



US006911471B2

(12) **United States Patent**
Zingaro et al.

(10) **Patent No.:** US 6,911,471 B2
(45) **Date of Patent:** Jun. 28, 2005

(54) **S-DIMETHYLARSINO-THIOSUCCINIC ACID S-DIMETHYLARSINO-2-THIOBENZOIC ACID S-(DIMETHYLARSINO) GLUTATHIONE AS TREATMENTS FOR CANCER**

(75) **Inventors:** Ralph A. Zingaro, College Station, TX (US); Emil J. Freireich, Houston, TX (US); Hatice Duzkale, Houston, TX (US); Hagop Kantarjian, Bellaire, TX (US); Srdan Verstovsek, Houston, TX (US); Merida Sotelo-Lerma, Sonora (MX)

(73) **Assignees:** The Texas A&M University System, College Station, TX (US); Board of Regents, The University of Texas System, Austin, TX (US)

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** 10/337,969

(22) **Filed:** Jan. 7, 2003

(65) **Prior Publication Data**

US 2004/0034095 A1 Feb. 19, 2004

Related U.S. Application Data

(60) Provisional application No. 60/346,492, filed on Jan. 7, 2002.

(51) **Int. Cl.⁷** A61K 31/285; A61K 33/36; C07F 9/70

(52) **U.S. Cl.** 514/504; 556/71; 556/77; 556/78; 556/80; 424/629

(58) **Field of Search** 424/629; 514/504; 556/71, 77, 78, 80

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,191,123 B1 2/2001 Uckun et al. 514/150
2004/0028750 A1 2/2004 Lu 424/629

FOREIGN PATENT DOCUMENTS

EP 1002537 10/1998
WO WO 99/24029 11/1998

OTHER PUBLICATIONS

American Conference of Governmental Industrial Hygienists, Inc. (ACGIH). Arsenic and soluble compounds, including arsine. Documentation of the Threshold Limit Values and Biological Exposure Indices, sixth edition, 1991.
Bachleitner-Hofmann et al., "Arsenic trioxide and ascorbic acid: synergy with potential implications for the treatment of acute myeloid leukaemia," *Br. J. Haematol.*, 112(3):783-786, 2001.
Banks et al., "Biomolecules bearing the S- or SeAsMe2 function: amino acid and steroid derivatives," *J. Medicinal Chem.* 22:572-575, 1979.

Beliles, "The Metals," In *Patty's Industrial Hygiene and Toxicology, fourth edition* G. D. Clayton and F.E. Clayton, eds. John Wiley & Sons, Inc.: New York. pp. 1913-1925, 1994.

Calleja and Warrell, "Differentiating agents in pediatric malignancies: all-trans-retinoic acid and arsenic in acute promyelocytic leukemia," *Curr. Oncol. Rep.*, 2:519-523, 2000.

Chen et al., "6-thio- and -seleno-alpha-D-glucose esters of dimethylarsinous acid," *Carb. Res.* 50:53-62, 1976.

Chen et al., "Synthesis of 1- and 6-S- and 1- and 6-Se-derivatives of 2-amino-2-deoxy-alpha/beta-D-glucopyranose," *J. Chemical Soc, Perkin Trans.* 1: 2287-2293, 1980.

Cuzick et al., "Medicinal arsenic and internal malignancies," *Br. J. Cancer*, 45:904-911, 1982.

Daniel and Zingaro "Dimethylarsinous acid esters of 1-thio- and -seleno- galactose. A new class of potential carcinostatic agents," *Phosphorus and Sulfur* 4:179-185, 1978.

Forkner and McNair-Scott, "Arsenic as a therapeutic agent in chronic myeloid leukemia," *JAMA* 97(1):3-6, 1931.

Geissler et al., "In vivo effects of arsenic trioxide in refractory acute myeloid leukemia other than acute promyelocytic leukemia," *Blood* 94:4230a, 1999.

Goyer, "Toxic effects of metals" In *Casarett and Doull's Toxicology: The Basic Science of Poisons, 5th edition*. C.D. Klassen, ed. McGraw-Hill: New York. pp. 691-698, 1996.

Grignani et al., "The acute promyelocytic leukemia-specific PML-RAR alpha fusion protein inhibits differentiation and promotes survival of myeloid precursor cells," *Cell*, 74:423-431, 1993.

Hughes and Kenyon, "Dose-dependent effects on the disposition of monomethylarsonic acid and dimethylarsinic acid in the mouse after intravenous administration," *J. Toxicol. Environ. Health*, 53(2):95-112, 1998.

IARC. Some metals and metallic compounds. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. vol. 23:39-141, 1980.

(Continued)

Primary Examiner—Porfirio Nazario-Gonzalez
(74) *Attorney, Agent, or Firm*—Ropes & Gray LLP

(57) **ABSTRACT**

Arsenic trioxide, an inorganic compound, is commercially available anti-cancer agent but it carries significant toxicity. Organic arsenicals, on the other hand, are much less toxic, to the extent that the methylation of inorganic arsenic in vivo into organic arsenicals has been considered a detoxification reaction. New organic arsenic derivatives have been synthesized, including S-dimethylarsino-glutathione, S-dimethylarsino-thiosuccinic acid and S-dimethylarsino-thiobenzoic acid, and established its potent in vitro cytotoxic activity against numerous human tumor cell lines, both of solid and hematological origin, as well as against malignant blood cells from patients with leukemia. Results form a basis for the development of S-dimethylarsino-glutathione, S-dimethylarsino-thiosuccinic acid, S-dimethylarsino-thiobenzoic acid, and other organic arsenicals as an anti-cancer therapy, combining high efficacy with very low, if any, toxicity.

40 Claims, 32 Drawing Sheets-

OTHER PUBLICATIONS

- Kitamura et al., "New retinoids and arsenic compounds for the treatment of refractory acute promyelocytic leukemia: clinical and basic studies for the next generation," *Cancer Chemother Pharmacol.*, 40 (Suppl): S36–S41, 1997.
- Knock et al., "The use of selected sulfhydryl inhibitors in a preferential drug attack on cancer," *Surg. Gynecol. Obstet.* 133:458–466, 1971.
- König, A. et al., "Comparative activity of melarsoprol and arsenic trioxide in chronic B-cell leukemia lines," *Blood* 90:562–570, 1997.
- Lallemand-Breitenbach et al., "Retinoic acid and arsenic synergize to eradicate leukemic cells in a mouse model of acute promyelocytic leukemia," *J. Exp. Med.*, 189:1043–1052, 1999.
- Mountain et al., "Chemotherapy studies in an animal tumor spectrum: II. Sensitivity of tumors to fourteen antitumor chemicals," *Cancer Res.* 26:181–206, 1966.
- Rivi et al., "Organic arsenical melarsoprol shows growth suppressive activity via programmed cell death on myeloid and lymphoid leukemia derived cell lines," *Blood* (Suppl.) 88:68a, 1996.
- Rosenthal and Zingaro, "The synthesis and characterization of thio sugar esters of diorganylarsinous acids," *Phosphorus and Sulfur* 9:107–116, 1980.
- Rousselot et al., "Use of arsenic trioxide (As_2O_3) in the treatment of chronic myelogenous leukemia: In vitro and in vivo studies," *Blood* 94:4457a, 1999.
- Soignet et al., "Clinical study of an organic arsenic melarsoprol, in patients with advanced leukemia," *Cancer Chemother. Pharmacol.* 44:471–421, 1999.
- Soignet et al., "Dose-ranging and clinical pharmacologic study of arsenic trioxide in patients with advanced hematologic cancers," *Blood* 94: 1247a, 1999.
- Tallman, "Therapy of acute promyelocytic leukemia: all-tans retinoic acid and beyond," *Leukemia*, 12 (Suppl 1):S37–S40, 1998.
- Wiernik et al., "Phase II trial of arsenic trioxide (As_2O_3) in patients with relapsed/refractory acute myeloid leukemia, blast crisis of CML or myelodysplasia," *Blood* 94:2283a, 1999.
- Zhang et al., "Arsenic trioxide treated 72 cases of acute promyelocytic leukemia," *Chin. J. Hematol.* 17:58–62, 1996.
- King and Ludford, "Relation between the constitution of arsenicals and their action on cell division," *Journal of the Chemical Society Abstracts*, 2086–2088, 1950.
- Lam et al., "Spectroscopic studies of arsenic (III) binding to *Escherichia coli* RI methyltransferase and to two mutants, C223S and W183F," *Biochemistry*, 31(43):10438–10442, 1992.
- Lin et al., "Methylarsenicals and arsinothiols are potent inhibitors of mouse liver thioredoxin reductase," *Chemical Research in Toxicology*, 12(10):924–930, 1999.
- Mester et al., "Specification of dimethylarsinic acid and monomethylarsonic acid by gas chromatography-mass spectrometry," *Journal of Chromatography*, 832(1+2):183–190, 1999.
- Schoene et al., "Speciation of arsenic-containing chemical warfare agents by gas chromatographic analysis after derivatization with thioglycolic acid methyl ester," *Journal of Chromatography*, 605(2):257–262, 1992.
- Scott et al., "Reactions of arsenic (III) and arsenic (V) species with glutathione," *Chemical Research in Toxicology*, 6(1):102–106, 1993.
- Styblo et al., "Comparative inhibition of yeast glutathione reductase by arsenicals and arsenothiols," *Chemical Research in Toxicology*, 10(1):27–33, 1997.
- Tslev et al., "Flow-injection hydride generation atomic absorption spectrometric study of the automated on-line pre-reduction of arsenate, methylarsonate and dimethylarsinate and high-performance liquid chromatographic separation of their L-cysteine complexes," *Talanta*, 51(6):1059–1068, 2000.
- Tsao et al., "Optically detected magnetic resonance study of the interaction of an arsenic (III) derivative of cacodylic acid with EcoRI methyltransferase," *Biochemistry*, 30(18):4565–4572, 1991.
- Vega et al., "Differential effects of trivalent and pentavalent arsenicals on cell proliferation and cytokine secretion in normal human epidermal keratinocytes," *Toxicology and Applied Pharmacology*, 172(3):225–232, 2001.
- Aslanidis et al., "Methylarsino-substituted hydroxy carboxylate esters," *Chemiker-Zeitung*, 112(4):125–127, 1988.
- Beckermann and Bernhard, "Determination of monomethylarsonic acid and dimethylarsinic acid by derivatization with thioglycolic acid methyl ester and gas-liquid chromatographic separation," *Analytica Chimica Acta*, 135(1):77–84, 1982.
- Cullen et al., "The metabolism of methylarsine oxide and sulfide," *Applied Organometallic Chemistry*, 3(1):71–78, 1989.
- Cullen et al., "The reaction of methylarsinicals with thiols: some biological implications," *J. Inorganic Biochemistry*, 21(3):179–194, 1984.
- Emran et al., "Synthesis and biodistribution of radioarsenic labeled dimethylarsinothiols: derivatives of penicillamine and mercaptoethanol," *International Journal of Nuclear Medicine and Biology*, 11(3–4):259–261, 1984.
- Hosain et al., "Synthesis of radioarsenic labeled dimethylchloroarsine for derivation of a new group of radiopharmaceuticals," *International Journal of Applied Radiation and Isotopes*, 33(12):1477–1478, 1982.
- Ionov et al., "Reaction of tertiary arsine sulfides with alkyl chlorocarbonates," *Zhurnal Obshchei Khimii*, 46(11):2555–2558, 1976.
- Kala et al., "The MRP2/cMOAT transporter and arsenic-glutathione complex formation are required for biliary excretion of arsenic," *J. Biol. Chem.*, 275(43):33404–33408, 2000.

FIG. 1
3 day MTT assay - NB4 cells





FIG. 2
3 day MTT assay - AML2 cells






FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F




FIG. 3G




FIG. 3H





FIG. 3I

FIG. 4
SAL1: trypan blue assay - HL60 cells






FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F




FIG. 5G




FIG. 5H




FIG. 5I

FIG. 6
3 day MTT assay - NB4 cells





FIG. 7
5 day clonogenic assay - HL60 cells






FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 8E




FIG. 8F




FIG. 8G




FIG. 8H




FIG. 8I

FIG. 9
3 day trypan blue assay - AML patient




FIG. 10 - 4 day trypan blue assay -
AML patient




FIG. 11 - MER1: 5 day trypan blue assay
- AML patient




FIG. 12
3 day trypan blue assay - CML-BP patient





FIG. 13 - MER1: 4 day trypan blue assay
- ALL patient






FIG. 15 - 8 day clonogenic assay - normal donor





FIG. 16
5 day trypan blue assay - CLL patient






FIG. 17




FIG. 19
MER1: trypan blue assay - HL60 cells




FIG. 20

MER1 - Annexin V assay - HL60 cells






FIG. 21

MER1 - propidium iodide assay - HL60 cells








FIG. 23A




FIG. 23B




FIG. 23C




FIG. 23D




FIG. 23E




FIG. 23F




FIG. 24




FIG. 25A




FIG. 25B




FIG. 25C




FIG. 26A




FIG. 26B




FIG. 26C

1

**S-DIMETHYLARSINO-THIOSUCCINIC ACID
S-DIMETHYLARSINO-2-THIOBENZOIC
ACID S-(DIMETHYLARSINO)
GLUTATHIONE AS TREATMENTS FOR
CANCER**

BACKGROUND OF THE INVENTION

The present application claims priority to co-pending provisional U.S. Application Ser. No. 60/346,492 filed Jan. 7, 2002. The entire text of the above referenced application is incorporated herein by reference and without disclaimer.

I. Field of the Invention

The present invention relates generally to the field of anti-cancer therapy. More particularly, it provides organic arsenic compounds and methods for their use in treating cancers such as leukemia.

II. Description of Related Art

Despite progress in leukemia therapy, most adult patients with leukemia still die from disease progression and an estimated 31,500 new cases and 21,500 deaths are expected in the year 2001. Arsenic trioxide, an inorganic compound, has recently been approved for the treatment of patients with relapsed or refractory acute promyelocytic leukemia (APL) and is being evaluated as therapy for other leukemia types. However, its use is limited by its toxicity.

Arsenic was used as a medicinal agent more than 2400 years ago in Greece and Rome, and arsenic still comprises the active ingredient in certain folk remedies, particularly in Central and Southern Asia (Bainbridge et al., 1914). The history and folklore of arsenic prompted intensive studies by many early pharmacologists. The foundations of many modern concepts of chemotherapy derive from Ehrlich's early work with arsenicals (e.g. the "silver bullet" for syphilis), and such drugs were once a mainstay of cancer chemotherapy. For example, in the early 1930's, Fowler's solution (inorganic arsenic dissolved in water) was used for controlling elevated leukocyte counts in chronic myelogenous leukemia (Forkner et al., 1931). In fact, clinical improvement of the leukemia, including control of fever, reduction of white cell count and splenomegaly, and improvement in anemia could be observed. Various arsenicals were briefly examined in early preclinical cancer screening studies at the U.S. National Cancer Institute (Tamowski et al., 1966). However, the clinical use of arsenicals in the treatment of cancer in the U.S. virtually ceased in the early 1970's with the advent of now traditional cytotoxic drugs and radiotherapy along with reports of arsenic poisoning from chronic low-dose ingestion (Knock et al., 1971; Cuzick et al., 1987). Still, there is large body of knowledge regarding effects of arsenicals on human subjects from both medicinal and toxicological standpoints. In current therapeutics, arsenicals are important only for the treatment of certain tropical diseases, such as the use of melarsoprol, an organic compound, against African trypanosomiasis (Investigational Drug Brochure, 1987). However, Chinese reports of the striking activity of arsenic trioxide (As₂O₃) (Zhang et al., 1996), an inorganic compound, against APL have engendered considerable interest. This resulted in the recent approval of arsenic trioxide for treatment of patients with relapsed or refractory APL. Preliminary data from China and the recent experience in the U.S., however, suggest a role for arsenic trioxide in the other hematologic cancers as well. Consequently, the activity of arsenic trioxide as an anti-leukemic agent is currently being investigated in many types of leukemia. Although the results look favorable in terms of the response rate of some of the


2

leukemia types that are being investigated, systemic toxicity of arsenic trioxide is a problem (Soignet et al., 1999; Wiemiket et al., 1999; Geissler et al., 1999; Rousselot et al., 1999).

The only organic arsenical (OA) still manufactured for human use, melarsoprol, has been evaluated for its antileukemic (WO9924029, EP1002537) properties and showed significant activity. However, this compound is excessively toxic to patients with leukemia at drug concentration and schedule used previously for treatment of trypanosomiasis. Therefore, there is a need to identify arsenic derivatives that can be used for the treatment of hematologic malignancies and cancer in general, that have similar or greater activity and lower toxicity than arsenic trioxide. Organic arsenic derivatives may fulfill this promise as they should be less toxic than the inorganic arsenic trioxide.


SUMMARY OF THE INVENTION

The present invention overcomes these and other defects in the art and provides organic arsenical compounds with anti-cancer properties. In some embodiments, the present invention comprises compounds having anti-cancer activity comprising the structure:



wherein R₁ and R₂ are independently alkyls with 1–10 carbon atoms; X is S or Se; R₃ is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH; n is 0 or 1; R₄ is —OH, —H, —CH₃, or a glutamine substituent; R₃, R₄, C₁ and C₂ all independently comprise part of an aromatic ring or substituted aromatic ring; and R₅ is a —OH, or glycine substituent; or a pharmaceutically acceptable salt or formulation thereof.

In particular embodiments, the compound can have the formula:




or is a pharmaceutically acceptable salt or formulation thereof.

Thus, the invention also comprises compositions and pharmaceutical compositions comprising the compounds described above.


3

In other embodiments, the invention comprises of a pharmaceutical composition having anti-cancer activity comprising a pharmaceutical carrier and an organic arsenical compound. In some embodiments, such a composition has the formula:



wherein R_1 and R_2 are independently alkyls with 1–10 carbon atoms; X is S or Se; R_3 is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH; n is 0 or 1; R_4 is —OH, —H, —CH₃, or a glutamine substituent; R_3 , R_4 , C_1 and C_2 all independently comprise part of an aromatic ring or substituted aromatic ring; and R_5 is a —OH, or glycine substituent; or a pharmaceutically acceptable salt thereof.

In yet other embodiments, the invention comprises a method of treating a patient with cancer comprising administering a composition comprising a therapeutically effective amount of a compound having the formula:




wherein R_1 and R_2 are independently alkyls with 1–10 carbon atoms; X is S or Se; R_3 is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH; n is 0 or 1; R_4 is —OH, —H, —CH₃, or a glutam substituent; R_3 , R_4 , C_1 and C_2 all independently comprise part of an aromatic ring or substituted aromatic ring; and R_5 is a —OH, or a glycine substituent; or a pharmaceutically acceptable salt or formulation thereof to the patient. The therapeutically effective amount of a compound could be 0.1–1000 mg/kg or 1–500 mg/kg, or 10–100 mg/kg.

In particular embodiments, the method may comprise administering the composition daily. It is further contemplated that treatment methods may involve multiple administrations. On other embodiments, the method further comprises administering one or more additional agents to the patient. The additional agent may be all-trans-retinoic acid, 9-cis retinoic acid, Am-80 or ascorbic acid. The use of other adjunct cancer therapies, such as chemotherapy, radiotherapy, gene therapy, hormone therapy and other cancer therapies known in the art are also contemplated in conjunction with the methods of the present invention.

Various methods of administration are contemplated, including regional, systemic, direct administration and by perfusion. Such methods include administration by injection, oral routes, intravenous, intraarterial, intratumoral, administration to tumoral vasculature, intraperitoneal, intratracheal, intramuscular, endoscopic, intralesional, percutaneous, subcutaneous, topical, nasal, buccal, mucosal, anogenital, rectal and the like.

In particular embodiments, the method of treating a patient with a cancer comprises administering a therapeutically effective amount of a compound having the formula:

4



or a pharmaceutically acceptable salt or formulation thereof.

The methods of the invention may be used to treat any cancer, including but not limited, to a solid tumor, such as brain, lung, liver, spleen, kidney, lymph node, small intestine, pancreas, blood cells, bone, colon, stomach, breast, endometrium, prostate, testicle, ovary, central nervous system, skin, head and neck, esophagus, or bone marrow cancer. Furthermore, the cancer could be a hematological cancer, such as leukemia, acute promyelocytic leukemia, lymphoma, multiple myeloma, myelodysplasia, myeloproliferative disease, or refractory anemia.

The method can comprise of administering the compound daily such as by injection. Alternative routes and methods of administration described in the specification may also be used and the mode of administration will mainly depend on the type and location of the cancer. Furthermore, the method can comprise administering one or more additional agents to the patient. The additional agent may be all-trans-retinoic acid, 9-cis retinoic acid, Am-80 or ascorbic acid. However, the use of other agents that are normally used in the therapy of cancer are also contemplated. This includes the use of chemotherapeutic agents, radiation, surgery, gene-therapy, cytokines, hormonal therapy and a vast variety of other anti-cancer therapies known in the art.

As used herein the specification, “a” or “an” may mean one or more. As used herein in the claim(s), when used in conjunction with the word “comprising”, the words “a” or “an” may mean one or more than one. As used herein “another” may mean at least a second or more.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1. The human leukemia cell line NB4 was incubated for 3 days with indicated concentrations of

5

S-dimethylarsino-thiosuccinic acid (MER1) or arsenic trioxide. Cell survival was assessed by the MTT assay.

FIG. 2. The human leukemia cell line AML2 was incubated for 3 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by the trypan-blue exclusion method.

FIGS. 3A, 3B, 3C, 3D, 3E, 3F, 3G, 3H, & 3I. Percent growth for 60 human cell lines when treated with MER1. Various human cancer cells were incubated on a microtiter plate with indicated concentrations of MER1 for 48 hours. End point determinations were made with sulforhodamine B, a protein binding dye. Results are reported as the percentage of growth of treated cells when compared to untreated control cells. Negative data indicate cell kill. FIG. 3A. Leukemia cell lines, FIG. 3B. CNS cell lines, FIG. 3C. Renal cancer cell lines, FIG. 3D. Non-small cell lung cancer cell lines, FIG. 3E. Melanoma cell lines, FIG. 3F. Prostate cancer cell lines, FIG. 3G. Colon cancer cell lines, FIG. 3H. Ovarian cancer cell lines, FIG. 3I. Breast cancer cell lines.

FIG. 4. HL60 human leukemia cells were incubated for 3 days with indicated concentrations of S-dimethylarsino-2-thiobenzoic acid (SAL1). Cell survival was assessed by trypan-blue exclusion method.

FIGS. 5A, 5B, 5C, 5D, 5E, 5F, 5G, 5H, & 5I. Percentage growth for 60 human cell lines when treated with SAL1. Various human cancer cells were incubated on a microtiter plate with indicated concentrations of SAL1 for 48 hours. End point determinations were made with sulforhodamine B, a protein binding dye. Results are reported as the percentage of growth of treated cells when compared to untreated control cells. Negative data indicate cell kill. FIG. 5A. Leukemia cell lines, FIG. 5B. CNS cell lines, FIG. 5C. Renal cancer cell lines, FIG. 5D. Non-small cell lung cancer cell lines, FIG. 5E. Melanoma cell lines, FIG. 5F. Prostate cancer cell lines, FIG. 5G. Colon cancer cell lines, FIG. 5H. Ovarian cancer cell lines, FIG. 5I. Breast cancer cell lines.

FIG. 6. NB4 cells were incubated for 3 days with indicated concentrations of S-dimethylarsino-glutathione (SGLU1) or arsenic trioxide. Cell survival was assessed by the MTT assay.

FIG. 7. A five day clonogenic assay was performed using HL60 cells and SGLU1 or arsenic trioxide. Cell aggregates composed of more than 50 cells were counted as one colony and growth inhibition was evaluated as a percentage of colony growth as compared to colony growth in control (no drug) samples.

FIGS. 8A, 8B, 8C, 8D, 8E, 8F, 8G, 8H, & 8I. Percent growth for 60 human cell lines when treated with SGLU1. Various human cancer cells were incubated on a microtiter plate with indicated concentrations of SGLU1 for 48 hours. End point determinations were made with sulforhodamine B, a protein binding dye. Results are reported as the percentage of growth of treated cells when compared to untreated control cells. Negative data indicate cell kill. FIG. 8A. Leukemia cell lines, FIG. 8B. CNS cell lines, FIG. 8C. Renal cancer cell lines, FIG. 8D. Non-small cell lung cancer cell lines, FIG. 8E. Melanoma cell lines, FIG. 8F. Prostate cancer cell lines, FIG. 8G. Colon cancer cell lines, FIG. 8H. Ovarian cancer cell lines, FIG. 8I. Breast cancer cell lines.

FIG. 9. Mononuclear cells from acute myeloid leukemia (AML) patient were incubated for 3 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

FIG. 10. Mononuclear cells from AML patient were incubated for 4 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

6

FIG. 11. Mononuclear cells from AML patient were incubated for 5 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

FIG. 12. Mononuclear cells from chronic myeloid leukemia—blastic phase (CML-BP) patient were incubated for 3 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

FIG. 13. Mononuclear cells from acute lymphoblastic leukemia (ALL) patient were incubated for 4 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

FIG. 14. Mononuclear cells from normal donor were incubated for 5 days with indicated concentrations of MER1 or arsenic trioxide. Cell survival was assessed by trypan-blue exclusion method.

FIG. 15. An 8 day clonogenic assay was performed using normal donor cells and MER1 or arsenic trioxide. Cell aggregates composed of more than 50 cells were counted as one colony and growth inhibition was evaluated as a percentage of colony growth as compared to colony growth in control (no drug) samples.

FIG. 16. Mononuclear cells from chronic lymphocytic leukemia (CLL) patient were incubated for 5 days with indicated concentrations of SGLU1 or arsenic trioxide. Cell survival was assessed by trypan blue assay.

FIG. 17. An 8 day clonogenic assay was performed using mononuclear cells from AML patient with SGLU1 or arsenic trioxide. Cell aggregates composed of more than 50 cells were counted as one colony and growth inhibition was evaluated as a percentage of colony growth as compared to colony growth in control (no drug) samples.

FIG. 18. An 8 day clonogenic assay was performed using normal donor cells, and SGLU1 or arsenic trioxide. Cell aggregates composed of more than 50 cells were counted as one colony and growth inhibition was evaluated as a percentage of colony growth as compared to colony growth in control (no drug) samples.

FIG. 19. Stability of MER-1 formulation. Trypan blue assays were performed using HL60 cells and MER-1 1–7 weeks since the preparation of MER-1. HL60 human leukemia cells were incubated for 3 days with indicated concentrations of MER1. Cell survival was assessed by trypan-blue exclusion method. Time of test (i.e., weeks from the time MER1 was prepared) is indicated.

FIG. 20. Apoptosis was assessed by annexin V assay in HL-60 cells treated with MER1 for 1, 2, or 3 days.

FIG. 21. Dead cell number was assessed by the propidium iodide assay on HL-60 cells treated with MER1 for 1, 2, or 3 days.

FIG. 22. Cell survival was assessed by trypan-blue exclusion method in HL60 cells treated with MER1 for 1, 2, or 3 days.

FIGS. 23A, 23B, 23C, 23D, 23E, & 23F. Apoptosis was assessed in HL-60 cells treated with MER1, SGLU1 or arsenic trioxide, as indicated, by the annexin V assay and propidium iodide staining at 48 hours (FIG. 23A) and at 72 hours (FIG. 23D); by the caspase assay (phi-phi-lux staining) at 48 hours (FIG. 23B) and 72 hours (FIG. 23E); and by the CMXRos/MT-Green Assays at 48 hours (FIG. 23C) and 72 hours (FIG. 23F).

FIG. 24. NB4 cells assayed for the effects of arsenic trioxide, SGLU1 and MER1 on maturation by assaying the expression of CD11b marker on the surface of leukemic

cells by flow cytometer (after 3 day incubation). Data shows that, in contrast to arsenic trioxide, SGLU1 and MER1 do not induce maturation.

FIGS. 25A, 25B & 25C. HL60 cells assayed for the effects of MER1 (FIG. 25A), SGLU1 (FIG. 25B) & arsenic trioxide (FIG. 25C) on the cell cycle after 3 day incubation with indicated concentrations.

FIGS. 26A, 26B & 26C. Three day MTT assay in U937/9PR cells treated with arsenic trioxide (FIG. 26A), SGLU1 (FIG. 26B) and MER1 (FIG. 26C) with and without zinc to analyze the role of the PML/RARalpha gene. Zinc activates the the PML/RARalpha gene. The data shows that the presence of functional PML/RARalpha gene is pre-requisite for cells to be sensitive to arsenic trioxide but has no influence on the sensitivity of the cells to SGLU1 and MER1.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

I. The Present Invention

The present invention overcomes deficiencies in the art by providing a number of organic arsenic compounds for the treatment of cancer having similar or greater activity and lower toxicity than current treatment with arsenic trioxide. More particularly, the present invention provides S-dialkylarsino-thio carboxylic acids, including S-dimethylarsino-thiosuccinic acid and S-dimethylarsino-2-

II. Organic Arsenicals

Twenty years ago a large number of organic arsenicals (OA) derivatives were synthesized by Prof. Dr. Ralph A. Zingaro at Texas A&M University, a co-inventor in this application, and their physicochemical properties determined (Chen et al., 1976; Rosenthal et al., 1980; Chen et al., 1980; Daniel et al., 1978; Banks et al., 1979, the entire contents of all these references are incorporated herein by reference in their entirety). After it was shown that one of the compounds displayed in vitro activity against cancer cells, many of the newly synthesized OA were submitted to the National Institute of Health (NIH) for the evaluation of their anticancer activity. The compounds were tested in vivo in mice bearing P388 lymphocytic leukemia cells. The activity of these compounds was assessed by recording the survival time of groups of six mice treated with different drugs (intraperitoneally, daily for 5 days), as well as a control group, as shown in Table 1. The NIH criterion for significant activity is the percentage treatment/control (T/C)>125. This means that the group of animals receiving the drug survives at least 25% longer than the control group. A number of compounds displayed significant antileukemic activity, and some even reached % T/C of 180. Furthermore, the compounds were non-toxic as some were used at the dose of 200 mg/kg (LD₅₀ for arsenic trioxide is 10 mg/kg).

TABLE 1


In vivo activity of OA derivatives in mice bearing P388 lymphocytic leukemia cells, as reported by the NIH.		
Compound	Dose (mg/kg)	% T/C
1. 2,3,4,6-tetra-O-acetyl-1-S-dibutylarsino-1-thio-β-D-glucopyranose	100	117
2. 2,3,4,6-tetra-O-acetyl-1-S-dipropylarsino-1-thio-β-D-glucopyranose	200	124
3. 2,3,4,6-tetra-O-acetyl-1-S-diethylarsino-1-thio-β-D-glucopyranose	100	180
4. 2,3,4,6-tetra-O-acetyl-1-S-hydroxyethylmethyl-1-thio-β-D-glucopyranose	100	116
5. 2,3,4,6-tetra-O-acetyl-1-S-dihexadecylarsino-1-thio-β-D-glucopyranose	50	125
6. 2,3,4,6-tetra-O-acetyl-1-S-dicyclohexylarsino-1-thio-β-D-glucopyranose	400	Toxic
7. 1,2,3,4,-tetra-O-acetyl-6-S-methyl-n-propylarsino-6-thio-β-D-glucopyranose	200	118
8. 1,2,3,4,-tetra-O-acetyl-6-S-dihexadecylarsino-6-thio-β-D-glucopyranose	100	125
9. 1,2,3,4,-tetra-O-acetyl-6-S-dicyclohexylarsino-6-thio-β-D-glucopyranose	25	120
10. S-dimethylarsino-DL-cysteine	50	110
11. S-dimethylarsino-DL-penicillamine	25	117
12. Bis [S-dimethylarsino-homocysteine]	100	144
13. S-dimethylarsino-glutathione	100	164
14. Bis [S-dimethylarsino-α-dihydrolypoic acid]	50	128
15. Se-dimethylarsino-selenocholesterol	200	129
16. 2,3,4,6-tetra-O-acetyl-1-S-dimethylarsino-1-thio-β-D-galactopyranose	100	129
17. 1-S-dimethylarsino-1-thio-β-D-galactopyranose	50	125
18. 2-acetamido-3,4,6-tri-O-acetyl-2-deoxy-1-S-dimethylarsino-1-thio-β-D-glucopyranose	50	136
19. 2-acetamido-2-deoxy-1-S-dimethylarsino-1-thio-β-D-glucopyranose	100	125
20. 2-acetamido-1,3,4-tri-O-acetyl-2-deoxy-6-S-dimethylarsino-6-thio-α-D-glucopyranose	100	136
21. 2-acetamido-2-deoxy-6-S-dimethylarsino-6-thio-α-D-glucopyranose	100	125

thiobenzoic acid, and methods for use in treating cancers. The present invention also provides S-dimethylarsino-glutathione and methods for its use in treating cancers.


Melarsoprol (Arsobal) is the only organic arsenical still manufactured for human use, and is not commercially available in the United States. Melarsoprol is accepted for

use for trypanosomiasis, or African sleeping sickness, and has been evaluated for its antileukemic properties (WO9924029, EP1002537). Studies unexpectedly showed that melarsoprol had activity at least equivalent to that of arsenic trioxide against both APL and non-APL cell lines (Konig et al., 1997; Rivi et al., 1996). A limited clinical study of melarsoprol was then initiated in the U.S. in patients with advanced leukemia (Soignet et al, 1999). A total of 8 patients were treated on a 3 days per week schedule for 3 consecutive weeks (a dosing schedule that had previously been used for treatment of central nervous system trypanosomiasis). Only one patient (with chronic lymphocytic leukemia) displayed antitumor effects while most experienced neurological side effects. These results suggest that the dosing schedule developed for treatment of trypanosomiasis is excessively toxic in patients with leukemia and that further preclinical experiments, particularly in animal model of leukemia, is needed for melarsoprol.

Other organic arsenicals have been synthesized and include carboxylic acid and dicarboxylic acid arsenicals. These arsenicals have trivalent arsenic of the form:



wherein R_1 and R_2 are independently alkyls with 1–10 carbon atoms and are preferably methyl, ethyl or hydroxyethyl and most preferably methyl. X is S or Se, and is preferably S. Q is an organic grouping, usually of biochemical origin such as a sugar, peptide, amino acid, or steroid. However, Q can also be a non-biochemical moiety, such as a carboxylic acid moiety. The organic arsenicals of the present invention have at least one carboxylic acid group separated by one or two carbon atoms from X. These compounds can be described by:



wherein R_1 & R_2 are the same as in the previous structure; R_3 is $-COOH$, $-CH_2-COOH$, $-CH_2-CH_2-COOH$, $-CH(CH_3)-COOH$, $-CH(CH_2-CH_3)-COOH$, $-CH_2-CH_2-CH_2-COOH$, or R_3 forms an aromatic ring or substituted aromatic ring comprising R_4 , C, and C_2 ; n is 0 or 1; and R_4 is $-H-CH_3$, or is part of an aromatic ring. Part of an aromatic ring is defined herein as an atom that is bonded to two or more other atoms in an aromatic ring system.

Compounds of particular relevance to the present invention include S-dimethylarsino-thiosuccinic acid (MER1), S-dimethylarsino-2-thiobenzoic acid (SAL-1), and S-(dimethylarsino) glutathione (SGLU1). The present inventors have shown that MER1, SAL-1, and SGLU1 exert significant anti-cancer activity against a panel of human leukemia cell lines. This observation has been confirmed and extended to the activity against human solid tumor cell lines as well (>60 cell lines in total) by the experiments done at the National Institute of Health. In addition, MER1 and SGLU1 showed significant activity against blood tumor cells from patients with leukemia. When compared to the activity of the arsenic trioxide, MER1, SAL1, and SGLU1 showed similar efficacy. MER1 and SGLU1 also showed

low toxicity against nonmalignant blood mononuclear cells (obtained from normal donors). Furthermore, MER1 and SGLU1 showed less toxicity toward normal blood mononuclear cells than arsenic trioxide.

5 III. Toxicity of Inorganic vs. Organic Arsenicals

The use of arsenic trioxide is limited by its toxicity. OA, on the other hand, are much less toxic, to the extent that the methylation of inorganic arsenic in vivo into OA has been considered a detoxification reaction. The OA monomethylarsinic acid and dimethylarsinic acid are the primary metabolites of inorganic arsenic (Hughes et al., 1998). Inorganic arsenicals, including arsenic trioxide, have varied effects on many organ systems, including cardiovascular system, gastrointestinal tract, kidneys, skin, nervous system, and blood. Inorganic arsenicals are particularly toxic to the liver, causing infiltration, central necrosis, and cirrhosis (IARC, 1980; ACGIH, 1991; Beliles et al, 1994; Goyer et al., 1996). There is now sufficient evidence that inorganic arsenic compounds are skin and lung carcinogens in humans (Goyer et al, 1996).

The toxicity of a given arsenical is related to the rate of its clearance from the body and to the extent of its tissue accumulation (Beliles et al., 1994). In general, toxicity increases in the following sequence: organic arsenicals $<As^{5+} < As^{3+}$ (including arsenic trioxide) $<$ arsine. Unlike inorganic arsenicals, no deaths or serious cases of toxicity due to OA have been reported in the literature. Consequently, in mammals the methylation of inorganic arsenic has been considered a detoxification reaction because of the lower toxicity of methylated OA, and their fast excretion and low retention (Beliles et al., 1994; Goyer et al., 1996). A good example is that of dimethylarsinic acid, an organic compound, the predominant urinary metabolite excreted by most mammals after exposure to inorganic arsenic, including arsenic trioxide. In in vivo toxicity studies in mice, after intraperitoneal administration of arsenic trioxide, the LD_{50} (a dose at which 50% of animals die due to acute toxicity) was 10 mg/kg, (Investigator's Brochure, 1998), while after administration of dimethylarsinic acid, the LD_{50} was 500 mg/kg (MSDS, 1998).

40 IV. Cancer Treatment

The organic arsenicals of the current invention may be used to treat a variety of cancers, including all solid tumors and all hematological cancers, including leukemia, lymphoma, multiple myeloma, myelodysplasia, or myeloproliferative disorders. The OA can also be used to treat hematological cancers that have become refractory to other forms of treatment.

Leukemia is a malignant neoplasm of blood-forming tissues, characterized by abnormal proliferation of leukocytes and is one of the four major types of cancer. Leukemias are classified according to the type of leucocyte most prominently involved. Acute leukemias are predominantly undifferentiated cell populations and chronic leukemias have more mature cell forms (WO9924029).

The acute leukemias are divided into lymphoblastic (ALL) and non-lymphoblastic (ANLL) types and may be further subdivided by morphologic and cytochemical appearance according to the French-American-British classification or according to their type and degree of differentiation. Specific B- and T-cell, as well as myeloid cell surface markers/antigens are used in the classification too. ALL is predominantly a childhood disease while ANLL, also known as acute myeloid leukemia, is a more common acute leukemia among adults.

Chronic leukemias are divided into lymphocytic (CLL) and myeloid (CML) types. CLL is characterized by the

increased number of mature lymphocytes in blood, bone marrow, and lymphoid organs. Most CLL patients have clonal expansion of lymphocytes with B cell characteristics. CLL is a disease of older persons. In CML, the granulocytic cells predominate at all stages of differentiation in blood and bone marrow, but may also affect liver, spleen, and other organs. Other malignant hematological disease that may be treated with the OA of the current invention, include, but are not limited to: myelodysplasia, myeloproliferative diseases, lymphomas, and multiple myeloma.

V. Pharmaceutical Preparations

Pharmaceutical compositions of the present invention comprise an effective amount of one or more organic arsenic derivative, or more particularly *s*-dialkyl-thio acetic acids such as MER-1, SAL1 or SGLU1 or a salt or a derivative of MER1, SAL1 or SGLU1, or additional agent dissolved or dispersed in a pharmaceutically acceptable carrier. The phrases "pharmaceutical or pharmacologically acceptable" refers to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, such as, for example, a human, as appropriate. The preparation of a pharmaceutical composition that contains at least one organic arsenical or additional active ingredient will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference. Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329, incorporated herein by reference). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated.

The organic arsenical may comprise different types of carriers depending on whether it is to be administered in solid, liquid or aerosol form, and whether it need to be sterile for such routes of administration as injection. The present invention can be administered intravenously, intradermally, intraarterially, intraperitoneally, intralesionally, intracranially, intraarticularly, intraprostatically, intrapleurally, intratracheally, intranasally, intravitreally, intravaginally, intrarectally, topically, intratumorally, intramuscularly, intraperitoneally, subcutaneously, subconjunctival, intravesicularlly, mucosally, intrapericardially, intraumbilically, intraocularly, orally, topically, locally, injection, infusion, continuous infusion, localized perfusion bathing target cells directly, via a catheter, via a lavage, in lipid compositions (e.g., liposomes), or by other method or any combination of the forgoing as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference).

The actual dosage amount of a composition of the present invention administered to a patient can be determined by

physical and physiological factors such as body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the patient and on the route of administration. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject.

In certain embodiments, pharmaceutical compositions may comprise, for example, at least about 0.1% of an organic arsenical compound. In other embodiments, the an active compound may comprise between about 2% to about 75% of the weight of the unit, or between about 25% to about 60%, for example, and any range derivable therein. In other non-limiting examples, a dose may also comprise from about 0.1 mg/kg/body weight, 0.5 mg/kg/body weight, 1 mg/kg/body weight, about 5 mg/kg/body weight, about 10 mg/kg/body weight, about 20 mg/kg/body weight, about 30 mg/kg/body weight, about 40 mg/kg/body weight, about 50 mg/kg/body weight, about 75 mg/kg/body weight, about 100 mg/kg/body weight, about 200 mg/kg/body weight, about 350 mg/kg/body weight, about 500 mg/kg/body weight, about 750 mg/kg/body weight, to about 1000 mg/kg/body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 10 mg/kg/body weight to about 100 mg/kg/body weight, etc., can be administered, based on the numbers described above.

In any case, the composition may comprise various antioxidants to retard oxidation of one or more component. Additionally, the prevention of the action of microorganisms can be brought about by preservatives such as various antibacterial and antifungal agents, including, but not limited to parabens (e.g., methylparabens, propylparabens), chlorobutanol, phenol, sorbic acid, thimerosal or combinations thereof.

The organic arsenical may be formulated into a composition in a free base, neutral or salt form. Pharmaceutically acceptable salts include the salts formed with the free carboxyl groups derived from inorganic bases such as for example, sodium, potassium, ammonium, calcium or ferric hydroxides; or such organic bases as isopropylamine, trimethylamine, histidine or procaine.

In embodiments where the composition is in a liquid form, a carrier can be a solvent or dispersion medium comprising, but not limited to, water, ethanol, polyol (e.g., glycerol, propylene glycol, liquid polyethylene glycol, etc.), lipids (e.g., triglycerides, vegetable oils, liposomes) and combinations thereof. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin; by the maintenance of the required particle size by dispersion in carriers such as, for example liquid polyol or lipids; by the use of surfactants such as, for example hydroxypropylcellulose; or combinations thereof such methods. In many cases, it will be preferable to include isotonic agents, such as, for example, sugars, sodium chloride or combinations thereof.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount of the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and/or the other ingredients. In the case of sterile powders for the preparation of sterile injectable solutions, suspensions or emulsion, the preferred methods of preparation are vacuum-drying or

freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered liquid medium thereof. The liquid medium should be suitably buffered if necessary and the liquid diluent first rendered isotonic prior to injection with sufficient saline or glucose. The preparation of highly concentrated compositions for direct injection is also contemplated, where the use of DMSO as solvent is envisioned to result in extremely rapid penetration, delivering high concentrations of the active agents to a small area.

The composition must be stable under the conditions of manufacture and storage, and preserved against the contaminating action of microorganisms, such as bacteria and fungi. It will be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein.

In particular embodiments, prolonged absorption of an injectable composition can be brought about by the use in the compositions of agents delaying absorption, such as, for example, aluminum monostearate, gelatin or combinations thereof.

VI. Combination Therapy

It is an aspect of this invention that the organic arsenical can be used in combination with another agent or therapy method, preferably another cancer treatment. The organic arsenical may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agent and expression construct are applied separately to the cell, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and expression construct would still be able to exert an advantageously combined effect on the cell. For example, in such instances, it is contemplated that one may contact the cell, tissue or organism with two, three, four or more modalities substantially simultaneously (i.e., within less than about a minute) with the organic arsenical. In other aspects, one or more agents may be administered within about 1 minute, about 5 minutes, about 10 minutes, about 20 minutes about 30 minutes, about 45 minutes, about 60 minutes, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 25 hours, about 26 hours, about 27 hours, about 28 hours, about 29 hours, about 30 hours, about 31 hours, about 32 hours, about 33 hours, about 34 hours, about 35 hours, about 36 hours, about 37 hours, about 38 hours, about 39 hours, about 40 hours, about 41 hours, about 42 hours, about 43 hours, about 44 hours, about 45 hours, about 46 hours, about 47 hours, to about 48 hours or more prior to and/or after administering the organic arsenical. In certain other embodiments, an agent may be administered within of from about 1 day, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 7 days, about 8 days, about 9 days, about 10 days, about 11 days, about 12 days, about 13 days, about 14 days, about 15 days, about 16 days, about 17 days, about 18 days, about 19 days, about 20, to about 21 days prior to and/or after administering the organic arsenical. In some situations, it may be desirable to extend the time period for treatment significantly, however, where several weeks (e.g., about 1, about 2, about 3, about 4, about 5, about 6, about 7 or about 8 weeks or more) lapse between the respective administrations.

Various combinations may be employed, the organic arsenical is "A" and the secondary agent, which can be any other therapeutic agent, is "B":

A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B/B B/A/B/B
 B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A
 B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

Administration of the therapeutic compositions of the present invention to a patient will follow general protocols for the administration of chemotherapeutics, taking into account the toxicity, if any. It is expected that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies or adjunct cancer therapies, as well as surgical intervention, may be applied in combination with the described arsenical agent. These therapies include but are not limited to chemotherapy, radiotherapy, immunotherapy, gene therapy and surgery. The section below describes some adjunct cancer therapies:

a. Chemotherapy

Cancer therapies also include a variety of combination therapies with both chemical and radiation based treatments. Combination chemotherapies include, for example, cisplatin (CDDP), carboplatin, procarbazine, mechlorethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatinum, 5-fluorouracil, vincristin, vinblastin and methotrexate, or any analog or derivative variant of the foregoing.

b. Radiotherapy

Other factors that cause DNA damage and have been used extensively include what are commonly known as γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also contemplated such as microwaves and UV-irradiation. It is most likely that all of these factors effect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells. The terms "contacted" and "exposed," when applied to a cell, are used herein to describe the process by which a therapeutic construct and a chemotherapeutic or radiotherapeutic agent are delivered to a target cell or are placed in direct juxtaposition with the target cell. To achieve cell killing or stasis, both agents are delivered to a cell in a combined amount effective to kill the cell or prevent it from dividing.

c. Immunotherapy

Immunotherapeutics, generally, rely on the use of immune effector cells and molecules to target and destroy cancer cells. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionucleotide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve merely as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells.

Immunotherapy, thus, could be used as part of a combined therapy, in conjunction with gene therapy. The general approach for combined therapy is discussed below. Generally, the tumor cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumor markers exist and any of these may be suitable for targeting in the context of the present invention. Common tumor markers include carcinoembryonic antigen, prostate specific antigen, urinary tumor associated antigen, fetal antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, estrogen receptor, laminin receptor, erb B and p155.

d. Gene Therapy

In yet another embodiment, the secondary treatment is a secondary gene therapy in which a therapeutic polynucleotide is administered before, after, or at the same time a first therapeutic agent. Delivery of the therapeutic agent in conjunction with a vector encoding a gene product will have a combined anti-hyperproliferative effect on target tissues.

e. Surgery

Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative and palliative surgery. Curative surgery is a cancer treatment that may be used in conjunction with other therapies, such as the treatment of the present invention, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy and/or alternative therapies. Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, andoscopically controlled surgery (Mohs' surgery). It is further contemplated that the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

VII. EXAMPLES

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Synthesis of S-dimethylarsino-thiosuccinic Acid (MER1), S-dimethylarsino-salicylic Acid (SAL1), and S-(dimethylarsino) glutathione (SGLU1)

MER-1: Mercaptosuccinic acid, 4.5 g, was placed in 100 ml of glyme (1,2-dimethoxyethane) in a 250 ml round-bottom flask. Four ml of dimethylchloroarsine (0.03 mol) was added drop-wise, followed by 4 ml of diethylamine (0.04 mol), again drop-wise. The reaction mixture was stirred for 20 h at room temperature. A white precipitate of diethylamine hydrochloride was formed and was separated by filtration. The solution of MER1 in the glyme was greatly reduced in volume by evaporation at reduced pressure. White crystals of MER1 were separated by filtration and washed with cold distilled water. The colorless crystalline product was then recrystallized from ethanol-water to a constant melting point of 150° C.

SAL-1: In a 100 ml flask 5 g of 2-mercapto benzoic acid (thiosalicylic acid), 75 ml of glyme, 5 ml of dimethylchloroarsine, and 5 ml diethylamine were placed. The mixture was refluxed for 1 hour under an atmosphere of nitrogen and stirred at room temperature overnight. The precipitate of diethylamine hydrochloride was separated by filtration. The filtrate was evaporated slowly under reduced pressure until crystals of the product separate. The evaporated solution containing the product was chilled in ice and the cold solution was filtered. Crystals of the product were recrystallized from ethanol to a constant melting point of 97° C.

SGLU-1: Glutathione (14.0 g, 45.6 mmol) was stirred rapidly in glyme while dimethylchloroarsine (6.5 g, 45.6 mmol) was added dropwise. Pyridine (6.9 g, 91.2 mmol) was then added to the slurry and the mixture was subsequently heated to reflux. The heat was removed immediately and the mixture stirred at room temperature for 4 h. Isolation of the resultant insoluble solid and recrystallization from ethanol afforded 4 as the pyridinium hydrochloride (75% yield): mp 115–118° C.; NMR (D20) δ 1.35 (s, 6H), 1.9–4.1 (m's, 10H), 7.8–9.0 (m, 5H); mass spectrum (m/e) 140, 125, 110, 105, 79, 52, 45, 36.

The studies that led to the synthesis of MER-1, SGLU-1 and SAL-1 were funded by the Robert A. Welch foundation of Houston, Tex., in a grant to inventor Ralph Zingaro.

Example 2

Assay for In Vitro Evaluation

A variety of in vitro assays were used to determine the response of cancer cells to the arsenical compounds, compositions, and/or formulations of the present invention. Some of the responses assayed included cell survival, cell cycle, apoptosis, and maturation. The present inventors also designed an assay to evaluate the requirement of the PML/RARalpha gene in cancer cells for sensitivity to the arsenical compositions of the invention. Provided below is a description of these assays:

Sulforhodamine B Assay. Various human cancer cells were incubated on a microtiter plate with or without indicated concentrations of MER1, SAL1 or SGLU1 for 48 hours, and then sulforhodamine B dye was added to the cultures. The sulforhodamine B dye is a protein binding dye and labels live cells. Results are reported as the percent growth of treated cells when compared to untreated control cells (negative data indicate cell kill).

MTT and Trypan Blue Assays. For these assays mononuclear cells from peripheral blood samples of leukemia patients and normal donors were separated by Ficoll Hipaque fractionation and resuspended in DMEM complete medium. Alternatively, cell line cells were used in some cases. Malignant cells from various human cell lines (usually at 5×10^4 cells/ml) or mononuclear cells from peripheral blood of leukemia patients and healthy donors (1×10^6 cells/ml) were incubated in either alpha MEM or RPMI 1640 with or without various concentrations of MER1, SAL1 or SGLU1. Each experimental condition was done in triplicate. After the indicated number of days (usually 3 days) of exposure to MER1, SAL1 or SGLU1, cell survival was assessed by the addition of a dye to the wells (either MTT or trypan-blue). The MTT dye changes its color depending on the presence of live cells in the well. Survival of cells under MTT treatment was evaluated as a percentage of control cell growth. The trypan-blue dye penetrates dead cells and live cells can be counted under the microscope and percentage survival estimated.

Clonogenic Assay. Clonogenicity or colony formation was analyzed by obtaining peripheral blood mononuclear cells (from normal donors or leukemia patients) which were resuspended in semisolid medium containing recombinant cytokines and plated in quadruplicate, 0.1 ml/well, in 96-well microtiter plates at 4×10^4 cells/0.1 ml density. Cell aggregates composed of more than 50 cells are counted as one colony after ~10 days of incubation at 37° C. in 5% CO₂ humidified atmosphere. Growth inhibition was evaluated as a percentage of colony growth as compared to colony growth in control (no drug) samples.

Analysis of Apoptosis. Three different methods were used to analyze apoptosis by assaying different events in the apoptotic pathways. Percentages of apoptotic cells induced by arsenic derivatives of the invention were evaluated using flow cytometer. Different methods of staining cells for apoptosis were utilized to assess different aspects of apoptotic cascade.

1. Annexin V and Propidium Iodide (PI) Staining. Annexin V binds to cells that express phosphatidylserine on the outer layer of the cell membrane, while propidium iodide stains the cellular DNA of cells with a compromised cell membrane. This allows live cells (unstained with either fluorochrome) to be discriminated from apoptotic cells (stained only with annexin V) and necrotic cells (stained with both annexin and PI).

Following treatment of cells in culture with indicated arsenicals of the invention for the indicated time, cells were washed in phosphate-buffered saline (PBS) and resuspended in 100 μ l of binding buffer containing annexin V-FITC (Travigene) and incubated for 15 minutes in the dark. Cells were analyzed on flow cytometer after the addition of PI.

2. Cytofluorometric Analysis of the Mitochondrial Membrane Potential. To evaluate the changes in the potential of mitochondrial membrane, following treatment with arsenic derivatives for indicated time, cells were incubated in sub-micromolar concentrations of MitoTracker probes. MitoTracker probes passively diffuse across the plasma membrane and accumulate in active mitochondria. Cells were stained with two colors: MitoTracker Red CMXRos (Molecular Probes) and MitoTracker Green FM (Molecular Probes). Cells were washed in PBS, stained with MitoTracker dyes and incubated for 1 hour at 37° C. in the dark. CMXRos is incorporated into mitochondria driven by the mitochondrial membrane potential and reacts with thiol residues to form covalent thiol ester bonds. MitoTracker Green FM dye preferentially accumulates in mitochondria regardless of mitochondrial membrane potential, making it a useful tool for determining mitochondrial mass.

3. Detection of Caspase Activity. In order to monitor caspase activity by flow cytometry, the fluorogenic substrate PhiPhiLux G1D1 (Oncoimmunin) was used. PhiPhiLux G1D1 is a substrate for the detection and measurement of caspase 3 and caspase 3-like activities in living cells. Following treatment with the arsenic derivatives of the invention for indicated time, cells were washed in PBS, resuspended in 5 μ l substrate solution and incubated for 1 hour at 37° C. in the dark. After incubation cells were washed, and few minutes before flow cytometry analysis, PI was added to exclude necrotic cells during analysis.

Cell Cycle Analysis. Cell cycle was analyzed as follows: After 72 h of incubation with the different arsenical compounds of the invention, cells (1×10^6) were washed twice in PBS. Cell pellet was resuspended in staining solution that contained hypotonic solution (RNase solution, Triton X-100, sodium citrate, PEG) and PI (25 μ g/ml). Cells were

incubated 15 minutes in dark at room temperature and then they were analyzed by flow cytometer using CellQuest program (Becton-Dickinson).

5 Maturation Analysis. Human acute prolymphocytic leukemia cell line NB4 was used to test the effect of the arsenicals of the invention on the maturation of leukemic cells. Phycoerythrin-conjugated anti-CD11b monoclonal antibody (Becton-Dickinson) was used as a marker of mature myelocytes. After 72 h of incubation with drugs, cells were washed in PBS. Cells in the density of 1×10^6 cells/ml were then incubated with monoclonal antibody in dilution 1:10 in dark at room temperature for 15 minutes. After incubation cells were washed in PBS and the pellet was resuspended in 500 μ l of PBS. To exclude nonspecific binding appropriate isotypic control was prepared in the same manner. Cells were sorted using a flow cytometer and analyzed using CellQuest Document Analysis.

15 Role of PML/RARalpha Protein. Arsenic trioxide is approved as a treatment for acute prolymphocytic leukemia and it kills APL cells in large part due to their expression of PML/RARalpha gene and protein. To establish whether the presence of PML/RARalpha fusion protein in the leukemic cells contributes to the observed sensitivity of leukemic cells to SGLU and MER1 the present inventors used the following system: U937 cells, known to be resistant to arsenic trioxide, were transfected with PML/RARalpha gene. The transfected cells are called U937/PR9 and were kindly provided by Dr. Michael Andreeff (M. D. Anderson Cancer Center). The PML/RARalpha gene becomes functional in the presence of zinc. Zn²⁺-inducible expression of the PML/RARalpha gene in the U937/PR9 cell line is described in Grignani et al. (1993). In order to establish the PML/RARalpha expression, cells were treated with 0.1 mM ZnSO₄ for 3 h before the addition of arsenic compounds for 72 h. PML/RAR α expression is typically established at about 3 h following zinc addition to the cells and is stable for 48 hours.

Example 3

In Vitro Evaluation of Anticancer Activity of MER1, SAL1, and SGLU1

40 The anti-leukemic activity of MER1 has been evaluated by 3 day MTT assay/trypan blue exclusion method against 6 different human leukemia cell lines: AML2, AML3 and HL60 (an AML derived cell line), NB4 (an APL derived cell line), K562 (a CML-BP derived cell line), and KBM7 (an AML derived cell line). MER1 was most effective against NB4 cells with an IC₅₀ (the concentration that results in 50% survival of cells, as compared to untreated control cells) at 1 μ M (FIG. 1). MER1 treatment of other cell lines, including the analysis of AML2 cells and KBM7 cells by the MTT assay and AML2 cells (see FIG. 2), AML3 cells, K562 cells, and HL60 cells by the trypan blue assay showed IC₅₀ between 1.5–4 μ M. This activity was similar to the activity of arsenic trioxide against these cell lines (examples of arsenic trioxide activity are shown in FIG. 1 and FIG. 2). MER1 was also tested for anticancer activity by the National Institute Of Health (NIH), in vitro against a panel of 60 tumor cell lines using sulforhodamine B assay (FIG. 3). The compound showed evidence of activity at low concentrations against a variety of tumor cell lines, but particularly against leukemia cells tested. At the concentration of 11M of MER1, the growth of all 6 leukemia cell lines tested was significantly retarded (<20% growth; FIG. 3, first panel).

65 The antileukemic activity of SAL1 has been evaluated by 3 day trypan blue assays against 2 human cell lines: HL60

19

cells (depicted in FIG. 4), and Z138 (an ALL cell line). SAL1 was also tested for anticancer activity by the NIH in vitro against a panel of 60 tumor cell lines using sulforhodamine B assay (FIG. 5). The compound showed evidence of activity at low concentrations against a variety of tumor cell lines.

The antileukemic activity of SGLU1 has been evaluated by 3 day MTT assay against 9 human leukemia cell lines: NB4, CAG (multiple myeloma cell line), JURKAT and RAJI (lymphoma cell line), HL60, AML2, AML3, KBM5 (CML-BP derived cell line, and KBM7. The results for the cell line NB4 are depicted in FIG. 6, as an example. The antileukemic activity of SGLU1 has also been evaluated by 3 day trypan blue exclusion method against 6 human leukemia cell lines: NB4, CAG, JURKAT, HL60, KBM3 (AML cell line), and Z119 (an ALL cell line), showing similar results. The activity was similar to the activity of arsenic trioxide as depicted in FIG. 6, as an example. The antileukemic activity of SGLU1 has also been evaluated by 5 day clonogenic assay against HL60 human leukemia cells (FIG. 7). SGLU1 was also tested for anticancer activity by the NIH in vitro against a panel of 60 tumor cell lines using sulforhodamine B assay (FIG. 8). The compound showed evidence of activity at low concentrations against a variety of tumor cell lines.

Example 4

Toxicity Determination of MER1 and SGLU1
Against Malignant and Normal Blood Cells

The present inventors also tested MER1 against blood mononuclear cells (>80% blasts) from 5 leukemia patients (3 with AML, one with CML-BP, and one with ALL; FIGS. 9–13). In short-term cell cultures MER1 was as effective as arsenic trioxide (an example is shown in FIGS. 9, 10, and 12). In addition, toxicity of MER1 against normal peripheral blood mononuclear cells was evaluated in samples from 4 healthy donors. In short-term cell suspension cultures by MTT assay MER1 was less toxic to normal cells than malignant cells from leukemia patients (FIG. 14). Most importantly, in long-term clonogenic assay MER1 was less toxic to normal cells than arsenic trioxide (FIG. 15).

SGLU1 was tested against blood mononuclear cells from 3 leukemia patients, including a patient with CLL (FIG. 16 which shows a comparison to arsenic trioxide), and 2 patients with AML (FIG. 17). In long-term clonogenic assay SGLU1 was less toxic than arsenic trioxide to normal cells (FIG. 18). In addition to the 8 day clonogenic assay shown in FIG. 18, 9 day and 13 day clonogenic assays have also been performed.

Example 5

Formulation and Stability of MER1

Data has been obtained that shows that MER1 is stable for at least 2 months when dissolved in phosphate buffered saline, as the solutions have maintained cytotoxic activity at the same level in in vitro experiments done during this time period (FIG. 19). In addition detailed pharmaceutical evaluation of MER1 and SGLU1 were performed.

I. Pharmaceutical Evaluation of MER-1

MER-1 was found to have a sufficient solubility and stability to be acceptable for administration in a clinical

20

setting (see data below). It is also sufficiently stable that solutions can be extemporaneously compounded for use in animal testing and possibly an early Phase I study. However, the solution stability is not adequate for manufacturing of larger batches of a liquid dosage form for use in larger clinical trials and distribution in the commercial marketplace where long-term storage is required. A lyophilized dosage form that is reconstituted at the time of use is contemplated for these applications. Preparation of such lyophilized compositions are well known in the art.

A. Solubility

MER-1 has an aqueous solubility of about 15 mg/mL. Higher MER-1 concentrations up to about 150 mg/mL can be achieved by the use of 0.1 N sodium hydroxide to adjust to pH 6. In ethanol, MER-1 has a solubility of more than 100 mg/mL.

B. Solution pH

The natural pH values of aqueous solutions of MER-1 are as follows:

0.1 mg/mL	pH 3.7
1 mg/mL	pH 3.1
10 mg/mL	pH 2.3

C. Solution Stability

The effects of various pH values were evaluated at a concentration of 10 mg/mL in 0.9% sodium chloride injection. Samples having a pH of 2.3 (natural pH) and also adjusted with sodium hydroxide to pH 5, 7.1, and 8.5 have been evaluated over a period of 3 months under refrigeration. The samples at pH 5 demonstrated better stability retaining about 89% of the initial concentration after 3 months. The solutions at pH 7.1 and 8.5 retained about 92% and 96%, respectively, after 14 days but fell below 90% after that time. The pH 2.3 samples were stable for 7 days but developed a precipitate after that time. See Table 2.

MER-1 is less stable in aqueous solutions at lower concentrations, but is increasingly stable at higher concentrations. At 0.1 mg/mL in water, about 40% of the drug was lost in as little as one hour. As concentrations increased from 1 to 10 mg/mL in 0.9% sodium chloride injection, the drug was stable for increasingly longer periods. The 10-mg/mL concentration was stable for up to 3 months under refrigeration, but unacceptable decomposition occurred after that time. See Table 3.

TABLE 2

pH Stability Profile of MER-1 10 mg/mL in 0.9% Sodium Chloride Injection				
Assay Interval (Days)	Remaining MER-1 (%)			
	pH 2.3	pH 5.0	pH 7.1	pH 8.5
0	100	100	100	100
7	102	105	96	97
14	ppt	101	92	96
30		100	80	82
60		91		
90		91		
180		87		

21

TABLE 3

MER-1 Solution Stability at Varying Concentrations in 0.9% Sodium Chloride Injection				
Assay Interval (days)	Remaining MER-1 (%)			
	0.1 mg/mL ^a	1 mg/mL	2 mg/mL	10 mg/mL
0	60 ^b	100	100	100
7		94	99	105
14		89	102	101
21		81	96	102
30		79	98	100
60			88	91
90			84	91
180				87

^aIn water.^bAbout 40% loss occurred in 60 minutes.^cNot determined at this interval.

II. Pharmaceutical Evaluation of SGLU-1

SGLU-1 is found to have sufficient solubility and stability to be acceptable for administration in a clinical setting. It is also sufficiently stable that solutions can be extemporaneously compounded for use in animal testing and possibly an early Phase I study. However, the solution stability is not adequate for manufacturing of larger batches of a liquid dosage form for use in larger clinical trials and distribution in the commercial marketplace where long-term storage is required. A lyophilized dosage form that is reconstituted at the time of use is contemplated for these applications.

A. Solubility

SGLU-1 has an aqueous solubility of about 60 mg/mL. Higher SGLU-1 concentrations can be achieved by the use of 0.1 N sodium hydroxide to raise the solution pH. However, the drug appears to be unstable in an alkaline environment. SGLU-1 is insoluble in ethanol.

B. Solution pH

The natural pH values of aqueous solutions of SGLU-1 are:

0.1 mg/mL	pH 3.9
1 mg/mL	pH 3.2
2.5 mg/mL	pH 3.0
60 mg/mL	pH 2.7

C. Solution Stability

The effects of various pH values were evaluated at a concentration of 2.5 mg/mL in 0.9% sodium chloride injection. Samples having a pH of 3 (natural pH) and also adjusted with sodium hydroxide to pH 5 and 7 were evaluated over 30 days under refrigeration. The samples at pH 5 demonstrated slightly better stability retaining about 90% concentration after 30 days. The solutions at pH 3 and 7 retained about 84% and 82%, respectively. See Table 4.

SGLU-1 at concentrations of 20 mg/mL and 50 mg/mL in 0.9% sodium chloride injection adjusted to pH 5 have undergone stability testing. Less than 10% loss occurred through 60 days of storage under refrigeration. The stability results are shown in Table 5.

SGLU-1 is less stable at lower concentrations. At 0.1 mg/mL in water, more than 10% decomposition occurred in 24 hours at room temperature.

22

TABLE 4

pH Stability Profile of SGLU-1 2.5 mg/mL in 0.9% Sodium Chloride Injection			
Assay Interval (Days)	Remaining SGLU-1 (%)		
	pH 3.0	pH 5.0	pH 7.0
0	100	100	100
3	104	101	97
5	100	99	93
7	100	97	91
14	97	97	91
21	87	93	84
30	84	91	82

TABLE 5

Stability if SGLU-1 20 mg/mL and 50 mg/mL in 0.9% Sodium Chloride Injection at 4° C.		
Assay Interval (Days)	Remaining SGLU-1 (%)	
	20 mg/mL	50 mg/mL
7	101	97
14	98	98
30	94	94
60	92	93
90	85	87

Example 6

Mechanisms for MER1, SAL1, and SGLU1

The induction of apoptosis, effects on the cell cycle, induction of maturation, and degradation of aberrant PML/RARalpha fusion protein, have all shown to be mechanisms of action of arsenic trioxide. The present inventors have examined the potential of MER1 to induce apoptosis in HL60 human leukemia cells (assay time 1–3 days). The induction of apoptosis followed closely diminished percentage of surviving cells (FIGS. 20, 21 & 22). Additional studies using both MER1 and SGLU have established that the induction of apoptosis (annexin V staining) by these compounds involves change of the potential of mitochondrial membrane (CMXRos staining) and activation of caspases (PhiPhiLux staining) see FIGS. 23A, 23B, 23C, 23D, 23E, & 23F.

It has been reported that arsenic trioxide induced maturation of cells expressing PML/RARalpha gene. To test whether SGLU and MER1 have similar capability NB4 cells (expressing the PML/RARalpha gene) were used and, after 3 days of exposure to arsenicals, the expression of CD11b on the surface of the cells was measured by flow cytometer. CD11b is a maturation marker for myeloid cells. Data is presented in FIG. 24 indicating that SGLU and MER1 do not induce maturation. Possible cell cycle disturbance in HL60 cells treated with the different arsenicals of the invention was assessed using flow cytometry and staining with propidium iodide. It was found that SGLU caused marked accumulation of cells in S-phase of cell cycle, with MER1 causing a similar effect to lesser degree (FIGS. 25A, 25B). FIG. 25C describes S-phase accumulation of cells in response to arsenic trioxide.

To establish whether the presence of PML/RARalpha fusion protein in the leukemic cells contributes to the observed sensitivity of leukemic cells to SGLU and MER1

23

the following system was used: U937 cells, known to be resistant to arsenic trioxide, were transfected with PML/RARalpha gene. This gene becomes functional in the presence of zinc. Thus, transfected U937 cells (U937/PR9) were treated with different arsenical with or without zinc. Results are shown in FIGS. 26A, 26B & 26C which indicate that the presence of functional PML/RARalpha gene is pre-requisite for cells to become sensitive to arsenic trioxide but have no influence on the sensitivity of the cells to SGLU and MER1.

Example 7

In Vivo Evaluation of the Therapeutic Potential of MER1, SAL1, and SGLU1

The animal model of human leukemia is represented by severe combined immunodeficient (SCID) mouse bearing human leukemia cells. This model is unique in that it allows growth of human leukemia in animals in a mode similar to that seen in patients. It offers an opportunity to rapidly test the in vivo efficacy of new drugs at different dose levels and schedules. Furthermore, not only can animal survival be monitored, but also the effect of treatment on the dissemination pattern of the disease. Treatment of SCID mice typically starts 2 days after inoculation with human leukemic cells. Initial in vivo experiments in SCID mice injected with one human leukemia cell line will determine dose and schedule of MER1, SAL1 or SGLU1 for the other mice model as well as for initial human trials.

Animals are monitored daily and sacrificed when moribund or at the completion of the study (usually double the survival time of the control group). Necropsy is performed on animals that survive for long time and tissues are analyzed for the presence of human DNA by polymerase chain reaction (PCR) using primers specific for DNA sequences of HLA-DQ α . Since leukemia is a systemic disease, the presence of minimal residual disease is studied by checking for HLA-DQ α in DNA from different mouse tissues. This data may help predict whether the drug is active in selective compartments for leukemia, e.g. bone marrow, solid organs, or central nervous system.

Prerequisites for in vivo therapeutic experimentation in SCID mice are 1) the verification of leukemia cells engraftment in animals and 2) the determination of acute toxicity of tested compounds (definition of maximally tolerated dose).

I. Verification of Leukemia Cells Engraftment in Animals. The first in vivo experiment involved 4 groups of SCID mice. Five mice per group were inoculated intraperitoneally with human leukemia cells of different types: HL60 (AML), KBM5 (CML-BP), KBM7-acute myeloid leukemia, and Z119 (ALL). HL60 and KBM5 cells showed excellent engraftment: in HL60 group all mice died within days 31 and 36 after inoculation, while in the KBM5 group mice died within 34th and 36th day. The engraftment was verified by performing PCR for DNA sequences of human HLA-DQ α (test was positive in all tissues from all the mice). At day 100, there were still 4 of 5 mice alive in KBM7 and 5 of 5 mice alive in Z119 group. At that day all mice were sacrificed and tissues analyzed by PCR for HLA-DQ α . Testing was negative indicating lack of leukemia cell engraftment. Alternative cell lines of same type will be required for planned therapeutic studies.

II. Determination of Acute Toxicity of Tested Compounds. For toxicology testing, immunocompetent Swiss Webster mice were used. The present inventors have confirmed that LD₅₀ concentration for arsenic trioxide is 10 mg/kg.

24

A. Brief Toxicity Testing Of SGLU1 in Swiss-Webster Mice. Two studies were performed on Swiss-Webster mice to test the toxicity of SGLU1. In the first study, SGLU1 was administered at doses of 178 mg/kg; 285 mg/kg; and 357 mg/kg via the IP route. The toxicity was measured by the mortality of mice. It was found that the mice tolerated the 178 mg/kg and the 285 mg/kg doses of SGLU1 well. The data of this study are summarized in Table 6.

TABLE 6

Brief Toxicity Testing Of SGLU1			
	Dose (mg/kg)		
Administered IP	178	285	357
Dead/Total Number	0/5	1/5	5/5

In the second study the toxicity with each mouse weighted were studies for administration of SGLU1 by both the IP and IV routes at doses of 318 mg/kg and 375 mg/kg. Thus, inventors have established that LD₅₀ concentration for SGLU1 is 350 mg/kg. The results are summarized in Table 7.

TABLE 7

Better Performance With Each Mouse Weighted		
	Dose (mg/kg)	
	318	375
Dead/Total Number for IP Administration	0/5	4/5
Dead/Total Number for IV Administration	1/5	5/5

B. Brief Toxicity Testing Of MER-1 in Swiss-Webster Mice. Two studies were performed on Swiss-Webster mice to test the toxicity of MER-1. In the first study, MER-1 was administered at doses of 71 mg/kg; 107 mg/kg; and 143 mg/kg via the IP route. The toxicity was measured by the mortality of mice. It was found that the mice tolerated the 71 mg/kg and the 107 mg/kg doses of MER-1 well with no mortality. The data of this study are summarized in Table 8.

TABLE 8

Brief Toxicity Testing Of MER-1			
	Dose (mg/kg)		
Administered IP	71	107	143
Dead/Total Number	0/5	0/5	5/5

In the second study the toxicity with each mouse weighted were studies for administration of MER-1 by both the IP and IV routes at doses of 125 mg/kg; 156 mg/kg; and 170 mg/kg. Thus, inventors have established that LD₅₀ concentration for MER1 is 150 mg/kg. The results are summarized in Table 9.

TABLE 10-continued

MER1 Single Dose: 3 Day, 14 Day, 42 Day Toxicology								
Serum Chem-14 day								
Serum Chem-42 day								
Comment:		•						
Gross Path - 3 day	Male	Female	Male	Female	Male	Female	Male	Female
Gross Path - 14 day								
Gross Path - 42 day								
Organ wts - 3 day (rel. to brain)								
Organ wts - 14 day (rel. to brain)								
Organ wts - 42 day (rel. to brain)								
Comment:		•						
Histopathology	Low		Low-intermediate		High-intermediate		High	
Sex	F	M	F	M	F	M	F	M
Dose: mg/kg/day	0	57	80	86	120	120	150	150
	<u>Lung/Heart: Arteritis/Vasculitis/thrombosis</u>							
72 h	0/5	0/5	0/5	1/5	0/5	0/2	1/5	0/5
14 d						2/5	1/5	1/5
40/42 d								
Total								
	<u>Ovary: increased apoptosis of follicular cells</u>							
72 h								
14 d								
40/42 d								
Total								
Comment:	• Vascular lesions in heart and lungs. To date, effect level in male arm = 86 mg/kg and female arm = 150 mg/kg.							

Unless indicated otherwise, values = treatment group mean + control group mean. "—" indicates no compound-related finding.

B. Multiple-Dose Toxicology Studies

Further studies were performed to determine the dose-limiting toxicity associated with the administration of

repeated doses in groups of mice. Tables 11–16 below describe the results of the multiple-dose toxicology studies for MER-1 and Tables 17–21 describe results of the multiple-dose toxicology of SGLU-1.

TABLE 11

MER-1 Multiple Dose Toxicology in Vivo				
Multiple Dose 28 d Toxicology				
Strain: Swiss Webster		Dosing Frequency: Daily x 5		
Age at Start: ~6 wks		Dosing Volume: 10 mL/kg		
Group Size: 3/grp, male only		Vehicle: Saline, pH w/NaOH to 5.0		
Groups: 0, 30, 40, 50, 60, 70, 80, 90 and 100 mg/kg/d		Lot: Merida Sotelo notebook, p43, May 10, 2001, M. P. 151 degrees		
Operator: Julie Miller		Dose Date: 10 MAR 02–14 MAR 02		
		Sacrifice Date: 12 APR 02		
Objective: To investigate the potential toxicity and maximally-tolerated cumulative dose of MER1 when given intravenously (tail vein) daily x 5 to mice. Total cumulative dose = 150, 200, 250, 300, 350, 400, 450, and 500 mg/kg.				
Parameters Evaluated: Clinical signs (daily), body weight (pre dose and twice weekly) x 4 weeks. Necropsy moribund/dead animals. Necropsy surviving animals (d 28).				
To process and read: Heart, lung, liver, kidney (d 28) in control (0 mg/kg) and high dose (100 mg/kg) groups. Depending on whether lesions are found, next lower dose groups may be evaluated.				
Conclusions:				
	<u>MER1-Related Findings</u>			
mg/kg/day	30	40	50	60
Death	—	—	—	—
Comment:	Initial period of hyperactivity after dosing, followed by slightly decreased activity. Normal activity resumed 2–3 hours after dosing.			
Clinical Signs				
Comments	Animal tolerance of multiple doses of pH adjusted MER1 —MER1_02 study. There is no			

TABLE 11-continued

MER-1 Multiple Dose Toxicology in Vivo								
evidence of tail necrosis after 4 days of daily dosing in the 0–80 mg/kg/day groups; no evidence of tail necrosis after 3 days of daily dosing in the 90–100 mg/kg/day groups. All animals recover within several hours from short period of moderately decreased activity. Tail lesions and necrosis first noted one day after daily × 5 dosing is completed (Day 6) in all dose groups from 30–100 mg/kg/d. Animals are monitored daily and tails amputated as needed.								
Fd Consump	N/A		N/A		N/A		N/A	
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female
	35 g	—	39 g	—	34 g	—	35 g	—
Histopathology								
28 d								

Unless indicated otherwise, values = treatment group mean + control group mean.
 “—” indicates no compound-related finding.

TABLE 12

Analysis of Organ Lesions in Response to MER-1					
Animal No.	Dose Group	Heart	Lung	Liver	Kidneys
MER1-02-1001	0 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-1003	0 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-1005	0 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-2001	30 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-2003	30 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-2005	30 mg/kg/day	No significant lesion	Vasculitis, 1+ Fibrin thrombi, 1+	No significant lesion	Inflammation, 1+
MER1-02-3001	40 mg/kg/day	No significant lesion	Fibrin thrombi, 1+	No significant lesion	No significant lesion
MER1-02-3003	40 mg/kg/day	No significant lesion	No significant lesion	Infarct, 1+	No significant lesion
MER1-02-3005	40 mg/kg/day	No significant lesion	Vasculitis, 1+ Fibrin thrombi, 1+	No significant lesion	No significant lesion
MER1-02-4001	50 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-4003	50 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-4005	50 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-5001	60 mg/kg/day	Vasculitis, 1+	Cardiomyopathy, 1+	No significant lesion	No significant lesion
MER1-02-5003	60 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-5005	60 mg/kg/day	Perivasculitis, 1+	No significant lesion	No significant lesion	No significant lesion
MER1-02-6001	70 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-6003	70 mg/kg/day	Perivasculitis, 1+	No significant lesion	No significant lesion	No significant lesion
MER1-02-6005	70 mg/kg/day	Vasculitis, 1+ Fibrin thrombi, 1+	No significant lesion	No significant lesion	No significant lesion
MER1-02-7001	80 mg/kg/day	No significant lesion	No significant lesion	No significant lesion	No significant lesion
MER1-02-7003	80 mg/kg/day	No significant lesion	No significant lesion	CL Hypertrophy, 1+	No significant lesion
MER1-02-7005	80 mg/kg/day	No significant lesion	No significant lesion	CL Hypertrophy, 1+ Hemorrhage, 1+	No significant lesion
MER1-02-8001	90 mg/kg/day	Perivasculitis, 1+	No significant lesion	CL Hypertrophy, 1+	No significant lesion
MER1-02-8003	90 mg/kg/day	No significant lesion	No significant lesion	CL Hypertrophy, 1+	No significant lesion
MER1-02-8005	90 mg/kg/day	No significant lesion	No significant lesion	CL Hypertrophy, 1+	No significant lesion
MER1-02-9001	100 mg/kg/day	No significant lesion	No significant lesion	CL Hypertrophy,	No significant lesion

TABLE 12-continued

Analysis of Organ Lesions in Response to MER-1									
Lesion	30 mg/kg	40 mg/kg	50 mg/kg	60 mg/kg	70 mg/kg	80 mg/kg/d	90 mg/kg/d	100 mg/kg/d	
Vasculitis/perivasculitis	1/3	1/3	0/3	2/3	2/3	0/3	1/3	0/3	
Fibrin thrombi	1/3	2/3	0/3	0/3	1/3	0/3	0/3	1/3	
CL Hypertrophy	0/3	0/3	0/3	0/3	0/3	2/3	3/3	3/3	

15

The vasculitis/perivasculitis/and fibrin thrombi are considered part of the same process, and are related to IV administration of MER1. In this study, there is not a no-observed-effect-level for the vasculitis. At the higher does (80 mg/kg/d and above) there was central lobular

hypertrophy of hepatocytes in the liver (CL Hypertrophy). This finding is not uncommon when a xenobiotic is metabolized in the liver, and the enlarged cells are indicative of smooth endoplasmic reticulum increase. This could be confirmed by measuring enzyme induction in the liver.

TABLE 13

MER-1 Multiple Dose Toxicology									
MER1 Multiple Dose 28 d Toxicology					Dosing Frequency: Daily x 5				
Strain: Swiss Webster					Dosing Volume: ~10 mL/kg				
Age at Start: ~16 wks					Vehicle: Saline, pH w/NaOH to 5.0				
Group Size: 5/sex, male and female					Lot: Merida Sotelo notebook, p43, May 10, 2001, M.P. 151 degrees				
Groups: 110, 120, 130, 140, 150 mg/kg/d					Objective: To investigate the potential toxicity and maximally-tolerated cumulative dose of MER1 when given intravenously (tail vein) daily x 5 to mice. Total cumulative dose = 500, 550, 600, 650, 700, 750 mg/kg.				
					Parameters Evaluated: Clinical signs (daily), body weight (pre dose and twice weekly) x 4 weeks.				
					Necropsy moribund/dead animals. Necropsy surviving animals (d 27, Friday JUN 14, 2002).				
					To Process and read: Heart, lung, kidney, liver in moribund/dead animals and in surviving animals (d 28) in control (0 mg/kg) and high does groups.				
Conclusions:									
•									
MER1-Related Findings									
mg/kg/day	0		110		120		130		
Death			2F 2M d2	3M d3	1F d1,	2F 5M d 2	1F d3, 1 F d4	1F d5	
Comment:									
Clinical Signs	—		—		—		Decr. activity, rough hair coat, tail necrosis.		
Comment:									
Fd Consump	N/A		N/A		N/A		N/A		
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female	
Average Weight→	51 g	40 g	50 g	38 g	49 g	36 g	48 g	35 g	
Standard Deviation→	0.5477	0	0.4472	1.3038	0.5477	0.5477	0.5477	0.5477	
MER1-Related Findings									
mg/kg/day	140		150						
Death	2F 2 M d1,		3F 5M d1,		2F d2				
Comment:									
Clinical Signs	—		—		Decr. activity & rough hair coat.				
Comment:									
Fd Consump	N/A		N/A		N/A		N/A		
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female	
Average Weight→	47 g	34 g	46 g	33 g	—	—	—	—	
Standard Deviation→	0	0.4472	0.5477	0					
MER1-Related Findings									
mg/kg/day	70		80		90		100		
Death	—		—		—		—		
Comment:									
Clinical Signs	Initial period of hyperactivity after dosing, followed by moderately decreased activity. Normal activity resumed 2-3 hours after dosing.								

TABLE 13-continued

MER-1 Multiple Dose Toxicology									
Fd Consump Body Weights	N/A		N/A		N/A		N/A		
	Male	Female	Male	Female	Male	Female	Male	Female	
	35 g	—	35 g	—	29 g	—	31 g	—	
Histopathology 28 d									
mg/kg/day	0								
Death	—								
Comment:									
Clinical Signs									
Fd Consump	—								
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female	
		—		—		—		—	
Histopathology 28 d									
mg/kg/day	—								
Gross Path	—								
	00		MER1-Related Findings				130		
	Male	Female	Male	Female	Male	Female	Male	Female	
Organ wts (rel. to brain)	—	—	—	—	—	—	—	—	
Comment:	—								
Histopathology	—								
Sex	•								
Dose: mg/kg/day									
Lung/Heart:	F	M	F	M	F	M	F	M	
28 d									
Total									
Ovary:									
28 d									
Total									
Comment:	•								
mg/kg/day	—								
Gross Path	—								
	140		150						
	Male	Female	Male	Female	Male	Female	Male	Female	
Organ wts (rel. to brain)	—	—	—	—	—	—	—	—	
Comment:	—								
Histopathology	—								
Sex	•								
Dose: mg/kg/day									
Lung/Heart:	F	M	F	M	F	M	F	M	
28 d									
Total									
Ovary:									
28 d									
Total									
Comment:									

Unless indicated otherwise, values = treatment group mean ± control group mean.
 “—” indicates no compound-related finding.

TABLE 14

50

Mortality and Number of Doses Tolerated for MER1										
No. of doses	110 mg/kg		120 mg/kg		130 mg/kg		140 mg/kg		150 mg/kg	
	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose
	F	M	F	M	F	M	F	M	F	M
1	1/5		2/5	2/5			4/5	4/5	4/5	5/5
2	1/5	2/5	1/5	3/5	2/5	1/5	1/5	1/5		
3		3/5			1/5					

TABLE 14-continued

Mortality and Number of Doses Tolerated for MER1										
No. of doses	110 mg/kg		120 mg/kg		130 mg/kg		140 mg/kg		150 mg/kg	
	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose	Number of animals/dose
	F	M	F	M	F	M	F	M	F	M
4			1/5		1/5					
5	3/5		1/5		1/5					

Based on mortality, there is not a no-observable-effect level (NOEL) for this study.

TABLE 15

<u>Summary of Histopathological Findings for MER1 Treated Females</u>			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
<u>0 mg/kg/day females</u>			
MER1-03-1002	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Modest, 1+ focal lymphoid aggregate
MER1-03-1004	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Modest, 1+ multifocal lymphoid aggregates
MER1-03-1006	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Modest, 1+ focal subacute/chronic inflammation
MER1-03-1008	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Modest, 1+ focal subacute/chronic inflammation
MER1-03-1010	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
<u>110 mg/kg/day females</u>			
MER1-03-2002	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-2004	1	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Mild, 2+ cell death, lymphoid aggregates
MER1-03-2006	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions Tail: Severe, 4+ acute necrosis with vasculitis, thrombosis, and ulcerative dermatitis
MER1-03-2008	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: modest, 1+ tubular necrosis, medulla
MER1-03-2010	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: Modest, 1+ hypertrophy/hyperplasia, alveolar macrophages Liver: no significant lesions Kidney: no significant lesions
<u>120 mg/kg/day females</u>			
MER1-03-3002	1	S-May 19, 2002	Heart: no significant lesions Lungs: Modest, 1+ fibrin thrombus Liver: no significant lesions Kidney: no significant lesions
MER1-03-3004	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-3006	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: Modest, 1+ focal lymphoid aggregate Tail: Severe, 4+ acute necrosis with vasculitis, thrombosis, and ulcerative dermatitis
MER1-03-3008	4	S-May 22, 2002	Heart: no significant lesions Lungs: Modest, 1+ fibrin thrombus Liver: Mild, 2+ centrilobular hepatocyte hypertrophy w/ vacuolar degeneration Kidney: no significant lesions Tail: Moderate, 3+ acute necrosis with vasculitis, thrombosis, and ulcerative dermatitis
MER1-03-3010	1	S-May 20, 2002	Heart: no significant lesions Lungs: Modest, 1+ thrombosis, veins

TABLE 15-continued

<u>Summary of Histopathological Findings for MER1 Treated Females</u>			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
			<u>Liver: Moderate, 3+ hypertrophy panlobular hepatocytes w/vacuolar degeneration Kidney: Mild, 2+ tubular necrosis, C-M junction 130 mg/kg/day females</u>
MER1-03-4002	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-4004	4	S-May 22, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions Tail: Severe, 4+ acute necrosis with vasculitis, thrombosis, and ulcerative dermatitis
MER1-03-4006	5	S-May 23, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-4008	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: 1. Moderate, 3+ hypertrophy panlobular hepatocytes w/vacuolar degeneration; 2. Modest, 1 + focal subacute/chronic inflammation, portal area Kidney: no significant lesions
MER1-03-4010	3	S-May 21, 2002	Heart: no significant lesions Lungs: Mild, 2+ acute thrombophlebitis, veins Liver: no significant lesions Kidney: no significant lesions Tail: Moderate, 3+ acute necrosis with vasculitis, thrombosis, and hemorrhage <u>140 mg/kg/day females</u>
MER1-03-5002	2	S-May 20, 2002	Heart: no significant lesions Lungs: Modest, 1+ fibrin thrombus Liver: no significant lesions Kidney: no significant lesions
MER1-03-5004	1	S-May 19, 2002	Heart: no significant lesions Lungs: Moderate, 3+ acute congestion & hemorrhage Liver: no significant lesions Kidney: no significant lesions
MER1-03-5006	1	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-5008	1	D-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: Mild to moderate, 2-3+ acute hepatocellular necrosis Kidney: no significant lesions
MER1-03-5010	1	S-May 19, 2002	Heart: no significant lesions Lungs: Modest, 1+ vasculitis & fibrin thrombus Liver: no significant lesions Kidney: no significant lesions <u>150 mg/kg/day females</u>
MER1-03-6002	1	D-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-6004	1	S-May 19, 2002	Heart: no significant lesions Lungs: Moderate, 3+ acute fibrin thrombosis, alveolar capillaries Liver: no significant lesions Kidney: no significant lesions
MER1-03-6006	1	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-6008	1	S-May 19, 2002	Heart: no significant lesions Lungs: Mild, 2+ acute fibrin thrombosis, alveolar capillaries

TABLE 15-continued

Summary of Histopathological Findings for MER1 Treated Females			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
MER1-03-6010	1	S-May 19, 2002	Liver: Moderate, 3+ hypertrophy panlobular hepatocytes w/vacuolar degeneration Kidney: no significant lesions Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions

TABLE 16

Summary of Histopathological Findings for MER1 Treated Males			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
<u>0 mg/kg/day males</u>			
MER1-03-1001	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: 1. Moderate 3+, increased cytoplasmic glycogen, hepatocytes 2. Modest 1+, focal acute necrosis Kidney: no significant lesions
MER1-03-1003	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-1005	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-1007	5	S-Jun. 14, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-1009	5	S-Jun. 14, 2002	Heart: 1. Moderate, 3+ medial hyperplasia, coronary arteries 2. Mild, 2+ polyarteritis, coronary arteries Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
<u>110 mg/kg/day males</u>			
MER1-03-2001	3	S-May 21, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-2003	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-2005	3	S-May 21, 2002	Heart: 1. Mild, 2+ acute subendocardial myocarditis 2. Mild, 2+ myocyte vacuolation Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-2007	2	S-May 20, 2002	Heart: no significant lesions Lungs: Mild, 2+ acute hemorrhage Liver: Modest, 1+ centrilobular hepatocyte hypertrophy w/vacuolar degeneration (microvesicular) Kidney: no significant lesions
MER1-03-2009	3	S-May 21, 2002	Heart: no significant lesions Lungs: Mild, 2+ acute hemorrhage Liver: ,1+ centrilobular hepatocyte hypertrophy Kidney: Mild, 2+ lymphoid hyperplasia, pelvis & perivascular
<u>120 mg/kg/day males</u>			
MER1-03-3001	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions

TABLE 16-continued

Summary of Histopathological Findings for MER1 Treated Males			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
MER1-03-3003	2	S-May 20, 2002	Liver: Mild, 2+ cystic hyaline degeneration of hepatocytes Kidney: no significant lesions Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions
MER1-03-3005	2	S-May 20, 2002	Kidney: no significant lesions Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions
MER1-03-3007	1	S-May 20, 2002	Kidney: Modest, 1+ hyaline casts tubules, medulla Heart: no significant lesions Lungs: Modest, 1+ acute thrombophlebitis Liver: no significant lesions
MER1-03-3009	1	D-May 20, 2002	Kidney: no significant lesions Heart: no significant lesions Lungs: Modest, 1+ vasculitis Liver: no significant lesions Kidney: no significant lesions <u>130 mg/kg/day males</u>
MER1-03-4001	1	D-May 20, 2002	Heart: no significant lesions Lungs: 1. Mild, 2+ acute hemorrhage; 2. Modest, 1 + fibrin thrombus Liver: no significant lesions
MER1-03-4003	1	D-May 20, 2002	Kidney: Mild, 2+ acute tubular necrosis Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions
MER1-03-4005	1	D-May 20, 2002	Kidney: no significant lesions Heart: no significant lesions Lungs: Mild, 2+ acute hemorrhage Liver: no significant lesions
MER1-03-4007	2	S-May 20, 2002	Kidney: Mild, 2+ acute tubular necrosis Heart: no significant lesions Lungs: Modest, 1+ acute hemorrhage Liver: Mild, 2+ fatty change hepatocytes
MER1-03-4009	1	D-May 20, 2002	Kidney: Modest, 1+ acute tubular necrosis Heart: no significant lesions Lungs: Modest, 1+ fibrin thrombus Liver: no significant lesions Kidney: no significant lesions <u>140 mg/kg/day males</u>
MER1-03-5001	2	S-May 20, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions
MER1-03-5003	1	S-May 20, 2002	Kidney: no significant lesions Heart: no significant lesions Lungs: Modest, 1+ acute venous thrombosis Liver: Moderate, 3+ hypertrophy panlobular hepatocytes
MER1-03-5005	1	D-May 20, 2002	Kidney: Mild, 2+ lymphoid hyperplasia, pelvis Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions
MER1-03-5007	1	D-May 19, 2002	Kidney: no significant lesions Heart: Moderate, 3+ medial hyperplasia, coronary arteries Lungs: Moderate, 3+ atelectasis Liver: no significant lesions
MER1-03-5009	1	D-May 19, 2002	Kidney: no significant lesions Heart: no significant lesions Lungs: Modest, 1+ acute hemorrhage Liver: Mild, 2+ hypertrophy panlobular hepatocytes Kidney: Mild, 2+ infarct kidney <u>150 mg/kg/day males</u>
MER1-03-6001	1	S-May 19, 2002	Heart: Mild, 2+ medial hyperplasia, coronary arteries Lungs: Mild, 2+ acute hemorrhage Liver: Mild, 2+ hypertrophy panlobular hepatocytes Kidney: no significant lesions

TABLE 16-continued

Summary of Histopathological Findings for MER1 Treated Males			
Animal ID	# doses	Sac. or Death-date	Diagnoses and/observations
MER1-03-6003	1	S-May 19, 2002	Heart: no significant lesions Lungs: Modest, 1+ thrombosis, veins & capillaries Liver: Mild, 2+ hypertrophy panlobular hepatocytes Kidney: no significant lesions
MER1-03-6005	1	S-May 19, 2002	Heart: no significant lesions Lungs: Modest, 1+ thrombosis, veins & capillaries Liver: no significant lesions Kidney: no significant lesions
MER1-03-6007	1	S-May 19, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions
MER1-03-6009	1	S-May 19, 2002	Heart: no significant lesions Lungs: no significant lesions Liver: no significant lesions Kidney: no significant lesions

The results of the MER1 Multiple Dose I.V. toxicology study evaluating 110, 120, 130, 140, and 150 mg/kg daily x5, are summarized as follows: The study focused on the microscopic examination of heart, lung liver, and kidney. Medial hyperplasia of coronary arteries is usually a spontaneous lesion that is more common in male mice than in female mice. Inflammatory lesions, e.g. lymphocyte aggregates, in the kidney and liver are concluded to be incidental findings unrelated to MER1. Hepatocyte hypertrophy and acute necrosis of renal tubules are lesions of uncertain significance encountered inconsistently in treated mice of both sexes. The panlobular hypertrophy observed in

males at the higher doses was sometimes associated with a microvesicular vacuolation suggesting possible hepatic toxicity. These animals generally were sacrificed in extremis after a single dose. Vascular lesions in lungs of these multi-dose mice are not consistent or striking when present although vasculopathies were common in the single-dose study of MER1. The tolerance of fewer doses and greater mortality in male mice suggests that the male gender is more sensitive to toxic effects of this compound. Analysis for a difference in metabolism in the liver between the sexes may be helpful.

TABLE 17

SGLU1 - Multiple Dose Toxicology In Vivo									
SGLU1					SGLU1-Related Findings				
Multiple - Dose Toxicity					Dose Dates Females: JUN 19, 02-JUN 23, 02				
Dose Dates Males: JUN 18, 02-JUN 22, 02					Dosing Frequency: Daily x 5				
Strain: Swiss Webster					Dosing Volume: 10 mL/kg				
Age at Start: ~6 wks					Vehicle: Saline				
Group Size: 5/grp, male and female					Lot: xxxxxxxxx				
Groups: 0, 50, 100, 150, 200, 250, 300, and 350 mg/kg/d					Objective: To investigate the potential toxicity and LD10-LD90 of SGLU1 when given intravenously (tail vein) daily x 5 to mice. Total cumulative dose = 250, 500, 750, 1000, 1250, 1500 and 1750 mg/kg. (IV single dose MTD = 350 mg/kg)				
Parameters Evaluated: Clinical signs (daily), body weight (pre dose and twice weekly) x 4 weeks. Necropsy moribund/dead animals.					To process and read: Heart, lung, kidney, liver in moribund/dead animals and in surviving animals (d 28) in control (0 mg/kg) and high dose groups.				
Conclusions: •									
mg/kg/day	50		100		150		200		
Death	—		—		—		—		
Comment:									
Clinical Signs	None.		None.		None.		2F tail necrosis: amputate		
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female	
Avg. in grams	25	25	23	24	23	25	24	24	
Std Dev.	0.707	0.837	1.304	0.837	0.707	0.894	0.894	0.447	
Comment:	•								
mg/kg/day	250		300		350		0		
Death	1F d4		3F d2, 1M d3		2M d1, 4F d1 1M d2, 1F d2		—		
Comment:									
Clinical Signs	2M tail necr.: amputate		Slight decr. activity 1F tail necr.: amputate		Moderate decr. activity 2M tail necr.: amputate		None.		
Body Weights	Male	Female	Male	Female	Male	Female	Male	Female	

TABLE 20-continued

		Compound-Related Lesions															
		Incidences of Test Substance-Related Microscopic Changes in Female and Male Mice															
		Group Designation															
		Concentration (mg/kg/day)															
		Number of Mice in Group															
		1		2		3		4		5		6		7		8	
		0		50		100		150		200		250		300		350	
		5		5		5		5		5		5		5		5	
		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<u>Liver:</u>																	
Hepatocyte hypertrophy																	
Centrilobular (compatible with SER)																	
Panlobular w/eosinophilia																	
Periportal w/vacuolation																	
<u>Thymus:</u>																	
Apoptosis, increased																	
<u>Testes:</u>																	
Seminiferous tubule degeneration																	
Epididymis:																	
Oligospermia																	

^aIndicates number of mice with microscopic lesion.

TABLE 21

		Summary Incidences of All Microscopic Observations															
		Group 1		Group 2		Group 3		Group 4		Group 5		Group 6		Group 7		Group 8	
		0 mg/kg		50 mg/kg		100 mg/kg		150 mg/kg		200 mg/kg		250 mg/kg		300 mg/kg		350 mg/kg	
		Incidence		Incidence		Incidence		Incidence		Incidence		Incidence		Incidence		Incidence	
		Per organ		Per organ		Per organ		Per organ		Per organ		Per organ		Per organ		Per organ	
		Examined		Examined		Examined		Examined		Examined		Examined		Examined		Examined	
Organ:		F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M
<u>Heart:</u>																	
Cardiomyopathy																	
Interstitial fat, increased																	
Medial hypertrophy, coronary artery																	
<u>Lung:</u>																	
Congestion																	
Fibrin Thrombi																	
Thrombus, organized																	
Vasculitis																	
Hemorrhage, focal																	
Pleural adhesion, focal																	
Atelectasis																	
Inflammation, focal																	
<u>Liver:</u>																	
Pyogranuloma, focal																	
Inflammation, portal																	
Increased mitoses																	
Hepatocyte hypertrophy:																	
Centrilobular:																	
Panlobular w/eosinophilia:																	
Periportal w/vacuolation:																	
Focus of Alteration, eosinophilic																	
Ito cell hypertrophy																	
Necrosis, focal																	
<u>Kidney:</u>																	
Murine progressive glomerulonephropathy																	
Inflammation, focal																	
Congestion																	

TABLE 21-continued

Organ:	Summary Incidences of All Microscopic Observations																
	Group 1 0 mg/kg		Group 2 50 mg/kg		Group 3 100 mg/kg		Group 4 150 mg/kg		Group 5 200 mg/kg		Group 6 250 mg/kg		Group 7 300 mg/kg		Group 8 350 mg/kg		
	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	Incidence Per organ Examined	
Microscopic Observation	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M	
<u>Thymus:</u>																	
Apoptosis, increased	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5/5	2/5
<u>Spleen:</u>																	
Extramedullary hematopoiesis, increased	1/5	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1/5
Congestion	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5/5	2/5
<u>Mandibular L. N.:</u>																	
Apoptosis, lymphocyte	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1/5	—
<u>Testes:</u>																	
Seminiferous tubular degeneration	—	—	—	—	—	—	—	1/5	—	1/5	—	2/5	—	2/5	—	5/5	—
<u>Epididymus:</u>																	
Oligospermia	—	—	—	—	—	—	—	—	—	1/5	—	1/5	—	2/5	—	5/5	—
<u>Bone Marrow:</u>																	
Myeloid hyperplasia	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1/5
<u>Tail:</u>																	
Acute necrosis with vasculitis, thrombosis, and hemorrhage	—	—	—	—	—	—	—	1/5	4/5	—	—	2/5	1/5	1/5	—	—	1/5
Ulcerative dermatitis	—	—	—	—	—	—	—	1/5	2/5	—	—	1/5	—	1/5	—	—	—

Example 10

HPLC Analytical Method Development and Validation

HPLC will be used in the methods development and validation for the use of organic arsenicals. The HPLC methods will include: standard curve and linearity, reproducibility (10 injections minimum), sensitivity (minimum quantifiable concentration; minimum detectable concentration), accuracy (such as using three independently prepared solutions of 0.025 mg/mL, 0.1 mg/mL, 1 mg/mL), intentional degradation from heat, basic solutions acidic solutions and H₂O₂, and peak definition for intact drug, bulk impurities and starting materials, and degradation products. Bulk raw drug will be analyzed in a reference standard lot through HPLC analysis of purity, loss on drying, optical rotation, melting point, and visual appearance.

Example 11

Dosage Forms Development

The dosage of organic arsenicals will be developed following the formulation solvent system developed by Pharmacology Laboratory. This includes determining the stability in potential aqueous vehicles and to filtration, selecting target concentration for further development, testing the osmolality and pH and adjusting if necessary, selecting package and closure configuration, determining the thermal stability (autoclaving), testing the visual appearance and particulate burden and determining the target pH values and acceptable range for target concentration.

Example 12

Clinical Trials

This example is concerned with the development of human treatment protocols using the arsenical compounds,

MER1, SGLU and SAL-1, and compositions of the invention or the pharmaceutical formulations thereof. These compositions will be of use in the clinical treatment of various cancers including leukemias and other forms of solid cancers and tumors.

The various elements of conducting a clinical trial, including patient treatment and monitoring, will be known to those of skill in the art in light of the present disclosure. The following information is being presented as a general guideline for use in establishing clinical trials using the compositions of the present invention.

Candidates for the phase 1 clinical trial will be patients on which all conventional therapies have failed. Pharmaceutical formulations of MER1, SAL-1 or SGLU-1 will be administered to them intravenously on a tentative schedule of 5 days every 4 weeks. One of skill in the art will appreciate that one may administer the therapeutic formulation of the invention by any alternative route that is suitable depending on the nature of the lesion including administration by any method including local, regional, or systemic administration. Oral and topical applications are also contemplated. A composition of the present invention is typically administered orally or parenterally in dosage unit formulations containing standard, well known non-toxic physiologically acceptable carriers, adjuvants, and vehicles as desired. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intra-arterial injection, or infusion techniques.

To monitor disease course and evaluate the anti-tumor responses, it is contemplated that the patients should be examined for appropriate tumor markers every month. To assess the effectiveness of the drug, the following parameters will be monitored: tumor size and/or bone marrow infiltration of the cancer cells. Tests that will be used to

monitor the progress of the patients and the effectiveness of the treatments may include: physical exam, X-ray, blood work and other clinical laboratory methodologies. The doses given in the phase 1 study will be escalated as is done in standard phase 1 clinical phase trials, i.e. doses will be

escalated until maximal tolerable ranges are reached. Clinical responses may be defined by acceptable measure. For example, a complete response may be defined by complete disappearance of evidence of cancer cells for at least 2 months. Whereas a partial response may be defined by a 50% reduction of cancer cells for at least 2 months.

The clinical trials may be performed with the therapeutic agents of the invention alone or in combination with other anti-cancer drugs and other standard cancer therapies used in the art. The therapeutic compositions of the invention may be delivered to the patient before, after or concurrently with the other anti-cancer agents.

The typical course of treatment will vary depending upon the individual patient and disease being treated in ways known to those of skill in the art. For example, a patient with leukemia might be treated in four week cycles, although longer duration may be used if adverse effects are observed with the patient, and shorter terms of treatment may result if the patient does tolerate the treatment as hoped. Each cycle will consist of 5 individual doses, although this too may be varied depending on the clinical situation. Upon election by the clinician the regimen may be continued with 5 doses every three weeks or on a less frequent basis. Of course, these are only exemplary times for treatment, and the skilled practitioner will readily recognize that many other time-courses are possible.

Patients may, but need not, have received previous chemo-, radio- or gene therapeutic treatments. Optimally the patient will exhibit adequate bone marrow function (defined as peripheral absolute granulocyte count of $>2,000/\text{mm}^3$ and platelet count of $100,000/\text{mm}^3$, adequate liver function (bilirubin 1.5 mg/dl) and adequate renal function (creatinine 1.5 mg/dl).

In one embodiment, administration simply entails injection of the therapeutic composition into the tumor. In another embodiment, a catheter is inserted into the site of the tumor and the cavity may be continuously perfused for a desired period of time.

Of course, the above-described treatment regimes may be altered in accordance with the knowledge gained from preclinical trials. Those of skill in the art will be able to take the information disclosed in this specification and optimize treatment regimes based on the clinical trials described in the specification.

All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

American Conference of Governmental Industrial Hygienists, Inc. (ACGIH). Arsenic and soluble compounds, including arsine. Documentation of the Threshold Limit Values and Biological Exposure Indices, sixth edition, 1991.

Bainbridge, W. S. in *The Cancer Problem*, Macmillan, N.Y., pp. 271-276, 1914.

Banks, C. H. et al., "Biomolecules bearing the S— or SeAsMe₂ function: amino acid and steroid derivatives," *J. Medicinal Chem.* 22:572-575, 1979.

Beliles, R. P. "The Metals," In *Patty's Industrial Hygiene and Toxicology, fourth edition* G. D. Clayton and F. E. Clayton, eds. John Wiley & Sons, Inc.: New York. pp. 1913-1925, 1994.

Chen, G. C. et al., "6-thio- and -seleno-alpha-D-glucose esters of dimethylarsinous acid," *Carb. Res.* 50:53-62, 1976.

Chen, G. C. et al., "Synthesis of 1- and 6-S— and 1- and 6-Se-derivatives of 2-amino-2-deoxy-alpha/beta-D-glucopyranose," *J. Chemical Soc, Perkin Trans.* 1, 2287-2293, 1980.

Cuzick, J. et al., "Medicinal arsenic and internal malignancies," *Br. J. Cancer* 45:904-911, 1982.

Daniel, J. R. and Zingaro, R. A. "Dimethylarsinous acid esters of 1-thio- and -seleno-galactose. A new class of potential carcinostatic agents," *Phosphorus and Sulfur* 4:179-185, 1978.

EP1002537

Forkner, C. and McNair-Scott, T. F. "Arsenic as a therapeutic agent in chronic myeloid leukemia," *JAMA* 97:305, 1931.

Geissler, K. et al., "In vivo effects of arsenic trioxide in refractory acute myeloid leukemia other than acute promyelocytic leukemia," *Blood* 94:4230a, 1999.

Goyer, R. A. "Toxic effects of metals" In *Casarett and Doull's Toxicology: The Basic Science of Poisons, 5th edition*. C. D. Klassen, ed. McGraw-Hill: New York. pp. 691-698, 1996.

Grignani et al., "The acute promyelocytic leukemia-specific PML-RAR alpha fusion protein inhibits differentiation and promotes survival of myeloid precursor cells," *Cell*, 74:423-431, 1993.

Hughes, M. F.; Kenyon, E. M. "Dose-dependent effects on the disposition of monomethylarsonic acid and dimethylarsinic acid in the mouse after intravenous administration" *J. Toxicol. Environ. Health A* 23:53(2) 95-112, 1998.

IALRC. Some metals and metallic compounds. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. Volume 23:39-141, 1980.

Investigational Drug Brochure: Informational Material for Physicians: Melarsoprol (Mel B) (Arsobal). Centers for Disease Control, Atlanta, Ga., 1987.

Investigator's Brochure: Arsenic Trioxide, PolaRx Biopharmaceuticals Inc., New York, N.Y., 1998.

Knock, F. E. et al., "The use of selected sulfhydryl inhibitors in a preferential drug attack on cancer," *Surg. Gynecol. Obstet.* 133:458-466, 1971.

Konig, A. et al., "Comparative activity of melarsoprol and arsenic trioxide in chronic B-cell leukemia lines," *Blood* 90:562-570, 1997.

Material Safety Data Sheet: Dimethylarsinic acid, Strem Chemicals Inc., Newburyport, Mass., 1998.

Rivi, R. et al., "Organic arsenical melarsoprol shows growth suppressive activity via programmed cell death on myeloid and lymphoid leukemia derived cell lines," *Blood (Suppl.)* 88:68a, 1996.

53

Rosenthal, M. V. and Zingaro, R. A. "The synthesis and characterization of thio sugar esters of diorganylarsinous acids," *Phosphorus and Sulfur* 9:107-116, 1980.

Rousselot, P. et al., "Use of arsenic trioxide (As₂O₃) in the treatment of chronic myelogenous leukemia: In vitro and in vivo studies," *Blood* 94:4457a, 1999.

Soignet, S. L. et al., "Clinical study of an organic arsenic melarsoprol, in patients with advanced leukemia," *Cancer Chemother. Pharmacol.* 44:471-421, 1999.

Soignet, S. L. et al., "Dose-ranging and clinical pharmacologic study of arsenic trioxide in patients with advanced hematologic cancers," *Blood* 94:1247a, 1999.

Tarnowski, G. S. et al., "Chemotherapy studies in an animal tumor spectrum: II. Sensitivity of tumors to fourteen antitumor chemicals," *Cancer Res.* 26:181-206, 1966.


Wiernik, P. H. et al., "Phase II trial of arsenic trioxide (As₂O₃) in patients with relapsed/refractory acute myeloid leukemia, blast crisis of CML or myelodysplasia," *Blood* 94:2283a, 1999.

WO9924029

Zhang, P. et al., "Arsenic trioxide treated 72 cases of acute promyelocytic leukemia," *Chin. J. Hematol.* 17:58-62, 1996.

What is claimed is:

1. A compound having anti-cancer activity comprising:



wherein

R₁ and R₂ are independently alkyls with 1-10 carbon atoms;

X is S or Se;

R₃ is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;


R₄ is —OH, —H, or —CH₃;

R₃ and R₄ form an unsubstituted or substituted aromatic ring with C₁ and C₂; and

R₅ is a —OH, or Glycine substituent;

or a pharmaceutically acceptable salt thereof.


2. The compound of claim 1, wherein said compound has the formula:



or a pharmaceutically acceptable salt thereof.


3. The compound of claim 1, wherein said compound has the formula:

54



or a pharmaceutically acceptable salt thereof.

4. A pharmaceutical composition having anti-cancer activity comprising a pharmaceutical carrier and an organic arsenical compound having the formula:



wherein

R₁ and R₂ are independently alkyls with 1-10 carbon atoms;

X is S or Se;

R₃ is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;


R₄ is —OH, —H, or —CH₃;

or R₃ and R₄ form an unsubstituted or substituted aromatic ring with C₁ and C₂; and

R₅ is a —OH, or Glycine substituent;

or a pharmaceutically acceptable salt thereof.

5. A method of treating a patient with solid tumor cancer comprising administering a composition comprising a therapeutically effective amount of a compound having the formula:



wherein

R₁ and R₂ are independently alkyls with 1-10 carbon atoms;

X is S or Se;

R₃ is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;

R₄ is —OH, —H, —CH₃, or a Glutamine substituent;


R₃ and R₄ form an unsubstituted or substituted aromatic ring with C₁ and C₂; and

R₅ is a —OH, or a Glycine substituent;

or a pharmaceutically acceptable salt thereof to said patient.


6. The method of claim 5, wherein said compound has the formula:

55




or a pharmaceutically acceptable salt thereof.

7. The method of claim 5, wherein said compound has the formula:



or a pharmaceutically acceptable salt thereof.


8. The method of claim 5, wherein said compound has the formula:



or a pharmaceutically acceptable salt thereof.

9. The method of claim 5, wherein said solid tumor cancer is brain, lung, liver, spleen, kidney, lymph node, small intestine, pancreas, bone, colon, stomach, breast, endometrium, prostate, testicle, ovary, central nervous system, skin, head and neck, esophagus, or bone marrow cancer.

10. The compound of claim 1, wherein the compound has the formula:



wherein

R_1 , and R_2 are independently alkyls with 1–10 carbon atoms;

X is S or Se;

R_3 is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;

R_4 is —OH, —H, or —CH₃; and

R_5 is a —OH, or Glycine substituent;

or a pharmaceutically acceptable salt thereof.

11. The method of claim 5, wherein said pharmaceutically effective amount is 0.1–1000 mg/kg.

56

12. The method of claim 11, wherein said pharmaceutically effective amount is 1–500 mg/kg.

13. The method of claim 12, wherein said pharmaceutically effective amount is 10–100 mg/kg.

14. The method of claim 5, wherein said compound is administered daily.

15. The method of claim 5, wherein said compound is administered by injection.

16. The method of claim 5, wherein an additional agent is administered to said patient.

17. The method of claim 16, wherein said additional agent is all-trans-retinoic acid, 9-cis retinoic acid, Am-80 or ascorbic acid.

18. The compound of claim 1, wherein R_5 is —OH.

19. The compound of claim 1, wherein R_3 and R_4 form an unsubstituted or substituted aromatic ring with C_1 and C_2 .

20. The compound of claim 1, wherein X is S.

21. The compound of claim 1, wherein R_1 and R_2 are both methyl.

22. The compound of claim 1, wherein R_5 is a Glycine substituent.


23. The pharmaceutical composition of claim 4, wherein R_5 is —OH.

24. The pharmaceutical composition of claim 4, wherein R_3 and R_4 form an unsubstituted or substituted aromatic ring with C_1 and C_2 .

25. The pharmaceutical composition of claim 4, wherein X is S.

26. The pharmaceutical composition of claim 4, wherein R_1 and R_2 are both methyl.

27. A pharmaceutical composition having anti-cancer activity comprising a pharmaceutical carrier and an organic arsenical compound having the formula:



wherein

R_1 and R_2 are independently alkyls with 1–10 carbon atoms;

X is S or Se;

R_3 is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;

R_4 is —OH, —H, —CH₃ or a Glutamine substituent; or R_3 and R_4 form an unsubstituted or substituted aromatic ring with C_1 and C_2 ; and

R_5 is a —OH, or Glycine substituent;


or a pharmaceutically acceptable salt thereof, wherein said pharmaceutical composition has a pH of 5 to 7.

28. The pharmaceutical composition of claim 27, wherein said pharmaceutical composition has a pH of 5.

29. The pharmaceutical composition of claim 27, wherein said pharmaceutical composition has a pH of 7.

30. The pharmaceutical composition of claim 27, wherein said compound has the formula:

57




or a pharmaceutically acceptable salt thereof.

31. The pharmaceutical composition of claim 30, wherein said compound is complexed with pyridine hydrochloride.

32. The method of claim 8, wherein said compound is complexed with pyridine hydrochloride.

33. A method of treating a patient with lymphoma, multiple myeloma, myelodysplasia, myeloproliferative disease, or refractory anemia comprising administering a composition comprising a therapeutically effective amount of a compound having the formula:



wherein

R_1 and R_2 are independently alkyls with 1–10 carbon atoms;

X is S or Se;

R_3 is —H, —COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;


n is 0 or 1;

R_4 is —OH, —H, —CH₃, or a Glutamine substituent; or R_3 and R_4 form an unsubstituted or substituted aromatic ring with C₁ and C₂; and

R_5 is a —OH, or a Glycine substituent;

or a pharmaceutically acceptable salt thereof, to said patient.

34. The method of claim 33, wherein said compound has the formula:




or a pharmaceutically acceptable salt thereof.

58

35. The method of claim 34, wherein said compound is complexed with pyridine hydrochloride.

36. A method of treating a patient with cancer comprising administering a composition comprising a therapeutically effective amount of a compound having the formula:



wherein

R_1 and R_2 are independently alkyls with 1–10 carbon atoms;

X is S or Se;

R_3 is —H, —COOH, —CH₂—COOH, —CH₂—CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;

R_4 is —OH, —H, or —CH₃;

or R_3 and R_4 form an unsubstituted or substituted aromatic ring with C₁ and C₂; and


R_5 is a —OH, or a Glycine substituent;

or a pharmaceutically acceptable salt thereof, to said patient.

37. The method of claim 36, wherein said cancer is acute promyelocytic leukemia.

38. The compound of claim 1, wherein R_3 and R_4 form an unsubstituted or substituted aromatic ring with C₁ and C₂.

39. The method of claim 5, wherein the compound has the formula:



wherein

R_1 and R_2 are independently alkyls with 1–10 carbon atoms;

X is S or Se;

R_3 is —H, —COOH, —CH₂—COOH, —CH₂—COOH, —CH(CH₃)—COOH, —CH(CH₂—CH₃)—COOH, or —CH₂—CH₂—CH₂—COOH;

n is 0 or 1;

R_4 is —OH, —H, or —CH₃; and

R_5 is a —OH, or Glycine substituent;

or a pharmaceutically acceptable salt thereof.

40. The method of claim 5, wherein R_3 and R_4 form an unsubstituted or substituted aromatic ring with C₁ and C₂.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,911,471 B2
DATED : June 28, 2005
INVENTOR(S) : Zingaro et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 53,

Line 47, please insert -- or -- before "R₃ and R₄";

Column 54,

Line 60, please insert -- or -- before "R₃ and R₄";

Column 56,

Line 16, please replace "from" with -- form -- after "R₃ and R₄";

Column 57,

Lines 36-38, please replace "R₃ is -H, -COOH, -CH₂-CH₂-COOH, -CH (CH₃)-COOH, -CH(CH₂-CH₃)-COOH, or -CH₂-CH₂-CH₂-COOH;" with -- R₃ is -H, -COOH, -CH₂-COOH, -CH₂-CH₂-COOH, -CH (CH₃)-COOH, -CH(CH₂-CH₃)-COOH, or -CH₂-CH₂-CH₂-COOH; --.

Column 58,

Line 50, please replace "R₃ is -H, -COOH, -CH₂-COOH, -CH₂-COOH," with -- R₃ is -H, -COOH, -CH₂-COOH, -CH₂-CH₂-COOH, --.

Signed and Sealed this

Thirteenth Day of September, 2005



JON W. DUDAS

Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,911,471 B2
APPLICATION NO. : 10/337969
DATED : June 28, 2005
INVENTOR(S) : Zingaro et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In column 4, line 31, please replace “refractory anemia” with --refractory leukemia--.

In claim 33, column 57, line 20, please replace “refractory anemia” with --refractory leukemia--.

Signed and Sealed this

Nineteenth Day of June, 2007

A handwritten signature in black ink on a light gray dotted background. The signature reads "Jon W. Dudas" in a cursive style.

JON W. DUDAS

Director of the United States Patent and Trademark Office