Spectrum of Higher Derivative 6D Chiral Supergravity on Minkowski $\times S^2$

Y. Pang, C.N. Pope and E. Sezgin

George P. & Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

ABSTRACT

Gauged off-shell Maxwell-Einstein supergravity in six dimensions with N = (1,0) supersymmetry has a higher derivative extension afforded by a supersymmetrized Riemann squared term. This theory admits a supersymmetric Minkowski $\times S^2$ compactification with a U(1) monopole of unit charge on S^2 . We determine the full spectrum of the theory on this background. We also determine the spectrum on a non-supersymmetric version of this compactification in which the monopole charge is different from unity, and we find the peculiar feature that there are massless gravitini in a representation of the S^2 isometry group determined by the monopole charge.

Contents

1	1 Introduction						
2	2 The Theory						
3	Spectrum in Supersymmetric Minkowski $_4 \times S^2$ Background3.1Supersymmetric Minkowski $_4 \times S^2$ background3.2Bosonic Sector3.3Fermionic sector3.4The supermultiplet structure and stability	7 7 8 14 18					
4	Spectrum in Non-Supersymmetric Minkowski $_4 \times S^2$ Background4.1Non-supersymmetric Minkowski $_4 \times S^2$ background4.2Bosonic sector4.3Fermionic sector4.4Remarks on the non-supersymmetric spectrum	 20 20 21 23 27 					
5	Conclusions						
\mathbf{A}	Conventions						
в	Spin-weighted Harmonics on S^2 B.1Scalar spin-weighted harmonicsB.2Vector spin-weighted harmonicsB.3Spin- $\frac{1}{2}$ spin-weighted harmonicsB.4Spin- $\frac{3}{2}$ spin-weighted harmonics	 29 29 31 33 35 					
\mathbf{C}	Spin Projection Operators	35					

1 Introduction

Higher-derivative supergravities are of considerable interest, especially when they arise as lowenergy effective actions of string theories with higher-derivative corrections proportional to powers of the slope parameter α' . However, their construction is notoriously difficult, in part due to the fact that supergravities exist only on-shell in ten dimensions. In view of this difficulty, the compactifications of these theories are rarely studied. In order to gain insights into the compactification of higher-derivative theories, it is instructive to investigate the issue in the simpler situation of lowerdimensional supergravities with higher-derivative terms, postponing for the present the question of how they may arise from ten dimensions. An important technical advantage is that in some lower-dimensional cases, off-shell formulations of the supergravity theories exist. This leads us to consider in particular $\mathcal{N} = (1,0)$ supergravity in six dimensions, which is the highest dimension, and the highest degree of supersymmetry, for which a supergravity with an off-shell formulation is known. The off-shell formulation of this supergravity was constructed in [1, 2], and a higherderivative extension with an off-shell supersymmetrized Riemann-squared term was obtained in [3, 4]. The gauging of the U(1) R-symmetry in the presence of this higher-derivative extension has also recently been obtained [5]. The model has two parameters, namely an overall coefficient M^{-2} in front of the higher-derivative superinvariant in the action, and the gauge-coupling constant q.

In the present paper, we shall study the six-dimensional gauged $\mathcal{N} = (1,0)$ theory with the Riemann-squared term constructed in [5]. In the absence of the curvature-squared terms the model is an (off-shell) version of the Salam-Sezgin theory constructed long ago [6]. It was shown in [6] that the model had the unusual feature of admitting a supersymmetric Minkowski₄ × S^2 vacuum, in which there is a U(1) monopole flux with charge $q = \pm 1$ on the S^2 internal space. A remarkable feature of the theory with the Riemann-squared extension is that the Minkowski₄ × S^2 background continues to be a supersymmetric solution [5]. It also admits non-supersymmetric Minkowski₄ × S^2 backgrounds in which the quantised monopole charge q is larger than 1.

Our focus in this paper is to study the spectrum of the Kaluza-Klein states in the fluctuations around the Minkowski₄ \times S² background. As far as we are aware, such a Kaluza-Klein spectral analysis of a higher-derivative supergravity around a background with non-abelian symmetries has not previously been carried out. Even in the much simpler S^2 reduction of the Salam-Sezgin model discussed in [6], the situation is of considerable interest because of the very unusual feature of obtaining non-abelian symmetries from a sphere reduction, whilst obtaining a Poincaré rather than AdS supergravity in the lower dimension. As expected, the states assemble into $\mathcal{N}=1$ four-dimensional supermultiplets. In the model constructed in [5] with the higher-order Riemannsquared extension, we find a number of novel features associated with the occurrence of higher-order wave operators, and the fact that certain fields that were purely auxiliary prior to the inclusion of the higher-order terms now become dynamical. In particular, we find that certain four-dimensional vector supermultiplets have wave operators that give rise to masses m that are determined by a non-trivial polynomial of fourth order in m^2 . This leads to mass-squared values that are not simply linear in the eigenvalues of the Laplace operators on the internal space, but, rather, involve non-trivial roots of the associated quartic equation. One consequence of this is that the values of m^2 can be negative or even complex, thus implying that there will be instabilities.

The occurrence of such states might at first sight seem surprising in a supersymmetric vacuum. A standard argument for positive semi-definiteness of the energy, first given in [7], uses the fact that if a state $|\psi\rangle$ is annihilated by the supercharge Q, then the superalgebra $\{Q, Q\} \sim P$ implies that $P_0 \geq 0$. However, a crucial ingredient in this argument is that the norm on the states $|\psi\rangle$ is positive definite [8] In our case, the higher-derivative terms in the six-dimensional theory lead to ghost modes in the spectrum, and thus the assumptions required for the positivity result in [7] are violated.

The detailed structure of the quartic polynomial in m^2 for the vector multiplets implies that two of the four roots are always real and positive, while the remaining two can be complex. The conditions under which this occurs are governed by the ratio M^2/g^2 and by the Kaluza-Klein level number ℓ of the harmonics on S^2 . As M^2 becomes larger, the non-positivity and complexity of the two roots sets in at larger and larger values of the level number ℓ . M^2 must at least satisfy $M^2 \geq 8(5 + 2\sqrt{6})g^2$ in order for the roots to be real and positive even at the lowest level $\ell = 0$.

We also study the spectrum of the modes in the non-supersymmetric Minkowski₄ × S^2 vacua that arise for S^2 monopole charges q greater than 1. An interesting feature in these cases is that the spectrum includes an SU(2) multiplet of massless spin- $\frac{3}{2}$ fields at level $\ell = \frac{1}{2}(|q| - 3)$.

The organisation of the paper is as follows. In section 2 we review the six-dimensional gauged $\mathcal{N} = (1,0)$ off-shell $R + |\text{Riem}|^2$ supergravity that was recently constructed in [5]. In section 3 we study the complete linearised spectrum of Kaluza-Klein modes in the supersymmetric Minkowski₄ × S^2 vacuum, which has a monopole charge q = 1 on the S^2 internal space, and exists for any value of the coupling M^{-2} of the Riemann-squared invariant. In section 4 we repeat the analysis for the non-supersymmetric Minkowski₄ × S^2 vacua, which have arbitrary integer monopole charge $|q| \geq 2$, and which exist only for a special value of the ratio g^2/M^2 . For this analysis we need many results on the properties of spin-weighted spherical harmonics on S^2 , since these are needed for the expansions in the monopole background of the fermion fields and certain vector fields that carry charges. We present a detailed discussion of these harmonics in appendix B. In appendix A we give our spinor conventions, and in appendix C we summarise some results for spin projection operators in four dimensions.

2 The Theory

The off-shell 6D(1,0) supergravity multiplet consists of the fields [1]

$$(e_{\mu}{}^{a}, V_{\mu}^{\prime i j}, V_{\mu}, B_{\mu\nu}, L, C_{\mu\nu\rho\sigma}, \psi_{\mu}^{i}, \chi^{i})$$
 (2.1)

where V_{μ}^{ij} is symmetric and traceless in its Sp(1) doublet indices, B and C are antisymmetric tensor fields, L is a real scalar, and the spinors are symplectic Majorana-Weyl. The above fields have (15, 12, 5, 10, 1, 5, 40, 8) degrees of freedom. In addition, we shall consider the off-shell Maxwell multiplet consisting of the fields

$$\left(A_{\mu}, Y^{ij}, \lambda^{i}\right) , \qquad (2.2)$$

where Y^{ij} is symmetric in its indices and the fermion is symplectic Majorana Weyl. These fields have (5,3,8) degrees of freedom.

The total Lagrangian we shall study is given by

$$\mathcal{L} = \mathcal{L}_R - \frac{1}{8M^2} \mathcal{L}_{R^2} , \qquad (2.3)$$

where the $U(1)_R$ gauged off-shell supergravity Lagrangian, up to quartic fermion terms, is $[1, 5]^1$

$$e^{-1}\mathcal{L}_{R} = \frac{1}{2}LR + \frac{1}{2}L^{-1}\partial_{\mu}L\partial^{\mu}L + 2\sqrt{2}gL\delta^{ij}Y_{ij} - \frac{1}{24}LH_{\mu\nu\rho}H^{\mu\nu\rho} + LV_{\mu}^{\prime ij}V^{\prime \mu}{}_{ij} - \frac{1}{4}L^{-1}E^{\mu}E_{\mu} + \frac{1}{\sqrt{2}}E^{\mu}\left(V_{\mu} + 2gA_{\mu}\right) + Y^{ij}Y_{ij} - \frac{1}{8}F_{\mu\nu}F^{\mu\nu} - \frac{1}{16}\varepsilon^{\mu\nu\rho\sigma\lambda\tau}B_{\mu\nu}F_{\rho\sigma}F_{\lambda\tau} - \frac{1}{2}L\bar{\psi}_{\rho}\gamma^{\mu\nu\rho}D_{\mu}\psi_{\nu} - \sqrt{2}\bar{\chi}_{i}\gamma^{\mu\nu}D_{\mu}\psi_{\nu j}\delta^{ij} + L^{-1}\bar{\chi}D\chi - \frac{1}{2}\bar{\psi}^{\mu}\gamma^{\nu}\psi_{\nu}\partial_{\mu}L - \frac{1}{\sqrt{2}}\delta_{ij}\bar{\psi}_{\nu}^{i}\gamma^{\mu}\gamma^{\nu}\chi^{j}L^{-1}\partial_{\mu}L - 2\sqrt{2}gL\bar{\lambda}_{i}\gamma^{\mu}\psi_{\mu j}\delta^{ij} + 2g\bar{\lambda}\chi + \frac{1}{2}V_{\mu}^{\prime ij}\left(2\sqrt{2}\bar{\chi}^{k}\psi_{i}^{\mu}\delta_{jk} - 3L^{-1}\bar{\chi}_{i}\gamma^{\mu}\chi_{j}\right) - \frac{1}{48}LH_{\mu\nu\rho}\left(\bar{\psi}^{\lambda}\gamma_{[\lambda}\gamma^{\mu\nu\rho}\gamma_{\tau]}\psi^{\tau} + 2\sqrt{2}L^{-1}\bar{\psi}_{\lambda}\gamma^{\lambda\mu\nu\rho}\chi_{j}\delta^{ij} - 2L^{-2}\bar{\chi}\gamma^{\mu\nu\rho}\chi\right) - \frac{1}{4\sqrt{2}}E_{\rho}\left(\psi_{\mu}^{i}\gamma^{\rho\mu\nu}\psi_{\nu}^{j}\delta_{ij} - 2\sqrt{2}L^{-1}\bar{\psi}_{\sigma}\gamma^{\rho}\gamma^{\sigma}\chi + 2L^{-2}\bar{\chi}_{i}\gamma^{\rho}\chi_{j}\delta^{ij}\right) - 2\bar{\lambda}D\lambda + \frac{1}{12}H_{\mu\nu\rho}\bar{\lambda}\gamma^{\mu\nu\rho}\lambda + \frac{1}{2\sqrt{2}}F_{\mu\nu}\bar{\lambda}\gamma^{\rho}\gamma^{\mu\nu}\psi_{\rho},$$
(2.4)

where $H_{\mu\nu\rho} = 3\partial_{[\mu}B_{\nu\rho]}$ and

$$E^{\mu} = \frac{1}{24} \varepsilon^{\mu\nu_1 \cdots \nu_5} \partial_{[\nu_1} C_{\nu_2 \cdots \nu_5]} .$$
 (2.5)

$$D_{\mu}\psi_{\nu}^{i} = (\partial_{\mu} + \frac{1}{4}\omega_{\mu}{}^{ab}\gamma_{ab})\psi_{\nu}^{i} - \frac{1}{2}V_{\mu}\delta^{ij}\psi_{\nu j} , \qquad (2.6)$$

$$D_{\mu}\chi^{i} = (\partial_{\mu} + \frac{1}{4}\omega_{\mu}{}^{ab}\gamma_{ab})\chi^{i} - \frac{1}{2}V_{\mu}\delta^{ij}\chi_{j} + V_{\mu}{}^{'i}{}_{j}\chi^{j} .$$
(2.7)

Note the presence of arbitrary coupling constant in \mathcal{L}_R . In fact, the sum of all the terms in this Lagrangian that depend on g separately have the off-shell supersymmetry. Thus, the total Lagrangian is a sum of three separately off-shell supersymmetric pieces.

The Lagrangian for the supersymmetrized Riemann squared term, up to quartic fermion terms, is given by [3, 4]

$$e^{-1}\mathcal{L}_{\mathbf{R}^{2}} = R_{\mu\nu}{}^{ab}(\omega_{-})R^{\mu\nu}{}_{ab}(\omega_{-}) - 2G^{ab}G_{ab} - 4G'^{ij}_{\mu\nu}G'^{ij\nu}, + \frac{1}{4}\varepsilon^{\mu\nu\rho\sigma\lambda\tau}B_{\mu\nu}R_{\rho\sigma ab}(\omega_{-})R_{\lambda\tau ab}(\omega_{-}) + 2\bar{\psi}^{ab}(\omega_{+})\gamma^{\mu}D_{\mu}(\omega,\omega_{-})\psi_{ab}(\omega_{+}) - R_{\nu\rho}{}^{ab}(\omega_{-})\bar{\psi}_{ab}(\omega_{+})\gamma^{\mu}\gamma^{\nu\rho}\psi_{\mu} - 8G^{ij}_{\mu\nu}\left(\bar{\psi}^{\mu}_{i}\gamma_{\lambda}\psi^{\lambda\nu}_{j}(\omega_{+}) + \frac{1}{6}\bar{\psi}^{\mu}_{i}\gamma\cdot H\psi^{\nu}_{j}\right) - \frac{1}{12}\bar{\psi}^{ab}(\omega_{+})\gamma\cdot H\psi_{ab}(\omega_{+}) - \frac{1}{2}\Big[D_{\mu}(\omega_{-},\Gamma_{+})R^{\mu\nu ab}(\omega_{-}) - 2H_{\mu\nu}{}^{\rho}R^{\mu\nu ab}(\omega_{-})\Big]\bar{\psi}^{a}\gamma_{\rho}\psi_{b}, \qquad (2.8)$$

¹We have let $g \to 4g$ and $A_{\mu} \to A_{\mu}/\sqrt{2}$ in the results of [5].

where $G_{\mu\nu}^{'ij}$ and $G_{\mu\nu}$ are the field strengths associated with $V_{\mu}^{'ij}$ and V_{μ} , which can be combined as $V_{\mu}^{ij} = V_{\mu}^{'ij} + \frac{1}{2}\delta^{ij}V_{\mu}$. Furthermore $\psi_{\mu\nu}(\omega_{+}) = 2D_{[\mu}(\omega_{+})\psi_{\nu]}$ and

$$D_{\mu}(\omega,\omega_{-})\psi^{abi} = (\partial_{\mu} + \frac{1}{4}\omega_{\mu}{}^{cd}\gamma_{cd})\psi^{abi} + 2\omega_{\mu-}{}^{c[a}\psi^{b]i}{}_{c} + V^{i}_{\mu j}\psi^{abi} ,$$

$$\omega_{\mu\pm}{}^{ab} = \omega_{\mu}{}^{ab} \pm \frac{1}{2}H_{\mu}{}^{ab} , \qquad \Gamma^{\rho}_{\mu\nu\pm} = \Gamma^{\rho}_{\mu\nu} \pm \frac{1}{2}H_{\mu}{}^{\nu\rho} .$$
(2.9)

The off-shell resulting supersymmetry transformations of the Poincaré multiplet, up to cubic fermion terms, are [1, 3, 5]

$$\begin{split} \delta e_{\mu}{}^{a} &= \frac{1}{2} \bar{\epsilon} \gamma^{a} \psi_{\mu} ,\\ \delta \psi_{\mu}{}^{i} &= (\partial_{\mu} + \frac{1}{4} \omega_{\mu a b} \gamma^{a b}) \epsilon^{i} + V_{\mu}{}^{i}{}_{j} \epsilon^{j} + \frac{1}{8} H_{\mu \nu \rho} \gamma^{\nu \rho} \epsilon^{i} ,\\ \delta B_{\mu \nu} &= -\bar{\epsilon} \gamma_{[\mu} \psi_{\nu]} ,\\ \delta \chi^{i} &= \frac{1}{2\sqrt{2}} \gamma^{\mu} \delta^{i j} \partial_{\mu} L \epsilon_{j} - \frac{1}{4} \gamma^{\mu} E_{\mu} \epsilon^{i} + \frac{1}{\sqrt{2}} \gamma^{\mu} V'_{\mu k}^{(i} \delta^{j) k} L \epsilon_{j} - \frac{1}{12\sqrt{2}} L \delta^{i j} \gamma \cdot H \epsilon_{j} ,\\ \delta L &= \frac{1}{\sqrt{2}} \bar{\epsilon}^{i} \chi^{j} \delta_{i j} ,\\ \delta C_{\mu \nu \rho \sigma} &= L \bar{\epsilon}^{i} \gamma_{[\mu \nu \rho} \psi_{\sigma]}^{j} \delta_{i j} - \frac{1}{2\sqrt{2}} \bar{\epsilon} \gamma_{\mu \nu \rho \sigma} \chi ,\\ \delta V_{\mu}{}^{i j} &= \frac{1}{2} \bar{\epsilon}^{(i} \gamma^{\rho} \psi_{\mu \rho}^{j)} + \frac{1}{12} \bar{\epsilon}^{(i} \gamma \cdot H \psi_{\mu}^{j)} + \frac{1}{8} \sigma^{-1} \bar{\epsilon}^{(i} \gamma^{\rho} \left(H_{[\mu}{}^{a b} \gamma_{a b} \psi_{\rho]}^{j)} \right) \end{split}$$
(2.10)

and the off-shell supersymmetry transformations of the vector multiplet are

$$\begin{split} \delta A_{\mu} &= -\bar{\epsilon}\gamma_{\mu}\lambda ,\\ \delta \lambda^{i} &= \frac{1}{8\sqrt{2}}\gamma^{\mu\nu}F_{\mu\nu}\epsilon^{i} - \frac{1}{2}Y^{ij}\epsilon_{j} ,\\ \delta Y^{ij} &= -\bar{\epsilon}^{(i}\gamma^{\mu}D_{\mu}\lambda^{j)} + \frac{1}{8}\bar{\epsilon}^{(i}\gamma^{\mu}\gamma \cdot H\psi^{j)}_{\mu} - \frac{1}{24}\bar{\lambda}^{i}\gamma \cdot H\lambda^{j} - \frac{1}{2}Y^{k(i}\bar{\epsilon}^{j)}\gamma^{\mu}\psi_{\mu k} . \end{split}$$
(2.11)

Of the auxiliary fields of the Poincaré supergravity, $V_{\mu}^{'ij}$ and V_{μ} can no longer be eliminated algebraically due to the presence of the Riemann squared invariant but Y^{ij} and $C_{\mu\nu\rho\sigma}$ can still be eliminated by means of their field equations as

$$Y^{ij} = -\sqrt{2}gL\delta^{ij} , \qquad E_{\mu} = \sqrt{2}L\left(V_{\mu} + 2gA_{\mu}\right) .$$
 (2.12)

The total Lagrangian we shall study here is given by

$$\mathcal{L} = L_{\rm R} - \frac{1}{8M^2} \mathcal{L}_{\rm R^2} , \qquad (2.13)$$

where M is an arbitrary mass parameter.

3 Spectrum in Supersymmetric Minkowski₄ \times S² Background

3.1 Supersymmetric Minkowski₄ \times S² background

We shall study the compactification on the one half supersymmetric vacuum solution with the geometry of Minkowski₄ × S². From here on, the 6D coordinates will be denoted by x^M and they will be split as (x^{μ}, y^m) to denote the coordinates of 4D spacetime and the internal two-dimensional space. The supersymmetric Minkowski₄ × S² vacuum solution given by [5]

$$\bar{R}_{\mu\nu\lambda\rho} = 0 , \qquad \bar{R}_{mn} = \alpha^2 \bar{g}_{mn} , \qquad \bar{L} = 1 ,
\bar{F}_{\mu\nu} = 0 , \qquad \bar{F}_{mn} = 4g\epsilon_{mn} ,
\bar{G}_{\mu\nu} = 0 , \qquad \bar{G}_{mn} = -\alpha^2 \epsilon_{mn} ,$$
(3.1)

where $\alpha^2 \equiv 8g^2$, \bar{g}_{mn} is the metric on S^2 with radius $1/\alpha$, and ϵ_{mn} is the Levi-Civita tensor on the same S^2 . We define the complex vectors

$$\hat{Z}_M = \hat{V}_M^{\prime 11} + i \hat{V}_M^{\prime 12} , \qquad (3.2)$$

and parametrize the linearized fluctuations around above background as follows

$$\hat{g}_{MN} = \bar{g}_{MN} + \hat{h}_{MN} , \qquad \hat{L} = 1 + \hat{\phi} , \qquad \hat{A}_M = \bar{A}_M + \hat{a}_M ,
\hat{V}_M = \bar{V}_M + \hat{v}_M , \qquad \hat{Z}_M = \hat{z}_M , \qquad \hat{B}_{MN} = \hat{b}_{MN} ,$$
(3.3)

where we use "hat" to stand for six dimensional quantities and "bar" to denote quantities evaluated in the vacuum background. In the background specified above, the linearized six dimensional bosonic and fermionic gauge symmetries are expressed as²

$$\begin{split} \delta \hat{h}_{MN} &= \bar{\nabla}_M \hat{\xi}_N + \bar{\nabla}_N \hat{\xi}_M , \qquad \delta \hat{a}_M = \hat{\xi}^N \bar{F}_{NM} + \partial_M \hat{\Lambda}, \\ \delta \hat{v}_M &= \hat{\xi}^N \bar{G}_{NM} - 2g \partial_M \hat{\Lambda} , \qquad \delta \hat{b}_{MN} = \partial_M \hat{\Lambda}_N - \partial_N \hat{\Lambda}_M , \\ \delta \hat{\psi}_M &= \bar{D}_M \hat{\epsilon}, \qquad \delta \hat{\lambda} = \frac{1}{16} \bar{\Gamma}^{MN} \bar{F}_{MN} \hat{\epsilon} + \frac{i}{2} g \hat{\epsilon} , \qquad \delta \hat{\chi} = 0 . \end{split}$$
(3.4)

This background preserves half supersymmetry because it admits a Killing spinor $\hat{\eta}$ which has the following properties

$$\delta\hat{\psi}_M = \bar{D}_M\hat{\eta} = 0, \qquad \delta\hat{\chi} = 0, \qquad \delta\hat{\lambda} = \left(\frac{1}{16}\bar{\Gamma}^{MN}\bar{F}_{MN} + \frac{i}{2}g\right)\hat{\eta} = 0 , \qquad (3.5)$$

and by choosing the six dimensional gamma matrices as in Appendix, it can be shown that

$$\hat{\eta} = \epsilon \otimes \eta, \qquad \eta = \begin{pmatrix} 0\\ 1 \end{pmatrix},$$
(3.6)

where ϵ is a constant four dimensional Weyl spinor with appropriate chirality inherited from six dimensions.

²For later convenience, starting from the USp(2) symplectic-Majorana-Weyl spinors we have defined Weyl spinors by complexifying as $\psi = \psi_1 + i\psi_2$ and rescaled $\hat{\chi}$ and $\hat{\lambda}$ used in [5] by $\hat{\chi} \to \sqrt{2}\hat{\chi}$, $\hat{\lambda} \to \sqrt{2}\hat{\lambda}$.

3.2 Bosonic Sector

In this section, we shall drop the "bar" on the covariant derivatives for simplicity in notation. The linearized bosonic field equations are given as follows

$$(\hat{R}_{MN}^{(L)} + \hat{\phi}\bar{R}_{MN}) = \hat{\nabla}_{M}\hat{\nabla}_{N}\hat{\phi} + \alpha^{2}\bar{g}_{MN}\hat{\phi} + \frac{1}{2}(\hat{F}_{MP}^{(L)}\bar{F}_{N}^{P} + \hat{F}_{NP}^{(L)}\bar{F}_{M}^{P} - \bar{F}_{MP}\bar{F}_{NQ}\hat{h}^{PQ})$$
(3.7)

$$+ \frac{1}{2}\hat{h}_{MN}(\alpha^{2} - \frac{1}{4}\bar{F}^{PQ}\bar{F}_{PQ}) - \frac{1}{4}\bar{g}_{MN}(\hat{F}_{PQ}^{(L)}\bar{F}^{PQ} - \bar{F}_{P}^{Q}\bar{F}^{PT}\hat{h}_{QT}),$$
$$- \frac{1}{8M^{2}}S_{MN}^{(L)} ,$$
(3.8)

$$\hat{R}^{(L)} = 2\alpha^{2}\hat{\phi} + 2\hat{\Box}\hat{\phi} ,$$
(3.8)

$$\hat{\nabla}^{P}\hat{H}_{PMN}^{(L)} = \frac{1}{2}\varepsilon_{MN}^{PQST}(\frac{1}{2}\hat{F}_{PQ}^{(L)}\bar{F}_{ST} - \frac{1}{2M^{2}}\tilde{R}^{(L)J}_{KPQ}\bar{R}_{JST}^{K}) + \frac{1}{M^{2}}\hat{\nabla}^{P}\hat{\Box}\hat{H}_{PMN}^{(L)},$$
$$+ 3M^{2}\hat{\nabla}^{P}(\hat{H}_{+}^{(L)ST}\bar{F}_{R}\bar{R}_{MN}]_{ST}) ,$$
(3.9)

$$0 = \hat{\nabla}^{P} \hat{F}_{PM}^{(L)} - \hat{\nabla}^{P} \hat{h}_{PQ} \bar{F}_{M}^{Q} - \hat{\nabla}_{P} \hat{h}_{QM} \bar{F}^{PQ} + \frac{1}{2} \hat{\nabla}_{P} \hat{h} \bar{F}_{M}^{P} + 4g(\hat{v}_{M} + 2g\hat{a}_{M}) - \frac{1}{2} * \hat{H}_{MPQ}^{(L)} \bar{F}^{PQ} ,$$

$$(3.10)$$

$$0 = \hat{\nabla}^{P} \hat{G}_{PM}^{(L)} - \hat{\nabla}^{P} \hat{h}_{PQ} \bar{G}_{M}^{Q} - \hat{\nabla}_{P} \hat{h}_{QM} \bar{G}^{PQ} + \frac{1}{2} \hat{\nabla}_{P} \hat{h} \bar{G}_{M}^{P} - M^{2} (\hat{v}_{M} + 2g\hat{a}_{M}) , \qquad (3.11)$$

$$0 = \left(\hat{\nabla}^{P} - i\bar{V}^{P}\right)\hat{G}_{PM}^{'(L)} - i\bar{G}_{MN}\hat{z}^{N} - M^{2}\hat{z}_{M} , \qquad (3.12)$$

where

$$\hat{G}_{MN}^{(L)} = 2\hat{D}_{[M}\hat{z}_{N]}, \qquad \hat{D}_{M}\hat{z}_{N} \equiv (\hat{\nabla}_{M} - i\bar{V}_{M})\hat{z}_{N},
R^{(L)P}_{MNQ} = \hat{R}^{(L)P}_{MNQ} - \hat{\nabla}_{[N}H^{(L)P}_{Q] M},
S^{(L)}_{MN} = 8\left(\hat{G}^{(L)}_{MP}\bar{G}_{N}^{P} + \hat{G}^{(L)}_{NP}\bar{G}_{M}^{P} - \bar{G}_{M}^{P}\bar{G}_{N}^{Q}\hat{h}_{PQ}\right) - 4\bar{g}_{MN}(\hat{G}^{(L)}_{PQ}\bar{G}^{PQ} - \bar{G}_{P}^{Q}\bar{G}^{PT}\hat{h}_{QT})
+ 4(\tilde{R}^{(L)S}_{QMP}\bar{R}^{Q}_{SN}^{P} + \bar{R}^{S}_{QM}^{P}\tilde{R}^{(L)Q}_{SNP} - \hat{h}^{PQ}\bar{R}^{S}_{TMP}\bar{R}^{T}_{SNQ}) - 2\hat{h}_{MN}\bar{G}^{PQ}\bar{G}_{PQ}
+ \hat{h}_{MN}\bar{R}^{PQST}\bar{R}_{PQST} - 2\bar{g}_{MN}(\tilde{R}^{(L)S}_{TPQ}\bar{R}^{T}_{S}^{PQ} + \bar{R}_{JKSP}\bar{R}^{JKS}_{Q}\hat{h}^{PQ})
+ 8(\hat{\nabla}^{P}\tilde{\nabla}^{Q}\tilde{R}_{P(MN)Q})^{(L)} + 8\hat{\nabla}^{S}(\bar{R}_{S(M}^{PQ}\hat{H}^{+(L)}_{N)PQ}), \qquad (3.13)$$

and the penultimate term takes the form

$$(\hat{\nabla}^{P} \tilde{\nabla}^{Q} \tilde{R}_{P(MN)Q})^{(L)} = \bar{R}^{P}_{(MN)} (\bar{R}^{S}_{P} \hat{h}_{SQ} - \bar{R}^{S}_{P} {}^{T}_{Q} \hat{h}_{ST} - \frac{1}{2} \hat{\Box} \hat{h}_{PQ}) + \hat{\nabla}_{P} \hat{\nabla}_{Q} \hat{h}_{S(M} \bar{R}^{P}_{N)} {}^{QS} - \frac{1}{2} (\hat{\nabla}^{P} \hat{\nabla}_{(M} \hat{h}^{QS} \bar{R}_{N)SPQ} + \hat{\nabla}^{P} \hat{\nabla}^{S} \hat{h}^{Q}_{(M} \bar{R}_{N)SPQ} - \frac{1}{2} \bar{R}^{PQS}_{T} \hat{h}^{T}_{(M} \bar{R}_{N)SPQ} + \frac{1}{2} \bar{R}^{PQ}_{T(M} \bar{R}_{N)SPQ} \hat{h}^{ST}) + \frac{1}{2} \bar{R}_{PMNQ} \hat{\nabla}^{P} \hat{\nabla}^{Q} \hat{h} + \hat{\nabla}_{P} \hat{\nabla}^{Q} \tilde{R}^{(L)P}_{(MN)Q} - \frac{1}{2} (\hat{\nabla}_{P} \hat{H}^{(L)}_{QS(M} \bar{R}^{PQS}_{N)} + \hat{\nabla}_{P} \hat{H}^{(L)}_{QS(M} \bar{R}^{PQS}_{N)}).$$

$$(3.14)$$

The covariant derivative $\tilde{\nabla}_M$ is defined with respect to the connection $\tilde{\Gamma}^{\rho}_{\mu\nu}$ containing bosonic torsion as

$$\widetilde{\Gamma}^{\rho}_{\mu\nu} = \left\{ \begin{array}{c} \rho \\ \mu\nu \end{array} \right\} + \frac{1}{2} H_{\mu\nu}^{\rho} .$$
(3.15)

Note that we are using $\hat{G}_{MN}^{\prime(L)}$ to denote the covariant field strength of the complex vector field \hat{z}_M , and $G_{MN}^{(L)}$ to denote the field strength of the real vector v_M . There are no transverse traceless spin-2 harmonics on S^2 , and the transverse spin-1 harmonics

are related to spin-0 harmonics by

$$Y_m^{(\ell)} = \epsilon_m {}^n \nabla_n Y^{(\ell)} .$$
(3.16)

We can expand the six-dimensional bosonic fields in terms of S^2 harmonics as follows

$$\begin{aligned}
\hat{h}_{\mu\nu} &= \sum_{\ell \ge 0} h_{\mu\nu}^{(\ell)} Y^{(\ell)} , \\
\hat{h}_{mn} &= \sum_{\ell \ge 2} \left(L^{(\ell)} \nabla_{\{m} Y_{n\}}^{(\ell)} + \tilde{L}^{(\ell)} \nabla_{\{m} \nabla_{n\}} Y^{(\ell)} \right) + \bar{g}_{mn} \sum_{\ell \ge 0} N^{(\ell)} Y^{(\ell)} , \\
\hat{h}_{\mu m} &= \sum_{\ell \ge 1} (k_{\mu}^{(\ell)} Y_{m}^{(\ell)} + \tilde{k}_{\mu}^{(\ell)} \nabla_{m} Y^{(\ell)}) , \\
\hat{\phi} &= \sum_{\ell \ge 0} \phi^{(\ell)} Y^{(\ell)} , \\
\hat{a}_{\mu} &= \sum_{\ell \ge 0} a_{\mu}^{(\ell)} Y^{(\ell)} , \quad \hat{a}_{m} = \sum_{\ell \ge 1} \left(a^{(\ell)} Y_{m}^{(\ell)} + \tilde{a}^{(\ell)} \nabla_{m} Y^{(\ell)} \right) , \\
\hat{v}_{\mu} &= \sum_{\ell \ge 0} v_{\mu}^{(\ell)} Y^{(\ell)} , \quad \hat{v}_{m} = \sum_{\ell \ge 1} \left(v^{(\ell)} Y_{m}^{(\ell)} + \tilde{v}^{(\ell)} \nabla_{m} Y^{(\ell)} \right) , \\
\hat{z}_{\mu} &= \sum_{\ell \ge 0} z_{\mu}^{(\ell)} - 1 Y_{m}^{(\ell)} + \sum_{\ell \ge 2} \left(z^{(\ell)} D_{m-1} Y^{(\ell)} + i \tilde{z}^{(\ell)} \epsilon_{m}^{n} D_{n-1} Y^{(\ell)} \right) , \\
\hat{b}_{\mu\nu} &= \sum_{\ell \ge 0} b_{\mu\nu}^{(\ell)} Y^{(\ell)} , \quad \hat{b}_{mn} = \epsilon_{mn} \sum_{\ell \ge 0} b^{(\ell)} Y^{(\ell)} , \\
\hat{b}_{\mu m} &= \sum_{\ell \ge 0} \left(b_{\mu}^{(\ell)} Y_{m}^{(\ell)} + \tilde{b}_{\mu}^{(\ell)} \nabla_{m} Y^{(\ell)} \right) ,
\end{aligned}$$
(3.17)

where the notation $\{mn\}$ means "symmetric and traceless," and in the \hat{z}_m expansion $_{-1}V_m^{(0)}$ and $_{-1}V_m^{(1)}$ are level $\ell = 0$ and $\ell = 1$ complex anti-self dual vector harmonics with charge -1 on the 2-sphere, whose explicit forms are given in Appendix B.2. D_m is the U(1) covariant derivative on the 2-sphere, and $_{-1}Y^{(\ell)}$ are the charged harmonics which are described in some detail in Appendix B.1. Furthermore, the scalar harmonics $Y^{(\ell)}$ employed above satisfy

$$\Box_2 Y^{(\ell)} = -\alpha^2 c_{\ell} Y^{(\ell)} , \qquad (3.18)$$

where \Box_2 is the d'Alembertian on S^2 with radius $1/\alpha$ and

$$c_{\ell} \equiv \ell(\ell+1) , \qquad \alpha^2 \equiv 8g^2 . \tag{3.19}$$

We have also used the spin-1 harmonics $Y_m^{(\ell)}$ which satisfy the relations

$$\Box_2 Y_n^{(\ell)} = -(c_\ell - 1)\alpha^2 Y_n^{(\ell)} , \qquad \epsilon^{mn} \nabla_m Y_n^{(\ell)} = \alpha^2 c_\ell Y^{(\ell)} . \qquad (3.20)$$

Utilizing the six dimensional gauge symmetries (3.4), we impose the following gauge condition on the linearized fields [9]

$$\hat{\nabla}^{m} \hat{h}_{\{mn\}} = 0, \qquad \hat{\nabla}^{m} \hat{h}_{m\mu} = 0,
\hat{\nabla}^{m} \hat{a}_{m} = 0, \qquad \hat{\nabla}^{m} \hat{b}_{mM} = 0.$$
(3.21)

Upon the use of these gauge conditions, the harmonic expansions (3.17) simplify to

$$\hat{h}_{\mu\nu} = \sum_{\ell \ge 0} h_{\mu\nu}^{(\ell)} Y^{(\ell)} , \qquad \hat{h}_{\mu m} = \sum_{\ell \ge 1} k_{\mu}^{(\ell)} Y_{m}^{(\ell)} ,
\hat{h}_{mn} = \bar{g}_{mn} \sum_{\ell \ge 0} N^{(\ell)} Y^{(\ell)} , \qquad \hat{\phi} = \sum_{\ell \ge 0} \phi^{(\ell)} Y^{(\ell)} ,
\hat{a}_{\mu} = \sum_{\ell \ge 0} a_{\mu}^{(\ell)} Y^{(\ell)} , \qquad \hat{a}_{m} = \sum_{\ell \ge 1} a^{(\ell)} Y_{m}^{(\ell)} ,
\hat{v}_{\mu} = \sum_{\ell \ge 0} v_{\mu}^{(\ell)} Y^{(\ell)} , \qquad \hat{v}_{m} = \sum_{\ell \ge 1} \left(v^{(\ell)} Y_{m}^{(\ell)} + \tilde{v}^{(\ell)} \nabla_{m} Y^{(\ell)} \right) ,
\hat{z}_{\mu} = \sum_{\ell \ge 0} z_{\mu}^{(\ell)} - 1 Y^{(\ell)} ,
\hat{z}_{m} = \sum_{\ell \ge 0} z_{\mu}^{(\ell)} - 1 Y_{m}^{(\ell)} + \sum_{\ell \ge 2} \left(z^{(\ell)} D_{m-1} Y^{(\ell)} + i \tilde{z}^{(\ell)} \epsilon_{m}^{\ n} D_{n-1} Y^{(\ell)} \right) ,
\hat{b}_{\mu\nu} = \sum_{\ell \ge 0} b_{\mu\nu}^{(\ell)} Y^{(\ell)} , \qquad \hat{b}_{\mu m} = \sum_{\ell \ge 1} b_{\mu}^{(\ell)} Y_{m}^{(\ell)} , \qquad \hat{b}_{mn} = \epsilon_{mn} b^{(0)} Y^{(0)} .$$
(3.22)

The de Donder-Lorentz gauge (3.21) does not fix all the gauge symmetries, and consequently there are some residual ones generated by harmonic zero modes, S^2 Killing vector $Y_m^{(1)}$ and conformal Killing vectors $\nabla_m Y^{(1)}$. Specifically, these residual gauge symmetries are:

• The four dimensional coordinate transformation generated by $\hat{\xi}_{\mu} = \xi^{(0)}_{\mu} Y^{(0)}$

$$\delta h_{\mu\nu}^{(0)} = \partial_{\mu}\xi_{\nu}^{(0)} + \partial_{\nu}\xi_{\mu}^{(0)} . \qquad (3.23)$$

• The Stueckelberg shift symmetries generated by $\hat{\xi}_m = \xi^{(1)} \nabla_m Y^{(1)}$

$$\delta h^{(1)}_{\mu\nu} = -\partial_{\mu}\partial_{\nu}\xi^{(1)}, \qquad \delta N^{(1)} = -2\xi^{(1)}
\delta a^{(1)} = 4g\xi^{(1)}, \qquad \delta v^{(1)} = -\alpha^{2}\xi^{(1)}.$$
(3.24)

• Linearized SU(2) symmetry generated by $\hat{\xi}_m = \xi'^{(1)} Y_m^{(1)}$ and $\hat{\Lambda} = -4g\xi'^{(1)}Y^{(1)}$

$$\delta k_{\mu}^{(1)} = \partial_{\mu} \xi^{\prime(1)}$$
 (3.25)

• Four dimensional $U(1)_{\rm R}$ symmetry generated by $\hat{\Lambda} = \Lambda^{(0)} Y^{(0)}$

$$\delta a^{(0)}_{\mu} = \partial_{\mu} \Lambda^{(0)} . \qquad (3.26)$$

• Abelian 2-form symmetry generated by $\hat{\Lambda}_{\mu} = \Lambda_{\mu}^{(0)} Y^{(0)}$

$$\delta b^{(0)}_{\mu\nu} = \partial_{\mu} \Lambda^{(0)}_{\nu} - \partial_{\nu} \Lambda^{(0)}_{\mu} . \qquad (3.27)$$

We shall take into account these symmetries in the analysis of the spectrum below, where we treat the spin-2, spin-1 and spin-0 sectors separately. In doing so we shall encounter the following wave operators

$$\mathcal{O}_{1} \equiv \hat{\Box}_{0} + \alpha^{2} - M^{2} ,$$

$$\mathcal{O}_{2} \equiv \hat{\Box}_{0}^{2} - M^{2}\hat{\Box}_{0} - \alpha^{4}c_{\ell} ,$$

$$\mathcal{O}_{4} \equiv \hat{\Box}_{0}^{4} + (\alpha^{2} - M^{2})\hat{\Box}_{0}^{3} - 2\alpha^{2}(\alpha^{2}c_{\ell} - M^{2})\hat{\Box}_{0}^{2} - 4c_{\ell}\alpha^{4}(\alpha^{2} - M^{2})\hat{\Box}_{0} - 2\alpha^{8}c_{\ell}^{2} , \quad (3.28)$$

where

$$\hat{\Box}_0 \equiv \Box - \alpha^2 c_\ell \ . \tag{3.29}$$

In particular, the operator \mathcal{O}_4 has the property that for $\ell = 1$ it factorizes as

$$\mathcal{O}_4|_{\ell=1} = \Box \mathcal{O}_3 ,$$

$$\mathcal{O}_3 \equiv \Box^3 - (M^2 + 7\alpha^2)\Box^2 + 2\alpha^2 (4M^2 + 7\alpha^2)\Box - 12\alpha^4 (M^2 + \alpha^2) .$$
(3.30)

In the $\ell = 0$ sector, we will encounter the wave operator

$$\widetilde{\mathcal{O}}_2 = \Box(\Box + \alpha^2) - M^2(\Box - 2\alpha^2) .$$
(3.31)

Spin-2 sector

The spin-2 sector contains only the transverse and traceless gravitons, which upon the use of the spin projector operators provided in the Appendix C, and for $\ell \geq 1$, satisfy the following equation

$$\ell \ge 1$$
: $\mathcal{O}_2 \left(\mathcal{P}^2 h \right)^{(\ell)}_{\mu\nu} = 0$, (3.32)

where P^2 is the spin-2 projector defined in Appendix C. This equation describes two massive gravitons with mass squared

$$\ell \ge 1$$
: $m_{\pm}^2(\ell) = \frac{1}{2} \left(M^2 + 2\alpha^2 c_\ell \pm \sqrt{M^4 + 4\alpha^4 c_\ell} \right)$. (3.33)

The $\ell = 0$ needs to be treated separately, and in this case the gravitons satisfy

$$(\Box - M^2) R^{L(0)}_{\mu\nu} = -M^2 \left(\partial_\mu \partial_\nu S^{(0)} + \alpha^2 \eta_{\mu\nu} S^{(0)} \right) + \partial_\mu \partial_\nu (\Box + \alpha^2) S^{(0)}, \qquad (3.34)$$

where $S^{(0)} = \phi^{(0)} + N^{(0)}$. The solutions of this equation can be expressed as $h^{(0)}_{\mu\nu} = h'^{(0)}_{\mu\nu} + h''^{(0)}_{\mu\nu}$, where $h''^{(0)}_{\mu\nu}$ is completely determined by $S^{(0)}$ while $h'^{(0)}_{\mu\nu}$ is the solution to the following equations modulo the gauge symmetry (3.23):

$$\ell = 0: \qquad \Box (\Box - M^2) h_{\mu\nu}^{\prime(0)} = 0, \qquad R^{\prime(0)} = 0 , \qquad (3.35)$$

which describe a massless graviton and massive graviton with $m^2 = M^2$.

Spin-1 sector

Let $\ell \geq 2$. Then, the spin-1 sector consists of eight vectors $(P^1h)_{\mu\nu}, \partial^{\nu}b_{\mu\nu}, z_{\mu}^T, k_{\mu}^T, a_{\mu}^T, v_{\mu}^T, b_{\mu\nu}^T, b_{\mu}^T)$, where "T" indicates the transverse part and $(P^1h)_{\mu\nu} = P^1_{\mu\nu}{}^{\rho\sigma}h_{\rho\sigma}$ (see Appendix B). Of these eight vectors, $(k_{\mu}^T, a_{\mu}^T, v_{\mu}^T, b_{\mu\nu}^T, b_{\mu}^T)$ have mixing with each other through the following equations

$$0 = \mathcal{O}_2 b_{\mu\nu}^{T(\ell)} + 4g M^2 \star F_{\mu\nu}^{(\ell)}(a) - \alpha^4 c_\ell \left(\star F_{\mu\nu}^{(\ell)}(k) - \star F_{\mu\nu}^{(\ell)}(b) \right) , \qquad (3.36)$$

$$0 = \mathcal{O}_1 \hat{\Box}_0 b^{T(\ell)}_\mu + \frac{1}{2} \alpha^2 \epsilon^{\nu \lambda \rho}_\mu \partial_\nu b^{T(\ell)}_{\lambda \rho} , \qquad (3.37)$$

$$0 = (\hat{\Box}_0 + \alpha^2) a_{\mu}^{T(\ell)} - 4g \alpha^2 c_{\ell} k_{\mu}^{T(\ell)} + 4g v_{\mu}^{T(\ell)} - 2g \epsilon_{\mu}^{\nu \lambda \rho} \partial_{\nu} b_{\lambda \rho}^{T(\ell)} , \qquad (3.38)$$

$$0 = (\hat{\Box}_0 - M^2) v_{\mu}^{T(\ell)} + \alpha^2 c_{\ell} k_{\mu}^{T(\ell)} + \frac{1}{4} g M^2 a_{\mu}^{(\ell)} , \qquad (3.39)$$

$$0 = \left(\mathcal{O}_{2} + \alpha^{2}(\hat{\Box}_{0} - \alpha^{2}c_{\ell})\right)k_{\mu}^{T(\ell)} + 4gM^{2}a_{\mu}^{T(\ell)} + \frac{1}{2}\alpha^{2}\left(4v_{\mu}^{T(\ell)} - \epsilon_{\mu}^{\nu\lambda\rho}\partial_{\nu}b_{\lambda\rho}^{T(\ell)}\right) .$$
(3.40)

Diagonalising the associated 5×5 operator-valued matrix, we find that the modes are annihilated by $\mathcal{O}_1^2 \mathcal{O}_2 \mathcal{O}_4$. In particular, the linear combinations with coefficients $(-1, -4g\alpha^2 c_{\ell}/M^2, 0, 0, 1)$ and $(2, 4g\alpha^2 c_{\ell}/M^2, 1, 0, 0)$ are annihilated by \mathcal{O}_1 . The remaining vectors, namely, $((P^1h)_{\mu\nu}, \partial^{\nu}b_{\mu\nu}, z_{\mu})$ are separately annihilated by \mathcal{O}_1 as well. In summary, for $\ell \geq 2$ the total wave operator can be denoted by

$$\ell \ge 2: \qquad \mathcal{O}^{(1)} = \mathcal{O}_1^6 \, \mathcal{O}_2 \, \mathcal{O}_4 \,, \tag{3.41}$$

implying six massive vectors with mass squared

$$\ell \ge 2$$
: $m^2(\ell) = M^2 + \alpha^2(c_\ell - 1)$, (3.42)

two massive vectors with mass squared defined in Eq.(3.33) and four massive vectors whose squared masses are given by the roots of the polynomial

$$\begin{aligned} x^{4} + ax^{3} + bx^{2} + cx + d &= 0 , \\ a &= -M^{2} - (4\ell^{2} + 4\ell - 1)\alpha^{2} , \\ b &= \alpha^{2} \left[2M^{2} + \ell(\ell + 1) \left((6\ell^{2} + 6\ell - 5)\alpha^{2} + 3M^{2} \right) \right] , \\ c &= -\ell(\ell + 1)\alpha^{2} \left\{ \alpha^{2} \left[4 + \ell(\ell + 1)(4\ell^{2} + 4\ell - 7) \right] \alpha^{2} + 3\ell(\ell + 1)M^{2} \right\} , \\ d &= \ell^{2}(\ell + 1)^{2}(\ell - 1)(\ell + 2)\alpha^{6} \left[(\ell^{2} + \ell - 1)\alpha^{2} + M^{2} \right] . \end{aligned}$$
(3.43)

Next, consider the case $\ell = 1$. Recalling the factorization result given in (3.30), the total wave operator becomes

$$\ell = 1: \qquad \mathcal{O}^{(1)} = \mathcal{O}_1^6 \mathcal{O}_2|_{\ell=1} \Box \mathcal{O}_3 .$$
 (3.44)

In particular the massless vector is a linear combination of $(k_{\mu}^{T(1)}, a_{\mu}^{T(1)}, v_{\mu}^{T(1)}, b_{\mu\nu}^{T(1)}, b_{\mu\nu}^{T(1)})$ with mixing coefficients $(1, -4g, \alpha^2, 1, 0)$. The squared masses associated with $\mathcal{O}_1^6 \mathcal{O}_2|_{\ell=1}$ can be read of from (3.42) and (3.33) by setting $\ell = 1$, and those associated with \mathcal{O}_3 are the roots of the following polynomial

$$x^{3} - (M^{2} + 7\alpha^{2})x^{2} + 2\alpha^{2}(4M^{2} + 7\alpha^{2})x - 12\alpha^{4}(M^{2} + \alpha^{2}) = 0.$$
(3.45)

There remains the case of $\ell = 0$. The only vector fluctuations at this level are $(b_{\mu\nu}^{T(0)}, a_{\mu}^{T(0)}, v_{\mu}^{T(0)})$. Upon diagonalising the associated 3×3 operator-valued matrix, we find that the modes are annihilated by the following partially-factorising operator polynomial

$$\ell = 0: \qquad \mathcal{O}^{(1)} = \Box (\Box - M^2) \widetilde{\mathcal{O}}_2 , \qquad (3.46)$$

where the would-be massless vector annihilated by \Box is eaten by the two form and the operator \tilde{O}_2 is defined in (3.31). Thus, for $\ell = 0$ there are no massless vector modes, a massive vector with mass M and two massive vectors with squared masses given by

$$\widetilde{m}_{\pm}^2 = \frac{1}{2} \left(M^2 - \alpha^2 \pm \sqrt{M^4 - 10M^2\alpha^2 + \alpha^4} \right) .$$
(3.47)

The Spin-0 sector

We start with the case $\ell \geq 2$. Defining $\tilde{\varphi} = \omega_{\mu\nu}h^{\mu\nu}$ and $\varphi = \frac{1}{3}\theta_{\mu\nu}h^{\mu\nu}$ (see Appendix C), this sector consists of thirteen scalars $(\phi, N, \varphi, \tilde{\varphi}, a, v, \partial^{\mu}k_{\mu}, \partial^{\mu}b_{\mu}, \partial^{\mu}a_{\mu}, \partial^{\mu}v_{\mu}, \partial^{\mu}z_{\mu}, z, \tilde{z})$. The first six scalars $(\phi, N, \varphi, \tilde{\varphi}, a, v)$ mix as follows

$$\ell \ge 2: \quad 0 = 2(\hat{\Box}_0 + \alpha^2)\phi^{(\ell)} + (2\hat{\Box}_0 + 2\alpha^2 + \alpha^2 c_\ell)N^{(\ell)} + 3\hat{\Box}_0\varphi^{(\ell)} - \alpha^2 c_\ell \tilde{\varphi}^{(\ell)} , \qquad (3.48)$$

$$0 = \mathcal{O}_2 \varphi^{(\ell)} + 2\alpha^2 M^2 \left[\phi^{(\ell)} - 2gc_\ell a^{(\ell)} \right] - 2\alpha^2 (M^2 - \alpha^2 c_\ell) N^{(\ell)} + 2\alpha^3 c_\ell v^{(\ell)} , \qquad (3.49)$$

$$0 = \left(M^2 - (\alpha^2 + \Box)\right)\tilde{\varphi}^{(\ell)} + 3(M^2 - \alpha^2)\varphi^{(\ell)} + 2M^2\phi^{(\ell)} + (2\hat{\Box}_0 + 2\alpha^2 + \alpha^2 c_\ell)N^{(\ell)} ,$$
(3.50)

$$0 = (\hat{\square}_0 + \alpha^2)a^{(\ell)} + 4gN^{(\ell)} - 2g(\tilde{\varphi}^{(\ell)} + 3\varphi^{(\ell)}) + 4gv^{(\ell)} , \qquad (3.51)$$

$$0 = (\hat{\Box}_0 - M^2)v^{(\ell)} - \alpha^2 N^{(\ell)} + \frac{1}{2}\alpha^2 (\tilde{\varphi}^{(\ell)} + 3\varphi^{(\ell)}) - 2gM^2 a^{(\ell)} , \qquad (3.52)$$

$$0 = \left(M^2 - (\alpha^2 - \Box)\right)N^{(\ell)} + 3M^2\varphi^{(\ell)} + 2M^2\phi^{(\ell)} - 4gM^2a^{(\ell)} - 2\alpha^2v^{(\ell)} - \alpha^2(c_\ell - 1)\tilde{\varphi}^{(\ell)} .$$
(3.53)

Diagonalising the associated 6×6 operator-valued matrix, we find that the modes are annihilated by $\mathcal{O}_1^3 \mathcal{O}_4$. Of the remaining scalars, $(\partial^{\mu}k_{\mu}, \partial^{\mu}b_{\mu}, \partial^{\mu}a_{\mu}, \partial^{\mu}v_{\mu}, \tilde{v})$ mix but only three of them are dynamical. We choose these to be $(\partial^{\mu}k_{\mu}, \partial^{\mu}b_{\mu}, \tilde{v})$ which are separately annihilated by \mathcal{O}_1 , while $(\partial^{\mu}a_{\mu}, \partial^{\mu}v_{\mu})$ are determined by

$$\partial^{\mu} a_{\mu}^{(\ell)} = \frac{\alpha^2}{2g} \left(\tilde{v}^{(\ell)} - \partial^{\mu} k_{\mu}^{(\ell)} \right) ,$$

$$\partial^{\mu} v_{\mu}^{(\ell)} = \alpha^2 (c_{\ell} - 1) \tilde{v}^{(\ell)} + \alpha^2 \partial^{\mu} k_{\mu}^{(\ell)} . \qquad (3.54)$$

Finally, the remaining scalars (z, \tilde{z}) are annihilated by \mathcal{O}_1 , and the longitudinal modes $\partial^{\mu} z_{\mu}$ are given in terms of z and \tilde{z} , by virtue of the equation $\hat{D}^M \hat{z}_M = 0$. Thus, for $\ell \geq 2$ the total wave operator is given by

$$\ell \ge 2: \qquad \mathcal{O}^{(0)} = \mathcal{O}_1^{10} \, \mathcal{O}_4 \; . \tag{3.55}$$

Of these, the three linear combinations of $(\phi, N, \varphi, \tilde{\varphi}, a, v)$ with coefficients $(2+\alpha^2 c_{\ell}/M^2, -2, 0, 0, 0, 2),$ $(-2-\alpha^2 c_{\ell}/M^2, 2, 0, 0, -8g/M^2, 0)$ and $(-2+\alpha^2 c_{\ell}/M^2, 2, 0, 4, 0, 0)$ are annihilated by \mathcal{O}_1 .

In the case of $\ell = 1$, utilizing the residual symmetry (3.24) and (3.25), one can eliminate $N^{(1)}$ and $\partial^{\mu}k^{(1)}_{\mu}$. Taking into account the fact that the harmonic expansion of \hat{z}_m contributes only one complex scalar for $\ell = 1$, namely $z^{(1)}$, we find that for $\ell = 1$ the total wave operator for the scalar fields is given by

$$\ell = 1: \qquad \mathcal{O}^{(0)} = \mathcal{O}_1^6|_{\ell=1}\mathcal{O}_3 .$$
 (3.56)

There remains the case of $\ell = 0$. Of the remaining scalars, $(\phi^{(0)}, N^{(0)}, b^{(0)})$ satisfy the equations $\widetilde{\mathcal{O}}_2 S^{(0)} = 0$ where $S^{(0)} = \phi^{(0)} + N^{(0)}$, $\Box \mathcal{O}_1|_{\ell=0} b^{(0)} = 0$ and $\Box \mathcal{O}_1|_{\ell=0} N^{(0)} = 0$. Finally, there is a complex scalar $z^{(0)}$ annihilated by $\mathcal{O}_1|_{\ell=0}$. Thus, for $\ell = 0$ the total wave operator or the scalar fields is given by

$$\ell = 0: \qquad \mathcal{O}^{(0)} = \Box^2 \, \mathcal{O}^4_1|_{\ell=0} \, \widetilde{\mathcal{O}}_2 \,\,, \tag{3.57}$$

implying two massless and six massive scalars in this sector.

3.3 Fermionic sector

In terms of the complex spinor, the linearized equations of fermions around the background (3.1) are given by

$$0 = \bar{\Gamma}^{MPQ}\bar{D}_P\psi_Q + 2i\bar{\Gamma}^{NM}\bar{D}_N\chi - 4gi\bar{\Gamma}^M\lambda - \frac{1}{2}\bar{F}_{PQ}\bar{\Gamma}^{PQ}\bar{\Gamma}^M\lambda - \frac{1}{8M^2}\Theta^M , \qquad (3.58)$$

$$0 = \bar{\Gamma}^{MN} \bar{D}_M \psi_N - 2i \bar{\Gamma}^M \bar{D}_M \chi - 8gi\lambda, \qquad (3.59)$$

$$0 = 2gi\bar{\Gamma}^M\psi_M + 8g\chi - 4\bar{\Gamma}^M\bar{D}_M\lambda + \frac{1}{4}\bar{F}_{PQ}\bar{\Gamma}^M\bar{\Gamma}^{PQ}\psi_M , \qquad (3.60)$$

where

$$\bar{D}_{M}\psi = (\partial_{M} + \frac{1}{4}\bar{\omega}_{M}^{AB}\bar{\Gamma}_{AB})\psi - \frac{i}{2}\bar{V}_{M}\psi ,$$

$$\Theta^{M} = 8\bar{\Gamma}^{P}\bar{D}_{Q}\bar{D}_{P}\psi^{QM} - 2\bar{R}^{PM}_{ST}\bar{\Gamma}^{Q}\bar{\Gamma}^{ST}\bar{D}_{P}\psi_{Q} + 2\bar{R}^{PQ}_{ST}\bar{\Gamma}^{ST}\bar{\Gamma}^{M}\bar{D}_{P}\psi_{Q}$$

$$+ 8i\bar{G}^{P[M}\bar{\Gamma}^{Q]}\bar{D}_{Q}\psi_{P} - 8i\bar{G}^{M[P}\bar{\Gamma}^{Q]}\bar{D}_{Q}\psi_{P} .$$
(3.61)

In the remainder of this section, we shall drop the "bar" on the covariant derivatives as well as the Γ -matrices for simplicity in notation. Since the (3.61) contains gauge field, we adopt the spinweighted harmonics $_{s-\frac{1}{2}}\eta^{(\ell)}$, which are described in detail in appendix B, as the expansion basis. In this section we will need the harmonics for s = 0, namely $-\frac{1}{2}\eta^{(\ell)}$ which we will denote as $\eta^{(\ell)}$ for brevity in notation. These harmonics satisfy the relations,

$$\eta_{-}^{(0)} = \eta , \qquad \eta_{+}^{(\ell)} = \frac{1}{i\alpha\sqrt{c_{\ell}}} \nabla_n Y^{(\ell)} \sigma^n \eta , \qquad \eta_{-}^{(\ell)} = Y^{(\ell)} \eta, \qquad \ell \ge 1 , \qquad (3.62)$$

and have the following properties

$$\begin{aligned}
\sigma_{3}\eta_{\pm}^{(\ell)} &= \pm \eta_{\pm}^{(\ell)} , & \sigma^{n} D_{n} \eta_{\pm}^{(\ell)} &= i\alpha \sqrt{c_{\ell}} \eta_{\mp}^{(\ell)} , \\
[D_{m}, D_{n}]\eta_{-}^{(\ell)} &= 0 , & [D_{m}, D_{n}]\eta_{+}^{(\ell)} &= i\alpha^{2} \epsilon_{mn} \eta_{+}^{(\ell)} , \\
D^{n} D_{n} \eta_{-}^{(\ell)} &= -\alpha^{2} c_{\ell} \eta_{-}^{(\ell)} , & D^{n} D_{n} \eta_{+}^{(\ell)} &= -\alpha^{2} (c_{\ell} - 1) \eta_{+}^{(\ell)} . \end{aligned} \tag{3.63}$$

The Killing spinor η also has the property $\sigma_3\eta = -\eta$. Furthermore, given that $\Gamma_7 = \gamma_5 \times \sigma_3$ (see Appendix A), the chirality property of a spinor in 6D correlates the 4D and σ_3 chiralities.

Since there is no gamma traceless and transverse spin-3/2 harmonics on the S^2 , generically, the harmonic expansion are carried out as

$$\hat{\psi}_{\mu} = \psi_{\mu-}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\psi_{\mu+}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \psi_{\mu-}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) ,$$

$$\hat{\psi}_{m} = \Gamma_{m} \psi_{+}^{(0)} \otimes \eta^{(0)} + \Gamma_{m} \sum_{\ell \ge 1} \left(\psi_{-}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \psi_{+}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) ,$$

$$+ \sum_{\ell \ge 1} \left(\tilde{\psi}_{+}^{(\ell)} \otimes D_{\{m\}} \eta_{+}^{(\ell)} + \tilde{\psi}_{-}^{(\ell)} \otimes D_{\{m\}} \eta_{-}^{(\ell)} \right) ,$$

$$\hat{\chi} = \chi_{+}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\chi_{-}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \chi_{+}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) ,$$

$$\hat{\lambda} = \lambda_{-}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\lambda_{+}^{(\ell)} \eta_{+}^{(\ell)} + \lambda_{-}^{(\ell)} \eta_{-}^{(\ell)} \right) ,$$
(3.64)

where $D_{\{m\}}$ is the gamma traceless covariant derivative and the \pm subscripts denote chirality property under γ_5 . Using the 6D linearized fermionic gauge symmetry (3.4), one can impose the following gauge condition

$$\hat{\psi}_{\{m\}} = 0$$
, (3.65)

where $\{m\}$ means Γ -traceless. As a consequence, the expansion takes the following simpler forms

$$\hat{\psi}_{\mu} = \psi_{\mu-}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\psi_{\mu+}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \psi_{\mu-}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) , \qquad (3.66)$$

$$\hat{\psi}_m = \Gamma_m \psi_+^{(0)} \otimes \eta^{(0)} + \Gamma_m \sum_{\ell \ge 1} \left(\psi_-^{(\ell)} \otimes \eta_+^{(\ell)} + \psi_+^{(\ell)} \otimes \eta_-^{(\ell)} \right) , \qquad (3.67)$$

$$\hat{\chi} = \chi_{+}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\chi_{-}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \chi_{+}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) , \qquad (3.68)$$

$$\hat{\lambda} = \lambda_{-}^{(0)} \otimes \eta^{(0)} + \sum_{\ell \ge 1} \left(\lambda_{+}^{(\ell)} \otimes \eta_{+}^{(\ell)} + \lambda_{-}^{(\ell)} \otimes \eta_{-}^{(\ell)} \right) .$$

$$(3.69)$$

The gauge choice (3.65) does not fix all the gauge symmetries, we find the following residual symmetry transformations

• Generated by $\hat{\epsilon}^{(0)} = \epsilon^{(0)} \eta^{(0)}$:

$$\delta\psi^{(0)}_{\mu} = \partial_{\mu}\epsilon^{(0)}, \qquad (3.70)$$

• Generated by $\hat{\epsilon} = \epsilon_+^{(1)} \eta_+^{(1)}$:

$$\delta\psi_{\mu+}^{(1)} = \partial_{\mu}\epsilon_{+}^{(1)} + \frac{i}{\sqrt{2}}\,\alpha\,\epsilon_{+}^{(1)} \,, \qquad (3.71)$$

$$\delta \lambda_+^{(1)} = ig \epsilon_+^{(1)} . \tag{3.72}$$

We shall take into account these symmetries in the analysis of the spectrum below, where we treat the spin-3/2, spin-1/2 sectors separately.

Spin-3/2 sector

Let us begin with the restriction $\ell \geq 1$. This sector contains only the gravitino fields which satisfy the following equations

$$\ell \ge 1: \qquad \vartheta \Big((\hat{\Box}_0 + \alpha^2) - M^2 \Big) P^{3/2} \psi_{\mu+}^{(\ell)} + i\alpha \sqrt{c_\ell} \Big((\hat{\Box}_0 + \alpha^2) - M^2 \Big) P^{3/2} \psi_{\mu-}^{(\ell)} = 0 , \\ i\alpha \sqrt{c_\ell} \Big((\hat{\Box}_0 + \alpha^2) - M^2 \Big) P^{3/2} \psi_{\mu+}^{(\ell)} - \vartheta \Big(\hat{\Box}_0 - M^2 \Big) P^{3/2} \psi_{\mu-}^{(\ell)} = 0 , \qquad (3.73)$$

where $P^{3/2}$ is the spin-3/2 projector operator defined in Appendix C. Diagonalising the associated 2×2 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of sixth order in ∂ , given by

$$\ell \ge 1: \qquad \mathcal{O}^{(3/2)} = \mathcal{O}_1 \mathcal{O}_2 \ . \tag{3.74}$$

Next, consider the case of $\ell = 0$. By choosing the gauge $\gamma^{\mu}\psi^{(0)}_{\mu} = 0$, the gravitino equation can be written as

$$\partial \left(\Box - M^2\right)\psi^{(0)\mu} = -\left(\partial \partial^{\mu} - M^2\gamma^{\mu}\right)\Psi^{(0)} - 2M^2\left(\gamma^{\mu\nu}\partial_{\nu}\psi^{(0)} - i\gamma^{\mu\nu}\partial_{\nu}\chi^{(0)}\right) .$$
(3.75)

The solutions of above equation can be expressed as $\psi_{\mu}^{(0)} = \psi_{\mu}^{\prime(0)} + \psi_{\mu}^{\prime\prime(0)}$ where $\psi_{\mu}^{\prime\prime(0)}$ is completely determined by $\psi^{(0)}$ and $\chi^{(0)}$ while $\psi_{\mu}^{\prime(0)}$ is the solution to the following equations modular gauge symmetry (3.70)

$$\ell = 0: \qquad \not \partial(\Box - M^2)\psi_{\mu}^{\prime(0)} = 0 , \qquad \gamma^{\mu}\psi_{\mu}^{\prime(0)} = 0 , \qquad \partial^{\mu}\psi_{\mu}^{\prime(0)} = 0 . \tag{3.76}$$

It describes a massless and two massive gravitini.

Spin-1/2 sector

The $\ell \geq 2$ sector consists of ten spin 1/2 fields $(\Lambda_+, \Psi_+, \Lambda_-, \Psi_-, \psi_+, \psi_-, \chi_+, \chi_-, \lambda_+, \lambda_-)$, where $\Psi^{(\ell)} \equiv \partial^{\mu} \psi^{(\ell)}_{\mu}$ and $\gamma^{\mu} \psi^{(\ell)}_{\mu} \equiv \Lambda^{(\ell)}$. The linearized equations describing their mixing are

$$0 = \partial \Lambda_{-}^{(\ell)} - \Psi_{+}^{(\ell)} + i\alpha \sqrt{c_{\ell}} \psi_{+}^{(\ell)} - i\alpha \sqrt{c_{\ell}} \Lambda_{+}^{(\ell)} + 2\partial \psi_{-}^{(\ell)} - 2i\partial \chi_{-}^{(\ell)} + 2\alpha \sqrt{c_{\ell}} \chi_{+}^{(\ell)} - 8gi\lambda_{+}^{(\ell)} , \qquad (3.77)$$

$$0 = \partial \Lambda_{+}^{(\ell)} - \Psi_{-}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\psi_{-}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\Lambda_{-}^{(\ell)} - 2\partial \psi_{+}^{(\ell)} + 2i\partial \chi_{+}^{(\ell)} + 2\alpha\sqrt{c_{\ell}}\chi_{-}^{(\ell)} - 8gi\lambda_{-}^{(\ell)} , \quad (3.78)$$

$$0 = ig\Lambda_{-}^{(\ell)} + 2g\chi_{-}^{(\ell)} - \partial \lambda_{+}^{(\ell)} - i\alpha\sqrt{c_{\ell}}\lambda_{-}^{(\ell)} , \qquad (3.79)$$

$$0 = 2ig\psi_{+}^{(\ell)} + 2g\chi_{+}^{(\ell)} + \partial \lambda_{-}^{(\ell)} - i\alpha\sqrt{c_{\ell}}\lambda_{+}^{(\ell)} , \qquad (3.80)$$

$$0 = i\alpha\sqrt{c_{\ell}}\vartheta(\Box + M^{2})\psi_{+}^{(\ell)} - iM^{2}\alpha\sqrt{c_{\ell}}\vartheta\Lambda_{+}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\left(M^{2} - \alpha^{2} + \alpha^{2}c_{\ell}\right)\Psi_{-}^{(\ell)} + 2M^{2}\alpha\sqrt{c_{\ell}}\vartheta\chi_{+}^{(\ell)} - 8igM^{2}\vartheta\lambda_{+}^{(\ell)} + \alpha^{2}(c_{\ell} - 1)\vartheta\Psi_{+}^{(\ell)} + \alpha^{2}(2 - c_{\ell})\Box\psi_{-}^{(\ell)} , \qquad (3.81)$$

$$0 = -i\partial \left(\Box + M^{2} - \alpha^{2}\right) \psi_{-}^{(\ell)} - iM^{2}\partial \Lambda_{-}^{(\ell)} + i\left(\alpha^{2} + \alpha^{2}c_{\ell} - M^{2}\right) \Psi_{+}^{(\ell)} - 2M^{2}\partial \chi_{-}^{(\ell)} - \alpha\sqrt{c_{\ell}} \left(\partial \Psi_{-}^{(\ell)} + \Box \psi_{+}^{(\ell)}\right) , \qquad (3.82)$$

$$0 = \vartheta \left(\Box + \alpha^{2} - \alpha^{2} c_{\ell} + 2M^{2} \right) \Lambda_{-}^{(\ell)} - \left(\Box + 2\alpha^{2} - 2\alpha^{2} c_{\ell} + 2M^{2} \right) \Psi_{+}^{(\ell)} + i\alpha \sqrt{c_{\ell}} (\Box + 4M^{2}) \psi_{+}^{(\ell)} + \vartheta \left(6M^{2} + 2\alpha^{2} - \alpha^{2} c_{\ell} \right) \right) \psi_{-}^{(\ell)} - i\alpha \sqrt{c_{\ell}} \left(\Box + \alpha^{2} - \alpha^{2} c_{\ell} + 3M^{2} \right) \Lambda_{+}^{(\ell)} - 6iM^{2} \vartheta \chi_{-}^{(\ell)} + 8\alpha M^{2} \sqrt{c_{\ell}} \chi_{+}^{(\ell)} - 32igM^{2} \lambda_{+}^{(\ell)} + i\alpha \sqrt{c_{\ell}} \vartheta \Psi_{-}^{(\ell)} , \qquad (3.83)$$

$$0 = -\vartheta \left(\Box - \alpha^{2} c_{\ell} + 2M^{2} \right) \Lambda_{+}^{(\ell)} + \left(\Box - 2\alpha^{2} c_{\ell} + 2M^{2} \right) \Psi_{-}^{(\ell)} - i\alpha \sqrt{c_{\ell}} \left(\Box - 4\alpha^{2} + 4M^{2} \right) \psi_{-}^{(\ell)} + \vartheta (6M^{2} - \alpha^{2} c_{\ell}) \psi_{+}^{(\ell)} - i\alpha \sqrt{c_{\ell}} \left(\Box + \alpha^{2} - \alpha^{2} c_{\ell} + 3M^{2} \right) \Lambda_{-}^{(\ell)} - 6iM^{2} \vartheta \chi_{+}^{(\ell)} - 8\alpha M^{2} \sqrt{c_{\ell}} \chi_{-}^{(\ell)} + i\alpha \sqrt{c_{\ell}} \vartheta \Psi_{+}^{(\ell)} , \qquad (3.84)$$

$$0 = M^{2} \left(-\Lambda_{-}^{(\ell)} + 2i\chi_{-}^{(\ell)} \right) - \left(2\Box - \alpha^{2}c_{\ell} \right)\psi_{-}^{(\ell)} + \partial \Psi_{+}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\Psi_{-}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\partial \psi_{+}^{(\ell)} , \qquad (3.85)$$

$$0 = (\alpha^{2} - M^{2})\Lambda_{+}^{(\ell)} - 2iM^{2}\chi_{+}^{(\ell)} + (2\Box + 2\alpha^{2} - \alpha^{2}c_{\ell})\psi_{+}^{(\ell)} - i\alpha\sqrt{c_{\ell}}\Psi_{+}^{(\ell)} + \partial\Psi_{-}^{(\ell)} + i\alpha\sqrt{c_{\ell}}\partial\psi_{-}^{(\ell)}.$$
(3.86)

Diagonalising the associated 10×10 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of eighteenth order in ∂ , given by

$$\ell \ge 2: \qquad \mathcal{O}^{1/2} = \mathcal{O}_1^5 \, \mathcal{O}_4 \; . \tag{3.87}$$

Next we consider the case of $\ell = 1$. In this case, one can use the fermionic shift symmetry (3.72) to eliminate $\psi_{+}^{(1)}$. Consequently we get 9 by 9 mixing and we find that the modes are annihilated by the partially-factorising operator polynomial, of fifteenth order in ∂ , given by

$$\ell = 1: \qquad \mathcal{O}^{1/2} = \mathcal{O}_1^4|_{\ell=1} \not \partial \mathcal{O}_3 , \qquad (3.88)$$

where \mathcal{O}_3 is defined in (3.30). The factor ∂ demonstrates that there is massless spin-1/2 mode. This massless mode corresponds to linear combinations of $(\Lambda_+^{(1)}, \Psi_+^{(1)}, \Lambda_-^{(1)}, \Psi_-^{(1)}, \psi_-^{(1)}, \chi_+^{(1)}, \chi_-^{(1)}, \lambda_+^{(1)}, \lambda_-^{(1)})$ with mixing coefficients (8, 0, 0, 0, -2, -2*i*, 0, 0, 1).

There remains the case of $\ell = 0$. In this case, we have

$$\ell = 0: \qquad 0 = -\Psi^{(0)} - 2\partial \psi^{(0)} + 2i\partial \chi^{(0)} - 8gi\lambda^{(0)} , \qquad (3.89)$$

$$0 = 2gi\psi^{(0)} + 2g\chi^{(0)} + \partial \lambda^{(0)} , \qquad (3.90)$$

$$0 = (1 + \frac{1}{2M^2} \Box) \Psi^{(0)} + 3 \partial \!\!\!/ \psi^{(0)} - 3i \partial \!\!/ \chi^{(0)} , \qquad (3.91)$$

$$0 = -\Psi^{(0)} - \partial \left(1 + \frac{1}{M^2} (\Box + \alpha^2) \right) \psi^{(0)} + 2i \partial \chi^{(0)} - 8gi\lambda^{(0)} .$$
 (3.92)

Diagonalising the associated 4×4 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of seventh order in ∂ , given by

$$\ell = 0: \qquad \mathcal{O}^{(1/2)} = \mathcal{O}_1|_{\ell=0} \, \partial \widetilde{\mathcal{O}}_2 \,. \tag{3.93}$$

Thus, at the $\ell = 0$ level, there is only one massless spin-1/2 modes given by $\Psi^{(0)} = 0, \lambda^{(0)} = 0, i\psi^{(0)} + \chi^{(0)} = 0.$

3.4 The supermultiplet structure and stability

In arranging the full spectrum described above into a collection of supermultiplet structure, it is useful to recall that following massive supermultiplets:

massive supergravity multiplet :
$$(h_{\mu\nu}, A_{\mu}, \Psi_{\mu})$$
,massive gravitino multiplet : $(\psi_{\mu}, Z_{\mu}, \chi)$,massive vectormultiplet multiplet : (A_{μ}, ϕ, λ) ,massive scalar multiplet : (Z, ψ) ,

where A_{μ} is a real and Z_{μ} is a complex vector, ϕ is a real and Z is a complex scalar, and Ψ_{μ} is Dirac and ψ_{μ}, χ, ψ are Majorana. The Dirac gravitino can be written as $\Psi = \psi_{\mu+}^1 + \psi_{\mu-}^2$, where the two terms represent Weyl spinors that are independent of each other, and consequently Ψ_{μ} on-shell describes 8 real degrees of freedom. The Majorana gravitino, on the other hand can be written as $\psi_{\mu} = \psi_{\mu+} + \psi_{\mu-}$ where $\psi_{\mu-} = (\psi_{\mu+})^*$. Thus, on shell ψ_{μ} describes 4 real degrees of freedom. With this information at hand, we can now tabulate the supermultiplet structure of the full spectrum. It is convenient to do so by specifying the wave operators for different spin fields and consider the cases of $\ell = 0$, $\ell = 1$ and $\ell \geq 2$ separately. The results are given in Table 1, Table 2 and Table 3.

Having established the full spectrum of states in the four-dimensional theory, we may now examine the question of stability, which is governed by the mass values for the massive fields. We begin with the $\ell = 0$ level, given in Table 1. In addition to the massless graviton and scalar

s=2	s=2 $s=3/2$		s = 1/2	s = 0
	ð			
$\Box - M^2$	$\Box - M^2$	$\Box - M^2$		
		$\widetilde{\mathcal{O}}_2$	$\widetilde{\mathcal{O}}_2$	$\widetilde{\mathcal{O}}_2$
			ð	\square^2
			\mathcal{O}_1	\mathcal{O}_1^4

Table 1: The spectrum of wave operators for $\ell = 0$. The operator \mathcal{O}_1 is to be evaluated for $\ell = 0$. There is one massless spin-2 and one massless spin-0 multiplet, a massive spin-2 multiplet with mass M, two spin-0 multiplets with squared mass $m^2 = M^2 - \alpha^2$ and two massive spin-1 multiplets with mass² given in (3.47).

s = 2	s = 3/2	s = 1	s = 1/2	s = 0
\mathcal{O}_2	\mathcal{O}_2	\mathcal{O}_2		
	\mathcal{O}_1	\mathcal{O}_1^4	\mathcal{O}_1	
		\mathcal{O}_3	\mathcal{O}_3	\mathcal{O}_3
		\mathcal{O}_1^2	\mathcal{O}_1^2	\mathcal{O}_1^2
			ð	
			\mathcal{O}_1	\mathcal{O}_1^4

Table 2: The spectrum of wave operators for $\ell = 1$. The operators \mathcal{O}_1 and \mathcal{O}_2 are to be evaluated for $\ell = 1$. There are two spin-2 multiplets with squared masses given in (3.33) for $\ell = 1$, two spin-3/2, two spin-1 multiplets and two spin-0 multiplets with squared mass $m^2 = M^2 + \alpha^2$, three spin-1 multiplets with squared masses given by the roots of the polynomial given in (3.45) and a massless vector multiplet.

multiplets, there are massive graviton, vector and scalar multiplets at this level. The massive graviton multiplet has $m^2 = M^2$, the scalar multiplet has $m^2 = M^2 - \alpha^2$, and the vector multiplet has masses given by (3.47). These imply, respectively, that stability requires $M^2 > 0$, $M^2 > \alpha^2$ and $M^2 \ge (5 + 2\sqrt{6})\alpha^2 \approx 9.89898\alpha^2$.

At the level $\ell = 1$, in addition to the massless vector multiplet, there are massive graviton, gravitino, vector and scalar multiplets. The gravitino multiplet, two of the massive vector multiplets and the scalar multiplet have masses given by the operator \mathcal{O}_1 , implying $m^2 = M^2 + \alpha^2$, which therefore impose no new conditions. The massive graviton multiplet with mass operator \mathcal{O}_2 has m^2 given by equation (3.33) with ℓ set equal to 1. This implies $m^2 = (M^2 + 4\alpha^2 \pm \sqrt{M^4 + 8\alpha^4})/2$, and hence gives no further restriction. There remains the massive vector multiplet with mass operator \mathcal{O}_3 given in (3.30). This gives a cubic polynomial in m^2 , and we find that this has three real (and positive) roots for m^2 provided that $\mu \equiv M^2/\alpha^2$ satisfies the condition

$$4\mu^4 - 64\mu^3 + 153\mu^2 - 26\mu - 139 \ge 0.$$
(3.95)

This implies we must have $\mu \ge \mu_{\min}$, where $\mu_{\min} \approx 13.1425$. In other words, at level $\ell = 1$ stability requires that M^2 should exceed approximately $13.1425\alpha^2$.

For levels $\ell \geq 2$, the multiplets are given in Table 3. The gravitino, vector and scalar multiplets with mass operator \mathcal{O}_1 have m^2 given in equation (3.42), and these are always positive for all values of $\ell \geq 2$. Likewise, for the graviton multiplet with mass operator \mathcal{O}_2 , m^2 , given in (3.33),

s = 2	s = 3/2	s = 1	s = 1/2	s = 0
\mathcal{O}_2	\mathcal{O}_2	\mathcal{O}_2		
	\mathcal{O}_1	\mathcal{O}_1^4	\mathcal{O}_1	
		\mathcal{O}_4	\mathcal{O}_4	\mathcal{O}_4
		\mathcal{O}_1^2	\mathcal{O}_1^2	\mathcal{O}_1^2
			\mathcal{O}_1^2	\mathcal{O}_1^8

Table 3: The spectrum of wave operators for $\ell \geq 2$. For each integer ℓ , there are two spin-2 multiplets with squared mass $m_{\pm}^2(\ell)$ given in (3.33), two spin-3/2, two spin-1 multiplets and four spin-0 multiplets with squared mass $m^2(\ell)$ given in (3.42), and four spin-1 multiplets with squared masses given by the roots of the polynomial given in (3.43).

is positive for all $\ell \geq 2$. There remains the vector multiplet with mass operator \mathcal{O}_4 . This leads to a quartic polynomial in m^2 , which can be read off from (3.28) and (3.29). One can show that this polynomial necessarily has at least two real roots, which are positive, and that if the four roots for m^2 are real then they are also positive. The condition for having four real roots is that a rather complicated discriminant of sixth order in $\mu = M^2/\alpha^2$ should be positive. This discriminant also depends on the level ℓ . For a few representative values of ℓ , we find the requirement $\mu \geq \mu_{\min}(\ell)$:

$\ell =$	2	3	5	10	100	1000	10000
$\mu_{\rm min} \approx$	16.9381	20.869	28.8614	49.0439	414.215	4067.07	40595.8

Table 4: Minimum values of $\mu = M^2/\alpha^2$ necessary to achieve real positive mass-squared values for the \mathcal{O}_4 vector multiplet at level ℓ .

In the limit of large ℓ , we find that to leading order, $\mu_{\min}(\ell)$ grows linearly with ℓ , with

$$\mu_{\min}(\ell) \sim 4.05874\,\ell + \cdots \tag{3.96}$$

This implies that for any given ratio $\mu = M^2/\alpha^2$, there is a critical level ℓ_{max} beyond which the Kaluza-Klein tower must be truncated in order not to have modes with complex masses, which would be associated with instabilities.

4 Spectrum in Non-Supersymmetric Minkowski₄ \times S² Background

4.1 Non-supersymmetric Minkowski₄ \times S² background

In addition to supersymmetric vacuum solution discussed in previous section, the theory [5] also possesses non-supersymmetric Minkowski₄ × S² vacua when $M^2 = \alpha^2$, with the curvature and flux given by

$$\bar{R}_{\mu\nu\lambda\rho} = 0 , \qquad \bar{R}_{mn} = \alpha^2 \bar{g}_{mn} , \qquad \bar{L} = 1 ,
\bar{F}_{\mu\nu} = 0 , \qquad \bar{F}_{mn} = 4qg\epsilon_{mn} ,
\bar{G}_{\mu\nu} = 0 , \qquad \bar{G}_{mn} = -q\alpha^2\epsilon_{mn} ,$$
(4.1)

where q plays the role of monopole charge and is quantized to be $q = 0, \pm 1, \pm 2...$ The supersymmetric vacua correspond to $q = \pm 1$.

4.2 Bosonic sector

We will perform a similar spectrum analysis around the non-supersymmetric background. The harmonic expansion (3.22) for the uncharged fields after the gauge fixing is still valid, and the residual gauge symmetries are almost the same except that some terms related to background flux should be multiplied by the monopole charge. We present the results for the spectrum below. While we shall use the same notation for operators such as $\hat{\Box}_0$ and others, it is understood that they are to be evaluated for $M^2 = \alpha^2$.

Spin-2 sector

The equations of motion satisfied by graviton for $\ell \geq 1$ is

$$\ell \ge 1: \qquad (\hat{\Box}_0^2 - \alpha^2 \hat{\Box}_0 - \alpha^4 c_\ell) (\mathcal{P}^2 h)_{\mu\nu}^{(\ell)} = 0 , \qquad (4.2)$$

describing massive gravitons with square masses

$$m_{\pm}^{2}(\ell) = \frac{1}{2}\alpha^{2} \left(1 + c_{\ell} \pm \sqrt{1 + 4c_{\ell}}\right) .$$
(4.3)

For $\ell = 0$, the linearized field equation is

$$\ell = 0: \qquad (\Box - \alpha^2) R^{L(0)}_{\mu\nu} = -\alpha^2 \partial_\mu \partial_\nu S^{(0)} - \alpha^4 \eta_{\mu\nu} S^{(0)} + \partial_\mu \partial_\nu (\Box + \alpha^2) S^{(0)} , \qquad (4.4)$$

where $S^{(0)} = \phi^{(0)} + N^{(0)}$. It describes a massless graviton and massive graviton with squared mass $m^2 = \alpha^2$.

Spin-1 sector

For $\ell \geq 2$ the mixing among the five vector fields $(k_{\mu}^{T}, a_{\mu}^{T}, v_{\mu}^{T}, b_{\mu\nu}^{T}, b_{\mu}^{T})$ now have the following form

$$\ell \ge 2: \quad 0 = (2c_{\ell}\alpha^4 - \hat{\Box}_0^2)k_{\mu}^{T(\ell)} - 4g\alpha^2 q a_{\mu}^{T(\ell)} - \frac{\alpha^2}{2}(4qv_{\mu}^{T(\ell)} - \epsilon_{\mu}^{\nu\lambda\rho}\partial_{\nu}b_{\lambda\rho}^{T(\ell)}) , \quad (4.5)$$

$$0 = (\alpha^2 \Box - \hat{\Box}_0^2) b_{\mu\nu}^{T(\ell)} - 4qg\alpha^2 \star F_{\mu\nu}^{(\ell)}(a) + \alpha^4 c_\ell (\star F_{\mu\nu}^{(\ell)}(k) - \star F_{\mu\nu}^{(\ell)}(b)) , \qquad (4.6)$$

$$0 = \hat{\Box}_0^2 b_\mu^{T(\ell)} + \frac{1}{2} \alpha^2 \epsilon_\mu^{\nu\lambda\rho} \partial_\nu b_{\lambda\rho}^{T(\ell)} , \qquad (4.7)$$

$$0 = (\hat{\Box}_0 + \alpha^2) a_{\mu}^{T(\ell)} - 4qg\alpha^2 c_{\ell} k_{\mu}^{T(\ell)} + 4gv_{\mu}^{T(\ell)} - 2qg\epsilon_{\mu}^{\ \nu\lambda\rho}\partial_{\nu} b_{\lambda\rho}^{T(\ell)} , \qquad (4.8)$$

$$0 = (\hat{\Box}_0 - \alpha^2) v_{\mu}^{T(\ell)} + \alpha^4 q c_{\ell} k_{\mu}^{T(\ell)} - 2g \alpha^2 a_{\mu}^{(\ell)} , \qquad (4.9)$$

Diagonalising the associated 5×5 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of eighth order in $\hat{\Box}_0$, given by $\hat{\Box}_0^2 \mathcal{O}_6$. The explicit form of \mathcal{O}_6 can be obtained straightforwardly from (4.9), and one finds that it is symmetric under $q \rightarrow -q$ meaning that vector spectrum is symmetric under the sign change of monopole charge. Of the remaining vectors $((\mathcal{P}^1 h)_{\mu\nu}, \partial^{\mu} b_{\mu\nu})$ are annihilated by $\hat{\Box}_0$. Thus, apart from the charged vectors which will be treated separately below, the total wave operator for $\ell \geq 2$ is given by

$$\ell \ge 2: \qquad \mathcal{O}^{(1)} = \hat{\Box}_0^4 |_{M^2 = \alpha^2} \mathcal{O}_6 , \qquad (4.10)$$

implying four massive vectors with squared masses $m^2 = \alpha^2 c_\ell$, and six massive vectors whose squared masses m^2 correspond to the roots \mathcal{O}_6 in which \Box is to be replaced by m^2 .

In the case of $\ell = 1$, again excluding the charged vector, we find that a massless vector appears since for $c_{\ell} = 2$, the operator \mathcal{O}_6 factorizes as $\mathcal{O}_6 = \Box \mathcal{O}_5$, and the total wave operator becomes

$$\ell = 1: \qquad \mathcal{O}^{(1)} = \hat{\Box}_0^4|_{\ell=1} \Box \mathcal{O}_5 .$$
 (4.11)

The massless vector is composed from a linear combination of $(k_{\mu}^{T(1)}, a_{\mu}^{T(1)}, v_{\mu}^{T(1)}, b_{\mu\nu}^{T(1)}, b_{\mu\nu}^{T(1)})$ with

mixing coefficients $\left(\frac{2}{1+q^2}, -\frac{8qg}{1+q^2}, \frac{2q\alpha^2}{1+q^2}, 1, 0\right)$. In the uncharged vector sector, there remains the case of $\ell = 0$, for which the relevant vector fields are $(b_{\mu\nu}^{T(0)}, a_{\mu}^{T(0)}, v_{\mu}^{T(0)})$. Upon diagonalising the associated 3×3 operator-valued matrix, we find that the modes are annihilated by the following partially-factorising operator polynomial

$$\ell = 0:$$
 $\mathcal{O}^{(1)} = \Box (\Box - \alpha^2) (\Box^2 + 2\alpha^4 q^2).$ (4.12)

As before, we find that the would-be massless modes annihilated by \Box is eaten by the two form. Thus there are no massless vector modes at $\ell = 0$.

Finally, we turn to the treatment of the complex vector \hat{z}_{μ} . This field \hat{z}_{μ} is expanded in terms of charge "-q" scalar harmonics starting from $\ell = |q|$ as follows:

$$\hat{z}_{\mu} = \sum_{\ell \ge q} z_{\mu}^{(\ell)} \,_{-q} Y^{(\ell)} \,, \tag{4.13}$$

The resulting linearized field equation is

$$\ell \ge |q|:$$
 $(\Box - \alpha^2 c_\ell + \alpha^2 q^2 - M^2) z_\mu^{T(\ell)} = 0.$ (4.14)

Spin-0 sector

For $\ell \geq 2$, the equations describing the mixing between $(\phi, N, \varphi, \tilde{\varphi}, a, v)$ take the following form

$$\ell \ge 2: \quad 0 = 2(\hat{\Box}_0 + \alpha^2)\phi^{(\ell)} + (2\hat{\Box}_0 + 2\alpha^2 + \alpha^2 c_\ell)N^{(\ell)} + 3\hat{\Box}_0\varphi^{(\ell)} - \alpha^2 c_\ell \tilde{\varphi}^{(\ell)} , \qquad (4.15)$$

$$0 = \alpha^2 (3\varphi^{(\ell)} + 2\phi^{(\ell)} - 4gqa^{(\ell)} - 2qv^{(\ell)}) + \alpha^2 (1 - c_\ell)\tilde{\varphi}^{(\ell)} + \Box N^{(\ell)} , \qquad (4.16)$$

$$0 = 2\alpha^2 \phi^{(\ell)} + (2\hat{\Box}_0 + 2\alpha^2 + \alpha^2 c_\ell) N^{(\ell)} - \Box \tilde{\varphi}^{(\ell)} , \qquad (4.17)$$

$$0 = (\hat{\square}_0 + \alpha^2)a^{(\ell)} + 4gqN^{(\ell)} - 2gq(3\varphi^{(\ell)} + \tilde{\varphi}^{(\ell)}) + 4gv^{(\ell)} , \qquad (4.18)$$

$$0 = (\hat{\Box}_0 - \alpha^2)v^{(\ell)} - \alpha^2 q N^{(\ell)} + \frac{\alpha^2}{2} q (3\varphi^{(\ell)} + \tilde{\varphi}^{\ell}) - 2g\alpha^2 a^{(\ell)} , \qquad (4.19)$$

$$0 = (\alpha^2 \Box - \hat{\Box}_0^2) \varphi^{(\ell)} + 2\alpha^4 \phi^{(\ell)} + \alpha^4 (2 - c_\ell) N^{(\ell)} - 4qgc_\ell \alpha^4 a^{(\ell)} - 2c_\ell \alpha^4 qv^{(\ell)} . (4.20)$$

Diagonalising the associated 6×6 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of seventh order in $\hat{\Box}_0$, given by $\mathcal{O}_4 \hat{\Box}_0^3|_{M^2=\alpha^2}$ Of the remaining scalars, $(\partial^{\mu}k_{\mu}, \partial^{\mu}b_{\mu}, \partial^{\mu}a_{\mu}, \partial^{\mu}v_{\mu}, \tilde{v})$, three of them namely $(\partial^{\mu}k_{\mu}, \partial^{\mu}b_{\mu}, \tilde{v})$ are annihilated by $\hat{\Box}_0$, and the remaining two are determined in terms of them. Thus, in total, apart from the complex scalars which will be treated separately below, the wave operator for the scalar fields is given by

$$\mathcal{O}^{(0)} = \hat{\Box}_0^6 \,\mathcal{O}_4 \,\,. \tag{4.21}$$

Next, consider the case $\ell = 1$. Utilizing the residual symmetry (3.24) and (3.25), one can eliminate $N^{(1)}$ and $\partial^{\mu}k^{(1)}_{\mu}$. The mass operator coming from the mixing among $(\phi, \varphi, \tilde{\varphi}, a, v)$ takes the form $\hat{\Box}^2_0|_{\ell=1}\mathcal{O}_3$. Taking into account $\partial^{\mu}b^{(1)}_{\mu}$ and $\tilde{v}^{(1)}$, the total wave operator, again, excluding the complex scalar sector, is given by

$$\ell = 1: \qquad \mathcal{O}^{(0)} = \hat{\Box}_0^4 \,\mathcal{O}_3 \;.$$
(4.22)

There remaining the case of $\ell = 0$. In this case, the relevant scalar fields are $(\phi^{(0)}, N^{(0)}, b^{(0)})$, and they satisfy the following equations respectively

$$\ell = 0: \qquad (\Box^2 + 2\alpha^4) S^{(0)} = 0 , \qquad S^{(0)} = \phi^{(0)} + N^{(0)} ,$$
$$\Box^2 b^{(0)} = 0 ,$$
$$\Box^2 N^{(0)} = 0 . \qquad (4.23)$$

Thus besides two massless modes, we also have modes with linear time coordinate dependence. Finally, we discuss the complex scalars originating from \hat{z}_m . For positive monopole charge, the harmonic expansion of \hat{z}_m is given by

$$\hat{z}_m = z^{(q-1)}{}_{-q}V_m^{(q-1)} + z^{(q)}{}_{-q}V_m^{(q)} + \sum_{\ell > q} (z^{(\ell)}D_{m-q}Y^{(\ell)} + \tilde{z}^{(\ell)}\epsilon_m^{\ n}D_{n-q}Y^{(\ell)}) \ .$$
(4.24)

Thus we have

$$\ell > q: \qquad z^{(\ell)} , \ \tilde{z}^{(\ell)} \qquad \text{with} \qquad m^2 = \alpha^2 c_\ell - \alpha^2 q + M^2 , \\ \ell = q: \qquad z^{(q)} , \qquad \text{with} \qquad m^2 = \alpha^2 q + M^2 , \\ \ell = q - 1: \qquad z^{(q-1)} , \qquad \text{with} \qquad m^2 = M^2 - \alpha^2 q .$$
(4.25)

4.3 Fermionic sector

The analysis of the fermionic spectrum in a non-supersymmetric background (4.1) is more subtle than that in supersymmetric background. Since the non-supersymmetric background do not posses Killing spinor, we will use spin-weighted harmonics $s-\frac{1}{2}\eta^{(\ell)}$, described in detail in appendix B, as basis of expansion. For brevity, we shall use the notation

$$_{s-\frac{1}{2}}\eta^{(\ell)} \equiv \tilde{\eta}^{(\ell)} , \qquad (4.26)$$

where $s = \frac{1}{2}(1-q)$. These harmonics satisfy the relations

$$\tilde{\eta}_{+}^{(\ell)} = \eta_{+} \left({}_{s-1}Y^{(\ell)} \right) , \qquad \tilde{\eta}_{-}^{(\ell)} = \eta_{-} \left({}_{s}Y^{(\ell)} \right) , \qquad (4.27)$$

$$\left(\frac{d}{d\theta} + m\csc\theta + s\cot\theta\right)\left({}_{s}Y^{(\ell)}\right) = \sqrt{(\ell+s)(\ell+1-s)}\left({}_{s-1}Y^{(\ell)}\right) , \qquad (4.28)$$

$$\left(\frac{d}{d\theta} - m\csc\theta - (s-1)\cot\theta\right)\left(_{s-1}Y^{(\ell)}\right) = -\sqrt{(\ell+s)(\ell+1-s)}\left(_{s}Y^{(\ell)}\right) .$$
(4.29)

The lowest level would have definite chirality when $\ell = -s$ for q > 0 and $\ell = s - 1$ for q < 0. The spin weighted harmonics satisfy the following properties

$$\begin{aligned}
\sigma_{3}\tilde{\eta}_{\pm}^{(\ell)} &= \pm \tilde{\eta}_{\pm}^{(\ell)} , & \sigma^{n} D_{n} \tilde{\eta}_{\pm}^{(\ell)} &= i\alpha \sqrt{\tilde{c}_{\ell}} \tilde{\eta}_{\mp}^{(\ell)} , \\
[D_{m}, D_{n}]\tilde{\eta}_{-}^{(\ell)} &= -is\alpha^{2} \epsilon_{mn} \tilde{\eta}_{-}^{(\ell)} , & [D_{m}, D_{n}]\tilde{\eta}_{+}^{(\ell)} &= i(1-s)\alpha^{2} \epsilon_{mn} \eta_{+}^{(\ell)} , \\
D^{n} D_{n} \tilde{\eta}_{-}^{(\ell)} &= -\alpha^{2} (\tilde{c}_{\ell} - s) \tilde{\eta}_{-}^{(\ell)} , & D^{n} D_{n} \tilde{\eta}_{+}^{(\ell)} &= \alpha^{2} (1-s-\tilde{c}_{\ell}) \tilde{\eta}_{+}^{(\ell)} ,
\end{aligned} \tag{4.30}$$

where

$$\tilde{c}_{\ell} = \left(c_{\ell} - s(s-1)\right) \,. \tag{4.31}$$

The harmonic expansion for 6D spin-1/2 fields follows the same procedure as in supersymmetric case by using the spin weighted harmonics, while it is more subtle when expanding the 6D gravitini.

It can be checked that the linearized equations have the following discret symmetry

$$q \to -q , \qquad \psi_{\mu+} \to -\psi_{\mu-}, \qquad \psi_{\mu-} \to \psi_{\mu+} ,$$

$$\psi_{-} \to \psi_{+} , \qquad \psi_{+} \to -\psi_{-} , \qquad \chi_{-} \to \chi_{+} ,$$

$$\chi_{+} \to -\chi_{-} , \qquad \lambda_{+} \to -\lambda_{-} , \qquad \lambda_{-} \to \lambda_{+} ,$$
(4.32)

which implies that the spectrum keeps the same under the sign change of monopole charge. In the following, we will focus on the case with positive monopole charge and use |s| to denote $\frac{1}{2}(q-1)$.

Spin-3/2 Sector

The gravitini satisfy

$$\ell \ge |s|+1: \qquad \qquad \oint \left(\Box + \alpha^2 |s| - \alpha^2 \tilde{c}_\ell\right) \psi_{\mu+}^{(\ell)} + i\alpha \sqrt{\tilde{c}_\ell} \left(\Box - \alpha^2 \tilde{c}_\ell\right) \psi_{\mu-}^{(\ell)} = 0 ,$$
$$i\alpha \sqrt{\tilde{c}_\ell} \left(\Box - \alpha^2 \tilde{c}_\ell\right) \psi_{\mu+}^{(\ell)} - \oint \left(\Box + (|s|+1-\tilde{c}_\ell)\alpha^2\right) \psi_{\mu-}^{(\ell)} = 0 . \qquad (4.33)$$

Diagonalising the associated 2×2 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of third order in \Box , given by

$$\mathcal{O}^{(3/2)} = \tilde{\mathcal{O}}_3 , \qquad (4.34)$$

where the explicit form of $\tilde{\mathcal{O}}_3$ can be deduced from (4.33).

Next, we consider the case of $\ell = |s|$. In this case, we find that the quadratic action for the lowest level fermionic fields is proportional to

$$\mathcal{L}^{(2)} \propto -i\bar{\chi}\gamma^{\mu\nu}\partial_{\mu}\psi_{\nu} - i\bar{\psi}_{\mu}\gamma^{\mu\nu}\partial_{\nu}\chi + 2i\bar{\chi}\partial\!\!\!/\psi - 2i\bar{\psi}\partial\!\!/\chi + 2\bar{\chi}\partial\!\!/\chi -8g\bar{\chi}\lambda - 8g\bar{\lambda}\chi - 4\bar{\lambda}\partial\!\!/\lambda - 4ig|s|\bar{\lambda}\gamma^{\mu}\psi_{\mu} - 4ig|s|\bar{\psi}_{\mu}\gamma^{\mu}\lambda -8g(1+|s|)i\bar{\lambda}\psi + 8g(1+|s|)i\bar{\psi}\lambda - \frac{1}{2}\bar{\psi}_{\mu}\gamma^{\mu\nu\lambda}\partial_{\nu}\psi_{\lambda} + \bar{\psi}\partial\!\!/\psi + \bar{\psi}_{\mu}\gamma^{\mu\nu}\partial_{\nu}\psi - \psi\gamma^{\mu\nu}\partial_{\mu}\psi_{\nu} - \frac{1}{\alpha^{2}}\left(\frac{1}{4}\bar{\psi}_{\mu\nu}\partial\!\!/\psi^{\mu\nu} - \bar{\psi}\partial\!\!/\Box\psi + \frac{1}{2}|s|\alpha^{2}\bar{\psi}_{\mu}\partial\!\!/\psi^{\mu} - |s|\alpha^{2}\bar{\psi}\partial_{\mu}\psi^{\mu} + |s|\alpha^{2}\bar{\psi}_{\mu}\partial^{\mu}\psi - (1+2|s|)\alpha^{2}\bar{\psi}\partial\!\!/\psi\right).$$
(4.35)

Unlike the supersymmetric case, we see the appearance of the terms $\bar{\lambda}\gamma^{\mu}\psi_{\mu}$, $\bar{\psi}_{\mu}\partial\psi^{\mu}\psi^{\mu}$ and $\bar{\psi}_{\mu}\partial^{\mu}\psi$ which break the fermionic gauge symmetry. The homogeneous solutions for gravitini satisfy

$$\mathscr{D}(\Box - \alpha^2 - |s|\alpha^2)\psi_{\mu}^{(|s|)} = 0, \qquad \gamma^{\mu}\psi_{\mu}^{(|s|)} = 0, \qquad \partial^{\mu}\psi_{\mu}^{(|s|)} = 0.$$
(4.36)

Thus, due to the lack of fermionic gauge symmetry, the longitudinal mode $\psi_{\mu} \propto p_{\mu} e^{ipx}$ with $p^2 = 0$ becomes a dynamical degree of freedom.

Spin-1/2 Sector

The $\ell \geq |s| + 2$ sector consists of ten spin-1/2 fields $(\Lambda_+, \Psi_+, \Lambda_-, \Psi_-, \psi_+, \psi_-, \chi_+, \chi_-, \lambda_+, \lambda_-)$. The linearized equations describing their mixing are

$$0 = \partial \Lambda_{-}^{(\ell)} - \Psi_{+}^{(\ell)} + i\alpha \sqrt{\tilde{c}_{\ell}} \psi_{+}^{(\ell)} - i\alpha \sqrt{\tilde{c}_{\ell}} \Lambda_{+}^{(\ell)} + 2\partial \psi_{-}^{(\ell)} - 2i\partial \chi_{-}^{(\ell)} + 2\alpha \sqrt{\tilde{c}_{\ell}} \chi_{+}^{(\ell)} - 8gi\lambda_{+}^{(\ell)} , \qquad (4.37)$$

$$0 = \partial \Lambda_{+}^{(\ell)} - \Psi_{-}^{(\ell)} + i\alpha \sqrt{\tilde{c}_{\ell}} \psi_{-}^{(\ell)} + i\alpha \sqrt{\tilde{c}_{\ell}} \Lambda_{-}^{(\ell)} - 2\partial \psi_{+}^{(\ell)} + 2i\partial \chi_{+}^{(\ell)} + 2\alpha \sqrt{\tilde{c}_{\ell}} \chi_{-}^{(\ell)} - 8gi\lambda_{-}^{(\ell)} , \qquad (4.38)$$

$$0 = g(1+|s|)i\Lambda_{-}^{(\ell)} + 2g\chi_{-}^{(\ell)} - \partial \lambda_{+}^{(\ell)} - i\alpha\sqrt{\tilde{c}_{\ell}}\lambda_{-}^{(\ell)} , \qquad (4.39)$$

$$0 = 2ig|s|\psi_{+}^{(\ell)} + 2g\chi_{+}^{(\ell)} + \partial \lambda_{-}^{(\ell)} - i\alpha\sqrt{\tilde{c}_{\ell}}\lambda_{+}^{(\ell)}, \qquad (4.40)$$

$$0 = i\sqrt{\tilde{c}_{\ell}}\vartheta(\Box + \alpha^{2} + |s|\alpha^{2})\psi_{+}^{(\ell)} - i\alpha^{2}\sqrt{\tilde{c}_{\ell}}\vartheta\Lambda_{+}^{(\ell)} + i\alpha^{2}\tilde{c}_{\ell}^{3/2}\Psi_{-}^{(\ell)} + 2\alpha^{2}\sqrt{\tilde{c}_{\ell}}\vartheta\chi_{+}^{(\ell)} -8g(1+|s|)\alpha\,i\vartheta\lambda_{+}^{(\ell)} + \alpha(\tilde{c}_{\ell} - 1 - |s|)\vartheta\Psi_{+}^{(\ell)} + \alpha(2+2|s| - \tilde{c}_{\ell})\Box\psi_{-}^{(\ell)} , \qquad (4.41)$$

$$0 = i\sqrt{\tilde{c}_{\ell}}\vartheta(\Box - |s|\alpha^2)\psi_{-}^{(\ell)} + i\alpha^2\sqrt{\tilde{c}_{\ell}}\vartheta\Lambda_{-}^{(\ell)} - i\alpha^2\tilde{c}_{\ell}^{3/2}\Psi_{+}^{(\ell)} + 2\alpha^2\sqrt{\tilde{c}_{\ell}}\vartheta\chi_{-}^{(\ell)} + 8ig|s|\alpha\vartheta\lambda_{-}^{(\ell)} + \alpha(\tilde{c}_{\ell} + |s|)\vartheta\Psi_{-}^{(\ell)} + \alpha(\tilde{c}_{\ell} + 2|s|)\Box\psi_{+}^{(\ell)} , \qquad (4.42)$$

$$0 = \vartheta(\Box + 3\alpha^{2} + |s|\alpha^{2} - \alpha^{2}\tilde{c_{\ell}})\Lambda_{-}^{(\ell)} - (\Box + 4\alpha^{2} + 2|s|\alpha^{2} - 2\alpha^{2}\tilde{c_{\ell}})\Psi_{+}^{(\ell)} + i\alpha\sqrt{\tilde{c_{\ell}}}(\Box + 4\alpha^{2} + 4|s|\alpha^{2})\psi_{+}^{(\ell)} + \alpha^{2}\vartheta(8 + 2|s| - \tilde{c_{\ell}})\psi_{-}^{(\ell)} - i\alpha\sqrt{\tilde{c_{\ell}}}(\Box + 4\alpha^{2} - \alpha^{2}\tilde{c_{\ell}})\Lambda_{+}^{(\ell)} - 6i\alpha^{2}\vartheta\chi_{-}^{(\ell)} + 8\alpha^{3}\sqrt{\tilde{c_{\ell}}}\chi_{+}^{(\ell)} - 32ig\alpha^{2}(1 + |s|)\lambda_{+}^{(\ell)} + i\alpha\sqrt{\tilde{c_{\ell}}}\vartheta\Psi_{-}^{(\ell)},$$

$$(4.43)$$

$$0 = -\emptyset(\Box + 2\alpha^{2} - |s|\alpha^{2} - \alpha^{2}\tilde{c}_{\ell})\Lambda_{+}^{(\ell)} + (\Box + 2\alpha^{2} - 2|s| - 2\alpha^{2}\tilde{c}_{\ell})\Psi_{-}^{(\ell)} - i\alpha\sqrt{\tilde{c}_{\ell}}(\Box - 4|s|\alpha^{2})\psi_{-}^{(\ell)} + \alpha^{2}\emptyset(6 - 2|s| - \tilde{c}_{\ell})\psi_{+}^{(\ell)} - i\alpha\sqrt{\tilde{c}_{\ell}}(\Box + 4\alpha^{2} - \alpha^{2}\tilde{c}_{\ell})\Lambda_{-}^{(\ell)} - 6i\alpha^{2}\vartheta\chi_{+}^{(\ell)} - 8\alpha^{3}\sqrt{\tilde{c}_{\ell}}\chi_{-}^{(\ell)} - 32ig|s|\alpha^{2}\lambda_{-}^{(\ell)} + i\alpha\sqrt{\tilde{c}_{\ell}}\vartheta\Psi_{+}^{(\ell)},$$

$$(4.44)$$

$$0 = (1+|s|)\alpha^{2}\Lambda_{-}^{(\ell)} - 2i\alpha^{2}\chi_{-}^{(\ell)} + (2\Box - 2|s|\alpha^{2} - \alpha^{2}\tilde{c}_{\ell})\psi_{-}^{(\ell)} - \partial\!\!\!/ \Psi_{+}^{(\ell)} - i\alpha\sqrt{\tilde{c}_{\ell}}\vartheta\psi_{+}^{(\ell)} ,$$

$$(4.45)$$

$$0 = |s|\alpha^{2}\Lambda_{+}^{(\ell)} - 2i\alpha^{2}\chi_{+}^{(\ell)} + (2\Box + 2\alpha^{2} + 2|s|\alpha^{2} - \alpha^{2}\tilde{c_{\ell}})\psi_{+}^{(\ell)} - i\alpha\sqrt{\tilde{c_{\ell}}}\Psi_{+}^{(\ell)} + \partial\!\!\!/\Psi_{-}^{(\ell)} + i\alpha\sqrt{\tilde{c_{\ell}}}\partial\!\!\!/\psi_{-}^{(\ell)} .$$

$$(4.46)$$

Diagonalising the associated 10×10 operator-valued matrix, we find that the modes are annihilated by the partially-factorising operator polynomial, of ninth order in \Box , given by

$$\ell \ge |s| + 2:$$
 $\mathcal{O}^{(0)} = \hat{\Box}_0^3 \tilde{\mathcal{O}}_6 ,$ (4.47)

where the explicit form of the operator \mathcal{O}_6 can be determined from the linearized spin 1/2 field equations listed above.

$$\ell = |s| + 1$$

At this level, since $D_m \eta_+^{(|s|+1)} = \frac{i}{2} \alpha \sqrt{\tilde{c_\ell}} \sigma_m \eta_-^{(|s|+1)}$, there emerges a fermionic gauge symmetry generated by $\hat{\epsilon} = \epsilon_+^{(|s|+1)} \eta_+^{(|s|+1)}$

$$\delta \psi_{\mu+}^{(|s|+1)} = \partial_{\mu} \epsilon_{+}^{(|s|+1)} + i\alpha \sqrt{\frac{(1+|s|)}{2}} \epsilon_{+}^{(|s|+1)} ,$$

$$\delta \lambda_{+}^{(|s|+1,m)} = ig(1+|s|) \epsilon_{+}^{(|s|+1)} .$$
(4.48)

Using this gauge symmetry, one can eliminate $\psi_{+}^{(|s|+1)}$, such that the 10 by 10 mixing becomes 9 by 9 mixing. Diagonalising this system, we find that the modes are determined by the partially-factorising operator polynomial, of fifteenth order in ∂ , given by

$$\ell = |s| + 1: \qquad \partial \left(\Box - (|s| + 2)\alpha^2 \right)^2 \tilde{\mathcal{O}}_5 , \qquad (4.49)$$

where the explicit form of $\tilde{\mathcal{O}}_5$ can be deduced from the mixing equations. It is clear that is a massless spin-1/2 mode. Explicitly, it is a linear combination of $(\Lambda_+^{(|s|+1)}, \Psi_+^{(|s|+1)}, \Lambda_-^{(|s|+1)}, \Psi_-^{(|s|+1)}, \psi_-^{(|s|+1)}, \chi_-^{(|s|+1)}, \chi_+^{(|s|+1)}, \Lambda_+^{(|s|+1)}, \chi_-^{(|s|+1)})$ with mixing coefficients

$$(8(1+2|s|), 0, 0, 0, -2, -2i(1+2|s|), 0, 0, \sqrt{1+|s|}(1+4|s|).$$

$$(4.50)$$



The coupled system of linearized field equations for the spin-1/2 fields $(\Lambda_{+}^{(|s|)}, \Psi_{-}^{(|s|)}, \psi_{+}^{(|s|)}, \chi_{+}^{(|s|)}, \lambda_{-}^{(|s|)})$ are

$$0 = \partial \Lambda_{+}^{(|s|)} - \Psi_{-}^{(|s|)} - 2 \partial \psi_{+}^{(|s|)} + 2i \partial \chi_{+}^{(|s|)} - 8gi\lambda_{-}^{(|s|)}, \qquad (4.51)$$

$$0 = ig|s|\Lambda_{+}^{(|s|)} + 2g(1+|s|)i\psi_{+}^{(|s|)} + 2g\chi_{+}^{(|s|)} + \partial \lambda_{-}^{(|s|)} = 0, \qquad (4.52)$$

$$0 = 8 ci \partial \lambda_{+}^{(|s|)} + \partial W_{+}^{(|s|)} + 2 \Box c \psi_{+}^{(|s|)} = 0 \qquad (4.52)$$

$$0 = 8gi\vartheta\lambda_{-}^{(l)} + \vartheta\Psi_{-}^{(l)} + 2\Box\psi_{+}^{(l)} = 0,$$

$$0 = (\Box + 2\alpha^{2} - |s|\alpha^{2})\partial\Lambda_{+}^{(|s|)} - (\Box + 2\alpha^{2} - 2|s|\alpha^{2})\Psi_{-}^{(|s|)} + (2|s| - 6)\alpha^{2}\partial\psi_{+}^{(|s|)} + 6\alpha^{2}i\partial\chi_{+}^{(|s|)}$$

$$(4.53)$$

$$+32g|s|\alpha^{2} i\lambda_{-}^{(|s|)}, \qquad (4.54)$$

$$0 = \alpha^{2} \partial \Lambda_{+}^{(|s|)} - (1+|s|)\alpha^{2} \Psi_{-}^{(|s|)} - \partial (\Box + 2\alpha^{2} + 2|s|\alpha^{2})\psi_{+}^{|s|} - 8g(1+|s|)\alpha^{2} i\lambda_{-}^{(|s|)} + 2\alpha^{2} i\partial \chi_{+}^{(|s|)}. \qquad (4.55)$$

Diagonalising this system, we find two massless modes which satisfy the relations

$$\Psi_{-}^{(|s|)} = 0 , \qquad \lambda_{-}^{(|s|)} = 0 , \qquad |s|\Lambda_{+}^{(|s|)} - 2i\chi_{+}^{(|s|)} + 2(1+|s|)\psi_{+}^{(|s|)} = 0 .$$

$$\boxed{\ell = |s| - 1}$$
(4.56)

When monopole charge $q \geq 3$, there exist charge "-q/2" vector-spinor harmonics η_m on S^2 possessing following properties

$$D^m \left({}_{s-\frac{1}{2}} \eta^{(|s|-1)} \right)_m = 0 , \qquad \sigma^m \left({}_{s-\frac{1}{2}} \eta^{(|s|-1)} \right)_m = 0 , \qquad (4.57)$$

where s = (1 - q)/2. It follows that

$$\sigma^m D_m \left({}_{s-\frac{1}{2}} \eta^{(|s|-1)} \right)_n = 0 .$$
(4.58)

Associated to this harmonics, there is a new spin-1/2 field $\tilde{\psi}^{(|s|-1)}$ satisfying

$$\partial \Box \tilde{\psi}^{(|s|-1)} = 0 . \tag{4.59}$$

4.4 Remarks on the non-supersymmetric spectrum

We saw in the supersymmetric vacuum that even at the $\ell = 0$ level in the Kaluza-Klein harmonic expansions, avoiding tachyons in the four-dimensional spectrum required imposing the condition $M^2 \ge (5 + 2\sqrt{6})\alpha^2$ on the parameter M in the six-dimensional Lagrangian. In the nonsupersymmetric vacua we necessarily have $M^2 = \alpha^2$, and in fact having larger values for the background monopole charge q then the $q = \pm 1$ supersymmetric case only makes the tachyon problem worse, as can be seen from the vector mass operator (4.12). For this reason, we shall not explore further the precise details of the occurrence of tachyonic states in the non-supersymmetric backgrounds.

One feature of interest that we shall, however, comment on is the occurrence of massless fermions in the non-supersymmetric backgrounds. As can be seen from the spin- $\frac{3}{2}$ operator in (4.36), there will be massless spin- $\frac{3}{2}$ fields at level $\ell = |s| = \frac{1}{2}(q-1)$; thus these will occur in an SU(2) multiplet of dimension $2\ell + 1 = q$. At $\ell = |s| + 1$, $\ell = |s|$ and $\ell = |s| - 1$ there will also be massless spin- $\frac{1}{2}$ modes, as was discussed in the spin- $\frac{1}{2}$ section above.

5 Conclusions

In this paper we have studied the complete linearised spectrum in the S^2 Kaluza-Klein reduction of off-shell six-dimensional $\mathcal{N} = (1,0)$ gauged supergravity extended by a Riemann-squared superinvariant. The higher-derivative terms in the six-dimensional theory can be expected to imply the occurrence of ghosts. As discussed in the introduction, the usual argument for the positivity of the energies of states in a supersymmetric background breaks down, and indeed we found that states in the Kaluza-Klein spectrum could now have complex energies, thus implying instabilities.

One way to understand the occurrence of complex masses is that there are mixings between fourdimensional ghostlike and non-ghostlike modes. This may be illustrated by the following simple example. Consider a set of fields ϕ_i with the Lagrangian

$$\mathcal{L} = -\frac{1}{2} K_{ij} \,\partial\phi_i \,\partial\phi_j - \frac{1}{2} V_{ij} \phi_i \phi_j \,, \tag{5.1}$$

where K_{ij} and V_{ij} are constant symmetric matrices. If the eigenvalues of K_{ij} are all positive, then K_{ij} and V_{ij} can be simultaneously diagonalised, by means of orthogonal transformations combined with rescalings of the fields. However, if K_{ij} has negative as well as positive eigenvalues, then the rescalings will introduce factors of $\sqrt{-1}$ and the diagonalised mass matrix will be complex.

One way to avoid the ghost problems of the higher-derivative theory is to treat it not as an exact model in its own right, but rather as an effective theory valid at energy scales $\sqrt{\Lambda}$ much smaller than M. In this case the propagators are governed by the leading-order theory without the higher-derivative terms, and these terms are treated as interactions. The Kaluza-Klein spectrum in

the reduced four-dimensional theory would then be simply that of the original Salam-Sezgin model corresponding to the $M^2 \to \infty$ limit. This spectrum has been given in [6] and our results agree in the $M^2 \to \infty$ limit. However, the Kaluza-Klein level number ℓ would have to be restricted to lie below some maximum value, in order to satisfy the $\Lambda \ll M^2$ limit. Interestingly, this condition is sufficient to ensure that in the full, extended, theory, the m^2 values of the retained modes would all be real and positive. This can be seen from (3.96), which indicates that the m^2 values will all be real and positive if ℓ is less than about $M^2/(4\alpha^2)$.

A couple of remarks about the consistency of the Kaluza-Klein reduction are in order. Although we have restricted ourselves to a linearised analysis of the four-dimensional spectrum, it should be emphasised that provided one is keeping all the infinite towers of modes, then even at the full non-linear order the reduction would still be consistent. The truncation of the spectrum at some maximum value of the level number ℓ that we discussed in the previous paragraph would not, of course, be consistent beyond the linear order, since the higher modes that were being set to zero would be excited by sources involving the modes that are being retained.

Another more subtle question of consistency arises in this model also. It was shown in [11] that the Salam-Sezgin theory admits a non-trivial consistent Pauli reduction on S^2 , in which a finite subset of fields including the $\ell = 1$ triplet of Yang Mills gauge bosons are retained. It would be interesting to see whether such a Pauli reduction is still possible in the theory with the higher-derivative extension that we have been considering in this paper.

Another interesting question is whether the six dimensional model, with the auxiliary fields eliminated in an order by order expansion in inverse powers of M^2 , can be embedded into the tendimensional heterotic string. In the gauged theory where $g \neq 0$, this continues to be a challenging problem even before the higher-derivative terms are considered (although some progress was made in a restricted sector of the theory in [12]). For g = 0, on the other hand, it was conjectured in [13] that there is a relation with the 4-torus reduction of the heterotic theory with Riemann-squared corrections that were constructed in [14]. This relation holds upon making a suitable truncation and performing an S-duality transformation. This conjecture was tested to lowest order in the bosonic sector in [13].

It would also be interesting to study exact solutions of the higher-derivative six-dimensional supergravity. While many solutions of the Salam-Sezgin theory are known, exact solutions of the higher-derivative theory, beyond the vacuum solutions we have discussed in this paper, are scarce. As far as we are aware, the only further example, which exists only in the ungauged theory, is the self-dual string that was found in [15].

A further question is whether there exist other quadratic-curvature superinvariants over and above the Riemann-squared invariant of the theory we have been considering. This may have consequences for the embedding of the theory in ten dimensions.

Acknowledgements

We are very grateful to F. Coomans and A. van Proeyen for many helpful discussions on the six-dimensional model. The research of C.N.P. is supported in part by DOE grant DE-FG03-95ER40917i, and that of E.S. is supported by in part by NSF grant PHY-0906222.

A Conventions

We choose the 6D gamma matrices to be

$$\Gamma^{0} = \gamma^{0} \otimes \sigma_{3} , \qquad \Gamma^{1} = \gamma^{1} \otimes \sigma_{3} , \qquad \Gamma^{2} = \gamma^{2} \otimes \sigma_{3} , \Gamma^{3} = \gamma_{3} \otimes \sigma_{3} , \qquad \Gamma^{4} = \mathbf{1}_{4 \times 4} \otimes \sigma_{1} , \qquad \Gamma^{5} = \mathbf{1}_{4 \times 4} \otimes \sigma_{2} .$$
(A.1)

One can check that

$$\Gamma^{0}\Gamma^{\mu\dagger}\Gamma^{0} = \Gamma^{\mu} , \qquad B = \Gamma^{3}\hat{\Gamma}^{5} ,
B^{*}B = -1 , \qquad B\Gamma^{\mu}B^{-1} = \Gamma^{\mu*} .$$
(A.2)

The SU(2) symplectic-Majorana-Weyl spinor is defined by

$$\psi^{*i} = (\psi_i)^* = \epsilon^{ij} B \psi_j. \tag{A.3}$$

A useful formula related to the SU(2) symplectic-Majorana-Weyl spinor is

$$\bar{\lambda}^{i}\Gamma^{(n)}\psi^{j} = t_{n}\bar{\psi}^{j}\Gamma^{(n)}\lambda^{i} , \qquad t_{n} = \begin{cases} +, & n = 1, 2, 5, 6; \\ -, & n = 0, 3, 4. \end{cases}$$
(A.4)

B Spin-weighted Harmonics on S^2

In this appendix, we give an elementary construction of the spin-weighted spherical harmonics. This is based on a specialisation of results for the analogous harmonics in the complex projective space CP^n , which were discussed in [10]. Since the azimuthal label m on the spin-weighted spherical harmonics ${}_{s}Y_{\ell m}$ plays an important role in the derivations in this appendix, we shall suspend our convention used in the body of the paper of suppressing the m label. In order to avoid confusion with coordinate indices, we shall use i, j,... for coordinate indices on S^2 in this appendix.

B.1 Scalar spin-weighted harmonics

The scalar spin-weighted spherical harmonics ${}_{s}Y_{\ell m}$ are the eigenfunctions of the *charged* scalar Laplacian $\Box_{(s)}$ on the unit S^2 , carrying electric charge s, in the presence of a Dirac monopole with potential $A = -\cos\theta \, d\phi$:

$$\Box_{(s)} \equiv \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta}\right) + \frac{1}{\sin^2\theta} \left(\frac{\partial}{\partial\phi} + is \cos\theta\right)^2.$$
(B.1)

In the language of differential forms, the charged Laplacian operator on the unit S^2 with metric $d\Omega_2^2 = d\theta^2 + \sin^2\theta \, d\phi^2$ may be written in terms of the charged Hodge-de Rham operator

$$\Delta \equiv *D*D + D*D*, \tag{B.2}$$

as

$$-\Box_{(s)} = \Delta \,, \tag{B.3}$$

where D is the charge-s gauge-covariant exterior derivative

$$D = d + is\cos\theta d\phi\,.\tag{B.4}$$

The spin-weighted harmonics may be constructed by starting with the four-dimensional scalar Laplacian on \mathbb{C}^2 , and then embedding the unit S^3 , viewed as a U(1) bundle over S^2 , in \mathbb{C}^2 . Introducing complex coordinates Z^a on \mathbb{C}^2 , the four-dimensional Laplacian is

$$\Box_4 = 4 \frac{\partial^2}{\partial Z^a \partial \bar{Z}_a} \,. \tag{B.5}$$

Clearly, if we define functions

$$f = T_{a_1 \cdots a_p}{}^{b_1 \cdots b_q} Z^{a_1} \cdots Z^{a_p} \bar{Z}_{b_1} \cdots \bar{Z}_{b_q} , \qquad (B.6)$$

where $T_{a_1\cdots a_p}{}^{b_1\cdots b_q}$ is symmetric in its upper and its lower indices, and traceless with respect to any contraction of upper and lower indices, then they will satisfy

$$\Box_4 f = 0. \tag{B.7}$$

Writing

$$Z^{1} = r e^{i(\psi + \phi)/2} \cos \frac{1}{2}\theta, \qquad Z^{2} = r e^{i(\psi - \phi)/2} \sin \frac{1}{2}\theta, \qquad (B.8)$$

the Euclidean metric on \mathbb{C}^2 is expressible as

$$ds_4^2 = dr^2 + r^2 \, d\Omega_3^2 \,, \tag{B.9}$$

where

$$d\Omega_3^2 = \frac{1}{4} (d\psi + \cos\theta \, d\phi)^2 + \frac{1}{4} (d\theta^2 + \sin^2\theta \, d\phi^2)$$
(B.10)

is the metric on the unit 3-sphere. The four-dimensional Laplacian is given by

$$\Box_4 = \frac{1}{r^3} \frac{\partial}{\partial r} \left(r^3 \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \Box_3, \tag{B.11}$$

where \Box_3 is the Laplacian on the unit S^3 . Noting from (B.6) and (B.8) that f takes the form

$$f = r^{p+q} e^{i(p-q)\psi/2} Y(\theta, \phi),$$
 (B.12)

then (B.7) and (B.11) imply that

$$\Box_3\left(e^{i(p-q)\psi/2} Y(\theta,\phi)\right) = -(p+q)(p+q+2) e^{i(p-q)\psi/2} Y(\theta,\phi).$$
(B.13)

From (B.10), \square_3 is given by

$$\frac{1}{4}\Box_3 = \frac{1}{\sin\theta} \frac{\partial}{\partial\theta} \left(\sin\theta \frac{\partial}{\partial\theta}\right) + \frac{1}{\sin^2\theta} \left(\frac{\partial}{\partial\phi} - \cos\theta \frac{\partial}{\partial\psi}\right)^2 + \frac{\partial^2}{\partial\psi^2},\tag{B.14}$$

and so if we define

$$p = \ell - s, \qquad q = \ell + s, \tag{B.15}$$

then (B.13) implies that $Y(\theta, \phi)$ satisfies

$$-\Box_{(s)} Y = [\ell(\ell+1) - s^2] Y, \qquad (B.16)$$

where $\Box_{(s)}$ is the charged scalar Laplacian on S^2 that we defined in equation (B.1). Up to an overall conventional normalisation, we see that $Y(\theta, \phi)$ constructed from (B.6) and (B.12) is nothing but a

spin-weighted spherical harmonic ${}_{s}Y_{\ell m}$. Since p and q in (B.6) are non-negative integers, and they are related to ℓ and s by (B.15), it follows that

$$\ell \ge |s| \,. \tag{B.17}$$

It is easily seen that the number of independent traceless symmetric tensors $T_{a_1 \cdots a_p}{}^{b_1 \cdots b_q}$ in (B.6) is equal to 1 + p + q, and hence we have constructed the $2\ell + 1$ spin-weighted spherical harmonics ${}_{s}Y_{\ell m}$ at level ℓ satisfying

$$-\Box_{(s)\ s}Y_{\ell m} = \left[\ell(\ell+1) - s^2\right]_s Y_{\ell m}, \qquad \ell \ge |s|, \qquad -\ell \le m \le \ell.$$
(B.18)

Note that s, ℓ and m are either all integers, or else all half-integers.

With the conventional normalisation, the spin-weighted spherical harmonics satisfy the relations

$$\mathcal{D}_{-s}Y_{\ell m} \equiv \left(\frac{\partial}{\partial\theta} + m\csc\theta + s\cot\theta\right)_{s}Y_{\ell m} = \sqrt{(\ell+s)(\ell+1-s)}_{s-1}Y_{\ell m},$$
$$\mathcal{D}_{+s-1}Y_{\ell m} \equiv \left(\frac{\partial}{\partial\theta} - m\csc\theta - (s-1)\cot\theta\right)_{s-1}Y_{\ell m} = -\sqrt{(\ell+s)(\ell+1-s)}_{s}Y_{\ell m}.$$
(B.19)

B.2 Vector spin-weighted harmonics

The spin-weighted vector harmonics are the eigenfunctions of the charged Hodge-de Rham operator (B.2) acting on 1-forms:

$$\Delta V = \lambda V , \qquad V = dy^i V_i . \tag{B.20}$$

Generically, these eigenfunctions can be constructed from the scalar spin-weighted harmonics ${}_{s}Y_{\ell m}$ (denoted simply as Y below) by writing

$$V = DY + \mu * DY, \tag{B.21}$$

where $D = d + i s \cos \theta \, d\phi$ is the gauge-covariant exterior derivative. We shall write the eigenvalues for the scalar spin-weighted harmonics, given by (B.18), simply as λ , so that

$$\Delta Y = \lambda Y, \qquad \lambda = \ell(\ell+1) - s^2,. \tag{B.22}$$

Noting that

$$D^2 = -is\Omega_2, \qquad (B.23)$$

where $\Omega_2 = \sin \theta \, d\theta \wedge d\phi$ is the volume form on the unit S^2 , that $D * D * (DY) = D\Delta Y = \lambda DY$, $*D*D(DY) = -is*D(*\Omega_2 Y) = -is*DY$ and that $\Delta * = *\Delta$, we see that

$$\Delta V = (\lambda + i\mu s)DY + (\mu\lambda - is)*DY.$$
(B.24)

Thus V is an eigenfunction, satisfying (B.20), if $\mu = \pm i$, and so generically we get two distinct vector eigenfunctions V^{\pm} with corresponding eigenvalues $\tilde{\lambda}_{\pm}$ from each scalar eigenfunction Y, where

$$V^{\pm} = DY \mp i * DY, \qquad \tilde{\lambda}_{\pm} = \lambda \pm s.$$
(B.25)

In terms of ℓ and s, these eigenvalues are given by

$$\tilde{\lambda}_{+} = (\ell + s)(\ell + 1 - s), \qquad \tilde{\lambda}_{-} = (\ell - s)(\ell + 1 + s).$$
(B.26)

Note that the vector harmonics V^{\pm} obey the complex duality conditions

$$*V^{\pm} = \pm iV^{\pm} \,. \tag{B.27}$$

A special case arises if the scalar eigenvalue λ is equal to s or -s. (Since λ is necessarily non-negative, the former can only arise if s is positive, and it implies $\ell = s$, while the latter arises if s is negative, and implies $\ell = -s$.) Calculating the norm of V^- , we find

$$\int *\bar{V}^- \wedge V^- = \int (*D\bar{Y} - i*D\bar{Y}) \wedge (DY + i*DY) = 2 \int (*DY \wedge D\bar{Y} - iDY \wedge D\bar{Y})$$
$$= 2 \int ((D*DY)\bar{Y} - i(D^2Y)\bar{Y}) = 2(\lambda - s) \int |Y|^2, \qquad (B.28)$$

and so $V^- = 0$ if $\lambda = s$. A similar calculation shows $V^+ = 0$ if $\lambda = -s$. Thus if $\lambda = s$ then the mode V^- , which from (B.25) would have had eigenvalue $\tilde{\lambda}_- = 0$, is absent. Similarly, if $\lambda = -s$ then V^+ , which would likewise have had eigenvalue $\tilde{\lambda}_+ = 0$, is absent.

In fact vector spin-weighted zero modes of Δ do arise, but they cannot be constructed from scalar harmonics in the manner described above. If $\Delta V = 0$ then integrating $*\bar{V}\Delta V$ over the sphere implies

$$\int (|D*V|^2 + |DV|^2) = 0 \tag{B.29}$$

and hence V is (gauge) closed and co-closed,

$$DV = 0, \qquad D*V = 0.$$
 (B.30)

We can project into the self-dual and anti-self-dual subspaces, and thus seek 1-forms V satisfying

$$*V = \pm iV, \qquad DV = 0. \tag{B.31}$$

Making the ansatz

$$V = e^{im\phi} \left(f d\theta + g d\phi \right), \tag{B.32}$$

where f and g are functions of θ , we find $*V = e^{im\phi} (-f\sin\theta \, d\phi + g\, \csc\theta \, d\theta)$ and hence the duality condition implies

$$g = \pm i f \sin \theta \,. \tag{B.33}$$

The condition DV = 0 implies $g' = if(m + s\cos\theta)$, and hence we obtain

$$f = c(\sin\theta)^{-1\pm s} \left(\tan\frac{1}{2}\theta\right)^{\pm m}.$$
(B.34)

Thus if $s \ge 1$ we obtain regular self-dual harmonics $(*V^+ = +iV^+)$ given by

$$V^{+} = (\sin\frac{1}{2}\theta)^{s-1+m} (\cos\frac{1}{2}\theta)^{s-1-m} e^{im\phi} (d\theta + i\,\sin\theta\,d\phi), \qquad -(s-1) \le m \le (s-1), \quad (B.35)$$

while if $s \leq -1$ we obtain regular anti-self-dual harmonics $(*V^- = -iV^-)$ given by

$$V^{-} = \left(\sin\frac{1}{2}\theta\right)^{-s-1-m} \left(\cos\frac{1}{2}\theta\right)^{-s-1-m} e^{im\phi} \left(d\theta - i\,\sin\theta\,d\phi\right), \qquad s+1 \le m \le -s-1.$$
(B.36)

In each case, these charge-s vector harmonics form an $\ell = |s| - 1$ representation of SU(2), as evidenced by the (2|s| - 1)-fold multiplet of m values.

In summary, each spin-weighted scalar harmonic Y with eigenvalue $\lambda = \ell(\ell + 1) - s^2$ and with with $\ell \ge |s| + 1$ gives rise to two spin-weighted vector harmonics, namely a self-dual harmonic V^+ with eigenvalue $\tilde{\lambda}_+ = (\ell + s)(\ell + 1 - s)$ for the charge-s Hodge-de Rham operator Δ , and an anti-self dual harmonic V^- with eigenvalue $\tilde{\lambda}_- = (\ell - s)(\ell + 1 + s)$. However, the lowest-level spin-weighted scalar harmonic, with $\ell = |s|$, gives rise to only one spin-weighted vector harmonic, namely V^+ if s is positive, or V^- if s is negative. The "missing" vector harmonic when $\ell = |s|$ would have been a zero-mode of Δ . In its place, a zero-mode harmonic satisfying $\Delta V = 0$ does occur, but it cannot be constructed from the scalar spin-weighted harmonics. It corresponds to $\ell = |s| - 1$, and therefore has multiplicity 2|s| - 1.

B.3 Spin- $\frac{1}{2}$ spin-weighted harmonics

We may define the spin-weighted spinor harmonics to be charged solutions of the Dirac equation in the monopole background. They may, in general, be constructed from the scalar spin-weighted harmonics, as we now describe. We first note that there exist two charged gauge-covariantly constant spinors on S^2 with the monopole background, satisfying $D\eta = 0$ where we now add a spin connection term to the gauge-covariant exterior derivative,

$$D = \nabla + i s \cos \theta \, d\phi \,, \qquad \nabla \equiv d + \frac{1}{4} \omega_{ab} \sigma^{ab} \tag{B.37}$$

and, when acting on η^{\pm} , $s = \pm \frac{1}{2}$. This can be seen from the integrability condition

$$0 = [D_i, D_j]\eta = \frac{1}{4}R_{ijk\ell}\sigma^{k\ell}\eta - i\,s\,\epsilon_{ij}\,\eta = i\,\epsilon_{ij}\,(\frac{1}{2}\sigma_3 - s)\eta \tag{B.38}$$

from which we see that there exist two solutions:

$$s = \frac{1}{2}: \quad D\eta_{+} = 0, \qquad \sigma_{3}\eta_{+} = \eta_{+}, s = -\frac{1}{2}: \quad D\eta_{-} = 0, \qquad \sigma_{3}\eta_{-} = -\eta_{-}.$$
(B.39)

Using the standard basis for the Pauli matrices σ_i , the solutions, we find that the gauge-covariantly constant spinors η_{\pm} are given by

$$\eta_{+} = \begin{pmatrix} 1\\0 \end{pmatrix}, \qquad \eta_{-} = \begin{pmatrix} 0\\1 \end{pmatrix}. \tag{B.40}$$

From these spinors, which are normalised so that $\bar{\eta}_+\eta_+ = \bar{\eta}_-\eta_- = 1$, we may construct the gaugecovariantly constant vector

$$U = \bar{\eta}_{-}\sigma^{i}\eta_{+}\partial_{i} = \frac{\partial}{\partial\theta} + i\csc\theta \frac{\partial}{\partial\phi}, \qquad (B.41)$$

which has charge s = 1. Its complex conjugate $\overline{U} = \partial/\partial\theta - i \csc\theta \partial/\partial\phi$ has charge s = -1. In fact U is the holormorphic (1,0)-form on the Kähler manifold S^2 , satisfying $J_i^j U_j = i U_i$, where $J_{ij} = \epsilon_{ij}$ is the Kähler form. Note that

$$\sigma^{i}\eta_{+} = U^{i}\eta_{-}, \qquad \sigma^{i}\eta_{-} = \bar{U}^{i}\eta_{+}.$$
(B.42)

The operators $U^i D_i$ and $\overline{U}^i D_i$ give precisely \mathcal{D}_+ and \mathcal{D}_- , defined in (B.19), when acting on ${}_{s}Y_{\ell m}$ and ${}_{s-1}Y_{\ell m}$ respectively. It is now clear why \mathcal{D}_+ and \mathcal{D}_- raise and lower the charge of the spinweighted scalar harmonics by one unit, since U^i and \overline{U}^i are gauge-covariantly constant vectors carrying +1 and -1 charge respectively. The solutions of the charged Dirac equation can be expressed in terms of chiral spinors ψ_+ and anti-chiral spinors ψ_- , satisfying

$$\sigma^i D_i \psi_+ = i \,\lambda_+ \,\psi_- \,, \qquad \sigma^i D_i \psi_- = i \,\lambda_- \,\psi_+ \,. \tag{B.43}$$

Since $(\sigma^i D_i)^2 \psi = D^i D_i \psi + (\tilde{s}\sigma_3 - \frac{1}{2})\psi$ where ψ is any spinor with charge \tilde{s} , we have

$$-D^{i}D_{i}\psi_{+} = (\lambda_{+}\lambda_{-} + \tilde{s} - \frac{1}{2})\psi_{+}, \qquad -D^{i}D_{i}\psi_{-} = (\lambda_{+}\lambda_{-} - \tilde{s} - \frac{1}{2})\psi_{-}.$$
(B.44)

The product $\lambda_+\lambda_-$ is therefore uniquely determined, as a function of \tilde{s} and ℓ , for each spinor eigenfunction of the second-order operator $D^i D_i$. The values of λ_+ and λ_- separately are not determined, but depend upon the choice of relative normalisation for ψ_+ and ψ_- in (B.43).

It is convenient to consider spinor eigenfunctions ψ_+ and ψ_- with charge $\tilde{s} = s - \frac{1}{2}$. We may construct these from the scalar spin-weighted harmonics ${}_{s}Y_{\ell m}$ discussed earlier by writing

$$\psi_{-} = \eta_{-} (_{s}Y_{\ell m}) , \qquad \psi_{+} = \eta_{+} (_{s-1}Y_{\ell m}) .$$
 (B.45)

Note these ψ_{\pm} are denoted by $\tilde{\eta}_{\pm}^{(\ell)}$ in (4.26). For brevity in notation, however, we shall continue to use the notation ψ_{\pm} instead in this section. Acting on these with the Dirac operator, we find, using (B.19), (B.41) and (B.42), that

$$\sigma^{i}D_{i}\psi_{-} = \sigma^{i}\eta_{-}D_{i\,s}Y_{\ell m} = \eta_{+}\bar{U}^{i}D_{i\,s}Y_{\ell m}$$

$$= \eta_{+}\left(\frac{\partial}{\partial\theta} + m\csc\theta + s\cot\theta\right)_{s}Y_{\ell m} = \eta_{+}\sqrt{(\ell+s)(\ell+1-s)}_{s-1}Y_{\ell m},$$

$$\sigma^{i}D_{i}\psi_{+} = \sigma^{i}\eta_{+}D_{i\,s-1}Y_{\ell m} = \eta_{-}U^{i}D_{i\,s-1}Y_{\ell m} \qquad (B.46)$$

$$= \eta_{-}\left(\frac{\partial}{\partial\theta} - m\csc\theta - (s-1)\cot\theta\right)_{s-1}Y_{\ell m} = -\eta_{-}\sqrt{(\ell+s)(\ell+1-s)}_{s}Y_{\ell m},$$

and hence

$$\sigma^{i} D_{i} \psi_{-} = \sqrt{(\ell+s)(\ell+1-s)} \psi_{+}, \qquad \sigma^{i} D_{i} \psi_{+} = -\sqrt{(\ell+s)(\ell+1-s)} \psi_{-}.$$
(B.47)

It is worth remarking that there is an alternative procedure that in general constructs the charged spin- $\frac{1}{2}$ harmonics from scalar harmonics, in which only one of the gauge-covariantly constant spinors is required. For example, using only η_{-} we can construct the negative-chirality spin- $\frac{1}{2}$ harmonics ψ_{-} as in the first equation in (B.45), while for the positive-chirality harmonics we take

$$\psi'_{+} = \sigma^{i} \eta_{-} D_{i s} Y_{\ell m} \,. \tag{B.48}$$

A straightforward calculation shows that

$$\sigma^{i} D_{i} \psi'_{+} = -(\ell + s)(\ell + 1 - s) \eta_{-s} Y_{\ell m} .$$
(B.49)

The harmonics ψ'_+ are in general proportional to the harmonics ψ_+ given in (B.45). However, the charge $\frac{1}{2}$ harmonic $\psi_+ = \eta_+$ itself (which is a zero mode of the Dirac operator) cannot be constructed using (B.48), since it would require taking s = 1 and $\ell = 0$, for which ${}_{s}Y_{\ell m}$ does not exist.

It might also seem that charge $-s - \frac{1}{2}$ zero modes ψ'_+ would be obtained if s were negative and $\ell = -s$. However, calculating the norm of ψ'_+ , we find

$$\int_{S^2} |\psi'_+|^2 \sqrt{g} d^2 x = (\ell + s)(\ell + 1 - s) \int_{S^2} |_s Y_{\ell m}|^2 \sqrt{g} d^2 x , \qquad (B.50)$$

and thus ψ'_+ would actually be identically zero if $\ell = -s$. These putative zero modes are in fact not obtained by the construction for ψ_+ in (B.45) either, since this would require the use of scalar harmonics with ℓ smaller than the magnitude of their spin weight.

B.4 Spin- $\frac{3}{2}$ spin-weighted harmonics

The general spin- $\frac{3}{2}$ harmonics η_i can be decomposed into chiral and antichiral projections η_i^{\pm} satisfying

$$\sigma^i D_i \eta_j^+ = \lambda_+ \eta_j^-, \qquad \sigma^i D_i \eta_j^- = \lambda_- \eta_j^+.$$
(B.51)

Each chiral projection admits a decomposition of the form

$$\eta_i^{\pm} = \sigma_i \psi^{\mp} + \eta_{\{i\}}^{\pm} + \tilde{\eta}_i^{\pm} \,, \tag{B.52}$$

where $\eta_{\{i\}}^{\pm}$ is longitudinal and gamma traceless, $\sigma^i \eta_{\{i\}}^{\pm} = 0$, and $\tilde{\eta}_i^{\pm}$ is transverse and gamma traceless, satisfying $D^i \tilde{\eta}_i^{\pm} = 0$ and $\sigma^i \tilde{\eta}_i^{\pm} = 0$. We can write $\eta_{\{i\}}^{\pm}$ in terms of spin- $\frac{1}{2}$ modes η^{\pm} as

$$\eta_{\{i\}}^{\pm} = 2D_i\psi^{\pm} - \sigma_i\sigma^j D_j\psi^{\pm} = (D_i \mp i\,\epsilon_i{}^j\,D_j)\psi^{\pm}\,. \tag{B.53}$$

In fact $\eta_{\{i\}}^{\pm}$ can alternatively be written in terms of the vector harmonics V^{\pm} constructed from scalar harmonics as in (B.25), by taking

$$\eta_{\{i\}}^{\pm} = V_i^{\pm} \, \eta^{\pm} \,. \tag{B.54}$$

The gamma-tracelessness of $\eta_{\{i\}}^{\pm}$ follows immediately from the fact that V_i^{\pm} and $\sigma^i \eta^{\pm}$ are either both self-dual or both anti-self dual. The charge carried by $\eta_{\{i\}}^{\pm}$ will, of course, be equal to $s \pm \frac{1}{2}$, where s is the charge of V_i^{\pm} .

The transverse traceless spin- $\frac{3}{2}$ harmonics $\tilde{\eta}_i$ can be constructed in the same way, and are given by (B.54) except that now, V_i^{\pm} are the self-dual vector harmonics (B.35) or the anti-self dual harmonics (B.36) that cannot be constructed from scalar harmonics. Since such V_i^{\pm} vectors arise only when $s \ge 1$ or $s \le -1$ respectively, the transverse traceless spin- $\frac{3}{2}$ harmonics $\tilde{\eta}_i^{\pm}$ arise only for charges $\tilde{s} \ge \frac{3}{2}$ or $\tilde{s} \le -\frac{3}{2}$ respectively.

All necessary properties of the spin- $\frac{3}{2}$ harmonics follow from the properties of the lower-spin harmonics that we discussed previously.

C Spin Projection Operators

The well known spin projector operators associated with a second rank symmetric tensor field are given by [16]

$$\mathcal{P}^{2}_{\mu\nu,\rho\sigma} = \frac{1}{2} (\theta_{\mu\rho}\theta_{\nu\sigma} + \theta_{\mu\sigma}\theta_{\nu\rho} - \frac{2}{3}\theta_{\mu\nu}\theta_{\rho\sigma}) ,$$

$$\mathcal{P}^{1}_{\mu\nu,\rho\sigma} = \frac{1}{2} (\theta_{\mu\rho}\omega_{\nu\sigma} + \theta_{\mu\sigma}\omega_{\nu\rho} + \theta_{\nu\rho}\omega_{\mu\sigma} + \theta_{\nu\sigma}\omega_{\mu\rho}) ,$$

$$\mathcal{P}^{(0,s)}_{\mu\nu,\rho\sigma} = \frac{1}{3}\theta_{\mu\nu}\theta_{\rho\sigma} ,$$

$$\mathcal{P}^{(0,\omega)}_{\mu\nu,\rho\sigma} = \omega_{\mu\nu}\omega_{\rho\sigma} ,$$

(C.1)

where

$$\theta_{\mu\nu} = \eta_{\mu\nu} - \Box^{-1}\partial_{\mu}\partial_{\nu} , \qquad \omega_{\mu\nu} = \Box^{-1}\partial_{\mu}\partial_{\nu} .$$
(C.2)

Similarly, the spin projector operators associated with vector-spinor field take the form

$$P_{\mu\nu}^{3/2} = \theta_{\mu\nu} - \frac{1}{3}\theta_{\mu}\theta_{\nu} ,$$

$$(P_{11}^{1/2})_{\mu\nu} = \frac{1}{3}\theta_{\mu}\theta_{\nu} , \qquad (P_{12}^{1/2})_{\mu\nu} = \frac{1}{\sqrt{3}}\theta_{\mu}\omega_{\nu} ,$$

$$(P_{21}^{1/2})_{\mu\nu} = \frac{1}{\sqrt{3}}\omega_{\mu}\theta_{\nu} , \qquad (P_{22}^{1/2})_{\mu\nu} = \omega_{\mu}\omega_{\nu} ,$$
(C.3)

where

$$\theta_{\mu} = \theta_{\mu\nu}\gamma^{\nu} , \qquad \omega_{\mu} = \omega_{\mu\nu}\gamma^{\nu} .$$
(C.4)

References

- E. Bergshoeff, E. Sezgin and A. Van Proeyen, Superconformal tensor calculus and matter couplings in six-dimensions, Nucl. Phys. B 264 (1986) 653, Erratum-ibid. B 598 (2001) 667.
- [2] F. Coomans and A. Proeyen, Off-shell N = (1,0), D = 6 supergravity from superconformal methods, JHEP **1102** (2011) 049, Erratum-ibid. **1201** (2012) 119, arXiv:1101.2403 [hep-th].
- [3] E. Bergshoeff, A. Salam and E. Sezgin, Supersymmetric R² actions, conformal invariance and Lorentz Chern-Simons term in six-dimensions and ten-dimensions, Nucl. Phys. B 279 (1987) 659.
- [4] E. A. Bergshoeff, J. Rosseel and E. Sezgin, *Off-shell* D = 5, N = 2 *Riemann squared super*gravity, Class. Quant. Grav. **28** (2011) 225016, arXiv:1107.2825 [hep-th].
- [5] E. Bergshoeff, F. Coomans, E. Sezgin and A. Van Proeyen, *Higher derivative extension of 6D chiral gauged supergravity*, arXiv:1203.2975 [hep-th].
- [6] A. Salam and E. Sezgin, Chiral compactification on Minkowski× S^2 of N = 2 Einstein-Maxwell supergravity in six-dimensions, Phys. Lett. B 147 (1984) 47.
- [7] S. Deser and C. Teitelboim, Supergravity has positive energy, Phys. Rev. Lett. **39** (1977) 249.
- [8] D. G. Boulware, S. Deser and K. S. Stelle, Energy and supercharge in higher derivative gravity, Phys. Lett. B 168 (1986) 336.
- [9] H. J. Kim, L. J. Romans and P. van Nieuwenhuizen, The mass spectrum of chiral N = 2, D = 10 supergravity on S^5 , Phys. Rev. D **32** (1985) 389.
- [10] P. Hoxha, R. R. Martinez-Acosta and C. N. Pope, Kaluza-Klein consistency, Killing vectors, and Kahler spaces, Class. Quant. Grav. 17, 4207 (2000), hep-th/0005172.
- [11] G. W. Gibbons and C. N. Pope, Consistent S² Pauli reduction of six-dimensional chiral gauged Einstein-Maxwell supergravity, Nucl. Phys. B 697 (2004) 225, hep-th/0307052.
- [12] M. Cvetic, G. W. Gibbons and C. N. Pope, A String and M-theory origin for the Salam-Sezgin model, Nucl. Phys. B 677 (2004) 164, hep-th/0308026.

- [13] H. Lu, C. N. Pope and E. Sezgin, Massive three-dimensional supergravity from $R + R^2$ action in six dimensions, JHEP **1010** (2010) 016, arXiv:1007.0173 [hep-th].
- [14] E. A. Bergshoeff and M. de Roo, The quartic effective action of the heterotic string and supersymmetry, Nucl. Phys. B 328 (1989) 439.
- [15] H. Lü and Y. Pang, On hybrid (topologically) massive supergravity in three dimensions, JHEP 1103, 050 (2011), arXiv:1011.6212 [hep-th].
- [16] P. Van Nieuwenhuizen, Supergravity, Phys. Rept. 68 (1981) 189.