

INDEX-CATALOGUE OF MEDICAL AND VETERINARY ZOOLOGY


SUPPLEMENT 22, PART 4

PARASITE-SUBJECT CATALOGUE
PARASITES: NEMATODA AND ACANTHOCEPHALA



UNITED STATES
DEPARTMENT OF
AGRICULTURE

PREPARED BY
SCIENCE AND
EDUCATION
ADMINISTRATION



SINCLAIR COMPARATIVE MEDICINE RESEARCH
ROUTE 3
COLUMBIA, MISSOURI

INDEX-CATALOGUE OF MEDICAL AND VETERINARY ZOOLOGY

SUPPLEMENT 22, PART 4

PARASITE-SUBJECT CATALOGUE
PARASITES: NEMATODA AND ACANTHOCEPHALA

By

SHIRLEY J. EDWARDS, In Charge

MARTHA W. HOOD, Zoologist

JUDITH H. SHAW, Zoologist

JANE D. RAYBURN, Technical Information Specialist

MARGIE D. KIRBY, Technical Information Specialist

DEBORAH T. HANFMAN, Technical Information Specialist

JUDITH A. ZIDAR, Technical Information Specialist



UNITED STATES
DEPARTMENT OF
AGRICULTURE

PREPARED BY
SCIENCE AND
EDUCATION
ADMINISTRATION

November 1980

THE UNIVERSITY OF CHICAGO
DEPARTMENT OF CHEMISTRY
GENERAL CHEMISTRY

LECTURE NOTES
BY

PROFESSOR
JAMES H. COOPER
AND
ASSISTANT PROFESSOR
ROBERT M. WAYNE

PREFACE

The Index-Catalogue of Medical and Veterinary Zoology is an index to the world's literature on animal parasites of animals, including man. The Catalogue is distributed to qualified individuals and libraries throughout the world without charge. It has been maintained in cumulative files since 1892. Only the Author Catalogue has been published in its entirety. A revision of the Author Catalogue of the Index-Catalogue of Medical and Veterinary Zoology, consisting of Parts 1 to 18, was published during the period 1932-52. Beginning in 1953, a series of supplements designed to publish the backlog was initiated. This was completed with Supplement 6, published in 1956. From 1956 to 1964, supplements covering authors A to Z were issued on an annual basis.

Beginning with Supplement 15, the Parasite-Subject Catalogues, containing indices to the author references, have been issued. The Author Catalogues of Supplements 15-21 continued the format of previous supplements. Users should note that for each reference in the Author Catalogues of these supplements the author(s) plus the date and letter (e.g., Smith, J.; and Doe, L., 1978 b) are the key to all items in the Parasite-Subject Catalogues derived from that reference. In other words, when using the Parasite-Subject Catalogues of Supplements 15-21, it is necessary to consult the Author Catalogue of the corresponding supplement for complete bibliographic information.

Commencing with Supplement 22, basic bibliographic information is included with each entry in Parts 2-7. It should be emphasized, however, that it will still be useful to consult the Author Catalogue for a variety of other information that may be found there: Title of the reference, translated title, language of text and summaries, issue date, library from which the original may be obtained, published corrections, related references by the same author, and other miscellaneous information.

Each supplement consists of the following parts:

- Part 1, Authors: A-Z
- Part 2, Parasite-Subject Catalogue: Parasites: Protozoa
- Part 3, Parasite-Subject Catalogue: Parasites: Trematoda and Cestoda
- Part 4, Parasite-Subject Catalogue: Parasites: Nematoda and Acanthocephala
- Part 5, Parasite-Subject Catalogue: Parasites: Arthropoda and Miscellaneous Phyla
- Part 6, Parasite-Subject Catalogue: Subject Headings and Treatment
- Part 7, Parasite-Subject Catalogue: Hosts

Users should bear in mind that this is an Index-Catalogue, not a treatise, and should not expect to find reasons for any given entry. Nor does citing of synonymy mean that it is necessarily correct. The same statement holds for hosts, locations, localities, authorship of taxa, designation of new taxa, etc. These items are cited as given by the author(s) of the publication being indexed.

The information included in any given supplement represents only the publications that have been indexed in that supplement; and therefore, exclusion of, or limited entries for, any given author or parasite has no significance. No pretension is made for completeness, and assistance in correcting errors or obtaining additional information is appreciated. Reprints of papers on parasitology are requested.


EXPLANATORY NOTE

Author Catalogue

The Author Catalogue (Part 1 of each supplement) contains full bibliographic information for each publication indexed during the compilation of that supplement. A symbol for the library from which the original publication may be obtained is given at the end of each entry, e.g., Wa, Wm, Wc, etc. A key to these library symbols may be found in Supplements 10 and 20. A list of serial abbreviations new to our files is published at the beginning of each Author Catalogue.

Parasite Catalogues

The Parasite Catalogues (Parts 2-5 of each supplement) are divided by parasite phyla (Protozoa, Trematoda, etc.). They are arranged alphabetically by genera, parasitic diseases, and higher taxa and then alphabetically by species within genera. Entries under each heading are in turn arranged alphabetically by authors and then chronologically for each author. Each entry consists of the name of the parasite or parasitic disease, the author(s) of the publication, date, abbreviated title of the publication, volume, number, inclusive pages, and a subheading. Illustrations of parasites are indicated by the word *illus.* following the name of the parasite.



A variety of information is found indented beneath the author line(s) of each entry: Classification, hosts, synonymy, keys, treatment, etc. Subheadings are guides to the subject matter of the publication.

- (1) **Classification:** In entries based on systematic articles, the subheading may give the higher taxa in which the taxon has been placed or it may list the lower taxa included in a higher taxon.
- (2) **Hosts:** The only hosts recorded are those that pertain directly to the author's own work. Scientific host names are used unless the author gives only common names, in which case the host names are given exactly as in the original publication.

However, when host common names are in Cyrillic alphabet languages, host Latin names are assigned and listed instead of the common name; these are in square brackets [].

Locations of parasites in or on hosts are given in parentheses (). Where a host-parasite relationship is well known, a host may be given under a parasite name and not in the Host Catalogue; this applies particularly to parasites of medical and veterinary importance and of worldwide distribution. A + before the host name on the parasite entry means that no host entry was made for this particular reference.

- (3) **Synonymy:** Usually only those synonyms which the author indicates as new, or which are new to the files of the Index-Catalogue of Medical and Veterinary Zoology, are given.
- (4) **Keys:** The subheading "key" indicates that the name is included in a taxonomic key.
- (5) **Treatment:** When there are several antiparasitic agents mentioned in a publication, a general term is used in the subheading, e.g., anthelmintics, insecticides, protozoacides. However, in the Treatment Catalogue, all agents tested by the investigator(s) are listed.
- (6) **Geographic Distribution:** When there are multiple hosts and geographic localities, the appropriate locality is

recorded after each host name; when the hosts of a parasite are all from one locality, they are recorded as "all from" this locality.

- (7) **Other Subject Matter:** Phrases indicate other subject matter discussed (e.g., immunity, metabolism, morphology, etc.).

Subject Headings Catalogue

The Subject Headings Catalogue (the first section of Part 6 of each supplement) is an alphabetic arrangement of entry terms from a controlled list of subject headings. Each entry consists of the subject heading, bibliographic information, and a subheading reflecting the information contained in the paper. Subject headings with numerous entries are separated into alphabetized subdivisions, e.g.,

Immunity
Immunity, Agglutination
Immunity, Allergy

Treatment Catalogue

In the Treatment Catalogue (a section of Part 6 of each supplement), all entries referring to one antiparasitic agent are grouped under one heading (regardless of the name used by the investigator) and are then listed alphabetically by author. Other names for the same agent are cross-referenced to the name used for filing. When generic and chemical names are available, preference is given to those names as headings rather than to trade names or code numbers and letters. Code number designations for compounds are entered in the Number Index in numerical order and cross-referenced to the name under which they are listed in the alphabetical section. Salts of a compound are usually grouped together, e.g., piperazine adipate, piperazine citrate, etc., are all listed under Piperazine. Sometimes verifying synonymy of drug names is impossible; consequently, groupings and cross-references are not always authenticated although as many as possible have been checked with reliable sources. In some instances, the cross-references are based entirely on information in papers indexed and verification was not possible. Foreign language terminology has been anglicized

where feasible. Chemosterilants, Molluscicides, and Repellents are entered under these three collective headings and not under the individual chemical. The format is the same as the parasite entries: Entry term (in this case, drug name), bibliographic information, and subheading.

Host Catalogue

The Host Catalogue (Part 7 of each supplement) is arranged alphabetically by genera, common names, and higher taxa and then alphabetically by species within genera. Nominate subspecies are interfiled with the species. Entries under each heading are in turn arranged alphabetically by author(s) and then chronologically for each author. The format is the same as in the other Catalogues, i.e., entry term (in this case, host name), bibliographic information, and subheading. Indented beneath the author line(s) of each host entry are all the parasites of a particular phylum that were reported from this host in the paper in question. Body locations of these parasites will be found in parentheses () either in the subheading or with the host name. Experimental infection is reported as such. When there are multiple parasites and geographic localities, the appropriate locality is recorded after each parasite name; when the parasites from this host are all from one locality, they are recorded as "all from" this locality. When authors use only common names of hosts, scientific names are cautiously supplied from authoritative sources after careful consideration. Cross-references from the common name used by the author to the scientific name supplied by the Index-Catalogue are filed among the host entries. Such supplied names are given in square brackets []. If a scientific name cannot be supplied, English common names are used. Scientific names or English common names are always supplied for common names in Cyrillic alphabet languages, and no cross-references are made. Surveys of parasites of humans and domestic animals are often indexed under geographic headings and entered in Part 6, Subject Headings, in addition to appearing in the Host Catalogue. In this case, all parasite phyla are grouped under the same host entry, and individual parasite entries are not included in the Parasite Catalogue.

Visitors are welcome to come to the Animal Parasitology Institute to use the cumulative files. Arrangements should be made in advance for lengthy visits.

All correspondence should be addressed to:

Index-Catalogue of Medical and Veterinary Zoology
Animal Parasitology Institute
USDA, SEA-AR, BARC-East, Building 1180
Beltsville, Maryland 20705 U.S.A.

It is hoped that these Catalogues will serve as a useful tool to workers in the field of parasitology. Users are requested to preserve the Catalogues, since they are not designed for general distribution and the edition is limited.

The compilers thank the staffs of the Technical Information Systems of the Science and Education Administration, the National Library of Medicine, and all other libraries who have aided us invaluablely by making publications available to us.

Trade names are used in this publication solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture or an endorsement by the Department over other products not mentioned.

- Abbreviata Travassos, 1920, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key
- Abbreviata sp. 2, larvae Chabaud, 1954
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Pachyscelis banghaasi
Stalagmoptera incostata
Pelorocnemis punctata
Somocoelia pinguis
Bioramix constricta
Zophosis punctata deflexa
Dissonomus sp.
all from Tadzhik SSR [and/or] Uzbek SSR
- Abbreviata sp.
King, S. R.; and Babero, B. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 241-248
Dipodomys merriami (stomach): Nevada
- Abbreviata sp.
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Pisterotarsa gigantea subsp. zoubkoffi:
Turkmenia
- Abbreviata caucasica (von Linstow 1902)
File, S. K.; McGrew, W. C.; and Tutin, C. E. G., 1976, J. Parasitol., v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces):
Gombe National Park, Tanzania
- Abbreviata caucasica, illus.
McConnell, E. E.; et al., 1974, Onderstepoort J. Vet. Research, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (stomach, small intestine):
Kruger National Park, Transvaal
- Abbreviata caucasica, illus.
Seureau, C., 1977, J. Invert. Path., v. 29 (2), 240-241
Abbreviata caucasica-infected Locusta migratoria, cytopathological accumulation of microtubules in gut epithelial cells, structure protects nematode from host reaction so that it can complete development when ingested by final host
- Abbreviata caucasica, illus.
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in Locusta migratoria (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Abbreviata (Abbreviata) turkomanica Andruschko et Markov, 1956, illus.
Annaev, Dzh.; and Mushkambarova, M. G., 1975, Izvest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk (5), 81-87
Abbreviata turkomanica, description, life cycle
Pisterotarsa gigantea zoubkoffi: Central Karakum (Ak-Molla)
Phrynocephalus helioscopus (stomach mucous membrane) (exper.)
Eremias velox (stomach) (exper.)
Agama sanguinolenta (mucous membrane of esophagus and stomach) (exper.)
Phrynocephalus mystaceus (mucous membrane of esophagus and stomach) (exper.)
Varanus griseus (stomach mucous membrane) (exper.)
- Acanthocheilidae Wuelker, 1929
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridoidea
key; key to genera
includes: Acanthocheilus; Metanisakis; Pseudanisakis
- Acanthocheilonema [? n. rank]
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of Dipetalonema; key
tod: D. (A.) dracunculoides (Cobbold, 1870)
- Acanthocheilonema perstans, illus.
Charters, A. D.; et al., 1972, Med. J. Australia, v. 1 (6), 268-271
Loa loa and Acanthocheilonema perstans adult worms recovered from calabar swellings in persons who previously had been employed in Nigeria, public health importance in possible transfer of disease entity from endemic areas: Western Australia
- Acanthocheilonema perstans
Katamine, D.; et al., 1967, Nettare Igaku (Trop. Med.), v. 9 (3), 143-157
Acanthocheilonema perstans, Wuchereria bancrofti, prevalence survey in area residents: Tanzania
- Acanthocheilonema perstans (Manson, 1891)
Sasa, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 3-48
human filariasis in the Americas, extensive review, epidemiology, geographic distribution, mosquito vectors, control measures, literature review
- Acanthocheilus Molin, 1858
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Acanthocheilidae
key
- Acanthospiculum jakutensis
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
also given in Addendum as O. jakutensis; possibly misprint or possibly transfer to Onchocerca

- Ackertia* Vaz, 1934
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
key
- Ackertia dorsti* Bain et Hocquet, 1968
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
as syn. of *Ackertia finlayi* (Mazza et Fiora, 1932)
- Ackertia finlayi* (Mazza et Fiora, 1932), illus.
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
synonymy
- Ackertia globulosa* sp. n., illus.
Muller, R. L.; and Nelson, G. S., 1975, *J. Parasitol.*, v. 61 (4), 606-609
Lemniscomys striatus striatus (pulmonary arteries)
Haemaphysalis leachi (nat. and exper.)
Otomys angoniensis (pulmonary arteries)
Aethomys kaiseri (pulmonary arteries)
Tatera robusta (pulmonary arteries)
all from Machakos and Nairobi areas of Kenya
- Ackertia globulosa*
Bianco, A. E., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (4), 429 [Demonstration]
Ackertia globulosa in *Meriones unguiculatus*, a new rodent filaria as possible model for studying diagnostic techniques and chemotherapy for onchocerciasis, *Haemaphysalis leachi* apparently natural vector for *Ackertia globulosa*
Meriones unguiculatus (exper.)
Haemaphysalis leachi (nat. and exper.)
Lemniscomys striatus
Tatera robusta
Otomys angoniensis
Aethomys kaiseri
all from Machakos, Kenya
- Ackertia globulosa*
Bianco, A. E.; and Muller, R. L., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (5), 383 [Demonstration]
Haemaphysalis leachii leachii as vector of *Ackertia globulosa* for rodent hosts, tick attachment and adaptations in both tick and nematode life cycles which enable tick to serve as vector
- Acuaria Bremser*, 1811, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Acuaria* sp.
Kayton, R. J.; and Schmidt, G. D., 1975, *J. Helminth.*, v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Acuaria anthuris*
Seureau, C.; and Quentin, J. C., 1977, *Ann. Parasitol.*, v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Acuaria anthuris*
Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Acuaria attenuata* (Rud., 1819) Railliet, Henry et Sisoff, 1912, illus.
Jaron, W., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 137-152
description, helminth fauna of adult swallows just returning from migration compared with young birds; dynamics of infection, species composition of helminths, various stages of nesting season
Hirundo rustica
Delichon urbica
Riparia riparia
(underneath the horny lining of the gizzard of all): all from Poland
- Acuaria depressa*
Seureau, C.; and Quentin, J. C., 1977, *Ann. Parasitol.*, v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Acuaria gruvelli* Gendre, 1913
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 267-271
Bubo bubo
Garrulus glandarius
all from Huesca, Alto Aragon
- Acuaria (Cheilospirura) hamulosa* (Diesing, 1851) Railliet et al., 1912
Bali, H. S.; and Kalra, I. S., 1975, *J. Research, Punjab Agric. Univ.*, v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Acuaria (Cheilospirura) hamulosa* (Diesing, 1851)
Fabiyyi, J. P., 1972, *Bull. Epizoot. Dis. Africa*, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria

- Acuaria (Cheilospirura) hamulosa Diesing, 1851
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagridis galeata (under gizzard lining): Vom area, Benue Plateau State, Nigeria
- Acuaria hamulosa
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Adesmia servillei schatzmayri
Trigonoscelis punctipleuris
all from Turkmenia
- Acuaria quiscula Williams, 1929
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coerulescens (under gizzard lining): Florida
- Acuaria (Dispharynx) spiralis (Molin, 1858)
Railliet, Henry and Sizoff, 1912
Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Acuaria (Dispharynx) spiralis (Molin, 1858)
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Acuaria (Dispharynx) spiralis Molin, 1858
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagridis galeata (proventriculus): Vom area, Benue Plateau State, Nigeria
- Acuaria spiralis
Gogoi, A. R., 1975, Kerala J. Vet. Sc., v. 5 (2), 131-134
fowl: Assam
- Acuaria spiralis
Singh, N. B.; Tewari, H. C.; and Deo, P. G., 1975, Ceylon Vet. J., v. 23 (3-4), 61 [Letter]
Acuaria spiralis, poultry, histopathology, pathogenicity in proventriculus and gizzard
- Acuariidae (Railliet, Henry & Sisoff, 1912, subfam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuarioidea
key to subfams.
includes: Acuariinae; Seuratiinae; Schistophinae
- Acuariinae Railliet, Henry, & Sisoff, 1912
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariidae
key; key to genera
includes: Paracuaria; Skrjabinocerca; Acuaria; Cheilospirura; Chordatortilis; Synacuaria; Pectinospirura; Chordocephalus; Chevreuxia; Skrjabinoclava; Echinuria; Stammerinema; Synhimantus; Desportesius; Cosmocephalus; Sexansocara
- Acuarioidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Acuarioidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
- Acuarioidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirurida
includes: Acuariidae
- Adenophorea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Nematoda
key
includes: Enoplida
- Adenophorea (=Aphasmidia) Chitwood, 1958
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Nematoda
includes: Enoplia; Chromadoria
- Adolpholutzia Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Aelurostrongylus abstrusus larvae
Ash, L. R., 1976, Rev. Biol. Trop., v. 24 (1), 163-174
Vaginulus plebeius: market gardens, near Noumea, New Caledonia
- Aelurostrongylus abstrusus, illus.
Clinton, R. L.; Stump, F. J.; and Wiggers, K. L., 1976, Feline Pract., v. 6 (2), 45-46
Aelurostrongylus abstrusus, 9-week-old kitten, levamisole hydrochloride, good results; sodium iodide, no effect on larvae; diethyl-carbamazine citrate, reduced number of larvae
- Aelurostrongylus abstrusus
Gregory, G. G.; and Munday, B. L., 1976, Austral. Vet. J., v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island

- Aelurostrongylus abstrusus*, *illus.*
Romero-Rodriguez, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 373-375
description of larval form
Felis catus domestica (nodulos subpleurales):
Granada (Spain)
- Aelurostrongylus abstrusus*, *illus.*
Scott, D. W., 1973, *Cornell Vet.*, v. 63 (3), 483-500
Aelurostrongylus abstrusus, cats (feces),
case reports, successful treatment with
L-tetramisole; literature review
- Aelurostrongylus abstrusus*
West, B.; Wilson, P.; and Hatch, C., 1977,
J. Helminthol., v. 51 (3), 210-211
Acinonyx jubatus (faeces): gardens of the
Royal Zoological Society of Ireland, Dublin
- Aelurostrongylus falciformis* (Schlegel, 1933)
Wetzel, 1938
Shakhmatova, V. I., 1966, *Trudy Gel'mint.*
Lab., *Akad. Nauk SSSR*, v. 17, 277-289
Meles meles
Gulo gulo
Martes martes
(lungs of all): all from Karelia
- Afrochona* nov. gen.
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Rhabdochoniidae, Rhabdochoninae; *tod*: *A. camerounensis* sp. n.
- Afrochona camerounensis* sp. n. (*tod*), *illus.*
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Aphyosemion camerounensis (*intestin*): Cameroon, Olounou
- ? *Agamerms saldulae* sp. n., *illus.*
Rubzov, I. A., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 97-100
"The question mark before the generic name has been put because no Mermithidae larvae from bugs were cultivated to the adult helminths and the inclusion of our larvae to the genus *Agamerms* might arise some doubt." *Saldula saltatoria*, larva: valley of the Zomnica river, Poland
- Agamerms pachysoma* (v. Linstow), *illus.*
Poinar, G. O., jr., 1976, *J. Parasitol.*, v. 62 (5), 843-844
infection in *Gumaga griseolum* (gut wall, skeletal muscles, fat body) represents paratenic host in life cycle
Culex pipiens (exper.)
- Agamospirura* sp. I
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Blaps fausti bactriana
Pachyscelis laevicollis
Adesmia planidorsis
Pachyscelis banghaasi
Trigonoscelis gemmulata
all from Tadzhik SSR [and/or] Uzbek SSR
- Agamospirura* sp. II
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Adesmia gebleri
A. planidorsis
Blaps fausti bactriana
Pachyscelis laevicollis
all from Tadzhik SSR [and/or] Uzbek SSR
- Agamospirura* sp. III
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Prosodes vincens: Tadzhik SSR
- Agamospirura* sp. IV
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Pisterotarsa kiritschenkoi
Adesmia planidorsis
all from Tadzhik SSR [and/or] Uzbek SSR
- Agamospirura* sp. 1
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Pisterotarsa gigantea subsp. *zoubkoffi*: Turkmenia
- Agamospirura* sp. 2
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Pisterotarsa gigantea subsp. *zoubkoffi*: Turkmenia
- Agamospirura* sp. 3
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Trigonoscelis punctipleuris: Turkmenia
- Agamospirura* sp. 4
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Cyphostete komarovi: Turkmenia
- Agamospirura* sp. 5
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Pisterotarsa gigantea subsp. *zoubkoffi*
Trigonoscelis gigas
all from Turkmenia
- Agamospirura* sp. 6
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Adesmia servillei schatzmayri
Trigonoscelis punctipleuris
Pisterotarsa gigantea subsp. *zoubkoffi*
P. kessleri
all from Turkmenia
- Agamospirura* sp. 7
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Adesmia gebleri: Turkmenia
- Agamospirura* sp. 8
Mushkambarova, M. G., 1973, *Ekol. Nasekom. Turkmen.* (Tashliev), 20-35
Pisterotarsa gigantea subsp. *zoubkoffi*: Turkmenia

- Agamospirura sp. 9
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Pisterotarsa kessleri: Turkmenia
- Agamospirura sp.
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes (stomach): Karelia
- Agamospirura, nec Skarbilovitsch, 1946, Henry et Sisoff, 1912, n. syn.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
as syn. of Physocephalus sexualatus (Molin, 1860) larvae
- Agrachanus Tichomirova in Skrjabin et al., 1971 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of Skrjabillanus Shigin & Shigina, 1958
- Agriostomum equidentatum
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 195-198
Antidorcas marsupialis (colon): Mountain Zebra National Park near Cradock, Cape Province
- Agriostomum gorgonis Le Roux, 1929
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of Syncerus caffer, some pathological findings, age of host
Syncerus caffer (small and large intestine): Kruger National Park
- Agriostomum gorgonis Le Roux
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Alcelaphus buselaphus cokei (digestive tract)
Connochaetes taurinus (large intestine)
all from Kenya
- Agriostomum vryburgi Railliet, 1902, illus.
Graber, M.; and Turpin, M., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (1), 23-30
Agriostomum vryburgi, description, localization, pathological effects
N'Dama cattle (colon): imported from Senegal, reared in Louila Ranch (Popular Congo Republic)
- Agriostomum vryburgi Railliet, 1902, illus.
Neto, M. P.; Grisi, L.; and Amato, J. F. R., 1975, Rev. Brasil. Biol., v. 35 (4), 639-643
Agriostomum vryburgi, incidence and intensity by seasons, cattle (Zebu-friesland) (large intestine): state of Rio de Janeiro, Brazil
- Alaeuris, illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Syn.: Thelastomoides
- Alaeuris caballeroi n. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraisemblablement" G. agasizii: desert de Sonora, Mexique (colon of all)
- Alaeuris gopheri n. sp.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
- Alaeuris gopheri gopheri n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraisemblablement" G. agasizii: desert de Sonora, Mexique (colon of all)
- Alaeuris gopheri macrolabiata n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus polyphemus (colon): sud de Lake Placid, comte de Highlands, Floride
- Alaeuris gopheri pudica n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. agasizii (colon): desert de Sonora, Mexique
- Alaeuris kinsellai n. sp.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
- Alaeuris kinsellai kinsellai n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. flavo-marginatus (colon): desert de Coahuila, Mexique
- Alaeuris kinsellai sonorae n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. agasizii (colon): desert de Sonora, Mexique
- Alaeuris longicollis (Walton, 1927) [? n. comb.], illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
description
Syn.: Thelastomoides longicollis Walton, 1927
Gopherus sp., "vraisemblablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraisemblablement" G. agasizii: desert de Sonora, Mexique (colon of all)

- Alaeuris mazzottii* n. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976,
Bull. Mus. National Hist. Nat., Paris, 3. s.
(389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" *G. flavo-*
marginatus (colon): desert de Coahuila,
Mexique
- Alaeuris paramazzottii* n. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976,
Bull. Mus. National Hist. Nat., Paris, 3. s.
(389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" *G. agas-*
sizii: desert de Sonora, Mexique
Gopherus polyphemus: sud de Lake Placid,
comte de Highlands, Floride
(colon of all)
- Alaimina* Clark, 1961
Maggenti, A. R., 1976, Organ. Nematodes
(Croll), 1-10
Dorylaimida
- Alaimus primitivus* De Man, 1880, illus.
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972,
Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera
beetles, possible importance in biological
control
Pimelia angulata: Abu-Rawash, Cairo, Egypt
Blaps polychresta: Cairo, Egypt
(intestine of all)
- Alainchabaudia* Mawson, 1968
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Hartertiidae
key
- Alcedospirura* Oshmarin, 1959
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Aviculariella* Wehr, 1931
- Alfortia*
Reinecke, R. K.; and le Roux, D. J., 1972, J.
South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys
and modified critical tests on horses using
mebendazole, highly effective
- Alfortia edentatus*, illus.
Loseva, N. G., 1975, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 25, 95-97
Alfortia edentatus, *Delafondia vulgaris*,
histochemistry of intestine, low glycogen
content related to blood feeding; quantity
and distribution of nucleic acids
- Alfortia edentata*
de Matos, P. F.; and Costa, J. O., 1976, Arq.
Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2),
173-180
gastrointestinal helminths, horses, levami-
sole, haloxon, crufomate, anthelmintic effi-
ciency
- Alfortia edentatus* (Looss, 1900) Skrjabin, 1933,
illus.
Puliaevaksia, N. V., 1969, Trudy Gel'mint.
Lab., Akad. Nauk SSSR, v. 20, 123-125
Alfortia edentatus females, morphology of
different sections of the genital tract
- Alfortia edentata* (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am.
Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Aliascaridinea (Kalyankar, 1971, subfam.)
Hartwich, G., 1974, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Raphidascaridinae
key
includes: *Aliascaris*
- Aliascaris* Kalyankar, 1971
Hartwich, G., 1974, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Aliascaridinea
key
- Aliella* Ali, 1968
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Dispharynx* subgen. of *Synhimantus*
- Alinema*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
subgen. of *Philometra*
key
- Alinema Jogis*, 1968, nec Rasheed, 1963
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Rusguniella* Seurat, 1919
- Allantonematidae n. sp., illus.
Laumond, C.; and Lyon, J. P., 1975, Acta
Trop., v. 32 (4), 334-339
Allantonematidae n. sp., probably belonging to
Iotonchium, morphology and life cycle
Helophilus trivittatus
H. pendulus
(hemocoel of all): all from Sud de la
France
- Allantonematidae
Poinar, G. O., jr.; and Nelson, B. C., 1973,
J. Med. Entom., v. 10 (4), 349-354
"should be re-instated"
- Allintoschius* Chitwood, 1937 [lapsus for *Allin-*
toshius]
Durette-Desset, M. C.; and Chabaud, A. G.,
1975, Ann. Parasitol., v. 50 (3), 303-337
Syn.: *Parallintoschius* [sic] Araujo, 1940

- Allintoshius Chitwood, 1957
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Ornithostrongylinae
Syn.: Parallintoshius Araujo, 1940
- Allintoschius [sic] dunni n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Myotis mystacinus (intestin): Bukit Mangol, Province de Selangor, Malaisie
Pipistrellus nanus: Maboke, Republique Centrafricaine
- Allodapa suctoria
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in Locusta migratoria (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Amidostomatidae (Travassos, 1919, sub. fam.), Baylis et Daubney, 1926
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
includes: Amidostomatinae; Inglamidinae; Epomidostomatinae; Mackerrastrongylinae
- Amidostomatidae
Mawson, P. M., 1976, Tr. Roy. Soc. South Australia, v. 100 (3), 121-123
Woodwardstrongylus transferred to family Amidostomatidae
- Amidostomatinae Travassos, 1919
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae
includes: Amidostomum (type genus); Amphibiophilus; Austrostrongylus; Batrachonema; Dromaeostrongylus; Herpetostrongylus; Nicolina; Paramidostomum; Paraastrostrongylus; Patricialina; Woolleya
- Amidostomum Railliet et Henry, 1909 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Amidostomum sp.
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (under gizzard lining): Florida
- Amidostomum sp.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Ceryle rudis: Israel
- Amidostomum acutum (Lundahl, 1848)
Bishop, C. A.; and Threlfall, W., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 25-35
Somateria mollissima (below gizzard lining, junction with proventriculus and duodenum): insular Newfoundland and/or southern Labrador
- Amidostomum acutum Lundahl, 1848
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anser albifrons
Casarca ferruginea
Anas platyrhynchos
A. strepera
A. penelope
A. clypeata
A. acuta
A. crecca
A. querquedula
Aythya ferina
A. nyroca
Netta rufina
(under cuticle of muscular stomach of all): all from Bulgaria
- Amidostomum acutum, illus.
Orlandi, M.; and Colombani, B., 1975, Ann. Fac. Med. Vet. Pisa, v. 27, 1974, 113-128
histopathology
Anas querquedula (gizzard): Tombolo (Pisa)
- Amidostomum acutum (Lundahl, 1848)
Turner, B. C.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 157-169
parasites of Anas crecca and A. discors, incidence and intensity, age and sex of host
Anas crecca
A. discors
(keratinous lining of gizzard of all): all from eastern Canada
- Amidostomum acutum (Lundahl, 1848), illus.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
brief description
Anas crecca (intestin grele): Kefar Ruppim, Israel
- Amidostomum anseris
Cervenka, J.; Zajicek, D.; and Nydl, J., 1975, Veterinarstvi, v. 25 (6), 263-264
helminths, geese, Mebendazole
- Amidostomum anseris Zeder, 1800
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anser anser
A. albifrons
A. erythropus
Anas platyrhynchos
A. querquedula
(under cuticle of muscular stomach of all): all from Bulgaria
- Amidostomum anseris (Zeder, 1800), illus.
Letonja, T., 1972, Bol. Chileno Parasitol., v. 27 (3-4), 131-133
Anser albifrons (gizzard): Chile
- Amidostomum anseris (Zeder, 1800), illus.
Phuc, D. V.; and Varga, I., 1975, Acta Vet., Budapest, v. 25 (2-3), 231-239
Amidostomum anseris, comparison of worm development and pathology in experimentally infected chickens and ducklings with those of goslings, stored larvae from ducklings proved infective to goslings and ducklings

- Amidostomum anseris*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Amidostomum anseris*, illus.
Vetesi, F.; Phuc, D. V.; and Varga, I., 1976, Acta Vet., Budapest, v. 26 (1), 113-127
Amidostomum anseris, goslings (ventriculus) (exper.), histopathology, comparison with infected ducklings and chickens
- Amidostomum fulicae*
Eley, T. J., jr., 1976, Calif. Fish and Game, v. 62 (2), 156-157
Fulica americana (gizzard-under lining): lower Colorado River
- Amidostomum fulicae* (Rudolphi, 1819)
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Porphyryla martinica (gizzard lining): Florida
- Amidostomum fulicae* (Rudolphi, 1819), illus.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
brief description
Fulica atra (intestin grele): Hula and Qishon, Israel
- Amidostomum quasifulicae* Macko, 1966
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans (gizzard lining): Florida
- Amphibiophilus* Skrjabin, 1916
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Amphicaecum* Walton, 1927
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Contracecum* Railliet & Henry, 1912
- Amphimermis elegans* (Hagmeier 1912) Welch 1963
Kaiser, H., 1974, Mitt. Naturw. Ver. Steiermark, v. 104, 177-181
synonymy
Steiermark, Osterreich
- Amplicaecum* Baylis, 1920
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Angusticaecinae
key
? Syn.: *Orneoscaris* Skrjabin, 1916, gen. dub.
- Amplicaecum alii* n. sp., illus.
Ilyas, R., 1974, Riv. Parassitol., Roma, v. 35 (2), 125-128
Milvus migrans (gizzard): Aurangabad, Maharashtra, India
- Amplicaecum robertsi*, illus.
Archer, G. T., 1969, Pathology, v. 1 (2), 133-140
Amplicaecum robertsi in rats (exper.), antigen-antibody precipitate was chemotactic to eosinophils, phagocytosis of precipitate by eosinophils occurred and was followed by lysis of eosinophil granules and discharge of granular material outside cells, mast cell changes followed eosinophilia and occurred at sites of eosinophil accumulation
- Amplicaecum robertsi*
Mesina, J. E.; et al., 1974, Tropenmed. u. Parasitol., v. 25 (1), 116-127
Rattus spp. (liver): North Queensland, Australia
- Anacanthocheilus* Wulker, 1930
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis* Layman & Borovkova, 1926
- Anacanthocheilus* Wuelker, 1930
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Pseudanisakis* (Layman & Borovkova, 1926, subgen.) *Mozgovoi*, 1951
- Anacanthocheilus australis* of Johnston & Mawson (1945), to *Acanthocheilus* [comb. not made]
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
". . . *Anacanthocheilus australis* belongs to the genus *Acanthocheilus* and may be identical with *A. rotundatus* (Rud., 1819) Hartwich, 1957 (= *A. bicuspis*; = *A. quadridentatus*)"
- Anacanthocheilus rajae* (Yamaguti, 1941) of Johnston & Mawson (1945)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis rajae* Yamaguti, 1941, sensu nov.
- Anacanthocheilus rotundatus* (Rud.) of Wulker (1930), of Baylis (1939)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis tricupola* nom. nov.
- Anatrichosoma buccalis*
Kinsella, J. M.; and Winegarner, C. E., 1975, J. Parasitol., v. 61 (4), 779-781
Anatrichosoma buccalis in *Didelphis virginiana*, incidence of lesions, host age, course of infection in field: Archbold Biological Station, Highlands County, Florida

- Anatrichosoma cynamolgi Smith and Chitwood 1954, illus.
Long, G. G.; Lichtenfels, J. R.; and Stookey, J. L., 1976, J. Parasitol., v. 62 (1), 111-115
synonymy, prevalence in Macaca mulatta, limited inflammatory response: U. S. Army Medical Research Institute of Infectious Diseases, Frederick, Maryland
- Anatrichosoma nacepobi Conrad and Wong 1973
Long, G. G.; Lichtenfels, J. R.; and Stookey, J. L., 1976, J. Parasitol., v. 62 (1), 111-115
as syn. of Anatrichosoma cynamolgi Smith and Chitwood 1954
- Anatrichosoma rhina Conrad and Wong 1973
Long, G. G.; Lichtenfels, J. R.; and Stookey, J. L., 1976, J. Parasitol., v. 62 (1), 111-115
as syn. of Anatrichosoma cynamolgi Smith and Chitwood 1954
- Ancylostoma
Biagi, F.; Smyth, J.; and Gonzalez, C., 1974, Prensa Med. Mexicana, v. 39 (1-2), 51-53
human intestinal helminths, successful clinical trials using mebendazole, drug well tolerated with minimal side effects: Mexico
- Ancylostom[a]
Genchi, C.; and Locatelli, A., 1974, Atti Soc. Ital. Sc. Vet., v. 28, 862-863
dog helminth eggs, contaminated samples from public parks, potential source of infection for humans and domestic animals: Milan
- Ancylostomum
Ockens, N., 1977, Medlemsbl. Danske Dyrægeforen., v. 60 (4), 147-148
Ancylostomum, Uncinaria, dogs, symptoms, good results with banminth
- Ancylostoma spp.
Galliard, H., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 295-304
visceral and cutaneous larva migrans, review
- Ancylostoma sp. (presumably A. tubaeforme)
Guterbock, W. M.; and Levine, N. D., 1977, J. Am. Vet. Med. Ass., v. 170 (12), 1411-1413
cats (feces): east central Illinois
- Ancylostoma [sp.]
Hass, D. K.; and Chitwood, M. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 255
dog (feces): west central Ohio
- Ancylostoma spp.
Lacuata, A. Q.; and Corpuz, Z. C., 1975, Philippine J. Vet. Med., v. 13 (1-2), 242-250
Ancylostoma spp., dogs, citarin administered subcutaneously, drug efficacy, fair results: Philippines
- Ancylostoma sp.
Papasarathorn, T.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 82-92
human hookworm, epidemiologic survey, relationship of climatic conditions and soil to disease spread, suggested control measures: Soong Nern District, Korat Province, Thailand
- Ancylostoma sp.
Pattanayak, P. C., 1974, Indian J. Animal Research, v. 8 (2), 56-58
Ancylostoma sp., toxocarasis, pups, clinical trial with tetramisole, effective safe medication
- Ancylostoma braziliense De Faria, 1910, illus.
Areekul, S.; and Tipayamontri, U., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 31-38
Ancylostoma braziliense, egg counts, sex, number and size of developing worms, measurements, hematologic changes in hosts, comparative study of findings in dogs and cats (both exper.) infected by oral and cutaneous routes
- Ancylostoma braziliense
Areekul, S.; Tipayamontri, U.; and Ukoskit, K., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 230-235
Ancylostoma braziliense, dogs and cats (both exper.), measurements of blood loss through gastrointestinal tract with finding that A. braziliense caused relatively insignificant loss in these animals
- Ancylostoma braziliense, illus.
Peiretti, H. A., 1975, Gac. Vet., Buenos Aires (306), v. 37, 649-654
incidence, measurements
dogs
cats
all from Corrientes, Argentina
- Ancylostoma braziliense
Rep, B. H.; and Heinemann, D. W., 1976, Trop. and Geogr. Med., v. 28 (2), 104-110
Ancylostoma spp., prevalence and geographical distribution survey of hookworm in dogs and cats, follow-up of 1967 survey
cat
dog
all from Surinam
- Ancylostoma braziliense
Robinson, M.; Hooke, F. G.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (2), 104-108
pyrantel pamoate, critical trials against roundworms and hookworms in naturally and experimentally infected dogs, drug efficacy compared with piperazine citrate: Australia
- Ancylostoma braziliense, illus.
Setasuban, P.; and Vajrasthira, S., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 505-509
scanning electron microscopy showing anterior end
- Ancylostoma braziliense
Setasuban, P.; and Waddell, A. H., 1973, Austral. Vet. J., v. 49 (2), 110
dog
cat
all from Cairns, North Queensland
- Ancylostoma braziliense
Tharaldsen, J., 1973, Norwegian J. Zool., v. 21 (4), 327-328 [Abstract]
cats (feces): quarantine station, Oslo, Norway

- Ancylostoma braziliense*
Vetter, J. C. M.; and Klaver-Wesseling, J. C.,
M., 1977, *Trop. and Geogr. Med.*, v. 29 (3),
319 [Abstract]
hookworms, dogs, detection of antibodies
using indirect fluorescent antibody tech-
nique was successful against cryostat sec-
tions of infective larvae but not against
living infective larvae
- Ancylostoma braziliense*, *illus.*
Yoshida, Y.; et al., 1973, *Southeast Asian J.*
Trop. Med. and Pub. Health, v. 4 (4), 498-503
prevalence survey for hookworms in stray
dogs, domestic and wild cats
dogs
cats
all from Kuala Lumpur, Malaysia
- Ancylostoma buckleyi*, Le Roux and Biocca, 1957,
illus.
Setasuban, P., 1976, *Southeast Asian J. Trop.*
Med. and Pub. Health, v. 7 (1), 45-49
Ancylostoma buckleyi, morphometric data,
scanning electron microscopy
dogs (intestine): Cairns, North Queensland,
Australia
- Ancylostoma caninum* (Ercolani 1859)
Acholonu, A. D., 1977, *J. Parasitol.*, v. 63
(4), 757-758
dog: Ponce, Puerto Rico
- Ancylostoma caninum*
Areekul, S., 1975, *Southeast Asian J. Trop.*
Med. and Pub. Health, v. 6 (1), 144-145
Ancylostoma caninum in dogs (exper.), estima-
tion of blood loss caused by parasitemia
using ⁵¹Cr-labelled red cells
- Ancylostoma caninum*
Areekul, S.; et al., 1971, *Southeast Asian J.*
Trop. Med. and Pub. Health, v. 2 (2), 201-209
Ancylostoma ceylanicum, *A. caninum*, hookworm
anemia in dogs (exper.) influenced by their
iron reserve and dietary iron, no difference
between effects of hookworm infection on
iron metabolism in dogs with normal and
deficient iron reserves
- A[*ncylostoma*] *caninum*
Areekul, S.; Viravan, C.; and Radomyos, P.,
1973, *Southeast Asian J. Trop. Med. and Pub.*
Health, v. 4 (1), 138
A[*ncylostoma*] *ceylanicum*, *A. caninum*, unsuc-
cessful attempts to infect volunteer subjects
by administering filariform larvae orally in
small quantities of water, larvae possibly
killed by high acidity of gastric juices
- Ancylostoma caninum* or *A. tubaeforme* larvae
Ash, L. R., 1976, *Rev. Biol. Trop.*, v. 24 (1),
163-174
Vaginulus plebeius
Laevicaulus alte
all from market gardens, near Noumea, New
Caledonia
- Ancylostoma caninum*
Banerjee, D.; Mandal, A.; and Prakash, O.,
1971, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v.
65 (5), 685-686
marked inhibitory action of mebendazole on
development of larval stages from eggs of
Ancylostoma duodenale, *A. caninum*, and
Necator americanus, no effect on infective
larval stages, *in vitro* trials
- Ancylostoma caninum*
Banerjee, D.; Prakash, O.; and Nayar, E.,
1971, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 65
(5), 681-682 [Letter]
Ancylostoma caninum in cyclops (exper.),
larvae remain viable for 2-3 days, possible
natural vector
- Ancylostoma caninum*
Bhopale, M. K.; and Johri, G. N., 1975, *J.*
Helminth., v. 49 (3), 179-185
Ancylostoma caninum larvae, migration and
distribution in various organs and muscles
of mice after oral infection with various
doses
- Ancylostoma caninum*
Bhopale, M. K.; and Johri, G. N., 1976, *J.*
Hyg., Epidemiol., Microbiol. and Immunol., v.
20 (4), 464-470
Ancylostoma caninum, Swiss albino mice
(exper.), analysis of serum protein compo-
nents in the presence of infection, signifi-
cant decrease in albumin and gamma globulin
with increase in beta globulin, most signifi-
cant changes occurred on the 9th day after
infection
- Ancylostoma caninum*
Buick, T. D.; Campbell, R. S. F.; and Hutch-
inson, G. W., 1977, *Austral. Vet. J.*, v. 53
(12), 602-603 [Letter]
Ancylostoma caninum, dog (cervical spinal
cord), first case in which partial maturity
of the nematode was achieved with intense
involvement of central nervous system:
Queensland
- Ancylostomum caninum*, *illus.*
Bwangamoi, O., 1973, *Bull. Epizoot. Dis.*
Africa, v. 21 (4), 363-370
recovery from dogs using Lindsey's method;
case report, dog (caecum, colon, rectum):
Uganda
- Ancylostoma caninum*
Chauhan, H. V. S.; Dwivedi, P.; and Kalra,
D. S., 1974, *Haryana Vet.*, v. 13 (1),
5-21
protozoan and helminth parasites, transmit-
ted through milk to newborn animals, review
- Ancylostoma caninum*
Genchi, C., 1976, *Arch. Vet. Ital.*, v. 27
(3-4), 98-99
dog intestinal helminth ova, incidence in
soil samples in public parks, potential
public health problem: Milan

- Ancylostoma caninum**
Hass, D. K., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 100-103
Ancylostoma caninum in beagles (exper.), vincofos effective against late third, fourth, early fifth, and juvenile stages, elimination of pathologic responses in host
- Ancylostoma caninum**
Hass, D. K.; and Collins, J. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 135-137
helminths, dogs, comparative efficacy of vincofos, ticarbodine, mebendazole
- Ancylostoma caninum**
Kelly, J. D.; et al., 1976, N. Zealand Vet. J., v. 24 (5), 93-94 [Letter]
larval Ancylostoma caninum, arrested development in gastro-intestinal tract of dogs, complication of treatment, diagnosis and control, significance in transmission
- Ancylostoma caninum**
Khuller, G. K.; Jain, S. K.; and Vinayak, V.K., 1977, Experientia, v. 33 (12), 1585
Ancylostoma caninum adults, phospholipid composition
- Ancylostoma caninum**
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, Vet. Rec., v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects
- Ancylostoma caninum**
Klein, J. B.; and Bradley, R. E., sr., 1976, Vet. Med. and Small Animal Clin., v. 71 (5), 598-599
dogs, cats, sansalid, critical testing, good results
- Ancylostoma caninum**
Krawokwa, S., 1977, J. Am. Vet. Med. Ass., 1977, v. 171 (8), 750-753
transplacentally acquired parasitic diseases of dogs, diagnostic features, review
- Ancylostoma caninum**
Lamas, R.; et al., 1970, Bol. Chileno Parasitol., v. 25 (1-2), 74-77
cutaneous larva migrans (possible Ancylostoma caninum origin) in city garbage collector successfully treated with thiabendazole, transmission of infection probably from fecal-contaminated material associated with occupation: Santiago, Chile
- Ancylostoma caninum**
Lee, K. T.; Little, M. D.; and Beaver, P. C., 1975, J. Parasitol., v. 61 (4), 589-598
Ancylostoma caninum larvae, persistence and intracellular (muscle-fiber) habitat in vertebrate paratenic hosts and nature of host responses studied in mouse, cat, and rhesus monkey (all exper.); larvae also recovered by tissue digestion from muscles of naturally infected dogs
- Ancylostoma caninum**
McCurdy, H. D.; and Guerrero, J., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1731-1733
helminths, dogs, mebendazole powder, controlled critical studies, good results against all helminths except Dipylidium caninum: Kansas; New Jersey; Texas
- Ancylostoma caninum**
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1975, Indian Vet. J., v. 52 (6), 451-456
Ancylostoma caninum, dogs (nat. and exper.), nitroxylnil subcutaneously, drug efficacy, good results; nitroxylnil not effective against Taenia sp., Dipylidium sp., Toxocara sp.
- Ancylostoma caninum**
Mark, D. L., 1975, J. Parasitol., v. 61 (3), 484-488
Ancylostoma caninum, infective larvae, survival on outdoor bluegrass plots in 40 experiments over 1 year
- Ancylostoma caninum**
Migasena, S.; Gilles, H. M.; and Maegraith, B. G., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 534-547
Ancylostoma caninum, experimental infections in dogs, study of interrelationships between larval infective dose, egg count and worm load; pathologic and hematologic changes
- Ancylostoma caninum, illus.**
Oyerinde, J. P. O., 1976, Ann. Trop. Med. and Parasitol., v. 70 (4), 455-462
Ancylostoma caninum, ability of Musca domestica to carry hookworm eggs and larvae externally and internally, viability of eggs and survival of larvae after such exposure, potential role of house fly in dissemination of hookworm
- Ancylostoma caninum, illus.**
Peiretti, H. A., 1975, Gac. Vet., Buenos Aires (306), v. 37, 649-654
incidence, measurements
dogs: Corrientes, Argentina
- Ancylostomum caninum**
Ramachandran, C. P., 1975, Kajian Vet., v. 7 (1), 31-38
helminths, immunization with radiation attenuated vaccines, review
- Ancylostoma caninum**
Ray, D. K.; Negi, S. K.; and Srivastava, P. S., 1975, Indian J. Animal Research, v. 9 (2), 75-78
jackals
wild cat
all from Tarai area, Uttar Pradesh
- Ancylostoma caninum**
Rep, B. H.; and Heinemann, D. W., 1976, Trop. and Geogr. Med., v. 28 (2), 104-110
Ancylostoma spp., prevalence and geographical distribution survey of hookworm in dogs and cats, follow-up of 1967 survey
dog: Surinam

- Ancylostoma caninum*
Roberson, E. L.; and Ager, A. L., 1976, Am. J. Vet. Research, v. 37 (12), 1479-1482
cestodes, nematodes, dogs, natural infections, uredofos highly effective, no toxicosis
- Ancylostoma caninum*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, Am. J. Vet. Research, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections
- Ancylostoma caninum*
Robinson, M.; Hooke, F. G.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (2), 104-108
pyrantel pamoate, critical trials against roundworms and hookworms in naturally and experimentally infected dogs, drug efficacy compared with piperazine citrate: Australia
- Ancylostoma caninum*
Robinson, M.; Hooke, F.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (3), 173-176
Trichuris vulpis, *Ancylostoma caninum*, *Toxocara canis*, dogs, oxtel pamoate separately and combined with pyrantel pamoate, drug efficacy, good results: Australia
- Ancylostoma caninum*
Ronald, N. C.; and Bell, R. R., 1976, Southwest. Vet., v. 29 (3), 217-218
dogs, critical evaluation of butamisol hydrochloride, highly effective against *Ancylostoma caninum* and *Trichuris vulpis*, not effective against *Toxocara canis*; *Dipylidium caninum* unaffected
- Ancylostoma caninum*
Schantz, P. M.; and Prezioso, U., 1976, Am. J. Vet. Research, v. 37 (5), 619-620
immature *Echinococcus granulosus*, dogs, efficacy of divided doses of fospirate (70-94%); also active against *Ancylostoma caninum* and *Toxocara canis*
- Ancylostoma caninum*
Schmelzle, H. M.; and Stoye, M., 1976, Ztschr. Parasitenk., v. 50 (2), 219-220
Ancylostoma caninum, lactating dogs infected percutaneously, orally or intravenously, chronology and extent of larvae in milk secretion, localization in mammary gland
- Ancylostoma caninum* (Ercolani, 1859), *illus.*
Setasuban, P., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (4), 519-523
infective larvae of *Ancylostoma tubaeforme* and *A. caninum* differentiated by scanning electron microscopy
- Ancylostoma caninum*
Setasuban, P., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 114-115
Ancylostoma tubaeforme, differentiation from *A. caninum* using Sarles phenomenon: precipitation around openings of third stage larvae of *A. tubaeforme* in presence of homologous antisera, no precipitation with *A. caninum*
- Ancylostoma caninum* (Ercolani, 1859), *illus.*
Setasuban, P., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 190-194
attempted experimental cross-breeding of *A. tubaeforme* and *A. caninum* with failure to produce progeny even though the two strains copulated, egg production observed only in identical single-strain combinations supporting assumption that the two species are genetically separate
- Ancylostoma caninum*, *illus.*
Setasuban, P.; Vajrasthira, S.; and Muennoo, C., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (4), 534-539
Ancylostoma ceylanicum, *A. caninum*, statistics of prevalence survey of parasitism in stray cats in Prachin Buri Province, Thailand
- Ancylostoma caninum*
Setasuban, P.; and Waddell, A. H., 1973, Austral. Vet. J., v. 49 (2), 110
dog
cat
all from Cairns and Brisbane, North Queensland
- Ancylostoma caninum*
Sinha, A. K.; and Sahai, B. N., 1976, Indian J. Med. Research, v. 64 (8), 1192-1196
morphological determinations of eggs and infective larvae recovered from dogs and human sources to establish *Ancylostoma duodenale* and *A. caninum* as sources of infections in Patna, Bihar, India
- Ancylostoma caninum*
Smith, C. F.; and Hooke, F. G., 1976, N. Zealand Vet. J., v. 24 (5), 95-96 [Letter]
Ancylostoma caninum, control measures in colostrum infection in dogs, potential problems: imported from Australia to New Zealand
- Ancylostoma caninum*
Stoye, M.; and Krause, J., 1976, Zentralbl. Vet.-Med., Reihe B, v. 23 (10), 822-839
Ancylostoma caninum, impatiently infected lactating ovariectomized dog, reactivation of inhibited larvae, oestradiol and progesterone induced larval excretion in milk
- Ancylostoma caninum*
Stromberg, B. E.; and Soulsby, E. J. L., 1977, Vet. Parasitol., v. 3 (2), 169-175
Ascaris suum, guinea pigs, heterologous resistance induced by *Toxocara canis* and *Ancylostoma caninum* but not by *Haemonchus contortus*, *Caenorhabditis briggsae*, or *Turbatrix aceti*

- Ancylostoma caninum*
Tharaldsen, J., 1973, Norwegian J. Zool., v. 21 (4), 327-328 [Abstract]
dogs (feces): quarantine station, Oslo, Norway
- Ancylostoma caninum*
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Ancylostoma caninum*
Thornton, J. E.; Bell, R. R.; and Reardon, M. J., 1974, *J. Wildlife Dis.*, v. 10 (3), 232-236
Canis latrans: Nueces County, Texas
- Ancylostoma caninum*
Todd, K. S.; and Yates, R. L., 1976, *Am. J. Vet. Research*, v. 37 (11), 1329-1330
Ancylostoma caninum, *Trichuris vulpis*, dogs (exper.), diuredosan, drug efficacy, good results
- Ancylostoma caninum*
Vetter, J. C. M.; and Klaver-Wesseling, J. C., M., 1977, *Trop. and Geogr. Med.*, v. 29 (3), 319 [Abstract]
hookworms, dogs, detection of antibodies using indirect fluorescent antibody technique was successful against cryostat sections of infective larvae but not against living infective larvae
- Ancylostoma caninum*
Williams, B. M., 1976, *Brit. Vet. J.*, v. 132 (3), 309-312
Vulpes vulpes (intestine): southwest Wales
- Ancylostoma caninum*
Yoshida, Y.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 498-503
dogs: Kuala Lumpur, Malaysia
- Ancylostoma caninum* or *A. tubaeforme*
Yoshida, Y.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 498-503
Felis marmorata: Kuala Lumpur, Malaysia
- Ancylostoma ceylanicum*
Areekul, S.; et al., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (2), 201-209
Ancylostoma ceylanicum, *A. caninum*, hookworm anemia in dogs (exper.) influenced by their iron reserve and dietary iron, no difference between effects of hookworm infection on iron metabolism in dogs with normal and deficient iron reserves
- Ancylostoma ceylanicum*
Areekul, S.; et al., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (4), 604-605
unable to establish experimental infection in *Macaca mulatta*
- A[ncylostoma] ceylanicum*
Areekul, S.; Viravan, C.; and Radomyos, P., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (1), 138
A[ncylostoma] ceylanicum, *A. caninum*, unsuccessful attempts to infect volunteer subjects by administering filariform larvae orally in small quantities of water, larvae possibly killed by high acidity of gastric juices
- Ancylostoma ceylanicum*
Rep, B. H.; and Heinemann, D. W., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 104-110
Ancylostoma spp., prevalence and geographical distribution survey of hookworm in dogs and cats, follow-up of 1967 survey
cat
dog
all from Surinam
- Ancylostoma ceylanicum*, *illus.*
Setasuban, P.; and Vajrasthira, S., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (4), 505-509
scanning electron microscopy showing anterior end
- Ancylostoma ceylanicum*, *illus.*
Setasuban, P.; Vajrasthira, S.; and Muennoo, C., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (4), 534-539
Ancylostoma ceylanicum, *A. caninum*, statistics of prevalence survey of parasitism in stray cats in Prachin Buri Province, Thailand
- Ancylostoma ceylanicum*
Vetter, J. C. M.; and Klaver-Wesseling, J. C. M., 1977, *J. Parasitol.*, v. 63 (4), 700
Ancylostoma ceylanicum, unimpaired infectivity of larvae after storage in liquid nitrogen for one year
- Ancylostoma ceylanicum*
Vetter, J. C. M.; and Klaver-Wesseling, J. C. M., 1977, *Trop. and Geogr. Med.*, v. 29 (3), 319 [Abstract]
hookworms, dogs, detection of antibodies using indirect fluorescent antibody technique was successful against cryostat sections of infective larvae but not against living infective larvae
- Ancylostoma ceylanicum*, *illus.*
Yoshida, Y.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 498-503
prevalence survey for hookworms in stray dogs, domestic and wild cats
dogs
cats
all from Kuala Lumpur, Malaysia
- Ancylostoma ceylanicum*
Zuidema, P. J.; Rep, B. H.; and Meuzelaar, H. L. C., 1971, *Nederl. Tijdschr. Geneesk.*, v. 115 (45), 1886-1889
Ancylostoma ceylanicum and *Necator americanus*, mild infections in Dutch soldiers who had served in Surinam, clinical aspects: Netherlands

- Ancylostoma conepati Solanet, 1911, illus.
Ibancz, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 142-145
description of male and female
Conepatus inca (tubo digestivo): San Juan, Cajamarca, Peru
- Ancylostoma duodenale
Adickman, M.; and Tuthill, T. M., 1976, Postgrad. Med., v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Ancylostoma duodenale
Arfaa, F.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 230-233
Ascaris, Trichostrongylus spp., Ancylostoma duodenale, evaluation of different methods of control of soil transmitted helminths (sanitation measures, mass-therapy, combined mass-therapy and sanitation) in villagers of Khuzestan, southwest Iran
- Ancylostoma duodenale
Asmera, J.; et al., 1975, Ceskoslov. Pediat., v. 30 (4), 171-172
human intestinal helminths, mebendazole used as successful wide-spectrum anthelmintic in clinical trials: Czechoslovakia
- Ancylostoma duodenale
Banerjee, D.; Mandal, A.; and Prakash, O., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 685-686
marked inhibitory action of mebendazole on development of larval stages from eggs of Ancylostoma duodenale, A. caninum, and Necator americanus, no effect on infective larval stages, in vitro trials
- Ancylostoma duodenale
Basset, M.; et al., 1975, Medecine Trop., v. 35 (6), 475-476
mixed Strongyloides infections apparent after treatment of Necator americanus and Ancylostoma duodenale with alcopar or didakene, case reports, statistics
- Ancylostoma duodenale
Bhandari, B.; and Shrimali, L. N., 1975, Progr. Drug Research, v. 19, 6-9
Ancylostoma duodenale, bitoscanate a promising safe and effective anthelmintic for children, clinical trials: Udaipur, India
- Ancylostoma duodenale
Biagi, F.; Lopez, R.; and Viso, J., 1975, Progr. Drug Research, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review
- Ancylostoma duodenale
Biroum-Noerjasin, 1973, Clin. and Exper. Immunol., v. 13 (4), 545-551
hookworm, human, IgE levels in relation to anti-helminthic treatment, to numbers of eggs/g of stool, to states of infestation (with and without reinfection, with and without superinfection), and to time post-treatment, changes in differential lymphocyte and eosinophil counts: East Java, Indonesia
- Ancylostoma duodenale
Brown, R. C.; and Girardeau, M. H. F., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 215-219
Strongyloides fuelleborni, Necator americanus, Ancylostoma duodenale, prevalence survey and study of possible transmammmary passage, presence of Strongyloides sp. larvae in milk of one nursing mother suggests that S. fuelleborni may be transmitted via milk in humans: Bulape, Zaire
- A[ncylostoma] duodenale
Bruch, K.; and Haas, J., 1976, Ann. Trop. Med. and Parasitol., v. 70 (2), 205-211
Ascaris, hookworm, Trichuris, human, single dose fenbendazole compared with pyrantel: Liberia
- Ancylostoma duodenale
Burke, G. J., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 402-405
10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Ancylostoma duodenale
Camerlynck, P.; and Benmansour, N., 1973, Maroc Med. (571), v. 53, 517-519
mass therapy of human infection with L-tetramisole: Maroc
- Ancylostoma duodenale
de Carneri, I., 1974, Riv. Parassitol., Roma, v. 35 (4), 277-284
Necator americanus, human, geographic distribution in northern band extending from Portugal to Iran, prevailing over Ancylostoma duodenale
- Ancylostoma duodenale
de Carvalho, P., 1973, Pediat. Prat., S. Paulo, v. 44 (3-4), 93-98
Ancylostoma duodenale, acute infection with anemia in infant, diagnostic difficulties, therapy of blood transfusions and tetrachloroethylene: Sao Paulo, Brasil
- Ancylostoma duodenale
Cerva, L., 1976, Ceskoslov. Epidemiol., Mikrobiol., Immunol., v. 25 (6), 365-367
helminths, diagnostic staining method for eggs and larvae in fresh feces or those fixed by formalin

- Ancylostoma duodenale*, *illus.*
Cerva, L., 1976, *Immun. u. Infekt.*, v. 4 (6), 279-282
intestinal helminths, diagnostic method for staining of eggs and larvae in smears of fresh and fixed stool samples
- Ancylostoma duodenale*
Chuttani, H. K.; and Misra, R. C., 1975, *Progr. Drug Research*, v. 19, 44-47
Ancylostoma duodenale, humans, effects of parasitism on function and structure of host small bowel, review
- Ancylostoma duodenale*
Doshi, J. C.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 636-639
human hookworm infections, clinical trials of dosage searching for tolerance and efficacy using C.9333-Go/CGP 4540, compound well tolerated with few side effects, active also in mixed infections with *Ascaris lumbricoides*: Bombay, India
- Ancylostoma duodenale*
Farahmandian, I.; et al., 1977, *Chemotherapy*, v. 23 (2), 98-105
human intestinal *Ascaris*, hookworm and *Trichostrongylus* spp., statistics of comparative trials using pyrantel, levamisole, mebendazole, thiabendazole and bephenium: Iran
- Ancylostoma duodenale*
Farid, Z.; et al., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (3), 425-426 [Letter]
Ancylostoma duodenale, *Ascaris lumbricoides*, male farmers, tetramisole extremely effective against both parasites, no side-effects: Egypt
- Ancylostoma duodenale*
Farid, Z.; et al., 1977, *J. Trop. Med. and Hyg.*, v. 80 (5), 107-108
human *Ascaris lumbricoides* and *Ancylostoma duodenale*, effectiveness of levamisole, pyrantel and bephenium compared in clinical trials, all drugs well tolerated: Egypt
- Ancylostoma duodenale*
Gemési, G., 1971, *Parasitol. Hungar.*, v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Ancylostoma duodenale*
Herrero Alonso, C., 1969, *Med. Trop.*, Madrid, v. 45 (1-2), 11-20
thiabendazole drug of choice in treating human infections of *Ancylostoma duodenale*, *Trichuris trichiura* and *Ascaris lumbricoides*, clinical trials
- Ancylostoma duodenale*
Hsieh, H. C., 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 568 [Demonstration]
Ancylostoma duodenale, *Necator americanus* and *Strongyloides stercoralis*-related spp., differentiation of species using a modified test-tube filter-paper culture
- Ancylostoma duodenale*
Iannucci, J. G., 1974, *Med. Times*, New York, v. 102 (10), 43-49
gastrointestinal hemorrhage as presenting symptom of human hookworm disease, clinical management, need for diagnostic awareness in non-endemic areas
- Ancylostoma duodenale*
Jose, D. G.; and Welch, J. S., 1970, *Med. J. Australia*, v. 1 (8), 349-356
possible role of intestinal parasitism in growth-retarded, anemic and malnourished Australian Aboriginal children, comparison with normal Aboriginal children: Queensland
- Ancylostoma duodenale*
Kaul, T. N.; and Mahajan, R. C., 1977, *Indian J. Med. Research*, v. 66 (3), 413-416
human hookworm infection, immunofluorescence using *Ancylostoma duodenale* antigen, useful immunodiagnostic tool especially in early stages of infections
- Ancylostoma duodenale*
Layrisse, M.; and Vargas, A., 1975, *Progr. Food and Nutrition Sc.*, v. 1 (10), 645-667
human intestinal parasites, mechanisms by which parasites interfere with host nutrition (competition for nutrients, malabsorption, blood loss, excess nutrient utilization), extensive review
- Ancylostoma duodenale*
Levai, J.; Lengyel, A.; and Banki, G., 1972, *Med. Universalis*, v. 5 (1), 31-34
clinical trials using levamisole to treat human intestinal helminths, good results against *Ascaris*, *Ancylostoma*, *Trichostrongylus*, ineffective against *Trichuris trichiura*: Hungary
- Ancylostoma duodenale*
Lozoff, B.; Warren, K. S.; and Mahmoud, A. A. F., 1975, *J. Infect. Dis.*, v. 132 (5), 606-610
Necator americanus, *Ancylostoma duodenale*, human, algorithms in diagnosis and management, mebendazole
- Ancylostoma duodenale*
Otsuru, M., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Ancylostoma duodenale*
Patricia, A. H.; et al., 1975, *Progr. Drug Research*, v. 19, 90-95
Necator americanus, *Ancylostoma duodenale*, humans, efficacy of bitoscanate compared with tetrachlorethylene and bephenium hydroxynaphthoate, found to be safe and effective

- Ancylostoma duodenale*
Radermecker, M.; et al., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months
- Ancylostoma duodenale*
Rajasekaran, P.; Dutt, P. R.; and Pisharoti, K. A., 1977, *Indian J. Med. Research*, v. 66 (2), 189-199
human intestinal parasites, survey of correlation between infection rate and source of water supply (well, street tap, home with tap water) as indication of control of water-borne diseases by public water supplies: Madurai district, Tamil Nadu, India
- Ancylostoma duodenale*
Samuel, M. R., 1975, *Progr. Drug Research*, v. 19, 96-107
human intestinal helminths, review of clinical experiences world wide comparing the efficacy and tolerance of bitoscanate with that of bephenium hydroxynaphthoate and tetrachlorethylene; found to be most useful against hookworm with results against other helminths still inconclusive
- Ancylostoma duodenale*
Schad, G. A.; et al., 1973, *Science* (4085), v. 180, 502-504
Ancylostoma duodenale, human, seasonal variation in egg counts and larval abundance and distribution, evidence for occurrence of arrested development as an adaptation to a seasonally unfavorable external environment: West Bengal, India
- Ancylostoma duodenale*, *illus.*
Setasuban, P.; and Vajrasthira, S., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (4), 505-509
scanning electron microscopy showing anterior end
- Ancylostoma duodenale*
Sinha, A. K.; and Sahai, B. N., 1976, *Indian J. Med. Research*, v. 64 (8), 1192-1196
morphological determinations of eggs and infective larvae recovered from dogs and human sources to establish *Ancylostoma duodenale* and *A. caninum* as sources of infections in Patna, Bihar, India
- Ancylostoma duodenale*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii kraï
- Ancylostoma duodenale*
Tarabini Castellani, G., 1970, *Minerva Gastroenterol.*, v. 16 (1), 45-49
human intestinal parasites, clinical trials using FI 6518 with and without thymol, acceptable drug for mass therapy: Somalia
- Ancylostoma duodenale*, *illus.*
Titus, H., 1974, *Medecine Afrique Noire*, v. 21 (10), 685-692
Ancylostoma duodenale, human, epidemiologic and clinical aspects, case reports: Haiti
- Ancylostoma duodenale*
Torres, P.; Figueroa, L.; and Navarrete, N., 1972, *Bol. Chileno Parasitol.*, v. 27 (1-2), 52-55
Trichostrongylus spp. survey in humans, morphological differentiation for diagnosis: Chile
- Ancylostoma duodenale*
Vakil, B. J.; et al., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (3), 247-250
human hookworm, clinical evaluation of C.9333 Go./CGP 4540, promising anthelmintic with few side effects: India
- Ancylostoma duodenale*
Vinayak, V. K.; Singh, T.; and Naik, S. R., 1977, *Indian J. Med. Research*, v. 66 (5), 737-744
human ancylostomiasis, intradermal skin test using *Ancylostoma duodenale* larval antigen, useful and rapid screening method for epidemiologic surveys, also recommended as adjunct to fecal examination in individual case diagnosis
- Ancylostoma duodenale*
Walker, A. C.; and Bellmaine, S. P., 1975, *Med. J. Australia*, v. 1 (24), 751-752
severe digestive tract hemorrhage in Aboriginal infants apparently associated with massive infestation of *Ancylostoma duodenale* with cessation of bleeding after administration of anthelmintic: Northern Territory, Australia
- Ancylostoma duodenale* (Dubini, 1843), *illus.*
Yoshida, Y.; et al., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (4), 515-518
Ancylostoma duodenale, scanning electron microscopic study of adult surface structures and infective-stage larvae
- Ancylostoma kusimaense*
Yoshida, Y.; and Arizono, N., 1976, *J. Parasitol.*, v. 62 (5), 766-770
Nyctereutes procyonoides: Kyoto and Miyazaki Prefectures, Japan
- Ancylostoma malayanum*, Alessandrini, 1905, *illus.*
Setasuban, P.; and Vajrasthira, S., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (4), 505-509
Ancylostoma malayanum, scanning electron microscopy, comparative morphometric measurements
Helarctos malayanus (small intestine): Nakhon Sri Thammarat Province, Southern Thailand
- Ancylostoma tubaeforme* (Zeder 1800)
Acholonu, A. D., 1977, *J. Parasitol.*, v. 63 (4), 757-758
cat: Ponce, Puerto Rico

- Ancylostoma tubaeforme* or *A. caninum* larvae
Ash, L. R., 1976, Rev. Biol. Trop., v. 24 (1), 163-174
Vaginulus plebeius
Laevicaulus alte
all from market gardens, near Noumea, New Caledonia
- Ancylostoma tubaeforme*
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, Vet. Rec., v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects
- Ancylostoma tubaeforme*
Matthews, B. E.; and Croll, N. A., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (4), 608 [Letter]
Ancylostoma tubaeforme, no consistent trends established between 'biological performance' of larvae and age of adult parasites
- Ancylostoma tubaeformae*
Nwosu, A. B. C., 1977, Bull. Animal Health and Prod. Africa, v. 25 (1), 73-78
Ancylostoma tubaeformae third-stage larvae, changes in lipid content, activity rate, and penetrability with time and at various temperatures, evaluation of infectivity using these parameters
- Ancylostoma tubaeforme*, illus.
Nwosu, A. B. C.; and Croll, N. A., 1977, Exper. Parasitol., v. 43 (1), 100-106
Ancylostoma tubaeforme, infective larvae, movement, mechanism of wave propagation
- Ancylostoma tubaeforme*, illus.
Peiretti, H. A., 1975, Gac. Vet., Buenos Aires (306), v. 37, 649-654
incidence, measurements
cats: Corrientes, Argentina
- Ancylostoma tubaeforme* (Zeder, 1800), illus.
Power, L. A., 1974, Rev. Med. Vet. y Parasitol., Maracay, v. 25 (1-8), 1973-1974, 104-111
gatos (intestino delgado, feces): Maracay, Venezuela
- Ancylostoma tubaeforme*
Rep, B. H.; and Heinemann, D. W., 1976, Trop. and Geogr. Med., v. 28 (2), 104-110
Ancylostoma spp., prevalence and geographical distribution survey of hookworm in dogs and cats, follow-up of 1967 survey
cat: Surinam
- Ancylostoma tubaeforme* (Zeder, 1800), illus.
Setasuban, P., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (4), 519-523
infective larvae of *Ancylostoma tubaeforme* and *A. caninum* differentiated by scanning electron microscopy
- Ancylostoma tubaeforme*
Setasuban, P., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 608-609
Ancylostoma tubaeforme, demonstration of transmammary transmission in baby mice (exper.); prenatal infection not established
- Ancylostoma tubaeforme*, illus.
Setasuban, P., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 114-115
Ancylostoma tubaeforme, differentiation from *A. caninum* using Sarles phenomenon: precipitation around openings of third stage larvae of *A. tubaeforme* in presence of homologous antisera, no precipitation with *A. caninum*
- Ancylostoma tubaeforme* (Zeder, 1800), illus.
Setasuban, P., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 190-194
attempted experimental cross-breeding of *A. tubaeforme* and *A. caninum* with failure to produce progeny even though the two strains copulated, egg production observed only in identical single-strain combinations supporting assumption that the two species are genetically separate
- Ancylostoma tubaeforme*
Setasuban, P.; and Waddell, A. H., 1973, Austral. Vet. J., v. 49 (2), 110
cat: Cairns and Brisbane, North Queensland
- Ancylostoma tubaeforme*, illus.
Smith, J. M., 1976, Internat. J. Parasitol., v. 6 (1), 9-13
Ancylostoma tubaeforme and *Necator americanus* third stage larvae, ultrastructure of oesophageal glands compared before and after penetration through rabbit skin
- Ancylostoma tubaeforme*
Tharaldsen, J., 1973, Norwegian J. Zool., v. 21 (4), 327-328 [Abstract]
cats (feces): quarantine station, Oslo, Norway
- Ancylostoma tubaeforme* or *A. caninum*
Yoshida, Y.; et al., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 498-503
Felis marmorata: Kuala Lumpur, Malaysia
- Ancylostomatoidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Strongylida
key
- Ankylostomes*
Dewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Ancylostomes*
Krubwa, F.; et al., 1974, Medecine Trop., v. 34 (5), 679-687
trial administration of mebendazole to school children on quarterly basis for attempted eradication of intestinal nematodes, recommendations for use as mass control measure: Zaire

- Ankylostomes**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ancylostomiasis**
Ahmed, S. H.; and Vaishnav, S., 1975, Progr. Drug Research, v. 19, 2-5
ancylostomiasis, children, clinical trials with bitoscanate successful, few transient side effects: New Delhi, India
- Ancylostomiasis**
Camargo, P. A. M.; et al., 1974, Pediat. Prat., S. Paulo, v. 45 (10-12), 189-194
human intestinal helminths, comparative treatment trials using mebendazole and pyrantel pamoate, efficacy of drugs checked using the McMaster technique to determine reduction in egg production
- Ancylostomiasis**
Capdevielle, P.; et al., 1975, Medecine Trop., v. 35 (5), 426-428
human ancylostomiasis, decreased prothrombin time as possible indicator of presence of infection
- Ancylostomiasis**
de Carneri, I.; and Amorosi, D., 1972, Parasitologia, v. 14 (1), 109-112
human, absence from horticultural area formerly endemic; absence attributed to nematocides used for plant parasitic nematodes: Tronto; Tesino
- Ancylostomiasis**
de Carneri, I.; and Comaschi, D., 1972, Parasitologia, v. 14 (1), 105-106
human, no longer found in former endemic area: Vigevano (Pavia)
- Ancylostomiasis**
de Carneri, I.; and Penna, R., 1972, Parasitologia, v. 14 (1), 97-101
human, no longer found in former endemic zone: Provincia di Alessandria
- Ancylostomiasis**
de Carneri, I.; and Penna, R., 1972, Parasitologia, v. 14 (1), 103-104
human, no longer found in former endemic area: Provincia di Vercelli
- Ancylostomiasis**
Colette, J.; Garrigue, G.; and Sellin, B., 1976, Nouv. Presse Med., v. 5 (24), 1540 [Letter]
Schistosoma mansoni and ancylostomiasis in humans, study of possible relationships between helminthiasis and presence of HBS antigens in blood in infected persons in Haute-Volta
- Ancylostomiasis**
Diop, B.; and Bao, O., 1974, Medecine Afrique Noire, v. 21 (1), 31-40
human intestinal helminths, clinical indications for treatment, suggested dosage, efficacy, tolerances, possible toxicities
- Ancylostomiasis**
El-Antably, S. A.; and El-Hoshy, M. H., 1975, Bull. Ophth. Soc. Egypt, v. 68, 275-280
human ascariasis, ancylostomiasis and schistosomiasis in association with soft cataracts, clinical aspects, possible relationships with nutritional deficiencies: Egypt
- Ancylostomiasis**
Gougerot, M. A.; et al., 1975, Bull. Soc. Path. Exot., v. 68 (3), 297-303
elevated IgE, useful indicator of possible human parasitism in absence of allergic type conditions
- Ancylostomiasis**
Goulart, E. G.; et al., 1977, J. Helminth., v. 51 (2), 131-132
ancylostomiasis, Strongyloides stercoralis, human, field trials of control by means of mass anthelmintic treatment combined with introduction of plants inhibitory to the free-living larval stages, reduction in prevalence: Ilha Do Governador, Rio de Janeiro, Brazil
- Ancylostomiasis**
Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
anthropozoonotic helminthiasis associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Ancylostomiasis**
Molet, B.; and Derr-Harf, C., 1975, Medecine Trop., v. 35 (4), 336-337
ancylostomiasis, strongyloidiasis, improved coproculture technique for diagnosis
- Ancylostomiasis**
Pasticier, A., 1975, Nouv. Presse. Med., v. 4 (37), 2659-2660
single and mixed parasitic infections in humans, good results with pyrantel pamoate
- Ancylostomiasis**
Pawlowski, Z.; and Chodera, L., 1975, Polski Tygod. Lekar., v. 30 (24), 1035-1037
ascariasis, trichuriasis, ancylostomiasis, humans, clinical trials show mebendazole to be effective drug for all 3 infections: Poznan, Poland
- Ancylostomiasis**
Pene, P.; and Delmont, J., 1973, Medecine Afrique Noire, v. 20 (1), 41-43
human ancylostomiasis, clinical trials using L-tetramisole: Mali

- Ancylostomiasis**
Poirot, J. L.; et al., 1976, *Medecine et Malad. Infect.*, v. 6 (1), 32-35
survey of origination of patients infected with strongyloidiasis and ancylostomiasis in Paris hospital
- Ancylostomiasis**
Rodrigo Gomez, J. M.; et al., 1971, *Med. Espan.* (386), an. 34, v. 65, 355-358
man with long-standing case of duodenal ancylostomiasis, clinical aspects, treatment with tetrachlorethylene, deficiency in intestinal absorption
- Ancylostomiasis**
Sankale, M.; Diop, B.; and Bao, O., 1974, *Medecine Afrique Noire*, v. 21 (2), 149-152
ascariasis, ancylostomiasis, human, successful trials with pyrantel pamoate: Senegal
- Ancylostomiasis**
Shah, P. M.; Junnarkar, A. R.; and Khare, R. D., 1975, *Progr. Drug. Research*, v. 19, 136-146
impact of periodic deworming on nutritional status of preschool children, evaluation of including periodic deworming in National Nutrition Programmes of India
- Ancylostomiasis**
Sohi, B. K., 1975, *Pediat. Clin. India*, v. 10 (3), 230
mixed strongyloidiasis and ancylostomiasis in young child, presenting symptom of acute abdomen, cure with mintezole: Pathankot, India
- Ancylostomiasis**
Sutanto, A. H.; Sembiring, L.; and Simatupang, J., 1976, *Paediat. Indonesiana*, v. 16 (11-12), 453-463
field survey of ancylostomiasis in school children, no evidence of effect on height or weight, tetrachlorethylene therapy only 69% successful: Indonesia
- Ancylostomiasis**
Thomas, J.; et al., 1971, *Medecine Trop.*, v. 31 (2), 177-181
bephenium hydroxynaphthoate, piperazine and thiabendazole in comparative clinical tests, institutionalized children with protein deficiency
- Ancylostomiasis**
Tjoa Tjhoeng Kie; et al., 1970, *Paediat. Indonesiana*, v. 10 (5), 179-189
human ancylostomiasis, poor treatment results with both alcopar and chenopodium oil: North Sumatra
- Ancylostomiasis**
Vinayak, V. K.; Singh, T.; and Naik, S. R., 1977, *Indian J. Med. Research*, v. 66 (5), 737-744
human ancylostomiasis, intradermal skin test using *Ancylostoma duodenale* larval antigen, useful and rapid screening method for epidemiologic surveys, also recommended as adjunct to fecal examination in individual case diagnosis
- Ancylostomidae**
de Figueiredo, N.; et al., 1968, *Bol. Chileno Parasitol.*, v. 23 (1-2), 57-59
intestinal histopathology as ascertained by oral biopsies, common human intestinal parasites
- Ancylostomidae**
Setasuban, P., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (3), 390-394
Bathmostomum sangeri in *Elephas indicus*, light and scanning electron microscopy, morphometric data, confirmation that parasite is probably a Strongyloidea rather than Ancylostomidae: circus elephants in Brisbane, Australia (primary origin unknown)
- Ancylostomum.** See *Ancylostoma*.
- Ancyracanthinae** Yorke & Maplestone, 1926
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatidae
key
includes: *Ancyracanthus*
- Ancyracanthopsis** Diesing, 1861, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Schistorophinae
key; synonymy
- Ancyracanthopsis** sp., illus.
Jaron, W., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 137-152
description, helminth fauna of adult swallows just returning from migration compared with young birds; dynamics of infection, species composition of helminths, various stages of nesting season
Hirundo rustica (underneath the horny lining of the gizzard): Poland
- Ancyracanthopsis buckleyi** sp. n., illus.
Ali, M. M., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 357-363
Amaurornis phoenicurus (gizzard): Hyderabad, Andhra Pradesh, India
- Ancyracanthopsis coronata**
Bush, A. O.; and Forrester, D. J., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Ancyracanthus** Diesing, 1858, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Ancyracanthinae
Syn.: *Piyayussunema* [sic] Kohn, Gomes & Motta, 1968
- Angiostrongylosis**
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiasis associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand

- Angiostrongylus*
Fontan, R.; Beauchamp, F.; and Beaver, P. C., 1975, Bull. Soc. Path. Exot., v. 68 (6), 557-566
endemic foci of eosinophilic meningitis in humans suggestive of *Angiostrongylus* origins: Laos
- Angiostrongylus* n. sp.
Larson, O. R., 1976, Proc. North Dakota Acad. Sc., v. 30, Part I, 29
Peromyscus maniculatus
Clethrionomys gapperi
(pulmonary arteries, bronchial lumina, feces of all): all from along Red River, eastern North Dakota
- Angiostrongylus* sp.
Picot, H.; et al., 1975, Acta Trop., v. 32 (4), 381-383
2 case reports of human eosinophilic meningitis indicate possible presence of *Angiostrongylus cantonensis*, larvae from *Achatina fulica* (hepatopancreas, muscles pedieux) used to infect rats produced 4th stage larvae of *Angiostrongylus* sp.: Ile de la Reunion
- Angiostrongylus cantonensis*
Ash, L. R., 1976, Rev. Biol. Trop., v. 24 (1), 163-174
Angiostrongylus cantonensis in mollusks and planarians occurring in market gardens supplying fresh produce; marked seasonal occurrence of planarians during cool months corresponds to peak vegetable growing season, important source of human infection
Laevicaulus alte
Vaginulus plebeius
Deroceras laeve
Bradybaena similaris
Geoplana forsterorum
Helix aspera (exper.)
all from market gardens, near Noumea, New Caledonia
- Angiostrongylus cantonensis*
Bisseru, B., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 523-526
Angiostrongylus cantonensis, prevalence survey for evidence of infection in the giant African snail (*Achatina fulica*): West Malaysia and Singapore
- Angiostrongylus cantonensis*
Bisseru, B.; Gill, S. S.; and Lucas, J. K., 1972, Med. J. Malaya, v. 26 (3), 164-167
first reported case of human eosinophilic meningoencephalitis in West Malaysia, history of eating raw prawns and salad greens, diagnosis, case report: Kuala Lumpur
- Angiostrongylus cantonensis*, illus.
Brockelman, C. R.; Chusatayanond, W.; and Baidikul, V., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 30-37
Angiostrongylus cantonensis, development and localization within the vector snail *Achatina fulica*
- Angiostrongylus cantonensis*
Chen, S.-N., 1975, Bull. Inst. Zool., Acad. Sinica, v. 14 (2), 109-113
Angiostrongylus cantonensis as possible cause of human cases of eosinophilic meningitis, immunoglobulins and leucocytes in blood and cerebrospinal fluid, antibody to *A. cantonensis*, evidence for specific immune response, history of eating raw *Achatina fulica*
- Angiostrongylus cantonensis*
Crook, J. R.; Fulton, S. E.; and Supanwong, K., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 602-605
Angiostrongylus cantonensis, chlorine or iodine treatment of water used for culinary purposes did not completely attenuate larvae shed into the water by drowned *Achatina fulica* snail vectors, possible source of human infection: Thailand
- Angiostrongylus cantonensis*
Dobson, C.; and Welch, J. S., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 223-228
survey for antibodies against *Dirofilaria immitis*, *Toxocara canis*, *Ascaris suum*, *Angiostrongylus cantonensis*, *A. mackerrasae*, in patients with eosinophilia using fluorescent antibody test and passive reversed Arthus test in guinea pigs; *D. immitis* implicated as etiologic agent of human eosinophilic meningitis: Australia
- Angiostrongylus cantonensis*
Heyneman, D.; and Lim, B. L., 1965, Med. J. Malaya, v. 20 (1), 67-68
correlation of habitat to rodent susceptibility to *Angiostrongylus cantonensis* infection
- Angiostrongylus cantonensis*
Intermill, R. W.; et al., 1972, Jap. J. Exper. Med., v. 42 (4), 355-359
statistical survey of potential hosts of *Angiostrongylus cantonensis* in Okinawa, implications for human public health
rat, albino (exper.)
Suncus murinus riukiuanus
Rattus rattus
R. norvegicus
Achatina fulica
Bradybaena circulus
Fruticicola despecta
Satsuma mercatoria
Vaginulus plebeius
Deroceras laeve
Philomycus (*Meghimatum*) *bilineatus*
all from Okinawa
- Angiostrongylus cantonensis*
Jindrak, K.; Mansukhani, M. G.; and Freiberg, A., 1977, J. Parasitol., v. 63 (6), 1132-1133
[See J. Parasitol., 1978, v. 64 (1), p. 44.]
Angiostrongylus cantonensis, use of commercially available edible snails *Otala lactea* and *Helix aperta* as laboratory hosts

- Angiostrongylus cantonensis*
John, D. T., 1974, Proc. Helminth. Soc. Wash-
ington, v. 41 (2), 121-126
Angiostrongylus cantonensis, mice (exper.),
acquired immunity, weight loss occurred to
a lesser degree in immunized mice, they ex-
perienced a greater leukocytosis
- Angiostrongylus cantonensis*
John, D. T., 1977, Virginia J. Sc., v. 28 (2),
64 [Abstract]
Angiostrongylus cantonensis, mice (exper.),
course of primary infection, pathology
- Angiostrongylus cantonensis*
Kamiya, M., 1970, Southeast Asian J. Trop. Med.
and Pub. Health, v. 1 (4), 570-571 [Demonstra-
tion]
Angiostrongylus cantonensis, positive indi-
rect hemagglutination titers in experimen-
tally infected rats after early stages of infec-
tions
- Angiostrongylus cantonensis*
Kamiya, M.; and Klongkamnuankarn, K., 1970,
Southeast Asian J. Trop. Med. and Pub. Health,
v. 1 (4), 571-572 [Demonstration]
Angiostrongylus cantonensis, rats (exper.),
hemagglutination activity after transfer of
adult worms to abdominal cavity of noninfec-
ted rats
- Angiostrongylus cantonensis*
Kamiya, M.; and Klongkamnuankarn, K., 1970,
Southeast Asian J. Trop. Med. and Pub. Health,
v. 1 (4), 572-573 [Demonstration]
Angiostrongylus cantonensis, fractionation
of serum of rats (exper.) with sephadex G-200
chromatography, hemagglutination tests of
each fraction
- Angiostrongylus cantonensis*
Kamiya, M.; and Klongkamnuankarn, K., 1970,
Southeast Asian J. Trop. Med. and Pub. Health,
v. 1 (4), 573 [Demonstration]
Angiostrongylus cantonensis, rabbits as
experimental host model, immunological re-
sponse to hemagglutination test
- Angiostrongylus cantonensis*
Kamiya, M.; Klongkamnuankarn, K.; and Bunnag,
D., 1970, Southeast Asian J. Trop. Med. and
Pub. Health, v. 1 (4), 574 [Demonstration]
Angiostrongylus cantonensis, evaluation of
hemagglutination test for diagnosis of infec-
tion in man
- Angiostrongylus cantonensis*
Kamiya, M.; Klongkamnuankarn, K.; and Harina-
suta, C., 1971, Southeast Asian J. Trop. Med.
and Pub. Health, v. 2 (2), 233-236
Angiostrongylus cantonensis, comparison of
oral, subcutaneous and intraperitoneal routes
for infecting experimental rats, results show
that infection could also be established in
nature by penetration through wounds or skin
abrasions thus putting at risk laboratory
workers caring for infected vector snails
- Angiostrongylus cantonensis*
Kamiya, M.; Tharavanij, S.; and Harinasuta,
C., 1973, Southeast Asian J. Trop. Med. and
Pub. Health, v. 4 (2), 187-194
Angiostrongylus cantonensis, fractionation
of male and female antigen extracts, anti-
genicity of each fraction determined by in-
direct hemagglutination and immunoelectro-
phoresis tests
- Angiostrongylus cantonensis*
Kamiya, M.; and Wongpisal, T., 1970, Southeast
Asian J. Trop. Med. and Pub. Health, v. 1 (4),
569-570 [Demonstration]
Angiostrongylus cantonensis, rats, compari-
son of oral, subcutaneous and intraperitoneal
methods of experimental infections, classifi-
cation of types of lung lesions
- Angiostrongylus cantonensis*
Kocan, A. A., 1974, Proc. Helminth. Soc. Wash-
ington, v. 41 (2), 237-241
influence of *Nippostrongylus brasiliensis*
on establishment of *Angiostrongylus cantonensis*,
role of various organs, laboratory
rat
- Angiostrongylus cantonensis* Chen, 1935, illus.
Lim, B. L., 1975, Southeast Asian J. Trop.
Med. and Pub. Health, v. 6 (3), 376-381
Angiostrongylus cantonensis, redescription,
morphologic comparisons with *A. malaysiensis*
and with *A. cantonensis* from Formosa
Rattus tiomanicus (pulmonary arteries):
East Coast of Peninsular Malaysia
- Angiostrongylus cantonensis*
Lim, B. L., 1976, Med. J. Malaysia, v. 30 (3),
207-211
clinical aspects and presenting symptoms of
human eosinophilic meningoencephalitis
caused by rat lungworm *Angiostrongylus*
cantonensis and speculation as to similar
infection by *Angiostrongylus malaysiensis*
indigenous to Malaysia, probable disease
transmission through raw or inadequately
cooked food
- Angiostrongylus cantonensis*
Lim, B. L.; and Omar-Ahmad, U. D., 1969, Med.
J. Malaya, v. 23 (3), 208-213
survey for *Angiostrongylus cantonensis* in-
fected wild rodents and land molluscs and
contaminated leaf lettuce, probable sources
of human meningoencephalitis in Malaysia
Rattus argentiventer
Rattus jalorensis
Rattus exulans
Macrochlamys resplendens
Achatina fuliva
Microparmarion malayanus
Laevicaulis alte
all from Pahang, West Malaysia
- Angiostrongylus cantonensis*
Loison, G.; Jardin, C.; and Crosnier, J., 1973,
Medicine Trop., v. 33 (2), 143-161
human eosinophilic meningitis caused by
Angiostrongylus cantonensis, possible trans-
mission through contaminated food, public
health program, improved sanitation as con-
trol measures: South Pacific Islands

- Angiostrongylus cantonensis*
Margono, S. S.; and Ilahude, H. D., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 236-240
Angiostrongylus cantonensis, measurements, distribution survey of definitive and intermediate hosts
Rattus rattus diardi
R. argentiventer
R. norvegicus
Achatina fulica
Laevicaulis alte
Pila scutata
all from Jakarta and its vicinity
- Angiostrongylus cantonensis*
Mason, K. V.; et al., 1976, Austral. Vet. J., v. 52 (6), 295 [Letter]
Angiostrongylus cantonensis, dogs, naturally occurring granulomatous encephalomyelitis: Queensland
- Angiostrongylus cantonensis*
Moreau, J. P. J.; and Outin-Fabre, D., 1972, Medecine Afrique Noire, v. 19 (1), 41-42
Angiostrongylus cantonensis, life cycle, cause of human eosinophilic meningitis
- Angiostrongylus cantonensis*
Nhuan, D. T.; and Hendricks, J. R., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 29-30
Rattus norvegicus
R. exulans
all from greater Saigon areas, Vietnam
- Angiostrongylus cantonensis*, illus.
Nitidandhaprabhas, P.; Harnsomburana, K.; and Thepsitthar, P., 1975, Am. J. Trop. Med. and Hyg., v. 24 (4), 711-712
Angiostrongylus cantonensis living 5th-stage female discovered in spinal fluid of man with eosinophilic meningitis: Thailand
- Angiostrongylus cantonensis*
Ottolenghi, A.; et al., 1977, Infect. and Immun., v. 15 (1), 13-18
Angiostrongylus cantonensis, nonsensitized and sensitized rats after challenge, phospholipase B activity in lungs and brains, eosinophilia in bone marrow, results support hypothesis that inflammation, elevated phospholipase B activity, and reduction in worm burden are causally related
- Angiostrongylus cantonensis*
Picot, H.; et al., 1975, Acta Trop., v. 32 (4), 381-383
2 case reports of human eosinophilic meningitis indicate possible presence of *Angiostrongylus cantonensis*, larvae from *Achatina fulica* (hepatopancreas, muscles pedieus) used to infect rats produced 4th stage larvae of *Angiostrongylus* sp.: Ile de la Reunion
- Angiostrongylus cantonensis*
Pirame, Y.; et al., 1973, Medecine Trop., v. 33 (2), 131-141
human eosinophilic meningitis caused by *Angiostrongylus cantonensis*, case reports, clinical aspects, dehydroemetine treatment: Nouvelle-Caledonie
- Angiostrongylus cantonensis*
Punyagupta, S.; Juttijudata, P.; and Bunnag, T., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 921-931
review of 484 clinical cases of human eosinophilic meningitis probably caused by *Angiostrongylus cantonensis*: Thailand
- Angiostrongylus cantonensis*
Rachford, F. W., 1975, J. Parasitol., v. 61 (6), 1117-1119
Angiostrongylus cantonensis, *Lymnaea palustris* (exper.) as potential paratenic as well as intermediate host, *Cambarus* sp. (exper.) as potential paratenic host
- Angiostrongylus cantonensis*
Rachford, F. W., 1976, Exper. Parasitol., v. 39 (3), 377-381
Angiostrongylus cantonensis, larval growth and development in *Lymnaea palustris*
- Angiostrongylus cantonensis*, illus.
Rachford, F. W., 1976, Exper. Parasitol., v. 39 (3), 382-392
Angiostrongylus cantonensis, histopathologic changes in *Lymnaea palustris*
- Angiostrongylus cantonensis*
Rachford, F. W., 1976, Exper. Parasitol., v. 39 (3), 393-400
Angiostrongylus cantonensis-infected *Lymnaea palustris* vs. non-infected controls, oxygen uptake at 25, 30, and 35 C over 34-day periods
- Angiostrongylus cantonensis*
Saltos, N.; Ghosh, H. K.; and Gan, A., 1975, Med. J. Australia, v. 1 (18), 561-562
case report of eosinophilic meningitis probably resulting from *Angiostrongylus cantonensis*, condition improved after course of thiabendazole: New South Wales
- Angiostrongylus cantonensis*, illus.
Sauerlaender, R., 1976, Ztschr. Parasitenk., v. 49 (3), 263-280
Angiostrongylus vasorum, *A. cantonensis*, histopathology of experimentally infected *Achatina fulica*, localization within host at various times after infection, cellular defense mechanisms
- Angiostrongylus cantonensis*
Saugrain, J., 1971, Medecine Trop., v. 31 (2), 233-236
filariasis, *Angiostrongylus cantonensis*, amoebiasis, human parasitic diseases of medical interest and their etiology: French Polynesia
- Angiostrongylus cantonensis*
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus rattus argentiventer
R. jalorensis
R. sabanus
all from Malaysia

- Angiostrongylus cantonensis*
Stafford, E. E.; et al., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 490-491
Rattus diardii (heart and lungs)
R. jaloriensis "
R. exulans "
Achatina fulica
all from Sumatra, Indonesia
- Angiostrongylus cantonensis*
Stafford, E. E.; Sukeri, S.; and Sutanti, T., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 41-44
survey for possible definitive and intermediate hosts
Bandingia indica setifera
Rattus rattus diardii
Achatina fulica
laboratory rats (exper.)
all from Ancol, Jakarta, Indonesia
- Angiostrongylus cantonensis*
Watts, M. B., 1969, Med. J. Malaya, v. 24 (2), 89-93
first 5 case reports of eosinophilic meningitis in Sarawak, *Angiostrongylus cantonensis* probable etiologic agent
- Angiostrongylus cantonensis*, illus.
Widagdo; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (1), 72-74
Angiostrongylus cantonensis male worm removed from eye of woman, some residual visual impairment, history of eating raw vegetables possibly contaminated by snails and of residence in rat-infested area: Semarang, Central Java
- Angiostrongylus cantonensis*
Wioreno, W., 1975, Bul. Kebun Raya, v. 2 (3), 81-84
Angiostrongylus cantonensis, morphology, life cycle, epidemiology, presence in Indonesia, review
- Angiostrongylus cantonensis* (Chen, 1935)
Wioreno, W., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 136-138
Rattus rattus diardi (lung): Bogar, West Java, Indonesia
- Angiostrongylus cantonensis*
Yii, C. Y., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 233-249
Angiostrongylus cantonensis, human, eosinophilic meningitis and meningoencephalitis, extensive clinical study of 125 patients with 16 case reports: Taiwan
- Angiostrongylus cantonensis*
Yii, C. Y.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 447-454
epidemiologic characteristics of human eosinophilic meningitis and meningoencephalitis probably caused by *Angiostrongylus cantonensis* being inadvertently ingested during preparation for consumption: Taiwan
Rattus rattus subsp.
Rattus norvegicus
Bandingia indica nemorivaga
rat, unidentified
Achatina fulica
all from Taiwan
- Angiostrongylus cantonensis*
Yoshimura, K.; et al., 1976, Japan. J. Vet. Sc., v. 38 (6), 579-593
Angiostrongylus cantonensis, guinea pigs, rats, evolution of cellular (macrophage migration inhibitory factor; delayed-type skin reactivity) and humoral (hemagglutinating and precipitating antibodies) immune responses
- Angiostrongylus cantonensis*
Yoshimura, K.; and Soulsby, E. J. L., 1976, Am. J. Trop. Med. and Hyg., v. 25 (1), 99-107
Angiostrongylus cantonensis, rats, lymphoid cell responsiveness, antibody production (reaginic and haemagglutinating)
- Angiostrongylus cantonensis*
Yoshimura, K.; and Yamagishi, T., 1976, Japan. J. Vet. Sc., Tokyo, v. 38 (1), 33-40
Angiostrongylus cantonensis, rabbits and rats, productions of reaginic and indirect hemagglutinating antibodies, reinfections, course of infection
- Angiostrongylus cantonensis*
Yousif, F.; and Laemmler, G., 1975, Ztschr. Parasitenk., v. 47 (3), 191-201
Angiostrongylus cantonensis, factors influencing infectivity of first stage larvae to *Biomphalaria glabrata*, size of snails, number of larvae, age of larvae, individual or mass exposure, length of exposure, temperature, light
- Angiostrongylus cantonensis*
Yousif, F.; and Laemmler, G., 1975, Ztschr. Parasitenk., v. 47 (3), 203-210
Angiostrongylus cantonensis, infectivity to various species of snails under standardized conditions measured by rate of infection, length of development, snail survival rate, rate of recovery of larvae; formula developed for "Capacity Index"
Biomphalaria glabrata
Biomphalaria alexandrina
Planorbis planorbis
Planorbis intermixtus
Bulinus truncatus
Bulinus contortus
Bulinus africanus
Bulinus tropicus
Helisoma sp.
Lymnaea natalensis
Lymnaea tomentosa
Lymnaea stagnalis
Stagnicola elodes
Physa acuta
Marisa cornuarietis
Lanistes carinatus
(exper. in all)
- Angiostrongylus costaricensis* Morera y Cespedes, 1971
Morera, P., [1971], Bol. Chileno Parasitol., v. 25, (3-4), 1970, 133-134
search for definitive host
Rattus rattus (mesenteric arteries of cecum, intestinal wall, feces)
Sigmodon hispidus (mesenteric arteries of cecum, intestinal wall, feces)
all from Costa Rica

- Angiostrongylus costaricensis* (Morera y Cespedes, 1971)
Morera, P.; and Ash, L. R., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 135
establishment of slug (*Vaginulus plebeius*) as intermediate host of *Angiostrongylus costaricensis* in Costa Rica
Vaginulus plebeius
Rattus norvegicus (exper.)
- Angiostrongylus costaricensis*
Sauerbrey, M., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, part 1), 1156-1158
Angiostrongylus costaricensis, human abdominal infections diagnosed using the gel-double diffusion precipitin reaction using as antigen sera from infected cotton rats
- Angiostrongylus* (*P[arastrongylus]*) *dujardini*
Drozdz et Doby, 1970, illus.
Meszaros, F., 1972, Parasitol. Hungar., v. 5, 163-176
Angiostrongylus dujardini, morphometric data, survey of incidence in rodents
Apodemus flavicollis
Clethrionomys glareolus
A. sylvaticus
Pitymys subterraneus
(lungs, right half of heart of all): all from Hungary
- Angiostrongylus mackerrasae*
Dobson, C.; and Welch, J. S., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 223-228
survey for antibodies against *Dirofilaria immitis*, *Toxocara canis*, *Ascaris suum*, *Angiostrongylus cantonensis*, *A. mackerrasae*, in patients with eosinophilia using fluorescent antibody test and passive reversed Arthus test in guinea pigs; *D. immitis* implicated as etiologic agent of human eosinophilic meningitis: Australia
- Angiostrongylus malaysiensis* n. sp., illus.
Bhaibulaya, M.; and Cross, J. H., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 527-533
Rattus jalorensis (pulmonary arteries, right side of heart)
Bradybaena similaris
Microparmarion malayanus
Laevicaulus alte
rats (exper.)
all from Malaysia
- Angiostrongylus malaysiensis*
Dondero, T. J., jr.; and Lim, B. L., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 38-40
Angiostrongylus malaysiensis, *Lymnaea rubiginosa* (common fresh-water snail in Peninsular Malaysia) a capable experimental intermediate host
- Angiostrongylus malaysiensis*
Lim, B. L., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 275-277 [Demonstration]
Angiostrongylus malaysiensis, comparisons and variations in adult worms from 3 regions of Western Malaysia
- Angiostrongylus malaysiensis*
Lim, B. L., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 376-381
Angiostrongylus cantonensis, redescription, morphologic comparisons with *A. malaysiensis* and with *A. cantonensis* from Formosa: East Coast of Peninsular Malaysia
- Angiostrongylus malaysiensis*
Lim, B. L., 1976, Med. J. Malaysia, v. 30 (3), 207-211
clinical aspects and presenting symptoms of human eosinophilic meningoencephalitis caused by rat lungworm *Angiostrongylus cantonensis* and speculation as to similar infection by *Angiostrongylus malaysiensis* indigenous to Malaysia, probable disease transmission through raw or inadequately cooked food
- Angiostrongylus malaysiensis*
Lim, B. L.; et al., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 384-389
survey for rodent and molluscan hosts and for evidence of human infections
laboratory albino rats (exper.)
Rattus rattus diardii
R. exulans
R. argentiventer
Pila scutata
Achatina fulica
Laevicaulus alte
Microparmarion malayanus
all from Tuaran, Sabah
- Angiostrongylus malaysiensis*
Lim, B. L.; Yap, L. F.; and Krishnansamy, M., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (1), 27-35
Angiostrongylus malaysiensis, survey for naturally infected freshwater snails and rodents in rice fields; laboratory trials to establish possible experimental molluscan hosts
Pila scutata (nat. and exper.)
Bellamyia ingallsiana (nat. and exper.)
Rattus tiomanicus
R. argentiventer
R. exulans
R. rattus diardii
albino Norway rats (exper.) (feces)
Microparmarion malayanus
all from ricefields of Peninsular Malaysia
- Angiostrongylus malaysiensis*
Stafford, E. E.; et al., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 490-491
laboratory rats (exper.)
Rattus diardii (heart and lungs)
R. jalorensis "
R. exulans "
Achatina fulica
all from Sumatra, Indonesia
- Angiostrongylus vasorum* Raillet, 1866, illus.
Bwangamoi, O., 1974, Bull. Epizoot. Dis. Africa, v. 22 (1), 55-68
Angiostrongylus vasorum, dogs, pathology

- Angiostrongylus vasorum*
Lynch, V., 1977, *Vet. Rec.*, v. 101 (2), 41-42
[Letter]
dogs (right ventricle, pulmonary artery),
clinical signs, case reports: United Kingdom
- Angiostrongylus vasorum, illus.*
Sauerlaender, R., 1976, *Ztschr. Parasitenk.*,
v. 49 (3), 263-280
Angiostrongylus vasorum, A. cantonensis,
histopathology of experimentally infected
Achatina fulica, localization within host at
various times after infection, cellular de-
fense mechanisms
- Angiostrongylus vasorum* (Railliet, 1866)
Smith, F. R.; and Threlfall, W., 1973, *Am.*
Midland Naturalist, v. 90 (1), 215-218
Vulpes fulva: insular Newfoundland
- Anguillicola Yamaguti*, 1935, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Anguillicolinae
- Anguillicolidae Yamaguti, 1935
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Dracunculoidea
key; key to subfamilies
includes: Anguillicolinae; Skrjabillaninae
- Anguillicolinae (Yamaguti, 1935, fam.)
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Anguillicolidae
key
includes: Anguillicola
- Angulocirrus Biocca et Le Roux*, 1956
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Molineinae
- Angusticaecinae Skrjabin & Karokhin, 1945
Hartwich, G., 1974, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Ascarididae
key; key to tribes
includes: Angusticaecinae; Ophidascaridinea
- Angusticaecinae Chabaud, 1965
Hartwich, G., 1974, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Angusticaecinae
key; key to genera
includes: Angusticaecum; Amplicaecum
- Angusticaecum Baylis*, 1920, *illus.*
Hartwich, G., 1974, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Angusticaecinae
key
- Anisakiasis
Sapunar, J.; Doerr, E.; and Letonja, T., 1976,
Bol. Chileno Parasitol., v. 31 (3-4), 79-83
human anisakiasis, increasing world wide
problem with increased use of fish for food,
suggested control measures; case report of
woman who expelled *Anisakis* sp. from throat
after eating raw fish, elimination of second
worm in feces after mebendazole therapy:
Santiago, Chile
- Anisakidae (Railliet & Henry, 1912, subfam.)
Skrjabin & Karokhin, 1945
Hartwich, G., 1974, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Ascaridoidea
key; key to subfamilies; synonymy
includes: Goeziinae; Anisakinae; Raphidas-
caridinae
- Anisakidae
Horwitz, M. A.; and Hughes, J. M., 1976, *J.*
Infect. Dis., v. 134 (3), 306-312
Trichinella spiralis, *Toxoplasma gondii*,
Anisakidae, implicated in human food borne
diseases in United States, compiled for 1974
by Center for Disease Control, Atlanta
- Anisakidae
Myers, B. J., 1976, *Tr. Am. Micr. Soc.*, v. 95
(2), 137-142
human anisakiasis, historical review
- Anisakidae [sp.]
Bussieras, J.; and Baudin-Laurencin, F.,
1973, *Rev. Elevage et Med. Vet. Pays Trop.*,
n. s., v. 26 (4), 13a-19a
Thunnus albacares
T. obesus
Katsuwonus pelamis
Euthynnus alleteratus
(peritoinae of all): all from tropical
Atlantic
- Anisakidae gen. sp.
Deliamure, S. L.; and Popov, V. N., 1975,
Biol. Nauk., Min. Vyssh. i Sredn. Spetsial.
Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (stomach,
intestine): Sakhalin Bay
- Anisakidae gen. sp.
Popov, V. N., 1976, *Biol. Nauk., Min. Vyssh.*
i Sredn. Spetsial. Obrazovan. SSSR (145), year
19, (1), 49-53
Histiophoca fasciata (stomach, intestine):
northern shore of Okhotsk Sea from Lisiansk
peninsula to Iamsk island
- Anisakid larvae
Lichtenfels, J. R.; et al., 1976, *Tr. Am. Micr.*
Soc., v. 95 (2), 265-266 [Abstract]
anisakid larvae, resembles *Paranisakiopsis*,
from commercially important shellfish, de-
scription of 4th stage, nearly 100% hyper-
parasitized by haplosporidan
Spisula solidissima
Busycon canaliculata
Lunatia heros
all from coastal waters from New Jersey to
North Carolina

- Anisakid worms, possibly *Paranisakiopsis pectinis*
Perkins, F. O.; Madden, P. A.; and Sawyer, T.
K., 1977, Tr. Am. Micr. Soc., v. 96 (3), 376-382
Spisula solidissima: coastal waters of Maryland and Virginia
- Anisakids, similar to *Paranisakiopsis pectinis* (Cobb 1930) Mosgovoy 1951, illus.
Perkins, F. O.; Zwerner, D. E.; and Dias, R. K., 1975, J. Parasitol., v. 61 (5), 944-949
Urosporidium spisuli sp. n., hyperparasite of anisakids (pseudocoel) in surf clams, no potential health hazard from ingesting clams since they are temperature treated during commercial processing: vicinity Chesapeake Light, off Cape Henry, Virginia, N. Atlantic Ocean
- Anisakiden-type, illus.
Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Anisakinae Railliet & Henry, 1912
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakidae
key; key to tribes; synonymy
includes: Anisakinea; Contracaecinea
- Anisakine nematodes
Stern, L.; et al., 1976, Tr. Am. Micr. Soc., v. 95 (2), 264 [Abstract]
Eopsetta jordani
Gadus macrocephalus
Microstomus pacificus
Oncorhynchus kisutch
O. nerka
Ophiodon elongatus
Parophrys vetulus
Sebastes alutus
S. brevispinis
S. caurinus
S. elongatus
S. entomelas
S. flavidus
S. melanops
S. paucispinis
S. pinniger
all from Washington state waters
- Anisakinea Chabaud, 1965
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinae
key; key to genera
includes: Anisakis; Phocanema; Terranova; Sulcascaris
- Anisakis Dujardin, 1845, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinea
key; synonymy
- Anisakis (Skrjabinisakis) Mozgovoi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakis Dujardin, 1845*
- Anisakis
Myers, B. J., 1976, Tr. Am. Micr. Soc., v. 95 (2), 137-142
human anisakiasis, historical review
- Anisakis
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Anisakis sp.
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Phalacrocorax ussuriensis
Larus crassirostris
Oceanodroma monorchis
(esophagus of all): all from Rimsko-Korskov islands
- Anisakis sp.
Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 80-88
helminth distribution among age groups of *Pleurogrammus azonus* (body cavity, muscles): Peter the Great Bay, Sea of Japan
- Anisakis larvae, illus.
Andreassen, J.; and Jørring, K., 1970, Nord. Med., Stockholm, v. 84 (48), 1492-1495
anisakiasis in human presenting as acute appendicitis, at surgical intervention white worm larvae discovered in intestinal wall, other larvae found in patient's home-salted herring: Denmark
- Anisakis sp.
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (3), 185-189
measurements
Hydrobates pelagicus (ventriculus): island of Rost, Norway
Fulmarus glacialis: off Greenland
- Anisakis sp.
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus argentatus
Lunda cirrhata
all from coast of Sea of Okhotsk
- Anisakis larvae Type I
Beverley-Burton, M.; Nyman, O. L.; and Pippy, J. H. C., 1977, J. Fish. Research Bd. Canada, v. 34 (1), 105-112
Anisakis simplex larvae, description, morphology with particular reference to excretory system; comparative morphology of larvae from *Clupea harengus harengus* and *Salmo salar* in widely separated areas of North Atlantic suggest that *Anisakis* larvae Type I is *A. simplex*, findings substantiated by acid phosphatase polymorphism studies
- Anisakis sp.
Bier, J. W.; et al., 1976, Tr. Am. Micr. Soc., v. 95 (2), 264-265 [Abstract]
Anisakis sp. and *Phocanema* sp. in pigs (exper.) (stomach), gross and microscopic pathology

- Anisakis sp.
Bier, J. W.; Jackson, G. J.; and Gerding, T. A., 1976, Tr. Am. Micr. Soc., v. 95 (2), 265 [Abstract]
Anisakis sp., Phocanema sp., blood analysis of experimentally infected pigs showed mild eosinophilia and increased levels of amylase, lactic dehydrogenase, and bilirubin
- Anisakis sp.
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus (stomach): European waters
- Anisakis Type I
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 227-232
incidence, intensity, host diet, habitat; ecological relationships of larval ascarioids from marine fishes
Carangoides fulvoguttatus
Euthynnus alletteratus
Kishinoella tonggol
Lutjanus amabilis
L. sebae
Plectropomus maculatus
Pranesus ogilbyi
Scomberomorus commerson
S. nipponius
all from south-eastern Queensland
- Anisakis sp. (Type I), illus.
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 233-243
description, key
- Anisakis sp. larvae
Cattan, P. E.; and Videla, N. N., 1976, Bol. Chileno Parasitol., v. 31 (3-4), 71-74
Anisakis sp., survey of parasitized Trachurus murphyi (cavidad celomatica, mesenterios, estomago, intestino, gonadas), relationship between size of fish and frequency of parasitism, potential for human infection through fish consumption: puertos de Arica e Iquique, Chile
- Anisakis sp., larva
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Belone belone: Finland
- Anisakis sp.
Hauck, A. K., 1977, J. Parasitol., v. 63 (3), 515-519
Anisakis sp. larvae in Clupea harengus palasi from Yaquina Bay, Oregon, effects of various methods of handling and processing (fresh, frozen, brine, cold smoked, and cold smoked-gibbed) on migration into and survival in flesh of fish, implications for transmission to humans
- Anisakis larvae, illus.
Hsieh, H. C.; and Chen, E. R., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 567-568 [Demonstration]
Anisakis larvae, penetration into stomach wall of experimentally infected Macaca cyclopis within 6 hours of infection per os
- Anisakis sp.
Jackson, G. J.; Bier, J. W.; and Payne, W. L., 1976, Tr. Am. Micr. Soc., v. 95 (2), 264 [Abstract]
Anisakis sp. and Phocanema sp. in miniature pigs (exper.) (stomach), course of infection
- Anisakis spp.
Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Anisakis sp.
Korotaeva, V. D., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 89-96
Enophrys diceraus
Icelus spiniger
(body cavity of all): all from Sea of Japan
- Anisakis sp.
McVicar, A. H., 1977, J. Helminth., v. 51 (1), 11-21
intestinal helminths of Raja naevus, incidence, intensity, pattern of infection with host age and sex, geographical differences in composition of parasite burden
Raja naevus (cyst in intestine wall): off Aberdeen
- Anisakis sp.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Thunnus thynnus
Euthynnus affinis
Auxis thazard
all from South China Sea
- Anisakis sp.
Matthews, B. E., 1977, Parasitology, v. 75 (2), xii-xiii [Abstract]
Anisakis sp. larvae produce secretions which contain one fraction with enzymic activity, this proteolytic activity along with body movements may account for mechanism of migration
- Anisakis sp. (Dujardin 1845)
Mehl, J. A. P., 1970, N. Zealand J. Marine and Freshwater Research, v. 4 (3), 241-247
Thyrsites atun (flesh): eastern Cook Strait, New Zealand
- Anisakis sp.
Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (1), 267-277
helminths, incidence and intensity of infection in young and adult Oncorhynchus nerka, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Anisakis sp.
Popov, V. N., 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
age dynamics of infection
Histriophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island

- Anisakis sp. (Type I)
 Sakaguchi, Y.; and Katamine, D., 1971, *Nettai Igaku (Trop. Med.)*, v. 13 (4), 159-169
 anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
Coryphaena hippurus: sea near Nagasaki
Sillago japonica: "
Todarodes pacificus: "
Decapterus kuroides: "
Saurida tumbil: East and South China Seas
Muraenesox cinereus: East China Sea
Argyrosomus argentatus: "
Decapterus maruadsi: East and South China Seas
Pseudosciaena crocea: East China Sea
Zeus japonicus: "
- Caranx equula*: East and South China Seas
Fugu vermicularis vermicularis: East China Sea
Uranoscopus japonicus: "
Inegocia meerdervoorti: "
Upeneus bensasi: East and South China Seas
Trachurus japonicus: East China Sea
Taius tumifrons: "
Lepidotrigla microptera: "
Branchiostegus japonicus japonicus: "
Bembras japonicus: "
Nemipterus virgatus: South China Sea
Lutjanus sebae: "
Rachycentron canadum: "
Priacanthus sp.: "
Abalistes stellatus: "
Lutjanus basmira: "
- Pristipomoides sieboldi*: "
Megalopsis cordyla: "
Epinephelus septemfasciatus: "
Clidoderma asperrimum: "
Plectorhynchus pictus: "
Ilisha elongata: "
Lethrinus haematopterus: "
Tachysurus falcaurus: "
Siganus fuscens: "
Lepidotus brama: "
- Anisakis sp., *illus.*
 Sapunar, J.; Doerr, E.; and Letonja, T., 1976, *Bol. Chileno Parasitol.*, v. 31 (3-4), 79-83
 human anisakiasis, increasing world wide problem with increased use of fish for food, suggested control measures; case report of woman who expelled *Anisakis sp.* from throat after eating raw fish, elimination of second worm in feces after mebendazole therapy: Santiago, Chile
- Anisakis sp.
 Shakhmatova, V. I., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 277-289
Martes martes (intestine): Karelia
- Anisakis sp.
 Smith, J. W., 1971, *Nature* (5330), v. 234, 478
Thysanoessa inermis
Thysanoessa longicauda
 (haemocoel of all): all from northern North Sea
- Anisakis sp. larvae Type I
 Sommerville, R. I.; and Davey, K. G., 1976, *Internat. J. Parasitol.*, v. 6 (5), 433-439
Anisakis sp. larva, cuticle formation and ecdysis in vitro, development restarted by physico-chemical stimuli (effect of different media, carbon dioxide, storage, temperature), feeding does not occur until after moulting
- Anisakis spp. larvae
 van Thiel, P. H., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 75-85
 human anisakiasis, current status, review (diagnosis, treatment, epidemiology, distribution, definitive and intermediate hosts)
- Anisakis sp. probably *Anisakis simplex* (Rudolphi)
 Vooren, C. M.; and Tracey, D., 1976, *N. Zealand J. Marine and Freshwater Research*, v. 10 (3), 499-509
 incidence, intensity, age of host
Cheilodactylus macropterus (mesenteries, liver, intestinal wall, stomach wall): New Zealand
- Anisakis marina
 Farstad, L., 1975, *Norsk Vet.-Tidsskr.*, v. 86 (4), 247-253
Anisakis marina, life cycle, morphology, clinical aspects of human disease, diagnosis, control, review
- Anisakis marina (L.), *illus.*
 Reichenbach-Klinke, H. H., 1975, *Fisch u. Umwelt* (1), 113-121
 Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Anisakis physeteris
 van Thiel, P. H., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 75-85
Kogia breviceps: Shinnecock Inlet, Long Island, USA
- Anisakis (*Pseudanisakis*) *rotundata* (Rud.) of
 Layman & Borovkova (1926)
 Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
 as syn. of *Pseudanisakis tricupola* nom. nov.
- Anisakis simplex (Rudolphi, 1809), *illus.*
 Bakke, T. A.; and Barus, V., 1975, *Norwegian J. Zool.*, v. 23 (3), 183-191
 description, seasonal occurrence, sex of host
Larus canus (alimentary canal): Agdenes area, Norway
- Anisakis simplex, larvae
 Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (1), 7-31
 nematodes of *Larus canus*, age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway

- Anisakis simplex*, *illus.*
Beverley-Burton, M.; Nyman, O. L.; and Pippy, J. H. C., 1977, *J. Fish. Research Bd. Canada*, v. 34 (1), 105-112
Anisakis simplex larvae, description, morphology with particular reference to excretory system; comparative morphology of larvae from *Clupea harengus harengus* and *Salmo salar* in widely separated areas of North Atlantic suggest that *Anisakis* larvae Type I is *A. simplex*, findings substantiated by acid phosphatase polymorphism studies
- Anisakis simplex*
Bonner, W. N., 1972, *Oceanogr. and Marine Biol. Ann. Rev.*, v. 10, 461-507
Halichoerus grypus (stomach): European waters
- Anisakis simplex* (Rudolphi)
Cannon, L. R. G., 1977, *Austral. J. Marine and Freshwater Research*, v. 28 (6), 717-722
Peponocephala electra (stomach): Moreton Island; Tweed Heads
- Anisakis simplex* (Rudolphi, 1809, det. Krabbe, 1878)
Dailey, M. D.; and Perrin, W. F., 1973, *Fish. Bull.*, National Oceanic and Atmos. Admin., v. 71 (2), 455-471
incidence related to age of host
Stenella graffmani
S. cf. *S. longirostris*
(stomachs of all): all from eastern tropical Pacific
- Anisakis simplex*, *illus.*
D'Aubert, S.; Cattaneo, P.; and Ardemagni, A., 1976, *Arch. Vet. Ital.*, Milano, v. 27 (1-2), 16-18
possible methods for control, massive infestation, *Scomber scombrus* (muscle, abdominal cavity): imported from Danimarca (porto di Frederikshaven, Aalbaeksgskagen)
- Anisakis simplex*
Margolis, L.; and Beverley-Burton, M., 1977, *Internat. J. Parasitol.*, v. 7 (4), 269-273
response of *Mustela vison* to experimentally administered larvae collected from *Merluccius productus*, non-anaesthetized mink react violently with vomiting and/or defecation and eliminate all larvae, anaesthetized mink underwent violent 'heaving' but retained larvae for longer and some larval penetration was seen, recommended that marine fish be cooked or frozen before feeding to ranch mink
- Anisakis simplex* (=marina)
van Thiel, P. H., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 75-85
human anisakiasis, current status, review (diagnosis, treatment, epidemiology, distribution, definitive and intermediate hosts)
Halichoerus grypus: Rockhall, Scotland
Phocaena phocaena: Montrose Bay, eastern Scotland
Tursiops truncatus: Frisian isle of Ter-schelling
- Anisakis typica* (Diesing)
Cannon, L. R. G., 1977, *Austral. J. Marine and Freshwater Research*, v. 28 (6), 717-722
Peponocephala electra (stomach): Moreton Island; Tweed Heads
- Anisakis typica* (Rudolphi 1809 det. Krabbe 1878)
Forrester, D. J.; and Robertson, W. D., 1975, *J. Parasitol.*, v. 61 (5), 922
Steno bredanensis (forestomach, intestine): sandbar 6 miles southeast of the mouth of the Suwannee River in the Gulf of Mexico
- Anisakis typica*
van Thiel, P. H., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 75-85
human anisakiasis, current status, review (diagnosis, treatment, epidemiology, distribution, definitive and intermediate hosts)
- Ankylostomes. See Ancylostomes.
- Annulofilaria* G. N. Hsu, 1957
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Thelazia* Bosc, 1819
- Annulospira* Jairajpuri & Siddiqi, 1969
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Ceratospira* Schneider, 1866
- Anoplostrongylineae Chandler, 1938
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae
includes: *Anoplostrongylus* (type genus); *Adolpholutzia*; *Biacantha*; *Bidigiticauda*; *Bradypostrongylus*; *Brevigraphidium*; *Caenostrongylus*; *Cheiropteronema*; *Dasypostrongylus*; *Delicata*; *Didactyluris*; *Filicapitis*; *Fontesia*; *Graphidiops*; *Histiostrongylus*; *Maciela*; *Moennigia*; *Neohistiostrongylus*; *Paragraphidium*; *Parahistiostrongylus*; *Spinestrongylus*; *Torrestrongylus*; *Trichohelix*; *Tricholeiperia*; *Trichotravassosia*; *Trifurcata*; *Tupaiostrongylus*
- Anoplostrongylus* Boulenger, 1926 (type genus of subfam.)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Anoplostrongylineae
- Anoplostrongylus* (*Tupaiostrongylus*) *liei* (Dunn, 1963) n. comb., *illus.*
Durette-Desset, M. C.; and Chabaud, A. G., 1975, *Ann. Parasitol.*, v. 50 (2), 173-185
redescription
Tupaia glis (intestin, poumons): Ulu Gombak FR Selangor, Malaisie; 15 miles de Sandakan-Sepilok, Borneo
Tupaia minor (intestin): Ulu Gombak FR Selangor, Malaisie
T. tana (intestin): 15 miles de Sandakan-Sepilok, Borneo

- Antennocara Wassilkowa, 1926
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Schistorophus Railliet, 1916
- Aplectana acuminata
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Rana temporaria
Bufo viridis
Bufo bufo
all from Macedonia, Yugoslavia
- Aplectana acuminata (Schränk, 1788) Railliet et Henry, 1916
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Aplectana itzocanensis Bravo, 1943
Caballero Deloya, J., 1974, An. Inst. Biol., Univ. Nac. Mexico, s. Zool., v. 45 (1), 45-50
as syn. of Oxyssomatium itzocanensis (Bravo, 1943) Skrjabin, 1961
- Aplectana lynae n. sp., illus.
Kennedy, M. J., 1977, Canad. J. Zool., v. 55 (3), 630-634
Rana aurora aurora (large intestine): Stave Lake, British Columbia, Canada
- Aplectana schneideri (Travassos, 1931), illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Bufo bufo (debelo crijevo): Yugoslavia
- Aplectana uropeltidarum sp. nov., illus.
Cruz, H.; and Ching, C. C., 1975, Ann. Parasitol., v. 50 (3), 339-349
Rhinophis drummondhayi (rectum): above Watawala railway station; Namunukula area
Uropeltis melanogaster (rectum): hills above Kandy
U. phillipsi (rectum, small intestine): Gammaduwa
Rhinophis philippinus (rectum): Pallatenne near Gammaduwa
R. blythi (rectum): Talawakele
Teretrurus sanguineus (rectum): Nalumukku Estate, Mandjölai
- Aprocta sp.
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coeruleus (air sacs): Florida
- Aprocta cylindrica Linstow, 1883, illus.
Quentin, J. C.; Troncy, P. M.; and Barre, N., 1976, Ann. Parasitol., v. 51 (1), 83-93
Aprocta cylindrica, description of adult, life cycle, growth, development, larval morphogenesis
Quelea quelea quelea (cavite orbitaire): Tchad
Ploceus capitalis (cavite orbitaire): Tchad
P. cucullatus (cavite orbitaire): Tchad
Euplectes orix (cavite orbitaire): Tchad
Locusta migratoria (exper.)
- Aprocta turgidae Stossich, 1902
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus: Azov Sea
- Aproctella stoddardi
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (body cavity)
Colinus virginianus
all from Florida
- Aproctidae Skrjabin et Schikobalova, 1945
Jurasek, V., 1977, Biologia, Bratislava, s. B, Zool. (1), v. 32 (2), 105-109
diagnosis
- Aproctidae gen. sp. 2 Oschmarin, 1963
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 3-6
as syn. of Sarconema pseudolabiata nov. sp.
- Aproctinae gen. sp. Sonin et Borgarenko, 1965
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 3-6
as syn. of Sarconema pseudolabiata nov. sp.
- Aproctoidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Spirurina
key
- Aproctoidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
- Apteragia Jansen, 1958
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
synonymy
- Araeolaimida DeConinck & Schuurmans Stekhoven, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Chromadoria
includes: Araeolaimina; Tripyloidina
- Araeolaimina DeConinck, 1965
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Araeolaimida
- Arduenna Railliet & Henry, 1911
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Ascarops van Beneden, 1873

- Arthrocephaloides longespiculum* (Maplestone, 1931) Yamaguti, 1961
Balasingam, E., 1965, *Med. J. Malaya*, v. 20 (2), 165-166
development of provisional and definitive buccal capsule in *Arthrocephaloides longespiculum*
- Arthrocephalus lotoris* (Schwartz, 1925) Chandler, 1942
Barnstable, R. W.; and Dyer, W. G., 1974, *Tr. Illinois State Acad. Sc.*, v. 67 (4), 451-460
as syn. of *Placoconus lotoris* (Schwartz, 1925) Webster, 1956
- Arthrocephalus lotoris*
Bartsch, R. C.; and Ward, B. C., 1976, *Vet. Path.*, v. 13 (4), 241-249
raccoons (jejunum, ileum): southeastern Florida
- Arthrocephalus lotoris*
Georgi, J. R.; et al., 1976, *Cornell Vet.*, v. 66 (3), 309-323
Procyon lotor: North Rose, Wayne County, New York
- Arthrostroma Cameron* 1927
Yoshida, Y.; and Arizono, N., 1976, *J. Parasitol.*, v. 62 (5), 766-770
key to species, measurements, includes: *Arthrostroma spatulatum* Jansen, 1968; *A. vampira* Schmidt and Kuntz, 1968; *A. cheni* Kou, 1958; *A. felineum* Cameron, 1927; *A. longespiculum* (Maplestone, 1931); *A. tunkanati* Inglis and Ogden, 1965; *A. miyazakiense* (Nagayosi, 1955) comb. n.
- Arthrostroma longespiculum* (Maplestone, 1931), *illus.*
Setasuban, P., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (3), 382-385
Arthrostroma longespiculum, scanning electron microscopic study of parasite surfaces
- Arthrostroma longespiculum* (Maplestone, 1931), *illus.*
Setasuban, P.; and Vajrasthira, S., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (2), 202-207
morphometric data
Paradoxurus hermaphroditus (small intestine): Nakorn Nayok, Thailand
- Arthrostroma miyazakiense* (Nagayosi 1955) comb. n., *illus.*
Yoshida, Y.; and Arizono, N., 1976, *J. Parasitol.*, v. 62 (5), 766-770
redescription, key, measurements
Syn.: *Necator miyazakiensis* Nagayosi 1955
Nyctereutes procyonoides (small intestine): Miyazaki and Kyoto Prefectures, Japan
- Artionema indica* Dutt, 1963
Subramanian, G.; and Srivastava, V. K., 1973, *Riv. Parassitol.*, Roma, v. 34 (1), 59-62
as syn. of *Setaria cervi* (Rudolphi, 1819)
- Arundelia* n. gen.
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
Trichonematidae: Cloacininae
tod: *A. dissimilis* (Johnston & Mawson)
n. comb.
- Arundelia dissimilis* (Johnston & Mawson) n. comb. (tod), *illus.*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
description, syn.: *Macropostrongylus dissimilis* Johnston & Mawson, 1939
Wallabia bicolor: Keyneton, Bemm River, Yarra Valley, and Dartmouth, Victoria
- Ascariasis
Angate, Y.; et al., 1974, *Medecine Afrique Noire*, v. 21 (1), 61-65
symptoms of acute abdomen resulting from human intestinal parasites, medical and surgical care, case reports: Abidjan, Ivory Coast
- Ascariasis
Ashizawa, H.; et al., 1975, *Bull. Fac. Agric. Miyazaki Univ.*, v. 22 (2), 221-229
ascariasis, swine, pathology, biliary tract
- Ascariasis
Baljozovic, A.; and Popovic, M., 1971, *Med. Glasnik*, v. 25 (11-12), 392-393
perforation of man's appendix caused by ascariasis, case report: Beograd
- Ascariasis
Beheshti, G.; et al., 1974, *Pahlavi Med. J.*, v. 5 (1), 94-99
biliary ascariasis in 2-year-old boy, fatal illness after complication of generalized peritonitis, case report: Iran
- Ascariasis
Bessieres-Cathala, M. H.; et al., 1975, *Medicine et Malad. Infect.*, v. 5 (12), 592-596
micromethod of complement fixation in diagnosis of various human parasites
- Ascariasis
Biagi, F., 1975, *Progr. Drug Research*, v. 19 23-27
human hookworm and trichuriasis successfully treated with bitoscanate in clinical trials; mixed results when used to treat ascariasis in comparison trials with dithiazanine and piperazine
- Ascariasis
Biagi, F.; Alvarez, R.; and Delgado, R., 1972, *Prensa Med. Mexicana*, v. 37 (5-6), 221-223
human intestinal helminths, clinical trials with pyrantel pamoate, good results
- Ascariasis
Biagi, F.; Lopez, R.; and Viso, J., 1975, *Progr. Drug Research*, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review

- Ascariasis**
Bomb, B. S.; and Bedi, H. K., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 358 [Letter]
ascariasis, 12-year-old girl treated with piperazine citrate at dosage recommended by manufacturer, neurotoxic side effects
- Ascariasis**
Bulucea, D., 1974, Med. Int., Bucuresti, v. 26 (2), 249-256
ascariasis and Trichocephalus as causes of massive eosinophilia
- Ascariasis**
Bulucea, D.; and Rarinca, A., 1972, Microbiol., Parazitol., Epidemiol., Bucuresti, v. 17 (2), 169-174
mixed ascariasis and Trichocephalus infection in woman complicated by pleuro-pulmonary staphylococcal pathology, probable parasitic allergic reactions involved in etiology, case report
- Ascariasis**
Diop, B.; and Bao, O., 1974, Medecine Afrique Noire, v. 21 (1), 31-40
human intestinal helminths, clinical indications for treatment, suggested dosage, efficacy, tolerances, possible toxicities
- Ascariasis**
Diouf, A. B.; et al., 1975, Medecine Afrique Noire, v. 22 (6), 453-460
human helminthiasis, statistics of 103 surgical parasitic cases over 10-year period
- Ascariasis**
Egorov, V. I., 1968, Botan. i Zool. Issled. Dal'nem Vostoke, v. 2, 250-255
ascariasis, trichuriasis, oesophagostomiasis, swine, single and mixed infections in various combinations, control studies, best system used dehelminthization 4 times a year
- Ascariasis**
Egorov, V. I., 1968, Botan. i Zool. Issled. Dal'nem Vostoke, v. 2, 256-260
ascariasis, trichuriasis, swine, epizootiology, seasonal distribution of depositing eggs and larvae in barns and pastures: Primor'e
- Ascariasis**
El-Antably, S. A.; and El-Hoshy, M. H., 1975, Bull. Ophth. Soc. Egypt, v. 68, 275-280
human ascariasis, ancylostomiasis and schistosomiasis in association with soft cataracts, clinical aspects, possible relationships with nutritional deficiencies: Egypt
- Ascariasis**
Grigorescu-Sido, P.; et al., 1972, Pediatria, Bucuresti, v. 21 (2), 115-124
human gastrointestinal parasitic diseases, possible relationships with lactase deficiency in children
- Ascariasis**
Gument, J. M., 1973, Nouv. Presse Med., v. 2 (32), 2125 [Letter]
human ascariasis, possible associations with bronchial asthma and rhinitis
- Ascariasis**
Ismail, D.; et al., 1976, Paediat. Indonesiana, v. 16 (9-10), 391-395
ascariasis in children, pyrantel pamoate more effective than either tetramisole or piperazine as therapy but all drugs reduced egg counts in feces by 80%: Indonesia
- Ascariasis**
Joshi, V. S.; et al., 1975, Pediat. Clin. India, v. 10 (3), 171-173
children with ascariasis or enterobiasis, successful therapy with vanpar: Pune, India
- Ascariasis**
Kim, C. W., 1975, Progr. Clin. Path., v. 6, 267-288
extensive review of techniques used to diagnose human parasitic diseases
- Ascariasis**
Kovalenko, I. I.; and Sakachina, V. I., 1976, Veterinariia, Kiev (43), 86-88
ascariasis and heterakiasis, chickens, mixed infection; furidine, furidine + phenothiazine, and piperazine + phenothiazine, good results; phenothiazine or furidine alone, poor results
- Ascariasis**
Meakins, R. H.; Carswell, F.; and Harland, P. S. E. G., 1977, Parasitology, v. 75 (2), xxxii [Abstract]
helminthiasis in school children, association with malnutrition: rural Tanzania
- Ascariasis**
Meister, A.; Tschaepe, M.; and Schroetter, E., 1977, Pharmazie, v. 32 (3), 174-177
human ascariasis, structure-activity relationships of phenol derivatives, possible ovicides
- Ascariasis**
Nemeseri, L., 1972, Parasitol. Hungar., v. 5, 217-224
coccidiosis and ascaridiasis in domestic animals and poultry, furidin and rodanin assayed for possible prophylactic use in conditions of large scale animal husbandry
- Ascariasis**
Nitzulescu, V.; et al., 1971, Pediatria, Bucuresti, v. 20 (2), 169-174
human ascariasis, L-tetramisole successfully used for mass therapy
- Ascariasis**
Nitzulescu, V.; and Cherman, I., 1974, Pediatria, Bucuresti, v. 23 (3), 271-272
mixed human infections of ascariasis and oxyuriasis, combantrin an effective anthelmintic, limited clinical trial

- Ascariasis**
Ogawa, K. M.; and Mathias, A. L., 1967, *Maternidade e Infancia*, v. 26 (4), 381-385
human infection of liver and biliary tract, clinical review, case reports
- Ascariasis**
Pasticier, A., 1975, *Nouv. Presse. Med.*, v. 4 (37), 2659-2660
single and mixed parasitic infections in humans, good results with pyrantel pamoate
- Ascariasis**
Patterson, R.; et al., 1975, *Pediatrics*, *Am. Acad. Pediat.*, v. 56 (3), 417-420
differentiation of visceral larva migrans caused by toxocariasis from that caused by ascariasis using radioimmunoassay to demonstrate serum immunoglobulins
- Ascariasis**
Pawlowski, Z.; and Chodera, L., 1975, *Polski Tygod. Lekar.*, v. 30 (24), 1035-1037
ascariasis, trichuriasis, ancylostomiasis, humans, clinical trials show mebendazole to be effective drug for all 3 infections: Poznan, Poland
- Ascariasis**
Radermecker, M.; et al., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months
- Ascariasis**
Sankale, M.; Diop, B.; and Bao, O., 1974, *Medecine Afrique Noire*, v. 21 (2), 149-152
ascariasis, ancylostomiasis, human, successful trials with pyrantel pamoate: Senegal
- Ascariasis**
Savanat, T.; et al., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 149-154
determinations of total serum IgE levels in humans with amoebic liver abscess or other parasitic infections
- Ascariasis**
Shah, P. M.; Junnarkar, A. R.; and Khare, R. D., 1975, *Progr. Drug. Research*, v. 19, 136-146
impact of periodic deworming on nutritional status of preschool children, evaluation of including periodic deworming in National Nutrition Programmes of India
- Ascariasis**
Singh, O. P.; and Rai, P., 1975, *Indian J. Animal Research*, v. 9 (1), 1-7
ascariasis, seasonal incidence, zebu and buffalo calves: India
- Ascariasis**
Spay, G., 1974, *Medecine Afrique Noire*, v. 21 (1), 55-58
surgical emergencies and manifestations in course of human intestinal ascariasis, differential diagnosis: Caboul, Afganistan
- Ascariasis**
Tarabini Castellani, G., 1970, *Minerva Gastroenterol.*, v. 16 (1), 45-49
human intestinal parasites, clinical trials using FI 6518 with and without thymol, acceptable drug for mass therapy: Somalia
- Ascariasis**
Thomas, J.; et al., 1971, *Medecine Trop.*, v. 31 (2), 177-181
bephenium hydroxynaphthoate, piperazine and thiabendazole in comparative clinical tests, institutionalized children with protein deficiency
- Ascariasis**
Trifonov, T.; and Velkov, D., 1975, *Vet. Sbirka*, v. 73 (11), 16-17
ascariasis, trichuriasis, oesophagostomiasis, pigs, dehelminthization with suiverm or hygromix in feed, economically effective
- Ascariasis**
Vakil, B. J.; and Dalal, N. J., 1975, *Progr. Drug Research*, v. 19, 166-175
comparative efficacy of newer anthelmintics in treating various human intestinal helminths, review
- Ascariasis**
Vanneste, J. A. L.; et al., 1975, *Nederl. Tijdschr. Geneesk.*, v. 119 (48), 1899-1901
ascariasis, child treated with piperazine showed signs of neurologic toxicity: Netherlands
- Ascariasis**
Weinand, H. A.; Broicher, K.; and Lieb, W., 1974, *Med. Welt*, v. 25 (31-32), 1283-1285
ascariasis, child, differential diagnosis by laparoscopy and histologic examination of liver
- Ascarides**
Bussieras, J., 1977, *Rev. Med. Vet.*, Toulouse, v. 128 (3), 343-348
strongyles, ascarids, horses, pyrantel pamoate orally or by naso-oesophageal catheter, no bad effects
- Ascarides**
Duewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Ascarides**
Genchi, C.; and Locatelli, A., 1974, *Atti Soc. Ital. Sc. Vet.*, v. 28, 862-863
dog helminth eggs, contaminated samples from public parks, potential source of infection for humans and domestic animals: Milan

Ascarides

- Lange, E.; and Tomsen, K., 1976, Tierarztl. Umschau, v. 31 (10), 441-443
 Strongyloides, strongyles, ascarides, horses, fenbendazole (aqueous suspension, granulate), field trials

Ascarides

- Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
 fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date

Ascaridia

- Jacobs, D.; and Schulze, H. W., 1977, Prakt. Tierarzt, v. 58 (1), 46-48
 pig parasites, natural infections, ivermectin and dichlorvos effective in field testing

Ascaridia

- Khanum, Z.; Farooq, M.; and Zuberi, H. B., 1976, Norwegian J. Zool., v. 24 (3), 191-194
 key to species from *Columba livia*

Ascaridia-type, illus.

- Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
 helminths of pet birds, diagnosis of eggs in fecal examination

Ascaridia

- Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
 anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report

Ascaridia columbae

- Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
 fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology

Ascaridia columbae (Gmelin, 1790) Travassos, 1913

- Khanum, Z.; Farooq, M.; and Zuberi, H. B., 1976, Norwegian J. Zool., v. 24 (3), 191-194
 key

Ascaridia compar

- Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix
Tetrao urogallus
 all from CSSR

Ascaridia cylindrica

- Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix
Tetrao urogallus
 all from CSSR

Ascaridia dissimilis

- Colglazier, M. L., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 60-61
Ascaridia dissimilis, *Heterakis gallinarum*, and *Capillaria obsignata*, turkeys, levamisole administered in drinking water effective as anthelmintic

Ascaridia dissimilis

- Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (duodenum; lower small intestine): Florida

Ascaridia dissimilis (Vigueras, 1931), illus.

- Mutafova, T., 1976, Ztschr. Parasitenk., v. 48 (3-4), 239-245
Ascaridia dissimilis, *A. galli*, gametogony, spermatogenesis, mitotic and male meiotic karyotypes, both species similar

Ascaridia dissimilis

- Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: southeastern United States

Ascaridia dissimilis

- Vasilev, I.; et al., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 59-72
Ascaridia galli, *A. dissimilis*, turkey poults, course of infection, pathogenesis (hematology, histology, histochemistry, biochemistry)

Ascaridia galli

- Amer, A. A.; El-Amrousi, S.; and El-Hammady, H. Y., 1976, Vet.-Med. Nachr. (1), 18-31
Ascaridia galli, chickens (nat. and exper.), tetramisole, niclosamide, efficacy of drugs administered singly and in combination, drug trials

Ascaridia galli

- Ansari, A. A.; Khan, M. A.; and Ghatak, S., 1976, Exper. Parasitol., v. 39 (1), 74-83
Ascaridia galli extracts, isolation of trypsin and chymotrypsin inhibitors, partial purification and some physical and chemical properties

Ascaridia galli, illus.

- Araujo, P.; and Bressan, M. C. R. V., 1977, Ann. Parasitol., v. 52 (5), 531-537
Ascaridia galli larvae undergo two molts before hatching, infective larva is therefore third stage

Ascaridia galli (Schrank, 1788) Freeborn, 1923

- Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
 fowl, domestic
 fowl, desi
 all from Punjab State, India

- Ascaridia galli*
Birova-Volosinovicova, V., 1974, *Biologia*, Bratislava, s. B, Zool. (1), v. 29 (2), 139-149
Ascaridia galli in chicks treated with hormones (ACTH and cortisone), low doses produced higher levels of parasitism, increased larval burden in first 5 days after infection and eliminated usual sudden decrease of larval count in chick intestine; high dose of cortisone produced low level of parasitism
- A[*Ascaridia*] *galli*
Birova-Volosinovicova, V., 1974, *Biologia*, Bratislava, s. B, Zool. (1), v. 29 (2), pp. 151-157
A[*Ascaridia*] *galli* in chicks treated with hormones (ACTH and cortisone), blood values (erythrocytes, leucocytes, haemoglobin, leucogram) same as normal chicks or infected, untreated chicks
- Ascaridia galli*
Bogoiavlenskii, Iu. K.; and Koroleva, N. A., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 21-29
Ascaridia galli, *Hystrichis tricolor*, comparison of micromorphology and histochemistry of hypodermal-muscular sac during pre-imaginal development
- Ascaridia galli*
Chakraborty, A. K.; Mehta, R. K.; and Datta, I. C., 1976, *Indian J. Exper. Biol.*, v. 14 (5), 585-587
Ascaridia galli, tetramisole, carbohydrate utilization, neuro-transmission, enzymatic activity
- Ascaridia galli*
Cruthers, L. R.; al-Khateeb, G. H.; and Hansen, M. F., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 119-121
Ascaridia galli, *Heterakis gallinarum*, *Capillaria obsignata*, chickens, levamisole in drinking water
- Ascaridia galli*
Dick, J. W.; Leland, S. E., jr.; and Hansen, M. F., 1973, *Tr. Am. Micr. Soc.*, v. 92 (2), 225-230
Ascaridia galli, 4 hatching techniques for embryonated eggs compared, subsequent cultivation in medium Ae and supplemented medium Ac
- Ascaridia galli*
Dubinsky, P.; Lestan, P.; and Rybos, M., 1976, *Biologia*, Bratislava, s. B, Zool. (4), v. 31 (11), 829-838
Ascaridia galli larvae, chickens fed with amino acid-deficient cereal diet, effect on components of blood serum
- Ascaridia galli*
Gogoi, A. R., 1975, *Kerala J. Vet. Sc.*, v. 5 (2), 131-134
fowl: Assam
- Ascaridia galli*
Hurwitz, S.; Shamir, N.; and Bar, A., 1973, *Isotopes and Radiation Parasitol.* III, 61-65
Ascaridia galli-infected chicks, absorption and digestion of protein and absorption of phosphate from intestine
- Ascaridia galli*
Kaushik, R. K., 1977, *Indian J. Animal Sc.*, v. 45 (11), 1975, 911-912
Ascaridia galli, chicks (exper.), serum protein changes
- Ascaridia galli*
Kovalenko, I. I., 1975, *Veterinariia*, Kiev (42), 92-94
Ascaridia galli, *Heterakis gallinarum*, length of development studied as basis for timing of anthelmintic treatment
- Ascaridia galli* (Schrank, 1788)
Kulkarni, D.; Rao, Y. V. B. G.; and Rao, B. V., 1977, *Indian Poultry Rev.*, v. 8 (17), 14-16
Ascaridia galli, chickens (exper.), tetramisole, morantel tartarate, wopell, anthelmintic efficacy, laboratory trial, tetramisole and morantel tartarate most effective against immature and mature stages, wopell only partially effective
- Ascaridia galli*
Lal, Jawahar; et al., 1976, *Indian J. Physiol. and Pharmacol.*, v. 20 (2), 64-68
Ascaridia galli, in vitro anthelmintic activity of 10 medicinal plants, comparison with piperazine hexahydrate
- Ascaridia galli* Schrank
Lal, J.; Subramanian, G.; and Sabir, M., 1977, *Indian J. Animal Sc.*, v. 45 (10), 1975, 807-808
Ascaridia galli, chicks (exper.), mebendazole, highly effective against intestinal larvae, ineffective against tissue-phase larvae
- Ascaridia galli*
Lee, Y. C.; et al., 1976, *J. Chinese Soc. Vet. Sc.*, v. 2 (1), 40-43
Ascaridia galli eggs, *Citrobacter freundii* isolated, failure to recover *Salmonella pullorum* from worm eggs after artificial infection of chickens
- Ascaridia galli*
Lee, Y. C.; Liu, C. K.; and Liu, C. C., 1975, *J. Chinese Soc. Vet. Sc.*, v. 1 (2), 79-82
Ascaridia galli eggs did not transmit *Salmonella pullorum* from chickens
- Ascaridia galli*
Lesin'sh, K. P.; et al., 1975, *Latvijas PSR Zinat. Akad. Vestis* (340) (11), 27-30
helminths, chickens, effect of host age and method of rearing on infestation: Latvian SSR
- Ascaridia galli*
Leutskaia, Z. K., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 88-90
Ascaridia galli antigen-immunized chickens, content of various forms of vitamin A in mitochondria and microsomes of spleen cells, possible role of vitamin A in antibody formation and immune process
- Ascaridia galli*
Leutskaia, Z. K., 1975, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 25, 71-90
Ascaridia galli in chicks, role of vitamin A in immunogenesis (increase of resistance, specific immunity, antibody formation), specifically, role in immunization

- Ascaridia galli*
Long, P. L., 1977, *Vet. Rec.*, v. 100 (16), 342
Ascaridia galli, chickens, incidence in earth
floored broiler house, piperazine citrate,
good results: Britain
- Ascaridia galli*, *illus.*
Mayaudon T., H.; and Bendjoya Cases, J.,
1974, *Rev. Med. Vet. y Parasitol.*, Maracay,
v. 25 (1-8), 1973-1974, 32-66
prevalence in *Gallus (Gallus) domesticus*:
Estado Aragua, Venezuela
- Ascaridia galli*
Mirzayans, A., 1975, *J. Vet. Fac. Univ. Tehran*,
v. 30 (4), 5
chickens: area of Tehran, Iran
- Ascaridia galli* (Schrank, 1788), *illus.*
Mutafova, T., 1976, *Ztschr. Parasitenk.*, v. 48
(3-4), 239-245
Ascaridia dissimilis, *A. galli*, gametogony,
spermatogenesis, mitotic and male meiotic
karyotypes, both species similar
- Ascaridia galli* (Schrank, 1788)
Nauriyal, D. C.; and Rai, P., 1976, *Agra Univ.*
J. Research, v. 23 (3), 1974, 75-84
Ascaridia galli, immature stages, chickens
(exper.), serum enzymes (alkaline and acid
phosphatase, cholinesterase) and gamma glob-
ulin levels, histochemical changes, diagnos-
tic value, results indicate significant rise
of serum alkaline phosphatase and parallel
histochemical changes; acid phosphatase,
cholinesterase, and gamma globulin not use-
ful in diagnosis
- Ascaridia galli*
Nol, M.; et al., 1975, *Rev. Avicult.*, v. 19
(3), 351-352
Ascaridia galli, *Heterakis gallinae*, poultry,
anemia, changes in leucocyte types, elevated
sedimentation rate, decrease in albumin, in-
crease in alpha and beta globulin
- Ascaridia galli*
Okon, E. D., 1976, *Research Vet. Sc.*, v. 21
(1), 104
Ascaridia galli third stage larvae, fowls,
pyrantel tartrate, drug efficacy, good re-
sults
- Ascaridia galli*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1),
23-30
ethyl-6-ethoxybenzothiazole-2-carbamate,
evaluation of anthelmintic activity in pon-
ies, swine, lambs, and chickens
- Ascaridia galli*
Pav, J.; and Zajicek, D., 1974, *Veterinarstvi*,
v. 24 (11), 517-520
Lyrus tetrix
Tetrao urogallus
all from CSSR
- Ascaridia galli*
Pavlicek, J.; and Dykova, I., 1976, *Vet. Med.*,
Praha, v. 49, v. 21 (6), 353-358
Ascaridia galli, chickens of different age
groups, effectiveness of piperazine, te-
tramisole and metrifphonate, metrifphonate
entirely ineffective; tissue reaction of
host and number of worms as measures of
effectiveness
- Ascaridia galli* (Schrank, 1788)
Peirce, M. A.; and Bevan, B. J., 1973, *Vet.*
Rec., v. 92 (10), 261 [Letter]
Neophema elegans (small intestine): bird
farm in England
Platyercus eximius ceciliae (duodenum):
bred in Holland and imported into the
United Kingdom
- Ascaridia galli*
Radhakrishnan, C. V.; and Ebrahimina, A.,
1975, *J. Vet. Fac. Univ. Tehran*, v. 30 (4), 1-4
chickens (small intestine): Darab, Fars
Province, Iran
- Ascaridia galli* Schrank, 1788
Ramon Vericad, J.; and Sanchez Acedo, C.,
1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 267-
271
Buteo buteo
Alectoris rufa
all from Huesca, Alto Aragon
- Ascaridia galli*
Rybos, M.; Lestan, P.; and Dubinsky, P., 1974,
Biologia, Bratislava, s. B, *Zool.* (1), v. 29
(2), 129-132
Ascaridia galli, hydrocarbon metabolism ex-
cretory products, principally acetic acid in
23 day-old worms, proprionic acid in mature
worms
- Ascaridia galli* Schrank, 1788, *illus.*
Segida, G. V., 1969, *Trudy Gel'mint. Lab.*,
Akad. Nauk SSSR, v. 20, 133-145
Ascaridia galli, fine structure of the
peripheral sensory nervous system
- Ascaridia galli*
Shikhobalova, N. P.; and Korsak-Paruzhinskaia,
L. S., 1975, *Trudy Gel'mint. Lab.*, Akad. Nauk
SSSR, v. 25, 204-214
Ascaridia galli, *Ganguleterakis spumosa*,
Trichocephalus muris, ionizing radiation of
eggs, postradiation effect of high and low
temperatures on radiosensitivity as evidenced
by development; radiation as possible control
of helminths
- Ascaridia galli*
Stepanian, S. G.; et al., 1976, *Biol. Zhurnal*
Armenii, v. 29 (10), 82-86
Ascaridia galli, chickens, role of colamine
phosphate in heightening natural host re-
sistance (strengthened hemopoiesis, increased
vitamin C in blood plasma and tissue, in-
creased organic action of acid soluble
fraction of phosphate in blood)

- Ascaridia galli*
Subramanian, G.; and Singh, K. S., 1973, *Iso-
topes and Radiation Parasitol.* III, 67-71
Ascaridia galli, one-week old chicks,
attempted immunization with irradiated vs.
normal eggs
- Ascaridia galli*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26),
216-230
fenbendazole (available in 5 forms), ef-
ficacy against nematodes in various animals,
well tolerated with no apparent effects on
fertility or fetus, extensive summary of re-
sults to date
- Ascaridia galli*
Torres, P.; et al., 1974, *Bol. Chileno Parasi-
tol.*, v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Ascaridia galli*
Torres, P.; and Barriga, O. O., 1974, *Bol.
Chileno Parasitol.*, v. 29 (3-4), 79-85
Ascaris suum, *A. lumbricoides*, *Toxocara
mystax*, *Ascaridia galli*, comparative anti-
genic analysis by gel double diffusion and
immunoelectrophoresis
- Ascaridia galli*
Vasilev, I.; et al., 1973, *Izvest. Tsentral.
Khelmint. Lab.*, v. 16, 43-58
Ascaridia galli, chicks, course of infection,
pathogenesis (hematology, histology, histo-
chemistry, biochemistry)
- Ascaridia galli*
Vasilev, I.; et al., 1973, *Izvest. Tsentral.
Khelmint. Lab.*, v. 16, 59-72
Ascaridia galli, *A. dissimilis*, turkey
poults, course of infection, pathogenesis
(hematology, histology, histochemistry, bio-
chemistry)
- Ascaridia galli*
Walker, T. R.; and Farrell, D. J., 1976,
Brit. Poultry Sc., v. 17 (1), 63-77
Ascaridia galli, significant reduction in
metabolisability of dietary energy and
nitrogen retention of infected chickens;
inadequate amounts of vitamin A reduced
size of larvae and perhaps decreased their
numbers
- Ascaridia galli*
Young, P. L.; and Babero, B. B., 1975, *Proc.
Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an
important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
chickens (small intestine)
(all exper.)
- Ascaridia hermaphrodita*
Robinson, P. T.; and Richter, A. G., 1977, *J.
Zoo Animal Med.*, v. 8 (1), 23-26
Poicephalus robustus robustus, successful
treatment with levamisole phosphate paren-
terally: [San Diego Zoo]
- Ascaridia lucknowensis* sp. nov., illus.
Kumar, P.; and Gupta, S. P., 1975, *Current
Sc.*, Bangalore, v. 44 (24), 891
Centropus sinensis (intestine): Lucknow
- Ascaridia magnipapilla*
Pav, J.; and Zajicek, D., 1974, *Veterinarstvi*,
v. 24 (11), 517-520
Lyrus tetrix
Tetrao urogallus
all from CSSR
- Ascaridia numidae* Leiper, 1908
Fabiyl, J. P., 1972, *Bull. Epizoot. Dis.
Africa*, v. 20 (3), 235-238
Numida meleagridis galeata (intestine):
Vom area, Benue Plateau State, Nigeria
- Ascaridia qadrii* sp. n., illus.
Khanum, Z.; Farooq, M.; and Zuberi, H. B.,
1976, *Norwegian J. Zool.*, v. 24 (3), 191-194
key
Columba livia (intestine): Karachi Univer-
sity Campus, Karachi, Pakistan
- Ascaridia razia* Akhtar, 1937
Khanum, Z.; Farooq, M.; and Zuberi, H. B.,
1976, *Norwegian J. Zool.*, v. 24 (3), 191-194
key
- Ascaridia styplocerca* (Stossich, 1904)
Fabiyl, J. P., 1972, *Bull. Epizoot. Dis.
Africa*, v. 20 (3), 229-234
survey of helminths of chickens, comparison
of techniques of management (native exten-
sive, deep-litter (intensive) and semi-in-
tensive systems) on worm burden; suggested
preventive measures and treatment with
piperazine: Vom area, Benue-Plateau State,
Nigeria
- Ascaridiasis*. See *Ascariasis*.
- Ascaridida*
Chabaud, A. G., 1974, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and Will-
mott)(1), 6-17
Secernentea
key; key to superfamilies
includes: *Cosmocercoidea*; *Seuratoidea*; *Het-
erakoidea*; *Ascaridoidea*; *Subuluroidea*
- Ascaridida* Railliet & Henry, 1915
Maggenti, A. R., 1976, *Organ. Nematodes*
(Croll), 1-10
Secernentea
- Ascarididae* Baird, 1853
Hartwich, G., 1974, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Ascaridoidea
key; key to subfamilies
includes: *Toxocarinae*; *Multicaecinae*; *As-
caridinae*; *Angusticaecinae*
- Ascarididae* [sp.], larvae
Andrews, S. E.; and Threlfall, W., 1975, *Proc.
Helminth. Soc. Washington*, v. 42 (1), 24-28
Corvus brachyrhynchos (mid- and posterior
section of small intestine): insular New-
foundland

- Ascaridinae (Baird, 1853)
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascarididae
key; key to genera
includes: Parascaris; Ascaris; Lagochilascaris; Baylisascaris; Toxascaris
- Ascaridoidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Ascaridida
key
- Ascaridoidea
Gibson, D. I.; and Taylor, A. L., 1976, Parasitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Ascaridoidea
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
key to families
includes: Crossophoridae; Heterocheilidae; Anisakidae; Acanthocheilidae; Ascarididae
- Ascaridoidea
Petter, A. J., 1977, Ann. Parasitol., v. 52 (2), 151-158
evolutionary aspects of distribution of Ascaridoidea in mammals
- Ascaridoidea
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Ascarids, illus.
Church, E. M.; Wyand, D. S.; and Lein, D. H., 1975, Am. J. Vet. Research, v. 36 (3), 331-335
cerebrospinal nematodiasis, experimentally induced in *Oryctolagus cuniculus* with *Ascaris columbiana*, *A. suum*, or *Toxocara canis*, naturally occurring in *Sylvilagus floridanus* and *O. cuniculus*, clinical signs, gross and microscopic changes, duration of infection and parasite morphology and distribution in CNS, potential of rabbits as intermediate or paratenic hosts for ascarids of carnivorous origins
- Ascarids
Dolnikov, A. E.; and Melnikova, K. V., 1976, Gig. Sanitariia (10), 115-116
ascarid eggs, attempted dehelminthization of liquids by electrohydraulic effect
- Ascarids
Fagot, G.; Clery, P.; and Pascal, D., 1977, Prat. Vet. Equine, v. 9 (3), 169-170
ascarids, strongyles, *Oxyuris* in race horses, clinical trials with strongid-P most successful in eradicating strongyles: France
- Ascarids
Genchi, C., 1976, Arch. Vet. Ital., v. 27 (3-4), 98-99
dog intestinal helminth ova, incidence in soil samples in public parks, potential public health problem: Milan
- Ascarids
Greve, J. H.; and Paul, J. W., 1976, Vet. Med. and Small Animal Clin., v. 71 (12), 1737-1740, 1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Ascarids
Howes, H. L., jr., 1972, Proc. Soc. Exper. Biol. and Med., v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that *T. muris*-mouse infection could be test model for antiwhipworm studies
- Ascarids
Lazdynia, M. A., 1975, Latvijas PSR Zinat. Akad. Vestis (340) (11), 31-33
ascarid larvae, effect of worm metabolic products on establishment of infection, chickens administered volatile fatty acids had 2-3 times more larvae established compared to chickens not administered volatile fatty acids
- Ascariden
Manz, D.; and Dingeldein, W., 1974, Prakt. Tierarzt, v. 55 (8), 422-425
nematodes of European and exotic herbivores, Banminth, good results
Equus quagga
- Ascarids
Pavlov, A. V.; and Koshkina, L. A., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 106-109
ascarids, chicks, increased ATP-ase and sodium and chloride ions in body fluid of worms from hosts vaccinated before infection, possible relationships to cuticle permeability and transport system
- Ascarids
Petric, S. W., 1977, J. South African Vet. Ass., v. 48 (2), 105-107
Spirocerca lupi, *Filaroides osleri*, ascarids, dogs, gastrointestinal fibroscope, useful diagnostic aid
- Ascarid, illus.
Schueler, R. L., 1973, J. Wildlife Dis., v. 9 (1), 58-60
larval ascarids, probably *Ascaris columbiana*, in *Sciurus granatensis* (brain, lung) with clinical signs of neurological disease
- Ascarids
Sewell, M. M. H.; and Urquhart, H. R., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 287 [Demonstration]
technique for quantitative recovery of ascarid and *Trichuris* eggs from 100g samples of soil

Ascarids

- Shchukin, P. I., 1977, Izvest. Akad. Nauk SSSR, s. Biol. (4), 589-601
 ascarids, toxic substances in metabolic products producing changes in blood serum proteins in various strains of rats as reflecting genotypical differences among strains

Ascariden

- Witteborg, K., 1976, Tierarztl. Umschau, v. 31 (2), 72-74
 ascariasis, puppies of dogs of large breeds, pyrantel pamoate at two standard dosages, not successful, comparison with piperazine citrate, discussion of dosages

Ascaris

- Arfaa, F.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 230-233
 Ascaris, Trichostrongylus spp., Ancylostoma duodenale, evaluation of different methods of control of soil transmitted helminths (sanitation measures, mass-therapy, combined mass-therapy and sanitation) in villagers of Khuzestan, southwest Iran

Ascaris

- Aung-Thun-Batu; et al., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 167 [Letter]
 Ascaris larvae and ova in core of human bile-duct stones: Rangoon, Burma

Ascaris, illus.

- Bennet-Clark, H. C., 1976, Organ. Nematodes (Croll), 313-342
 mechanics of nematode feeding: anatomy of nematode oesophageal pump; feeding performance; theory of filling and emptying of nematode oesophagus; a model for filling cycle of oesophagus of Ascaris

Ascaris

- Biagi, F.; and Medina, J. A., 1974, Prensa Med. Mexicana, v. 39 (9-10), 440-443
 human intestinal parasites, pyrantel pamoate, metabolism and pharmaceutical action

Ascaris

- Biagi, F.; Smyth, J.; and Gonzalez, C., 1974, Prensa Med. Mexicana, v. 39 (1-2), 51-53
 human intestinal helminths, successful clinical trials using mebendazole, drug well tolerated with minimal side effects: Mexico

Ascaris, illus.

- Bradley, R. H.; and Burghardt, R. C., 1976, Proc. 34. Ann. Meet. Electron Microsc. Soc. America (Miami Beach, Florida, Aug. 9-13), 158-159
 Ascaris spermatozoa, immunocytochemistry of surface changes during maturation, specific antigenic differences between inactive and active, mature cells, studies by unlabeled antibody enzyme method; possible relationship to sperm's ability to recognize and/or penetrate oolemma of oocyte

Ascaris

- Bruch, K.; and Haas, J., 1976, Ann. Trop. Med. and Parasitol., v. 70 (2), 205-211
 Ascaris, hookworm, Trichuris, human, single dose fenbendazole compared with pyrantel: Liberia

Ascaris

- Bulucea, D., 1971, Pediatria, Bucuresti, v. 20 (2), 163-168
 mixed ascariasis and lambliaisis in young child with persistent eosinophilia which was not relieved by cure of parasitemia, clinical case report: Romania

Ascaris

- Cabrera, B. D.; Arambulo, P. V. III; and Portillo, G. P., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 510-518
 human ascariasis, statistical epidemiologic survey and development of working model for control and/or eradication of infection using a rural community in the Philippines

Ascaris

- Carswell, F.; Meakins, R. H.; and Harland, P. S. E. G., 1976, Lancet, London (7988), v. 2, 706-707
 Ascaris, no evidence that infection in humans suppresses development of asthma, clinical survey in endemic area: Tanzania

Ascaris

- Dafalla, A. A., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 146-147
 Toxocara canis, T. cati in humans, immunodiagnosis using the capillary-tube precipitin test, cross reaction with Ascaris could be eliminated by absorption with Ascaris antigen

Ascaris

- Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
 fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology

Ascaris

- Enigk, K.; Dey-Hazra, A.; and Batke, J., 1976, Tierarztl. Umschau, v. 31 (8), 360-362
 swine nematodes, mebendazole treatment

Ascaris

- Gougerot, M. A.; et al., 1975, Bull. Soc. Path. Exot., v. 68 (3), 297-303
 elevated IgE, useful indicator of possible human parasitism in absence of allergic type conditions

Ascaris

- Gupta, M. C.; et al., 1977, Lancet, London (8029), v. 2, 108-110
 ascariasis-infested preschool children, effect of periodic deworming with tetramisole on nutritional status and growth in children receiving government-supplied food supplements: Uttar Pradesh, India

Ascaris L., 1758

- Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
 Ascaridinae
 key; synonymy

Ascaris

- Ito, K.; et al., 1976, *Japan. J. Exper. Med.*, v. 46 (1), 79-85
levels of immunoglobulin E antibodies to *Ascaris* using *Ascaris suum* antigens higher in Papua New Guineans than in Japanese subjects

Ascaris

- Kagei, N., 1975, *Bull. Inst. Pub. Health, Tokyo*, v. 24 (3), 169-175
comparison of Kato thick smear and Tween 80 citric acid ether sedimentation methods for diagnosis of helminth ova

Ascaris

- Kelly, J. F.; and Patterson, R., 1973, *Internat. Arch. Allergy and Applied Immunol.*, v. 44 (4), 553-562
rhesus monkeys, characteristics of reverse passive respiratory reaction including relationship to *Ascaris* sensitivity

Ascaris

- Layrisse, M.; and Vargas, A., 1975, *Progr. Food and Nutrition Sc.*, v. 1 (10), 645-667
human intestinal parasites, mechanisms by which parasites interfere with host nutrition (competition for nutrients, malabsorption, blood loss, excess nutrient utilization), extensive review

Ascaris

- McFarlane, H., 1976, *Proc. Nutrition Soc.*, v. 35 (3), 263-272
parasitic infestation, preschool children, malnutrition and impaired immune response, brief review comment: Nigeria

Ascaris

- Monteoliva Hernandez, M., 1973, *Rev. Iber. Parasitol.*, v. 33 (4), 599-606
Ascaris, motility studies in vitro, motility longer with glucose than when fasting; longer at 25° C than at 37° C; younger worms more active than mature ones

Ascaris

- Neppert, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis

Ascaris

- Ong, H. C., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (3), 407-412
clinical observations during pregnancy in Malaysian aborigines living in deep jungle vs. those who have migrated "outside", includes observations on *Trichomonas vaginalis* and helminthic infection: Gombak, Selangor, Malaysia

Ascaris

- Patterson, R.; and Harris, K. E., 1975, *Internat. Arch. Allergy and Applied Immunol.*, v. 49 (3), 381-390
Ascaris-induced model of asthma in rhesus monkeys and dogs, arterial and muscle oxygen tension as additional parameter for assessment of model

Ascaris

- Patterson, R.; and Kelly, J. F., 1973, *Internat. Arch. Allergy and Applied Immunol.*, v. 45 (1-2), 98-109
rhesus monkeys, dogs, cellular and physiologic studies of immediate-type respiratory reactions including those to *Ascaris* antigen, review

Ascaris

- Patterson, R.; Suszko, I. M.; and Harris, K. E., 1975, *Clin. and Exper. Immunol.*, v. 19 (2), 335-342
dogs or rhesus monkeys with immediate-type allergy to *Ascaris*, potentiation of IgE-mediated cutaneous reactivity and blood leucocyte histamine release by deuterium oxide

Ascaris

- Pitchford, R. J.; and Visser, P. S., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 16 [Demonstration]
quantitative technique for the estimation of helminth eggs in urine and faeces

Ascaris

- Raynaud, J.-P.; Sennelier, J.; and Irisarri, E., 1975, *Folia Vet. Latina*, v. 5 (3), 412-429
gastrointestinal helminths, swine, post natal infection of piglets in contact with infected mothers, comparison of various methods of husbandry and hygiene, studies during pregnancy and lactation, routine daily hygiene recommended

Ascaris

- Samuel, M. R., 1975, *Progr. Drug Research*, v. 19, 96-107
human intestinal helminths, review of clinical experiences world wide comparing the efficacy and tolerance of bitoscanate with that of bephenium hydroxynaphthoate and tetrachlorethylene; found to be most useful against hookworm with results against other helminths still inconclusive

Ascaris

- Sanyal, R. K.; et al., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 50 (3), 291-296
adrenaline-induced increase in eosinophil count in patients with tropical pulmonary eosinophilia and in *Ascaris* larvae-fed guinea pigs, possible use as diagnostic test of tropical pulmonary eosinophilia

Ascaris

- Schroetter, E.; Tschaeppe, M.; and Meister, A., 1977, *Pharmazie*, v. 32 (3), 171-174
human ascariasis, synthesis of phenol derivatives for possible ovicidal effects

Ascaris

- Sfer, H.; Bollea, A.; and Leon, S., 1972, *Prensa Med. Argent.*, v. 59 (3), 116-118
human ascariasis of biliary tract with resulting obstruction, differential diagnosis, surgical management: Argentina

- Ascaris
Suberman, R. I., 1976, *Pediat. Ann.*, v. 5 (1), 32-48
Ascaris in children, dead worms in cecum after therapy as cause of intestinal obstruction and constipation, diagnosis by X-ray examination
- Ascaris
Tantengco, V. O.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 524-533
Ascaris, Trichuris, hookworm infections apparently not contributory cause of nutritional anemia in schoolchildren: Philippine Islands
- Ascaris
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Ascaris
Ugrinovic, N.; et al., 1972, *Med. Casop.*, v. 7 (1-2), 50-55
intestinal complications in children infected with Ascaris or Oxyuris, review of hospital cases: Yugoslavia
- Ascaris
Wadstroem, T.; et al., 1976, *Arch. Dis. Childhood*, v. 51 (11), 865-870
Ascaris, Entamoeba histolytica, Giardia lamblia, children with diarrheal disease, survey, no correlation with toxinogenic bacteria in stools: Ethiopia
- Ascaris sp.
Cabrera, B. D., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 50-55
Rattus rattus (feces): Leyte, Philippines
- Ascaris [sp.]
Chintanawongse, C.; et al., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 581 [Demonstration]
Ascaris fertilized eggs recovered in exudate of thoracoparacentesis performed on 4-year old girl with undiagnosed fever, upper gastrointestinal X-ray showed Ascaris worm shadows, 10 worms passed after piperazine therapy
- Ascaris sp.
Davidson, W. R., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Ascaris [sp.]
Guildal, J. A.; and Clausen, B., 1973, *Norwegian J. Zool.*, v. 21 (4), 329-330 [Abstract]
Vulpes vulpes (feces): Denmark
- Ascaris spp.
Hogarth-Scott, R. S.; and Feery, B. J., 1976, *Austral. J. Exper. Biol. and Med. Sc.*, v. 54 (4), 317-327
existence of cross-reacting antigens between Toxocara canis and Ascaris spp. and probably between T. canis and other nematodes confirmed by in vitro and in vivo tests, such cross-reactions compromise usefulness of skin tests in diagnosis
- Ascaris [sp.]
Rousset, J. J.; et al., 1976, *Nouv. Presse Med.*, v. 5 (28), 1760 [Letter]
human ascariasis in man diagnosed by x-ray discovery of calcified Ascaris in peritoneum: France
- Ascaris sp.
Valenzuela, G.; et al., 1976, *Bol. Chileno Parasitol.*, v. 31 (3-4), 87
Ascaris suum discovered at necropsy of sheep, fecal examination positive for Ascaris sp. eggs, confirmation of possibility of cross infection of gastrointestinal nematodes between sheep and swine: Pelchiquin, Provincia de Valdivia, Chile
- Ascaris sp.
Yang, J.; and Scholten, T., 1977, *Am. J. Clin. Path.*, v. 67 (3), 300-304
diagnosis of human intestinal parasites, fecal examination technique using Junod's fixative for concentration and permanent staining procedures, comparison with results using formalin-ether procedure
- Ascaris columnaris
Barnstable, R. W.; and Dyer, W. G., 1974, *Tr. Illinois State Acad. Sc.*, v. 67 (4), 451-460
Procyon lotor (small intestine): southern Illinois
- Ascaris columnaris, illus.
Church, E. M.; Wyand, D. S.; and Lein, D. H., 1975, *Am. J. Vet. Research*, v. 36 (3), 331-335
cerebrospinal nematodiasis, experimentally induced in Oryctolagus cuniculus with Ascaris columnaris, A. suum, or Toxocara canis, naturally occurring in Sylvilagus floridanus and O. cuniculus, clinical signs, gross and microscopic changes, duration of infection and parasite morphology and distribution in CNS, potential of rabbits as intermediate or paratenic hosts for ascarids of carnivorous origins
- Ascaris columnaris
Georgi, J. R.; et al., 1976, *Cornell Vet.*, v. 66 (3), 309-323
Procyon lotor: North Rose, Wayne County, New York
- Ascaris columnaris Leidy, 1856
Kozlov, D. P., 1969, *Trudy Gel'mint. Lab. Akad. Nauk SSSR*, v. 20, 71-78
Martes martes
Gulo gulo
all from Pechora river basin

- Ascaris columnaris*, *illus.*
Schueler, R. L., 1973, *J. Wildlife Dis.*, v. 9 (1), 58-60
larval ascarids, probably *Ascaris columnaris*, in *Sciurus granatensis* (brain, lung) with clinical signs of neurological disease
- Ascaris columnaris* Leidy, 1856
Young, P. L.; and Babero, B. B., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Ascaris dentata* Zeder
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 189-193
Barbus haibus (intestine): Vistula River near Warsaw
- Ascaris equorum*
Orr, J. P., 1972, *Vet. Rec.*, v. 90 (20), 571
Ascaris equorum in Thoroughbred foal (peritoneal cavity, stomach), probable cause of fatal perforated duodenal ulcer; accompanied by aneurysm at root of anterior mesenteric artery containing near-adult *Strongylus vulgaris*, case report
- Ascaris filiformis*, Goeze 1782
Rozman, M., 1971, *Acta Parasitol. Jugoslavica*, v. 2 (2), 67-77
as syn. of *Oswaldocruzia goezei* Skrjabin et Schulz, 1952
- Ascaris helicina* of Gedoelst, 1916 (in part)
Sprent, J. F. A., 1977, *J. Helminthol.*, v. 51 (3), 253-287
as syn. of *Dujardinascaris dujardini* (Travassos, 1920)
- Ascaris labiata* Rudolphi
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 189-193
Anguilla anguilla (intestine): Vistula River near Warsaw
- Ascaris lumbricoides*
Adickman, M.; and Tuthill, T. M., 1976, *Postgrad. Med.*, v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Ascaris lumbricoides*, *illus.*
Ahluwalia, H. S., 1965, *Med. J. Malaya*, v. 19 (4), 316-317
case report of acute peritonitis in man caused by granulomatous lesion of *Ascaris lumbricoides*, original diagnosis of tuberculosis had been made: Malaysia
- Ascaris lumbricoides*
Alcasid, M. L. S.; et al., 1973, *N. York State J. Med.*, v. 73 (13), 1786-1788
human intestinal parasites, no significant correlation with bronchial asthma, results of study with control and asthmatic groups: New York
- Ascaris lumbricoides*
Antunes, M. L.; and Delascio, D., 1967, *Maternidade e Infancia*, v. 26 (1), 63-67
laboratory diagnosis of vulvo-vaginitis of parasitic origin in children: Brasil
- Ascaris lumbricoides*
Arfaa, F.; and Ghadirian, E., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (5, part 1), 866-871
Ascaris lumbricoides, statistics of epidemiologic survey and mass therapy using pyrantel pamoate in 6 rural villages in central Iran
- Ascaris lumbricoides*
Arnaud, J. P.; and Danis, M., 1976, *Medecine Infant.*, v. 83 (1), 9-22
helminthiasis of digestive tract in children, differential diagnosis, current treatment methods
- Ascaris lumbricoides*
Asmera, J.; et al., 1975, *Ceskoslov. Pediat.*, v. 30 (4), 171-172
human intestinal helminths, mebendazole used as successful wide-spectrum anthelmintic in clinical trials: Czechoslovakia
- Ascaris lumbricoides*, *illus.*
Ausin, J. M.; Baquedano, F.; and Medina, A., 1966, *Bol. Chileno Parasitol.*, v. 21 (3), 88-91
Ascaris lumbricoides in 4-year-old child, expulsion of 2 living adult worms through the umbilicus, X-ray examination showed fistula into small bowel, case report: San Fernando, Chile
- Ascaris lumbricoides*
Baldini, I.; Pala, V.; and Ferro, M., 1974, *Pathologica* (959-960), v. 66, 339-349
Ascaris lumbricoides in humans, complement fixation, passive agglutination and latex agglutination compared with direct fecal examination as means of diagnosis
- Ascaris lumbricoides*
Baldo, B. A.; and Fletcher, T. C., 1973, *Nature* (5429), v. 246, 145-146
evidence of C-reactive protein-like precipitins in *Pleuronectes platessa* serum, reaction with *Ascaris lumbricoides* extract, possibility that these non-antibody precipitins form part of the fishes' humoral defenses against invasion by parasites
- Ascaris lumbricoides*
Ballo, T.; et al., 1975, *Med. Univ.*, v. 8 (3), 135-136
Ascaris lumbricoides, recommendations for diagnosis and treatment of children with levamisole: Hungary

- Ascaris lumbricoides*
Barrett, J., 1975, *J. Parasitol.*, v. 61 (3), 545-546
nucleosidediphosphate kinase, occurrence and intracellular distribution in 6 parasitic helminths
- Ascaris lumbricoides*
Barrett, J., 1976, *Parasitology*, v. 73 (1), 109-121
Ascaris lumbricoides eggs, studies on mechanism of induction of permeability, "no firm conclusions can be drawn"
- Ascaris lumbricoides*
Barrett, J., 1977, *Symposia Brit. Soc. Parasitol.*, v. 15, pp. 121-144
helminths, metabolic changes associated with infection, metabolism of adults vs. free-living stages compared, symposium presentation
- Ascaris lumbricoides*
Bell, W. J.; and Nassif, S., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 584-588
clinical trials comparing the efficacy of pyrantel pamoate and piperazine phosphate in the treatment of human *Ascaris lumbricoides* infections: Egypt
- Ascaris lumbricoides, illus.*
Bernstein, R. B., 1977, *Am. J. Digest. Dis.*, v. 22 (4), 391-394
Ascaris lumbricoides, biliary ascariasis in young girl with resulting chronic dilatation of the biliary system, diagnosis using intravenous cholangiography and ERCP radiography, clinical case report with emphasis on need for diagnostic awareness in persons originating from endemic areas: Oakland, California (native of Philippines)
- Ascaris lumbricoides, illus.*
Bevanger, L., 1974, *Tidsskr. Norske Laegefor.*, v. 94 (10), 651-652
single and mixed intestinal parasitic infections in adoptive children from Asiatic areas, need for control measures: Norway
- Ascaris lumbricoides*
Biagi, F.; Smyth, J.; and Gonzalez, C., 1975, *Prensa Med. Mexicana*, v. 40 (5-6), 189-192
human intestinal parasites, clinical trials with mebendazole show it to be useful drug against many parasites and therefore recommended for mass therapy in low socioeconomic areas where multiple parasitism is likely to be present: Mexico
- Ascaris lumbricoides*
Blumenthal, D. S.; and Schultz, M. G., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (5), 682-690
Ascaris lumbricoides, nutritional status of infected children compared with matched controls using anthropometric, clinical and laboratory criteria, adverse effects on serum albumin and vitamin C levels: Louisiana
- Ascaris lumbricoides*
Botero R., D., 1975, *Progr. Drug Research*, v. 19, 28-43
human intestinal soil transmitted helminths, epidemiology and public health importance, pathology, extensive review
- Ascaris lumbricoides*
Botero, D.; Orozco, H. I.; and Perez, A., 1966, *Bol. Chileno Parasitol.*, v. 21 (4), 111-114
Ascaris lumbricoides and *Enterobius vermicularis* in children, evaluation of combination syrup containing piperazine and pyriminium in treatment of infections
- Ascaris lumbricoides*
Brus, R.; et al., 1976, *Polski Tygod. Lekar.*, v. 31 (5), 195-197
Enterobius vermicularis in children, comparative mass clinical trials with vanquin and combantrin; combantrin indicated as drug of choice since it was slightly more effective than vanquin and also effective against *Ascaris lumbricoides*: Cieszyn County, Poland
- Ascaris lumbricoides*
Bryant, C., 1975, *Advances Parasitol.*, v. 13, 35-69
carbon dioxide utilisation and regulation of respiratory metabolic pathways in parasitic helminths, extensive review
- Ascaris lumbricoides*
Bull, F.; Oyarce, R.; and Stehr, I., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 10-15
prevalence and epidemiologic survey of human intestinal parasites in slum areas of Concepcion Province, Chile
- Ascaris lumbricoides, illus.*
Bustamante-Sarabia, J.; Martuscelli-Q., A.; and Tay, J., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (3), 568-569
Ascaris lumbricoides in young woman, massive intestinal infection with migration of parasites from colon through fistula to kidney and through subcutaneous abscess to exterior skin surfaces, clinical case report of fatal illness: State of Guerrero, Mexico
- Ascaris lumbricoides*
Cain, G. D., 1976, *Exper. Parasitol.*, v. 40 (1), 112-115
Ascaris lumbricoides gut, muscles, and eggs, coproporphyrinogen oxidase activity, results suggest that porphyrin synthesis may be maintained at low levels due to continual feedback inhibition by protoheme
- Ascaris lumbricoides*
Cain, G. D.; and Bassow, F., 1976, *Internat. J. Parasitol.*, v. 6 (1), 79-82
Ascaris lumbricoides, porphyrins in perienteric fluid

- Ascaris lumbricoides*
Camargo, P. A. M.; et al., 1974, *Pediat. Prat.*, S. Paulo, v. 45 (10-12), 189-194
human intestinal helminths, comparative treatment trials using mebendazole and pyrantel pamoate, efficacy of drugs checked using the McMaster technique to determine reduction in egg production
- Ascaris lumbricoides, illus.*
Cerva, L., 1976, *Immun. u. Infekt.*, v. 4 (6), 279-282
intestinal helminths, diagnostic method for staining of eggs and larvae in smears of fresh and fixed stool samples
- Ascaris lumbricoides*
Cervoni, W. A.; and Oliver-Gonzalez, J., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 589-591
pyrantel pamoate effective against human infection of *Ascaris lumbricoides* and *Necator americanus* but not *Trichuris trichiura* in clinical evaluations: Puerto Rico
- Ascaris lumbricoides*
Chen, P. C. Y., 1970, *Med. J. Malaya*, v. 24 (3), 176-182
survey of rural Malay community for beliefs and practices regarding *Ascaris lumbricoides* infections, poor sanitation and tribal practices aid in spread of infections: Malaysia
- Ascaris lumbricoides*
Daili, S.; et al., 1972, *Paediat. Indonesiana*, v. 12 (6), 271-275
Ascaris lumbricoides in children, ascaridil as single dose therapy, satisfactory results: West Sumatra
- Ascaris lumbricoides, illus.*
Denzler, T. B.; and Gunning, J.-J., 1976, *Arch. Int. Med.*, Chicago, v. 136 (9), 1044-1045
recently immigrated Vietnamese refugee with acute abdominal symptoms resulting from intestinal infection with *Ascaris lumbricoides*, radiologic diagnosis and piperazine treatment, case report: California
- Ascaris lumbricoides*
Doshi, J. C.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 636-639
human hookworm infections, clinical trials of dosage searching for tolerance and efficacy using C.9333-Go/CGP 4540, compound well tolerated with few side effects, active also in mixed infections with *Ascaris lumbricoides*: Bombay, India
- Ascaris lumbricoides, illus.*
Dutt, A. K.; Beasley, D.; and Sandosham, A. A., 1969, *Med. J. Malaya*, v. 24 (2), 158-160
case report of woman with eosinophilic granuloma of pancreas caused by *Ascaris lumbricoides* eggs, symptoms suggestive of cancer: Malaysia
- Ascaris lumbricoides*
Ehrlich, I.; and Hrszenjak, T., 1975, *Vet. Arhiv, Zagreb*, v. 45 (5-6), 129-132
Ascaris lumbricoides, thin layer chromatography of polar lipids from tissue homogenates, identification; globosides as possible antigen components
- Ascaris lumbricoides*
Evans, H. J.; Sullivan, C. E.; and Piez, K. A., 1976, *Biochemistry*, Washington, v. 15 (7), 1435-1439; Correction (11), 2500
Ascaris lumbricoides, cuticle collagen, resolution into three chain types by chromatography on phosphocellulose
- Ascaris lumbricoides*
Farahmandian, I.; et al., 1977, *Chemotherapy*, v. 23 (2), 98-105
human intestinal *Ascaris*, hookworm and *Trichostrongylus* spp., statistics of comparative trials using pyrantel, levamisole, mebendazole, thiabendazole and bephenium: Iran
- Ascaris lumbricoides*
Farid, Z.; et al., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (3), 425-426 [Letter]
Ancylostoma duodenale, *Ascaris lumbricoides*, male farmers, tetramisole extremely effective against both parasites, no side-effects: Egypt
- Ascaris lumbricoides*
Farid, Z.; et al., 1977, *J. Trop. Med. and Hyg.*, v. 80 (5), 107-108
human *Ascaris lumbricoides* and *Ancylostoma duodenale*, effectiveness of levamisole, pyrantel and bephenium compared in clinical trials, all drugs well tolerated: Egypt
- Ascaris lumbricoides*
Farzan, S., 1970, *N. England J. Med.*, v. 282 (22), 1273 [Letter]
Ascaris lumbricoides in asthmatic persons, generally a concomitant occurrence rather than cause of asthma
- Ascaris lumbricoides*
de Figueiredo, N.; et al., 1968, *Bol. Chileno Parasitol.*, v. 23 (1-2), 57-59
intestinal histopathology as ascertained by oral biopsies, common human intestinal parasites
- Ascaris lumbricoides*
Fischerova, H.; and Kubistova, J., 1968, *Physiol. Bohemoslov.*, v. 17 (4), 375-382
absence of phosphagen in *Ascaris lumbricoides* muscles and *Fasciola hepatica* tissues supports theory that phosphagen acts only as reserve of readily mobilized high energy phosphate bond in muscle cell
- Ascaris lumbricoides*
Forman, D. W.; et al., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 598-601
parasitic pathogens discovered during etiologic survey of causes of diarrheal disease in U.S. Marines in Vietnam
- Ascaris lumbricoides*
Fujimoto, D., 1975, *J. Biochem.*, Tokyo, v. 78 (5), 905-909
Ascaris lumbricoides, extent of hydrolysis of *Ascaris* cuticle collagen by bacterial collagenase under various conditions, amino acid composition of collagenase digests, results suggest CaCl_2 necessary for hydrolysis of certain regions in molecule of *Ascaris* collagen not present in mammalian collagens

- Ascaris lumbricoides*
Garrocho-Sandoval, C.; and Torres-Ruvalcaba, A., 1977, *Am. J. Clin. Path.*, v. 67 (6), 603-605
human intestinal parasites, comparison study of two different methods for collecting fecal samples for diagnostic purposes
- Ascaris lumbricoides*
Gemési, G., 1971, *Parasitol. Hungar.*, v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Ascaris lumbricoides*
Gietko, M.; and Zapart, W., 1975, *Pediat. Polska*, v. 50 (1), 61-68
visceral larva migrans in young children with severe eosinophilia, diagnosed by intradermal skin test antigens of *Ascaris lumbricoides* and *Toxocara canis*
- Ascaris lumbricoides*
Gold, B. M.; and Meyers, M. A., 1977, *Am. J. Roentgenol.*, v. 128 (3), 493-494
Taenia saginata, case report of tapeworm in butcher who routinely ate raw beef, radiologic manifestations and differentiation from *Ascaris lumbricoides* as seen on X-ray, atabrine: New York City
- Ascaris lumbricoides*
Gortazar Hajar, P.; and Cigala Cano, J. L., 1970, *Minerva Pediat.*, v. 22 (26), 1333-1335
Ascaris lumbricoides infection in child resulting in cholecystitis and hepatic abscess, case report, clinical management: Guadalajara
- Ascaris lumbricoides*
de Groot, J. F. M.; et al., 1976, *Nederl. Tijdschr. Geneesk.*, v. 120 (51), 2263-2267
Trichuris trichiura, *Ascaris lumbricoides* and *Entamoeba histolytica*, fecal survey of nursing staff and inmates of hospital for mentally retarded for presence of intestinal parasites: Groot Schuylenburg, Apeldoorn, Netherlands
- Ascaris lumbricoides*
Gut, C.; Zulauf, E.; and Tobler, H., 1977, *Experientia*, v. 33 (6), 819 [Abstract]
Ascaris lumbricoides, isolation of messenger RNA coding for egg shell protein
- Ascaris lumbricoides*
Hays, B. D., 1977, *J. Environ. Health*, v. 39 (6), 424-426
transmission of protozoan cysts and metazoan eggs from land application of sewage effluents and sludges, brief literature review, parasite survey from selected Pittsburgh area sludges, control measures
- Ascaris lumbricoides*
Hermoso Yanez, R.; and Monteoliva Hernandez, M., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 427-437
Ascaris lumbricoides, perivisceral fluid, glucose and trehalose content apparently independent of maturity of parasite or nutritive state of swine host
- Ascaris lumbricoides*
Hermoso, R.; and Monteoliva, M., 1974, *Rev. Iber. Parasitol.*, v. 34 (3-4), 295-304
Ascaris lumbricoides, content of trehalose, glucose, soluble glycogen and combined glycogen in sexual organs, body walls and perivisceral fluid, values before and after culture in solution without glucose, estimate of endogenous utilization
- Ascaris lumbricoides*
Hermoso, R.; Monteoliva, M.; and Sanchez, M., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 329-337
Ascaris lumbricoides, culture in media with added glucose or hemoglobin, relationship of nutrients to total carbohydrate content of certain tissues; no clear relationship to content of specific sugars (glucose, trehalose, glycogen)
- Ascaris lumbricoides*
Hermoso, R.; Sanchez, M.; and Monteoliva, M., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 355-360
Ascaris lumbricoides, culture in media with added glucose or hemoglobin, soluble protein and hemoglobin content of certain tissues, values for body wall most accurately reflect physiological state of parasite; Tyrode solution with 1.0 g glucose and 0.3 g pig hemoglobin per liter best medium for culture
- Ascaris lumbricoides*
Herrero Alonso, C., 1969, *Med. Trop., Madrid*, v. 45 (1-2), 11-20
thiabendazole drug of choice in treating human infections of *Ancylostoma duodenale*, *Trichuris trichiura* and *Ascaris lumbricoides*, clinical trials
- Ascaris lumbricoides*
Hiatt, W. R., 1969, *Oral Surg.*, v. 28 (4), 610-612
Ascaris lumbricoides, adult female, removed from submandibular mass on neck of small child, probable migration through thyroglossal duct via vomitus, clinical case report: Guam, Mariana Islands
- Ascaris lumbricoides*
Hoghooghi, N.; et al., 1973, *Pahlavi Med. J.*, v. 4 (1), 53-60
Ascaris lumbricoides, survey of soil contamination with *Ascaris* eggs, seasonal variations, possible control with improved sewage management: Isfahan, Iran
- Ascaris lumbricoides*
Hrzenjak, T.; and Ehrlich, I., 1976, *Vet. Arhiv, Zagreb*, v. 46 (9-10), 263-267
helminths, separation of polar lipids, comparative biochemistry
- Ascaris lumbricoides*
Hsieh, H. C.; and Chen, E. R., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (3), 362-364
Ascaris lumbricoides, humans, single dose therapy with combantrin resulted in high cure rates and reduced egg counts: Taiwan

- Ascaris lumbricoides*
Ishikawa, H., 1977, *Comp. Biochem. and Physiol.*, v. 56 (3B), 229-234
several protostomes (including *Ascaris lumbricoides* and *Fasciola hepatica*), ribosomal RNA's, thermal stability and molecular integrity, evolutionary implications
- Ascaris lumbricoides*
Ishizaki, T., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 65-81
human *Ascaris lumbricoides*, epidemiologic and historical review, nightsoil control measures, comparison of anthelmintics in current use: Japan
- Ascaris lumbricoides*
Islam, N.; and Chowdhury, N. A., 1976, *South-east Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 81-84
human intestinal parasites, comparative treatment trials using mebendazole and pyrantel pamoate, recommendations for use: Dacca, Bangladesh
- Ascaris lumbricoides*
Iwanczuk, I.; 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 139-145
human parasite incidence in water and surfaces of swimming pools; change of incidence in children using swimming pool for 6 week period: Poland
- Ascaris lumbricoides*
Iwanowski, H.; and Herman, T., 1977, *Pediat. Polska*, v. 52 (6), 649-654
Ascaris lumbricoides in children, Soprunova urine test used successfully to diagnose ascariasis and differentiate from infections of *Trichuris trichiura*, *Enterobius vermicularis* and *Lamblia intestinalis*
- Ascaris lumbricoides*
Jacobson, R. J., 1972, *N. England J. Med.*, v. 286 (25), 1369 [Letter]
Ascaris lumbricoides pneumonia and asthma occurring in "beer-drinking" Bantu natives, report of 2 outbreaks after consumption of home-brewed apparently *Ascaris* contaminated beer, resulting hypersensitivity reaction: South Africa
- Ascaris lumbricoides*
James, T., 1969, *Med. Proc., Johannesburg*, v. 15 (17), 304-305
factors connecting endemic ascariasis with endemic poor nutrition, world wide problem in child population
- Ascaris lumbricoides*
James, T., 1970, *Med. Proc., Johannesburg*, v. 16 (8), 127-131
Ascaris lumbricoides abscess of liver in young children, clinical symptoms and pathology
- Ascaris lumbricoides, illus.*
Jarman, M., 1976, *Organ. Nematodes (Croll)*, 293-312
Ascaris lumbricoides, neuromuscular physiology, review: morphology; working of nerve-muscle preparation; ionic fluxes across muscle cell membranes
- Ascaris lumbricoides*
Jones, H. I., 1976, *Papua N. Guinea Med. J.*, v. 19 (3), 153-155
pigs as important factor in spread of *Ascaris lumbricoides* by eating human feces containing *Ascaris* eggs which then pass unharmed through pig digestive system and are spread widely in human environment; human hookworm eggs largely destroyed in pig digestive system thereby effectively reducing human exposure to hookworm larvae: Papua New Guinea
- Ascaris lumbricoides*
Jones, H. I., 1976, *Papua N. Guinea Med. J.*, v. 19 (3), 165-172
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, epidemiologic survey of human helminths in natives of Kar Kar Island, *Ascaris* infections highest and heaviest in children and principal cause of high eosinophilia in the communities: Madang Province, Papua New Guinea
- Ascaris lumbricoides*
Jones, H. I., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (4), 523-529
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, statistical epidemiologic survey of human intestinal parasites with discussion of socio-cultural and environmental factors influencing spread of infections: Venilale District of East Timor
- Ascaris lumbricoides*
Jones, H. I., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (2), 219-226
use of haemagglutination test as aid in study of epidemiology of human *Ascaris lumbricoides* infections, A. suum used as antigen: Papua New Guinea and East Timor
- Ascaris lumbricoides*
Jose, D. G.; and Welch, J. S., 1970, *Med. J. Australia*, v. 1 (8), 349-356
possible role of intestinal parasitism in growth-retarded, anemic and malnourished Australian Aboriginal children, comparison with normal Aboriginal children: Queensland
- Ascaris lumbricoides*
Khan, S. M.; and Khairul Anuar, A., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 260-264
Ascaris lumbricoides, *Trichuris trichiura*, hookworms, prevalence survey of soil-transmitted intestinal helminths in patients admitted to the District Hospital in Balik Pulau, Penang, Malaysia
- Ascaris lumbricoides*
Kieninger, G.; and Madecki, O., 1972, *Med. Welt*, v. 23 (2), 59-61
Ascaris lumbricoides, human, intestinal obstruction and perforation, surgical aspects of treatment

- Ascaris lumbricoides*
Krasnonos, L. N.; Brudastov, A. N.; and Tadzhikbaev, A. T., 1976, *Gig. Sanitariia* (10), 114-115
Ascaris lumbricoides, extent of contamination of soil with eggs in relation to numbers of infected people in microfoci of ascariasis, hygienic factors: Namangansk oblast, Uzbek SSR
- Ascaris lumbricoides*
Krubwa, F.; et al., 1974, *Medecine Trop.*, v. 34 (5), 679-687
trial administration of mebendazole to school children on quarterly basis for attempted eradication of intestinal nematodes, recommendations for use as mass control measure: Zaire
- Ascaris lumbricoides*, *illus.*
Kuster, G.; Oyarce, R.; and Boero, D., 1966, *Bol. Chileno Parasitol.*, v. 21 (2), 48-50
Ascaris lumbricoides in human, case report of hepatic granuloma surrounding parasite eggs, discovered at surgery to correct biliary colic, piperazine therapy resulted in elimination of 3 adult worms: Chile
- Ascaris lumbricoides*
Lamina, J., 1976, *Deutsche Tierarztl. Wchnschr.*, v. 83 (4), 133-136
Ascaris lumbricoides, guinea-pigs (exper.), detection of humoral antibodies by microprecipitation test
- Ascaris lumbricoides*
Lang, F., 1973, *Nepegeszsegugy*, v. 54 (5), 313-315
Ascaris lumbricoides, successful elimination of infection through epidemiologic measures, health education and treatment of infected persons: Hungary
- Ascaris lumbricoides*
Lang, F.; and Woelfer, V., 1976, *Med. Universalis*, v. 9 (5), 255-256
Ascaris lumbricoides, levamisole in humans, good results in field trials: Gyor-Sopron County, Hungary
- Ascaris lumbricoides*
LaPorte, V. D.; and Gibbs, R. S., 1977, *Obst. and Gynec.*, v. 49 (1), suppl., 84-85
Ascaris lumbricoides, acute pancreatitis associated with ascariasis in 2 pregnant Vietnamese evacuees, clinical management: Virginia
- Ascaris lumbricoides*
Lengyel, A.; et al., 1973, *Orvosi Hetilap*, v. 114 (23), 1368-1371
clinical trials testing levamisole for therapy of human intestinal parasites, good results against *Ascaris lumbricoides*
- Ascaris lumbricoides*
Lie, K. J.; Kwo, E. H.; and Ow-Yang, C. K., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (2), 196-200
Ascaris lumbricoides, *Trichuris trichiura*, *Necator americanus*, fecal survey for evidence of soil-transmitted helminths in infants and children living near Kuala Lumpur, Malaysia
- Ascaris lumbricoides*
Loc-Chay, M., 1975, *Provence Med.*, v. 43 (6-7), 11-16
Ascaris lumbricoides, humans, clinical trials comparing the efficacy of piperazine and san-tonine
- Ascaris lumbricoides*, *illus.*
Mathur, S. C.; et al., 1975, *Med. and Surg.*, Bombay, v. 15 (7), 5-7
report of 3 cases of small bowel perforation resulting from infections with *Ascaris lumbricoides*: Jaipur, Rajasthan, India
- Ascaris lumbricoides* Linne, 1758, *illus.*
Maung, M., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (1), 41-45
Ascaris lumbricoides from man, *Ascaris suum* from pigs, morphological comparisons of denticles, cuticular striations, eggs and spicules
- Ascaris lumbricoides*
Monteoliva Hernandez, M., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 407-425
Ascaris lumbricoides, perivisceral fluid, separation and analysis of proteins and hemoglobin
- Ascaris lumbricoides*
Monteoliva, M.; Benito, M.; and Hermoso, R., 1973, *Rev. Iber. Parasitol.*, v. 33 (4), 515-524
Ascaris lumbricoides, perivisceral liquid, statistical analysis of content of proteins, hemoglobin and "adenine derivatives"; protein and hemoglobin content influenced by intestinal content and diet of host but not by maturity of parasite; types of proteins dependent on weight and maturity of parasite
- Ascaris lumbricoides*
Monteoliva, M.; and Hermoso, R., 1974, *Rev. Iber. Parasitol.*, v. 34 (3-4), 229-236
Ascaris lumbricoides, cultured in starvation conditions, soluble protein content decreased significantly in sexual organs and body wall, slightly in perivisceral fluid; hemoglobin decreased significantly in sexual organs and body wall, increased slightly in perivisceral fluid; estimates of endogenous utilization and elimination
- Ascaris lumbricoides*
Most, H., 1972, *N. England J. Med.*, v. 287 (10), 495-498; (14), 698-702
common parasitic infections of man encountered in the United States, recommendations for treatment, review
- Ascaris lumbricoides*
Mutalik, G. S.; Gulati, R. B.; and Igbal, A. K., 1975, *Progr. Drug Research*, v. 19, 81-85
human intestinal parasites, clinical trials with bitoscanate show it to be safe and useful anthelmintic especially against hook-worm infections: India
- Ascaris lumbricoides*
Nalin, D. R.; and McLaughlin, J., 1976, *J. Parasitol.*, v. 62 (5), 839-841
Ascaris lumbricoides adults found to be colonized by *Vibrio cholerae* when isolated from cholera patients: Bangladesh

- Ascaris lumbricoides*
Neppert, J.; and Warns, C.-M., 1974, Tropenmed. u. Parasitol., v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from *Ascaris*, hookworm, *Onchocerca*, *Dirofilaria immitis*, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination
- Ascaris lumbricoides*
Obtulowicz, K., 1972, Polski Tygod. Lekar., v. 27 (8), 296-298
human intestinal parasites, possible role in etiology of asthma and other allergic conditions
- Ascaris lumbricoides*
Okumura, M.; et al., 1975, Med. and Surg., Bombay, v. 15 (8), 8-13
acute intestinal obstruction resulting from *Ascaris lumbricoides*, review of 455 cases observed in young children, differential diagnosis, pathologic findings, clinical management: Sao Paulo
- Ascaris lumbricoides*
Orren, A.; and Dowdle, E. B., 1975, Internat. Arch. Allergy and Applied Immunol., v. 49 (6), 814-830
serum IgE concentrations and immediate skin hypersensitivity to common allergens analyzed with respect to ethnic group (whites, Cape Coloreds, Africans), sex, allergic status, and evidence of intestinal helminthic infestation: Western Cape Province, South Africa
- Ascaris lumbricoides*
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Ascaris lumbricoides*
Owsinski, J.; Hytros, M.; and Walaszek, K. M., 1977, Przegl. Lek., v. 34 (6), 563-565
ascariasis complicating a pleural-pericardial neoplastic cyst in young man, clinical case report: Krakow, Poland
- Ascaris lumbricoides*
Pasternak, J.; and Barrell, R., 1976, Exper. Cell Research, v. 102 (1), 191-199
Ascaris lumbricoides, quantitative changes in DNA, histone, and basic nuclear protein content during spermatogenesis and oogenesis, microspectrophotometry
- Ascaris lumbricoides*
Pasternak, J.; and Barrell, R., 1976, Genet. Research, v. 27 (2), 339-348
Ascaris lumbricoides, quantitation of nuclear DNA during gametogenesis and embryogenesis by Feulgen-microspectrophotometry, DNA constancy and chromatin diminution
- Ascaris lumbricoides*
Pasternak, J.; and Barrell, R., 1976, J. Exper. Zool., v. 198 (2), 217-223
Ascaris lumbricoides, patterns of nuclear DNA and protein changes during embryogenesis, microspectrophotometric analysis, comparison with free-living *Panagrellus silusiae*
- Ascaris lumbricoides*
Pawlowski, Z.; and Chodera, L., 1972, Polski Tygod. Lekar., v. 27 (41), 1605-1607
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, humans, levo-tetramisole clinical trials successful except in trichuriasis
- A[*scaris*] *lumbricoides*
Penna, R.; and Grassi, L., 1972, Parassitologia, v. 14 (2-3), 339-341
[*Trichuris trichiura*, A[*scaris*] *lumbricoides*, H[*ymenolepis*] *nana*, survey of prevalence of geohelminthiasis in school children, helminthiasis related to socio-economic and hygienic conditions, not to scholastic achievement, prevalence higher in plain country than in mountain country: Province of Alesandria (Italy)]
- Ascaris lumbricoides*
Podgorny, G., 1967, North Carolina Med. J., v. 28 (1), 21-23
Ascaris lumbricoides infection in 3-year-old child complicating recovery from accidental gunshot wounds penetrating intestines, clinical case report: Winston Salem, North Carolina
- Ascaris lumbricoides*
Purtilo, D. T.; et al., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 229-232
parasitized children with protein-calorie malnutrition, complete blood count, serum immunoglobulin concentration, significant relationship between intensity of parasitism and Ig levels: Brazil
- Ascaris lumbricoides*
Rajasekaran, P.; Dutt, P. R.; and Pisharoti, K. A., 1977, Indian J. Med. Research, v. 66 (2), 189-199
human intestinal parasites, survey of correlation between infection rate and source of water supply (well, street tap, home with tap water) as indication of control of water-borne diseases by public water supplies: Madurai district, Tamil Nadu, India
- Ascaris lumbricoides*
Ray, D. K.; Negi, S. K.; and Srivastava, P. S., 1976, Indian J. Animal Research, v. 10 (1), 53
Ratufa bicolor (feces): Calcutta Zoological Garden
- Ascaris lumbricoides*
Rees, P. H.; and Marsden, P. D., 1970, Brit. J. Clin. Pract., v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment

- A[scaris] lumbricoides
Ricci, M.; and Federico, A., 1972, Riv. Parasitol., Roma, v. 33 (3), 209-218
A[scaris] lumbricoides, human, control, mass treatment with piperazine adipate: Comune di Parenti, Cosenza
- Ascaris lumbricoides
Ripert, C.; and Avouac-Borzee, F., 1974, Medecine Afrique Noire, v. 21 (10), 713-715
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, human, epidemiology: Mirebalais, Haiti
- Ascaris lumbricoides
Roe, T. N. C.; Lal, K.; and Cawkell, W. A., 1968, Med. J. Malaya, v. 23 (1), 78-79
case reports of human Ascaris lumbricoides infections resulting in small bowel obstruction with volvulus: Malaya
- Ascaris lumbricoides
Rukmono, B.; Naumar, S. A.; and Talogo, R. W., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (2), 249-255
Ascaris lumbricoides, Entamoeba histolytica, Giardia lamblia implicated as possible pathogens in survey of infantile diarrhea in a population of low socio-economic group in Djakarta, Indonesia
- Ascaris lumbricoides
Sagua, H.; et al., 1973, Bol. Chileno Parasitol., v. 28 (3-4), 58-60
comparison of phenol-alcohol-formalin sedimentation and polyvinyl alcohol fixative tests in diagnosis of human intestinal helminths and protozoa
- Ascaris lumbricoides
Sapunar, J., 1973, Bol. Chileno Parasitol., v. 28 (1-2), 34-36
Ascaris lumbricoides causing chronic cholecystitis, 11 female worms found in woman's ductus choledochus at surgery: Chile
- Ascaris lumbricoides
Sasi, P. K.; and Kaleysaraj, R., 1975, Experimentia, v. 31 (11), 1261-1262
Ascaris lumbricoides, incubation in medium containing sub-lethal concentrations of piperazine, decreased phospholipid level, partial stimulation of phospholipase C activity, partial inhibition of choline kinase activity
- Ascaris lumbricoides
Schenone, H.; et al., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 118-122
single combined dose of pyrvinium and tetramisole effective in treating children with Enterobius vermicularis and mixed infections with Ascaris lumbricoides: Chile
- Ascaris lumbricoides
Schenone, H.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 86-89
pyrantel pamoate recommended for treatment of both Ascaris lumbricoides and Enterobius vermicularis, individual or mass therapy, clinical trials in children: Chile
- Ascaris lumbricoides
Schenone, H.; et al., 1976, Bol. Chileno Parasitol., v. 31 (3-4), 75-78
Enterobius vermicularis, Ascaris lumbricoides, and Trichuris trichiura in children, clinical trials of oral mebendazole highly successful, recommended for individual or mass therapy
- Ascaris lumbricoides
Schenone, H.; Galdames, M.; and Cabello, C., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 89-90
intestinal parasites, young girls, combined therapy with mebendazole and thiabendazole
- Ascaris lumbricoides, illus.
Schultz, M. G., 1977, Postgrad. Med., v. 62 (2), 121-125
- Ascaris lumbricoides
Seah, S. K. K., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 534-542
intestinal parasites, persons living in non-endemic areas who acquired infections while travelling or who have immigrated from endemic areas, pyrantel pamoate successful for Ascaris lumbricoides, results with other parasites varied: Montreal, Canada
- Ascaris lumbricoides
Seah, S. K. K., 1976, Canad. Med. Ass. J., v. 115 (8), 777-779
mebendazole suggested as drug of choice for Trichuris trichiura and mixed nematode infections after extensive clinical trials: Canada
- Ascaris lumbricoides
Signorello, G., 1973, Minerva Med., v. 64 (52), 2736-2740
Ascaris lumbricoides in children resulting in cutaneous allergic reactions, skin test diagnosis, frequent mixed infections with Trichuris trichiura also diagnosed
- Ascaris lumbricoides
Simon, K., 1972, Med. Welt, v. 23 (44), 1601-1602
Ascaris lumbricoides, therapy of bronchitis in children resulting from Ascaris infections: Germany
- Ascaris lumbricoides
Singson, C. N.; Banzon, T. C.; and Cross, J. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 932-934
Capillaria philippinensis, human clinical trials using mebendazole for intestinal capillariasis, additionally effective against Ascaris lumbricoides, Trichuris trichiura and hookworm: Philippines
- Ascaris lumbricoides
Sinios, A., 1972, Monatschr. Kinderh., v. 120 (9), 378-381
Ascaris lumbricoides causing interstitial eosinophilic pertussis-like pneumonia in newborn, diagnosis by direct and indirect precipitation reaction after emesis of adult Ascaris by mother

- Ascaris lumbricoides*
Smedresman, P., 1977, Clin. Pediat., Philadelphia, v. 16 (2), 197
Ascaris lumbricoides as cause of appendicitis in young girl, case report, medical management: New York City (lived previously in South Carolina)
- Ascaris lumbricoides*
Stankov, M.; Movsesijan, M.; and Kovandzic, D., 1975, Acta Parasitol. Iugoslavica, v. 6 (1), 31-36
Ascaris suum, pigs, antibodies form specific antigen-antibody complex; same antibodies are specifically bound to same organs and tissues of A. lumbricoides, sera of humans infected with A. lumbricoides form fluorescent complex with same organs of A. lumbricoides and A. suum
- Ascaris lumbricoides, illus.*
Stretton, A. O. W., 1976, J. Exper. Biol., London, v. 64 (3), 773-788
Ascaris lumbricoides, anatomy of muscle cells and their neuromuscular contacts, development of musculature from larval to adult forms, brief comparison with Oxyuris equi
- Ascaris lumbricoides*
Tantengco, V. O.; Marzan, A. M.; and de Castro, C. R., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 580-586
Ascaris lumbricoides, Trichuris trichiura, establishment of blood parameters in infected and control children, little significant differences except lowered albumin and elevated globulin levels in presence of infections: Philippine Islands
- Ascaris lumbricoides*
Terziiski, A., 1972, Izvest. Tsentral. Khel'mint. Lab., v. 15, 199-207
Ascaris suum, A. lumbricoides, Toxocara canis, migration in white mice, comparison
- Ascaris lumbricoides*
Torisu, M.; et al., 1975, Clin. Immunol. and Immunopathol., v. 4 (4), 467-477
Ascaris lumbricoides, patients with worm migration into biliary tree, skin tests, complement fixation, hemagglutination tests, immunoglobulin levels, pre- and post-surgical results, significant preoperative rise in IgE appears to be dependent on Ascaris infection, purified Ascaris antigen has high chemotactic effect on eosinophils
- Ascaris lumbricoides*
Torres, P.; and Barriga, O. O., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 79-85
Ascaris suum, A. lumbricoides, Toxocara mystax, Ascaridia galli, comparative antigenic analysis by gel double diffusion and immunoelectrophoresis
- Ascaris lumbricoides*
Turner, K. J.; Baldo, B. A.; and Anderson, H. R., 1975, Internat. Arch. Allergy and Applied Immunol., v. 48 (6), 784-799
humans, serum IgE levels, no significant correlation with faecal egg counts to hookworm, Ascaris lumbricoides, and Trichuris, incidence of IgE antibodies to Ascaris lumbricoides not correlated with incidence of asthma but significantly elevated in patients with chronic obstructive lung disease, hypersensitivity to Ascaris apparently not factor of importance in etiology of asthma in this area: Highland area of Papua-New Guinea
- Ascaris lumbricoides, illus.*
Ubelaker, J. E.; and Allison, V. F., 1975, J. Parasitol., v. 61 (5), 802-807
eggs, fine external morphology, scanning electron microscopy
- Ascaris lumbricoides*
Vega Franco, L.; et al., 1975, Prensa Med. Mexicana, v. 40 (7-8), 197-201
intestinal parasites, comparison of D-xylose intestinal absorption in infected children showed that only those with Giardia lamblia had statistically different absorption from non-infected children: Mexico
- Ascaris lumbricoides*
Vinayak, V. K.; and Sehgal, S. C., 1976, Indian J. Med. Research, v. 64 (9), 1347-1350
human helminthic and protozoan parasites, comparison of nigrosin-methylene blue diagnostic test with formol-ether method and direct examination
- Ascaris lumbricoides, illus.*
Wang, J. S.; and Fujita, J., 1976, J. Chinese Soc. Vet. Sc., v. 2 (1), 35-39
Ascaris suum, A. lumbricoides, scanning electron microscopy, especially head, lip, papillae, tail, cuticle, denticles, no morphological difference found
- Ascaris lumbricoides*
Warren, K. S.; and Mahmoud, A. A. F., 1977, J. Infect. Dis., v. 135 (5), 868-872
human ascariasis and toxocariasis, algorithms in diagnosis and clinical management
- Ascaris lumbricoides*
Weisblat, D. A.; Byerly, L.; and Russell, R. L., 1976, J. Comp. Physiol., A, v. 111 (2), 93-113
Ascaris lumbricoides, ionic mechanisms of electrical activity in somatic muscle, significance of results in understanding role of myogenic activity in nematode locomotion
- Ascaris lumbricoides*
Weisblat, D. A.; and Russell, R. L., 1976, J. Comp. Physiol., A, v. 107 (3), 293-307
Ascaris lumbricoides, propagation of electrical activity in nerve cord and muscle syncytium
- Ascaris lumbricoides, illus.*
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105

- Ascaris lumbricoides*, *illus.*
Zuidema, P. J., 1976, *Nederl. Tijdschr. Geneesk.*, v. 120 (43), 1849-1854
human intestinal helminths, differential diagnosis and clinical management of parasitic infestations seen in immigrants from Surinam: Netherlands
- Ascaris lumbricoides* var. *hominis*
Mukerji, K.; et al., 1976, *Indian J. Med. Research*, v. 64 (11), 1611-1619
Ascaris lumbricoides var. *hominis*, purification and protein properties of trypsin inhibitor located in muscular and cuticular layers of parasite, speculations on immunologic role
- Ascaris lumbricoides* var. *hominis*
Mukerji, K.; et al., 1977, *Indian J. Med. Research*, v. 66 (5), 745-755
Ascaris lumbricoides var. *hominis*, purification of parasite chymotrypsin inhibitor and properties of partially purified chymotrypsin and trypsin inhibitors
- Ascaris lumbricoides* var. *suis*
Aoki, T.; et al., 1975, *Proc. Japan Acad. Tokyo*, v. 51 (9), 733-736
Ascaris lumbricoides var. *suis* ovary, demonstration of glutamine-dependent carbamoyl-phosphate synthetase, catalytic and regulatory properties
- Ascaris lumbricoides* var. *suis*
Homandberg, G. A.; and Peanasky, R. J., 1976, *J. Biol. Chem.*, v. 251 (8), 2226-2233
Ascaris lumbricoides var. *suis*, carboxypeptidase inhibitors, purification and properties, evidence for atypical stoichiometry
- Ascaris lumbricoides* var. *suis*
Koehler, P.; and Saz, H. J., 1976, *J. Biol. Chem.*, v. 251 (8), 2217-2225
Ascaris lumbricoides var. *suis*, NADH:NAD⁺ transhydrogenase from muscle mitochondria, demonstration and possible physiological function in hydride ion translocation
- Ascaris lumbricoides* var. *suis*
Moritz, K. B.; and Roth, G. E., 1976, *Nature* (5538), v. 259, 55-57
Ascaris lumbricoides, *Parascaris equorum*, complexity of germline and somatic DNA
- Ascaris lumbricoides* var. *suis*
Saz, H. J.; and Lescure, O. L., 1968, *Molec. Pharm.*, v. 4 (4), 407-410
Ascaris lumbricoides var. *suis*, inhibition by anticestodal agents of mitochondrial ³²P-ATP exchange reaction indicates that selective toxicity of these compounds for cestodes is a result of differences in permeability between these two groups of helminths
- Ascaris lumbricoides* var. *suis*
Stefl, B.; and Kubistova, J., 1972, *Physiol. Bohemoslov.*, v. 21 (3), 261-263
Ascaris lumbricoides var. *suis*, low actomyosin content of body-wall muscles, possible role in metabolism
- Ascaris lumbricoides* var. *suum*
Coles, G. C.; East, J. M.; and Jenkins, S. N., 1975, *Gen. Pharmacol.*, v. 6 (4), 309-313
Nippostrongylus brasiliensis, *Ascaris lumbricoides* var. *suum*, levamisole, mechanism of action, paralysis and recovery of motility, resistance of recovered worms to paralysis by methyridine, pyrantel, or bephenium, results suggest levamisole acts as ganglion stimulant and that inhibition of fumarate reductase is not relevant to in vivo mode of action
- Ascaris lumbricoides* var. *suum*
Douch, P. G. C., 1976, *Xenobiotica*, v. 6 (9), 531-536
Ascaris lumbricoides var. *suum*, *Moniezia expansa*, azo- and nitro-reductase activities, absence of cytochromes P-450 and b₅, possible new approach for development of ant-helminthic drugs
- Ascaris lumbricoides* var. *suum*
Douch, P. G. C.; and Gahagan, H. M., 1977, *Xenobiotica*, v. 7 (5), 301-307
Moniezia expansa, *Ascaris lumbricoides* var. *suum*, reduction and/or hydrolysis of niclosamide and related compounds by intact helminths and by enzyme preparations from the helminths and from mouse and sheep liver homogenates, reduction of niclosamide inhibited by allopurinol, indicates that co-administration of niclosamide and allopurinol might improve efficacy of anthelmintic, hydrolysis of benzanilide and related compounds inhibited by anthelmintic organophosphates
- Ascaris lumbricoides* var. *suum*
Douch, P. G. C.; and Gahagan, H. M., 1977, *Xenobiotica*, v. 7 (5), 309-314
Ascaris lumbricoides var. *suum*, N-deacetylase, localization and some properties, inhibition by anthelmintic organophosphates indicates they have potential use as adjuvants for anthelmintics of other chemical classes
- Ascaris lumbricoides* var. *suum*, *illus.*
Fill, A.; Goldstein, P.; and Moens, P. B., 1977, *Chromosoma*, v. 65 (1), 21-35
Ascaris lumbricoides var. *suum*, precocious formation of synaptonemal-like polycomplexes and their subsequent fate
- Ascaris lumbricoides* var. *suum*
Fioravanti, C. F.; and Saz, H. J., 1976, *Arch. Biochem. and Biophys.*, v. 175 (1), 21-30
Hymenolepis diminuta, pyridine nucleotide transhydrogenases, comparison with *Ascaris lumbricoides* var. *suum*
- Ascaris lumbricoides* var. *suum*, *illus.*
Goldstein, P., 1977, *J. Morphol.*, v. 152 (2), 141-151
Ascaris lumbricoides var. *suum*, chromatin diminution in early embryogenesis, three characteristic types of mitoses (pre-diminution, diminution, and post-diminution mitosis)

- Ascaris lumbricoides* var. *suum*, *illus.*
Goldstein, P., 1977, *J. Morphol.*, v. 154 (3), 317-337
Ascaris lumbricoides var. *suum*, spermatogenesis and spermiogenesis, morphology of chromosomes
- Ascaris lumbricoides* var. *suum*, *illus.*
Goldstein, P.; and Moens, P. B., 1976, *Chromosoma*, v. 58 (2), 101-111
Ascaris lumbricoides var. *suum*, chromosome number determined from count of synaptonemal complexes, oocyte and spermatocyte pachytene nuclei
- Ascaris lumbricoides* var. *suum*
Grzywacz, M., 1975, *Mater. Med. Pol.* (25), v. 7 (4), 311-313
Ascaris lumbricoides var. *suum*, pharmacodynamic analysis of piperazine adipate penetration into parasite using scintigraphy and autoradiography
- Ascaris lumbricoides suum*
Kasuya, S.; Ohtomo, H.; and Ishizaki, T., 1977, *Japan. J. Med. Sc. and Biol.*, v. 30 (6), 297-307
suppressing effects of purified eosinophils derived from *Ascaris lumbricoides* suum-immunized guinea pigs on lymphocyte blast formation
- Ascaris lumbricoides* var. *suum*
Krell, R. D.; and Chakrin, L. W., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 51 (6), 641-655
fragmented canine lung preparation actively or passively sensitized to *Ascaris* antigen as useful in vitro model for study of immediate-type hypersensitivity reactions, biochemical and pharmacological characterization
- Ascaris lumbricoides* var. *suum*
Sanchez Rasero, F., 1973, *Rev. Iber. Parasitol.*, v. 33 (1), 65-79
Ascaris lumbricoides var. *suum*, determination of DNA and RNA, factors in technique affecting values; values for whole worms, males and females, various organs and tissues
- Ascaris lumbricoides* var. *suum* (Goeze, 1782)
Young, P. L.; and Babero, B. B., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Ascaris mucronata* Schrank
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 189-193
Esox lucius (intestine): Vistula River near Warsaw
- Ascaris ovis*
Vasilev, I.; and Mutafova, T., 1974, *Izvest. Tsentral. Khelmint. Lab.*, v. 17, 33-42
Ascaris suum, *A. ovis*, no difference in karyotype, successful experimental infection of pigs with egg cultures from lambs, *A. ovis* should be considered synonym of *A. suum*
- Ascaris rotundata* Rud. of von Linstow (1880), of Jagerskiold (1894)
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
as syn. of *Pseudanisakis tricupola* nom. nov.
- Ascaris suis*
Kepron, W.; et al., 1977, *Am. Rev. Resp. Dis.*, v. 115 (4, pt. 2), 61 [Abstract]
Ascaris suis, technique for repeated artificial infestation of dogs with infective ova to provide simple and reliable model for study of allergic bronchoconstriction
- Ascaris suis*
Kumagai, Y.; Kumagai, R.; and Mori, N., 1974, *Nihon Univ. J. Med.*, v. 16 (3), 229-240
Ascaris suis, effect of Co-gamma rays on cleavage and development
- Ascaris suum*
Ambler, J.; Miller, J. N.; and Orr, T. S., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (3), 351-361
Ascaris suum, allergen A, spectroscopic and fluorescence properties
- Ascaris suum*
Ambler, J.; Miller, J. N.; and Orr, T. S. C., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 46 (3), 427-437
Ascaris suum allergen A, characterization
- Ascaris suum*
Andersen, S., 1976, *Nord. Vet.-Med.*, v. 28 (6), 322-330
Ascaris suum, negative influence on growth rate of pigs (exper.)
- Ascaris suum*
Anisimov, A. P., 1976, *Tsitologiya*, v. 18 (4), 445-450
Ascaris suum, oesophageal glands, polyploidization and growth of giant nuclei during postnatal ontogenesis
- Ascaris suum*
Annabaeva, G. D.; and Soprunov, F. F., 1975, *Ang. Parasitol.*, v. 16 (3), 170-177
activity of phosphoenolpyruvate carboxykinase in subcellular fractions of *Ascaris suum* and *Fasciola hepatica* tissues, radioisotope and spectrophotometric methods, biological role in metabolism of helminths
- Ascaris suum, illus.*
Araujo, P., 1975, *Ann. Parasitol.*, v. 50 (2), 167-172
Ascaris suum, third-stage larvae, sexual dimorphism in genital primordium and in internal structure of tail (rectal glands)

Ascaris suum

- Archer, G. T.; Robson, J. E.; and Thompson, A. R., 1977, Pathology, v. 9 (2), 137-153
 Ascaris suum, Echinococcus granulosus, isolation from both parasites of a phospholipid capable of inducing eosinophilia and mast cell hyperplasia when injected into rats (exper.), phagocytosis found to be complement dependent and eosinophilia possibly resulted from stimulation of alternate complement pathway by the phospholipid

Ascaris suum

- Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, Vet. Rec., v. 99 (7), 119-122
 swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with pica-dex

Ascaris suum

- Barriga, O. O., 1977, J. Clin. Microbiol., v. 6 (3), 274-279
 Trichinella spiralis, different antigenic fractions, reactivity and specificity (tested for cross-reactions against Ascaris suum) in cutaneous (immediate and delayed) and serological (bentonite agglutination, hemagglutination, hemagglutination inhibition) tests, implications for clinical diagnosis of trichinellosis

Ascaris suum

- Benkova, M.; and Boroskova, Z., 1976, Vet. Med., Praha, v. 49, v. 21 (6), 369-373
 Ascaris suum, rabbit (non-specific host), demonstration of migration phase, sensitivity of complement-fixation and latex-fixation test

Ascaris suum

- Bindseil, E., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 313-318
 Ascaris suum migrating larvae, pathogenesis of liver and lung lesions, differences in pigs vs. mice and young vs. old pigs

Ascaris suum, illus.

- Bogdanov, Iu. F., 1977, Chromosoma, v. 61 (1), 1-21
 Ascaris suum, male meiosis, formation of cytoplasmic synaptonemal-like polycomplexes at leptotene and normal synaptonemal complexes at zygotene

Ascaris suum

- Boisvenue, R. J.; Emmick, T. L.; and Galloway, R. B., 1977, Exper. Parasitol., v. 42 (1), 67-72
 Haemonchus contortus, some compounds with juvenile hormone activity inhibited in vitro development of infective larvae, none of these compounds had anthelmintic properties against Ascaris suum or Nematospiroides dubius in mice

[Ascaris suum] Askaris suum

- Bonev, B.; et al., 1975, Vet. Sbirka, v. 73 (11), 14-15
 pigs, [Ascaris suum], [Trichocephalus suis], [Oesophagostomum dentatum], hygromycin B in feed, completely controls ascariasis

Ascaris suum

- Boroskova, Z.; and Benkova, M., 1974, Vet. Med., Praha, v. 47, v. 19 (5), 271-275
 Ascaris suum larvae, somatic antigen with greater immunogenic properties and serologic activity than antigens from delipidized extract and fraction from mature stages

Ascaris suum

- Boucher, R. C.; et al., 1977, J. Allergy and Clin. Immunol., v. 60 (2), 134-140
 Ascaris suum-sensitive Macaca mulatta, airway mucosal permeability

Ascaris suum

- Bradbury, S. M.; Percy, D. H.; and Strejan, G. H., 1974, Internat. Arch. Allergy and Applied Immunol., v. 46 (4), 498-511
 Ascaris suum, rats infected with embryonated eggs, characteristic pathological changes in liver and lungs, eosinophilia, production of reaginic antibodies, purified Ascaris antigen (Asc-1) present in all stages of parasite life cycle and directly involved in stimulating reagin production during migratory phase of infection

Ascaris suum

- Brown, A. R.; and Crandall, C. A., 1976, J. Immunol., v. 116 (4), 1105-1109
 mice, Ascaris suum-induced phosphorylcholine-binding component identified as IgM antibody having idiotypic determinants in common with PC-binding IgA myeloma TEPC 15, response not duplicated by immunization with dead Ascaris larvae or by infection with Heligmosomoides polygyrus or Trichinella spiralis

Ascaris suum

- Brown, A. R.; Crandall, C. A.; and Crandall, R. B., 1977, J. Parasitol., v. 63 (5), 950-952
 Ascaris suum in mice with X-linked B lymphocyte defect, immune response and acquired resistance

Ascaris suum

- Cappuccinelli, P., 1972, Parassitologia, v. 14 (2-3), 255-260
 Armillifer armillatus, antigens, complement fixation and immunodiffusion studies of antibody response in rabbit; identification of active fractions by immunoelectrophoresis; immunodiffusion tests against Echinococcus granulosus, Fasciola hepatica, Dicrocoelium dendriticum, Onchocerca volvulus and Ascaris suum, no common antigens found

Ascaris suum

- Casarosa, L.; Lugetti, G.; and Marconcini, A., 1973, Isotopes and Radiation Parasitol. III, 113-126
 Ascaris suum, guinea pigs vaccinated and then subjected to whole-body irradiation, enteric wall reactivity against challenge, relationship to in vitro adherence reaction

- Ascaris suum Goeze, 1782
Casarosa, L.; Lugetti, G.; and Marconcini, A., 1974, Ann. Fac. Med. Vet. Pisa, v. 26, 1973, 385-401
Ascaris suum-vaccinated guinea pigs, total body x-irradiation and challenge infection, enteric wall reactivity, globule leukocytes, immunoglobulin-containing cells; globule leukocytes depleted in challenged hosts; higher number of fluorescing mature plasma cells in lamina propria of vaccinated animals
- Ascaris suum
Church, E. M.; Wyand, D. S.; and Lein, D. H., 1975, Am. J. Vet. Research, v. 36 (3), 331-335
cerebrospinal nematodiasis, experimentally induced in *Oryctolagus cuniculus* with *Ascaris columnaris*, *A. suum*, or *Toxocara canis*, naturally occurring in *Sylvilagus floridanus* and *O. cuniculus*, clinical signs, gross and microscopic changes, duration of infection and parasite morphology and distribution in CNS, potential of rabbits as intermediate or paratenic hosts for ascarids of carnivorous origins
- Ascaris suum
Collins, R. F.; and Ivey, M. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 455-459
skin test responses in guinea pigs infected with small numbers of *Toxocara canis* or *Ascaris suum* and challenged intradermally with several adult and larval somatic antigenic preparations
- Ascaris suum
Collins, R. F.; and Ivey, M. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 460-464
passive cutaneous anaphylaxis responses of sensitized guinea pigs to various antigens of adult and larval stages of *Toxocara canis* or *Ascaris suum*; homologous reactions; *Ascaris* larval antigen reacted with *Toxocara* antiserum
- Ascaris suum
Connan, R. M., 1977, Vet. Rec., v. 100 (20), 421-422
Ascaris suum, time required for eggs to become infective in a pig house at different times of the year, epidemiological significance
- Ascaris suum
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
Sus scrofa domesticus x *Sus scrofa cristatus* (small intestine): Aransas National Wildlife Refuge, southern Texas
- Ascaris suum
Corwin, R. M., 1977, Am. J. Vet. Research, v. 38 (4), 465-467
mixed nematode infections, pigs, oxfendazole, critical evaluation: Missouri
- Ascaris suum
Cotton, D. J.; et al., 1977, J. Applied Physiol.: Respiratory, Environmental and Exercise Physiol., v. 42 (1), 101-106
Ascaris suum antigen inhalation by dogs, rapid shallow breathing, results indicate that vagal afferent pathways mediate antigen-induced tachypnea and this response does not primarily depend on bronchoconstriction
- Ascaris suum
Crandall, C. A., 1976, Exper. Parasitol., v. 39 (1), 69-73
Ascaris suum, mice, measurement of homocytotropic antibody response (IgG₁, IgE), infection did not potentiate reaginic response to ovalbumin, not promising model for study of reagin production in helminth infections
- Ascaris suum
Crandall, C. A.; and Crandall, R. B., 1976, Exper. Parasitol., v. 40 (3), 363-372
Ascaris suum, mice, definite but selective immunosuppression during acute infection
- Ascaris suum
Delespesse, G.; Ishizaka, K.; and Kishimoto, T., 1975, J. Immunol., v. 114 (3), 1065-1071
rabbit lymphocyte populations responding to haptenic and carrier determinants for DNA synthesis, DNP-*Ascaris suum* conjugate used as one of antigens
- Ascaris suum
Dey-Hazra, A., 1976, Ztschr. Parasitenk., v. 50 (2), 198
helminths, pigs, mode of pathogenicity, review
- Ascaris suum
Dobson, C.; Rockey, J. H.; and Soulsby, E. J. L., 1971, J. Immunol., v. 107 (5), 1431-1439
Ascaris suum, guinea pigs, characterization of IgE antibodies
- Ascaris suum
Dobson, C.; and Welch, J. S., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 223-228
survey for antibodies against *Dirofilaria immitis*, *Toxocara canis*, *Ascaris suum*, *Angiostrongylus cantonensis*, *A. mackerrasae*, in patients with eosinophilia using fluorescent antibody test and passive reversed Arthus test in guinea pigs; *D. immitis* implicated as etiologic agent of human eosinophilic meningitis: Australia
- Ascaris suum
Dorf, M. E.; et al., 1975, J. Immunol., v. 114 (6), 1717-1719
in vivo cooperative responses between lymphocytes are controlled by genes in K-end of H-2 complex, DNP-*Ascaris suum* used as antigen
- Ascaris suum
El-Moukdad, A.-R., 1977, Wien Tierarztl. Monatschr., v. 64 (3), 85-91
effect of disinfectants on eggs of *Ascaris suum*, *Toxascaris leonina*, small horse strongyles and coccidia oocysts

- Ascaris suum
Enigk, K.; et al., 1975, Zentralbl. Vet.-Med., Reihe B, v. 22 (8), 687-702
survival of resistant external stages of parasites during fermentation of liquid cattle manure at high temperatures
- Ascaris suum
Enigk, K.; and Dey-Hazra, A., 1976, Prakt. Tierarzt, v. 57 (4), 232-234
Strongyloides ransomi, prepatent Ascaris suum, experimental infection in swine, Cam-bendazol in feed, effective treatment
- Ascaris suum, illus.
Enigk, K.; and Dey-Hazra, A., 1976, Berl. u. Munchen. Tierarztl. Wchnschr., v. 89 (14), 276-281
Ascaris suum eggs, protein layer, surface structure and function
- Ascaris suum
Ershov, V. S.; et al., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 343-348
anaphylactic shock in guinea pigs after sensitization with free-living or plant-parasitic nematodes and challenge with various helminth antigens indicates antigenic components in common; intradermal tests using antigen from free-living nematode in cases of ascariasis, trichinellosis, and cysticercosis; possible use of free-living nematode to immunize against dictyocaulosis and ascariasis
- Ascaris suum
Fairchild, S. S.; and Malley, A., 1975, J. Immunol., v. 115 (2), 446-449
induction of mouse homocytotropic antibodies to timothy pollen antigen coupled to purified fraction of Ascaris suum
- Ascaris suum, illus.
Foor, W. E., 1976, J. Parasitol., v. 62 (6), 849-864
Ascaris suum, glandular vas deferens, structure and function
- Ascaris suum
Gaur, S. N. S.; and Dutt, S. C., 1975, Indian Vet. J., v. 52 (6), 474-477
mice fed with irradiated and non-irradiated Ascaris suum eggs, reduced size, motility, and longevity of resulting migrating A. suum juveniles recovered from liver and lungs
- Ascaris suum
Gaur, S. N. S.; and Dutt, S. C., 1976, Indian Vet. J., v. 53 (4), 305-306
immunizing effect of single dose of irradiated Ascaris suum eggs, mice, guinea pigs, pathologic changes
- Ascaris suum
Gaur, S. N. S.; and Dutt, S. C., 1977, Pantnagar J. Research, v. 2 (1), 80-84
Ascaris suum eggs, immunizing effect in mice, rats and guinea-pigs against migratory juveniles; double immunizing doses in mice produced stronger immunity but more tissue damage
- Ascaris suum
Gaur, S. N. S.; and Dutt, S. C., 1977, Pantnagar J. Research, v. 2 (2), 190-191
Ascaris suum, guinea pigs, successful immunization by subcutaneous administration of pooled immune sera, challenge by migrating juveniles of A. suum
- Ascaris suum
Gaur, S. N. S.; Dutt, S. C.; and Sokolic, A., 1975, Ceylon Vet. J., v. 23 (3-4), 46-48
Ascaris suum, mice, attenuation of eggs with various doses of gamma-rays, infectivity of developing larvae; preliminary to immunization studies
- Ascaris suum
Getler, K., 1972, Med. Wet., v. 28 (8), 476-477
nematodes, pigs on industrial swill feeding farm, Atgard
- Ascaris suum
Goldstein, P., 1977, J. Parasitol., v. 63 (4), 689
Ascaris suum, females, presence of extra reproductive tract in 23 of 2000 worms dissected
- Ascaris suum
Grunnet, K.; and Henriksen, S. A., 1977, Nord. Vet.-Med., v. 29 (10), 458-459
Ascaris suum eggs, elimination from sewage by air flotation (bubbling air-saturated water through sewage) after sedimentation, preliminary experiments
- Ascaris suum
Grzywinski, L.; et al., 1975, Medycyna Wet., v. 31 (9), 524-526
swine, Nilverm by injection for control
- Ascaris suum
Grzywinski, L.; Martynowicz, T.; and Kluczniok, P., 1976, Med. Wet., v. 32 (4), 227-229
Oesophagostomum, Ascaris suum, pigs, Cambendazole effective
- Ascaris suum, illus.
Guerrero, J.; and Mercadante, M. L. P., 1976, Rev. Fac. Med. Vet. e Zootec. Univ. S. Paulo, v. 13 (2), 391-400
Ascaris suum larvae, simplification of culture medium used for in vitro cultivation, results indicate that hemin plus any other protein supplement (replacing fetal calf serum) considerably increase larval development and survival
- Ascaris suum
Guerrero, J.; and Silverman, P. H., 1972, Rev. Invest. Pecuarias, v. 1 (2), 209-213
Ascaris suum, larval culture method, metabolic antigens immunizing mice
- Ascaris suum
Guerrero, J.; and Silverman, P. H., 1972, Rev. Invest. Pecuarias, v. 1 (2), 215-222
Ascaris suum larval culture, various media with gases; Eagles medium in air plus CO₂, best immunization of mice with metabolic antigens

- Ascaris suum*
Gupta, R. P., 1973, Indian Vet. J., v. 50 (1), 18-22
Ascaris suum, piglets (exper.), cambendazole, ineffective against migratory larvae
- Ascaris suum*, *illus.*
Gutman, G. A.; and Mitchell, G. F., 1977, Exper. Parasitol., v. 43 (1), 161-168
Ascaris suum, location of phosphorylcholine in lung larvae confined to internal membranous structures and lining of intestinal tract, outside of cuticle is negative, availability of phosphorylcholine-containing parasite antigens to host for immune induction may occur through intestinal excretions, damage to larvae, or at larval moulting stages
- Ascaris suum*
Hale, O. M.; Stewart, T. B.; and Johnson, J. C., jr., 1977, Research Bull. (203) Agric. Exper. Stations Univ. Georgia, 3-21
Strongyloides ransomi, *Ascaris suum*, naturally infected gilt and/or barrow crossbred pigs, superimposed *S. ransomi* infection, differences in performance with diets of varying levels of protein and vitamins not significant
- Ascaris suum*
Halliwell, R. E. W., 1973, J. Immunol., v. 110 (2), 422-430
canine skin, immunofluorescent staining for IgE and *Ascaris*-binding antibodies, association with mast cells, correlation with results of intradermal skin tests
- Ascaris suum*
Hamaoka, T.; et al., 1974, J. Immunol., v. 113 (3), 958-973
Ascaris suum, mice, establishment of parameters for generation of helper T cell function regulating primary and secondary responses of IgE and IgG B lymphocytes
- Ascaris suum*, *illus.*
Harpur, R. P., 1975, J. Parasitol., v. 61 (5), 881
Ascaris suum, specimen with three uteri
- Ascaris suum*
Harpur, R. P.; and Jackson, D. M., 1975, J. Parasitol., v. 61 (5), 808-814
Ascaris suum, pieces of gut tissue in vitro, oxygen consumption, production of fermentation acids, anaerobic synthesis of protein
- Ascaris suum*
Harpur, R. P.; and Jackson, D. M., 1976, Comp. Biochem. and Physiol., v. 54 (4B), 455-460
Ascaris suum, isolated intestine, glucose diminished oxygen uptake (Crabtree effect), glucose caused increases in tissue carbohydrate, CO₂ evolution, and acid production, no Pasteur effect demonstrated, glutamine increased oxygen uptake in presence and absence of glucose but did not abolish Crabtree effect
- Ascaris suum*
Hinck, L. W.; and Ivey, M. H., 1976, J. Parasitol., v. 62 (5), 771-774
Ascaris suum, proteinase activity in eggs, hatching fluid, and excretions-secretions of hatched larvae
- Ascaris suum*
Hirashima, M.; and Hayashi, H., 1976, Immunology, v. 30 (2), 203-212
two different chemotactic factors for eosinophils, isolation from allergic tissue lesions induced by DNP-*Ascaris suum* extract in guinea pigs, characterization
- Ascaris suum*
Hoegel, E.; and Schroetter, E., 1973, Pharmazie, v. 28 (10), 636-640
Ascaris suum, mice (exper.), laboratory synthesis and trials with substituted β -aminoethyl-[4-n-alkoxy-3-ethyl-phenyl]-ketonhydrochloride for possible anthelmintic activity
- Ascaris suum*
Hogarth-Scott, R. S., 1967, Internat. Arch. Allergy and Applied Immunol., v. 32 (2), 201-207
Toxocara canis, *T. cati*, *Toxascaris leonina*, *Ascaris suum*, rabbits (exper.), presence of reagin-like antibodies demonstrable by homologous passive cutaneous anaphylaxis, responsible allergens were common to all 4 nematode species
- Ascaris suum*
Hubert, J., 1977, Rec. Med. Vet., v. 153 (12), 923-929
Ascaris suum (exper.), *Trichuris suis* (exper.), *Oesophagostomum* spp., pigs, comparison of coproscopical method of count after dilution in dense solution and flotation in Mac Master Slides and count after sedimentation and flotation in dense solution in Mac Master Slides; various densities of solutions compared with both techniques
- Ascaris suum*
Hubert, J.; Yvore, P.; and Kerboeuf, D., 1976, Ann. Recherches Vet., v. 7 (1), 83-90
parasite survival in liquid manure, anti-parasitic action of xylene
- Ascaris suum*
Hung, C. H.; et al., 1977, J. Biol. Chem., v. 252 (11), 3995-4001
Ascaris suum, intestinal basement membrane, analysis of polypeptide components, amino acid and carbohydrate composition
- Ascaris suum*
Hussain, R.; Bradbury, S. M.; and Strejan, G., 1973, J. Immunol., v. 111 (1), 260-268
Ascaris suum, characterization of highly purified allergen
- Ascaris suum*
Hussain, R.; Strejan, G.; and Campbell, D. H., 1972, J. Immunol., v. 109 (3), 638-647
Ascaris suum allergen, isolation and partial characterization

- Ascaris suum*
Ince, A. J., 1976, *Internat. J. Parasitol.*, v. 6 (2), 127-128
Ascaris suum, analytical determinations of trace element levels by method of activation analysis, possible trace element interrelationships
- Ascaris suum*
Ishizaka, K.; and Kishimoto, T., 1972, *J. Immunol.*, v. 109 (1), 65-73
Ascaris suum, rabbits, formation of reaginic antibody
- Ascaris suum*
Ito, K.; et al., 1976, *Japan. J. Exper. Med.*, v. 46 (1), 79-85
levels of immunoglobulin E antibodies to Ascaris using Ascaris suum antigens higher in Papua New Guineans than in Japanese subjects
- Ascaris suum*
Jablonowski, Z., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 271-284
Ascaris suum, guinea pigs, vitamin C-deficient diet, haemogram level, serum total protein and protein fractions, vitamin C content of various organs
- Ascaris suum*
Jakovljevic, D. D., 1975, *Acta Vet. Beograd*, v. 25 (6), 315-325
Ascaris suum, method for obtaining embryonated eggs capable of infection; number of developed worms in intestines inversely related to number of administered eggs and age of suckling pigs; pathomorphological changes in liver and lungs; earthworms (*Allolobophora caliginosa*, *Octolasion transpadanum*, *Lumbricus rubellus*, *Allolobophora leoni*) have only a passive role as 'carriers' of embryonated eggs in transfer of infection to swine; incidence and economic importance: Yugoslavia
white mice (exper.)
guinea pigs (exper.)
- Ascaris suum*, *illus.*
Jeska, E. L., 1973, *J. Reticuloendothel. Soc.*, v. 14 (2), 146-157
Ascaris suum infective juveniles, cell adherence reactions of mouse peritoneal exudate system in vitro
- Ascaris suum*
Johnson, H. G.; and van Hout, C. A., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 50 (4), 454-462
Ascaris suum-sensitive monkeys, respiratory response after aerosol challenge with Ascaris antigen, poor correlation with cutaneous response, inhibition of allergic reaction by cromoglycate and by a new anti-allergic drug
- Ascaris suum*
Johnson, J. C., jr.; Stewart, T. B.; and Hale, O. M., 1975, *J. Parasitol.*, v. 61 (3), 517-524
responses of pigs to natural infections of *Strongyloides ransomi* and *Ascaris suum* and to superimposed artificial infection with *Strongyloides ransomi*: effects of breed (Duroc, Hampshire, Duroc-Hampshire crossbred), level of *Strongyloides ransomi* infection, and season (spring, fall) on performance of growing-finishing pigs
- Ascaris suum*
Jones, H. I., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (2), 219-226
use of haemagglutination test as aid in study of epidemiology of human *Ascaris lumbricoides* infections, A. suum used as antigen: Papua New Guinea and East Timor
- Ascaris suum*
Jorgensen, R. J.; et al., 1975, *Vet. Parasitol.*, v. 1 (2), 151-157
Ascaris suum, piglets (exper.), failure to establish patent infections with primary infection dose of 10,000 eggs, significant eosinophil response; patent infections established with dose of 50 eggs, no significant eosinophil response; mechanism of expulsion of high level infections not known
- Ascaris suum*
Katz, D. H.; et al., 1974, *J. Immunol.*, v. 113 (3), 974-983
Ascaris suum, mice, evidence for distinctive sensitivities of IgE and IgG B lymphocytes to regulatory influences of T cells
- Ascaris suum*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiasis associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Ascaris suum*
Kessler, G. F.; Frick, O. L.; and Gold, W. M., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (3), 313-328
Toxocara canis or *Ascaris suum*-sensitive dogs, experimental asthma, immunologic and physiologic characterization of role of reaginic antibodies
- Ascaris suum*
Khoury, P. B.; and Soulsby, E. J. L., 1977, *Exper. Parasitol.*, v. 41 (1), 141-159
Ascaris suum, guinea pigs, lymphoid cell responses during primary infections assessed by antigen-induced lymphocyte transformation, rosette-formation, and rosette-plaques techniques, progression of local lymphoid cell responses in lymphoid organs draining parasitized tissues
- Ascaris suum*
Khoury, P. B.; and Soulsby, E. J. L., 1977, *Exper. Parasitol.*, v. 41 (2), 432-445
Ascaris suum, immunized guinea pigs given challenge infections, lymphoid cell responses of draining lymph nodes and spleen, in vitro antigen-induced lymphocyte transformation, rosette formation, rosette inhibition, and rosette-plaques techniques
- Ascaris suum*
Khoury, P. B.; Stromberg, B. E.; and Soulsby, E. J. L., 1977, *Immunology*, v. 32 (4), 405-411
Ascaris suum, guinea pigs, passive transfer of immunity by cells or serum, significant protection with immune IgG2, IgE + IgG1 and whole immune serum or with lymphocytes from hepatic and mediastinal lymph nodes of immune animals, minimal protection with IgM and IgA, spleen lymphocytes enhanced rather than reduced degree of infection

- Ascaris suum**
Kimoto, M.; et al., 1977, *J. Immunol.*, v. 118 (3), 840-845
induction of in vitro IgE antibody response in murine spleen cells and demonstration of possible involvement of distinct T-helper cells in IgE and IgG antibody responses, *Ascaris suum* and DNP-A. suum used as antigens
- Ascaris suum**
Kishimoto, T.; et al., 1975, *J. Immunol.*, v. 115 (5), 1179-1184
effect of anti-immunoglobulin and enhancing soluble factor on differentiation and proliferation of B cells, *Ascaris suum* used as antigen
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1972, *J. Immunol.*, v. 109 (3), 612-622
Ascaris suum, rabbits, role of hapten-specific memory cells and carrier-specific helper cells on distribution of anti-hapten antibodies in IgG, IgM, and IgE classes
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1972, *J. Immunol.*, v. 109 (6), 1163-1173
Ascaris suum, rabbits, distribution of immunoglobulin heavy chain antigenic determinants on hapten-specific memory cells
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1973, *J. Immunol.*, v. 111 (1), 1-9
Ascaris suum, rabbits, effect of carrier-specific (T) helper cells on generation of hapten-specific (B) memory cells of different immunoglobulin classes
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1973, *J. Immunol.*, v. 111 (3), 720-732
Ascaris suum, used as priming or supplemental immunization in combination with various other antigens, results show that carrier-specific helper cells for IgE antibody formation are different from those for IgG/IgM antibody formation
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1974, *J. Immunol.*, v. 112 (5), 1685-1697
Ascaris suum, rabbits, multiplicity of soluble factors released from carrier-specific cells
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1975, *J. Immunol.*, v. 114 (2, pt. 1), 585-591
induction of secondary anti-hapten IgE antibody response by anti-immunoglobulin and enhancing soluble factor, DNP-*Ascaris suum* used as priming antigen
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1975, *J. Immunol.*, v. 114 (4), 1177-1184
immunologic and physicochemical properties of enhancing soluble factors for IgG and IgE antibody responses, DNP-*Ascaris suum* conjugate used as antigen
- Ascaris suum**
Kishimoto, T.; and Ishizaka, K., 1976, *J. Immunol.*, v. 116 (2), 534-541
biphasic effect of cyclic AMP on secondary anti-hapten antibody response to anti-immunoglobulin and enhancing soluble factor, DNP-*Ascaris suum* used as antigen
- Ascaris suum**
Klucas, C. A.; and Rhodes, M. B., 1976, *Proc. Nebraska Acad. Sc.*, 20
Ascaris suum, hatching fluid, immunization of pigs
- Ascaris suum**
Koehler, P., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 187
Ascaris suum, chemotherapy, effect on mitochondrial hydrogen transport
- Ascaris suum**
Koehler, P., 1977, *Internat. J. Biochem.*, v. 8 (2), 141-147
Ascaris suum, transport of dicarboxylates and some properties of fumarase in muscle mitochondria
- Ascaris suum**
Komandarev, S.; and Mikhov, L., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 111-118
Trichinella spiralis-immunized mice, immunity to homologous and heterologous (*Ascaris suum*) challenge
- Ascaris suum**
Krvavica, S.; Francetic, D.; and Zivkovic, D., 1976, *Vet. Arhiv, Zagreb*, v. 46 (9-10), 231-239
nematodes, trematodes, cestodes, activity, distribution and cofactor dependence of malic enzymes, majority are located in mitochondria in all investigated parasites
- Ascaris suum**
Kuo, C. Y.; and Yoo, T. J., 1977, *Internat. Arch. Allergy and Applied Immunol.*, v. 54 (4), 308-314
Ascaris suum, new allergen obtained from perienteric fluid through isoelectric focusing techniques
- Ascaris suum, illus.**
Leake, L. D., 1975, *Comparative histology. An introduction to the microscopic structure of animals.*, 738 pp.
parasites, comparative histology, textbook
- Ascaris suum**
Lee, Y. C.; and Huang, K. J., 1975, *J. Chinese Soc. Vet. Sc.*, v. 1 (1), 24-30
Ascaris suum, pigs, relationship to prevalence of infectious pneumonia caused by bacteria, more severe lesions with both present
- Ascaris suum**
Lee, Y. C.; and Liu, C. C., 1976, *J. Chinese Soc. Vet. Sc.*, v. 2 (2), 59-61
Ascaris suum eggs, isolation of *Escherichia coli*, *Alcaligenes faecalis* and *Pseudomonas aeruginosa*

Ascaris suum

Leventhal, R.; and Soulsby, E. J. L., 1976, Internat. J. Parasitol., v. 6 (3), 279-283
 Ascaris suum larvae, adhesion and degranulation of polymorphonuclear leukocytes on surface, evaluation of serum components which are responsible for opsonization

Ascaris suum, illus.

Leventhal, R.; and Soulsby, E. J. L., 1977, Exper. Parasitol., v. 41 (2), 423-431
 Ascaris suum early larval stages, cuticular binding of third component of complement

Ascaris suum

Mackler, B.; Malley, A.; and Amkraut, A. A., 1971, Internat. Arch. Allergy and Applied Immunol., v. 41 (5), 765-777
 in vitro lymphocyte transformation demonstrated using peripheral lymphocytes from rhesus monkeys showing 'pure' atopic hypersensitivity (Ascaris suum-induced)

Ascaris suum, illus.

Madden, P. A.; and Tromba, F. G., 1976, J. Parasitol., v. 62 (2), 265-271
 Ascaris suum adults of known ages recovered from pigs experimentally infected with eggs from the same stock, variation in number, size, and shape of lip denticles, concluded that denticles are functional and become worn through use

Ascaris suum

Malley, A.; et al., 1968, J. Immunol., v. 101 (2), 292-300
 Ascaris suum, Macaca mulatta inoculated with crude worm extract or electrophoretically purified fraction, atopic-type hypersensitivity demonstrated by direct active cutaneous anaphylaxis and by passive transfer experiments; second fraction showed toxic effects but did not sensitize normal M. mulatta nor cross-react with the sensitizing fraction

Ascaris suum

Matausic, M.; et al., 1976, Vet. Arhiv, Zagreb, v. 46 (7-8), 189-196
 Ascaris suum, extraction and identification of polar lipids from isolated organs and eggs in uterus

Ascaris suum Goeze, 1782, illus.

Maung, M., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (1), 41-45
 Ascaris lumbricoides from man, Ascaris suum from pigs, morphological comparisons of denticles, cuticular striations, eggs and spicules

Ascaris suum

Mazhuga, N. A., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 98-101
 Ascaris suum, proteolytic enzymes, optimal pH; action against synthetic substrate; activity distinct from that of trypsin of host

Ascaris suum

Mitchell, G. F., 1976, Internat. Arch. Allergy and Applied Immunol., v. 52 (1-4), 79-94
 Ascaris suum body fluid as antigen mixture, mice, effects on circulating reagent titers of manipulations such as T cell deprivation and reconstitution, lipopolysaccharide and cyclophosphamide injection, and altered route of administration of antigen

Ascaris suum

Mitchell, G. F.; et al., 1976, Internat. Arch. Allergy and Applied Immunol., v. 52 (1-4), 64-78
 Ascaris suum, mouse strain variation in susceptibility, resistance to second infection, comparison of susceptibility of normal vs. hypothyroid nu/nu mice, appearance of antibodies in infected mice reacting with phosphorylcholine

Ascaris suum

Mitchell, G. F.; and Lewers, H. M., 1976, Internat. Arch. Allergy and Applied Immunol., v. 52 (1-4), 235-240
 Ascaris suum, Nippostrongylus brasiliensis, mice, inhibition of an anti-DNP antibody response with DNP-Ficoll containing phosphorylcholine, results suggest that parasites may 'utilize' molecules such as phosphorylcholine to induce state of selective tolerance to parasite antigens as a mechanism for facilitating survival

Ascaris suum

Newburger, P. E.; Hamaoka, T.; and Katz, D. H., 1974, J. Immunol., v. 113 (3), 824-829
 effect of bacterial lipopolysaccharide on hapten-specific IgG and IgE response to DNP-Ascaris suum primed spleen cells, analysis in adoptive transfer system in mice shows that adjuvant effect is due to action on carrier-specific T cell function

Ascaris suum

Niel, G.; et al., 1972, Medecine et Malad. Infect., v. 2 (5), 193-202
 filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of Setaria labiatopapillosa as antigen, comparison with Dipetalone-ma vitae and Ascaris suum as antigens

Ascaris suum

Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
 Sus scrofa: Chile

Ascaris suum

Oelerich, S.; and Nwokolo, C., 1974, Tropenmed. u. Parasitol., v. 25 (2), 137-146
 Paragonimus uterobilateralis, sera from 27 patients, complement fixation, indirect hemagglutination, double gel diffusion, reactions with homologous antigen and cross-reactions with other helminth antigens, disc-electrophoretic analysis of P. uterobilateralis antigen: Nigeria

- Ascaris suum
Oelerich, S.; Umaly, R. C.; and Lederer, I.,
1974, Tropenmed. u. Parasitol., v. 25 (3),
318-326
Schistosoma mansoni, different developmental
stages. S. japonicum, Fasciola hepatica,
Ascaris suum, cross reactions in double gel
diffusion, Cerkarienhullenreaktion, comple-
ment fixation, indirect immunofluorescence,
indirect haemagglutination, mice, rabbits
- Ascaris suum
Okudaira, H.; and Ishizaka, K., 1973, J. Im-
munol., v. 111 (5), 1420-1428
spleen cells from mice immunized with dini-
trophenylated derivative of Ascaris suum
extract (DNP-Asc) cooperated with ovalbumin-
specific helper cells to form anti-DNP IgE
antibody after DNP-ovalbumin administration
- Ascaris suum
Okumura, K.; and Tada, T., 1971, J. Immunol.,
v. 107 (6), 1682-1689
Ascaris suum, rats, homocytotropic antibody
response, inhibition by thymocytes and
spleen lymphocytes from hyperimmunized
donors
- Ascaris suum
Okumura, K.; and Tada, T., 1974, J. Immunol.,
v. 112 (2), 783-791
Ascaris suum, rats, chemical and physico-
chemical characterization of antigen-speci-
fic inhibitory T cell factor in hapten-speci-
fic homocytotropic antibody response
- Ascaris suum
Okumura, K.; Tada, T.; and Ochiai, T., 1974,
Immunology, v. 26 (2), 257-268
Ascaris suum, rats, time and dose of anti-
thymocyte serum administration are crucial
factors in determining subsequent suppressed
or enhanced reaginic antibody responses
- Ascaris suum
Osikovski, E.; Koenigk, E.; and Schmidt, G.,
1972, Izvest. Tsentral. Khelmin. Lab., v. 15,
159-166
Ascaris suum, tRNA methylase, characteriza-
tion
- Ascaris suum
Panitz, E., 1977, J. Helminth., v. 51 (1),
23-30
ethyl-6-ethoxybenzothiazole-2-carbamate,
evaluation of anthelmintic activity in po-
nies, swine, lambs, and chickens
- Ascaris suum
Pare, P. D.; Michoud, M.-C.; and Hogg, J. C.,
1976, J. Applied Physiol., v. 41 (5, pt. 1),
668-676
Ascaris suum-sensitive Macaca mulatta, lung
mechanics following antigen challenge as an
animal model for human asthma
- Ascaris suum
Patterson, R.; et al., 1975, Pediatrics, Am.
Acad. Pediat., v. 56 (3), 417-420
differentiation of visceral larva migrans
caused by toxocariasis from that caused by
ascariasis using radioimmunoassay to demon-
strate serum immunoglobulins
- Ascaris suum
Patterson, R.; and Booth, B. H., 1971, N. York
State J. Med., v. 71 (7), 755-759
Ascaris suum-infected dogs and monkeys used
as animal model for immediate type human
respiratory allergy
- Ascaris suum
Patterson, R.; Irons, J. S.; and Harris, K.
E., 1975, Internat. Arch. Allergy and Applied
Immunol., v. 48 (3), 412-421
Ascaris-induced reagin-mediated model of
asthma in rhesus monkey, potentiating effect
of D₂O studied with double aerosolized anti-
gen challenge technique
- Ascaris suum
Patterson, R.; Talbot, C. H.; and Brandfon-
brener, M., 1971, Internat. Arch. Allergy and
Applied Immunol., v. 41 (4), 592-603
Ascaris suum-sensitive Macaca mulatta chal-
lenged with Ascaris antigen, effect of di-
sodium cromoglycate in respiratory and cuta-
neous reactions and on electrocardiograms
- Ascaris suum
Patterson, R.; Talbot, C. H.; and Roberts, M.,
1972, Clin. and Exper. Immunol., v. 10 (2),
267-274
reverse passive respiratory reactions due
to anti-IgE in Macaca mulatta both with and
without respiratory reactivity to Ascaris
suum antigen
- Ascaris suum
Pavlov, A. V.; Shishova-Kasatochkina, O. A.;
and Volynskaia, K. B., 1970, Parazitologiya,
Leningrad, v. 4 (3), 231-236
Ascaris suum, comparative transport L- vs.
D-isomers of arginine, tryptophan, phenyl-
alanine, and alanine in vitro
- Ascaris suum
Peczon, B. D.; et al., 1977, J. Biol. Chem.,
v. 252 (11), 4002-4006
Ascaris suum, intestinal basement membrane,
characterization of carbohydrate units
- Ascaris suum
Perper, R. J.; Sanda, M.; and Lichtenstein, L.
M., 1972, Internat. Arch. Allergy and Applied
Immunol., v. 43 (6), 837-844
Ascaris suum, rhesus monkeys, inhibition of
dermal allergic reaction by cAMP active
agents
- Ascaris suum
Phillips, J. L.; Sturman, G.; and West, G. B.,
1975, Gen. Pharmacol., v. 6 (4), 295-297
Ascaris suum, content and distribution of
histamine in tissues
- Ascaris suum
Phillips, J. L.; Sturman, G.; and West, G. B.,
1976, Brit. J. Pharmacol., v. 57 (3), 417-420
comparative study of the interaction between
histamine and piperazine and histamine and
bephenium in Ascaris suum

- Ascaris suum
Pinon, J. M.; and Dropsy, G., 1977, J. Immunol. Methods, v. 16 (1), 15-22
various human parasitic diseases, application of enzyme-linked-immuno-electro-diffusion assay (combination of immunoelectrodifusion and immunoenzyme method), sensitivity and specificity, enables class of immunoglobulins involved to be determined
- Ascaris suum
Pinon, J. M.; and Dropsy, G., 1977, Path. Biol., v. 25 (1), 23-27
human parasitic diseases, use of enzyme-linked-immuno-electro-diffusion-assay (ELIEDA) in diagnosis and immunologic studies
- Ascaris suum, illus.
Plotnikova, S. I.; Shishov, B. A.; and Kuz'mina, L. V., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 103-108
Ascaris suum, biogenous monoamines in nervous system
- Ascaris suum
Raynaud, J. P., 1976, Pathophysiol. Parasit. Infect., 99-104
Oesophagostomum spp., Hyostrogylus rubidus, Ascaris suum, young swine (exper.), multi-stage multiparasite model for pathological and anthelmintic studies
- Ascaris suum
Rhodes, M. B.; et al., 1977, Exper. Parasitol., v. 42 (2), 356-362
Ascaris suum embryonated eggs, hatching in orally inoculated pigs, in ligated intestinal segments, and in isolated intestinal loops of pigs, immune status of pig had no effect on hatching
- Ascaris suum
Richharia, V. S.; Jeska, E. L.; and Greve, J. H., 1975, J. Parasitol., v. 61 (6), 1113-1115
Ascaris suum, swine (exper.), demonstration of true delayed hypersensitivity responses
- Ascaris suum Goetze, 1782
Rivera, M. A.; and Gaafar, S. M., 1976, Vet. Parasitol., v. 2 (4), 341-353
Ascaris suum, pigs, sequential development of esophagogastric ulcers induced by infection
- Ascaris suum, illus.
Rodrick, G. E.; et al., 1977, Exper. Parasitol., v. 42 (1), 150-156
Ascaris suum, mitochondrial DNA in fertilized eggs and adult body muscle, observations agree with previous conclusion that egg is typically aerobic whereas energy metabolism of muscle is essentially anaerobic
- Ascaris suum, illus.
Roneus, O.; and Christensson, D., 1977, Vet. Parasitol., v. 3 (4), 371-375
calves (faeces, ductus choledochus, intestine): Sweden
- Ascaris suum, illus.
Rubin, H.; and Trelease, R. N., 1975, J. Parasitol., v. 61 (4), 577-588
Ascaris suum, developing larvae, correlation of ultrastructural changes in lipid body and glycogen patterns with certain biochemical events occurring during lipid to carbohydrate interconversion, elucidation of specific tissue sites and accompanying organelles associated with this metabolic conversion
- Ascaris suum
Rubin, H.; and Trelease, R. N., 1976, J. Cell Biol., v. 70 (2, pt. 1), 374-383
Ascaris suum larvae, evidence favors localization of glyoxylate cycle enzymes in mitochondria
- Ascaris suum
de Savigny, D. H., 1975, J. Parasitol., v. 61 (4), 781-782
larvae hatched and maintained in Eagle's Minimal Essential Medium with Hanks' salts (HMEM) did not survive longer than 21 days
- Ascaris suum
de Savigny, D. H.; and Tizard, I. R., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 501-507
Toxocara larva migrans, larval excretions and secretions from in vitro cultures used as antigen in passive hemagglutination and fluorescent antibody tests to diagnose visceral larva migrans in man and laboratory animals (exper.), preliminary evaluation for serodiagnostic purposes, no cross reactions with Ascaris suum infections
- Ascaris suum
Saz, H. J.; and Dunbar, G. A., 1975, J. Parasitol., v. 61 (5), 794-801
stibophen inhibition of phosphofructokinase
- Ascaris suum, illus.
Schaffert, R.; and Strauch, D., 1976, Berl. u. Munchen. Tierarztl. Wchnschr., v. 89 (20), 399-402
Ascaris suum eggs in municipal sewage and pig slurry, rotating aeration, temperatures above 50° C necessary for egg destruction, pH not significant
- Ascaris suum
Schroetter, E.; and Hoegel, E., 1972, Pharmazie, v. 27 (2), 93-94
Ascaris suum, in vitro testing of substituted phenylhydrazones as possible anthelmintics
- Ascaris suum
Schroetter, E.; Hoegel, E.; and Tschaeppe, M., 1975, Pharmazie, v. 30 (3), 147-151
Ascaris suum, 2-ethylphenol derivatives ovidical action in laboratory trials with mice, promising results
- Ascaris suum
Shivacheva, T.; and Terziiski, A., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 169-185
Ascaris suum, mice, dynamics of cellular reaction of lymphoid tissue in tunica propria mucosae of caecum

- Ascaris suum**
Shivacheva, T.; and Terziiski, A., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 187-195
Ascaris suum, mice, plasma cell response of mesenteric lymph nodes
- Ascaris suum**
Sibalic, S.; et al., 1976, Vet. Glasnik, v. 30 (7), 613-617
Ascaris suum, piglets (exper.), cambendazole most effective against migrating larvae when given 14 days after infection
- Ascaris suum** (Goeze, 1782)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Sus scrofa: insular Newfoundland
- Ascaris suum** Goeze, 1782, illus.
Sowinska, A., 1975, Przegł. Zool., v. 19 (2), 270-273
Ascaris suum, structural anomalies of female reproductive system
- Ascaris suum**
Stalheim, O. H. V., 1973, Am. J. Vet. Research, v. 34 (10), 1257-1260
failure to reproduce acute porcine leptospirosis by physiologic stress or concurrent infections (including one with Ascaris suum)
- Ascaris suum, illus.**
Stankov, M.; Movsesijan, M.; and Kovandzic, D., 1975, Acta Parasitol. Iugoslavica, v. 6 (1), 31-36
Ascaris suum, pigs, antibodies form specific antigen-antibody complex; same antibodies are specifically bound to same organs and tissues of A. lumbricoides, sera of humans infected with A. lumbricoides form fluorescent complex with same organs of A. lumbricoides and A. suum
- Ascaris suum**
Stephenson, L. S.; Georgi, J. R.; and Cleveland, D. J., 1977, Cornell Vet., v. 67 (1), 92-102
Ascaris suum, pigs (exper.), worm burden in weanling pigs fed low and high protein diets after infection with known numbers of larvae isolated from rabbits, production of worm burdens of consistent size, potential model for human Ascaris studies
- Ascaris suum**
Stevenson, P., 1977, Vet. Rec., v. 100 (23), 500 [Letter]
Ascaris suum eggs, development during summer months, possible importance in control
- Ascaris suum**
Stevenson, P.; and Jacobs, D. E., 1976, Parasitology, v. 73 (2), i-ii [Abstract]
pigs, milkspot lesion of liver, evidence from serological surveys supports view of Ascaris suum as major factor in aetiology and gives little indication that Toxocara spp. play any significant role: N. Wales; East Anglia
- Ascaris suum**
Stevenson, P.; and Jacobs, D. E., 1977, J. Helminth., v. 51 (2), 149-154
Toxocara canis, T. cati, Ascaris suum, Toxocara leonina, Parascaris equorum, pigs (exper.), in vitro larval precipitate test and indirect fluorescent antibody test using T. canis larvae as antigen, indirect fluorescent antibody test using A. suum larvae as antigen, specificity
- Ascaris suum**
Strejan, G. H.; et al., 1977, Internat. Arch. Allergy and Applied Immunol., v. 54 (6), 502-516
Ascaris suum, rats, influence of type of adjuvant and of carrier priming on induction of IgE and IgG antibodies to dinitrophenyl conjugates
- Ascaris suum**
Strejan, G. H.; and Surlan, D., 1977, Internat. Arch. Allergy and Applied Immunol. v. 54 (6), 487-501
Ascaris suum, rats, function of glutaraldehyde-polymerized antigen in induction of reaginic antibodies
- Ascaris suum**
Stromberg, B. E.; Khoury, P. B.; and Soulsby, E. J. L., 1977, Internat. J. Parasitol., v. 7 (2), 149-151
Ascaris suum, culture from third to fourth stage in chemically defined medium
- Ascaris suum**
Stromberg, B. E.; and Soulsby, E. J. L., 1976, Vet. Parasitol., v. 2 (2), 197-208
Ascaris suum, guinea pigs, capacity of various worm developmental stages to induce protective immune response using various routes of inoculation, antibody titer as assessed by indirect hemagglutination was not correlated with degree of protection
- Ascaris suum**
Stromberg, B. E.; and Soulsby, E. J. L., 1977, Vet. Parasitol., v. 3 (2), 169-175
Ascaris suum, guinea pigs, heterologous resistance induced by Toxocara canis and Ancylostoma caninum but not by Haemonchus contortus, Caenorhabditis briggsae, or Turbatrix aceti
- Ascaris suum**
Stromberg, B. E.; and Soulsby, E. J. L., 1977, Internat. J. Parasitol., v. 7 (4), 287-291
Ascaris suum, guinea pigs, immunization with soluble antigens (extracts or excretory-secretory products of adult and larval stages)
- Ascaris suum**
Tada, T.; et al., 1972, Internat. Arch. Allergy and Applied Immunol., v. 43 (2), 207-216
Ascaris suum, rats, lymphocytosis-promoting factor purified from culture fluid of Bordetella pertussis exerted strong adjuvant effect on production of reaginic antibody

Ascaris suum

Tada, T.; et al., 1975, *Internat. Arch. Allergy and Applied Immunol.*, v. 48 (1), 116-131
 Ascaris suum-sensitized rats, half-lives of two types of rat homocytotropic antibodies in circulation and in skin

Ascaris suum

Tada, T.; and Okumura, K., 1971, *J. Immunol.*, v. 107 (4), 1137-1145

Ascaris suum, anti-hapten homocytotropic antibody (HTA) formation induced in rats with dinitrophenylated Ascaris extracts, suppressive activity of anti-hapten and anti-carrier antibodies on HTA formation, results indicate cooperation of carrier-specific and hapten-specific recognition cells for induction of anti-hapten HTA

Ascaris suum

Tada, T.; Okumura, K.; and Taniguchi, M., 1972, *J. Immunol.*, v. 108 (6), 1535-1541
 Ascaris suum, rats, nature and activities of carrier-specific cells in induction and inhibition of homocytotropic antibody formation, regulator and helper cells may be identical

Ascaris suum

Tada, T.; Okumura, K.; and Taniguchi, M., 1973, *J. Immunol.*, v. 111 (3), 952-961
 Ascaris suum, rats, antigen-specific T cell factor that regulates anti-hapten homocytotropic antibody response

Ascaris suum

Taniguchi, M.; and Tada, T., 1974, *J. Immunol.*, v. 113 (6), 1757-1769
 Ascaris suum, rat, IgT-like molecule for induction of homocytotropic antibody response

Ascaris suum

Tarczyński, S.; Romaniuk, K.; and Szelagiewicz-Czosnek, M., 1972, *Med. Wet.*, v. 28 (4), 217-218
 intestinal nematodes, swine, Suiverm

Ascaris suum

Terziński, A., 1972, *Izvest. Tsentral. Khel'mint. Lab.*, v. 15, 199-207
 Ascaris suum, *A. lumbricoides*, *Toxocara canis*, migration in white mice, comparison

Ascaris suum

Terziński, A.; and Shivacheva, T., 1974, *Izvest. Tsentral. Khel'mint. Lab.*, v. 17, 145-152
 Ascaris suum, mice (immunized or not immunized, challenged per os with antigen), cell reaction in mesenteric lymph nodes

Ascaris suum

Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
 oxbendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction

Ascaris suum

Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
 anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report

Ascaris suum

Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
 fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date

Ascaris suum

Torres, P.; and Barriga, O. O., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 79-85
 Ascaris suum, *A. lumbricoides*, *Toxocara mystax*, *Ascaridia galli*, comparative antigenic analysis by gel double diffusion and immunoelectrophoresis

Ascaris suum, illus.

Trimble, J. J. III; and Thompson, S. A., 1975, *Ztschr. Parasitenk.*, v. 47 (2), 131-144
 Ascaris suum, intestinal epithelium, carbohydrate cytochemistry, microvilli surface, basal lamella, electron microscopy

Ascaris suum, illus.

Trimble, J. J. III; and Thompson, S. A., 1976, *Cell and Tissue Research*, v. 172 (3), 357-363
 Ascaris suum, *Parascaris equorum*, distribution of concanavalin A binding site on nematode intestinal epithelium

Ascaris suum

Tsuji, M.; et al., 1977, *Internat. Arch. Allergy and Applied Immunol.*, v. 55 (1-6), 78-81
 IgE antibodies to Ascaris antigens in serum of person experimentally sensitized with Ascaris suum antigens

Ascaris suum, illus.

Ubelaker, J. E.; and Allison, V. F., 1975, *J. Parasitol.*, v. 61 (5), 802-807
 eggs, fine external morphology, scanning electron microscopy

Ascaris suum

Umaly, R. C.; Oelerich, S.; and Haas, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 422-432
 Schistosoma haematobium, human, with and without other helminthic infections, serodiagnosis, various schistosome antigens plus Ascaris suum and Fasciola hepatica tested in Cercarien-hüllenreaktion, indirect immunofluorescence, indirect haemagglutination, complement fixation, and double gel diffusion tests, evaluation of sensitivity and specificity, attempt to correlate results of serologic tests with some clinical symptoms and with influence of chemotherapy

- Ascaris suum* Goeze, 1782, *illus.*
Valenzuela, G.; et al., 1976, *Bol. Chileno Parasitol.*, v. 31 (3-4), 87
Ascaris suum discovered at necropsy of sheep, fecal examination positive for *Ascaris* sp. eggs, confirmation of possibility of cross infection of gastrointestinal nematodes between sheep and swine: Pelchuquin, Provincia de Valdivia, Chile
- Ascaris suum*
Valenzuela, G.; et al., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (intestino delgado): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Ascaris suum*
Vasilev, I.; and Mutafova, T., 1974, *Izvest. Tsentral. Khelminth. Lab.*, v. 17, 33-42
Ascaris suum, *A. ovis*, no difference in karyotype, successful experimental infection of pigs with egg cultures from lambs, *A. ovis* should be considered synonym of *A. suum*
- Ascaris suum*, *illus.*
Wang, J. S.; and Fujita, J., 1976, *J. Chinese Soc. Vet. Sc.*, v. 2 (1), 35-39
Ascaris suum, *A. lumbricoides*, scanning electron microscopy, especially head, lip, papillae, tail, cuticle, denticles, no morphological difference found
- Ascaris suum*
Wong, H. S. W.; Embil, J. A.; and Ozere, R. L., 1976, *Exper. Parasitol.*, v. 40 (3), 421-426
Ascaris suum, *Toxocara canis*, guinea pigs sensitized with egg extract antigens, dermal reactivity, macrophage migration inhibition test, and lymphocyte transformation using homologous and heterologous antigens
- Ascaris suum*
Zimmermann, I.; and Ulmer, W. T., 1976, *Pneumonologie*, v. 153 (2), 95-103
Ascaris suum extract aerosol used in dogs (exper.) to produce asthma models for study of pharmacodynamics of disodium cromoglycate on asthmatic state
- Ascaroidea* [sp.]
Coggins, J. R., 1975, *J. Elisha Mitchell Scient. Soc.*, v. 91 (2), 73
parasitic fauna, effect of host diet and habitat
Turdus migratorius
Quiscalus quiscula
Agelaius phoeniceus
all from Kellogg Bird Sanctuary, Michigan
- Ascaroidea* [sp.] ova
Faust, B. S.; and Pappas, P. W., 1977, *J. Zoo Animal Med.*, v. 8 (1), 18-23
Rhea americana
Aix galericulata
Cygnus olor
(feces of all): all from Columbus (Ohio) Zoo
- Ascarophis van Beneden*, 1871, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key; synonymy
- Ascarophis* sp.
Fagerholm, H.-P., 1976, *Norwegian J. Zool.*, v. 24 (4), 466 [Abstract]
Finland
- Ascarophis* sp.
Moeller, H., 1976, *J. Marine Biol. Ass. United Kingdom*, v. 56 (3), 781-785
Gadus morhua (intestine): Kiel Fjord (western Baltic Sea)
- Ascarophis species A*, *illus.*
Poinar, G. O., jr.; and Thomas, G. M., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 28-33
description, infective juvenile stages
Callianassa californiensis (hemocoel): Bodega Bay, California
- Ascarophis species B*, *illus.*
Poinar, G. O., jr.; and Thomas, G. M., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 28-33
description, infective juvenile stages
Callianassa californiensis (hemocoel): Bodega Bay, California
- Ascarophis* sp.
Poinar, G. O., jr.; and Thomas, G. M., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 28-33
description, infective juvenile stages
Pagurus samuelis
P. granosimanus
(dorsal wall of abdomen of all): all from Bodega Bay, California
- Ascarophis* sp.
Poinar, G. O., jr.; and Thomas, G. M., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 28-33
Pugettia producta (pyloric stomach, mouthparts, under carapace): Bodega Bay, California; Hood Canal, Washington
Pachycheles pubescens (hepatopancreas): Bodega Bay, California
- Ascarophis argumentosus* nov. sp., *illus.*
Skriabina, E. S., 1966, *Trudy Gel'mint. Lab. Akad. Nauk SSSR*, v. 17, 169-182
Acipenser baeri (stomach mucosa): Yenisei and Lena Rivers
- Ascarophis ayalai* n. sp., *illus.*
Caballero R., G., 1975, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (301), *Zool.* (211), 649-652
Arius liropus (estomac): lagune d'Agua Brava, Nayarit et estuaire de Moroncarit, Sonora, Mexique
- Ascarophis ovotrichuria* Skrjabin, 1924, *illus.*
Skriabina, E. S., 1966, *Trudy Gel'mint. Lab. Akad. Nauk SSSR*, v. 17, 169-182
Acipenser baeri (stomach mucosa): Yenisei River

- Ascarophis pacificus* Zhukov, 1960
Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 80-88
helminth distribution among age groups of *Pleurogrammus azonus*: Peter the Great Bay, Sea of Japan
- Ascarophus pacificus* Zhukov, 1953
Korotaeva, V. D., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 89-96
Icelus spiniger
Hemilepidotus gilberti
Cottiusculus goner
Enophrys diceraus
all from Sea of Japan
- Ascarops van Beneden*, 1873, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascaropsinae
key
Syn.: *Arduenna* Railliet & Henry, 1911
- Ascarops strongylinea*
Peterson, P. M.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1778-1780
Ascarops strongylinea, *Physocephalus sexalatus*, *Hyostrongylus rubidus*, natural incidence, comparison with earlier surveys
hogs (stomachs): Georgia; Wisconsin
- Ascarops strongylinea* (Rudolphi, 1819), larvae, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Syn.: *A. strongylinea* Beneden, 1873
Myotis daubentoni
M. dasycneme
Nyctalus noctula
N. leisleri
Pipistrellus pipistrellus
Vespertilio murinus
Eptesicus serotinus
all from Moldavia
- Ascarops strongylinea* Beneden, 1873
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
as syn. of *A. strongylinea* (Rudolphi, 1819), larvae
- Ascarops strongylinea* (Rudolphi, 1819)
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Rhinolophus hipposideros
Myotis oxygnathus
M. daubentoni
M. mystacinus
Plecotus auritus
Nyctalus noctula
Eptesicus serotinus
all from Moldavia
- Ascarops strongylinea*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Aphodius pusillus
A. distinctus
all from Uzbekistan
- Ascarops strongylinea*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Ascarops strongylinea*
Varma, S.; Malik, P. D.; and Lal, S. S., 1976, Vet. Rec., v. 99 (13), 256
Balantidium coli, consumption of *Ascarops strongylinea* eggs observed during routine fecal examination, pig; possible role in epidemiology of *A. strongylinea*
- Ascarops strongylinea*
Varma, S.; Malik, P. D.; and Lal, S. S., 1977, J. Helminth., v. 51 (2), 134-135
Hybosorus orientalis (exper.)
Oniticellus pallipes (exper.)
Onitis philemon (exper.)
Onthophagus falsus (exper.)
O. mopsus (exper.)
O. ramosellus (exper.)
Trox granulatus (exper.)
Neocleonus sannio (exper.)
Sphenariopsis tristis (exper.)
Hister corax (nat. and exper.)
H. maindronni (nat. and exper.)
Onthophagus catta (nat. and exper.)
O. gazella (nat. and exper.)
O. quadridentatus (nat. and exper.)
all from Hissar, India
- Ascaropsinae Alicata & McIntosh, 1933
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirocercidae
key; key to genera
includes: *Ascarops*; *Streptopharagus*; *Simondsia*; *Leiuris*; *Tejeraia*; *Texicospirura*; *Pygarginema*; *Physocephalus*
- Ashworthius Le Roux*, 1930
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Haemonchinae*
- Ashworthius martinagliai* Ortepp, 1935
Hiregoudar, L. S., 1976, Indian Vet. J., v. 53 (3), 237
Boselaphus tragocamelus (duodenum, small intestine): Gir forest, Gujarat State, India
- Ashworthius tuyenquangi* sp. n., illus.
Drozd, J., 1970, Acta Parasitol. Polon., v. 17 (20-38), 253-258
Muntjac muntjak (abomasum): province Tuyen-Quang, Vietnam
- Aspiculuris*
Sharp, J. W.; and Wescott, R. B., 1976, Lab. Animal Sc., v. 26 (2, pt. I), 222-223
Aspiculuris, *Syphacia*, mice, mebendazole, good results

- Aspicularis dinniki* Schulz, 1927
Tenora, F.; Pfaller, K.; and Murai, E., 1971, *Parasitol. Hungar.*, v. 4, 157-167
Microtus nivalis (Enddarm): Obergurgl; Kuh-tai; Timmelsjoch (Tiroler Zentralalpen)
- Aspicularis pakistanica*
Nama, H. S.; and Parihar, A., 1976, *J. Helminth.*, v. 50 (2), 99-102
Rattus rattus rufescens (intestine): Jodhpur City area, India
- Aspicularis* (*Paraspicularis*) *pakistanica* Akhtar, 1955, illus.
Saxena, A.; and Nama, H. S., 1977, *Geobios*, v. 4 (6), 243-244
description
Rattus rattus (colon): Jodhpur, India
- Aspicularis ratti* Johnson 1969
Singhvi, A.; and Johnson, S., 1977, *J. Parasitol.*, v. 63 (5), 858-860
Aspicularis ratti, *Syphacia muris*, female to male ratio of nematodes in concurrent infections in *Rattus rattus*, no significant correlation with worm burden, possible explanations
- Aspicularis ratti* Johnson, 1969
Sood, M. L.; and Parshad, V. R., 1975, *Riv. Parasitol.*, Roma, v. 36 (2-3), 189-196
infections in *Millardia melitana*, survey of seasonal distribution, possible correlations between host diet and sex and incidence of infection
- Aspicularis tetraptera*
Anya, A. O., 1976, *Internat. J. Parasitol.*, v. 6 (2), 173-177
Aspicularis tetraptera, pattern of the sex attraction phenomenon, origin of attractant secretions
- Aspicularis tetraptera*
Behnke, J. M., 1975, *J. Helminth.*, v. 49 (2), 85-90
Aspicularis tetraptera and *Syphacia obvelata*, survey of levels of infection in wild *Mus musculus*, prevalence of infection of *A. tetraptera* greater in male than in female mice: London Zoo
- Aspicularis tetraptera*
Behnke, J. M., 1976, *J. Helminth.*, v. 50 (3), 197-202
Aspicularis tetraptera in wild *Mus musculus* of different ages, prevalence and level of infection decreased in older animals, either innate or acquired resistance could account for observations
- Aspicularis tetraptera*
Behnke, J. M.; et al., 1976, *Parasitology*, v. 73 (2), xv [Abstract]
Trichinella spiralis expulsion from mice, effect on concurrent helminth infections (*Hymenolepis diminuta*, *H. microstoma*, *Aspicularis tetraptera*)
- Aspicularis tetraptera*
Berenguer Puvia, F. J.; and Gallego Berenguer, J., 1973, *Rev. Iber. Parasitol.*, v. 33 (1), 81-106
Aspicularis tetraptera, *Syphacia obvelata*, *Nippostrongylus brasiliensis*, mice, piperazine and phenothiazine compared with 4 phenothiazine-piperazine derivatives; piperazine: good activity, phenothiazine: low activity against *N. brasiliensis*, low toxicity of both; derivatives: more toxic, no anthelmintic activity
- Aspicularis tetraptera*
Berenguer Puvia, F. J.; and Gallego Berenguer, J., 1973, *Rev. Iber. Parasitol.*, v. 33 (4), 573-598
Aspicularis tetraptera, *Syphacia obvelata*, natural infections in mice, treatment with phenothiazine, piperazine hydrate, piperazine anhydrate; at various doses, calculation of elimination index and statistical analysis
- Aspicularis tetraptera*
Dewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Aspicularis tetraptera*
Farell-Sala, A.; Berenguer-Puvia, F. J.; and Gallego-Berenguer, J., 1974, *Rev. Iber. Parasitol.*, v. 34 (3-4), 331-353
Aspicularis tetraptera, *Syphacia obvelata*, mice, piperazine treatment, comparison of methods of measuring drug activity (deparasitization index and elimination index); relationships of dosage and activity, statistical analysis
- Aspicularis tetraptera*
Gavaghan, A. D.; and Nunn, A. J., 1974, *Pharm. Acta Helveticae*, v. 49 (7-8), 250-258
bis-(6-indazolyloxy) alkanes, no schistosomicidal or other significant anthelmintic properties in laboratory trials with mice
- Aspicularis tetraptera*
McNair, D. M.; and Timmons, E. H., 1977, *Lab. Animal Sc.*, v. 27 (1), 38-42
Syphacia obvelata and *Aspicularis tetraptera*, effects on exploratory behavior of inbred mouse strain (exper.)
- Aspicularis tetraptera*
Martin, O. C., 1975, *Philippine Agric.*, v. 59 (3-4), 114-118
brief description
Mus musculus: Bureau of Research and Laboratories, Alabang, Rizal
- Aspicularis tetraptera*
Owen, D., 1976, *Lab. Animals*, v. 10 (3), 271-278
Mus musculus: Carshalton
- Aspicularis tetraptera*
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 214
Maus
Ratte

- Aspicularis tetraptera*
Rak, H., 1974, Rev. Fac. Vet. Univ. Teheran, v. 29 (4), 21-28
Mus musculus: Iran
- Aspicularis tetraptera*
Taffs, L. F., 1975, J. Helminth., v. 49 (3), 173-177
continuous feed medication with thiabendazole for removal of *Hymenolepis nana*, *Syphacia obvelata*, and *Aspicularis tetraptera* in naturally infected laboratory mice, unexplained deaths among inbred strain C3H/Hef Nimr mice
- Aspicularis tetraptera*
Taffs, L. F., 1976, Vet. Rec., v. 99 (8), 143-144
Hymenolepis nana, *Syphacia obvelata*, *Aspicularis tetraptera*, mice, efficacy of thiabendazole given in diet
- Aspicularis tetraptera*
Wescott, R. B.; Malczewski, A.; and Van Hoosier, G. L., 1976, Lab. Animal Sc., v. 26 (5), 742-745
filter top caging effective method for preventing pinworm infection in pathogen-free mice being introduced into laboratory colony where *Aspicularis tetraptera* and *Syphacia obvelata* were enzootic
- Asymmetracantha* Mawson, 1960
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Mackerrastrongyliinae
- Asymmetricostrongylus* Nagaty, 1932
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Filarinema* Moennig, 1929
- Atractis dactyluris*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Testudo graeca
Testudo haermani
all from Macedonia, Yugoslavia
- Aulonocephalus pennula*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (ceca): Florida
- Austrostrongylus* Chandler, 1924
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Avellaria* Freitas et Lent, 1934
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Viannella* Travassos, 1919
- Aviabronema* Ali, 1961
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Procyrnea* (Chabaud, 1958, subgen.)
- Aviculariella* Wehr, 1931, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key
Syn.: *Alcedospirura* Oshmarin, 1959
- Aviculariella collaricephala* (Oshmarin, 1959), illus.
Aleksseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
redescription
Alcedo attis: Rimsko-Korsakov islands
- Avioserpens* Wehr & Chitwood, 1934
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Dracunculidae
key; synonymy
- Avitellina*
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1974, Indian J. Animal Research, v. 8 (2), 75-78
Haemonchus contortus, other nematodes, experimentally or naturally infected sheep, nitroxynil highly effective, critical testing; in vitro testing against *H. contortus*

- Bancroftian filariasis. See [Wuchereria bancrofti]
- Bancroftinema Johnston & Mawson, 1941
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
"seem to belong to the Physalopteroidea but, as yet, are not well enough known to be classified"
- Barusispirura n. subg., illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of Oxyspirura
tod: O. (B.) rodriguesi Barus, 1968
key
"Yorkeispirura cannot be used and we prefer to follow the classification of Barus (1963) and propose, for the species that was called Yorkeispirura, a new subgenus Barusispirura n. subg."
- Bathmostomum sangeri Cobbold, 1879, illus.
Setasuban, P., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 390-394
Bathmostomum sangeri in Elephas indicus, light and scanning electron microscopy, morphometric data, confirmation that parasite is probably a Strongylidae rather than Ancylostomidae: circus elephants in Brisbane, Australia (primary origin unknown)
- Batrachonema Yuen, 1965
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Batrachostromylus Yuen, 1963
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Baylisascaris Sprent, 1968
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridinae
key
- Baylisascaris spp., illus.
Dade, A. W.; et al., 1977, J. Am. Vet. Med. Ass., v. 171 (9), 885-886
Baylisascaris spp. in Myocaster coypus (brain), complicated by nosematosis, case history, clinical and pathologic findings: city zoo, Grand Rapids, Michigan
- Baylisascaris procyonis
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (thoracic cavity): southeastern United States
- Baylisascaris procyonis
Jacobson, H. A.; et al., 1976, J. Wildlife Dis., v. 12 (3), 357-360
Baylisascaris procyonis, outbreak of cerebrospinal nematodiasis in Sylvilagus floridanus and Marmota monax following establishment of infected Procyon lotor population; laboratory transmission to Sylvilagus floridanus: Center Woods, Virginia
- Baylisascaris tasmaniensis
Munday, B. L.; and Gregory, G. G., 1974, J. Wildlife Dis., v. 10 (3), 241-242
infection of ascarid-free Tasmanian devils by feeding visceral granulomata containing nematode larvae from wombats
Vombatus ursinus: north and northeast Tasmania
Sarcophilus harrisi (nat. and exper.): northeast Tasmania
- Baylisascaris transfuga (Rudolphi, 1819)
Rogers, L. L., 1975, J. Wildlife Dis., v. 11 (2), 189-192
Ursus americanus (intestinal tracts): Minnesota
- Baylisascaris transfuga (Rudolphi, 1819) Sprent, 1968
Vercruyse, J.; et al., 1976, Acta Zool. et Path. Antverpiensia (64), 115-119
Baylisascaris transfuga, polar bears (Thalartos maritimus), mebendazole effective, control program discussed: Zoological Garden, Antwerp
- Befilaria puertoricensis n. sp., illus.
Bain, O.; and Chaniotis, B. N., 1975, Bull. Mus. National Hist. Nat., Paris, 3. s. (281), Zool. (191), 1-5
Anolis cristellus (tissu sous-cutane, sang): Puerto Rico (Caraibes)
- Belanisakis Maplestone, 1932
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Porrocaecum Railliet & Henry, 1912
- Belascaris Leiper, 1907
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Toxocara Stiles, 1905
- Bergheia Drozd, 1965
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Bhalfilaria badamii Bhalerao et Rao, 1944, illus.
Gupta, N. K.; and Acharya, A. K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 69-71
description of male
Arborophila torqueola (heart): Badhan (District Kangra) Himachal Pradesh
- Biacantha Wolfgang, 1954
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae

- Bicaulus sagittatus*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence:
Zehusice enclosure
- Bicaulus sagittatus*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence:
Zehusice enclosure
- Bicaulus sagittatus*
Wolf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere: Trebic District
- Bidigiticauda* Chitwood, 1938
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylineae
- Bigalkenema Ortlepp*, 1963
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Longistrongylus* Le Roux, 1931
- Bigalkenema Ortlepp*, 1963
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus* Le Roux, 1931
- Bigalkenema curvispiculum* Gibbons, 1973
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus curvispiculum* (Gibbons, 1973) n. comb.
- Bigalkenema curvispiculum* Gibbons, 1973
Gibbons, L. M.; and Khalil, L. F., 1976, Trop. Animal Health and Prod., v. 8 (3), 168
sheep
goats
(gut of all): all from Kajiado district, Kenya
- Bigalkenema curvispiculum* Gibbons
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut)
G. granti (abomasum)
all from Kenya
- Bigalkenema namaquensis* Ortlepp, 1963
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus namaquensis* (Ortlepp, 1963) n. comb.
- Bigalkenema neveulemairei* (Gutteres, 1947) Janzen, 1958
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
species inquirenda
- Bigalkenema sabie* (Moennig, 1932) nov. comb.
Ortlepp, R. J., 1963, Onderstepoort J. Vet. Research, v. 30 (1), 119-123 [For author reference see Supplement 16, Part 1]
- Bigalkenema sabie* (Monning, 1932) Ortlepp, 1963
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus sabie* (Monning, 1932) n. comb.
- Binema korsakowi* (Sergiev, 1923) Basir, 1956
Hristovski, N. D., 1972, Acta Parasitol. Iugoslavica, v. 3 (2), 109-115
Gryllotalpa gryllotalpa: Jugoslavija (Skopje); Grcija (Lerin; Solun)
- Binema korsakowi*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Gryllotalpa gryllotalpa: Macedonia, Yugoslavia
- Biogastranema Rohrbacher et Ehrenford*, 1954
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Haemonchinae
- Blatticola blattae* (Graeffe, 1860) Chitwood, 1932
Hristovski, N. D., 1972, Acta Parasitol. Iugoslavica, v. 3 (2), 109-115
Blattella germanica: Jugoslavija (Belgrad)
- Blatticola blattae*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Blattella germanica: Macedonia, Yugoslavia
- Boehmiella Gebauer*, 1932
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Haemonchinae
- Boehmiella wilsoni*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (stomach): southeastern United States
- Boreostrongylus Durette-Desset*, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Nippostrongyliinae
- Boreostrongylus minutus* (Dujardin, 1845)
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Microtus guentheri: Israel
- Boreostrongylus seurati* (Travassos et Darriba, 1929), illus.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
description of 4th stage larvae
Gerbillus pyramidum: Holon, Nahal Rubin, Rishon Le Ziyon, El-Arish-Romani Road and Be'er Sheva, dunes, Israel
G. allenbyi: Caesarea, Hadassim, Herzliyya, Holon, Nahal Rubin and Rishon le Ziyon, Israel
G. dasyurus: Rosh Ha-Niqra, Israel
Meriones tristrami: Mishmar Ha'emeq, Akko, Shave Ziyon and Nahariyya, Israel
M. sacramenti: Israel

- Bostrichodera freitaslenti* (Yeh, 1957) n. comb.
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
Syn.: *Deraiphoronema freitaslenti* Yeh, 1957
- Bostrichodera spiralis* (Molin, 1860) n. comb.
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
Syn.: *Dipetalonema spiralis* (Molin, 1860)
- Bourgelatia diducta*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii krai
- Bourgelatioides traguli* Chandler, 1931, illus.
Chabaud, A. G.; and Krishnasamy, M., 1976, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (388), *Zool.* (270), 721-727
brief description
Tragulus javanicus: Selangor, Subang, Subang Forest Reserve
- Bradyopstrongylus* Price, 1928
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
synonymy
- Breinlia Yorke* et Maplestone, 1926
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
key
includes subgenera: *Breinlia*; *Johnstonema*
- Breinlia*
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
subgen. of *Breinlia*; key
tod: *B. (B.) trichosuri* (Breinl, 1913)
- Breinlia (Johnstonema) sp.* (Spratt et Varughese, 1975)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia sp.*
Singh, M.; and Cheong Chee Hock, 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 516-521
Rattus rattus rumpia
R. mulleri
R. sabanus
all from Malaysia
- Breinlia (Johnstonema) andersoni* (Spratt et Varughese, 1975), illus.
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia (Johnstonema) annulipapillatum* (Johnston et Mawson, 1938) (tod of subgen.)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia (Breinlia) boltoni* (Spratt et Varughese, 1975)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia booliati* sp. n., Singh & Ho, 1972 (in press [nom. nud.])
Singh, M.; Ho, B. C.; and Lim, B. L., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 622
Breinlia booliati, experimental transmission from naturally infected *Rattus sabanus* to laboratory albino rats (exper.) using infective larvae from *Armigeres subalbatus* (exper.) infected vector mosquitoes
- Breinlia (Breinlia) booliati* Singh et Ho, 1973
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia booliati*
Ho, B.-C.; et al., 1976, *Internat. J. Parasitol.*, v. 6 (2), 113-116
Breinlia booliati, transmission to different strains of laboratory rats by various routes of inoculation, female rats more susceptible to infection than males, localization of adult worms in thoracic and abdominal cavities
- Breinlia booliati* Singh and Ho, 1973, illus.
Mak, J. W.; and Lim, B. L., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (1), 22-28
Breinlia booliati, morphometric data, host survey
Rattus sabanus
R. cremoriventer
R. muelleri
all from Sarawak, Indonesia
- Breinlia booliati* Singh and Ho, microfilariae, illus.
Miyata, A.; and Tsukamoto, M., 1975, *Nettai Igaku (Trop. Med.)*, v. 16 (3), 113-130
Rattus mulleri balabagensis: Palawan Island, the Philippines
- Breinlia booliati*
Singh, M.; et al., 1976, *J. Helminth.*, v. 50 (2), 103-110
Breinlia booliati, course of development in *Rattus sabanus* and in laboratory albino rat (both exper.), measurements of developing stages
- Breinlia booliati*
Yap, E.-H.; et al., 1975, *J. Helminth.*, v. 49 (4), 263-269
Breinlia booliati, nocturnal subperiodicity in *Rattus sabanus* vs. irregular fluctuations of microfilariae in albino rats, prepatent period, levels and patterns of microfilaremia during course of infection, cortisone without effect on microfilarial levels when administered near or at postpatency
- Breinlia (Breinlia) dasyuri* (Johnston et Mawson, 1938)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia (Breinlia) dendrolagi* Solomon, 1933
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397

- Breinlia* (*Breinlia*) *dantonensis* [i.e., *dentonensis*] (Spratt et Varughese, 1975)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *mackerrasae* (Walker et Mc Millan, 1974)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *macropi* Wahid, 1962
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *mundayi* (Spratt et Varughese, 1975)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *pseudocheiri* (Spratt et Varughese, 1975)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *rarum* (Johnston et Mawson, 1938)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* *ratti*
Viraboonchai, S.; et al., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (1), 143
[Demonstration]
Rattus jalorensis (nat. and exper.): Narathivas province, Thailand
R. rattus (exper.)
R. surifer "
hamsters "
- Breinlia* (*Breinlia*) *robertsi* (Johnston et Mawson, 1938)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *sergenti* (Mathis et Leger, 1909)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* *sergenti*
Ho, B. C.; and Kan, S. P., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (1), 86 [Demonstration]
Breinlia sergenti, development of filarial larva in fat body of *Aedes togoi* vectors, evidence that fat body provides some nutrition in the development
- Breinlia* *sergenti*
Ho, B. C.; Singh, M.; and Yap, E. H., 1974, *J. Med. Entom.*, v. 11 (5), 622-628
Brugia malayi, *Dirofilaria immitis*, *Breinlia sergenti*, migratory patterns in *Aedes togoi* (exper.), spontaneous escape of infective larvae from mosquitoes denied access to a blood meal; level of host microfilaremia does not seem to influence mortality rate or migratory patterns of infective larvae in mosquitoes
- Breinlia sergenti*, illus.
Ho, B. C.; Yap, E. H.; and Singh, M., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 741-750
Breinlia sergenti, larval development and migration in *Aedes togoi*
- Breinlia sergenti*
Kan, S. P.; and Ho, B. C., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (1), 90-91 [Demonstration]
Breinlia sergenti, mode of nutrition of infective larvae developing within the individual fat cells of *Aedes togoi* vector mosquitoes
- Breinlia sergenti*
Zaman, V., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 462-465
Breinlia sergenti, electron microscopic observations on development within the fat body of *Armigeres subalbatus* (exper.)
- Breinlia sergenti*
Zaman, V.; and Lim, E. P. C., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (4), 579-580
Breinlia sergenti larvae unable to complete development in *Armigeres subalbatus* vector mosquitoes that had been treated with levamisole
- Breinlia* (*Breinlia*) *spelaea* (Leidy, 1875)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *thylogali* (Mackerras, 1954)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Breinlia*) *trichosuri* (Breinl, 1913), illus. (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Breinlia* (*Johnstonema*) *woerli* (Spratt et Varughese, 1975)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Brevigraphidium* Freitas et Mendonca, 1960
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Anoplostrongylineae
- Brevispiculoides* Ortlepp, 1939
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
as syn. of *Viannella* Travassos, 1919
- Brevistriata* Travassos, 1937
Durette-Desset, M. C., 1976, *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation

- Brevistriata Travassos, 1937 (type genus of subfam.)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Brevistriata bergerardi Durette-Desset, 1970
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
as syn. of Brevistriata skrjabini (Schulz et Lubimov, 1932)
- Brevistriata brevispicula Ow Yang, 1967
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
as syn. of Fissicauda brevispicula (Ow Yang, 1967) n. comb.
- Brevistriata callosciuri Supperer et Kutzer, 1963
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
as syn. of Fissicauda callosciuri (Supperer et Kutzer, 1963) n. comb.
- Brevistriata malayensis Ow Yang, 1967
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
as syn. of Calypsostrongylus malayensis (Ow Yang, 1967) n. comb.
- Brevistriata skrjabini (Schulz et Lubimov, 1932), illus.
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
synophe
Syn.: Brevistriata bergerardi Durette-Desset, 1970
- Brevistriata sundasciuri Schmidt, Myers et Kuntz, 1967
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
as syn. of Fissicauda sundasciuri (Schmidt, Myers et Kuntz, 1967)
- Brevistriatinae
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Heligmosomidae
redefinition based on evolution of important characteristics (orientation of ridges, carene development, number and segmentation of crests), good correlation between morphological characters and distribution of species among hosts and geographical regions; key to genera
- Brevistriatinae Durette-Desset, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae
includes: Brevistriata (type genus); Calypsostrongylus; Cordicauda; Fissicauda; Kuala; Metheligmonella; Paraheligmonina; Quentinstongylus; Srivastavanema; Xericola
- Brugia
Laurence, B. R.; and Simpson, M. G., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 12 [Demonstration]
Brugia spp., autoradiography and ultrastructure of filarial larvae development and metabolism in mosquito hosts, uptake of amino and nucleic acids
- Brugia sp.
Mullin, S. W.; et al., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 548-551
Presbytis melalophos
P. obscura
P. cristata
all from Malaysia
- Brugia sp. "Timor filaria"
Partono, F.; et al., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 354-355 [Letter]
man (blood)
Anopheles barbirostris (nat. and exper.)
Aedes togoi (exper.)
Meriones unguiculatus (exper.) (testes, lungs, blood)
Felis domesticus (exper.) (blood)
- Brugia [sp.] "Timor filaria", illus.
Purnomo; et al., 1976, J. Parasitol., v. 62 (6), 881-885
Timor filaria, development in Aedes togoi (exper.), comparison with Brugia malayi, findings support view that Timor filaria is member of Brugia complex
- Brugia-like species, possibly B. beaveri, illus.
Schlesinger, J. J.; Dubois, J. G.; and Beaver, P. C., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 204-207
Brugia-like filarial infection in young soldier, mature male filarial worm removed from retro-auricular lymph node: New Jersey and Maryland army training areas
- Brugia malayi, illus.
Aoki, Y., 1971, Nettare Igaku (Trop. Med.), v. 13 (3), 134-140
Wuchereria bancrofti, Brugia malayi, B. pahangi, demonstration of exsheathing of microfilariae on thick blood film or on agar plate, effects of temperature on exsheathment
- Brugia malayi
Atomosoedjono, S.; van Peenen, P. F. D.; and Putrali, J., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (3), 259
Brugia malayi, continued identification of Anopheles barbirostris as vector of human infection: Central Sulawesi, Indonesia
- Brugia malayi
Barbee, W. C.; Ewert, A.; and Folse, D., 1977, Trop. and Geogr. Med., v. 29 (1), 65-73
mixed Brugia malayi-fungal infections of lymphatic system in cats (exper.) resulted in exacerbation of both infections
- Brugia malayi
Beckett, E. B., 1973, Ann. Trop. Med. and Parasitol., v. 67 (4), 455-466
Brugia pahangi in Aedes aegypti, B. malayi in Aedes aegypti and Mansonia uniformis, flight muscle damage, quantitative aspects

- Brugia malayi**
Beckett, E. B., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (3), 353-357
Brugia pahangi, B. malayi, pattern of flight muscle damage in relation to distribution of developing filarial larvae in *Aedes aegypti* and *Mansonia uniformis*
- Brugia malayi**
Bosworth, W.; and Chernin, E., 1976, *J. Parasitol.*, v. 62 (6), 1013-1014
Brugia malayi, oral transmission to anesthetized jirds, results resemble previous findings with B. pahangi
- Brugia malayi**
Bosworth, W.; and Ewert, A., 1973, *J. Med. Entom.*, v. 10 (2), 217-219
Brugia malayi, superinfection of *Aedes togoi* (exper.), more infective larvae collected after 2 spaced blood meals than after 1 blood meal
- Brugia malayi**
Bosworth, W.; and Ewert, A., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (1), 21-25
Brugia malayi in cats (exper.), added infection of beta haemolytic streptococcus to hind leg regions resulted in elephantiasis, patterns of infections reversible with collateral lymphatic vessels developing
- Brugia malayi**
Bosworth, W.; Ewert, A.; and Flores, A. E., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 255-259
Brugia malayi, cats (exper.), determination of antistreptolysin O titers in filariasis and comparison of titers in combined filariasis and streptococcal involvement, attempted correlation with observable pathology
- Brugia malayi**
Bosworth, W.; Sullivan, J. J.; and Chernin, E., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (5), 700-703
Brugia pahangi, B. malayi, jirds (exper.), viability and oral infectivity of third stage larvae kept in water or recovered from dead mosquitoes, rearward migration of Brugia from mosquito hosts, implications for naturally acquired infections
- Brugia malayi**
Bwangamoi, O.; and Isyagi, A. O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (1), 33-37
filariasis, dogs, incidence survey, morbidity rate per breed, age, and sex of host: Uganda
- Brugia malayi**
Cabrera, B. D., 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 496-504
Brugia malayi, *Mansonia uniformis* and *M. bonnea* incriminated as vector mosquitoes in Bunawan, Agusan, Philippines (nat. and exper.)
- Brugia malayi**
Chow, C. Y., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 404-418
Wuchereria bancrofti, *Brugia malayi*, up-to-date review of confirmed and suspected vectors in WHO Western Pacific Region, available data of natural and experimental infection and infective rates of most vector mosquitoes, remarks on ecology and control
- Brugia malayi**
Dondero, T. J., jr.; Mullin, S. W.; and Bala-singam, S., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 569-575
clinical features of acute Malaysian filariasis as established in 3 human experimental subjects
- Brugia malayi**
Dondero, T. J., jr.; Ramachandran, C. P.; and bin Yusoff, O., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 503-515
Brugia malayi, clinical and pathological findings and statistics of persons infected with Malayan filariasis: West Malaysia
- Brugia malayi**
Dondero, T. J., jr.; and Sivanandam, S., 1973, *Med. J. Malaysia*, v. 27 (4), 306-309
no evidence of human filariasis infection in "spot" blood survey of 4 localities, not public health problem in Kelantan River area of Malaysia
- Brugia malayi**
Dondero, T. J., jr.; Sivanandam, S.; and Lee, C.-C., 1971, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 65 (5), 691-693
Brugia malayi, diurnally sub-periodic microfilarial pattern found in region where nocturnal forms existed: West Malaysia
- Brugia malayi**
ElBihari, S.; and Ewert, A., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (2), 184-186
Brugia malayi, no correlation between prepatent period and either the size of B. malayi inoculum or the number of adult female worms recovered from infected *Meriones unguiculatus* (exper.)
- Brugia malayi**
Ewert, A.; and Bosworth, W., 1975, *J. Parasitol.*, v. 61 (4), 610-614
Brugia malayi, distribution and development in cats reinfected in 1 of 3 ways (on previously infected hind limb only, on contralateral infected limb only, or on both hind limbs simultaneously)
- Brugia malayi**
Ewert, A.; and Emerson, G. A., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (1), 71-73
effect of diethylcarbamazine on third stage Brugia malayi larvae in cats
- Brugia malayi**
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, *Medicine et Malad. Infect.*, v. 3 (8-9), 351-353
diagnostic review of human filariasis

- Brugia malayi**
Grove, D. I.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 220-229
antigen abstract prepared from subperiodic *Brugia malayi* compared with *Dirofilaria immitis* antigen in diagnosis of human filariasis, concluded that antigens from microfilariae, adult worms, and 3rd-stage larvae of *B. malayi* are more sensitive than *D. immitis* antigens and do not have a significantly higher number of false positive reactions: Philippines
- Brugia malayi**
Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, J. Infect. Dis., v. 132 (3), 340-352
algorithms in the diagnosis and management of human forms of filariases
- Brugia malayi**
Guent, J. M., 1972, Maroc Med. (555), v. 52 171-178
man, filariasis presenting as bronchial asthma and rhinosinusitis, case report, diethylcarbamazine: Maroc, previously from South Pacific area
- Brugia malayi**
Guptavanij, P.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 44-50
Brugia malayi, blood survey of humans, domestic animals and rodents for evidence of infections established microfilariae to be nocturnally subperiodic, *Mansonia bonneae* and *M. uniformis* established as mosquito vectors: Banduat Canton, southern Thailand
- Brugia malayi-like microfilariae**
Guptavanij, P.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 44-50
cat (blood): Banduat Canton, southern Thailand
- Brugia malayi**
Guptavanij, P.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 98-99 [Demonstration]
Brugia malayi, transmission of periodic infection from man to cats (exper.), comparative statistics of infections in both subjects
- Brugia malayi**
Guptavanij, P.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 100-101 [Demonstration]
Brugia malayi, prevalence of nocturnal subperiodic infections in areas near Thai-Malaysian border
- Brugia malayi**
Guptavanij, P.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 589-590
Brugia malayi, mosquito survey for vectors of periodic and subperiodic infections
Mansonia indiana
M. uniformis
M. bonneae
M. dives
all from Peninsula of Southern Thailand
- B[rugia] malayi**
Guptavanij, P.; et al., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (1), 42-52
Malayan filariasis, statistics of prevalence and distribution survey for microfilariae and elephantiasis in the provinces of peninsular Thailand
- Brugia malayi, illus.**
Guptavanij, P.; and Harinasuta, C., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 94-95 [Demonstration]
Brugia malayi, statistics of sheathed and unsheathed appearance of periodic and subperiodic microfilariae from southern Thailand
- Brugia malayi**
Guptavanij, P.; and Harinasuta, C., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 578 [Demonstration]
Brugia malayi, *Wuchereria bancrofti*, spontaneous disappearance of microfilariae from 3 human carriers after they were transferred from an endemic area to a non-endemic area in Bangkok
- Brugia malayi**
Guptavanij, P.; and Harinasuta, C., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 185-189
Brugia malayi, humans, comparative study of periodicity of 2 endemic areas of South Thailand (Pattani and Narathiwat Provinces)
- Brugia malayi**
Guptavanij, P.; Harinasuta, C.; and Vutikes, S., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 95-98 [Demonstration]
Brugia malayi, comparative measurements of male and female worms recovered from infected cat (exper.) (lymph glands)
- Brugia malayi**
Guptavanij, P.; Vutikes, S.; and Surathin, K., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 288-289 [Demonstration]
Brugia malayi, subperiodic form, biting cycles of vector mosquitoes (*Mansonia bonneae*, *M. dives*, *M. indiana* and *M. uniformis*): southern Thailand
- Brugia malayi**
Harbut, C. L., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 487-491
Brugia pahangi, *B. malayi*, evaluation of susceptibility and potential uses of white rats and golden hamsters (*Mesocricetus auratus*) as laboratory hosts; successful *B. pahangi* infections established in rats and hamsters; hamsters inoculated with *B. malayi* failed to develop patent infections although adult worms occurred within testes and/or heart and lungs, carcass
- Brugia malayi**
Harinasuta, C.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 103-105 [Demonstration]
Brugia malayi, bionomics of *Mansonia* mosquito vectors in southern Thailand

- B[rugia] malayi
 Harinasuta, C.; and Sucharit, S., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 179-184
 malayan filariasis, decreased incidence of infection in humans and vector mosquitoes assumed to be due to interruption of transmission of filariasis cycle in the mosquitoes resulting from DDT spraying operations of the Malaria Eradication Programme in South Thailand
- Brugia malayi
 Hawking, F., 1974, Progr. Drug Research, v. 18, 173-190
 Wuchereria bancrofti, Brugia malayi, review of present status of human infections in India with emphasis on public health issues and possible control measures
- Brugia malayi
 Ho, B.-C.; Singh, M.; and Yap, E.-H., 1974, J. Med. Entom., v. 11 (5), 622-628
 Brugia malayi, Dirofilaria immitis, Breinlia sergenti, migratory patterns in Aedes togoi (exper.), spontaneous escape of infective larvae from mosquitoes denied access to a blood meal; level of host microfilaremia does not seem to influence mortality rate or migratory patterns of infective larvae in mosquitoes
- Brugia malayi
 Husain, A.; and Kershaw, W. E., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 617-619
 Brugia malayi filariasis, possible effects on ability of Anopheles togoi vector mosquitoes to fly and feed and to transmit infections
- Brugia malayi
 Kapojos, D. S.; et al., 1976, Trop. and Geogr. Med., v. 28 (4), 359-361
 Brugia malayi in humans, prevalence survey: North Celebes, Indonesia
- Brugia malayi, illus.
 Kozek, W. J., 1977, J. Parasitol., v. 63 (6), 952-1000
 Brugia malayi, adults and all larval stages harbor intracytoplasmic bacterial organisms that appear to be transovarially transmitted and show special preference for lateral chords and for germinal tissues of females
- Brugia malayi
 Laurence, B. R., 1970, Med. History, v. 14 (4), 352-363
 Brugia malayi, early history of elephantiasis in India and its association with religious beliefs (St. Thomas' curse)
- Brugia malayi, illus.
 Leong, A. S. Y., 1976, Am. J. Trop. Med. and Hyg., v. 25 (4), 655-656
 Brugia malayi microfilaria discovered in routine cervical smear of healthy Malaysian aborigine: Kuala Lumpur
- Brugia malayi
 Mak, J. W.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 141 [Demonstration]
 Brugia malayi, complete adult male and gravid female recovered from small cyst from bulbar conjunctiva of woman's right eye, blood smear positive for B. malayi, case report: Johore, Malaysia
- Brugia malayi, illus.
 Mak, J. W.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 226-229
 Brugia malayi, infection of human conjunctiva with recovery of adult male and female worms from conjunctival cyst, woman's microfilaremia successfully treated with diethylcarbamazine: near Segamat, Johore, Indonesia
- Brugia malayi, illus.
 Mak, J. W.; and Sivanandam, S., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 21-29
 Brugia malayi, experimental infections in cats to determine whether human ocular lesions are due to site of entry of infective larvae, patent infections produced via ocular instillation, and subconjunctival and subcutaneous inoculation of infective larvae, results show that ocular lesions could be result of site of bite by vector mosquito and thus entry site of infective larvae
- Brugia malayi
 Mazaud, R.; Pelloux, H.; and Ferrus, R., 1974, Medecine Trop., v. 34 (1), 7-24
 Brugia malayi, Fasciola hepatica, humans, cardiovascular complications resulting from cell-mediated immunity
- Brugia malayi
 Muller, R. L.; and Denham, D. A., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 8-9 [Demonstration]
 filarial spp. larvae, field technique for recovery and preservation of infective larvae from their insect vectors, application to studies of transmission dynamics
- Brugia malayi
 Mullin, S. W.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (2), 263
 unsuccessful search for evidence of periodic infection in cats suggests that cats are not important host for periodic strain in Singkir district of central Kedah, Malaysia
- Brugia malayi
 Mullin, S. W.; et al., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 548-551
 Macaca fascicularis: Malaysia
- Brugia malayi
 Mullin, S. W.; and Dondero, T. J., jr., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 91 [Demonstration]
 Brugia malayi, infectivity successfully maintained when stored up to 12 hours at 4°C

- Brugia malayi*
Obiamiwe, B. A., 1977, Ann. Trop. Med. and Parasitol., v. 71 (4), 487-490
sub-periodic *Brugia malayi*, periodic *Wuchereria bancrofti*, influence of the sb gene in *Culex pipiens* vectors on susceptibility to parasite development
- Brugia malayi*, *illus.*
Omar, M. S., 1977, Tropenmed. u. Parasitol., v. 28 (1), 100-108
Wuchereria bancrofti, *Brugia malayi*, *B. pahangi*, *Dirofilaria immitis*, distribution of acid phosphatase activity in larval stages in the mosquito, presence or absence of enzymic activity in the excretory cell complex and amphids of developing larvae useful as adjunctive diagnostic method
- Brugia malayi*
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Brugia malayi*
Partono, F.; et al., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 537-547
Brugia malayi, epidemiologic survey to establish extent of human filariasis (nocturnally periodic), vectors (*Anopheles barbirostris*) and reservoir hosts (none found); trials with hetrazan resulted in marked decrease of microfilariae in carriers: Margolemo, South Sulawesi
- Brugia malayi*
Partono, F.; et al., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 484-486
epidemiologic survey in South Sulawesi for human malaria and filariasis conducted 8 and 22 months after arrival of transmigrants from Central Java showed decreased incidence of malaria and increased filariasis, discussion of associated problems and effects on populace: Indonesia
- Brugia malayi*
Petranyi, G.; Mieth, H.; and Leitner, I., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 328-337
Brugia malayi subperiodic form, statistics of study using *Mastomys natalensis* (exper.) as laboratory host
- Brugia malayi*
Ponnampalam, J. T., 1971, Med. J. Malaya, v. 26 (1), 62-64
review of human chronic endemic filariasis in West Malaysia, case reports of varying presenting symptoms
- Brugia malayi*
Purnomo; et al., 1976, J. Parasitol., v. 62 (6), 881-885
Timor filaria, development in *Aedes togoi* (exper.), comparison with *Brugia malayi*, findings support view that Timor filaria is member of *Brugia* complex
- Brugia malayi*
Purnomo; Dennis, D. T.; and Partono, F., 1977, J. Parasitol., v. 63 (6), 1001-1006
Brugia timori, morphologic description with comparison to *B. malayi*
- Brugia malayi*
Ramachandran, C. P., 1975, Kajian Vet., v. 7 (1), 31-38
helminths, immunization with radiation attenuated vaccines, review
- Brugia malayi*
Ramachandran, C. P.; et al., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 505-515
Brugia malayi, epidemiologic and entomologic survey
Mansonia bonneae
M. dives
M. uniformis
all from Trengganu State, West Malaysia
- Brugia malayi*
Ramachandran, C. P.; et al., 1971, Med. J. Malaya, v. 25 (4), 273-277
survey of workers on rubber plantation for evidence of *Brugia malayi* filariasis: Negeri Sembilan, West Malaysia
- Brugia malayi*
Ramachandran, C. P.; Sivanandam, S.; and Mullin, S. W., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 92-93 [Demonstration]
Brugia malayi, comparison of various methods and routes of inoculation of sub-periodic parasites into cats (exper.)
- Brugia malayi*
Ramachandran, C. P.; and Zaini, M. A., 1968, Med. J. Malaya, v. 22 (3), 198-203
laboratory studies of transmission of sub-periodic *Brugia malayi* by *Aedes togoi*, development of parasites to infective form, relationship of density of microfilariae in vertebrate host to number of mosquitoes infected
- Brugia malayi*
Ramachandran, P.; and Zaini, M. A., 1968, Med. J. Malaya, v. 22 (4), 323-329
survival pattern of *Aedes togoi* mosquitoes infected with sub-periodic *Brugia malayi* studies under laboratory conditions
- Brugia malayi*, *illus.*
Redington, B. C.; et al., 1975, Ann. Trop. Med. and Parasitol., v. 69 (4), 489-492
Brugia pahangi, sub-periodic *B. malayi*, microfilariae, differentiation on basis of specific distribution of acid phosphatase activity, superior to previously used morphologic and biologic methods for differentiating these 2 microfilarial spp.
- Brugia malayi*
Richet, P., 1973, Medecine Afrique Noire, v. 20 (11), 899-920
guidelines for mass therapy in human filarial infections, drugs in current use, review

Brugia malayi

Rodhain, F.; and Rodhain-Rebourg, F., 1976, *Medecine et Malad. Infect.*, v. 6 (3), 108-114
 Wuchereria bancrofti, *Brugia malayi*, review of geographic distribution of human filariasis on the Asian continent

Brugia malayi

Sasa, M.; et al., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (3), 370-376
Brugia malayi, *Wuchereria bancrofti*, field study to determine pattern of microfilarial periodicity conducted in locality where *Brugia malayi* was first described, both parasites of nocturnal periodic form: Bireuen, Indonesia

Brugia malayi

Sasa, M.; and Tanaka, H., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 518-536
Wuchereria bancrofti, *Brugia malayi*, humans, technique for statistical analysis of microfilarial periodicity survey data

Brugia malayi, illus.

Seo, B. S., 1975, *J. National Acad. Sc., Republic Korea, Nat. Sc. s.*, v. 14, 177-229
Brugia malayi, humans, survey, incidence and prevalence rates, mathematical analysis, clinical aspects, diethylcarbamazine citrate, mass treatment program at 3 dosage schedules, histopathology, adenolymphangitis in untreated and treated cases: Cheju Island, Korea

Brugia malayi

Sivanandam, S.; and Dondero, T. J., jr., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (1), 93-94 [Demonstration]
Brugia malayi, comparison of measurements of abnormally long microfilariae recovered from man and cat, probably represents abnormal, possible mutant form of periodic *B. malayi*

Brugia malayi

Sivanandam, S.; and Dondero, T. J., jr., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (3), 413 [Demonstration]
Brugia malayi, rapid Giemsa staining technique for differentiating microfilariae in thick blood films

Brugia malayi

Sivanandam, S.; and Fredericks, H. J., 1966, *Med. J. Malaya*, v. 20 (4), 337-338
 comparison of relative lengths of Innenkorper (central viscus) in differential diagnosis of *Brugia pahangi* and *Brugia malayi*

Brugia malayi, illus.

Sivanandam, S.; and Mak, J. W., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (3), 449-451 [Demonstration]
Brugia malayi, early embryonic forms of subperiodic infection seen in peripheral blood of jirds (exper.), possible abnormal host-parasite reaction

Brugia malayi

Sivanandam, S.; Mak, J. W.; and Lai, P. F., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (1), 68-73
Brugia malayi, experimental infections of *Rattus sabanus* and *R. muelleri* show that both rats can support full parasite development but are poor hosts and probably not important in transmission of subperiodic infection to man in Malaysia

Brugia malayi

Stiller, D.; Sivanandam, S.; and Abu Hassan, R. B., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (1), 132-133 [Demonstration]
Brugia malayi, *B. pahangi*, failure to develop beyond microfilarial stage in nymphs of *Haemaphysalis nadchatrami* fed on infected cats

Brugia malayi

Sucharit, S.; et al., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (4), 549-554
Wuchereria bancrofti, *Brugia* spp., attempted differentiation using morphological characteristics

Brugia malayi (Brug, 1927), illus.

Vincent, A. L.; Ash, L. R.; and Frommes, S. P., 1975, *J. Parasitol.*, v. 61 (3), 499-512
Brugia malayi, adults in pulmonary arteries of male *Meriones unguiculatus*, ultrastructure (cuticle, chords and intrachordal hypodermis, somatic musculature, basal laminae and pseudocoel, alimentary tract, reproductive systems)

Brugia malayi, illus.

Vincent, A. L.; Frommes, S. P.; and Ash, L. R., 1976, *Exper. Parasitol.*, v. 40 (3), 330-354
Brugia, 3 spp. in *Meriones unguiculatus*, pulmonary pathology, results suggest that localization in pulmonary arteries should not be considered an aberrant mode of development

Brugia malayi, illus.

Vincent, A. L.; Portaro, J. K.; and Ash, L. R., 1975, *J. Parasitol.*, v. 61 (3), 567-570
Brugia pahangi, midbody ultrastructure of cuticle, hypodermis, and somatic musculature of adults, compared to *B. malayi*

Brugia malayi

Wenceslao, J. M.; Oban, E.; and Cabrera, B. D., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 552-561
 epidemiologic survey established widespread endemicity of human filariasis in Eastern Samar province, survey for vector mosquitoes *Mansonia uniformis*: Republic of the Philippines

Brugia malayi

Wilson, T.; and Ramachandran, C. P., 1971, *Ann. Trop. Med. and Parasitol.*, v. 65 (4), 525-546
Brugia malayi in man and cats, *B. pahangi* in cats, review of data on infections: long-term observations on microfilaraemia and estimates of efficiency of transmission from mosquito vector to definitive host

- Brugia malayi**
Wong, M. M.; et al., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 265-273
Brugia malayi, summary of longitudinal studies of 28 rhesus monkeys experimentally infected with single, double or multiple inoculations of larvae: clinical aspects, antibody responses, pathology, treatment with diethylcarbamazine, host biochemical changes, microfilaremia
- Brugia malayi**
Wong, M. M.; and Lim, K. C., 1975, J. Parasitol., v. 61 (4), 598
successful experimental infection of Galago crassicaudatus panganiensis with Brugia malayi and B. pahangi, diurnal microfilarial periodicity of B. malayi, advantages for use as laboratory host compared to other known exper. primate hosts
- Brugia pahangi**
Abaru, D. E.; and Denham, D. A., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 367-369
Brugia pahangi, Dirofilaria immitis, no significant differences could be detected between nuclepore and millipore filter systems for collecting microfilariae, controlled laboratory trials
- Brugia pahangi**
Abaru, D. E.; and Denham, D. A., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 333-334
laboratory evaluation of new technique for counting microfilariae in blood, comparison with counting chamber method, possible advantages of new technique in field studies
- Brugia pahangi**
Aoki, Y., 1971, Nettare Igaku (Trop. Med.), v. 13 (3), 134-140
Wuchereria bancrofti, Brugia malayi, B. pahangi, demonstration of exsheathing of microfilariae on thick blood film or on agar plate, effects of temperature on exsheathment
- Brugia pahangi**
Aoki, Y., 1971, Nettare Igaku (Trop. Med.), v. 13 (4), 170-179
Wuchereria bancrofti, Brugia pahangi, exsheathing of microfilariae on thick blood film or on agar plate, effects of temperature, salinity, and pH
- Brugia pahangi**
Beckett, E. B., 1973, Ann. Trop. Med. and Parasitol., v. 67 (4), 455-466
Brugia pahangi in Aedes aegypti, B. malayi in Aedes aegypti and Mansonia uniformis, flight muscle damage, quantitative aspects
- Brugia pahangi**
Beckett, E. B., 1974, Ann. Trop. Med. and Parasitol., v. 68 (3), 353-357
Brugia pahangi, B. malayi, pattern of flight muscle damage in relation to distribution of developing filarial larvae in Aedes aegypti and Mansonia uniformis
- Brugia pahangi**
Benjamin, D. B.; and Soulsby, E. J. L., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 266-272
Brugia pahangi in Mastomys natalensis, homocytotropic and hemagglutinating antibody responses detected using Dirofilaria immitis as antigen
- Brugia pahangi**
Bosworth, W.; and Chernin, E., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 762-764
Brugia pahangi, oral transmission of third stage larvae to dogs (exper.) with penetration of mucosa in or near mouth, adult worms recovered from mandibular, retropharyngeal and axillary lymphatics, head, skin, shoulder-neck area
- Brugia pahangi**
Bosworth, W.; and Chernin, E., 1976, J. Parasitol., v. 62 (4), 654-655
Brugia pahangi, high susceptibility of orally exposed neonatal jirds, survival of larvae within stomach and intestinal tract and subsequent migration to other parts of body
- Brugia pahangi**
Bosworth, W.; Sullivan, J. J.; and Chernin, E., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 700-703
Brugia pahangi, B. malayi, jirds (exper.), viability and oral infectivity of third stage larvae kept in water or recovered from dead mosquitoes, rearward migration of Brugia from mosquito hosts, implications for naturally acquired infections
- Brugia pahangi**
Bryan, J. H.; et al., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 14 [Demonstration]
Brugia pahangi, effects of vector mosquito (Anopheles and Aedes spp.) pharyngeal armature on motility of microfilariae during parasite invasion of haemocoel
- Brugia pahangi**
Denham, D. A.; et al., 1976, J. Helminth., v. 50 (4), 243-250
Brugia pahangi, in vivo (Felis catus, Meriones unguiculatus), in vitro, filaricidal effect of compounds F151, HOE 33258, and their reaction product or mixture V5851 = E
- Brugia pahangi**
Denham, D. A.; McGreevy, P. B.; and Suswillo, R. R., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 13-14 [Demonstration]
hybrid microfilariae obtained by cross-mating Brugia pateri and B. pahangi
- Brugia pahangi**
Denham, D. A.; and Rogers, R., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (2), 173-176
Brugia pahangi in cats (exper.), structural and functional changes in lymphatic system during infection

- Brugia pahangi**
Fox, E. G.; and Schacher, J. F., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 523
Brugia pahangi-infected syngeneic laboratory rat-strains, permits differential studies of cellular response elicited by filarial infection and possible analysis of effect of histocompatibility type on immunopathologic picture of infection in man
- Brugia pahangi** (Buckley and Edeson)
Gaaboub, I. A.; and Busvine, J. R., 1975, Ann. Trop. Med. and Parasitol., v. 69 (4), 493-501
Brugia pahangi, vectorial capacity of Aedes aegypti in relation to its DDT-resistance status and effects of DDT exposure
- Brugia pahangi**
Gaaboub, I. A.; and Busvine, J. R., 1976, Ann. Trop. Med. and Parasitol., v. 70 (3), 355-360
Brugia pahangi, larval treatment of Aedes aegypti with insect development inhibitor, effect on vectorial capacity
- Brugia pahangi**
Gwadz, R. W.; and Chernin, E., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (6), 808-813
Brugia pahangi, exper. oral transmission to Meriones unguiculatus, suggests possibility that cryptic infection by mouth may represent component of epidemiology of filariasis and tropical pulmonary eosinophilia
- Brugia pahangi**
Gwadz, R. W.; and Chernin, E., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (6), 814-818
Brugia pahangi, escape of infective larvae from Aedes aegypti into water, implications for possibility of oral transmission
- Brugia pahangi**
Harbut, C. L., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 487-491
Brugia pahangi, B. malayi, evaluation of susceptibility and potential uses of white rats and golden hamsters (Mesocricetus auratus) as laboratory hosts; successful B. pahangi infections established in rats and hamsters; hamsters inoculated with B. malayi failed to develop patent infections although adult worms occurred within testes and/or heart and lungs, carcass
- Brugia pahangi**
Hedge, E. C.; and Ridley, D. S., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (4), 304-307
evaluation of microfilarial antigen for use with indirect immunofluorescent test in diagnosing human filariasis; best results obtained with sonicated microfilariae of Brugia pahangi with which both cytoplasmic and sheath antigens could be utilized simultaneously
- Brugia pahangi**
Ho, B. C.; and Lavoipierre, M. M. J., 1975, J. Helminth., v. 49 (1), 65-72
Brugia pahangi third-stage larvae, quantitative loss and rate of escape from Aedes togoi allowed to probe on a cat or a mouse at various time intervals or to feed on several hosts until fully engorged, implications for filariasis transmission
- Brugia pahangi**
Jaffe, J. J.; et al., 1977, J. Parasitol., v. 63 (3), 547-553
Brugia pahangi-infected and normal Aedes aegypti, comparative activity and properties of lactate dehydrogenase, xanthine dehydrogenase, and dihydrofolate reductase
- Brugia pahangi**
Javadian, E.; and Macdonald, W. W., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 477-481
Brugia pahangi, Dirofilaria repens, infection as cause of reduced egg-production of Aedes aegypti
- Brugia pahangi**
Lehane, M. J.; and Laurence, B. R., 1977, Parasitology, v. 74 (1), 87-92
Brugia pahangi, susceptible and refractory mosquitoes, flight muscle ultrastructure, changes during parasite melanization in 'resistant' Anopheles labranchiae atroparvus, comparison with lack of change in 'susceptible' Aedes togoi and 'resistant' Aedes aegypti, decrease of glycogen in all 3 spp.
- Brugia pahangi**
McGreevy, P. B.; et al., 1975, J. Helminth., v. 49 (2), 107-113
Brugia pahangi, larval and adult stages transferred from donor cats to jirds immunized against cats survived as well as in normal jirds, infective larvae transferred from mosquitoes to cats immunized against mosquitoes survived as well as in normal cats, host antigenic determinants not detected on parasite surface in substantial amounts using fluorescent antibody techniques
- Brugia pahangi**
McGreevy, P. B.; McClelland, G. A. H.; and Lavoipierre, M. M. J., 1974, Ann. Trop. Med. and Parasitol., v. 68 (1), 97-109
Dirofilaria immitis, susceptibility of Aedes aegypti controlled by sex-linked recessive gene which is distinct from those controlling development of Brugia pahangi or D. corynodes, variation in filarial infectivity as well as in mosquito susceptibility
- Brugia pahangi, illus.**
McLaren, D. J.; et al., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (5-6), 509-514
larval Dirofilaria immitis and Brugia pahangi, gram-negative micro-organisms within hypodermal tissue, possible adverse affect on development
- Brugia pahangi, illus.**
Malone, J. B.; Leininger, J. R.; and Chapman, W. L., jr., 1976, Exper. Parasitol., v. 40 (1), 62-73
Brugia pahangi, histopathology in Mesocricetus auratus, lymph vessels and nodes, testes, epididymis, and spermatic cord, lungs, liver, potential for model of human filariasis

- Brugia pahangi**
Malone, J. B.; Leininger, J. R.; and Thompson, P. E., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (2), 170-171 [Letter]
Brugia pahangi in golden hamster (exper.), pathology, possible use as laboratory model for human infections
- Brugia pahangi**
Muller, R. L.; and Denham, D. A., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 8-9 [Demonstration]
filarial spp. larvae, field technique for recovery and preservation of infective larvae from their insect vectors, application to studies of transmission dynamics
- Brugia pahangi**
Obiamiwe, B. A., 1977, Ann. Trop. Med. and Parasitol., v. 71 (3), 367-370
Brugia pahangi, susceptibility rates of 5 strains of Culex pipiens fatigans (exper.) to infections found to be low with no direct relationship to parasitemia but rather controlled by sex-linked recessive gene
- Brugia pahangi**
Obiamiwe, B. A., 1977, Ann. Trop. Med. and Parasitol., v. 71 (3), 371-374
Brugia pahangi, effects of anticoagulant on early migration of microfilariae from blood meals of susceptible and refractory strains of Culex pipiens (exper.)
- Brugia pahangi**
Obiamiwe, B. A., 1977, Ann. Trop. Med. and Parasitol., v. 71 (3), 375-377
Brugia pahangi, comparative estimates of mortality of parasites in mid-gut and thorax of susceptible and refractory strains of Culex pipiens and Aedes aegypti (both exper.)
- Brugia pahangi**
Obiamiwe, B. A., 1977, Ann. Trop. Med. and Parasitol., v. 71 (4), 491-500
Brugia pahangi, relationships between microfilarial density, number of microfilariae ingested by Culex pipiens mosquito vectors and the proportion of mosquitoes infected with larvae
- Brugia pahangi**
Obiamiwe, B. A.; and Macdonald, W. W., 1971, Ann. Trop. Med. and Parasitol., v. 65 (4), 547-554
Brugia pahangi microfilariae, technique for freezing and storing, subsequent development in mosquito host and final development in vertebrate host
- Brugia pahangi**
Obiamiwe, B. A.; and MacDonald, W. W., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 32-33 [Demonstration]
Brugia pahangi, effect of heparin on migration of microfilariae in Culex pipiens, uptake of microfilariae in C. pipiens and infectivity of C. pipiens in relation to microfilarial densities, evidence of sex-linked gene controlling susceptibility of C. pipiens to infection
- Brugia pahangi, illus.**
Omar, M. S., 1977, Tropenmed. u. Parasitol., v. 28 (1), 100-108
Wuchereria bancrofti, Brugia malayi, B. pahangi, Dirofilaria immitis, distribution of acid phosphatase activity in larval stages in the mosquito, presence or absence of enzymic activity in the excretory cell complex and amphids of developing larvae useful as adjunctive diagnostic method
- Brugia pahangi, illus.**
Omar, M. S.; and Gwadz, R. W., 1974, Tropenmed. u. Parasitol., v. 25 (2), 167-174
Brugia pahangi-infected Aedes aegypti, differential uptake and incorporation of tritiated thymidine and adenine by parasite and host
- Brugia pahangi**
Owen, R. R., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (2), 110-111 [Demonstration]
Brugia pahangi, analysis of pattern of migration of microfilariae over 3 hours following blood meal distinguishes susceptible and refractory members of the Aedes scutellaris complex
- Brugia pahangi (Buckley & Edeson)**
Paige, C. J.; and Craig, G. B., jr., 1975, J. Med. Entom., v. 12 (5), 485-493
Brugia pahangi, variation in susceptibility among 29 strains of Aedes aegypti, possible implications for role of A. aegypti as vector of human filariasis in East Africa
- Brugia pahangi, illus.**
Ponnampalam, J. T., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 511-517
Brugia pahangi, cats (exper.), histopathologic changes resulting from microfilarial infection
- Brugia pahangi**
Ponnudurai, T.; Denham, D. A.; and Rogers, R., 1975, J. Helminth., v. 49 (1), 25-30
Brugia pahangi microfilariae transfused from infected to other cats, wide variation in longevity in normal cats, not detected in immunized recipients after 18 hours
- Brugia pahangi**
Portaro, J. K.; et al., 1977, J. Parasitol., v. 63 (1), 172-174
differential response of Brugia pahangi-sensitized splenocytes to antigens from Brugia pahangi, Dirofilaria immitis, and Trichinella spiralis, possible diagnostic use
- Brugia pahangi**
Portaro, J. K.; Britton, S.; and Ash, L. R., 1976, Exper. Parasitol., v. 40 (3), 438-446
Brugia pahangi in Meriones unguiculatus, depressed reactivity of splenocytes to mitogens phytohemagglutinin and concanavalin A, data suggest this depression is cell-mediated

- Brugia pahangi**
Portaro, J. K.; Kowalski, J. C.; and Ash, L.R., 1977, *Exper. Parasitol.*, v. 43 (1), 122-127
adaptation of *Meriones unguiculatus* lymphocytes to an in vitro microassay system, use in study of cellular immune function with mitogens, mitogen reactivity decreased with jird age and was depressed by infection with *Brugia pahangi*
- Brugia pahangi, illus.**
Redington, B. C.; et al., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (4), 489-492
Brugia pahangi, sub-periodic *B. malayi*, microfilariae, differentiation on basis of specific distribution of acid phosphatase activity, superior to previously used morphologic and biologic methods for differentiating these 2 microfilarial spp.
- Brugia pahangi**
Rew, R. S.; and Saz, H. J., 1977, *J. Parasitol.*, v. 63 (1), 123-129
Litomosoides carinii, *Dipetalonema viteae*, and particularly *Brugia pahangi* microfilariae, oxygen requirements, carbohydrate metabolism, effect of levamisole
- Brugia pahangi**
Richey, T. J.; and Rodriguez, P. H., 1976, *J. Parasitol.*, v. 62 (4), 655-656
Brugia pahangi in *Aedes aegypti*, effects of gamma radiation on development
- Brugia pahangi**
Ridley, D. S.; and Hedge, E. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Brugia pahangi**
Robins, R. A.; et al., 1976, *J. Parasitol.*, v. 62 (1), 171-172
Brugia pahangi-infected and uninfected *Meriones unguiculatus*, development of markers for determining frequency of Fc- and complement-receptor lymphocyte subpopulations, usefulness for immunological evaluation of jird
- Brugia pahangi**
Rodriguez, P. H., 1973, *J. Med. Entom.*, v. 10 (2), 194-197
Brugia pahangi in *Aedes aegypti* (exper.), susceptibility of different strains to infection
- Brugia pahangi (Buckley & Edeson)**
Rodriguez, P. H., 1975, *J. Med. Entom.*, v. 12 (4), 447-450
Brugia pahangi, developing larvae in 3 susceptible strains of *Aedes aegypti*, decrease in susceptibility in high temperature stress
- Brugia pahangi**
Rogers, R.; et al., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (1), 77-84
Brugia pahangi, cats, histological changes in lymph nodes characteristic of cell-mediated and antibody-type immunological responses
- Brugia pahangi**
Rogers, R.; Davis, R.; and Denham, D. A., 1975, *J. Helminth.*, v. 49 (1), 31-32
new technique for visualizing changes in lymphatics caused by filarial worms, injection of Hypaque solution followed by Xero-radiography, use with *Brugia pahangi* in cats
- Brugia pahangi**
Rogers, R.; and Denham, D. A., 1976, *J. Helminth.*, v. 50 (1), 21-28
Brugia pahangi adults, 3rd stage infective larvae, microfilariae, action of levamisole in vitro and in vivo (cats and *Aedes aegypti*)
- Brugia pahangi, illus.**
Rogers, R.; Ellis, D. S.; and Denham, D. A., 1976, *J. Helminth.*, v. 50 (4), 251-257
Brugia pahangi, intrauterine development of embryos from after fertilization to birth with particular emphasis on origin and development of sheath of microfilaria and its possible role in nutrition of developing embryo, comparison with other filarial species
- Brugia pahangi**
Saz, H. J.; and Dunbar, G. A., 1975, *J. Parasitol.*, v. 61 (5), 794-801
stibophen, inhibition of phosphofructokinase and lactate formation, effect on internal hexose phosphate accumulation, inhibition of aldolase, comparison with potassium antimony tartrate (inhibits PFK at higher concentrations but not aldolase)
- Brugia pahangi**
Saz, H. J.; Dunbar, G. A.; and Bueding, E., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (3), 574-575
Brugia pahangi, small doses of 4-isothiocyanato-4'-nitrodiphenylamine given over 2-month period resulted in cure of infected jirds (exper.)
- Brugia pahangi (Buckley and Edeson, 1956)**
Schacher, J. F.; et al., 1973, *Ann. Trop. Med. and Parasitol.*, v. 67 (1), 81-94
Brugia pahangi, dogs, lymphographic studies at various intervals after infection up to 18 months, lymphatic system of male animals less able to accommodate to filarial insult than that of females
- Brugia pahangi**
Sivanandam, S.; and Fredericks, H. J., 1966, *Med. J. Malaya*, v. 20 (4), 337-338
comparison of relative lengths of Innenkorper (central viscus) in differential diagnosis of *Brugia pahangi* and *Brugia malayi*
- Brugia pahangi**
Sivanandam, S.; and Sandosham, A. A., 1965, *Med. J. Malaya*, v. 20 (1), 65
multiple filarial infections in domestic cat: East Pahang, Malaya

- Brugia pahangi*
 Stilller, D.; Sivanandam, S.; and Abu Hassan, R. B., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (1), 132-133 [Demonstration]
Brugia malayi, *B. pahangi*, failure to develop beyond microfilarial stage in nymphs of *Haemaphysalis nadchatrami* fed on infected cats
- Brugia pahangi*
 Strauss, J. M.; and Sivanandam, S., 1966, Med. J. Malaya, v. 20 (4), 336
 mixed *Brugia pahangi* and *Dirofilaria immitis* infections in *Panthera pardus*: National Zoo, Kuala Lumpur (captured in state of Pahang)
- Brugia pahangi*
 Sucharit, S., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 492-497
Brugia pahangi, differences in periodic patterns of microfilariae in experimental infections of rats and cats
- Brugia pahangi*
 Sucharit, S.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 549-554
Wuchereria bancrofti, *Brugia* spp., attempted differentiation using morphological characteristics
- Brugia pahangi*
 Sucharit, S.; and Macdonald, W. W., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 577-578 [Demonstration]
Brugia pahangi females, intra-species variations in measurements of vulva opening
- Brugia pahangi*
 Sucharit, S.; and Macdonald, W. W., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (1), 71-77
Brugia pahangi, rats (exper.), selection program in which rats susceptible to infection were selectively bred, increase in microfilaria rate by F4 generation, resistance to infection in older male rats seemed to be reduced
- Brugia pahangi*
 Sucharit, S.; Riganti, M.; and Harinasuta, C., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 286-287 [Demonstration]
Brugia pahangi, rats and gerbils, splenic granulomas and high eosinophilia in research animals used in selection process for increased susceptibility, possible application to research on tropical pulmonary eosinophilia
- Brugia pahangi*, illus.
 Sucharit, S.; Riganti, M.; and Harinasuta, C., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 223-225
Brugia pahangi in white rats and gerbils being used in laboratory studies to increase susceptibility to infection, evidence of splenic granulomas and high eosinophilia, possible application to research on human pulmonary eosinophilia
- Brugia pahangi*
 Sucharit, S.; and Vutikes, S., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 437-439
Brugia pahangi in cats (exper.), comparison of counting chamber, Sasa and standard smear methods used for counting microfilariae
- Brugia pahangi*
 Sullivan, J. J.; and Chernin, E., 1975, J. Parasitol., v. 61 (3), 572-573
Brugia pahangi, proportion and location of developing larvae recovered from male *Meriones unguiculatus* killed 10 or 11 days after oral or subcutaneous infection, useful for rapid determination of infectivity
- Brugia pahangi*
 Sullivan, J. J.; and Chernin, E., 1976, Internat. J. Parasitol., v. 6 (1), 75-78
Brugia pahangi and *Dipetalonema viteae* compared, differences in oral vs. subcutaneous infection of anaesthetized vs. unanaesthetized adult vs. neonatal *Meriones unguiculatus*
- Brugia pahangi*
 Suswillo, R. R.; and Denham, D. A., 1977, J. Parasitol., v. 63 (3), 591-592
Brugia pahangi, transplanted infections in *Meriones unguiculatus* as a new system of testing drugs for filaricidal activity
- Brugia pahangi*
 Terwedow, H. A., jr.; and Craig, G. B., jr., 1977, Exper. Parasitol., v. 41 (2), 272-282
Waltonella flexicauda, development controlled by genetic factor in *Aedes aegypti*, this factor for susceptibility did not control development of *Brugia pahangi* or *Dirofilaria immitis*
- Brugia pahangi*
 Townson, H., 1974, Ann. Trop. Med. and Parasitol., v. 68 (2), 239-240
Brugia pahangi, development in male *Aedes aegypti* of 'refractory' genotype, females remained refractory even when larvae were introduced by inoculation
- Brugia pahangi*
 Townson, H., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 12-13 [Demonstration]
Brugia pahangi, refined technique for inoculating mosquitoes with microfilariae
- Brugia pahangi*, illus.
 Vincent, A. L.; Frommes, S. P.; and Ash, L. R., 1976, Exper. Parasitol., v. 40 (3), 330-354
Brugia, 3 spp. in *Meriones unguiculatus*, pulmonary pathology, results suggest that localization in pulmonary arteries should not be considered an aberrant mode of development
- Brugia pahangi*, illus.
 Vincent, A. L.; Portaro, J. K.; and Ash, L. R., 1975, J. Parasitol., v. 61 (3), 567-570
Brugia pahangi, midbody ultrastructure of cuticle, hypodermis, and somatic musculature of adults, compared to *B. malayi*

- Brugia pahangi**
Wade, J. O., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 19 [Demonstration]
Brugia pahangi, improved method using membrane feeding unit with Aedes aegypti so that uptake of both erythrocytes and microfilariae remain constant over successive feeding intervals
- Brugia pahangi**
Wade, J. O., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 19 [Demonstration]
effect of heparin and crystamycin in blood food of membrane feeding units on survival and fecundity of vector Aedes aegypti and on migration and development of Brugia pahangi microfilariae
- Brugia pahangi**
Wilson, T.; and Ramachandran, C. P., 1971, Ann. Trop. Med. and Parasitol., v. 65 (4), 525-546
Brugia malayi in man and cats, B. pahangi in cats, review of data on infections: long-term observations on microfilaraemia and estimates of efficiency of transmission from mosquito vector to definitive host
- Brugia pahangi**
Wong, M. M.; and Lim, K. C., 1975, J. Parasitol., v. 61 (4), 598
successful experimental infection of Galago crassicaudatus panganiensis with Brugia malayi and B. pahangi, diurnal microfilarial periodicity of B. malayi, advantages for use as laboratory host compared to other known exper. primate hosts
- Brugia pahangi**
Zielke, E., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (1), 36-44
Culex pipiens fatigans, Aedes aegypti, selection of strains differing in susceptibility to Dirofilaria immitis, demonstrated that inheritance of susceptibility in Aedes aegypti is controlled by sex-linked recessive gene, also found that susceptibility to infection with Brugia pahangi is on same chromosome but in different locus
- Brugia patei**
Abaru, D. E.; and Denham, D. A., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 333-334
laboratory evaluation of new technique for counting microfilariae in blood, comparison with counting chamber method, possible advantages of new technique in field studies
- Brugia patei**
Bwangamoi, O.; and Isyagi, A. O., 1973, Bull. Epizoot. Dis. Africa, v. 21 (1), 33-37
filariasis, dogs, incidence survey, morbidity rate per breed, age, and sex of host: Uganda
- Brugia patei**
Denham, D. A.; McGreevy, P. B.; and Suswillo, R. R., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 13-14 [Demonstration]
hybrid microfilariae obtained by cross-mating Brugia patei and B. pahangi
- Brugia patei**
Laurence, B. R.; and Simpson, M. G., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 12 [Demonstration]
Brugia spp., autoradiography and ultrastructure of filarial larvae development and metabolism in mosquito hosts, uptake of amino and nucleic acids
- Brugia patei**
Vincent, A. L.; Frommes, S. P.; and Ash, L. R., 1976, Exper. Parasitol., v. 40 (3), 330-354
Brugia, 3 spp. in Meriones unguiculatus, pulmonary pathology, results suggest that localization in pulmonary arteries should not be considered an aberrant mode of development
- Brugia timori sp. n., illus.**
Partono, F.; et al., 1977, J. Parasitol., v. 63 (3), 540-546
man (lymphatics, blood)
Anopheles barbirostris
Meriones unguiculatus (exper.) (testes, lungs, heart, major vessels at base of heart)
Aedes togoi (exper.)
all from Ae Bubu, Flores Island, Southeast Indonesia
- Brugia timori**
Partono, F.; et al., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 155-157
Brugia timori, experimental infections in jirds (Meriones unguiculatus) with recovery of adult worms and in cats with development of patent infections but low level microfilaremia; attempted infection of Macaca fascicularis unsuccessful
- Brugia timori**
Partono, F.; and Idris, K. N., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 158-164
factors that influence loss of microfilariae from stained thick blood films, results of laboratory experiments
- Brugia timori, illus.**
Purnomo; Dennis, D. T.; and Partono, F., 1977, J. Parasitol., v. 63 (6), 1001-1006
Brugia timori, morphologic description with comparison to B. malayi
- Brugia tupaiae**
Sucharit, S.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 549-554
Wuchereria bancrofti, Brugia spp., attempted differentiation using morphological characteristics
- Buckleyella Sarwar, 1956**
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Trichostrongylus Looss, 1905
- Buckleyella Rasheed, 1963, illus.**
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key

- Bulbocephalus Rasheed*, 1966, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Proleptinae
key
Syn.: *Cestocephalus Rasheed*, 1966
- Bulbodacnitis alpinus* Mudry and McCart, 1974
Mudry, D. R.; and McCart, P. J., 1976, J. Fish. Research Bd. Canada, v. 33 (2), 271-275
Salvelinus alpinus (pyloric caeca): Alaska; Yukon
- Bulbodacnitis globosa*
Lockard, L. L.; Parsons, R. R.; and Schaplow, B. M., 1975, Great Basin Nat., v. 35 (4), 442-448
Salmo trutta (upper digestive tract), relationship of incidence and intensity of nematode infection to age and sexual maturity of host, higher infection rate in sexually mature trout due to aggressive feeding behavior: streams in southern and western Montana
- Bulbodacnitis truttae* (Fabricius, 1794)
Mudry, D. R.; and Anderson, R. S., 1977, J. Fish Biol., v. 11 (1), 21-33
Salmo gairdneri: Jasper National Park, Canada
Salvelinus namaycush: Jasper National Park, Canada
Prosopium williamsoni: Jasper and Waterton Lakes National Parks, Canada
- Bunostomum*
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Bun[ostomum]*
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Bunostomum*
Chhabra, R. C.; Bali, H. S.; and Toor, L. S., 1976, J. Research, Punjab Agric. Univ., v. 13 (3), 308-311
gastrointestinal strongyles in sheep, critical drug evaluation, thiabendazole (most effective), tetramisole (good results), morantel tartrate (fair results), clioxanide and methyridine (least effective): India
- Bunostomum*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Bunostomum*
Fudalewicz-Niemczyk, W.; et al., 1975, Med. Vet., v. 31 (11), 666-668
sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Bunostomum*
Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Bunostomum*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Bunostomum*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Bunostomum*
Klein Mori, J., 1972, Rev. Med. Vet. y Parasitol., Maracay, v. 24 (1-8), 1971-1972, 207-226
gastrointestinal nematodes, sheep, Neguvon, Ripercol, Thibenzoline, comparison, various management systems, all effective, Ripercol easiest to administer, Neguvon somewhat toxic
- Bunostomum*
Krishna Iyer, P. P.; and Peter C. T., 1975, Kerala J. Vet. Sc., v. 5 (2), 121-123
gastrointestinal nematodes, goats, methyridine
- Bunostomum*
Qadir, A. N. M. A., 1976, Indian Vet. J., v. 53 (11), 855-858
gastrointestinal nematodes, goats and calves, urea for pasture control of free-living stages
- Bunostomum*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Bunostomum*
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Bunostomum*
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment

- Bunostomum spp.**
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Bunostomum spp.**
Düewel, D.; et al., 1974, Prakt. Tierarzt, v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Bunostomum sp.**
Graves, I. L.; Adams, W. H.; and Pyakural, S., 1975, Am. J. Vet. Research, v. 36 (6), 843-846
Babesia bigemina, *Bos grunniens* moved from high to low altitude and challenged with influenza A viruses, hemolytic anemia, possible explanations, death due to *Fasciola hepatica* and *F. gigantica*, incidental finding of *Bunostomum* sp., *Trichuris* sp., *Neoscaris vitulorum*, *Dictyocaulus* sp., coccidia, some reasons for poor survival of yaks at low altitude: Nepal
- Bunostomum spp.**
Niec, R.; et al., 1976, Gac. Vet., Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary anthelmintic treatment that could prevent normal buildup of resistance
- Bunostomum sp.**
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (intestine): Sweden
- Bunostomum sp.**
Schneider, C. R.; et al., 1975, Ann. Trop. Med. and Parasitol., v. 69 (2), 227-232
Bubalus bubalis: Khong Island, Laos
- Bunostomum spp.**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Bunostomum [sp.]**
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Bunostomum phlebotomum**
Anderson, P. J. S.; and Marais, F. S., 1975, J. South African Vet. Ass., v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Bunostomum phlebotomum**
Bryan, R. P., 1976, Austral. Vet. J., v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Bunostomum phlebotomum**
Ciordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Bunostomum phlebotomum**
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Bunostomum phlebotomum**
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Bunostomum phlebotomum**
Randall, R. W.; and Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Bunostomum phlebotomum**
Reinecke, R. K., 1972, Onderstepoort J. Vet. Research, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Bunostomum phlebotomum**
Rowlands, D. ap T.; and Berger, J., 1977, J. South African Vet. Ass., v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Bunostomum phlebotomum**
Schroeder, J.; Honer, M. R.; and Louw, J. P., 1977, J. South African Vet. Ass., v. 48 (2), 95-97
trematodes, nematodes, cattle (exper.), rafoxanide, efficacy of subcutaneous injections against immature larvae and adults

- Bunostomum phlebotomum**
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Bunostomum phlebotomum**
Snijders, A. J.; and Horak, I. G., 1975, J. South African Vet. Ass., v. 46 (3), 265-267
cattle (exper.), rafoxanide, drug trials, efficacy, good results
- Bunostomum phlebotomum**
Tager-Kagan, P., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (4), 317-321
gastro-intestinal nematodes, zebu cattle (1 to 2 years old), cambendazole: Niger
- Bunostomum phlebotomum**
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Bunostomum phlebotomum**
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Bunostomum phlebotomum**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Bunostomum phlebotomum**
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Bunostomum phlebotomum**
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Bunostomum phlebotomum**
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Bunostomum trigenocephalum (Rudolphi, 1808)**
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Bunostomum trigenocephalum**
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Bunostomum trigenocephalum**
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Bunostomum trigenocephalum**
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisolé compared
- Bunostomum trigenocephalum**
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (tenue): Sardegna
- Bunostomum trigenocephalum**
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Bunostomum trigenocephalum**
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Bunostomum trigenocephalum Rudolphi, 1808, illus.**
Guttekova, A.; and Zmoray, I., 1977, Biologia, Bratislava, s. B, Zool. (2), v. 32 (5), 351-361
Bunostomum trigenocephalum, ultrastructure of intestinal cells; probably not haemato-phage, host anemia probably results from injuries to host gut followed by blood loss
- Bunostomum trigenocephalum (Rudolphi, 1808)**
Railliet, 1902
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Bunostomum trigenocephalum**
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia

- Bunostomum trigonocephalum*
 Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
 lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Bunostomum trigonocephalum*
 Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Ovis aries: Chile
- Bunostomum trigonocephalum*
 Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
 ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Bunostomum trigonocephalum* (Rudolphi, 1808)
 Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Bunostomum trigonocephalum*
 Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, Vet. Glasnik, v. 30 (1), 11-17
 sheep, morantel tartarate + diethylcarbazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Bunostomum trigonocephalum*
 Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
 nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Bursaphelenchus lignicolus* n. sp., illus.
 Mamiya, Y.; and Kiyohara, T., 1972, Nematologica, v. 18 (1), 120-124
- Bursaphelenchus sexdentati*
 Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
 nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR

- Caballeroispirura**
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Oxyspirura*
key
- Caenostromylus** Lent et Freitas, 1938
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostromylinae
- Calcaronema trifurcatum** Hovorka et Macko, 1959
Gundlach, J. L., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 83-89
Ciconia nigra (trachea): Lublin Palatinate
- Calypsostrongylus** Schmidt, Myers et Kuntz, 1967
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Calypsostrongylus** Schmidt, Myers et Kuntz, 1967
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Calypsostrongylus longipene** (Chabaud et Durette-Desset, 1967) [i. e. (Durette-Desset et Chabaud, 1967)] n. comb.
Kliks, M.; and Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 693-696
- Calypsostrongylus malayensis** (Ow Yang, 1967) n. comb., illus.
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
synophe
Syn.: *Brevistriata malayensis* Ow Yang, 1967
Aeromys tephromelas: Hill Lagong Kepong, Selangor
Callosciurus caniceps: Bukit Kedap, reserve forestiere, Ulu, Trenggiannu (intestin of all)
- Calypsostrongylus ogdeni** Schmidt, Myers et Kuntz, 1967, illus.
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
synophe
- Calypsostrongylus titasuthi** n. sp., illus.
Kliks, M.; and Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 693-696
Callosciurus flavimanus (intestin): Doi Pui, province de Chiang Mai, Thaïlande
- Camallanidae**
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanoidea
key to genera
includes: *Procamallanus*; *Malayocamallanus*; *Spirocamallanus*; *Paracamallanus*; *Camallanides*; *Oncophora*; *Serpinema*; *Camallanus*
- Camallanides** Baylis & Daubney, 1922, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanidae
key
- Camallanides prashadi** (Baylis et Daubney, 1922) Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, Dobuts. Zasshi, Tokyo, v. 84 (3), 258-261
seasonal differences of infection rate, worm burden
Naja hannah: West Bengal villages
- Camallanina**
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurida
key; key to superfamilies
includes: *Camallanoidea*; *Dracunculoidea*
- Camallanina**
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurida
includes: *Camallanoidea*; *Dracunculoidea*
- Camallanina Chitwood**, 1936
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Spirurida
- Camallanoidea**
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Camallanina
key
- Camallanoidea**
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanina
includes: *Camallanidae*
- Camallanus** Railliet & Henry, 1915, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanidae
key
Syn.: *Zeylanema* Yeh, 1960
- ?**Camallanus** sp.
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Gasterosteus aculeatus: Loch Leven, Scotland
- Camallanus** sp.-larve L3
Vassiliades, G., 1972, Bull. Inst. Fond. Afrique Noire, s. A, v. 34 (3), 529-533
Epiplatys senegalensis: Sangalkam, Senegal
- Camallanus atropusi** n. sp., illus.
Bashirullah, A. K. M.; and Khan, H. R., 1973, Riv. Parassitol., Roma, v. 34 (4), 291-294
Atropus atropus (fore intestine): Bay of Bengal (Cox's Bazar coast)

- Camallanus ctenopomae* Vassiliades et Petter (sous presse)
Vassiliades, G., 1972, Bull. Inst. Fond. Afrique Noire, s. A, v. 34 (3), 529-533
Ctenopoma kingsleyae: Senegal (Sangalkam; Lac Mbaouane)
- Camallanus dimitrovi* n. sp., illus.
Durette-Desset, M.-C.; and Batcharov, G., 1974, Ann. Parasitol., v. 49 (5), 567-576
Dicroglossus occipitalis (intestine grele, ampoule rectale): Lome, Togo
Rana galamensis (intestine grele): Klouto, Togo
- Camallanus dollfusi* n. sp., illus.
Bashirullah, A. K. M.; and Khan, H. R., 1973, Riv. Parassitol., Roma, v. 34 (4), 291-294
Lepturacanthus haumela (intestine): Bay of Bengal (Cox's Bazar coast)
- Camallanus fotedari* Raina et Dhar, 1972, illus.
Campana-Rouget, Y.; et al., 1976, Bull. Acad. Vet. France, v. 49 (2), 205-210
Camallanus fotedari, measurements, life cycle
Lebistes reticulatus (muqueuse anale)
Danio rerio (muqueuse anale)
Cyclops (exper.)
- Camallanus lacustris* (Zoega, 1776)
Dabrowska, Z., 1970, Acta Parasitol. Polon., v. 17 (20-38), 189-193
Esox lucius
Anguilla anguilla
Lota lota
Perca fluviatilis
(intestine of all): all from Vistula River near Warsaw
- Camallanus lacustris* (Zoega, 1776) Railliet et Henry, 1915
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 195-201
Lota l. lota (intestine, pyloric appendices, anterior portion of intestine)
Esox lucius
Silurus glanis
Perca fluviatilis
Anguilla anguilla
all from Poland
- Camallanus lacustris* (Zoega, 1776) Railliet et Henry, 1915
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
Silurus glanis (anterior portions of intestine): river Biebrza basin, Poland
- Camallanus lacustris* (Zoega 1776)
Lee, R. L. G., 1977, Lond. Naturalist (1976) (56), 57-70
Gobio gobio (gut)
Gymnocephalus cernua (gut)
Perca fluviatilis (gut)
Anguilla anguilla (mid-region of the intestine)
all from Serpentine lake, Hyde Park and Kensington Gardens, central London
- Camallanus lacustris*
Perłowska, R., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 27-32
Esox lucius
Perca fluviatilis
all from Zegrzynski Reservoir
- Camallanus mastacembeli* Agrawal, 1967
Sinha, A.; and Sahay, U., 1971, Indian J. Animal Research, v. 5 (2), 67-72
as syn. of *Zeylanema mastacembeli* Sahay et Sinha, 1966
- Camallanus moravecii* n. sp., illus.
Petter, A. J.; Cassone, J.; and France, B. M., [1975], Ann. Parasitol., v. 49 (6), 1974, 677-683
Camallanus moravecii n. sp. causing host mortality, histological study of lesions
Xiphophorus helleri
Mollienisia latipinna
(rectum of all): all from Singapur
- Camallanus oxycephalus* Ward and Magath, 1916
Baker, J. C.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 37-39
Ictalurus punctatus (intestines): island region of western Lake Erie
- Camallanus oxycephalus*
Combs, D. L.; Harley, J. P.; and Williams, J. C., 1977, Tr. Kentucky Acad. Sc., v. 38 (3-4), 128-131
Moxostoma erythrurum (gut): Kentucky River
- Camallanus oxycephalus* Ward and Magath, 1917
Cooper, C. L.; Ashmead, R. R.; and Crites, J. L., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 96
prevalence, comparison with previous years
Perca flavescens (intestine): western Lake Erie
- Camallanus oxycephalus* Ward and Magath 1916
Crites, J. L., 1976, J. Parasitol., v. 62 (1), 166
Camallanus oxycephalus, possible alternative pathway in life cycle, transfer of living encapsulated larvae from forage fish to predaceous fish
Aplodinotus grunniens (mesenteries of posterior intestine near urinary bladder): western Lake Erie
Micropterus dolomieu (exper.)
- Camallanus oxycephalus*
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Ictalurus punctatus
Aplodinotus grunniens
Micropterus salmoides
M. punctulatus
Lepomis gulosus
Pomoxis annularis
all from Eagle Mountain Lake, Texas
- Camallanus oxycephalus*
Harley, J. P., 1977, Tr. Kentucky Acad. Sci., v. 38 (3-4), 136-138
Pomoxis annularis (anus): Lake Wilgreen, Madison County, Kentucky

- Camallanus oxycephalus*
Lockard, L. L.; and Parsons, R. R., 1975, Great Basin Nat., v. 35 (4), 425-426
Polyodon spathula: Yellowstone River near Intake, Montana
- Camallanus patani* Sood
Sinha, A.; and Sahay, U., 1971, Indian J. Animal Research, v. 5 (2), 67-72
"Sood . . . has suggested the name *Camallanus patani* as a substitute name for Sahay and Sinha's worm *Z[eylanema] mastacembeli*. . . . The authors, however, . . . suggest that *Z. mastacembeli* should be recognised as a valid species . . ."
- Camallanus singhi* (Ali, 1956) Yeh, 1960, illus. Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
female redescribed
Trachinotus ovatus (intestine): Fish Harbour, Karachi, Pakistan
- Camallanus trichiuris* n. sp., illus.
Bashirullah, A. K. M.; and Rahman, H., 1972, Riv. Parassitol., Roma, v. 33 (4), 289-292
Lepturacanthus (*Trichiuris*) *savala* (small intestine): Bay of Bengal at Cox's Bazar, Bangladesh
- Camallanus trispinosus* (Leidy, 1851)
Platt, T. R., 1977, Ohio J. Sc., v. 77 (2), 97-98
Chrysemys picta marginata (stomach, small intestine)
Emydoidea blandingii (small intestine)
all from Ottawa National Wildlife Refuge, Ottawa Co., Ohio
- Camallanus truncatus* (Rudolphi, 1814)
Murai, E., 1971, Parasitol. Hungar., v. 4, 145-155
Anguilla anguilla (intestinal tract): Lake Balaton, Hungary
- Camallanus truncatus* Rud., 1814
Ponyi, J.; Biro, P.; and Murai, E., 1972, Parasitol. Hungar., v. 5, 383-408
internal helminths of *Acerina cernua* (intestine), incidence survey, seasonal variations and host growth and development in relationship to parasitic burden: Lake Balaton, Hungary
- Camallanus* (*Neocamallanus*) *vachai* (Wahid and Perveen, 1969) n. comb.
Bashirullah, A. K. M.; and Rahman, H., 1972, Riv. Parassitol., Roma, v. 33 (4), 289-292
- Camelostrongylus Orloff*, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae*
- Camelostrongylus mentulatus*
Beveridge, I.; et al., 1974, Austral. Vet. J., v. 50 (1), 36-37
Camelostrongylus mentulatus, experimentally infected sheep and associated gastritis
Camelus dromedarius (gastrointestinal tract): Melbourne Zoo (Australian origin)
- Camelostrongylus mentulatus*
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antilope cervicapra (abomasum): Texas
- Camerostrongylus* Wolfgang, 1951
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Travassostrongylus Orloff*, 1933
- Capillaria*
Forstner, M. J.; Kopp, H.; and Wiesner, H., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (9), 180-183
nematodes of ruminants, mebendazole, good results: Hellabrunn Zoo, Munich
- Capillaria*
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, Am. J. Vet. Research, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Capillaria*
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Capillaria*
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Capillaria*
Whitehead, R., 1973, Major Problems Path., v. 3, 105-110
human intestinal infection, diagnosis, pathological appearance of mucosal biopsy of gastrointestinal tract
- Capillaria*-like eggs
Ashford, R. W., 1977, Ann. Trop. Med. and Parasitol., v. 71 (1), 29-34
Vulpes vulpes (feces): Wales; Exmoor
- Capillaria* sp.
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (abomasum): vicinity of Nowy Targ, Carpathian Mountains
- Capillaria* sp.
Bishop, C. A.; and Threlfall, W., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 25-35
Somateria mollissima (small intestine): insular Newfoundland and/or southern Labrador

- Capillaria sp.
Bisseru, B.; and Lim, K. C., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 412 [Demonstration]
Corvus splendens protegatus (intestine): Klang, Selangor, Malaysia
- Capillaria sp., illus.
Brown, R. J.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (4), 599-600
Capillaria sp. in Cynopterus brachyotis and Emballonura alectro, liver tissue of bats revealed ova resembling Capillaria hepatica in general morphology, variations discussed: Indonesia
- Capillaria [sp.], illus.
Brown, R. J.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 599-601
Capillaria [sp.] causing hepatitis in Suncus murinus: West Java
- Capillaria sp.
Buck, O. D.; Cooper, C. L.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 233-234
Larus argentatus: Bass Island region of Lake Erie
- Capillaria sp.
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (small intestine): Florida
- Capillaria sp.
Cabrera, B. D., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 50-55
Rattus rattus (feces): Leyte, Philippines
- Capillaria sp.
Christensson, D.; and Rehbinder, C., 1975, Nord. Vet.-Med., v. 27 (10), 496-498
gastrointestinal parasites of reindeer calves, none found in first month of life, increasing infection with age: Norrbotten
- Capillaria [sp.], illus.
Cosgrove, G. E.; and Jared, D. W., 1977, Lab. Animal Sc., v. 27 (4), 526-527
Capillaria [sp.] in Xenopus laevis (skin), thiabendazole, good results with repeated therapy: Oak Ridge National Laboratory, Tennessee
- Capillaria sp. like contorta
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (esophagus, proventriculus): Florida; Louisiana
- Capillaria sp. like mergi
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (small intestine, ceca, cloaca): Florida
- Capillaria spp.
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Capillaria sp.
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 419-423
Ochotona roylei (intestins): Gasainhund, Nepal and Thare Pate a l'est de Gasainhund
- Capillaria sp.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Myotis blythii oxygnathus: Zagorska pec, Novi, Yougoslavie
- Capillaria sp.
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland
- Capillaria sp.
Faust, B. S.; and Pappas, P. W., 1977, J. Zoo Animal Med., v. 8 (1), 18-23
Gallus gallus (feces): Columbus (Ohio) Zoo
- Capillaria sp.
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (small intestine): Florida
- Capillaria sp.
Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, J. Parasitol., v. 61 (3), 547-548
Grus canadensis pratensis (lower small intestine): Florida
- Capillaria sp.
Gogoi, A. R., 1975, Kerala J. Vet. Sc., v. 5 (2), 131-134
fowl: Assam
- Capillaria sp.
Guerrero, C.; Rojas, M.; and Vargas, J., 1974, Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity of 1-tetramisole, significant body weight gain in treated animals
- Capillaria [sp.]
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes (feces): Denmark
- Capillaria spp.
Guterbock, W. M.; and Levine, N. D., 1977, J. Am. Vet. Med. Ass., v. 170 (12), 1411-1413
cats (feces): east central Illinois
- Capillaria spp., illus.
Henriksen, S. A., 1977, Medlemsbl. Danske Dyrlaegeforen., v. 60 (11), 482-485
morphology, life cycle, epidemiology, pathology, diagnosis, control, brief review

- Capillaria sp. 1
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (duodenum; lower small intestine): Florida
- Capillaria sp. 2
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (lower small intestine): Florida
- Capillaria sp., illus.
Huizinga, H. W.; Cosgrove, G. E.; and Sturrock, R. F., 1976, J. Wildlife Dis., v. 12 (1), 93-96
Capillaria sp. in Herpestes auropunctatus, kidney pathology: St. Lucia, West Indies
- Capillaria sp.
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans (gizzard lining): Florida
- Capillaria sp.
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Capillaria sp., illus.
McVicar, A. H., 1977, J. Helminth., v. 51 (1), 11-21
description, may be new species
intestinal helminths of Raja naevus, incidence, intensity, pattern of infection with host age and sex, geographical differences in composition of parasite burden
Raja naevus (spiral intestine): off Plymouth; off Aberdeen
- Capillaria sp.
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Ovis aries: Chile
- Capillaria sp., eggs
Pampiglione, S.; and Ricciardi, M. L., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 89-108
Capillaria sp. eggs, Pygmy (feces), discovered during parasitic prevalence survey, thought to be transit ones from ingested rodents: Cameroun
- Capillaria spp.
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Capillaria [sp.], illus.
Prestwood, A. K.; Nettles, V. F.; and Farrell, R. L., 1977, Am. J. Vet. Research, v. 38 (4), 529-532
pathology
Didelphis marsupialis (lung): Clarke County, Georgia
- Capillaria spec.
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Capillaria sp.
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (intestine): Sweden
- Capillaria-type, illus.
Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Capillaria spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Capillaria sp.
Torres, P.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Capillaria sp.
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 39-42
Rattus norvegicus (intestino delgado): Chile
- Capillaria sp.
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Capillaria (Thominx) aerophila, illus.
Aftandeliants, R.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (1), 64-71
Capillaria aerophila, granulomatous lesion containing worm removed from lung of child presenting with asthmatic symptoms and eosinophilia, treatment with diethylcarbamazine and thiabendazole relieved symptoms, clinical case report, possible transmission through cat-contaminated play area: Teheran, Iran
- Capillaria aerophila Miller & Harkema 1970 [et auct.] (not Creplin, 1839)
Butterworth, E. W.; and Beverley-Burton, M., 1977, Canad. J. Zool., v. 55 (3), 616-619
as syn. of Capillaria didelphis n. sp.

- Capillaria aerophila*
Endres, W. A., 1976, Vet. Med. and Small Animal Clin., v. 71 (11), 1553
Capillaria aerophila, cat, levamisole, good results; dichlorvos ineffective
- Capillaria aerophila*
Gregory, G. G.; and Munday, B. L., 1976, Austral. Vet. J., v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island
- Capillaria aerophila*
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Capillaria aerophila*
Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Capillaria aerophila*
Zeh, J. B.; Stone, W. B.; and Roscoe, D. E., 1977, N. York Fish and Game J., v. 24 (1), 91-93
red fox
gray fox
all from New York
- Capillaria americana*
Anderson, M. M.; and McDaniel, J. S., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
Peromyscus leucopus: eastern North Carolina
- Capillaria americana*
Bienek, G. K.; and Klikoff, L. G., 1974, Am. Midland Naturalist, v. 91 (1), 251-253
Dipodomys microps
- Capillaria americana*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Capillaria anatis* (Shrank, 1790)
Turner, B. C.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 157-169
parasites of *Anas crecca* and *A. discors*, incidence and intensity, age and sex of host
Anas crecca
A. discors
all from eastern Canada
- Capillaria annulata* Molin, 1858
Fabiyy, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (under crop lining): Vom area, Benue Plateau State, Nigeria
- Capillaria annulata*
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: southeastern United States
- Capillaria annulosa* (Dujardin, 1845) Travassos, 1915, illus.
Meszaros, F., 1977, Acta Zool. Acad. Scient. Hungar., v. 23 (1-2), 133-138
description
Cricetus cricetus (small intestine): Hungary
- Capillaria annulosa*
Owen, D., 1976, Lab. Animals, v. 10 (3), 271-278
Rattus norvegicus: Carshalton
- Capillaria anseris*
Cervenka, J.; Zajicek, D.; and Nydl, J., 1975, Veterinarstvi, v. 25 (6), 263-264
helminths, geese, Mebendazole
- Capillaria anseris* Madsen, 1945
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anas querquedula (small intestine): Bulgaria
- Capillaria bilobata*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Capillaria blarinae*
Anderson, M. M.; and McDaniel, J. S., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
Blarina brevicauda: eastern North Carolina
- Capillaria bovis* (Schnyder, 1906)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Capillaria bovis* (Schnyder, 1906), illus.
Goffredo, G.; and Sobrero, R., 1972, Parassitologia, v. 14 (1), 143-148
Dama dama (intestine): foresta Umbra (promontorio garganico, provincia di Foggia)
- Capillaria bovis*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (small intestine): Kentucky
- Capillaria bovis* (Schnyder, 1906) Ransom, 1911
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria

- Capillaria bovis*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence:
Zehusice enclosure
- Capillaria bovis*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Capillaria bovis*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites *Odocoileus virginianus*: Hardy County, West Virginia
- Capillaria bovis* (Schnyder, 1906)
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (small intestine): Cumberland County, New Jersey; Oklahoma
- Capillaria bovis*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Capillaria bovis*
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Capillaria bovis*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Capillaria bovis*
Wolf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Capillaria bovis* (Schnyder, 1906) Ransom, 1911, illus.
Wang, J.-S.; et al., 1975, Bull. Nippon Vet. and Zootech. Coll. (24), 54-56
Syn.: *C. longipes* Ransom, 1911
Capricornis crispus (posterior small intestine): Omachi Alpine Museum, Omachi, Nagano Prefecture, Japan
- Capillaria bursata*
Lesin'sh, K. P.; et al., 1975, Latvijas PSR Zinat. Akad. Vestis (340) (11), 27-30
helminths, chickens, effect of host age and method of rearing on infestation: Latvian SSR
- Capillaria capillaris* (von Linstow, 1882)
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
as syn. of *Capillaria incrassata* (Diesing, 1854)
- Capillaria carbonis* Rudolphi, 1819, illus.
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
description
Chlidonias nigra
Gelochelidon niloticae
Larus genei
all from Azov Sea
- Capillaria catenata* Van Cleave and Mueller, 1932
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Cyprinus carpio (intestine): Sacramento-San Joaquin Delta, California
- Capillaria caudinflata*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (heart, innominate artery): Franklin county, Ohio
- Capillaria caudinflata*
Lesin'sh, K. P.; et al., 1975, Latvijas PSR Zinat. Akad. Vestis (340) (11), 27-30
helminths, chickens, effect of host age and method of rearing on infestation: Latvian SSR
- Capillaria caudinflata*
Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrrix
Tetrao urogallus
all from CSSR
- Capillaria caudinflata*
Torres, P.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Capillaria contorta* (Creplin, 1839)
Andrews, S. E.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 24-28
Corvus brachyrhynchos (esophagus): insular Newfoundland
- Capillaria contorta* (Creplin, 1839)
Buck, O. D.; Cooper, C. L.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 233-234
Larus argentatus: Bass Island region of Lake Erie
- Capillaria contorta*
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (esophagus): Florida
- Capillaria contorta*
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (esophagus): South Bass Island, Ohio

- Capillaria contorta*
 Graber, M.; and Euzeby, J., 1976, *Ann. Parasitol.*, v. 51 (2), 199-205
Anas boschas: Guadeloupe
- Capillaria contorta*
 Kocan, A. A.; and Locke, L. N., 1974, *J. Wildlife Dis.*, v. 10 (1), 8-10
Haliaeetus leucocephalus: Iowa; Missouri; Minnesota; Wisconsin
- Capillaria contorta*
 Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, *Proc. 3. National Wild Turkey Symp.*, 27-32
Meleagris gallopavo silvestris: southeastern United States
- Capillaria contorta*
 Torres, P.; et al., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Capillaria contorta* (Creplin, 1839)
 Turner, B. C.; and Threlfall, W., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 157-169
 parasites of *Anas crecca* and *A. discors*, incidence and intensity, age and sex of host
Anas crecca
A. discors
 (esophageal mucosa of all): all from eastern Canada
- Capillaria didelphis* n. sp., illus.
 Butterworth, E. W.; and Beverley-Burton, M., 1977, *Canad. J. Zool.*, v. 55 (3), 616-619
 Syn.: *Capillaria aerephila* Miller & Harkema 1970 [et auct.] (not Creplin, 1839)
Didelphis virginiana (bronchioles and alveolar spaces of lung): Georgia
- Capillaria erinacei*
 Isenbuegel, E., 1976, *Prakt. Tierarzt*, v. 57, Sondernummer, 21-27
 Eustidil, Telmin, Neguvon, Citarin L, Levamisol
 Igel
- Capillaria exigua* (Dujardin, 1845)
 Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
Crocidura russula: Catalan Pyrenean Mountains
- Capillaria exilis*
 Cooper, C. L.; and Crites, J. L., 1974, *J. Wildlife Dis.*, v. 10 (4), 397-398
Turdus migratorius (intestine): South Bass Island, Ohio
- Capillaria exilis* (Dujardin, 1845)
 Cooper, C. L.; and Crites, J. L., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 233-237
Quiscalus quiscula versicolor (intestine): South Bass Island, Ottawa County, Ohio
- Capillaria exilis*
 Cooper, C. L.; and Crites, J. L., 1976, *J. Parasitol.*, v. 62 (1), 105-110
 similarity index of helminth faunas of 7 passerine bird species, index of association of 10 species of helminths identified as having foci of infection, competition for invertebrate food resources and aggregation into mixed feeding flocks maximizes transmission: South Bass Island, Ottawa County, Ohio
- Capillaria fulicae* (Pavlov and Borgarenko, 1959)
 Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, *Am. Midland Naturalist*, v. 89 (2), 467-473
 comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans
Porphyryla martinica
 (ceca of all): all from Florida
- Capillaria garfiai* n. sp., illus.
 Gallego, J.; and Mas-Coma, S., [1976], *Vie et Milieu*, s. C, *Biol. Terr.*, v. 25 (2), 1975, 237-248
 pathology
Sus scrofa (epitelio poliestratificado lateral de la lengua): Valle de Aran, provincia de Lerida (Espana)
- Capillaria gastrica* Baylis, 1926
 Beveridge, I.; and Barker, I. K., 1975, *J. Helminth.*, v. 49 (4), 211-227
 comparative measurements of *C. rickardi* sp. n. with *C. gastrica*
- Capillaria gastrica*
 Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
Rattus norvegicus (estomac): Tunisia, Tunisia
- Capillaria helenae* Layman, 1930
 Korotaeva, V. D., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 89-96
Icelus spiniger (intestine, pyloric caeca): Sea of Japan
- Capillaria hepatica*
 Anderson, M. M.; and McDaniel, J. S., 1975, *J. Elisha Mitchell Scient. Soc.*, v. 91 (2), 73
Blarina brevicauda
Peromyscus leucopus
 all from eastern North Carolina
- Capillaria hepatica*
 Brown, R. J.; et al., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (4), 599-600
Capillaria sp. in *Cynopterus brachyotis* and *Emballonura alectro*, liver tissue of bats revealed ova resembling *Capillaria hepatica* in general morphology, variations discussed: Indonesia
- Capillaria hepatica* (Bancroft 1893)
 Farhang-Azad, A., 1977, *J. Parasitol.*, v. 63 (1), 117-122
Capillaria hepatica in *Rattus norvegicus*, prevalence, intensity, aspects of rat population ecology and environmental factors which relate to parasite transmission and maintenance: Baltimore Zoo, Maryland

- Capillaria hepatica* (Bancroft 1893)
Farhang-Azad, A., 1977, *J. Parasitol.*, v. 63 (4), 701-706
Capillaria hepatica, egg-releasing mechanisms and transmission ecology among Norway rat populations, cannibalism serves as primary egg-releasing mechanism with secondary role played by predators and normal death and decomposition, minor role of carrion insects and soil invertebrates: Baltimore Zoo
- Capillaria hepatica*, *illus.*
Grigonis, G. J., jr.; and Solomon, G. B., 1976, *Exper. Parasitol.*, v. 40 (2), 286-297
Capillaria hepatica, egg shell in situ following freeze-dry fixation of infected mouse liver, fine structure, histochemistry
- Capillaria hepatica*
Hays, B. D., 1977, *J. Environ. Health*, v. 39 (6), 424-426
transmission of protozoan cysts and metazoan eggs from land application of sewage effluents and sludges, brief literature review, parasite survey from selected Pittsburgh area sludges, control measures
- Capillaria hepatica*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Capillaria hepatica*
Kinsella, J. M., 1974, *Am. Mus. Novitates* (2540), 1-12
Sigmodon hispidus (liver): Florida
- Capillaria hepatica*
Kutzer, E.; and Frey, H., 1976, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 89 (24), 480-483
Lepus europaeus: Austria
- Capillaria hepatica*
Laemmler, G.; et al., 1974, *Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol.* (Vienna, Austria, Sept. 18-20, 1973), 327-341
Capillaria hepatica in *Mastomys natalensis* as a model system, review: infection and early development, egg production, host reactions (pathological and pathophysiological changes, serologic response), implications for human infections
- Capillaria hepatica*
Laemmler, G.; and Gruener, D., 1976, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 89 (11), 222-225; (12), 229-233
Capillaria hepatica in *Mastomys natalensis*, comparative activity of 24 anthelmintics, possible model screening system
- Capillaria hepatica*
Meszaros, J.; and Varga, I., 1976, *Acta Vet., Budapest*, v. 26 (3), 377-383
Capillaria hepatica in *Cavia porcellus* and mice (both exper.), fewer liver lesions in *C. porcellus* than in mice, eventual disintegration of parasites in *C. porcellus*, apparently not a very susceptible host
- Capillaria hepatica*, *illus.*
Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
experimental development in domestic cat unsuccessful
Rattus norvegicus (foie): Tunis, Tunisia
- Capillaria hepatica*
Owen, D., 1976, *Lab. Animals*, v. 10 (3), 271-278
Rattus norvegicus: Carshalton
- Capillaria hepatica*
Reynolds, W. A.; and Gavutis, G., jr., 1975, *J. Wildlife Dis.*, v. 11 (1), 13
Marmota monax (liver): Great Swamp National Wildlife Refuge
- Capillaria hepatica*
Silveira, D.; et al., 1975, *Arq. Escola Vet. Univ. Fed. Minas Gerais*, v. 27 (2), 231-234
pathology
dog (liver): State of Minas Gerais, Brazil
- Capillaria hepatica* (Bancroft, 1893)
Singh, M.; and Cheong Chee Hock, 1971, *South-east Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 516-521
Rattus rattus argentiventer
R. r. jarak
R. r. rumpia
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
R. whiteheadi
all from Malaysia
- Capillaria hepatica* (Bancroft, 1893), *illus.*
Slais, J.; Sterva, J.; and Zikova, A., 1973, *Pizen. Lek. Sborn.* (39), 231-234
Capillaria hepatica manifesting as solitary hepatic granuloma in humans, pathologic findings
- Capillaria hepatica*, *illus.*
Solomon, G. B.; and Grigonis, G. J., jr., 1976, *Exper. Parasitol.*, v. 40 (2), 298-307
Capillaria hepatica, changes in egg shell structure following collection of eggs by physical methods or after passage through mouse gastrointestinal tract, relationship to origin and release of antigens contributing to immunological response during granuloma formation; hypothesis concerning exper. egg granuloma formation, maintenance of homeostasis of eggs in situ, and possible modes of action which trigger development
- Capillaria hepatica*
Stokes, R., 1973, *Austral. Vet. J.*, v. 49 (2), 109
dog (liver): Brisbane
- Capillaria hepatica*, *illus.*
Taniguchi, M.; et al., 1977, *Bull. Coll. Agric. and Vet. Med., Nihon Univ.* (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo

- Capillaria hepatica* (Bancroft, 1893) Travassos, 1915
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 39-42
Rattus norvegicus (higado): Chile
- Capillaria hepatica*
Vollerthun, R.; et al., 1976, *Zentralbl. Vet.-Med., Beihefte* (25), 161-163
Capillaria hepatica, pathophysiology, immunology, *Mastomys natalensis*, rabbits; animal models for human infection studies
- Capillaria hepatica* (Bancroft, 1893), illus.
Wiroren, W., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (1), 136-138
Rattus rattus diardi (liver surface): Bogar, West Java, Indonesia
- Capillaria hepatica* (Bancroft, 1893), illus.
Wobeser, G.; and Rock, T. W., 1973, *J. Wildlife Dis.*, v. 9 (3), 225-226
Capillaria hepatica, diagnosis in *Canis latrans* on basis of histopathology, no recognizable portions of adult parasites were seen: southern Saskatchewan
- Capillaria hepatica* Bancroft, 1893, illus.
Wright, K. A., 1976, *J. Nematol.*, v. 8 (1), 92-95
Capillaria hepatica, somatic centrioles, morphology, rare occurrence, incomplete structure, relation to eutely
- Capillaria hepatica*, illus.
Wright, K. A., 1976, *Organ. Nematodes* (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Capillaria hepatica*
Zahner, H.; et al., 1976, *Ztschr. Parasitenk.*, v. 49 (1), 41-61
Capillaria hepatica in *Mastomys natalensis* (liver) (exper.), development, prepatency, egg production (duration and dynamics dependent upon infective dose), necroses of liver as related to infection rate, liver and spleen weight increases
- Capillaria incrasata* (Diesing, 1854), illus.
Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
synonymy
Sorex araneus (vejiga urinaria): Catalan Pyrenean Mountains
- Capillaria italica* Ricci, 1949
Skvortsov, V. G., 1973, *Parazity Zhivot. i Rasten.*, *Akad. Nauk Moldavsk. SSR* (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Myotis daubentoni
M. bechsteini
all from Moldavia
- Capillaria kutori* Ruchljadewa, 1946
Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
synonymy
Sorex araneus
S. minutus
(tracto gastrointestinal of all): all from Catalan Pyrenean Mountains
- Capillaria linstowi* Travassos, 1914
Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
as syn. of *Capillaria incrasata* (Diesing, 1854)
- Capillaria longipes*
Boag, B.; and Thomas, R. J., 1977, *Research Vet. Sc.*, v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Capillaria longipes* Ransom, 1911
Wang, J.-S.; et al., 1975, *Bull. Nippon Vet. and Zotech. Coll.* (24), 54-56
as syn. of *C. bovis* (Schnyder, 1906)
Ransom, 1911
- Capillaria mergi* Madsen, 1945, illus.
Daia, G. G., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 49-53
redescription
Mergus merganser: Lena estuary
Bucephala clangula: Kamchatka
- Capillaria mergi*
Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Capillaria mucronata* (Molin, 1858) Travassos, 1915
Shakhmatova, V. I., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 277-289
Martes martes
Mustela lutreola
Mustela putorius
Mustela erminea
all from Karelia
- Capillaria nyrocinorum* (Madsen, 1945)
Bishop, C. A.; and Threlfall, W., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 25-35
Somateria mollissima (gizzard, cloaca, oviducts): insular Newfoundland and/or southern Labrador
- Capillaria obsignata*
Clarkson, M. J.; and Esfandiari, A., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 15-16 [Demonstration]
Capillaria obsignata in fowls (exper.), dynamics of infections, self cure and host immune response

- Capillaria obsignata*
Colglazier, M. L., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 60-61
Ascaridia dissimilis, Heterakis gallinarum, and *Capillaria obsignata*, turkeys, levamisole administered in drinking water effective as anthelmintic
- Capillaria obsignata*
Cruthers, L. R.; al-Khateeb, G. H.; and Hansen, M. F., 1975, Proc. Oklahoma Acad. Sc., v. 55, 119-121
Ascaridia galli, Heterakis gallinarum, *Capillaria obsignata*, chickens, levamisole in drinking water
- Capillaria obsignata*
Lesin'sh, K. P.; et al., 1975, Latvijas PSR Zinat. Akad. Vestis (340) (11), 27-30
helminths, chickens, effect of host age and method of rearing on infestation: Latvian SSR
- Capillaria obsignata*
Radhakrishnan, C. V.; and Ebrahimina, A., 1975, J. Vet. Fac. Univ. Tehran, v. 30 (4), 1-4
chickens (small intestine): Darab, Fars Province, Iran
- Capillaria obsignata*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Capillaria obsignata*
Torres, P.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Capillaria obsignata* (Madsen, 1945)
Ziegler, K., 1975, Acta Vet. Brno, v. 44 (1-2), 115-122
Capillaria obsignata, vaccination, chickens, X-irradiated embryonated eggs, safe, effective
- Capillaria oesophagicola* Soltys, 1952
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
Sorex araneus (esofago): Catalan Pyrenean Mountains
- Capillaria ovopunctatum*
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (intestine): South Bass Island, Ohio
- Capillaria ovopunctatum*
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 399-403
survey, helminths of red-winged blackbirds including a check list of previous findings
Agelaius phoeniceus (intestine): South Bass Island, Ohio
- Capillaria ovopunctatum* (von Linstow, 1873)
Cooper, C. L.; and Crites, J. L., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (intestine): South Bass Island, Ottawa County, Ohio
- Capillaria ovopunctatum*
Cooper, C. L.; and Crites, J. L., 1976, J. Parasitol., v. 62 (1), 105-110
similarity index of helminth faunas of 7 passerine bird species, index of association of 10 species of helminths identified as having foci of infection, competition for invertebrate food resources and aggregation into mixed feeding flocks maximizes transmission: South Bass Island, Ottawa County, Ohio
- Capillaria ovopunctatum*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (intestine): Ottawa county, Ohio
- Capillaria papuensis* sp. nov., illus.
Copland, J. W., 1975, J. Helminth., v. 49 (3), 187-190
prepatent period; pathological changes in infected epithelium
Sus scrofa papuensis (tongue): Kasena and Nupa Village, both villages within 20 miles of Goroka, Eastern Highlands of Papua New Guinea
- Capillaria perforans*, illus.
Itagaki, H.; et al., 1975, Bull. Azabu Vet. Coll. (30), 57-62
Capillaria perforans eggs, development, optimum conditions for culture
- Capillaria petruschewski* Shulman, illus.
Reichenbach-Klinke, H. H., 1975, Fisch u. Umwelt (1), 113-121
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Capillaria phalacrocoraxi* sp. nov., illus.
Borgarenko, L. F., 1975, Dokl. Akad. Nauk Tadzhiksk. SSR, v. 18 (10), 63-66
Phalacrocorax pygmaeus (bursa of Fabricius, cloaca): Tigrovaia Balka zapovednik, Piandzhskii region, Tadzhikistan
- Capillaria philippinensis*, illus.
Pradatsundarasar, A.; et al., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (1), 131-134
Capillaria philippinensis, intestinal capillariasis in 18-month-old child (feces), clinical case report, effective therapy with long-term administration of thiabendazole: Bangpree District of Samut Prakarn Province, Thailand
- Capillaria philippinensis*
Singson, C. N.; Banzon, T. C.; and Cross, J. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 932-934
Capillaria philippinensis, human clinical trials using mebendazole for intestinal capillariasis, additionally effective against Ascaris lumbricoides, Trichuris trichiura and hookworm: Philippines

- Capillaria philippinensis*, *illus.*
Sun, S. C.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (4), 524-533
Capillaria philippinensis, ultrastructural survey of pathologic changes in intestinal infections in humans and *Meriones unguiculatus* (exper.)
- Capillaria plica*
Anderson, M. M.; and McDaniel, J. S., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
Blarina brevicauda: eastern North Carolina
- Capillaria plica*
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Capillaria podicipitis* Yamaguti, 1941, *illus.*
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
redescription
Podiceps griseigena (caecum, small intestine): Rimsko-Korsakov islands
- Capillaria procyonis* sp. n., *illus.*
Pence, D. B., 1975, J. Parasitol., v. 61 (5), 815-818
Procyon lotor (stratified squamous epithelial lining of esophagus): LaPlace, St. John the Baptist Parish, Louisiana
- Capillaria putorii* (Rudolphi, 1819) Travassos, 1915
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela lutreola
Mustela putorius
Mustela erminea
Meles meles
Lutra lutra
all from Karelia
- ? *Capillaria rauschi* Read, 1949
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
as syn. of *Capillaria kutori* Ruchljadewa, 1946
- Capillaria recurvirostrae* Mawson, 1968
Ahern, W. B.; and Schmidt, G. D., 1976, Parasitology, v. 73 (3), 381-398
Recurvirostra americana: Kansas and/or Colorado
- Capillaria resecta* (Dujardin, 1843)
Andrews, S. E.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 24-28
Corvus brachyrhynchos (duodenum and small intestine): insular Newfoundland
- Capillaria rickardi* sp. n., *illus.*
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
pathology, comparative measurements with *C. gastrica*
Antechinus stuartii (stomach, intestine): Powelltown, Dartmouth, and Healesville, Victoria
- Capillaria rickardi* Beveridge & Barker
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (stomach, intestine): Powelltown, Victoria
- Capillaria romana* Ricci, 1949
Skvortsov, V. G., 1973, Parazyty Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Myotis dasycneme
M. daubentoni
M. mystacinus
all from Moldavia
- Capillaria salvelini* Polyanski, 1952
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Salmo trutta (intestine)
Perca fluviatilis
Esox lucius (intestine)
all from Loch Leven, Scotland
- Capillaria sunci* Chen, 1937
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
as syn. of *Capillaria incrassata* (Diesing, 1854)
- Capillaria tridens*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (intestine): Franklin county, Ohio
- Capillaria tridens* (Dujardin 1845) Travassos 1915
Davidson, W. R.; Kellogg, F. E.; and Doster, G. L., 1975, J. Parasitol., v. 61 (6), 1115
Meleagris gallopavo (small intestines): Clarke County, Alabama; Morehouse Parish, Louisiana; Berkeley County, South Carolina; Pocahontas County, West Virginia
- Capillaria vanelli*, *illus.*
Orlandi, M.; and Colombani, B., 1975, Ann. Fac. Med. Vet. Pisa, v. 27, 1974, 113-128
histopathology
Vanellus vanellus (gizzard): Tombolo (Pisa)
- Capillaria ventricola* Soltys, 1952
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
as syn. of *Capillaria kutori* Ruchljadewa, 1946
- Capillaria wioletti* Ruchljadewa, 1950
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Arvicola terrestris (stomach): Karelia

- Capillariidae gen. sp.
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Charadrius hiaticula
Tringa glareola
Numenius ph. phaeopus
Calidris temminckii
Phalaropus lobatus
Xenus cinereus
all from lower Yenisei [and/or] Keta lake
- Capillospirura Skrjabin, 1924
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Ascarophis van Beneden, 1871
- Capreocaulus capreoli
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (lungs): Czechoslovakia
- Capreocaulus capreoli
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of Ovis ammon musimon and Capreolus capreolus, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (lung): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Capreocaulus capreoli
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Capreocaulus capreoli (Stroh et Schmidt, 1938)
Schulz et Kadenazii, 1948
Maklakova, L. P., 1975, Trudy Gel'mint Lab., Akad. Nauk SSSR, v. 25, 102-106
Capreocaulus capreoli, infection in relation to age of mollusc hosts
Eulota fruticum (nat. and exper.): Medynsk region, Kaluzhsk oblast
Succinea putris (exper.)
- Capreologia Schulz, Andreeva et Kadenazii, 1953
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Capsularia Zeder, 1800, nec Modeer, 1793
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Anisakis Dujardin, 1845
- Capsulariinae Johnston & Mawson, 1943
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Anisakinae Railliet & Henry, 1912
- Cardiofilaria inornata (Anderson, 1956)
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coerulescens (air sacs):
Florida
- Cardiofilaria nuda (Hamann, 1940)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Gallinago media: lower Yenisei
- Cathematella Yamaguti, 1961
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Torquatoides (Williams, 1929, subgen.) Inglis, 1965
- Cathematella Yamaguti (1961)
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
as syn. of Torquatoides Williams, 1929
- Cephalobata Paramonov, 1956
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Rhabditida
- Cephalobus persegnis Bastian, 1865, illus.
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972, Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera beetles, possible importance in biological control
Scarabaeus sacer: Abu-Rawash, Cairo, Egypt
- Cephaluris Akhtar, 1947
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
revision, key to species, male, includes: Cephaluris hashmi Akhtar, 1956; C. alaskensis Akhtar, 1958; C. andrejevi Shul'ts, 1948; C. chabaudi Inglis, 1959; C. coloradensis Olsen, 1949; C. ochotonae Akhtar, 1947
- Cephaluris alaskensis Akhtar, 1958, illus.
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
redescription, measurements, key
Ochotona princeps: Gunnison County, Colorado; Rocky Mountains, Sheep River and east of Kananaskis River, southwest Alberta
O. collaris: Ogilvie Mountains, Yukon
- Cephaluris alaskensis
Seese, F. M., 1973, Am. Midland Naturalist, v. 89 (2), 257-265
key
- Cephaluris andrejevi Shul'ts, 1948
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
measurements, key
Ochotona rutila: Kazakhstan, USSR
O. hyperborea yesoensis: Daisetsusan National Park, Japan
- Cephaluris collaris Akhtar, 1958
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
as syn. of Cephaluris coloradensis Olsen, 1949

- Cephaluris collaris*
Seese, F. M., 1973, Am. Midland Naturalist, v. 89 (2), 257-265
key
- Cephaluris coloradensis* Olsen
Grundmann, A. W.; and Lombardi, P. S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 39-46
Ochotona princeps cinnemomea: Tushar Mountains, Utah
O. p. wasatchensis: Wasatch Mountains, Utah
O. p. uinta: Uinta Mountains, Utah
O. p. lasalensis: La Sal Mountains, Utah
O. p. fuscipes: Markagunt Plateau, Utah
O. p. barnsei: Fish Lake Mountains, Utah
O. p. nevadensis: Ruby Mountains, Nevada
- Cephaluris coloradensis* Olsen, 1949, illus.
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
redescription, measurements, key, syn.:
Cephaluris collaris Akhtar, 1958
Ochotona princeps: Colorado; Rocky Mountains, Sheep River and east of Kananaskis River, southwest Alberta
O. collaris: Ogilvie Mountains, Yukon
- Cephaluris coloradensis* Olsen, 1949
Seese, F. M., 1973, Am. Midland Naturalist, v. 89 (2), 257-265
key
Ochotona p. princeps (caecum, large intestine): St. Joe Baldy Mountain, Benewah County, Idaho
- Cephaluris coloradensis* Olsen 1949, illus.
Seese, F. M., 1977, J. Parasitol., v. 63 (3), 511-514
Cephaluris coloradensis, eggshell, morphology and histochemistry
- Cephaluris ochotonae* Akhtar, 1947
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
measurements, key, syn.: *Cephaluris vakhanica* Erhardova-Kotrla and Daniel, 1970
Ochotona rutila: Kazakhstan, USSR
- Cephaluris vakhanica* Erhardova-Kotrla and Daniel, 1970
Hobbs, R. P., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 151-160
as syn. of *Cephaluris ochotonae* Akhtar, 1947
- Cerascaris Cobb*, 1929
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Contraecum* Railliet & Henry, 1912
- Ceratospira* Schneider, 1866
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelaziinae
key
Syn.: *Annulospira* Jairajpuri & Siddiqi, 1969
- Cestocephalus Rasheed*, 1966
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Bulbocephalus* Rasheed, 1966
- Chabaudgolvania terdentatum* (Linstow, 1960) [i. e. 1890] Skrjabin, Schichobalova et Lagodoskaja, 1964
Hristovski, N. D., 1975, Acta Parasitol. Jugoslavica, v. 6 (1), 3-5
Rana graeca: Bitola district, Macedonia, Yugoslavia
- Chabaudstrongylus Durette-Desset et Denke*, 1978 [nomen nudum]
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Cooperiinae
- Chabertia*
Baines, D. M.; and Colegrave, A. J., 1977, Vet. Rec., v. 100 (11), 217-219
gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales
- Chabertia*
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Chab[ertia]*
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Chabertia*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Chabertia*
Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
sheep helminths, effective control with Nil-verb and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Chabertia*
Klein Mori, J., 1972, Rev. Med. Vet. y Parasitol., Maracay, v. 24 (1-8), 1971-1972, 207-226
gastrointestinal nematodes, sheep, Neguvon, Ripercol, Thibenzoline, comparison, various management systems, all effective, Ripercol easiest to administer, Neguvon somewhat toxic

- Chabertia**
Sewell, M. M. H., 1973, *Vet. Rec.*, v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Chabertia**
Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Chabertia**
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Chabertia sp.?**
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Chabertia spp.**
Searson, J. E.; and Doughty, F. R., 1977, *Austral. Vet. J.*, v. 55 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult *Ostertagia ostertagi* than larval forms): southern New South Wales
- Chabertia spp.**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Chabertia ovina**
Anderson, P. J. S.; and Marais, F. S., 1972, *J. South African Vet. Ass.*, v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Chabertia ovina** (Gmelin, 1790)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (large intestine): vicinity of Nowy Targ, Carpathian Mountains
- Chabertia ovina**
Boag, B.; and Thomas, R. J., 1973, *Research Vet. Sc.*, v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Chabertia ovina**
Boag, B.; and Thomas, R. J., 1977, *Research Vet. Sc.*, v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Chabertia ovina**
Chalmers, K., 1977, *N. Zealand Vet. J.*, v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Chabertia ovina**
Chroust, K.; and Dyk, V., 1975, *Deutsche Tierarztl. Wchnschr.*, v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Chabertia ovina**
Corticelli, B.; and Lai, M., 1972, *Parassitologia*, v. 14 (1), 95-96
Ovis musimon: Sardegna
- Chabertia ovina**
Downey, N. E., 1977, *Vet. Rec.*, v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Chabertia ovina**
Düewel, D.; et al., 1974, *Prakt. Tierarzt*, v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Chabertia ovina**
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Chabertia ovina**
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Chabertia ovina**
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia

- Chabertia ovina*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Chabertia ovina*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Chabertia ovina*
Foix, J., 1977, Rev. Med. Vet., Toulouse, v. 128 (8-9), 1111-1119
Dicrocoelium dendriticum, sheep, cambendazole, good results against flukes as well as gastrointestinal strongyles
- Chabertia ovina*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- [*Chabertia*] *ovina* "*Khabersia ovina*"
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, Vet. Sbirka, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of tetramisole, group and individual applications
- Chabertia ovina*
Horak, I. G.; Snijders, A. J.; and Louw, J.P., 1972, J. South African Vet. Ass., v. 43 (4), 397-403
trematodes and nematodes, sheep (exper.), rafoxanide, efficacy studies
- Chabertia ovina* (Fabricius, 1878) Railliet et Henry, 1909
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
Capreolus capreolus (large intestine): southern Bulgaria
- Chabertia ovina*
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Chabertia ovina*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Chabertia ovina*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Chabertia ovina*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence: Zehusice enclosure
- Chabertia ovina*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure
- Chabertia ovina*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Chabertia ovina*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Ovis aries: Chile
- Chabertia ovina*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Chabertia ovina*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Chabertia ovina*
Schweisgut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Chabertia ovina*
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Chabertia ovina*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Chabertia ovina*
Vlassoff, A., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand

- Chabertia ovina*
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Chabertia ovina*
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Chabertiella pesteri* Tadros
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Giraffa camelopardalis tippelskirchi (rectum, large intestine): Kenya
- Chandlerella Yorke* and Maplestone, 1926
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key to males of species
- Chandlerella* sp.
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (heart): Florida
- Chandlerella* sp.
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (lungs): Florida
- Chandlerella alii* Sultana, 1962, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella apusi* Sonin, 1963, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella bosei* (Chandler, 1924), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella braziliensis* Yeh, 1957, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella chitwoodae* Anderson, 1961, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella columbae* Sonin, 1966, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella columbigallinae* (Augustine, 1937), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella flexivaginalis* (Jones, 1961), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella hawkingi* Chatterjee, Sen, and Bhattacharya, 1965, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella himalayansis* Sultana, 1962, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella longicaudata* Sonin, 1966, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella periarterialis* (Cabellero, 1948), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella petrowi* (Tschertkow, 1946), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella quisicali* (von Linstow, 1904)
Cooper, C. L.; and Crites, J. L., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (brain):
South Bass Island, Ottawa County, Ohio
- Chandlerella quisicali*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (cerebrum): Franklin and Ottawa counties, Ohio
- Chandlerella quisicali* (von Linstow, 1904), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella robinsoni* n. sp., illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
Columba fasciata fasciata (blood vessels of liver; fascia of thighs): Colorado, U.S.A.
- Chandlerella sinensis* Li, 1933, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella singhi* Ali, 1956, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella skrjabini* (Petrov and Tschertkow, 1947), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key

- Chandlerella stantchinskyi* Gilbert, 1932, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella striatospicula* Hibler, 1964, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella sultana* Sonin, 1966, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella thapari* Rasheed, 1960, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Chandlerella travassosi* (Koroliowa, 1926), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Cheilobus* sp., illus.
Miles, P. M., 1977, Entom. Month. Mag. (1344-1347), v. 112, 169-172
Isotoma viridis: Ynyslas dune-slack, Nr. Borth, Ceredigion, Dyfed, Wales
- Cheilonematodum* Johnston & Mawson, 1941
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"poorly known but it may be a synonym of *Rusguniella*"
- Cheilospirura* Diesing, 1861
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Cheilospirura spinosa*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (gizzard lining): Florida
- Cheiropteranema* Sandground, 1929
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Cheiropteranema*
Durette-Desset, M. C.; and Tchepakoff, R., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1091-1094
Molineinae
systematic position and phyletic affinities
- Cheiropteranema globocephala* Sandground, 1929, illus.
Durette-Desset, M. C.; and Tchepakoff, R., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1091-1094
redescription
Artibeus jamaicensis lituratus (intestin): Guyane francaise
- Chelonidracunculus* Yamaguti, 1961
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Dracunculus* (Reichard, 1759)
- Cheniellospirura* Kou, 1962
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Gendrespirura* Chabaud, 1958
- Chenofilaria* [? n. rank]
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of *Dipetalonema*; key
tod: *D. (C.) filaria* (Kou, 1958)
- Chenofilaria filaria* Kou 1958
Esslinger, J. H., 1976, J. Parasitol., v. 62 (4), 527
Syn.: *Dipetalonema fausti* Esslinger 1966
- Chenospirura* Hsu, 1957, nec Kou, 1958
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cyrnea* Seurat, 1914
- Chenospirura* Kou, 1958, nec Hsu, 1957
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Gendrespirura* Chabaud, 1958
- Chevreauxia* Seurat, 1918, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Chevreauxia americana* Schmidt, 1968
Ahern, W. B.; and Schmidt, G. D., 1976, Parasitology, v. 73 (3), 381-398
Recurvirostra americana (under the koilon of the gizzard): Kansas; Colorado
- Chevreauxia revoluta* (Rud., 1819)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Charadrius hiaticula: Keta lake
- Chitwoodspirura* Chabaud & Rousselot, 1956, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
- Chlamydonema praeputiale* (Linstow 1889)
Acholonu, A. D., 1977, J. Parasitol., v. 63 (4), 757-758
cat: Ponce, Puerto Rico
- Chlamydopecta* Chandler, 1954
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"may be a synonym of *Vigisospirura*"

- Chordatortilis Mendonca & Rodrigues, 1965, illus.
- Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Chordocephalus [sic] Alegret, 1941, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
Syn.: Skrjabinocara Kurashvili, 1941; "We presume that Chordocephalus has priority but this is not certain."
- Chromadoria Pearse, 1942
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Adenophorea
includes: Araeolaimida; Monhysterida; Desmodorida; Chromadorida; Desmoscolecida
- Chromadorida Chitwood, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Chromadoria
includes: Chromadorina; Cyatholaimina
- Chromadorina Chitwood & Chitwood, 1937
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Chromadorida
- Chromonema n. gen.
Khan, A.; Brooks, W. M.; and Hirschmann, H., 1976, J. Nematol., v. 8 (2), 159-168
Steinernematidae
tod: C. heliothidis n. sp.
- Chromonema heliothidis n. gen., n. sp. (tod), illus.
Khan, A.; Brooks, W. M.; and Hirschmann, H., 1976, J. Nematol., v. 8 (2), 159-168
life cycle
Heliothis zea: soil, Clayton, North Carolina
Galleria mellonella (exper.)
Heliothis virescens (exper.)
Argyrotaenia velutinana (exper.)
Manduca sexta (exper.)
Estigmene acraea (exper.)
Maladera castanea (exper.)
Graphognathus sp. (exper.)
Culex pipiens quinquefasciatus (exper.)
Musca domestica (exper.)
Phormia regina (exper.)
- Chromonema heliothidis
Khan, A.; and Brooks, W. M., 1977, J. Invert. Pathol., v. 29 (3), 253-261
characterization of chromogenic bioluminescent bacterium associated with the entomophilic nematode Chromonema heliothidis, comparison of this bacterium with Achromobacter nematophilus
- Citellina marmotae Manter, 1930
Harley, J. P.; Thompson, M. P.; and Aubrey, D., 1973, Tr. Kentucky Acad. Sc., v. 34 (3, 4), 59
as syn. of C. triradiata Hall, 1916
- Citellina triradiata (Hall, 1915)
Babero, B. B., 1973, Tr. Am. Micr. Soc., v. 92 (2), 265-272
Spermophilus tereticaudus
Ammospermophilus leucurus
(caecum of all): all from Nevada
- Citellina triradiata Hall, 1916
Harley, J. P.; Thompson, M. P.; and Aubrey, D., 1973, Tr. Kentucky Acad. Sc., v. 34 (3, 4), 59
Syn.: C. marmotae Manter, 1930
Marmota monax (cecum): Washington County, Kentucky
- Citellina triradiata (Hall) Manter, 1930
Larson, O. R.; and Scharf, W. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 174-175
Marmota monax (large intestine): Itasca State Park, Minnesota
- Citellinema Hall, 1916
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
Syn.: Warrenius Hall, 1916
- Citellinema bifurcatum
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Citellinoides Dikmans, 1939
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
- Clavinema Yamaguti, 1935
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
? as syn. of Philometra Costa, 1845
- Cloacina australis (Yorke & Maplestone) [n. comb.]
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
Syn.: Macropostrongylus australis Yorke & Maplestone
- Cloacina australis Johnston & Mawson (1938), preoccupied, renamed C. daveyi nom. nov.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
- Cloacina daveyi nom. nov. for C. australis
Johnston & Mawson (1938), preoccupied
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
- Cloeoascaris Baylis, 1923
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Galeiceps Railliet, 1916

- Cloacascaris* sp., illus.
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
description
Martes martes (small intestine): Karelia
- Cnizostrongylus* Chabaud, Durette-Desset et Houin, 1967
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Libyostrongylinae
- Cobbostrongylus* [sic] Sarwar, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Trichostrongylus* Looss, 1905
- Cochlus* Zeder, 1803
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Goezia* Zeder, 1800
- Collarinema* Sey, 1970
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Pseudoproleptus* Khera, 1953
- Columbostrongylus* Puyllaert, 1968
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Comephoronema* Layman, 1933
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cystidicola* Fischer, 1798
- Cometeterakis* gen. nov.
Crusz, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
Heterakidae, Meteterakinae
mt: *C. lyriocephali* sp. nov.
- Cometeterakis lyriocephali* gen. et sp. nov. (mt), illus.
Crusz, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
Lyriocephalus scutatus (rectum): Godekande, Ceylon
- Conocephalus* Diesing, 1861, nec Thunberg, 1812
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakis* Dujardin, 1845
- Contortospiculum* rhea
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Contortylenchus pseudodiplogaster*
Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Contraeacinea* Mozgovoi & Shakhmatova, 1971
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinae
key; key to genera
includes: *Duplicaecum*; *Galeiceps*; *Contraeacum*; *Phocascaris*
- Contraeacum*
Courtney, C. H.; Forrester, D. J.; and White, F. H., 1977, J. Am. Vet. Med. Ass., v. 171 (9), 991-992
helminths in *Pelecanus occidentalis*, anthelmintic activity of arecoline hydrobromide, thiabendazole, niclosamide, 1-tetramisole: Bird Keys and Port Orange, Florida
- Contraeacum* Railliet & Henry, 1912, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Contraeacinea
key; synonymy
- Contraeacum* (*Ornitocaecum*) Mozgovoi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Contraeacum* Railliet & Henry, 1912
- Contraeacum* (*Synthetonema*) Kreis, 1952
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Contraeacum* Railliet & Henry, 1912
- Contraeacum* (*Erschowicaecum*) Mozgovoi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Thynnascaris* Dollfus, 1933
- Contraeacum* (*Simplexonema*) Kreis, 1952, nom. nud.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Thynnascaris* Dollfus, 1933
- Contraeacum* (*Thynnascaris*) Dollfus, 1935
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Thynnascaris* Dollfus, 1933
- Contraeacum*
Myers, B. J., 1976, Tr. Am. Micr. Soc., v. 95 (2), 137-142
human anisakiasis, historical review

- Contraecaecum
Soleim, O., 1976, Norwegian J. Zool., v. 24 (4), 464 [Abstract]
"It is concluded that the genus *Thynnascaris* should be maintained and that *Phocascaris* become a synonym of *Contraecaecum*."
- Contraecaecum sp.
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Cepphus carbo
Uria aalge
all from Rimsko-Korsakov islands
- Contraecaecum sp., probably *C. rudolphii*
Bakke, T. A.; and Barus, V., 1975, Norwegian J. Zool., v. 23 (3), 183-191
Larus canus (alimentary canal): Agdenes area, Norway
- Contraecaecum sp.
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
nematodes of *Larus canus*, age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Contraecaecum sp.
Beacham, B. E.; and Haley, A. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 232-233
Morone americana (mesenteries): Chesapeake Bay
- Contraecaecum sp.
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (esophagus): Florida
- Contraecaecum Type I
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 227-232
incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
Mugil cephalus
M. dussumieri
M. strongylocephalus
all from south-eastern Queensland
- Contraecaecum Type II
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 227-232
incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
Apogon fasciata
Platycephalus arenarius
Pseudorhombus arsius
P. jenynsii
all from south-eastern Queensland
- Contraecaecum sp. (Type I), illus.
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 233-243
description, key
- Contraecaecum sp. (Type II), illus.
Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 233-243
description, key
- Contraecaecum sp.
Edwards, R. W.; Harley, J. P.; and Williams, J. C., 1977, Tr. Kentucky Acad. Sc., v. 38 (3-4), 132-135
Ictalurus punctatus (mesenteries): Kentucky River drainage
- Contraecaecum sp.
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Ictalurus punctatus
Aplodinotus grunniens
Micropterus salmoides
Pomoxis annularis
Lepomis macrochirus
L. megalotis
L. microlophus
all from Eagle Mountain Lake, Texas
- Contraecaecum sp.
Kocan, A. A.; and Locke, L. N., 1974, J. Wildlife Dis., v. 10 (1), 8-10
Haliaeetus leucocephalus: New Jersey; Minnesota; South Dakota; Illinois; Wisconsin
- Contraecaecum sp.
Korotaeva, V. D., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 89-96
description, larva
Icelus spiniger (stomach, intestine): Sea of Japan
- Contraecaecum sp.
Lichtenfels, J. R., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 115
Polyorchis penicillatus (mesoglea near the radial canals): San Francisco Bay
- Contraecaecum sp.
Lockard, L. L.; and Parsons, R. R., 1975, Great Basin Nat., v. 35 (4), 425-426
Polyodon spathula: Yellowstone River near Intake, Montana
- Contraecaecum spp.
Lowe, P. O.; Ffolliott, P.; and Goodwin, J. G., 1977, Southwest. Nat., v. 22 (4), 537-538
Micropterus salmoides (mesenteries): Presa de Novillo Reservoir, Sonora, Mexico
- Contraecaecum sp.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Thunnus thynnus
Euthynnus affinis
Auxis thazard
Thunnus sp.
(body cavity of all): all from South China Sea
- Contraecaecum sp.
Miller, R. L.; Olson, A. C., jr.; and Miller, L. W., 1973, Calif. Fish and Game, v. 59 (3), 196-206
Lepomis macrochirus
Micropterus salmoides
Pomoxis annularis
P. nigromaculatus
all from southern California reservoirs

- Contracaecum sp.
 Niederkorn, J. Y., 1974, Tr. Missouri Acad. Sci., v. 7-8, 1973-1974, 160-163
 Lepomis cynellus: Johnson County, Missouri
- Contracaecum [sp.]
 Paperna, I., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 252
 brief description
 (pericardial cavity of all)
 Tilapia nilotica: Lake George; northern Lake Victoria
 Haplochromis spp.: Lake George; northern Lake Victoria
 Bagrus docmac: Lake George
- Contracaecum sp.
 Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (1), 267-277
 helminths, incidence and intensity of infection in young and adult Oncorhynchus nerka, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Contracaecum sp., larva
 Ponyi, J.; Biro, P.; and Murai, E., 1972, Parasitol. Hungar., v. 5, 383-408
 internal helminths of Acerina cernua (intestine), incidence survey, seasonal variations and host growth and development in relationship to parasitic burden: Lake Balaton, Hungary
- Contracaecum larvae
 Prudhoe, S.; and Hussey, C. G., 1977, Zoologica Africana, v. 12 (1), 113-147
 Clarias gariepinus (mesenteries, body wall, bile ducts): Transvaal, South Africa (Oliphants River; Kareepan, Wolmaranstad); Malkerns, Swaziland
- Contracaecum sp. (Type A)
 Sakaguchi, Y.; and Katamine, D., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 159-169
 anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
 Sillago japonica: sea near Nagasaki
 Argyrosomus argentatus: East China Sea
 Uranoscopus japonicus: "
 Rhinoplagusia japonica: "
 Upeneus bensasi: "
 Pleuronichthys cornutus: "
 Nemipterus virgatus: South China Sea
- Contracaecum sp. (Type E), illus.
 Sakaguchi, Y.; and Katamine, D., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 159-169
 anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
 Saurida tumbil: East China Sea
 Argyrosomus argentatus: "
 Zeus japonicus: "
 Fugu vermicularis vermicularis: "
 Uranoscopus japonicus: "
 Inegocia meerdervoorti: "
 Upeneus bensasi: East and South China Seas
 Pleuronichthys cornutus: East China Sea
 Taius tumifrons: "
 Lepidotrigla microptera: "
 Branchiostegus japonicus japonicus: "
- Contracaecum sp. (Type E), illus.-- Continued.
 Sakaguchi, Y.; and Katamine, D., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 159-169.-- Continued.
 Nemipterus virgatus: South China Sea
 Priacanthus sp.: "
 Abalistes stellatus: "
 Lutjanus basmira: "
 Pristipomoides sieboldi: "
 Epinephelus septemfasciatus: "
 Clidoderma asperrimum: "
 Plectorhynchus pictus: "
 Pseudalutarius [l.e. Pseudaluteres] nasicornis: "
 Lethrinus haematopterus: "
 Tachysurus falcarius: "
 Caranx equula: "
 Ilisha elongata: "
- Contracaecum sp. (Type F), illus.
 Sakaguchi, Y.; and Katamine, D., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 159-169
 anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
 Saurida tumbil
 Abalistes stellatus
 Clidoderma asperrimum
 all from South China Sea
- Contracaecum sp.
 Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
 Phocoena phocoena: insular Newfoundland and its adjacent waters
- Contracaecum sp.
 Smith, J. W., 1971, Nature (5330), v. 234, 478
 Thysanoessa raschii
 Nyctiphanes couchii
 Sagitta elegans
 all from off northeast coast of Scotland
- Contracaecum [sp.]
 Tasto, R. N., 1975, Fish Bull. (165), State Calif., Resources Agency, Dept. Fish and Game, 123-135
 Leptocottus armatus (coelom): Anaheim Bay
- Contracaecum sp., probably Contracaecum aduncum (Rudolphi)
 Vooren, C. M.; and Tracey, D., 1976, N. Zealand J. Marine and Freshwater Research, v. 10 (3), 499-509
 incidence, intensity
 Cheilodactylus macropterus (body cavity, stomach wall): New Zealand
- Contracaecum sp.
 White, G. E., 1974, Tr. Am. Micr. Soc., v. 93 (2), Apr., 280-282
 Catostomus commersoni: Kentucky River drainage system
- Contracaecum aduncum (Rudolphi, 1802)
 Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 80-88
 helminth distribution among age groups of Pleurogrammus azonus (intestine): Peter the Great Bay, Sea of Japan

- Contracaecum aduncum* (Rudolphi, 1802)
Korotaeva, V. D., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 89-96
Enophrys diceraus (body cavity): Sea of Japan
- Contracaecum aduncum*
Moeller, H., 1976, J. Marine Biol. Ass. United Kingdom, v. 56 (3), 781-785
intestinal helminths, elimination from host held in captivity, high rate of elimination of helminths unattached or slightly attached to host, lower elimination rate of helminths attached to host
Gadus morhua
Zoarces viviparus
Myoxocephalus scorpius
Platichthys flesus
(intestine of all): all from Kiel Fjord (western Baltic Sea)
- Contracaecum aduncum* (Rud.)
Soleim, Ø., 1976, Norwegian J. Zool., v. 24 (4), 319-323
as syn. of *Thynnascaris adunca* (Rud. 1802)
- Contracaecum brachyurum*
Harley, J. P., 1977, Tr. Kentucky Acad. Sci., v. 38 (3-4), 136-138
Pomoxis annularis (stomach, intestine): Lake Wilgreen, Madison County, Kentucky
- Contracaecum brachyurum* (Ward and Magath, 1917)
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Ictalurus catus
Morone saxatilis
Pomoxis nigromaculatus
Alosa sapidissima
all from Sacramento-San Joaquin Delta, California
- Contracaecum brachyurum* (Ward & Magath, 1917)
Mudry, D. R.; and Anderson, R. S., 1977, J. Fish. Biol., v. 11 (1), 21-33
Salvelinus namaycush
Lota lota
all from Waterton Lakes National Park, Canada
- Contracaecum brachyurum*
Rubertone, J. A.; and Hall, J. E., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 58-59
Micropterus dolomieu (intestine): Greenbrier River below Alderson, West Virginia
- Contracaecum microcephalum* Rudolphi, 1899
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmin. Lab., v. 15, 109-133
Anas platyrhynchos (small intestine): Bulgaria
- Contracaecum micropapillatum* (Stossich, 1890)
Baylis, 1920, illus.
Semenova, M. K., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 145-156
Contracaecum micropapillatum, detailed morphology of egg, four larval stages, juvenile and adult
Pelecanus crispus (nat. and exper.)
- Contracaecum multipapillatum*
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis: Florida and/or Louisiana
- Contracaecum* (C.) nehli Karokhin, 1949
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Podiceps griseigena: Rimsko-Korsakov islands
- Contracaecum osculatum* Rudolphi
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus
Phoca vitulina
(stomach of all): all from European waters
- Contracaecum osculatum* (Rudolphi, 1802)
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (stomach): Sakhalin Bay
- Contracaecum osculatum* (Rudolphi, 1802)
Popov, V. N.; 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
age dynamics of infection
Histriophoca fasciata (stomach, intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Contracaecum osculatum* (Rudolphi, 1802)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Pagophilus groenlandicus: insular Newfoundland and its adjacent waters
- Contracaecum osculatum*
Sweeney, J. C.; and Gilmartin, W. G., 1974, J. Wildlife Dis., v. 10 (4), 370-376
survey, diseases in California sea lions, diagnosis, treatment
Zalophus californianus: southern California beaches
- Contracaecum rudolphii* Hartwich, 1964
Bakke, T. A.; and Barus, V., 1975, Norwegian J. Zool., v. 23 (3), 183-191
Larus canus (alimentary canal): Agdenes area, Norway
- Contracaecum rudolphii*
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
nematodes of Larus canus, age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Contracaecum spheniscus*
Boero, J. J.; Led, J. E.; and Brandetti, E., 1972, Analecta Vet., v. 4 (1), 17-34
Spheniscus magellanicus
Eudyptes crestatus
(intestine of all): all from Argentine Republic

- Contraecaecum* (C.) *spiculigerum* (Rudolphi, 1809)
 Alekseev, V. M.; and Smetanina, Z. B., 1968,
 Gel'mint. Zhivot. Tikhogo Okeana (Skriabin),
 97-104
Phalacrocorax pelagicus
P. ussuriensis
Larus crassirostris
Cepphus carbo
Uria aalge
Botaurus stellaris
Nycticorax nycticorax
 all from Rimsko-Korsakov islands
- Contraecaecum spiculigerum* (Rudolphi, 1809)
 Belogurov, O. I.; Leonov, V. A.; and Zueva,
 L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana
 (Skriabin), 105-124
Larus argentatus
Sterna hirundo
Larus crassirostris
L. schistisagus
Stercorarius parasiticus
Uria lomvia
Cepphus carbo
Lunda cirrhata
 all from coast of Sea of Okhotsk
- Contraecaecum spiculigerum* (Rudolphi, 1819)
 Buck, O. D.; Cooper, C. L.; and Crites, J. L.,
 1976, Proc. Helminth. Soc. Washington, v. 43
 (2), 233-234
Larus argentatus: Bass Island region of
 Lake Erie
- Contraecaecum spiculigerum*
 Courtney, C. H.; and Forrester, D. J., 1974,
 Proc. Helminth. Soc. Washington, v. 41 (1),
 89-93
 prevalence and intensity, age of host
Pelecanus occidentalis: Florida and/or
 Louisiana
- Contraecaecum spiculigerum*
 Harley, J. P., 1977, Tr. Kentucky Acad. Sci.,
 v. 38 (3-4), 136-138
Pomoxis annularis (stomach, intestine):
 Lake Wilgreen, Madison County, Kentucky
- Contraecaecum spiculigerum* (Rudolphi, 1809)
 Hensley, G. H.; and Nahhas, F. M., 1975,
 Calif. Fish and Game, v. 61 (4), 201-208
Ictalurus nebulosus (mesentery): Sacramento-
 San Joaquin Delta, California
- Contraecaecum spiculigerum* (Rud. 1809)
 Sergeeva, T. P., 1969, Trudy Gel'mint. Lab.,
 Akad. Nauk SSSR, v. 20, 146-155
Larus ridibundus
Sterna hirundo
Larus ichthyaetus
 all from Tuva
- Cooperia*
 Bliss, D. H.; and Todd, A. C., 1977, Vet. Med.
 and Small Animal Clin., v. 72 (10), 1612-1617
 milk production in dairy cows exposed to
 mixed trichostrongylid larvae, results in-
 dicate that greatest milk loss occurs during
 the first 90 days of lactation, relationship
 between exposure time and stage of lactation
- Cooperia*
 Boag, B.; and Thomas, R. J., 1975, Research
 Vet. Sc., v. 19 (3), 293-295
 sheep nematodes, population dynamics, field
 studies, level of larval mortality may vary
 from year to year with prevailing climatic
 conditions, 'spring rise' in ewes is major
 source of pasture contamination causing wave
 of lamb infections in late August and Sep-
 tember
- Coop[eria]*
 Brunson, R. V., 1976, N. Zealand J. Exper.
 Agric., v. 4 (3), 275-279
 lambs, effectiveness of single thiabendazole
 drench at weaning in controlling build-up of
 trichostrongyle worm burdens, relative im-
 portance of various sources of pasture con-
 tamination (overwintered larvae; larvae de-
 posited by ewes and lambs in pre-weaning
 period; larvae deposited by lambs at weaning)
- Cooperia*
 Buerger, H.-J., 1976, Vet. Parasitol., v. 1
 (4), 359-366
Ostertagia, *Cooperia*, *Nematodirus*, signifi-
 cantly higher numbers of larvae on herbage
 samples collected from calf pastures vs. cow
 pastures, improved control of trichostrongyle
 infection during late summer and autumn might
 be achieved by transfer of calves to cow pas-
 tures
- Cooperia*
 Buerger, H. J., 1976, Ztschr. Parasitenk.,
 v. 50 (2), 219
 incidence on grass from cattle pasture:
 Niedersachsen
- Cooperia*
 Crowley, J. W., jr.; et al., 1977, Am. J. Vet.
 Research, v. 38 (5), 689-692
 lungworms, gastrointestinal parasites,
 cattle, 3 controlled critical trials, highly
 effective
- Cooperia*
 Downey, N. E.; and Moore, J. F., 1977, Vet.
 Rec., v. 101 (24), 487-488
Ostertagia, *Cooperia*, incidence in calves
 grazing on pastures which were fertilized
 with slurry: Ireland
- Cooperia*
 Duwel, D., 1977, Cahiers Bleus Vet. (26),
 201-215
 fenbendazole, efficacy against nematodes in
 various animals, useful as broad spectrum
 anthelmintic, mechanism of action, pharmaco-
 kinetics, metabolism, toxicology
- Cooperia Ransom*, 1907 (type genus)
 Durette-Desset, M. C.; and Chabaud, A. G.,
 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Cooperiinae*
- Cooperia*
 Eckert, J., 1972, Schweiz. Arch. Tierh., v.
 114 (12), 652-667
 control measures based on limited informa-
 tion available, pasture change and anthel-
 mintic treatment at end of June, review:
 Switzerland

- Cooperia**
Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Cooperia**
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Cooperia**
Henriksen, Sv. Aa.; et al., 1976, Vet. Parasitol., v. 2 (3), 259-272
gastro-intestinal nematodes, young calves during first grazing season, infection levels, blood findings, body weight gains, comparison of animals grazing same pasture entire season with those moved in early July and between levamisole-treated and untreated animals: Denmark
- Cooperia**
Jordan, H. E.; et al., 1977, Am. J. Vet. Research, v. 38 (8), 1157-1160
Ostertagia ostertagi, Cooperia, influence on energy efficiency in full-fed vs. maintenance-fed steers with high vs. low worm burdens (low worm burdens did not significantly effect energy utilization; in full-fed steers, energy retention was greater in steers with lower worm burdens; maintenance-fed steers were more heavily parasitized than full-fed steers)
- Cooperia**
Sewell, M. M. H., 1973, Vet. Rec., v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Cooperia**
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, Am. J. Vet. Research, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Cooperia**
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Cooperia**
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Cooperia**
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Cooperia**
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, Vet. Glasnik, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against Dictyocaulus filaria and most gastrointestinal helminths except Strongyloides papillosus, Trichuris ovis, and Moniezia sp.
- Cooperia**
Wallnoefer, E., 1977, Wien. Tierarztl. Monatsschr., v. 64 (4), 129-131
sheep parasites, Mebenvet, good results when treatment was repeated after 14 days: Austria
- Cooperia**
Whitlock, J. H.; and Georgi, J. R., 1976, Parasitology, v. 72 (3), 207-224
biological controls in mixed trichostrongylid infections (predominantly Haemonchus contortus cayugensis) in sheep, different ecosystems (barn vs. pasture) and different treatment groups, course of infections (erythrocyte loss, fecal egg counts, hematocrit values), "Anaphylactoid 'self-cure' did not occur in this experiment but something like premunition certainly did."
- Cooperia**
Zeakes, S. J.; et al., 1976, Am. J. Vet. Research, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos crumbles and naftalofos boluses
- Cooperia spp.**
Bryan, R. P., 1976, Austral. Vet. J., v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Cooperia spp.**
Bryan, R. P.; Bainbridge, M. J.; and Kerr, J. D., 1976, Austral. J. Zool., v. 24 (3), 417-421
Bubalus bubalis
cattle
(large and small intestine of all): all from Northern Territory, Australia
- Cooperia [sp.]**
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
ruminants, survey, treatment, economic importance: Kaedi area (Mauritania)
- Cooperia sp.**
Campbell, W. C.; and Thomson, B. M., 1973, Austral. Vet. J., v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of Trichostrongylus colubriformis proved uninfactive even if they had not been frozen

- Cooperia sp.
Canale, A.; et al., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 306-310
Fasciola hepatica, calves, light experimental infection alone or in combination with gastrointestinal nematodes, digestive function not impaired
- Cooperia spp.
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Cooperia sp.
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 526-532
cattle, morantel tartrate, good results against Cooperia sp., Ostertagia sp., and Trichostrongylus sp., increased weight gain in treated cattle, field trials: United Kingdom
- Cooperia spp.
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Vet. Rec., v. 92 (20), 551-554
control of clinical infections of gastrointestinal nematodes and lungworms in calves using morantel/diethylcarbazine solution, field trials, good results as measured by growth response and clinical symptoms; routine treatment economically sound under conditions of heavy infection
- Cooperia spp.
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxfendazole, drug efficacy
- Cooperia spp.
Curr, C., 1977, Austral. Vet. J., v. 53 (9), 425-428
nematodes, calves, levamisole, efficiency of pour-on formulation, drug trials, good results
- Cooperia spp.
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Cooperia spp.
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against Ostertagia spp., similar efficacy against other species), both drugs increased calves' weight gains
- Cooperia spp.
Duncan, J. L.; et al., 1976, Vet. Rec., v. 98 (17), 342
Ostertagia ostertagi (inhibited 4th stage larvae), Trichostrongylus axei, Cooperia spp., fenbendazole, good results, compared with levamisole
- Cooperia spp.
El-Abdin, Y. Z.; et al., 1975, Egypt. J. Vet. Sc., v. 12 (1), 31-43
serum constituents and serum enzyme activities, normal and nematode infested Camelus dromedarius: Cairo abattoir
- Cooperia [sp.]
Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Cooperia spp.
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Cooperia spp.
Lukovich, R.; et al., 1977, Gac. Vet., Buenos Aires (318), v. 39, 91-95
helminths, cattle, levamisole, results from injectable and dermal application similar
- Cooperia sp. 4th stage
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Cooperia spp.
McBain, D. G.; et al., 1977, Vet. Rec., v. 101 (14), 285-286
helminths, calves, fenbendazole in feed blocks
- Cooperia spp.
Niec, R.; et al., 1976, Gac. Vet., Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary antihelminthic treatment that could prevent normal buildup of resistance
- Cooperia sp.
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile
- Cooperia sp.
de Oliveira, A. R., 1976, Arq. Inst. Biol., Sao Paulo, v. 43 (1-2), 53-56
Oesophagostomum sp., Cooperia sp., Haemonchus sp., calves (exper.), no correlation between level of infestation and circulating eosinophils, may result from eosinophil migration to affected organs or bone marrow exhaustion

- Cooperia sp.
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Cooperia spp.
Ronald, N. C.; Bell, R. R.; and Craig, T. M., 1977, J. Am. Vet. Med. Ass., v. 170 (3), 317-319
gastrointestinal nematodes, calves, levamisole phosphate, effective at one-half recommended dosage
- Cooperia spp.
Searson, J. E.; and Doughty, F. R., 1977, Austral. Vet. J., v. 53 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult *Ostertagia ostertagi* than larval forms): southern New South Wales
- Cooperia spp.
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Cooperia spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Cooperia sp.
Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidian drug (Cozurone), good control of all except *Strongyloides*
- Cooperia spp.
Vlassoff, A., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand
- Cooperia [sp.]
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Cooperia sp.
Wikerhauser, T.; et al., 1974, Acta Parasitol. Iugoslavica, v. 5 (2), 79-81
trichostrongylids, cattle, fenbendazole compared with thiabendazole, good results from both
- Cooperia sp.
Williams, J. C.; et al., 1977, Vet. Rec., v. 101 (24), 484-486
Ostertagia ostertagi, cattle, albendazole, good results against inhibited fourth stage larvae as well as developing stages and adults, high efficacy against *Haemonchus* and *Cooperia* spp., no signs of toxicity
- Cooperia spp.
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Cooperia spp.
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Cooperia antidorca
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella granti (small intestine): Kenya
- Cooperia bisonis
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Cooperia bisonis
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Cooperia bisonis
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Cooperia curticei
Ahluwalia, J. S., 1976, Indian J. Animal Sc., v. 46 (5), 256-267
Cooperia curticei, survival, migration on soil and grass of infective larvae under natural conditions, various meteorological data
- Cooperia curticei
Ahluwalia, J. S., 1977, Indian J. Animal Sc., v. 45 (8), 1975, 592-593
Cooperia curticei, survival of infective larvae in water, under natural conditions, influence of temperature at different times of the year

- Cooperia curticei*
Ahluwalia, J. S., 1977, Indian J. Animal Sc., v. 45 (9), 1975, 706-708
Cooperia curticei, egg production in sheep measured by fecal counts and post mortem, can be as high as 2000/female/day
- Cooperia curticei*
Ahluwalia, J. S., 1977, Indian J. Animal Sc., v. 45 (12), 1975, 978-980
Cooperia curticei, sheep (exper.), gel-precipitin tests, results indicate that antibodies can be formed locally in alimentary tract, and circulating antibodies of serum detected earlier than mucous samples
- Cooperia curticei* (Railliet, 1893)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Cooperia curticei*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Cooperia curticei*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Cooperia curticei*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Cooperia curticei*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Cooperia curticei*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Cooperia curticei*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Cooperia curticei*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Cooperia curticei*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Cooperia curticei*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Cooperia curticei*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (small intestine): Kentucky
- Cooperia curticei*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Cooperia curticei*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Cooperia curticei*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Cooperia curticei*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter

- Cooperia curticei*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974,
Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus
Ovis aries
all from Chile
- Cooperia curticei*
Panitz, E., 1977, J. Helminth., v. 51 (1),
23-30
ethyl-6-ethoxybenzothiazole-2-carbamate,
evaluation of anthelmintic activity in po-
nies, swine, lambs, and chickens
- Cooperia curticei*
Prestwood, A. K.; Pursglove, S. R.; and Hayes,
F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-
385
survey of parasites of *Odocoileus virgini-
anus* and *Ovis aries* on common range, deer
unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Cooperia curticei*
Zajicek, D.; and Kozdon, O., 1977, Veterinarst-
vi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthi-
zation with pyrantel HCl, helmantac and nil-
verm to nematode incidence on pastures,
three-year study, overall decrease
- Cooperia fueelleborni* Hung, 1926
Basson, P. A.; et al., 1970, Onderstepoort J.
Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus
caffer*, some pathological findings, age of
host
Syncerus caffer (small intestine): Kruger
National Park
- Cooperia fueelleborni* Hung
Pester, F. R. N.; and Laurence, B. R., 1974,
J. Zool., London, v. 174 (3), 397-406
Alcelaphus buselaphus cokei (digestive tract)
Connochaetes taurinus (small intestine)
all from Kenya
- Cooperia mcmasteri*
Guerrero, C.; Rojas, M.; and Vargas, J., 1974,
Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity
of 1-tetramisole, significant body weight
gain in treated animals
- Cooperia mcmasteri*
Helle, O.; and Tharaldsen, J., 1976, Vet. Para-
sitol., v. 1 (4), 345-357
Ostertagia ostertagi and *Cooperia* spp. in
young cattle during their first grazing sea-
son, free-living stages overwintered in suf-
ficient numbers to cause reduced weight gain
and clinical disease in early spring, thia-
bendazole treatment and move to clean pasture
improved weight gain: Norway
- Cooperia mcmasterii*
Kelly, J. D.; et al., 1975, Research Vet.
Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole
against naturally acquired *Dictyocaulus
filaria* infection associated with concu-
rent infection of gastro-intestinal nema-
todes in sheep
- Cooperia mcmasteri*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974,
Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Cooperia mcmasteri*
Prestwood, A. K.; Pursglove, S. R.; and Hayes,
F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-
385
survey of parasites of *Odocoileus virgini-
anus* and *Ovis aries* on common range, deer
unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Cooperia mcmasteri*
Randall, R. W.; and Gibbs, H. C., 1977, Am.
J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle,
occurrence, degree of parasitism, and season-
al fluctuations: Maine
- Cooperia mcmasteri*
Vargas, J.; Guerrero, C.; and Rojas, M., 1972,
Rev. Invest. Pecuarias, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight
toxicity
- Cooperia neitzi* Moennig, 1930
Verster, A.; Imes, G. D., jr.; and Smit,
J. P. J., 1975, Onderstepoort J. Vet. Re-
search, v. 42 (1), 29-31
Damaliscus dorcas dorcas: captured at
Bontebok National Park, Swellendam and
transferred to the National Zoological
Gardens, Pretoria
- Cooperia oncophora*
van Adrichem, P. W. M.; and Shaw, J. C., 1977,
J. Animal Sc., v. 45 (3), 423-429
gastrointestinal nematodes, effects on
growth performance and milk production in
cambendazole-treated vs. non-treated mono-
zygous twin cattle naturally infected on
pasture during the first lactation period
- Cooperia oncophora*
Benz, G. W.; and Ernst, J. V., 1977, Am. J.
Vet. Research, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.),
albendazole significantly reduced infesta-
tions
- Cooperia oncophora*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25
(10), 266-269
gastrointestinal nematodes and cestodes,
sheep, oxfendazole, drug efficacy, good
results: New Zealand
- Cooperia oncophora*
Chroust, K.; and Dyk, V., 1975, Deutsche
Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of heifers,
efficacy of fenbendazole, thiabendazole
and tetramisole compared

- Cooperia oncophora*
Giordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH₄NO₃, prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Cooperia oncophora*
Coles, G. C.; and Simpkin, K. G., 1977, Research Vet. Sc., v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Cooperia oncophora*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Cooperia oncophora*
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 518-525
gastrointestinal nematodes and lungworms, calves (exper.), morantel tartrate, efficacy in 5 controlled trials, toxicity experiments demonstrate wide safety margin
- Cooperia oncophora*
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Cooperia oncophora*
Downey, N. E.; and O'Shea, J., 1977, Vet. Rec., v. 100 (13), 265-266
Dictyocaulus viviparus, *Ostertagia ostertagi*, *Cooperia oncophora*, calves (exper.), low dose levels of levamisole or morantel administered via drinking water, good results
- Cooperia oncophora*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Cooperia oncophora*
Eckert, J.; and Eisenegger, H., 1976, Zentralbl. Vet.-Med., Beihefte (25), 155-160
Dictyocaulus viviparus, cattle, program for management and control, vaccination with Dicotol, tetramisole treatment; *Ostertagia ostertagi*, *Cooperia oncophora*, pyrantel tartrate treatment to control concurrent infection limiting weight gain
- Cooperia oncophora*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Cooperia oncophora*
Fincher, G. T., 1975, J. Parasitol., v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Cooperia oncophora*
Goldberg, A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Cooperia oncophora*
Guerrero, C.; Rojas, M.; and Vargas, J., 1974, Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity of 1-tetramisole, significant body weight gain in treated animals
- Cooperia oncophora*
Helle, O.; and Tharaldsen, J., 1976, Vet. Parasitol., v. 1 (4), 345-357
Ostertagia ostertagi and *Cooperia* spp. in young cattle during their first grazing season, free-living stages overwintered in sufficient numbers to cause reduced weight gain and clinical disease in early spring, thiabendazole treatment and move to clean pasture improved weight gain: Norway
- Cooperia oncophora*
Henriksen, S. A.; Benthholm, B. R.; and Nielsen-Englyst, A., 1976, Nord. Vet.-Med., v. 28 (4-5) 201-209
gastro-intestinal strongyles, cattle, seasonal distribution on pastures
- Cooperia oncophora*
Herlich, H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.), oxfendazole, efficacy against adult and larval stages
- Cooperia oncophora*
Herlich, H., 1977, Am. J. Vet. Research, v. 38 (8), 1247-1248
efficacy of albendazole against gastrointestinal nematodes and *Fasciola hepatica* in cattle (exper.); comparison of critical vs. controlled tests
- Cooperia oncophora* (Railliet, 1898) Ransom, 1907
Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Cooperia oncophora*
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (small intestine): southwestern Iceland, slaughterhouse in Reykjavik

- Cooperia oncophora*
 Kelly, J. D.; et al., 1975, *Research Vet. Sc.*, v. 19 (1), 105-107
 anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Cooperia oncophora*
 Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
 nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Cooperia oncophora*
 Lancaster, M. B.; and Hong, C., 1977, *Vet. Rec.*, v. 101 (4), 81-82
Ostertagia ostertagi, *Cooperia oncophora*, calves, variable action of fenbendazole on arrested fourth stage larvae
- Cooperia oncophora*
 Lyons, E. T.; et al., 1975, *Am. J. Vet. Research*, v. 36 (6), 777-780
 calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Cooperia oncophora*
 Michel, J. F.; Lancaster, M. B.; and Hong, C., 1974, *J. Comp. Path.*, v. 84 (4), 539-554
Ostertagia ostertagi, *Cooperia oncophora*, evidence that arrested development is due to action of environmental factors, nature of environmental signals not precisely identified but not simple, changes which they induce in larvae are spontaneously reversed after a time
- Cooperia oncophora*
 Michel, J. F.; Lancaster, M. B.; and Hong, C., 1975, *J. Comp. Path.*, v. 85 (1), 133-138
Ostertagia ostertagi, *Cooperia oncophora*, arrested development, effect of temperature at free-living 3rd stage, larvae stored at 4° C. compared with storage at 15° C. and with a change of temperature after 12 weeks
- Cooperia oncophora*
 Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus
Ovis aries
 all from Chile
- Cooperia oncophora*
 Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
 survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Cooperia oncophora*
 Pursglove, S. R.; et al., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (9), 896-900
 intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Cooperia oncophora*
 Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
 gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Cooperia oncophora*
 Reinecke, R. K., 1972, *Onderstepoort J. Vet. Research*, v. 39 (3), 153-178
 gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Cooperia oncophora*
 Rose, J. H., 1971, *Symposia Brit. Soc. Parasitol.*, v. 9, 109-121
 gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Cooperia oncophora*
 Rose, J. H., 1973, *Research Vet. Sc.*, v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostrogylus rubidus*, culture from infective larva to adult worm in WAe medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Cooperia oncophora*
 Smeal, M. G.; et al., 1977, *Austral. Vet. J.*, v. 53 (12), 566-573
 nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Cooperia oncophora* (Railliet, 1898)
 Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Bos taurus: insular Newfoundland
- Cooperia oncophora*
 Smith, H. J., 1976, *Canad. J. Comp. Med.*, v. 40 (3), 320-321
 mixed *Ostertagia ostertagi* and *Cooperia oncophora* larvae in experimentally infected calves, no significant maturation requirement obtained, maximum infectivity possibly related to incubation temperature

- Cooperia oncophora*
Tharaldsen, J., 1976, Acta Vet. Scand., v. 17, Suppl. 61, 1-21
trichostrongylid infections, calves, survival of larvae on pasture, occurrence of larvae not influenced by artificial irrigation; treatment with thiabendazole did not effectively control infection due to overwintering larvae, neither improved weight gain nor reduced egg production: Norway
- Cooperia oncophora*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Cooperia oncophora*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Cooperia oncophora*
Vargas, J.; Guerrero, C.; and Rojas, M., 1972, Rev. Invest. Pecuarias, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight toxicity
- Cooperia pectinata*
Anderson, P. J. S.; and Marais, F. S., 1975, J. South African Vet. Ass., v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Cooperia pectinata*
Giordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Cooperia pectinata*
Dharsana, R. S.; Fabiyi, J. P.; and Hutchinson, G. W., 1976, Vet. Parasitol., v. 2 (4), 333-340
mixed gastro-intestinal nematode infections, calves, effects on host intestinal enzymes
- Cooperia pectinata*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Cooperia pectinata*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Cooperia pectinata*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Cooperia pectinata*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Cooperia pectinata*
Fincher, G. T., 1975, J. Parasitol., v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Cooperia pectinata*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Cooperia pectinata*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence: Zehusice enclosure
- Cooperia pectinata*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure
- Cooperia pectinata*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Cooperia pectinata* Ransom
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Alcelaphus buselaphus cokel (digestive tract): Kenya
- Cooperia pectinata*
Randall, R. W.; and Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine

- Cooperia pectinata**
Reinecke, R. K., 1972, Onderstepoort J. Vet. Research, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Cooperia pectinata**
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Cooperia pectinata**
Rowlands, D. ap T.; and Berger, J., 1977, J. South African Vet. Ass., v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Cooperia pectinata**
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Cooperia pectinata**
Tager-Kagan, P., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (4), 317-321
gastro-intestinal nematodes, zebu cattle (1 to 2 years old), cambendazole: Niger
- Cooperia pectinata**
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Cooperia pectinata**
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Cooperia punctata**
van Adrichem, P. W. M.; and Shaw, J. C., 1977, J. Animal Sc., v. 45 (3), 417-422
gastrointestinal nematodes, monozygous twin cattle, comparison of treated and untreated pairs infected naturally on pasture, growth performance, results indicate that the reduced growth may be long-lasting
- Cooperia punctata**
van Adrichem, P. W. M.; and Shaw, J. C., 1977, J. Animal Sc., v. 45 (3), 423-429
gastrointestinal nematodes, effects on growth performance and milk production in cambendazole-treated vs. non-treated monozygous twin cattle naturally infected on pasture during the first lactation period
- Cooperia punctata**
Anderson, P. J. S.; and Marais, F. S., 1975, J. South African Vet. Ass., v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Cooperia punctata**
Benz, G. W.; and Ernst, J. V., 1976, Am. J. Vet. Research, v. 37 (8), 895-899
Cooperia punctata and/or Eimeria bovis-infected calves, reduced alkaline phosphatase activities in intestinal mucosa
- Cooperia punctata**
Benz, G. W.; and Ernst, J. V., 1977, Am. J. Vet. Research, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.), albendazole significantly reduced infestations
- Cooperia punctata**
Ciordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Cooperia punctata**
Dharsana, R. S.; Fabiyi, J. P.; and Hutchinson, G. W., 1976, Vet. Parasitol., v. 2 (4), 333-340
mixed gastro-intestinal nematode infections, calves, effects on host intestinal enzymes
- Cooperia punctata**
Fincher, G. T., 1975, J. Parasitol., v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Cooperia punctata**
Goldberg, A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Cooperia punctata**
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil

- Cooperia punctata*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, *J. South African Vet. Ass.*, v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Cooperia punctata* (Linstow, 1906) Ransom, 1907
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Cooperia punctata*
Leland, S. E., jr.; et al., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 449-456
Cooperia punctata, in vitro-grown parasitic stages, evaluation of 28 anthelmintics using presumptive and confirmatory tests, comparison of in vitro potency with established in vivo activity against *Cooperia* and various other parasites, potential usefulness as preliminary screening method
- Cooperia punctata*
Lyons, E. T.; et al., 1975, *Am. J. Vet. Research*, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Cooperia punctata*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus: Chile
- Cooperia punctata*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries
Odocoileus virginianus
all from Hardy County, West Virginia
- Cooperia punctata*
Pursglove, S. R.; et al., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Cooperia punctata*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Cooperia punctata*
Reinecke, R. K., 1972, *Onderstepoort J. Vet. Research*, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Cooperia punctata*
Ridley, R. K.; Slonka, G. F.; and Leland, S. E., jr., 1977, *J. Parasitol.*, v. 63 (2), 348-356
Cooperia punctata, L₄ and adult stages grown in vitro, utilization of propionic acid, use of propionate by worms would result in depriving ruminant host of some of its necessary glucogenic precursors and could account for specific pathogenic mechanism attendant to heavy infections
- Cooperia punctata*
Rowlands, D. ap T.; and Berger, J., 1977, *J. South African Vet. Ass.*, v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Cooperia punctata*
Schweisgut, I., 1975, *Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75*, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Cooperia punctata*
Smeal, M. G.; et al., 1977, *Austral. Vet. J.*, v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Cooperia punctata*
Tager-Kagan, P., 1976, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 29 (4), 317-321
gastro-intestinal nematodes, zebu cattle (1 to 2 years old), cambendazole: Niger
- Cooperia punctata*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Cooperia punctata*
Todd, K. S., jr.; Levine, N. D.; and Wagher, B. N., 1977, *J. Parasitol.*, v. 63 (5), 956-957
Cooperia punctata, effects of repeated desiccation and rehydration on survival of infective larvae

- Cooperia punctata*
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Cooperia punctata*
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Cooperia punctata*
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Cooperia spatulata*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Cooperia surnabada*
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Cooperia verrucosa* Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Alcelaphus buselaphus cokei (digestive tract): Kenya
- Cooperiinae (Skrjabin et Schulz, 1937, tribu) Skrjabin et Schikhobalova, 1952
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae
includes: *Cooperia* (type genus); *Chabaudstrongylus* [nomen nudum]; *Cooperioides*; *Gazellostrongylus*; *Impalaia*; *Megacooperia*; *Paracooperia*
- Cooperioides* Daubney, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Cooperiinae
- Cooperioides* sp.
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Cooperioides antidorca* Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Cooperioides antidorci*
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 195-198
Antidorcas marsupialis (duodenum): Mountain Zebra National Park near Cradock, Cape Province
- Cordicauda Durette-Desset*, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Cordicauda Durette-Desset*, 1971
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Cordicauda magnabursa* n. sp., illus.
Durette-Desset, M.-C.; Diaw, O.; and Krishnasamy, M., 1975, Ann. Parasitol., v. 50 (4), 477-491
Trichys lipura (intestin): Malaisie
- Cordicauda malayensis* n. sp., illus.
Durette-Desset, M.-C.; Diaw, O.; and Krishnasamy, M., 1975, Ann. Parasitol., v. 50 (4), 477-491
Trichys lipura (intestin): Malaisie
- Cordicauda trichysi* n. sp., illus.
Durette-Desset, M.-C.; Diaw, O.; and Krishnasamy, M., 1975, Ann. Parasitol., v. 50 (4), 477-491
Trichys lipura (intestin): Malaisie
- Cordonema* Schmidt & Kuntz, 1971
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Skrjabinoclava* Sobolev, 1943
- Cordophilus sagittus*
Young, E.; and Basson, P. A., 1976, J. South African Vet. Med. Ass., v. 47 (1), 57
Cordophilus sagittus, *Haemonchus* spp., eland, pathology, levamisole hydrochloride, good results against gastro-intestinal parasites: Kruger National Park, translocated from Addo Elephant National Park
- Coregonema* Bauer, 1946
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
? as syn. of *Philonema* Kuitunen-Ekbaum, 1933
- Coronilla minuta* Beneden, 1871
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus acutus* Dujardin, 1845
- Coronilla robusta* Beneden, 1871
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus acutus* Dujardin, 1845

- Cosmocephalus* Molin, 1858, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Cosmocephalus* sp., larva
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Charadrius hiaticula
Philomachus pugnax
all from Keta lake
- Cosmocephalus australiensis* Johnston and Mawson, 1952
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
as syn. of Synhimantus australiensis (Johnston and Mawson, 1952) Yamaguti, 1961
- Cosmocephalus diesingi* Molin, 1858
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
species inquirenda
- Cosmocephalus firlothei* Rao, 1951
Keppner, E. J., 1973, Tr. Am. Micr. Soc., v. 92 (2), 288-291
Larus californicus (esophagus): city dump of Laramie, Wyoming
- Cosmocephalus obvelatus* (Creplin, 1825)
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Larus crassirostris
Cepphus carbo
all from Rimsko-Korsakov islands
- Cosmocephalus obvelatus* (Creplin, 1825)
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
nematodes of Larus canus (esophagus, ventriculus), age and sex of host, seasonal variations, distribution in alimentary canal:
Agdenes, Norway
- Cosmocephalus obvelatus*
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus argentatus
L. crassirostris
L. canus
Sterna hirundo
Larus ridibundus
all from coast of Sea of Okhotsk
- Cosmocephalus obvelatus* (Creplin, 1825)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Tringa glareola
Heteroscelus incanus brevipes
all from lower Yenisei [and/or] Keta lake
- Cosmocephalus obvelatus* (Creplin, 1825)
Buck, O. D.; Cooper, C. L.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 233-234
Larus argentatus: Bass Island region of Lake Erie
- Cosmocephalus obvelatus*
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (esophagus, proventriculus): Florida; Louisiana
- Cosmocephalus obvelatus* (Creplin, 1825)
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus genei: Azov Sea
L. ichthyaetus: Azov Sea; Tuva
L. argentatus: Azov Sea
L. ridibundus: Azov Sea; Tuva
L. minutus: Azov Sea
Sterna sandvinensis: Azov Sea
S. hirundo: Tuva
- Cosmocerca brasiliensis* Travassos 1925, illus.
Dyer, W. G.; and Altig, R., 1976, J. Parasitol., v. 62 (2), 262-264
redescription
Dendrobates parvulus
Eleutherodactylus sp.
Hamptophryne bolivianus
Hyla boans
Hyla geographica
Hyla lanciformis
Hyla leucophyllata
Hyla marmorata
Hyla sarayacuensis
Ischnocnema quixenis
Phyllomedusa tarsius
(lumen of large intestine of all): all from Santa Cecilia, Napo Province, Ecuador
- Cosmocerca brasiliensis* Travassos 1925
Dyer, W. G.; and Altig, R., 1977, Herpetologica, v. 33 (3), 293-296
Eleutherodactylus altamazonicus
E. lanthinites
(large intestine of all): all from Santa Cecilia, Napo Province, Ecuador
- Cosmocerca commutata* (Diesing, 1851)
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Rana esculenta: Samara river valley, Ukrainian SSR
- Cosmocerca commutata*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Rana ridibunda
Bufo viridis
Bufo bufo
all from Macedonia, Yugoslavia
- Cosmocerca commutata* (Diesing, 1851) Diesing, 1861, illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Bufo viridis
B. bufo
(debelo crijevo of all): all from Yugoslavia
- Cosmocerca kashmirensis* Fotedar 1959, illus.
Fotedar, D. N.; Duda, P. L.; and Raina, M. K., 1973, Chromosome Inform. Serv. (14), 16-18
Cosmocerca kashmirensis, chromosome number and morphology
Bufo viridis (intestines): Kashmir, India

- Cosmocerca ornata* Railliet et Henry
Bozhkov, D., 1974, *Izvest. Tsentral. Khelmint. Lab.*, v. 17, 25-31
8 helminth species in *Rana ridibunda* fed to *Natrix natrix* or *N. tessellata*, found that *Diplodiscus subclavatus*, *Opisthioglyphe ran-ae*, *Cephalogonimus retusus*, and *Cosmocerca ornata* can pass alive from body of ingested frog to intestine of *Natrix natrix*, and *D. subclavatus* to *N. tessellata*
- Cosmocerca ornata*
Hristovski, N. D., 1973, *Acta Parasitol. Iugoslavica*, v. 4 (2), 87-91
Rana ridibunda
Rana temporaria
Rana graeca
Hyla arborea
Pelobates syriacus balcanicus
Bombina variegata
Bufo viridis
Bufo bufo
Triturus vulgaris
all from Macedonia, Yugoslavia
- Cosmocerca ornata* (Dujardin, 1945) Railliet et Henry, 1916
Hristovski, N. D., 1975, *Acta Parasitol. Iugoslavica*, v. 6 (1), 3-5
Rana graeca: Bitola district, Macedonia, Yugoslavia
- Cosmocerca ornata* (Dujardin, 1845) Railliet et Henry, 1916
Hristovski, N. D.; and Lees, E., 1973, *Acta Parasitol. Iugoslavica*, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Cosmocerca ornata* (Dujardin, 1845), illus.
Milka, R., 1976, *Veterinaria, Sarajevo*, v. 25 (3), 449-476
Rana ridibunda
R. esculenta
R. temporaria
R. agilis
Bombina bombina
B. variegata
Bufo bufo
(debelo crijevo of all): all from Yugoslavia
- Cosmocerca ornata* (Dujardin, 1845; Railliet et Henry, 1916), illus.
Rozman, M., 1971, *Acta Parasitol. Iugoslavica*, v. 2 (2), 67-77
description
synonymy
Rana esculenta (debelo i tanko crijevo): environs of Novi Sad, Yugoslavia
- Cosmocercella polessiensis* Maguza, 1973
Antsyshkina, L. M.; et al., 1976, *Vestnik Zool. Akad. Nauk Ukrainsk. SSR, Inst. Zool.* (2), 82-84
Bombina bombina
Rana ridibunda
all from Samara river valley, Ukrainian SSR
- Cosmocercoidea*
Chabaud, A. G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (1), 6-17
Ascaridida
key
- Cosmocercoides* sp.
Koller, R. L.; and Gaudin, A. J., 1977, *Southwest. Nat.*, v. 21 (4), 503-509
helminth recovery at 2 sites with diverse climates, statistical analysis indicates correlations between incidence and/or intensity of infection and host species, locality, and sex and size of host
Hyla regilla
Bufo boreas
all from Los Angeles County, California (Malibu Creek; Big Tujunga Wash)
- Cosmocercoides dukae* (Holl, 1928) Wilkie, 1930
Dyer, W. G.; and Brandon, R. A., 1973, *Tr. Illinois Acad. Sc.*, v. 66 (1-2), 23-29
Plethodon dorsalis (large intestine)
Eurycea lucifuga (large intestine)
all from Equality Cave, southwest of Equality, Saline County, Illinois
- Cosmocercoides dukae* (Holl, 1928) Travassos, 1931
Lewis, P. D., jr., 1973, *Tr. Am. Micr. Soc.*, v. 92 (2), 286-287
Discus cronkhitei
Deroceras laeve
Stenotrema leai
Succinea ovalis
Triodopsis albolabris
Triodopsis multilinata
Cionella lubrica
all from Nebraska
- Cottocomephoronema* Layman, 1933
Arthur, J. R.; and Margolis, L., 1975, *Canad. J. Zool.*, v. 53 (6), 736-747
as syn. of *Haplonema* Ward and Magath, 1917
- Cottocomephoronema hamulatum* (Moulton, 1931) Sudarikov and Ryzhikov, 1952
Arthur, J. R.; and Margolis, L., 1975, *Canad. J. Zool.*, v. 53 (6), 736-747
as syn. of *Haplonema hamulatum* Moulton, 1931
- Cottocomephoronema problematica* Layman, 1933
Arthur, J. R.; and Margolis, L., 1975, *Canad. J. Zool.*, v. 53 (6), 736-747
as syn. of *Haplonema hamulatum* Moulton, 1931
- Cramispirura*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Oxyspirura*
key
- Crassicauda* Leiper & Atkinson, 1914, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Crassicaudinae
key
- Crassicauda* sp., illus.
Dailey, M. D.; and Perrin, W. F., 1973, *Fish. Bull., National Oceanic and Atmos. Admin.*, v. 71 (2), 455-471
Stenella cf. *S. longirostris* (abdominal muscle adjacent to mammary gland): eastern tropical Pacific

- Crassicauda-like [sp.]
Dailey, M. D.; and Perrin, W. F., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (2), 455-471
irreversible lesions in ventral skull region of spotted porpoise, incidence related to age of host
Stenella graffmani (air sinuses): eastern tropical Pacific
- Crassicaudinae Yorke & Maplestone, 1926
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Tetrameridae
key; key to genera
includes: Crassicauda; Placentonema
- Craterostomum
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Craterostomum
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Craterostomum acuticaudatum
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Craterostomum acuticaudatum
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Craterostomum acuticaudatum
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Craterostomum mucronatum
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Craterostomum mucronatum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Crenosoma hermani Anderson, 1962, illus. Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
description
Mustela erminea (lungs): Karelia
- Crenosoma petrowi Morozov, 1939
Craig, R. E.; and Borecky, R. A., 1976, Canad. J. Zool., v. 54 (5), 806-807
Martes pennanti (lungs): Ontario
- Crenosoma petrowi Marosow, 1939
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Martes martes: Pechora river basin
- Crenosoma petrowi Morozov, 1939
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes (lungs, trachea): Karelia
- Crenosoma schulzi Gagarin, 1956
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Meles meles (lungs): Karelia
- Crenosoma striatum
Isenbuegel, E., 1976, Prakt. Tierarzt, v. 57, Sondernummer, 21-27
Citarin, Laevamisol
Igel
- Crenosoma striatum (Zeder, 1800), illus.
Saupe, E., 1976, Vet.-Med. Nachr. (1), 91-96
Crenosoma striatum, hedgehog, tetramisole, good results
- Crenosoma vulpis
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Crenosoma vulpis (Duj., 1874)
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Alopes lagopus
Vulpes vulpes
all from Pechora river basin
- Crenosoma vulpis (Dujardin, 1845)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Canis familiaris
Vulpes fulva
all from insular Newfoundland
- Crenosoma vulpis
Stockdale, P. H. G.; and Smart, M. E., 1975, Research Vet. Sc., v. 18 (2), 178-181
Crenosoma vulpis, dogs (exper.), treatment with levamisole, diethylcarbamazine, thi-acetarsamide, and cyacetazide (from most to least effective), pulmonary lesions
- Crenosoma vulpis
Zeh, J. B.; Stone, W. B.; and Roscoe, D. E., 1977, N. York Fish and Game J., v. 24 (1), 91-93
red fox
gray fox
all from New York

- Cristiceps* Mawson, 1971
Mawson, P. M., 1976, Tr. Roy. Soc. South Australia, v. 100 (3), 121-123
as syn. of *Woodwardostrongylus* Wahid, 1964
- Cristitectus* Petter, 1970, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Crossocephalus viviparus* Linstow
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Equus burchelli (intestines): Kenya
- Crossophoridae (Baylis, 1920, subfam.) Hartwich, 1957
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridoidea
key; key to genera
includes: *Crossophorus*; *Dartevellenia*
- Crossophorus* Hemprich & Ehrenberg, 1828, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Crossophoridae
key
- Cruorifilaria* gen. n.
Eberhard, M. L.; Morales, G. A.; and Orihel, T. C., 1976, J. Parasitol., v. 62 (4), 604-607
Onchocercidae, tod: *C. tubero cauda* sp. n.
- Cruorifilaria tubero cauda* sp. n. (tod), illus.
Eberhard, M. L.; Morales, G. A.; and Orihel, T. C., 1976, J. Parasitol., v. 62 (4), 604-607
Hydrochoerus hydrochaeris (blood vessels of kidney, lungs, and heart, microfilaria in blood): Carimagua, Meta State, Colombia, South America
- Cruzia americana*
Prestwood, A. K.; Nettles, V. F.; and Farrell, R. L., 1977, Am. J. Vet. Research, v. 38 (4), 529-532
Didelphis marsupialis: Georgia
- Cruzia morleyi* Pearse, 1936
Caballero Deloya, J., 1974, An. Inst. Biol., Univ. Nac. Mexico, s. Zool., v. 45 (1), 45-50
as syn. of *Neocruzia morleyi* (Pearse, 1936)
Yamaguti, 1961
- Cryptaphelenchus minutus*
Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Ctenascarophis* gen. nov.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Rhabdochonidae, Rhabdochoninae
tod: *C. gastricus* sp. nov.
- Ctenascarophis* Mamaev, 1968, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Ctenascarophis gastricus* gen. et sp. nov. (tod), illus.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Auxis thazard
Euthynnus affinis
(stomach of all): all from South China Sea
- Cucullanellus minutus* (Rud., 1819)
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland
- Cucullanellus minutus*
Moeller, H., 1976, J. Marine Biol. Ass. United Kingdom, v. 56 (3), 781-785
Platichthys flesus: Kiel Fjord (western Baltic Sea)
- Cucullanidae
Ivashkin, V. M.; and Khromova, L. A., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 37-43
life cycles, taxonomy, brief review
- Cucullanus* Goeze, 1782, nec Mueller, 1777
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Goezia* Zeder, 1800
- Cucullanus* sp.
Rubertone, J. A.; and Hall, J. E., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 58-59
Pylodictus olivaris (intestine): Greenbrier River below Alderson, West Virginia
- Cucullanus annulatus* Margolis, 1960, illus.
Rehana, R.; and Bilqees, F. M., 1976, Agric. Pakistan, v. 26 (4), 1975, 521-528
Mystus cavasius (intestine): Kalri Lake, Sind area
- Cucullanus caballeroi* Petter, 1976
Petter, A. J.; Golvan, Y. J.; and Tcheprakoff, R., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 159-171
Eleotris pisonis
Gobiomorus maculatus
all from river of Guadeloupe
- Cucullanus cirratus*
Moeller, H., 1976, J. Marine Biol. Ass. United Kingdom, v. 56 (3), 781-785
Gadus morhua (intestine): Kiel Fjord (western Baltic Sea)
- Cucullanus diminutus*
Rehana, R.; and Bilqees, F. M., 1976, Agric. Pakistan, v. 26 (4), 1975, 521-528
- Cucullanus heterochrous* Rud., 1802
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland

- Cucullanus heterochrous*
Moeller, H., 1976, J. Marine Biol. Ass. United Kingdom, v. 56 (3), 781-785
intestinal helminths, elimination from host held in captivity, high rate of elimination of helminths unattached or slightly attached to host, lower elimination rate of helminths attached to host
Platichthys flesus (intestine): Kiel Fjord (western Baltic Sea)
- Cucullanus indentatus*
Rehana, R.; and Bilqees, F. M., 1976, Agric. Pakistan, v. 26 (4), 1975, 521-528
- Cucullanus lebedevi* E. Skriabina, 1966, illus.
Skriabina, E. S., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 169-182
description
Acipenser baeri (stomach, intestine): Yenisei and Lena Rivers
- Cucullanus theraponi*
Rehana, R.; and Bilqees, F. M., 1976, Agric. Pakistan, v. 26 (4), 1975, 521-528
- Cucullanus truttae* (Fabricius, 1794)
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
Silurus glanis (anterior portion of intestine): river Biebrza basin, Poland
- Cucullanus truttae* (Fabricius)
Lang, B. Z.; and Edson, S. A., 1976, J. Parasitol., v. 62 (1), 93
Rhinichthys osculus: Turnbull National Wildlife Refuge, Spokane County, Washington
- Cutifilaria* n. gen.
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 450-453
Onchocercinae
tod: *C. wenki* n. sp.
- Cutifilaria wenki* n. gen., n. sp. (tod), illus.
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 450-453
Cervus elaphus (intradermique; dos, ventre et cou): Schonbuch de Tubingen, Wurttemberg, Allemagne
- Cutifilaria wenki* Bain und Schulz-Key 1974, illus.
Schulz-Key, H., 1975, Tropenmed. u. Parasitol., v. 26 (3), 348-358
Filariidae spp., microfilariae infecting *Cervus elaphus*, morphological comparisons and localization patterns
Cervus elaphus (skin of trunk): southern Germany
- Cutifilaria wenki* Bain un Schulz-Key 1974
Schulz-Key, H., 1975, Tropenmed. und Parasitol., v. 26 (4), 494-498
development and distribution of adults and microfilariae intradermally and subcutaneously in Dama dama: Southern Germany
- Cyatholaimina* DeConinck, 1965
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Chromadorida
- Cyathospirura* Baylis, 1934, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirocercinae
key
- Cyathospirura dasyuridis*
Gregory, G. G.; and Munday, B. L., 1976, Austral. Vet. J., v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island
- Cyathospirura seurati* Gibbs, 1957, illus.
Quentin, J. C.; and Wertheim, G., 1975, Ann. Parasitol., v. 50 (1), 63-85
redescription, valid species
Acomys cahirinus
Gerbillus sp.
all from Israel
- Cyathostoma* Blanchard, 1849
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
Cyathostominae Nicoll, 1927
revision, keys to subgen. and species includes subgen.: *Cyathostoma* (Blanchard, 1849) subg. n.; *Ryzhikovistoma* subg. n.
- Cyathostoma* (Blanchard, 1849) subg. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
subgen. of *Cyathostoma*
key, key to species
diagnosis, tod: *Cyathostoma* (*Cyathostoma*) *lari* Blanchard, 1849 comb. n.
- Cyathostoma* (*Ryzhikovistoma*) sp. (Ali)
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* sp. 1
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (trachea, lungs): Florida
- Cyathostoma* sp. 2
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (trachea, lungs): Florida
- Cyathostoma* (*Ryzhikovistoma*) *americanum* Chapin, 1925 comb. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* (*Ryzhikovistoma*) *brantae* Cram, 1928 comb. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* (*Cyathostoma*) *bronchialis* (Muehling, 1884) Ghapin, 1925 n. comb.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
as syn. of *Cyathostoma* (*C.*) *variegatum* (Creplin, 1849) Chapin, 1925 n. comb.

- Cyathostoma bronchialis* (Muhlig, 1884) Chapin, 1925
Hernandez-Rodriguez, S.; et al., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 367-371
lesions and localization in *Anser anser* (traquea y bronquios): Parque Zoológico "Juan Barasona" de Córdoba, España
- Cyathostoma bronchialis*, *illus.*
Vasilev, I.; and Denev, I., 1972, Izvest. Tsentral. Khelminth. Lab., v. 15, 21-32
Cyathostoma bronchialis, development of eggs and larvae, importance of earthworms in life cycle in enhancing larval infectivity
Lumbricus terrestris (nat. and exper.)
L. rubellus
L. polyphemus
Bimastus tenuis
Octolasion rebeli
O. complanatum
O. lacteum
Allolobophora caliginosa
[*Anser anser*] (exper.)
[*Gallus gallus*] "
[*Meleagris gallopavo*] (exper.)
[*Anas platyrhynchos*] "
[*Alectoris graeca*] "
[*Numida meleagris*] "
[*Phasianus colchicus*] "
[Partridge] (exper.)
- Cyathostoma* (*Ryzhikovistoma*) *coscorobae* Chapin, 1925 comb. n. (tod of subgen.)
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* (*Cyathostoma*) *lari* Blanchard, 1849 comb. n. (tod of subgen.)
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma lari* Blanchard, 1849
Andrews, S. E.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 24-28
Corvus brachyrhynchos (nasal cavity): insular Newfoundland
- Cyathostoma lari* Blanchard, 1849, *illus.*
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
nematodes of *Larus canus* (cloaca, nasal and orbital cavities, trachea), age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Cyathostoma* (*Cyathostoma*) *phenisci* Baudet, 1937 comb. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma phenisci*
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (trachea, lungs): Florida
- Cyathostoma* (*Ryzhikovistoma*) *sarcidiornis* Kreis, 1953 comb. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* (*Cyathostoma*) *tadornae* Chatin, 1874 comb. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key
- Cyathostoma* (*Cyathostoma*) *variegatum* (Creplin, 1849) Chapin, 1925 comb. n., *illus.*
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
key, description
Syn.: *Cyathostoma* (*C.*) *bronchialis* (Muehling, 1884) Chapin, 1925 comb. n.
Pavo cristatus (trachea): India
- Cyathostomidae**
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Cyathostominae** Nicoll, 1927
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
host specificity, distribution, taxonomy, review
- Cyathostomum**
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Cyathostomum spp.**
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxbendazole
- Cyathostomum catinatum**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cyathostomum catinatum**
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum coronatum**, *illus.*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level

- Cyathostomum coronatum*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cyathostomum coronatum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cyathostomum coronatum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cyathostomum coronatum*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum labiatum, illus.*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cyathostomum labiatum*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cyathostomum labiatum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cyathostomum labiatum*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum pateratum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cyathostomum pateratum*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum tetracanthum*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cyathostomum labiatum*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum labratum, illus.*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cyathostomum labratum*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cyathostomum labratum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cyathostomum labratum*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cyathostomum labratum*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations

- Cyclodontostomum purvisi* (Adams, 1933), illus.
Bhaibulaya, M.; and Indrangarm, S., 1975,
Southeast Asian J. Trop. Med. and Pub. Health,
v. 6 (3), 391-394
Cyclodontostomum purvisi, accidental parasitism in man discovered during survey for incidence in rats, morphology, measurements man (feces)
Bandicota savilei (large intestine)
B. indica (large intestine)
all from Thailand
- Cyclodontostomum purvisi* (Adams, 1933)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus rattus argentiventer
R. bowersi
R. cremoriventer
R. mulleri
R. rajah subsp.
R. sabanus
R. whiteheadi
all from Malaysia
- Cyclodontostomum purvisi* (Adams, 1933), illus.
Varughese, G., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (1), 78-95
Cyclodontostomum purvisi, life cycle and developmental anatomy of free-living and parasitic stages
laboratory rats (exper.)
laboratory mice (exper.)
Rattus whiteheadi
R. muelleri
R. sabanus
(large intestine, caecum of all): all from Malaysia
- Cyclostrongylus* Johnston & Mawson
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
redefined, key to species
Syn.: *Oesophagonastes*
- Cyclostrongylus dissimilis*, to *Macropostrongyloides* [comb. not made]
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
- Cyclostrongylus gallardi* Johnston & Mawson, 1939
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
key
- Cyclostrongylus kartana* (Mawson 1955) [n. comb.]
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
key
Syns.: *Spirostrongylus kartana*; *Oesophagonastes kartana*: Mawson, 1965
- Cyclostrongylus leptos* (Mawson 1965) [n. comb.]
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
key, syn.: *Oesophagonastes leptos*
- Cyclostrongylus medioannulatus*
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
sp. inq.
- Cyclostrongylus parma* (Johnston & Mawson 1939) [n. comb.]
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
key, syns.: *Pharyngostrongylus parma*; *Spirostrongylus parma*: Mawson, 1955; *Oesophagonastes parma*: Mawson, 1965
- Cyclostrongylus wallabiae* Johnston & Mawson, 1939
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
synonymy, key
- Cyclozone Dogiel*, 1932, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Cylicobrachytus brevicapsulatus*, illus.
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicobrachytus brevicapsulatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocercus alveatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocercus catinatus*, illus.
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocercus catinatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocercus catinatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis

- Cylicocercus goldi*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocercus goldi*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocercus goldi*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocercus pateratus*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocercus pateratus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocercus pateratus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocyclus*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Cylicocyclus*
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Cylicocyclus* spp.
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Cylicocyclus ashworthi*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocyclus auriculatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocyclus brevicapsulatus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicocyclus elongatus, illus.*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocyclus elongatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocyclus elongatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocyclus elongatus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicocyclus elongatum* (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Cylicocyclus insigne, illus.*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level

- Cylicocycclus insigne*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocycclus insigne*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocycclus insigne*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicocycclus insigne*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicocycclus leptostomus*, *illus.*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocycclus leptostomus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocycclus leptostomus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicocycclus leptostomus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicocycclus nassatus*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicocycclus nassatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocycclus nassatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocycclus nassatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicocycclus nassatus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicocycclus nassatum* (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Cylicocycclus radiatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicocycclus radiatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicocycclus ultrajectinus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicodontophorus*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method

- Cylicodontophorus* spp.
Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxbendazole
- Cylicodontophorus bicoronatus*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicodontophorus bicoronatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicodontophorus bicoronatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicodontophorus bicoronatus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicodontophorus euproctus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicodontophorus euproctus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicodontophorus euproctus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicodontophorus mettami*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicodontophorus ultrajectinus*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicodontophorus ultrajectinus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicospirura*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (3), 29-58
subgen. of *Cylicospirura*
key
- Cylicospirura* Vevers, 1922, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (3), 29-58
Spirocercinae
key
Syn.: *Petrowospirura* Matschulsky, 1952
includes subgens.: *Cylicospirura*; *Gastrododus*; *Skrjabinocercina*
- Cylicospirura felineus*
Gregory, G. G.; and Munday, B. L., 1976, *Austral. Vet. J.*, v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island
- Cylicospirura skrjabini* Kozlov, Owsjukova et Radkewitch, 1964
Kozlov, D. P., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 71-78
Alopex lagopus: Pechora river basin
- Cylicostephanus*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Cylicostephanus* spp.
Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxbendazole
- Cylicostephanus asymmetricus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicostephanus asymmetricus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England

- Cylicostephanus calicatus*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicostephanus calicatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicostephanus calicatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicostephanus calicatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicostephanus calicatus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicostephanus goldi*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicostephanus goldi*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicostephanus hybridus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicostephanus longibursatus*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicostephanus longibursatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicostephanus longibursatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicostephanus longibursatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Cylicostephanus longibursatus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicostephanus minutus*
Braide, E. I.; and Georgi, J. R., 1974, Cornell Vet., v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicostephanus minutus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicostephanus minutus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicostephanus minutus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations

- Cylicostephanus minutus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicostephanus poculatus*
Braide, E. I.; and Georgi, J. R., 1974, *Cornell Vet.*, v. 64 (2), 233-239
equine cyathostomes, number of external leaf crown elements, unsuitable for generic classification but provides relatively constant character at species level
- Cylicostephanus poculatus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicostephanus poculatus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cylicostephanus poculatus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Cylicotetrapedon asymmetricus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Cylicotetrapedon asymmetricus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Cyrnea* Seurat, 1914, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key; synonymy
- Cyrnea*
Davidson, W. R.; Hon, L. T.; and Forrester, D. J., 1977, *J. Parasitol.*, v. 63 (2), 332-336
generic status discussed
- Cyrnea* (*Procyrnea*) sp.
Buscher, H. N.; and Tyler, J. D., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 108-111
Speotyto cunicularia (gizzard lining, gut mesentery): Oklahoma
- Cyrnea* (=Seurocyrnea) sp. [of] Maxfield et al. (1963)
Davidson, W. R.; Hon, L. T.; and Forrester, D. J., 1977, *J. Parasitol.*, v. 63 (2), 332-336
as syn. of *Cyrnea* (*Cyrnea*) neeli sp. n.
- Cyrnea* sp.
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 119-127
Meleagris gallopavo (gizzard lining): Florida
- Cyrnea* spp.
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, *Proc. 3. National Wild Turkey Symp.*, 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Cyrnea* (*Cyrnea*) colini Cram 1927, illus.
Davidson, W. R.; Hon, L. T.; and Forrester, D. J., 1977, *J. Parasitol.*, v. 63 (2), 332-336
number and arrangement of caudal papillae redescribed
"reexamination of the specimens reported by Hon et al. (1975) as *C. eurycerca* revealed them to be *C. colini*."
Meleagris gallopavo: Clarke County, Alabama; Crittenden and Desha counties, Arkansas; Alachua, Flagler, Gadsden, Leon, Taylor, Wakulla, Glades, and Osceola counties, Florida; Grady County, Georgia; Bolivar and Tunica counties, Mississippi; Stafford County, Virginia
Colinus virginianus: Leon County, Florida
- Cyrnea* (*Cyrnea*) eurycerca
Davidson, W. R.; Hon, L. T.; and Forrester, D. J., 1977, *J. Parasitol.*, v. 63 (2), 332-336
"reexamination of the specimens reported by Hon et al. (1975) as *C. eurycerca* revealed them to be *C. colini*."
- Cyrnea eurycerca*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 119-127
Meleagris gallopavo (gizzard lining): Florida
- Cyrnea eurycerca*
Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Cyrnea* (*Cyrnea*) neeli sp. n., illus.
Davidson, W. R.; Hon, L. T.; and Forrester, D. J., 1977, *J. Parasitol.*, v. 63 (2), 332-336
Syn.: *Cyrnea* (=Seurocyrnea) sp. [of] Maxfield et al. (1963)
Meleagris gallopavo (proventricular-gizzard isthmus): Clarke County, Alabama; Gadsden, Glades, Leon, Osceola, Taylor, and Wakulla counties, Florida

- Cyrnea seurati*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Cyrneinae* gen. sp., larvae
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Adesmia planidorsis
Pachyscelis laevicollis
Prosodes bactriana
all from Tadzhik SSR [and/or] Uzbek SSR
- Cyrtosomum heyneani*
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus graciosus (cecum): Great Basin and Upper Colorado Plateau, Utah
- Cyrtosomum penneri*
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus graciosus
Sceloporus magister
(cecum of all): all from Great Basin and Upper Colorado Plateau, Utah
- Cyrtosomum readi*
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus graciosus (cecum)
Sceloporus magister (cecum and small intestine)
all from Great Basin and Upper Colorado Plateau, Utah
- Cystidicola* Fischer, 1798, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key; synonymy
- Cystidicola* sp. larvae
Mudry, D. R.; and Anderson, R. S., 1977, J. Fish Biol., v. 11 (1), 21-33
Prosopium coulteri: Yoho National Park, Canada
- Cystidicola farionis* Fischer, 1798
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Salmo trutta (swim bladder): Loch Leven, Scotland
- Cystidicola farionis* Fischer, illus.
Otto, F.; and Koerting, W., 1973, Vet. Med. Rev. (2), 99-106
endoparasites, behavioral changes in infected rainbow trout (swim bladder), post mortem findings: fish farm, South Germany
- Cystidicola farionis* Fischer, illus.
Reichenbach-Klinke, H. H., 1975, Fisch u. Umwelt (1), 113-121
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Cystidicola stigmatura* (Leidy, 1886)
Mudry, D. R.; and Anderson, R. S., 1977, J. Fish Biol., v. 11 (1), 21-33
Salmo gairdneri: Jasper National Park, Canada
Salvelinus namaycush: Jasper and Waterton Lakes National Parks, Canada
Prosopium williamsoni: Jasper National Park, Canada
- Cystidicola stigmatura* Skinker, 1931
Mudry, D. R.; and McCart, P. J., 1976, J. Fish. Research Bd. Canada, v. 33 (2), 271-275
Salvelinus alpinus (swim bladder): Alaska; Yukon
- Cystidicolidae (Skrjabin, 1946, subfam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematoidea
key; key to genera
includes: *Crisitectus*; *Metabronema*; *Salvelinema*; *Cystidicoloides*; *Spinitectus*; *Ctenascrophis*; *Cyclozone*; *Pseudoproleptus*; *Ascarophis*; *Parascrophis*; *Cystidicola*; *Spinitectoides*
- Cystidicolidae
Chabaud, A. G.; and Krishnasamy, M., [1976], Ann. Parasitol., v. 50 (6), 1975, 813-820
Trichospirura, should be placed in Rhabdochoniidae, evolutionary position, host range, osmoexcretory apparatus, relations between Rhabdochoniidae and Cystidicolidae
- Cystidicoloides* Skinker, 1931, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
Syn.: *Sterliadochona* Skrjabin, 1946
- Cystidicoloides* sp.
Arvy, L.; and Sowa, R., 1976, Ann. Parasitol., v. 51 (1), 111-120
Ephemera danica: region de Cracovie, Pologne.
- Cystidicoloides salvelini*
Lockard, L. L.; Parsons, R. R.; and Schaplow, B. M., 1975, Great Basin Nat., v. 35 (4), 442-448
Salmo trutta (upper digestive tract), relationship of incidence and intensity of nematode infection to age and sexual maturity of host, higher infection rate in sexually mature trout due to aggressive feeding behavior: streams in southern and western Montana
- Cystidicoloides tenuissima* Zeder, 1800
Kakacheva-Avramova, D., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 89-107
Salmo trutta morpha fario (stomach): River Tundzha
- Cystidicoloides tenuissima* (Zeder, 1800)
Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 87-110
Salmo trutta morpha fario (stomach, intestine): Balkan Mountain river(s)

- Cystocaulus**
Nemeseri, L., 1976, Magy. Allat. Lapja, v. 98, v. 32 [i. e. 31] (7), 459-461
Dictyocaulus filaria, trichostrongylids, Fasciola hepatica, sheep, successful treatment with combination of diamphenetide and tetramisole; no effect on Protostrongylidae
- Cystocaulus**
Paul, I.; et al., 1975, Lucrar. Stiint. Inst. Agron. I. Ionescu Brad Iasi, II. Zootehn., Med. Vet., 71-72
Protostrongylus, Muellerius, Cystocaulus, bronchopneumonia, adult sheep, morphopathology, result of association of several species
- Cystocaulus ocreatus**
Aller, B.; and Aller, J. M., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 337-346
nematodes of sheep lungs, lower incidence of fungi in parasitized lungs: Leon
- Cystocaulus ocreatus, illus.**
Casarosa, L.; Lugetti, G.; and Orlandi, M., 1975, Ann. Fac. Med. Vet. Pisa, v. 27, 1974, 87-90
Cystocaulus ocreatus, first stage larvae sensitized with immune serum, adherence reaction with normal guinea pig peritoneal macrophages
- Cystocaulus ocreatus**
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (7), 129
Dictyocaulus filaria, Protostrongylus rufescens, and Cystocaulus ocreatus in sheep, fenbendazole effectively eliminated lung-worms
- Cystocaulus ocreatus**
Ferreira, L. D. B. B., 1975, Rev. Portug. Cien. Vet. (433-434), v. 70, 25-39
broncho-pulmonary nematodes of sheep, life cycles, epizootiology, instructive review for veterinarians
- Cystocaulus ocreatus**
Melikov, Iu. F., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 80-83
distribution in sheep of sheep farms of Kura-Araksinsk lowland, Azerbaidzhan
- Cystocaulus ocreatus**
Ramisz, A.; Urban, E.; and Piechocki, B., 1975, Med. Wet., v. 31 (11), 677-679
Protostrongylidae of sheep, tetramisole effective against all species except Muellerius capillaris
- Cystocaulus ocreatus**
Rojo Vazquez, J., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 51-101
Protostrongylinae, frequency in ovine lungs, primary nodes are infected by bacteria at a lower rate than non-parasitized lung areas: municipal slaughterhouse, Leon
- Cystocaulus ocreatus, illus.**
Sumnaliev, P., 1973, Izvest. Tsentral. Khel-mint. Lab., v. 16, 161-167
Neostongylus linearis, Muellerius capillaris, Cystocaulus ocreatus, Protostrongylus sp., differential diagnosis of infective larvae based on biometric studies
Helicella obvia
Zebrina detrita
all from Bulgaria
- Cystocaulus ocreatus (Railliet et Henry, 1907)**
Mikacic, 1939, illus.
Svarc, R., 1977, Biologia, Bratislava, s. B, Zool. (3), v. 32 (8), 575-584
Cystocaulus ocreatus, penetration of first stage larvae into snails, localization, morphological changes during maturation, moulting
Cepaea vindobonensis
Succinea putris
(all exper.) (mucous tubular glands of foot (typical site); penis sacwall, liver (atypical, accidental sites) of all)
- Cystocaulus ocreatus**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date

- Dacnitis truttae* (Fabricius, 1794)
Fagerholm, H.-P., 1976, Norwegian J. Zool.,
v. 24 (4), 466 [Abstract]
Finland
- Dacnitoidea cotylophora* Ward & Magath, 1916
Beacham, B. E.; and Haley, A. J., 1976, Proc.
Helminth. Soc. Washington, v. 43 (2), 232-233
Morone americana (intestine): Chesapeake
Bay
- Dacnitoidea cotylophora* (Ward and Magath, 1917)
Cooper, C. L.; Ashmead, R. R.; and Crites,
J. L., 1977, Proc. Helminth. Soc. Washington,
v. 44 (1), 96
prevalence, comparison with previous years
Perca flavescens (intestine): western Lake
Erie
- Dacnitoidea cotylophora*
Gruninger, T. L.; Murphy, C. E.; Britton, J.
C., 1977, Southwest. Nat., v. 22 (4), 525-535
Ictalurus punctatus (intestine): Eagle
Mountain Lake, Texas
- Dartevellenia*
Gibson, D. I.; and Taylor, A. L., 1976, Para-
sitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon
taxonomic significance and function
- Dartevellenia Ezzat*, 1954
Hartwich, G., 1974, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
Crossophoridae
key
- Dasyostrostrongylus Travassos*, 1935
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Daubneyia*
Schmidt, G. D.; and Kuntz, R. E., 1975, Proc.
Oklahoma Acad. Sc., v. 55, 163-165
"The finding of a species lacking a cervical
groove in this subgenus supports the re-
jection of *Daubneyia* as a valid generic
concept!"
- Decorataria Sobolev*, 1949
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Synquaria* Gilbert, 1927
- [*Delafondia*] *deliafondii*
Kadyrov, N. T., 1973, Vestnik Sel'skokhoz.
Nauki Kazakhstana (11), 68-70
[*Delafondia*] invasive larvae cultured from
uterine eggs, experimental infection of colts
- Delafondia*
Reinecke, R. K.; and le Roux, D. J., 1972, J.
South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys
and modified critical tests on horses using
mebendazole, highly effective
- Delafondia* (*Strongylus*) *vulgaris*
Avellini, G.; and Vitellozzi, G., 1977, Clin.
Vet., Milano, v. 100 (4), 268-271
Delafondia vulgaris, colts, two cases of
sudden death owing to hemorrhage, verminous
mesenteric aneurysm
- Delafondia vulgaris*
Loseva, N. G., 1975, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 25, 95-97
Alfortia edentatus, *Delafondia vulgaris*,
histochemistry of intestine, low glycogen
content related to blood feeding; quantity
and distribution of nucleic acids
- Delafondia vulgaris*
de Matos, P. F.; and Costa, J. O., 1976, Arq.
Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2),
173-180
gastrointestinal helminths, horses, levami-
sole, haloxon, crufomate, anthelmintic effi-
ciency
- Delafondia vulgaris* (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am.
Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Delafondiosis*
Kadyrov, N. T.; and Suleimenova, F. Z., 1975,
Vestnik Sel'skokhoz. Nauki Kazakhstana (12),
91-93
delafondiasis, infected and uninfected
horses, blood levels of lactic acid, serum
activity of alanine-aminotransferase, aspar-
tate-aminotransferase and aldolase
- Delamurella Gubanov* 1952
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
key
- Delicata Travassos*, 1935
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Delicata perronae* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and
Cassone, J., 1977, Bull. Mus. National Hist.
Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, di-
delphic forms of *Trichostrongyloidea* more
abundant than monodelphic ones
Tamandua tetradactyla (partie moyenne de
l'intestin): region de Belem, Province de
Para, Bresil
- Delicata soyeriae* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and
Cassone, J., 1977, Bull. Mus. National Hist.
Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, di-
delphic forms of *Trichostrongyloidea* more
abundant than monodelphic ones
Tamandua tetradactyla (deuxieme moitie de
l'intestin): region de Belem, Province de
Para, Bresil

- Dentomermis* gen. nov.
Rubtsov, I. A.; and Polishchuk, V. V., 1975, *Gidrobiol. Zhurnal*, v. 11 (2), 95-97
Mermithidae
tod: *D. markevitschi* sp. n.
- Dentomermis markevitschi* gen. et sp. n. (tod),
illus.
Rubtsov, I. A.; and Polishchuk, V. V., 1975, *Gidrobiol. Zhurnal*, v. 11 (2), 95-97
host unknown: Ukrainian SSR, south of city of Chuguev
- Deraiphoronema freitaslenti* Yeh, 1957
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
as syn. of *Bostrichodera freitaslenti* (Yeh, 1957) n. comb.
- Dermatoxys* sp.
Jacobson, H. A.; and Kirkpatrick, R. L., 1974, *J. Wildlife Dis.*, v. 10 (4), 384-391
comparison of selected physiological measurements in untreated parasitized cottontail rabbits and those treated with 1-tetramisole hydrochloride and 2,2-dichlorovinyl, dimethyl phosphate: Montgomery County, Virginia
- Dermatoxys* sp., illus.
Seese, F. M., 1973, *Am. Midland Naturalist*, v. 89 (2), 257-265
key
Ochotona p. princeps (caecum, large intestine): St. Joe Baldy Mountain, Benewah County, Idaho
- Dermatoxys talkeetnarius* [sic] Akhtar (1956)
Seese, F. M., 1973, *Am. Midland Naturalist*, v. 89 (2), 257-265
key
- Desmidocerca aerophila*
Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Desmidocercella* (D.) skrjabini Guschanskaja, 1949
Aleksiev, V. M.; and Smetanina, Z. B., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 97-104
Phalacrocorax pelagicus
P. ussuriensis
all from Rimsko-Korsakov islands
- Desmodorida* DeConinck, 1965
Maggenti, A. R., 1976, *Organ. Nematodes* (Croll), 1-10
Chromadoria
includes: *Draconematina*
- Desmoscolecida* Filipjev, 1934
Maggenti, A. R., 1976, *Organ. Nematodes* (Croll), 1-10
Chromadoria
- Desportesius* (Chabaud & Campana, 1949, subgen.)
Skrjabin et al., 1965, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Diaphanocephaloidea*
Chabaud, A. G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (1), 6-17
Strongylida
key
- Dicheilonema ciconiae* (Schrank, 1788)
Gundlach, J. L., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 83-89
Ciconia nigra (pectoral muscles): Lublin Palatinate
- Dichelyne robusta* (Van Cleave and Mueller, 1932)
Baker, J. C.; and Crites, J. L., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 37-39
Ictalurus punctatus (intestines): island region of western Lake Erie
- Dictyocaulidae*
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongyloidea
- Dictyocaulosis*
Epel'dimov, L. S.; and Shchelkanov, K. G., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 116-122
dictyocaulosis, sheep treated with anthelmintics, histological changes
- Dictyocaulosis*
Melikov, Iu. F., 1976, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (6), 81-86
distribution in sheep, cattle and buffalo: regions of Kura-Araksin lowland
- Dictyocaulosis*
Ohshima, K.; et al., 1976, *J. Fac. Agric., Iwate Univ.*, v. 13 (2), 161-176
viral disease in cattle previously or concurrently having fascioliasis, dictyocaulosis, and/or piroplasmiasis, histopathological study of mucosal lesions, differential diagnosis: Japan
- Dictyocaulus*
Fudalewicz-Niemczyk, W.; et al., 1975, *Med. Wet.*, v. 31 (11), 666-668
sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Dictyocaulus*
Guarino, C.; and Rivellini, P., 1972, *Atti Soc. Ital. Sc. Vet.*, v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- [*Dictyocaulus*] *diktiokauli*
Panchin, O. G.; et al., 1975, *Veterinariia*, Kiev (40), 100-104
helminths and coccidia, sheep, seasonal incidence on pastures, measures for control: Kalanchats'k region, Kherson oblast

- Dictyocaulus**
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Dictyocaulus spp.**
Euzeby, J., 1977, *Rev. Med. Vet.*, Toulouse, v. 128 (5), 605-622
respiratory strongylosis of domestic ruminants, review
- Dictyocaulus sp.**
Graves, I. L.; Adams, W. H.; and Pyakural, S., 1975, *Am. J. Vet. Research*, v. 36 (6), 843-846
Babesia bigemina, *Bos grunniens* moved from high to low altitude and challenged with influenza A viruses, hemolytic anemia, possible explanations, death due to *Fasciola hepatica* and *F. gigantica*, incidental finding of *Bu-nostomum* sp., *Trichuris* sp., *Neoscaris vitulorum*, *Dictyocaulus* sp., coccidia, some reasons for poor survival of yaks at low altitude: Nepal
- Dictyocaulus sp.**
Kirton, A. H.; et al., 1976, *N. Zealand Vet. J.*, v. 24 (4), 59-64
Dictyocaulus sp., lambs, seasonal incidence of enzootic pneumonia, effect on growth, little economic significance in present flock
- Dictyocaulus sp.**, probably *D. viviparus*
Low, W. A., 1976, *Canad. Field-Naturalist*, v. 90 (2), 189-191
Rangifer tarandus caribou (lung alveoli, bronchioles): Tweedsmuir Provincial Park, British Columbia
- Dictyocaulus sp.**
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Alcelaphus buselaphus cokei (nasal cavity) *Equus burchelli* (bronchi)
all from Kenya
- Dictyocaulus arnfieldi**
Dooley, D. E.; and King, E. M., 1977, *J. Zoo Animal Med.*, v. 8 (1), 16-17
Dictyocaulus arnfieldi, Hartmann's zebra (bronchi and bronchioles), possible reservoir host: Lion Country, Laguna Hills, California
- Dictyocaulus arnfieldi, illus.**
Rickard, M. D.; and James, D. E., 1976, *Austral. Vet. J.*, v. 52 (10), 487-488 [Letter]
Dictyocaulus arnfieldi, horses, mebendazole: Australia, imported from New Zealand
- Dictyocaulus arnfieldi (Cobbold 1884)**
Round, M. C., 1976, *Vet. Rec.*, v. 99 (20), 393-395
horses, donkeys, life cycle, prepatency, patency, resistance, pathogenicity, pathology, experimental infections, diagnosis, treatment, review
- Dictyocaulus arnfieldi**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Dictyocaulus (M.) eckerti** Skrjabin, 1931
Ianchev, I., 1973, *Izvest. Tsentral. Khelmin. Lab.*, v. 16, 205-220
Capreolus capreolus (bronchi): southern Bulgaria
- Dictyocaulus filaria**
Al-Sammarræ, S. A.; and Sewell, M. M. H., 1977, *Research Vet. Sc.*, v. 23 (3), 336-339
Dictyocaulus filaria, Blackface sheep, infection levels, seasonal distribution on lowland farm: south east Scotland
- Dictyocaulus filaria**
Al-Sammarræ, S. A.; and Sewell, M. M. H., 1977, *Research Vet. Sc.*, v. 23 (3), 383-384
Dictyocaulus filaria, Soay and Blackface sheep, comparison of resistance, Soay sheep more susceptible
- Dictyocaulus filaria (Rudolphi, 1809)**
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (trachea, bronchi): vicinity of Nowy Targ, Carpathian Mountains
- Dictyocaulus filaria (Rudolphi 1809), illus.**
Casarosa, L., 1976, *Ann. Fac. Med. Vet.*, Univ. Pisa, v. 28, 1975, 53-63
Dictyocaulus filaria, guinea pigs (exper.), 3 increasing doses followed by challenge, histopathology of mesenteric and bronchial lymph nodes, trapped larvae surrounded by leucocytes and macrophages
- Dictyocaulus filaria, illus.**
Casarosa, L.; et al., 1976, *Ann. Fac. Med. Vet.*, Univ. Pisa, v. 28, 1975, 71-77
Dictyocaulus filaria, third stage larvae, sensitized with immune sera, in vitro adherence reaction with eosinophils and pyroninophil cells from guinea pigs immunized with *D. filaria* somatic metabolic antigen
- Dictyocaulus filaria**
Casarosa, L.; and Lugetti, G., 1972, *Parassitologia*, v. 14 (1), 71-72
Dictyocaulus filaria, third stage larvae sensitized in immune serum, adherence reaction with guinea pig peritoneal macrophages

- Dictyocaulus filaria*, illus.
Casarosa, L.; Lugetti, G.; and Favati, V., 1974, Ann. Fac. Med. Vet. Pisa, v. 26, 1973, 373-384
adherence reaction between infective *Dictyocaulus filaria* larvae (sensitized with immune guinea pig serum) and guinea pig peritoneal macrophages; mixed antiglobulin reaction between infective *D. filaria* larvae and sheep red blood cells previously sensitized with photo-oxidized guinea pig anti-serum; adherence reaction induced by gamma globulin combining with cuticle of parasite
- Dictyocaulus filaria*
Denev, I.; and Kolev, M., 1975, Vet. Med. Nauki, v. 12 (1), 97-100
Dictyocaulus filaria, lambs (exper.), diagnosis, immunofluorescence
- Dictyocaulus filaria*
Denev, I.; Komandarev, S.; and Mikhov, L., 1975, Vet. Med. Nauki, v. 12 (5), 19-26
Dictyocaulus filaria, comparative study of somatic and functional antigens, disk electrophoresis
- Dictyocaulus filaria*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Dictyocaulus filaria*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Dictyocaulus filaria*
Ershov, V. S.; et al., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 343-348
anaphylactic shock in guinea pigs after sensitization with free-living or plant-parasitic nematodes and challenge with various helminth antigens indicates antigenic components in common; intradermal tests using antigen from free-living nematode in cases of ascariasis, trichinellosis, and cysticercosis; possible use of free-living nematode to immunize against dictyocaulosis and ascariasis
- Dictyocaulus filaria*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (7), 129
Dictyocaulus filaria, *Protostrongylus rufescens*, and *Cystocaulus ocreatus* in sheep, fenbendazole effectively eliminated lung-worms
- Dictyocaulus filaria*
Ferreira, L. D. B. B., 1975, Rev. Portug. Cien. Vet. (433-434), v. 70, 25-39
broncho-pulmonary nematodes of sheep, life cycles, epizootiology, instructive review for veterinarians
- D[ictyocaulus] filaria*
Fromunda, V., 1976, Rev. Crest. Animalelor, v. 26 (3), 86-90
helminthic diseases, sheep, prevention during grazing
- Dictyocaulus filaria*
Gallie, G. J.; and Nunns, V. J., 1976, J. Helminth., v. 50 (2), 79-89
Dictyocaulus filaria, different seasons, development and survival of free-living larvae on pasture, transmission of infection between lambs: North-East England
- Dictyocaulus filaria*
Gallie, G. J.; Thomas, R. J.; and Nunns, V. J., 1977, Research Vet. Sc., v. 22 (2), 251-256
Dictyocaulus filaria, sheep, prevalence and intensity of infection, seasonal pattern and weather conditions, sources of infection for lambs: north east England
- Dictyocaulus filaria*
Goda, Fawzy F. M., 1974, Bull. Epizoot. Dis. Africa, v. 22 (1), 75-78
sheep: Benghazi, Libya
- Dictyocaulus filaria*
Groeneveld, H. T.; and Reinecke, R. K., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 285-297
non-parametric statistical method for comparing worm burdens in two groups of sheep, application in interpreting results of anthelmintic tests
- Dictyocaulus filaria*
Grzywinski, L.; et al., 1975, Medycyna Wet., v. 31 (9), 524-526
sheep, Nilverm by injection for control
- Dictyocaulus filaria*
Jabir, M.H.; Kadhim, J.K.; and Singh, K.S., 1977, Vet. Parasitol., v. 3 (1), 67-73
Dictyocaulus filaria, Awasi sheep dosed at about 200 larvae/kg body weight are good producers of live larvae
- Dictyocaulus filaria*
Jedreas, A., 1976, Med. Wet., v. 32 (2), 73-75
Dictyocaulus filaria, sheep, immunization by vaccine of normal or X-irradiated larvae; animals vaccinated twice, highly resistant to challenge dose; however, vaccination by high doses of normal larvae sometimes caused death
- Dictyocaulus filaria*
Jedreas, A., 1976, Med. Wet., v. 32 (9), 525-529
Dictyocaulus filaria, sheep, immunization with irradiated and non-irradiated larvae, challenged with non-irradiated larvae
- Dictyocaulus filaria*
Jørgensen, R. J., 1975, Vet. Parasitol., v. 1 (1), 61-67
isolation of infective *Dictyocaulus* larvae from herbage, 3 steps: 1. isolation of debris from herbage 2. isolation of larvae from debris (migration in agar gel containing bile) 3. staining to differentiate from free-living nematodes

- Dictyocaulus filaria*
Kadhim, J. K.; Jabbar, M. H.; and Altaif, K. I., 1972, *Research Vet. Sc.*, v. 13 (6), 597-599
Dictyocaulus filaria, goats (exper.), comparison between efficacy of tetramisole vs. morantel/diethylcarbamazine
- Dictyocaulus filaria*
Kassai, T.; et al., 1973, *Isotopes and Radiation Parasitol.* III, 51-60
Dictyocaulus filaria, sheep and goats, laboratory investigation of immunity induced by irradiated vs. non-irradiated larvae, marked resistance following infection with normal larvae, irradiated larvae failed to prevent re-infection in sheep but did in 3 of 5 goats; sheep, field trials of radiation-attenuated vaccine compared to efficacy of anthelmintic treatment, no protection shown in vaccinated sheep
- Dictyocaulus filaria*
Kelly, J. D.; et al., 1975, *Research Vet. Sc.*, v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Dictyocaulus filaria*
Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Dictyocaulus filaria*
Kozdon, O.; and Zajicek, D., 1976, *Vet. Med.*, Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Dictyocaulus filaria*
Malik, P. D.; and Gupta, R. P., 1975, *Haryana Agric. Univ. J. Research*, v. 5 (1), 58-65
phenothiazine + piperazine adipate, methyridine, tetramisole, thiabendazole, drug efficacy, sheep: University Farm, Hissar
- D[*dictyocaulus*] *filaria*, *illus.*
Movsesijan, M.; and Lalic, R., 1971, *Acta Parasitol. Iugoslavica*, v. 2 (2), 57-65
D[*dictyocaulus*] *filaria*; sheep, diagnosis, localization of antigen-antibody complex
- Dictyocaulus filaria*
Movsesijan, M.; and Lalic, R., 1973, *Isotopes and Radiation Parasitol.* III, 33-42
Dictyocaulus filaria, sheep, radioimmunoassay using radioactively labelled immunoglobulins and whole parasites (12-14 days old) as antigen
- Dictyocaulus filaria*
Nemeseri, L., 1976, *Magy. Allat. Lapja*, v. 98, v. 32 [i. e. 31] (7), 459-461
Dictyocaulus filaria, trichostrongylids, *Fasciola hepatica*, sheep, successful treatment with combination of diamphenetide and tetramisole; no effect on Protostrongylidae
- Dictyocaulus filaria*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Ovis aries: Chile
- Dictyocaulus filaria*
Polley, L.; and Hoerning, B., 1977, *Rev. Suisse Zool.*, v. 84 (3), 675-680
Rupicapra rupicapra (tracheas and larger bronchi): Switzerland
- Dictyocaulus filaria*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Dictyocaulus filaria*
Ramachandran, C. P., 1975, *Kajian Vet.*, v. 7 (1), 31-38
helminths, immunization with radiation attenuated vaccines, review
- Dictyocaulus filaria*
Rose, J. H., 1971, *Symposia Brit. Soc. Parasitol.*, v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Dictyocaulus filaria*
Rybnikar, A., 1975, *Acta Vet. Brno*, v. 44 (4), 385-391
Dictyocaulus filaria, experimentally infected lambs used as donors for obtaining larvae for preparation of radiation vaccine; amount of larvae excreted dependent upon body mass and age of lambs, total dose of larvae and season of infestation
- Dictyocaulus filaria*
Sharma, D. N.; Rajya, B. S.; and Dwivedi, J. N., 1977, *Indian J. Animal Sc.*, v. 45 (5), 1975, 275-281
Dictyocaulus filaria in Jaagziekte and Maedi-infected sheep and goats (exper.), no effect of *Dictyocaulus filaria* in precipitating either disease
- Dictyocaulus filaria*
Shnain, A. H.; et al., 1973, *Vet. Rec.*, v. 92 (19), 499-500, 501
cannulation of thoracic duct for recovery of *Dictyocaulus filaria* from lymph, surgical technique, sheep
- Dictyocaulus filaria* (Rudolphi, 1809)
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Ovis aries: insular Newfoundland

- Dictyocaulus filaria*
Taranik, K. T., 1975, *Visnik Sil's'kogospod. Nauki* (7) (212), 101-102
Dictyocaulus filaria, pathogenicity studies of experimental bovine infections, light infections producing no clinical signs, heavy infections (50,000 larvae) producing pneumonia and emphysema
- Dictyocaulus filaria*
Taranik, K. T.; and Antonov, V. S., 1975, *Veterinariia*, Kiev (42), 95-98
Dictyocaulus filaria, calves immunized with nonspecific gammaglobulin, lambs immunized with anti-*Dictyocaulus* immunoglobulin from cattle, changes in blood proteins after infection
- Dictyocaulus filaria*
Tewari, H. C.; Dhar, D. N.; and Singh, K. S., 1973, *Isotopes and Radiation Parasitol.* III, 43-50
Dictyocaulus filaria, sheep, incidence, laboratory and field trials with gamma-irradiated vaccine, high degree of protection conferred, no correlation between antibody response and ability to withstand challenge: Kashmir, India
- Dictyocaulus filaria*
Theodorides, V. J.; Nawalinski, T.; and Chang, J., 1976, *Am. J. Vet. Research*, v. 37 (12), 1515-1516
gastrointestinal nematodes, *Moniezia* spp., sheep, albendazole highly effective
- Dictyocaulus filaria*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Dictyocaulus filaria*
Tomanek, J., 1974, *Acta Vet. Brno*, v. 43 (1), 47-52
guinea-pigs orally infected with irradiated and non-irradiated *Dictyocaulus filaria* larvae, development of resulting larvae recovered from lungs; decreased larval counts with increased X-ray exposure, greater survival of female worms, implications for use in immunization
- Dictyocaulus filaria*
Tomanek, J.; and Franek, M., 1975, *Acta Vet. Brno*, v. 44 (4), 393-399
Dictyocaulus filaria, presence of complement-fixing antibodies in immunoglobulin fractions of sera from lambs, during primary and secondary response to infection with non-irradiated and X-irradiated larvae
- Dictyocaulus filaria*
Tomanek, J.; and Sedlacek, M., 1975, *Acta Vet. Brno*, v. 44 (1-2), 123-129
Dictyocaulus filaria larvae, X-irradiation, radiosensitivity, larval concentration, fecal contamination, temperature, salinity, aeration
- Dictyocaulus filaria*
Turton, J. A., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (2), 207-213
Dictyocaulus filaria, lambs, levamisole, efficacy administered orally or subcutaneously at rate of 7.5 mg per kg against different life cycle stages
- Dictyocaulus filaria*
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, *Vet. Glasnik*, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Dictyocaulus filaria*
Wallnoefer, E., 1977, *Wien. Tierarztl. Monatsschr.*, v. 64 (4), 129-131
sheep parasites, Mebenvet, good results when treatment was repeated after 14 days: Austria
- Dictyocaulus filaria*
Zajicek, D.; and Kozdon, O., 1977, *Veterinarstvi*, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Dictyocaulus magna* (Monnig, 1932)
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, *Onderstepoort J. Vet. Research*, v. 42 (1), 29-31
Antidorcas marsupialis: Bontebok National Park, Swellendam
Damaliscus dorcas dorcas (feces): captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Dictyocaulus viviparus*
Baxter, J. T.; and Allan, D., 1977, *Vet. Rec.*, v. 101 (19), 394 [Letter]
Dictyocaulus viviparus, persistence of larvae on pasture herbage October to April, possibly due to gradual disintegration of feces: Northern Ireland
- Dictyocaulus viviparus*
Benitez-Usher, C.; Armour, J.; and Urquhart, G. M., 1976, *Vet. Parasitol.*, v. 2 (2), 209-222
Dictyocaulus viviparus, young calves, efficacy of immunization with Dictol below commercial recommendation of 8 weeks of age, concluded that it may be practical to vaccinate milk-fed and suckling calves from 3-4 weeks of age
- Dictyocaulus viviparus*
Bergstrom, R. C., 1975, *J. Wildlife Dis.*, v. 11 (1), 40-44
Dictyocaulus viviparus in *Cervus canadensis*, seasonal distribution, fecal analyses useful in prevalence survey: Teton County, Wyoming

- Dictyocaulus viviparus*
 Braca, G.; and Arispici, M., 1974, Atti Soc. Ital. Sc. Vet., v. 28, 708-710
Dictyocaulus viviparus, pathology of parasitic bronchitis in cattle, possible relationships to non-parasitic respiratory infections
- Dictyocaulus viviparus*
 Breeze, R. G.; et al., 1974, J. Comp. Path., v. 84 (4), 577-588
 no evidence that fog fever occurs as result of hypersensitivity reaction in sensitized cattle to oral reinfection with *D. viviparus* larvae
- Dictyocaulus viviparus*
 Breeze, R. G.; et al., 1975, Folia Vet. Latina, v. 5 (1), 95-128
Dictyocaulus viviparus, adult cattle, reinfection syndrome, respiratory distress, pulmonary lymphoid nodules: Britain
- Dictyocaulus viviparus*
 Christensson, D.; and Reh binder, C., 1975, Nord. Vet.-Med., v. 27 (10), 496-498
 gastrointestinal parasites of reindeer calves, none found in first month of life, increasing infection with age: Norrbotten
- Dictyocaulus viviparus*
 Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
 gastrointestinal nematodes of heifers, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Dictyocaulus viviparus*
 Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 518-525
 gastrointestinal nematodes and lungworms, calves (exper.), morantel tartrate, efficacy in 5 controlled trials, toxicity experiments demonstrate wide safety margin
- Dictyocaulus viviparus*
 Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Vet. Rec., v. 92 (20), 551-554
 control of clinical infections of gastrointestinal nematodes and lungworms in calves using morantel/diethylcarbazine solution, field trials, good results as measured by growth response and clinical symptoms; routine treatment economically sound under conditions of heavy infection
- Dictyocaulus viviparus*
 Crowley, J. W., jr.; et al., 1977, Am. J. Vet. Research, v. 38 (5), 689-692
 lungworms, gastrointestinal parasites, cattle, 3 controlled critical trials, highly effective
- Dictyocaulus viviparus*
 Curr, C., 1977, Austral. Vet. J., v. 53 (9), 425-428
 nematodes, calves, levamisole, efficiency of pour-on formulation, drug trials, good results
- Dictyocaulus viviparus*
 Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
 nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Dictyocaulus viviparus*, illus.
 Downey, N. E.; and O'Shea, J., 1977, Vet. Rec., v. 100 (13), 265-266
Dictyocaulus viviparus, *Ostertagia ostertagi*, *Cooperia oncophora*, calves (exper.), low dose levels of levamisole or morantel administered via drinking water, good results
- Dictyocaulus viviparus*
 Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
 fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Dictyocaulus viviparus*
 Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
 roe deer (lungs): Czechoslovakia
- Dictyocaulus viviparus*
 Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
 helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (lung): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Dictyocaulus viviparus*
 Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
 coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Dictyocaulus viviparus*
 Eckert, J., 1972, Schweiz. Arch. Tierh., v. 114 (12), 652-667
 basis for control measures including chemoprophylaxis, vaccination most promising, review: Switzerland
- Dictyocaulus viviparus*
 Eckert, J.; and Eisenegger, H., 1976, Zentralbl. Vet.-Med., Beihefte (25), 155-160
Dictyocaulus viviparus, cattle, program for management and control, vaccination with Diclol, tetramisole treatment; *Ostertagia ostertagi*, *Cooperia oncophora*, pyrantel tartrate treatment to control concurrent infection limiting weight gain
- D[*ictyocaulus*] *viviparus*
 Fromunda, V., 1976, Rev. Crest. Animalelor, v. 26 (3), 86-90
 helminthic diseases, sheep, prevention during grazing

- Dictyocaulus viviparus*
Goda, Fawzy F. M., 1974, Bull. Epizoot. Dis. Africa, v. 22 (1), 75-78
cattle: Benghazi, Libya
- Dictyocaulus viviparus* (Bloch, 1782)
Gupta, R. P., 1976, Indian J. Animal Sc., v. 45 (2), 1975, 66-70
Dictyocaulus viviparus, techniques to culture, store and test infectivity of larvae, storage at various temperatures; infectivity declined as storage time progressed
- Dictyocaulus viviparus* (Bloch, 1782)
Gupta, R. P.; Chauhan, H. V. S.; and Rao, U. R. K., 1974, Indian J. Animal Sc., v. 43 (8), 1973, 779-781
incidence
cattle (lungs): slaughterhouse at Jabalpur, India
- Dictyocaulus viviparus*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (bronchi): Kentucky
- Dictyocaulus viviparus*
Inderbitzin, F.; and Eckert, J., 1976, Ztschr. Parasitenk., v. 50 (2), 218
Dictyocaulus viviparus, calves infected with larvae previously maintained at 4°C, induced inhibited development
- Dictyocaulus viviparus*
Ito, S.; and Taniguchi, R., 1976, Nippon Zyuishi-Kai Zassi (J. Japan. Vet. Med. Ass.), v. 29 (11), 616-619
Dictyocaulus viviparus, calf, possible contact infection in calves living in same pen
- Dictyocaulus viviparus*, illus.
Jørgensen, R. J., 1973, Acta Vet. Scand., v. 14 (2), 341-343
Dictyocaulus viviparus, 3rd stage larvae, activation by bile, in vitro, technique for evaluating viability of larval suspensions
- Dictyocaulus viviparus*
Jørgensen, R. J., 1975, Vet. Parasitol., v. 1 (1), 61-67
isolation of infective *Dictyocaulus* larvae from herbage, 3 steps: 1. isolation of debris from herbage 2. isolation of larvae from debris (migration in agar gel containing bile) 3. staining to differentiate from free-living nematodes
- Dictyocaulus viviparus*, illus.
Kummeneje, K., 1977, Acta Vet. Scand., v. 18 (1), 86-90
Dictyocaulus viviparus, reindeer, verminous bronchopneumonia, pathology; endemic with healthy carriers; infestation probably on pasture in summer or fall, but clinical disease in spring, probable inhibited larval development in winter: Finnmark, northern Norway
- Dictyocaulus vivipara*
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Dictyocaulus viviparus*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence: Zehusice enclosure
- Dictyocaulus viviparus*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure
- Dictyocaulus viviparus*
Oakley, G. A., 1977, Vet. Rec., v. 101 (10), 187-188
Dictyocaulus viviparus, overwinter survival on pasture, possible mechanisms: Cheshire
- Dictyocaulus viviparus*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Dictyocaulus viviparus*, illus.
Parker, S.; and Croll, N. A., 1976, Exper. Parasitol., v. 40 (1), 80-85
Dictyocaulus viviparus, pepsin did cause exsheathment but was not an absolute requirement, exsheathment occurred in other proteases and in chitinase at appropriate pH optima, concluded that exsheathment in vivo is caused by host gut enzymes
- Dictyocaulus viviparus*
Petrovic, A. P., 1975, Vet. Glasnik, v. 29 (10), 771-774
cattle, local Zebu breed and Boran breed (lungs of all): all from export slaughterhouse, Tanzania
- Dictyocaulus viviparus*
Pfeiffer, H., 1976, Wien. Tierarztl. Monatschr., v. 63 (2), 54-55
Dictyocaulus viviparus, cattle, inhibition of development greater in animals experimentally infected with larvae maintained at 7° C during all or most of development than with larvae matured quickly at 25° C; dosing with anthelmintics during prepatent period may increase rate of inhibition
- Dictyocaulus viviparus* (Bloch, 1782)
Phillips, J. H.; Harley, J. P.; and Rudersdorf, W. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 250
Dama dama (lungs): western Kentucky

- Dictyocaulus viviparus*
 Presidente, P. J. A.; and Knapp, S. E., 1973, J. Wildlife Dis., v. 9 (1), 41-43
Dictyocaulus viviparus isolate from *Odocoileus hemionus columbianus*, bovine calves experimentally inoculated or naturally exposed on contaminated pasture, clinical response but no development of patent infections, implications for natural cross-transmission
- Dictyocaulus viviparus*
 Presidente, P. J. A.; Knapp, S. E.; and Dean, R. E., 1973, J. Wildlife Dis., v. 9 (1), 34-40
Dictyocaulus viviparus, captive *Odocoileus hemionus columbianus* fawns, treatment with cambendazole vs. levamisole hydrochloride, survival and infectivity of larvae on contaminated pasture
- Dictyocaulus viviparus*
 Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
 survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites *Odocoileus virginianus*: Hardy County, West Virginia
- Dictyocaulus viviparus*
 Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
 nematodes, seasonal dynamics in deer
- Dictyocaulus viviparus* (Bloch, 1782)
 Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (lungs): Cumberland County, New Jersey
- Dictyocaulus viviparus*
 Ramachandran, C. P., 1975, Kajian Vet., v. 7 (1), 31-38
 helminths, immunization with radiation attenuated vaccines, review
- Dictyocaulus viviparus*
 Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
 gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Dictyocaulus viviparus*
 Rowlands, D. ap T.; and Berger, J., 1977, J. South African Vet. Ass., v. 48 (2), 85-93
 nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Dictyocaulus viviparus*
 Rubin, R., 1972, Am. J. Vet. Research, v. 33 (2), 425-426
Dictyocaulus viviparus, cattle (exper.), efficacy of cambendazole
- Dictyocaulus viviparus*
 Saad, M. B. E.; and Rubin, R., 1977, Am. J. Vet. Research, v. 38 (9), 1427-1428
Dictyocaulus viviparus, calves (exper.), fenbendazole administered as a drench or as a feed additive highly effective against adult lungworms
- Dictyocaulus viviparus*
 Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
 helminths of *Alces alces*, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Dictyocaulus viviparus*
 Selman, I. E.; et al., 1977, Vet. Rec., v. 101 (14), 278-283
 cattle, change from poor to lush, lungworm-free pasture; while *D. viviparus* cannot be excluded absolutely from playing a role in fog fever, the disease cannot arise as the result of recent larval invasion
- Dictyocaulus viviparus*
 Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
 nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Dictyocaulus viviparus* (Bloch, 1782)
 Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Bos taurus: insular Newfoundland
- Dictyocaulus viviparus*
 Swietlikowski, M., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 101-115
Dictyocaulus viviparus, calves, immunization, normal or X-ray inactivated larvae, numbers of infective larvae, levels of complement fixing and precipitating antibodies, course of infection, precipitating antibodies appearing later than complement fixing antibodies and probably produced by mature parasites
- Dictyocaulus viviparus*
 Swietlikowski, M., 1969, Acta Parasitol. Polon., v. 17 (1-19), 89-94
Dictyocaulus viviparus, calves, immunization by subcutaneous injection of larvae; infective larvae produce immunity and antibodies in sera; uninformative larvae immunize but produce no antibodies
- Dictyocaulus viviparus*
 Swietlikowski, M., 1969, Acta Parasitol. Polon., v. 17 (1-19), 95-101
Dictyocaulus viviparus, calves infected orally by larvae refrigerated 3 or 8 months; young larvae produce more severe disease; both ages cause similar immunological response; implications for overwintering, epizootiology, and self-cure

- Dictyocaulus viviparus*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Dictyocaulus viviparus*
Wolf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere: Trebic District
- Dictyocaulus viviparus*
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Dictyocaulus viviparus*
Winters, J. B.; and Worley, D. E., 1975, Am. J. Vet. Research, v. 36 (3), 327-329
Dictyocaulus viviparus, occurrence in beef cattle of various age groups, seasonal cycle, potential role of wild ruminants in epizootiology of lungworms: Park, Gallatin, and Ravalli counties, western Montana
- Didelphonema Wolfgangi*, 1953, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirocerinae
key
- Didelphostrongylus* gen. n.
Prestwood, A. K., 1976, J. Parasitol., v. 62 (2), 272-275
Filaroididae, Angiostrongylinae, mt: D. hayesi sp. n.
- Didelphostrongylus hayesi* sp. n. (mt), illus.
Prestwood, A. K., 1976, J. Parasitol., v. 62 (2), 272-275
Didelphis marsupialis (nat. and exper.) (lungs, subpleural): Clarke, Jackson, and Madison counties, Georgia
Mesodon perigraptus (exper.)
Triodopsis albolabris (exper.)
- Didelphostrongylus hayesi*, illus.
Prestwood, A. K.; Nettles, V. F.; and Farrell, R. L., 1977, Am. J. Vet. Research, v. 38 (4), 529-532
pathology
Didelphis marsupialis (lung) (nat. and exper.): Georgia (Clarke County; Madison County)
- Diectophyma*. See *Diectophyme*.
- Diectophymatina* Skryabin, 1927
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Trichocephalida
- Diectophymatoidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Enoplida
key
- Diectophyme renale* (Goeze, 1782) illus.
Bogoiavlenskii, Iu. K.; and Khatkevich, L. M., 1970, Parazitologiya, Leningrad, v. 4 (3), 223-230
4 spp. of *Diectophymata*, fine structure of somatic musculature, distribution of DNA and RNA
- Diectophyme renale* Goetze
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus
Phoca vitulina
(kidney of all): all from European waters
- Diectophyma renale*
Mace, T. F., 1976, J. Wildlife Dis., v. 12 (1), 88-92
Diectophyma renale in *Mustela vison*, kidney lesions: Washago region, Ontario, Canada
- Diectophyma renale*, illus.
Taniguchi, M.; et al., 1977, Bull. Coll. Agric. and Vet. Med., Nihon Univ. (34), 202-217
Rattus norvegicus: Setagaya-ku area, Tokyo
- Dipetalonema*
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
Dipetalonema lineage, definition and limits of the line, morphological development, hypotheses on evolution, key to genera and subgenera
- Dipetalonema* Diesing, 1861
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
key
includes subgenera: *Orihelia* n. subgen.; *Acanthocheilonema*; *Molinema*; *Loxodontofilaria*; *Chenofilaria*; *Dipetalonema*
- Dipetalonema* [? n. rank]
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of *Dipetalonema*; key
tod: D. (D.) *caudispina* (Molin, 1858)
- Dipetalonema* sp., illus.
Alcaino, H.; and Rudolph, W., 1970, Bol. Chileno Parasitol., v. 25 (1-2), 89-90
perro (blood): Chile
- Dipetalonema* sp., illus.
Balbo, T.; and Abate, O., 1972, Parassitologia, v. 14 (2-3), 239-244
Dirofilaria immitis, D. repens, *Dipetalonema* sp., microfilaria from dogs, staining for localization of acid phosphatase, detailed procedure, basis for diagnostic differentiation

- Dipetalonema [sp.], perhaps *D. reconditum*
Beaucournu, J. C.; and Deunff, J., [1976],
Ann. Parasitol., v. 50 (6), 1975, 831-835
as hyperparasites of fleas, causing para-
sitic castration
Pulex irritans: France
- Dipetalonema sp.
Christensson, D., 1977, Svensk Vet.-Tidn.,
v. 29 (12), 521-522
incidence in dogs: Sweden, imported from
Spain
- Dipetalonema [sp.]
Jackson, R. F.; et al., 1975, Vet. Rec., v.
97 (24), 476-477
dogs (blood): Florida, imported from South-
ern Ireland
- Dipetalonema sp.
Kelly, J. D., 1973, Austral. Vet. J., v. 49
(1), 23-27
Dirofilaria immitis, *Dipetalonema* sp., detec-
tion and differentiation of microfilariae in
canine blood, comparison of techniques,
modified Knott technique and modified filter
technique most reliable and consistent for
detection; morphological and physiological
features considered collectively is recom-
mended for positive identification, striking
difference in acid phosphatase activity of-
fers most accurate method of differentiation
- Dipetalonema sp., *illus.*
King, N. W., jr., 1976, Scient. Publication
(317). Pan Am. Health Organ., 169-198
- Dipetalonema [sp.] microfilariae
Schillhorn van Veen, T.; and Blotkamp, J.,
1975, Ann. Trop. Med. and Parasitol., v. 69
(4), 517-518
dogs: * Zaria area, Nigeria
- Dipetalonema sp.
Sengbusch, H.; Sartori, P.; and Wade, S.,
1975, Am. J. Vet. Research, v. 36 (7), 1035-
1036
Dirofilaria immitis, 2 of 100 stray dogs in-
fected, none with *Dipetalonema* sp., first
report of heartworm in domesticated dogs in
the state: Buffalo, New York
- Dipetalonema (*Orihelia*) *anticiava* (Molin, 1858),
illus. (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema (*Molinema*) *arbuta* Highby, 1943
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema (*Molinema*) *bifida* (Molin, 1858)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema (*Dipetalonema*) *caudispina* (Molin,
1858) (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema caudispina*, tentative identifica-
tion, *illus.*
Chalifoux, L. V.; et al., 1973, Lab. Animal
Sc., v. 23 (2), 211-220
differentiation of 11 types of circulating
microfilariae in blood smears from 7 spp. of
New World monkeys based on differences in
histochemical localization of acid phosphatase
Saimiri sciureus
Ateles geoffroyi
Saguinus tamarinus
Cebus albifrons
all from New England Regional Primate Re-
search Center
- Dipetalonema *dessetae* Bain, 1973
Bain, O., 1976, Bull. World Health Organ.,
v. 54 (4), 397-401
human filariasis, number of developing and
infective larvae dependent upon number of
microfilariae penetrating into haemocoel of
vector, relationship based on proportionality,
facilitation and limitation, applica-
tion to disease control and treatment
methods
- Dipetalonema *dessetae*
Bain, O.; and Chabaud, A. G., 1975, Compt.
Rend. Acad. Sc., Paris, v. 281, s. D, Sc. Nat.
(16), 1199-1202
Dipetalonema dessetae, relation of number of
microfilariae ingested by *Aedes aegypti*
to number penetrating stomach wall, 'limi-
tation' phenomenon, apparently specific
lysis of stomach cells
- Dipetalonema *dessetae*
Bain, O.; and Chabaud, A.-G., 1977, Ann. Para-
sitol., v. 52 (1), 84-86
Dipetalonema dessetae in *Aedes aegypti*,
phenomenon of 'limitation' regulating num-
bers of microfilariae crossing stomach wall
of vector
- Dipetalonema (*Molinema*) *dessetae* Bain, 1973,
illus.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema *dessetae*
Petit, G.; Bain, O.; and Spitalier-Kaveh, H.,
1977, Ann. Parasitol., v. 52 (5), 585-586
Dipetalonema dessetae, factors favorable to
laboratory transmission in *Aedes aegypti*
and *Proechimys guyanensis*
- Dipetalonema (*Molinema*) *diacantha* (Freitas et
Lent, 1939), *illus.* (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Dipetalonema *dracunculoides*
Bwangamoi, O.; and Isyagi, A. O., 1973,
Bull. Epizoot. Dis. Africa, v. 21 (1),
33-37
filariasis, dogs, incidence survey, mor-
bidity rate per breed, age, and sex of
host: Uganda
- Dipetalonema (*Acanthocheilonema*) *dracunculoides*
(Cobbold, 1870), *illus.* (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397

- Dipetalonema evansi*
Michael, S. A.; and Saleh, S. M., 1977, Trop. Animal Health and Prod., v. 9 (4), 241-244
Dipetalonema evansi, camels, slide agglutination test for diagnosis, high number of positives from *D. evansi* infections, no positives in camels infected with *Trypanosoma evansi*
- Dipetalonema fausti* Esslinger, 1966
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
as syn. of *Dipetalonema* (*Chenofilaria*) *filaria* (Kou, 1958)
- Dipetalonema fausti* Esslinger 1966
Esslinger, J. H., 1976, J. Parasitol., v. 62 (4), 527
as syn. of *Chenofilaria filaria* Kou 1958
- Dipetalonema* (*Chenofilaria*) *filaria* (Kou, 1958), illus. (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
Syn.: *Dipetalonema fausti* Esslinger, 1966
- Dipetalonema finlayi*
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
as syn. of *Ackertia finlayi*)Mazza et Fiora, 1932)
- Dipetalonema* (*Loxodontofilaria*) *gossi* (Baylis, 1923)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Dipetalonema*) *gracile* (Rudolphi, 1809), illus.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema gracile*, illus.
Chalifoux, L. V.; et al., 1973, Lab. Animal Sc., v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saimiri sciureus
Ateles geoffroyi
all from New England Regional Primate Research Center
- Dipetalonema* (*Dipetalonema*) *graciliformis* Freitas, 1964
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Acanthocheilonema*) *grassii* Noe, 1907
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Loxodontofilaria*) *hippopotami* (Leiper, 1910)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema interstitium*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (subcutaneous): south-eastern United States
- Dipetalonema* (*Chenofilaria*) *johnstoni* (Spratt et Varughese, 1975) [i. e. (Mackerras, M. J., 1954)]
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Loxodontofilaria*) *loxodontis* (Berghe et Gillain, 1939) (tod of subgen.)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Acanthocheilonema*) *mansonbahri* Nelson, 1961
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema* (*Acanthocheilonema*) *mephitis* (Webster et Beauregard, 1964)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema obtusa*, illus.
Chalifoux, L. V.; et al., 1973, Lab. Animal Sc., v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Cebus albifrons: New England Regional Primate Research Center
- Dipetalonema* (*Acanthocheilonema*) *odendhali* Perry, 1967
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema odendhali* Perry, 1967, illus.
Kagei, N.; and Oda, T., 1975, Bull. Inst. Pub. Health, Tokyo, v. 24 (4), 203-205
Dipetalonema odendhali in *Callorhinus ursinus* (intermuscular fascia), description, measurements compared with *D. spirocauda*: Enoshima Marine Zoo, Kanagawa Prefecture, Japan, captured near Kesenuma, Hokkaido, Japan
- Dipetalonema odendhali* Perry, 1967, illus.
Machida, M., 1977, Bull. National Sc. Mus. Tokyo, s. A, Zool., v. 3 (2), 67-71
description
Callorhinus ursinus (cervical and thoracic region of subfascia): off Sanriku District, Japan
- Dipetalonema* (*Loxodontofilaria*) *okapiae* Fain, 1948
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397

- Dipetalonema* (*Acanthocheilonema*) *pachycephalum* (Ortlepp, 1961)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Dipetalonema* (*Chenofilaria*) *pearsoni* (Spratt et Varughese, 1975)
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Dipetalonema* *perstans*
Ambrose-Thomas, P.; and Kien Truong, T., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema viteae*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance
- Dipetalonema* *perstans*
Anderson, R. I.; and Buck, A. A., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 447-456
complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad
- Dipetalonema* *perstans*
Barclay, R., 1971, *Med. J. Zambia*, v. 5 (6), 201-203
night blood survey undertaken in area where abnormally high number of hydrocoele noticed in clinic outpatients, *Dipetalonema perstans* only microfilariae found: Luangwa Basin, Zambia
- Dipetalonema* *perstans*, *illus.*
Beaver, P. C.; Neel, J. V.; and Orihel, T. C., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (2), 263-265
Indians (blood): southern Venezuela
- Dipetalonema* *perstans*
Burke, G. J., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (4), 402-405
10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Dipetalonema* *perstans*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, *Medecine et Malad. Infect.*, v. 3 (8-9), 351-353
diagnostic review of human filariasis
- Dipetalonema* *perstans*
Niel, G.; et al., 1972, *Medecine et Malad. Infect.*, v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalonema viteae* and *Ascaris suum* as antigens
- Dipetalonema* *perstans*
Picq, J. J.; and Roux, J., 1973, *Medecine Trop.*, v. 33 (5), 451-461
Onchocerca volvulus, human, microfilaruria in relation to age and sex of host, other filarial diseases, geographic region, cutaneous microfilarial densities, albuminuria during suramin treatment, eggs of *Schistosoma haematobium* in urine, and diethylcarbamazine chemotherapy
- Dipetalonema* *perstans*
Ridley, D. S.; and Hedge, E. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Dipetalonema* *perstans*
Stuerchler, D.; and Degremont, A., 1976, *Schweiz. Med. Wchnschr.*, v. 106 (20), 682-688
extensive diagnostic and clinical review of filarial parasites frequently encountered by travelers to endemic tropical areas: Switzerland
- Dipetalonema* *perstans*
Weiss, N.; and Degremont, A., 1976, *Tropenmed. u. Parasitol.*, v. 27 (3), 377-384
filariasis in persons returning from endemic areas, comparison immunoserologic diagnostic tests (immunoelectrophoresis, indirect fluorescent antibody, indirect hemagglutination, two-dimensional gel diffusion tests) showed that highest sensitivity obtained with immunoelectrophoresis, combined tests gave best results
- Dipetalonema* *perstans*
White, G. B., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (2), 161-175
Onchocerca volvulus, *Loa loa*, and *Dipetalonema perstans*, survey of man-biting species of Chrysops, Culicoides and Simulium as potential vectors of human infection: Ethiopia
- Dipetalonema* (*Acanthocheilonema*) *procyonis* Price, 1955
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Dipetalonema* (*Molinema*) *proechimiyis* Esslinger, 1974
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Dipetalonema* (*Molinema*) *raposoensis* Esslinger, 1974
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397

- Dipetalonema reconditum*
Bwagamoi, O., 1973, Bull. Epizoot. Dis. Africa, v. 21 (4), 363-370
Dirofilaria repens, *Dipetalonema reconditum*, recovery from dogs using Lindsey's method: Uganda
- Dipetalonema reconditum*
Bwagamoi, O.; and Isyagi, A. O., 1973, Bull. Epizoot. Dis. Africa, v. 21 (1), 33-37
filariasis, dogs, incidence survey, morbidity rate per breed, age, and sex of host: Uganda
- Dipetalonema (Acanthocheilonema) reconditum* (Grassi, 1889)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema reconditum*, illus.
Christofferson, P. V., 1976, Vet. Med. and Small Animal Clin., v. 71 (4), 489-492
Dirofilaria immitis, *Dipetalonema reconditum*, diagnosis, Difil Test, possible problems and artifacts, necessary precautions in use
- Dipetalonema reconditum*
Evans, K., 1977, Austral. Vet. Practitioner, v. 7 (3-4), 224-226
canine heartworm disease, techniques in diagnosis, use of wet smear test and thorough clinical examination
- Dipetalonema reconditum*
Farnell, D. R., 1976, J. Alabama Acad. Sc., v. 47 (3), 173 [Abstract]
Dipetalonema reconditum, dogs, possibly antagonistic to *Dirofilaria immitis*
- Dipetalonema reconditum*
Hills, J., 1976, Austral. Vet. Practitioner, v. 6 (1), 7 [Letter]
Dirofilaria immitis, cat, dog; *Dipetalonema reconditum*, dog, diagnosis
- Dipetalonema reconditum*
Holmes, P. R.; Kelly, J. D., 1973, Austral. Vet. J., v. 49 (1), 55
Dirofilaria immitis, *Dipetalonema reconditum*, dogs, low incidence combined with low average number of worms per dog suggests that *D. immitis* is of lesser importance than first suspected: Sydney, Australia
- Dipetalonema reconditum*
Rabalais, F. C.; and Votava, C. L., 1972, J. Am. Vet. Med. Ass., v. 160 (2), 202-203
Dirofilaria immitis, *Dipetalonema reconditum*, dogs (blood), prevalence: northwestern Ohio (some dogs originally from southern states)
- Dipetalonema reconditum*
Streitel, R. H.; Stromberg, P. C.; and Dubey, J. P., 1977, J. Am. Vet. Med. Ass., v. 170 (7), 720-721
prevalence in dogs stable since 1964, no highly enzootic center: humane shelter, Columbus, Ohio
- Dipetalonema reconditum* Grassi
Todaro, W. S.; Morris, C. D.; and Heacock, N. A., 1977, Am. J. Vet. Research, v. 38 (8), 1197-1200
prevalence
dogs (blood): central New York
- Dipetalonema reconditum*
Watson, A. D. J.; Porges, W. L.; and Testoni, F. J., 1973, Austral. Vet. J., v. 49 (1), 31-34
dogs, survey of incidence of *Dipetalonema reconditum* and *Dirofilaria immitis* microfilariae and adult *D. immitis*: Sydney, Australia
- Dipetalonema reconditum*
Watson, A. D. J.; Testoni, F. J.; and Porges, W. L., 1973, Austral. Vet. J., v. 49 (1), 28-30
Dirofilaria immitis, *Dipetalonema reconditum*, comparison of microfilarial characteristics isolated by modified Knott and filter methods, results suggest consideration of mean microfilarial length and numbers on filter may permit accurate differentiation of microfilariae in canine blood
- Dipetalonema (Loxodontofilaria) ruandae* Fain et Herin, 1955, illus.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema (Acanthocheilonema) setariosa* (Monnig, 1927)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema spiralis* (Molin, 1860)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
as syn. of *Bostrichodera spiralis* (Molin, 1860) n. comb.
- Dipetalonema (Acanthocheilonema) spirocauda* (Leidy, 1858)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema spirocauda* (Leidy, 1858)
Kagei, N.; and Oda, T., 1975, Bull. Inst. Pub. Health, Tokyo, v. 24 (4), 203-205
Dipetalonema odendhali, measurements compared with *D. spirocauda*
- Dipetalonema spirocauda* (Leidy, 1858), illus.
Machida, M., 1977, Bull. National Sc. Mus., Tokyo, s. A, Zool., v. 3 (2), 67-71
description
Phoca vitulina (heart): Odaito, Hokkaido, Japan
- Dipetalonema (Molinema) sprengi* Anderson, 1953
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema streptocerca*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, Medecine et Malad. Infect., v. 3 (8-9), 351-353
diagnostic review of human filariasis

- Dipetalonema streptocerca*
McDougall, A. C., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (4), 360-361 [Letter]
onchocerciasis and streptocerciasis in persons with leprosy, possible relationships
- Dipetalonema streptocerca, illus.*
Meyers, W. M.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, part 1), 1153-1155
Dipetalonema streptocerca, identifying features of adult male worms excised from human dermal collagen during skin biopsies of patients after treatment for streptocerciasis: Bolonga-Scieurs, Equator Region, Republic of Zaire
- Dipetalonema streptocerca*
Meyers, W. M.; and Connor, D. H., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (5-6), 524-525
human *Onchocerca volvulus* and *Dipetalonema streptocerca* in persons with leprosy, altered Mazzotti reactions following administration of diethylcarbamazine, possible immunological implications
- Dipetalonema (Molinema) travassosi* Artigas et Pacheco, 1933
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema viteae*. See *Dipetalonema viteae*.
- Dipetalonema (Acanthocheilonema) weissi, illus.*
Bain, O.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (5), 569-575
Dipetalonema weissi, adult morphology, larval development in *Ornithodoros erraticus* (exper.), comparison with *D. viteae*
Elephantulus rozeti (peritoine, tissue sous-cutane, au niveau de la tete, du ventre et du dos): environs de Sfax, Tunisie
- Dipetalonema (Acanthocheilonema) weissi* (Seurat, 1914)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema viteae*
Abaru, D. E.; and Denham, D. A., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 333-334
laboratory evaluation of new technique for counting microfilariae in blood, comparison with counting chamber method, possible advantages of new technique in field studies
- Dipetalonema viteae*
Ambrose-Thomas, P.; and Kien Truong, T., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema viteae*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance
- Dipetalonema viteae*
Ambrose-Thomas, P.; Prod'hon, J.; and Venard, P., 1975, Bull. Soc. Path. Exot., v. 68 (6), 579-588
Wuchereria bancrofti, humans, epidemiologic prevalence survey comparing findings from microfilaremia determinations and those from indirect immunofluorescence studies using *Dipetalonema viteae* antigens: south-east coast of Madagascar
- Dipetalonema viteae*
Arafa, M. S.; Salit, A. M.; and Hilal, T., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 13 [Demonstration]
unsheathed microfilariae undistinguishable from *Dipetalonema viteae* found in blood of *Rattus* spp. in Egypt
- Dipetalonema (Acanthocheilonema) viteae, illus.*
Bain, O.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (5), 569-575
Dipetalonema weissi, adult morphology, larval development in *Ornithodoros erraticus* (exper.), comparison with *D. viteae*
- Dipetalonema (Acanthocheilonema) viteae* (Krepkogorskaya, 1953)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Dipetalonema viteae*
Dalesandro, D. A.; and Klei, T. R., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 534-535 [Letter]
Dipetalonema viteae-infected *Mesocricetus auratus* (exper.) and *Meriones unguiculatus* (exper.) showed immunodepression at 10 but not at 5 weeks post-inoculation, possible association between microfilaremia and immunodepression
- Dipetalonema viteae*
Diesfeld, H. J.; and Kirsten, C., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 435-438
filariasis, human, diagnosis, new embedding technique employing 'methacrylate' for preparation of antigen (*Dipetalonema viteae*) to be used in indirect fluorescent antibody test (tested on onchocerciasis sera from Togo), compared with usual frozen-section method
- Dipetalonema viteae, illus.*
Diesfeld, H. J.; and Kirsten, C., 1975, Tropenmed. und Parasitol., v. 26 (4), 499-502
Dipetalonema viteae, localization of antigen-antibody reactions in male and female using immunofluorescence and serum from human filariasis patients; possible implications for diagnosis of human infection
- D[*ipetalonema*] *viteae*
Dutta, S. N.; Diesfeld, H. J.; and Kirsten, C., 1976, Tropenmed. u. Parasitol., v. 27 (4), 479-482
immunofluorescent antibody test using D[*ipetalonema*] *viteae* as antigen applied to mothers' blood and to the umbilical cord blood of their newborns, results show that in a *Wuchereria bancrofti* endemic area maternal antibodies against filariae are passed via placenta to newborn, findings not demonstrable after 6th month of life: India

- Dipetalonema viteae**
Eberhard, M. L.; and Rabalais F. C., 1976, *Exper. Parasitol.*, v. 40 (1), 5-12
Dipetalonema viteae in *Meriones unguiculatus*, hypo- and hyperthermic stress resulted in increased numbers of microfilariae in peripheral blood, microfilariae do not exhibit cyclic periodicity
- Dipetalonema viteae**
Holdstock, R. P., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (1), 9 [Demonstration]
Dipetalonema viteae, *Mastomys natalensis* a successful exper. host with patent infections established in mature males and immature males and females, periodicity, parasite localization, peripheral microfilaremia in relation to presence of female worms
- Dipetalonema viteae**
Hutchison, W. F.; et al., 1976, *Comp. Biochem. and Physiol.*, v. 53 (4B), 495-497
Dirofilaria immitis adults, lipid analysis, compared with *Dipetalonema viteae* and *Litomosoides carinii*
- Dipetalonema viteae, illus.**
Londono M., I., 1976, *J. Parasitol.*, v. 62 (4), 589-595
Dipetalonema viteae larvae, distribution and migration in *Ornithodoros tartakowskyi*, as cause of injury to and disorganization of tick muscle fibers, failure of some ticks to feed after biting may protect from fatal hyperinfection
- Dipetalonema viteae**
Londono M., I., 1976, *J. Parasitol.*, v. 62 (4), 596-603
number and proportion of infective *Dipetalonema viteae* larvae moving from biting *Ornithodoros tartakowskyi* to *Meriones unguiculatus*, possible factors involved in this migration
- Dipetalonema viteae**
Londono M., I., 1976, *J. Parasitol.*, v. 62 (5), 786-788
Ornithodoros tartakowskyi, transmission of *Dipetalonema viteae* by small ticks feeding on engorged larger infected ticks, normal development with subsequent transmission to jirds; elimination of microfilariae in coxal fluid as a possible mechanism to prevent hyperinfection
- Dipetalonema viteae**
Neilson, J. T. M., 1975, *J. Parasitol.*, v. 61 (5), 785-793
Dipetalonema viteae, adults, soluble somatic extracts, extracts of solubilized cuticles and membranes, fractionation by Sephadex column chromatography and polyacrylamide gel electrophoresis, constituents of each preparation compared by immunodiffusion and immunoelectrophoresis
- Dipetalonema viteae**
Neilson, J. T. M., 1976, *Tropenmed. u. Parasitol.*, v. 27 (2), 233-237
a comparison of the acquired resistance to *Dipetalonema viteae* stimulated in hamsters by trickle versus tertiary infections
- Dipetalonema vitae**
Niel, G.; et al., 1972, *Medecine et Malad. Infect.*, v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalonema vitae* and *Ascaris suum* as antigens
- Dipetalonema vitae**
Pinon, J. M.; and Dropsy, G., 1977, *J. Immunol. Methods*, v. 16 (1), 15-22
various human parasitic diseases, application of enzyme-linked-immuno-electro-diffusion assay (combination of immunoelectrodiffusion and immunoenzyme method), sensitivity and specificity, enables class of immunoglobulins involved to be determined
- Dipetalonema vitae**
Pinon, J. M.; and Dropsy, G., 1977, *Path. Biol.*, v. 25 (1), 23-27
human parasitic diseases, use of enzyme-linked-immuno-electro-diffusion-assay (ELIEDA) in diagnosis and immunologic studies
- Dipetalonema viteae**
Rew, R. S.; and Saz, H. J., 1977, *J. Parasitol.*, v. 63 (1), 123-129
Litomosoides carinii, *Dipetalonema viteae*, and particularly *Brugia pahangi* microfilariae, oxygen requirements, carbohydrate metabolism, effect of levamisole
- Dipetalonema witei (= viteae)**
Saz, H. J.; and Dunbar, G. A., 1975, *J. Parasitol.*, v. 61 (5), 794-801
stibophen, inhibition of phosphofructokinase and lactate formation, effect on internal hexose phosphate accumulation, inhibition of aldolase, comparison with potassium antimony tartrate (inhibits PFK at higher concentrations but not aldolase)
- Dipetalonema viteae, illus.**
Simpson, C. F.; and Neilson, J. T. M., 1976, *Tropenmed. u. Parasitol.*, v. 27 (3), 349-354
Dipetalonema viteae in *Mesocricetus auratus* (exper.), comparison of pathologic changes in hamsters with single versus quadruple infections (hyperinfected hamsters developed more extensive lesions and subcutaneous nodules)
- Dipetalonema witei**
Striebel, H. P., 1976, *Experientia*, v. 32 (4) 457-458
4-isothiocyanato-4'-nitrodiphenylamine, an anthelmintic with an unusual spectrum of activity against intestinal nematodes, filariae and schistosomes
- Dipetalonema viteae**
Sullivan, J. J.; and Chernin, E., 1976, *Internat. J. Parasitol.*, v. 6 (1), 75-78
Brugia pahangi and *Dipetalonema viteae* compared, differences in oral vs. subcutaneous infection of anaesthetized vs. unanaesthetized adult vs. neonatal *Meriones unguiculatus*

- Dipetalonema viteae*
Turner, A. C.; et al., 1975, J. Mississippi Acad. Sc., v. 20, Suppl., 25
Dirofilaria immitis, dogs, lipid content of worms, total fatty acid methyl esters in Dirofilaria immitis, compared with Dipetalonema viteae and Litomosoides carinii, similar fatty acid patterns, gas chromatography
- Diplogasterata* Paramonov, 1956
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Rhabditida
- Diplogasteroides* sp.
Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Diplotrriaena* sp.
Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Diplotrriaena* sp.
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coerulescens (air sacs): Florida
- Diplotrriaena* sp.
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Porphyryla martinica (body cavity): Florida
- Diplotrriaena bargusina*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (air sacs): Ottawa county, Ohio
- Diplotrriaena darnaudii* sp. n., illus.
Webster, W. A.; and Speckmann, G., 1976, J. Parasitol., v. 62 (3), 451-452
Trachyphonus darnaudii (thoracic air sacs): Lake Baringo area, Kenya, died at Assiniboine Park Zoo, Winnipeg, Manitoba
- Diplotrriaena falconis*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Diplotrriaena obtusa* (Rud., 1802) Railliet et Henry, 1909
Jaron, W., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 137-152
description, helminth fauna of adult swallows just returning from migration compared with young birds; dynamics of infection, species composition of helminths, various stages of nesting season
Hirundo rustica
Riparia riparia
Hirundo savignyi
Delichon urbica
Junco hyemalis
Alauda arvensis
all from Poland
- Diplotrriaena obtusa* (Rud., 1802)
Webster, W. A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109
Progne subis (body cavity): Ottawa, Ontario
- Diplotrriaena tridens*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in Locusta migratoria (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Diplotrriaenoidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Diplotrriaenoidea*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
- Dirofilaria* [sp.], resembling *D. ursi*, illus.
Beaver, P. C.; and Samuel, W. M., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 329-330
Dirofilaria [sp.] resembling *D. ursi*, infertile female removed from subcutaneous nodule from upper arm near axilla of middle-aged woman: Kamloops, British Columbia, Canada
- Dirofilaria* (Nochtiella) sp., illus.
Meerovitch, E.; Faubert, G.; and Groulx, G., 1976, Canad. J. Pub. Health, v. 67 (4), 333-335
Dirofilaria sp. removed from superficial nodule on woman's breast, probable third documented finding of undetermined zoonotic filarial "Welty-type" nematode in human: rural Quebec Province
- Dirofilaria* sp., fig.
Thomas, D.; et al., 1976, Am. J. Ophth., Chicago, v. 82 (6), 931-933
Dirofilaria sp., man, eye orbit with presenting symptoms of conjunctivitis and inflammatory pseudotumor, sexually mature male worm excised: Florida

- Dirofilaria acutuscula* (D. repens), *illus.*
Stevanovic, R.; Savic-Cvetojevic, M.; and Petrovic, Z., 1971, *Acta Parasitol. Lugoslavica*, v. 2 (2), 93-97
case report in 10-year-old boy (under skin on spine): vicinity of Varvarin
- Dirofilaria* (Nochtiella) *conjunctivae*, *illus.*
Tasmouranis, N.; et al., 1976, *Arch. Inst. Pasteur Helvet.*, v. 22, 61-73
Dirofilaria conjunctivae, clinical case reports of subcutaneous nodules each containing immature female worm, excised from 2 women (neck region above left clavicle, abdominal wall): Greece
- Dirofilaria corynoides* (?=D. aethiops)
McGreavy, P. B.; McClelland, G. A. H.; and Lavogier, M. M. J., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (1), 97-109
Dirofilaria immitis, susceptibility of *Aedes aegypti* controlled by sex-linked recessive gene which is distinct from those controlling development of *Brugia pahangi* or *D. corynoides*, variation in filarial infectivity as well as in mosquito susceptibility
- Dirofilaria immitis*
Abaru, D. E.; and Denham, D. A., 1976, *South East Asian J. Trop. Med. and Pub. Health*, v. 7 (3), 367-369
Brugia pahangi, *Dirofilaria immitis*, no significant differences could be detected between nucleopore and millipore filter systems for collecting microfilariae, controlled laboratory trials
- Dirofilaria immitis*
Abaru, D. E.; and Denham, D. A., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (4), 333-334
laboratory evaluation of new technique for counting microfilariae in blood, comparison with counting chamber method, possible advantages of new technique in field studies
- Dirofilaria immitis*
Ambrose-Thomas, P.; and Kien Truong, T., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema vivax*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance
- Dirofilaria immitis*
Aoki, Y., 1971, *Nettai Igaku (Trop. Med.)*, v. 13 (1), 7-15
Dirofilaria immitis, dogs (exper.), seasonal distribution of microfilariae in peripheral circulation and effect of temperature on migration from lung capillaries to peripheral circulation
- Dirofilaria immitis*, *illus.*
Aoki, Y.; and Katamine, D., 1975, *Nettai Igaku (Trop. Med.)*, v. 17 (1), 27-34
Dirofilaria immitis, scanning electron morphology, morphometric data
- Dirofilaria immitis*, *illus.*
Babio, T.; and Abate, D., 1972, *Parassitologia*, v. 14 (2-3), 239-244
Dirofilaria immitis, D. repens, *Dipetalonema* sp., microfilaria from dogs, staining for localization of acid phosphatase, detailed procedure, basis for diagnostic differentiation
- Dirofilaria immitis*
Barranti, J. A.; Kristensen, F.; and Drumheller, F. B., 1977, *Am. J. Vet. Research*, v. 38 (7), 1055-1058
Dirofilaria immitis, dogs, analysis of serum proteins using agarose electrophoresis, relationship of differences in concentration of protein fractions to differences in age, sex, and infected vs. non-infected dogs
- Dirofilaria immitis*
Bewick, W. J.; and Moorhouse, D. E., 1968, *J. Med. Entomol.*, v. 5 (3), 269-272
Dirofilaria immitis, attempted experimental infections of potential vectors in laboratory or by feeding on infected dogs, positive results with *Culex pipiens fatigans*, *C. annulirostris*, *Aedes vigilax*, *A. notoscriptus*
- Dirofilaria immitis*
Bongis, R. G., 1975, *J. South African Vet. Ass.*, v. 46 (4), 373
Dirofilaria immitis, control, diagnosis, transmission of canine heartworm by imported dogs
- Dirofilaria immitis*
Benjamin, D. B.; and Soulsby, E. J. L., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (2), 266-272
Brugia pahangi in *Mastomys natalensis*, monocytotropic and hemagglutinating antibody responses detected using *Dirofilaria immitis* as antigen
- Dirofilaria immitis*
Sickle, W. E., 1976, *Mosquito News*, v. 36 (3), 366-367
failure of *Culex salinarius* to transmit *Dirofilaria immitis* from dog to dog
- Dirofilaria immitis*
Sickley, W. E.; et al., 1977, *Mosquito News*, v. 37 (1), 137-138
Dirofilaria immitis, dog-to-dog transmission by *Aedes canadensis*
- Dirofilaria immitis*
Bradley, R. E., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (3), 311-316
Dirofilaria immitis, dogs (nat. and exper.), thiocetarsamide sodium to destroy adult worms, levamisole resinate 6 weeks later at dosage of 11 mg/kg (but not 5.5 mg/kg) was effective as a microfilaricide, 2 cases of levamisole toxicosis in presence of adult worms; incidental observation on greater susceptibility of older dogs to infection
- Dirofilaria immitis*
Bradley, R. E.; Sr.; and Alford, B. T., 1977, *Mod. Vet. Pract.*, v. 58 (6), 518-520
Dirofilaria immitis, dogs, levamisole resinate, good results against microfilariae and adult male worms, less effective against adult female worms

- Dirofilaria immitis*, *ill.*.
Brightman, A. H.; Helper, L. C.; and Todd, K. S., Jr., 1977, *Vet. Med. and Small Animal Clin.*, v. 72 (1), 1021-1023
Dirofilaria immitis, dog (anterior chamber of eye), surgical removal
- Dirofilaria immitis*
Brino, J. A. S.; et al., 1971, *Med. J. Australia*, v. 2 (24), 1238-1241
woman, excised pulmonary lesion contained *Dirofilaria immitis*, case report: Western Australia
- Dirofilaria immitis*
Brown, S. R.; and Marshall, E. W. III, 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (8), 1078-1080
Dirofilaria immitis, dogs, survey: New Jersey (Fort Dix; McGuire Air Force Base)
- Dirofilaria immitis*
Bwangami, O.; and Isyagi, A. O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (1), 33-37
filariasis, dogs, incidence survey, morbidity rate per breed, age, and sex of host: Uganda
- Dirofilaria immitis*
Chellappah, W.T.; and Chellappah, G.R., Jr., 1968, *J. Med. Entom.*, v. 5 (3), 358-361
Dirofilaria immitis, attempted experimental infections of potential vectors by feeding on infected dogs, positive results with *Aedes aegypti*, *A. albopictus*, *Culex fatigans*, negative results with *Araucaria sub-sabatus*
- Dirofilaria immitis*
Christensen, B. M.; and Andrews, M. N., 1976, *J. Parasitol.*, v. 62 (2), 276-280
Aedes trivittatus (nat. and exper.) (Malpighian tubules, abdomen, thorax, head): Ames, Iowa
Anopheles punctipennis (Malpighian tubules): Ames, Iowa
- Dirofilaria immitis*
Christenson, D., 1977, *Svensk Vet.-Tidn.*, v. 29 (12), 521-522
incidence in dogs: Sweden (imported from Singapore, Tanzania, Kenya, Japan)
- Dirofilaria immitis*, *ill.*.
Christofferson, P. V., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (4), 489-492
Dirofilaria immitis, *Dipetalonema reconditum*, diagnosis, Difil Test, possible problems and artifacts, necessary precautions in use
- Dirofilaria immitis*
Church, E. M.; Georgi, J. R.; and Robson, D. S., 1976, *Cornell Vet.*, v. 66 (3), 333-336
Dirofilaria immitis, microfilarial periodicity, dogs, statistical analysis, variation in peak hours: emotional or environmental stresses may obscure periodicity
- Dirofilaria immitis*
Courtney, C. H.; and Nachreiner, R. F., 1976, *Am. J. Vet. Research*, v. 37 (5), 1095-1097
4 male dogs given diethylcarbamazine citrate for 6 mos. at twice daily dose recommended for heartworm prophylaxis, no significant deterioration in quantity, morphology, motility, or viability of sperm as compared to 4 unmedicated controls
- Dirofilaria immitis*
Crans, W. J.; and Feldhauser, M. F., 1974, *Proc. and Papers 42. Ann. Conf. Calif. Mosquito Control Ass. and 30. Ann. Meet. Am. Mosquito Control Ass. (Anaheim, Feb. 24-27), 168 [Abstract]*
survey of mosquito vectors of dog heartworm: coastal-area of southern New Jersey
- Dirofilaria immitis*
Cusick, P. K.; et al., 1976, *J. Am. Animal Hosp. Ass.*, v. 12 (4), 490-491
Dirofilaria immitis, cat (brain, heart), neuropathologic changes, case history: Massachusetts
- Dirofilaria immitis* (Leidy, 1856), *ill.*.
D'Alessandro, A. L., 1972, *Rev. Med. Vet. y Parasitol.*, Maracay, v. 25 (1-8), 1971-1972, 109-130
dogs, prevalence, statistical analysis, clinical aspects, geographic distribution: Estado Aragua, Venezuela
- Dirofilaria immitis*
Desowitz, R. S.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (4), 430 [Demonstration]
Dirofilaria immitis-infected dogs treated with diethylcarbamazine, severe drug reaction generally in dogs with high microfilaremia probably result of antigen and antibody combining on surface of serotonin-rich platelets, release of serotonin from damaged platelets having vascular permeability-increasing effect leading to shock
- Dirofilaria immitis*
Desowitz, R. S.; and Una, S. R., 1976, *J. Helminth.*, v. 50 (1), 53-57
Dirofilaria immitis-infected cats and dogs, *Wuchereria bancrofti*-infected humans, diagnosis, counterimmunoelectrophoresis using *D. immitis* adult and microfilarial antigens
- Dirofilaria immitis*
Dobson, C.; and Welch, J. S., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (3), 223-228
survey for antibodies against *Dirofilaria immitis*, *Toxocara canis*, *Ascaris suum*, *Angiostrongylus cantonensis*, *A. mackerrasae*, in patients with eosinophilia using fluorescent antibody test and passive reversed Arthus test in guinea pigs; *D. immitis* implicated as etiologic agent of human eosinophilic meningitis: Australia
- Dirofilaria immitis*
Donahoe, J. M. R., 1975, *J. Parasitol.*, v. 61 (4), 599-605
Dirofilaria immitis, cats (exper.), microfilaremia, immediate skin hypersensitivity, cutaneous nodules, chylothorax, low adult worm recoveries, one successful mosquito passage from-cat to dog

- Dirofilaria immitis*
Eberhard, M. L.; et al., 1977, J. Parasitol., v. 63 (6), 978
Dirofilaria immitis found in eye of microfilariae-negative dog, conversion to microfilariae-positive 105 days after discovery of this worm: Arkansas
- [*Dirofilaria*] *immitis*
Evans, K., 1977, Austral. Vet. Practitioner, v. 7 (3-4), 224-226
canine heartworm disease, techniques in diagnosis, use of wet smear test and thorough clinical examination
- Dirofilaria immitis*
Farnell, D. R., 1976, J. Alabama Acad. Sc., v. 47 (3), 175 [Abstract]
Dipetalonema reconditum, dogs, possibly antagonistic to *Dirofilaria immitis*
- Dirofilaria immitis* (Leidy, 1856)
Foil, L.; and Orihel, T. C., 1975, J. Parasitol., v. 61 (3), 433
Castor canadensis (right ventricle): near Raymond, Mississippi, held in captivity at Jackson Zoological Park, Jackson, Mississippi
- Dirofilaria immitis*
Foley, R. H., 1976, Canine Pract., Santa Barbara, v. 3 (6), 8-10 [Letter]
Dirofilaria immitis, microfilaruria, technique for differentiating urinary protein abnormalities in urine of dogs simultaneously having heartworm and renal and/or bladder disease: Florida
- Dirofilaria immitis*
Fowler, J. L.; Furusho, Y.; and Fernau, R. C., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 466-468
Dirofilaria immitis, dogs (exper.), fenthion used successfully as a prophylaxis against developing stages of parasite
- Dirofilaria immitis*
Franson, J. C.; Jorgenson, R. D.; and Bogges, E. K., 1976, J. Wildlife Dis., v. 12 (2), 165-166
Canis latrans (right ventricle): southwestern Iowa
- [*Dirofilaria immitis*] heartworm
Fujii, I., 1975, Bull. Azabu Vet. Coll. (30), 105-118
dog heartworm causing embolism in venae cavae, clinical signs, pathology, surgical treatment
- Dirofilaria immitis*, *illus.*
Fuller, W. J., 1976, Vet. Med. and Small Animal Clin., v. 71 (5), 632
Dirofilaria immitis, dog, clinical signs, digoxin, heartworm knotted around host pulmonary valves causing a severe valvular deficit, case history
- Dirofilaria immitis*
Garlick, N. L., 1976, Clin. Toxicol., v. 9 (6), 981-992
Dirofilaria immitis, dogs, drug-induced shock precipitated by microfilariae, proposed hypothesis
- Dirofilaria immitis*
Garlick, N. L., 1976, Canine Pract., Santa Barbara, v. 3 (2), 64-69
Dirofilaria immitis, dogs, drug trials, levamisole, good results against microfilariae and adult male heartworms, adult female heartworms resistant to drug, hematological and biochemical effects; levamisole + thiacetarsamide sodium + diethylcarbamazine, satisfactory results; levamisole followed by fenthion, good results
- Dirofilaria immitis*
Garlick, N. L.; Beck, A. M.; and Bryan, R. K., 1976, Canine Pract., Santa Barbara, v. 3 (4), 44-47
Dirofilaria immitis microfilariae and adults, dogs, dithiazanine iodide followed by thiacetarsamide sodium, good results, 547 cases, only one fatality
- Dirofilaria immitis*
Garlick, N. L.; and Christy, K. E., 1977, Vet. and Human Toxicol., v. 19 (1), 14-22
Dirofilaria immitis, microfilariae, dogs, biochemical lesions, pathology, treatment, toxic phenomena, blood serum chemistry, additive effects, review
- Dirofilaria immitis*
Garlick, N. L.; and Darby, T. E., 1976, Canine Pract., Santa Barbara, v. 3 (6), 29-30, 35-40
Dirofilaria immitis, dogs, electronic microfilaria counts using Coulter Counter, direct relationship between numbers of microfilariae and adult female worms, useful in selecting suitable treatment regimen
- Dirofilaria immitis*
Graham, J. M., 1975, J. Parasitol., v. 61 (3), 513-516
Dirofilaria immitis, adult heartworms found in 11 of 133 *Canis latrans*, host age and sex, discussion as reservoirs: Kansas; Colorado
- Dirofilaria immitis*
Gross, D. R.; et al., 1975, Southwest. Vet., v. 28 (3), 233-238
Dirofilaria immitis, dog, severe pulmonary hypertension, case report
- Dirofilaria immitis*
Grove, D. I.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 220-229
antigen abstract prepared from subperiodic *Brugia malayi* compared with *Dirofilaria immitis* antigen in diagnosis of human filariasis, concluded that antigens from microfilariae, adult worms, and 3rd-stage larvae of *B. malayi* are more sensitive than *D. immitis* antigens and do not have a significantly higher number of false positive reactions: Philippines

- Dirofilaria immitis*
Hedge, E. C.; and Ridley, D. S., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (4), 304-307
evaluation of microfilarial antigen for use with indirect immunofluorescent test in diagnosing human filariasis; best results obtained with sonicated microfilariae of *Brugia pahangi* with which both cytoplasmic and sheath antigens could be utilized simultaneously
- Dirofilaria immitis*
Hills, J., 1976, Austral. Vet. Practitioner, v. 6 (1), 7 [Letter]
Dirofilaria immitis, cat, dog; *Dipetalonema reconditum*, dog, diagnosis
- Dirofilaria immitis*
Ho, B.-C.; Singh, M.; and Yap, E.-H., 1974, J. Med. Entom., v. 11 (5), 622-628
Brugia malayi, *Dirofilaria immitis*, *Brelinia sergenti*, migratory patterns in *Aedes togoi* (exper.), spontaneous escape of infective larvae from mosquitoes denied access to a blood meal; level of host microfilaremia does not seem to influence mortality rate or migratory patterns of infective larvae in mosquitoes
- Dirofilaria immitis*
Holmes, P. R.; Kelly, J. D., 1973, Austral. Vet. J., v. 49 (1), 55
Dirofilaria immitis, *Dipetalonema reconditum*, dogs, low incidence combined with low average number of worms per dog suggests that *D. immitis* is of lesser importance than first suspected: Sydney, Australia
- Dirofilaria immitis*
Hutchison, W. F.; et al., 1976, Comp. Biochem. and Physiol., v. 53 (4B), 495-497
Dirofilaria immitis adults, lipid analysis, compared with *Dipetalonema viteae* and *Litomosoides carinii*
- Dirofilaria immitis*
Hutchison, W. F.; and Turner, A. C., 1976, J. Mississippi Acad. Sc., Suppl., v. 21, 47 [Abstract]
Dirofilaria immitis, hexokinase, characterization, kinetics
- Dirofilaria immitis*
Hutchison, W. F.; Turner, A. C.; and Oelshlegel, F. J., jr., 1977, Comp. Biochem. and Physiol., v. 58 (2B), 131-134
Dirofilaria immitis adults, hexokinase, properties
- Dirofilaria immitis*
Iwamoto, I., 1972, Nettai Igaku (Trop. Med.), v. 14 (3), 124-137
Dirofilaria immitis, *Wuchereria bancrofti*, in vitro survival in various media, effects of temperature; *D. immitis* survival in dogs (exper.), periodicity, migration and localization
- Dirofilaria immitis*, illus.
Jackson, R. F.; et al., 1977, J. Am. Vet. Med. Ass., v. 171 (10), 1065-1069
Dirofilaria immitis, dogs (heart), caval syndrome, surgical treatment
- Dirofilaria immitis*
Jankowski, T. J.; and Bickley, W. E., 1976, Ann. Entom. Soc. Am., v. 69 (5), 781-785
Dirofilaria immitis, maturation of larvae in *Aedes canadensis* and *A. vexans*, fed on infected dogs, high vector potential of both mosquitoes: Maryland
- Dirofilaria immitis* (Leidy 1856)
Johnson, C. A. III, 1975, J. Parasitol., v. 61 (5), 940
Ursus americanus (right ventricle, pulmonary artery): Carteret County, N.C.
- Dirofilaria immitis*
Johnson, S., 1975, Southwest. Vet., v. 28 (3), 263-265
Dirofilaria immitis, dog, complete diagnostic regime, case report
- Dirofilaria immitis*
Kan, S. P.; Rajah, K. V.; and Dissanaik, A. S., 1977, Vet. Parasitol., v. 3 (2), 177-181
Dirofilaria immitis, dogs, incidence and degree of infection in survey of 370 animals, differences in relation to pedigree vs. mixed-breed and short-haired vs. long-haired breeds, no age and sex differences: Seremban, Malaysia
- Dirofilaria immitis*
Katamine, D., 1969, Nettai Igaku (Trop. Med.), v. 11 (1), 1-10
Wuchereria bancrofti in humans, skin test diagnosis using purified antigen (FPT) prepared from *Dirofilaria immitis*, useful for tool in mass diagnostic survey
- Dirofilaria immitis*
Keegan, H. L., 1975, J. Mississippi Acad. Sc., v. 20, Suppl., 57
Dirofilaria immitis, dogs, survey, minor public health problem, economic impact of canine dirofilariasis is of considerable importance: Mississippi
- Dirofilaria immitis*
Keegan, H. L., 1977, J. Mississippi Acad. Sc., v. 22, 53-56
Dirofilaria immitis, dogs, prevalence, minor public health problem, considerable economic impact: Mississippi
- Dirofilaria immitis*
Kelly, J. D., 1973, Austral. Vet. J., v. 49 (1), 23-27
Dirofilaria immitis, *Dipetalonema* sp., detection and differentiation of microfilariae in canine blood, comparison of techniques, modified Knott technique and modified filter technique most reliable and consistent for detection; morphological and physiological features considered collectively is recommended for positive identification, striking difference in acid phosphatase activity offers most accurate method of differentiation
- Dirofilaria immitis*
Kimbell, J. W., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 80
prevalence, dogs: Fairfax County, Virginia

- Dirofilaria immitis*, illus.
Kipnis, R. M.; et al., 1976, *Canine Pract.*, Santa Barbara, v. 3 (4), 49-50, 52
Dirofilaria immitis, dog, case report, microfilariae detected in urine, differentiated from casts and mucous threads in urine sediment, thiacetarsamide and Dizan, good results
- Dirofilaria immitis*
Klein, J. B.; and Stoddard, E. D., 1977, *J. Am. Vet. Med. Ass.*, v. 171 (4), 354-355
horse, crossbred colt (pulmonary vessels, right ventricle of heart)
- Dirofilaria immitis*
Kotani, T.; Tomimura, T.; and Mochizuki, H., 1976, *Japan. J. Vet. Sc.*, v. 38 (5), 495-506
Dirofilaria immitis, dogs, postmortem pulmonary angiography, gross and histopathological observations
- Dirofilaria immitis*
Krakowka, S., 1977, *J. Am. Vet. Med. Ass.*, 1977, v. 171 (8), 750-753
transplacentally acquired parasitic diseases of dogs, diagnostic features, review
- Dirofilaria immitis*
Krzaczynski, J.; and Daehler, M. H., 1973, *J. Am. Vet. Med. Ass.*, v. 162 (5), 397-398
Dirofilaria immitis, dog (right atrium, pulmonary artery), contrast radiography as a diagnostic aid
- Dirofilaria immitis*
Lindemann, B. A., 1977, *Mosquito News*, v. 37 (2), 293-295
Dirofilaria immitis encapsulation in *Aedes aegypti*, relationship to larval type and location in mosquitoes
- Dirofilaria immitis*
McCall, J. W.; and Crouthamel, H. H., 1976, *J. Parasitol.*, v. 62 (5), 844-845
Dirofilaria immitis, dogs (exper.), mebendazole 100% effective against developing larvae, potential as prophylactic
- Dirofilaria immitis*
McGreevy, P. B.; McClelland, G. A. H.; and Lavoipierre, M. M. J., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (1), 97-109
Dirofilaria immitis, susceptibility of *Aedes aegypti* controlled by sex-linked recessive gene which is distinct from those controlling development of *Brugia pahangi* or *D. corynodes*, variation in filarial infectivity as well as in mosquito susceptibility
- Dirofilaria immitis*, illus.
McLaren, D. J.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (5-6), 509-514
larval *Dirofilaria immitis* and *Brugia pahangi*, gram-negative micro-organisms within hypodermal tissue, possible adverse affect on development
- Dirofilaria immitis*
Mandelker, L.; and Brutus, R. L., 1971, *J. Am. Vet. Med. Ass.*, v. 159 (6), 776
Dirofilaria immitis, cat (brain), dog (blood, brain), encephalitis
- Dirofilaria immitis*, illus.
Martire, J. R.; Liberman, I. L.; and Goldman, S. M., 1975, *Maryland State Med. J.*, v. 24 (4), 62-63
Dirofilaria immitis, human, cause of pulmonary infarction, case report, clinical diagnosis and management: Maryland
- Dirofilaria immitis*
Miyahara, A.; Chung, N. Y.; and Chung, G., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (10), 1429-1430
Dirofilaria immitis, dogs, increasing incidence, higher incidence in males: Oahu, Hawaii
- Dirofilaria immitis*
Moorhouse, D. E.; et al., 1971, *Med. J. Australia*, v. 2 (24), 1230-1233
Dirofilaria immitis, human infections presenting as pulmonary granulomata of unknown etiology, surgical excision, case reports: Queensland, Australia
- Dirofilaria immitis*
Muscoflat, C. C.; et al., 1977, *Am. J. Vet. Research*, v. 38 (12), 2095-2096
Dirofilaria immitis, dogs, density gradient method of separating lymphocytes, eosinophils, and microfilariae from blood
- Dirofilaria immitis*, illus.
Nayar, J. K.; and Sauerman, D. M., jr., 1975, *J. Insect Physiol.*, v. 21 (12), 1965-1975
Dirofilaria immitis, susceptibility of various species of mosquitoes, patterns of microfilarial development, sites of blockage or prevention of infection, possible physiological factors (substances in salivary gland secretion, melanization of microfilariae)
Aedes taeniorhynchus (exper.)
A. sollicitans (exper.)
A. aegypti (exper.)
Anopheles quadrimaculatus (exper.)
Culex nigripalpus (exper.)
C. pipiens quinquefasciatus (exper.)
Mansonia titillans (exper.)
- Dirofilaria immitis*
Neppert, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Dirofilaria immitis*
Neppert, J.; and Warns, C.-M., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from *Ascaris*, hookworm, *Onchocerca*, *Dirofilaria immitis*, closed hexagonal immunodiffusion, complement fixation reaction, indirect haemagglutination

- Dirofilaria immitis*, illus.
Omar, M. S., 1977, Tropenmed. u. Parasitol., v. 28 (1), 100-108
Wuchereria bancrofti, Brugia malayi, B. pahangi, *Dirofilaria immitis*, distribution of acid phosphatase activity in larval stages in the mosquito, presence or absence of enzymic activity in the excretory cell complex and amphids of developing larvae useful as adjunctive diagnostic method
- Dirofilaria immitis*
Oster, Z., 1976, J. Nuclear Med., v. 17 (5), 425-426
elderly man with supraclavicular mass visible on bone scan, biopsy of mass revealed nematode cysts, probably filariasis of *Dirofilaria immitis* origin
- Dirofilaria immitis*
Otto, G. F.; et al., 1976, J. Am. Vet. Med. Ass., v. 168 (7), 605-607
Dirofilaria immitis, dogs (exper.), number of circulating microfilariae is not an index of the number of adult heartworms or the severity of disease, reduced numbers of microfilariae per adult occur with increased numbers of adults, possible mechanisms
- Dirofilaria immitis*
Portaro, J. K.; et al., 1977, J. Parasitol., v. 63 (1), 172-174
differential response of *Brugia pahangi*-sensitized splenocytes to antigens from *Brugia pahangi*, *Dirofilaria immitis*, and *Trichinella spiralis*, possible diagnostic use
- Dirofilaria immitis*
Rabalais, F. C.; and Votava, C. L., 1972, J. Am. Vet. Med. Ass., v. 160 (2), 202-203
Dirofilaria immitis, *Dipetalonema reconditum*, dogs (blood), prevalence: northwestern Ohio (some dogs originally from southern states)
- Dirofilaria immitis*
Rawlings, C. A.; and Lewis, R. E., 1977, Am. J. Vet. Research, v. 38 (11), 1801-1805
Dirofilaria immitis, dogs with spontaneous infection, evaluation of size of right ventricle by thoracic radiography, electrocardiography, and right ventricular free wall weights; based on radiography, only 2 of 15 dogs had right ventricles of normal size
- Dirofilaria immitis*
Rawlings, C. A.; Losonsky, J. M.; and Lewis, R. E., 1977, Am. J. Vet. Research, v. 38 (9), 1365-1369
dogs infected with *Dirofilaria immitis* but without pulmonary hypertension showed exaggerated increase of pulmonary artery pressure when subjected to hypoxia, apparently due to increased pulmonary vascular resistance; possible implications for pulmonary hypertension in humans
- Dirofilaria immitis*
Rep, B. H.; and Heinemann, D. W., 1976, Trop. and Geogr. Med., v. 28 (2), 104-110
dog: Surinam
- Dirofilaria immitis*
Ridley, D. S.; and Hedge, E. C., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Dirofilaria immitis*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, Am. J. Vet. Research, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections
- Dirofilaria immitis*
Robinson, F. R.; and Garner, F. M., 1973, Am. J. Vet. Research, v. 34 (3), 437-442
dogs, histopathological survey (1964-1971), review of lesions in specimens
- Dirofilaria immitis*
Rodhain-Rebourg, F.; and Rodhain, F., 1972, Medecine et Malad. Infect., v. 2 (8-9), 309-313
review of human infection with *Dirofilaria immitis*, cause of solitary pulmonary nodules
- [*Dirofilaria immitis*] canine heartworm
Rosenberg, M. A.; et al., 1976, Vet. Med. and Small Animal Clin., v. 71 (4), 496, 498-503
[*Dirofilaria immitis*], evaluation of prognosis and treatment, mathematical formula
- Dirofilaria immitis*, illus.
Sarkar, P.; Basak, D. K.; and Bhattacharyya, H. M., 1976, Indian Vet. J., v. 53 (1), 55-57
Dirofilaria immitis, dogs (heart), pathology
- Dirofilaria immitis*
Sawada, T.; et al., 1975, Progr. Drug Research, v. 19, 128-135
human filariasis, diagnosis using *Dirofilaria immitis* adult worm antigen for skin tests, purification of antigen
- Dirofilaria immitis*
Scheib, C. W.; and Corwin, D. S., 1977, Vet. Med. and Small Animal Clin., v. 72 (10), 1594-1595
dog (right ventricle, pulmonary artery, lung parenchyma), clinical signs, no microfilariae found in blood, diagnostic difficulty: Maine
- Dirofilaria immitis* (Leidy)
Seeley, D. C., jr.; and Bickley, W. E., 1974, Proc. and Papers 42. Ann. Conf. Calif. Mosquito Control Ass. and 30. Ann. Meet. Am. Mosquito Control Ass. (Anaheim, Feb. 24-27), 87-92
Dirofilaria immitis in 3 strains of *Culex salinarius* (exper.), potential vector

- Dirofilaria immitis*
Sengbusch, H.; Sartori, P.; and Wade, S., 1975, *Am. J. Vet. Research*, v. 36 (7), 1035-1036
Dirofilaria immitis, 2 of 100 stray dogs infected, none with *Dipetalonema* sp., first report of heartworm in domesticated dogs in the state: Buffalo, New York
- Dirofilaria immitis*
Sivanandam, S.; and Sandosham, A. A., 1965, *Med. J. Malaya*, v. 20 (1), 65
multiple filarial infections in domestic cat: East Pahang, Malaya
- Dirofilaria immitis*, *illus.*
Slonka, G. F.; Castleman, W.; and Krum, S., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (7), 717-719
Dirofilaria immitis, dog, mature worms in femoral and other systemic arteries, ischemic necrosis to abdominal viscera and hindlimbs, posterior paresis and paralysis
- Dirofilaria immitis*
Smith, M. W., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 18 [Demonstration]
dogs: Trinidad
- Dirofilaria immitis*, *illus.*
Soifer, F. K., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (4), 484, 486
Dirofilaria immitis, cat, also positive for feline leukemia virus, possible depression of immune response mechanism: Houston area, Texas
- Dirofilaria immitis*
Strauss, J. M.; and Sivanandam, S., 1966, *Med. J. Malaya*, v. 20 (4), 336
mixed *Brugia pahangi* and *Dirofilaria immitis* infections in *Panthera pardus*: National Zoo, Kuala Lumpur (captured in state of Pahang)
- Dirofilaria immitis*
Streitel, R. H.; Stromberg, P. C.; and Dubey, J. P., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (7), 720-721
prevalence in dogs stable since 1964, no highly enzootic center: humane shelter, Columbus, Ohio
- Dirofilaria immitis*
Suenaga, O., 1972, *Nettai Igaku (Trop. Med.)*, v. 14 (1), 32-40
development of *Dirofilaria immitis* larvae in potential mosquito vectors, effects of temperature and season
Aedes togoi
Culex pipiens pallens
C. tritaeniorhynchus summorosus
Armigeres subalbatus
(all exper.)
- Dirofilaria immitis*
Suenaga, O., 1972, *Nettai Igaku (Trop. Med.)*, v. 14 (3), 144-150
Aedes albopictus (exper.)
Culex pipiens molestus (exper.)
- Dirofilaria immitis*
Suenaga, O., 1973, *Nettai Igaku (Trop. Med.)*, v. 15 (2), 84-91
Dirofilaria immitis, laboratory trials, development in local mosquitoes; high experimental infection rate in *Anopheles sinensis*, low in *Aedes vexans nipponii*: Nagasaki City
- Dirofilaria immitis*
Suenaga, O., 1975, *Nettai Igaku (Trop. Med.)*, v. 17 (1), 35-40
Dirofilaria immitis, prevalence survey in dogs and vector mosquitoes
Culex pipiens pallens
Aedes albopictus
Anopheles sinensis
Culex tritaeniorhynchus summorosus: all from Nagasaki Prefecture
- Dirofilaria immitis*
Suenaga, O.; and Itoh, T., 1973, *Nettai Igaku (Trop. Med.)*, v. 15 (3), 131-140
Dirofilaria immitis, field collections of possible vector mosquitoes to determine natural infections
Culex pipiens pallens
Aedes albopictus
Culex tritaeniorhynchus summorosus
Aedes vexans nipponii
all from Nagasaki City areas
- Dirofilaria immitis*
Suenaga, O.; and Itoh, T., 1976, *Nettai Igaku (Trop. Med.)*, v. 18 (1), 59-63
Dirofilaria immitis, possible importance of *Aedes albopictus* and *Culex pipiens pallens* in transmitting canine heartworm to man: Nagasaki City, Japan
- Dirofilaria immitis*
Suenaga, O.; Itoh, T.; and Nishioka, T., 1971, *Nettai Igaku (Trop. Med.)*, v. 12 (4), 169-178
Dirofilaria immitis, prevalence survey in domestic dogs, seasonal distribution: Nagasaki City, Japan
- Dirofilaria immitis*
Suenaga, O.; Kamahara, H.; and Shibata, M., 1974, *Nettai Igaku (Trop. Med.)*, v. 16 (2), 95-101
Dirofilaria immitis in house dogs, prevalence survey for microfilariae in peripheral blood: Omura City, Japan
- Dirofilaria immitis*
Takahashi, J.; and Sato, K., 1976, *Japan. J. Exper. Med.*, v. 46 (1), 7-13
fractionation and purification of *Dirofilaria immitis* antigens by column chromatography and disc electrophoresis, evaluation for use in diagnosis of human *Wuchereria bancrofti* by hemagglutination test
- Dirofilaria immitis*
Terwedow, H. A., jr.; and Craig, G. B., jr., 1977, *Exper. Parasitol.*, v. 41 (2), 272-282
Waltonella flexicauda, development controlled by genetic factor in *Aedes aegypti*; this factor for susceptibility did not control development of *Brugia pahangi* or *Dirofilaria immitis*

- Dirofilaria immitis*
Theis, J. H.; and McGreevy, P. B., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 32 [Demonstration]
description of film on development and transmission
- Dirofilaria immitis*
Thornton, J. E.; Bell, R. R.; and Reardon, M. J., 1974, J. Wildlife Dis., v. 10 (3), 232-236
Canis latrans (right ventricle and pulmonary artery): Nueces County, Texas
- Dirofilaria immitis* Leidy
Todaro, W. S.; Morris, C. D.; and Heacock, N. A., 1977, Am. J. Vet. Research, v. 38 (8), 1197-1200
Dirofilaria immitis, prevalence in dogs and mosquitoes, seasonal distribution, potential vectors
dogs (blood)
Aedes canadensis (exper.)
A. vexans (nat. and exper.)
A. triseriatus (exper.)
A. aegypti (exper.)
Anopheles quadrimaculatus (nat. and exper.)
Culex pipiens-restuans group (exper.)
Coquillettidia perturbans (exper.)
all from central New York
- Dirofilaria immitis*, illus.
Todd, K. S., jr.; et al., 1976, Feline Pract., v. 6 (2), 41-44
Dirofilaria immitis, 3 cats, case histories, pathology, literature review: Illinois
- Dirofilaria immitis*
Turner, A. C.; et al., 1975, J. Mississippi Acad. Sc., v. 20, Suppl., 25
Dirofilaria immitis, dogs, lipid content of worms, total fatty acid methyl esters in *Dirofilaria immitis*, compared with *Dipetalonema viteae* and *Litomosoides carinii*, similar fatty acid patterns, gas chromatography
- Dirofilaria immitis*
Watson, A. D. J.; Porges, W. L.; and Testoni, F. J., 1972, Med. J. Australia, v. 1 (3), 141
increasing incidence of *Dirofilaria immitis* in canines in Sydney, Australia area, need for awareness as possible zoonosis
- Dirofilaria immitis*
Watson, A. D. J.; Porges, W. L.; and Testoni, F. J., 1973, Austral. Vet. J., v. 49 (1), 31-34
dogs, survey of incidence of *Dipetalonema reconditum* and *Dirofilaria immitis* microfilariae and adult *D. immitis*: Sydney, Australia
- Dirofilaria immitis*
Watson, A. D. J.; Testoni, F. J.; and Porges, W. L., 1973, Austral. Vet. J., v. 49 (1), 28-30
Dirofilaria immitis, *Dipetalonema reconditum*, comparison of microfilarial characteristics isolated by modified Knott and filter methods, results suggest consideration of mean microfilarial length and numbers on filter may permit accurate differentiation of microfilariae in canine blood
- Dirofilaria immitis*
Weinmann, C. J.; and Garcia, R., 1975, Proc. and Papers 43. Ann. Conf. Calif. Mosquito Control Ass. (Redding, Jan. 26-29), 96 [Abstract]
Aedes sierrensis and *A. dorsalis* as vectors of *Dirofilaria immitis*
- Dirofilaria immitis*
Welch, J. S.; and Dobson, C., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (6), 466-472
comparative fluorescent antibody test survey of Aborigines and Caucasians for presence of antibodies to *Dirofilaria immitis* and correlations with canine filariasis; cross-reactions to *Toxocara canis* observed only in presence of eosinophilia: Queensland, Australia
- Dirofilaria immitis*
Wilkerson, M., 1976, J. Elisha Mitchell Scient. Soc., v. 92 (2), 49 [Abstract]
Dirofilaria immitis, dogs, prevalence; large, outside dogs have highest probability for infection; higher percentage positive for microfilaria during July-November than for entire year of 1975
- Dirofilaria immitis*
Williams, J. F.; et al., 1977, J. Am. Vet. Med. Ass., v. 170 (7), 714-716
Dirofilaria immitis, dogs, diagnostic techniques, evaluation of polycarbonate filter found superior to cellulose filter system
- Dirofilaria immitis*
Williams, J. F.; and Dade, A. W., 1976, J. Parasitol., v. 62 (1), 174-175
Dirofilaria immitis, severe vascular and pulmonary disease, *Gulo luscus* (right ventricle, right atrium, pulmonary artery, posterior vena cava): imported from Ontario, Canada to zoo, Lansing, Michigan
- Dirofilaria immitis*, illus.
Wong, M. M., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (4), 480-486
Dirofilaria immitis, dirofilariasis without microfilaremia produced in dogs (exper.), immunologic responses analogized to etiology of tropical eosinophilic lung in humans
- Dirofilaria immitis*, illus.
Wong, M. M., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (6), 479-490
Dirofilaria immitis in normal and immunosuppressed *Macaca* spp., histopathology, W.B.C. and eosinophil counts, radiological examination, serological examination by IFA test, recovery of adult worms only in immunosuppressed hosts indicates that host susceptibility rather than parasite infectivity was the factor concerned
Macaca fascicularis
M. arctoides
M. mulatta
(all exper.)
- Dirofilaria immitis*
Wong, M. M.; and Lim, K. C., 1975, J. Parasitol., v. 61 (3), 573-574
Dirofilaria immitis, development of intraperitoneally inoculated larvae in *Meriones unguiculatus*

- [*Dirofilaria immitis*] heartworm, *illus.*
Yamashita, H.; et al., 1975, *Bull. Nippon Vet. and Zootech. Coll.* (24), 94-101
surgical removal of heartworms by right auriculotomy and puncture of pulmonary artery, dogs
- Dirofilaria immitis*, *illus.*
Yoshihara, T.; et al., 1977, *Exper. Rep. Equine Health Lab.* (14), 1-12
Dirofilaria immitis, horses (heart, lung), parasite morphology and measurements, pathology: Japan
- Dirofilaria immitis*
Zielke, E., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (1), 32-35
Dirofilaria immitis, mechanism of transmission by *Aedes togoi* (exper.), route of invasion via puncture of mosquito bite
- Dirofilaria immitis*
Zielke, E., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (1), 36-44
Culex pipiens fatigans, *Aedes aegypti*, selection of strains differing in susceptibility to *Dirofilaria immitis*, demonstrated that inheritance of susceptibility in *Aedes aegypti* is controlled by sex-linked recessive gene, also found that susceptibility to infection with *Brugia pahangi* is on same chromosome but in different locus
- Dirofilaria immitis*
Zielke, E., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (2), 214-221
Dirofilaria immitis, quantitative aspects of transmission by *Anopheles atroparvus* and *Aedes togoi*, intake of microfilariae by mosquitoes is directly proportional to number of microfilariae in host blood and the amount of blood taken up by the insect, infectious potential considerably reduced by increased mortality rates of infected mosquitoes
- Dirofilaria immitis*
Zielke, E., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (2), 243-244
Dirofilaria immitis, preliminary trials show that parasite can be transplanted from dog and survive for some time in peritoneal cavity of laboratory rodents (rat, mouse, guinea-pigs, *Meriones*, *Mastomys*) with some migration outside peritoneal cavity
- Dirofilaria immitis*
Zymet, C. L., 1977, *Vet. Med. and Small Animal Clin.*, v. 72 (12), 1848-1852
Dirofilaria immitis, dog, shock syndrome resulting from administration of diethyl-carbamazine citrate, case history, tentative diagnosis of Addisonian crisis
- Dirofilaria magnilarvata*
Cheong, W. H.; and Omar, A. H. B., 1966, *Med. J. Malaya*, v. 20 (4), 332
some early findings on the possibility of using *Aedes togoi* as a vector of *Dirofilaria magnilarvata* and *Dirofilaria repens* in the laboratory
- Dirofilaria magnilarvatum*
Mullin, S. W.; et al., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 548-551
Presbytis obscura
P. cristata
Macaca fascicularis
all from Malaysia
- Dirofilaria repens*
Bain, O., 1976, *Bull. World Health Organ.*, v. 54 (4), 397-401
human filariasis, number of developing and infective larvae dependent upon number of microfilariae penetrating into haemocyte of vector, relationship based on proportionality, facilitation and limitation, application to disease control and treatment methods
- Dirofilaria repens*, *illus.*
Balbo, T.; and Abate, O., 1972, *Parassitologia*, v. 14 (2-3), 239-244
Dirofilaria immitis, *D. repens*, *Dipetalonema* sp., microfilaria from dogs, staining for localization of acid phosphatase, detailed procedure, basis for diagnostic differentiation
- Dirofilaria repens*
Bwangamoi, O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (4), 363-370
Dirofilaria repens, *Dipetalonema reconditum*, recovery from dogs using Lindsey's method: Uganda
- Dirofilaria repens*
Bwangamoi, O.; and Isyagi, A. O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (1), 33-37
filariasis, dogs, incidence survey, morbidity rate per breed, age, and sex of host: Uganda
- Dirofilaria repens*
Cheong, W. H.; and Omar, A. H. B., 1966, *Med. J. Malaya*, v. 20 (4), 332
some early findings on the possibility of using *Aedes togoi* as a vector of *Dirofilaria magnilarvata* and *Dirofilaria repens* in the laboratory
- Dirofilaria repens*
Javadian, E.; and Macdonald, W. W., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (4), 477-481
Brugia pahangi, *Dirofilaria repens*, infection as cause of reduced egg-production of *Aedes aegypti*
- Dirofilaria repens*
Schillhorn van Veen, T.; and Blotkamp, J., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (4), 517-518
"evidence strongly suggests that they [microfilariae] are *D. repens*"
dogs: Zaria area, Nigeria

- Dirofilaria repens*, *illus.*
Vodovozov, A. M.; Jarulin, G. R.; and Djakonova, S. W., 1973, *Ophthalmologica*, Basel, v. 166 (2), 88-93
Dirofilaria repens, extraction of living intra-ocular nematode from man's eye with preservation of vision, clinical case report: Wolgograd (UdSSR)
- Dirofilaria repens*, *illus.*
Wong, M. M., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (1), 88-93
Dirofilaria repens, healthy and immunosuppressed macaques (exper.), larvae and adult worms recovered in 10 of 13 but microfilaria seen only in prednisolone-treated animals, host responses (eosinophilia and filarial antibodies)
Macaca fascicularis (exper.)
Macaca nemestrina (exper.)
Macaca mulatta (exper.)
Macaca arctoides (exper.)
- Dirofilaria repens*, *illus.*
Zeybek, H.; and Oge, A., 1977, *Vet. Hekim. Dernegi Dergisi*, v. 47 (1), 52-54
human, woman (eye): Bafra county, Samsun, Turkey
- Dirofilaria tenuis*, *illus.*
Christie, R. W., 1977, *N. England J. Med.*, v. 297 (13), 706-707
Dirofilaria tenuis infection in woman presenting as periorbital swelling and nodule over right eye, prevalence of mosquitoes and raccoons in area suggests raccoon-to-human transmission via mosquito vector: Holland, Vermont
- Dirofilaria tenuis*, probably, *illus.*
Davies, P.; et al., 1973, *N. York State J. Med.*, v. 73 (15), 1999-2001
probable *Dirofilaria tenuis* infection in woman with presenting symptom of tender, painful swelling over right temple, apparent cure after surgical biopsy and follow-up diethylcarbamazine, history of extensive bites from mosquitoes and association with dog that had recently been in southern states: Albany, New York
- Dirofilaria tenuis*, *illus.*
Wong, M. M.; and Lim, K. C., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (1), 94-98
Dirofilaria tenuis, healthy and immunosuppressed macaques (exper.), larvae and adult worms recovered from 7 of 11 but microfilaria seen only in a prednisolone-treated animal, host responses (eosinophilia and filarial antibodies)
Macaca arctoides (exper.)
Macaca mulatta (exper.)
- Dirofilaria timidi* nov. sp., *illus.*
Gubanov, N. M.; and Fedorov, K. P., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 47-48
Lepus timidus (thoracic cavity): Central Yakut
- Dirofilaria ursi* Yamaguti, 1941
Rogers, L. L., 1975, *J. Wildlife Dis.*, v. 11 (2), 189-192
Ursus americanus (connective tissue, peritoneal cavities, blood): Michigan; Minnesota
- Dirofilariaeformia pulmonis*
Davidson, W. R., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (pulmonary arteries): southeastern United States
- Dirofilariasis*
de Carneri, I.; Sacchi, S.; and Pazzaglia, A., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (6), 887-888 [Letter]
subcutaneous dirofilariasis in man, more widespread than previously believed, definitely zoonotic
- Dirofilariasis*
Humphries, J. P.; and Goodnight, D. B., 1977, *Southwest. Vet.*, v. 30 (1), 40-51
reference outline to common heart disease syndromes in dogs
- Dirofilariasis*
Ishihara, K.; et al., 1977, *Japan. J. Vet. Sc.*, v. 39 (3), 255-264
dirofilariasis, canine, serum lipoproteins in relation to stage and intensity of infection, possible clinical and diagnostic value
- Dirofilariasis*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiasis associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Diserratosomus Mirza*, 1933
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
"corresponds possibly to the subgenus *Mesopectines* Quentin, 1969 but since this is uncertain we retain the taxon *Mesopectines*."
- Dispharynx*, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Synhimantus*
key
Syn.: *Aliella Ali*, 1968
- Dispharynx* sp.
Alekseev, V. M.; and Smetanina, Z. B., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 97-104
description
Larus crassirostris (muscles): Rimsko-Korsakov islands

- Dispharynx nasuta (Rudolphi, 1819) Henry and Sisoff 1912
Christensen, Z. D.; and Pence, D. B., 1977, J. Parasitol., v. 63 (5), 830
Ortalis vetula maccalli: near San Benito, Cameron Co., Texas
- Dispharynx nasuta
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (proventriculus): South Bass Island, Ohio
- Dispharynx nasuta
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 399-403
survey, helminths of red-winged blackbirds including a check list of previous findings
Agelaius phoeniceus (proventriculus): South Bass Island, Ohio
- Dispharynx nasuta (Rudolphi, 1819)
Cooper, C. L.; and Crites, J. L., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (proventriculus): South Bass Island, Ottawa County, Ohio
- Dispharynx nasuta
Cooper, C. L.; and Crites, J. L., 1976, J. Parasitol., v. 62 (1), 105-110
similarity index of helminth faunas of 7 passerine bird species, index of association of 10 species of helminths identified as having foci of infection, competition for invertebrate food resources and aggregation into mixed feeding flocks maximizes transmission: South Bass Island, Ottawa County, Ohio
- Dispharynx nasuta
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (proventriculus): Franklin and Ottawa counties, Ohio
- Dispharynx nasuta (Rudolphi, 1819)
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (proventriculus): Florida
- Dispharynx nasuta
Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, J. Parasitol., v. 61 (3), 547-548
Grus canadensis pratensis (proventriculus): Florida
- Dispharynx nasuta
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (proventriculus)
Colinus virginianus
Corvus brachyrhynchos
Cyanocitta cristata
all from Florida
- Dispharynx nasuta (Rudolphi, 1819)
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coerulescens (proventriculus): Florida
- Dispharynx nasuta (Rudolphi, 1819), illus.
Led, J. E.; and Brandetti, E., 1972, *Analecta Vet.*, v. 4 (1), 37-40
pavo (estomago glandular): environs of La Plata City
- Dispharynx nasuta
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: southeastern United States
- Dispharynx nasuta (Rudolphi, 1819)
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus genei: Azov Sea
- Diximermis Nickle
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
"Diximermis peterseni and Perutilimermis culicis must be regarded as genera et species inquirendarum."
- Diximermis peterseni Nickle
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
"Diximermis peterseni and Perutilimermis culicis must be regarded as genera et species inquirendarum."
- Diximermis peterseni
Woodard, D. B.; and Fukuda, T., 1977, Mosquito News, v. 37 (2), 192-195
Diximermis peterseni as biological control agent for *Anopheles quadrimaculatus*, laboratory resistance, mechanism is behavioural (avoidance of attack and snapping at nematodes by mosquito larvae during exposure)
- Dogielina Sobolev, 1949
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
"seem to belong to the Physalopteroidea but, as yet, are not well enough known to be classified"
- Dogielina Sobolev, 1949
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Physalopteridae, Physalopterinae
key
- Dollfusstrongylus Quentin, 1970
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Dorcopsinema n. gen.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
Trichonematidae; Zoniolaiminae
tod: *D. dorcopsis* (Baylis) n. comb.
- Dorcopsinema dorcopsis* (Baylis) n. comb. (tod), illus.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
Syn.: *Macropostrongylus dorcopsis* Baylis, 1940

- Dorylaimida Pearse, 1942
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Enoplia
includes: Dorylaimina; Alaimina
- Dorylaimina Pearse, 1936
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Dorylaimida
- Dorylaimoidea, illus.
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
- Dorylaimus sp. (?) Dujardin, 1845
Amin, O. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 81-88
accidental infection
Catostomus commersoni (intestine): southeastern Wisconsin
- Dorylaimus sp. (?)
Amin, O. M., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 43-46
Catostomus commersoni: southeastern Wisconsin
- Draconematina DeConinck, 1965
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Desmodorida
- Dracunculidae (Stiles, 1907, subfam.) Leiper, 1912
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Dracunculoidea
key; key to genera
includes: Avioserpens; Dracunculus
- Dracunculoidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Camallanina
key
- Dracunculoidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanina
key to families
includes: Anguillicolidae; Guyanemidae; Philometridae; Micropleuridae; Dracunculidae
- Dracunculosis, illus.
Dorfmann, H.; and de Seze, C., 1972, Nouv. Presse Med., v. 1 (15), 1013-1016
dracunculosis in man, filarial arthritic infection in man's knee joint diagnosed by arthroscopy, clinical case report: France (had resided in Mali)
- Dracunculosis
Niel, G.; et al., 1972, Medecine et Malad. Infect., v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalonema vitae* and *Ascaris suum* as antigens
- Dracunculosis
Pinon, J. M.; and Gentilini, M., 1973, Nouv. Presse Med., v. 2 (19), 1283-1287
human filariasis, application of cellular immunologic tests (rosette formation, macrophage migration) in diagnosis and comparison with serologic tests (fluorescent antibody, passive hemagglutination, gel diffusion)
- Dracunculosis
Richet, P., 1973, Medecine Afrique Noire, v. 20 (11), 899-920
guidelines for mass therapy in human filarial infections, drugs in current use, review
- Dracunculosis
Ricosse, J. H.; and Picq, J. J., 1973, Medecine Afrique Noire, v. 20 (11), 877-897
human filariasis, clinical indications for treatment, drugs in current use, review
- Dracunculus (Reichard, 1759), illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Dracunculidae
key; synonymy
- Dracunculus insignis
Beverley-Burton, M.; and Crichton, V.F.J., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 152 [Letter]
infection of rhesus monkey with *Dracunculus insignis* obtained originally from wild-caught raccoon and subsequently cycled in mink (exper.), "This preliminary observation suggests that *D. medinensis* and *D. insignis* may be conspecific. . ."
- Dracunculus insignis
Beverley-Burton, M.; and Crichton, V. F. J., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 704-708
Dracunculus insignis, *D. medinensis*, variations in susceptibility of mammalian hosts and migration routes and sites of emergence of parasites, may represent physiological strains of single species or may in fact be two distinct species
Macaca mulatta (exper.)
Mustela putorius furo (exper.)
- Dracunculus insignis
Georgi, J. R.; et al., 1976, Cornell Vet., v. 66 (3), 309-323
Procyon lotor: North Rose, Wayne County, New York

- [*Dracunculus medinensis*] guinea worm
Belcher, D. W.; Wurapa, F. K.; and Ward, W. B., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (3), 444-446
thiabendazole and metronidazole unsatisfactory in treatment of human guinea worm in exper. trials comparing effectiveness, cost and side-effects: Accra, Ghana
- Dracunculus medinensis*
Beverley-Burton, M.; and Crichton, V.F.J., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (1), 152 [Letter]
infection of rhesus monkey with *Dracunculus insignis* obtained originally from wild-caught raccoon and subsequently cycled in mink (exper.), "This preliminary observation suggests that *D. medinensis* and *D. insignis* may be conspecific. . ."
- Dracunculus medinensis*
Beverley-Burton, M.; and Crichton, V. F. J., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (5), 704-708
Dracunculus insignis, *D. medinensis*, variations in susceptibility of mammalian hosts and migration routes and sites of emergence of parasites, may represent physiological strains of single species or may in fact be two distinct species
- Dracunculus medinensis*
Bourrel, P.; et al., 1972, *Medecine Trop.*, v. 32 (2), 169-174
Dracunculus medinensis, localized in human scrotum, diagnosis and clinical management, surgical treatment
- Dracunculus medinensis*, *illus.*
Bourrel, P.; Cerutti, J.; and Damas, R., 1975, *Medecine Trop.*, v. 35 (1), 55-66
Dracunculus medinensis, aberrant and typical migration patterns, differential diagnosis, complications, clinical aspects, prognosis and medical management
- Dracunculus medinensis*
Bourrel, P.; and Delatte, P., 1972, *Medecine Trop.*, v. 32 (3), 291-294
calcified filaria of *Dracunculus medinensis* localized in boney areas and joints, differential diagnosis from osteo-arthritis, humans
- Dracunculus medinensis*
Giordano, C.; et al., 1976, *Medecine Afrique Noire*, v. 23 (2), 83-87
case reports of human spinal cord compression by *Dracunculus medinensis*, clinical aspects, medical management: Ivory coast
- [*Dracunculus medinensis*] guinea worm
Kale, O. O., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (1), 91-95
dracontiasis, efficacy of niridazole and metronidazole, controlled field trial: southeast of Ibadan, Western State, Nigeria
- Dracunculus medinensis*
Kale, O. O., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (4), 600-605
Dracunculus medinensis, mebendazole field trials, humans, safe and effective for mass therapy, eliminates adult worms and prevents clinical relapses but not significantly effective in eliminating symptoms and healing ulcers
- Dracunculus medinensis*
Kale, O. O., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (2), 208-214
Dracunculus medinensis, clinico-epidemiologic survey of guinea worm infection in native population, economic and occupational importance, possible control measures by provision of wholesome water supplies: Ibadan district, Nigeria
- [*Dracunculus medinensis*] guinea worm
Kulkarni, D. R.; and Nagalotimath, S. J., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 169-170 [Letter]
guinea worm infection in humans, trials with metronidazole unsuccessful: India
- Dracunculus medinensis*, *illus.*
Muller, R., 1976, *Pathophysiol. Parasit. Infect.*, 133-147
Dracunculus medinensis in rhesus monkeys, pathological processes associated with emergence of female worms, chemotherapy (some drugs had no effect on pre-emergent worms but markedly reduced host tissue reaction thus allowing easier extraction)
+*Macacca mulatta* (exper.)
Cyclops vernalis americanus (exper.)
+cats (exper.)
- [*Dracunculus medinensis*] guinea worm
Odei, M. A., 1975, *Ghana J. Sc.*, v. 15 (2), 219-224
Schistosoma haematobium and guinea worm infections in humans, prospects for increased disease incidence with construction of Weiija Dam and suggested methods for control: Ghana
- Dracunculus medinensis* (Linn, 1758), *illus.*
Pardanani, D. S.; et al., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (1), 45-52
Dracunculus medinensis in humans, 85% cure rate in double blind clinical study using metronidazole, drug assessed with infections in various stages, treatment well tolerated and without serious side effects: India
- Dracunculus medinensis*, *illus.*
St. George, J., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (3), 383-386
Dracunculus medinensis, retroplacental localization as cause of bleeding in pregnancy and possible cause of habitual abortion, 2 case reports: Nigeria
- Dracunculus* [*medinensis*]
Shafei, A. Z., 1976, *J. Trop. Med. and Hyg.*, v. 79 (9), 197-200
Dracunculus [*medinensis*] adult worms, humans, mebendazole effective in clinical trials: Nigeria

- Dracunculus medinensis*
Subrahmanyam, B.; Reddy, Y. R.; and Paul, S., 1976, Indian Vet. J., v. 53 (8), 637-639
Dracunculus medinensis, dog, case report, Flagyl, good results: Kurnool town
- Draschearia* Skrjabin, Sobolev & Ivashkin, 1965
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"cannot be retained . . . probably a synonym of *Procyrnea*"
- Draschia* Chitwood & Wehr, 1934, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
- Draschia megastoma*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Draschia megastoma*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Draschia megastoma*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Draschia megastoma*
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxfendazole
- Dromaeostrongylus* Lubimov, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Dudekemia longicarpus* n. sp., illus.
Rao, V. J., 1973, Riv. Parassitol., Roma, v. 34 (4), 287-290
Spirostreptus sp. (intestine): Mananoor, Andhra Pradesh, India
- Dujardinascaris* Campana-Rouget, 1960
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Multicaecinae
key; key to genera
includes: *Hartwichia*; *Dujardinascaris*
- Dujardinascaris* Baylis, 1927
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Dujardinascaridinea
key; synonymy
- Dujardinascaris* Baylis, 1947
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
redescription
- Dujardinascaris alata*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris antipini*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris australiensis*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris chabaudi*, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
"there is a probability that paulista and chabaudi are identical"
Caiman crocodilus (= scleros) (stomach): Barquisimeto, Venezuela
- Dujardinascaris dujardini* (Travassos, 1920), illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
synonymy
- Dujardinascaris dujardini madagascariensis* Chabaud & Caballero, 1966
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
as syn. of *D. madagascariensis* Chabaud & Caballero, 1966 [? n. rank]
- Dujardinascaris gedoelsti* n. sp., illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Syn.: *D. helicina* of Gedoelst, 1916 (in part)
Crocodylus niloticus (stomach, intestine): Leopoldville; Djoko-Punda, Zaire; Dolo, Zaire
- Dujardinascaris helicina* of Gedoelst, 1916 (in part)
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
as syn. of *D. gedoelsti* n. sp.
- Dujardinascaris helicina* (Molin, 1860) non *D. helicina* of Gedoelst, 1916; Baylis, 1923; Walton, 1927, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Crocodylus acutus (stomach): Amsterdam Zoo; San Diego Zoo

- Dujardinascaris longispicula* Travassos, 1933, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Caiman crocodilus (= sclerops) (stomach): Rio S. Laurencio, Brazil; Barquisimeto, Venezuela
- Dujardinascaris madagascariensis* Chabaud & Caballero, 1966 [? n. rank], illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Syn.: *D. dujardini* madagascariensis Chabaud & Caballero, 1966
Crocodylus niloticus (stomach): Mairgodo, Madagascar
C. cataphractus (stomach)
- Dujardinascaris mawsonae* n. sp., illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Crocodylus novaeguineae: Lake Murray and Central Province, Papua New Guinea
C. porosus: Arnhem Land, Northern Australia
C. johnstoni?: Herbert River, Queensland (stomach of all)
- Dujardinascaris paulista* Travassos, 1933, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
"there is a probability that paulista and chabaudi are identical"
- Dujardinascaris puylaerti*, n. sp., illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Crocodylus niloticus (stomach): Boma, Zaire; Zambia
- Dujardinascaris qadrii* n. sp., illus.
Zubairi, H. B.; and Farooq, M., 1976, Pakistan J. Zool., v. 8 (1), 73-76
Sciaena sp. (intestine): Karachi coast, Pakistan
- Dujardinascaris ritai* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Rita rita (mesentery): Lahore, Pakistan
- Dujardinascaris salomonis*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris tasmani*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris taylorae* n. sp., illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Crocodylus porosus: Arnhem Land, Australia
C. novaeguineae: Papua New Guinea (stomach of all)
- Dujardinascaris vandenbrandeni*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
excluded from genus *Dujardinascaris*
- Dujardinascaris waltoni* n. sp. non *Dujardinascaris helicina* of Walton, 1927, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Alligator mississippiensis (stomach, intestine): Victoria County, Texas, U.S.A.; Florida, U.S.A.; Louisiana U.S.A.
- Dujardinascaris woodlandi* Baylis, 1923, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (3), 253-287
Gavialis gangeticus (stomach): River Ganges, India
- Dujardinia* Gedoelst, 1916, in part, nec Quatre-fages, 1844
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Dujardinascaris* Baylis, 1927
- Dujardinia* Gedoelst, 1916, in part, nec Quatre-fages, 1844
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Paradujardinia* Travassos, 1933
- Duplicaecum* Majumdar & Chakravarty, 1963
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Contraecinea
key

- Echinocephalus* Molin, 1858, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatinae
key
- Echinocephalus* sp.
Anantaraman, S., 1963, *J. Marine Biol. Ass. India*, v. 5 (1), 137-139
Hemifusus pugilinus: Madras Coast
- Echinocephalus sinensis* Ko, 1975, *illus.*
Ko, R. C., 1976, *Canad. J. Zool.*, v. 54 (4), 597-609
Echinocephalus sinensis, seasonal variation in incidence and intensity in *Crassostrea gigas*, seasonal variation in infectivity to exper. mammal hosts possibly dependent upon ambient temperature, pathology in mammals, implications for possible human infection from eating poorly cooked oysters
Crassostrea gigas (adductor muscle, genital duct, Leydig tissue): Hong Kong
Kittens (exper.)
puppies (exper.)
Macacca mulatta (exper.)
M. nemestrini (exper.)
- Echinocephalus spinosissimus* (Linstow, 1905)
Shafee, M. S.; and Natarajan, R., 1976, *Current Sc.*, Bangalore, v. 45 (9), 339-340
[Letter]
brief description
Aetobatis narinari (inner wall of intestine): Bay of Bengal, Porto Novo
- Echinocephalus uncinatus*, *illus.*
Bhaibulaya, M.; and Indra-ngarm, S., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 118-119
Echinocephalus uncinatus, morphometric data, electron microscopy
Pentapodus setosus (cyst on surface of intestinal wall): Gulf of Thailand
- Echinocephalus uncinatus* (Molin, 1858)
Shafee, M. S.; and Natarajan, R., 1976, *Current Sc.*, Bangalore, v. 45 (9), 339-340
[Letter]
Aetobatis narinari (inner wall of stomach): Bay of Bengal, Porto Novo
- Echinuria* Soloviev, 1912, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
Syn.: Hamannia Railliet, Henry & Sisoff, 1912
- Echinuria borealis* (Mawson, 1956)
Bishop, C. A.; and Threlfall, W., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 25-35
Somateria mollissima: insular Newfoundland and/or southern Labrador
- Echinuria heterobrachiata* Wehr, 1937
Keppner, E. J., 1973, *Tr. Am. Micr. Soc.*, v. 92 (2), 288-291
Larus californicus: city dump of Laramie, Wyoming
- Echinuria uncinata* (Rudolphi, 1819)
Bishop, C. A.; and Threlfall, W., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 25-35
Somateria mollissima (proventriculus, gizzard): insular Newfoundland and/or southern Labrador
- Echinuria uncinata*
George, R. R.; and Bolen, E. G., 1975, *J. Wildlife Dis.*, v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas
- Echinuria uncinata* (Rudolphi, 1819) Soloviev, 1912
Kamburov, P.; and Vasilev, I., 1972, *Izvest. Tsentral. Khelmit. Lab.*, v. 15, 109-133
Anas platyrhynchos
Aythya nyroca
Mergus serrator
(small intestine of all): all from Bulgaria
- Echinuria uncinata*, *illus.*
Misiura, M., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 247-251
Echinuria uncinata, larval development in *Daphnia pulex*, *D. magna* and *Heterocypris incongruens* (all exper.) (hemocoel of all)
Cygnus olor (proventriculus): Zoological Garden in Warszawa
- Edesonfilaria malayensis* Yeh, 1960, *illus.*
Mullin, S. W., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (2), 256
Edesonfilaria malayensis, morphologic description of microfilariae
- Edesonfilaria malayensis*
Mullin, S. W.; et al., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 548-551
Macaca fascicularis: Malaysia
M. nemestrina
- Elaeophora poeli*, *illus.*
Prasad, M. C.; and Bhalla, N. P., 1977, *Indian Vet. J.*, v. 54 (2), 97-101
Elaeophora poeli, *Bubalus bubalis*, pathology of parasitic aortitis
- Elaeophora schneideri* Wehr and Dikmans, 1935
Clark, G. G.; and Hibler, C. P., 1973, *J. Wildlife Dis.*, v. 9 (1), 21-25
Hybomitra laticornis considered to be most important horse fly vector
H. phaenops
H. tetrica rubrilata
Tabanus abditus
T. eurycerus
T. gilanus
T. punctifer
all from Gila National Forest, New Mexico
- Elaeophora schneideri*
Davies, R. B.; and Clark, G. G., 1974, *J. Wildlife Dis.*, v. 10 (1), 63-65
Hybomitra laticornis: Gila National Forest, New Mexico

- Elaeophora schneideri* Wehr and Dikmans, 1935
Hibler, C. P., 1974, *J. Wildlife Dis.*, v. 10 (1), 44-46
Elaeophora schneideri, successful experimental infections in neonatal and yearling *Odocoileus hemionus*, no evidence of disease indicates mule deer are normal definitive hosts
- Elaeophora schneideri* Wehr & Dikmans, 1935, *illus.*
Hibler, C. P.; and Metzger, C. J., 1974, *J. Wildlife Dis.*, v. 10 (4), 361-369
Elaeophora schneideri, morphology and development of larval stages in horse flies and *Odocoileus hemionus*, pathogenesis in abnormal definitive hosts (*Cervus canadensis*, *Ovis aries*)
- Elaeophora schneideri*
Weinmann, C. J.; et al., 1973, *J. Wildlife Dis.*, v. 9 (3), 213-220
Wehrdikmansia cervipedis, *Elaeophora schneideri*, *Setaria yehi*, prevalence of infection in various age classes of *Odocoileus hemionus columbianus*: northern California
- Elaeophora schneideri* Wehr and Dikmans, 1935
Worley, D. E., 1975, *J. Wildlife Dis.*, v. 11 (4), 486-488
Alces alces
Odocoileus hemionus
(arteries of all): all from Montana
- Elaphostrongylus* sp., *illus.*
Lankester, M. W.; Crichton, V. J.; and Timmermann, H. R., 1976, *Canad. J. Zool.*, v. 54 (5), 680-684
1st stage protostrongylid larvae in *Rangifer tarandus caribou* (feces) may be *Elaphostrongylus* sp., lack of pathogenic effects seems to rule out *Parelaphostrongylus tenuis*: northwestern Ontario and Manitoba
- Elaphostrongylus cervi*, *illus.*
Sutherland, R. J., 1976, *N. Zealand Vet. J.*, v. 24 (11), 263-266
Elaphostrongylus cervi, *Cervus elaphus*, gross and histological lesions
- Elaphostrongylus rangiferi*
Halvorsen, O.; et al., 1976, *Norwegian J. Zool.*, v. 24 (4), 461 [Abstract]
occurrence of possible intermediate gastropod hosts in summer pasture of a flock of domesticated reindeer: Finnmark, north Norway
- Elaphostrongylus rangiferi*
Hansen, S. H.; and Halvorsen, O., 1976, *Norwegian J. Zool.*, v. 24 (4), 462 [Abstract]
reaction of intermediate host to infection *Arianta arbustorum* (muscular part of foot) (exper.)
Eobania vermiculata (muscular part of foot) (exper.)
- Elaphostrongylus rangiferi*
Lorentzen, G.; and Halvorsen, O., 1976, *Norwegian J. Zool.*, v. 24 (4), 461 [Abstract]
Elaphostrongylus rangiferi free living first stage larvae, overwintering, temperature tolerance; this stage not likely to be a factor limiting success of parasite: north Norway
- Elaphostrongylus rangiferi*, probably
Rehbinder, C.; and Christensson, D., 1977, *Nörd. Vet.-Med.*, v. 29 (12), 556-557
reindeer (faeces): Sweden
- Elaphostrongylus rangiferi*
Wissler, K.; and Halvorsen, O., 1976, *Norwegian J. Zool.*, v. 24 (4), 462-463 [Abstract]
Elaphostrongylus rangiferi, reindeer (feces), seasonal distribution, age of host: north Scandinavia
- Enoplia* Pearse, 1942
Maggenti, A. R., 1976, *Organ. Nematodes* (Croll), 1-10
Adenophorea
includes: *Enoplida*; *Dorylaimida*; *Mononchida*; *Isolaimida*; *Trichosyringida*; *Trichocephalida*
- Enoplida*
Chabaud, A. G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (1), 6-17
Adenophorea
key to superfamilies
includes: *Diectophymatoidea*; *Trichuroidea*; *Muspiceoidea*
- Enoplida* Schuurmans Stekhoven & DeConinck, 1933
Maggenti, A. R., 1976, *Organ. Nematodes* (Croll), 1-10
Enoplia
includes: *Enoplina*; *Oncholaimina*
- Enoplina* Chitwood & Chitwood, 1937
Maggenti, A. R., 1976, *Organ. Nematodes* (Croll), 1-10
Enoplia
- Enterobiasis*
Biagi, F.; Alvarez, R.; and Delgado, R., 1972, *Prensa Med. Mexicana*, v. 37 (5-6), 221-223
human intestinal helminths, clinical trials with pyrantel pamoate, good results
- Enterobiasis*
Carney, D. E.; O'Reilly, B. J.; and Tweddell, E. D., 1971, *Med. J. Australia*, v. 2 (5), 254-256
human enterobiasis, comparative clinical trials using pyrantel embonate and vipryinium embonate: Australia
- Enterobiasis*
Grudzien, M.; and Krauze, M., 1970, *Polski Tygod. Lekar.*, v. 25 (3), 1133-1134
enterobiasis, survey of pregnant women for evidence of infection, recommendations for prophylactic measures and routine examinations in pregnancy: Poland
- Enterobiasis*
Joshi, V. S.; et al., 1975, *Pediat. Clin. India*, v. 10 (3), 171-173
children with ascariasis or enterobiasis, successful therapy with vanpar: Pune, India

- Enterobiasis**
Palicka, P., 1971, *Ceskoslov. Pediat.*, v. 26 (11), 563-565
enterobiasis, long-term follow-up of children treated with a single dose of pyrvinium embonate and recommendations for therapy: Czechoslovakia
- Enterobiasis**
Quirolgico, E. O., 1973, *Philippine J. Pediat.*, v. 22 (2), 70-73
enterobiasis in children, 1-year comparative study, perianal inspection most practical and accurate method of diagnosis for use in private practice
- Enterobiasis**
Radermecker, M.; et al., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months
- Enterobius**
Biagi, F.; and Medina, J. A., 1974, *Prensa Med. Mexicana*, v. 39 (9-10), 440-443
human intestinal parasites, pyrantel pamoate, metabolism and pharmaceutical action
- Enterobius**
Biagi, F.; Smyth, J.; and Gonzalez, C., 1974, *Prensa Med. Mexicana*, v. 39 (1-2), 51-53
human intestinal helminths, successful clinical trials using mebendazole, drug well tolerated with minimal side effects: Mexico
- Enterobius bipapillatus**
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 214
Rhesusaffe
- Enterobius sciuri**
Davidson, W. R., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (cecum, large intestine); southeastern United States
- Enterobius vermicularis**
Alcasid, M. L. S.; et al., 1973, *N. York State J. Med.*, v. 73 (13), 1786-1788
human intestinal parasites, no significant correlation with bronchial asthma, results of study with control and asthmatic groups: New York
- Enterobius vermicularis**
al-Allaf, G. A.; and Hayatee, Z. G., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (4), 351
Enterobius vermicularis, recto-urethral migration of gravid female worms reported in 2 adult males (urethra) who admitted to being habitual sodomists, each had had recent contact with an underage (under 14) partner: Iraq
- Enterobius vermicularis, illus.**
Alvarez San Cristobal, A.; and de Mundi, A., 1976, *Acta Cytol.*, v. 20 (3), 190-192
Enterobius vermicularis larvae detected in vaginal smear of young girl using Papanicolaou and acridine orange staining methods
- Enterobius vermicularis**
Antunes, M. L.; and Delascio, D., 1967, *Maternidade e Infancia*, v. 26 (1), 63-67
laboratory diagnosis of vulvo-vaginitis of parasitic origin in children: Brasil
- Enterobius vermicularis**
Arnaud, J. P.; and Danis, M., 1976, *Medecine Infant.*, v. 83 (1), 9-22
helminthiasis of digestive tract in children, differential diagnosis, current treatment methods
- Enterobius vermicularis**
Asmera, J.; et al., 1971, *Ceskoslov. Pediat.*, v. 26 (9), 456-458
Enterobius vermicularis in young children, comparative clinical trials using pyrvinium suspension (Spofa) and vanquin (Parke-Davis), no significant difference between drugs: Czechoslovakia
- Enterobius vermicularis**
Asmera, J.; et al., 1975, *Ceskoslov. Pediat.*, v. 30 (4), 171-172
human intestinal helminths, mebendazole used as successful wide-spectrum anthelmintic in clinical trials: Czechoslovakia
- Enterobius vermicularis**
Bailenger, J.; et al., 1977, *Pharmacie Biol.* (109), v. 11, 267-277
human intestinal parasites, fecal examination using floatation or diphasic concentration, principles involved in both diagnostic procedures, changes in results when mercury or sodium merthiolate is added during procedure, special application of Janeckso and Urbanyi reaction
- Enterobius vermicularis**
Biagi, F.; Lopez, R.; and Viso, J., 1975, *Progr. Drug Research*, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review
- Enterobius vermicularis**
Biagi, F.; Smyth, J.; and Gonzalez, C., 1975, *Prensa Med. Mexicana*, v. 40 (5-6), 189-192
human intestinal parasites, clinical trials with mebeticlol show it to be useful drug against many parasites and therefore recommended for mass therapy in low socioeconomic areas where multiple parasitism is likely to be present: Mexico

- Enterobius vermicularis*
Botero, D.; Orozco, H. I.; and Perez, A., 1966, Bol. Chileno Parasitol., v. 21 (4), 111-114
Ascaris lumbricoides and Enterobius vermicularis in children, evaluation of combination syrup containing piperazine and pyrvinium in treatment of infections
- Enterobius vermicularis*
Brus, R.; et al., 1976, Polski Tygod. Lekar., v. 31 (5), 195-197
Enterobius vermicularis in children, comparative mass clinical trials with vanquin and combantrin; combantrin indicated as drug of choice since it was slightly more effective than vanquin and also effective against Ascaris lumbricoides: Cieszyn County, Poland
- Enterobius vermicularis*
de Carneri, I.; de Dominici, M.; and Carozzi, A., 1973, Riv. Parassitol., Roma, v. 34 (3), 213-218
Enterobius vermicularis, school children, incidence, no correlation with age or sex or infection with trichuriasis, some correlation with socio-economic status: Pavia Province, Italy
- Enterobius vermicularis, illus.*
Cavallo, G., 1974, Pediatria, Napoli, v. 82 (4), 628-653
Enterobius vermicularis, extensive morphologic and clinical review of infection in humans
- Enterobius vermicularis*
Cerva, L., 1976, Ceskoslov. Epidemiol., Mikrobiol., Immunol., v. 25 (6), 365-367
helminths, diagnostic staining method for eggs and larvae in fresh feces or those fixed by formalin
- Enterobius vermicularis, illus.*
Cerva, L., 1976, Immun. u. Infekt., v. 4 (6), 279-282
intestinal helminths, diagnostic method for staining of eggs and larvae in smears of fresh and fixed stool samples
- Enterobius vermicularis*
Cervova, H., 1973, Ceskoslov. Pediat., v. 28 (1), 43
Enterobius vermicularis in kindergarten-age children successfully treated with pyrvinium pamoate suspension: Czechoslovakia
- Enterobius vermicularis, illus.*
Chandrasoma, P. T.; and Mendis, K. N., 1977, Am. J. Trop. Med. and Hyg., v. 26 (4), 644-649
Enterobius vermicularis in humans, case reports of parasite migration to ectopic sites with resulting granulomata and accompanying pathology, sex of parasite, relationship to area of migration
- Enterobius vermicularis*
Copeman, R.; Pashen, D.; and Burger, G., 1975, Med. J. Australia, v. 1, suppl. 2, 8-13
association of common intestinal parasites to growth, nutrition and living situation of Aboriginal children: Cunnamulla, Western Queensland
- Enterobius (Oxyuris) vermicularis*
Cuevas, R.; et al., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 121-123
boarding school children, survey indicates 87.8% infection rate after multiple Scotch tape tests: Santiago, Chile
- Enterobius vermicularis*
Desser, S. S.; and Yang, Y. J., 1976, Canad. Med. Ass. J., v. 114 (4), 290-291
Dientamoeba fragilis infection in child resulting in severe intestinal symptoms, diagnosis only after multiple fecal examinations, diphetarsone treatment; possible role of Enterobius vermicularis as vector of Dientamoeba fragilis in human infections
- Enterobius vermicularis, illus.*
Di Guardo, G.; and Pampiglione, S., 1972, Parassitologia, v. 14 (1), 115-119
Enterobius vermicularis, Taenia sp., prevalence in appendices surgically excised: Luino
- Enterobius vermicularis*
Dufek, M.; and Rakosnikova, M., 1973, Ceskoslov. Pediat., v. 28 (5), 264-265
Enterobius vermicularis in children, successful clinical trials with pyrvinium suspension, 91.5% cure rate: Praha
- Enterobius vermicularis*
Gemesi, G., 1971, Parasitol. Hungar., v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Enterobius vermicularis*
Ghee, C. G., 1970, Med. J. Malaya, v. 24 (3), 218-220
evaluation of tetramisole as effective antihelminthic against Enterobius vermicularis in children
- Enterobius vermicularis*
Hays, B. D., 1977, J. Environ. Health, v. 39 (6), 424-426
transmission of protozoan cysts and metazoan eggs from land application of sewage effluents and sludges, brief literature review, parasite survey from selected Pittsburgh area sludges, control measures
- Enterobius vermicularis*
Iwanczuk, I., 1969, Acta Parasitol. Polon., v. 17 (1-19), 139-145
human parasite incidence in water and surfaces of swimming pools; change of incidence in children using swimming pool for 6 week period: Poland
- Enterobius vermicularis*
Iwanowski, H.; and Herman, T., 1977, Pediat. Polska, v. 52 (6), 649-654
Ascaris lumbricoides in children, Soprunova urine test used successfully to diagnose ascariasis and differentiate from infections of Trichuris trichiura, Enterobius vermicularis and Lamblia intestinalis

- Enterobius vermicularis**
Jaskiewicz, W.; and Soroczan, W., 1971, *Pediat. Polska*, v. 46 (1), 57-61
Enterobius vermicularis infection in children (80%) and personnel (50%) of institution for children with impaired vision, samples of dust and air and bed linens also showed contamination, clinical report: Lublin, Poland
- Enterobius vermicularis**
Jose, D. G.; and Welch, J. S., 1970, *Med. J. Australia*, v. 1 (8), 349-356
possible role of intestinal parasitism in growth-retarded, anemic and malnourished Australian Aboriginal children, comparison with normal Aboriginal children: Queensland
- Enterobius vermicularis**
Kochanska, E.; Kochanski, M.; and Stehlik, A., 1972, *PolSKI Tygod. Lekar.*, v. 27 (37), 1453-1455
Enterobius vermicularis, children in group home, clinical trials using combantrin, good results: Poland
- Enterobius vermicularis**
Krasa, E.; and Kaupeny, M., 1977, *Paediat. u. Paedol.*, v. 12 (1), 73-75
children with Enterobius vermicularis and associated asthma or broncho-pulmonary conditions, mebendazole therapy highly effective
- Enterobius vermicularis**
Kvasz, L., 1972, *Ceskoslov. Pediat.*, v. 27 (11), 555-557
Enterobius vermicularis in children, Graham-Brumpt method superior to other method used for diagnostic surveys: Bratislava
- Enterobius vermicularis**
Lessig, P. M., 1973, *Maternal and Child Health*, v. 5 (11), 12-13
Enterobius vermicularis, diagnosis, control measures, eradication, humans
- Enterobius vermicularis**
Most, H., 1972, *N. England J. Med.*, v. 287 (10), 495-498; (14), 698-702
common parasitic infections of man encountered in the United States, recommendations for treatment, review
- Enterobius vermicularis, illus.**
Ockert, G.; and Schmidt, T., 1976, *J. Hyg., Epidemiol., Microbiol. and Immunol.*, v. 20 (1), 76-81
Enterobius vermicularis, confirmation as vector of *Dientamoeba fragilis*
- Enterobius vermicularis**
Otsuru, M., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Enterobius vermicularis**
Pawlowski, Z.; Chwirot, E.; and Zizlowski, J., 1973, *PolSKI Tygod. Lekar.*, v. 28 (8), 291-293
Enterobius vermicularis, adults and children, efficacy of combantrin compared with that of pyrinium embonate
- Enterobius vermicularis**
Pazdziora, E., 1976, *Ceskoslov. Epidemiol., Mikrobiol., Immunol.*, v. 25 (1), 50-55
Enterobius vermicularis, children's convalescent home, mass treatment with pyrinium pamoate, 17% not cured could serve as permanent source of disease and must be treated with particular attention if enterobiasis is to be eliminated
- Enterobius vermicularis**
Purtilo, D. T.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (2), 229-232
parasitized children with protein-calorie malnutrition, complete blood count, serum immunoglobulin concentration, significant relationship between intensity of parasitism and Ig levels: Brazil
- Enterobius vermicularis**
Rajasekaran, P.; Dutt, P. R.; and Pisharoti, K. A., 1977, *Indian J. Med. Research*, v. 66 (2), 189-199
human intestinal parasites, survey of correlation between infection rate and source of water supply (well, street tap, home with tap water) as indication of control of water-borne diseases by public water supplies: Madurai district, Tamil Nadu, India
- Enterobius vermicularis**
Rees, P. H.; and Marsden, P. D., 1970, *Brit. J. Clin. Pract.*, v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment
- Enterobius vermicularis**
Ruitenbergh, E. J.; et al., 1976, *Nederl. Tijdschr. Geneesk.*, v. 120 (15), 645-649
Toxocara canis, survey of 253 children for complement-fixing antibodies against *Toxocara* shows low incidence; eosinophilia attributed to presence of *Enterobius vermicularis*: Netherlands
- Enterobius vermicularis, illus.**
Saffos, R.; and Rhatigan, R. M., 1977, *Am. J. Clin. Path.*, v. 67 (3), 296-299
Enterobius vermicularis in woman resulting in perforated unilateral salpingitis, surgical excision of granular mass containing ova, case report: Florida
- Enterobius vermicularis**
Sagua, H.; and Poblete, H., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 6-7
Enterobius vermicularis in boys in boarding schools, diagnostic comparisons of Graham's scotch tape method and a modified Graham's technique: Antofagasta, Chile

- Enterobius vermicularis*
Schenone, H.; et al., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 113-117
Enterobius vermicularis, human, comparison of anal swabs and direct fecal examination in diagnosis, value of serial examinations
- Enterobius vermicularis*
Schenone, H.; et al., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 118-122
single combined dose of pyrinium and tetramisole effective in treating children with *Enterobius vermicularis* and mixed infections with *Ascaris lumbricoides*: Chile
- Enterobius vermicularis*
Schenone, H.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 86-89
pyrantel pamoate recommended for treatment of both *Ascaris lumbricoides* and *Enterobius vermicularis*, individual or mass therapy, clinical trials in children: Chile
- Enterobius vermicularis*
Schenone, H.; et al., 1976, Bol. Chileno Parasitol., v. 31 (3-4), 75-78
Enterobius vermicularis, *Ascaris lumbricoides*, and *Trichuris trichiura* in children, clinical trials of oral mebendazole highly successful, recommended for individual or mass therapy
- Enterobius vermicularis*
Schenone, H.; Galdames, M.; and Cabello, C., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 89-90
intestinal parasites, young girls, combined therapy with mebendazole and thiabendazole
- Enterobius vermicularis*
Seah, S. K. K., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 534-542
intestinal parasites, persons living in non-endemic areas who acquired infections while travelling or who have immigrated from endemic areas, pyrantel pamoate successful for *Ascaris lumbricoides*, results with other parasites varied: Montreal, Canada
- Enterobius vermicularis*
Sehgal, S. C.; Vinayak, V. K.; and Gupta, U., 1977, Indian J. Med. Research, v. 65 (4), 509-512
human helminthic ova in feces, diagnosis using the Kato thick smear technique more successful than commonly used techniques, recommended for epidemiologic surveys: Chandigarh, India
- Enterobius vermicularis*
Shipton, E. A.; McInerney, R. J. F.; and Hulbert, L., 1973, Med. J. Australia, v. 1 (20), 1014
Enterobius vermicularis ova discovered in vaginal smear of adult woman, routine wet films from vaginal discharge recommended for diagnosis: Australia
- Enterobius vermicularis*
Tanowitz, H. B., 1974, Med. Aspects Human Sexual., v. 8 (9), 45-65
human parasitic gynecologic diseases, clinical aspects, epidemiology, sexual transmission, review
- Enterobius vermicularis*
Vinayak, V. K.; and Sehgal, S. C., 1976, Indian J. Med. Research, v. 64 (9), 1347-1350
human helminthic and protozoan parasites, comparison of nigrosin-methylene blue diagnostic test with formol-ether method and direct examination
- Enterobius vermicularis*
Virgala, J.; Spaldonova, R.; and Vzentkova, O., 1973, Ceskoslov. Pediat., v. 28 (6), 325-326
Enterobius vermicularis in humans, clinical trials with mintage, good results with some side effects: Czechoslovakia
- Enterobius vermicularis*
Warren, K. S.; and Mahmoud, A. A. F., 1975, J. Infect. Dis., v. 132 (2), 229-232
Enterobius vermicularis, life cycle, epidemiology, disease syndromes, diagnosis, management (sanitary measures of no avail, mebendazole effective)
- Enterobius vermicularis*
Yang, J.; and Scholten, T., 1977, Am. J. Trop. Med. and Hyg., v. 26 (1), 16-22
Dientamoeba fragilis in humans, extensive epidemiologic survey, pathology, fecal examination diagnostic methods, periodicity, frequent occurrence in presence of *Enterobius vermicularis* suggests possibility of helminths as vectors: Toronto, Canada
- Epomidiostomatinae Skrjabin et Schulz, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae
includes: *Epomidiostomum* (type genus); *Pseudamidostomum*
- Epomidiostomum* Skrjabin, 1915 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Epomidiostomatinae
- Epomidiostomum anatinum* Skrjabin, 1915
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmin. Lab., v. 15, 109-133
Anser anser
A. erythropus
A. albifrons
Casarca ferruginea
Anas platyrhynchos
A. penelope
A. clypeata
A. acuta
A. crecca
A. querquedula
Aythya nyroca
Netta rufina
(under cuticle of muscular stomach of all): all from Bulgaria
- Epomidiostomum uncinatum* (Lundahl, 1848)
Turner, B. C.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 157-169
parasites of *Anas crecca* and *A. discors*, incidence and intensity, age and sex of host
Anas crecca
A. discors
(beneath the gizzard lining of all): all from eastern Canada

- Esslingeria* subgen. n.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of *Tetrapetalonema*; key
tod of subgen.: *Tetrapetalonema* (*Esslingeria*)
perstans (Manson, 1891)
- Eucoleus annulatus* (Molin, 1858) Lopez-Neyra, 1946
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelminth. Lab., v. 15, 109-133
Anas platyrhynchos (esophagus): Bulgaria
- Eucoleus corvicola* Vasilkova, 1930
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
as syn. of *Thominx contorta* (Creplin, 1839)
- Eucoleus laricola* Vasilkova, 1930
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
possible syn. of *Thominx contorta* (Creplin, 1839)
- Eucoleus laricola* Wassilkowa, 1930, illus.
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus canus
Larus ridibundus
L. crassirostris
Sterna hirundo
(esophagus of all): all from coast of Sea of Okhotsk
- Eucoleus laricola* Wassilkowa, 1930
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus: Azov Sea
L. ridibundus: Azov Sea; Tuva
L. genei: Azov Sea
L. ichthyaetus: Tuva
Sterna hirundo: Tuva
- Eucoleus pachyderma*
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
as syn. of *Thominx contorta* (Creplin, 1839)
- Eucoleus trilobus* (Linstow, 1875) Lopez-Neyra, 1946
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Philomachus pugnax
Eudromias morinellus
Charadrius hiaticula
Calidris temminckii
Heteroscelus incanus brevipes
Gallinago media
all from lower Yenisei [and/or] Keta lake
- Eucyathostomum* Molin (1861)
Pursglove, S. R., jr., 1976, J. Parasitol., v. 62 (4), 574-578
taxonomy
- Eucyathostomum webbi* sp. n., illus.
Pursglove, S. R., jr., 1976, J. Parasitol., v. 62 (4), 574-578
Odocoileus virginianus (large intestine):
Blackbeard Island, McIntosh County, Georgia;
Pushmataha County, Oklahoma; Florida; South Carolina
- Eucyathostomum webbi* Pursglove, 1976
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (large intestine):
Oklahoma
- Eucyathostomum webbi*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Eufilaria lari* Yamaguti, 1935
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus argentatus
L. canus
(connective tissue between trachea and esophagus of all): all from coast of Sea of Okhotsk
- Eugenuris* Schulz 1948
Grundmann, A. W.; and Lombardi, P. S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 39-46
diagnosis
- Eugenuris utahensis* n. sp., illus.
Grundmann, A. W.; and Lombardi, P. S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 39-46
Ochotona princeps cinnemomea: Mt. Delano, Tushar Mountains, Utah
O. p. uinta: Uinta Mountains, Utah
O. p. lasalensis: La Sal Mountains, Utah
O. p. barnsei: Fish Lake Mountains, Utah
O. p. nevadensis: Ruby Mountains, Nevada
O. p. fuscipes: Markagunt Plateau, Utah
(cecum of all)
- Eurymermis* in part: Rubtsov 1972
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
as syn. of *Romanomermis* Coman, 1961
- Eurymermis boschkoi* sp. n., illus.
Rubtsov, I. A., 1976, Zool. Zhurnal, v. 55 (9), 1292-1298
Chrysops relictus: Kiev obl.
- Eustoma* van Beneden, 1871
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis* Layman & Borovkova, 1926
- Eustoma* van Beneden, 1870, nec Piette, 1855
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Pseudanisakis* (Layman & Borovkova, 1926, subgen.) *Mozgovoi*, 1951
- Eustoma rajae* (Yamaguti, 1941) of Hartwich (1957)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis rajae* Yamaguti, 1941, *sensu nov.*

- Eustoma rotundata* (Rud.) of Polyanski (1955), of Campana-Rouget (1955), of Kreis (1958, adults only), of Myers (1959), of Berland (1961), and of Threlfall (1969)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as possible syn. of *Pseudanisakis tricupola* nom. nov.
- Eustoma rotundatum* (Rud.) of Punt (1941)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis baylisi* nom. nov.
- Eustoma truncata* van Beneden of Hartwich (1957) (in part)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis tricupola* nom. nov.
- Eustoma truncata* van Beneden, 1871, of Hartwich (1957) (in part)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of *Pseudanisakis baylisi* nom. nov.
- Eustrongylides Jagerskiold*, 1909
Fastzkie, J. S.; and Crites, J. L., 1977, J. Parasitol., v. 63 (4), 707-712
key to species, including: *E. perpapillatus* Jagerskiold 1909; *E. phalacrocoracis* Johnston and Mawson 1941; *E. plotinus* Johnston and Mawson 1941; *E. excisus* Jagerskiold 1909; *E. formosensis* Sugimoto 1931; *E. mergorum* Rudolphi 1809; *E. tubifex* Jagerskiold 1909; *E. ignotus* Jagerskiold 1909; *E. tricolor* Sugimoto 1931; *E. africanus* Jagerskiold 1909
- Eustrongylides* sp.
Abram, J. B.; and Lichtenfels, J. R., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 253
Lutra canadensis (intestine): Anne Arundel County, Maryland
- Eustrongylides* sp., tentatively identified as *E. tubifex* (Nitzsch, 1819)
Buck, O. D.; Cooper, C. L.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 233-234
Larus argentatus: Bass Island region of Lake Erie
- Eustrongylides* spp. Jagerskiold, 1909
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Salmo trutta: Loch Leven, Scotland
- Eustrongylides* sp.
Fastzkie, J. S.; and Crites, J. L., 1977, J. Parasitol., v. 63 (4), 707-712
Mergus serrator: Lake Erie
- Eustrongylides* sp.
Kennedy, C. R.; and Lie, S. F., 1976, J. Fish Biol., v. 8 (4), 293-302
Eustrongylides sp., distribution, pathogenicity of larvae in fish host, preference for female fish and larger fish, little accumulation in older fish, no seasonal changes in infection levels nor pathological effects
Salmo trutta (stomach, body wall, intestine, liver, ovary, swim bladder): Fernworthy Reservoir, Devon Galloway; Orkney
Gymnocephalus cernua: Serpentine
Perca fluviatilis: Serpentine
- Eustrongylides* sp.
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Bot. Africaines, v. 87 (2), 209-248
Protopterus aethiopicus (intestinal wall): Lake Victoria, Uganda
Engraulicypris argenteus (body cavity): Lake Victoria, Uganda
Bagrus docmac (body cavity): Lake Victoria, Uganda
Haplochromis sp. (body cavity): Lake Victoria, Uganda
H. angustifrons (body cavity): Lake George, Uganda
- Eustrongylides* sp. (? *excisus*) (Jagerskiold 1909)
Lee, R. L. G., 1977, Lond. Naturalist (1976) (56), 57-70
Gymnocephalus cernua
Perca fluviatilis
(wall of gut within the body cavity of all): all from Serpentine lake, Hyde Park and Kensington Gardens, central London
- Eustrongylides* sp.
Lichtenfels, J. R.; and Lavies, B., 1976, Lab. Animal Sc., v. 26 (3), 465-467
larval *Eustrongylides* sp., case reports, cause of death, surgical treatment
Thamnophis sirtalis parietalis (beneath skin, lungs, body cavity, along spinal column): 100 miles north of Winnipeg, Manitoba (transported to Annapolis, Maryland)
- Eustrongylides* sp.
Lockard, L. L.; Parsons, R. R.; and Schaplow, B. M., 1975, Great Basin Nat., v. 35 (4), 442-448
Salmo trutta (upper digestive tract), relationship of incidence and intensity of nematode infection to age and sexual maturity of host, higher infection rate in sexually mature trout due to aggressive feeding behavior: streams in southern and western Montana
- Eustrongylides* sp.
Miller, R. L.; Olson, A. C., jr.; and Miller, L. W., 1973, Calif. Fish and Game, v. 59 (3), 196-206
Lepomis macrochirus (muscles): southern California reservoirs
- Eustrongylides ignotus, illus.*
Winterfield, R. W.; and Kazacos, K. R., 1977, Avian Dis., v. 21 (3), 448-451
case report, mortality, morbidity, cause of death in young birds
Ardea herodias (proventriculus, gizzard): Madison County, Indiana

- Eustrongylides mergorum* (Rudolphi, 1809), illus. Bogoiavlenskii, Iu. K.; and Khatkevich, L. M., 1970, *Parazitologiya*, Leningrad, v. 4 (3), 223-230
4 spp. of Diactophymata, fine structure of somatic musculature, distribution of DNA and RNA
- Eustrongylides papillosus* Rudolphi, 1802
Kamburov, P.; and Vasilev, I., 1972, *Izvest. Tsentral. Khelmit. Lab.*, v. 15, 109-133
Mergus serrator (esophagus): Bulgaria
- Eustrongylides tubifex* Jagerskiold, 1909
Baker, J. C.; and Crites, J. L., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 37-39
Ictalurus punctatus (mesenteries): island region of western Lake Erie
- Eustrongylides tubifex* (Nitzsch, 1819)
Cooper, C. L.; Ashmead, R. R.; and Crites, J. L., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 96
prevalence, comparison with previous years
Perca flavescens (mesenteries): western Lake Erie
- Eustrongylides tubifex* (Nitzsch 1819) Jagerskiold 1909, illus.
Fastzkie, J. S.; and Crites, J. L., 1977, *J. Parasitol.*, v. 63 (4), 707-712
redescription, key
Perca flavescens (mesentery): western basin of Lake Erie
Anas platyrhynchos (proventriculus) (exper.)
- Evandroia Travassos*, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, Pudicinae
- Excisa Gendre*, 1928, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key

- Filaria* sp. Serkowa, 1948
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 3-6
as syn. of *Sarconema pseudolabiata* nov. sp.
- Filaria* [sp.], tentatively identified as immature *Dirofilaria immitis*, illus.
Dissanaike, A. S.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, part 1), 1143-1147
filaria in vitreous of man's eye, worm not recovered but apparently died after diethyl-carbamazine therapy, probable source of infection was pet dogs, 3 of which were infected with *Dirofilaria immitis*: Peninsular Malaysia
- Filaria* [sp.]
Kanda, T.; et al., 1976, Mosquito News, v. 36 (3), 364-365
biting cycles of possible vectors of "Timor filaria"
Anopheles barbirostris: Timor Island, Indonesia
- Filaria bubali*
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
re-examination of specimens of Rudolphi 1819, referred to in legend to fig. as *Setaria bubali*
- Filaria inflexocaudata* von Siebold 1842
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Halocercus invaginatus* (Quekett 1841) Dougherty 1943
- Filaria malayi* (*Brugia malayi*)
Zuidema, P. J., 1972, Nederl. Tijdschr. Geneesk., v. 116 (1), 21-23
Filaria malayi, historical review of first research on human filariasis
- Filaria medinensis*, illus.
Fedchenko, A. P., 1971, Am. J. Trop. Med. and Hyg., v. 20 (4), 511-523
English translation of Russian article published in 1870 on the structure and reproduction of the guinea worm (*Filaria medinensis*)
- Filaria neglecta* Diesing, 1851
Rozman, M., 1971, Acta Parasitol. Iugoslavica, v. 2 (2), 67-77
as syn. of *Icosiella neglecta* (Diesing, 1851)
- Filaria rubella* Rudolphi, 1819 sensu Nablas et Sabrazes, 1892; sensu Blanchard, 1895; sensu Porta, 1912; sensu SKRjabin, 1923, illus.
Rozman, M., 1971, Acta Parasitol. Iugoslavica, v. 2 (2), 67-77
as syn. of *Icosiella neglecta* (Diesing, 1851)
- Filariiae
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Filarial worms
Sivanandam, S.; and Sandosham, A. A., 1965, Med. J. Malaya, v. 20 (1), 65
multiple filarial infections in domestic cat: East Pahang, Malaya
- Filariasis, bancroftian. See [*Wuchereria bancrofti*]
- Filariasis
Aalberse, R. C.; Brummelhuis, H. G. J.; and Reerink-Brongers, E. E., 1973, Immunochemistry, v. 10 (5), 295-303
plasma from patients with *Schistosoma mansoni* and tropical eosinophilia probably due to a microfilaria infection, purification of polyclonal IgE by immunosorption
- Filariasis
Ambroise-Thomas, P.; and Kien Truong, T., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema viteae*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance
- Filariasis
Arrighi, E.; and Artignan, P., 1972, Medecine Trop., v. 32 (3), 305-310
surgical procedure for the treatment of human scrotal elephantiasis caused by filariasis
- Filariasis
Ayachit, S. D., 1972, Oriental Arch. Ophth., v. 10 (2), 92-96
adult filarial worm removed from eye of man being treated for tropical eosinophilia, clinical case report: Madhya Pradesh, India
- Filariasis
Basio, R. G., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 435-436
human filariasis, possible biological control of *Mansonia uniformis* mosquito vectors as larvae are ideal food source for mudfish (*Ophicephalus striatus*) in Philippines
- Filariasis
Bloch-Michel, H.; and Waltzing, P., 1973, Nouv. Presse Med., v. 2 (6), 382 [Letter]
human filariasis, application of rosette phenomenon for diagnosis
- Filariasis
Charters, A. D.; et al., 1972, Med. J. Australia, v. 2 (21), 1195-1198
tropical pulmonary eosinophilia in 2 youths who recently migrated into Western Australia, case reports, clinical management, need for awareness of possible disease transmission from endemic areas
- Filariasis
Chernin, E., 1977, Am. J. Trop. Med. and Hyg., v. 26 (5, part 2), 1065-1070
human filariasis, historical review of Patrick Manson's investigations on transmission

Filariasis

Dennis, D. T.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (6), 797-802

Timor filariasis, human, epidemiology, clinical manifestations, age and sex stratification of microfilaremia and symptoms: Ae Bubu, southeastern Flores, Southeast Indonesia

Filariasis

Diesfeld, H. J.; and Kirsten, C., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 435-438

filariasis, human, diagnosis, new embedding technique employing 'methacrylate' for preparation of antigen (*Dipetalonema viteae*) to be used in indirect fluorescent antibody test (tested on onchocerciasis sera from Togo), compared with usual frozen-section method

Filariasis

Dumas, M.; Girard, P. L.; and Gentilini, M., 1976, *Medecine Afrique Noire*, v. 23 (2), 89-93

human filariasis, invasion of central nervous system by microfilaria, diagnosis by immunofluorescence and finding of antibodies in spinal fluid

Filariasis

Elslager, E. F., 1974, *Progr. Drug Research*, v. 18, 99-172

human malaria and filariasis, new perspectives on chemotherapy, extensive review

Filariasis

Ezigo, J. C.; et al., 1976, *Parasitology*, v. 73 (2), xxxii [Abstract]

filariasis in normal and protein-deficient cotton rats, serum enzyme levels

Filariasis

Hedge, E. C.; and Ridley, D. S., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (4), 304-307

evaluation of microfilarial antigen for use with indirect immunofluorescent test in diagnosing human filariasis; best results obtained with sonicated microfilariae of *Brugia pahangi* with which both cytoplasmic and sheath antigens could be utilized simultaneously

Filariasis

Ismail, M. M.; and Nagaratnam, N., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (3), 405-409

filariasis as possible cause of arthritis, clinical features and laboratory findings in 33 cases, age distribution, diethylcarbamazine treatment gave good results: Sri Lanka

Filariasis

Katiyar, J. C.; et al., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (2), 169-170 [Letter]

human filariasis, persons undergoing diethylcarbamazine therapy for *Wuchereria bancrofti*, interference with skin test reactions when *W. bancrofti* used as antigen, review of possible mechanisms

Filariasis

Kerdipibule, V.; et al., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (1), 150-151 [Demonstration]

possible biological control of *Mansonia uniformis* vectors of human filariasis through nematode parasitism with *Reesimermis nielsenii*, preliminary laboratory trials

Filariasis

Kershaw, W. E.; et al., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (4), 438 [Demonstration]

filariasis, cotton rats, first infection retards development of worms of second infection, second infection may cause abrupt fall in circulating microfilariae

Filariasis

Kim, C. W., 1975, *Progr. Clin. Path.*, v. 6, 267-288

extensive review of techniques used to diagnose human parasitic diseases

Filariasis

Kurokawa, K., 1976, *Nippon Zyuisi-Kai Zasshi (J. Japan Vet. Med. Ass.)*, v. 29 (4), 190-197

canine filariasis, surgical treatment

Filariasis

Laurens, A.; et al., 1973, *Medecine et Armees*, v. 1 (4), 61-64

woman, hemato-chyluria of filarial origin, diagnosis by lymphography, case report, diethylcarbamazine

Filariasis

Macdonald, W. W., 1976, *Symposia Brit. Soc. Parasitol.*, v. 14, 1-24

Aedes scutellaris complex, *Culex pipiens*, genetics, relevance to possible control of filariasis vector populations, symposium presentation

Filariasis

McMillan, B., 1968, *Med. J. Australia*, v. 2 (2), 63-64

use of masticated lawang bark for relief of pain of human filarial lymphangitis and associated ulcerations: northern New Guinea

Filariasis

Mazaud, R.; and Ferrus, R., 1973, *Medecine Trop.*, v. 33 (2), 177-185

human malaria and filariasis as possible causes of idiopathic endomyocardial fibrosis in tropical areas

Filariasis

Murakami, F.; et al., 1972, *Nettai Igaku (Trop. Med.)*, v. 14 (3), 138-143

successful treatment of human filarial chyluria with medium chain triglyceride and low fat diet

Filariasis

Nagaratnam, N.; and Ismail, M. M., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (6), 803-807

comparative study of 101 cases of rheumatic fever and arthritis believed to be of filarial origin, borderline cases where aetiology in doubt

Filariasis

Nelson, G. S., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 273-285
schistosomiasis, filariasis, zoonophylaxis ("the prevention or amelioration of disease in man as the result of previous exposure to heterologous infections of animal origin")

Filariasis

Niel, G.; et al., 1972, Medecine et Malad. Infect., v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalone-ma vitae* and *Ascaris suum* as antigens

Filariasis

Oster, Z., 1976, J. Nuclear Med., v. 17 (5), 425-426
elderly man with supraclavicular mass visible on bone scan, biopsy of mass revealed nematode cysts, probably filariasis of *Dirofilaria immitis* origin

Filariasis

Ottesen, E. A.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, Pt. 2), 134-141
eosinophils from normal humans and from patients with schistosomiasis or filariasis, immunoglobulin and complement receptors, role in cellular adherence to *Schistosoma mansoni* schistosomules, workshop report

Filariasis

Pedinielli, L.; et al., 1970, Marseille Chir., v. 22 (5), 504-508
filariasis, man, scrotal elephantiasis and chyluria, successful surgical intervention, case report, clinical review: France

Filariasis

Pieron, R.; et al., 1974, Poumon et Coeur, v. 30 (4), 313-318
lymphatic filariasis with associated intrathoracic cystic lymphangioma, clinical case report: Paris (native of Africa)

Filariasis

Pinon, J. M.; and Dropsy, G., 1977, J. Immunol. Methods, v. 16 (1), 15-22
various human parasitic diseases, application of enzyme-linked-immuno-electro-diffusion assay (combination of immunoelectrodiffusion and immunoenzyme method), sensitivity and specificity, enables class of immunoglobulins involved to be determined

Filariasis

Pinon, J. M.; and Gentilini, M., 1973, Nouv. Presse Med., v. 2 (19), 1283-1287
human filariasis, application of cellular immunologic tests (rosette formation, macrophage migration) in diagnosis and comparison with serologic tests (fluorescent antibody, passive hemagglutination, gel diffusion)

Filariasis

Radermecker, M.; et al., 1974, Internat. Arch. Allergy and Applied Immunol., v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months

Filariasis

Rajagopalan, P. K.; Yasuno, M.; and Menon, P. K. B., 1976, Indian J. Med. Research, v. 64 (5), 688-708
density and survival studies on *Culex pipiens fatigans*, vector of human Bancroftian filariasis: Delhi, India

Filariasis

Razig, S. A., 1976, Vet.-Med. Nachr. (1), 75-84
microfilariae, horses (blood), stibophen, 9 cases tested, good results

Filariasis

Ree, G. H.; et al., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 542-543
filariasis, onchocerciasis, blood plasma levels of diethylcarbamazine after oral administration to humans, results show that drug is rapidly absorbed from gastrointestinal tract and reaches peak levels at 1-2 hours after administration

Filariasis

Ricosse, J. H.; and Picq, J. J., 1973, Medecine Afrique Noire, v. 20 (11), 877-897
human filariasis, clinical indications for treatment, drugs in current use, review

Filariasis

Rodhain, F.; and Rodhain-Rebourg, F., 1975, Medecine Trop., v. 35 (4), 267-273
importance of knowledge of animal filariasis in diagnosing human infections, review

Filariasis

Rogers, R.; Davis, R.; and Denham, D. A., 1975, J. Helminth., v. 49 (1), 31-32
new technique for visualizing changes in lymphatics caused by filarial worms, injection of Hypaque solution followed by Xero-radiography, use with *Brugia pahangi* in cats

Filariasis

Sasa, M., 1974, Progr. Drug Research, v. 18, 259-268
human filariasis, historical review of anti-filariasis control campaigns using mass drug therapy, vector control, and epidemiologic surveys

Filariasis

Sasa, M., 1976, Human filariasis. A global survey of epidemiology and control, 819 pp., illus., maps
human filariasis, global survey, epidemiology and control

Filariasis

Sasa, M.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 415 [Demonstration]
 filariasis, possible use of poeciliid fishes for biological control of mosquito vectors

Filariasis

Saugrain, J., 1971, Medecine Trop., v. 31 (2), 233-236
 filariasis, *Angiostrongylus cantonensis*, amoebiasis, human parasitic diseases of medical interest and their etiology: French Polynesia

Filariasis

Sawada, T.; et al., 1975, Progr. Drug Research, v. 19, 128-135
 human filariasis, diagnosis using *Dirofilaria immitis* adult worm antigen for skin tests, purification of antigen

Filariasis

Singh, H.; Chaudhuri, P. C.; and Kumar, A., 1976, Indian Vet. J., v. 53 (6), 467-468
 intra-ocular filariasis, horses, surgical removal using paracentesis oculi technique

Filariasis

Sinha, V. P., 1977, Mosquito News, v. 37 (2), 300-301
 impact of industrialization and urbanization on the transmission of filariasis, very brief review

Filariasis

Sivaloganathan, V.; and Cheah, W., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 279-280 [Demonstration]
 human filariasis, case report of surgical repair of elephantiasis of legs: Kepala Batas, Province Wellesley, Malaysia

Filariasis

Sivanandam, S.; and Dondero, T. J., jr., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 415 [Demonstration]
 description of simplified, non-breakable Sinton pipette useful in filariasis surveys and research where measured thick blood smears are required

Filariasis

Southgate, B. A., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 172-178
 human filariasis, field trial of counting-chamber technique for determination of microfilarial rates and densities, applicable for epidemiologic surveys

Filariasis

Southgate, B. A., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 177-186
 human bancroftian filariasis, simultaneous trials using 4 known field techniques to diagnose microfilaremia in order to obtain comparative epidemiologic profiles: Fiji

Filariasis

Stuerchler, D.; and Degremont, A., 1976, Schweiz. Med. Wchnschr., v. 106 (20), 682-688
 extensive diagnostic and clinical review of filarial parasites frequently encountered by travelers to endemic tropical areas: Switzerland

Filariasis

Velimirovic, B.; and Clarke, J. L., 1975, Tropenmed. und Parasitol., v. 26 (4), 503-506
 possible use of larvivorous fishes in wells as biological control against culicine vectors of human filariasis: Maldives Republic

Filariasis

Webber, R. H., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 544-548
 filariasis, theoretical considerations in the vector control of human infections, construction of model using retrospective data

Filariasis

White, G. B., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (4), 278-301
 malaria and filariasis in humans, extensive review of *Anopheles gambiae* complex vector mosquitoes, their morphologic identification and relationship to disease spread, possible control measures: Africa

Filariasis

Yap, H. H.; and Hanapi, S., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (4), 575-580
 human Malayan filariasis, laboratory insecticide susceptibility tests against *Mansonia* larvae

Filariasis

Zaman, V.; and Fung, W. P., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 144-145 [Letter]
 eosinophilic lung (tropical eosinophilia), successful treatment with levamisole, case report

Filarids

Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
 comparison of helminth fauna of common and purple gallinules
Porphyrula martinica (body cavity): Florida

Filarinema Moennig, 1929

Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Amidostomatidae, Mackerrastrongylinae
 Syn: *Asymmetricostrongylus Nagaty*, 1932

Filarioidea

Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
 Spirurina
 key

- Filarioidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
- Filarissima Chabaud, 1974
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
key
- Filaroides gordius, probably, *illus.*
King, N. W., jr., 1976, Scient. Publication (317). Pan Am. Health Organ., 169-198
- Filaroides hirthi
Georgi, J. R., 1976, Science (4266), v. 194, 735
Filaroides hirthi, dogs (exper.), transmission through ingestion of first-stage larvae, theoretical possibility of autoinfection
- Filaroides hirthi Georgi and Anderson, 1975, *illus.*
Georgi, J. R.; et al., 1976, Cornell Vet., v. 66 (3), 309-323
Filaroides hirthi, life history in beagles, occurrence as function of age, intermediate or alternate definitive hosts not found, levamisole not effective: commercial breeding establishment, North Rose, Wayne County, New York
- Filaroides hirthi, *illus.*
Georgi, J. R.; Georgi, M. E.; and Cleveland, D. J., 1977, Parasitology, v. 75 (2), 251-257
Filaroides hirthi, dogs (nat. and exper.), diagnosis, zinc sulphate flotation more efficient than Baermann technique in concentrating larvae from feces, larvae recovered from feces proved infective and it was concluded that infection can be transmitted directly and immediately by fresh fecal contamination, mongrel dogs as well as beagles can be infected, finding of larvae in mesenteric lymph nodes long after single exposure to exogenous infection supports hypothesis of autogenous re-infection of host by proportion of larvae migrating from lungs to anus
- Filaroides hirthi
Kagei, N.; et al., 1976, Bull. Inst. Pub. Health, Tokyo, v. 25 (3), 140-144
Filaroides hirthi, Strongyloides sp., Toxocara canis, and Trichuris vulpis in imported dogs, inability to experimentally infect other dogs with Strongyloides stercoralis of human origin: Japan, imported from U.S.A.
- Filaroides martis (Werner, 1782)
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela lutreola
Mustela putorius
Mustela erminea
Gulo gulo
(lungs of all): all from Karelia
- Filaroides milksi, *illus.*
Yasuda, N.; Sakamoto, H.; and Kono, I., 1976, Bull. Fac. Agric. Kagoshima Univ. (26), 113-118
dog, Pomeranian (lungs): Kagoshima city, Japan
- Filaroides osleri
Darke, P. G. G., 1976, Vet. Rec., v. 99 (15), 293-294
Filaroides osleri-infected dogs, tracheo-bronchitis, levamisole, good results
- Filaroides osleri
Hill, B. L.; and McChesney, A. E., 1976, J. Am. Animal Hosp. Ass., v. 12 (4), 487-489
Filaroides osleri, dog, clinical findings, diagnosis, thiabendazole, case history
- Filaroides osleri, *illus.*
Jones, B. R.; et al., 1977, N. Zealand Vet. J., v. 25 (4), 103-104
Filaroides osleri, dog (trachea, lung parenchyma), clinical findings, thiabendazole, levamisole, and fenbendazole suppressed larval production but failed to kill adults, pathology: New Zealand
- Filaroides osleri, *illus.*
Luengo, M.; and Arata, N., 1970, Bol. Chileno Parasitol., v. 25 (1-2), 87-88
chronic respiratory distress in dog caused by tracheal polyps containing Filaroides osleri, dogs (perro): Chile
- Filaroides osleri
Petrick, S. W., 1977, J. South African Vet. Ass., v. 48 (2), 105-107
Spirocerca lupi, Filaroides osleri, ascarids, dogs, gastrointestinal fibroscope, useful diagnostic aid
- Filaroides osleri
Polley, L.; and Creighton, S. R., 1977, Vet. Rec., v. 100 (7), 136-137
Filaroides osleri, dogs, experimental direct transmission
- Filaroides osleri
Thornton, J. E.; Bell, R. R.; and Reardon, M. J., 1974, J. Wildlife Dis., v. 10 (3), 232-236
Canis latrans (submucosal tissue at the tracheal bifurcation): Nueces County, Texas
- Filicapitis Travassos, 1949
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Filocapsularia Deslongchamps, 1824, nom. oblit.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Anisakis Dujardin, 1845
- Filocapsularia sp., probably F. marina
White, J. R., 1976, Florida Scientist., v. 39 (1), 37-41
Feresa attenuata (stomach): Lake Worth, Florida

- Filocapsulariinae* Yamaguti, 1961, in part
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakinae* Railliet & Henry, 1912
- Filochona*, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Rhabdochona*
key
- Fissicauda* n. gen.
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Heligmosomidae, *Brevistriatinae*
tod: *F. fissicauda* n. gen. n. sp.
- Fissicauda*
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
[genus described as new in Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710]
- Fissicauda* Durette-Desset et Krishnasamy, 1976
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Fissicauda* Durette-Desset et Krishnasamy, 1977
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, *Brevistriatinae*
- Fissicauda brevispicula* (Ow Yang, 1967) n. comb.
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Syn.: *Brevistriata brevispicula* Ow Yang, 1967
- Fissicauda callosciuri* (Supperer et Kutzer, 1963), n. comb., *illus.*
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
description, synlophe
Syn.: *Brevistriata callosciuri* Supperer et Kutzer, 1963
Callosciurus notatus (intestin grele):
Malaisie
- Fissicauda callosciuri* (Supperer et Kutzer, 1963)
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Tragulus javanicus (intestin grele): Selangor, Subang, Forest Reserve
- Fissicauda fissicauda* n. gen. n. sp. (tod), *illus.*
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Iomys horsfieldi: Selangor, Tg. Duablas, Tg. Rabok
Aeromys tephromelas: Pahang, Gunang, Benom
Ratufa bicolor: Selangor, Tg. Duablas, Tg. Rabok
(intestin grele of all)
- Fissicauda schmidti* n. sp., *illus.*
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Tragulus javanicus (intestin grele): Selangor, Subang, Forest Reserve
- Fissicauda sonini* n. sp., *illus.*
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Ratufa bicolor (intestin grele): Selangor, Jinjang, Bt. Legong, Forest Reserve
- Fissicauda sundasciuri* (Schmidt, Myers et Kuntz, 1967) n. comb., *illus.*
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 685-692
description, synlophe
Sundasciurus steeri juvenis: Philippines
- Fissicauda sundasciuri* (Schmidt, Myers et Kuntz, 1967)
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Syn.: *Brevistriata sundasciuri* Schmidt, Myers et Kuntz, 1967
- Fontesia Travassos*, 1928
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Anoplostrongylinae*
- Fontesia fontesi* Travassos, 1928, *illus.*
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
redescription, frequency and distribution in host gut, didelphic forms of *Trichostrongyloidea* more abundant than monodelphic ones
Tamandua tetradactyla (estomac, intestin): region de Belem, Province de Para, Bresil
- Freitasia Barus & Coy Otero*, 1968, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochonidae
key
- Fusaria Zeder*, 1800
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Ascaris* L., 1758

- Gaigeria* sp.
Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidian drug (Cozurone), good control of all except *Strongyloides*
- Gaigeria pachyscelis* Railliet & Henry, 1910
Hart, R. J.; and Wagner, A. M., 1971, Onderstepoort J. Vet. Research, v. 38 (2), 111-116
Gaigeria pachyscelis, sheep (exper.), pathological physiology (macrocytic normochromic anaemia, hypoproteinaemia, hypocalcaemia, hyperglycaemia, eosinophilia); some fatalities, due primarily to loss of blood
- Gaigeria pachyscelis*
Horak, I. G.; Snijders, A. J.; and Louw, J. P., 1972, J. South African Vet. Ass., v. 43 (4), 397-403
trematodes and nematodes, sheep (exper.), rafoxanide, efficacy studies
- Gaigeria pachyscelis*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Galeiceps* Railliet, 1916
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Contraceptiva
key
Syn.: *Cloeoascaris* Baylis, 1923
- Gallegostrongylus* n. gen.
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (6), 637-642
Metastrongylidae, Angiostrongylinae
tod: *Gallegostrongylus ibicensis* n. gen., n. sp.
- Gallegostrongylus andersoni* (Petter, 1972) n. comb.
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (6), 637-642
Syn.: *Morerastrongylus andersoni* (Petter, 1972)
- Gallegostrongylus ibicensis* n. gen., n. sp., illus. (tod)
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (6), 637-642
Mus musculus (poumons): Ile d'Ibiza, Balears, Espagne
- Ganguleterakis* sp. 1
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
description
Halcyon pileata (muscular stomach): Rimsko-Korsakov islands
- Ganguleterakis* sp. 2
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
description
Ixobrychus eurhythmus (intestine): Rimsko-Korsakov islands
- Ganguleterakis* dispar
Cervenka, J.; Zajicek, D.; and Nydl, J., 1975, Veterinarstvi, v. 25 (6), 263-264
helminths, geese, Mebendazole
- Ganguleterakis* dispar Schrank, 1790
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anser anser
A. albifrons
(caecum of all): all from Bulgaria
- Ganguleterakis* spumosa
Shikhobalova, N. P.; and Korsak-Paruzhinskaia, L. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 204-214
Ascaridia galli, *Ganguleterakis spumosa*, *Trichocephalus muris*, ionizing radiation of eggs, postradiation effect of high and low temperatures on radiosensitivity as evidenced by development; radiation as possible control of helminths
- Ganguleterakis tenuicauda*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Gastromermis* sp.
Mondet, B.; Pendriez, B.; and Bernadou, J., 1976, Cahiers O.R.S.T.O.M., s. Entom. Med., v. 14 (2), 141-149
Simulium vorax: sud-ouest de la Haute-Volta (riviere Yanaon)
S. hargreavesi: sud-ouest de la Haute-Volta (riviere Yanaon); sud-est du Mali (riviere Farako)
S. cervicornutum: sud-est du Mali (riviere Farako)
S. damnosum s. l.: sud-est du Mali (riviere Farako)
- Gastromermis* sp.
Mondet, B.; Pendriez, B.; and Bernadou, J., 1976, Cahiers O.R.S.T.O.M., s. Entom. Med., v. 14 (2), 141-149
Simulium damnosum
S. vorax
S. adersi
S. alcocki
S. cervicornutum
S. unicornutum
all from Coted'Ivoire (riviere Mounongo)
- Gastromermis boophthorae* Welch and Rubz., illus.
Rubtsov, I. A., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 128-156
ontogenesis of mermithids, illustrated description of structure of body, cuticle, amphids, longitudinal fields, stichosome, osmosome, trophosome and reproductive organs; technique of preparing material

- Gastromermis likhovosi* sp. n., illus.
Rubtsov, I. A., 1976, Zool. Zhurnal, v. 55 (9), 1292-1298
Wilhelmia equina: Rovensk obl., r. Ust'e
- Gastromermis viridis* Welch
Ebsary, B. A.; and Bennett, G. F., 1975, Canad. J. Zool., v. 53 (8), 1058-1062
Simulium corbis
Simulium vittatum
all from insular Newfoundland
- Gastronodus*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Cylicospirura*
key
- Gazellostrongylus* Yeh, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Cooperiinae
- Gazellostrongylus* Yeh, 1956
Khalil, L. F.; and Gibbons, L. M., 1975, J. Helminth., v. 49 (4), 271-279
"no justification for the synonymy of the genus *Gazellostrongylus* with *Paracooperia*." placed in tribe *Ostertagia*
- Gazellostrongylus lerouxi* Yeh, 1956, illus.
Khalil, L. F.; and Gibbons, L. M., 1975, J. Helminth., v. 49 (4), 271-279
redescription
Gazella thomsoni
G. granti
(abomasum of all): all from Kajiado district, Kenya
- Gelanostrongylus* Popova (1952)
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
genus suppressed
- Gelanostrongylus irma*: Popova, 1952
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
as syn. of *Macrostrongylus irma* Johnston & Mawson, 1940
- Gelanostrongylus lesouefi*: Popova, 1952
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
as syn. of *Macrostrongylus lesouefi* Johnston & Mawson, 1939
- Gelanostrongylus macrostoma*: Popova, 1952
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
as syn. of *Macrostrongylus macrostoma* Davey & Wood, 1938
- Gelanostrongylus wallabiae*: Popova, 1952
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
as syn. of *Popovastrongylus wallabiae* (Johnston & Mawson) [n. comb.]
- Gendrespirura* Chabaud, 1958, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key; synonymy
- Gendria leberrei* Bain et Philippon, 1969, illus.
Durette-Desset, M.-C.; and Batcharov, G., 1974, Ann. Parasitol., v. 49 (5), 567-576
female described, male redescribed
Dicroglossus occipitalis: Lome, Togo
- Geopetitia* Chabaud, 1951, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Geopetitiinae
- Geopetitiinae Chabaud, 1951
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Tetrameridae
key
includes: *Geopetitia*
- Gilesia* Sarwar, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Trichostrongylus* Looss, 1905
- Gilsonia* Gedoelst, 1919
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Hadjelia* Seurat, 1916
- Globocephaloides*
Durette-Desset, M. C.; Denke, M. A.; and Murua, R., 1976, Ann. Parasitol., v. 51 (4), 453-460
placed in *Globocephalinae*, excluded from *Amidostomatidae*
- Globocephalus connorfilii* (Lane, 1922)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. (4), 516-521
Rattus sabanus: Malaysia
- Globocephalus urosubulatus*
Bussieras, J., 1976, Rec. Med. Vet., v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Globocephalus urosubulatus*
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
Sus scrofa domesticus x *Sus scrofa cristatus* (small intestine): Aransas National Wildlife Refuge, southern Texas
- Globochona*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Rhabdochona*
key

- Gnathostoma Owen, 1836, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatinae
key
- Gnathostoma spp.
Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Gnathostoma sp.
Thuraisingam, V.; Tan Ewe Aik, P.; and Sandosham, A. A., 1969, Med. J. Malaya, v. 24 (2), 107-112
presumptive case of gnathostomiasis in Chinese woman probably caused by Gnathostoma spinigerum, history of consumption of raw fish: Malaysia
- Gnathostoma procyonis
Barnstable, R. W.; and Dyer, W. G., 1974, Tr. Illinois State Acad. Sc., v. 67 (4), 451-460
Procyon lotor (stomach): southern Illinois
- Gnathostoma procyonis or Physaloptera rara
Bartsch, R. C.; and Ward, B. C., 1976, Vet. Path., v. 13 (4), 241-249
"may have been either Gnathostoma procyonis or Physaloptera"
raccoons (gastric mucosa): southeastern Florida
- Gnathostoma procyonis Chandler, 1942
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (small intestine): Florida
- Gnathostoma spinigerum
Daengsvang, S., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 95-101
review of natural sources of infection and methods of disease transmission
- Gnathostoma spinigerum
Daengsvang, S.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 359-361
Gnathostoma spinigerum, cats (nat. and exper.), successful elimination of adult worms from gastrointestinal tract using ancylool disopenol but larvae seemed resistant to therapy
- Gnathostoma spinigerum, illus.
Daengsvang, S.; Sangsingkeo, P.; and Senivong-Na-Ayudhaya, B., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 260-262
Gnathostoma spinigerum larva excised from finger of man who had symptoms of migratory swelling, pain and itching over left forearm for 2 years, possible transmission by consumption of partially cooked chicken: Thailand
- Gnathostoma spinigerum, illus.
Fontan, R.; Beauchamp, F.; and Beaver, P. C., 1975, Bull. Soc. Path. Exot., v. 68 (6), 557-566
human case reports of human intestinal and pulmonary gnathostomiasis: Laos
- Gnathostoma spinigerum Owen, 1836, illus.
Lim, B. L., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (4), 530-533
Gnathostoma spinigerum adult worms removed from naturally infected Prionodon linsang (stomach) during survey for possible infections in civet and wild cats, morphologic statistics, discussion of civet cat dietary habits in relationship to infection: Bukit Mandol Forest Reserve, Selangor, Kuala Lumpur
- Gnathostoma spinigerum, illus.
Nitidandhaprabhas, P.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (1), 49-51
fourth reported case of human urinary tract gnathostomiasis, woman with history of eating raw fish passed adult male Gnathostoma spinigerum in urine, probable parasite migration through lung and spinal column to bladder: Thailand
- Gnathostoma spinigerum, illus.
Nitidandhaprabhas, P.; Hanchansin, S.; and Vongsloesvidhya, Y., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 547-548
living adult male Gnathostoma spinigerum coughed up by Thai woman, probably infected from eating pork, case report: Korat, Thailand
- Gnathostoma spinigerum
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Gnathostoma spinigerum
Prasansuk, S., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 260 [Letter]
Gnathostoma spinigerum in man, neuro-otological symptoms produced by mature male worm later removed from right external acoustic meatus, residual facial palsy, case report: Bangkok
- Gnathostoma spinigerum
Thuraisingam, V.; Tan Ewe Aik, P.; and Sandosham, A. A., 1969, Med. J. Malaya, v. 24 (2), 107-112
presumptive case of gnathostomiasis in Chinese woman probably caused by Gnathostoma spinigerum, history of consumption of raw fish: Malaysia
- Gnathostoma spinigerum
Trueman, K. F.; and Ferris, P. B. C., 1977, Austral. Vet. J., v. 53 (10), 498-499
Gnathostoma spinigerum, cats, 3 case reports, histopathology, clinical signs, fatal case with perforated stomach: Townsville, Queensland, Australia

- Gnathostoma vietnamicum Le-Van-Hoa 1965, illus. Daengsvang, S., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (1), 63-70
Gnathostoma vietnamicum, morphology and morphometric measurements show variation of cephalic hooklet rows on the head bulbs and few body cuticular spines, gross pathology in naturally infected *Aonyx cinerea* (urinary system) and preliminary experimental life cycle in *Mesocyclops leuckarti* and *Trichopsis vittatus*: Thailand
- Gnathostomatidae Railliet, 1895
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatoidea
key to subfamilies
includes: Spiroxyinae; Ancyracanthinae; Gnathostomatinae
- Gnathostomatinae (Railliet, 1895, fam.) Baylis & Lane, 1920
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatidae
key; key to genera
includes: Tanqua; Echinocephalus; Gnathostoma
- Gnathostomatoidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Gnathostomatoidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
includes: Gnathostomatidae
- Gnathostomiasis
Thuraisingam, V.; Tan Ewe Aik, P.; and Sando-sham, A. A., 1969, Med. J. Malaya, v. 24 (2), 107-112
presumptive case of gnathostomiasis in Chinese woman probably caused by *Gnathostoma spinigerum*, history of consumption of raw fish: Malaysia
- Goezia Zeder, 1800, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Goeziinae
synonymy
- Goezia Zeder, 1800
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
amended key to subgenera
- Goezia subgen.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
key
- Goezia (*Pseudogoezia*) sp. Rai, 1967
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
as syn. of *G. (P.) chitali* nom. nov. [i.e., n. sp.]
- Goezia (*Pseudogoezia*) *chitali* nom. nov. [i.e., n. sp.], illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Syn.: *Goezia (Pseudogoezia)* sp. Rai, 1967
Notopterus chitala (stomach): Taunsa Barrage, Pakistan
- Goezia (*Pseudogoezia*) *heteropneusti* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Heteropneustes fossilis (intestine): Taunsa Barrage, Pakistan
- Goezia (*Goezia*) *taunsa* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Channa gachua (intestine)
Notopterus chitala (stomach)
all from Taunsa Barrage, Pakistan
- Goeziinae Travassos, 1919
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakidae
key
includes: *Goezia*
- Gongylonema
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Gongylonema*
key
- Gongylonema Molin, 1857, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Gongylonematidae
key to subgens.
includes: *Gongylonema*; *Gongylonemoides*
- Gongylonema
Isenbuegel, E., 1976, Prakt. Tierarzt, v. 57, Sondernummer, 21-27
carbon tetrachloride, unsuccessful
Igel
- Gongylonema sp. Molin, 1857
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (under crop lining): Vom area, Benue Plateau State, Nigeria
- Gongylonema sp., illus.
King, N. W., jr., 1976, Scient. Publication (317). Pan Am. Health Organ., 169-198

- Gongylonema* sp.
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (stomach): Florida
- Gongylonema* spec.
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Gongylonema* sp.
Schulte, J. W.; Klimstra, W. D.; and Dyer, W. G., 1976, J. Wildlife Management, v. 40 (3), 579-581
Odocoileus virginianus clavium (feces): Big Pine Key, Florida
- Gongylonema* congolense (Fain, 1955)
Fabiyl, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Gongylonema* congolense Fain, 1955
Fabiyl, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (under crop lining): Vom area, Benue Plateau State, Nigeria
- Gongylonema* dipodomys Kruidenier and Peebles, 1953
King, S. R.; and Babero, B. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 241-248
Dipodomys merriami: Nevada
- Gongylonema* ingluvicola
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Gongylonema* neoplasticum
Mishra, G. S.; and Gonzalez, J. P., 1975, Arch. Inst. Pasteur Tunis, v. 52 (1-2), 71-87
Rattus norvegicus (estomac, oesophage): Tunis, Tunisia
- Gongylonema* neoplasticum Fibiger et Ditlevsen, 1914, illus.
Quentin, J. Cl.; and Krishnasamy, M., [1976], Ann. Parasitol., v. 50 (6), 1975, 795-812
Tupaia glis (paroi oesophagienne): station biologique de Sepilok, Sandakan, Province de Sabah
- Gongylonema* neoplasticum (Fibiger and Ditlevsen, 1914)
Singh, M.; and Cheong Chee Hock, 1971, South-east Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus r. jarak: Malaysia
- Gongylonema* pithyusensis n. sp., illus.
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (1), 13-18
Eliomys quercinus ophiusae (muqueuse de l'oesophage): Ile de Formentera (Balears, Espagne)
- Gongylonema* problematicum Schulz, 1924
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Blaps fausti bactriana
B. deplanata reichardtii
Trigonoscelis gemmulata
T. ceromatica
Pisterotarsa kiritschenkoi
Pseudeuthripta tadzshikistana
all from Tadzshik SSR [and/or] Uzbek SSR
- Gongylonema* problematicum
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Trigonoscelis gigas
Blaps fausti
B. scutellata
B. seriata
all from Turkmenia
- Gongylonema* pulchrum Molin, 1857, illus.
Amaszta, M.; et al., 1972, Parasitol. Hungar., v. 5, 239-246
Gongylonema pulchrum, 3 specimens removed within one month from oral areas of young woman (lips, oral fold, oral mucous membrane), associated complaints of gastrointestinal disturbances, apparent cure after administration of levamisole, source of infection could not be identified: Debrecen, Hungary
- Gongylonema* pulchrum, illus.
Amaszta, M.; et al., 1973, Orvosi Hetilap, v. 114 (37), 2237-2238
Gongylonema pulchrum, case report of human infection, levamisole therapy: Hungary
- Gongylonema* pulchrum
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
Sus scrofa domesticus x Sus scrofa cristatus (tongue, esophagus): Aransas National Wildlife Refuge, southern Texas
- Gongylonema* pulchrum
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (esophagus): southeastern United States
- Gongylonema* pulchrum Molin, 1857
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Prosodes biformis: environs of Dushanbe town
- Gongylonema* pulchrum
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (esophagus): Kentucky

- Gongylonema pulchrum* Molin, 1857
Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
Capreolus capreolus (esophagus): southern Bulgaria
- Gongylonema pulchrum*
Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
anthropozoonotic helminthiasis associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Gongylonema pulchrum*
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Trigonoscelis punctipleuris
Blaps fausti
all from Turkmenia
- Gongylonema pulchrum*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries
Odocoileus virginianus
all from Hardy County, West Virginia
- Gongylonema pulchrum* Molin, 1857
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (esophagus): Cumberland County, New Jersey; Oklahoma
- Gongylonema pulchrum*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Aphodius lugens
A. makowskyi
A. fimetaris
A. granarius
A. erraticus
Gymnopleurus morsus
G. aciculatus
Geotrupes impressus
Chironitis hungaricus
C. fuscifer
Copris hispanus
Copris lunaris
Onitis humerosus
Oniticellus pallipes
Oniticellus fulvus
Onthophagus amyntas
Onthophagus pugurgus
Onthophagus koshantschikoff
Onthophagus nuchicornis
Onthophagus gibbulus
Onthophagus taurus
Onthophagus fracticornis
Onthophagus transcaspicus
Scarabaeus sacer
Blaps oblonga
B. deplanata
all from Uzbekistan
- Gongylonema sumani* Bhalerao, 1933
Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
Fowl, domestic
Fowl, desi
all from Punjab State, India
- Gongylonema verrucosum* (Giles, 1892)
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (rumen): Oklahoma
- Gongylonematidae (Hall, 1916, subfam.) Sobolev, 1949
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruroidea
key
includes: *Gongylonema*
- Gongylonemoides*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Gongylonema*
key
- Gopheruris n. g.*
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Pharyngodonidae
tod: *G. aspicula n. sp.*
- Gopheruris aspicula n. sp.* (tod), illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" *G. agassizii* (colon): desert de Sonora, Mexique
- Gordiacea sp.*, proche du *Neochordodes occidentalis*, illus.
Arvy, L.; and Sowa, R., 1976, Ann. Parasitol., v. 51 (1), 111-120
Baetis muticus: region de Cracovie, Pologne
- Gordius sp.*
Zalom, F. G., 1977, Am. Midland Naturalist, v. 97 (1), 229-230
Notonecta undulata (abdomen): Lake Mountain Lookout, NE of McNary, Apache Co., Arizona
- Gordius aquaticus*, illus.
Arvy, L.; and Sowa, R., 1976, Ann. Parasitol., v. 51 (1), 111-120
- Gordius panighettensis*, illus.
Lora Lamia Donin, C.; and Cotelli, F., 1977, J. Ultrastructure Research, v. 61 (2), 193-200
Gordius panighettensis, ultrastructure of spermatozoon; aberrant immotile, rod-like sperm unlike any other in zoological scale
- Grammocephalus hybridatus* van der Westhuisen, 1938, illus.
Pellegrini, N.; et al., 1976, Ann. Fac. Med. Vet., Univ. Pisa, v. 28, 1975, 139-149
Grammocephalus hybridatus, *Elephas maximus* (bile ducts), histopathological changes in liver and pancreas: died in zoo, Tuscany
- Grammocephalus hybridatus*
Pillay, K. R. Sukumara; et al., 1976, Indian Vet. J., v. 53 (9), 670-671
Grammocephalus hybridatus, Indian elephant, case history, pathogenesis: Tithmathy, Coorg District, Karnataka State

- Graphidiella Olsen, 1948
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Graphidiinae
- Graphidiella ochotonae
Seesee, F. M., 1973, Am. Midland Naturalist,
v. 89 (2), 257-265
key
Ochotona p. princeps (stomach): St. Joe
Baldy Mountain, Benewah County, Idaho
- Graphidiella szechuanensis sp. nov., illus.
Wu, C., 1977, Tung Wu Hsueh Pao (Acta Zool.
Sinica), v. 23 (3), 283-285
Ochotona sp. (stomach): Shiqu and Seda,
Sichuan Province
- Graphidiinae Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae
includes: Graphidium (type genus); Graphi-
diella; Hyostrongylus; Parostertagia
- Graphidioides Cameron, 1923
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Trichostrongylinae
- Graphidioides taglei n. sp., illus.
Babero, B. B.; and Cattán, P. E., 1975, Bol.
Chileno Parasitol., v. 30 (3-4), 68-76
Octodon degus (estomago e intestino delgado):
Quebrada de la Plata, Santiago, Chile
- Graphidiops Lent et Freitas, 1938
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Graphidiops costalimai Lent et Freitas, 1938,
illus.
Durette-Desset, M. C.; Chabaud, A. G.; and
Cassone, J., 1977, Bull. Mus. National Hist.
Nat., Paris, 3. s. (428), Zool. (298), 133-158
redescription of male, frequency and dis-
tribution in host gut, didelphic forms of
Trichostrongyloidea more abundant than mono-
delphic ones
Tamandua tetradactyla (estomac): region de
Belem, Province de Para, Bresil
- Graphidiops cyclopi n. sp., illus.
Diaw, O. T., [1977], Bull. Mus. National Hist.
Nat., Paris, 3. s. (405), 1976, Zool. (282),
1065-1089
Cyclopes didactylus (duodenum): Belem,
province de Para, Bresil
- Graphidium Railliet et Henry, 1909 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Graphidiinae
- Graphidium strigosum
Cremers, H. J. W. M.; Jansen, J.; and Swier-
stra, D., 1975, Tijdschr. Diergeneesk.,
v. 100 (22), 1209-1211
Oryctolagus cuniculus: Netherlands
- Graphidium strigosum
Kutzer, E.; and Frey, H., 1976, Berl. u. Mün-
chen. Tierarztl. Wchnschr., v. 89 (24), 480-
483
Lepus europaeus: Austria
- Graphidium strigosum
Kutzer, E.; and Frey, H., 1976, Ztschr. Para-
sitenk., v. 50 (2), 213-214
Lepus europaeus
- Graphinema gen. n.
Guerrero, C. A.; and Rojas, J. E., 1969, Bol.
Chileno Parasitol., v. 24 (3-4), 127-129
Graphidiinae
tod: G. aucheniae n. sp.
- Graphinema [sp.]
Leguia, G.; and Bendezu, P., 1974, Rev. In-
vest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in
fecal egg counts, 2 year period, pregnant
Lama pacos: Central Sierra of Peru (Dept.
Pasco)
- Graphinema aucheniae n. g., n. sp. (tod), illus.
Guerrero, C. A.; and Rojas, J. E., 1969, Bol.
Chileno Parasitol., v. 24 (3-4), 127-129
Lama pacos
Vicugna vicugna
Lama glama
(abomasum of all): all from Cuzco y Puno,
Peru, Sud America
- Graphinema aucheniae
Guerrero, C.; Rojas, M.; and Vargas, J., 1974,
Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity
of 1-tetramisole, significant body weight
gain in treated animals
- Graphinema aucheniae
Vargas, J.; Guerrero, C.; and Rojas, M., 1972,
Rev. Invest. Pecuarias, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight
toxicity
- Grosspiculagia (Orloff, 1933) Sarwar, 1956
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Guinea worm. See [Dracunculus medinensis]
- Guyanema Petter, 1975, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Guyanemidae
- Guyanemidae Petter, 1975
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Dracunculoidea
key
includes: Guyanema

Gyalocephalus

Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method

Gyalocephalus

Reinecke, R. K.; and le Roux, D. J., 1972, *J. South African Vet. Ass.*, v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective

Gyalocephalus capitatus

Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method

Gyalocephalus capitatus

Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole

Gyalocephalus capitatus

Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England

Gynaecophila

Pence, D. B.; Mollhagen, T.; and Forrester, D. J., 1975, *J. Parasitol.*, v. 61 (3), 494-498
subgenus of *Tetrameres*
subgeneric status discussed, probable revision indicated

- Habronema Diesing, 1861, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
- Habronema
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, not effective
- Habronema spp.
Duncan, J. L.; et al., 1977, Equine Vet. J., v. 9 (3), 146-149
immature strongyles, ponies, fenbendazole
- Habronema sp., illus.
Finazzi, M.; Lattanzio, G.; and Malnati, G., 1977, Clin. Vet., Milano, v. 100 (1), 23-31
Habronema sp., stallion (prepuce, glans), histopathology, nodules surrounded by hyperplastic connective tissue
- Habronema spp.
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Habronema spp.
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxbendazole
- Habronema americanum
Kocan, A. A.; and Locke, L. N., 1974, J. Wildlife Dis., v. 10 (1), 8-10
Haliaeetus leucocephalus: Minnesota; Iowa; Illinois; Wisconsin
- Habronema megastoma
Reddy, A. B.; Gaur, S. N. S.; and Sharma, U. K., 1976, Indian J. Animal Sc., v. 46 (4), 207-210
Habronema muscae, Habronema megastoma, horses, gross and microscopic pathological changes, stomach and other organs
- Habronema microstoma
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Habronema muscae
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Habronema muscae
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Habronema muscae
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Habronema muscae
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Habronema muscae
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Habronema muscae
Reddy, A. B.; Gaur, S. N. S.; and Sharma, U. K., 1976, Indian J. Animal Sc., v. 46 (4), 207-210
Habronema muscae, Habronema megastoma, horses, gross and microscopic pathological changes, stomach and other organs
- Habronema numidae Ortlepp, 1938
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (under gizzard lining): Vom area, Benue Plateau State, Nigeria
- Habronematidae (Chitwood & Wehr, 1932) Ivaschkin, 1961
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematoidea
key; key to subfams.
includes: Habronematinae; Parabronematinae; Histioccephalinae
- Habronematinae Chitwood & Wehr, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematidae
key; key to genera
includes: Odontospirura; Draschia; Chitwoodspirura; Sicarius; Gendrespirura; Excisa; Procyrnea; Cyrnea; Metacyrnea; Habronema
- Habronematoidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Habronematoidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina

- Habronematoidea**
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 Spirurida
 key to families
 includes: Hedruridae; Tetrameridae; Habronematidae; Cystidicolidae
- Hadjelia Seurat, 1916, illus.**
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 Histiocéphalinae
 key; synonymy
- Hadjelia neglecta (Lent and Frietas, 1939)**
 Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 39 (2), 467-473
 comparison of helminth fauna of common and purple gallinules
 Porphyryla martinica (proventriculus):
 Florida
- Hadjelia pyrhnota sp. n., illus.**
 Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
 Petrochelidon pyrhnota (proventriculus):
 north of Rockport, Colorado, U.S.A.
- Hadjelia truncata (Creplin, 1825)**
 Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
 role of Tenebrionidae as intermediate hosts
 Pachyscelis banghaasi
 Trigonoscelis gemmulata
 Dila bucharica
 all from Tadzhik SSR
- Hadjelia truncata**
 Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
 Adesmia servillei schatzmayri
 Trigonoscelis gigas
 Pisterotarsa gigantea subsp. zoubkoffi
 P. kessleri
 all from Turkmenia
- Haemonchinae (Skrjabin et Schulz, 1937, tribu)**
 Skrjabin et Schulz, 1952
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Trichostrongylidae
 includes: Haemonchus (type genus); Ashworthius; Biogastranema; Boehmiella; Leiperiatus; Mecistocirrus
- Haemonchus**
 Baines, D. M.; and Colegrave, A. J., 1977, Vet. Rec., v. 100 (11), 217-219
 gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole:
 England and Wales
- Haemonchus**
 Bliss, D. H.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (10), 1612-1617
 milk production in dairy cows exposed to mixed trichostrongylid larvae, results indicate that greatest milk loss occurs during the first 90 days of lactation, relationship between exposure time and stage of lactation
- Haemonchus**
 Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
 sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Haem[onchus]**
 Brunson, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
 lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Haemonchus**
 Chhabra, R. C.; Bali, H. S.; and Toor, L. S., 1976, J. Research, Punjab Agric. Univ., v. 13 (3), 308-311
 gastrointestinal strongyles in sheep, critical drug evaluation, thiabendazole (most effective), tetramisole (good results), morantel tartrate (fair results), clioxanide and methyridine (least effective): India
- Haemonchus**
 Chowanec, W.; et al., 1975, Med. Wet., v. 31 (12), 741-743
 Fasciola hepatica, Haemonchus, Trichostrongylus, Nematodirus, cattle, Nilzan, Zanil, field trials, good results
- Haemonchus**
 Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
 nematodes, cattle, oxbendazole, drug efficacy
- Haemonchus**
 Crowley, J. W., jr.; et al., 1977, Am. J. Vet. Research, v. 38 (5), 689-692
 lungworms, gastrointestinal parasites, cattle, 3 controlled critical trials, highly effective
- Haemonchus**
 Duwel, D., 1977, Cahiers Bleus Vet. (26), 201-215
 fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Haemonchus Cobb, 1898 (type genus)**
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Trichostrongylidae, Haemonchinae
- Haemonchus**
 Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
 sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district

- Haemonchus**
Fudalewicz-Niemczyk, W.; et al., 1976, Acta Zootech., Bratislava (32), 5-19
gastrointestinal helminths, mountain sheep, nilverm and zani, favorable influence on body weight and wool production of treated animals, no influence of treatment on fertility: Poland
- Haemonchus**
Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Haemonchus**
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Haemonchus**
Klein Mori, J., 1972, Rev. Med. Vet. y Parasitol., Maracay, v. 24 (1-8), 1971-1972, 207-226
gastrointestinal nematodes, sheep, Neguvon, Ripercol, Thibenzoline, comparison, various management systems, all effective, Ripercol easiest to administer, Neguvon somewhat toxic
- Haemonchus**
Misra, S. C.; and Ruprah, N. S., 1974, Indian Vet. J., v. 51 (2), 147-148
Haemonchus larvae, survival on 2 pastures under changing temperature and relative humidity: India
- Haemonchus**
Qadir, A. N. M. A., 1976, Indian Vet. J., v. 53 (11), 855-858
gastrointestinal nematodes, goats and calves, urea for pasture control of free-living stages
- Haemonchus**
Sewell, M. M. H., 1973, Vet. Rec., v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Haemonchus**
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, Am. J. Vet. Research, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Haemonchus**
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcvi
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Haemonchus**
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Haemonchus**
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Haemonchus**
Tsolov, B.; and Tsanov, G., 1975, Vet. Sbirka, v. 73 (9), 20-22
strongyloids of sheep, economics of tetramisole treatment, body weight, milk production, not effective against Trichocephalus
- Haemonchus**
Vashishta, M. S.; and Singh, R. P., 1977, Livestock Advis., v. 2 (7), 35-38
Haemonchus epidemic in sheep and goats, banminth-II, fenbendazole, helmatac, thibenzole, good results; development of geophagia in survivals or treated groups: Haryana
- Haemonchus**
Zeakes, S. J.; et al., 1976, Am. J. Vet. Research, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos crumbles and naftalofos boluses
- Haemonchus [sp.]**
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
ruminants, survey, treatment, economic importance: Kaedi area (Mauritania)
- Haemonchus spp.**
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against Ostertagia spp., similar efficacy against other species), both drugs increased calves' weight gains
- Haemonchus spp.**
El-Abdin, Y. Z.; et al., 1975, Egypt. J. Vet. Sc., v. 12 (1), 31-43
serum constituents and serum enzyme activities, normal and nematode infested Camelus dromedarius: Cairo abattoir
- Haemonchus sp.**
Krishna Iyer, P. P.; and Peter C. T., 1975, Kerala J. Vet. Sc., v. 5 (2), 121-123
gastrointestinal nematodes, goats, methyridine
- Haemonchus sp.**
Lara, S. I. M.; de Oliveira, C. M. B.; and Porto, J. C. A., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 93-99
Haemonchus sp., sheep (exper.), cobalt sulfate diet supplement, increased production and size of eggs, lower number of worms in autopsy

- Haemonchus* spp.
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Haemonchus* sp.
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Haemonchus* sp.
de Oliveira, A. R., 1976, Arq. Inst. Biol., Sao Paulo, v. 43 (1-2), 53-56
Oesophagostomum sp., *Cooperia* sp., *Haemonchus* sp., calves (exper.), no correlation between level of infestation and circulating eosinophils, may result from eosinophil migration to affected organs or bone marrow exhaustion
- Haemonchus* sp.
Qadir, A. N. M. A., 1976, Indian Vet. J., v. 53 (6), 448-450
Haemonchus sp., *Trichostrongylus* sp., *Oesophagostomum* sp., larvicidal action of 4 chemical compounds on infective nematode larvae in experimental outdoor plots; urea most effective
- Haemonchus* spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Haemonchus* sp.
Tongson, M. S.; and Montenegro, M. M., 1975, Philippine J. Vet. Med., v. 13 (1-2), 170-182
purified microfine phenothiazine + lead arsenate, anthelmintic efficiency, good results, goats: Philippines
- Haemonchus* sp.
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, *Camelus dromedarius*, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Haemonchus* sp.
Van Geldorp, P. J. A.; and Schillhorn van Veen, T. W., 1976, Vet. Parasitol., v. 1 (3), 265-269
periparturient rise in faecal helminth egg counts of Udah sheep, suggested that increase was due to helminths (mainly *Haemonchus* sp.) which had been inhibited during dry season and resumed development at beginning of rainy season, rapid decline in egg counts 5-6 weeks after lambing considered to be due to self cure associated with high rainfall: Zaria area of Nigeria
- Haemonchus* sp.
Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidian drug (Cozurone), good control of all except *Strongyloides*
- Haemonchus* sp.
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damaliscus dorcas dorcas (abomasum): captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Haemonchus* sp.
Williams, J. C.; et al., 1977, Vet. Rec., v. 101 (24), 484-486
Ostertagia ostertagi, cattle, albendazole, good results against inhibited fourth stage larvae as well as developing stages and adults, high efficacy against *Haemonchus* and *Cooperia* spp., no signs of toxicity
- Haemonchus* spp.
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Haemonchus* sp.
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 77-81
Connochaetes gnou
Redunca fulvorufula
(abomasum of all): all from Mountain Zebra National Park
- Haemonchus* spp.
Young, E.; and Basson, P. A., 1976, J. South African Vet. Med. Ass., v. 47 (1), 57
Cordophilus sagittus, *Haemonchus* spp., eland, pathology, levamisole hydrochloride, good results against gastro-intestinal parasites: Kruger National Park, translocated from Addo Elephant National Park
- Haemonchus bedfordi* Le Roux, 1929
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (abomasum): Kruger National Park
- Haemonchus bedfordi*
Gibbons, L. M.; and Khalil, L. F., 1976, Trop. Animal Health and Prod., v. 8 (3), 168
sheep
goats
(gut of all): all from Kajiado district, Kenya
- Haemonchus bedfordi* Le Roux
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya

- Haemonchus bedfordi*, illus.
Sachs, R.; Gibbons, L. M.; and Lweno, M. F., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 467-475
Syncerus caffer
Kobus defassa
Hippotragus niger
Connochaetes taurinus
Damaliscus korrigum
Alcelaphus buselaphus cokei
Alcelaphus buselaphus lichtensteini
Aepyceros melampus
Gazella granti
G. thomsonii
Giraffa camelopardalis
(abomasum of all): all from Tanzania
- Haemonchus contortus*
van Adrichem, P. W. M.; and Shaw, J. C., 1977, J. Animal Sc., v. 45 (3), 423-429
gastrointestinal nematodes, effects on growth performance and milk production in cambendazole-treated vs. non-treated monozygous twin cattle naturally infected on pasture during the first lactation period
- Haemonchus contortus*
Ajayi, J. A.; and Todd, A. C., 1973, Am. J. Vet. Research, v. 34 (3), 449-450
Haemonchus contortus, populations of differing pathogenicity used to preimmunize and challenge sheep, ability to distinguish 2 worm populations in sheep exposed to both by separation in curve of distribution frequencies of parasite lengths
- Haemonchus contortus*
Allonby, E. W.; and Urquhart, G. M., 1975, Vet. Parasitol., v. 1 (2), 129-143
Haemonchus contortus, Merino ewes and their lambs, epidemiology and pathogenic significance, faecal egg counts, worm burdens, haematological indices, clinical signs, levels of infective larvae on pasture, classical acute haemonchosis occurred during high rainfall periods, self-cure confirmed as flock phenomenon, importance of moderate infections: Naivasha, Kenya
- Haemonchus contortus*
Allonby, E. W.; and Urquhart, G. M., 1976, Research Vet. Sc., v. 20 (2), 212-214
Haemonchus contortus, Merino sheep, possible relationship between haemoglobin type and resistance to haemonchosis: Kenya
- Haemonchus contortus*
Anderson, P. J. S.; and Marais, F. S., 1972, J. South African Vet. Ass., v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Haemonchus contortus*
Bali, M. K.; and Singh, R. P., 1976, Indian J. Animal Research, v. 10 (2), 111-112
Haemonchus contortus, *Oesophagostomum* sp., *Trichostrongylus* sp., *Trichuris* sp., sheep, goats, morantel tartrate, good results against all parasites except for *Trichuris* sp.
- Haemonchus contortus* (Rudolphi, 1803)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (abomasum): Kruger National Park
- Haemonchus contortus*
Benitez-Usher, C.; et al., 1977, Vet. Parasitol., v. 3 (4), 327-342
Haemonchus contortus, Scottish Blackface lambs, immunization with gamma-irradiated larvae, roles of host age, size of immunizing dose, previous exposure to infection, and anthelmintic (thiabendazole) therapy
- Haemonchus contortus*
Benz, G. W.; and Ernst, J. V., 1977, Am. J. Vet. Research, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.), albendazole did not significantly reduce infestation
- Haemonchus contortus*
Berger, J., 1975, J. South African Vet. Ass., v. 46 (4), 369-372
sheep, laboratory strain and parbendazole resistant field strain of *Haemonchus contortus*, resistant to five benzimidazoles, susceptible to levamisole and haloxon
- Haemonchus contortus* (Rudolphi, 1803)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Haemonchus contortus*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Haemonchus contortus*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Haemonchus contortus*
Boisvenue, R. J.; Emmick, T. L.; and Galloway, R. B., 1977, Exper. Parasitol., v. 42 (1), 67-72
Haemonchus contortus, some compounds with juvenile hormone activity inhibited in vitro development of infective larvae, none of these compounds had anthelmintic properties against *Ascaris suum* or *Nematospiroides dubius* in mice

- Haemonchus contortus**
Campbell, W. C.; and Thomson, B. M., 1973, Austral. Vet. J., v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus colubriformis* proved uninfecive even if they had not been frozen
- Haemonchus contortus**
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Haemonchus contortus**
Chen, P.; and Soulsby, E. J. L., 1976, Internat. J. Parasitol., v. 6 (2), 135-141
Haemonchus contortus infections in ewes during pregnancy, parturition, and lactation, blastogenic responses of peripheral blood leukocytes to non-specific mitogen, non-helminth antigens, and specific 3rd stage larval antigen, relationship to 'spring-rise' and 'self-cure' phenomena, possible hormonal factors
- Haemonchus contortus**
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Haemonchus contortus**
Coadwell, W. J.; and Ward, P. F. V., 1977, Parasitology, v. 74 (2), 121-132
Haemonchus contortus, sheep (exper.), suggested that cyclic change in parasite growth pattern and arrested development is controlled by seasonal variation in concentration of substance(s) in host blood, sex of host and duration of infection had no effect on parasite length, age of host did relate to parasite length but relationship may have been an artifact
- Haemonchus contortus**
Coles, G. C.; and Simpkin, K. G., 1977, Research Vet. Sc., v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Haemonchus contortus**
Colglazier, M. L.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with *Trichuris* spp. and *Moniezia expansa*
- Haemonchus contortus**
Colglazier, M. L.; Kates, K. C.; and Enzie, F. D., 1975, J. Parasitol., v. 61 (4), 778-779
Haemonchus contortus, cambendazole-resistant strain, cross-resistance to thiabendazole, mebendazole, and oxbendazole, fully sensitive to levamisole
- Haemonchus contortus**
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Haemonchus contortus**
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 518-525
gastrointestinal nematodes and lungworms, calves (exper.), morantel tartrate, efficacy in 5 controlled trials, toxicity experiments demonstrate wide safety margin
- Haemonchus contortus**
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (abomaso): Sardegna
- Haemonchus contortus**
Dalton, S. E., 1977, Parasitology, v. 75 (2), xvi [Abstract]
Haemonchus contortus, *Nematodirus spathiger*, effect of thiophanate on egg output, hatchability, and worm burden, sheep
- Haemonchus contortus**
Dargie, J. D., 1975, Symposia Brit. Soc. Parasitol., v. 13, 1-26
helminth diseases of sheep, red cell and plasma protein metabolism, anaemia, applications of radioisotopic methods, extensive review with particular emphasis on *Fasciola hepatica* and *Haemonchus contortus*
- Haemonchus contortus**
Daskalov, P., 1972, Izvest. Tsentral. Khelmin. Lab., v. 15, 49-56
Haemonchus contortus, morphological variation of linguiform females, cuticular inflations develop with increasing worm age but depend on worm reproductive activity rather than directly on age, occurrence of various forms of linguiform females does not depend on season, cuticular inflations should not be used as taxonomic characters since they are nonhereditary morphological variations
- Haemonchus contortus**
Daskalov, P., 1974; Izvest. Tsentral. Khelmin. Lab., v. 17, 51-57
Haemonchus contortus, genetic studies of female polymorphism
- Haemonchus contortus**
Daskalov, P.; Komandarev, S.; and Mikhov, L., 1972, Izvest. Tsent. Khelmin. Lab., v. 15, 57-67
Haemonchus contortus, males and three morphological types of females, comparison of protein fractions by disc electrophoresis in polyacrylamide gel, differences found between sexes but none between types of females, nonhereditary morphological variations
- Haemonchus contortus, illus.**
Daskalov, P.; and Mutafova, T., 1972, Izvest. Tsentral. Khelmin. Lab., v. 15, 69-80
Haemonchus contortus, effect of ultraviolet and gamma rays and refrigeration on infective larvae of different morphological types of females

- Haemonchus contortus**
Dennis, R. D. W., 1977, *Internat. J. Parasitol.*, v. 7 (3), 181-188
Haemonchus contortus, extracts, amount of ecdysone-like material
- Haemonchus contortus**
Dewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Haemonchus contortus**
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Haemonchus contortus**
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Haemonchus contortus**
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Haemonchus contortus**
Dyk, V.; and Chroust, K., 1975, *Veterinarstvi*, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Haemonchus contortus**
Eichler, D. A., 1973, *Brit. Vet. J.*, v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Haemonchus contortus**
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*, v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Haemonchus contortus**
Eve, J. H.; and Kellogg, F. E., 1977, *J. Wildlife Management*, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (*Odocoileus virginianus*) density: south-eastern United States
- Haemonchus contortus**
Folz, S. D.; Rector, D. L.; and Ceng, S., 1976, *J. Parasitol.*, v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Haemonchus contortus**
Georgieva, D.; and Vladimirova, A., 1975, *Vet. Med. Nauk.*, v. 12 (9), 61-65
Haemonchus contortus, lambs, drop in albumin and rise in globulin fractions
- Haemonchus contortus**
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, *Vet. Med. Nauki*, v. 12 (1), 69-75
lambs (exper.), course of infection
- [**Haemonchus contortus**] **Khemonkhous kontortus**
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, *Vet. Sbirka*, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of tetramisole, group and individual applications
- Haemonchus contortus**
Ghosh, S. S.; et al., 1976, *Indian J. Animal Health*, v. 15 (1), 81-82
Haemonchus contortus, goats, high mortality rate, histopathology: Union Territory of Mizoram
- Haemonchus contortus**
Gibbons, L. M.; and Khalil, L. F., 1976, *Trop. Animal Health and Prod.*, v. 8 (3), 168
sheep
goats
(gut of all): all from Kajiado district, Kenya
- Haemonchus contortus**
Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (4), 533-534
Haemonchus contortus, *Ostertagia* sp., sheep, "spring rise" in fecal egg counts, higher incidence in bred sheep than in unbred sheep, role of contaminated bedding: Maine
- Haemonchus contortus**
Gibson, T. E.; and Everett, G., 1976, *Brit. Vet. J.*, v. 132 (1), 50-59
Haemonchus contortus, development and survival of free-living stages on pasture studied over period of 3 years, only in July, August, and September were climatic conditions favorable, concluded that climate in Southern England is not ideal for development and survival of preparasitic stages of *Haemonchus contortus*
- Haemonchus contortus**
Goldberg, A., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Haemonchus contortus**
Guimaraes, M. P.; et al., 1976, *Arq. Escola Vet. Univ. Fed. Minas Gerais*, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil

- Haemonchus contortus*, *illus.*
Guttekova, A.; and Zmoray, I., 1975, *Biologia, Bratislava, s. B, Zool.*, v. 30 (8), 605-614
Haemonchus contortus, ultrastructure of intestine, relationship to diet and metabolism; possibly phylogenetically young parasite in adaptation to host
- Haemonchus contortus*, *illus.*
Guttekova, A.; and Zmoray, I., 1977, *Biologia, Bratislava, v. 32 (8), s. B, Zool.* (3), 585-590
Haemonchus contortus, ultrastructure of body wall, strong musculature and many mitochondria as adaptation for continual undulation while attached to host abomasum wall
- Haemonchus contortus*
Herlich, H., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.), oxbendazole, efficacy against adult and larval stages
- Haemonchus contortus*
Herlich, H., 1977, *Am. J. Vet. Research*, v. 38 (8), 1247-1248
efficacy of albendazole against gastrointestinal nematodes and *Fasciola hepatica* in cattle (exper.); comparison of critical vs. controlled tests
- Haemonchus contortus*
Hogarth-Scott, R. S.; et al., 1976, *Research Vet. Sc.*, v. 21 (2), 232-237
Haemonchus contortus, *Trichostrongylus colubriformis*, thiabendazole-resistant strains, efficacy of fenbendazole, sheep, implications for management of helminth disease in grazing animals with respect to benzimidazole-resistance
- Haemonchus contortus*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, *J. South African Vet. Ass.*, v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Haemonchus contortus*
Horak, I. G.; Snijders, A. J.; and Louw, J. P., 1972, *J. South African Vet. Ass.*, v. 43 (4), 397-403
trematodes and nematodes, sheep (exper.), rafoxanide, efficacy studies
- Haemonchus contortus*
Hsu, C. K.; and Levine, N. D., 1977, *Am. J. Vet. Research*, v. 38 (8), 1115-1119
Haemonchus contortus, *Trichostrongylus colubriformis*, development of infective larvae under cyclic vs. constant conditions of temperature and humidity, degree-day concept appears to be applicable
- Haemonchus contortus*
Hubert, J.; Yvone, P.; and Kerboeuf, D., 1976, *Ann. Recherches Vet.*, v. 7 (1), 83-90
parasite survival in liquid manure, anti-parasitic action of xylene
- Haemonchus contortus*, *illus.*
Hutchinson, G. W.; and Slocombe, J. O. D., 1976, *J. Helminth.*, v. 50 (3), 143-152
Haemonchus contortus in rabbits (exper.), parasitological findings, developmental morphology, concluded that laboratory rabbit could be useful experimental host
- Haemonchus contortus* (Rudolphi, 1803) Cobb, 1898
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus (rennet): southern Bulgaria
- Haemonchus contortus*
Kelly, J. D.; et al., 1975, *Research Vet. Sc.*, v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Haemonchus contortus*
Kelly, J. D.; et al., 1977, *Research Vet. Sc.*, v. 22 (2), 161-168
Haemonchus contortus, *Trichostrongylus colubriformis*, resistant or susceptible to thiabendazole, sheep, effect of route of administration (oral, intra-ruminal or intra-abomasal) on efficacy of benzimidazole anthelmintics
- Haemonchus contortus*
Kennedy, T. J.; and Todd, A. C., 1975, *Am. J. Vet. Research*, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Haemonchus contortus*
Kerboeuf, D., 1977, *Ann. Recherches Vet.*, v. 8 (3), 257-266
Haemonchus contortus, lambs, changes in serum pepsinogen, protein and lipid levels
- Haemonchus contortus*
Kerboeuf, D.; and Leimbacher, F., 1977, *Rec. Med. Vet.*, v. 153 (1), 19-25
Ostertagia circumcincta, *Trichostrongylus axei*, *Haemonchus contortus*, lambs, use of serum pepsinogen measurements to assess average worm burden in a herd
- Haemonchus contortus*
Knight, R. A.; and Rodgers, D., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 116
Haemonchus contortus, lambs, age resistance to single inoculation
- Haemonchus contortus*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, *Am. J. Vet. Research*, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska

- Haemonchus contortus**
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Haemonchus contortus**
Kutzer, E.; and Frey, H., 1976, Berl. u. Munchen. Tierarztl. Wchnschr., v. 89 (24), 480-483
Lepus europaeus: Austria
- Haemonchus contortus**
Kutzer, E.; and Frey, H., 1976, Ztschr. Parasitenk., v. 50 (2), 213-214
Lepus europaeus
- Haemonchus contortus**
Le Jambre, L. F., 1976, Vet. Parasitol., v. 2 (4), Dec., 385-391
Haemonchus contortus, *Ostertagia circumcincta*, technique for assay of thiabendazole resistance by hatching eggs in solutions of thiabendazole
- Haemonchus contortus**
Le Jambre, L. F.; and Royal, W. M., 1976, Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (fecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except *Nematodirus* resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales
- Haemonchus contortus**
Le Jambre, L. F.; and Royal, W. M., 1977, Internat. J. Parasitol., v. 7 (6), 481-487
Haemonchus contortus from New South Wales, genetics of vulvar morph types, dominance hierarchy of these characters found to be smooth > knobbed > linguiform, linguiform phenotype most common in wild type population
- Haemonchus contortus**
Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1976, Internat. J. Parasitol., v. 6 (3), 217-222
Haemonchus contortus, dose mortality response of strains selected by thiabendazole or by thiabendazole plus morantel tartrate to thiabendazole, morantel tartrate, or levamisole, results indicate that resistance to thiabendazole is due to a single gene and resistance to morantel tartrate is polygenic in nature
- Haemonchus contortus**
Levine, N. D.; et al., 1975, Am. J. Vet. Research, v. 36 (10), 1459-1464
lambs grazing with their ewes under 2 pasture rotation systems, lambs under rotation had more nematodes and gained less weight than nonrotated control lambs, rotation is not recommended to control nematode parasitism of sheep in Illinois
- Haemonchus contortus**
Levine, N. D.; Mansfield, M. E.; and Todd, K. S., jr., 1977, J. Parasitol., v. 63 (5), 954-956
Haemonchus contortus, effects of photoperiod during development and of storage on maturation of larvae in lambs
- Haemonchus contortus**
McCallister, G. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 89-90
Haemonchus contortus, free-living stages, haloxon, thiabendazole; haloxon ineffective against all free-living stages and exsheathed third-stage larvae, thiabendazole had noticeable effect only on unembryonated egg
- Haemonchus contortus**
McKenna, P. B., 1973, Research Vet. Sc., v. 14 (3), 312-316
Haemonchus contortus, two morphologically and geographically distinct strains, effect of storage at 5 or 21 C for varying intervals on infectivity and parasitic development of third-stage larvae in sheep, evidence that small proportion of infective larvae may be innately 'inhibition-prone' and that effect of season on host or ageing of infective larvae or both may be contributory to changes in level of inhibition
- Haemonchus contortus**
McKenna, P. B., 1976, N. Zealand J. Exper. Agric., v. 4 (2), 235-237
post-mortem recovery of *Haemonchus contortus*, *Ostertagia* spp., *Trichostrongylus axei*, sheep, peptic digestion of ovine abomasum unlikely to be of diagnostic value for field-submitted specimens
- Haemonchus contortus**
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1974, Indian J. Animal Research, v. 8 (2), 75-78
Haemonchus contortus, other nematodes, experimentally or naturally infected sheep, nitroxylin highly effective, critical testing; in vitro testing against *H. contortus*
- Haemonchus contortus**
Mansfield, M. E.; Todd, K. S., jr.; and Levine, N. D., 1977, Am. J. Vet. Research, v. 38 (6), 803-806
Haemonchus contortus larvae, effects of various temperatures and storage conditions on developmental arrest in lambs (exper.)
- Haemonchus contortus**
Mapes, C. J.; and Gallie, G. J., 1977, Parasitology, v. 74 (3), 235-242
Haemonchus contortus, development and growth in stomach of laboratory rabbits (exper.), effect of exsheathment on infectivity, effect of dose size, no adult worms recovered and no eggs found in feces
- Haemonchus contortus**
Misra, S. C., 1972, Indian J. Animal Research, v. 6 (2), 95-96
parasitic gastro-enteritis, goats, epidemiology, seasonal incidence: Orissa

- Haemonchus contortus*, *illus.*
Munn, E. A., 1977, *Tissue and Cell*, v. 9 (1), 23-34
Haemonchus contortus, structure of intestinal cells, helical polymeric extracellular protein associated with luminal surface for which name contortin is proposed, *Ostertagia circumcincta* also contained contortin-like material but *Nippostrongylus brasiliensis* and *Syphacia obvelata* contained material associated with outer surface of microvilli which was quite distinct from contortin
- Haemonchus contortus*
Mutafova, T., 1972, *Izvest. Tsentral. Khelminth. Lab.*, v. 15, 143-150
Haemonchus contortus, *Ostertagia* sp., *Trichostrongylus* sp., sheep, seasonal changes in egg-production, spring rise observed irrespective of age or sex of host
- Haemonchus contortus*
Mwegoha, W. M.; and Jørgensen, R. J., 1977, *Acta Vet. Scand.*, v. 18 (3), 293-299
Haemonchus contortus, *Ostertagia ostertagi*, recovery of infective 3rd stage larvae by migration in agar gel, with and without addition of ox bile to agar gel, technique for recovery of larvae from field samples of pasture herbage
- Haemonchus contortus*
Niec, R.; et al., 1976, *Gac. Vet.*, Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary anthelmintic treatment that could prevent normal buildup of resistance
- Haemonchus contortus*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Haemonchus contortus*
Obergh, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Ovis aries: Chile
- Haemonchus contortus*
Okon, E. D.; and Enyenihi, U. K., 1975, *Bull. Office Internat. Epizoot.*, v. 83 (11-12), 1139-1144
Haemonchus contortus, *Trichostrongylus colubriformis*, lambs grazed on contaminated pastures, control by pasture rotation
- Haemonchus contortus*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Haemonchus contortus* Rudolphi
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Gazella thomsonii (gut)
G. granti (abomasum)
Alcelaphus buselaphus cokei (digestive tract)
Connochaetes taurinus (small intestine)
all from Kenya
- Haemonchus contortus*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Haemonchus contortus*
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Haemonchus contortus* (Rudolphi, 1803)
Pursglove, S. R., jr., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Oklahoma
- Haemonchus contortus*
Rahman, M. S.; et al., 1977, *N. Zealand Vet. J.*, v. 25 (4), 79-83
metabolic changes in *Moniezia expansa*, *Haemonchus contortus*, and *Fasciola hepatica* from mebendazole-treated sheep, total nucleotide concentrations, ATP levels, ATP/ADP ratios; detachment of *Fasciola hepatica* from host tissue diminished its contact with the drug
- Haemonchus contortus* Rudolphi, 1803
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 267-271
Sus scrofa: Huesca, Alto Aragon
- Haemonchus contortus*
Rogers, W. P.; and Brooks, F., 1976, *Internat. J. Parasitol.*, v. 6 (4), 315-319
Haemonchus contortus, suggested that exsheathing fluid contains a zinc metalloenzyme (probably leucine aminopeptidase) which is involved in process of exsheathment
- Haemonchus contortus*, *illus.*
Rogers, W. P.; and Brooks, F., 1977, *Internat. J. Parasitol.*, v. 7 (1), 61-65
Haemonchus contortus, egg hatching, presence of leucine aminopeptidase and lipase in hatching fluid, inhibition of hatching by 1,10-phenanthroline reversed by Zn^{2+}
- Haemonchus contortus*
Romanowski, R. D.; et al., 1975, *J. Parasitol.*, v. 61 (4), 777-778
Haemonchus contortus, cambendazole-resistant vs. -sensitive strains, effect on fumarate reductase of cambendazole, thiabendazole, and levamisole

- Haemonchus contortus*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Haemonchus contortus*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Haemonchus contortus*, *illus.*
Sachs, R.; Gibbons, L. M.; and Lweno, M. F., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 467-475
Ovis aries
Capra hircus
Bos indicus
Syncerus caffer
Redunca redunca
Kobus vardonii
K. defassa
Hippotragus equinus
Damaliscus korrigum
Alcelaphus buselaphus cokei
A. b. lichtensteini
Aepyceros melampus
Gazella granti
G. thomsonii
Giraffa camelopardalis
(abomasum of all): all from Tanzania
- Haemonchus contortus*
Schillhorn van Veen, T.; and Brinckman, W. L., 1975, Samaru Agric. Newsletter, v. 17 (2), 70-74
lambs, regular drenching with thiabendazole at regular intervals during rainy season, better weight gain, cost/benefit; possible influence of resistance and breed of sheep
- Haemonchus contortus* (Rudolphi, 1803) Cobbold, 1898
Sharma, L. D.; Bahga, H. S.; and Srivastava, P. S., 1971, Indian J. Animal Research, v. 5 (1), 33-38
Haemonchus contortus, sheep, goats, screening medicinal plant extracts
- Haemonchus contortus*
Sinclair, K. B.; and Prichard, R. K., 1975, Research Vet. Sc., v. 19 (2), 232-234
Haemonchus contortus, sheep, disophenol administered to sheep to prevent development of populations of adult worms and to study pathogenicity of arrested 4th-stage larvae which remained, concluded that they may cause damage to abomasal mucosa
- Haemonchus contortus*
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Haemonchus contortus*
Smith, W. D., 1977, Research Vet. Sc., v. 22 (1), 128-129
Haemonchus contortus, sheep immunized with larval antigens, stimulation of serum and mucus IgG antibody response, no IgA antibody response, no protection against challenge infection
- Haemonchus contortus*
Smith, W. D., 1977, Research Vet. Sc., v. 22 (3), 334-338
sheep hyperinfected with *Haemonchus contortus*, anti-larval antibody levels in serum and abomasal mucus, detected by radioimmunoassay, no immunological memory observed following challenge infection, presence of IgA antibodies in abomasal mucus thought to be locally produced while IgG antibodies largely derived from blood
- Haemonchus contortus*
Snijders, A. J.; and Horak, I. G., 1972, J. South African Vet. Ass., v. 43 (3), 295-297
Gedoelestia haessleri and *Haemonchus contortus* in naturally infected blesbuck, treatment with rafoxanide, highly effective
Damaliscus dorcas phillipsi: Republic of South Africa
- Haemonchus contortus*
Sommerville, R. I., 1976, J. Parasitol., v. 62 (2), 242-246
Haemonchus contortus, development and ecdysis in vitro, effects of changes in both ionic composition and osmotic pressure, potassium as necessary component of salt solution
- Haemonchus contortus*
Sommerville, R. I., 1977, J. Parasitol., v. 63 (2), 344-347
Haemonchus contortus, development in vitro, effect of rumen fluid and of a succession of media which incorporated changes in pH, pCO₂, and pO₂ likely to be encountered in transition from rumen to abomasum
- Haemonchus contortus*, *illus.*
Sommerville, R. I., 1977, J. Parasitol., v. 63 (5), 952-954
Haemonchus contortus, development to fourth stage in laboratory mice (gastric mucosa)
- Haemonchus contortus* (Rud., 1803), *illus.*
Sood, M. L.; and Kalra, S., 1977, Ztschr. Parasitenk., v. 51 (3), 265-273
Haemonchus contortus, histochemistry of body wall, comparison with plant parasitic nematode
- Haemonchus contortus*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Haemonchus contortus*
Stromberg, B. E.; and Soulsby, E. J. L., 1977, Vet. Parasitol., v. 3 (2), 169-175
Ascaris suum, guinea pigs, heterologous resistance induced by *Toxocara canis* and *Ancylostoma caninum* but not by *Haemonchus contortus*, *Caenorhabditis briggsae*, or *Turbatrix aceti*

- Haemonchus contortus**
Tager-Kagan, P., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (4), 317-321
gastro-intestinal nematodes, zebu cattle (1 to 2 years old), cambendazole: Niger
- Haemonchus contortus**
Tarazona, J. M., 1975, An. Inst. Nac. Invest. Agrar., s. Hig. y San. Animal (2), 11-17
trichostrongylidosis, ovine, seasonal incidence, 1971-1975: Spain
- Haemonchus contortus**
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcvi
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Haemonchus contortus**
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Haemonchus contortus**
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Haemonchus contortus**
Theodorides, V. J.; Nawalinski, T.; and Chang, J., 1976, Am. J. Vet. Research, v. 37 (12), 1515-1516
gastrointestinal nematodes, *Moniezia* spp., sheep, albendazole highly effective
- Haemonchus contortus**
Thomas, R. J.; Waller, P. J.; and Cottrill, B. R., 1975, Research Vet. Sc., v. 19 (1), 113-114
Haemonchus contortus larvae used as source of antigen, decrease in antigenic potency following storage for 2 months at 5°C, no such decline in larvae killed by freezing and stored at -15°C, suggested that loss of potency with ageing may be partly responsible for increased worm populations in sheep in spring
- Haemonchus contortus**
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antelope cervicapra (abomasum): Texas
- Haemonchus contortus**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Haemonchus contortus**
Todd, K. S., jr.; Levine, N. D.; and Boatman, P. A., 1976, Am. J. Vet. Research, v. 37 (8), 991-992
Haemonchus contortus, free-living stages, survival in sheep fecal pellets, various constant temperatures
- Haemonchus contortus**
Todd, K. S., jr.; Levine, N. D.; and Boatman, P. A., 1976, J. Parasitol., v. 62 (2), 247-249
Haemonchus contortus, survival of desiccated and undesiccated infective larvae at various constant temperatures
- Haemonchus contortus**
Tripathi, J. C., 1977, Indian J. Animal Sc., v. 47 (11), 739-742
Haemonchus contortus, effect of different temperatures on infective larvae in water and in faecal medium; desiccation of infective larvae in diffused light and sunlight compared
- Haemonchus contortus**
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Haemonchus contortus**
Varshney, T. R.; and Singh, Y. P., 1976, Indian Vet. J., v. 53 (9), 672-676
Haemonchus contortus, trichostrongyles, Trichuris, lambs, critical trial, efficacy and economic value, various anthelmintics: Central Sheep and Wool Research Station, Pashulok Rishikesh (Dehradun)
- Haemonchus contortus**
Varshney, T. R.; and Singh, Y. P., 1977, Indian J. Animal Sc., v. 46 (12), 666-668
Haemonchus contortus, sheep, resistance to phenothiazine and thiabendazole, critical and controlled conditions: Pashulok (Dehra Dun)
- Haemonchus contortus** (Rudolphi, 1803)
Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karroo
- Haemonchus contortus**
Virat, M.; and Gevrey, T., 1976, Ztschr. Parasitenk., v. 48 (3-4), 299 [Abstract]
Haemonchus contortus, infectious larvae, trapping activity of several species of predacious fungi, adhesive networks more effective than sticky knobs or constricting rings, *Arthrobotrys oligospora* and *Dactylaria thausasia* more effective species, invasion of worms
- Haemonchus contortus**
Virat, M.; and Pelouille, M., 1977, Ann. Recherches Vet., v. 8 (1), 51-58
predatory activity of fungus *Arthrobotrys oligospora* against larvae of **Haemonchus contortus**, optimal temperatures, larval density
- Haemonchus contortus**
Vlassoff, A., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand

- Haemonchus contortus**
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Haemonchus contortus, illus.**
Weise, R. W., 1977, J. Parasitol., v. 63 (5), 854-857
Haemonchus contortus, dorsal buccal lancet, light and electron microscopy, functional relation of morphologic findings to feeding mechanism
- Haemonchus contortus**
Wikerhauser, T.; et al., 1974, Acta Parasitol. Iugoslavica, v. 5 (2), 79-81
trichostrongylids, cattle, fenbendazole compared with thiabendazole, good results from both
- Haemonchus contortus**
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Haemonchus contortus**
Wilson, P. A. G., 1976, Ztschr. Parasitenk., v. 49 (3), 243-252
Haemonchus contortus, Nippostrongylus brasiliensis, infective larvae, carbohydrate content and ageing process contrasted; carbohydrate level variation in H. contortus due to capacity to synthesize glycogen during ageing, low level in N. brasiliensis remains constant
- Haemonchus contortus**
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 77-81
Damaliscus dorcas phillipsi (abomasum): Mountain Zebra National Park
- Haemonchus contortus**
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 195-198
Antidorcas marsupialis (abomasum): Mountain Zebra National Park near Cradock, Cape Province
- Haemonchus contortus**
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Haemonchus cont[ortus]**
Zielinski, J., 1972, Med. Wet., v. 28 (9), 566-567
parasites, sheep, Nilverm, copper sulfate
- Haemonchus contortus cayugensis**
Le Jambre, L. F., 1977, Internat. J. Parasitol., v. 7 (1), 9-14
Haemonchus contortus cayugensis females, genetics of vulvar morph types, dominance hierarchy found to be linguiform > knobbed > smooth
- Haemonchus contortus cayugensis**
Le Jambre, L. F.; and Ractliffe, L. H., 1976, Parasitology, v. 73 (2), 213-222
Haemonchus contortus cayugensis, lambs, infection with selected strain of smooth or of linguiform worms and subsequent grazing on same pasture, seasonal changes in phenotypes in relation to population density (affects frequencies of linguiform A vs. B but not of smooth vs. linguiform), "It appears therefore that the proportion of smooth to linguiform worms is a stable equilibrium maintained by natural selection."
- Haemonchus contortus cayugensis**
Le Jambre, L. F.; and Whitlock, J. H., 1976, Parasitology, v. 73 (2), 223-238
Haemonchus contortus cayugensis (New York State), Haemonchus contortus contortus (Ohio), vulvar phenotypes and hatch rate of eggs over a range of temperatures
- Haemonchus contortus cayugensis**
Whitlock, J. H.; and Georgi, J. R., 1976, Parasitology, v. 72 (3), 207-224
biological controls in mixed trichostrongylid infections (predominantly Haemonchus contortus cayugensis) in sheep, different ecosystems (barn vs. pasture) and different treatment groups, course of infections (erythrocyte loss, fecal egg counts, hematocrit values), "Anaphylactoid 'self-cure' did not occur in this experiment but something like premunition certainly did."
- Haemonchus contortus contortus**
Le Jambre, L. F.; and Whitlock, J. H., 1976, Parasitology, v. 73 (2), 223-238
Haemonchus contortus cayugensis (New York State), Haemonchus contortus contortus (Ohio), vulvar phenotypes and hatch rate of eggs over a range of temperatures
- Haemonchus dinniki n. sp., illus.**
Sachs, R.; Gibbons, L. M.; and Lweno, M. F., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 467-475
Aepyceros melampus
Gazella granti
G. thomsonii
Rhinchotragus kirkii
(abomasum of all): all from Serengeti National Park, Tanzania
- Haemonchus longistipes**
Lodha, K. R.; Raisinghani, P. M.; and Karwar, R. S., 1977, Indian J. Animal Sc., v. 47 (10), 677-682
helminths, camels, promintic and banminth II effective, nilverm inconsistent in action, thiabendazole ineffective
- Haemonchus mitchelli**
Gibbons, L. M.; and Khalil, L. F., 1976, Trop. Animal Health and Prod., v. 8 (3), 168
goat (gut): Kajiado district, Kenya

- Haemonchus mitchelli*, *illus.*
Sachs, R.; Gibbons, L. M.; and Lweno, M. F., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 467-475
Taurotragus oryx
Gazella granti
G. thomsonii
Rynchotragus kirkii
Giraffa camelopardalis
(abomasum of all): all from Tanzania
- Haemonchus placei*
Anderson, P. J. S.; and Marais, F. S., 1975, *J. South African Vet. Ass.*, v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Haemonchus placei*
Bryan, R. P., 1976, *Austral. Vet. J.*, v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Haemonchus placei*
Bryan, R. P.; Bainbridge, M. J.; and Kerr, J. D., 1976, *Austral. J. Zool.*, v. 24 (3), 417-421
Bubalus bubalis
cattle
(large and small intestine of all): all from Northern Territory, Australia
- Haemonchus placei*
Campbell, W. C.; and Thomson, B. M., 1973, *Austral. Vet. J.*, v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus colubriformis* proved uninfected even if they had not been frozen
- Haemonchus placei*
Ciordia, H.; et al., 1977, *Am. J. Vet. Research*, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Haemonchus placei*
Curr, C., 1977, *Austral. Vet. J.*, v. 53 (9), 425-428
nematodes, calves, levamisole, efficiency of pour-on formulation, drug trials, good results
- Haemonchus placei*
Dharsana, R. S.; Fabiyi, J. P.; and Hutchinson, G. W., 1976, *Vet. Parasitol.*, v. 2 (4), 333-340
mixed gastro-intestinal nematode infections, calves, effects on host intestinal enzymes
- Haemonchus placei*
Dorn, H.; and Federmann, M., 1976, *Vet.-Med. Nachr.* (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Haemonchus placei*
Fincher, G. T., 1975, *J. Parasitol.*, v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Haemonchus placei*
Kelly, J. D.; et al., 1975, *Research Vet. Sc.*, v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Haemonchus placei*
Lukovich, R.; et al., 1977, *Gac. Vet., Buenos Aires* (318), v. 39, 91-95
helminths, cattle, levamisole, results from injectable and dermal application similar
- Haemonchus placei*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Haemonchus placei*
Reinecke, R. K., 1972, *Onderstepoort J. Vet. Research*, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Haemonchus placei*
Ronald, N. C.; Bell, R. R.; and Craig, T. M., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (3), 317-319
gastrointestinal nematodes, calves, levamisole phosphate, effective at one-half recommended dosage
- Haemonchus placei*
Rothwell, T. L. W.; et al., 1976, *Vet. Parasitol.*, v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-ACHE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Haemonchus placei*
Rowlands, D. ap T.; and Berger, J., 1977, *J. South African Vet. Ass.*, v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration

- Haemonchus placei*
Schroeder, J.; Honer, M. R.; and Louw, J. P.,
1977, J. South African Vet. Ass., v. 48 (2),
95-97
trematodes, nematodes, cattle (exper.),
rafoxanide, efficacy of subcutaneous in-
jections against immature larvae and adults
- Haemonchus placei*
Smeal, M. G.; et al., 1977, Austral. Vet. J.,
v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal
distribution, poor relationship between
faecal egg counts and worm burdens: North
Coast and Tableland regions of New South
Wales
- Haemonchus placei*, *illus.*
Smith, K., 1975, J. Microscopy, v. 105 (2),
229-232
method for concentrating nematode larvae
from fecal material, procedure for processing
for transmission electron microscopy
- Haemonchus placei*
Snijders, A. J.; and Horak, I. G., 1975, J.
South African Vet. Ass., v. 46 (3), 265-267
cattle (exper.), rafoxanide, drug trials,
efficacy, good results
- Haemonchus placei*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26),
216-230
fenbendazole (available in 5 forms), ef-
ficacy against nematodes in various animals,
well tolerated with no apparent effects on
fertility or fetus, extensive summary of re-
sults to date
- Haemonchus placei*
Williams, J. C.; Sheehan, D.; and Fuselier,
R. H., 1977, Am. J. Vet. Research, v. 38
(12), 2037-2038
gastrointestinal nematodes, tapeworms,
cattle, efficacy of albendazole (oral drench)
- Haemonchus similis*, *illus.*
Sachs, R.; Gibbons, L. M.; and Lweno, M. F.,
1973, Ztschr. Tropenmed. u. Parasitol., v. 24
(4), 467-475
Bos indicus (abomasum): Tanzania
- Haemonchus vegliai*, *illus.*
Sachs, R.; Gibbons, L. M.; and Lweno, M. F.,
1973, Ztschr. Tropenmed. u. Parasitol., v. 24
(4), 467-475
Taurotragus oryx (abomasum): Tanzania
- Halocercus* Baylis and Daubney 1925
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
key; key to species in *Phocoena phocoena*
- Halocercus* sp.
Smith, F. R.; and Threlfall, W., 1973, Am.
Midland Naturalist, v. 90 (1), 215-218
Phocoena phocoena: insular Newfoundland
and its adjacent waters
- Halocercus delphini* Baylis & Daubney, 1925,
illus.
Dailey, M. D.; and Perrin, W. F., 1973, Fish.
Bull., National Oceanic and Atmos. Admin.,
v. 71 (2), 455-471
Stenella graffmani
S. cf. *S. longirostris*
(bronchiole region of all): all from
eastern tropical Pacific
- Halocercus gymnurus* Railliet
Bonner, W. N., 1972, Oceanogr. and Marine Biol.
Ann. Rev., v. 10, 461-507
Phoca vitulina (lung): European waters
- Halocercus inflexocaudatus* (von Siebold) Baylis
and Daubney 1925
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Halocercus invaginatus* (Quekett
1841) Dougherty 1943
- Halocercus inflexocaudatus* (of Schmidt-Ries
1939)
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Halocercus taurica* Delyamure 1942
- Halocercus inflexocaudatus* (von Siebold, 1842)
Smith, F. R.; and Threlfall, W., 1973, Am.
Midland Naturalist, v. 90 (1), 215-218
Phocoena phocoena: insular Newfoundland
and its adjacent waters
- Halocercus invaginatus* (Quekett 1841) Dougherty
1943, *illus.*
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
synonymy; redescription; key
Phocoena phocoena: Bay of Fundy, Canada;
Vancouver Is., British Columbia, Canada
- Halocercus ponticus* Delyamure 1946
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Halocercus invaginatus* (Quekett
1841) Dougherty 1943
- Halocercus taurica* Delyamure 1942, *illus.*
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
redescription; key
Syn.: *Halocercus inflexocaudatus* (of
Schmidt-Ries 1939)
Phocoena phocoena: Bay of Fundy, Canada;
Vancouver Is., British Columbia, Canada
- Hamannia* Railliet, Henry & Sisoff, 1912
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Echinuria* Soloviev, 1912
- Hamatospiculum cylindricum* (Zeder, 1803)
Buscher, H. N.; and Tyler, J. D., 1975, Proc.
Oklahoma Acad. Sc., v. 55, 108-111
Speotyto cunicularia: Oklahoma

- Hammerschmidtella diesingi*
Hominick, W. M., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (5), 383 [Demonstration]
Thelastoma sp., bacterial infection of cuticle of pinworms inhabiting hindgut of laboratory reared *Periplaneta americana*, bacterial preference for *Thelastoma* sp. over *Hammerschmidtella diesingi* possibly related to structure of cuticle
- Hammerschmidtella diesingi*
Leong, L.; and Paran, T. P., 1966, Med. J. Malaya, v. 20 (4), 349
Periplaneta americana
Blatta orientalis
all from Singapore
- Hamulofilaria*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Oxyspirura*
key
Syn.: *Skrjabinispirura* subgen. of *Oxyspirura*
- Haplonema Ward and Magath, 1917*
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
synonymy; diagnosis; revision of genus
- Haplonema aditum* Mueller, 1934
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
"transferred to *Paraquimperia* by Moravec (1966)"
- Haplonema hamulatum* Moulton, 1931, illus.
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
synonymy; redescription
Lota lota: Aishihik Lake, Yukon Territory, Canada
- Haplonema immutatum* Ward and Magath, 1917, illus.
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
redescription
Amia calva: New Orleans area, Louisiana
- Haplonema orthocephalum* Dogiel and Akhmerov, 1959
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
species inquirenda
- Haplonema sinensis* (Hsu, 1933) Chitwood and Wehr, 1934
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
"The generic independence of *Pingus* is no longer questioned and the species is well known under its original name, *Pingus sinensis*."
- Haplonema tenerrima* (von Linstow, 1878) Mueller, 1934
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
"von Linstow's species has been reestablished in the genus *Paraquimperia* (see Moravec 1966)"
- Hartertia* Seurat, 1914, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Hartertiidae
key
- Hartertiidae* (Quentin, 1970, subfam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruroidea
key; key to genera
includes: *Alainchabaudia*; *Hartertia*
- Hartwichia* Chabaud & Bain, 1966
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Dujardinascariidea
key
- Hassalstrongylus Durette-Desset, 1971*
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, *Nippostrongylineae*
- Hassalstrongylus aduncus*
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (small intestine): Florida
- Hassalstrongylus chabaudi* n. sp., illus.
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Wiedomys pyrhorinus (duodenum): Exu, Pernambuco, Bresil
- Hassalstrongylus echalieri* n. sp., illus.
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Oryzomys sp. (duodenum et intestin): Guyane française
- Heartworm. See [*Dirofilaria immitis*]
- Hedruridae* (Railliet, 1916, subfam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematoidea
key
includes: *Hedruris*
- Hedruris* Nitzsch, 1812, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Hedruridae

- Hedruris spinigera* Baylis
Griffiths, W. E., 1976, N. Zealand J. Marine and Freshwater Research, v. 10 (3), 533-536
Hedruris spinigera in *Perca fluviatilis* (stomach), high rate of infestation when perch fed on *Retropinna retropinna*, rapid decline when this food source no longer available
Perca fluviatilis (stomach): Selwyn River, New Zealand
Paracalliope fluviatilis
- Heliconema* Travassos, 1919, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Proleptinae
key
Syn.: *Ortleppina* Schulz, 1927
- Heliconema* Travassos, 1919
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Physalopteridae, Physalopterinae
key
- Heliconema urolophi* (Johnston and Mawson, 1951) comb. nov.
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Syn.: *Proleptus urolophi* Johnston and Mawson, 1951
- Heligmobaylisia* Mawson, 1961
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Paraheligmomina* (Ortlepp, 1939)
- Heligmodendrium* Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Heligmostrongylus* Travassos, 1917
- Heligmodendrium hassalli*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Heligmonella* Moennig, 1927 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Heligmonellinae
- Heligmonella* sp. Wertheim et Nevo, 1971
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
as syn. of *Heligmonina nevoi* n. sp.
- Heligmonella dremomys* (Yen, 1973) n. comb.
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
Syn.: *Impalaia dremomys* Yen, 1973
- Heligmonella dremomysi* Durette-Desset, 1974, homonym of *H. dremomys* (Yen, 1973)
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
renamed *Heligmonella moreli* nom. nov.
- Heligmonella limbooliati* n. sp., illus.
Durette-Desset, M.-C.; Diaw, O.; and Krishna-samy, M., 1975, Ann. Parasitol., v. 50 (4), 477-491
Trichys lipura (intestin): Malaisie
- Heligmonella moreli* nom. nov.
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
for *Heligmonella dremomysi* Durette-Desset, 1974, homonym of *Heligmonella dremomys* (Yen, 1973) n. comb.
- Heligmonellidae (Skrjabin et Schikhobalova, 1952, tribu), nov. fam.
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
includes: Heligmonellinae; Pudicinae; Brevistriatinae; Nippostrongylineae
- Heligmonellinae (Skrjabin et Schikhobalova, 1952, tribu) Durette-Desset, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae
includes: *Heligmonella* (type genus); *Paraheligmonella*; *Tricholinstowia*
- Heligmonina* Baylis, 1928
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Nippostrongylineae
synonymy
- Heligmonina nevoi* n. sp., illus.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Syn.: *Heligmonella* sp. Wertheim et Nevo, 1971
Spalax ehrenbergi (intestin grele): Maor, pres de Tiberias, Ein Zetim, Ma'alot, Sasa, Ramat Ishai, Shefar'am, El-Al, Majdal shams, Quneitra, Gabri, Elroi, Yogune'am, Tiberias, Anza, Jenin, Schechem (Nablus), Nashonim, Atarat, Ramallah, Bet-El, and Lahav, Israel
- Heligmonoides* Baylis, 1928
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Nippostrongylineae
- Heligmoskrjabinia* Freitas et Lent, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Viannaiinae
- Heligmosomidae (Travassos, 1914, sub. fam.), Cram, 1927
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
includes: *Ornithostrongylineae*; *Viannaiinae*; *Heligmosominae*

- Heligmosominae Travassos, 1914
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae
includes: Heligmosomum (type genus); Citelinema; Citellinoides; Heligmosomoides; Longistriata; Ohbayashinema; Suncinema
- Heligmosomoides Hall, 1916
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
synonymy
- Heligmosomoides glareoli Baylis, 1928, illus.
Mishra, G. S.; Durette-Desset, M. C.; and Bercovier, H., 1976, Ann. Parasitol., v. 51 (1), 157-160
description, differentiation from H. taticus
Clethrionomys glareolus: Villegouin (Indre)
- Heligmosomoides glareoli Baylis, 1928
Wiger, R.; Lien, L.; and Tenora, F., 1976, Norwegian J. Zool., v. 24 (2), 133-135
Clethrionomys glareolus (small intestine): Kviteseid, Norway
- Heligmosomoides polygyrus (Nematospiroides dubius)
Brown, A. R.; and Crandall, C. A., 1976, J. Immunol., v. 116 (4), 1105-1109
mice, Ascaris suum-induced phosphorylcholine-binding component identified as IgM antibody having idiotypic determinants in common with PC-binding IgA myeloma TEPC 15, response not duplicated by immunization with dead Ascaris larvae or by infection with Heligmosomoides polygyrus or Trichinella spiralis
- Heligmosomoides polygyrus
Brown, A. R.; Crandall, R. B.; and Crandall, C. A., 1976, J. Parasitol., v. 62 (1), 169-171
Heligmosomoides polygyrus-infected mice, increased IgG catabolism as possible factor in observed suppression of circulating antibody levels following immunization to sheep erythrocytes
- Heligmosomoides polygyrus
Cypess, R. H.; et al., 1977, Exper. Parasitol., v. 42 (1), 34-43
Heligmosomoides polygyrus, temporal, spatial, and morphological population characteristics in LAF₁ vs. A/He mice examined in order to determine possible mechanisms responsible for differences in expression of resistance between these two mouse strains
- Heligmosomoides polygyrus
Cypess, R. H.; Ebersole, J. L.; and Molinari, J. A., 1977, Internat. Arch. Allergy and Applied Immunol., v. 55 (1-6), 496-503
Heligmosomoides polygyrus-infected mice, intestinal perfusates, radial immunodiffusion analysis, alteration in amount and class of immunoglobulins as well as anti-parasitic antibody
- Heligmosomoides polygyrus (= Nematospiroides dubius)
Cypess, R. H.; and Zidian, J. L., 1975, J. Parasitol., v. 61 (5), 819-824
Heligmosomoides polygyrus, development of self-cure and/or protection, influence of host genetic background (several inbred and outbred mouse strains) and various experimental conditions (route, dose, larval preparation)
- Heligmosomoides polygyrus (= Nematospiroides dubius)
Forrester, D. J., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 171-174
Heligmosomoides polygyrus, blood changes in irradiated experimentally infected mice
- Heligmosomoides polygyrus (Dujardin, 1845)
Wiger, R.; Lien, L.; and Tenora, F., 1976, Norwegian J. Zool., v. 24 (2), 133-135
Apodemus sylvaticus (small intestine): Oslo, Norway
- Heligmosomoides polygyrus polygyrus (Dujardin, 1845)
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Apodemus mystacinus: Mt. Miron, Carmel, caves, Mas'ada, Ein Hemed, Jerusalem, Saffat, Israel
A. sylvaticus: Dan, Sasa, Israel
- Heligmosomoides taticus (Erhardova, 1955)
Tenora, F.; Pfaller, K.; and Murai, E., 1971, Parasitol. Hungar., v. 4, 157-167
Microtus nivalis (Dunndarm): Obergurgl; Kuhtai; Schwarzee (Tiroler Zentralalpen)
- Heligmosomoides travassosi Schulz, 1926, illus.
Meszaros, F., 1977, Acta Zool. Acad. Scient. Hungar., v. 23 (1-2), 133-138
description
Cricetus cricetus (small intestine): Hungary
- Heligmosomum Railliet et Henry, 1909 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
- Heligmosomum borealis
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela lutreola
all from Karelia
- Heligmosomum costellatum Railliet et Henry, 1909
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Clethrionomys glareolus (small and large intestine): Karelia
- Heligmosomum costellatum Dujardin, 1845
Tenora, F.; and Meszaros, F., 1972, Parasitol. Hungar., v. 5, 159-161
Pitymys duodecimcostatus (small intestine): Burguete, western Pyrenees, Spain

- Heligmosomum costellatum* (Dujardin, 1845)
Tenora, F.; Pfaller, K.; and Murai, E., 1971, Parasitol. Hungar., v. 4, 157-167
Microtus nivalis (Dunndarm): Obergurgl; Kuhtai; Timmelsjoch (Tiroler Zentralalpen)
- Heligmosomum dubinini* Gvozdev, 1966
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 415-418
as syn. of *Ohbayashinema dubinini* (Gvozdev, 1966) n. comb.
- Heligmosomum glareoli* (Baylis, 1928) Skrjabin et Schulz, 1952
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Clethrionomys glareolus
Clethrionomys sp.
(small intestine of all): all from Karelia
- Heligmosomum halli* Schulz, 1926
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Arvicola terrestris (small intestine): Karelia
- Heligmosomum mixtum*
Merkusheva, I. V., 1975, Vestsi Akad. Navuk BSSR, s. Biial. Navuk (6), 82-86
helminths of rodents as model for quantitative indices in analysis of faunistic and ecological studies
- Heligmosomum polygyrum* (Dujardin, 1845) Railliet et Henry, 1909
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Arvicola terrestris
Ondatra zibethica
(intestine of all): all from Karelia
- Heligmospiroides* Ortlepp, 1939
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Heligmonina* Baylis, 1928
- Heligmostrongylus Travassos*, 1917
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, *Pudicinae*
synonymy
- Heligmostrongylus echimyos* n. sp., *illus.*
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Echimys armatus (intestin): Guyane française
- Heligmus* Dujardin, 1845
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
"excluded from the key since . . . incompletely described"
- Hempelia* Vaz, 1937
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelaziinae
key
- Hepaticola* sp. Yamaguti, 1935
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola* sp.
Kakacheva-Avramova, D., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 89-107
Barbus tauricus cyclolepis (liver): River Tundzha
- Hepaticola* sp., *illus.*
Reichenbach-Klinke, H. H., 1975, Fisch u. Umwelt (1), 113-121,
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Hepaticola cholidicola* (Soltys, 1952)
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola fagei* (Arvy, 1951)
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola hepatica* (Bancroft, 1893) Hall, 1916
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Arvicola terrestris (liver): Karelia
- Hepaticola hepaticola* (Bancroft, 1893) Hall, 1916
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola hepatophila* (Babos, 1954) comb. nov.
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola petruschewskii* Schulman, 1948
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola soricicola* Yokogawa in Nischigori, 1924
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepaticola tritonis-cristati* (Diesing, 1861)
Ivashkin, V. M.; and Shmytova, G. Ia., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 64-65
- Hepatinema* Rasheed, 1964, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochoniidae
key
- Hepatojarakus* Yeh, 1955
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Molineinae*

- Hepatojarakus bandicoti Sood et Parshad, 1973, illus.
Parshad, V. R.; and Sood, M. L., 1976, Ztschr. Parasitenk., v. 49 (2), 139-144
Hepatojarakus bandicoti, prevalence in rodents, pathological changes of liver and biliary system
Banicota bengalensis
Tatera indica
Millardia meltada
Golunda elliotti
Mus musculus bactrianus
M. booduga
- Hepatojarakus bandicoti Sood et Parshad, 1973
Sood, M. L.; and Parshad, V. R., 1975, Riv. Parassitol., Roma, v. 36 (2-3), 189-196
infections in Millardia meltada, survey of seasonal distribution, possible correlations between host diet and sex and incidence of infection
- Hepatojarakus malayae Yeh, 1955, illus.
Durette-Desset, M.-C.; and Chabaud, A.-G., 1975, Ann. Parasitol., v. 50 (2), 173-185
redescription
Tupaia tana: Ranan. Kg Nalapak, Sabah, Borneo
Aeromys thomasi (intestin grele): Kinabalu, Borneo
- Hepatojarakus malayae (Yeh, 1955)
Singh, M.; and Cheong Chee Hock, 1971, South-east Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus rattus argentiventer
R. r. jarak
R. bowersi
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
R. whiteheadi
all from Malaysia
- Heptochona Rasheed, 1965, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochoniidae
key
Syn.: Pontochona Mamaev, 1968
- Heptochona yamagutii n. sp., illus.
Kalyankar, S. D., 1972, Riv. Parassitol., Roma, v. 33 (4), 281-288
Stromateus niger (body cavity): Bombay, Maharashtra, India
- Herpestostrongylus Khera, 1956
Deshmukh, P. G., 1973, Riv. Parassitol., Roma, v. 34 (1), 63-65
generic diagnosis emended
- Herpestostrongylus herpestis Khera, 1956, illus.
Deshmukh, P. G., 1973, Riv. Parassitol., Roma, v. 34 (1), 63-65
description
Herpestes edwardsii (lung): Aurangabad (Maharashtra) India
- Herpestostrongylus Baylis, 1931
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Heterakiasis
Kovalenko, I. I.; and Sakachina, V. I., 1976, Veterinariia, Kiev (43), 86-88
ascariasis and heterakiasis, chickens, mixed infection; furidine, furidine + phenothiazine, and piperazine + phenothiazine, good results; phenothiazine or furidine alone, poor results
- Heterakis
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Heterakis
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Heterakis sp.
Kemp, R. L.; and Franson, J. C., 1975, Avian Dis., v. 19 (4), 741-744
Histomonas meleagridis, experimental transmission to chickens and turkeys by earthworms recovered from pheasant yard soil; earthworms may serve as transport hosts for Heterakis eggs and thus ultimately for H. meleagridis
- Heterakis beramporia Lane, 1914
Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Heterakis brevispiculum Gendre, 1911
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Heterakis brevispiculum Gendre, 1911
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagridis galeata (caeca): Vom area, Benue Plateau State, Nigeria
- Heterakis caudebrevis
Mack, J. K.; and Birova, V., 1976, Biologia, Bratislava, s. B, Zool. (1), v. 31 (2), 91-103
as syn. of Heterakis gallinarum

- Heterakis gallinae*
Alcaino, H.; and Gorman, T., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 157-159
Gallus gallus domesticus (ciego): Santiago, Chile
- Heterakis gallinae*
Gogoi, A. R., 1975, Kerala J. Vet. Sc., v. 5 (2), 131-134
fowl: Assam
- Heterakis gallinae*
Graber, M.; and Euzeby, J., 1976, Ann. Parasitol., v. 51 (2), 199-205
Anas boschas: Guadeloupe
- Heterakis gallinae*, illus.
Mayaudon T., H.; and Bendjaya Cases, J., 1974, Rev. Med. Vet. y Parasitol., Maracay, v. 25 (1-8), 1973-1974, 32-66
prevalence in Gallus (Gallus) domesticus: Estado Aragua, Venezuela
- Heterakis gallinae*
Nol, M.; et al., 1975, Rev. Avicult., v. 19 (3), 351-352
Ascaridia galli, Heterakis gallinae, poultry, anemia, changes in leucocyte types, elevated sedimentation rate, decrease in albumin, increase in alpha and beta globulin
- Heterakis gallinae*
Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix
Tetrao urogallus
all from CSSR
- Heterakis gallinae*
Torres, P.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 115-117
Gallus gallus domesticus: Chile
- Heterakis gallinarum* (Schrank, 1788) Madsen, 1949
Alavi, K. A.; and Ansari, J. A., 1973, Indian J. Zool., v. 1 (1), 13-16
Heterakis gallinarum, fowl (intestinal contents, caeca), incidence and seasonal variation: Aligarh and its suburbs
- Heterakis gallinarum* (Gmelin, 1790) Freeborn, 1923
Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Heterakis gallinarum*
Birova, V.; and Macko, J. K., 1976, Biologia, Bratislava, s. B, Zool. (3), v. 31 (8), 583-593
Heterakis gallinarum, short-, shortened-, and normal-tailed forms, survival and viability of descendants in three consecutive generations
- Heterakis gallinarum*
Cervenka, J.; Zajicek, D.; and Nydl, J., 1975, Veterinarstvi, v. 25 (6), 263-264
helminths, geese, Mebendazole
- Heterakis gallinarum*
Chute, A. M.; and Lund, E. E., 1972, J. Protozool., v. 19 (4), 639-643
Histomonas meleagridis in Numida meleagris (exper.), severe infections following inoculation by rectal route, much less severe response to natural route of infection (ingestion of Heterakis gallinarum eggs or earthworms carrying H. gallinarum and Histomonas meleagridis)
- Heterakis gallinarum*
Colglazier, M. L., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 60-61
Ascaridia dissimilis, Heterakis gallinarum, and Capillaria obsignata, turkeys, levamisole administered in drinking water effective as anthelmintic
- Heterakis gallinarum*
Cremers, H. J. W. M.; Jansen, J.; and Swierstra, D., 1975, Tijdschr. Diergeneesk., v. 100 (22), 1209-1211
Numida meleagris: Netherlands
- Heterakis gallinarum*
Cruthers, L. R.; al-Khateeb, G. H.; and Hansen, M. F., 1975, Proc. Oklahoma Acad. Sc., v. 55, 119-121
Ascaridia galli, Heterakis gallinarum, Capillaria obsignata, chickens, levamisole in drinking water
- Heterakis gallinarum* (Shrank, 1788) Madsen, 1949
Fine, P.E.M., 1975, J. Helminth., v. 49 (4), 229-243
Heterakis gallinarum in chickens with concomitant Parahistomonas wenrichi infections, worm fertility (egg production), worm mortality, inter-caecal distribution and migration tendencies of worms with reference to probability of pairing of males and females
- Heterakis gallinarum*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (ceca): Florida
- Heterakis gallinarum*
Kovalenko, I. I., 1975, Veterinariia, Kiev (42), 92-94
Ascaridia galli, Heterakis gallinarum, length of development studied as basis for timing of anthelmintic treatment
- Heterakis gallinarum*
Lesin'sh, K. P.; et al., 1975, Latvijas PSR Zinat. Akad. Vestis (340) (11), 27-30
helminths, chickens, effect of host age and method of rearing on infestation: Latvian SSR

- Heterakis gallinarum*
Long, P. L., 1971, Symposia Brit. Soc. Parasitol., v. 9, 65-75
maintenance of *Eimeria* and *Histomonas* in vivo, extensive review: maintenance of *Eimeria* (purification of species, exper. infections in chickens, factors affecting susceptibility of chickens and chicken embryos to infection; freeze-preservation); maintenance of *Histomonas meleagridis* (infection with ground-up infected tissues; use of in vitro cultures to obtain in vivo infections; use of embryonated *Heterakis gallinarum* ova to induce histomoniasis in vivo)
- Heterakis gallinarum*
Lund, E. E.; and Chute, A. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 73-76
larval *Heterakis gallinarum*, reciprocal transfer between *Phasianus colchicus* and *Coturnix coturnix japonica*, influence of early development in incompatible host on subsequent development; effect on liberation of *Histomonas meleagridis* and *Parahistomonas wenrichi* and transmission of *H. meleagridis*
- Heterakis gallinarum*
Lund, E. E.; and Chute, A. M., 1975, J. Protozool., v. 22 (3), 27A-28A [Abstract]
Parahistomonas wenrichi, factors in transmission by *Heterakis gallinarum*
- Heterakis gallinarum* (Schrank, 1788), *illus.*
Mack, J. K.; and Birova, V., 1976, Biologia, Bratislava, s. B, Zool. (1), v. 31 (2), 91-103
Syn.: *H. caudebrevis* Popova, 1949
Heterakis gallinarum, variations in tail length, numbers of caudal papillae, positions of caudal papillae, configuration of alae; various phenotypes, significance in helminthological systematics
- Heterakis gallinarum*
Mirzayans, A., 1975, J. Vet. Fac. Univ. Tehran, v. 30 (4), 5
chickens: area of Tehran, Iran
- Heterakis gallinarum*
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Heterakis gallinarum* (Schrank, 1788)
Pence, D. B.; and Bickel, S., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 104-105
Meleagris gallopavo intermedia: near Paint Rock, Concho County, Texas
- Heterakis gallinarum*
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Heterakis gallinarum*
Rudek, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 225-235
Heterakis gallinarum, chickens, various size doses, clinical course of disease, host weight gains, haemograph, pathomorphological changes, possible economic loss
- Heterakis gallinarum*
Stoimenov, K., 1975, Vet. Med. Nauki, v. 12 (1), 45-53
Heterakis gallinarum, chickens (exper.), recuperation processes after treatment with phenothiazine, early treatment most beneficial
- Heterakis gallinarum*
Stoimenov, K., 1976, Vet. Med. Nauki, v. 13 (7), 48-54
Heterakis gallinarum, chickens, parasite survival rate and pathogenicity increased by lower dose of infection and in 3-4 month old hosts; effects of blood factors; more female worms in older birds
- Heterakis gallinarum*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Heterakis gallinarum, illus.*
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Heterakis gallinarum* (Schrank 1788) Madsen 1949, *illus.*
Wright, K. A., 1977, J. Parasitol., v. 63 (3), 528-539
Heterakis gallinarum, labial sense organs, scanning and transmission electron microscopy
- Heterakis gallinarum, illus.*
Wright, K. A.; and Hui, N., 1976, J. Parasitol., v. 62 (4), 579-584
Heterakis gallinarum, post-labial sense organs, structure, probable function as chemo- and mechano-receptors
- Heterakis gallinarum* (Schrank, 1788)
Young, P. L.; and Babero, B. B., 1975, Proc. Oklahoma Acad. Sc., v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Heterakis hispidum*
Owen, D., 1976, Lab. Animals, v. 10 (3), 271-278
Rattus norvegicus: Carshalton

- Heterakis isolonche*, *illus.*
Griner, L. A.; et al., 1977, *Vet. Path.*, v. 14 (6), 582-590
Heterakis isolonche in gallinaceous birds, cecal nodular granulomas, pathology, case reports
Crossoptilon mantchurium
C. aritum
Agriocharis ocellata
Chrysolophus pictus
C. amherstiae
(ceca of all): all from San Diego Zoological Garden
- Heterakis spumosa*
Duewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Heterakis spumosa*
Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
Rattus norvegicus (gros intestin, caecum): Tunis, Tunisia
- Heterakis spumosa* Schneider, 1866, *illus.*
Palomino, H.; and Barriga, O. O., 1967, *Bol. Chileno Parasitol.*, v. 22 (2), 79
Heterakis spumosa infection in *Rattus norvegicus* (colon, ciego) also harboring *Hymenolepis nana* var. *fraterna*: Chile
- Heterakis spumosa* (Schneider, 1866)
Singh, M.; and Cheong Chee Hock, 1971, *South-east Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 516-521
Rattus bowersi: Malaysia
- Heterakis spumosa* Schneider, 1866
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Rattus norvegicus: insular Newfoundland
- Heterakis spumosa*, *illus.*
Taniguchi, M.; et al., 1977, *Bull. Coll. Agric. and Vet. Med.*, Nihon Univ. (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo
- Heterakis spumosa* Schneider, 1886
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 39-42
Rattus norvegicus (intestino grueso): Chile
- Heterakis tragopanis* Lal, 1942
Bali, H. S.; and Kalra, I. S., 1975, *J. Research, Punjab Agric. Univ.*, v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Heterakoidea*
Chabaud, A. G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (1), 6-17
Ascaridida
key
- Heterocheilidae* Railliet & Henry, 1915, in part
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakidae* (Railliet & Henry, 1912, subfam.) Skrjabin & Karokhin, 1945
- Heterocheilidae* (Railliet & Henry, 1912, subfam.) Railliet & Henry, 1915
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridoidea
key
includes: *Heterocheilus*
- Heterocheilus* Diesing, 1839
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Heterocheilidae
synonymy
- Heteromyoxyuris deserti*
Garner, H. W.; Richardson, L. W.; and Felts, L. A., 1976, *Southwest. Nat.*, v. 21 (3), 327-334
monthly percentages of animals parasitized
Dipodomys ordii (caecum): western Texas
- Heteromyoxyuris deserti* (Read and Millemann, 1953)
King, S. R.; and Babero, B. B., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 241-248
Dipodomys merriami
D. deserti
(cecum of all): all from Nevada
- Heterotylenchus* sp. Nicholas and Hughes, 1970
Slobodianiuk, O. V., 1975, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 25, 156-168
as syn. of *Paraiotonchium nicholasi* sp. nov.
- Heterotylenchus autumnalis*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, *J. Parasitol.*, v. 62 (6), 877-880
Musca autumnalis: Kentucky
- Heterotylenchus autumnalis* Nickle
Robinson, J. V.; and Combs, R. L., jr., 1976, *J. Econom. Entom.*, v. 69 (6), 722-724
Heterotylenchus autumnalis in *Musca autumnalis*, seasonal incidence, host longevity: Monroe and Chicksaw Counties, Mississippi
- Heterotylenchus autumnalis* Nickle, 1967
Slobodianiuk, O. V., 1975, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 25, 156-168
as syn. of *Paraiotonchium autumnalis* (Nickle, 1967) comb. n.
- Heterotylenchus pawlowskyi* Kurochkin, 1960
Poinar, G. O., jr.; and Nelson, B. C., 1973, *J. Med. Entom.*, v. 10 (4), 349-354
as syn. of *Psyllotylenchus pawlowskyi* (Kurochkin, 1960) n. comb.
- Heterotyplum* Spaul, 1927
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinea
key

- Heteroxyinema chiliensis*
 Babero, B. B.; and Cattán, P. E., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 68-76
Octodon degus (ciego e intestino grueso): Quebrada de la Plata, Santiago, Chile
- Heteroxyinema chiliensis* (Quentin, 1975)
 Cattán, P. E.; George-Nascimento, M.; and Rodríguez, J., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 16-20
 prevalence survey of helminths of *Octodon degus*, seasonal variations, age and sex of hosts: Chile
- Hexameris albicans* (Siebold)
 Drea, J. J.; et al., 1977, Entomophaga, v. 22 (2), 141-146
Hexameris albicans in *Lymantria dispar* and *Stilpnotia salicis*, percentage parasitism, emergence period of nematodes, biological control
Lymantria dispar: Burgenland, Austria; Würzburg, West Germany
Stilpnotia salicis: Burgenland, Austria
- Hexameris albicans* (Siebold 1848) Polozhentsev & Artiukhovskiy 1959
 Kaiser, H., 1974, Mitt. Naturw. Ver. Steiermark, v. 104, 177-181
 synonymy
 Steiermark, Österreich
- Hexameris brevis* (Hagmeier 1912) Polozhentsev & Artiukhovskiy 1959
 Kaiser, H., 1974, Mitt. Naturw. Ver. Steiermark, v. 104, 177-181
 synonymy
 Steiermark, Österreich
- Hexametra Travassos*, 1919
 Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ophidascaridinea
 key
- Hexametra* sp., illus.
 Clarke, M. D.; Chang, T. Y.; and Cross, J. H., jr., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 565-567 [Demonstration] morphology, life cycle
Trimeresurus stejneri: Taiwan, Republic of China
 mouse (exper.) (liver and abdominal cavity tissues)
- Hexametra quadricornis* Wedl 1862
 Pinnell, J. L.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 337-340
Psammodynastes pulverulentus
Trimeresurus albolabris
 all from Komodo Island, Indonesia
- Hexametrinae* Mozgovoi & Shakhmatova, 1971
 Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
 as syn. of *Ophidascaridinea* (Hartwich, 1954, subfam.) Chabaud, 1965
- Histiocephalinae* Gendre, 1922
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematidae
 key; key to genera
 includes: *Hadjelia*; *Histiocephalus*; *Viguiera*; *Stellocaronema*; *Torquatoides*
- Histiocephalus* Diesing, 1851, illus.
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Histiocephalinae
 key
- Histiostrongylus* Molin, 1861
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Anoplostrongylinae*
- Hoineffia* n. gen.
 Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Heligmosomidae, *Viannaiinae*
 tod: *H. cayennensis* n. gen. n. sp.
- Hoineffia* Diaw, 1977
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, *Viannaiinae*
- Hoineffia cayennensis* Diaw, 1976 [? nom. nud.]
 Diaw, O. T., 1976, Ann. Parasitol., v. 51 (3), 355-363
 trichostrongyloid nematode fauna of *Didelphis marsupialis* compared to that of *Metachirops opossum*, localization within intestine
Didelphis marsupialis (intestin): Guyane francaise
- Hoineffia cayennensis* n. gen. n. sp. (tod), illus.
 Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Didelphis marsupialis (dernier quart de l'intestin): Guyane francaise
- Hookworm
 Anderson, R. I.; and Buck, A. A., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 447-456
 complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad
- Hookworm
 Areekul, S., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 310-312
 human hookworm, changes in blood volume, red cell volume and plasma volume in infected persons compared with normal subjects
- Hookworm
 Areekul, S.; et al., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 519-523
 human hookworm anemia, assessment of blood loss, iron absorption and iron reabsorption in infected humans: Thailand

- Hookworms**
Areekul, S.; et al., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 568-569 [Demonstration]
hookworms, humans and dogs, inverse relationship between number of worms and blood loss per worm per day
- Hookworms**
Areekul, S.; et al., 1970, Southeast Asian J. Trop. Med. and Pub. Health, v. 1 (4), 569 [Demonstration]
human hookworm infection, measurement of gastro-intestinal protein loss
- Hookworm**
Areekul, S.; et al., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 107 [Demonstration]
comparison of serum vitamin B₁₂ levels in patients with hepatic amoebic abscess, opisthorchiasis or hookworm infections
- Hookworm**
Areekul, S.; Devakul, K.; and Boonyananta, C., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 585 [Demonstration]
iron storage in infected humans compared with that of normal controls, results showed that iron excretion in urine of infected persons was less than that of controls
- Hookworm**
Areekul, S.; Devakul, K.; and Kanakakorn, K., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 414 [Demonstration]
hookworm, amoebiasis, humans, effects of anemia on iron absorption compared in infected and control subjects
- Hookworm**
Areekul, S.; Pinyawatana, W.; and Kitkornphan, S., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 386-390
hookworm, human, red cell and serum folate levels and folic acid absorption, impairment of folate absorption and iron deficiency anemia were probably primary and secondary causes of low serum folate content in these patients
- Hookworm**
Biagi, F., 1975, Progr. Drug Research, v. 19 23-27
human hookworm and trichuriasis successfully treated with bitoscanate in clinical trials; mixed results when used to treat ascariasis in comparison trials with dithiazanine and piperazine
- Hookworm**
Briggs, M.; Wenlock, R. W.; and Briggs, M. H., 1972, Med. J. Zambia, v. 6 (2), 39-41
comparison of serum proteins of healthy subjects and those of persons with untreated malaria, hookworm or schistosomiasis, possible diagnostic value: Zambia
- Hookworm**
Bruce-Tagoe, A. A.; et al., 1977, Trop. and Geogr. Med., v. 29 (3), 237-244
human malaria and hookworm, correlations with hematological values and anemia in survey of rural population in Ghana
- Hookworm**
Burke, G. J., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 402-405
10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Hookworm sp.**
Cabrera, B. D., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (1), 50-55
Rattus rattus (feces): Leyte, Philippines
- Hookworm**
Chamorro, H.; and Moizeszowicz, J., 1973, Bol. Chileno Parasitol., v. 28 (1-2), 24-30
hookworm anemia, clinical trials testing iron therapy: Argentina
- Hookworm**
Douglas, J.; and Tamanika, M., 1973, Papua N. Guinea Med. J., v. 16 (1), 54-58
human hookworm anemia, positive correlation between degree of hookworm infection and degree of anemia: Sogeri rubber tappers, Papua New Guinea
- Hookworm**
Forman, D. W.; et al., 1971, Am. J. Trop. Med. and Hyg., v. 20 (4), 598-601
parasitic pathogens discovered during etiologic survey of causes of diarrheal disease in U.S. Marines in Vietnam
- Hookworm**
Garby, L.; and Areekul, S., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 467-476
iron-deficiency anemia caused by hookworm, possible prevention by iron supplementation of fish-sauce (a widely used food product): Thailand
- Hookworm**
Goulart, E. G.; et al., 1977, J. Helminth., v. 51 (2), 131-132
ancylostomiasis, Strongyloides stercoralis, human, field trials of control by means of mass anthelmintic treatment combined with introduction of plants inhibitory to the free-living larval stages, reduction in prevalence: Ilha Do Governador, Rio de Janeiro, Brazil
- Hookworm**
Grove, D. I.; Burston, T. O.; and Forbes, I. J., 1974, Clin. and Exper. Immunol., v. 18 (4), 565-569
hookworm-infested population, high serum IgE levels, serum IgE and blood eosinophil levels fell after treatment with pyrantel: Papua New Guinea
- Hookworm**
Gupta, M. C.; et al., 1977, Indian J. Med. Research, v. 66 (2), 244-252
health and nutrition survey of men doing heavy manual labor, mild anemia in group but hookworm and other parasites apparently non-contributory: Dumka region in Bihar and Garhwal, India

- Hookworm (egg), illus.
Hira, P. R.; and Patel, B. G., 1977, Am. J. Trop. Med. and Hyg., v. 26 (4), 640-643
- Hookworm
Hoves, H. L., jr., ., 1972, Proc. Soc. Exper. Biol. and Med., v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that T. muris-mouse infection could be test model for antiwhipworm studies
- Hookworm
Islam, N.; and Chowdhury, N. A., 1976, South-east Asian J. Trop. Med. and Pub. Health, v. 7 (1), 81-84
human intestinal parasites, comparative treatment trials using mebendazole and pyrantel pamoate, recommendations for use: Dacca, Bangladesh
- Hookworm
Johnson, S., 1975, Progr. Drug Research, v. 19, 70-74
human hookworm, comparative field trials with bitoscanate, bephenium hydroxynaphthoate, and tetrachlorethylene, results most favorable with bitoscanate, transient side effects: Vellore, Tamilnadu, India
- Hookworm
Kagei, N., 1975, Bull. Inst. Pub. Health, Tokyo, v. 24 (3), 169-175
comparison of Kato thick smear and Tween 80 citric acid ether sedimentation methods for diagnosis of helminth ova
- Hookworm
Kaplan, M. H.; and Bernstein, L. S., 1974, Mil. Med., v. 139 (6), 444-448
Plasmodium vivax, U.S. soldiers in Vietnam, improved care with primaquine combined with chloroquine therapy; survey of servicemen shows frequent mixed hookworm infections, increased serum IgM levels and decreased cholesterol levels
- Hookworm
Kaul, T. N.; and Mahajan, R. C., 1977, Indian J. Med. Research, v. 66 (3), 413-416
human hookworm infection, immunofluorescence using Ancylostoma duodenale antigen, useful immunodiagnostic tool especially in early stages of infections
- Hookworms
Khan, S. M.; and Khairul Anuar, A., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (2), 260-264
Ascaris lumbricoides, Trichuris trichiura, hookworms, prevalence survey of soil-transmitted intestinal helminths in patients admitted to the District Hospital in Balik Pulau, Penang, Malaysia
- Hookworms
Langer, A.; and Hung, C. T., 1973, Obst. and Gynec., v. 42 (4), 564-567
human hookworm anemia associated with pregnancy, clinical management with tetrachlorethylene or with tetrachlorethylene and thia-bendazole in Trichuris trichiura-associated infections
- Hookworm
Legendre, A. M., 1973, J. Am. Vet. Med. Ass., v. 163 (2), 149-150
disophenol toxicosis in a dog treated for hookworm
- Hookworm
Mahmud-Durrani, A.; Desai, M. H.; and Tembo, D., 1970, Med. J. Zambia, v. 4 (4), 121-123
commonest intestinal parasite in Zambia, with most people suffering only light infections
- Hookworm
Meakins, R. H.; Carswell, F.; and Harland, P. S. E. G., 1977, Parasitology, v. 75 (2), xxxii [Abstract]
helminthiasis in school children, association with malnutrition: rural Tanzania
- Hookworms
Most, H., 1972, N. England J. Med., v. 287 (10), 495-498; (14), 698-702
common parasitic infections of man encountered in the United States, recommendations for treatment, review
- Hookworm
Mutalik, G. S.; and Gulati, R. B., 1975, Progr. Drug Research, v. 19, 86-89
human hookworm, comparative study of bitoscanate, bephenium hydroxynaphthoate, and tetrachlorethylene; bitoscanate most effective with mild and transient side effects
- Hookworm
Mutalik, G. S.; Gulati, R. B.; and Igbal, A. K., 1975, Progr. Drug Research, v. 19, 81-85
human intestinal parasites, clinical trials with bitoscanate show it to be safe and useful anthelmintic especially against hookworm infections: India
- Hookworm
Naik, S. R.; et al., 1976, Digestion, v. 14 (2), 133-141
human gastric acid secretory responses to continuous infusion of histamine, laboratory trials to assess maximal acid output in patients with hookworm iron deficiency anemia
- Hookworms
Ndiritu, C. G.; and Al-Sadi, H. I., 1977, J. Small Animal Practice, v. 18 (3), 199-205
hookworms, dogs, age and sex incidence, seasonal distribution, clinical picture, pathology: Nairobi, Kenya
- Hookworm
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Hookworm
Neppert, J.; and Warns, C.-M., 1974, Tropenmed. u. Parasitol., v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from Ascaris, hookworm, Onchocerca, Dirofilaria immitis, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination

- Hookworm**
Nhonoli, A. M.; and Chukwuemeka, A. C., 1971, *Med. J. Zambia*, v. 5 (3), 95-101
analysis of electrocardiographic changes in patients with severe hookworm anemia before and after treatment with alcopar and tetrachlorethylene: Zambia
- Hookworm**
Ohlrogge, R.; and Schlipkoeter, H. W., 1973, *J. Microbiol. and Immunol.*, v. 158 (4), 267-274
deviations in erythrogram before and after treatment of human malaria infections with resochoin, no such changes in hookworm and other worm infections: Liberia; Togo
- Hookworm**
Oomi, H.; Watanabe, T.; and Yoshida, Y., 1976, *J. Agric. Sc., Tokyo Nogyo Daigaku*, v. 20 (3-4), 206-212
hookworm, whipworm, dogs, subcutaneous injection, methyridine + disophenol, good results
- Hookworm**
Pitchford, R. J.; and Visser, P. S., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 16 [Demonstration]
quantitative technique for the estimation of helminth eggs in urine and faeces
- Hookworm**
Purtilo, D. T.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (2), 229-232
parasitized children with protein-calorie malnutrition, complete blood count, serum immunoglobulin concentration, significant relationship between intensity of parasitism and Ig levels: Brazil
- Hookworm**
Rees, P. H.; and Marsden, P. D., 1970, *Brit. J. Clin. Pract.*, v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment
- Hookworm**
Reid, B. D.; Reid, T. E.; and Ullstrop, G., 1971, *Med. J. Zambia*, v. 5 (2), 61-69
study of health of schoolchildren, parasitic survey and possible associations with nutritional status: Zambia
- Hookworm**
Saraya, A. K.; and Tandon, B. N., 1975, *Progr. Drug Research*, v. 19, 108-118
human hookworm infection, study of associated anemia and intestinal malabsorption
- Hookworm**
Savanat, T.; et al., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 149-154
determinations of total serum IgE levels in humans with amoebic liver abscess or other parasitic infections
- Hookworm**
Seah, S. K. K., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 534-542
intestinal parasites, persons living in non-endemic areas who acquired infections while travelling or who have immigrated from endemic areas, pyrantel pamoate successful for *Ascaris lumbricoides*, results with other parasites varied: Montreal, Canada
- Hookworm**
Seah, S. K. K., 1976, *Canad. Med. Ass. J.*, v. 115 (8), 777-779
mebendazole suggested as drug of choice for *Trichuris trichiura* and mixed nematode infections after extensive clinical trials: Canada
- Hookworm**
Sehgal, S. C.; Vinayak, V. K.; and Gupta, U., 1977, *Indian J. Med. Research*, v. 65 (4), 509-512
human helminth ova in feces, diagnosis using the Kato thick smear technique more successful than commonly used techniques, recommended for epidemiologic surveys: Chandigarh, India
- Hookworm**
Singson, C. N.; Banzon, T. C.; and Cross, J. H., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (6, pt. 1), 932-934
Capillaria philippinensis, human clinical trials using mebendazole for intestinal capillariasis, additionally effective against *Ascaris lumbricoides*, *Trichuris trichiura* and hookworm: Philippines
- Hookworm**
Tantengco, V. O.; et al., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (2), 210-221
hookworm, analysis of serum and red cell folate activity and its relationship to hemoglobin concentration in infected and hookworm free children
- Hookworm**
Tantengco, V. O.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 524-533
Ascaris, *Trichuris*, hookworm infections apparently not contributory cause of nutritional anemia in schoolchildren: Philippine Islands
- Hookworm**
Turner, K. J.; Baldo, B. A.; and Anderson, H. R., 1975, *Internat. Arch. Allergy and Applied Immunol.*, v. 48 (6), 784-799
humans, serum IgE levels, no significant correlation with faecal egg counts to hookworm, *Ascaris lumbricoides*, and *Trichuris*, incidence of IgE antibodies to *Ascaris lumbricoides* not correlated with incidence of asthma but significantly elevated in patients with chronic obstructive lung disease, hypersensitivity to *Ascaris* apparently not factor of importance in etiology of asthma in this area: Highland area of Papua-New Guinea

- Hookworm
 Wakil, B. J.; and Dalal, N. J., 1975, Progr. Drug Research, v. 19, 166-175
 comparative efficacy of newer anthelmintics in treating various human intestinal helminths, review
- Hookworm
 Vinayak, V. K.; and Sehgal, S. C., 1976, Indian J. Med. Research, v. 64 (9), 1347-1350
 human helminthic and protozoan parasites, comparison of nigrosin-methylene blue diagnostic test with formol-ether method and direct examination
- Hookworm
 Wenlock, R. W., 1977, Brit. J. Nutrition, v. 38 (2), 239-243
 efficiency of urinary hydroxyproline index as indicator of nutritional status in mass surveys evaluated in presence of schistosomiasis, hookworm and malaria; in endemic malaria areas index probably of little value without prior evaluation of malarial status of all subjects
- Hookworm
 Whitehead, R., 1973, Major Problems Path., v. 3, 105-110
 human intestinal infection, diagnosis, pathological appearance of mucosal biopsy of gastrointestinal tract
- Hookworm
 Yang, J.; and Scholten, T., 1977, Am. J. Clin. Path., v. 67 (3), 300-304
 diagnosis of human intestinal parasites, fecal examination technique using Junod's fixative for concentration and permanent staining procedures, comparison with results using formalin-ether procedure
- Hookworm
 Zigas, V., 1973, Papua N. Guinea Med. J., v. 16 (1), 51-53
 human hookworm anemia, positive correlation between hemoglobin levels and degree of infection: Southern Highlands of Papua New Guinea
- Hooperstrongylus Lie et Kong, 1963
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Molineidae, Molineinae
- Hovorkonema gen. n.
 Jurasek, V., 1977, Biologia, Bratislava, s. B, Zool. (1), v. 32 (2), 105-109
 Aprocridae
 tod: *H. gastrofilaria* sp. n.
- Hovorkonema gastrofilaria sp. n. (tod), illus.
 Jurasek, V., 1977, Biologia, Bratislava, s. B, Zool. (1), v. 32 (2), 105-109
Sus scrofa atilla (stomach): Lucenec (Baglas), Slovak Socialist Republic (CSSR)
- Howardula species
 Elsey, K. D., 1977, J. Invert. Pathol., v. 29 (3), 384-385
Chaetocnema confinis
C. pulicaria
Colaspis brunnea
Epitrix fuscula
 all from North Carolina
- Howardula species, possibly *H. phyllotretae*
 Elsey, K. D., 1977, J. Invert. Pathol., v. 29 (3), 384-385
Phyllotreta striolata
P. zimmermanni
 all from North Carolina
- Howardula sp.
 Elsey, K. D.; and Pitts, J. M., 1976, Environment. Entom., v. 5 (4), 707-711
Howardula sp., incidence and importance in natural control of tobacco flea beetle
Epitrix hirtipennis: Oxford and Clayton, North Carolina
- Howardula sp., illus.
 Miles, P. M., 1977, Entom. Month. Mag. (1344-1347), v. 112, 169-172
Parasitus sp.: Tarlton, Gloucestershire
- Howardula acris n. sp., illus.
 Remillet, M.; and van Waerebeke, D., 1977, Cahiers O.R.S.T.O.M., s. Biol., v. 11 (3), 1976, 219-224
 life cycle
 Hydrophilidae: parc de Tsimbazaza, Tananarive
- Howardula benigna
 Elsey, K. D., 1977, J. Invert. Pathol., v. 29 (3), 384-385
Diabrotica undecimpunctata howardi
Acalymma vittata
 all from North Carolina
- Howardula husseyi Richardson, Hesling & Riding
 Richardson, P. N.; and Hesling, J. J., 1977, Ann. Applied Biol., v. 86 (3), 321-327
Howardula husseyi, winter decline in parasitism of *Megaselia halterata*, major mushroom pest; beakers in field test
- Humpsore. See Stephanofilariasis.
- Hydromermis
 Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
 key
- Hydromermis contorta
 Richter, S., 1975, Ztschr. Parasitenk., v. 47 (1), 69 [Correction for 1971 a]
Hydromermis contorta, preparasitic larvae, stoma armament, orientation
- Hydromermis contorta, illus.
 Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105

- Hyostromylus*
Dey-Hazra, A., 1976, Ztschr. Parasitenk., v. 50 (2), 198
helminths, pigs, mode of pathogenicity, review
- Hyostromylus* Hall, 1921
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Graphidiinae
- Hyostromylus*
Jacobs, D.; and Schulze, H. W., 1977, Prakt. Tierarzt, v. 58 (1), 46-48
pig parasites, natural infections, vermitin and dichlorvos effective in field testing
- Hyostromylus rubidus*
Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, Vet. Rec., v. 99 (7), 119-122
swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with pica-dex
- Hyostromylus rubidus*
Bussieras, J., 1976, Rec. Med. Vet., v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Hyostromylus rubidus*
Connan, R., 1977, Vet. Rec., v. 100 (12), 242-243
declining prevalence in pigs (stomachs): abattoir [Great Britain]
- Hyostromylus rubidus*
Düwel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Hyostromylus rubidus* (Hassall et Stiles, 1892)
Hall, 1921, illus.
Ianchev, I., 1973, Izvest. Tsentral. Khel'mint. Lab., v. 16, 205-220
Capreolus capreolus (rennet): southern Bulgaria
- Hyostromylus rubidus*
Kirsch, R.; and Düwel, D., 1975, Research Vet. Sc., v. 19 (3), 327-329
Hyostromylus rubidus, *Oesophagostomum* spp., pigs (exper.), efficacy of fenbendazole
- Hyostromylus rubidus*
Kutzer, E.; and Frey, H., 1976, Berl. u. Münch. Tierärztl. Wchnschr., v. 89 (24), 480-483
Lepus europaeus: Austria
- Hyostromylus rubidus*
Kutzer, E.; and Frey, H., 1976, Ztschr. Parasitenk., v. 50 (2), 213-214
Lepus europaeus
- Hyostromylus rubidus*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile
- Hyostromylus rubidus*
Peterson, P. M.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1778-1780
Ascarops strongylina, *Physocephalus sexalatus*, *Hyostromylus rubidus*, natural incidence, comparison with earlier surveys
hogs (stomachs): Georgia; Wisconsin
- Hyostromylus rubidus*
Pfeiffer, A., 1977, Prakt. Tierarzt, v. 58 (1), 32-38
Hyostromylus rubidus, *Oesophagostomum* spp., sows treated for improved weight gain of weanling pigs
- Hyostromylus rubidus* (Hassall and Stiles, 1892)
Probert, A. J.; Smith, B. D. S.; and Herbert, I. V., 1973, Vet. Rec., v. 93 (11), 302, 303-306
Hyostromylus rubidus, pigs (exper.), levamisole administered orally and subcutaneously, more effective against adults than larvae with selective elimination of male larvae compared with females
- Hyostromylus rubidus*
Raynaud, J. P., 1976, Pathophysiol. Parasit. Infect., 99-104
Oesophagostomum spp., *Hyostromylus rubidus*, *Ascaris suum*, young swine (exper.), multi-stage multiparasite model for pathological and anthelmintic studies
- Hyostromylus rubidus*
Raynaud, J.-P.; Sennelier, J.; and Irisarri, E., 1975, Folia Vet. Latina, v. 5 (3), 412-429
gastrointestinal helminths, swine, post natal infection of piglets in contact with infected mothers, comparison of various methods of husbandry and hygiene, studies during pregnancy and lactation, routine daily hygiene recommended
- Hyostromylus rubidus*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Hyostromylus rubidus*
Rose, J. H., 1973, Research Vet. Sc., v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostromylus rubidus*, culture from infective larva to adult worm in WAe medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Hyostromylus rubidus* (Hassall and Stiles, 1892)
Smith, H. V.; and Herbert, I. V., 1976, Vet. Parasitol., v. 1 (4), 327-335
Hyostromylus rubidus, pigs, primary infection, antibody response, time course and kinetics as shown by passive haemagglutination test

- Hyostrongylus rubidus*
Smith, H. V.; and Herbert, I. V., 1976, *Immunology*, v. 30 (2), 213-219
Hyostrongylus rubidus, passive transfer of humoral immunity from infected sows to their offspring via colostrum, demonstration that agglutinating antibodies mainly of the IgG class were associated with protection
- Hyostrongylus rubidus* (Hassall and Stiles, 1892)
Smith, H. V.; Herbert, I. V.; and Davis, A. J., 1976, *Vet. Parasitol.*, v. 1 (4), 337-344
Hyostrongylus rubidus, pigs, multiple infections and reinfections, antibody response, passive haemagglutination test
- Hyostrongylus rubidus*
Stewart, T. B.; Hale, O. M.; and Marti, O. G., 1975, *Am. J. Vet. Research*, v. 36 (6), 771-772
Hyostrongylus rubidus, pigs (exper.), treatment with coated vs. uncoated formulations of dichlorvos, efficacious against adult worms but little or no activity against 5- or 15-day-old worms, not as effective in sows as in barrows and gilts
- Hyostrongylus rubidus*
Taffs, L. F., 1976, *Brit. Vet. J.*, v. 132 (1), 105-111
cambendazole, little or no effect against 10-day-old *Hyostrongylus rubidus* or *Oesophagostomum* spp. larvae when given to pigs at oral dose rates of 15, 20, and 25 mg/kg bodyweight
- Hyostrongylus rubidus*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Hyostrongylus rubidus*
Valenzuela, G.; et al., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago): *Planta Faenadora de Carnes Socoagro*, Valdivia, Chile
- Hypocristata Durette-Desset*, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, *Nippostrongylinae*
- Hypodontus*
Durette-Desset, M. C.; Denke, M. A.; and Murua, R., 1976, *Ann. Parasitol.*, v. 51 (4), 453-460
placed in *Uncinariinae*, excluded from *Amidostomatidae*
- Hypodontus macropi* Monnig, 1929, *illus.*
Setasuban, P.; and Arundel, J. H., 1976, *South-east Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 102-105
Hypodontus macropi, morphometric data, comparison of statistics of males and females, scanning electron microscopy
Macropus rufogrisea (intestine): Tasmania, Australia
- Hysteracrum (Oesophagostomum) venulosum* (Rudolphi, 1809), *illus.*
Goffredo, G.; and Sobrero, R., 1972, *Parassitologia*, v. 14 (1), 143-148
Dama dama (intestine): foresta Umbra (promontorio garganico, provincia di Foggia)
- Hysterothylacium* Ward & Magath, 1916
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Raphidascaris* Railliet & Henry, 1915
- Hystrichis tricolor* (Dujardin, 1845), *illus.*
Bogoiavlenskii, Iu. K.; and Khatkevich, L. M., 1970, *Parazitologiya*, Leningrad, v. 4 (3), 223-230
4 spp. of *Diectophymata*, fine structure of somatic musculature, distribution of DNA and RNA
- Hystrichis tricolor*
Bogoiavlenskii, Iu. K.; and Koroleva, N. A., 1969, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 20, 21-29
Ascaridia galli, *Hystrichis tricolor*, comparison of micromorphology and histochemistry of hypodermal-muscular sac during pre-imaginal development
- Hystrichis tricolor* Dujardin, 1845
Kamburov, P.; and Vasilev, I., 1972, *Izvest. Tsentral. Khelmint. Lab.*, v. 15, 109-133
Anas platyrhynchos
A. penelope
A. querquedula
Aythya nyroca
Mergus serrator
(wall of crop of all): all from Bulgaria
- Hystrichis tricolor* Dujardin, 1845
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, *Am. Midland Naturalist*, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans
Porphyryula martinica
(proventriculus of all): all from Florida

- Ichthyanisakis Gendre, 1928
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Paranisakis Baylis, 1923
- Ichthyascaris Wu, 1949
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Raphidascaris Railliet & Henry, 1915
- Ichthyobronema Gnedina and Savina, 1930
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
as syn. of Haplonema Ward and Magath, 1917
- Ichthyobronema Gnedina & Ssavina, 1930
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
? as syn. of Rhabdochona Railliet, 1916
- Ichthyobronema conoura (von Linstow, 1885) sensu Gnedina and Savina, 1930
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
as syn. of Haplonema hamulatum Moulton, 1931
- Ichthyobronema gnedini Sudarikov and Ryzhikov, 1952
Arthur, J. R.; and Margolis, L., 1975, Canad. J. Zool., v. 53 (6), 736-747
as syn. of Haplonema hamulatum Moulton, 1931
- Ichthyofilaria Yamaguti, 1935, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key
- Ichthyonema cylindraceum Ward and Magath, 1916
Molnar, K.; and Fernando, C. H., 1975, J. Helminth., v. 49 (1), 19-24
as syn. of Philometra cylindracea (Ward and Magath, 1916)
- Ichthyospirura Skrjabin, 1917
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of Rhabdochona Railliet, 1916
- Ichthyobronema. See Ichthyobronema.
- Icosiella neglecta (Diesing, 1851) Seurat, 1917
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Icosiella neglecta (Diesing, 1851), illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Rana ridibunda
R. esculenta
(misici straznjih ekstremiteta of all): all from Yugoslavia
- Icosiella neglecta (Diesing, 1851) Seurat, 1917, illus.
Rozman, M., 1971, Acta Parasitol. Iugoslavica, v. 2 (2), 67-77
description
synonymy
Rana esculenta (muskulatura zadnjih ekstremiteta): environs of Novi Sad, Yugoslavia
- Icosiella neglecta Diesing 1851, illus.
Sanchez-Acedo, C.; and Vericad, J. R., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 197-203
Buteo buteo: Aragon Pyrenees
- Iheringascaris Pereira, 1935
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Thynnascaris Dollfus, 1933
- Impalaia Moennig, 1923
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Cooperiinae
synonymy
- Impalaia aegyptiaca Soliman, 1958
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
as syn. of Impalaia tuberculata Moennig, 1923
- Impalaia dremomys Yen, 1973
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
as syn. of Heligmonella dremomys (Yen, 1973) n. comb.
- Impalaia nudicollis of Daubney, 1933; Yeh, 1956; and Pande et al., 1962
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
tentatively as syn. of Impalaia tuberculata Moennig, 1923
- Impalaia nudicollis Moennig, 1931, illus.
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, Ann. Parasitol., v. 52 (4), 435-446
redescription, key
Damaliscus albifrons: South Africa
- Impalaia nudicollis Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Impalaia nudicollis
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Impalaia nudicollis
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, Camelus dromedarius, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad

- Impalaia okapiae* van den Berghe, 1937
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, *Ann. Parasitol.*, v. 52 (4), 435-446
valid species, key
- Impalaia taurotragi* (Le Roux, 1936) Travassos, 1937, *illus.*
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, *Ann. Parasitol.*, v. 52 (4), 435-446
redescription, key
Taurotragus oryx (duodenum): Northern Rhodesia
- Impalaia tuberculata* Moennig, 1923, *illus.*
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, *Ann. Parasitol.*, v. 52 (4), 435-446
synonymy, redescription, key
Aepyceros melampus (intestine): Transvaal, South Africa; Suguroi Estate, Kenya
Taurotragus oryx (small intestine): Serengeti region, Tanzania
Gazella thomsonii (small intestine): Gilgil, Kenya
Giraffa camelopardalis (small intestine): Maxwell Zoological Park, England
Camelus dromedarius: Agadis, Niger; slaughter house of Negele Borana, Addis-Abeba, Ethiopia
- Impalaia tuberculata*
Wilson, D. E.; and Hirst, S. M., 1977, *Wildlife Monogr.* (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Impalaia tuberculata* var. *longispiculata* Wetzel and Fortmeyer, 1960
Gibbons, L.; Durette-Desset, M. C.; and Daynes, P., 1977, *Ann. Parasitol.*, v. 52 (4), 435-446
as syn. of *Impalaia tuberculata* Moennig, 1923
- Indocamallanus Chakravarty et al.*, 1963
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Procamallanus* Baylis, 1923
- Indocucullanus calcariferii* n. sp., *illus.*
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
Lates calcarifer (intestine): Karachi Coast, Pakistan
- Indocucullanus karachii* n. sp., *illus.*
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
Engraulis indica (intestine): Fish Harbour, Karachi, Pakistan
- Indocucullanus longispiculum* Khan, 1969, *illus.*
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
male and female redescribed
Lates calcarifer (stomach): Karachi Coast, Pakistan
- Inglamidinae n. subfam.
Durette-Desset, M. C.; Denke, M. A.; and Murua, R., 1976, *Ann. Parasitol.*, v. 51 (4), 453-460
Amidostomatidae
type genus: *Inglamidum* n. gen.
- Inglamidinae Durette-Desset, Diaw et Murua, 1976
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae
includes: *Inglamidum* (type genus)
- Inglamidum* n. gen. (type genus of subfam.)
Durette-Desset, M. C.; Denke, M. A.; and Murua, R., 1976, *Ann. Parasitol.*, v. 51 (4), 453-460
Amidostomatidae, *Inglamidinae*
tod: *I. akodon* n. sp.
- Inglamidum* Durette-Desset, Diaw et Murua, 1976 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae, *Inglamidinae*
- Inglamidum akodon* n. sp. (tod), *illus.*
Durette-Desset, M. C.; Denke, M. A.; and Murua, R., 1976, *Ann. Parasitol.*, v. 51 (4), 453-460
Akodon olivaceus (intestine grele): San Martin, Province de Valdivia, Chile; Esc. Normal, Province de Valdivia, Chile
Akodon sanborni: Picada, Province d'Osorno, Chile
- Inglamidum akodon*
Denke, M. A.; and Murua, R., 1977, *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (428), *Zool.* (298), 127-131
Akodon olivaceus: Province de Valdivia, Chile
A. sanborni: Province d'Orsono, Chile
- Ingliseria* Gibson, 1968, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key
- Itonchium* n. sp., *illus.*
Laumond, C.; and Lyon, J. P., 1975, *Acta Trop.*, v. 32 (4), 334-339
Allantonematidae n. sp., probably belonging to *Itonchium*, morphology and life cycle
Helophilus trivittatus
H. pendulus
(hemocoel of all): all from Sud de la France
- Irukanema dalli* Yamaguti 1951
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Torynurus dalli* (Yamaguti 1951)
Delyamure 1972
- Irukanema dalli* Yamaguti, 1951
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Phocoena phocoena: insular Newfoundland and its adjacent waters

- Isolaimida Timm, 1969
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Enoplia
- Isomermis sp., illus.
Rubtsov, I. A., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 128-156
ontogenesis of mermithids, illustrated description of structure of body, cuticle, amphids, longitudinal fields, stichosome, osmosome, trophosome and reproductive organs; technique of preparing material
- Isomermis wisconsinensis Welch
Ebsary, B. A.; and Bennett, G. F., 1975, Canad. J. Zool., v. 53 (8), 1058-1062
Simulium venustum
Simulium vittatum
all from insular Newfoundland
- Johnstonema [? n. rank]
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of Breinlia; key
tod: B. (J.) annulipapillatum (Johnston et Mawson, 1938)
- Johnstonmawsonia Campana-Rouget, 1955, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochoniidae
key
Syn.: Prosungulonema Roytman, 1963
- Johnstonmawsonia Campana-Rouget, 1955
Puylaert, F. A., 1973, Rev. Zool. et Botan. Africaines, v. 87 (4), 647-665
discussion of systematic position
- Johnstonmawsonia sp., illus.
Petter, A. J.; Goivan, Y. J.; and Tcheprakoff, R., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 159-171
description
Anguilla rostrata (tube digestif): riviere Sarcelles, Guadeloupe
- Johnstonmawsonia campanae sp. n., illus.
Puylaert, F. A., 1973, Rev. Zool. et Botan. Africaines, v. 87 (4), 647-665
Aphyosemion cameronensis (intestin, estomac): Cameroun, Olounou
- Johnstonmawsonoides n. g.
Machida, M., 1975, Bull. National Sc. Mus., Tokyo, s. A, Zool., v. 1 (1), 1-4
Rhabdochoniidae, Ascarophidinae
tod: J. nemichthyos n. sp.
- Johnstonmawsonoides nemichthyos n. g., n. sp. (tod), illus.
Machida, M., 1975, Bull. National Sc. Mus., Tokyo, s. A, Zool., v. 1 (1), 1-4
Nemichthys scolopaceus (intestine): Suruga Bay, Japan
- Kalicephalus
Fernandes, M. P. M.; and Artigas, P. T., 1975, Mem. Inst. Butantan, v. 39, 103-121
key to species, includes: *Kalicephalus inermis*; *K. subulatus*; *K. rectiphilus rectiphilus*; *K. appendiculatus*; *K. costatus*
- Kalicephalus indicus* (Molin, 1861)
Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, Dobuts. Zasshi, Tokyo, v. 84 (3), 258-261
seasonal differences of infection rate, worm burden
Naja naja: West Bengal villages
- Kalicephalus longior* (Molin, 1861)
Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, Dobuts. Zasshi, Tokyo, v. 84 (3), 258-261
seasonal differences of infection rate, worm burden
Naja hannah: West Bengal villages
- Kalicephalus subulatus* Molin, 1861, illus.
Fernandes, M. P. M.; and Artigas, P. T., 1975, Mem. Inst. Butantan, v. 39, 103-121
Kalicephalus subulatus, key, redescription, valid species, geographical dispersion in serpents, host specificity
Epicrates cenchria cenchria (intestino medio): Vale do Guapore (Estado do Acre); Maraba (Estado do Para)
Boa constrictor constrictor (intestino delgado, intestino medio): Maraba (Estado do Para); Manaus (Estado do Amazonas)
Corallus caninus (intestino medio): Maraba (Estado do Para)
- Kalicephalus willeyi* Linstow 1904
Pinnell, J. L.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 337-340
Vipera russelli: Komodo Island, Indonesia
- Kathlania leptura* (Rud. 1819), illus.
Boero, J. J.; and Led, J. E., 1974, Rev. Agron. y Vet., v. 3 (1), 16-17
description
Thalassochelys caretta (intestino grueso): Jardin Zoologico, La Plata, Argentina (captured near Mar del Plata)
- Kathleena Leiper & Atkinson, 1914
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Contraecaecum* Railliet & Henry, 1912
- Kobusinema Ortlepp*, 1963
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Longistrongylus* Le Roux, 1931
- Kobusinema Ortlepp*, 1963
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus* Le Roux, 1931
- Kobusinema banagiense* Gibbons, 1972
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
as syn. of *Longistrongylus banagiense* (Gibbons, 1972) n. comb.

- Kobusinema schrenki (Ortlepp, 1939) Ortlepp, 1963
 Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
 as syn. of Longistrongylus schrenki (Ortlepp, 1939) n. comb.
- Koriakinema Oshmarin, 1949
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 "too poorly known to be classified but it may be a synonym of Streptocara"
- Krusadia Sanwal, 1952
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 as syn. of Schistorophus Railliet, 1916
- Kuala n. gen.
 Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
 Heligmosomidae, Brevistriatinae
 tod: K. chaii n. gen. n. sp.
- Kuala Durette-Desset et Krishnasamy, 1976
 Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
 Brevistriatinae
 key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Kuala Durette-Desset et Krishnasamy, 1977
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Heligmonellidae, Brevistriatinae
- Kuala chaii n. gen. n. sp. (tod), illus.
 Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
 Tragulus javanicus (intestin grele): Selangor, Subang, Forest Reserve
- Kuala gola (Inglis et Ogden, 1965) n. comb.
 Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
 Syn.: Longistriata gola Inglis et Ogden, 1965

- Labiostomum* sp., *illus.*
Seese, F. M., 1973, *Am. Midland Naturalist*, v. 89 (2), 257-265
key
Ochotona p. princeps (caecum, large intestine): St. Joe Baldy Mountain, Benewah County, Idaho
- Labiostomum coloradensis* Leiby
Grundmann, A. W.; and Lombardi, P. S., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 39-46
Ochotona princeps cinnemomea: Tushar Mountains, Utah
O. p. wasatchensis: Wasatch Mountains, Utah
O. p. uinta: Uinta Mountains, Utah
O. p. fuscipes: Markagunt Plateau, Utah
O. p. barnsei: Fish Lake Mountains, Utah
O. p. nevadensis: Ruby Mountains, Nevada
- Labiostomum coloradensis*
Seese, F. M., 1973, *Am. Midland Naturalist*, v. 89 (2), 257-265
key
- Labiostomum rauschi*
Seese, F. M., 1973, *Am. Midland Naturalist*, v. 89 (2), 257-265
key
- Labiostromylus eugenii*, *illus.*
Bird, A. F., 1976, *Organ. Nematodes (Croll)*, 107-137
skeletal structures in nematodes (copulatory spicules, cuticle, egg shell): structure, chemical composition, ontogeny, function, review
- Labiostromylus eugenii*, *illus.*
Smales, L. R., 1977, *Internat. J. Parasitol.*, v. 7 (6), 449-456
Labiostromylus eugenii, life history: embryogenesis, larval development within egg, hatching process, second and third stage larval morphology and development, optimal temperatures
- Labiostromylus eugenii*, *illus.*
Smales, L. R., 1977, *Internat. J. Parasitol.*, v. 7 (6), 457-461
Labiostromylus eugenii, life history: morphology and development of parasitic stages; failure to establish experimental infections in *Macropus eugenii*
- Labiostromylus eugenii*, *illus.*
Smales, L. R.; and Sommerville, R. I., 1977, *Internat. J. Parasitol.*, v. 7 (3), 205-209
Labiostromylus eugenii, exsheathment, important components of stimulus were pCO₂, pH, and temperature, similar to trichostrongylids
- Lagochilascaris Leiper*, 1909
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridinae
key
- Lagochilascaris minor*
Oostburg, B. F. J., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 580-583
thiabendazole therapy of *Lagochilascaris minor* infection in youth causing painful tumor in neck region, case report: Surinam
- Laimaphelenchus* sp.
Vosilite, B. S., 1975, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Lamanema Becklund*, 1963
Durette-Desset, M. L.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Nematodirinae
- Lamanema chavezii*, *illus.*
Guerrero, C. A.; et al., 1973, *Rev. Invest. Pecuarias*, v. 2 (1), 29-42
Lamanema chavezii, Lama pacos, liver and intestine, pathology; enterohepatic life cycle
- Lamanema chavezii*
Guerrero, C.; Rojas, M.; and Vargas, J., 1974, *Rev. Invest. Pecuarias*, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity of 1-tetramisole, significant body weight gain in treated animals
- Lamanema chavezii*
Leguia, G.; and Bendezu, P., 1974, *Rev. Invest. Pecuarias*, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Lamanema chavezii*
Sillau, H.; et al., 1973, *Rev. Invest. Pecuarias*, v. 2 (1), 103-105
Lamanema chavezii-infected alpacas, liver function tests
- Lamanema chavezii*
Vargas, J.; Guerrero, C.; and Rojas, M., 1972, *Rev. Invest. Pecuarias*, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight toxicity
- Lappetascaridinea* (Rasheed, 1965, subfam.)
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinae
key
includes: *Lappetascaris*
- Lappetascaris Rasheed*, 1965
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Lappetascaridinea
key

- Larva migrans
Bada, J. L., 1971, *Med. Trop.*, Madrid, v. 47 (2), 124-133
clinical review of 100 cases of human cutaneous larva migrans, pathologic findings, body locations of lesions, medical management: Monrovia
- Larva migrans
Bolio Cicero, A.; Zavala Velazquez, J.; and Patron Sanchez, F., 1970, *Prensa Med. Mexicana*, v. 35 (5-6), 218-219
visceral larva migrans, case report of larval granuloma of scrotum in young child: Mexico
- Larva migrans
Cypess, R. H.; and Glickman, L. T., 1976, *Mod. Vet. Pract.*, v. 57 (6), 462-464
zoonosis of visceral larva migrans, dogs to humans, review
- Larva migrans
Dafalla, A. A.; Satti, M. H.; and Abdel Nur, O. M., 1977, *J. Trop. Med. and Hyg.*, v. 80 (3), 63-64
cutaneous larva migrans, preliminary epidemiologic survey shows high prevalence in children especially during rainy seasons, etiology unknown but high incidence of *Strongyloides* spp. in soil samples, suggested control measures: Northern Kordofan, Sudan
- Larva migrans
Gietko, M.; and Zapart, W., 1975, *Pediat. Polska*, v. 50 (1), 61-68
visceral larva migrans in young children with severe eosinophilia, diagnosed by intradermal skin test antigens of *Ascaris lumbricoides* and *Toxocara canis*
- Larva migrans
Harland, P. S. E. G.; Meakins, R. H.; and Harland, R. H., 1977, *Brit. Med. J.* (6089), v. 2, 772 [Letter]
human cutaneous larva migrans, successful use of topical applications of thiabendazole rather than systemic therapy
- Larva migrans
Hayden, D. W.; and Van Kruiningen, H. J., 1973, *J. Am. Vet. Med. Ass.*, v. 162 (5), 379-384
dogs, eosinophilic gastroenteritis occurring simultaneously with visceral larva migrans and manifesting as chronic diarrhea, clinico- and histopathology, case reports, nematode larvae found in lesions of 3 of 5 cases, identified as *Toxocara canis* in 1
- Larva migrans
Kapur, S.; et al., 1976, *Neurol. India*, v. 24 (2), 104-107
human visceral larva migrans, nematode larva demonstrated in brain biopsy specimen of man presenting with severe neurologic impairment and stupor, neurologic symptoms attributed to allergic reaction to parasite presence, case report, biopsy material injected into mice (exper.), nematode larva later observed in mouse lung: India
- Larva migrans
Karel, I.; et al., 1977, *Ophthalmologica*, Basel, v. 174 (1), 14-19
Toxocara larva migrans, woman with granulomatous ocular lesion and active larva in the pupillary area, intolerance to mintezol therapy, case report: Czechoslovakia
- Larva migrans
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Larva migrans
Kim, C. W., 1975, *Progr. Clin. Path.*, v. 6, 267-288
extensive review of techniques used to diagnose human parasitic diseases
- Larva migrans
Lamas, R.; et al., 1970, *Bol. Chileno Parasitol.*, v. 25 (1-2), 74-77
cutaneous larva migrans (possible *Ancylostoma caninum* origin) in city garbage collector successfully treated with thiabendazole, transmission of infection probably from fecal-contaminated material associated with occupation: Santiago, Chile
- Larva migrans
Most, H., 1972, *N. England J. Med.*, v. 287 (10), 495-498; (14), 698-702
common parasitic infections of man encountered in the United States, recommendations for treatment, review
- Larva migrans
Sadana, J. R.; and Kalra, D. S., 1977, *Indian J. Animal Sc.*, v. 44 (11), 1974, 865-868
Stephanurus dentatus, visceral larva migrans, pigs (lungs), pathology: abattoirs at Hissar and Delhi
- Larva migrans
de Savigny, D. H.; and Tizard, I. R., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 501-507
Toxocara larva migrans, larval excretions and secretions from in vitro cultures used as antigen in passive hemagglutination and fluorescent antibody tests to diagnose visceral larva migrans in man and laboratory animals (exper.), preliminary evaluation for serodiagnostic purposes, no cross reactions with *Ascaris suum* infections
- Larva migrans
Styles, T. J.; and Evans, D. S., 1971, *N. York State J. Med.*, v. 71 (23), 2755-2757
visceral larva migrans syndrome present in dogs and cats, area survey, possible transfer to young children, need for public health
- Larva migrans
Thomas, J.; et al., 1969, *Marseille Med.*, v. 106 (9), 717-721
larva migrans, human infection, successful treatment with thiabendazole: France (had returned from Africa)

- Lecanocephalus* Diesing, 1839
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Goezia* Zeder, 1800
- Leidynema appendiculata* (Leidy, 1850) Chitwood, 1932
Hristovski, N. D., 1972, Acta Parasitol. Iugoslavica, v. 3 (2), 109-115
Blatta orientalis: Jugoslaviya (Skopje; Belgrad; Split; Zagreb); Grcija (Solun)
- Leidynema appendiculata*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Blatta orientalis: Macedonia, Yugoslavia
- Leidynema appendiculata*
Leong, L.; and Paran, T. P., 1966, Med. J. Malaya, v. 20 (4), 349
Periplaneta americana
Blatta orientalis
all from Singapore
- Leiperiatus* Sandground, 1930
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Haemonchinae
- Leipernema* n. g.
Singh, S. N., 1976, J. Helminth., v. 50 (4), 267-274
Strongyloididae
tod: *Leipernema leiperi* n. g., n. sp.
- Leipernema leiperi* n. g., n. sp. (tod), illus.
Singh, S. N., 1976, J. Helminth., v. 50 (4), 267-274
Manis pentadactyla (intestine): Hyderabad, A.P., India
- Leiuris* Leuckart, 1850, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascarosinae
key
Syn.: *Spirocercella* Thwaite, 1928
- Leprirophus* Skrjabin, Sobolev & Ivaschkin, 1965
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Torquatoides* (Williams, 1929, subgen.) Inglis, 1965
- Libyostrongylinae* n. sub. fam.
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae
includes: *Libyostrongylus* (type genus); *Cnizostrongylus*; *Obeliscoides*; *Paralibyostrongylus*; *Pararhabdonema*; *Pseudostertagia*
- Libyostrongylus* Lane, 1923 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Libyostrongylinae*
- Litomosa* sp. 1
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Vespertilio murinus: Bellerive, Geneve, Suisse; Col de Jaman, Vaud, Suisse
- Litomosa* sp. 2
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Tadarina teniotis: Col de Bretolet, Valais, Suisse
- Litomosa chiropterorum*
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Miniopterus natalensis: grotte de Baya, Katanga
- Litomosa desportesii* Bain, 1967
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Myotis myotis: grotte de Pertuis, Neuchatel, Suisse
- Litomosa filaria*
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Plecotus auritus: grotte Chaudiere d'Enfer, Vaud, Suisse; Col de Jaman, Vaud, Suisse
P. austriacus: Geneve, Suisse
- Litomosa filaria* Beneden, 1873, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Plecotus auritus
P. austriacus
Myotis myotis
M. daubentoni
M. oxygnathus
M. emarginatus
Rhinolophus ferrumequinum
Miniopterus schreibersi
Vespertilio murinus
(body cavity of all): all from Moldavia
- Litomosa filaria* Beneden, 1873
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Plecotus auritus: Moldavia
- Litomosoides carinii*
Ah, H. S.; and Burke, T. M., 1976, J. Parasitol., v. 62 (4), 651-652
Litomosoides carinii, recovery of substantial numbers of developing 4th stage larvae from pleural cavity of infected jirds 10 and 30 days after exposure to infected *Ornithonyssus bacoti*, subsequent use of these larvae in establishing quantitatively controlled infections in other jirds
- Litomosoides carinii*
Al-Baldawi, F. A. K.; et al., 1976, Parasitology, v. 73 (2), xviii [Abstract]
Litomosoides carinii in protein-deficient cotton rats, immune response assessed by measuring IgG, IgM, and anaphylactic antibody level

- Litomosoides carinii*
Barton, S. P.; Storey, D. M.; and Kershaw, W. E., 1976, *Parasitology*, v. 73 (2), xviii-xix [Abstract]
Litomosoides carinii from protein-deficient cotton rats, little if any detectable difference in structure as compared to worms from well-fed hosts, large structures within uteri of worms from both host groups may be 'abnormal' embryos of other authors
- Litomosoides carinii*
Bingham, A.; et al., 1976, *Parasitology*, v. 73 (2), xxxii-xxxiii [Abstract]
Litomosoides carinii in cotton rats and fast-growing white rats fed a vitamin E deficient diet, plasma enzyme activity, size of worms
- Litomosoides carinii*
Chand, B.; Ramachandran, M.; and Hussain, O. Z., 1977, *Indian J. Exper. Biol.*, v. 15 (8), 667-668
Litomosoides carinii-infected albino rats, respiratory activity of liver, spleen, kidney, and lung tissues, results indicate severe damage to liver and spleen
- Litomosoides carinii*
Emslie, V. W.; and Kershaw, W. E., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (1), 7 [Demonstration]
mice infected with *Trypanosoma brucei*, treated with antrypol and then infected with *Litomosoides carinii*, decreased host resistance to subsequent infection and to relapse from first infection, few visible immunologic changes observed
- Litomosoides carinii*
Enayat, M. S., 1976, *Pahlavi Med. J.*, v. 7 (3), 352-364
Litomosoides carinii, 2 specific antigenic components differentiated using gel diffusion and immunoelectrophoresis
- Litomosoides carinii*
Hendow, H. T.; Storey, D. M.; and Kershaw, W. E., 1976, *Parasitology*, v. 73 (2), xiii [Abstract]
Litomosoides carinii and *Trypanosoma brucei*, hooded rats, effect of combined infections (in some of which trypanosomiasis was cured with Berenil and in some allowed to run its course)
- Litomosoides carinii*
Hutchison, W. F.; et al., 1976, *Comp. Biochem. and Physiol.*, v. 53 (4B), 495-497
Dirofilaria immitis adults, lipid analysis, compared with *Dipetalonema viteae* and *Litomosoides carinii*
- Litomosoides carinii*
Illgen, B.; Wenk, P.; and Seitz, H. M., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 178-179
Litomosoides carinii female in vitro, embryonic development and liberation of microfilariae
- Litomosoides carinii*
Kershaw, W. E.; et al., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (1), 31-32 [Demonstration]
Litomosoides carinii in vitamin A deficient cotton rats and in vitamin E deficient cotton and white rats, antigen extracts from adults and microfilariae; *Nippostrongylus brasiliensis*, mast cell population in lungs of infected rats
- Litomosoides carinii*
Kershaw, W. E.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 11-12 [Demonstration]
Litomosoides carinii, effect of host dietary fat content on growth and development of parasite and cotton rat hosts
- Litomosoides carinii*
Kershaw, W. E.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 11-12 [Demonstration]
Litomosoides carinii, effect of protein-deficient diets on growth and development of parasites and cotton rat hosts
- Litomosoides carinii*
Kershaw, W. E.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 11-12 [Demonstration]
Litomosoides carinii, serum immunoglobulin levels in cotton rats on protein-deficient diets
- Litomosoides carinii*
Kershaw, W. E.; and Storey, D. M., 1976, *Ann. Trop. Med. and Parasitol.*, v. 70 (3), 303-312
Litomosoides carinii infection in cotton rats, comparison of 5 methods of calculating number of infective larvae transmitted to hosts by *Liponyssus bacoti* vectors, relationship between transmission intensities and worms recovered, role of heavy infections in vectors
- Litomosoides carinii*
Kinsella, J. M., 1974, *Am. Mus. Novitates* (2540), 1-12
Sigmodon hispidus (pleural cavity): Florida
- Litomosoides carinii*
Laemmler, G.; and Gruener, D., 1975, *Tropenmed. u. Parasitol.*, v. 26 (3), 359-369
Litomosoides carinii in *Mastomys natalensis* (exper.), 12 organophosphorus compounds tested for efficacy against micro- and macrofilarial infections
- Litomosoides carinii*
Laemmler, G.; Gruener, D.; and Zahner, H., 1975, *Tropenmed. u. Parasitol.*, v. 26 (1), 98-110
Litomosoides carinii in *Mastomys natalensis* (exper.), peripheral blood composition before and after therapy with diethylcarbamazine or suramin (or combination) or HOE 258 V, effects of therapy on pathologic changes

- Litomosoides carinii**
Laemmler, G.; and Herzog, H., 1974, Tropenmed. u. Parasitol., v. 25 (1), 78-83
Litomosoides carinii in Mastomys natalensis, suramin showed high suppressive activity and macrofilaricidal effect on various parasite stages from beginning of infection until end of prepatent period but failed to show prophylactic activity when administered prior to infection, results indicate necessity to confirm larvicidal activity of suramin in Onchocerca volvulus-infected chimpanzees and should initiate chemoprophylactic use of the drug in small clinical trials in human patients
- Litomosoides carinii**
Laemmler, G.; and Wolf, E., 1977, Tropenmed. u. Parasitol., v. 28 (2), 205-225
Litomosoides carinii in Mastomys natalensis (exper.), 17 anthelmintics and chemotherapeutics comparatively tested for their chemoprophylactic activity against various larval stages of the parasites
- Litomosoides carinii**
McCall, J. W., 1976, J. Parasitol., v. 62 (4), 585-588
Litomosoides carinii, simple method for collecting substantial quantities of infective larvae from pelts of Meriones unguiculatus exposed for only a few hours to large number of infected Ornithonyssus bacoti; jirds inoculated with larvae collected in this manner exhibited high mean % recoveries in a narrow range
- Litomosoides carinii**
Mohan, R. N., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (6), 883-884 [Letter]
Litomosoides carinii, white rats, pathological changes
- Litomosoides carinii, illus.**
Nelson, D. S.; et al., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (3), 254-255
Litomosoides carinii-infected rats, pleural exudate cellular morphology
- Litomosoides carinii**
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Litomosoides carinii**
Pringle, G., 1974, Ann. Trop. Med. and Parasitol., v. 68 (2), 205-224
Litomosoides carinii in Praomys (Mastomys) natalensis as laboratory host, course of infection, effects of infection and superinfection on host and on cells of pleural exudate, effect of splenectomy
- Litomosoides carinii**
Rao, Y. V. B. G.; Mehta, K.; and Subrahmanyam, D., 1977, Exper. Parasitol., v. 43 (1), 39-44
Litomosoides carinii, albino rats, effects of irradiation on development of infective larvae, immunization with irradiated infective larvae, immune response in vaccinated animals, effect of irradiation on microfilariae, effect of immunization with irradiated microfilariae
- Litomosoides carinii**
Rew, R. S.; and Saz, H. J., 1977, J. Parasitol., v. 63 (1), 123-129
Litomosoides carinii, Dipetalonema viteae, and particularly Brugia pahangi microfilariae, oxygen requirements, carbohydrate metabolism, effect of levamisole
- Litomosoides carinii**
Saxena, J. K.; et al., 1977, Exper. Parasitol., v. 43 (1), 239-243
Litomosoides carinii, microfilariae and adults, presence and concentration of biogenic amines
- Litomosoides carinii**
Saz, H. J.; and Dumbar, G. A., 1975, J. Parasitol., v. 61 (5), 794-801
stibophen, inhibition of phosphofructokinase and lactate formation, effect on internal hexose phosphate accumulation, inhibition of aldolase, comparison with potassium antimony tartrate (inhibits PFK at higher concentrations but not aldolase)
- Litomosoides carinii**
Sharma, S.; Iyer, R. N.; and Anand, N., 1975, Indian J. Chem., v. 13 (5), 468-472
synthesis of 4,8-dialkyl-1,4,8-triazabicyclo[4.4.0]decan-2-one, 3,7-dialkyl-1,3,7-triazabicyclo[4.3.0]nonan-2-one and related oxygen analogs, no noteworthy activity in Litomosoides carinii, cotton rats
- Litomosoides carinii**
Sharma, S. C.; and Ramachandran, M., 1976, Indian J. Exper. Biol., v. 14 (2), 188-189
Litomosoides carinii-infected albino rats, significantly lower agglutinin titers to sheep erythrocytes
- Litomosoides carinii**
Siddiqui, M. A.; and Kershaw, W. E., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 19-20 [Demonstration]
Litomosoides carinii, experimental laboratory trials show Delhi white rats to be poor hosts and un-irradiated and irradiated golden hamsters highly susceptible hosts when quantitative infections were established using Liponyssus bacoti in transmission
- Litomosoides carinii**
Siddiqui, M. A.; and Kershaw, W. E., 1976, Ann. Trop. Med. and Parasitol., v. 70 (3), 313-322
Litomosoides carinii, laboratory transmission to Delhi and Carworth strains of Rattus norvegicus, relationships between transmission intensity and worm recovery, possible age resistance
- Litomosoides carinii**
Soulsby, E. J. L.; et al., 1976, Pathophysiol. Parasit. Infect., 149-159
Litomosoides carinii in Mastomys natalensis, homocytotropic antibody response, passive and active cutaneous anaphylaxis
- Litomosoides carinii**
Storey, D. M.; and Court, J. P., 1977, Parasitology, v. 75 (2), ix [Abstract]
Litomosoides carinii, host or host-like antigens demonstrated in adult worms and their presence inferred in blood microfilariae

- Litomosoides carinii*
Striebel, H. P., 1976, *Experientia*, v. 32 (4), 457-458
4-isothiocyanato-4'-nitrodiphenylamine, an anthelmintic with an unusual spectrum of activity against intestinal nematodes, filariae and schistosomes
- Litomosoides carinii*
Sturm, P. A.; et al., 1977, *J. Med. Chem.*, v. 20 (10), 1327-1333
Litomosoides carinii in gerbils (exper.), laboratory trials with cis- and trans-1,2-cyclobutanediamines as analogues of diethylcarbamazine showed no activity as antifilarial agents
- Litomosoides carinii*
Sturm, P. A.; et al., 1977, *J. Med. Chem.*, v. 20 (10), 1333-1337
Litomosoides carinii in gerbils (exper.), 3-aminopyrrolidines and 1,4-diazabicyclo [3.2.1]octane derivatives acted as potent microfilaricides but showed no activity against adult worms in laboratory trials comparing activity with that of diethylcarbamazine
- Litomosoides carinii*
Subrahmanyam, D.; et al., 1976, *Nature*, London (5551), v. 260, 529-530
Litomosoides carinii, mechanism of leukocyte adhesion in vitro, mediated by serum factor, accompanied by cytotoxic effect on microfilariae, results implicate both humoral and cellular factors in destruction of microfilariae
- Litomosoides carinii*
Tanaka, H.; et al., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (1), 19-26
Litomosoides carinii in *Sigmodon hispidus* (exper.), suppression of microfilaricidal activity of diethylcarbamazine by anti-lymphocyte and anti-thymocyte serum establishes role of lymphocytes in mechanism of drug action
- Litomosoides carinii*
Turner, A. C.; et al., 1975, *J. Mississippi Acad. Sc.*, v. 20, Suppl., 25
Dirofilaria immitis, dogs, lipid content of worms, total fatty acid methyl esters in *Dirofilaria immitis*, compared with *Dipetalonema viteae* and *Litomosoides carinii*, similar fatty acid patterns, gas chromatography
- Litomosoides carinii*
Wegerhof, P. H.; and Wenk, P., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 179-180
Litomosoides carinii, cotton rats, subcutaneous, intravenous or intra-peritoneal injection of embryonal or blood microfilariae, course of microfilaraemia
- Litomosoides carinii*, illus.
Weiner, D. J.; and Soulsby, E. J. L., 1976, *J. Parasitol.*, v. 62 (6), 886-893
Litomosoides carinii, host response to adult worms intrapleurally or intraperitoneally transplanted into infected vs. naive *Mastomys natalensis*, concluded that preparation period is necessary for successful residence of adult worms
- Litomosoides carinii*
Wenk, P.; and Wegerhof, P. H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 180
Litomosoides carinii, cotton rats, effect of previous injection of microfilariae on challenge infection
- Litomosoides carinii*
Worms, M. J.; and McLaren, D. J., 1977, *Parasitology*, v. 75 (2), xxxvii [Abstract]
Litomosoides carinii in *Meriones* spp., pathogenesis, local response of tissues with special reference to pleural surface
- Litomosoides carinii*
Zahner, H.; et al., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 224
Litomosoides carinii in *Mastomys natalensis*, diethylcarbamazine and haloxon treatment, reduction of microfilaremia, immobilization of microfilariae in various organs
- Litomosoides carolliae venezuelensis* ssp. n.
Gracia Rodrigo, A., 1964, *An. 2. Cong. Latino-Am. Zool.*, v. 1, 261-263
Carollia perspicillata perspicillata (seros- as): El Caura, Edo. Bolivar, Venezuela
- Loa* [sp.], illus.
Poltera, A. A., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (6), 819-829
retrospective study of 40 conjunctival biopsies, portions of nematode identified as belonging to *Loa* group found in 18: Uganda
- Loa loa*
Abdalla, R. E., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (1), 53-55
Wuchereria bancrofti and *Onchocerca volvulus* as cause of human filariasis with hydrocele and elephantiasis, newly reported cases in previously non-endemic areas, need for differentiation from *Loa loa*: Sudan
- Loa loa*
Ambrose-Thomas, P.; and Kien Truong, T., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema viteae*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance
- Loa loa*
Anderson, R. I.; and Buck, A. A., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 447-456
complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad
- Loa loa*
Bendelac, J., 1971, *Maroc Med.* (544), v. 51, 174-175
human infections, case reports, notezine: Khouribga

- Loa loa**
 Burke, G. J., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 402-405
 10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Loa loa, illus.**
 Callihan, T. R.; Oertel, Y. C.; and Mendoza, M., 1977, Am. J. Trop. Med. and Hyg., v. 26 (3), 572-573
 Loa loa microfilaria found in routine cervico-vaginal smear of young woman, possibility of chronic genital filariasis: Washington, D. C. (immigrant from Nigeria)
- Loa loa, illus.**
 Charters, A. D.; et al., 1972, Med. J. Australia, v. 1 (6), 268-271
 Loa loa and Acanthocheilonema perstans adult worms recovered from calabar swellings in persons who previously had been employed in Nigeria, public health importance in possible transfer of disease entity from endemic areas: Western Australia
- Loa loa**
 Duke, B. O. L., 1972, Zool. J. Linn. Soc., London, v. 51, Suppl. 1, 97-107
 Loa loa, behavioral aspects of human and simian strains which have contributed to divergent adaptive evolution with 2 separate host-vector complexes that seldom result in parasite interchange
- Loa loa**
 Fritel, D.; Bariety, J.; and Gentilini, M., 1970, Medecine Afrique Noire, v. 17 (1), 13-15
 kidney pathology in association with Loa loa, possible relationships, case reports
- Loa loa**
 Garin, J. P.; Rougier, J.; and Mojon, M., 1975, Acta Trop., v. 32 (4), 384-388
 Loa loa, human (40 year old missionary), case report, severe ocular manifestations, eventual cure with diethylcarbamazine: Italy (had lived 5 years in Cameroon)
- Loa loa**
 Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, Medecine et Malad. Infect., v. 3 (8-9), 351-353
 diagnostic review of human filariasis
- Loa loa**
 Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, J. Infect. Dis., v. 132 (3), 340-352
 algorithms in the diagnosis and management of human forms of filariases
- Loa loa**
 Gruentzig, J., 1975, Med. Welt., v. 26 (46), 2110-2113
 Loa loa, de Bruy engraving of 1598 presents probable excision of eye worm
- Loa loa**
 Gruentzig, J.; and Jennes, B., 1977, Am. J. Trop. Med. and Hyg., v. 26 (4), 679-683
 reinterpretation of historical references to presence of Loa loa on Island of Ormus in the Persian Gulf
- Loa loa**
 Hagen, H., 1973, Med. Welt, v. 24 (18), 751-752
 Loa loa, human infection acquired while traveling in Cameroun, case report: Germany
- Loa loa**
 Hedge, E. C.; and Ridley, D. S., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (4), 304-307
 evaluation of microfilarial antigen for use with indirect immunofluorescent test in diagnosing human filariasis; best results obtained with sonicated microfilariae of Brugia pahangi with which both cytoplasmic and sheath antigens could be utilized simultaneously
- Loa loa**
 Lucot, J.; and Chovet, M., 1972, Medecine Trop., v. 32 (4), 523-525
 Loa loa, conjunctival manifestations of human loasis
- Loa loa**
 Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
 cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Loa loa**
 Niel, G.; et al., 1972, Medecine et Malad. Infect., v. 2 (5), 193-202
 filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of Setaria labiatopapillosa as antigen, comparison with Dipetalonema vitae and Ascaris suum as antigens
- Loa loa**
 Ogunba, E. O., 1977, Trop. and Geogr. Med., v. 29 (1), 51-55
 Loa loa, survey of Nigerian population shows low microfilaria levels in half of those surveyed, indications that Mansonia africana mosquitoes could become important vector
- Loa loa, illus.**
 Orihel, T. C.; and Lowrie, R. C., jr., 1975, Am. J. Trop. Med. and Hyg., v. 24 (4), 610-615
 Loa loa, entire life cycle maintained in laboratory outside endemic area using Chrysops atlanticus, potential as vectors Chrysops atlanticus (exper.) patas monkeys (exper.)
- Loa loa**
 Orihel, T. C.; and Moore, P. J., 1975, Am. J. Trop. Med. and Hyg., v. 24 (4), 606-609
 Loa loa, baboon and patas monkey successfully infected, possible use as laboratory hosts
 Papio anubis (exper.)
 Erythrocebus patas (exper.)

- Loa loa**
Pinon, J. M.; and Gentilini, M., 1973, *Nouv. Presse Med.*, v. 2 (19), 1283-1287
human filariasis, application of cellular immunologic tests (rosette formation, macrophage migration) in diagnosis and comparison with serologic tests (fluorescent antibody, passive hemagglutination, gel diffusion)
- Loa loa**
Richet, P., 1973, *Medecine Afrique Noire*, v. 20 (11), 899-920
guidelines for mass therapy in human filarial infections, drugs in current use, review
- Loa loa**
Ricosse, J. H.; and Picq, J. J., 1973, *Medecine Afrique Noire*, v. 20 (11), 877-897
human filariasis, clinical indications for treatment, drugs in current use, review
- Loa loa**
Ridley, D. S.; and Hedge, E. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Loa loa**
Rodhain, F.; and Dodin, A., 1971, *Medecine et Malad. Infect.*, v. 1 (4), 185-188
Wuchereria bancrofti, *Loa loa*, variations in human antistreptolysin O titers before and after treatment for filariasis, possible antigenic immune reaction between filariae and *Streptococcus*
- Loa loa**
Rodhain, F.; and Rodhain-Rebourg, F., 1973, *Medecine et Malad. Infect.*, v. 3 (11), 429-436
geographic distribution of *Loa loa* in African population south of the Sahara in equatorial rain-forest
- Loa loa, illus.**
Sacks, H. N.; Williams, D. N.; and Eifrig, D. E., 1976, *Arch. Int. Med.*, Chicago, v. 136 (8), 914-915
Loa loa adult female excised from upper eyelid of Nigerian residing in Minnesota, follow-up therapy with diethylcarbamazine, case report of variations in clinical features from those observed in endemic areas
- Loa loa**
Stuerchler, D.; and Degremont, A., 1976, *Schweiz. Med. Wchnschr.*, v. 106 (20), 682-688
extensive diagnostic and clinical review of filarial parasites frequently encountered by travelers to endemic tropical areas: Switzerland
- Loa loa**
Vedy, J.; Cahuzac, G.; and Labegorre, J., 1975, *Medecine et Armees*, v. 3 (9), 739-746
Loa loa, atypical ocular manifestations in human filariasis, clinical aspects, case reports: Gabon
- Loa loa**
Weiss, N.; and Degremont, A., 1976, *Tropenmed. u. Parasitol.*, v. 27 (3), 377-384
filariasis in persons returning from endemic areas, comparison immunoserologic diagnostic tests (immuno-electrophoresis, indirect fluorescent antibody, indirect hemagglutination, two-dimensional gel diffusion tests) showed that highest sensitivity obtained with immunoelectrophoresis, combined tests gave best results
- Loa loa**
White, G. B., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (2), 161-175
Onchocerca volvulus, *Loa loa*, and *Dipetalonema perstans*, survey of man-biting species of Chrysops, Culicoides and Simulium as potential vectors of human infection: Ethiopia
- Loa loa**
Woodruff, A. W., 1971, *Brit. J. Clin. Pract.*, v. 25 (12), 529-535
achievements in tropical medicine during past 25 years, control, prophylaxis, treatment
- Loa loa loa**
Suswillo, R. R.; et al., 1977, *J. Helminth.*, v. 51 (2), 132-134
failure to experimentally infect *Meriones unguiculatus*
- Loainae gen. sp.** Lubimow, 1926
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 3-6
as syn. of *Sarconema pseudolabiata* nov. sp.
- Lobocephalus** Diesing, 1838, nom. nud.
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Heterocheilus* Diesing, 1839
- Lombricoides** Merat, 1821
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Ascaris* L., 1758
- Longistriata** Schulz, 1926 (? *Neoheligionoides* Sadvovskaja, 1952)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, *Heligmosominae*
- Longistriata** sp.
Anderson, M. M.; and McDaniel, J. S., 1975, *J. Elisha Mitchell Scient. Soc.*, v. 91 (2), 73
Peromyscus leucopus: eastern North Carolina

- Longistriata adunca*
Coggins, J. R.; and McDaniel, J. S., 1975, Proc. Oklahoma Acad. Sc., v. 55, 112-118
helminths of cotton rat, seasonal variation, host size, higher incidence in males, no significant difference in number or kind of parasite in pregnant females
Sigmodon hispidus komareki: Greenville, Pitt County, North Carolina
- Longistriata degusi* n. sp., illus.
Babero, B. B.; and Cattán, P. E., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 68-76
Octodon degus (intestino delgado): Quebrada de la Plata, Santiago, Chile
- Longistriata degusi* (Babero y Cattán, 1975)
Cattán, P. E.; George-Nascimento, M.; and Rodríguez, J., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 16-20
prevalence survey of helminths of *Octodon degus*, seasonal variations, age and sex of hosts: Chile
- Longistriata gola* Inglis et Ogden, 1965
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
as syn. of *Kuala gola* (Inglis et Ogden, 1965) n. comb.
- Longistriata musasabi* Yamaguti, 1941
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
as syn. of *Srivastavanema musasabi* (Yamaguti, 1941) n. comb.
- Longistriata* (B.) *myopotami*
Sadykhov, I. A., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (1), 74-78
influence of ecological factors (age and sex of host, wild or caged animals, season of year) on parasitism
[*Myocastor coypus*]: Azerbaidzhan
- Longistriata noviberiae* Dikmans, 1935
Williams, P. B., jr., 1975, J. Alabama Acad. Sc., v. 46 (3-4), 97 [Abstract]
Longistriata noviberiae, life cycle, domestic rabbit, no intermediate host necessary to complete development
- Longistriata pseudodidas* Vaucher & Durette-Desset, 1973
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
Sorex araneus
S. minutus
(intestino of all): all from Catalan Pyrenean Mountains
- Longistriata wolgensis* Schulz, 1926
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Arvicola terrestris
Microtus agrestis
(small intestine of all): all from Karelia
- Longistrongylea*
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
tribe cannot be accepted
- Longistrongylus* Le Roux, 1931
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae* synonymy
- Longistrongylus* Le Roux, 1931
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
Ostertagia, *Trichostrongylinae* revision; syn.: *Bigalkenema* Ortlepp, 1973; *Kobusinema* Ortlepp, 1963; key to species
- Longistrongylus albifrontis* (Monnig, 1931) Travassos, 1937, illus.
Gibbons, L. M.; 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Antidorcas marsupialis (abomasum): Theunissen
- Longistrongylus banagiense* (Gibbons, 1972) n. comb., illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Syn.: *Kobusinema banagiense* Gibbons, 1972
Damaliscus korrigum: Serengeti, Tanzania
Aepyceros melampus: Kenya
Alcelaphus buselaphus cokei: Kenya
Gazella granti: Kenya
G. thomsonii: Kenya; Serengeti region, Tanzania
Oryx gazella: Kenya
Redunca sp.: Kenya; Serengeti region, Tanzania
Alcelaphus buselaphus jacksoni: Uganda (abomasum of all)
- Longistrongylus curvispiculum* (Gibbons, 1973) n. comb., illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Syn.: *Bigalkenema curvispiculum* Gibbons, 1973
Gazella granti (abomasum): Serengeti region, Tanzania; Kenya
Aepyceros melampus: Serengeti region, Tanzania
Damaliscus korrigum: Serengeti region, Tanzania
Connochaetes taurinus: Serengeti region, Tanzania
Gazella thomsonii: Serengeti region, Tanzania; Kenya
- Longistrongylus curvispiculum*
Gibbons, L. M.; and Khalil, L. F., 1977, J. Helminthol., v. 51 (3), 209-210
Oryx tao (abomasum): Marwell Zoological Park, England

- Longistrongylus meyeri* Le Roux, 1931, illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Alcelaphus caama: South West Africa
Aepyceros melampus (abomasum): Serengeti region, Tanzania
Connochaetes taurinus (abomasum): Serengeti region, Tanzania
Gazella thomsonii (abomasum): Serengeti region, Tanzania
G. granti (abomasum): Kenya
Alcelaphus buselaphus (abomasum): Kenya
unknown host: Kenya
- Longistrongylus meyeri* Le Roux, 1931
Gibbons, L. M.; and Khalil, L. F., 1976, Trop. Animal Health and Prod., v. 8 (3), 168
goat (gut): Kajiado district, Kenya
- Longistrongylus meyeri* Le Roux
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Longistrongylus muraschkinzevi* (Shulz and Kadentsii, 1950) Jansen, 1958
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
species inquirenda
- Longistrongylus namaquensis* (Ortlepp, 1963) n. comb., illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Syn.: Bigalkenema namaquensis Ortlepp, 1963
Ovis aries (abomasum): Namakwaland, South Africa
- Longistrongylus sabie* (Monnig, 1932) n. comb., illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Syn.: Bigalkenema sabie (Monnig, 1932) Ortlepp, 1963
Aepyceros melampus (duodenum, pylorus, abomasum): Serengeti region, Tanzania; Kenya
Gazella thomsonii (abomasum): Kenya
- Longistrongylus schrenki* (Ortlepp, 1939) n. comb., illus.
Gibbons, L. M., 1977, J. Helminth., v. 51 (1), 41-62
redescription, key, summary of hosts and geographic distribution
Syn.: Kobusinema schrenki (Ortlepp, 1939) Ortlepp, 1963
Kobus kob (abomasum): West Acholi District, Uganda; Semliki, Uganda; Tanzania
Ourebia ourebi: West Acholi District, Uganda
Kobus defassa (abomasum): Semliki, Uganda
Kobus sp. (abomasum): Kenya; Serengeti region, Tanzania
Redunca sp. (abomasum): Kenya
Kobus vardoni (abomasum): Serengeti region, Tanzania
- Loxodontofilaria* [? n. rank]
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of Dipetalonema; key
tod: D. (L.) loxodontis (Berghe et Gillain, 1939)
- Lungworms
Nelson, M., 1977, Vet. Rec., v. 101 (12), 248 [Letter]
lungworms, winter survival, evidence for possible migration into and overwintering in soil
- Lungworm
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Alcelaphus buselaphus cokei: Kenya
- Lutzinema Lent et Freitas, 1934
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Ornithostromylinae

- Macdonaldius Khanna*, 1933
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
key
Syn.: *Saurofilaria Caballero*, 1954
- Maciela* [sic] Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Mackerrastrongylinae* Inglis, 1968
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae
includes: *Mackerrastrongylus* (type genus); *Asymmetracantha*; *Filarinema*; *Peramelistrongylus*; *Tetrabothriostongylus*; *Woodwardostongylus*
- Mackerrastrongylus* Mawson, 1960 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae, *Mackerrastrongylinae*
- Macropostrongylus* Yorke & Maplestone
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
Trichonematidae
revised generic diagnosis, key to species (excluding *M. irma*)
- Macropostrongylus* Yorke & Maplestone p.p.
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Popovastrongylus* n. gen.
- Macropostrongylus australis* Yorke & Maplestone
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Cloacina australis* (Yorke & Maplestone) [n. comb.]
- Macropostrongylus baylisi* Wood, 1930, to *Macropostrongyloides* [comb. not made]
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
- Macropostrongylus dissimilis* Johnston & Mawson, 1939
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Arundelia dissimilis* (Johnston & Mawson) n. comb.
- Macropostrongylus dorcopsis* Baylis, 1940
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Dorcopsinema dorcopsis* (Baylis) n. comb.
- Macropostrongylus irma* Johnston & Mawson, 1940
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
Syn.: *Gelanostrongylus irma*: Popova, 1952
- Macropostrongylus lesouefi* Johnston & Mawson, 1939, *illus.*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
key, description, syn.: *Gelanostrongylus lesouefi*: Popova, 1952
- Macropostrongylus macropostrongylus* Yorke & Maplestone, *illus.*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
key, description
Macropus agilis
Thylogale brunii
all from Weam, Papua New Guinea
- Macropostrongylus macrostoma* Davey & Wood, 1938, *illus.*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
synonymy, key, description
- Macropostrongylus pearsoni* Johnston & Mawson, 1940
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Popovastrongylus pearsoni* (Johnston & Mawson) [n. comb.]
- Macropostrongylus wallabiae* Johnston & Mawson, 1939
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Popovastrongylus wallabiae* (Johnston & Mawson) [n. comb.]
- Macropostrongylus yorkei* Baylis, 1927, *illus.*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
key, description
Macropus agilis (stomach): Tipperary Stn., N. T.
- Macropostrongylus yorkei* (non Baylis): Johnston & Mawson, 1939
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (2-4), 51-62
as syn. of *Macropostrongylus macrostoma* Davey & Wood, 1938
- Madochotera* Bain et Brunhes, 1968
Bain, O.; and Prod'hon, J., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 721-739
Onchocercidae; Waltonellinae n. subfam.
- Madochotera landauae* Prod'hon et Bain, 1973
Bain, O.; and Prod'hon, J., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 721-739
- Madochotera pichoni* n. sp., *illus.*
Bain, O.; and Prod'hon, J., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 721-739
Racophorus sp. (cavite generale): Andringitra, Madagascar
- Malayocamallanus* Jothy & Fernando, 1971
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanidae
key
- Mammanidula* Sadovskaja, 1952
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, Nippostrongylinae
synonymy

- Mammomonogamus laryngeus*
Euzéby, J.; et al., 1977, Bull. Acad. Vet. France, v. 50 (2), 267-273
cattle and man, review: Central America; Antilles
- Mammomonogamus nasicola*
Euzéby, J.; et al., 1977, Bull. Acad. Vet. France, v. 50 (2), 267-273
cattle in South America and Antilles, review; unsuccessful attempt to study life cycle
bovins (larynx, pharynx): Guadeloupe
mouton (larynx): Martinique
- Mammomonogamus nasicola*
Magdeleine, J.; et al., 1974, Medecine Afrique Noire, v. 21 (8-9), 651-655
Mammomonogamus nasicola, human infections, clinical findings, diagnosis by bronchoscopy, possible reservoir hosts, epidemiology, relative frequency in Martinique
- Manistrongylus Baer*, 1959
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Trichocheenia Kou*, 1958
- Manistrongylus Cameron et Myers*, 1960
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Trichocheenia Kou*, 1958
- Mansonella ozzardi*, illus.
Beaver, P. C.; Neel, J. V.; and Orihel, T. C., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 263-265
Indians (blood): southern Venezuela
- Mansonella ozzardi*
Boorman, J.; and Mellor, P. S., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 438 [Demonstration]
Mansonella ozzardi, technique for infecting small insects as *Aedes aegypti*, *Culicoides riethi* and *C. variipennis* for laboratory study
- Mansonella ozzardi*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, Medecine et Malad. Infect., v. 3 (8-9), 351-353
diagnostic review of human filariasis
- Mansonella ozzardi*
Mellor, P. S., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 353 [Letter]
Culicoides variipennis (exper.)
Culicoides nubeculosus (exper.)
Culicoides riethi (exper.)
Aedes aegypti (exper.)
- Mansonella ozzardi*
Miller, M. J.; Ratard, R. C.; and McNeeley, D. F., 1976, J. Parasitol., v. 62 (5), 845-847
Wuchereria bancrofti, human, nocturnal microfilarial periodicity; presence of *Mansonella ozzardi* also reported: Haiti
- Mansonella ozzardi*
Moraes, M. A. P., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 16 [Demonstration]
Mansonella ozzardi in humans, diagnosis comparing microfilariae revealed in bloodless skin snips and in blood: Brazil
- Mansonella ozzardi*
Nelson, G. S.; and Davies, J. B., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 16-17 [Demonstration]
Mansonella ozzardi, humans, epidemiologic survey, concentration of microfilariae in superficial capillaries, mixed infections with *Wuchereria bancrofti* differentiated using stained filters: Trinidad
- Mansonella ozzardi*
Raccurt, C.; and Hodges, W., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (5), 452-453 [Letter]
human epidemiologic survey of filariasis in Haiti
- Mansonella ozzardi* (Manson, 1897)
Sasa, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 3-48
human filariasis in the Americas, extensive review, epidemiology, geographic distribution, mosquito vectors, control measures, literature review
- Mansonella ozzardi*
Shelley, A. J., 1975, Ann. Trop. Med. and Parasitol., v. 69 (3), 407-412
Mansonella ozzardi, human, epidemiological survey: rural communities on river Purus, state of Amazonas, Brazil
- Mansonella ozzardi*
Shelley, A. J.; and Shelley, A., 1976, Ann. Trop. Med. and Parasitol., v. 70 (2), 213-217
Mansonella ozzardi, conformation of *Simulium amazonicum* (nat. and exper.) as vector: Brazil
- Mansonella ozzardi*
Suswillo, R. R.; et al., 1977, J. Helminth., v. 51 (2), 132-134
failure to experimentally infect *Meriones unguiculatus*
- Marimermis* gen. n. (type genus)
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
Marimermithidae fam. n., tod: *M. maritima* sp. n.
- Marimermis kergelensis* sp. n., illus.
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
Hippasteria hyadesi: Kerguelen Islands
- Marimermis litoralis* sp. n., illus.
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
host unknown: Kerguelen Islands

- Marimermis maritima* sp. n. (tod), illus.
Rubtsov, I. A.; and Platonova, T. A., 1974,
Zool. Zhurnal, v. 53 (10), 1445-1458
host unknown: Pacific Ocean coast, Simushir
Island, Roka Cape
- Marimermithidae* fam. n.
Rubtsov, I. A.; and Platonova, T. A., 1974,
Zool. Zhurnal, v. 53 (10), 1445-1458
Enoplida
includes: *Marimermis* gen. n. (type genus);
Trophomera gen. n.; *Thalassonema* Ward, 1933
- Marshallagia* (Orloff, 1933) Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- [*Marshallagia*] *marshalagii*
Nurtazin, A. T., 1975, *Vestnik Sel'skokhoz.*
Nauki Kazakhstana (4), 84-86
sheep, trichostrongyles, anthelmintic ef-
ficacy of banminth, good results
- Marshallagia marshalli* Orloff, 1933
Bergstrom, R. C., 1975, *Proc. Helminth. Soc.*
Washington, v. 42 (1), 61-63
Marshallagia marshalli, other trichostrongy-
lylids, incidence and intensity in Ovis
aries under 3 types of management (farm
flocks, fenced range, and seasonally herded
range) and in wild *Antilocapra americana*;
evidence for transmission of *M. marshalli*
from antelope to sheep where range is shared:
Wyoming
- Marshallagia marshalli* (Orlov, 1933)
Bergstrom, R. C., 1975, *Proc. Oklahoma Acad.*
Sc., v. 55, 101-102
elk (feces)
mule deer (feces)
Antilocapra americana (feces, gastrointesti-
nal tract)
Ovis canadensis (feces, gastrointestinal
tract)
all from Wyoming
- Marshallagia marshalli*
Bergstrom, R. C.; Maki, L. R.; and Werner,
B. A., 1976, *Proc. Helminth. Soc. Washington*,
v. 43 (2), 171-174
trichostrongylid eggs in cattle or sheep
feces, dung beetles (*Aphodius* spp.; *Canthon*
practicola) as possible biological control
agents, laboratory studies showed decreased
eggs in feces when beetles were present
- Marshallagia marshalli*
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*,
v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenben-
dazole, satisfactory results: Iran
- Marshallagia marshalli*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1),
23-30
ethyl-6-ethoxybenzothiazole-2-carbamate,
evaluation of anthelmintic activity in pon-
ies, swine, lambs, and chickens
- Mastigonema* gen. n.
Dailey, M. D.; and Perrin, W. F., 1973, *Fish.*
Bull., National Oceanic and Atmos. Admin.,
v. 71 (2), 455-471
Spiuroidea, Ascaropsinae; mt: *M. stenellae*
sp. n.
- Mastigonema stenellae* sp. n. (mt), illus.
Dailey, M. D.; and Perrin, W. F., 1973, *Fish.*
Bull., National Oceanic and Atmos. Admin.,
v. 71 (2), 455-471
Stenella graffmani
S. cf. *S. longirostris*
(forestomachs of all): all from eastern
tropical Pacific
- Mastophorinae Quentin, 1970
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Spiroceridae
key
includes: *Mastophorus*
- Mastophorus* Diesing, 1853, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Mastophorinae
- Mastophorus*
Dyer, W. G., 1972, *Tr. Illinois Acad. Sc.*, v.
65 (1-2), 23-25
Mastophorus numidica, morphology, presence
or absence of pseudolabial teeth, shape of
stoma and form of spicules are most reliable
characters for separating species of *Masto-*
phorus
- Mastophorus-Protospirura* spp.
Singh, M.; and Cheong Chee Hock, 1971, *South-*
east Asian J. Trop. Med. and Pub. Health, v. 2
(4), 516-521
Rattus rattus argentiventer
R. r. rumpia
R. bowersi
R. canus
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
R. whiteheadi
all from Malaysia
- Mastophorus dipodomis* Read and Millemann, 1953
King, S. R.; and Babero, B. B., 1974, *Proc.*
Helminth. Soc. Washington, v. 41 (2), 241-248
Dipodomys merriami: Nevada
- Mastophorus muris*, illus.
Beaucournu, J. C.; and Deunff, J., [1976],
Ann. Parasitol., v. 50 (6), 1975, 831-835
Ctenophthalmus baeticus arvernus (cavite
generale): France
- Mastophorus muris*
Garner, H. W.; Richardson, L. W.; and Felts,
L. A., 1976, *Southwest. Nat.*, v. 21 (3), 327-
334
monthly percentages of animals parasitized
Dipodomys ordii (stomach): western Texas

- Mastophorus muris*, encapsulated third stage larva
Killick-Kendrick, R.; et al., 1976, Tr. Roy.
Soc. Trop. Med. and Hyg., v. 70 (1), 22 [Demon-
stration]
Phlebotomus ariasi: Gard, southern France
- Mastophorus muris*
Kinsella, J. M., 1974, Am. Mus. Novitates
(2540), 1-12
Sigmodon hispidus (stomach): Florida
- Mastophorus muris*
Mishra, G. S.; and Gonzalez, J. P., 1975,
Arch. Inst. Pasteur Tunis, v. 52 (1), 71-87
partial life cycle study, experimental de-
velopment in domestic cat unsuccessful
Rattus norvegicus (estomac): Tunisia
Blatella germanica (exper.)
- Mastophorus muris*
Seureau, C.; and Quentin, J. C., 1977, Ann.
Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subu-
lurid and spirurid nematodes in *Locusta mi-*
gratoria (exper.), course and duration of
migration, histopathologic consequences,
brief discussion of relation to phylogeny
of nematodes and host hemocytic defense
reaction
- Mastophorus muris*
Shakhmatova, V. I., 1966, Trudy Gel'mint.
Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes: Karelia
- Mastophorus muris*, *illus.*
Wertheim, G.; and Chabaud, A. G., 1977, Ann.
Parasitol., v. 52 (6), 647-657
- Mastophorus muris ascaroides*
Coggins, J. R.; and McDaniel, J. S., 1975,
Proc. Oklahoma Acad. Sc., v. 55, 112-118
helminths of cotton rat, seasonal variation,
host size, higher incidence in males, no
significant difference in number or kind
of parasite in pregnant females
Sigmodon hispidus komareki: Greenville,
Pitt County, North Carolina
- Mastophorus numidica* Seurat 1914
Bienek, G. K.; and Klikoff, L. G., 1974, Am.
Midland Naturalist, v. 91 (1), 251-253
Dipodomys merriami vulcani: Dixie State
Park, Washington Co., Utah
- Mastophorus numidica* (Seurat, 1914) Read and
Millemann, 1953
Dyer, W. G., 1972, Tr. Illinois Acad. Sc., v.
65 (1-2), 23-25
Mastophorus numidica, morphology, presence
or absence of pseudolabial teeth, shape of
stoma and form of spicules are most reliable
characters for separating species of *Masto-*
phorus
- Mazamostrongylus* Cameron, 1935
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Mazzia Khalil & Vogelsang, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
"cannot be classified . . . probably a valid
genus of Spirocerinae"
- Mecistocirrus* Railliet et Henry, 1912
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Haemonchinae
- Mecistocirrus*
Qadir, A. N. M. A., 1976, Indian Vet. J., v.
53 (11), 855-858
gastrointestinal nematodes, goats and
calves, urea for pasture control of free-
living stages
- Mecistocirrus* sp.
Schneider, C. R.; et al., 1975, Ann. Trop.
Med. and Parasitol., v. 69 (2), 227-232
Bubalus bubalis: Khong Island, Laos
- Mecistocirrus digitatus* (von Linstow, 1906)
Railliet & Henry, 1912
Gaur, S. N. S.; and Dutt, S. C., 1977, Indian
J. Animal Sc., v. 45 (7), 1975, 446-450
Mecistocirrus digitatus, exper. infection of
cattle and sheep, seasonal incidence in cat-
tle and buffaloes: India
- Megacooperia* Khalil et Gibbons, 1976
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Cooperiinae
- Mehdiella inflatocervix* Akhtar, 1937, *illus.*
Hanuskova, Z.; and Tilc, K., 1975, Acta Vet.
Brno, v. 44 (4), 407-412
nematodes, incidence in *Agrionemys horsfieldi*
(intestine) with regard to unsuitable con-
ditions and food: Czechoslovakia, imported
from USSR
- Mehdiella microstoma* Drasche, 1884
Hanuskova, Z.; and Tilc, K., 1975, Acta Vet.
Brno, v. 44 (4), 407-412
nematodes, incidence in *Agrionemys horsfieldi*
(intestine) with regard to unsuitable con-
ditions and food: Czechoslovakia, imported
from USSR
- Mehdiella uncinata*
Hristovski, N. D., 1973, Acta Parasitol. Iugo-
slavica, v. 4 (2), 87-91
Testudo graeca
Testudo haermani
all from Macedonia, Yugoslavia
- Mermis* sp.
Buettiker, W.; and Nicolet, J., 1975, Rev.
Elevage et Med. Vet. Pays Trop., n. s., v. 28
(3), 319-329
Arcyophora patricula: Minankro pres de
Bouake

- Mermis sp.
Jakhmola, S. S.; and Yadav, H. S., 1975,
Indian J. Entom., v. 35 (2), 170-172
Mermis sp., degree of parasitism correlated
with rainfall, biological control of cater-
pillars
Antigastra catalaunalis (haemocoele):
Madhya Pradesh, India
- Mermis sp., illus.
Sharma, V. K.; Singh, J. M.; and Chaudhary,
R. N., 1976, Pantnagar J. Research, v. 1 (2),
144-145
Chilo partellus (haemocoel): India
- Mermis myrmecophila Baylis, to Pheromermis n.
gen. [comb. not made]
Poinar, G. O., jr.; Lane, R. S.; and Thomas, G.
M., 1976, Nematologica, v. 22 (3), 360-370
- Mermis nigrescens
Condon, W. J.; and Gordon, R., 1977, Canad. J.
Zool., v. 55 (4), 690-692
Mermis nigrescens in Locusta migratoria,
hemolymph uric acid increased, fecal uric
acid decreased, host protein turnover
- Mermis nigrescens Duj.
Denner, M. W., 1976, Proc. Indiana Acad. Sc.,
v. 85, 1975, 406 [Abstract]
Mermis nigrescens, oviposition may occur in
absence or presence of sunlight, presence
of haemoglobin may not act as chromotropic
stimulus for egg laying
- Mermis nigrescens
Denner, M. W., 1976, Proc. Indiana Acad. Sc.,
v. 85, 1975, 406-407 [Abstract]
Mermis nigrescens, grasshoppers, strong host
preference not indicated
Cyrthacanthacridinae
Oediponinae
Conocephalinae
- Mermis paranigrescens sp. n., illus.
Rubtsov, I. A., 1976, Zool. Zhurnal, v. 55 (9),
1292-1298
host unknown: Chinkentsk obl., Dzhebagly
preserve, Kish Kaiandy
- Mermithid larva
Barrett, T. V., 1977, Tr. Roy. Soc. Trop. Med.
and Hyg., v. 71 (5), 382 [Demonstration]
Triatoma sordida: Bahia, Brazil
- Mermithid, illus.
Poinar, G. O., jr., 1976, J. Parasitol., v. 62
(5), 843-844
infection in Limnodrilus silvani (intestinal
tissue) represents paratenic host in life
cycle
- Mermithid
Wuelker, W., 1976, Ztschr. Parasitenk., v. 50
(2), 196-197
mermithid-infected Chironomus (exper.), in-
tersexual patterns of adult antennae
- Mermithidae
Charnov, E. L.; and Bull, J., 1977, Nature,
London (5605), v. 266, 828-830
environmental sex determination, hypothesis
with mention of Mermithidae and parasitic
copepods as examples
- Mermithidae
Ross, J. F.; and Smith, S. M., 1976, Canad. J.
Zool., v. 54 (7), 1084-1102
key to adults of mermithid genera infecting
mosquitoes in North America; comments on
useful characters in taxonomy
- Mermithidae [sp.]
Carter, J. B., 1976, J. Applied Ecol., v. 13
(1), 103-122
Mermithidae as possible biological control
agent of Tipula spp.
Tipula paludosa
T. marmorata or alpium?
Ptychoptera (albimana?)
all from north-east England
- Mermithidae [sp.], illus.
Dundee, D. S., 1977, J. Parasitol., v. 63 (3),
590
Veronicella ameghini (body cavity): Mobile,
Alabama
- Mermithidae [sp.]
Garris, G. I.; and Noblet, R., 1975, J. Med.
Entom., v. 12 (4), 481-482
Mermithidae and Microsporidia as possible
natural control agents of Simuliidae
Simulium venustum
S. verecundum
S. tuberosum
Simulium sp.
Cnephia sp.
all from South Carolina
- Mermithidae n. gen., n. spp.
Mondet, B.; Pendriez, B.; and Bernadou, J.,
1976, Cahiers O.R.S.T.O.M., s. Entom. Med.,
v. 14 (2), 141-149
Simulium damnosum
S. vorax
S. adersi
S. alcocki
S. cervicornutum
S. unicornutum
all from Coted'Ivoire (riviere Mounongo)
- Mermithidae [spp.]
Rubzov, I. A., 1969, Acta Parasitol. Polon.,
v. 16 (1-19), 1968-1969, 97-100
Elasmucha betula: Siberia
Eurygaster austriaca: Bulgaria
- Mermithides
Beaucournu, J. C.; and Deunff, J., [1976],
Ann. Parasitol., v. 50 (6), 1975, 831-835
as hyperparasites of fleas, causing para-
sitic castration
siphonapteres: France

- Mermithoidea**
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Mesomermis bilateralis** sp. n., *illus.*
Rubtsov, I. A., 1976, Zool. Zhurnal, v. 55 (9), 1292-1298
host unknown: Komi ASSR, r. Shchugor
- Mesomermis melusinae** sp. nov., *illus.*
Rubtsov, I. A., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 128-156
ontogenesis of mermithids, illustrated description of structure of body, cuticle, amphids, longitudinal fields, stichosome, osmosome, trophosome and reproductive organs; technique of preparing material
[Simuliidae]: Luzhsk region, Leningrad oblast
- Mesomermis subtilis** Coman, *illus.*
Rubtsov, I. A., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 128-156
ontogenesis of mermithids, illustrated description of structure of body, cuticle, amphids, longitudinal fields, stichosome, osmosome, trophosome and reproductive organs; technique of preparing material
- Mesopectines**, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of Pterygodermatites
key
- Mesorhabditis spiculigera** (Steiner, 1936), Dougherty, 1953, *illus.*
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972, Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera beetles, possible importance in biological control
Scarabaeus sacer: Abu-Rawash, Cairo, Egypt
- Metabronema Yorke & Maplestone**, 1926, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Metabronema salvelini** (Fujita, 1922)
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Pogonichthys macrolepidotus (intestine): Sacramento-San Joaquin Delta, California
- Metacyrnea** (Chabaud, 1960, subgen.), *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
- Metangusticaecum** Mozgovi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of Terranova Leiper & Atkinson, 1914
- Metanisakis** Mosgovi, 1950
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of Pseudanisakis Layman & Borovkova, 1926
- Metanisakis** Mozgovi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Acanthocheilidae
key
Syn.: Pseudanisakis Yamaguti, 1941, *nec* Layman & Borovkova, 1926
- Metanisakis rajae** (Yamaguti, 1941) *of* Mosgovi (1950, 1953)
Gibson, D. I., 1973, J. Nat. Hist., v. 7 (3), 319-340
as syn. of Pseudanisakis rajae Yamaguti, 1941, *sensu nov.*
- Metascaris** Schuurmans-Stekhoven, 1950
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
"excluded from the key since . . . incompletely described"
- Metastrongyloidea**
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Strongylida
key
- Metastrongylus** spp.
Barratt, M. E. J., 1972, Immunology, v. 22 (4), 615-623
Metastrongylus spp., pigs, immediate hypersensitivity, partial characterization of allergens, suggested that cross reactions so commonly found when using nematode antigens in wheal and erythema reactions can be eliminated by suitable dilution of the allergen
- Met[aststrongylus]** sp.
Getler, K., 1972, Med. Wet., v. 28 (8), 476-477
nematodes, pigs on industrial swill feeding farm, Atgard
- Metastrongylus** spp.
Poeschel, G. P.; and Emro, J. E., 1972, J. Am. Vet. Med. Ass., v. 160 (12), 1637-1640
Metastrongylus spp., hogs (lungs), levamisole hydrochloride and levamisole resinate, good results
- Metastrongylus** spp.
Preston, K. S.; and Switzer, W. P., 1976, Vet. Microbiol., v. 1 (1), 15-18
failure of Metastrongylus spp.-infected earthworms to transmit mycoplasmal pneumonia to swine; failure of Metastrongylus to produce antibody titers against mycoplasma pneumonia or to produce pneumonic lesions

- Metastrongylus* spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Metastrongylus apri*
Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, Vet. Rec., v. 99 (7), 119-122
swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with pica-dex
- Metastrongylus apri*
Barratt, M. E. J., 1972, Immunology, v. 22 (4), 601-614
Metastrongylus spp., pigs, immediate hypersensitivity, production and partial characterization of homocytotropic antibody, passive transfer of skin sensitivity to uninfected recipients, homocytotropic activity closely associated with but does not parallel distribution of IgA and may be mediated by another immunoglobulin
- Metastrongylus apri*
Bussieras, J., 1976, Rec. Med. Vet., v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Metastrongylus apri*
Chatterjee, A.; Das, S. K.; and Bhattacharyya, H. M., 1975, Indian J. Animal Health, v. 14 (2), 167-168
Metastrongylus apri, *M. salmi*, giant cell pneumonia, histopathology
pigs (diaphragmatic lobes of lungs): West Bengal
- Metastrongylus apri*
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
Sus scrofa domesticus x *Sus scrofa cristatus* (lungs): Aransas National Wildlife Refuge, southern Texas
- Metastrongylus apri*
Duewel, D.; Hajdu, P.; and Damm, D., 1975, Berl. u. Munchen. Tierarztl. Wchnschr., v. 88 (21), 418-419
Metastrongylus apri, pigs, fenbendazol, therapeutic dosage, serum levels, fast elimination and low levels required for effectiveness
- Metastrongylus apri*
Estudillo C., J. J., 1976, Veterinaria, Mexico, v. 7 (1), 15-16
Metastrongylus apri, *M. pudendotectus*, pigs (lungs), sex ratio, role of earthworm intermediate host in parasite prevalence and distribution: slaughterhouse in Veracruz, Mexico
- Metastrongylus apri*
Koga, M.; et al., 1976, Japan. J. Vet. Sci., v. 38 (6), 611-618
Metastrongylus apri, thymectomized guinea pigs exposed to whole-body X-irradiation or anti-thymocyte serum, vaccination and challenge, results suggest important role for T-cells in defense mechanism
- Metastrongylus apri*
Krvavica, S.; et al., 1976, Vet. Arhiv, Zagreb, v. 46 (11-12), 271-287
Metastrongylus apri, activity of enzymes taking part in glucose fermentation; aerobic metabolism; tricarboxylic acid cycle
- Metastrongylus apri*
Krvavica, S.; Francetic, D.; and Zivkovic, D., 1976, Vet. Arhiv, Zagreb, v. 46 (9-10), 231-239
nematodes, trematodes, cestodes, activity, distribution and cofactor dependence of malic enzymes; majority are located in mitochondria in all investigated parasites
- Metastrongylus apri, illus.*
Kumar, V.; and Mortelmans, J., 1974, Riv. Parasitol., Roma, v. 35 (2), 149-151
Metastrongylus apri, occurrence of 2 distinct morphological forms of embryonated eggs, differential hatching behavior
- Metastrongylus apri*
Kumar, V.; and Mortelmans, J., 1976, Parasitology, v. 72 (1), 13-18
Metastrongylus apri, guinea pigs, levamisole-terminated prepatent infection, stimulation of strong immunity to challenge, increase in serum gamma-globulin levels
- Metastrongylus apri*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile
- Metastrongylus elongatus*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Metastrongylus elongatus* (Dujardin, 1846) Raillet et Henry, 1911
Sobieszewski, K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 91-95
Metastrongylus elongatus, *M. pudendotectus*, pigs (lungs), incidence, mixed infection, pathological changes: Lublin Palatinate, Poland
- Metastrongylus elongatus*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (26), 123-128
[pig, wild]: Primorskii krai

- Metastrongylus pudendotectus*
Barratt, M. E. J., 1972, *Immunology*, v. 22 (4), 601-614
Metastrongylus spp., pigs, immediate hypersensitivity, production and partial characterization of homocytotropic antibody, passive transfer of skin sensitivity to uninfected recipients, homocytotropic activity closely associated with but does not parallel distribution of IgA and may be mediated by another immunoglobulin
- Metastrongylus pudendotectus*
Bussieras, J., 1976, *Rec. Med. Vet.*, v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Metastrongylus pudendotectus*
Coombs, D. W.; and Springer, M. D., 1974, *J. Wildlife Dis.*, v. 10 (4), 436-441
Sus scrofa domesticus x *Sus scrofa cristatus* (lungs): Aransas National Wildlife Refuge, southern Texas
- Metastrongylus pudendotectus*
Corwin, R. M., 1977, *Am. J. Vet. Research*, v. 38 (4), 465-467
mixed nematode infections, pigs, oxfendazole, critical evaluation: Missouri
- Metastrongylus pudendotectus*
Estudillo C., J. J., 1976, *Veterinaria, Mexico*, v. 7 (1), 15-16
Metastrongylus apri, *M. pudendotectus*, pigs (lungs), sex ratio, role of earthworm intermediate host in parasite prevalence and distribution: slaughterhouse in Veracruz, Mexico
- Metastrongylus pudendotectus*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Sus scrofa: Chile
- Metastrongylus pudendotectus* Wostokow, 1905
Sobieszewski, K., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 91-95
Metastrongylus elongatus, *M. pudendotectus*, pigs (lungs), incidence, mixed infection, pathological changes: Lublin Palatinate, Poland
- Metastrongylus pudendotectus*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii krai
- Metastrongylus salmi*
Chatterjee, A.; Das, S. K.; and Bhattacharyya, H. M., 1975, *Indian J. Animal Health*, v. 14 (2), 167-168
Metastrongylus apri, *M. salmi*, giant cell pneumonia, histopathology
pigs (diaphragmatic lobes of lungs): West Bengal
- Metastrongylus salmi*
Coombs, D. W.; and Springer, M. D., 1974, *J. Wildlife Dis.*, v. 10 (4), 436-441
Sus scrofa domesticus x *Sus scrofa cristatus* (lungs): Aransas National Wildlife Refuge, southern Texas
- Metastrongylus salmi, illus.*
Pulviaevskaia, N. V., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 119-122
Metastrongylus salmi and *Spirocerca lupi* females, morphology of genital tract
- Metastrongylus salmi*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii krai
- Metathelazia* Skinker, 1931, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Pneumospiruridae
key, synonymy
- Metathelazia*
Pence, D. B.; and Stone, J. E., 1977, *J. Parasitol.*, v. 63 (6), 979-991
systematics of family *Pneumospiruridae*
- Metathelazia*
Wertheim, G.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (6), 647-657
Thelazioidea, Pneumospiruridae
cephalic structures, scanning electron microscopy
Syn.: *Pneumospirura*
- Metathelazia acomysi, illus.*
Wertheim, G.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (6), 647-657
cephalic structures, scanning electron microscopy
- Metathelazia californica* Skinker 1931, *illus.*
Pence, D. B.; and Stone, J. E., 1977, *J. Parasitol.*, v. 63 (6), 979-991
redescription, pathology
Felis rufus (alveolar spaces and terminal bronchioles of lung parenchyma): Benjamin, Knox Co., Texas; Adams Ranch, King Co., Texas; Pitchfork and Beggs Ranches, Dickens Co., Texas
- Metathelazia capsulata, illus.*
Wertheim, G.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (6), 647-657
cephalic structures, scanning electron microscopy
- Metathelazia exilis* (Biocca and Chabaud, 1952) n. comb.
Wertheim, G.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (6), 647-657
- Metathelazia felis* (Vogel 1928) Dougherty 1943
Pence, D. B.; and Stone, J. E., 1977, *J. Parasitol.*, v. 63 (6), 979-991
as syn. of *Vogeloides felis* (Vogel 1928)
Davitian 1933

- Metathelazia felis* (Vogel, 1928) n. comb.
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
- Metathelazia hainanensis* (Wu and Hu, 1938) n. comb.
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
- Metathelazia multipapillata*, illus.
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
cephalic structures, scanning electron microscopy
- Metathelazia oesophageus*
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
as syn. of *Vogeloides oesophageus* (Gerichter, 1948), Dougherty, 1952
- Metathelazia rodentium* (Wertheim and Giladi, 1977) n. comb., illus.
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
cephalic structures, scanning electron microscopy
Syn.: *Pneumospirura rodentium*
- Meteterakinae
Crusz, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
definition emended
- Meteterakis sinharajensis* sp. nov., illus.
Crusz, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
Lyriocephalus scutatus (duodenum, rectum): Godekande in Hiniduma, bordering on Sinharaja rain forest, Ceylon
Pseudotyphlops philippinus (rectum): Dewatura Estate below Namunukula, Ceylon
- Metheligmonella Durette-Desset*, 1971
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Metheligmonella Durette-Desset*, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Mikoletzkyia* sp.
Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Mikoletzkyia pinicola*
Vosilite, B. S., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Microfilaria*
Aikat, T. K.; and Das, M., 1977, Indian J. Med. Research, v. 65 (1), 58-64
Wuchereria bancrofti, modified statistical method for analysis of periodicity of microfilaria using harmonic wave equation
- Microfilaria*
Arafa, M. S.; Salit, A. M.; and Hilal, T., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 13 [Demonstration]
unsheathed microfilariae undistinguishable from *Dipetalonema witeae* found in blood of *Rattus* spp. in Egypt
- Microfilaria* D, illus.
Chalifoux, L. V.; et al., 1973, Lab. Animal Sc., v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saguinus oedipus
Cebus apella
all from New England Regional Primate Research Center
- Microfilaria*
Greiner, E. C.; et al., 1975, Canad. J. Zool., v. 53 (12), 1762-1787
avian hematozoa, prevalence with reference to distribution by geographic region, by host family, by vertical stratification of nesting sites and by feeding behavior of known vectors: North America north of Mexico
[Checklist includes 388 bird species and contains both published and unpublished records. For records from specific hosts, see entries in Supplement 22, Part 7, Hosts.]
- Microfilaria*
Webber, R. H., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 537-538 [Letter]
human microfilaria, comparison of use of counting chamber and measured blood films for epidemiologic estimations, survey of *Wuchereria bancrofti*-endemic area in the Solomon Islands
- Microfilaria* [sp.]
Bennett, G. F.; et al., 1974, J. Wildlife Dis., v. 10 (4), 442-451
survey, prevalence of hematozoa in anatids, infection rate increases with host age, seasonal distribution
Anas rubripes
Anas platyrhynchos
Anas platyrhynchos x *A. rubripes*
Anas carolinensis
Aix sponsa
(blood of all): all from Massachusetts

- Microfilaria* [sp.]
 Bennett, G. F.; and Borrero H., J. I., 1976, J. Wildlife Dis., v. 12 (3), 454-458
Pionopsitta haematotis
P. pulchra
Tangara xanthogastra
Thraupis palmarum
T. virens
Coeligena wilsoni
Eutoxeres aquila
Catharus ustulatus
Myadestes ralloides
Tyrannus melancholicus
 (blood of all): all from Colombia
- Microfilaria* [sp.]
 Bennett, G. F.; and Herman, C. M., 1976, J. Wildlife Dis., v. 12 (1), 59-65
 (blood of all)
Streptopelia senegalensis: Kenya
Pytelia afra: Zaire
Dryoscopus cubla: Zaire
Cyanomitra olivacea: Tanzania
Nectarinia amethystina: Zaire
N. kilimensis: Kenya
Numida meleagris: Zaire
Oriolus auratus: Zaire
Cisticola robusta: Kenya
Zoothera oberlaenderi: Tanzania
Pycnonotus xanthopygus: "
- Microfilaria* [sp.]
 Bennett, G. F.; Okia, N. O.; and Cameron, M. F., 1974, J. Wildlife Dis., v. 10 (4), 458-465
 survey, avian hematozoa, seasonal prevalence
Ceuthmochares aereus
Pycnonotus barbatus
Zosterops senegalensis
 (blood of all): all from Uganda
- Microfilaria* [sp.]
 Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus
Phoca vitulina
 (blood of all): all from European waters
- Microfilaria* [sp.]
 Choudhury, A.; and Misra, K. K., 1973, J. Protozool., v. 20 (4), 514
Lanius schach collaris (blood): India
- Microfilaria* [sp.], illus.
 Dissanaïke, A. S., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 142-143 [Demonstration]
 morphometric measurements of sheathed microfilaria from blood of *Rattus muelleri*: Johore Lobis Forest Reserve
- Microfilaria* [sp.], type 2
 Dissanaïke, A. S.; and Fernando, M. A., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 138 [Demonstration]
Gallus gallus spadiceus (blood and lung smears): Malaysia
- Microfilaria* [sp.], type 3
 Dissanaïke, A. S.; and Fernando, M. A., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 138 [Demonstration]
Gallus gallus spadiceus (blood and lung smears): Malaysia
- Microfilaria* [sp.], unidentified
 Miyata, A.; and Tsukamoto, M., 1975, Nettai Igaku (Trop. Med.), v. 16 (3), 113-130
Callosciurus juvencus: Palawan Island, the Philippines
- Microfilaria* [sp.], unidentified sheathed microfilaria, illus.
 Miyata, A.; and Tsukamoto, M., 1975, Nettai Igaku (Trop. Med.), v. 16 (3), 113-130
Varanus salvator: Palawan Island, the Philippines
- Microfilaria* [sp.]
 Oduye, O. O.; and Dipeolu, O. O., 1976, J. Small Animal Practice, v. 17 (5), 331-337
 blood parasites of dogs, single and mixed infections, correlation between incidence and rainfall, degree of parasitaemia, infectivity rate within age groups, no significant difference in host susceptibility of local and exotic breeds to infection: Ibadan, Nigeria
- Microfilaria* sp.
 Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrurus tetricus
Tetrao urogallus
 all from USSR
- Microfilaria* [sp.], illus.
 Peirce, M. A.; and Bevan, B. J., 1977, Vet. Rec., v. 100 (14), 282-283
Amazona aestiva
Cacatua galerita
C. sulphurea
Psittacula eupatria
P. k. krameri
 (blood of all): all imported to London from Indonesia
- Microfilaria* [sp.]
 Stabler, R. M.; Kitzmiller, N. J.; and Braun, C. E., 1977, J. Wildlife Management, v. 41 (1), 128-130
Columba fasciata (blood): Colorado; California; Mexico
- Microfilaria* [sp.]
 Thomas, S. E.; and Dobson, L. D., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 67-68
Netta erythrophthalma
Streptopelia senegalensis
 (blood of all): all from vicinity of Onderstepoort, Republic of South Africa
- Microfilaria* [sp.]
 Wood, S. F., 1975, J. Parasitol., v. 61 (5), 969-970
Peromyscus boylii rowleyii (blood): Tonto National Monument, near Roosevelt, Gila County, Arizona
P. truei (blood): Chaco Canyon National Monument, San Juan County, New Mexico
P. maniculatus (blood): Chaco Canyon National Monument, San Juan County, New Mexico
- Microfilariae*, Bancroftian. See [*Wuchereria bancrofti*]
- Microfilariae*
 Andrews, S. E.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 24-28
Corvus brachyrhynchos: insular Newfoundland

Microfilariidae

- Ashford, R. W.; et al., 1976, J. Wildlife Dis., v. 12 (3), 409-426
- Anhinga rufa
 Necrosyrtes monachus
 Actophilornis africana
 Tringa hypoleucos
 Columba guinea
 Oena capensis
 Aplopelia larvata
 Clamator jacobinus
 Otus leucotis
 Alcedo cristata
 Halcyon senegalensis
 H. leucocephala
 Merops apiaster
 M. nubicus
 M. albicollis
 M. lafresnayii
 Upupa epops
 Tockus alboterminatus
 Lybius leucomelas
 Riparia riparia
 R. paludicola
 Hirundo smithii
 Motacilla flava
 Pycnonotus barbatus
 Prionops plumata
 Dryoscopus gambensis
 Tchagra minuta
 T. senegala
 Laniarius aethiopicus
 Cossypha heuglini
 Luscinia megarhynchos
 Turdus pelios
 Turdoides rubiginosus
 Acrocephalus gracilirostris
 Sphenoeacus mentalis
 Sylvia atricapilla
 S. curruca
 S. mystacea
 Cisticola erythrops
 C. brachyptera
 Phyllolais pulchella
 Muscicapa striata
 Terpsiphone viridis
 Anthreptes platurus
 Nectarinia senegalensis
 Zosterops senegalensis
 Emberiza tahapisi
 Serinus mozambicus
 S. citrinelloides
 Hypochera chalybeata
 Pytelia phoenicopter
 Estrilda rhodopyga
 E. astrild
 Uraeginthus ianthinogaster
 U. bengalus
 Lagonosticta larvata
 L. senegala
 L. rubricata
 Lonchura malabarica
 Amblyospiza albifrons
 Ploceus luteolus
 P. taeniopterus
 P. velatus
 P. cucullatus
 P. ocularis
 Malimbus rubriceps
 Euplectes macrourus
 Dinemellia dinemelli
 Passer griseus
 P. eminibey
 Lamprotornis chalybaeus
 Creatophora cinerea
 Dicrurus adsimilis
 all from Ethiopia

Microfilariidae

- Brooks, M. A., 1976, Invert. Tissue Cult. Research Applic. (Maramorosch), 181-199
 parasite transmission, applications of insect tissue culture, review and prospects
- Microfilariidae B, illus.
 Chalifoux, L. V.; et al., 1973, Lab. Animal Sc., v. 23 (2), 211-220
 differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
 Saimiri sciureus
 Saguinus oedipus
 all from New England Regional Primate Research Center
- Microfilariidae K, illus.
 Chalifoux, L. V.; et al., 1973, Lab. Animal Sc., v. 23 (2), 211-220
 differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
 Saguinus tamarinus: New England Regional Primate Research Center
- Microfilariidae
 Cheke, R. A.; Hassall, M.; and Peirce, M. A., 1976, J. Wildlife Dis., v. 12 (2), 133-138
 Parus caeruleus (blood): Great Britain
- Microfilariidae, morphologically similar to Dipetalonema interstitium microfilariae
 Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
 epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
 Sciurus carolinensis (blood): Maryland; North Carolina; South Carolina; Georgia; Mississippi; Alabama; Arkansas; Tennessee; Kentucky; Virginia
- Microfilariidae
 El-Bishlawi, O., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (2), 307-308 [Letter]
 Schistosoma haematobium, adhesion of red blood cells to ova as cause of hematuria; similar adhesion phenomenon observed with microfilariae in blood-tinged hydrocoele fluid
- Microfilariidae, possibly Aproctella stoddardi or Diplotriaenoides sp.
 Eve, J. H.; and Davidson, W. R., 1976, J. Parasitol., v. 62 (1), 142-144
 Bonasa umbellus (blood): Kentucky; Michigan; West Virginia
- Microfilariidae
 Fallis, A. M.; Jacobson, R. L.; and Raybould, J. N., 1973, J. Protozool., v. 20 (3), 438-442
 Gutтера pucherani (blood): Amani, Tanzania

- Microfilariae**
 Jones, T. C.; Mott, K.; and Pedrosa, L. C., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (2), 243-246
 technique for isolating and concentrating microfilariae from peripheral blood using gradient centrifugation
- Microfilariae**
 Manwell, R. D.; Allen, C. S.; and Kuntz, R. E., 1976, *J. Protozool.*, v. 23 (4), 571-576
Bambusicola thoracica sonorivox
Turnix suscitator rostrata
 (blood of all): all from Taiwan
- Microfilariae**
 Manwell, R. D.; and Rossi, G. S., 1975, *J. Protozool.*, v. 22 (1), 124-127
Cyanothorax affinis (lungs and air sacs): imported to U. S., origin unknown
- Microfilariae, resembling Onchocerca sp.**
 Partono, F.; et al., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 537-547
 domestic cows (blood): Margolembo, South Sulawesi
- Microfilariae, resembling Setaria sp.**
 Partono, F.; et al., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 537-547
 domestic cows (blood): Margolembo, South Sulawesi
- Microfilariae**
 Partono, F.; and Idris, K. N., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 158-164
 factors that influence loss of microfilariae from stained thick blood films, results of laboratory experiments
- Microfilariae**
 Peirce, M. A.; and Cheke, A. S., 1977, *J. Protozool.*, v. 24 (2), Suppl., 33A [Abstract]
Turdus bewsheri moheliensis: Comoro Islands
- Microfilariae, illus.**
 Perera, P. A. C., 1977, *Med. Lab. Sc.*, v. 34 (2), 127-129
 method of staining the sheath of microfilariae in human microfilarial granulomata of the breast
- Microfilariae**
 Rajapaksa, N.; and Garnham, P. C. C., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (1), 4 [Demonstration]
Arborophila charltoni (blood)
- Microfilariae**
 Sasa, M., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (2), 197-210
 human microfilariae, statistical technique for estimating efficiency of detection of parasites in varying volumes of blood samples taken during epidemiologic surveys
- Microfilariae**
 Schillhorn van Veen, T.; and Blotkamp, J., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (4), 517-518
 dogs: Zaria area, Nigeria
- Microfilariae**
 Schulz-Key, H., 1975, *Tropenmed. und Parasitol.*, v. 26 (4), 494-498
 skin-inhabiting microfilariae of unidentified filarial worm discovered in *Dama dama*: Southern Germany
- Microfilariae**
 Telford, S. R., jr., 1977, *Internat. J. Parasitol.*, v. 7 (4), 299-314
 microfilariae distribution and zoogeography, taxonomic problems, incidence by locality and season, implications for vector searches, mixed infections, comparative effectiveness of initial vs. repeated examinations of blood smears for detection: Middle America
Anolis capito: Costa Rica; Panama
Ameiva undulata: Costa Rica
Thecadactylus rapicaudus: Panama
Gonatodes albogularis: Panama
Anolis limifrons: Panama
A. lionotus: Panama
A. poecilopus: Panama
A. biporcatus: Panama
Corytophanes cristatus: Panama
Ameiva ameiva: Panama
Mabuya mabouya: Panama
- Microfilariae**
 Townson, H., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 12-13 [Demonstration]
Brugia pahangi, refined technique for inoculating mosquitoes with microfilariae
- Microfilariae**
 Worms, M. J., 1972, *Zool. J. Linn. Soc., London*, v. 51, Suppl. 1, 53-67
 microfilariae, *Plasmodium* spp., trypanosomes, rhythmic behavior, significance in relation to transmission, review
- Microhadjelia Jogis, 1965, illus.**
 Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
 Tetramerinae
 key
- Microhadjelia**
 Quentin, J. C.; and Wertheim, G., 1975, *Ann. Parasitol.*, v. 50 (1), 63-85
 Tetrameridae, redefinition
- Microhadjelia multipapillata Mogis, 1965, illus.**
 Quentin, J. C.; and Wertheim, G., 1975, *Ann. Parasitol.*, v. 50 (1), 63-85
 redescription
Lanius senator: Bet Guvrin, Israel
- Micronema deletrix, illus.**
 Powers, R. D.; and Benz, G. W., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (2), 175-177
Micronema deletrix, central nervous system of horse, pathology
- Micropleura Linstow, 1906**
 Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
 Micropleuridae

- Micropleuridae (Baylis & Daubney, 1926, subfam.)
Travassos, 1960
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Dracunculoidea
key
includes: Micropleura
- Microtetrameres, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of Tetrameres
key
- Microtetrameres sp.
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (proventriculus): Franklin county, Ohio
- Microtetrameres sp.
Ellis, C. J.; and Calderwood, G., 1977, Proc. Iowa Acad. Sc., v. 84 (1), 30-31
Catharus ustulata
Dendroica pensylvanica
Seiurus auropallidus
Oporornis philadelphia
Passerina cyanea
(proventriculi of all): all from Iowa
- Microtetrameres corax
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Cyphostete komarovi
Adesmia servillei schatzmayri
A. gebleri
Trigonoscelis gigas
T. punctipleuris
Pisterotarsa gigantea subsp. zoubkoffi
P. kessleri
all from Turkmenia
- Microtetrameres helix
Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 399-403
survey, helminths of red-winged blackbirds including a check list of previous findings
Agelaius phoeniceus (proventriculus): South Bass Island, Ohio
- Microtetrameres inermis Travassos, 1914
Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Microtetrameres (Microtetrameres) platales sp. n., illus.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 315-327
Platalea leucorodia (proventriculus): Hyderabad, Andhra Pradesh, India
- Microtetrameres pusilla Travassos
Ellis, C. J.; and Calderwood, G., 1977, Proc. Iowa Acad. Sc., v. 84 (1), 30-31
Mniotilta varia (proventriculi): Iowa
- Microtetrameres spiculata Boyd, 1956
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma coerulescens coerulescens (proventriculus): Florida
- Mikoletzky. See Micoletzkyka.
- Mirandaia Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Stilestrongylus Freitas, Lent et Almeida, 1937
- Mirzaiella asiatica Basir, 1942
Hristovski, N. D., 1972, Acta Parasitol. Iugoslavica, v. 3 (2), 109-115
Gryllotalpa gryllotalpa: Armenia-SSSR (s. Samagar)
- Moennigia Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
synonymy
- Moennigia
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
Syn.: Pintonema
- Moennigia sp. Durette-Desset, 1974
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
as syn. of Moennigia dessetae n. sp.
- Moennigia sp., illus.
Durette-Desset, M. C., 1974, Ann. Parasitol., v. 49 (5), 555-566
description, 4 spp. of Trichostrongyloidea, localization in intestine, larval and adult synlophes compared, implications for taxonomy and evolution
Metachirops opossum (intestin): Guyane Francaise
- Moennigia alonsoi n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, didelphic forms of Trichostrongyloidea more abundant than monodelphic ones
Tamandua tetradactyla (intestin): region de Belem, Province de Para, Bresil
- Moennigia dessetae Diaw, 1976 [? nom. nud.]
Diaw, O. T., 1976, Ann. Parasitol., v. 51 (3), 355-363
trichostrongyloid nematode fauna of Didelphis marsupialis compared to that of Metachirops opossum, localization within intestine
Didelphis marsupialis
Metachirops opossum
(intestin of all): all from Guyane francaise
- Moennigia dessetae n. sp., illus.
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Syn.: Moennigia sp. Durette-Desset, 1974
Didelphis marsupialis (intestin anterieur): Guyane francaise

- Moennigia lentinae* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, didelphic forms of *Trichostrongyloidea* more abundant than monodelphic ones
Tamandua tetradactyla (intestin): region de Belem, Province de Para, Bresil
- Moennigia levyi* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, didelphic forms of *Trichostrongyloidea* more abundant than monodelphic ones
Tamandua tetradactyla (intestin): region de Belem, Province de Para, Bresil
- Moennigia michelae* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, didelphic forms of *Trichostrongyloidea* more abundant than monodelphic ones
Tamandua tetradactyla (intestin): region de Belem, Province de Para, Bresil
- Moennigia obelsi* n. sp., illus.
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
frequency and distribution in host gut, didelphic forms of *Trichostrongyloidea* more abundant than monodelphic ones
Tamandua tetradactyla (intestin): region de Belem, Province de Para, Bresil
- Moguranema*
Durette-Desset, M. C., 1977, Ann. Parasitol., v. 52 (5), 583-584
placed in *Molineidae*, *Molineinae*
- Moguranema Yamaguti*, 1941
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Molineinae*
- Moguranema japonicum Yamaguti*, 1941, illus.
Durette-Desset, M. C., 1977, Ann. Parasitol., v. 52 (5), 583-584
Mogera wogura: Japon
- Molineidae* (Skrjabin et Schulz, 1937, sub. fam.), n. fam.
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
includes: *Molineinae*; *Anoplostrongylinae*; *Nematodirinae*
- Molineinae* Skrjabin et Schulz, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae
includes: *Molineus* (type genus); *Angulocirrus*; *Batrachostromylus*; *Columbostromylus*; *Dollfusstromylus*; *Hepatojarakus*; *Hooperstromylus*; *Moguranema*; *Molinostrongylus*; *Nochtia*; *Nycteridostromylus*; *Ortleppstrongylus*; *Oswaldocruzia*; *Pithecostrongylus*; *Poekilostromylus*; *Schulzia*; *Trichochenia*; *Trichoskrjabinia*
- Molinema* [? n. rank]
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
subgen. of *Dipetalonema*; key
tod: *D. (M.) diacantha* (Freitas et Lent, 1939)
- Molineus* Cameron, 1923 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Molineinae*
synonymy
- Molineus* sp., probably *M. barbatus*
Hass, D. K.; and Chitwood, M. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 255
dog (small intestine): west central Ohio
- Molineus patens*
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Molineus patens* (Duj., 1845) Petrov, 1928
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Martes martes
Mustela erminea
all from Pechora river basin
- Molineus patens* (Dujardin, 1845) Petrov, 1928
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela lutreola
Meles meles
Mustela putorius
all from Karelia
- Molineus torulosis*, illus.
King, N. W., jr., 1976, Scient. Publication (317). Pan Am. Health Organ., 169-198
- Molinospirura*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Oxyspirura*
key
- Molinostrongylus* Skarbilowitsch, 1934
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, *Molineinae*
- Molinostrongylus* Skarbilovitsch, 1934
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
differential diagnosis of *Molinostrongylus* spp. from *Microchiroptera*, includes: *M. skrjabini*; *M. alatus*; *M. delicatus*; *M. heydoni*; *M. ornatus*; *M. rhinolophi*; *M. tipula*; *M. panousei*; *M. pseudoornatus*; *M. vespertilionis*; *M. dollfusi*; *M. longispicula*; *M. spasskii*
- Molinostrongylus* sp.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Nyctalus leisleri
Pipistrellus nathusii
all from Col de Jaman, Vaud, Suisse

- Molinostrongylus* sp. Andreiko et Skvortsov, 1966
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
as syn. of *M. spasskii* Andreiko, Pinchuk, Skvortsov, 1968
- Molinostrongylus aelleni* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Nyctalus lasiopterus (intestin grele):
Col de Bretolet, Valais, Suisse
- Molinostrongylus alatus* (Ortlepp, 1932), illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
description
Myotis myotis: Trelex, Vaud, Suisse
M. bechsteini: Commugny, Vaud, Suisse
M. blythii oxygnathus: Zagorska pec, Novi, Yougoslavie
M. nattereri: Commugny, Vaud, Suisse
- Molinostrongylus alatus* (Ortlepp, 1932) Skarbilovitsch, 1934, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
synonymy, description, geographic distribution
Miniopertus schreibersi
M. natalensis
Myotis oxygnathus
M. mystacinus
M. dasycneme
M. nattereri
Pipistrellus pipistrellus
Rhinolophus ferrumequinum
R. hipposideros
Nyctalus noctula
(small intestine of all): all from Moldavia
- Molinostrongylus alatus* (Ortlepp, 1932) Skarbilovitsch, 1934
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Rhinolophus hipposideros
Myotis oxygnathus
all from Moldavia
- Molinostrongylus bauchoti* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Chaerophon sp. (intestin): Tananarive, Madagascar
- Molinostrongylus benexae* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Chaerophon limbatus (intestin): Perinet, Madagascar
- Molinostrongylus colleyi* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Tylonycteris pachypus (intestin)
T. robustula
all from Ulu Gombak, District de Selangor, Malaisie
- Molinostrongylus daubentoni* Zdzitowiecki, 1970
syn. n.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 57-75
as syn. of *Molinostrongylus spasskii* Andreiko, Pinchuk, Skvortsov, 1968
- Molinostrongylus delicatus* (Schwartz, 1927)
Travassos, 1937
Cain, G. D.; and Studier, E. H., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 113-114
Molossus ater: Sinaloa, Mexico
- Molinostrongylus delicatus*
Martin, D. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 85-86
Tadarida brasiliensis: Texas; Louisiana
- Molinostrongylus ovyangi* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Glischropus tylopus (intestin): Ulu Gombak, District de Selangor, Malaisie
- Molinostrongylus panousei* Dollfus, 1954, illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
description
Miniopertus schreibersi: grotte de Pouade (Pyrenees-Orientales), France
- Molinostrongylus richardae* n. sp., illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Mormopterus albiventer (intestin): Anjiro, Madagascar
Eptesicus pusillus: Tananarive, Madagascar
- Molinostrongylus skrjabini* Skarbilovitsch, 1934, illus.
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
description
Nyctalus noctula: Uster, Zurich, Suisse;
Col de Bretolet, Valais; Col de Jaman, Vaud, Suisse
- Molinostrongylus skrjabini* Skarbilovitsch, 1934, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Rhinolophus ferrumequinum
R. hipposideros
Myotis bechsteini
M. daubentoni
M. oxygnathus
M. myotis
M. mystacinus
M. dasycneme
Plecotus auritus
Nyctalus noctula
Vespertilio murinus
Miniopertus schreibersi
Chiroptera sp.
(small intestine of all): all from Moldavia

- Molinostrongylus skrjabini* Skarbilovitsch, 1934
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Myotis oxygnathus
M. myotis
M. daubentoni
M. bechsteini
M. mystacinus
Plecotus auritus
Nyctalus noctula
all from Moldavia
- Molinostrongylus spasskii* Andreiko, Pinchuk, Skvorzov, 1968
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 57-75
Syn.: *Molinostrongylus daubentoni* Zdzitowiecki, 1970 syn. n.
- Molinostrongylus spasskii* Andreiko, Pinchuk, Skvorzov, 1968, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Syn.: *Molinostrongylus* sp. Andreiko et Skvorzov, 1966
Myotis myotis
M. daubentoni
M. mystacinus
(small intestine of all): all from Moldavia
- Molinostrongylus spasskii* Andreiko, Pinchuk, Skvorzov, 1968
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Myotis myotis
M. daubentoni
M. mystacinus
all from Moldavia
- Molinostrongylus vespertilionis* Morosov et Spassky, 1961, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Myotis oxygnathus
M. myotis
M. dasycneme
M. daubentoni
M. mystacinus
M. bechsteini
Vespertilio pipistrellus
Nyctalus leisleri
N. noctula
(small intestine of all): all from Moldavia
- Molinostrongylus vespertilionis* Morosov et Spassky, 1961
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Myotis oxygnathus
M. myotis
M. dasycneme
M. daubentoni
M. bechsteini
M. mystacinus
Nyctalus leisleri
N. noctula
Pipistrellus pipistrellus
all from Moldavia
- Molnaria* Moravec, 1968, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Skrjabillaninae
key
- Monanema Anteson*, 1968
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
key
- Monanema globulosa* (Muller et Nelson, 1975) n. comb.
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Monanema marmotae* (Webster, 1967), illus.
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Monanema nilotica* n.sp., illus.
ElBihari, S.; Hussein, H. S.; and Muller, R., 1977, J. Helminthol., v. 51 (4), 317-321
Arvicanthus niloticus testicularis (adults in right auricle of heart and pulmonary artery; microfilariae in skin): Shambat, Khartoum-North, Sudan
- Monhysterida* DeConinck & Schuurmans Stekhoven, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Chromadoria
- Monodontella giraffae* Yorke & Maplestone Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Giraffa camelopardalis tippelskirchi (liver): Kenya
- Monodontus louisianensis*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Mononchida* Jairajpuri, 1969
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Enoplia
- Morerastrongylus* Chabaud, 1972 emend.
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (6), 637-642
diagnosis
- Morerastrongylus andersoni* (Petter, 1972)
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (6), 637-642
as syn. of *Gallegostrongylus andersoni* (Petter, 1972) n. comb.
- Morerastrongylus andersoni* (Petter, 1972), illus.
Petter, A. J.; and Cassone, J., 1975, Ann. Parasitol., v. 50 (4), 469-475
Morerastrongylus andersoni, mode of entry (ingestion) and localization in *Limnaea stagnalis* and *Planorbium corneum*

- Morganiella Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Tricholinstowia Travassos, 1937
- Morganostrogylus Fahmy, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Tricholinstowia Travassos, 1937
- Muellerius
Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
sheep helminths, effective control with Nil-
verm and Zanil, increased weight gains and
shearing yields: Hanczowa, Gorlice district
- Muellerius
Nemeseri, L., 1976, Magy. Allat. Lapja, v. 98,
v. 32 [i. e. 31] (7), 459-461
Dictyocaulus filaria, trichostrongylids,
Fasciola hepatica, sheep, successful treat-
ment with combination of diamphenetide and
tetramisole; no effect on Protostrongylidae
- Muellerius
Paul, I.; et al., 1975, Lucrar. Stiint. Inst.
Agron. I. Ionescu Brad Iasi, II. Zootehn.,
Med. Vet., 71-72
Protostrongylus, Muellerius, Cystocaulus,
bronchopneumonia, adult sheep, morphopathol-
ogy, result of association of several species
- Muellerius [sp.]
Helle, O., 1976, Norwegian J. Zool., v. 24
(4), 463 [Abstract]
goats (feces): Norway
- Muellerius capillaris
Aller, B.; and Aller, J. M., 1973, Rev. Iber.
Parasitol., v. 33 (2-3), 337-346
nematodes of sheep lungs, lower incidence of
fungi in parasitized lungs: Leon
- Muellerius capillaris
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno,
v. 43 (2), 123-131
helminths and coccidians of Ovis ammon mus-
imon and Capreolus capreolus, intensity
variation with age of host, lack of evidence
for parasite exchange between mouflons and
roe deer
Ovis ammon musimon (lung): School Forest
Enterprise, University of Agriculture Brno,
Krtiny
- Muellerius capillaris
Dyk, V.; and Chroust, K., 1975, Vet. Parasi-
tol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe
deer, incidence and intensity, possible cross
transmission, implications for game manage-
ment
Ovis ammon musimon: Czechoslovakia
- Muellerius capillaris
Dyk, V.; and Chroust, K., 1975, Veterinarstvi,
v. 25 (7), 315-317
helminths, incidence by age of host, problem
in mouflon husbandry: Brno oblast
- Muellerius capillaris
Ferreira, L. D. B. B., 1975, Rev. Portug.
Cien. Vet. (433-434), v. 70, 25-39
broncho-pulmonary nematodes of sheep, life
cycles, epizootiology, instructive review
for veterinarians
- Muellerius capillaris
Goda, Fawzy F. M., 1974, Bull. Epizoot. Dis.
Africa, v. 22 (1), 75-78
sheep: Benghazi, Libya
- Muellerius capillaris
Helle, O., 1976, Norwegian J. Zool., v. 24
(4), 463 [Abstract]
goats (feces): Norway
- Muellerius capillaris
Kozdon, O.; and Zajicek, D., 1976, Vet. Med.,
Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field con-
ditions, seasonal distribution as modified
by dehelminthization, possible management
strategies for effective timing of dehel-
minthization: Western Bohemia
- Muellerius capillaris, illus.
Leguia, G., 1972, Rev. Invest. Pecuarias, v. 1
(2), 233-234
Muellerius capillaris, verminous pneumonia,
case report, Corriedale sheep: Huancayo
- Muellerius capillaris
Melikov, Iu. F., 1975, Izvest. Akad. Nauk
Azerbaidzhan. SSR, s. Biol. Nauk (3), 80-83
distribution in sheep of sheep farms of
Kura-Araksinsk lowland, Azerbaidzhan
- Muellerius capillaris
Prestwood, A. K.; Pursglove, S. R.; and Hayes,
F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-
385
survey of parasites of Odocoileus virgini-
anus and Ovis aries on common range, deer
unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Muellerius capillaris
Ramisz, A.; Urban, E.; and Piechocki, B.,
1975, Med. Wet., v. 31 (11), 677-679
Protostrongylidae of sheep, tetramisole
effective against all species except Mueller-
ius capillaris
- Muellerius capillaris
Rojo Vazquez, J., 1977, An. Fac. Vet. Leon,
Oviedo, v. 21 (21), 1975, 51-101
Protostrongylinae, frequency in ovine lungs,
primary nodes are infected by bacteria at a
lower rate than non-parasitized lung areas:
municipal slaughterhouse, Leon
- Muellerius capillaris
Rose, J. H., 1971, Symposia Brit. Soc. Parasi-
tol., v. 9, 109-121
gastrointestinal nematodes and lungworms of
farm animals, isolation and maintenance in
vivo, extensive review

- Muellerius capillaris* (Mueller, 1899)
Sultanov, M. A.; Azimov, D. A.; and Ubaidullaev, Ia. U., 1975, *Uzbek. Biol. Zhurnal* (6), 39-42
Muellerius capillaris, development in molluscs as first and second intermediate hosts, sheep and goats as final hosts
Helicella candaharica (nat. and exper.):
Uzbekistan
Succinea pfeifferi (exper.)
Subzebrinus albiplicatus (exper.)
Subzebrinus sogdianus (exper.)
Bradybaena dichrozona (exper.)
B. phaeozona (exper.)
[*Ovis aries*] (exper.)
[*Capra hircus*] (exper.)
- Muellerius capillaris*, *illus.*
Sumnaliev, P., 1973, *Izvest. Tsentral. Khel'mint. Lab.*, v. 16, 161-167
Neostromylus linearis, *Muellerius capillaris*, *Cystocaulus ocreatus*, *Protostrongylus* sp., differential diagnosis of infective larvae based on biometric studies
Helicella obvia
Zebrina detrita
all from Bulgaria
- Muellerius capillaris*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Muellerius capillaris*
Volf, K.; and Volfova, M., 1974, *Veterinarstvi*, v. 24 (3), 125-126
srnci zvere: Trebic District
- Muellerius capillaris*
Zajicek, D.; and Kozdon, O., 1977, *Veterinarstvi*, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Muellerius minutissimus*
Kirkpatrick, C. M.; et al., 1976, *Research Bull. Purdue Univ., Agric. Exper. Station* (932), 1-42
Odocoileus virginianus: Naval Weapons Support Center, Crane, Indiana
- Muellerius tenuispiculatus* Gebauer, 1932, *illus.*
Svarc, R.; and Zmoray, I., 1974, *Biologia, Bratislava*, s. B, *Zool.* (1), v. 29 (2), 121-127
Muellerius tenuispiculatus larvae, penetration into and development in soles of feet of *Cepaea vindobonensis* and *Succinea putris*, localization near sole glands, acid mucopolysaccharides and phospholipids in this site probably have role in larval nutrition
- Muflonagia* Schulz et Coll., 1953
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae*
- Multicaecinae* (Hartwich, 1954, fam.) Campana-Rouget, 1960
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascarididae
key; key to tribes
includes: *Multicaecinae*; *Dujardinascariidae*
- Multicaecinae* Campana-Rouget, 1960
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Multicaecinae
key; key to genera
includes: *Typhlophorus*; *Multicaecum*
- Multicaecum* Baylis, 1923, *illus.*
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Multicaecinae
key; synonymy
- Multicaecum* (*Brevimulticaecum*) Mozgovoi, 1951
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Multicaecum* Baylis, 1923
- Multiplectines*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Pterygodermatites*
key
- Murielea* Skrjabin et Schikhobalova, 1952
Durette-Desset, M. C., 1974, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (216), *Zool.* (144), 419-423
elevation of tribe *Murielea* Skrjabin et Schikhobalova, 1952 to subfamily *Murielinae*
- Murielinae* [new rank]
Durette-Desset, M. C., 1974, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (216), *Zool.* (144), 419-423
elevation of tribe *Murielea* Skrjabin et Schikhobalova, 1952 to subfamily *Murielinae*
- Murielus* Dikmans, 1939
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, *Nematodirinae*
- Murielus abei* n. sp., *illus.*
Durette-Desset, M. C., 1974, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (216), *Zool.* (144), 419-423
Ochotona roylei (intestins): Gasainhund, Nepal and Thare Pate a l'est de Gasainhund

Murielus harpeticulus Dikmans, 1939, illus.
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 419-423
description

Murielus harpeticulus Dikmans
Grundmann, A. W.; and Lombardi, P. S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 39-46

Ochotona princeps cinnemomea: Tushar Mountains, Utah

O. p. wasatchensis: Wasatch Mountains, Utah

O. p. uinta: Uinta Mountains, Utah

O. p. fuscipes: Markagunt Plateau, Utah

O. p. barnsei: Fish Lake Mountains, Utah

O. p. nevadensis: Ruby Mountains, Nevada

Murielus harpeticulus Dikmans, 1939
Seesee, F. M., 1973, Am. Midland Naturalist, v. 89 (2), 257-265

key

Ochotona p. princeps (small intestine):

St. Joe Baldy Mountain, Benewah County, Idaho

Muspiceoidea

Chaubaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chaubaud, and Willmott)(1), 6-17

Enoplida

key

- Necator**
Hawley, T. G., 1973, N. Zealand Med. J. (489), v. 77, 95-97
Necator infestation in rural dwelling Fijians and Indians, relationship to mean hemoglobin levels
- Necator sp.**
File, S. K.; McGrew, W. C.; and Tutin, C. E. G., 1976, J. Parasitol., v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces): Gombe National Park, Tanzania
- Necator sp., illus.**
Ghadirian, E.; and Biocca, E., 1972, Parassitologia, v. 14 (1), 137-142
Necator sp., description, distinction from Necator americanus
human (intestino (tenue)): Iran
- Necator americanus**
Adickman, M.; and Tuthill, T. M., 1976, Postgrad. Med., v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Necator americanus**
Bailenger, J.; et al., 1977, Pharmacia Biol. (109), v. 11, 267-277
human intestinal parasites, fecal examination using floatation or diphasic concentration, principles involved in both diagnostic procedures, changes in results when mercury or sodium merthiolate is added during procedure, special application of Janeckso and Urbanyi reaction
- Necator americanus**
Banerjee, D.; Mandal, A.; and Prakash, O., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 685-686
marked inhibitory action of mebendazole on development of larval stages from eggs of Ancylostoma duodenale, A. caninum, and Necator americanus, no effect on infective larval stages, in vitro trials
- Necator americanus**
Basset, M.; et al., 1975, Medecine Trop., v. 35 (6), 475-476
mixed Strongyloides infections apparent after treatment of Necator americanus and Ancylostoma duodenale with alcopar or didakene, case reports, statistics
- Necator americanus**
Beltran-H., F.; Gutierrez-Q., M.; and Navarrete-C., E., 1972, Prensa Med. Mexicana, v. 37 (5-6), 224-227
Necator americanus, humans, successful clinical trials with HOE 16842, mild side effects: Mexico
- Necator americanus**
Biagi, F.; Lopez, R.; and Viso, J., 1975, Progr. Drug Research, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review
- Necator americanus**
Biroum-Noerjasin, 1973, Clin. and Exper. Immunol., v. 13 (4), 545-551
hookworm, human, IgE levels in relation to anti-helminthic treatment, to numbers of eggs/g of stool, to states of infestation (with and without reinfestation, with and without superinfestation), and to time post-treatment, changes in differential lymphocyte and eosinophil counts: East Java, Indonesia
- Necator americanus**
Borda, C. E., 1973, Bol. Chileno Parasitol., v. 28 (1-2), 19-23
Necator americanus, epidemiologic survey of island families, favorable conditions of soil in area favor continued development: San Mateo Island, Argentina
- Necator americanus**
Borda, C. E.; Rea, M. J.; and Borda, R. W., 1976, Prensa Med. Argent., v. 63 (2), 55-58
Necator americanus, humans, clinical trials with mebendazole, recommended for mass therapy because of easy administration, good tolerance and few side effects: Argentina
- Necator americanus**
Botero R., D., 1975, Progr. Drug Research, v. 19, 28-43
human intestinal soil transmitted helminths, epidemiology and public health importance, pathology, extensive review
- Necator americanus**
Brown, R. C.; and Girardeau, M. H. F., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 215-219
Strongyloides fuelleborni, Necator americanus, Ancylostoma duodenale, prevalence survey and study of possible transmammmary passage, presence of Strongyloides sp. larvae in milk of one nursing mother suggests that S. fuelleborni may be transmitted via milk in humans: Bulape, Zaire
- Necator americanus**
Bruch, K.; and Haas, J., 1976, Ann. Trop. Med. and Parasitol., v. 70 (2), 205-211
Ascaris, hookworm, Trichuris, human, single dose fenbendazole compared with pyrantel: Liberia
- Necator americanus**
Cantor, D. S.; et al., 1966, Bol. Chileno Parasitol., v. 21 (3), 70-76
Necator americanus, small bowel studies of humans with single and mixed parasitic infections to correlate pathology and fecal fat excretion: Argentina
- Necator americanus**
de Carneri, I., 1974, Riv. Parassitol., Roma, v. 35 (4), 277-284
Necator americanus, human, geographic distribution in northern band extending from Portugal to Iran, prevailing over Ancylostoma duodenale

- Necator americanus*
de Carneri, I.; and Biasin, G., 1972, *Parassitologia*, v. 14 (1), 107-108
Necator americanus, human, endemic focus in rural area, local population attributes introduction of parasite to visit by local family to coffee plantation in Sao Paulo, Brazil 39 years earlier: Lusia (Rovigo)
- Necator americanus*
Cerva, L., 1976, *Ceskoslov. Epidemiol., Mikrobiol., Imunol.*, v. 25 (6), 365-367
helminths, diagnostic staining method for eggs and larvae in fresh feces or those fixed by formalin
- Necator americanus*
Cervoni, W. A.; and Oliver-Gonzalez, J., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 589-591
pyrantel pamoate effective against human infection of *Ascaris lumbricoides* and *Necator americanus* but not *Trichuris trichiura* in clinical evaluations: Puerto Rico
- Necator americanus*
Chinery, W. A.; et al., 1973, *Ann. Trop. Med. and Parasitol.*, v. 67 (1), 75-80
Necator americanus, human, wide variation in response to bephenium hydroxynaphthoate may be due to specific tolerance of the drug: Accra
- Necator americanus*
Chodera, L.; et al., 1973, *Polski Tygod. Lekar.*, v. 28 (9), 338-340
Necator americanus, *Trichuris trichiura*, evaluation of bitoscanate as therapy; poor results in trichuriasis and frequent side effects in hookworm
- Necator americanus*
Doshi, J. C.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 636-639
human hookworm infections, clinical trials of dosage searching for tolerance and efficacy using C.9333-Go/CGP 4540, compound well tolerated with few side effects, active also in mixed infections with *Ascaris lumbricoides*: Bombay, India
- Necator americanus*
Gemsi, G., 1971, *Parasitol. Hungar.*, v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Necator americanus*
Ghadirian, E.; and Biocca, E., 1972, *Parassitologia*, v. 14 (1), 137-142
Necator sp., description, distinction from *Necator americanus*
- Necator americanus*
Grove, D. I.; Burston, T. O.; and Forbes, I. J., 1974, *Clin. and Exper. Immunol.*, v. 18 (4), 565-569
hookworm-infested population, high serum IgE levels, serum IgE and blood eosinophil levels fell after treatment with pyrantel: Papua New Guinea
- Necator americanus*
Grove, D. I.; and Forbes, I. J., 1975, *Med. J. Australia*, v. 1 (11), 336-338
study suggests increased resistance to hookworm infestation in asthmatic and atopic population: Papua New Guinea
- Necator americanus*
Harinasuta, T.; and Bunnag, D., 1975, *Progr. Drug Research*, v. 19, 64-69
Necator americanus, humans, clinical field trials with bitoscanate testing various regimens and dosages: South Thailand
- Necator americanus*
Hsieh, H. C., 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 568 [Demonstration]
Ancylostoma duodenale, *Necator americanus* and *Strongyloides stercoralis*-related spp., differentiation of species using a modified test-tube filter-paper culture
- Necator americanus*
Iannucci, J. G., 1974, *Med. Times, New York*, v. 102 (10), 43-49
gastrointestinal hemorrhage as presenting symptom of human hookworm disease, clinical management, need for diagnostic awareness in non-endemic areas
- Necator* [americanus]
Inglessis, G., 1973, *Arch. Inst. Cardiol. Mexico*, v. 43 (1), 87-101
Necator [americanus] in children, evidence of cardiac pathology associated with parasitic anemia, reversal of symptoms when anemia treated: Venezuela
- Necator* [americanus]
Inglessis, G., 1973, *Arch. Inst. Cardiol. Mexico*, v. 43 (4), 602-608
Necator [americanus], alterations in hemodynamic values and blood pressure associated with hookworm anemia in children: Venezuela
- Necator americanus*
Jones, H. I., 1976, *Papua N. Guinea Med. J.*, v. 19 (3), 153-155
pigs as important factor in spread of *Ascaris lumbricoides* by eating human feces containing *Ascaris* eggs which then pass unharmed through pig digestive system and are spread widely in human environment; human hookworm eggs largely destroyed in pig digestive system thereby effectively reducing human exposure to hookworm larvae: Papua New Guinea
- Necator americanus*
Jones, H. I., 1976, *Papua N. Guinea Med. J.*, v. 19 (3), 165-172
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, epidemiologic survey of human helminths in natives of Kar Kar Island, *Ascaris* infections highest and heaviest in children and principal cause of high eosinophilia in the communities: Madang Province, Papua New Guinea

- Necator americanus*
Jones, H. I., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (4), 523-529
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, statistical epidemiologic survey of human intestinal parasites with discussion of socio-cultural and environmental factors influencing spread of infections: Venilale District of East Timor
- Necator americanus*
Junod, C., 1972, Medecine et Malad. Infect., v. 2 (2), 55-60
Necator americanus, predominant intestinal parasite of natives of Antilles now residing in France, statistics of infection, means of detection, approved treatment with didakene, alcopar or mintagezole
- Necator americanus*
Layrisse, M.; and Vargas, A., 1975, Progr. Food and Nutrition Sc., v. 1 (10), 645-667
human intestinal parasites, mechanisms by which parasites interfere with host nutrition (competition for nutrients, malabsorption, blood loss, excess nutrient utilization), extensive review
- Necator americanus*
Lie, K. J.; Kwo, E. H.; and Ow-Yang, C. K., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (2), 196-200
Ascaris lumbricoides, Trichuris trichiura, Necator americanus, fecal survey for evidence of soil-transmitted helminths in infants and children living near Kuala Lumpur, Malaysia
- Necator americanus*
Lozoff, B.; Warren, K. S.; and Mahmoud, A. A. F., 1975, J. Infect. Dis., v. 132 (5), 606-610
Necator americanus, Ancylostoma duodenale, human, algorithms in diagnosis and management, mebendazole
- Necator americanus*
Nepfert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Necator americanus*
Nepfert, J.; and Warns, C. M., 1974, Tropenmed. u. Parasitol., v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from Ascaris, hookworm, Onchocerca, Dirofilaria immitis, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination
- Necator americanus*
O'Holohan, D. R.; Hugoe-Matthews, J.; and Kanagasabai, K., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (1), 51-55
Necator americanus, patients with hookworm and anemia treated with jonit, drug well accepted and side effects minimal, effective hookworm agent with reduction in anemia levels and reduced fecal egg counts: Malaysia
- Necator americanus*
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Necator americanus*
Papasarathorn, T.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 82-92
human hookworm, epidemiologic survey, relationship of climatic conditions and soil to disease spread, suggested control measures: Soong Nern District, Korat Province, Thailand
- Necator americanus*
Pasticier, A., 1975, Nouv. Presse. Med., v. 4 (37), 2659-2660
single and mixed parasitic infections in humans, good results with pyrantel pamoate
- Necator americanus*
Patricia, A. H.; et al., 1975, Progr. Drug Research, v. 19, 90-95
Necator americanus, Ancylostoma duodenale, humans, efficacy of bitoscanate compared with tetrachlorethylene and bephenium hydroxynaphthoate, found to be safe and effective
- Necator americanus*
Pawlowski, Z.; and Chodera, L., 1972, Polski Tygod. Lekar., v. 27 (41), 1605-1607
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, humans, levo-tetramisole clinical trials successful except in trichuriasis
- Necator americanus*
Ripert, C.; and Avouac-Borzee, F., 1974, Medecine Afrique Noire, v. 21 (10), 713-715
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, human, epidemiology: Mirebalais, Haiti
- Necator americanus*
Rogers, S.; and Goldsmid, J. M., 1977, Ann. Trop. Med. and Parasitol., v. 71 (4), 503-504
Ternidens deminutus, man, indirect fluorescent antibody test evaluated for possible diagnostic use, some cross reactions with Necator americanus, promising epidemiologic tool
- Necator americanus*
Samuel, M. R., 1975, Progr. Drug Research, v. 19, 96-107
human intestinal helminths, review of clinical experiences world wide comparing the efficacy and tolerance of bitoscanate with that of bephenium hydroxynaphthoate and tetrachlorethylene; found to be most useful against hookworm with results against other helminths still inconclusive

- Necator americanus*
Senewiratne, B.; et al., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (2), 233-239
Necator americanus, human, tetrachlorethylene single dose compared with bephenium hydroxynaphthoate and pyrantel pamoate single and multiple doses, tetrachlorethylene remains drug of choice: Ceylon
- Necator americanus*, illus.
Smith, J. M., 1976, *Internat. J. Parasitol.*, v. 6 (1), 9-13
Ancylostoma tubaeforme and *Necator americanus* third stage larvae, ultrastructure of oesophageal glands compared before and after penetration through rabbit skin
- Necator americanus*
Sroczyński, J., 1977, *Polski Tygod. Lekar.*, v. 32 (16), 589-591
Necator americanus, [Wuchereria] bancrofti, Schistosoma mansoni, studies on hospitalized Africans to assess variations in blood picture during infections showed anemia in hookworm to be mainly iron deficiency while schistosomiasis caused protein deficiency, eosinophilia of peripheral origin rather than correlated with changes in bone marrow
- Necator americanus*
Taylor, M. M.; and Turton, J. A., 1976, *Tropenmed. u. Parasitol.*, v. 27 (1), 89-92
Necator americanus, cell-mediated immunity in man demonstrated by antigen-induced lymphocyte blastogenesis
- Necator americanus*
Turton, J. A., 1977, *Parasitology*, v. 75 (2), xxxvi [Abstract]
Necator americanus, human (exper.), 4 infections in one person over 2-year period, protective response not elicited (using egg count as criterion), anthelmintic activity of levamisole exceptionally varied (from 0 to 99%) but mebendazole always 100% effective, symptoms, blood picture, antibody production, lesions produced during dermal penetration
- Necator americanus*
Vakil, B. J.; et al., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (3), 247-250
human hookworm, clinical evaluation of C.9333 Go./CGP 4540, promising anthelmintic with few side effects: India
- Necator americanus*
Yeates, R. A.; and Ogilvie, B. M., 1976, *J. Immunol. Methods*, v. 12 (1-2), 57-65
Nippostrongylus brasiliensis, *Necator americanus*, measurement of antibodies to an unpurified enzyme (acetylcholinesterase) using an active-site directed radiolabel, cross-reactions between the two species
- Necator americanus* (Stiles, 1902), illus.
Yoshida, Y.; et al., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (4), 510-514
Necator americanus, scanning electron microscopic study of adult surface structures and infective-stage larvae
- Necator americanus*
Zuidema, P. J., 1971, *Nederl. Tijdschr. Geneesk.*, v. 115 (29), 1209-1212
man, case history of mixed infections of Entamoeba histolytica, *Necator americanus*, Plasmodium falciparum and Schistosoma mansoni, medical management: Netherlands (native of Surinam)
- Necator americanus*
Zuidema, P. J.; Rep, B. H.; and Meuzelaar, H. L. C., 1971, *Nederl. Tijdschr. Geneesk.*, v. 115 (45), 1886-1889
Ancylostoma ceylanicum and *Necator americanus*, mild infections in Dutch soldiers who had served in Surinam, clinical aspects: Netherlands
- Necator miyazakiensis* Nagayosi 1955
Yoshida, Y.; and Arizono, N., 1976, *J. Parasitol.*, v. 62 (5), 766-770
as syn. of Arthrostoma miyazakiense (Nagayosi 1955) comb. n.
- Necatoriasis
Lanari Zubiaur, F. J. B.; and Benavento de Beneventano, N. Y., 1974, *Medicina*, Buenos Aires, v. 34 (3), 249-252
electrocardiographic correlation between the double Master test and voluntary hyperventilation in patients with severe chronic hookworm anemia
- Nematod[a]
Ali, M. M., 1975, *Riv. Parassitol.*, Roma, v. 36 (2-3), 223-225
parasitic nematodes, comments on evolution and adaptation, abstract of thesis
- Nematoda
Andreiko, O. F., 1973, [Parasites of mammals of Moldavia], 184 pp., illus.
parasites of mammals, parasite lists, descriptions, host lists, ecology, geographic distribution, epidemiological and epizootiological distribution, monographic review: Moldavian SSR
- Nematoda
Atkinson, H. J., 1976, *Organ. Nematodes* (Croll), 243-272
respiratory physiology of nematodes, review: diffusion of oxygen into nematodes; factors influencing oxygen demand; oxygen availability; respiration in low oxygen regimes; respiratory function of haemoglobin in nematodes
- Nematoda
Barrett, J., 1976, *Organ. Nematodes* (Croll), 11-70
energy metabolism in nematodes, extensive review
- Nematoda
Bennet-Clark, H. C., 1976, *Organ. Nematodes* (Croll), 313-342
mechanics of nematode feeding: anatomy of nematode oesophageal pump; feeding performance; theory of filling and emptying of nematode oesophagus; a model for filling cycle of oesophagus of Ascaris

- Nematoda
Bird, A. F., 1976, Organ. Nematodes (Croll), 107-137
skeletal structures in nematodes (copulatory spicules, cuticle, egg shell): structure, chemical composition, ontogeny, function, review
- Nematoda
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
key to subclasses, orders, and superfamilies includes: Adenophorea; Secernentea
- Nematoda
Chitwood, B. G.; and Chitwood, M. B., 1974, Introduction to nematology, 334 pp., illus. introduction to nematology
- Nematod[a], larval
Croll, N. A., 1972, Zool. J. Linn. Soc., London, v. 51, Suppl. 1, 31-52
larval nematodes, behavior, review (nematode senses, locomotion, movement patterns as tracked on agar, mechanisms of orientation)
- Nematoda
Croll, N. A., 1975, Advances Parasitol., v. 13, 71-122
behavioral analysis of nematode movement, extensive review
- Nematoda
Croll, N. A., 1976, Organ. Nematodes (Croll), 343-364
behavioral coordination of nematodes, review: movements of the body and parts of the body; models of behavioral integration; sequential analysis; neurotransmitters and coordination
- Nematoda
Croll, N. A.; and Matthews, B. E., 1977, Biology of nematodes, 201 pp.
biology of nematodes
- Nematoda
Davey, K. G., 1976, Organ. Nematodes (Croll), 273-291
hormones in nematodes, review: neurosecretion; hormonal control of development; effect of exogenous hormones
- Nematoda
Evans, A. A. F.; and Perry, R. N., 1976, Organ. Nematodes (Croll), 383-424
survival strategies in nematodes, review: quiescence with special reference to cryptobiosis; diapause (in unhatched larvae; in larvae outside the egg; in adult stages; induction and termination; morphological and behavioral correlates)
- Nematoda
Evans, K.; and Forder, D., 1976, Nematologica, v. 22 (4), 475-476
automatic nematode counter
- Nematoda
Gabrisch, K., 1976, Prakt. Tierarzt, v. 57, Sondernummer, 37-40
parasites of reptiles, diagnosis, treatment, brief review
- Nematoda
Kazakov, B. E., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 43-52
helminths of vertebrates of tundra zones, biological peculiarities related to habitat, review
- Nematoda
Kunz, P.; and Klíngler, J., 1976, Nematologica, v. 22 (4), 477-479
nematode tracks, method for direct or microscopic observation and photography
- Nematoda
Lee, D. L.; and Atkinson, H. J., 1976, Physiology of nematodes. 2. ed. 215 pp., illus. physiology of nematodes, textbook
- Nematoda
Lindquist, W. D., 1970, Dis. Swine (Dunne), 3. ed., 708-744
swine, pathology, diagnosis, control, textbook
- Nematoda
Lindquist, W. D., 1975, Dis. Swine (Dunne), 4. ed., 780-815
helminths of swine, emphasis on nematodes, morphology, pathology, life cycle, diagnosis, treatment and control, review
- Nematoda
Lomakin, V. V., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 90-95
nematodes of fishes, zoogeographic analysis, review: Caspian Sea
- Nematoda
McLaren, D. J., 1976, Organ. Nematodes (Croll), 139-161
nematodes, morphology of sense organ/secretory cell relationship (amphidial and phasmidial sense organs, papillae), mechanics of secretion, experimental evidence for secretion, review
- Nematoda Cobb, 1919
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
taxonomic position among pseudocoelomate bilateria, should be considered as independent phylum
includes: Adenophorea; Secernentea
- Nematod[a]
Mueller, N. H. G., 1976, Gac. Vet., Buenos Aires (315), v. 38, 440-449
parasitic nematodes, inhibition of larval development, possible significance in cattle and sheep production; possibility of orienting anthelmintic treatment by bioclimatograph- and photoperiod-based forecast of onset of hypobiosis, review: Argentina

- Nematoda
Ollenschlaeger, B., 1975, *Fisch u. Umwelt* (1), 35-44
blood parasites of economically important fishes, species, importance, recommendations for therapy, review
- Nematoda
Otto, G. F., 1970, *Immun. Parasitic Animals* (Jackson, Herman and Singer), v. 2, 963-980
insect-borne nematodes, immunology, review
- Nematod[a]
Pouplard, L., 1976, *Ann. Med. Vet.*, v. 120 (8), 515-529
nematodes, anthelmintics in veterinary medicine, review
- Nematoda
Priadko, E. I., 1976, [Helminths of Cervidae] [Russian text], 228 pp., illus., maps
helminths of Cervidae, systematics, faunistics, parasite and host lists, zoogeographic and epizootiological aspects, control, extensive worldwide review
- Nematoda
Ramisz, A.; Urban, E.; and Surma, F., 1976, *Med. Wet.*, v. 32 (5), 291-293
Fasciola hepatica, gastro-intestinal nematodes, sheep, mixture of ranide and thiabendazole more effective than ranide alone
- Nematoda
Reichenbach-Klinke, H. H., 1975, *Fisch u. Umwelt* (1), 113-121
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Nematoda
Ryzhikov, K. M., 1975, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 25, 124-135
helminths of birds in Russia, number of species in each class of helminths, comparison with numbers worldwide, review of literature
- Nematoda
Smith, K., 1975, *J. Microscopy*, v. 105 (2), 229-232
method for concentrating nematode larvae from fecal material, procedure for processing for transmission electron microscopy
- Nematoda
Sood, M. L., 1977, *Research Bull. Panjab Univ.*, v. 24 (1-2), 1973, 33-41
classified list of nematodes from Indian freshwater fishes, review
- Nematoda
Thorson, R. E., 1970, *Immun. Parasitic Animals* (Jackson, Herman and Singer), v. 2, 913-961
direct-infection nematodes, immunology, review
- Nematoda
Wright, D. J.; and Newall, D. R., 1976, *Organ. Nematodes* (Croll), 163-210
nematodes, nitrogen catabolism and excretory products, osmotic and ionic regulation, excretory structures, review
- Nematoda
Wright, K. A., 1976, *Organ. Nematodes* (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Nematoda
Zuckerman, B. M., 1976, *Organ. Nematodes* (Croll), 211-241
free-living nematodes as models for aging studies, may have implications for parasitology
- Nematod[a sp.]
Anderson, M. M.; and McDaniel, J. S., 1975, *J. Elisha Mitchell Scient. Soc.*, v. 91 (2), 73
Peromyscus leucopus: eastern North Carolina
- Nematod[a sp.]
Arvy, L.; and Sowa, R., 1976, *Ann. Parasitol.*, v. 51 (1), 111-120
Ephemera danica: region de Cracovie, Pologne
- Nematod[a sp.]
Ashford, R. W., 1974, *J. Med. Entom.*, v. 11 (5), 605-616
Phlebotomus orientalis (terminal abdominal region, hemocoel, mouthparts): Ethiopia
- Nematod[a sp.] (undescribed)
Beveridge, I.; and Barker, I. K., 1976, *Austral. J. Zool.*, v. 24 (2), 265-272
helminths and arthropods, Antechinus stuartii, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (paraocloacal glands): Powelltown, Victoria
- Nematod[a spp.], illus.
Brinck-Lindroth, G.; and Smit, F. G. A. M., 1973, *Entom. Scand.*, v. 4 (4), 302-322
Amphipsylla sibirica sibirica from small mammals, castration and intersexuality due to hyperparasitism by nematodes, morphological changes
Amphipsylla sibirica sibirica: Stora Sjöfallet National Park and Vittangi area, Sweden; Straumsnes, Norway; Czechoslovakia
Palaeopsylla soricis: Stora Sjöfallet National Park, Sweden; Straumsnes, Norway
P. soricis rosickyi: Czechoslovakia
Rhadinopsylla integella: Stora Sjöfallet National Park, Sweden
R. pentacantha: Czechoslovakia
Malaraeus penicilliger: Stora Sjöfallet National Park, Sweden; Straumsnes, Norway
M. penicilliger kratochvili: Czechoslovakia
Megabothris rectangulatus: Stora Sjöfallet National Park, Sweden; Straumsnes, Norway
M. turbidus: Czechoslovakia
Amphipsylla sibirica hetera: Czechoslovakia
Parapsyllus cardinis: Antipodes Island
- Nematod[a sp.], illus.
Gwahaba, J. J., 1975, *J. East Africa Nat. Hist. Soc. and National Mus.* (153), 1-14
Corvus albus (abdominal cavity, synovial fluid cavity of tibiotarsal joint): Kampala, Uganda

- Nematod[a sp.]
 Irwin, S. W. B.; and Prentice, H. J., 1976, Irish Naturalists' J., v. 18 (9), 281-282
 Larus argentatus (digestive tract): Roe Island, Strangford Lough, County Down
- Nematod[a sp.], illus.
 Kapur, S.; et al., 1976, Neurol. India, v. 24 (2), 104-107
 human visceral larva migrans, nematode larva demonstrated in brain biopsy specimen of man presenting with severe neurologic impairment and stupor, neurologic symptoms attributed to allergic reaction to parasite presence, case report, biopsy material injected into mice (exper.), nematode larva later observed in mouse lung: India
- Nematod[a sp.] larva
 Lainson, R.; et al., 1976, Acta Amazonica, v. 6 (4), 55-60
 Psychodopygus complexus: Municipio of Aripuana, Mato Grosso State, Brazil
- Nematod[a sp.], unidentified
 Lank, D. R., jr., 1971, Proc. Indiana Acad. Sc., v. 81 (2), 359-364
 Rana catesbeiana: Indiana
- Nematod[a sp.]
 Lockard, L. L.; and Parsons, R. R., 1975, Great Basin Nat., v. 35 (4), 425-426
 Polyodon spathula (surface of stomach, pyloric caecum, intestine): Yellowstone River near Intake, Montana
- Nematoda [sp.]
 Maklakova, L. P., 1975, Trudy Gel'mint Lab., Akad. Nauk SSSR, v. 25, 102-106
 Succinea putris
 Cochlicopa lubrica
 Columella edentula
 Eulota fruticum
 all from Medynsk region, Kaluzhsk oblast
- Nematod[a sp.], adult worm, illus.
 Miyata, A.; and Tsukamoto, M., 1975, Nettai Igaku (Trop. Med.), v. 16 (3), 113-130
 Callosciurus juvencus (peripheral blood): Palawan Island, the Philippines
- Nematod[a sp.] larvae
 Mudry, D. R.; and Anderson, R. S., 1977, J. Fish Biol., v. 11 (1), 21-33
 Salmo gairdneri: Jasper and Banff National Parks, Canada
 Prosopium williamsoni: Waterton Lakes National Park, Canada
- Nematoda sp., larva
 Ponyi, J.; Biro, P.; and Murai, E., 1972, Parasitol. Hungar., v. 5, 383-408
 internal helminths of Acerina cernua (intestine), incidence survey, seasonal variations and host growth and development in relationship to parasitic burden: Lake Balaton, Hungary
- Nematod[a sp.]
 Smit, F. G. A. M., 1974, Senckenbergiana Biol., v. 55 (4-6), 357-398
 Palaeopsylla tauberi (abdominal cavity): Nepal
- Nematod[a sp.] nematode (lungs)
 Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
 Lepus americanus: insular Newfoundland
- Nematod[a sp.], possibly second or third stage filariae, illus.
 Stiller, D.; Sivanandam, S.; and Abu Hassan, R. B., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 447-448 [Demonstration] larval nematodes from Ornithodoros batuensis (hemocoele), a tick host-specific for the cave fruit bat (Eonycteris spelaea), search for possible transmission to bats unsuccessful: Dark Cave, Batu Caves, Kuala Lumpur
- Nematodiasis
 Stackhouse, L. L., 1977, J. Am. Vet. Med. Ass., v. 171 (9), 987-988
 cerebral nematodiasis in Alces alces, histopathologic features in brain compatible with cerebrospinal nematodiasis resulting from infection by Parelaphostrongylus tenuis: New Hampshire
- Nematodirella Yorke et Maplestone, 1926
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Molineidae, Nematodirinae
 synonymy
- Nematodirella alcidis (Dickmans, 1935) Ivaschkin, 1954
 Drozd, J.; and Bylund, G., 1970, Acta Parasitol. Polon., v. 17 (20-38), 259-260
 Alces alces (abomasa): Poland
- Nematodirella longispiculata
 Bergstrom, R. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 61-63
 Antilocapra americana: Wyoming
- Nematodirella longispiculata
 Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
 helminths of Alces alces, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Nematodirella longissimespiculata (= N. 1. longispiculata)
 Low, W. A., 1976, Canad. Field-Naturalist, v. 90 (2), 189-191
 Rangifer tarandus caribou (small intestine): Tweedsmuir Provincial Park, British Columbia
- Nematodirinae Skrjabin et Orloff, 1934
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Molineidae
 includes: Nematodirus (type genus); Lama-nema; Murielus; Nematodirella; Nematodiroides

- Nematodiroides Bernard, 1967
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Nematodirinae
- Nematodirus
Baines, D. M.; and Colegrave, A. J., 1977, Vet. Rec., v. 100 (11), 217-219
gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales
- Nematodirus
Bliss, D. H.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (10), 1612-1617
milk production in dairy cows exposed to mixed trichostrongylid larvae, results indicate that greatest milk loss occurs during the first 90 days of lactation, relationship between exposure time and stage of lactation
- Nematodirus
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Nematodirus
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Nematodirus
Buerger, H. J., 1976, Vet. Parasitol., v. 1 (4), 359-366
Ostertagia, Cooperia, Nematodirus, significantly higher numbers of larvae on herbage samples collected from calf pastures vs. cow pastures, improved control of trichostrongyle infection during late summer and autumn might be achieved by transfer of calves to cow pastures
- Nematodirus
Buerger, H. J., 1976, Ztschr. Parasitenk., v. 50 (2), 219
incidence on grass from cattle pasture: Niedersachsen
- Nematodirus
Chowaniec, W.; et al., 1975, Med. Wet., v. 31 (12), 741-743
Fasciola hepatica, Haemonchus, Trichostrongylus, Nematodirus, cattle, Nilzan, Zanil, field trials, good results
- Nematodirus
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxbendazole, drug efficacy
- Nematodirus
Dewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Nematodirus Ransom, 1907 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Nematodirinae
- Nematodirus
Forstner, M. J.; Kopp, H.; and Wiesner, H., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (9), 180-183
nematodes of ruminants, mebendazole, good results: Hellabrunn Zoo, Munich
- Nematodirus
Fudalewicz-Niemczyk, W.; et al., 1976, Acta Zootech., Bratislava (32), 5-19
gastrointestinal helminths, mountain sheep, nilverm and zanil, favorable influence on body weight and wool production of treated animals, no influence of treatment on fertility: Poland
- Nematodirus
Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Nematodirus
Henriksen, Sv. Aa.; et al., 1976, Vet. Parasitol., v. 2 (3), 259-272
gastro-intestinal nematodes, young calves during first grazing season, infection levels, blood findings, body weight gains, comparison of animals grazing same pasture entire season with those moved in early July and between levamisole-treated and untreated animals: Denmark
- Nematodirus
Lodha, K. R.; Raisinghani, P. M.; and Karwar, R. S., 1977, Indian J. Animal Sc., v. 47 (10), 677-682
helminths, camels, promintic and banminth II effective, nilverm inconsistent in action, thiabendazole ineffective
- Nematodirus
Malczewski, A.; et al., 1975, Med. Wet., v. 31 (12), 728-731
helminths, sheep, treatment with Nilverm and Nilzan more effective in May and November than in May and September, higher economic profit, increased weight gains and shearing yields: Olsztyn province

- [Nematodirus] nematodirusi
Nurtazin, A. T., 1975, Vestnik Sel'skokhoz. Nauki Kazakhstana (4), 84-86
sheep, trichostrongyles, anthelmintic efficacy of banminth, good results
- [Nematodirus] nematodiri
Panchin, O. G.; et al., 1975, Veterinariia, Kiev (40), 100-104
helminths and coccidia, sheep, seasonal incidence on pastures, measures for control: Kalanchats'k region, Kherson oblast
- Nematodirus
Sewell, M. M. H., 1973, Vet. Rec., v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Nematodirus
Tharaldsen, J., 1976, Acta Vet. Scand., v. 17, Suppl. 61, 1-21
trichostrongylid infections, calves, survival of larvae on pasture, occurrence of larvae not influenced by artificial irrigation; treatment with thiabendazole did not effectively control infection due to overwintering larvae, neither improved weight gain nor reduced egg production: Norway
- Nematodirus
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Nematodirus
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Nematodirus
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Nematodirus
Tsolov, B.; and Tsanov, G., 1975, Vet. Sbirka, v. 73 (9), 20-22
strongyloids of sheep, economics of tetramisole treatment, body weight, milk production, not effective against Trichocephalus
- Nematodirus
Zielinski, J., 1972, Med. Wet., v. 28 (9), 566-567
parasites, sheep, Nilverm, copper sulfate
- Nematodirus spp., 4th stage larvae
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, Trichostrongylus, and Nematodirus, oxfendazole highly effective against adult stages in sheep
- Nematodirus sp.
Bergstrom, R. C.; Maki, L. R.; and Werner, B. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 171-174
trichostrongylid eggs in cattle or sheep feces, dung beetles (Aphodius spp.; Canthon practicola) as possible biological control agents, laboratory studies showed decreased eggs in feces when beetles were present
- Nematodirus spp.
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Nematodirus spp.
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Nematodirus spp.
Düewel, D.; et al., 1974, Prakt. Tierarzt, v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Nematodirus sp.
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, Vet. Sbirka, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of tetramisole, group and individual applications
- Nematodirus spp.
Henriksen, S. A.; Bentholt, B. R.; and Nielsen-Englyst, A., 1976, Nord. Vet.-Med., v. 28 (4-5) 201-209
gastro-intestinal strongyles, cattle, seasonal distribution on pastures
- Nematodirus spp.
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Nematodirus sp.
Le Jambre, L. F.; and Royal, W. M., 1976, Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (fecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except Nematodirus resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales
- Nematodirus sp.
Low, W. A., 1976, Canad. Field-Naturalist, v. 90 (2), 189-191
Rangifer tarandus caribou (small intestine): Tweedsmuir Provincial Park, British Columbia

- Nematodirus* spp.
Lukovich, R.; et al., 1977, *Gac. Vet.*, Buenos Aires (318), v. 39, 91-95
helminths, cattle, levamisole, results from injectable and dermal application similar
- Nematodirus* sp.
Lyons, E. T.; et al., 1975, *Am. J. Vet. Research*, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Nematodirus* sp.
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Equus caballus: Chile
- Nematodirus* spp.
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Nematodirus* spp.
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Nematodirus* spp.
Vlassoff, A., 1976, *N. Zealand J. Exper. Agric.*, v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand
- Nematodirus* [sp.]
Volf, K.; and Volfova, M., 1974, *Veterinarstvi*, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Nematodirus* spp.
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, *Vet. Glasnik*, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Nematodirus* sp.
Williams, J. C.; and Knox, J. W., 1976, *Am. J. Vet. Research*, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Nematodirus* sp.
Young, E.; et al., 1973, *Research J. National Parks Republic South Africa* (16), 195-198
Antidorcas marsupialis (duodenum): Mountain Zebra National Park near Cradock, Cape Province
- Nematodirus abnormalis*
Corticelli, B.; and Lai, M., 1972, *Parassitologia*, v. 14 (1), 95-96
Ovis musimon (tenue): Sardegna
- Nematodirus abnormalis* May, 1920
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Nematodirus abnormalis*
Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Nematodirus abnormalis*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, *Am. J. Vet. Research*, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Nematodirus abnormalis*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Nematodirus abnormalis* May, 1920
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Nematodirus battus*
Boag, B.; and Thomas, R. J., 1973, *Research Vet. Sc.*, v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Nematodirus battus*
Boag, B.; and Thomas, R. J., 1975, *Research Vet. Sc.*, v. 19 (3), 263-268
Nematodirus battus vs. *N. filicollis*, sheep, epidemiological studies over a 3 year period under field conditions starting from clean pasture, annual increase in infection levels, persistence of infection despite pasture rest, consistent difference in larval pattern between the two species, possibility of control by plowing and reseeded or by alternate grazing with cattle

- Nematodirus battus**
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Nematodirus battus**
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastro-intestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Nematodirus battus**
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Nematodirus battus**
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Nematodirus battus**
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Nematodirus battus, illus.**
Martin, J.; and Lee, D. L., 1976, Parasitology, v. 72 (1), 75-80
Nematodirus battus, appearance of large hexagonal crystals blocking intestine, lipoprotein in composition, apparently associated with development of immunity to this nematode in lambs
- Nematodirus battus**
Parkin, J. T., 1976, Parasitology, v. 73 (3), 343-354
Nematodirus battus, egg development and hatching, effect of variations in humidity and osmotic pressure
- Nematodirus battus, illus.**
Perry, R. N., 1977, Parasitology, v. 74 (2), 133-137
Nematodirus battus larvae, reassessment of variations in water content during hatching process
- Nematodirus battus**
Reid, J. F. S., 1976, Vet. Rec., v. 98 (25), 496-499
gastrointestinal nematodes, coccidiosis, diarrhea of sheep, age and seasonal factors: Britain
- Nematodirus battus**
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Nematodirus battus**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Nematodirus europaeus**
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Nematodirus filicollis**
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, Trichostrongylus, and Nematodirus, oxfendazole highly effective against adult stages in sheep
- Nematodirus filicollis (Rudolphi, 1802)**
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Nematodirus filicollis**
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Nematodirus filicollis**
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 263-268
Nematodirus battus vs. N. filicollis, sheep, epidemiological studies over a 3 year period under field conditions starting from clean pasture, annual increase in infection levels, persistence of infection despite pasture rest, consistent difference in larval pattern between the two species, possibility of control by plowing and reseeded or by alternate grazing with cattle
- Nematodirus filicollis**
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species

- Nematodirus filicollis*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Nematodirus filicollis*
Corticelli, B.; and Lai, M., 1972, Parasitologia, v. 14 (1), 95-96
Ovis musimon (tenue): Sardegna
- Nematodirus filicollis*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Nematodirus filicollis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Nematodirus filicollis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Nematodirus filicollis*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitology, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Nematodirus filicollis*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Nematodirus filicollis*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Nematodirus filicollis*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Nematodirus filicollis*
Gibson, T. E.; and Everett, G., 1976, Research Vet. Sc., v. 20 (2), 158-161
Nematodirus filicollis, development and survival of eggs placed on grass plots over a period of a year, extraordinary persistence of eggs and larvae under weather conditions of southern England makes control difficult
- Nematodirus filicollis*
Gonzalez, H.; and Plaza, J., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 134-137
gastrointestinal nematodes of sheep, comparative therapeutic trials using banminth, phenothiazine, and thiabendazole
- Nematodirus filicollis* (Rudolphi, 1802) Ransom, 1907
Ianchev, I., 1973, Izvest. Tsentral. Khelmin. Lab., v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Nematodirus filicollis*
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Nematodirus filicollis*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Nematodirus filicollis*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Nematodirus filicollis*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Nematodirus filicollis*
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Nematodirus filicollis*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure

- Nematodirus filicollis*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Nematodirus filicollis*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Ovis aries: Chile
- Nematodirus filicollis*
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Nematodirus filicollis*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Nematodirus filicollis*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Nematodirus filicollis*
Ramajo Martin, V.; and Simon Vicente, F., 1975, Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C., v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Nematodirus filicollis*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Nematodirus filicollis*
Schweigsut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Nematodirus filicollis*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Nematodirus filicollis*
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Nematodirus helvetianus*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Nematodirus helvetianus*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Nematodirus helvetianus*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Nematodirus helvetianus*
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Nematodirus helvetianus*
McBain, D. G.; et al., 1977, Vet. Rec., v. 101 (14), 285-286
helminths, calves, fenbendazole in feed blocks
- Nematodirus helvetianus*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Nematodirus helvetianus*
Pfeiffer, H.; and Supperer, R., 1976, Berl. u. Munchen. Tierarztl. Wchnschr., v. 89 (13), 249-252
Nematodirus helvetianus, cattle, Fenbendazol effective against adults and growing larvae but not against inhibited fourth stage larvae
- Nematodirus helvetianus*
Randall, R. W.; and Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- [*Nematodirus*] *helvetianus*
Reinecke, R. K., 1972, Onderstepoort J. Vet. Research, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure

- Nematodirus helvetianus*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Nematodirus helvetianus*
Rose, J. H., 1973, Research Vet. Sc., v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostromylus rubidus*, culture from infective larva to adult worm in WAe medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Nematodirus helvetianus*
Rose, J. H., 1975, Research Vet. Sc., v. 18 (2), 175-177
Nematodirus helvetianus, eggs which survive on pastures throughout winter are able to transmit infection to calves turned on to pasture in early summer but such calves did not exhibit symptoms of clinical disease, concluded that such eggs are unlikely to be associated with disease outbreaks
- Nematodirus helvetianus*
Wikerhauser, T.; et al., 1974, Acta Parasitol. Iugoslavica, v. 5 (2), 79-81
trichostrongylids, cattle, fenbendazole compared with thiabendazole, good results from both
- Nematodirus lamae*
Guerrero, C.; Rojas, M.; and Vargas, J., 1974, Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity of 1-tetramisole, significant body weight gain in treated animals
- Nematodirus lamae*
Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Nematodirus lamae*
Vargas, J.; Guerrero, C.; and Rojas, M., 1972, Rev. Invest. Pecuarias, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight toxicity
- Nematodirus odocoilei*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Nematodirus roscidus*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Nematodirus spathiger*
Anderson, P. J. S.; and Marais, F. S., 1972, J. South African Vet. Ass., v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Nematodirus spathiger*
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, *Trichostrongylus*, and *Nematodirus*, oxfendazole highly effective against adult stages in sheep
- Nematodirus spathiger* (Railliet, 1896)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Nematodirus spathiger*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Nematodirus spathiger*
Ciordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Nematodirus spathiger*
Coles, G. C.; and McNeillie, R. M., 1977, J. Helminth., v. 51 (4), 323-326
dietary feeding of drugs for 5 days to mice infected with *Nematospiroides dubius* and *Hymenolepis nana*, detected all modern anthelmintics examined except stilbazium; simple test using *Nematodirus spathiger* eggs and *Nippostrongylus brasiliensis* adults to detect anthelmintics in vitro
- Nematodirus spathiger*
Coles, G. C.; and Simpkin, K. G., 1977, Research Vet. Sc., v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Nematodirus spathiger*
Colglazier, M. L.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with *Trichuris* spp. and *Moniezia expansa*

- Nematodirus spathiger*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastro-intestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Nematodirus spathiger*
Dalton, S. E., 1977, Parasitology, v. 75 (2), xvi [Abstract]
Haemonchus contortus, *Nematodirus spathiger*, effect of thiophanate on egg output, hatchability, and worm burden, sheep
- Nematodirus spathiger*
Dineen, J. K.; et al., 1977, Internat. J. Parasitol., v. 7 (3), 211-215
Trichostrongylus colubriformis-vaccinated sheep, high level of protection against single-species homologous challenge, lowered level of protection against single-species challenge with *T. vitrinus*, no protection against single-species challenge with *Nematodirus spathiger*, high level of protection against all 3 species to simultaneous challenge with all 3 species, latter suggests that terminal effectors of resistance are immunologically non-specific
- Nematodirus spathiger*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Nematodirus spathiger*
Groeneveld, H. T.; and Reinecke, R. K., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 285-297
non-parametric statistical method for comparing worm burdens in two groups of sheep, application in interpreting results of anthelmintic tests
- Nematodirus spathiger* (Railliet, 1896) Railliet et Henry, 1909
Ianchev, I., 1973, Izvest. Tsentral. Khelmin. Lab., v. 16, 205-220
Capreolus capreolus (small intestine): southern Bulgaria
- Nematodirus spathiger*
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Nematodirus spathiger*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Nematodirus spathiger*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Nematodirus spathiger*
Knight, R. A., 1977, J. Parasitol., v. 63 (5), 957-958
Trichostrongylus affinis, *Nematodirus spathiger*, rabbits (exper.), effect of dexamethasone (higher egg counts but no difference in numbers of worms that developed, did not prevent worm expulsion)
- Nematodirus spathiger*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Nematodirus spathiger*
Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Nematodirus spathiger*
Martinez Gomez, F.; and Hernandez Rodriguez, S., 1973, Rev. Iber. Parasitol., v. 33 (1), 11-20
Ovis aries (duodenum): Cordoba, Spain
- Nematodirus spathiger*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Nematodirus spathiger*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Nematodirus spathiger*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Nematodirus spathiger*
Ramajo Martin, V.; and Simon Vicente, F., 1975, Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C., v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Nematodirus spathiger* (Railliet, 1896), *illus.*
Rivellini, P.; Guarino, C.; and Signoretta, N., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 492-494
Ovis aries: Campania

- Nematodirus spathiger*
Simpkin, K. G.; and Coles, G. C., 1976, Parasitology, v. 73 (2), iv [Abstract]
Nematospiroides dubius, *Nippostrongylus brasiliensis*, *Nematodirus spathiger*, modes of action of thiabendazole and mebendazole apparently different from those reported on other species
- Nematodirus spathiger* (Railliet, 1896)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Nematodirus spathiger*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Nematodirus spathiger*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Nematodirus spathiger*
Theodorides, V. J.; Nawalinski, T.; and Chang, J., 1976, Am. J. Vet. Research, v. 37 (12), 1515-1516
gastrointestinal nematodes, *Moniezia* spp., sheep, albendazole highly effective
- Nematodirus spathiger*
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antelope cervicapra (small intestine): Texas
- Nematodirus spathiger*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Nematodirus spathiger*
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, *Camelus dromedarius*, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Nematodirus spathiger* (Railliet, 1896)
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damaliscus dorcas dorcas (small intestine): captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Nematodirus spathiger* (Railliet, 1896)
Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karoo
- Nematodirus spathiger*
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 77-81
Redunca fulvorufula
Damaliscus dorcas phillipsi (duodenum of all): all from Mountain Zebra National Park
- Nematodirus triangularis* Boughton, 1932
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Lepus americanus: insular Newfoundland
- Nematomorpha, illus.*
Lanzavecchia, G.; Valvassori, R.; and De Eguileor, M., 1977, J. Molecular Biol., v. 111 (3), 371-374
Nematomorpha, existence of bipolarity in thick myofilaments, possible implications for mechanism of muscle contraction
- Nematomorpha* sp.
Arvy, L.; and Sowa, R., 1976, Ann. Parasitol., v. 51 (1), 111-120
Ecdyonurus starmachi: region de Cracovie, Pologne
- Nematospira* Walton, 1923
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Heligmosomoides* Hall, 1916
- Nematospiroides* Baylis, 1926
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Heligmosomoides* Hall, 1916
- Nematospiroides dubius*
Behnke, J. M., 1977, Parasitology, v. 75 (2), xv [Abstract]
Nematospiroides dubius, inhibition of larval development in immune mice, transfer of immunity by immune serum and syngeneic mesenteric lymph node cells
- Nematospiroides dubius*
Behnke, J. M.; and Wakelin, D., 1977, J. Helminthol., v. 51 (3), 167-175
Nematospiroides dubius, stimulation of acquired immunity in inbred strains of mice
- Nematospiroides dubius*
Behnke, J. M.; Wakelin, D.; and Wilson, M. M., 1977, Parasitology, v. 75 (2), xxxiv-xxxv [Abstract]
interactions between intestinal phase of *Trichinella spiralis* and *Nematospiroides dubius*
- Nematospiroides dubius*
Boisvenue, R. J.; Emmick, T. L.; and Galloway, R. B., 1977, Exper. Parasitol., v. 42 (1), 67-72
Haemonchus contortus, some compounds with juvenile hormone activity inhibited in vitro development of infective larvae, none of these compounds had anthelmintic properties against *Ascaris suum* or *Nematospiroides dubius* in mice

- Nematospiroides dubius*
Chaicumpa, S.; et al., 1977, Austral. J. Exper. Biol. and Med. Sc., v. 55 (4), 393-400
Nematospiroides dubius, mice, immunization with live third-stage larvae given orally, intravenously, intraperitoneally, or subcutaneously
- Nematospiroides dubius*
Chaicumpa, V.; Jenkin, C. R.; and Fischer, H., 1977, Austral. J. Exper. Biol. and Med. Sc., v. 55 (5), 561-570
Nematospiroides dubius, effect in vivo of peritoneal exudate cells of immune and normal mice on infectivity of third stage larvae
- Nematospiroides dubius*
Chaicumpa, V.; Jenkin, C. R.; and Rowley, D., 1976, Austral. J. Exper. Biol. and Med. Sc., v. 54 (3), 245-252
Nematospiroides dubius, no passive transfer of immunity to normal mice by serum from immune mice, passive transfer of immunity from immunized mice to their offspring, immunity dependent on intake of immunoglobulin via milk for period longer than 24 hours, passive transfer of immunity from immune mothers to neonatal mice does not appear to be dependent on a specific class of immunoglobulins
- Nematospiroides dubius*
Coles, G. C.; and McNeillie, R. M., 1977, J. Helminth., v. 51 (4), 323-326
dietary feeding of drugs for 5 days to mice infected with *Nematospiroides dubius* and *Hymenolepis nana*, detected all modern anthelmintics examined except stibazium; simple test using *Nematodirus spathiger* eggs and *Nippostrongylus brasiliensis* adults to detect anthelmintics in vitro
- Nematospiroides dubius*
Cremers, H. J. W. M.; Jansen, J.; and Swierstra, D., 1975, Tijdschr. Diergeneesk., v. 100 (22), 1209-1211
Mus musculus: Netherlands
- Nematospiroides dubius*
Cypess, R. H.; et al., 1974, J. Infect. Dis., v. 130 (5), 534-538
Nematospiroides dubius, influence of parasitic infection in mouse (exper.) on enteric colonization and immune response to *Escherichia coli*
- Nematospiroides dubius*
Damian, R. T., 1976, J. Parasitol., v. 62 (1), 168-169
Nematospiroides dubius, separation and cleaning of infective larvae and eggs from gross fecal contaminants, centrifugation on Ficoll-Isopaque cushion
- Nematospiroides dubius*
Della Bruna, C.; and Xenia, B., 1976, J. Parasitol., v. 62 (3), 490-491
Nippostrongylus brasiliensis: reduced worm burden and prolonged infection in mice harboring *Nematospiroides dubius*
- Nematospiroides dubius*
Dobson, C.; and Owen, M. E., 1977, Internat. J. Parasitol., v. 7 (6), 463-466
Nematospiroides dubius, influence of serial passage on infectivity and immunogenicity in mice
- Nematospiroides dubius*
Duwel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Nematospiroides dubius*
Gavaghan, A. D.; and Nunn, A. J., 1972, Pharm. Acta Helveticae, v. 47 (11-12), 719-726
Nippostrongylus brasiliensis, *Strongyloides ratti*, *Nematospiroides dubius*, synthesis of thiocyanine and hemithiocyanine derivatives of benzothiazole, significant anthelmintic activity in laboratory trials with mice
- Nematospiroides dubius*
Hepler, D. I.; and Lueker, D. C., 1976, Abst. Ann. Meet. Am. Soc. Microbiol., 76
Nematospiroides dubius, mice (exper.), cellular and humoral response after oral immunization
- Nematospiroides dubius*
Hepler, D. I.; and Lueker, D. C., 1976, Experientia, v. 32 (3), 386-387
Nematospiroides dubius, high degree of native resistance of *Peromyscus maniculatus* to infection, infection only established with use of steroid treatment
- Nematospiroides dubius*
Hepler, D. I.; Lueker, D. C.; and Rubin, R., 1976, J. Parasitol., v. 62 (3), 491-492
Nematospiroides dubius, vaccination, immune response of outbred mouse strains stronger than inbred ones, oral route of administering larvae superior to subcutaneous route, steroid hormones blocked expression of immunity in subcutaneously vaccinated mice but not in orally vaccinated ones
- Nematospiroides dubius*
Hosier, D. W.; and Durning, J. P., 1975, J. Parasitol., v. 61 (3), 564-566
Nematospiroides dubius, male vs. female ICR mice challenged with 200 larvae after receiving one stimulating infection of 400 larvae, effect of gonadectomy or supplemental sex hormone treatment on worm burden
- Nematospiroides dubius*
Howes, H. L., jr., 1972, Proc. Soc. Exper. Biol. and Med., v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that *T. muris*-mouse infection could be test model for antiwhipworm studies

- Nematospiroides dubius*
Husain, M. I.; and Agarwal, S. K., 1975, Indian J. Chem., v. 13 (11), 1238-1239
N-(4-N,N-diethylaminobenzylidene)-p-(4-methyl-1-piperazino)aniline shows significant activity in laboratory screening trials with *Nematospiroides dubius* infection in mice, comparison with tetramisole
- Nematospiroides dubius*
Jenkins, D. C., 1977, Exper. Parasitol., v. 41 (2), 335-340
Nematospiroides dubius, course of primary and challenge infections in male and female *Meriones unguiculatus*, rate of establishment, morphology, sex ratio and distribution within host intestine, expulsion in primary infections, resistance to challenge infections, lactating jirds with depressed immunocompetence were significantly more susceptible to reinfection than nulliparous jirds of same age
- Nematospiroides dubius*
Jenkins, S. N.; and Behnke, J. M., 1977, Parasitology, v. 75 (1), 71-78
Trichuris muris, mice, primary immune expulsion markedly delayed by concurrent infection with *Nematospiroides dubius*, possible relevance in pathogenesis of concurrent tropical diseases
- Nematospiroides dubius*
Jenkins, S. N.; and Behnke, J. M., 1977, Parasitology, v. 75 (2), xxxiv [Abstract]
Trichuris muris, delay of primary expulsion in mice concurrently infected with *Nematospiroides dubius*
- Nematospiroides dubius* (Baylis, 1926)
Lewis, J. W.; and Bryant, V., 1976, J. Helminth., v. 50 (3), 163-171
Nematospiroides dubius, distribution within small intestine of mice up to 60 days post-infection, relation of establishment and pattern of distribution to host age and sex, degree of aggregation of worm populations with respect to host intestine and to each other
- Nematospiroides dubius*
Lewis, J. W.; and Shava, F. H. M., 1977, Parasitology, v. 75 (2), iv [Abstract]
Syphacia obvelata, *Nematospiroides dubius*, differences in periodicity of egg deposition can be correlated with differences in transmission of infective stages to definitive host
- Nematospiroides dubius*
Long, R. A.; Ellis, W. L.; and Taylor, G. R., 1976, Texas J. Sc., v. 27 (1), 163-172
Nematospiroides dubius, response to deep space environment of Apollo 16 manned space-flight, reduced hatching rate of eggs, unchanged infectivity to mice
- Nematospiroides dubius* Baylis, 1962
Manger, B. R., 1976, Parasitology, v. 73 (2), xiii-xiv [Abstract]
Nematospiroides dubius, mice, anthelmintic-terminated immunizing infections (Cambendazole selected as most active), variation in degree of protection between host strains, timing of termination of immunizing infection indicated exsheathment per se not essential in production of resistance, protection could not be transferred with serum alone
- Nematospiroides dubius*
Murua, R., 1975, J. Helminth., v. 49 (4), 293-296
Nematospiroides dubius larvae, hatching time, rate of development to third stage, optimum hatching and development temperature
- Nematospiroides dubius*
Roberts, T. M.; and Thorson, R. E., 1977, J. Parasitol., v. 63 (4), 764-766
Nippostrongylus brasiliensis, pairing between adults and other species of nematodes in vitro, comparison of intra and interspecific response patterns, results suggest that pairing is not related entirely to sexual attraction
- Nematospiroides dubius*
Scofield, A. M., 1975, Comp. Biochem. and Physiol., v. 52 (4A), 685-689
Nematospiroides dubius, rats, transient malabsorption of amino acids by small intestine during infection
- Nematospiroides dubius*
Sharpe, M. J.; and Lee, D. L., 1977, Parasitology, v. 75 (2), xvi [Abstract]
Trichostrongylus colubriformis, *Nematospiroides dubius*, effect of levamisole on adenylate energy charge and on levels of acetylcholinesterase
- Nematospiroides dubius*
Simpkin, K. G.; and Coles, G. C., 1976, Parasitology, v. 73 (2), iv [Abstract]
Nematospiroides dubius, *Nippostrongylus brasiliensis*, *Nematodirus spathiger*, modes of action of thiabendazole and mebendazole apparently different from those reported on other species
- Nematospiroides dubius*
Striebel, H. P., 1976, Experientia, v. 32 (4), 457-458
4-isothiocyanato-4'-nitrodiphenylamine, an anthelmintic with an unusual spectrum of activity against intestinal nematodes, filariae and schistosomes
- Nematospiroides dubius*
Takats, C., 1972, Parasitol. Hungar., v. 5, 203-216
human helminths, 171 compounds tested on laboratory animals for possible anthelmintic action, techniques of screening procedures described

- Nematospiroides dubius*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Nematospiroides dubius*, illus.
Wright, E. J.; and Sommerville, R. I., 1977, Internat. J. Parasitol., v. 7 (5), 353-359
Nematospiroides dubius, spermatozoa, description and location, movement in vitro, speed, locomotion not considered to be amoeboid but resembles movement of monopodial neutrophils
- Neoapectana*
Mracek, Z., 1977, J. Invert. Path., v. 30 (1), 87-94
diagnostic characters distinguishing *Steinernema* from *Neoapectana*
- Neoapectana* sp.
Finney, J. R.; and Walker, C., 1977, J. Invert. Path., v. 29 (1), 7-9
Neoapectana sp., DD-136 strain, possible biological control agent for European elm bark beetle, *Scolytus scolytus*, logs sprayed with nematodes in solutions of either distilled water, glycerin, or P.B.I. Wetting agent
- Neoapectana carpocapsae*
Andreadis, T. G.; and Hall, D. W., 1976, Exper. Parasitol., v. 39 (2), 252-261
Neoapectana carpocapsae, encapsulation and melanization in *Aedes aegypti*, changes in host hemocytes and hemolymph proteins, distribution of DOPA-oxidase within hemocytes, hemocyte changes as possible pathological condition created by parasite, possible function of protein in host defense reactions
- Neoapectana carpocapsae*
Beresky, M. A.; and Hall, D. W., 1977, J. Invert. Path., v. 29 (1), 74-80
Neoapectana carpocapsae-infected *Aedes aegypti* larvae, phenylthiourea treatment reduced nematode encapsulation and melanization and mosquito mortality, possible explanations
- Neoapectana carpocapsae* Weiser
Kamionek, M., 1977, Bull. Acad. Polon. Sc., Cl. II., s. Sc. Biol., v. 25 (4), 243-246
Neoapectana carpocapsae in *Hylemya antiqua*, *Pegomya hyoscyami*, *Hyponomeuta malinellus*, and *Aphis fabae* (all exper.), host susceptibility, percentage and rate of mortality, intensity of infection
- Neoapectana carpocapsae* Weiser
Kamionek, M.; and Sandner, H., 1977, Bull. Acad. Polon. Sc., Cl. II., s. Sc. Biol., v. 25 (4), 247-249
Neoapectana carpocapsae in *Sitophilus oryzae*, *Tribolium castaneum*, and *Trogoderma granarium* (all exper.), increased pathogenicity when 2 insect species were placed together as compared to those kept singly
- Neoapectana carpocapsae*
Kamionek, M.; and Seryczynska, H., 1976, Bull. Acad. Polon. Sc., Cl. II., s. Sc. Biol., v. 24 (8), 483-485
Neoapectana carpocapsae, changes in hemocyte ultrastructure of *Galleria mellonella*
- Neoapectana carpocapsae* Weiser, illus.
de Luca, Y., 1976, Rev. Zool. Agric. et Path. Veg., v. 75 (4), 127-131
Neoapectana carpocapsae, biological control of *Acanthoscelides obtectus* adults
- Neoapectana carpocapsae*, illus.
Triggiani, O.; and Poinar, G. O., jr., 1976, J. Invert. Path., v. 27 (3), 413-414
entry of *Neoapectana carpocapsae* in *Phryganidia californica* and *Galleria mellonella* via host tracheal system
- Neoapectana dutkyi*
Prakasa Rao, P. S.; Das, P. K.; and Padhi, G., 1977, Indian J. Agric. Sc., v. 45 (6), 1975, 275-277
Neoapectana dutkyi, compatibility with some insecticides and fertilizers, useful in control of rice insects
- Neoscaris* Travassos, 1927
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Toxocara* Stiles, 1905
- Neoscaris* sp.
Schulte, J. W.; Klimstra, W. D.; and Dyer, W. G., 1976, J. Wildlife Management, v. 40 (3), 579-581
Odocoileus virginianus clavium (feces): Big Pine Key, Florida
- Neoscaris vitulorum*
Chauhan, H. V. S.; Dwivedi, P.; and Kalra, D. S., 1974, Haryana Vet., v. 13 (1), 5-21
protozoan and helminth parasites, transmitted through milk to newborn animals, review
- Neoscaris vitulorum* (Goeze, 1782) Travassos, 1927, illus.
Chauhan, P. P. S.; et al., 1977, Indian J. Animal Sc., v. 44 (10), 1974, 801-804
Neoscaris vitulorum in albino rats and chickens (both exper.), migratory behaviour, histopathology
- Neoscaris vitulorum*
El-Abdin, Y. Z.; Mossalam, I.; and Hamza, S. M., 1975, Egypt. J. Vet. Sc., v. 12 (2), 143-152
Neoscaris vitulorum, buffalo calves, blood picture, biochemical blood constituents, enzyme activities, before and after treatment with Concurat
- Neoscaris vitulorum*
Gautam, O. P.; Malik, P. D.; and Singh, D. K., 1976, Current Sc., Bangalore, v. 45 (9), 350-351 [Letter]
Neoscaris vitulorum larvae, transmammmary infection, buffalo calves; larvae found in milk and colostrum

- Neoscaris vitulorum*
Graves, I. L.; Adams, W. H.; and Pyakural, S., 1975, *Am. J. Vet. Research*, v. 36 (6), 843-846
Babesia bigemina, Bos grunniens moved from high to low altitude and challenged with influenza A viruses, hemolytic anemia, possible explanations, death due to Fasciola hepatica and F. gigantica, incidental finding of Bunostomum sp., Trichuris sp., Neoscaris vitulorum, Dictyocaulus sp., coccidia, some reasons for poor survival of yaks at low altitude: Nepal
- Neoscaris vitulorum*
Gupta, G. C.; Joshi, B. P.; and Rai, P., 1976, *Indian Vet. J.*, v. 53 (6), 438-441
Neoscaris vitulorum, buffalo calves, pathogenesis, biochemical alterations in host blood suggest severe hepatic insufficiency
- Neoscaris vitulorum*
Ismail, E.; Tawfik, A. A.; and El-Ebrashi, N. M. A., 1977, *Arzneimittel-Forsch.*, v. 27 (7), 1393-1394
6-hydroxy-4-methoxy-5-(p-methoxy-cinnamoyl)-benzofuran, in vitro broad spectrum anthelmintic activity against livestock helminths, promising results indicate need for future research
- Neoscaris vitulorum*
Krvavica, S.; et al., 1976, *Vet. Arhiv*, Zagreb, v. 46 (9-10), 215-229
Paramphistomum microbothrium, Neoscaris vitulorum, activity of enzymes taking part in glucose decomposition; anaerobic metabolism; metabolic pathways
- Neoscaris vitulorum*
Krvavica, S.; Francetic, D.; and Zivkovic, D., 1976, *Vet. Arhiv*, Zagreb, v. 46 (9-10), 231-239
nematodes, trematodes, cestodes, activity, distribution and cofactor dependence of malic enzymes, majority are located in mitochondria in all investigated parasites
- Neoscaris vitulorum*
Mozgovoi, A. A.; and Shakhmatova, V. I., 1969, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 20, 97-101
Neoscaris vitulorum, description of larval stages, life cycle, development and migration
[Mus musculus]
[Cavia porcellus]
[Rat, white]
[Ovis aries] (liver, lungs)
[Bos] (liver, lungs, amniotic fluid)
[Leporidae]
(all exper.)
- Neoscaris vitulorum*
Schneider, C. R.; et al., 1975, *Ann. Trop. Med. and Parasitol.*, v. 69 (2), 227-232
Bubalus bubalis: Khong Island, Laos
- Neoscaris vitulorum*, illus.
Tawfik, M. A. A., 1975, *Egypt. J. Vet. Sc.*, v. 12 (2), 85-88
Neoscaris vitulorum, white rats, pathological changes induced by migration (ileum, caecum, liver and lung)
- Neoscaris vitulorum*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Neoscaris vitulorum*
Troncy, P. M.; and Oumate, O., 1973, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 26 (2), 199-202
Neoscaris vitulorum, Strongyloides papillosus, Zebu calves, morantel tartrate: Tchad
- Neoscarophis n. g.*
Machida, M., 1976, *Bull. National Sc. Mus.*, Tokyo, s. A, Zool., v. 2 (1), 1-6
Rhabdochonidae, Ascarophidinae
tod: N. yarihige n. sp.
- Neoscarophis bathygadi n. sp.*, illus.
Machida, M., 1976, *Bull. National Sc. Mus.*, Tokyo, s. A, Zool., v. 2 (1), 1-6
Bathygadus garretti (intestine): Suruga Bay, Japan
- Neoscarophis yarihige n. g.*, n. sp. (tod), illus.
Machida, M., 1976, *Bull. National Sc. Mus.*, Tokyo, s. A, Zool., v. 2 (1), 1-6
Coelorrhynchus multispinulosus (intestine): Suruga Bay, Japan
- Neocamallanus* Ali, 1957
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of Paracamallanus Yorke & Maplestone, 1928
- Neocamallanus sensu* Chakravarty et al., 1962, not Ali, 1957
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of Procammallanus Baylis, 1923
- Neocamallanus atridentus* Khera, 1954
Sahay, U., 1972, *Indian J. Animal Research*, v. 6 (2), 97-98
Syn.: N. bengalensis Soota & Chaturvedi, 1971
- Neocamallanus bengalensis* Soota & Chaturvedi, 1971
Sahay, U., 1972, *Indian J. Animal Research*, v. 6 (2), 97-98
as syn. of Neocamallanus atridentus Khera, 1954
- Neocamallanus sindensis n. sp.*, illus.
Akram, M., 1976, *Pakistan J. Zool.*, v. 8 (1), 93-95
Channa punctatus (stomach, intestine): Sind region

- Neocruzia morleyi* (Pearse, 1936) Yamaguti, 1961, illus.
Caballero Deloya, J., 1974, An. Inst. Biol., Univ. Nac. Mexico, s. Zool., v. 45 (1), 45-50
redescription, syn.: *Cruzia morleyi* Pearse, 1936
Bufo horribilis (intestino, cloaca): Laguna de Catemaco, Veracruz, Mexico
- Neoditylenchinae* subfam. n.
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Sychnotylenchidae Paramonov, 1967
key, type gen.: *Neoditylenchus* Meyl, 1960
- Neoditylenchus* Meyl, 1960 (type gen.)
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Sychnotylenchidae, *Neoditylenchinae* subfam. n.
key
- Neoditylenchus dalei* nov. sp., illus.
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Cerambyx cerdo acuminatus (under elytra)
- Neogoezia* Kreis, 1937
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
? as syn. of *Raphidascaris* Railliet & Henry, 1915
- Neogoezia* subgen.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
key
- Neoheligmone* Durette-Desset, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmoneidae, Nippostrongylineae
- Neohistiostrongylus* Barus et Rysavy, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylineae
- Neoleptus* gen. n.
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Physalopteridae, Physalopterinae
key, tod: *Neoleptus australis* (Johnston and Mawson, 1943) comb. nov.
- Neoleptus australis* (Johnston and Mawson, 1943) comb. nov. (tod)
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
[Syn.]: *Paraleptus australis* Johnston and Mawson, 1943
- Neomermis* sp.
Mondet, B.; Pendriez, B.; and Bernadou, J., 1976, Cahiers O.R.S.T.O.M., s. Entom. Med., v. 14 (2), 141-149
Simulium damnosum
S. unicornutum
all from Coted'Ivoire (riviere N'zi)
- Neomesomermis flumenalis*
Condon, W. J.; and Gordon, R., 1977, J. Invert. Path., v. 29 (1), 56-62
Neomesomermis flumenalis, effects on neuro-endocrine systems and storage of fat body glycogen in larval *Prosimulium mixtum fuscum* and *Simulium venustum*
- Neomesomermis flumenalis* (Welch)
Ebsary, B. A.; and Bennett, G. F., 1975, Canad. J. Zool., v. 53 (8), 1058-1062
Prosimulium fuscum/mixtum
Simulium venustum
Simulium latipes
Simulium tuberosum
all from insular Newfoundland
- Neomesomermis flumenalis*
Molloy, D.; and Jambback, H., 1977, Mosquito News, v. 37 (1), 104-108
Neomesomermis flumenalis, control of *Simulium venustum*, *S. vittatum*, cost too great for intermediate stage field trials
- Neopaucipectines*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Pterygodermatites*
key
- Neorailletnema praeputiale* Skrjabin, 1916
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Rana ridibunda
R. esculenta
all from Samara river valley, Ukrainian SSR
- Neospinitectus* Kalyankar, 1971
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"too poorly known to be classified at present"
- Neospirocerca* Johnson, 1968
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"unclassifiable but it may be a juvenile female of *Protospirura*"
- Neostongylus*
Nemeseri, L., 1976, Magy. Allat. Lapja, v. 98, v. 32 [i. e. 31] (7), 459-461
Dictyocaulus filaria, trichostrongylids, *Fasciola hepatica*, sheep, successful treatment with combination of diamphenetide and tetramisole; no effect on Protostrongylidae
- Neostongylus linearis*
Aller, B.; and Aller, J. M., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 337-346
nematodes of sheep lungs, lower incidence of fungi in parasitized lungs: Leon

- Neostromylus linearis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (lung): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Neostromylus linearis*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Neostromylus linearis*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Neostromylus linearis*
Ramisz, A.; Urban, E.; and Piechocki, B., 1975, Med. Wet., v. 31 (11), 677-679
Protostrongylidae of sheep, tetramisole effective against all species except *Muellerius capillaris*
- Neostromylus linearis* (Marotel, 1913) Gebaur, 1932, illus.
Rojo-Vazquez, F. A.; and Cordero del Campillo, M., [1975], Ann. Parasitol., v. 49 (6), 1974, 685-699
Neostromylus linearis, life cycle in mollusc intermediate hosts and sheep, morphology of larval stages
Ovis aries
Helicella variabilis
H. neglecta
H. apicina
Cepaea nemoralis
Helix aspersa
Zebrina sp.
(exper. in all)
- Neostromylus linearis*
Rojo Vazquez, J., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 51-101
Protostrongylinae, frequency in ovine lungs, primary nodes are infected by bacteria at a lower rate than non-parasitized lung areas: municipal slaughterhouse, Leon
- Neostromylus linearis* (Marotel, 1913) Gebauer, 1932, illus.
del Rosario Marcos Martinez, M., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 103-174
Neostromylus linearis, histological reactions in *Ceratomyxa* (*Xeromagna*) *cespitem ari-gonis* and *C. (C.) virgata* (exper.); review of literature on morpho-eco-biology and systematics of mollusks studied and method of infestation and culturing intermediate hosts
- Neostromylus linearis*, illus.
Sumnaliev, P., 1973, Izvest. Tsentral. Khel-mint. Lab., v. 16, 161-167
Neostromylus linearis, *Muellerius capillar-is*, *Cystocaulus ocreatus*, *Protostrongylus* sp., differential diagnosis of infective larvae based on biometric studies
Helicella obvia
Zebrina detrita
all from Bulgaria
- Neoxysomatium brevicaudatum*
Hristovski, N. D., 1973, Acta Parasitol. Iugo-slavica, v. 4 (2), 87-91
Rana temporaria
Bufo bufo
Testudo graeca
Angius fragilis
all from Macedonia, Yugoslavia
- Neoxysomatium brevicaudatum* (Zeder, 1800) Ballesteros Marquez, 1945
Hristovski, N. D., 1975, Acta Parasitol. Iugoslavica, v. 6 (1), 3-5
Rana graeca: Bitola district, Macedonia, Yugoslavia
- Neoxysomatium brevicaudatum* (Zeder, 1800)
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Nicollina Baylis*, 1930
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
synonymy
- Nicollina antechini* sp. n., illus.
Beveridge, I.; and Barker, I. K., 1975, J. Parasitol., v. 61 (3), 489-493
Nicollina antechini sp. n. in *Antechinus stuartii* (nasal cavity), pathology: Victoria, Australia (Powelltown; Mt. Sabine, Otway Ranges)
- Nicollina antechini* Beveridge & Barker
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (nasal cavity): Powelltown, Victoria
- Nilonema Khalil*, 1960, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key
- Nippostrongylinae* Durette-Desset, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae
includes: *Nippostrongylus* (type genus); *Boreostromylus*; *Hassalstrongylus*; *Heligmonina*; *Heligmonoides*; *Hypocristata*; *Mammanidula*; *Neoheligmonella*; *Odilia*; *Oriostromylus*; *Stilestrongylus*; *Tenorastrongylus*

- Nippostrongylus* Lane, 1923 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, *Nippostrongylinae*
synonymy
- Nippostrongylus* sp.
Durette-Desset, M. C.; and Lim, B. L., [1975],
Ann. Parasitol., v. 49 (6), 1974, 669-675
Tomys horsfieldi (intestin grele): Kuala
Langat Forest Reserve, Bukit Mandol Aborig-
ine village, Klang, Selangor, Malaisie
- Nippostrongylus brasiliensis*
Ambler, J.; and Orr, T. S. C., 1972, *Immuno-
chemistry*, v. 9 (3), 263-272
Nippostrongylus brasiliensis adults, extrac-
tion and characterization of allergenic com-
ponent which bears striking resemblance in
many general physicochemical properties to
purified allergens of clinical importance
which have been extracted from other animal
and plant sources
- Nippostrongylus brasiliensis*
Berenguer Puvia, F. J.; and Gallego Berenguer,
J., 1973, *Rev. Iber. Parasitol.*, v. 33 (1), 81-
106
Aspiculuris tetraptera, *Syphacia obvelata*,
Nippostrongylus brasiliensis, mice, piper-
azine and phenothiazine compared with 4 pheno-
thiazine-piperazine derivatives; piperazine:
good activity, phenothiazine: low activity
against *N. brasiliensis*, low toxicity of
both; derivatives: more toxic, no anthelmin-
tic activity
- Nippostrongylus brasiliensis*
Bloch, K. J.; Towle, C.; and Mills, J. A.,
1977, *Cellular Immunol.*, v. 28 (1), 181-189
Nippostrongylus brasiliensis, rats, mesen-
teric lymph node and spleen cells, stimula-
tion by worm metabolic antigen and by con A
of tritiated thymidine incorporation
- Nippostrongylus brasiliensis*
Bone, L. W.; et al., 1977, *Exper. Parasitol.*,
v. 42 (1), 82-86
Nippostrongylus brasiliensis, pheromone
production and response, effect of para-
site age, time of day periodicity not demon-
strated
- Nippostrongylus brasiliensis*
Bone, L. W.; and Shorey, H. H., 1977, *J. Para-
sitol.*, v. 63 (5), 845-848
Nippostrongylus brasiliensis, interactive
influences of male- and female-produced
pheromones on male attraction to female
- Nippostrongylus brasiliensis*
Bone, L. W.; and Shorey, H. H., 1977, *Science*
(4304), v. 197, 694-695
Nippostrongylus brasiliensis, males main-
tained in environment permeated with female
pheromone have greatly reduced subsequent
ability to orient to gradient of pheromone
emanating from living females for periods up
to 2 hours, possible use of this phenomenon
in control
- Nippostrongylus brasiliensis*
Bone, L. W.; Shorey, H. H.; and Gaston, L. K.,
1977, *J. Parasitol.*, v. 63 (2), 364-367
Nippostrongylus brasiliensis, mutual hetero-
sexual attraction in vitro, pheromonal dos-
age response, homosexual trials
- Nippostrongylus brasiliensis*, illus.
Bonner, T. P.; and Buratt, M., 1976,
Internat. J. Parasitol., v. 6 (4), 289-
294
Nippostrongylus brasiliensis, inhibition
of development and infectivity by actino-
mycin-D, supports hypothesis that infec-
tion is dependent upon new gene expression,
i.e., transcription
- Nippostrongylus brasiliensis*
Bonner, T. P.; Evans, K.; and Kline, L., 1976,
Internat. J. Parasitol., v. 6 (6), 473-477
Nippostrongylus brasiliensis, role of gene
expression in regulating cuticle formation
during second molt, results strongly suggest
that messenger RNA specific for molting was
synthesized at 90 hours
- Nippostrongylus braziliensis*
Carr, J. B.; Durham, H. G.; and Hass, D. K.,
1977, *J. Med. Chem.*, v. 20 (7), 934-939
activity of isoxazole anthelmintics against
Nippostrongylus braziliensis in rats, labora-
tory screening of selected compounds
- Nippostrongylus braziliensis*
Carson, D.; Metzger, H.; and Bazin, H., 1975,
J. Immunol., v. 115 (2), 561-563
Nippostrongylus braziliensis-infected rats,
simple radioimmunoassay for measurement of
IgE levels by ammonium sulfate precipitation
- Nippostrongylus brasiliensis*
Carson, D.; Metzger, H.; and Bloch, K. J.,
1975, *J. Immunol.*, v. 114 (1, Pt.2), 521-523
Nippostrongylus brasiliensis-infected rats,
serum IgE levels during potentiated reagin
response to egg albumin, results suggest
N. brasiliensis functions as an IgE-specific
adjuvant in the rat
- Nippostrongylus brasiliensis*
Church, M. K., 1975, *Immunology*, v. 29 (3),
527-534
Nippostrongylus brasiliensis-sensitized, re-
sensitized, or passively sensitized rats,
interrelationships of anaphylactic broncho-
constriction, active cutaneous anaphylaxis,
and circulating reagenic antibody level
- Nippostrongylus brasiliensis*
Coles, G. C.; East, J. M.; and Jenkins, S. N.,
1975, *Gen. Pharmacol.*, v. 6 (4), 309-313
Nippostrongylus brasiliensis, *Ascaris lum-
bricoides* var. *levisole*, mechanism
of action, paralysis and recovery of motili-
ty, resistance of recovered worms to paraly-
sis by methyridine, pyrantel, or bephenium,
results suggest levamisole acts as ganglion
stimulant and that inhibition of fumarate
reductase is not relevant to in vivo mode of
action

- Nippostrongylus brasiliensis*
Coles, G. C.; and McNeillie, R. M., 1977, *J. Helminth.*, v. 51 (4), 323-326
dietary feeding of drugs for 5 days to mice infected with *Nematospiroides dubius* and *Hymenolepis nana*, detected all modern anthelmintics examined except stilbazium; simple test using *Nematodirus spathiger* eggs and *Nippostrongylus brasiliensis* adults to detect anthelmintics in vitro
- Nippostrongylus brasiliensis*
Croll, N. A., 1976, *Internat. J. Parasitol.*, v. 6 (5), 441-448
Nippostrongylus brasiliensis, dispersion in rat intestine related to host feeding and diet
- Nippostrongylus brasiliensis*
Croll, N. A., 1977, *Internat. J. Parasitol.*, v. 7 (3), 201-204
Nippostrongylus brasiliensis, rats, relative importance of active behavioral factors and passive carriage in larval migration between host's tissues and organs
- Nippostrongylus brasiliensis*
Croll, N. A., 1977, *Parasitology*, v. 75 (2), iii [Abstract]
Nippostrongylus brasiliensis, dispersion in rat intestine, effect of several manipulations (diet and surgical)
- Nippostrongylus brasiliensis*
Croll, N. A.; and Ma, K., 1977, *Internat. J. Parasitol.*, v. 7 (1), 21-26
Nippostrongylus brasiliensis, localization in rats, influence of surgical manipulation of intestine and mesenteric blood supply on dispersion
- Nippostrongylus brasiliensis*, illus.
Croll, N. A.; and Smith, J. M., 1977, *Internat. J. Parasitol.*, v. 7 (3), 195-200
Nippostrongylus brasiliensis in fed vs. fasted rats, 10 vs. 15 day old infections, activity rates, postures, and temporal movement patterns observed in vivo, differences correlated with feeding regime of host and could thus lead to habitat selection
- Nippostrongylus brasiliensis*
Della Bruna, C.; and Xenia, B., 1976, *J. Parasitol.*, v. 62 (3), 490-491
Nippostrongylus brasiliensis: reduced worm burden and prolonged infection in mice harboring *Nematospiroides dubius*
- Nippostrongylus brasiliensis*
Dineen, J. K.; et al., 1974, *Prostaglandins*, v. 5 (2), 209-220
Nippostrongylus brasiliensis-infected rats (exper.), inhibitor of prostaglandin action detected in residue of ram and human semen after extraction of prostaglandins; inhibitor prevented expulsion of adult worms from rat small intestine, possibly plays important role in reproductive physiology
- Nippostrongylus brasiliensis*
Dineen, J. K.; and Kelly, J. D., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 51 (4), 429-440
Nippostrongylus brasiliensis, primary and secondary infections in rats, intestinal prostaglandin levels in relation to worm expulsion
- Nippostrongylus brasiliensis*
Dineen, J. K.; Kelly, J. D.; and Love, R. J., 1973, *Internat. Arch. Allergy and Applied Immunol.*, v. 45 (4), 504-512
Nippostrongylus brasiliensis, mesenteric lymph node cells from immunized donors caused expulsion of transplanted damaged worms or adult worms developed from larval infection in both irradiated or non-irradiated syngeneic recipients but cells from unimmunized donors failed to affect the parasite
- Nippostrongylus brasiliensis*
Dineen, J. K.; Ogilvie, B. M.; and Kelly, J. D., 1973, *Immunology*, v. 24 (3), 467-475
Nippostrongylus brasiliensis, expulsion from intestine of rats, collaboration between humoral and cellular components of immune response
- Nippostrongylus brasiliensis*
Gavaghan, A. D.; and Nunn, A. J., 1972, *Pharm. Acta Helvetiae*, v. 47 (11-12), 719-726
Nippostrongylus brasiliensis, *Strongyloides ratti*, *Nematospiroides dubius*, synthesis of thiocyanine and hemithiocyanine derivatives of benzothiazole, significant anthelmintic activity in laboratory trials with mice
- Nippostrongylus brasiliensis*
Gavaghan, A. D.; and Nunn, A. J., 1974, *Pharm. Acta Helvetiae*, v. 49 (7-8), 250-258
bis-(6-indazolyloxy) alkanes, no schistosomicidal or other significant anthelmintic properties in laboratory trials with mice
- Nippostrongylus brasiliensis*
Gerber, H. C.; Hoerchner, F.; and Zander, B., 1976, *Ztschr. Parasitenk.*, v. 50 (1), 57-66
Nippostrongylus brasiliensis, studies in vitro with serum or secretory antibodies from rats, no effect with antibodies from either infected or uninfected rats; degranulation of mast cells from infected rats but no effect on *N. brasiliensis*
- Nippostrongylus brasiliensis*
Golenser, J.; Spira, D. T.; and Shmuel, Z., 1976, *Parasitology*, v. 73 (2), xiii [Abstract]
rats, mutual influence of infection with *Plasmodium berghei* and *Nippostrongylus brasiliensis*
- Nippostrongylus brasiliensis*
Goose, J., 1977, *Parasitology*, v. 75 (2), xxxv [Abstract]
Fasciola hepatica-infected rats, depressed immune response to sheep red blood cells, greater numbers of peritoneal macrophages and increased phagocytosis, increased resistance to infection with *Trypanosoma congolense* and *Nippostrongylus brasiliensis* not serum-transferable, concluded that *Fasciola hepatica* stimulates certain T-cell populations as well as reticulo-endothelial system

- Nippostrongylus brasiliensis*, *illus.*
Greenberg, Z.; and Wertheim, G., 1973, *Immunology*, v. 24 (3), 531-543
Nippostrongylus brasiliensis, rats, cellular responses to intraperitoneal inoculation of larvae, sequence of cell-types adhering to larvae and subsequent formation of granulomas around cell-coated larvae, initial neutrophilia in rats infected with L2 or L3 larvae, pronounced eosinophilia seen only in rats inoculated with L3 larvae
- Nippostrongylus brasiliensis*
Henney, R. W.; MacLean, J. M.; and Mulligan, W., 1973, *Isotopes and Radiation Parasitol.* III, 3-10
Nippostrongylus brasiliensis, decline in uptake of ^{32}P -phosphate and ^{75}Se -selenomethionine from host's tissue fluids from day 7 of infection onward, recovery upon transfer to fresh rat, results imply significant interference with parasites' metabolism several days before expulsion occurs
- Nippostrongylus brasiliensis*
Hindsbo, O., 1973, *Norwegian J. Zool.*, v. 21 (4), 328 [Abstract]
anterior migration of *Nippostrongylus brasiliensis* in small intestine of rats during primary and secondary infections correlated to self-cure; experiments with prednisolone-treated rats showed that anterior migration is independent of worm expulsion
- Nippostrongylus brasiliensis*
Hogarth-Scott, R. S., 1973, *Immunology*, v. 24 (3), 503-509
Nippostrongylus brasiliensis, rats, peripheral circulating allergens as cause of loss of homologous passive cutaneous anaphylaxis reactivity, reaction appeared to be immunologically specific
- Nippostrongylus brasiliensis*
Ishizaka, T.; et al., 1975, *J. Immunol.*, v. 115 (4), 1078-1083
Nippostrongylus brasiliensis-infected rats, increase in serum IgE, most receptors for IgE on mast cells of infected animals are occupied by their own IgE, no differences in mast cells of normal vs. infected rats with respect to histamine content or intracellular levels of cyclic nucleotides
- Nippostrongylus brasiliensis*
Ishizaka, T.; et al., 1976, *J. Allergy and Clin. Immunol.*, v. 58 (4), 523-538
Nippostrongylus brasiliensis-infected rats, nature of IgE synthesis, role of T and B cells, characteristic properties of T cells primed by infection
- Nippostrongylus brasiliensis*
Ishizaka, T.; Urban, J. F., jr.; and Ishizaka, K., 1976, *Cellular Immunol.*, v. 22 (2), 248-261
Nippostrongylus brasiliensis, rats, kinetics of IgE synthesis, selective proliferation of IgE-bearing lymphocytes and differentiation of these to IgE-forming cells
- Nippostrongylus brasiliensis*, *illus.*
Jacobson, R. H.; and Reed, N. D., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 52 (1-4), 160-168
Nippostrongylus brasiliensis, mice, thymus-dependent lymphocytes are required to elicit both humoral and cell-mediated steps of worm expulsion
- Nippostrongylus brasiliensis*
Jacobson, R. H.; Reed, N. D.; and Manning, D. D., *Immunology*, v. 32 (6), 867-874
Nippostrongylus brasiliensis, mice experiencing immunosuppressive effects of anti- μ antibodies, worm expulsion unaffected even though antibody production potential had been eliminated, suggests that anti-worm antibodies may not be requisite in mechanism of expulsion
- Nippostrongylus brasiliensis*
Jarrett, E. E. E., 1972, *Immunology*, v. 22 (6), 1099-1101
potentiation of reaginic (IgE) antibody to ovalbumin by infection of rats with *Fasciola hepatica* and re potentiation by subsequent infection with *Nippostrongylus brasiliensis*
- Nippostrongylus brasiliensis*
Jarrett, E. E. E.; Haig, D. M.; and Bazin, H., 1976, *Clin. and Exper. Immunol.*, v. 24 (2), 346-351
Nippostrongylus brasiliensis, rats, relationship in time between elevation of total serum IgE, the parasite-specific IgE response, and the potentiated IgE response to unrelated antigen
- Nippostrongylus brasiliensis*
Jarrett, E. E. E.; Orr, T. S. C.; and Riley, P., 1971, *Clin. and Exper. Immunol.*, v. 9 (5), 585-594
inhibition of allergic reactions due to competition for mast cell sensitization sites by two reagins (egg albumin reagins and *Nippostrongylus brasiliensis* reagins), practical importance
- Nippostrongylus brasiliensis*
Jarrett, E. E. E.; and Stewart, D. C., 1973, *Immunology*, v. 24 (1), 37-45
Nippostrongylus brasiliensis, rats, no direct quantitative relationship between size of skin test reactions and level of specific circulating reaginic antibody
- Nippostrongylus brasiliensis*
Jones, V. E.; and Ogilvie, B. M., 1967, *Internat. Arch. Allergy and Applied Immunol.*, v. 31 (5), 490-504
Nippostrongylus brasiliensis, attempts to transfer protective immunity and reagins passively to young rats in 3 different ways: by maternal milk, by feeding antiserum, and by antiserum given parenterally
- Nippostrongylus brasiliensis* (Travassos, 1914)
Katiyar, J. C.; and Sen, A. B., 1969, *Indian J. Helminth.*, v. 21 (1), 81-85
Nippostrongylus brasiliensis, susceptibility of 3 strains of white rats to infection

- Nippostrongylus brasiliensis*
Kazacos, K. R., 1975, *Vet. Parasitol.*, v. 1 (2), 165-174
Trichinella spiralis-immunized rats, increased resistance to *Nippostrongylus brasiliensis*, heterologous and homologous tests of immune precipitate formation on infective larvae
- Nippostrongylus brasiliensis*, illus.
Kazacos, K. R.; and Thorson, R. E., 1975, *J. Parasitol.*, v. 61 (3), 525-529
rats, immunization with *Nippostrongylus brasiliensis* or *Strongyloides ratti* protected against homologous and heterologous challenge; precipitates formed on infective larvae incubated in vitro in homologous or heterologous immune globulins
- Nippostrongylus brasiliensis*
Keller, R., 1973, *Clin. and Exper. Immunol.*, v. 13 (1), 139-147
incubation of sensitized cells from *Nippostrongylus brasiliensis*-infected rats with concanavalin A in vitro induces release of histamine, release mechanism is similar to that elicited by specific worm allergen, suggests that release mechanism is triggered by interaction with cell-bound IgE-type antibody
- Nippostrongylus brasiliensis*
Keller, R.; and Beeger, I., 1971, *Internat. Arch. Allergy and Applied Immunol.*, v. 41 (2-3), 278-285
Nippostrongylus brasiliensis-infected rats, increase in sensitivity to histamine but hyperglycemic response to histamine and epinephrine is unchanged, blockade of adrenergic function is not important feature in increase in histamine lethality of nematode-infected rat
- Nippostrongylus brasiliensis*
Keller, R.; Ogilvie, B. M.; and Simpson, E., 1971, *Lancet*, London (7701), v. 1, 678-680
Nippostrongylus brasiliensis, rats, mice, effect of infection on tumor growth, results raise possibility that helminths and tumors may share antigens and suggest that helminth infections have non-specific effect on tumor growth
- Nippostrongylus brasiliensis*
Kelly, J. D.; et al., 1974, *Internat. Arch. Allergy and Applied Immunol.*, v. 47 (3), 458-465
Nippostrongylus brasiliensis, rats, comparative capacity of synthetic prostaglandins and pharmacologically active amines to initiate worm expulsion
- Nippostrongylus brasiliensis*
Kelly, J. D.; and Dineen, J. K., 1972, *Immunology*, v. 22 (2), 199-210
Nippostrongylus brasiliensis, rats, successful adoptive immunization with mesenteric lymph node cells from immune donors
- Nippostrongylus brasiliensis*
Kelly, J. D.; and Dineen, J. K., 1973, *Immunology*, v. 24 (3), 551-558
Nippostrongylus brasiliensis, suppression of rejection in castrated male rats treated with ovine prolactin, main effect of prolactin is directed towards the immunologically specific lymphoid phase of the worm rejection mechanism
- Nippostrongylus brasiliensis*
Kelly, J. D.; and Dineen, J. K., 1976, *Austral. Vet. J.*, v. 52 (9), 391-397
gastrointestinal helminths, role of prostaglandins in worm expulsion, review
- Nippostrongylus brasiliensis*
Kershaw, W. E.; et al., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (1), 31-32 [Demonstration]
Litomosoides carinii in vitamin A deficient cotton rats and in vitamin E deficient cotton and white rats, antigen extracts from adults and microfilariae; *Nippostrongylus brasiliensis*, mast cell population in lungs of infected rats
- Nippostrongylus brasiliensis*
Kocan, A. A., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 237-241
influence of *Nippostrongylus brasiliensis* on establishment of *Angiostrongylus cantonensis*, role of various organs, laboratory rat
- Nippostrongylus brasiliensis*
Kojima, S.; and Ovary, Z., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 50 (1), 81-86
rats immunized with dinitrophenylated *Nippostrongylus brasiliensis* protein 2 weeks after initial infection, induction of carrier-specific enhancing effect on primary and secondary antihapten IgE antibody response
- Nippostrongylus brasiliensis*
Larsh, J. E., jr.; and Weatherly, N. F., 1975, *Advances Parasitol.*, v. 13, 183-222
principles of delayed (cellular) hypersensitivity, cell-mediated immunity against parasitic worms, extensive review
- Nippostrongylus brasiliensis*
Love, R. J.; and Ogilvie, B. M., 1977, *Exper. Parasitol.*, v. 41 (1), 124-132
Nippostrongylus brasiliensis, *Trichinella spiralis*, rats, increased localization of thoracic duct lymphoblasts in small intestine, not correlated with presence of antigen but related to factors associated with inflammation, no increase in blast localization in lactating vs. nonlactating rats but increased localization in lactating mammary gland
- Nippostrongylus brasiliensis*
Love, R. J.; Ogilvie, B. M.; and McLaren, D. J., 1976, *Immunology*, v. 30 (1), 7-15
Trichinella spiralis, duration of infections in rats (young, adult, and lactating), rapidity of expulsion from previously infected rats, transfer of immunity with antiserum and lymph node cells, ultrastructural signs of antibody damage to worms, results suggest that mechanism of immune expulsion requires both antibody and cells, comparison with *Nippostrongylus brasiliensis*

- Nippostrongylus brasiliensis*
Mayrhofer, G.; Bazin, H.; and Gowans, J. L., 1976, *European J. Immunol.*, v. 6 (8), 537-545
- Nippostrongylus brasiliensis*, immunized rats, nature of cells binding anti-IgE, IgE synthesis in regional lymph nodes and concentration in mucosal mast cells, results do not support suggestion that IgE is secretory immunoglobulin with physiology analogous to that of IgA
- Nippostrongylus brasiliensis*
Meacock, S. C. R.; and Marsden, C. H., 1976, *Immunology*, v. 30 (4), 491-496
- Nippostrongylus brasiliensis*, rats, potentiating effects of infection on IgE antibody production after two injections of egg albumin, IgG_a antibody not potentiated
- Nippostrongylus brasiliensis*
Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
- Rattus norvegicus* (intestin grele): Tunisia
- Nippostrongylus brasiliensis*
Mitchell, G. F.; et al., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 52 (1-4), 95-104
- Nippostrongylus brasiliensis* infections in hypothyroid nu/nu mice
- Nippostrongylus brasiliensis*
Mitchell, G. F.; and Lewers, H. M., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 52 (1-4), 235-240
- Ascaris suum*, *Nippostrongylus brasiliensis*, mice, inhibition of an anti-DNP antibody response with DNP-Ficoll containing phosphorylcholine, results suggest that parasites may 'utilize' molecules such as phosphorylcholine to induce state of selective tolerance to parasite antigens as a mechanism for facilitating survival
- Nippostrongylus brasiliensis*
Mohan, J.; and Pujari, H. K., 1975, *Indian J. Chem.*, v. 13 (5), 528-529
- synthesis of 3-aryl-5,6-dimethylimidazo[2,1-b]thiazoles, no activity in anthelmintic tests with *Hymenolepis nana* and *Nippostrongylus brasiliensis*
- Nippostrongylus brasiliensis*
Munn, E. A., 1977, *Tissue and Cell*, v. 9 (1), 23-34
- Haemonchus contortus*, structure of intestinal cells, helical polymeric extracellular protein associated with luminal surface for which name contortin is proposed, *Ostertagia circumcincta* also contained contortin-like material but *Nippostrongylus brasiliensis* and *Syphacia obvelata* contained material associated with outer surface of microvilli which was quite distinct from contortin
- Nippostrongylus brasiliensis*
Murray, P. K.; et al., 1974, *Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol.* (Vienna, Austria, Sept. 18-20, 1973), 133-150
- Trypanosoma brucei* in rodents, pathology of immune system, role of mononuclear phagocytic system in immunosuppression, humoral and T cell responsiveness, effect of infection on *Nippostrongylus brasiliensis* infection, recovery of immune response after trypanocidal therapy, review
- Nippostrongylus brasiliensis*
Ogilvie, B. M.; et al., 1977, *Immunology*, v. 32 (4), 521-528
- Nippostrongylus brasiliensis*, rats, cellular requirement for worm expulsion, results suggest that following antibody damage this nematode is expelled by nonimmunoglobulin-bearing lymphocytes which are effective in the absence of newly formed cells derived from the cell recipients
- Nippostrongylus brasiliensis*
Ogilvie, B. M.; and Love, R. J., 1974, *Transplant. Rev.*, v. 19, 147-169
- immune mechanisms in *Nippostrongylus brasiliensis*-rat model, co-operation between antibodies and cells in immune expulsion, review
- Nippostrongylus brasiliensis*
Ogilvie, B. M.; Mackenzie, C. D.; and Love, R. J., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (6, Pt. 2), 61-67
- Nippostrongylus brasiliensis*, rats, lymphocytes and eosinophils in immune response to initial and subsequent infections, workshop report
- Nippostrongylus brasiliensis*
Ohman, J. L., jr.; and Bloch, K. J., 1972, *J. Immunol.*, v. 108 (6), 1637-1646
- Nippostrongylus brasiliensis*, rats, effect of IgE on passive cutaneous anaphylactic reaction mediated by IgG_a antibodies
- Nippostrongylus brasiliensis*
Orr, T. S. C.; Riley, P. A.; and Doe, J. E., 1972, *Immunology*, v. 22 (2), 211-217
- Nippostrongylus brasiliensis*-infected rats, time course of potentiated reagin response to egg albumin
- Nippostrongylus brasiliensis*
Parrott, D. M. V.; and Ferguson, A., 1974, *Immunology*, v. 26 (3), 571-588
- mesenteric lymph node cells from donors infected with *Nippostrongylus brasiliensis* used in investigating selective migration of lymphocytes within mouse small intestine
- Nippostrongylus brasiliensis*
Petit, A.; et al., 1977, *Ann. Immunol.*, v. 128C (4-5), 937 [Abstract]
- Nippostrongylus brasiliensis*, rats, circulating immune complexes in serum

- Nippostrongylus brasiliensis*
Poulain, J.; et al., 1977, *Ann. Immunol.*, v. 128C (1-2), 245-247
Nippostrongylus brasiliensis, rats, synthesis of hemagglutinating antibodies in intestinal secretions, detected before serum antibodies, possible role of local antibodies in mechanism of worm expulsion
- Nippostrongylus brasiliensis*
Poulain, J.; Luffau, G.; and Pery, P., 1975, *Compt. Rend. Acad. Sc.*, Paris, v. 281, s. D, Sc. Nat. (23), 1933-1934
Nippostrongylus brasiliensis infected rats, formation of hemagglutinating antibodies in serum and intestinal secretions; local antibodies, new immune mechanism
- Nippostrongylus brasiliensis*
Poulain, J.; Luffau, G.; and Pery, P., 1976, *Ann. Immunol.*, v. 127C (2), 215-224
Nippostrongylus brasiliensis, rats, kinetics of haemagglutinin production in serum vs. intestinal secretions, antibody response to low doses of infecting larvae, nature of immunoglobulin classes involved
- Nippostrongylus brasiliensis*
Poulain, J.; Pery, P.; and Luffau, G., 1976, *Ann. Immunol.*, v. 127C (2), 209-213
Nippostrongylus brasiliensis, rats, vaccination with worm homogenates, whole fixed worms, or metabolic products administered orally, metabolic products induced highest level of protection
- Nippostrongylus brasiliensis*
Premvati, G.; and Chopra, A. K., 1977, *Indian J. Zoot.*, v. 16 (1), 1975, 1-5
Nippostrongylus brasiliensis, infective larvae exposed to various doses of ultraviolet rays, development, sex ratio of adult worms, no resistance in rats challenged with lethal dose of untreated larvae subsequent to lethal dose of irradiated larvae
- Nippostrongylus brasiliensis*
Preston, P. J.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 17 [Demonstration]
Nippostrongylus brasiliensis, in vitro and in vivo, isotope scanning, potential for study of parasite migration and behavior in living host
- Nippostrongylus brasiliensis*
Preston, P. J.; et al., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (2), 289-290 [Letter]
Nippostrongylus brasiliensis, isotopic scanning technique whereby life cycle and migration of nematode parasites can be studied within living host
- Nippostrongylus brasiliensis*
Przyjalkowski, Z.; and Wescott, R. B., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 265-269
Nippostrongylus brasiliensis, conventional and germfree mice, some with monocultures of three species of bacteria, differential leucocyte counts, packed cell volume; higher worm burdens in male mice and in conventional mice, little differences in blood values
- Nippostrongylus brasiliensis*, illus.
Richards, A. J.; et al., 1977, *Internat. J. Parasitol.*, v. 7 (2), 153-158
Nippostrongylus brasiliensis, in vitro incubation with prostaglandin E₁, effect on glycolysis, on worm morphology, and on survival in vivo, results support view that prostaglandins play vital role in mechanism of worm expulsion
- Nippostrongylus brasiliensis*
Roberts, T. M.; and Thorson, R. E., 1977, *J. Parasitol.*, v. 63 (2), 357-363
Nippostrongylus brasiliensis, chemical attraction between adults both within host intestine and in vitro with particular regard to effect of host immunity on parasite behavior
- Nippostrongylus brasiliensis*
Roberts, T. M.; and Thorson, R. E., 1977, *J. Parasitol.*, v. 63 (4), 764-766
Nippostrongylus brasiliensis, pairing between adults and other species of nematodes in vitro, comparison of intra and interspecific response patterns, results suggest that pairing is not related entirely to sexual attraction
- Nippostrongylus brasiliensis*
Roberts, T. M.; and Thorson, R. E., 1977, *J. Parasitol.*, v. 63 (5), 849-853
Nippostrongylus brasiliensis, characterization of pheromone which attracts females (lipid fraction of excretory and secretory products of both males and females), that from immune-damaged females was not attractive
- Nippostrongylus brasiliensis*
Sanderson, B. E.; Jenkins, D. C.; and Phillipson, R. F., 1976, *Internat. J. Parasitol.*, v. 6 (2), 99-102
Nippostrongylus brasiliensis, relation between changes in host immunity and worm acetylcholinesterase levels, results indicate that immunity-associated cholinesterase increase is to some extent reversible
- Nippostrongylus brasiliensis*
Scofield, A. M., 1977, *Internat. J. Parasitol.*, v. 7 (2), 159-165
Nippostrongylus brasiliensis-infected rats, intestinal absorption of hexoses, possible relation to immune reaction
- Nippostrongylus brasiliensis*, illus.
Seese, F. M.; Wescott, R. B.; and Gorham, J. R., 1976, *Exper. Parasitol.*, v. 39 (2), 214-221
Nippostrongylus brasiliensis, mice, indirect fluorescent antibody studies on localization of antigenic sites within worm and on classes of antibody involved in humoral response
- Nippostrongylus brasiliensis*, illus.
Seese, F. M.; Wescott, R. B.; and Gorham, J. R., 1977, *J. Parasitol.*, v. 63 (6), 1135-1137
Nippostrongylus brasiliensis, scanning electron microscopy, suitable method of specimen preparation, visualization of surface topography in situ

- Nippostrongylus brasiliensis*
Simpkin, K. G.; and Coles, G. C., 1976, Parasitology, v. 73 (2), iv [Abstract]
Nematospiroides dubius, *Nippostrongylus brasiliensis*, *Nematodirus spathiger*, modes of action of thiabendazole and mebendazole apparently different from those reported on other species
- Nippostrongylus braziliensis* (Travassos, 1914)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus canus
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
all from Malaysia
- Nippostrongylus brasiliense*
Singhal, K. C., 1976, Indian J. Exper. Biol., v. 14 (3), 345-347
berberine hydrochloride, in vivo activity against *Syphacia obvelata*, *Nippostrongylus brasiliense*, and *Hymenolepis nana*, mice; elimination of *S. obvelata* only, drug considered equipotent to piperazine citrate
- Nippostrongylus brasiliensis*
Sinski, E.; and Holmes, P. H., 1977, Exper. Parasitol., v. 43 (2), 382-389
Nippostrongylus brasiliensis, rats, systemic and local antibody response, specific anti-parasite binding capacity of IgA and IgG in intestinal mucosa and serum measured by radioimmunoassay, results compared with total levels of IgA and IgG and hemagglutinating and precipitating antibody titers, indirect evidence that local IgA may be part of effector mechanism
- Nippostrongylus brasiliensis*
Smith, S. R.; et al., 1973, Internat. Arch. Allergy and Applied Immunol., v. 44 (3), 382-395
potentiation of IgG and IgE reaginic responses to alum-precipitated dinitrophenylated bovine gamma-globulin with *Nippostrongylus brasiliensis*, rats
- Nippostrongylus brasiliensis*
Smith, S. R.; Petillo, J.; and Hwang, A., 1972, Internat. Arch. Allergy and Applied Immunol., v. 43 (1), 145-157
high titers of reagins against egg albumin obtained in rats by initial immunization with egg albumin and *Bordetella pertussis* and subsequent infection with *N. brasiliensis*, IgE type antibodies against *N. brasiliensis* are also produced in these same animals but not until titers of IgE against egg albumin have decreased
- Nippostrongylus brasiliensis*
Stafford, E. E.; and Jachowski, L. A., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 184-185
Nippostrongylus brasiliensis, method of quantitative intestinal worm recovery, rapid and accurate
- Nippostrongylus brasiliensis*, illus.
Symons, L. E. A., 1976, Internat. J. Parasitol., v. 6 (2), 107-111
Nippostrongylus brasiliensis-infected rat jejunum, scanning electron microscopy
- Nippostrongylus brasiliensis*
Takats, C., 1972, Parasitol. Hungar., v. 5, 203-216
human helminths, 171 compounds tested on laboratory animals for possible anthelmintic action, techniques of screening procedures described
- Nippostrongylus brasiliensis*, illus.
Taniguchi, M.; et al., 1977, Bull. Coll. Agric. and Vet. Med., Nihon Univ. (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo
- Nippostrongylus brasiliensis* (Travassos, 1914)
Travassos y Darriba, 1929
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 39-42
Rattus norvegicus (intestino delgado): Chile
- Nippostrongylus brasiliensis*
Urban, J. F., jr.; Ishizaka, T.; and Ishizaka, K., 1977, J. Immunol., v. 118 (6), 1982-1986
Nippostrongylus brasiliensis, rats, normal and neonatally thymectomized, proliferation of IgE-bearing cells
- Nippostrongylus brasiliensis*
Urban, J. F., jr.; Ishizaka, T.; and Ishizaka, K., 1977, J. Immunol., v. 119 (2), 583-590
Nippostrongylus brasiliensis, rats, soluble factor for generation of IgE-bearing lymphocytes
- Nippostrongylus brasiliensis*, illus.
Urquhart, G. M.; et al., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (4), 528-535
Trypanosoma brucei-infected rats, failure of humoral and immediate-type responses to superimposed *Nippostrongylus brasiliensis* infection; *T. brucei*-infected mice, cell-mediated immunity as measured by oxazolone sensitization still occurred to significant extent although less than in uninfected mice
- Nippostrongylus brasiliensis*
Watanabe, N.; et al., 1977, J. Immunol., v. 118 (2), 485-488
Nippostrongylus brasiliensis, suppression of IgE antibody production in SJL mice, expression of Ly-1 antigen on helper and non-specific suppressor T cells
- Nippostrongylus brasiliensis*
Watanabe, N.; Kojima, S.; and Ovary, Z., 1977, J. Immunol., v. 118 (1), 251-255
Nippostrongylus brasiliensis, mice, tolerizing effect of DNP-Ficoll on IgE antibody production

- Nippostrongylus brasiliensis*
Wells, P. D., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 15 [Demonstration]
Nippostrongylus brasiliensis in rats (exper.), mast cells in lungs after super-challenge infection, no evidence of sudden expulsion of larvae from lungs or adult worms from intestine •
- Nippostrongylus brasiliensis*
Wells, P. D., 1977, Exper. Parasitol., v. 43 (2), 326-335
Nippostrongylus brasiliensis, repeatedly inoculated rats, lung mast cell populations and larval and adult worm populations
- Nippostrongylus brasiliensis* (Travassos, 1914)
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Rattus norvegicus
R. rattus
all from Israel
- Nippostrongylus brasiliensis*
Whur, P.; and White R. G., 1970, Internat. Arch. Allergy and Applied Immunol., v. 38 (2), 185-195
fluorescence microscopy of small intestines of commercially slaughtered sheep and of *Nippostrongylus brasiliensis*-infected rats, concluded that globule leucocytes do not contain immunoglobulin
- Nippostrongylus brasiliensis*
Wilson, P. A. G., 1976, Ztschr. Parasitenk., v. 49 (3), 243-252
Haemonchus contortus, *Nippostrongylus brasiliensis*, infective larvae, carbohydrate content and ageing process contrasted; carbohydrate level variation in *H. contortus* due to capacity to synthesize glycogen during ageing, low level in *N. brasiliensis* remains constant
- Nippostrongylus brasiliensis*
Wilson, P. A. G.; Gentle, M.; and Scott, D. S., 1976, Parasitology, v. 72 (3), 355-360
rats, milk-borne infection, *Strongyloides ratti* (real and important vehicle for infection) vs. *Nippostrongylus brasiliensis* (possible but quantitatively insignificant)
- Nippostrongylus brasiliensis*
Wilson, P. A. G.; Gentle, M.; and Scott, D. S., 1976, Parasitology, v. 73 (3), 399-406
Strongyloides ratti, dynamic determinants of route of larval migration in lactating rats; *Nippostrongylus brasiliensis*, *Strongyloides ratti*, control of exper. error in quantitative studies of milk transmission of skin-penetrating roundworms
- Nippostrongylus brasiliensis* (Travassos, 1914)
Travassos and Darriba 1929, illus.
Wright, K. A., 1975, Canad. J. Zool., v. 53 (8), 1131-1146
Nippostrongylus brasiliensis, cephalic sense organs, form and function
- Nippostrongylus brasiliensis*, illus.
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Nippostrongylus brasiliensis*
Yeates, R. A.; and Ogilvie, B. M., 1976, J. Immunol. Methods, v. 12 (1-2), 57-65
Nippostrongylus brasiliensis, *Necator americanus*, measurement of antibodies to an unpurified enzyme (acetylcholinesterase) using an active-site directed radiolabel, cross-reactions between the two species
- Nippostrongylus brasiliensis*
Zander, B.; and Hoerchner, F., 1976, Pathophysiol. Parasit. Infect., 241-246
Nippostrongylus brasiliensis in vitro, effect of rat macrophages and lymphocytes vs. effects of serum, results suggest that cell dependent reactions rather than humoral components are involved in immune response
- Nippostrongylus muris*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Nippostrongylus muris*
Striebel, H. P., 1976, Experientia, v. 32 (4), 457-458
4-isothiocyanato-4'-nitrodiphenylamine, an anthelmintic with an unusual spectrum of activity against intestinal nematodes, filariae and schistosomes
- Nippostrongylus witenbergi* Greenberg, 1972
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Nesokia indica: Israel
- Nochitia Travassos et Vogelsang*, 1929
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Nochitia nochti*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Notopteroides Chakravarty & Majumdar*, 1962
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Pseudopropleptus Khera*, 1953
- Nycteridostrongylus*, Baylis, 1930
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Nycteridostrongylus (Petiellus) petersi* n. sub. gen., n. sp. (tod of subgen.)
Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (2), 173-185
Tupaia glis (intestin, poumons)
T. tana (intestin)
all from Sepilok, 15 miles de Sandakan, Borneo

- Obeliscoides* Graybill, 1924
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, Libyostrongylinae
synonymy
- Obeliscoides cuniculi*
Bemis, D. A.; and Georgi, J. R., 1972, *Cornell Vet.*, v. 62 (4), 680-685
Obeliscoides cuniculi, rabbits (exper.), estimated erythrocyte loss, concluded that *O. cuniculi* is not a significant bloodsucker
- Obeliscoides cuniculi*
Fox, J. C., 1976, *Vet. Parasitol.*, v. 1 (3), 209-220
Obeliscoides cuniculi, inhibited development in rabbits: effects of active and passive immunization and resumption of larval development (fewer males developed than females), data indicate that host immune responses contribute to inhibition and that worm egg production is also responsive to immunologic control
- Obeliscoides cuniculi*
Hutchinson, G. W.; and Fernando, M. A., 1976, *J. Parasitol.*, v. 62 (5), 841-843
Obeliscoides cuniculi, various life cycle stages, activity of enzymes of pentose phosphate pathway and glycolysis
- Obeliscoides cuniculi*
Jacobson, H. A.; and Kirkpatrick, R. L., 1974, *J. Wildlife Dis.*, v. 10 (4), 384-391
comparison of selected physiological measurements in untreated parasitized cottontail rabbits and those treated with 1-tetramisole hydrochloride and 2,2-dichlorovinyl, dimethyl phosphate: Montgomery County, Virginia
- Obeliscoides cuniculi*
Yonders, P. C.; and Dixon, C. F., 1976, *J. Alabama Acad. Sc.*, v. 47 (3), 115 [Abstract]
Obeliscoides cuniculi, domestic rabbit, ⁵⁹Fe absorption in small intestine, higher in blood of infected hosts
- Ochoterenella Caballero*, 1944
Bain, O.; and Prod'hon, J., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 721-739
Onchocercidae; Waltonellinae n. subfam.
- Ochoterenella digiticauda Caballero* 1944
Dyer, W. G.; and Altig, R., 1977, *Herpetologica*, v. 33 (3), 293-296
Hyla lanciformis (peritoneal cavity): Santa Cecilia, Napo Province, Ecuador
- Ochoterenella guibei* n. sp., illus.
Bain, O.; and Prod'hon, J., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 721-739
Mantidactylus redimitus (cavite generale): Marojezy, Madagascar
- Octodontoxys* n. gen.
Quentin, J. C.; Courtin, S. S.; and Fontecilla, J. A., 1975, *Bol. Chileno Parasitol.*, v. 30 (1-2), 21-25
Oxyurinae
tod: *O. gigantea* n. sp.
- Octodontoxys gigantea* n.gen., n. sp. (tod), illus.
Quentin, J. C.; Courtin, S. S.; and Fontecilla, J. A., 1975, *Bol. Chileno Parasitol.*, v. 30 (1-2), 21-25
Octodon degus: Parque Nacional de Fray Jorge, Provincia de Coquimbo, Chile
- Octodontoxys gigantea*
Babero, B. B.; and Cattán, P. E., 1975, *Bol. Chileno Parasitol.*, v. 30 (3-4), 68-76
Octodon degus (ciego e intestino grueso): Quebrada de la Plata, Santiago, Chile
- Octodontoxys gigantea* (Quentin y col., 1975)
Cattán, P. E.; George-Nascimento, M.; and Rodríguez, J., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 16-20
prevalence survey of helminths of Octodon degus, seasonal variations, age and sex of hosts: Chile
- Octomyomeris*
Ross, J. F.; and Smith, S. M., 1976, *Canad. J. Zool.*, v. 54 (7), 1084-1102
key
- Octomyomeris muspratti* (Obiamiwe and Macdonald, 1973) n. comb.
Ross, J. F.; and Smith, S. M., 1976, *Canad. J. Zool.*, v. 54 (7), 1084-1102
Syn.: *Reesimeris muspratti* Obiamiwe and Macdonald, 1973
- Octomyomeris troglodytis* sp. n., illus.
Poinar, G. O., jr.; and Sanders, R. D., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 37-41
bionomics; possible biological control agent for treehole mosquitoes
Aedes sierrensis (body cavity): near Novato, Marin County, California
- Odilia Durette-Desset*, 1973
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, Nippostrongylinae
synonymy
- Odontospirura* Wehr, 1933, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
Syn.: *Vaznema Freitas & Lent*, 1947
- Oesophagodontus*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Oesophagodontus*
Reinecke, R. K.; and le Roux, D. J., 1972, *J. South African Vet. Ass.*, v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective

- Oesophagodontus robustus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Oesophagodontus robustus*
Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxbendazole
- Oesophagodontus robustus*
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
horses (large intestine): south-west England
- Oesophagodontus robustus* Giles
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Equus burchelli (intestines): Kenya
- Oesophagonastes*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (1), 19-20
as syn. of *Cyclostrongylus Johnston* & Mawson
- Oesophagonastes gallardi*: Mawson, 1965
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (1), 19-20
as syn. of *Cyclostrongylus wallabiae Johnston* & Mawson, 1939
- Oesophagonastes kartana*: Mawson, 1965
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (1), 19-20
as syn. of *Cyclostrongylus kartana* (Mawson 1955) [n. comb.]
- Oesophagonastes leptos*
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (1), 19-20
as syn. of *Cyclostrongylus leptos* (Mawson 1965) [n. comb.]
- Oesophagonastes parma*: Mawson, 1965
Mawson, P. M., 1977, *Tr. Roy. Soc. South Australia*, v. 101 (1), 19-20
as syn. of *Cyclostrongylus parma* (Johnston & Mawson 1939) [n. comb.]
- Oesophagostomiasis*
Egorov, V. I., 1968, *Botan. i Zool. Issled. Dal'nem Vostokey*, v. 2, 250-255
ascariasis, trichuriasis, oesophagostomiasis, swine, single and mixed infections in various combinations, control studies, best system used dehelminthization 4 times a year
- Oesophagostomiasis*
Rahman, A.; Uddin Ahmed, M.; and Mia, A. S., 1975, *Trop. Animal Health and Prod.*, v. 7 (3), 164
goats: slaughterhouses in Bangladesh
- Oesophagostomiasis*
Trifonov, T.; and Velkov, D., 1975, *Vet. Sbirka*, v. 73 (11), 16-17
ascariasis, trichuriasis, oesophagostomiasis, pigs, dehelminthization with suiverm or hygromix in feed, economically effective
- Oesophagostomiasis*
Wojcik, A. R.; and Grzywinski, L., 1975, *Medycyna Wet.*, v. 31 (10), 597-598
incidence in sheep, economic losses: Torun slaughterhouse
- Oesophagostomoides traguli* Maplestone, 1932
Chabaud, A. G.; and Krishnasamy, M., 1976, *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (388), *Zool.* (270), 721-727
as syn. of *Rhabditostomum traguli* (Maplestone, 1932) n. gen. n. comb.
- Oesophagostomum*
Baines, D. M.; and Colegrave, A. J., 1977, *Vet. Rec.*, v. 100 (11), 217-219
gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales
- Oes[ophagostomum]
Brunsdon, R. V., 1976, *N. Zealand J. Exper. Agric.*, v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Oesophagostomum*
Chhabra, R. C.; Bali, H. S.; and Toor, L. S., 1976, *J. Research, Punjab Agric. Univ.*, v. 13 (3), 308-311
gastrointestinal strongyles in sheep, critical drug evaluation, thiabendazole (most effective), tetramisole (good results), morantel tartrate (fair results), clixanide and methyridine (least effective): India
- Oesophagostomum*
Crowley, J. W., jr.; et al., 1977, *Am. J. Vet. Research*, v. 38 (5), 689-692
lungworms, gastrointestinal parasites, cattle, 3 controlled critical trials, highly effective
- Oesophagostomum*
Dey-Hazra, A., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 198
helminths, pigs, mode of pathogenicity, review
- Oesophagostomum*
Dewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology

- Oesophagostomum
Fudalewicz-Niemczyk, W.; et al., 1975, *Med. Wet.*, v. 31 (11), 666-668
sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Oesophagostomum
Fudalewicz-Niemczyk, W.; et al., 1976, *Acta Zootech.*, Bratislava (32), 5-19
gastrointestinal helminths, mountain sheep, nilverm and zanil, favorable influence on body weight and wool production of treated animals, no influence of treatment on fertility: Poland
- Oesophagostomum
Grzywinski, L.; Martynowicz, T.; and Kluczniok, P., 1976, *Med. Wet.*, v. 32 (4), 227-229
Oesophagostomum, *Ascaris suum*, pigs, Cambendazole effective
- Oesophagostomum
Guarino, C.; and Rivellini, P., 1972, *Atti Soc. Ital. Sc. Vet.*, v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Oesophagostomum
Guimaraes, M. P.; et al., 1976, *Arq. Escola Vet. Univ. Fed. Minas Gerais*, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Oesophagostomum
Jacobs, D.; and Schulze, H. W., 1977, *Prakt. Tierarzt*, v. 58 (1), 46-48
pig parasites, natural infections, vermitin and dichlorvos effective in field testing
- Oesophagostomum
Klein Mori, J., 1972, *Rev. Med. Vet. y Parasitol.*, Maracay, v. 24 (1-8), 1971-1972, 207-226
gastrointestinal nematodes, sheep, Neguvon, Ripercol, Thibenzoline, comparison, various management systems, all effective, Ripercol easiest to administer, Neguvon somewhat toxic
- Oesophagostomum
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1974, *Indian J. Animal Research*, v. 8 (2), 75-78
Haemonchus contortus, other nematodes, experimentally or naturally infected sheep, nitroxylin highly effective, critical testing; in vitro testing against *H. contortus*
- Oesophagostomum
Qadir, A. N. M. A., 1976, *Indian Vet. J.*, v. 53 (11), 855-858
gastrointestinal nematodes, goats and calves, urea for pasture control of free-living stages
- Oesophagostomum
Sewell, M. M. H., 1973, *Vet. Rec.*, v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Oesophagostomum
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, *Am. J. Vet. Research*, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Oesophagostomum
Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Oesophagostomum
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Oesophagostomum
Todd, A. C.; et al., 1976, *Am. J. Vet. Research*, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Oesophagostomum
Zeakes, S. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos crumbles and naftalofos boluses
- Oesophagostomum spp.
van Adrichem, P. W. M.; and Shaw, J. C., 1977, *J. Animal Sc.*, v. 45 (3), 417-422
gastrointestinal nematodes, monozygous twin cattle, comparison of treated and untreated pairs infected naturally on pasture, growth performance, results indicate that the reduced growth may be long-lasting
- Oesophagostomum spp.
van Adrichem, P. W. M.; and Shaw, J. C., 1977, *J. Animal Sc.*, v. 45 (3), 423-429
gastrointestinal nematodes, effects on growth performance and milk production in cambendazole-treated vs. non-treated monozygous twin cattle naturally infected on pasture during the first lactation period
- Oesophagostomum sp.
Bali, M. K.; and Singh, R. P., 1976, *Indian J. Animal Research*, v. 10 (2), 111-112
Haemonchus contortus, *Oesophagostomum* sp., *Trichostrongylus* sp., *Trichuris* sp., sheep, goats, morantel tartrate, good results against all parasites except for *Trichuris* sp.
- Oesophagostomum [sp.]
Cabaret, J., 1976, *Rev. Elevage et Med. Vet. Pays Trop.* v. 29 (3), 221-226
cattle, survey, treatment, economic importance: Kaedi area (Mauritania)

- Oesophagostomum (Hysteracrum) sp. 1, illus.
Chabaud, A. G.; and Krishnasamy, M., 1976,
Bull. Mus. National Hist. Nat., Paris, 3. s.
(388), Zool. (270), 721-727
description
Tragulus javanicus: Selangor, Subang, Su-
bang Forest Reserve
- Oesophagostomum (Hysteracrum) sp. 2, illus.
Chabaud, A. G.; and Krishnasamy, M., 1976,
Bull. Mus. National Hist. Nat., Paris, 3. s.
(388), Zool. (270), 721-727
brief description
Tragulus javanicus (intestin grele): Selan-
gor, Jinjang, Bt. Legong Forest Reserve
- Oesophagostomum sp.
File, S. K.; McGrew, W. C.; and Tutin, C. E.
G., 1976, J. Parasitol., v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces):
Gombe National Park, Tanzania
- [Oesophagostomum] sp. "Oezofagostomum"
Georgieva, D.; Vladimirova, A.; and Monov, M.,
1975, Vet. Sbirka, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of
tetramisole, group and individual appli-
cations
- Oesophagostomum sp.
Grzywinski, L.; et al., 1975, Medycyna Wet.,
v. 31 (9), 524-526
swine, Nilverm by injection for control
- Oesophagostomum sp.
Grzywinski, L.; and Poznanski, W., 1976, Med.
Wet., v. 32 (12), 737-739
Oesophagostomum sp., piglets from sows treat-
ed 2 weeks before farrowing with thibendazole
had higher live weight gain rate than those
from sows treated with nilverm
- Oesophagostomum spp.
Hubert, J., 1977, Rec. Med. Vet., v. 153 (12),
923-929
Ascaris suum (exper.), Trichuris suis
(exper.), Oesophagostomum spp., pigs, com-
parison of coproscopical method of count
after dilution in dense solution and flota-
tion in Mac Master Slides and count after
sedimentation and flotation in dense solution
in Mac Master Slides; various densities of
solutions compared with both techniques
- Oesophagostomum sp. illus.
Kaminsky, R. G.; and Ndinya-Achola, J. O.,
1977, East African Med. J., v. 54 (5), 296-297
immature male worm excised from abscess cav-
ity during biopsy of abdominal wall and
peritoneum of 14 year old boy from Kenya
- Oesophagostomum sp.
Krishna Iyer, P. P.; and Peter C. T., 1975,
Kerala J. Vet. Sc., v. 5 (2), 121-123
gastrointestinal nematodes, goats, methyri-
dine
- Oesophagostomum spp.
Levine, N. D.; et al., 1975, Am. J. Vet. Re-
search, v. 36 (10), 1459-1464
lambs grazing with their ewes under 2 pas-
ture rotation systems, lambs under rotation
had more nematodes and gained less weight
than nonrotated control lambs, rotation is
not recommended to control nematode para-
sitism of sheep in Illinois
- Oesophagostomum sp. 4th stage
Lyons, E. T.; et al., 1975, Am. J. Vet. Re-
search, v. 36 (6), 777-780
calves, natural infections of gastrointest-
inal parasites and lungworms, controlled
test of activity of levamisole administered
via drinking water, subcutaneous injection,
or alfalfa pellet premix
- Oesophagostomum spp.
Maksimovic, A.; and Cvetkovic, L., 1976, Vet.
Glasnik, v. 30 (6), 537-541
Oesophagostomum spp., use of piperazine and
thiabendazole on sows before farrowing, pre-
vention of infection in piglets by preventing
post parturient egg rise in sows
- Oesophagostomum sp.
de Oliveira, A. R., 1976, Arq. Inst. Biol.,
Sao Paulo, v. 43 (1-2), 53-56
Oesophagostomum sp., Cooperia sp., Haemonchus
sp., calves (exper.), no correlation between
level of infestation and circulating eosino-
phils, may result from eosinophil migration
to affected organs or bone marrow exhaustion
- Oesophagostomum spp.
Pfeiffer, A., 1977, Prakt. Tierarzt, v. 58 (1),
32-38
Hyostrogylus rubidus, Oesophagostomum spp.,
sows treated for improved weight gain of
weanling pigs
- Oesophagostomum sp.
Qadir, A. N. M. A., 1976, Indian Vet. J., v.
53 (6), 448-450
Haemonchus sp., Trichostrongylus sp.,
Oesophagostomum sp., larvicidal action of 4
chemical compounds on infective nematode
larvae in experimental outdoor plots; urea
most effective
- Oesophagostomum [sp.]
Rawlings, C. A.; and Splitter, G. A., 1973,
Lab. Animal Sc., v. 23 (2), 259-261
Pneumonyssus simicola in Macaca mulatta
causing multiple bronchopleural fistulas and
resultant tension pneumothorax, Oesophago-
stomum contributing to chronic debilitating
condition, fatal outcome, case report
- Oesophagostomum spp.
Raynaud, J. P.; Sennelier, J.; and Irisarri,
E., 1975, Folia Vet. Latina, v. 5 (3), 412-429
gastrointestinal helminths, swine, post
natal infection of piglets in contact with
infected mothers, comparison of various
methods of husbandry and hygiene, studies
during pregnancy and lactation, routine
daily hygiene recommended

- Oesophagostomum* sp.
Schillhorn van Veen, T.; and Brinckman, W. L., 1975, Samaru Agric. Newsletter, v. 17 (2), 70-74
lambs, regular drenching with thiabendazole at regular intervals during rainy season, better weight gain, cost/benefit; possible influence of resistance and breed of sheep
- Oesophagostomum* sp.
Schulte, J. W.; Klimstra, W. D.; and Dyer, W. G., 1976, J. Wildlife Management, v. 40 (3), 579-581
Odocoileus virginianus clavium (feces): Big Pine Key, Florida
- Oesophagostomum* sp.
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Oesophagostomum* sp.
Tarczynski, S.; Romaniuk, K.; and Szelagiewicz-Czosnek, M., 1972, Med. Wet., v. 28 (4), 217-218
intestinal nematodes, swine, Suiverm
- Oesophagostomum* sp.
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antelope cervicapra (large intestine): Texas
- Oesophagostomum* spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Oesophagostomum* sp.
Tongson, M. S.; and Montenegro, M. M., 1975, Philippine J. Vet. Med., v. 13 (1-2), 170-182
purified microfine phenothiazine + lead arsenate, anthelmintic efficiency, good results, goats: Philippines
- Oesophagostomum* sp.
Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidian drug (Cozurone), good control of all except *Strongyloides*
- Oesophagostomum* spp.
Vegad, J. L.; Kolte, G. N.; and Awadhiya, R.P., Vet. Rec., v. 95 (10), 207-208
Oesophagostomum spp., Rambouillet ram, clinical observations, histopathology, fatal termination due to rupture of necrotic nodule into peritoneal cavity: Adhartal
- Oesophagostomum* [sp.]
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Oesophagostomum* sp.
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 77-81
Connochaetes gnou (colon): Mountain Zebra National Park
- Oesophagostomum aculeatum*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Oesophagostomum asperum* Railliet and Henry, 1913, illus.
Rao, S. H.; and Venkataratnam, A., 1977, Indian Vet. J., v. 54 (1), 14-20
Oesophagostomum asperum, goats (large intestine), morphology and development of eggs and free living larval stages, measurements of larvae compared with *O. venulosum* and *O. columbianum*
- Oesophagostomum asperum* Railliet and Henry, 1913, illus.
Rao, S. H.; and Venkataratnam, A., 1977, Indian Vet. J., v. 54 (2), 102-107
Oesophagostomum asperum in goats, life history study, detailed description of morphology and development of 4th and 5th larval stages
- Oesophagostomum asperum* Railliet and Henry, 1931, illus.
Sathianesan, V.; and Peter, C. T., 1976, Kerala J. Vet. Sc., v. 7 (1), 43-47
Oesophagostomum asperum, development of free living stages, measurements of 3rd stage larva compared with *O. venulosum*
- Oesophagostomum bifurcum*
McConnell, E. E.; et al., 1974, Onderstepoort J. Vet. Research, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (large intestine): Kruger National Park, Transvaal
- Oesophagostomum bifurcum*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Oesophagostomum cervi*
Baker, M. R.; and Pursglove, S. R., jr., 1976, J. Parasitol., v. 62 (1), 166-168
previous reports of *Oesophagostomum cervi* from white-tailed deer probably refer to *O. venulosum*, "The present study suggests that *O. cervi* is a synonym of *O. venulosum*."
- Oesophagostomum columbianum*
Anderson, P. J. S.; and Marais, F. S., 1972, J. South African Vet. Ass., v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Oesophagostomum columbianum*
Campbell, W. C.; and Thomson, B. M., 1973, Austral. Vet. J., v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus columbiformis* proved uninfective even if they had not been frozen

- Oesophagostomum columbianum*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Oesophagostomum columbianum*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Oesophagostomum columbianum*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Oesophagostomum columbianum*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazine, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Oesophagostomum columbianum*
Gerber, H. M., 1975, J. South African Vet. Ass., v. 46 (3), 273-275
percutaneous infestation of pure strains of *Oesophagostomum radiatum* and *O. columbianum* respectively in calves and lambs, fecal egg counts, worm development and recovery
- Oesophagostomum columbianum*
Groeneveld, H. T.; and Reinecke, R. K., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 285-297
non-parametric statistical method for comparing worm burdens in two groups of sheep, application in interpreting results of anthelmintic tests
- Oesophagostomum columbianum*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Oesophagostomum columbianum*
Gupta, O. P.; et al., 1976, Indian J. Exper. Biol., v. 14 (3), 356-357
in vitro anthelmintic activity of embelin disalts, *Paramphistomum cervi*, *Oesophagostomum columbianum*, *Trichuris ovis*, *Dipylidium caninum*, good results
- Oesophagostomum columbianum*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Oesophagostomum columbianum*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21⁹ (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Oesophagostomum columbianum*
Misra, S. C., 1972, Indian J. Animal Research, v. 6 (2), 95-96
parasitic gastro-enteritis, goats, epidemiology, seasonal incidence: Orissa
- Oesophagostomum columbianum*
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Oesophagostomum columbianus* Curtice
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Connochaetes taurinus (large intestine): Kenya
- Oesophagostomum columbianum*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Oesophagostomum columbianum*
Rao, S. H.; and Venkataratnam, A., 1977, Indian Vet. J., v. 54 (1), 14-20
Oesophagostomum asperum, goats (large intestine), morphology and development of eggs and free living larval stages, measurements of larvae compared with *O. venulosum* and *O. columbianum*
- Oesophagostomum columbianum*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific

- Oesophagostomum columbianum*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Oesophagostomum columbianum*
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, Camelus dromedarius, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Oesophagostomum columbianum* Curtice, 1890
Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karoo
- Oesophagostomum columbianum*
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Oesophagostomum columbianum*
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Oesophagostomum dentatum*
Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, Vet. Rec., v. 99 (7), 119-122
swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with picedex
- [*Oesophagostomum dentatum*] *Ezofagostomum dentatum*
Bonev, B.; et al., 1975, Vet. Sbirka, v. 73 (11), 14-15
pigs, [Ascaris suum], [Trichocephalus suis], [*Oesophagostomum dentatum*], hygromycin B in feed, completely controls ascariasis
- Oesophagostomum dentatum*
Bussieras, J., 1976, Rec. Med. Vet., v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Oesophagostomum dentatum*
Corwin, R. M., 1977, Am. J. Vet. Research, v. 38 (4), 465-467
mixed nematode infections, pigs, oxfendazole, critical evaluation: Missouri
- Oesophagostomum dentatum*
Getler, K., 1972, Med. Wet., v. 28 (8), 476-477
nematodes, pigs on industrial swill feeding farm, Atgard
- Oesophagostomum dentatum*
Grzywinski, L.; et al., 1975, Medycyna Wet., v. 31 (9), 524-526
swine, Nilverm by injection for control
- Oesophagostomum dentatum* (Rudolphi, 1803) Molin, 1861, illus.
Kendall, S. B.; Small, A. J.; and Phipps, L. P., 1977, J. Comp. Path., v. 87 (2), 223-229
pigs (exper.)
- Oesophagostomum dentatum*
Kirsch, R.; and Duewel, D., 1975, Research Vet. Sc., v. 19 (3), 327-329
Hyostrongylus rubidus, *Oesophagostomum* spp., pigs (exper.), efficacy of fenbendazole
- Oesophagostomum dentatum*
Oakley, G. A., 1977, Vet. Rec., v. 100 (15), 310-312
Oesophagostomum dentatum, pigs (exper.), levamisole hydrochloride, controlled and critical trials, good results
- Oesophagostomum dentatum*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile
- Oesophagostomum dentatum* Rudolphi, 1885
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 267-271
Sus scrofa: Huesca, Alto Aragon
- Oesophagostomum dentatum*
Raynaud, J. P., 1976, Pathophysiol. Parasit. Infect., 99-104
Oesophagostomum spp., *Hyostrongylus rubidus*, *Ascaris suum*, young swine (exper.), multi-stage multiparasite model for pathological and anthelmintic studies
- Oesophagostomum dentatum*
Rose, J. H., 1973, Research Vet. Sc., v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostrongylus rubidus*, culture from infective larva to adult worm in WAE medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Oesophagostomum dentatum*
Taffs, L. F., 1976, Brit. Vet. J., v. 132 (1), 105-111
cambendazole, little or no effect against 10-day-old *Hyostrongylus rubidus* or *Oesophagostomum* spp. larvae when given to pigs at oral dose rates of 15, 20, and 25 mg/kg bodyweight

- Oesophagostomum dentatum*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (intestino grueso): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Oesophagostomum granatensis*
Raynaud, J. P., 1976, Pathophysiol. Parasit. Infect., 99-104
Oesophagostomum spp., *Hyostrogylus rubidus*, *Ascaris suum*, young swine (exper.), multi-stage multiparasite model for pathological and anthelmintic studies
- Oesophagostomum quadrispinulatum*
Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, Vet. Rec., v. 99 (7), 119-122
swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with pica-dex
- Oesophagostomum quadrispinulatum*
Bussieras, J., 1976, Rec. Med. Vet., v. 152 (3), 219-222
strongyloses of swine, immunological phenomena, clinical manifestations, applications in diagnosis, prophylaxis and treatment, review
- Oesophagostomum quadrispinulatum*
Hubert, J.; Yvore, P.; and Kerboeuf, D., 1976, Ann. Recherches Vet., v. 7 (1), 83-90
parasite survival in liquid manure, anti-parasitic action of xylene
- Oesophagostomum quadrispinulatum* (Marccone, 1901)
Alicata, 1935, illus.
Kendall, S. B.; Small, A. J.; and Phipps, L. P., 1977, J. Comp. Path., v. 87 (2), 223-229
Oesophagostomum quadrispinulatum, pigs (exper.), description, life cycle, pathology
- Oesophagostomum quadrispinulatum*
Kendall, S. B.; Small, A. J.; and Phipps, L. P., 1977, J. Comp. Path., v. 87 (4), 551-555
Oesophagostomum quadrispinulatum, pigs (exper.), repeated infection with 500 larvae induced a solid resistance in about 50 days, barrier to reinfection develops against third stage larvae
- Oesophagostomum quadrispinulatum*
Kirsch, R.; and Duewel, D., 1975, Research Vet. Sc., v. 19 (3), 327-329
Hyostrogylus rubidus, *Oesophagostomum* spp., pigs (exper.), efficacy of fenbendazole
- Oesophagostomum quadrispinulatum*
Raynaud, J. P., 1976, Pathophysiol. Parasit. Infect., 99-104
Oesophagostomum spp., *Hyostrogylus rubidus*, *Ascaris suum*, young swine (exper.), multi-stage multiparasite model for pathological and anthelmintic studies
- Oesophagostomum quadrispinulatum*
Small, A. J.; and Kendall, S. B., 1977, Parasitology, v. 75 (2), x [Abstract]
Oesophagostomum quadrispinulatum, rate of development in pigs, resistance of pigs to re-infection
- Oesophagostomum quadrispinulatum*
Taffs, L. F., 1976, Brit. Vet. J., v. 132 (1), 105-111
cambendazole, little or no effect against 10-day-old *Hyostrogylus rubidus* or *Oesophagostomum* spp. larvae when given to pigs at oral dose rates of 15, 20, and 25 mg/kg bodyweight
- Oesophagostomum quadrispinulatum*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (intestino grueso): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Oesophagostomum radiatum*
Anderson, P. J. S.; and Marais, F. S., 1975, J. South African Vet. Ass., v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Oesophagostomum radiatum* (Rudolphi, 1803)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (large intestine): Kruger National Park
- Oesophagostomum radiatum*
Benz, G. W.; and Ernst, J. V., 1977, Am. J. Vet. Research, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.), albendazole significantly reduced infestations
- Oesophagostomum radiatum*
Bremner, K. C.; Keith, R. K.; and Winks, R., 1976, Research Vet. Sc., v. 20 (3), 350-351
Oesophagostomum radiatum, castrated male calves, resistance to initial infection increases with age
- Oesophagostomum radiatum*
Bryan, R. P., 1976, Austral. Vet. J., v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Oesophagostomum radiatum*
Bryan, R. P.; Bainbridge, M. J.; and Kerr, J. D., 1976, Austral. J. Zool., v. 24 (3), 417-421
Bubalus bubalis
cattle
(large and small intestine of all): all from Northern Territory, Australia
- Oesophagostomum radiatum*
Campbell, W. C.; and Thomson, B. M., 1973, Austral. Vet. J., v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus colubriformis* proved uninfected even if they had not been frozen

- Oesophagostomum radiatum*
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of heifers, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Oesophagostomum radiatum*
Ciordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Oesophagostomum radiatum*
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxbendazole, drug efficacy
- Oesophagostomum radiatum*
Curr, C., 1977, Austral. Vet. J., v. 53 (9), 425-428
nematodes, calves, levamisole, efficiency of pour-on formulation, drug trials, good results
- Oesophagostomum radiatum*
Dharsana, R. S.; Fabiyi, J. P.; and Hutchinson, G. W., 1976, Vet. Parasitol., v. 2 (4), 333-340
mixed gastro-intestinal nematode infections, calves, effects on host intestinal enzymes
- Oesophagostomum radiatum*
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Oesophagostomum radiatum*
Fincher, G. T., 1975, J. Parasitol., v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Oesophagostomum radiatum*
Gerber, H. M., 1975, J. South African Vet. Ass., v. 46 (3), 273-275
percutaneous infestation of pure strains of *Oesophagostomum radiatum* and *O. columbianum* respectively in calves and lambs, fecal egg counts, worm development and recovery
- Oesophagostomum radiatum*
Goldberg, A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Oesophagostomum radiatum*
Herlich, H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.), oxbendazole, efficacy against adult and larval stages
- Oesophagostomum radiatum*
Herlich, H., 1977, Am. J. Vet. Research, v. 38 (8), 1247-1248
efficacy of albendazole against gastrointestinal nematodes and *Fasciola hepatica* in cattle (exper.); comparison of critical vs. controlled tests
- Oesophagostomum radiatum* Rudolphi, 1803
Hiregoudar, L. S., 1976, Indian Vet. J., v. 53 (3), 237
Axis axis (large intestine): Gir forest, Gujarat State, India
- Oesophagostomum radiatum*
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Oesophagostomum radiatum*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure
- Oesophagostomum radiatum*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Oesophagostomum radiatum*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Oesophagostomum radiatum*
Randall, R. W.; and Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Oesophagostomum radiatum*
Reinecke, R. K., 1972, Onderstepoort J. Vet. Research, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Oesophagostomum radiatum*
Ronald, N. C.; Bell, R. R.; and Craig, T. M., 1977, J. Am. Vet. Med. Ass., v. 170 (3), 317-319
gastrointestinal nematodes, calves, levamisole phosphate, effective at one-half recommended dosage

- Oesophagostomum radiatum*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Oesophagostomum radiatum*
Rowlands, D. ap T.; and Berger, J., 1977, J. South African Vet. Ass., v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Oesophagostomum radiatum*
Schroeder, J.; Honer, M. R.; and Louw, J. P., 1977, J. South African Vet. Ass., v. 48 (2), 95-97
trematodes, nematodes, cattle (exper.), rafoxanide, efficacy of subcutaneous injections against immature larvae and adults
- Oesophagostomum radiatum*
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Oesophagostomum radiatum*
Searson, J. E.; and Doughty, F. R., 1977, Austral. Vet. J., v. 53 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult *Ostertagia ostertagi* than larval forms): southern New South Wales
- Oesophagostomum radiatum*
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Oesophagostomum radiatum* (Rudolphi, 1803)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Bos taurus: insular Newfoundland
- Oesophagostomum* (*Bosicola*) *radiatum*
Tager-Kagan, P., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (4), 317-321
gastro-intestinal nematodes, zebu cattle (1 to 2 years old), cambendazole: Niger
- Oesophagostomum radiatum*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcvi
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Oesophagostomum radiatum*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Oesophagostomum radiatum*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Oesophagostomum radiatum*
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Oesophagostomum radiatum*
Willadsen, P., 1977, Austral. J. Biol. Sc., v. 30 (5), 411-419
Oesophagostomum radiatum, isolation and partial characterization of α -chymotrypsin inhibitor in the cuticle, hypodermis, and musculature, affinity chromatography compared with ion-exchange and molecular sieve chromatography
- Oesophagostomum radiatum*
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Oesophagostomum radiatum*
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Oesophagostomum* (*Conoweberia*) *selfi* sp. n., *illus.*
Schmidt, G. D.; and Kuntz, R. E., 1975, Proc. Oklahoma Acad. Sc., v. 55, 163-165
Rattus coxinga coxinga (small intestine): Chuei-feng, Nan-tou Hsien, Taiwan, Republic of China
- Oesophagostomum* (*Proteracum*) *synceri* n. sp., *illus.*
Troncy, P. M.; and Thal, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 173-176
Bubalus caffer aequinoxialis (gros intestin): Republique Centrafricaine and Tchad
- Oesophagostomum venulosum* (Rudolphi 1809) *Raillet* et Henry 1913
Baker, M. R.; and Pursglove, S. R., jr., 1976, J. Parasitol., v. 62 (1), 166-168
previous reports of *Oesophagostomum cervi* from white-tailed deer probably refer to *O. venulosum*, "The present study suggests that *O. cervi* is a synonym of *O. venulosum*."
Odocoileus virginianus: Norfolk County, Ontario; Warren County, Mississippi; Adair County, Oklahoma

- Oesophagostomum venulosum* (Rudolphi, 1809)
 Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
 brief description
 sheep (large intestine, cecum): vicinity of Nowy Targ, Carpathian Mountains
- Oesophagostomum venulosum*
 Boag, B.; and Thomas, R. J., 1977, *Research Vet. Sc.*, v. 22 (1), 62-67
 gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Oesophagostomum venulosum*
 Chalmers, K., 1977, *N. Zealand Vet. J.*, v. 25 (10), 266-269
 gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Oesophagostomum venulosum*
 Chroust, K.; and Dyk, V., 1975, *Deutsche Tierarztl. Wchnschr.*, v. 82 (12), 487-491
 gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Oesophagostomum venulosum*
 Colglazier, M. L.; et al., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 145-150
 gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with *Trichuris* spp. and *Moniezia expansa*
- Oesophagostomum venulosum*
 Corticelli, B.; and Lai, M., 1972, *Parassitologia*, v. 14 (1), 95-96
Ovis musimon: Sardegna
- Oesophagostomum venulosum*
 Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (1), 65-77
 roe deer (digestive tract): Czechoslovakia
- Oesophagostomum venulosum*
 Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
 helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
 (digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Oesophagostomum venulosum*
 Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
 coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
 all from Czechoslovakia
- Oesophagostomum venulosum*
 Dyk, V.; and Chroust, K., 1975, *Veterinarstvi*, v. 25 (7), 315-317
 helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Oesophagostomum venulosum*
 Gonzalez, H.; and Plaza, J., 1968, *Bol. Chileno Parasitol.*, v. 23 (3-4), 134-137
 gastrointestinal nematodes of sheep, comparative therapeutic trials using banminth, phenothiazine, and thiabendazole
- Oesophagostomum venulosum*
 Heuer, D. E.; et al., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 141-143
Odocoileus virginianus (cecum): Kentucky
- Oesophagostomum venulosum* (Rudolphi 1809),
 Railliet, 1896
 Hiregoudar, L. S., 1976, *Indian Vet. J.*, v. 53 (3), 237
Boselaphus tragocamelus (large intestine): Gir forest, Gujarat State, India
- Oesophagostomum* (H.) *venulosum* (Rudolphi, 1809)
 Railliet et Henry, 1913
 Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus (large intestine): southern Bulgaria
- Oesophagostomum venulosum*
 Kelly, J. D.; et al., 1975, *Research Vet. Sc.*, v. 19 (1), 105-107
 anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Oesophagostomum venulosum*
 Kennedy, T. J.; and Todd, A. C., 1975, *Am. J. Vet. Research*, v. 36 (10), 1465-1467
 gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Oesophagostomum venulosum*
 Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
 nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Oesophagostomum venulosum*
 Le Jambre, L. F.; and Royal, W. M., 1976, *Austral. Vet. J.*, v. 52 (4), 181-183
 nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (faecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except *Nematodirus* resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales

- Oesophagostomum venulosum*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Oesophagostomum venulosum*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Ovis aries: Chile
- Oesophagostomum venulosum*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries
Odocoileus virginianus
all from Hardy County, West Virginia
- Oesophagostomum venulosum*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Oesophagostomum venulosum* (Rudolphi, 1809)
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (large intestine): Oklahoma
- Oesophagostomum venulosum*
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Oesophagostomum venulosum*
Rao, S. H.; and Venkataratnam, A., 1977, Indian Vet. J., v. 54 (1), 14-20
Oesophagostomum asperum, goats (large intestine), morphology and development of eggs and free living larval stages, measurements of larvae compared with *O. venulosum* and *O. columbianum*
- Oesophagostomum venulosum*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Oesophagostomum venulosum*
Sathianesan, V.; and Peter, C. T., 1976, Kerala J. Vet. Sc., v. 7 (1), 43-47
Oesophagostomum asperum, development of free living stages, measurements of 3rd stage larva compared with *O. venulosum*
- Oesophagostomum venulosum*
Schweisgut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Oesophagostomum venulosum*
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Oesophagostomum venulosum* (Rudolphi, 1809)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Oesophagostomum walkeri* Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii: Kenya
- Ohbayashinema* n. gen.
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 415-418
Heligmosomidae, Heligmosominae
tod: *Ohbayashinema ochotoni* n. gen., n. sp. evolution
- Ohbayashinema* Durette-Desset, 1974
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
- Ohbayashinema dubinini* (Gvozdev, 1966) n. comb.
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 415-418
Syn.: *Heligmosomum dubinini* Gvozdev, 1966
- Ohbayashinema ochotoni* n. gen., n. sp. (tod), illus.
Durette-Desset, M. C., 1974, Bull. Mus. National Hist. Nat., Paris, 3. s. (216), Zool. (144), 415-418
Ochotona macrotis (intestin): Gasainhund, Nepal
- Okapinema* Ivaschkin, 1960, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Parabronematinae
key
- Ollulanidae
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
- Ollulanus tricuspis*
Gregory, G. G.; and Munday, B. L., 1976, Austral. Vet. J., v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island

- Ollulanus tricuspis* (Leuckart 1865), illus.
Haenichen, T.; and Hasslinger, M. A., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (3), 59-62
Ollulanus tricuspis, case history of chronic gastritis in cat, pathogenicity and incidence discussed
- Ollulanus tricuspis* Leuckart, 1865, illus.
Hinaidy, H. K., 1976, Zentralbl. Vet.-Med., Reihe B, v. 23 (1), 66-73
synonymy, description of spicule
Vulpes vulpes: Osterreich
- Onchocerca*
Vargas, L., 1976, Rev. Invest. Salud Pub., Mexico, v. 36 (3), 185-186
human onchocerciasis, possible diagnosis of onchocercoma using xeroradiography
- Onchocerca* sp., illus.
Ali-Khan, Z., 1977, Ann. Trop. Med. and Parasitol., v. 71 (4), 469-482
Onchocerca, comparative morphology, histology and measurements of one male and one female specimen of *Onchocerca* sp. excised from the wrist of an Ontario, Canada resident, 2 bovine spp. from Canada and *O. cervicalis* from the ligamentum nuchae of a horse from Georgia, U.S.A.
- Onchocerca* spp., illus.
Prasad, M. C.; Rajya, B. S.; and Mohanty, G. C., 1977, Indian J. Animal Sc., v. 45 (5), 1975, 270-274
Onchocerca spp., goats (aortas), incidence, pathology: India
- Onchocerca* sp., illus.
Rehbinder, C.; Christensson, D.; and Glatthard, V., 1975, Nord. Vet.-Med., v. 27 (10), 499-507
Onchocerca sp. in subperitoneal and subcutaneous granulomas and *Setaria tundrae* in encapsulations in peritoneum of reindeer, increasing incidence, association with liver lesions caused by *Corynebacteria*, found in forest herds but not mountain herds
- Onchocerca* sp.
Scholtens, R. G.; and Adams, S. R., 1977, Am. J. Vet. Research, v. 38 (7), 1099-1101
microfilariae, cattle (skin), microbiopsy technique, survey over two years, evidence of local transmission: near Lawrenceville, Georgia
- Onchocerca armillata* Railliet et Henry, 1909, illus.
Bain, O., [1976], Ann. Parasitol., v. 50 (6), 1975, 763-788
redescription, differentiation from other *Onchocerca* spp.
- Onchocerca armillata*, illus.
Chauhan, P. P. S.; and Pande, B. P., 1972, Indian J. Animal Sc., v. 42 (10), 809-813
Onchocerca armillata, lesions in aortae of buffaloes and bullocks, histopathological studies: abattoirs, Uttar Pradesh
- Onchocerca armillata*
ElBihari, S.; and Hussein, H. S., 1975, J. Parasitol., v. 61 (4), 656
Onchocerca armillata, location of microfilariae in rat (exper.) and cow, found in skin not blood
- Onchocerca armillata* (Railliet and Henry, 1909)
Schillhorn-Van-Veen, T. W.; Bello, S. I.; and Folaranmi, D. O. B., 1976, Rev. Elevage et Med. Vet. Pays Trop., v. 29 (3), 227-228
Camelus dromedari (aorta): Kano abattoir, northern Nigeria
- Onchocerca armillata*
Schillhorn van Veen, T.; and Robl, M. G., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (3), 305-310
Onchocerca armillata, cattle (White Fulani and N'bororo breeds), incidence and description of lesions in aortic wall: Nigeria
- Onchocerca armillata*
Zak, F., 1975, Pathol. et Microbiol., v. 43 (2-3), 150 [Abstract]
aortic calcifications in cattle resulting from *Onchocerca armillata* infection: West Sudan
- Onchocerca armillata*
al-Zubaidy, A. J., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (4), 436 [Demonstration]
Onchocerca armillata, cattle, parasitic aortitis: Iraq
- Onchocerca bohmi* (Supperer, 1953) n. comb.
Bain, O.; et al., 1976, J. Helminth., v. 50 (4), 287-293
- Onchocerca caecutiens*
Hanna, L. S.; Abboud, I. A.; and Ragab, H. A. A., 1973, Bull. Ophth. Soc. Egypt, v. 66, 575-586
Onchocerca caecutiens living and dead microfilariae injected sub-conjunctivally and into anterior chamber of hamster and guinea pigs, pathology of developing lesions
- Onchocerca cervicalis*, illus.
Ali-Khan, Z., 1977, Ann. Trop. Med. and Parasitol., v. 71 (4), 469-482
Onchocerca, comparative morphology, histology and measurements of one male and one female specimen of *Onchocerca* sp. excised from the wrist of an Ontario, Canada resident, 2 bovine spp. from Canada and *O. cervicalis* from the ligamentum nuchae of a horse from Georgia, U.S.A.
- Onchocerca cervicalis* Railliet et Henry, 1910, illus.
Bain, O., [1976], Ann. Parasitol., v. 50 (6), 1975, 763-788
redescription, differentiation from other *Onchocerca* spp.
- Onchocerca cervicalis* microfilariae
McCullough, C.; et al., 1977, J. Parasitol., v. 63 (6), 1065
horses: Maryland

- Onchocerca cervicalis*, illus.
Marcoux, M.; Frechette, J. L.; and Morin, M., 1977, *Canad. Vet. J.*, v. 18 (4), 108-110
cheval (peau), clinical signs, diagnosis: Quebec
- Onchocerca cervicalis*
Maurer, N. D.; and Bell, R. R., 1977, *Southwest. Vet.*, v. 30 (2), 176-179
Onchocerca cervicalis, naturally infected-ponies, injectable trichlorfon, no effect: Texas
- Onchocerca cervicalis* Railliet and Henry 1910
Mellor, P. S., 1975, *J. Helminth.*, v. 49 (1), 33-42
Onchocerca cervicalis, migration and development of larvae in *Culicoides nubeculosus* and *C. variipennis* (both exper.)
- Onchocerca cervicalis*
Schulz-Key, H.; Bain, O.; and Wenk, P., 1976, *Tropenmed. u. Parasitol.*, v. 27 (2), 229-232
distribution pattern of *Onchocerca garmsi* microfilariae in the skin of *Cervus elaphus* and comparisons with distribution of adults and microfilariae of *Onchocerca tubingensis*, *O. cervicalis*, *O. tarsicola* and *Wehrdikmansia rugosicauda*: Suddeutschland
- Onchocerca cervicalis*
Stannard, A. A.; and Cello, R. M., 1975, *Am. J. Vet. Research*, v. 36 (7), 1029-1031
Onchocerca cervicalis, horses, prevalence, more common in older animals, occurred in both sexes equally, distribution and concentration of microfilariae within skin, cutaneous pathologic changes, invasion of eye with microfilariae in 60% of cases
- Onchocerca cervipedis* Wehr et Dikmans, 1935
Bain, O.; and Schulz-Key, H., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 437-449
Wehrdikmansia cervipedis returned to *Onchocerca*
- Onchocerca dermatata* Bain, Bussieras et Amegee, 1974
Bain, O.; Bussieras, J.; and Amegee, E., 1976, *Ann. Parasitol.*, v. 51 (4), 461-471
as syn. of *O. ochengi* Bwngamoi, 1969
- Onchocerca dewittei* n. sp., illus.
Bain, O.; et al., 1977, *Ann. Parasitol.*, v. 52 (4), 471-479
Sus scrofa jubatus (pied, niveau des metatarses (tendons et conjonctif sous-cutane)): reserve de Krau, Etat de Pahang, Malaisie
- Onchocerca dukei* Bain, Bussieras et Amegee, 1974, illus.
Bain, O.; Bussieras, J.; and Amegee, E., 1976, *Ann. Parasitol.*, v. 51 (4), 461-471
redescription
Bos taurus (tissus conjonctifs sous-cutane et musculaire, niveau des muscles de la poitrine): Togo (Lama Kara)
- Onchocerca flexuosa* (Wedl, 1856), illus.
Bain, O.; and Schulz-Key, H., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 437-449
redescription
Cervus elaphus (nodules sous-cutanes du tronc, generalement dorsaux): Allemagne Occidentale
- Onchocerca flexuosa* (Wedl, 1856), illus.
Schulz-Key, H., 1975, *Tropenmed. u. Parasitol.*, v. 26 (1), 60-69
Onchocerca flexuosa, development of nodules on hide of *Cervus elaphus*, relationship to age and sex of parasites, parasitic reproduction within nodules
- Onchocerca flexuosa* (Wedl, 1856), illus.
Schulz-Key, H., 1975, *Tropenmed. u. Parasitol.*, v. 26 (3), 348-358
Filariidae spp., microfilariae infecting *Cervus elaphus*, morphological comparisons and localization patterns
Cervus elaphus (posterior part of body mainly on inner hind legs): southern Germany
- Onchocerca garmsi* Bain and Schulz-Key, 1976 [nom. nud.]
Schulz-Key, H.; Bain, O.; and Wenk, P., 1976, *Tropenmed. u. Parasitol.*, v. 27 (2), 229-232
distribution pattern of *Onchocerca garmsi* microfilariae in the skin of *Cervus elaphus* and comparisons with distribution of adults and microfilariae of *Onchocerca tubingensis*, *O. cervicalis*, *O. tarsicola* and *Wehrdikmansia rugosicauda*: Suddeutschland
- Onchocerca garmsi* n. sp., illus.
Bain, O.; and Schulz-Key, H., 1976, *Tropenmed. u. Parasitol.*, v. 27 (4), 474-478
Cervus elaphus (tissu sous-cutane du poitrail): Allemagne Occidentale
- Onchocerca gutturosa*
Amakiri, S. F., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (2), 123-128
Bos indicus (skin from horn and hump regions): Nigeria
- Onchocerca gutturosa* Neumann, 1910, illus.
Bain, O., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 763-788
redescription, differentiation from other *Onchocerca* spp.
- Onchocerca gutturosa*
Bartlett, A.; Bidwell, D. E.; and Voller, A., 1975, *Tropenmed. u. Parasitol.*, v. 26 (3), 370-374
Onchocerca volvulus, evaluation of enzyme immunoassay (ELISA) for the diagnosis of human infections, use of *O. gutturosa* antigens more promising than antigens prepared from various other nematodes
- Onchocerca gutturosa*, illus.
Chauhan, P. P. S.; and Pande, B. P., 1972, *Indian J. Animal Sc.*, v. 42 (10), 809-813
Onchocerca gutturosa, lesions in ligamentum nuchae of buffaloes and bullocks, histopathological studies: abattoirs, Uttar Pradesh
- Onchocerca gutturosa*
Denham, D. A.; and Mellor, P., 1976, *J. Helminth.*, v. 50 (1), 49-52
Onchocerca gutturosa, cattle, anthelmintic effects of compound "E", possible tertiary screening system for drug action against *O. volvulus* in man

- Onchocerca gutturosa*
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Onchocerca gutturosa*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Simuliidae sp.: Uzbekistan
- Onchocerca jakutensis*
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
given in Addendum as *O. jakutensis*; possibly misprint or possibly transfer from *Acanthospiculum*
- Onchocerca lienalis*, *illus.*
Scholtens, R. G.; Adams, S. R.; and Broderson, J. R., 1977, Am. J. Vet. Research, v. 38 (7), 1093-1097
prevalence in *Bos taurus* (skin, fascia of nuchal ligaments), possible animal model for study of human *Onchocerca*: slaughterhouse, Talmo, Georgia
- Onchocerca ochengi* Bwagamoi, 1969, *illus.*
Bain, O.; Bussieras, J.; and Amegee, E., 1976, Ann. Parasitol., v. 51 (4), 461-471
redescription
Syn.: *O. dermatata* Bain, Bussieras et Amegee, 1974
Bos taurus (intradermique): Togo
Bos indicus (intradermique): provenant de Haute-Volta, abattus au Togo
bovins (fragments de peau, tissu mammaire): Afrique Orientale
- Onchocerca raillieti* sp. n., *illus.*
Bain, O.; et al., 1976, J. Helminth., v. 50 (4), 287-293
Equus asinus (niveau du garrot, kyste sous-cutane du penis, conjonctif perimusculaire du flanc): Egypte; Nigeria; Kenya (Mombasa)
- Onchocerca reticulata*, Diesing, 1841, *illus.*
Bain, O., [1976], Ann. Parasitol., v. 50 (6), 1975, 763-788
redescription, differentiation from other *Onchocerca* spp.
- Onchocerca synceri* Sandground, 1938, *illus.*
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (subcutis): Kruger National Park
- Onchocerca tarsicola* n. sp., *illus.*
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
Cervus elaphus (sous-cutanee, au niveau des articulations tibiotarsale et radiocarpale, sur les tendons flechisseurs et partiellement entre les os et la peau): Allemagne Occidentale
- Onchocerca tarsicola* Bain und Schulz-Key 1974, *illus.*
Schulz-Key, H., 1975, Tropenmed. u. Parasitol., v. 26 (3), 348-358
Filariidae spp., microfilariae infecting *Cervus elaphus*, morphological comparisons and localization patterns
Cervus elaphus (skin of ears and nose): southern Germany
- Onchocerca tarsicola*
Schulz-Key, H.; Bain, O.; and Wenk, P., 1976, Tropenmed. u. Parasitol., v. 27 (2), 229-232
distribution pattern of *Onchocerca garmsi* microfilariae in the skin of *Cervus elaphus* and comparisons with distribution of adults and microfilariae of *Onchocerca tubingensis*, *O. cervicalis*, *O. tarsicola* and *Wehrdikmansia rugosicauda*: Suddeutschland
- Onchocerca tubingensis* n. sp. [lapsus p. 444 for *O. tubingensis* n. sp.]
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
- Onchocerca tubingensis* n. sp., *illus.*
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
[lapsus p. 444 as *O. tubingensis* n. sp.]
Syn.: *Wehrdikmansia cervipedis sensu* Tarczyński, 1954
Cervus elaphus (nodules sous-cutanes du tronc, generalement dorsaux): Allemagne Occidentale
- Onchocerca tubingensis* Bain et Schulz-Key, 1974, *illus.*
Bain, O.; and Schulz-Key, H., 1976, Tropenmed. u. Parasitol., v. 27 (4), 474-478
Onchocerca tubingensis, description of female and morphologic comparisons with *O. garmsi* n. sp.
- Onchocerca tubingensis* Bain und Schulz-Key 1974, *illus.*
Schulz-Key, H., 1975, Tropenmed. u. Parasitol., v. 26 (3), 348-358
Filariidae spp., microfilariae infecting *Cervus elaphus*, morphological comparisons and localization patterns
Cervus elaphus (ventral part of body mostly on sternum and inner sides of hindlegs): southern Germany
- Onchocerca tubingensis*
Schulz-Key, H.; Bain, O.; and Wenk, P., 1976, Tropenmed. u. Parasitol., v. 27 (2), 229-232
distribution pattern of *Onchocerca garmsi* microfilariae in the skin of *Cervus elaphus* and comparisons with distribution of adults and microfilariae of *Onchocerca tubingensis*, *O. cervicalis*, *O. tarsicola* and *Wehrdikmansia rugosicauda*: Suddeutschland
- Onchocerca volvulus*
Abdalla, R. E., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 53-55
Wuchereria bancrofti and *Onchocerca volvulus* as cause of human filariasis with hydrocele and elephantiasis, newly reported cases in previously non-endemic areas, need for differentiation from *Loa loa*: Sudan

Onchocerca volvulus

Ambroise-Thomas, P.; Kien Truong, T., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 435-452

filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (*Dipetalonema viteae*, *Dirofilaria immitis*, *Wuchereria bancrofti*, *Loa loa*, *Onchocerca volvulus*), possible application to epidemiological surveys and post-therapeutic surveillance

Onchocerca volvulus

Anderson, J.; et al., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 190-208

Onchocerca volvulus, survey of total population 5 years old and over in 22 village groups using standardized techniques to assess eye and skin lesions, comparison with persons not infected with onchocerciasis: United Cameroon Republic

Onchocerca volvulus

Anderson, J.; et al., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 209-222

Onchocerca volvulus, survey of total populations aged 5 years and older in 16 villages of rain-forest and savanna zones, standard techniques used to assess intensity of infection, clinical manifestations; differences thought to be influenced by hormonal factors, strain pathogenicity, transmission patterns: United Cameroon Republic

Onchocerca volvulus

Anderson, J.; and Fuglsang, H., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (4), 544-548

Onchocerca volvulus, human, variation in numbers of microfilariae in anterior chamber of eye

Onchocerca volvulus

Anderson, J.; and Fuglsang, H., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (5), 710-717

Onchocerca volvulus, human, effects of 3% diethylcarbamazine eye drops on microfilariae in cornea and anterior chamber, numbers reduced but not eliminated and reinvasion occurred soon after treatment stopped, some severe reactions particularly in heavily infected subjects: Cameroon

Onchocerca volvulus

Anderson, J.; Fuglsang, H.; and Bird, A. C., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 378-395

Onchocerca volvulus, comparative pathologic study of posterior segment ocular lesions of infected villagers from savanna and rain-forest regions using fluorescein fundus angiography: Cameroon

Onchocerca volvulus

Anderson, J.; Fuglsang, H.; and Marshall, T. F. de C., 1976, Tropenmed. u. Parasitol., v. 27 (3), 263-278

Onchocerca volvulus, human eye lesions, trials with diethylcarbamazine with and without added effects of corticosteroids (betamethazone), value of therapy varied with type of eye lesions: north Cameroon

Onchocerca volvulus

Anderson, J.; Fuglsang, H.; and Marshall, T. F. de C., 1976, Tropenmed. u. Parasitol., v. 27 (3), 279-296

Onchocerca volvulus, human eye lesions, trials with suramin alone or followed by course of diethylcarbamazine, value of therapy varied with type of eye lesions, severe toxic reactions (2 fatalities) in some persons treated with suramin: north Cameroon

Onchocerca volvulus

Anderson, J.; Fuglsang, H.; and Marshall, T. F. de C., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 362-373

Onchocerca volvulus, comparative follow-up epidemiologic study of infected villagers from savannah and rain-forest areas, relationships between development of eye lesions and high concentrations of microfilariae in skin, particularly around shoulders, as well as eye, implications for prevention of blindness: Cameroon

Onchocerca volvulus

Anderson, R. I., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (3), 418 [Letter]

Onchocerca volvulus, human, location of microfilariae, in addition to skin and eye also found in urine which suggests invasion of renal system and possibly even other internal organs: Chad

Onchocerca volvulus

Anderson, R. I.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (1), 66-70

prevalence survey for microfilaruriae and other manifestations of human *Onchocerca volvulus* infection in rural area, correlation with microfilariae in skin snips, age and sex of hosts: Cameroon

Onchocerca volvulus

Anderson, R. I.; and Buck, A. A., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 447-456

complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad

Onchocerca volvulus

Anderson, R. I.; Fazen, L. E.; and Buck, A. A., 1975, Am. J. Trop. Med. and Hyg., v. 24 (1), 58-61

migration of *Onchocerca volvulus* into urine, blood and sputum after medication of infected humans with diethylcarbamazine: Guatemala

Onchocerca volvulus

Anderson, R. I.; Fazen, L. E.; and Buck, A. A., 1975, Am. J. Trop. Med. and Hyg., v. 24 (1), 62-65

daytime periodicity of *Onchocerca volvulus* established in Guatemala with maximal density of microfilariae in skin coinciding with most active feeding time of principal *Simulium* vector in area

- Onchocerca volvulus*
Asibey, E. O. A., 1977, Environment. Conservation, v. 4 (4), 291-195
Onchocerca volvulus, control of blackfly vector, problems and effectiveness of mass spraying, urgent need for more information concerning distribution of disease, possible environmental and economic effects of spraying and other control measures: West Africa
- Onchocerca volvulus* (Leuckart, 1893), illus.
Bain, O., [1976], Ann. Parasitol., v. 50 (6), 1975, 763-788
redescription, differentiation from other *Onchocerca* spp.
- Onchocerca volvulus*
Bain, O., 1976, Bull. World Health Organ., v. 54 (4), 397-401
human filariasis, number of developing and infective larvae dependent upon number of microfilariae penetrating into haemocoel of vector, relationship based on proportionality, facilitation and limitation, application to disease control and treatment methods
- Onchocerca volvulus*
Bain, O.; et al., 1976, Compt. Rend. Acad. Sc., Paris, v. 283, s. D (4), 391-392
Onchocerca volvulus, correlation between thickness of vector's (*Simulium damnosum*) peritrophic membrane and number of microfilariae ingested explains 'limitation phenomenon'
- Onchocerca volvulus*
Bartlett, A.; Bidwell, D. E.; and Voller, A., 1975, Tropenmed. u. Parasitol., v. 26 (3), 370-374
Onchocerca volvulus, evaluation of enzyme immunoassay (ELISA) for the diagnosis of human infections, use of *O. gutturosa* antigens more promising than antigens prepared from various other nematodes
- Onchocerca volvulus*
Bradley, A. K., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (3), 225-229
Onchocerca volvulus, relationship of prevalence of onchocerciasis and resulting blindness to abandonment of native villages, statistical survey of various geographic localities: Middle Hawal Valley, Nigeria
- Onchocerca volvulus*
Braun-Munzinger, R. A.; Scheiber, P.; and Southgate, B. A., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 548-549
Onchocerca volvulus, humans, simplified modifications of the microtitration plate technique for determining microfilariae densities in skin snips, use in epidemiologic surveys
- Onchocerca volvulus*
Braun-Munzinger, R. A.; and Southgate, B. A., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (5), 385 [Demonstration]
Onchocerca volvulus, 4 distinct types differentiated in 4 native villages of Togo
- Onchocerca volvulus*
Brinkmann, U. K., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 397-403
Onchocerca volvulus, human, diagnosis, dimensions of skin snips obtained with modified scleral punch (weight, surface size, largest diameter), pattern of emergence of microfilariae: Liberia
- Onchocerca volvulus*
Brinkmann, U. K.; et al., 1976, Bull. World Health Organ., v. 54 (6), 708-709
Onchocerca volvulus microfilariae, transplacental transmission with microfilariae found in skin and umbilical cords of newborn infants examined at time of delivery: Upper Region, Ghana
- Onchocerca volvulus*
Bryceson, A. D. M., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 397-399
Onchocerca volvulus in humans, evidence that host immune response underlies pathogenesis of infection and that variations in responses contribute to geographical differences in disease patterns, review
- Onchocerca volvulus*
Bryceson, A. D. M.; et al., 1976, Clin. and Exper. Immunol., v. 24 (1), 168-176
Onchocerca volvulus, antigenic diversity among worms from one village in Nigeria, consistent differences in worm antigen patterns and antibody response with worms originating from forest vs. savanna zones of United Cameroon Republic, relevance of findings to pathology and prevention of onchocerciasis
- Onchocerca volvulus*
Buck, A. A., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (3), 336-338
Onchocerca volvulus, human, microfilaruria, review and research recommendations: Africa
- Onchocerca volvulus*
Buck, A. A.; Anderson, R. I.; and MacRae, A. A., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (1), 21-31
serum immunoglobulin levels in five villages compared, comparative studies of IgG, IgA, IgM, and IgD levels between *Onchocerca volvulus* patients with and without microfilaruria, different age and sex patterns, effect of infection intensity, IgE and combined infection with *Schistosoma mansoni*: Chad
- Onchocerca volvulus*
Budden, F. H., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (3), 265-266
Onchocerca volvulus, route of entry of microfilariae into human eye

- Onchocerca volvulus*
Budden, F. H., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (5-6), 1976, 484-491
Onchocerca volvulus, human ocular involvement, assessment of new eye lesions and progression of existing lesions 14-15 years after single course of suramin, comparison with control group shows slightly less deterioration in treated group and no latent effects of suramin therapy: Hawal Valley, Nigeria
- Onchocerca volvulus*
Cappuccinelli, P., 1972, Parassitologia, v. 14 (2-3), 255-260
Armillifer armillatus, antigens, complement fixation and immunodiffusion studies of antibody response in rabbit; identification of active fractions by immunoelectrophoresis; immunodiffusion tests against Echinococcus granulosus, Fasciola hepatica, Dicrocoelium dendriticum, Onchocerca volvulus and Ascaris suum, no common antigens found
- Onchocerca volvulus*
Chlebowski, H. O.; and Zielke, E., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (2), 181-182 [Letter]
Wuchereria bancrofti, Onchocerca volvulus in humans, diagnosis of microfilaruria using a modified membrane filter technique; application also to filtration of eggs from urine of persons suspected to have schistosomal infections: Liberia, West Africa
- Onchocerca volvulus*
Collins, R. C.; et al., 1977, Tropenmed. u. Parasitol., v. 28 (2), 235-243
Onchocerca volvulus, quantitative aspects of infection in Simulium vectors after feeding on humans with moderate numbers of microfilariae, statistical study
- Onchocerca volvulus*
Denham, D. A.; and Mellor, P., 1976, J. Helminth., v. 50 (1), 49-52
Onchocerca gutturosa, cattle, anthelmintic effects of compound "E", possible tertiary screening system for drug action against *O. volvulus* in man
- Onchocerca volvulus*
Discamps, G.; et al., 1975, Medecine Trop., v. 35 (4), 338-344
Plasmodium falciparum, Onchocerca volvulus, world trends and recent findings of 1974 epidemiologic surveys
- Onchocerca volvulus*
Duke, B. O. L., 1973, Ann. Trop. Med. and Parasitol., v. 67 (1), 95-99
Onchocerca volvulus, infective 'forest' Simulium damnosum feeding on man, nearly 80% of infective larvae escaped from flies, 40% of flies shed all larvae and became negative, considerable number of remaining 60% survived in infective state and were potentially able to transmit the parasite at subsequent blood-meal(s)
- Onchocerca volvulus*
Duke, B. O. L., 1974, Ann. Trop. Med. and Parasitol., v. 68 (2), 241-242
Onchocerca volvulus, chimpanzee (exper.), trichlorophone is no more than microfilaricide and shows no promise as macrofilaricide for West African forest strain
- Onchocerca volvulus*
Duke, B. O. L., 1974, Proc. 1. Internat. Cong. Ecol., 323-328
life cycles of Onchocerca volvulus and its vector, Simulium spp.; ecology of human onchocerciasis, epidemiology, control, distribution, review
- Onchocerca volvulus*
Duke, B. O. L., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 71 [Letter]
Onchocerca volvulus, acute infection in exper. chimpanzee given large intramuscular formulation of levamisole, probable action on adult worms as well as microfilariae
- Onchocerca volvulus*
Duke, B. O. L., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (2), 172 [Letter]
Onchocerca volvulus-infected chimpanzee, mebendazole ineffective treatment when administered orally
- Onchocerca volvulus*
Duke, B. O. L., 1974, Tropenmed. u. Parasitol., v. 25 (1), 84-88
Onchocerca volvulus in chimpanzees, suramin lethal to all immature worm stages, implications for human treatment
- Onchocerca volvulus*
Duke, B. O. L., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (2), 287-288 [Letter]
Onchocerca volvulus, levamisole, no evidence of action on either microfilariae or adult worms at a maximum dosage tolerated by humans
- Onchocerca volvulus*
Duke, B. O. L., 1975, Tropenmed. u. Parasitol., v. 26 (1), 88-97
Onchocerca volvulus, dispersal of nulliparous and parous Simulium damnosum vectors, highest number carrying infective larvae found near banks of breeding rivers: Sudan-savanna zone of Cameroon
- Onchocerca volvulus*
Duke, B. O. L., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 90-91 [Letter]
Onchocerca volvulus in man, speculation that route of entry of microfilariae into eye is along sheaths of ciliary vessels and nerves: Cameroon
- Onchocerca volvulus*
Duke, B. O. L.; and Garner, A., 1975, Tropenmed. u. Parasitol., v. 26 (4), 435-448
Onchocerca volvulus, rabbits (exper.) pre-immunized with live or freeze-killed microfilariae and later challenged by subconjunctival inoculation of live microfilariae, severe ocular pathology resulted with live microfilariae, minimal reaction with killed; possible immunological bearing on human infection

- Onchocerca volvulus*
Duke, B. O. L.; and Garner, A., 1976, Tropenmed. u. Parasitol., v. 27 (1), 3-17
pathologic changes in eyes of rabbits inoculated in posterior chamber with *Onchocerca volvulus* microfilariae, comparisons with normal controls
- Onchocerca volvulus*
Duke, B. O. L.; and Moore, P. J., 1974, Tropenmed. u. Parasitol., v. 25 (2), 153-159
Onchocerca volvulus, Guatemalan strain in chimpanzees, periodicity of concentrations of microfilariae in skin, peak coincided with peak wave of biting density of main *Simulium* vectors in Guatemala, pattern of circadian undulation differs from West African strain
- Onchocerca volvulus*
Duke, B. O. L.; Moore, P. J.; and Vincelette, J., 1975, Tropenmed. u. Parasitol., v. 26 (4), 449-468
Onchocerca volvulus in humans, factors influencing passage of microfilariae into urine (extra fluid intake, pyrexial episodes, dosage with diethylcarbamazine, host activity), probably from reservoir in kidney glomerular capillaries, implications for epidemiologic surveys
- Onchocerca volvulus*
Duke, B. O. L.; Moore, P. J.; and Vincelette, J., 1976, Tropenmed. u. Parasitol., v. 27 (2), 133-144
the population dynamics of *Onchocerca volvulus* microfilariae during treatment with suramin and diethylcarbamazine
- Onchocerca volvulus*
Duke, B. O. L.; Vincelette, J.; and Moore, P. J., 1976, Tropenmed. u. Parasitol., v. 27 (2), 123-132
microfilariae in cerebrospinal fluid of patients heavily infected with *Onchocerca volvulus* and resulting neurological complications, association between these complications and diethylcarbamazine therapy, implications for treatment programs with diethylcarbamazine and/or suramin: northern Cameroon
- Onchocerca volvulus*
Fazen, L. E.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (1), 52-57
epidemiologic study of human microfilaruria in Guatemala, its frequency of occurrence and association with microfilariae in skin and its relationship to presence of subcutaneous nodules
- Onchocerca volvulus*
Fazen, L. E.; et al., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 250-256
Onchocerca volvulus, humans with light infections, single oral dose of diethylcarbamazine, consequent clinical and laboratory changes, distribution of microfilariae, corticosteroid appeared to reduce symptoms of DEC reaction: Guatemala
- Onchocerca volvulus*
Figuerola M., H.; Collins, R. C.; and Kozek, W. J., 1977, Am. J. Trop. Med. and Hyg., v. 26 (1), 75-79
technique for post-feeding transportation and maintenance of *Simulium ochraceum* infected with *Onchocerca volvulus* from area of infection in Guatemala to research laboratories in California
- Onchocerca volvulus*
Frentzel-Beyme, R., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (3), 339-357
Onchocerca volvulus, human, epidemiological survey, parasitological and clinical findings, age and sex specific prevalence rates, relation to rate of blindness: Bong Range, Liberia
- Onchocerca volvulus*
Frentzel-Beyme, R., 1975, Tropenmed. u. Parasitol., v. 26 (1), 70-87
Onchocerca volvulus, extensive incidence survey showing geographic patterns, age and sex differences in infected population of Liberia
- O[onchocerca] volvulus*
Frentzel-Beyme, R. R., 1975, Tropenmed. u. Parasitol., v. 26 (4), 469-488
Onchocerca volvulus, epidemiologic survey of inhabitants of 121 communities over 3-year period for incidence and risk of blindness associated with onchocerciasis infection, importance as public health problem, results show 50% increased risk associated with exposure to parasites: Liberia
- Onchocerca volvulus*
Fuglsang, H.; et al., 1976, Tropenmed. u. Parasitol., v. 27 (3), 365-369
Onchocerca volvulus in humans, variations in concentrations of microfilariae in diagnostic skin snips suggests seasonal variation corresponding to climatic changes or to biting activity of *Simulium* vectors
- Onchocerca volvulus*
Fuglsang, H.; and Anderson, J., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 72-73 [Letter]
Onchocerca volvulus in humans, respiratory distress and collapse in several persons receiving diethylcarbamazine as part of mass therapy trials, possible association with unusual presence of microfilariae in circulation: Cameroon
- Onchocerca volvulus*
Fuglsang, H.; and Anderson, J., 1977, Tropenmed. u. Parasitol., v. 28 (1), 63-67
Onchocerca volvulus in humans, incidence survey using corneoscleral punch-microtiter plate quantitative skin snip technique, concentration of microfilariae in skin near eye used as simple measure of severity of infection and indicator of people at risk in developing eye pathology: Cameroon

- Onchocerca volvulus*
Fuglsang, H.; Anderson, J.; and Marshall, T. F. de C., 1976, *Tropenmed. u. Parasitol.*, v. 27 (3), 355-364
Onchocerca volvulus in humans, comparative survey of natives of rain-forest and savannah areas for presence of head nodules containing adult worms, possible associations between presence of nodules and ocular onchocerciasis: Cameroon
- Onchocerca volvulus*
Garms, R., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (2), 222-231
Simulium damnosum, distribution, infections with *Onchocerca volvulus* in different cytological categories of species complex: Liberia
- Onchocerca volvulus*
Garms, R., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (3), 358-372
Onchocerca volvulus, quantitative aspects of transmission by *Simulium damnosum*, daily and annual cycles of biting densities, age composition and infection rates of vector population, transmission potential: Bong Range, Liberia
- Onchocerca volvulus, illus.*
Garner, A., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (5-6), 1976, 374-377
Onchocerca volvulus, pathology of ocular infections in humans and experimental rabbits, granulomatous lesions resulting from direct microfilarial invasion and inflammatory lesions a probable response to free microfilarial antigens, review of current research
- Onchocerca volvulus*
Garner, A.; and Duke, B. O. L., 1976, *Tropenmed. u. Parasitol.*, v. 27 (1), 19-29
Onchocerca volvulus, rabbits, fundus lesions following inoculation of microfilariae into posterior segment of eye
- Onchocerca volvulus, illus.*
Garner, A.; Duke, B. O. L.; and Anderson, J., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 385-396
Onchocerca volvulus, rabbits, comparison of lesions produced in cornea of eye by microfilariae of forest vs. Sudan-savanna strains from Cameroon, observations confirm greater pathogenicity of Sudan-savanna strain
- Onchocerca volvulus*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, *Medecine et Malad. Infect.*, v. 3 (8-9), 351-353
diagnostic review of human filariasis
- Onchocerca volvulus, illus.*
Gibson, D. W.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (1), 74-87
Onchocerca volvulus, humans with severe onchocercal dermatitis, ultrastructure of microfilariae and host skin tissues before and after diethylcarbamazine treatment: Cameroon
- Onchocerca volvulus*
Gougerot, M. A.; et al., 1975, *Bull. Soc. Path. Exot.*, v. 68 (3), 297-303
elevated IgE, useful indicator of possible human parasitism in absence of allergic type conditions
- Onchocerca volvulus*
Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, *J. Infect. Dis.*, v. 132 (3), 340-352
algorithms in the diagnosis and management of human forms of filariases
- Onchocerca volvulus*
Grunewald, J., 1976, *Tropenmed. u. Parasitol.*, v. 27 (4), 438-454
Onchocerca volvulus, *Simulium damnosum* complex vectors, analysis of physical and hydro-chemical conditions at breeding sites in Upper Volta, Liberia, Cameroon, Tanzania, and Kenya
- Onchocerca volvulus*
Hansen, E. L.; and Hansen, J. W., 1976, *IRCS J. Med. Sc.*, v. 4 (11), 508
Romanomeris culicivox, experimental parasitism of *Simulium damnosum* vectors of *Onchocerca volvulus*, possible use as biological control agent and as laboratory model for developing similar techniques with other merithids
- Onchocerca volvulus*
Hunponu-Wusu, O. O.; and Somorin, A. O., 1977, *J. Trop. Med. and Hyg.*, v. 80 (6), 129-131
Onchocerca volvulus, evaluation of geographical basis of endemicity of human onchocerciasis in Nigeria
- Onchocerca volvulus, illus.*
Iwamoto, I.; Tada, I.; and Wonde, T., 1973, *Nettai Igaku (Trop. Med.)*, v. 15 (1), 36-45
Onchocerca volvulus in humans, epidemiologic survey, clinical manifestations, hetrazan: Ilubabor Province, Ethiopia
- Onchocerca volvulus, illus.*
Kozek, W. J.; and Figueroa Marroquin, H., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 663-678
Onchocerca volvulus, intracellular organisms similar to chlamydiae found within lateral chords of adult worms and larval stages, also in oogonia, oocysts, and developing eggs of females and in microfilariae, significance unknown
- Onchocerca volvulus*
Laemmler, G.; and Herzog, H., 1974, *Tropenmed. u. Parasitol.*, v. 25 (1), 78-83
Litosomoides carinii in *Mastomys natalensis*, suramin showed high suppressive activity and macrofilaricidal effect on various parasite stages from beginning of infection until end of prepatent period but failed to show prophylactic activity when administered prior to infection, results indicate necessity to confirm larvicidal activity of suramin in *Onchocerca volvulus*-infected chimpanzees and should initiate chemoprophylactic use of the drug in small clinical trials in human patients

- Onchocerca volvulus*
Limbos, P., 1971, *Medecine Afrique Noire*, v. 18 (3), 269-271
Onchocerca volvulus, pruritus and skin manifestations in human onchocerciasis, variations between Europeans and Africans, differential diagnosis
- Onchocerca volvulus*
McDougall, A. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (4), 360-361 [Letter]
onchocerciasis and streptocerciasis in persons with leprosy, possible relationships
- Onchocerca volvulus*
MacRae, A. A.; Anderson, R. I.; and Fazen, L. E., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 658-662
Onchocerca volvulus in humans, evaluation of complement fixation for diagnosis; not good index of degree of infection (suppression of titer in persons with large numbers of microfilariae in skin) but useful diagnostic tool in recently acquired infection and suspected cases without microfilariae in skin and no apparent nodules
- Onchocerca volvulus*
Marcoullis, G.; and Graesbeck, R., 1976, *Tropenmed. u. Parasitol.*, v. 27 (3), 314-322
Onchocerca volvulus, antigen extracts, preliminary identification and characterization; cross-reactions in immunodiffusion using other helminth antigens and sera from patients with other parasitic diseases
- Onchocerca volvulus*, *illus.*
Martinez Baez, M., 1976, *Rev. Invest. Salud Pub.*, Mexico, v. 36 (3), 139-150
Onchocerca volvulus microfilariae, extensive morphological description with emphasis on cephalic and caudal areas: Chiapas, Mexico
- Onchocerca volvulus*, *illus.*
Martinez-Palomo, A.; and Martinez-Baez, M., 1977, *J. Parasitol.*, v. 63 (6), 1007-1018
Onchocerca volvulus, microfilariae from patients in Mexico, morphology (light and scanning and transmission electron microscopy) for comparison with African strains
- Onchocerca volvulus*
Merino, F.; and Brand, A., 1977, *Tropenmed. u. Parasitol.*, v. 28 (2), 229-234
Onchocerca volvulus in humans living in endemic area, determinations of immunological profile in comparison with normal controls showed little alteration in immunoglobulin levels, third component of complement or in response of peripheral blood lymphocytes to mitogens
- Onchocerca volvulus*
Meyers, W. M.; and Connor, D. H., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (5-6), 524-525
human *Onchocerca volvulus* and *Dipetalonema streptocerca* in persons with leprosy, altered Mazzotti reactions following administration of diethylcarbamazine, possible immunological implications
- Onchocerca volvulus*, *illus.*
Meyers, W. M.; Neafie, R. C.; and Connor, D. H., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 650-657
human onchocerciasis, 2 autopsy case reports of microfilariae found in kidney, liver, pancreas, lung, and wall of thoracic aorta, associated pathology: Republic of Zaire
- Onchocerca volvulus*
Mougey, Y.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (1), 95-110
5 filaria-vector sets, stochastic models for assessing relation between numbers of microfilariae ingested and numbers of microfilariae passing into hemocoel, limitation phenomenon
- Onchocerca volvulus*
Muller, R. L.; and Denham, D. A., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (1), 8-9 [Demonstration]
filarial spp. larvae, field technique for recovery and preservation of infective larvae from their insect vectors, application to studies of transmission dynamics
- Onchocerca volvulus*
Neppert, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Onchocerca volvulus*
Neppert, J.; and Warns, C. M., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from Ascaris, hookworm, *Onchocerca*, *Dirofilaria immitis*, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination
- Onchocerc[a] volvulus*, *illus.*
Ngu, J. L.; and Blackett, K., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 111-120
Onchocerc[a] volvulus in humans, immunologic studies attempting to delineate role of humoral and cellular immune responses in the heterogeneity of onchocercal lesions
- Onchocerca volvulus*
Noamesi, G. K., 1971, *Ann. Trop. Med. and Parasitol.*, v. 65 (4), 555-565
Simulium damnosum, dry season survival, implications for control of *Onchocerca volvulus*: northern Ghana, West Africa
- Onchocerca volvulus*, *illus.*
Omar, M. S.; and Garms, R., 1977, *Tropenmed. u. Parasitol.*, v. 28 (1), 109-119
Onchocerca volvulus Guatemalan strain, high intakes of microfilariae by *Simulium metallicum* vectors proved lethal to large proportion of flies within 24 hours of taking infected blood, gross histopathological changes
- Onchocerca volvulus*
Onuigbo, W. I. B., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (4), 708-709
Onchocerca volvulus, statistical survey of biopsied nodules in the Igbos of Nigeria

- Onchocerca volvulus*
Picq, J. J.; and Roux, J., 1973, *Medecine Trop.*, v. 33 (5), 451-461
Onchocerca volvulus, human, microfilaruria in relation to age and sex of host, other filarial diseases, geographic region, cutaneous microfilarial densities, albuminuria during suramin treatment, eggs of *Schistosoma haematobium* in urine, and diethylcarbamazine chemotherapy
- Onchocerca volvulus*
Pinon, J. M.; and Gentilini, M., 1973, *Nouv. Presse Med.*, v. 2 (19), 1283-1287
human filariasis, application of cellular immunologic tests (rosette formation, macrophage migration) in diagnosis and comparison with serologic tests (fluorescent antibody, passive hemagglutination, gel diffusion)
- Onchocerca volvulus*
Poux, J.; and Picq, J. J., 1974, *Medecine et Armees*, v. 2 (10), 877-882
Onchocerca volvulus, occurrence of microfilaria during treatment with diethylcarbamazine, possible correlation with infection intensity: Haute-Volta
- Onchocerca volvulus*, illus.
Rassi, E.; et al., 1975, *Bull. Pan Am. Health Organ.*, v. 9 (1), 10-12
Onchocerca volvulus, newly discovered vector of human onchocerciasis in South America *Simulium amazonicum* (thorax muscle tissue, stomach): Toototobi River, Amazonas State, Brazil
- Onchocerca volvulus*, illus.
Rassi, E.; et al., 1977, *Bull. Pan Am. Health Organ.*, v. 11 (1), 41-64
human, incidence survey, new focus discovered *Simulium pintoii*
S. amazonicum
all from Federal Territory of Amazonas, Venezuela
- Onchocerca volvulus*
Ree, G. H., 1977, *Brit. J. Dermat.*, v. 97 (5), 551-554
Onchocerca volvulus, humans, assessment of diethylcarbamazine for therapy, mixed results in clinical trials with 48% cure rate after one course of drug; criteria for cure still uncertain
- Onchocerca volvulus*
Richet, P., 1973, *Medecine Afrique Noire*, v. 20 (11), 899-920
guidelines for mass therapy in human filarial infections, drugs in current use, review
- Onchocerca volvulus*
Ridley, D. S.; and Hedge, E. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Onchocerca volvulus*, illus.
Rougemont, A.; et al., 1974, *Medecine Trop.*, v. 34 (4), 508-522
Onchocerca volvulus, histopathology of cutaneous reactions and microfilarial response in persons treated with diethylcarbamazine or suramine
- Onchocerca volvulus*
Scheiber, P.; et al., 1976, *Bull. World Health Organ.*, v. 53 (4), 472-475
human *Onchocerca volvulus*, membrane filter concentration technique for epidemiologic field studies, comparison with standard techniques showed that new technique resulted in increased incidence of observed prevalence and density of microfilariae
- Onchocerca volvulus*
Scheiber, P.; Braun-Munzinger, R. A.; and Southgate, B. A., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (1), 19 [Demonstration]
Onchocerca volvulus, humans, comparison of the millipore membrane-concentration technique and the standard skin snip technique for use as epidemiologic tools in field surveys: Togo
- Onchocerca volvulus*
Smit, A. M.; and Zuidema, P. J., 1973, *Nederl. Tijdschr. Geneesk.*, v. 117 (33), 1225-1230
Onchocerca volvulus in European travelers to West or Central Africa, presentation mainly as a skin disease, 100 case reports, diethylcarbamazine drug of choice with follow-up of suramine if needed: Netherlands
- Onchocerca volvulus*
Somorin, A. O.; Heiner, D. C.; and Ajugwo, R. E., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (5, part 1), 872-876
Onchocerca volvulus, Nigerian patients, total serum IgE measured by radioimmunoassay, comparison with IgE levels in uninfected and atopic groups in Nigeria and California
- Onchocerca volvulus*
Stuerchler, D.; and Degremont, A., 1976, *Schweiz. Med. Wchnschr.*, v. 106 (20), 682-688
extensive diagnostic and clinical review of filarial parasites frequently encountered by travelers to endemic tropical areas: Switzerland
- Onchocerca volvulus*
Suswillo, R. R.; et al., 1977, *J. Helminth.*, v. 51 (2), 132-134
failure to experimentally infect *Meriones unguiculatus*
- Onchocerca volvulus*
Tada, I.; Iwamoto, I.; and Wonde, T., 1973, *Nettai Igaku (Trop. Med.)*, v. 15 (2), 121-122
Onchocerca volvulus, humans, evaluation of current skin snip method used for diagnosis and recommendation for improvement
- Onchocerca volvulus*
Thompson, B. H., 1976, *Ann. Trop. Med. and Parasitol.*, v. 70 (3), 329-341
Onchocerca volvulus, intervals between blood-meals of *Simulium damnosum* vectors

- Onchocerca volvulus*
Thompson, B. H., 1976, Ann. Trop. Med. and Parasitol., v. 70 (3), 343-354
Onchocerca volvulus, flight range and dispersal of Simulium damnosum vectors: Cameroon rain forests
- Onchocerca volvulus*
Torroella, J. J., 1976, Rev. Invest. Salud Pub., Mexico, v. 36 (3), 167-170
Onchocerca volvulus, report in 1875 by A.J.P. Da Silva Aranzo entitled "Filaria-dermatemica" may be first report of onchocerciasis in Brazil
- Onchocerca volvulus*
Vajime, C. G.; and Dunbar, R. W., 1975, Tropenmed. u. Parasitol., v. 26 (1), 111-138
Onchocerca volvulus, chromosomal studies of black fly vector revealed 8 species rather than just Simulium damnosum previously thought to be the single vector of human onchocerciasis
- Onchocerca volvulus*
Vargas, D. L., 1974, Prensa Med. Mexicana, v. 39 (3-4), 215-219
Onchocerca volvulus, humans, immunologic analysis of serum immunoglobulins, finding that parasite produces heterophilic antigen of A type blood groups
- Onchocerca volvulus*
Vedy, J.; et al., 1975, Medecine et Armees, v. 3 (10), 851-858
Onchocerca volvulus, human bilateral onchocercal papillitis, clinical management, diagnosis, aggravation of condition with diethylcarbamazine treatment: Tchad
- Onchocerca volvulus*
Verin, P., 1974, Medecine Afrique Noire, v. 21 (12), 981-984
human infections, current concepts on epidemiology, immunity
- Onchocerca volvulus*
Weiss, N.; and Degremont, A., 1976, Tropenmed. u. Parasitol., v. 27 (3), 377-384
filariasis in persons returning from endemic areas, comparison immunoserologic diagnostic tests (immunolectrophoresis, indirect fluorescent antibody, indirect hemagglutination, two-dimensional gel diffusion tests) showed that highest sensitivity obtained with immunoelectrophoresis, combined tests gave best results
- Onchocerca volvulus*
White, G. B., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (2), 161-175
Onchocerca volvulus, Loa loa, and Dipetalonema perstans, survey of man-biting species of Chrysops, Culicoides and Simulium as potential vectors of human infection: Ethiopia
- Onchocerca volvulus*
Wilton, D. P.; and Collins, R. C., 1977, Tropenmed. u. Parasitol., v. 28 (3), 384-386
Onchocerca volvulus, comparison study of recovery of parasites from vector black flies by individual dissection and by mass separation; poor results using mass separation technique
- Onchocerca volvulus*
Zielke, E., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 546-547
Onchocerca volvulus, attempted development of larvae in mosquito hosts for purposes of laboratory study; Aedes aegypti selected Liverpool strain supported development with third-stage larvae found in the mosquito abdomen and, in one instance, in the head
- Onchocerca volvulus*
Zielke, E.; Schulz-Key, H.; and Albiez, E. J., 1977, Tropenmed. u. Parasitol., v. 28 (2), 254-257
Onchocerca volvulus, unsuccessful attempts to infect Anopheles gambiae and Aedes aegypti for use in laboratory studies; A. gambiae refractory and larvae developed only to third stage with some abnormalities in A. aegypti
- Onchocerciasis*
Anderson, J.; Fuglsang, H.; and al Zubaidy, A., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 30-31 [Demonstration]
onchocerciasis, human, clinical and pathological features with special reference to 'sowda', distribution of microfilariae: Yemen
- Onchocerciasis*
Bianco, A. E., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 429 [Demonstration]
Ackertia globulosa in Meriones unguiculatus, a new rodent filaria as possible model for studying diagnostic techniques and chemotherapy for onchocerciasis, Haemaphysalis leachi apparently natural vector for Ackertia globulosa
- Onchocerciasis*
Brinkmann, U. K., 1974, Tropenmed. u. Parasitol., v. 25 (2), 160-166
onchocerciasis, human, method for assessing microfilarial densities in skin snips under field conditions
- Onchocerciasis*
Burke, G. J., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 402-405
10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Onchocerciasis*
Diesfeld, H. J.; and Kirsten, C., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 435-438
filariasis, human, diagnosis, new embedding technique employing 'methacrylate' for preparation of antigen (Dipetalonema viteae) to be used in indirect fluorescent antibody test (tested on onchocerciasis sera from Togo), compared with usual frozen-section method
- Onchocerciasis*
Enarson, D. A., 1976, Ann. Trop. Med. and Parasitol., v. 70 (4), 481-482
onchocerciasis, prevalence amongst 4 ethnic groups, appears to be widely distributed disease with significant endemicity in certain areas: Sudan Republic

Onchocerciasis

Enarson, D. A., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (4), 465-468

human onchocerciasis, epidemiologic survey for endemicity, intensity of infection and clinical features: Sudan Republic

Onchocerciasis

Fuglsang, H.; and Anderson, J., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (5-6), 1976, 534 [Letter]

human onchocerciasis, unsuccessful trials using deseril to alleviate side effects of diethylcarbamazine therapy: Cameroon

Onchocerciasis

Granbacka, P., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (1), 150 [Letter]

onchocerciasis, human, discovery of endemic area: southern Tanzania

Onchocerciasis

Hamilton, P. J. S.; et al., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (3), 187-189

human onchocerciasis, observer variations in relation to ocular and general clinical signs, need for standardization for purposes of assessment of control measures: United Cameroon Republic

Onchocerciasis

Imperato, P. J.; and Sow, O., 1971, *Trop. and Geogr. Med.*, v. 23 (4), 385-389

human onchocerciasis, clinical survey of incidence of infection and survey of local beliefs and attitudes concerning transmission, signs and symptoms and blindness resulting from this infection: Senegal River basin areas

Onchocerciasis

Metge, P.; et al., 1974, *Medecine Trop.*, v. 34 (5), 625-632

human ocular onchocerciasis, localization in paracentral temporal area characteristic of chorioretinal involvement and probable point of microfilarial penetration, diagnosis by angiofluorography

Onchocerciasis

Mondet, B.; Pendriez, B.; and Bernadou, J., 1976, *Cahiers O.R.S.T.O.M.*, s. Entom. Med., v. 14 (2), 141-149

onchocerciasis, Simulium vectors, mermithid parasites, potential biological control: West Africa

Onchocerciasis

Nebout, N., 1971, *Medecine Trop.*, v. 31 (2), 229-232

important focus of human onchocerciasis in the southern area of Tchad near Baibokoum

Onchocerciasis

Niel, G.; et al., 1972, *Medecine et Malad. Infect.*, v. 2 (5), 193-202

filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalonema vitae* and *Ascaris suum* as antigens

Onchocerciasis

Onuigbo, W. I. B., 1975, *Brit. J. Clin. Pract.*, v. 29 (11), 290-291

differential diagnosis of nodules of onchocerciasis, frequently misdiagnosed as cancers, benign tumors or lymph nodes: Nigeria

Onchocerciasis

Ree, G. H.; et al., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 542-543

filariasis, onchocerciasis, blood plasma levels of diethylcarbamazine after oral administration to humans, results show that drug is rapidly absorbed from gastrointestinal tract and reaches peak levels at 1-2 hours after administration

Onchocerciasis

Ricosse, J. H.; and Picq, J. J., 1973, *Medicine Afrique Noire*, v. 20 (11), 877-897

human filariasis, clinical indications for treatment, drugs in current use, review

Onchocerciasis

Rodger, F. C., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (2), 225-237

ocular onchocerciasis, human, prevalence, evaluation of effectiveness of Simulium control scheme at Abuja, North Nigeria

Onchocerciasis

Rougemont, A.; et al., 1976, *Bull. World Health Organ.*, v. 54 (4), 403-410

human onchocerciasis, large-scale mass treatment trials (native villagers over 10 years of age) using oral diethylcarbamazine, advantages and disadvantages of program and follow-up studies: Bamako, Mali

Onchocerciasis

Torroella, J. J., 1976, *Rev. Invest. Salud Pub., Mexico*, v. 36 (3), 171-173

human ocular onchocerciasis, possible link between release of serotonin and ocular pathology

Onchocerciasis

Torroella, J. J.; and Portillo, C., 1976, *Rev. Invest. Salud Pub., Mexico*, v. 36 (3), 175-183

human onchocerciasis, past and current research projects, review: Chiapas, Mexico

Onchocerciasis

Vedy, J., 1972, *Medecine Trop.*, v. 32, Special number, 427-433

onchocerciasis, review of causes of human blindness in Tchad

Onchocerciasis

Vedy, J.; and Sirol, J., 1971, *Medecine Trop.*, v. 31 (6), 559-564

youth with sub-acute bilateral edematous papillitis associated with anodular onchocerciasis, macrofilarial treatment with suramin, followed by diethylcarbamazine resulted in degenerative chorioretinopathy probably caused by diethylcarbamazine treatment: Tchad

- Onchocerciasis**
 Vedy, J.; Sirol, J. C.; and Coulm, J., 1972, *Medecine Trop.*, v. 32, Special number, 403-409
 human onchocerciasis infection of retina resulting in white degenerating intra-retinal spots, only young people affected, no functional impairment
- Onchocerciasis**
 Waddy, B. R., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 39-50
 human parasitism and public health problems associated with building of new lakes and dams, review of several water projects with emphasis on Africa
- Onchocerciasis**
 Woodruff, A. W., 1971, *Brit. J. Clin. Pract.*, v. 25 (12), 529-535
 achievements in tropical medicine during past 25 years, control, prophylaxis, treatment
- Onchocerciasis**
 Woodruff, A. W.; et al., 1977, *J. Trop. Med. and Hyg.*, v. 80 (4), 68-73
 human onchocerciasis in savanna and rain-forest areas, comparison of epidemiologic aspects and pathology: Nigeria and Cameroon
- Onchocerciasis**
 Woodruff, A. W.; et al., 1977, *J. Trop. Med. and Hyg.*, v. 80 (4), 74-77
 clinicopathologic picture of human onchocerciasis in Ethiopia as compared to that in Nigeria and Cameroon
- Onchocerciasis**
 Wyatt, G. B., 1971, *Ann. Trop. Med. and Parasitol.*, v. 65 (4), 513-523
 onchocerciasis, human, prevalence, age and sex rates, microfilaremia, subcutaneous nodules, symptoms, skin changes, visual acuity, corneal opacities, hemoglobin genotype, entomology: Ibarapa, Western State, Nigeria
- Oncholaimina DeConinck, 1965**
 Maggenti, A. R., 1976, *Organ. Nematodes (Croll)*, 1-10
 Enoplida
- Oncophora Diesing, 1851, illus.**
 Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (3), 1-27
 Camallanidae
 key
 Syn.: *Piscillania* [sic] Yeh, 1960
- Oncophora albacarensis Baudin-Laurencin, 1972**
 Bussieras, J.; and Baudin-Laurencin, F., 1973, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 26 (4), 13a-19a
Thunnus albacares (organe pylorique): tropical Atlantic
- Ophidascaridinea (Hartwich, 1954, subfam.) Chabaud, 1965**
 Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (2), pp. 1-15
 Angusticaecinae
 synonymy
 key; key to genera
 includes: *Ophidascaris*; *Polydelphis*; *Hexametra*
- Ophidascaris Baylis, 1921**
 Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (2), pp. 1-15
 Ophidascaridinea
 key
- Ophidascaris filaria (Dujardin 1845), illus.**
 Elbihari, S.; and Hussein, M. F., 1973, *J. Wildlife Dis.*, v. 9 (2), 171-173
Ophidascaris filaria in *Python sebae* (stomach), associated granulomatous pre-neoplastic gastric lesions: Sudan
- Ophidascaris najae (Baylis, 1921)**
 Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, *Dobuts. Zasshi, Tokyo*, v. 84 (3), 258-261
 seasonal differences of infection rate, worm burden
Naja naja: West Bengal villages
- Ophidascaris wui Freitas 1967**
 Pinnell, J. L.; and Schmidt, G. D., 1977, *J. Parasitol.*, v. 63 (2), 337-340
Naja naja sputatrix: Komodo Island, Indonesia
- Ophiidracunculus Yamaguti, 1961**
 Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (3), 1-27
 as syn. of *Dracunculus* (Reichard, 1759)
- Orientostrongylus Durette-Desset, 1970**
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
 Heligmonellidae, Nippostrongylinae
- Orientostrongylus dendrogali n. sp., illus.**
 Durette-Desset, M. C.; and Lim, B. L., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 669-675
Dendrogale melanura (intestin grele): Mount Kinabalu, Bundu Tahan Reserve, Sabah, Nord Borneo
- Orientostrongylus krishnansamyi n. sp., illus.**
 Durette-Desset, M. C.; and Lim, B. L., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 669-675
Iomys horsfieldi (intestin grele): Kuala Langat Forest Reserve, Bukit Mandol Aborigine village, Klang, Selangor, Malaisie
- Orihelia n. subgen.**
 Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
 subgen. of *Dipetalonema*; key
 tod of subgen.: *Dipetalonema* (*Orihelia*)
 anticlava (Molin, 1858)

- Orloffia Drozdz*, 1965
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Orneoascaris* Skrjabin, 1916, gen. dub.
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
? as syn. of *Amplificaecum* Baylis, 1920
- Ornithofilaria* sp. Borgarenko, 1960
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 3-6
as syn. of *Sarconema pseudolabiata* nov. sp.
- Ornithofilaria fallisensis* Anderson
Bennett, G. F.; et al., 1975, *J. Wildlife Dis.*, v. 11 (2), 280-289
hematozoa of Anatidae, prevalence survey
Anas rubripes rubripes
Anas platyrhynchos platyrhynchos
Anas crecca carolinensis
Anas acuta
Aix sponsa
Mareca americana
all from Maritime provinces of Canada
- Ornithonema* Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, Ornithostrongylinae
- Ornithostrongylinae* Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae
includes: *Ornithostrongylus* (type genus);
Allintoshius; *Lutznema*; *Ornithonema*; *Oswaldostromylus*
- Ornithostrongylus* Travassos, 1914 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, Ornithostrongylinae
- Ortleppina* Schulz, 1927
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Heliconema* Travassos, 1919
- Ortleppstrongylus* Durette-Desset, 1970
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Molineinae
- Ortoanisakis* (Mozgovoi, 1951, subgen.) Yamaguti, 1961
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Paranisakis* Baylis, 1923
- Oshimaia* Sugimoto, 1934
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Avioserpens* Wehr & Chitwood, 1934
- Osleroides* Orlov, Davtian & Lubimov, 1933
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Vogeloides* Orlov, Davtian & Lubimov in Skrjabin, 1933
- Osleroides* Orlov, Davtian & Lubimov in Skrjabin, 1933
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Metathelazia* Skinner, 1931
- Osleroides* (*Osleroides*) *felis* (Vogel 1928) Orlov, Davtian, and Liubimov 1933
Pence, D. B.; and Stone, J. E., 1977, *J. Parasitol.*, v. 63 (6), 979-991
as syn. of *Vogeloides felis* (Vogel 1928) Davtian 1933
- Oslerus* Hall, 1971 (in part)
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Metathelazia* Skinner, 1931
- Oslerus felis* Vogel 1928
Pence, D. B.; and Stone, J. E., 1977, *J. Parasitol.*, v. 63 (6), 979-991
as syn. of *Vogeloides felis* (Vogel 1928) Davtian 1933
- Ostertagia*
Baines, D. M.; and Colegrave, A. J., 1977, *Vet. Rec.*, v. 100 (11), 217-219
gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales
- Ostertagia*
Bliss, D. H.; and Todd, A. C., 1977, *Vet. Med. and Small Animal Clin.*, v. 72 (10), 1612-1617
milk production in dairy cows' exposed to mixed trichostrongylid larvae, results indicate that greatest milk loss occurs during the first 90 days of lactation, relationship between exposure time and stage of lactation
- Ostertagia*
Boag, B.; and Thomas, R. J., 1975, *Research Vet. Sc.*, v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Ost[ertagia]*
Brunsdon, R. V., 1976, *N. Zealand J. Exper. Agric.*, v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)

- Ostertagia**
Buerger, H. J., 1976, *Vet. Parasitol.*, v. 1 (4), 359-366
Ostertagia, Cooperia, Nematodirus, significantly higher numbers of larvae on herbage samples collected from calf pastures vs. cow pastures, improved control of trichostrongyle infection during late summer and autumn might be achieved by transfer of calves to cow pastures
- Ostertagia**
Buerger, H. J., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 219
incidence on grass from cattle pasture: Niedersachsen
- Ostertagia**
Crowley, J. W., jr.; et al., 1977, *Am. J. Vet. Research*, v. 38 (5), 689-692
lungworms, gastrointestinal parasites, cattle, 3 controlled critical trials, highly effective
- Ostertagia**
Downey, N. E.; and Moore, J. F., 1977, *Vet. Rec.*, v. 101 (24), 487-488
Ostertagia, Cooperia, incidence in calves grazing on pastures which were fertilized with slurry: Ireland
- Ostertagia**
Dewel, D., 1977, *Cahiers Bleus Vet.* (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Ostertagia Ransom**, 1907 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
synonymy
- Ostertagia**
Eckert, J., 1972, *Schweiz. Arch. Tierh.*, v. 114 (12), 652-667
control measures based on limited information available, pasture change and anthelmintic treatment at end of June, review: Switzerland
- Ostertagia**
Ford, G. E., 1976, *Pathophysiol. Parasit. Infect.*, 83-97
trichostrongylid parasitism of ruminants (with emphasis on Ostertagia in cattle), blood pepsinogen estimations: seasonal variation in relation to climate; clinical significance (diagnosis, pathogenesis, response to anthelmintic treatment); parasitological significance; production significance; review
- Ostertagia, illus.**
Goffredo, G.; and Sobrero, R., 1972, *Parassitologia*, v. 14 (1), 143-148
- Ostertagia**
Guarino, C.; and Rivellini, P., 1972, *Atti Soc. Ital. Sc. Vet.*, v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Ostertagia**
Henriksen, Sv. Aa.; et al., 1976, *Vet. Parasitol.*, v. 2 (3), 259-272
gastro-intestinal nematodes, young calves during first grazing season, infection levels, blood findings, body weight gains, comparison of animals grazing same pasture entire season with those moved in early July and between levamisole-treated and untreated animals: Denmark
- Ostertagia**
Qadir, A. N. M. A., 1976, *Indian Vet. J.*, v. 53 (11), 855-858
gastrointestinal nematodes, goats and calves, urea for pasture control of free-living stages
- Ostertagia**
Sewell, M. M. H., 1973, *Vet. Rec.*, v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Ostertagia**
Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Ostertagia**
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Ostertagia**
Todd, A. C.; et al., 1976, *Am. J. Vet. Research*, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Ostertagia**
Tsolov, B.; and Tsanov, G., 1975, *Vet. Sbirka*, v. 73 (9), 20-22
strongylids of sheep, economics of tetramisole treatment, body weight, milk production, not effective against *Trichocephalus*
- Ostertagia**
Wallnoefer, E., 1977, *Wien. Tierarztl. Monatsschr.*, v. 64 (4), 129-131
sheep parasites, Mebenvet, good results when treatment was repeated after 14 days: Austria

- Ostertagia spp.
Whitlock, J. H.; and Georgi, J. R., 1976, Parasitology, v. 72 (3), 207-224
biological controls in mixed trichostrongylid infections (predominantly Haemonchus contortus cayugensis) in sheep, different ecosystems (barn vs. pasture) and different treatment groups, course of infections (erythrocyte loss, fecal egg counts, hematocrit values), "Amaphylactoid 'self-cure'" did not occur in this experiment but something like premunition certainly did."
- Ostertagia spp.
Zeakes, S. J.; et al., 1976, Am. J. Vet. Research, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos-crumbles and nalfalofos boluses
- Ostertagia spp.
Zilinski, J., 1972, Med.-Wet., v. 28 (9), 566-567
parasites, sheep, Nilverm, copper sulfate
- Ostertagia sp.
Bergstrom, R. C.; Maki, L. R.; and Werner, B. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 171-174
Trichostrongylid eggs in cattle or sheep feces, dung beetles (Aphodius spp.; Canthon praticolus) as possible biological control agents, laboratory studies showed decreased eggs in feces when beetles were present
- Ostertagia sp.
Canale, A.; et al., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 306-310
Fasciolis hepatica, calves, light experimental infection alone or in combination with gastrointestinal nematodes, digestive function not impaired
- Ostertagia sp.
Cheljina, S. N., 1977, Vet. Rec., v. 100 (6), 120
Ostertagia sp., adult cattle, diagnosis based on plasma pepsinogen levels, discrepancies, unreliable for diagnosis
- Ostertagia spp.
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs and heifers, efficacy of fenbendazole, thiabendazole and tetranisole compared
- Ostertagia sp.
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 526-532
cattle, morantel tartrate, good results - against Cooperia sp., Ostertagia sp., and Trichostrongylus sp., increased weight gain in treated cattle, field trials: United Kingdom
- Ostertagia spp.
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Vet. Rec., v. 92 (20), 551-554
control of clinical infections of gastrointestinal nematodes and lungworms in calves using morantel/diethylcarbamazine solution, field trials, good results as measured by growth response and clinical symptoms; routine treatment economically sound under conditions of heavy infection
- Ostertagia spp.
Carr, C., 1977, Austral. Vet. J., v. 53 (9), 425-428
nematodes, calves, Ievamisole, efficiency of pour-on formulation, drug trials, good results
- Ostertagia spp.
Dorr, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citirizin spot-on, application on skin, good results
- Ostertagia spp.
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with Ievamisole (oxfendazole showed higher efficacy than Ievamisole against Ostertagia spp., similar efficacy against other species), both drugs increased calves' weight gains
- Ostertagia larvae
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-265
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with Ievamisole
- Ostertagia spp.
Düwel, D.; et al., 1974, Prakt. Tierarztl. v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Ostertagia spp.
El-Abdin, Y. Z.; et al., 1975, Egypt. J. Vet. Sc., v. 12 (1), 31-43
serum constituents and serum enzyme activities, normal and nematode infested Camelus dromedarius: Cairo abattoir
- Ostertagia sp.
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, Vet. Sbirka, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of tetranisole, group and individual applications
- Ostertagia sp.
Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (5), 533-534
Haemonchus contortus, Ostertagia sp., sheep, "spring rise" in fecal egg counts, higher incidence in bred sheep than in unbred sheep, role of contaminated bedding: Maine
- Ostertagia sp.
Gonzalez, H.; and Plaza, J., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 134-137
gastrointestinal nematodes of sheep, comparative therapeutic trials using banminth, phenothiazine, and thiabendazole

- Ostertagia spp.
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Ostertagia [sp.]
Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Ostertagia spp.
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Ostertagia sp. 4th stage
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Ostertagia spp.
McKenna, P. B., 1976, N. Zealand J. Exper. Agric., v. 4 (2), 235-237
post-mortem recovery of Haemonchus contortus, Ostertagia spp., Trichostrongylus axei, sheep, peptic digestion of ovine abomasa unlikely to be of diagnostic value for field-submitted specimens
- Ostertagia sp.
Mutafova, T., 1972, Izvest. Tsentral. Khelminth. Lab., v. 15, 143-150
Haemonchus contortus, Ostertagia sp., Trichostrongylus sp., sheep, seasonal changes in egg-production, spring rise observed irrespective of age or sex of host
- Ostertagia sp.
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Ostertagia sp.
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Ostertagia spp.
Reid, J. F. S., 1976, Vet. Rec., v. 98 (25), 496-499
gastrointestinal nematodes, coccidiosis, diarrhea of sheep, age and seasonal factors: Britain
- Ostertagia spp.
Reid, J. F. S.; Duncan, J. L.; and Bairden, K., 1976, Vet. Rec., v. 98 (21), 426-427
Ostertagia spp., pregnant sheep, efficacy of levamisole against inhibited larvae, good results: Scotland
- Ostertagia spp.
Romero Rodriguez, J.; and Valero Lopez, A., 1975, Rev. Iber. Parasitol., v. 35 (1-2), 147-156
Ostertagia spp., measurements, incidence in sheep
Ovis aries: Granada (Espana)
- Ostertagia sp.
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Ostertagia spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ostertagia type II
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ostertagia spp.
Titchen, D. A.; and Anderson, N., 1977, Austral. Vet. J., v. 53 (8), 369-373
Ostertagia, sheep and cattle, physiopathology of gastritis, secretory changes of parasitized and non-parasitized mucosa, review
- Ostertagia sp.
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Ostertagia spp.
Vlassoff, A., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand
- Ostertagia [sp.]
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Ostertagia sp.
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus equinus: Percy Fyfe Nature Reserve, South Africa

- Ostertagia antipini* Matschulski, 1950
Drozd, J.; and Bylund, G., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 259-260
Alces alces (abomasa): Poland
- Ostertagia circumcincta*
Anderson, N.; Blake, R.; and Titchen, D. A., 1976, *Parasitology*, v. 72 (1), 1-12
Ostertagia circumcincta, sheep, repeated infections, food intake, total acid output of fundic pouches, pH of abomasal contents, plasma pepsinogen levels, effects reversed by thiabendazole treatment, secretory capacity of fundic pouches tested with pharmacologic agents and feeding
- Ostertagia circumcincta*
Anderson, P. J. S.; and Marais, F. S., 1972, *J. South African Vet. Ass.*, v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Ostertagia circumcincta*
Baker, N. F.; and Fisk, R. A., 1977, *Am. J. Vet. Research*, v. 38 (9), 1315-1316
Ostertagia, *Trichostrongylus*, and *Nematodirus*, oxfendazole highly effective against adult stages in sheep
- Ostertagia circumcincta*
Boag, B.; and Thomas, R. J., 1973, *Research Vet. Sc.*, v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Ostertagia circumcincta*
Boag, B.; and Thomas, R. J., 1977, *Research Vet. Sc.*, v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Ostertagia circumcincta*
Campbell, W. C.; and Thomson, B. M., 1973, *Austral. Vet. J.*, v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus colubriformis* proved uninfective even if they had not been frozen
- Ostertagia circumcincta*
Chalmers, K., 1977, *N. Zealand Vet. J.*, v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Ostertagia circumcincta*
Coles, G. C.; and Simpkin, K. G., 1977, *Research Vet. Sc.*, v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Ostertagia circumcincta*
Colglazier, M. L.; et al., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with *Trichuris* spp. and *Moniezia expansa*
- Ostertagia circumcincta*
Coop, R. L.; Sykes, A. R.; and Angus, K. W., 1976, *Parasitology*, v. 73 (2), xxxii [Abstract]
Ostertagia circumcincta, lambs, subclinical infection, effect on food intake and utilization and skeletal growth
- Ostertagia circumcincta*
Coop, R. L.; Sykes, A. R.; and Angus, K. W., 1977, *Research Vet. Sc.*, v. 23 (1), 76-83
Ostertagia circumcincta, sheep (exper.), continuous small infections, clinical observations, body weight and food intake, worm populations and faecal egg count, pathology, concentrations of serum constituents and assessment of their value in diagnosis of sub-clinical ostertagiasis
- Ostertagia circumcincta*
Cornwell, R. L., 1975, *Research Vet. Sc.*, v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Ostertagia circumcincta*
Corticelli, B.; and Lai, M., 1972, *Parassitologia*, v. 14 (1), 95-96
Ovis musimon (abomasa, tenue): Sardegna
- Ostertagia circumcincta*, *illus.*
Daskalov, P., 1974, *Izvest. Tsentral. Khelminth. Lab.*, v. 17, 59-72
Ostertagia circumcincta, *O. trifurcata*, *Teladorsagia davtianii*, males crossed with different morphological types of female *Ostertagia*, no reproductive barriers between them, proposed that they be considered the *Ostertagia circumcincta* complex
- Ostertagia circumcincta*
Downey, N. E., 1977, *Vet. Rec.*, v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole

- Ostertagia circumcincta*
Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Ostertagia circumcincta*
Durham, P. J. K.; and Elliott, D. C., 1976, Vet. Parasitol., v. 2 (2), 157-166
Ostertagia spp., sheep (exper.), development of worm populations and lesions at different larval dose-levels and at different times following infection
- Ostertagia circumcincta*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Ostertagia circumcincta*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Ostertagia circumcincta*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Ostertagia circumcincta*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Ostertagia circumcincta*
Elliott, D. C.; and Durham, P. J. K., 1976, Vet. Parasitol., v. 2 (2), 167-175
Ostertagia spp., challenge infections in previously exposed sheep (exper.), greatly reduced worm numbers, rate of worm development, and pathological effects when compared to infections in previously worm-free animals
- Ostertagia circumcincta*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Ostertagia circumcincta*
Foix, J., 1977, Rev. Med. Vet., Toulouse, v. 128 (8-9), 1111-1119
Dicrocoelium dendriticum, sheep, cambendazole, good results against flukes as well as gastrointestinal strongyles
- Ostertagia circumcincta*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1975, Vet. Parasitol., v. 1 (1), 77-83
Ostertagia circumcincta infection in lambs originating from larvae which survived the winter, pastures with high vs. low residual larval infection, differences in worm burdens between 2 groups but no significant difference in weight gain, implications for worm control
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1975, Vet. Parasitol., v. 1 (1), 85-89
Ostertagia circumcincta, ewes and lambs exposed to experimentally simulated postparturient rise of faecal egg count on pastures, compared to similar group not exposed to postparturient rise, worm burdens and weight gains, implications for control
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1976, J. Comp. Path., v. 86 (2), 269-274
Ostertagia circumcincta, lambs, effect of different levels of larval intake on faecal egg counts and weight gain, no significant acquired resistance demonstrated
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1977, Brit. Vet. J., v. 133 (4), 360-364
Ostertagia circumcincta, lambs, different levels of larval intake to simulate seasonal pasture conditions, effect on fecal egg output, possible grazing management regimes to reduce worm infections
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1977, Brit. Vet. J., v. 133 (6), 559-563
Ostertagia circumcincta, sheep, pasture larval infection and fecal egg output lower in resistant animals (those with previous exper. infection and anthelmintic treatment), possible use of adult animals to graze contaminated paddocks as control measure
- Ostertagia circumcincta*
Gibson, T. E.; and Everett, G., 1977, Research Vet. Sc., v. 23 (2), 191-195
Ostertagia circumcincta, lambs (exper.), contribution of residual pasture larvae and the spring rise as sources of infection, weather conditions as useful tool in predicting patterns of infection and most effective preventive measures
- Ostertagia circumcincta*
Gibson, T. E.; and Parfitt, J. W., 1976, J. Comp. Path., v. 86 (4), 547-555
Ostertagia circumcincta, sheep, phenothiazine and thiabendazole treatment, no influence on host resistance, slight effect on weight gain

- Ostertagia circumcincta
Gibson, T.E.; and Parfitt, J.W., 1977, Vet. Parasitol., v. 3 (1), 61-66
Ostertagia circumcincta, lambs, pattern of faecal egg counts after experimental infection with three different numbers of larvae
- Ostertagia circumcincta
Groeneveld, H. T.; and Reinecke, R. K., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 285-297
non-parametric statistical method for comparing worm burdens in two groups of sheep, application in interpreting results of anthelmintic tests
- Ostertagia circumcincta
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and Oestrus ovis, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Ostertagia circumcincta (Stadelman, 1894) Ransom, 1907
Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
Capreolus capreolus (rennet): southern Bulgaria
- Ostertagia circumcincta
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (abomasum): southwestern Iceland, slaughterhouse in Reykjavik
- Ostertagia circumcincta
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired Dictyocaulus filaria infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Ostertagia circumcincta
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Ostertagia circumcincta
Kerboeuf, D.; and Leimbacher, F., 1977, Rec. Med. Vet., v. 153 (1), 19-25
Ostertagia circumcincta, Trichostrongylus axei, Haemonchus contortus, lambs, use of serum pepsinogen measurements to assess average worm burden in a herd
- Ostertagia circumcincta
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Ostertagia circumcincta
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Ostertagia circumcincta
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Ostertagia circumcincta
Le Jambre, L. F., 1976, Vet. Parasitol., v. 2 (4), Dec., 385-391
Haemonchus contortus, Ostertagia circumcincta, technique for assay of thiabendazole resistance by hatching eggs in solutions of thiabendazole
- Ostertagia circumcincta
Le Jambre, L. F.; and Royal, W. M., 1976, Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (fecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except Nematodirus resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales
- Ostertagia circumcincta
Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1977, Internat. J. Parasitol., v. 7 (6), 473-479
Ostertagia circumcincta, resistance of selected lines to thiabendazole, morantel tartrate, and levamisole following 8 generations of selection in the laboratory and up to 4 years selection in the field: Australia
- Ostertagia circumcincta
Low, W. A., 1976, Canad. Field-Naturalist, v. 90 (2), 189-191
Rangifer tarandus caribou: Tweedsmuir Provincial Park, British Columbia
- Ostertagia circumcincta, illus.
Martinez Gomez, F.; and Hernandez Rodriguez, S., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 295-313
Ostertagia circumcincta, ultrastructure of body wall and intestinal epithelium
- Ostertagia circumcincta
Mulligan, W., 1972, Proc. Nutrition Soc., v. 31 (1), 47-51
gastrointestinal parasites, domestic animals (nat. and exper.) and rabbits (exper.), effects of infections on protein metabolism

- Ostertagia circumcincta*, *illus.*
Munn, E. A., 1977, *Tissue and Cell*, v. 9 (1), 23-34
Haemonchus contortus, structure of intestinal cells, helical polymeric extracellular protein associated with luminal surface for which name contortin is proposed, *Ostertagia circumcincta* also contained contortin-like material but *Nippostrongylus brasiliensis* and *Syphacia obvelata* contained material associated with outer surface of microvilli which was quite distinct from contortin
- Ostertagia circumcincta*
Niec, R.; et al., 1976, *Gac. Vet.*, Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary anthelmintic treatment that could prevent normal buildup of resistance
- Ostertagia circumcincta*
Obergh, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus
Ovis aries
Equus caballus
all from Chile
- Ostertagia circumcincta*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Ostertagia circumcincta*
Parkins, J. J.; Holmes, P. H.; and Bremner, K. C., 1973, *Research Vet. Sc.*, v. 14 (1), 21-28
Ostertagia circumcincta, effects of different infection levels on feed intake, apparent digestibility, and nitrogen balance in Blackface sheep when fed rations of different nitrogen content
- Ostertagia circumcincta*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Ostertagia circumcincta*
Ramajo Martín, V.; and Simon Vicente, F., 1975, *Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C.*, v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Ostertagia circumcincta*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Ostertagia circumcincta*
Reh binder, C.; and Christensson, D., 1977, *Nord. Vet.-Med.*, v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Ostertagia circumcincta*
Reid, J. F. S.; and Armour, J., 1975, *J. Comp. Path.*, v. 85 (1), 163-170
Ostertagia circumcincta, breeding ewes, degree by which immune status varied on a seasonal basis (remained substantially immune to challenge), situation not altered by thiabendazole treatment
- Ostertagia circumcincta*, Stadelmann, 1894, *illus.*
Romero Rodriguez, J.; and Valero Lopez, A., 1975, *Rev. Iber. Parasitol.*, v. 35 (1-2), 147-156
Ostertagia spp., measurements, incidence in sheep
Ovis aries: Granada (España)
- Ostertagia circumcincta*
Rose, J. H., 1971, *Symposia Brit. Soc. Parasitol.*, v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Ostertagia circumcincta*, *illus.*
Rose, J. H., 1973, *Research Vet. Sc.*, v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostrongylus rubidus*, culture from infective larva to adult worm in WAe medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Ostertagia circumcincta*
Rose, J. H., 1976, *Research Vet. Sc.*, v. 21 (1), 76-78
immunization of lambs using metabolites from *Ostertagia circumcincta* grown in vitro and living worms at various stages of development, degree of protection shown by worm burdens, worm lengths, and faecal worm egg counts
- Ostertagia circumcincta*
Rothwell, T. L. W.; et al., 1976, *Vet. Parasitol.*, v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Ostertagia circumcincta* (Stadelmann, 1894)
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Ovis aries: insular Newfoundland

- Ostertagia circumcincta*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Ostertagia circumcincta*
Sykes, A. R.; and Coop, R. L., 1976, Proc. Nutrition Soc., v. 35 (1), 13A-14A [Abstract]
Trichostrongylus colubriformis, *Ostertagia circumcincta*, sheep (exper.), effects of extensive chronic parasitism on food intake and utilization by growing lambs, deposition of nitrogen and fat severely impaired and food intake reduced by 9%
- Ostertagia circumcincta*
Sykes, A. R.; and Coop, R. L., 1977, J. Agric. Sc., v. 88 (3), 671-677
Ostertagia circumcincta larvae, daily dosing of growing sheep, reduction of food intake and utilization resulting from abomasal damage; reduced weight gain
- Ostertagia circumcincta*
Sykes, A. R.; Coop, R. L.; and Angus, K. W., 1977, J. Comp. Path., v. 87 (4), 521-529
Ostertagia circumcincta, chronic sub-clinical parasitism of lambs, significantly reduces skeletal growth
- Ostertagia circumcincta*
Tarazona, J. M., 1975, An. Inst. Nac. Invest. Agrar., s. Hig. y San. Animal (2), 11-17
trichostrongylidosis, ovine, seasonal incidence, 1971-1975: Spain
- Ostertagia circumcincta*
Thomas, R. J.; and Waller, P. J., 1975, Vet. Rec., v. 97 (24), 468-471
Ostertagia circumcincta, lambs naturally infected on pasture from spring to autumn, faecal egg counts, worm counts, serum pepsinogen levels, body weights, correlations; serum pepsinogen estimations as possible diagnostic test
- Ostertagia circumcincta*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ostertagia circumcincta*
Titchen, D. A.; and Anderson, N., 1977, Austral. Vet. J., v. 53 (8), 369-373
Ostertagia, sheep and cattle, physiopathology of gastritis, secretory changes of parasitized and non-parasitized mucosa, review
- Ostertagia circumcincta*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Ostertagia circumcincta* (Stadelmann, 1894)
Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karroo
- Ostertagia circumcincta*
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, Vet. Glasnik, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Ostertagia circumcincta*
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Ostertagia crimensis*
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Ostertagia dikmansii*
Eve, J. H.; and Kellogg, F. E., 1977, J. Wildlife Management, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (*Odocoileus virginianus*) density: southeastern United States
- Ostertagia dikmansii*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites *Odocoileus virginianus*: Hardy County, West Virginia
- Ostertagia dikmansii* Becklund and Walker, 1968
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Cumberland County, New Jersey; Oklahoma
- Ostertagia hamata* Monnig, 1932
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damaliscus dorcas dorcas (abomasum): captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Ostertagia lasensis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia

- Ostertagia lasensis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Ostertagia lasensis*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Ostertagia leptospicularis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Ostertagia leptospicularis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Ostertagia leptospicularis*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Ostertagia leptospicularis* Asadov, 1953
Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
synonymy
Capreolus capreolus (rennet): southern Bulgaria
- Ostertagia leptospicularis*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Ostertagia leptospicularis*
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Ostertagia leptospicularis*
Schweigsut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Ostertagia lyrata*
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Ostertagia lyrata*
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (abomasum): southwestern Iceland, slaughterhouse in Reykjavik
- Ostertagia lyrata*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Ostertagia lyrata*
Randall, R. W.; and Gibbs, H. C., 1977, Am. J. Vet. Research, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Ostertagia lyrata*
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Ostertagia* (*Marshallagia*) *marshalli*, Ransom, 1907, illus.
Romero Rodriguez, J.; and Valero Lopez, A., 1975, Rev. Iber. Parasitol., v. 35 (1-2), 147-156
Ostertagia spp., measurements, incidence in sheep
Ovis aries: Granada (España)
- Ostertagia* (=Marshallagia) *marshalli*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ostertagia mossi*
Eve, J. H.; and Kellogg, F. E., 1977, J. Wildlife Management, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (*Odocoileus virginianus*) density: southeastern United States
- Ostertagia mossi*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (abomasum): Kentucky
- Ostertagia mossi* Dikmans, 1931
Phillips, J. H.; Harley, J. P.; and Rudersdorf, W. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 250
Dama dama (abomasum): western Kentucky

- Ostertagia mossi*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites *Odocoileus virginianus*: Hardy County, West Virginia
- Ostertagia mossi* Dikmans, 1931
Pursglove, S. R., jr., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Cumberland County, New Jersey; Oklahoma
- Ostertagia occidentalis*
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*, v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Ostertagia ostertagi*
van Adrichem, P. W. M.; and Shaw, J. C., 1977, *J. Animal Sc.*, v. 45 (3), 417-422
gastrointestinal nematodes, monozygous twin cattle, comparison of treated and untreated pairs infected naturally on pasture, growth performance, results indicate that the reduced growth may be long-lasting
- Ostertagia ostertagi*
van Adrichem, P. W. M.; and Shaw, J. C., 1977, *J. Animal Sc.*, v. 45 (3), 423-429
gastrointestinal nematodes, effects on growth performance and milk production in cambendazole-treated vs. non-treated monozygous twin cattle naturally infected on pasture during the first lactation period
- Ostertagia ostertagi*
Anderson, N., 1977, *Research Vet. Sc.*, v. 23 (3), 298-302
Ostertagia ostertagi, *Trichostrongylus axei*, cattle, efficacy of levamisole, thiabendazole, and fenbendazole
- Ostertagia ostertagi*
Anderson, P. J. S.; and Marais, F. S., 1975, *J. South African Vet. Ass.*, v. 46 (4), 325-329
adult gastrointestinal nematodes, calves, controlled trials with morantel tartrate
- Ostertagia ostertagi*
Anderson, R. M.; and Michel, J. F., 1977, *Internat. J. Parasitol.*, v. 7 (4), 321-329
Ostertagia ostertagi in calves, survival characteristics of worm populations, two simple mathematical models proposed to describe density-dependent survival
- Ostertagia ostertagi*
Bayly, W. M., 1977, *Southwest. Vet.*, v. 30 (2), 171-175
Ostertagia ostertagi, cattle, Type II ostertagiasis, life cycle, epidemiology, immunity, pathophysiology, treatment, review
- Ostertagia ostertagi*
Benz, G. W.; and Ernst, J. V., 1977, *Am. J. Vet. Research*, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.), albendazole significantly reduced infestations
- Ostertagia ostertagi* (Stiles, 1892)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (abomasum): vicinity of Nowy Targ, Carpathian Mountains
- Ostertagia ostertagi*
Bryan, R. P., 1976, *Austral. Vet. J.*, v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Ostertagia ostertagi*
Campbell, W. C.; and Thomson, B. M., 1973, *Austral. Vet. J.*, v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathing; exsheathed larvae of *Trichostrongylus colubriformis* proved ineffective even if they had not been frozen
- Ostertagia ostertagi*
Canale, A.; et al., 1977, *Folia Vet. Latina*, v. 7 (1), 82-90
Ostertagia ostertagi, calves (exper.), digestive utilization of host diet, results indicate that the diminished digestibility is not sufficient to account for the reduced growth
- Ostertagia ostertagi*
Giordia, H.; et al., 1977, *Am. J. Vet. Research*, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Ostertagia ostertagi*
Coles, G. C.; and Simpkin, K. G., 1977, *Research Vet. Sc.*, v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Ostertagia ostertagi*
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, *Brit. Vet. J.*, v. 129 (6), 518-525
gastrointestinal nematodes and lungworms, calves (exper.), morantel tartrate, efficacy in 5 controlled trials, toxicity experiments demonstrate wide safety margin

- Ostertagia ostertagi*
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxfendazole, drug efficacy
- Ostertagia ostertagi*
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Ostertagia ostertagi*
Dotta, U.; et al., 1977, Clin. Vet., Milano, v. 100 (10), 685-696
Ostertagia ostertagi, calves (exper.), clinical manifestations and physiopathological observations
- Ostertagia ostertagi*
Douvres, F. W.; and Malakatis, G. M., 1977, J. Parasitol., v. 63 (3), 520-529
Ostertagia ostertagi, in vitro cultivation, development from infective larvae to egg-laying adults, 70-minute exsheathing process and two-step roller culture system
- Ostertagia ostertagi*
Downey, N. E., 1976, Vet. Rec., v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Ostertagia ostertagi*
Downey, N. E.; and O'Shea, J., 1977, Vet. Rec., v. 100 (13), 265-266
Dictyocaulus viviparus, *Ostertagia ostertagi*, *Cooperia oncophora*, calves (exper.), low dose levels of levamisole or morantel administered via drinking water, good results
- Ostertagia ostertagi*
Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Ostertagia ostertagi*
Duncan, J. L.; et al., 1976, Vet. Rec., v. 98 (17), 342
Ostertagia ostertagi (inhibited 4th stage larvae), *Trichostrongylus axei*, *Cooperia* spp., fenbendazole, good results, compared with levamisole
- Ostertagia ostertagi*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Ostertagia ostertagi*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Ostertagia ostertagi*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Ostertagia ostertagi*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Ostertagia ostertagi*
Eckert, J.; and Eisenegger, H., 1976, Zentralbl. Vet.-Med., Beihefte (25), 155-160
Dictyocaulus viviparus, cattle, program for management and control, vaccination with Dictol, tetramisole treatment; *Ostertagia ostertagi*, *Cooperia oncophora*, pyrantel tartrate treatment to control concurrent infection limiting weight gain
- Ostertagia ostertagi*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Ostertagia ostertagi*
Fincher, G. T., 1975, J. Parasitol., v. 61 (4), 759-762
numbers of nematode parasites acquired by parasite-free calves grazing contaminated pastures containing dung beetle populations of different densities, worm counts reduced with increased dung beetle populations
- Ostertagia ostertagi*
Forsyth, B. A.; and Shepherd, G. E., 1977, Austral. Vet. J., v. 53 (12), 598-599 [Letter]
Ostertagia ostertagi, cattle, levamisole, fenbendazole, clinical responses and weight gains showed that both drugs were highly effective in controlling Type II ostertagiosis outbreak: Western District of Victoria
- Ostertagia ostertagi* (Stiles, 1907), illus.
Goffredo, G.; and Sobrero, R., 1972, Parassitologia, v. 14 (1), 143-148
Dama dama (intestine): foresta Umbra (promontorio garganico, provincia di Foggia)

- Ostertagia ostertagi
Goldberg, A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Ostertagia ostertagi
Helle, O.; and Tharaldsen, J., 1976, Vet. Parasitol., v. 1 (4), 345-357
Ostertagia ostertagi and Cooperia spp. in young cattle during their first grazing season, free-living stages overwintered in sufficient numbers to cause reduced weight gain and clinical disease in early spring, thiabendazole treatment and move to clean pasture improved weight gain: Norway
- Ostertagia ostertagi, illus.
Henriksen, Sv. Aa., 1975, Medlemsbl. Danske Dyrlaegeforen., v. 58 (21), 833-842
Ostertagia ostertagi, morphology, life cycle, comparison with other nematodes, pathogenicity, clinical aspects, immunity, diagnosis, control, brief review
- Ostertagia ostertagi
Henriksen, Sv. Aa.; and Aagaard, K., 1976, Nord. Vet.-Med., v. 28 (7-8), 392-397
Ostertagia ostertagi, flotation technique and McMaster method, fecal examination
- Ostertagia ostertagi
Henriksen, S. A.; Benthholm, B. R.; and Nielsen-Englyst, A., 1976, Nord. Vet.-Med., v. 28 (4-5) 201-209
gastro-intestinal strongyles, cattle, seasonal distribution on pastures
- Ostertagia ostertagi
Herlich, H., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 52-54
Ostertagia ostertagi, lambs, no patent infections produced after oral inoculation of infective larvae but larvae did exsheath, enter gastric pits, undergo development to adolescent stage, and produce abomasal lesions; adults and mixture of larvae and adolescents recovered from calves and transferred to lambs orally did produce patent infections; mixture of larvae and adolescents recovered from lambs and transferred orally to lambs and calves did not produce patent infections
- Ostertagia ostertagi
Herlich, H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.), oxicendazole, efficacy against adult and larval stages
- Ostertagia ostertagi
Herlich, H., 1977, Am. J. Vet. Research, v. 38 (8), 1247-1248
efficacy of albendazole against gastrointestinal nematodes and Fasciola hepatica in cattle (exper.); comparison of critical vs. controlled tests
- Ostertagia ostertagi
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (abomasum): Kentucky
- Ostertagia ostertagi
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (abomasum): southwestern Iceland, slaughterhouse in Reykjavik
- Ostertagia ostertagi
Jørgensen, R. J.; et al., 1976, Nord. Vet.-Med., v. 28 (4-5), 210-216
Ostertagia ostertagi, cattle, serum pepsinogen analysis
- Ostertagia ostertagi
Jordan, H. E.; et al., 1977, Am. J. Vet. Research, v. 38 (8), 1157-1160
Ostertagia ostertagi, Cooperia, influence on energy efficiency in full-fed vs. maintenance-fed steers with high vs. low worm burdens (low worm burdens did not significantly effect energy utilization; in full-fed steers, energy retention was greater in steers with lower worm burdens; maintenance-fed steers were more heavily parasitized than full-fed steers)
- Ostertagia ostertagii
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired Dictyocaulus filaria infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Ostertagia ostertagi
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Ostertagia ostertagi
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Ostertagia ostertagi
Lancaster, M. B.; and Hong, C., 1977, Vet. Rec., v. 101 (4), 81-82
Ostertagia ostertagi, Cooperia oncophora, calves, variable action of fenbendazole on arrested fourth stage larvae
- Ostertagia ostertagi
Lukovich, R.; et al., 1977, Gac. Vet., Buenos Aires (318), v. 39, 91-95
helminths, cattle, levamisole, results from injectable and dermal application similar
- Ostertagia ostertagi
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix

- Ostertagia ostertagi*
McBain, D. G.; et al., 1977, *Vet. Rec.*, v. 101 (14), 285-286
helminths, calves, fenbendazole in feed blocks
- Ostertagia ostertagi*
Martin, B., 1977, *Vet. Rec.*, v. 101 (1), 11-13
cattle, clinical signs, diagnosis, epidemiology, immunity, control, review
- Ostertagia ostertagi*
Michel, J. F., 1976, *Advances Parasitol.*, v. 14, 355-397
nematode infections in grazing animals, epidemiology and control, extensive review: free-living stages (bionomics, transport); parasitic stages (population regulation, immunity, host differences, arrested development, post-parturient rise); parasitic gastro-enteritis in sheep and cattle; parasitic bronchitis in cattle
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1973, *J. Comp. Path.*, v. 83 (3), 351-356
Ostertagia ostertagi, variation within a population with respect to inhibited development, progeny of worms whose development had been arrested had a greater propensity for interrupted development than progeny of worms whose development had not been arrested, cattle
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1974, *J. Comp. Path.*, v. 84 (4), 539-554
Ostertagia ostertagi, *Cooperia oncophora*, evidence that arrested development is due to action of environmental factors, nature of environmental signals not precisely identified but not simple, changes which they induce in larvae are spontaneously reversed after a time
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1975, *J. Comp. Path.*, v. 85 (1), 133-138
Ostertagia ostertagi, *Cooperia oncophora*, arrested development, effect of temperature at free-living 3rd stage, larvae stored at 4° C. compared with storage at 15° C. and with a change of temperature after 12 weeks
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1976, *J. Comp. Path.*, v. 86 (1), 73-80
Ostertagia ostertagi, resumed development of arrested nematodes in naturally infected yearling cattle
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1976, *J. Comp. Path.*, v. 86 (4), 615-619
Ostertagia ostertagi, development arrested by cold storage, calves infected repeatedly, rate of resumed larval development, worm burden
- Ostertagia ostertagi*
Michel, J. F.; Lancaster, M. B.; and Hong, C., 1976, *Internat. J. Parasitol.*, v. 6 (1), 83-86
Ostertagia ostertagi females, variation in form of vulval flap, effect of genetic factors much smaller than effect of host resistance
- Ostertagia ostertagi*
Musila, V., 1976, *Veterinarstvi*, v. 26 (6), 264
helminths of fallow deer, incidence: Zehusice enclosure
- Ostertagia ostertagi*
Mwegoha, W. M.; and Jørgensen, R. J., 1977, *Acta Vet. Scand.*, v. 18 (3), 293-299
Haemonchus contortus, *Ostertagia ostertagi*, recovery of infective 3rd stage larvae by migration in agar gel, with and without addition of ox bile to agar gel, technique for recovery of larvae from field samples of pasture herbage
- Ostertagia ostertagi*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Ostertagia ostertagi*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus
Sus scrofa
all from Chile
- Ostertagia ostertagi*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Ostertagia ostertagi*
Raynaud, J. P.; and Bouchet, A., 1976, *Ann. Recherches Vet.*, v. 7 (3), 253-280
bovine ostertagiosis, analysis of types and syndromes, total worm counts, post mortem examinations, survey of 74 cattle: France
- Ostertagia ostertagi*
Raynaud, J. P.; and Bouchet, A., 1977, *Rev. Med. Vet.*, Toulouse, v. 128 (2), 155-189
cattle, post-mortem examinations and total worm counts, analysis of types and syndromes: France
- Ostertagia ostertagi*
Rehbinder, C.; and Christensson, D., 1977, *Nord. Vet.-Med.*, v. 29 (12), 556-557
reindeer (abomasum): Sweden

- Ostertagia ostertagi*
Reinecke, R. K., 1972, Onderstepoort J. Vet. Research, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Ostertagia ostertagi*
Ronald, N. C.; Bell, R. R.; and Craig, T. M., 1977, J. Am. Vet. Med. Ass., v. 170 (3), 317-319
gastrointestinal nematodes, calves, levamisole phosphate, effective at one-half recommended dosage
- Ostertagia ostertagi*
Rose, J. H., 1971, Symposia Brit. Soc. Parasitol., v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Ostertagia ostertagi*
Rose, J. H., 1973, Research Vet. Sc., v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostromylus rubidus*, culture from infective larva to adult worm in WAE medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Ostertagia ostertagi*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Ostertagia ostertagi*
Rowlands, D. ap T.; and Berger, J., 1977, J. South African Vet. Ass., v. 48 (2), 85-93
nematodes, calves (exper.), levamisole, dermal application, efficacy against third and fourth larval stages and fifth stage larvae/adult worms, results equivalent to those achieved by orthodox methods of drug administration
- Ostertagia ostertagi*
Searson, J. E.; and Doughty, F. R., 1977, Austral. Vet. J., v. 53 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult *Ostertagia ostertagi* than larval forms): southern New South Wales
- Ostertagia ostertagi*
Selman, I. E.; et al., 1976, Vet. Rec., v. 99 (3), 141-143
Ostertagia ostertagi, outbreaks of ostertagiasis affecting adult beef cattle, clinical, biochemical, haematological, parasitological and pathological findings characteristic of type II ostertagiasis in immature cattle, diagnosis
- Ostertagia ostertagi*
Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Ostertagia ostertagi*
Smith, H. J., 1976, Canad. J. Comp. Med., v. 40 (3), 320-321
mixed *Ostertagia ostertagi* and *Cooperia oncophora* larvae in experimentally infected calves, no significant maturation requirement obtained, maximum infectivity possibly related to incubation temperature
- Ostertagia ostertagi*
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, Am. J. Vet. Research, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Ostertagia ostertagi*, illus.
Stringfellow, F., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 76-81
monospecific and dual species infections of *Ostertagia ostertagi* and *Trichostrongylus axei*, calves, histochemical studies of abomasal tissue
- Ostertagia ostertagi*
Tharaldsen, J., 1976, Acta Vet. Scand., v. 17, Suppl. 61, 1-21
trichostrongylid infections, calves, survival of larvae on pasture, occurrence of larvae not influenced by artificial irrigation; treatment with thiabendazole did not effectively control infection due to overwintering larvae, neither improved weight gain nor reduced egg production: Norway
- Ostertagia ostertagi*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Ostertagia ostertagi*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Ostertagia ostertagi*
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole

- Ostertagia ostertagi*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Ostertagia ostertagi*
Wikerhauser, T.; et al., 1974, Acta Parasitol. Jugoslavica, v. 5 (2), 79-81
trichostrongylids, cattle, fenbendazole compared with thiabendazole, good results from both
- Ostertagia ostertagi*
Williams, J. C.; et al., 1977, Vet. Rec., v. 101 (24), 484-486
Ostertagia ostertagi, cattle, albendazole, good results against inhibited fourth stage larvae as well as developing stages and adults, high efficacy against *Haemonchus* and *Cooperia* spp., no signs of toxicity
- Ostertagia ostertagi*
Williams, J. C.; and Bilkovich, F. R., 1973, Am. J. Vet. Research, v. 34 (10), 1337-1344
Ostertagia ostertagi, distribution of infective larvae on pasture herbage, rainfall and temperature: Louisiana
- Ostertagia ostertagi*
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Ostertagia ostertagi*
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Ostertagia ostertagi*
Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmintac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Ostertagia pinnata*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Ostertagia pinnata*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Ostertagia pinnata*
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (abomaso): Sardegna
- Ostertagia pinnata*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Ostertagia* (*Muflonagia*) *podjapolskyi*, Schulz, Andreeva and Kadezii, 1954, illus.
Romero Rodriguez, J.; and Valero Lopez, A., 1975, Rev. Iber. Parasitol., v. 35 (1-2), 147-156
Ostertagia spp., measurements, incidence in sheep
Ovis aries: Granada (Espana)
- Ostertagia thalae* n. sp., illus.
Troncy, P. M.; and Graber, M., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 221-224
Alcelaphus buselaphus
Hippotragus equinus
(abomasum of all): all from Republique Centrafricaine
- Ostertagia trifurcata*
Anderson, P. J. S.; and Marais, F. S., 1972, J. South African Vet. Ass., v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Ostertagia trifurcata*
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, *Trichostrongylus*, and *Nematodirus*, oxfendazole highly effective against adult stages in sheep
- Ostertagia trifurcata*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Ostertagia trifurcata*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Ostertagia trifurcata*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand

- Ostertagia trifurcata*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Ostertagia trifurcata*
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (abomaso): Sardegna
- Ostertagia trifurcata*, illus.
Daskalov, P., 1974, Izvest. Tsentral. Khel'mint. Lab., v. 17, 59-72
Ostertagia circumcincta, *O. trifurcata*, *Teladorsagia davtiani*, males crossed with different morphological types of female *Ostertagia*, no reproductive barriers between them, proposed that they be considered the *Ostertagia circumcincta* complex
- Ostertagia trifurcata*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Ostertagia trifurcata*
Durham, P. J. K.; and Elliott, D. C., 1976, Vet. Parasitol., v. 2 (2), 157-166
Ostertagia spp., sheep (exper.), development of worm populations and lesions at different larval dose-levels and at different times following infection
- Ostertagia trifurcata*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Ostertagia trifurcata*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Ostertagia trifurcata*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Ostertagia trifurcata*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Ostertagia trifurcata*
Elliott, D. C.; and Durham, P. J. K., 1976, Vet. Parasitol., v. 2 (2), 167-175
Ostertagia spp., challenge infections in previously exposed sheep (exper.), greatly reduced worm numbers, rate of worm development, and pathological effects when compared to infections in previously worm-free animals
- Ostertagia trifurcata*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Ostertagia trifurcata*
Foix, J., 1977, Rev. Med. Vet., Toulouse, v. 128 (8-9), 1111-1119
Dicrocoelium dendriticum, sheep, cambendazole, good results against flukes as well as gastrointestinal strongyles
- Ostertagia trifurcata*
Groeneveld, H. T.; and Reinecke, R. K., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 285-297
non-parametric statistical method for comparing worm burdens in two groups of sheep, application in interpreting results of anthelmintic tests
- Ostertagia trifurcata*
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (abomasum): southwestern Iceland, slaughterhouse in Reykjavik
- Ostertagia trifurcata*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Ostertagia trifurcata*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Ostertagia trifurcata*
Low, W. A., 1976, Canad. Field-Naturalist, v. 90 (2), 189-191
Rangifer tarandus caribou: Tweedsmuir Provincial Park, British Columbia
- Ostertagia trifurcata*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus
Ovis aries
all from Chile

- Ostertagia trifurcata*
Oberg, C.; and Valenzuela, G., 1976, Bol. Chileno Parasitol., v. 31 (3-4), 85-86
differentiation from *Teladorsagia davtiani* by examination of genital cone
- Ostertagia trifurcata*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Ostertagia trifurcata*
Ramajo Martin, V.; and Simon Vicente, F., 1975, Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C., v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Ostertagia trifurcata*
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Ostertagia trifurcata*, Ransom, 1907, illus.
Romero Rodriguez, J.; and Valero Lopez, A., 1975, Rev. Iber. Parasitol., v. 35 (1-2), 147-156
Ostertagia spp., measurements, incidence in sheep
Ovis aries: Granada (España)
- Ostertagia trifurcata*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Ostertagia Dikov*, 1963
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae*
- Ostertagiasis*
Clements, O.; Hamilton, A. F.; and Redahan, E., 1977, Vet. Rec., v. 100 (9), 187 [Letter]
ostertagiasis, calves, importance of serum pepsinogen levels in diagnosis
- Ostertagiella Andreeva*, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Ostertagia* Ransom, 1907
- [*Ostertagiella*] *ostertagielly*
Nurtazin, A. T., 1975, Vestnik Sel'skokhoz. Nauki Kazakhstana (4), 84-86
sheep, trichostrongyles, anthelmintic efficacy of banminth, good results
- Ostertagiinae* (Skrjabin et Schulz, 1937, tribu) Sarwar, 1956
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae
includes: *Ostertagia* (type genus); *Aptergia*; *Bergheia*; *Camelostrongylus*; *Capreolagia*; *Grosspiculaglia*; *Longistrongylus*; *Marshallagia*; *Mazamostrongylus*; *Muflonagia*; *Orloffia*; *Ostertagia*; *Ostertamia*; *Paramecistocirrus*; *Rinadia*; *Sarwaria*; *Skrjabina*; *Spiculoptergia*; *Teladorsagia*
- Ostertamia* Dikov, 1963
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae*
- Oswaldocruzia* Travassos, 1917
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Oswaldocruzia* sp.
Koller, R. L.; and Gaudin, A. J., 1977, South-west. Nat., v. 21 (4), 503-509
helminth recovery at 2 sites with diverse climates, statistical analysis indicates correlations between incidence and/or intensity of infection and host species, locality, and sex and size of host
Hyla regilla
Bufo boreas
all from Los Angeles County, California (Malibu Creek; Big Tujunga Wash)
- Oswaldocruzia bialata* (Molin, 1860)
Plasota, K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 47-60
helminths of frogs, comparison of aquatic and terrestrial hosts, relation of parasite fauna to environment, food supplies and food habits, host life cycle, temperature, rainfall, season, age and sex of host, competition between species of parasite, localization within host
Rana esculenta
R. terrestris
all from Kampinos National Park, Poland
- Oswaldocruzia filiformis* (Goeze, 1782) Travassos, 1917
Hristovski, N. D., 1975, Acta Parasitol. Iugoslavica, v. 6 (1), 3-5
Rana graeca: Bitola district, Macedonia, Yugoslavia
- Oswaldocruzia filiformis* (Goeze, 1782)
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Oswaldocruzia gansi* sp. nov., illus.
Crusz, H.; and Ching, C. C., 1975, Ann. Parasitol., v. 50 (3), 339-349
Rhinophis drummondhayi: above Watawala railway station; Namunukula area
R. philippinus: Pallatenne near Gammaduwa
Uropeltis melanogaster: hills above Kandy (small intestine of all)

- Oswaldocruzia goezei* Skrjabin et Schulz, 1952
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Pelobates fuscus
Rana terrestris
all from Samara river valley, Ukrainian SSR
- Oswaldocruzia goezei* Skrjabin et Schulz, 1952
Markov, G. S.; and Mozgovoi, A. A., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 91-96
low level of helminth infection in *Vipera berus* influenced by temperature, humidity and peculiarities of its geographic distribution and biotic origin
Vipera berus (small intestine, stomach): Karelian ASSR
- Oswaldocruzia goezei* (Skrjabin et Suljc 1952), illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Rana ridibunda
R. esculenta
R. temporaria
R. agilis
Bufo bufo
B. viridis
Bombina bombina
B. variegata
Hyla arborea
all from Yugoslavia
- Oswaldocruzia goezei* Skrjabin et Schulz, 1952, illus.
Rozman, M., 1971, Acta Parasitol. Iugoslavica, v. 2 (2), 67-77
description
synonymy
Rana esculenta (Tanko crijevo): environs of Novi Sad, Yugoslavia
- Oswaldocruzia leidy* Travassos, 1917
Ernst, E. M.; and Ernst, C. H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 176-178
Terrapene carolina: College Park, Maryland
- Oswaldocruzia mazzai* Travassos 1934
Dyer, W. G.; and Altig, R., 1977, Herpetologica, v. 33 (3), 293-296
Bufo typhonius
Eleutherodactylus altamazonicus
Leptodactylus mystaceus
L. pentadactylus
(large intestine of all): all from Santa Cecilia, Napo Province, Ecuador
- Oswaldofilaria belemensis* Bain et Sulahian, 1975, illus.
Bain, O.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (2), 209-221
Oswaldofilaria 3 spp., morphology and development in mosquitoes
Aedes aegypti (exper.)
- Oswaldofilaria petersi* Bain et Sulahian, 1975, illus.
Bain, O.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (2), 209-221
Oswaldofilaria 3 spp., morphology and development in mosquitoes
Culex fatigans (exper.)
C. pipiens (exper.)
- Oswaldofilaria spinosa* Bain et Sulahian, 1975, illus.
Bain, O.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (2), 209-221
Oswaldofilaria 3 spp., morphology and development in mosquitoes
Culex fatigans (exper.) (abdomen)
C. pipiens (exper.) (abdomen, thorax)
- Oswaldonema* Travassos, 1927
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Viannaiinae
- Oswaldostrongylus* Lent et Freitas, 1934
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Ornithostrongylinae
- Otophocaenus*
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus* Leuckart 1848
- Otophocaenus oserskoi* Skrjabin 1942
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden 1870) n. comb.
- Otostrongylus circumlitus* Railliet
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus
Phoca vitulina
(lung of all): all from European waters
- Otostrongylus circumlitus* (Railliet, 1899) Braun, 1933
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (lungs): Sakhalin Bay
- Oxysomatium* sp.
Dyer, W. G.; and Brandon, R. A., 1973, Tr. Illinois Acad. Sc., v. 66 (1-2), 23-29
Eurycea lucifuga (large intestine): Equality Cave, southwest of Equality, Saline County, Illinois
- Oxysomatium* sp.
Plasota, K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 47-60
helminths of frogs, comparison of aquatic and terrestrial hosts, relation of parasite fauna to environment, food supplies and food habits, host life cycle, temperature, rainfall, season, age and sex of host, competition between species of parasite, localization within host
Rana esculenta (intestine): Kampinos National Park, Poland
- Oxysomatium brevicaudatum* (Zeder, 1800), illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Bufo bufo
Rana temporaria
(straznji dio tankog crijeva of all): all from Yugoslavia

- Oxysomatium brevicaudatum* (Zeder, 1800)
Plasota, K., 1969, *Acta Parasitol. Polon.*,
v. 16 (1-19), 1968-1969, 47-60
helminths of frogs, comparison of aquatic
and terrestrial hosts, relation of parasite
fauna to environment, food supplies and food
habits, host life cycle, temperature, rain-
fall, season, age and sex of host, competi-
tion between species of parasite, localiza-
tion within host
Rana esculenta (intestine): Kampinos
National Park, Poland
- Oxysomatium itzocanensis* (Bravo, 1943) Skrjabin,
1961
Caballero Deloya, J., 1974, *An. Inst. Biol.,
Univ. Nac. Mexico, s. Zool.*, v. 45 (1), 45-50
Syn.: *Aplectana itzocanensis* Bravo, 1943
Bufo horribilis (intestino, cloaca): Laguna
de Catemaco, Veracruz, Mexico
- Oxysomatium longicaudata*
Lank, D. R., jr., 1971, *Proc. Indiana Acad.
Sc.*, v. 81 (2), 359-364
Rana catesbeiana: Indiana
- Oxyspirura* Drasch in Stossich, 1897
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
Oxyspirurinae
key to subgenera
includes: *Molinospirura*; *Cramispirura*;
Barusispirura n. subg.; *Caballeroispirura*;
Oxyspirura; *Hamulofilaria*
- Oxyspirura*, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
subgen. of *Oxyspirura*
key
Syn.: *Yorkeispirura* subgen. of *Oxyspirura*
- Oxyspirura petrowi*
Cooper, C. L.; Troutman, E. L.; and Crites,
J. L., 1973, *Ohio J. Sc.*, v. 73 (6), 376-380
Molothrus a. ater (under nictitating mem-
brane of eye): Ottawa county, Ohio
- Oxyspirura* (*Oxyspirura*) *petrowi* Skrjabin, 1929
Pence, D. B., 1975, *Proc. Helminth. Soc. Wash-
ington*, v. 42 (2), 181-183
checklist of previously reported North Ameri-
can bird hosts
Callipepla squamata: Presidio County, Texas
Cyrtonyx montezumae: Davis Mountains near
Fort Davis, Texas
(eyes of all)
- Oxyspirura pusillae* Wehr and Hwang, 1957
Kinsella, J. M., 1974, *Proc. Helminth. Soc.
Washington*, v. 41 (2), 127-130
Aphelocoma c. coerulescens (under nictitat-
ing membrane): Florida
- Oxyspirura* (*Barusispirura*) *rodriguesi* Barus,
1968 (tod of subgen.)
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
- Oxyspirurinae (Skrjabin, 1916, fam.) Yamaguti,
1961
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
Thelaziidae
key
includes: *Oxyspirura*
- Oxyures
Düewel, D., 1977, *Cahiers Bleus Vet.* (26),
201-215
fenbendazole, efficacy against nematodes in
various animals, useful as broad spectrum
anthelmintic; mechanism of action, pharmaco-
kinetics, metabolism, toxicology
- Oxyures
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26),
216-230
fenbendazole (available in 5 forms), ef-
ficacy against nematodes in various animals,
well tolerated with no apparent effects on
fertility or fetus, extensive summary of re-
sults to date
- Oxyuriasis
Angate, Y.; et al., 1974, *Medecine Afrique
Noire*, v. 21 (1), 61-65
symptoms of acute abdomen resulting from
human intestinal parasites, medical and sur-
gical care, case reports: Abidjan, Ivory
Coast
- Oxyuriasis
Diop, B.; and Bao, O., 1974, *Medecine Afrique
Noire*, v. 21 (1), 31-40
human intestinal helminths, clinical indi-
cations for treatment, suggested dosage,
efficacy, tolerances, possible toxicities
- Oxyuriasis
Diouf, A. B.; et al., 1975, *Medecine Afrique
Noire*, v. 22 (6), 453-460
human helminthiasis, statistics of 103
surgical parasitic cases over 10-year
period
- Oxyuriasis
Levin, M. B., 1969, *Pediatrics, Am. Acad.
Pediat.*, v. 44 (1), 125-126
oxyuriasis in young females, pain and rest-
lessness during night time sleep attributed
to migration of female pinworm into vaginal
area from anal area in order to deposit eggs
- Oxyuriasis
Nitzulescu, V.; and Cherman, I., 1974, *Pedi-
atria, Bucuresti*, v. 23 (3), 271-272
mixed human infections of ascariasis and
oxyuriasis, combantrin an effective anthel-
mintic, limited clinical trial
- Oxyuriasis
Privitera, P.; and Privitera, U., 1972, *Minerva
Chir.*, v. 27 (16), 899-906
human oxyuriasis resulting in appendicitis,
case reports, clinical management: Italy

- Oxyuriasis**
 Tarabini Castellani, G., 1970, *Minerva Gastroenterol.*, v. 16 (1), 45-49
 human intestinal parasites, clinical trials using FI 6518 with and without thymol, acceptable drug for mass therapy: Somalia
- Oxyurida**
 Chabaud, A. G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott)(1), 6-17
 Secernentea
 key
 includes: Oxyuroidea
- Oxyuridae, illus.**
 Toft, J. D. II; Schmidt, R. E.; and De Paoli, A., 1976, *J. Med. Primatol.*, v. 5 (6), 360-364
 Oxyuridae in Pan troglodytes, multiple intestinal polyps, histologic characteristics
- Oxyuridae [sp.]**
 Thornton, J. E.; Bell, R. R.; and Reardon, M. J., 1974, *J. Wildlife Dis.*, v. 10 (3), 232-236
 Canis latrans (stomach): Nueces County, Texas
- Oxyurid, larvae**
 Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
 Equus burchelli (rectum): Kenya
- Oxyurids**
 Petter, A. J.; and Douglass, J. F., 1976, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (389), *Zool.* (271), 731-768
 comparative study of populations from Gopherus spp. in 3 locations of North America and in other regions of the world (frequency, distribution, variation, speciation)
- Oxyuris**
 Blanchard, J., 1971, *Prat. Vet. Equine*, v. 3 (1), 29-31
 gastrointestinal nematodes in race horses, clinical trials with equigard, good immediate results especially with ascarids; rapid reinfection rates (2 months after therapy), suggests contaminated paddocks; recommendations therefore for program of routine treatment: France
- Oxyuris**
 Fagot, G.; Clery, P.; and Pascal, D., 1977, *Prat. Vet. Equine*, v. 9 (3), 169-170
 ascarids, strongyles, Oxyuris in race horses, clinical trials with strongid-P most successful in eradicating strongyles: France
- Oxyuris**
 Farag, H. H.; Youssef, A. F.; and Omran, L. A., 1977, *J. Pharm. Sc.*, v. 66 (3), 423-425
 Oxyuris, rats, 2-acetylrimino-3-[(N-phenylcarbamoyl)methyl]-2,3,4,5-tetrahydrothiazole
- Oxyuris**
 Reinecke, R. K.; and le Roux, D. J., 1972, *J. South African Vet. Ass.*, v. 43 (3), 287-294
 adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Oxyuris**
 Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
 anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Oxyuris**
 Ugrinovic, N.; et al., 1972, *Med. Casop.*, v. 7 (1-2), 50-55
 intestinal complications in children infected with Ascaris or Oxyuris, review of hospital cases: Yugoslavia
- Oxyuris sp., iflus.**
 Petter, A. J.; and Douglass, J. F., 1976, *Bull. Mus. National Hist. Nat., Paris*, 3. s. (389), *Zool.* (271), 731-768
 dimensions
 Gopherus polyphemus (colon): sud de Lake Placid, comte de Highlands, Floride
- Oxyuris equi**
 Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
 gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Oxyuris equi**
 Drudge, J. H.; and Lyons, E. T., 1977, *Am. J. Vet. Research*, v. 38 (10), 1581-1586
 internal parasites, horses, antiparasitic drugs, methods of evaluation, critical tests, controlled tests and clinical trials, review of methods and criteria
- Oxyuris equi**
 Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, *Am. J. Vet. Research*, v. 36 (4), Part 1, 435-439
 cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Oxyuris equi**
 Furmaga, S.; Gundlach, J. L.; and Patyra, J., 1976, *Med. Wet.*, v. 32 (12), 734-737
 roundworms, horses, fenbendazol and cambendazol very effective
- Oxyuris equi**
 Greve, J. H.; and Paul, J. W., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (12), 1737-1740, 1742
 nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Oxyuris equi**
 Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 128-135
 internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis

- Oxyuris equi
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, Am. J. Vet. Research, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and Gastrophilus (inactive against latter)
- Oxyuris equi
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 721-723
helminths and bots in horses, thiabendazole and trichlorfon sequentially administered via stomach tube, critical testing, drug efficacies, good results
- Oxyuris equi
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Oxyuris equi
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Oxyuris equi
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Oxyuris equi
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Oxyuris equi
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Oxyuris equi
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Oxyuris equi
Smith, H. J., 1976, Canad. J. Comp. Med., v. 40(4), 327-333
strongyles, Oxyuris equi, Parascaris equorum, naturally-infected ponies, thiabendazole, results support regular repeated treatment
- Oxyuris equi
Stretton, A. O. W., 1976, J. Exper. Biol., London, v. 64 (3), 773-788
Ascaris lumbricoides, anatomy of muscle cells and their neuromuscular contacts, development of musculature from larval to adult forms, brief comparison with Oxyuris equi
- Oxyuris equi
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Oxyuris vermicularis
Bos, P., 1967, Ceskoslov. Pediat., v. 22 (9), 857-858
Oxyuris vermicularis, drug reaction and coma in child taking behavior modification drugs and then treated for oxyuriasis with helmirazin, clinical case report: Czechoslovakia
- Oxyuris vermicularis, illus.
Czorniuk, A.; and Stopinska-Gluszak, U., 1975, Patol. Polska, v. 26 (2), 299-302
Oxyuris vermicularis, woman with symptoms of abdominal tumor, which when removed surgically was inflamed oviduct with ova of pinworms in its wall and lumen, clinical case report: Warsaw, Poland
- Oxyuroidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Oxyurida
- Oxyuroidea [sp.]
Coggins, J. R., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
parasitic fauna, effect of host diet and habitat
Quiscalus quiscula
Agelaius phoeniceus
all from Kellogg Bird Sanctuary, Michigan
- Ozolaimus megatyphlon (Rudolphi, 1819) Dujardin, 1845
Acholonu, A. D., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 106-116
synonymy
Ameiva exsul (small intestine): San German, Puerto Rico

- Panagrolaimus wichmanni* Ruhm, 1955, illus.
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972,
Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera
beetles, possible importance in biological
control
Scarabaeus sacer: Abu-Rawash, Cairo, Egypt
- Pancreatonema torriensis*
McVicar, A. H., 1977, *J. Helminth.*, v. 51 (1),
11-21
intestinal helminths of *Raja naevus*, inci-
dence, intensity, pattern of infection with
host age and sex, geographical differences
in composition of parasite burden
Raja naevus (pancreatic duct): Loch Ewe;
off Aberdeen
- Papilloslerus Khera*, 1944
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
? as syn. of *Metathelazia* Skinner, 1931
- Parabronema Baylis*, 1921, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
Parabronematinae
key
Syn.: *Squamanema* van Thiel, 1925
- Parabronema longispiculatum* n. sp., illus.
Graber, M., 1975, *Rev. Elevage et Med. Vet.
Pays Trop.*, n. s., v. 28 (4), 473-479
[given as *Parabronema rhodesiense longispicu-
latum* on pp. 474, 475]
Loxodonta africana (stomach): N'Djamena
(Fort-Lamy), Chad
- Parabronema rhodesiense longispiculatum*, illus.
Graber, M., 1975, *Rev. Elevage et Med. Vet.
Pays Trop.*, n. s., v. 28 (4), 473-479
[given as *Parabronema longispiculatum* n. sp.
in title, and on pp. 473, 474, 477, 478]
- Parabronema skrjabini* Rassowska, 1924
Basson, P. A.; et al., 1970, *Onderstepoort J.
Vet. Research*, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus
caffer*, some pathological findings, age of
host
Syncerus caffer (abomasum): Kruger National
Park
- Parabronematinae *Skrjabin*, 1941
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
Habronematidae
key; key to genera
includes: *Okapinema*; *Parabronema*
- Paracamallanus Yorke & Maplestone*, 1928, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
Camallanidae
key
Syn.: *Neocamallanus* Ali, 1957
- Paracamallanus senegalensis* Vassiliades, 1970
Vassiliades, G., 1972, *Bull. Inst. Fond. Af-
rique Noire*, s. A, v. 34 (3), 529-533
Clarias senegalensis: Sangalkam, Senegal
- Paracanthocheilus* Kreis, 1952
Hartwich, G., 1974, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (2), pp. 1-15
? as syn. of *Terranova* Leiper & Atkinson,
1914
- Paracooperia Travassos*, 1935
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, *Cooperiinae*
synonymy
- Paracooperia Travassos*, 1935
Khalil, L. F.; and Gibbons, L. M., 1975, *J.
Helminth.*, v. 49 (4), 271-279
"no justification for the synonymy of the
genus *Gazellostrongylus* with *Paracooperia*."
- Paracooperia lerouxi* Yeh
Pester, F. R. N.; and Laurence, B. R., 1974,
J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Paracooperia matoffi*
Chauhan, P. P. S.; and Pande, B. P., 1972,
Indian J. Animal Sc., v. 42 (11), 919-929
as syn. of *P. nodulosa*
- Paracooperia nodulosa*, illus.
Chauhan, P. P. S.; and Pande, B. P., 1972,
Indian J. Animal Sc., v. 42 (11), 919-929
Paracooperia nodulosa, third- and fourth-
stage larvae (future males and females)
described, preadult and adult males and fe-
males described, morphological variations,
buffalo calves
Syn.: *P. matoffi*
- Paracooperia nodulosa*, illus.
Chauhan, P. P. S.; and Pande, B. P., 1972,
Indian J. Animal Sc., v. 42 (11), 930-934
Paracooperia nodulosa, buffalo calves,
histopathology of nodules in host small
intestine
- Paracooperia nodulosa*, illus.
Chauhan, P. P. S.; and Pande, B. P., 1972,
Indian J. Animal Sc., v. 42 (12), 1033-1038
Paracooperia nodulosa, buffalo calves, sea-
sonal incidence of numbers and types of
nodules, numbers of free worms, and numbers
of juvenile worms correlated, development
and description of pre-parasitic stages
Antilope cervicapra (wall of small intes-
tine)
buffaloes (small intestine)
- Paracooperia raphiceri* Ortler
Pester, F. R. N.; and Laurence, B. R., 1974,
J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya

- Paracooperia serrata* (Monnig, 1931) Travassos, 1935, *illus.*
Khalil, L. F.; and Gibbons, L. M., 1975, *J. Helminth.*, v. 49 (4), 271-279
redescription
Gazella thomsoni (small intestine): Kenya
- Paracooperia serrata* Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Gazella thomsonii (gut): Kenya
- Paracooperia serrata*
Young, E.; et al., 1973, *Research J. National Parks Republic South Africa* (16), 195-198
Antidorcas marsupialis (duodenum): Mountain Zebra National Park near Cradock, Cape Province
- Paracrenosoma combesi* n. sp., *illus.*
Mas-Coma, S., 1977, *Ann. Parasitol.*, v. 52 (4), 447-456
Crocidura russula (poumons): Espagne (Catalogne; Tona; Aiguafreda, Prades)
- Paracuaria Krishna Rao*, 1951, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Paracuaria* sp., larvae
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Adesmia gebleri
A. planidorsis
Pisterotarsa kiritschenkoi
P. kessleri
Allotadzhikistania comata
Stalagmoptera incostata
Somocoelia pinguis
Prosodes bactriana
P. biformis
Trigonoscelis gemmulata
all from Tadzhik SSR [and/or] Uzbek SSR
- Paracuaria macdonaldi* Rao, 1951
Aleksiev, V. M.; and Smetanina, Z. B., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 97-104
Larus crassirostris (muscular stomach, proventriculus): Rimsko-Korsakov islands
- Paracuaria tridentata* (Linstow, 1877)
Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (1), 7-31
synonymy, nematodes of *Larus canus* (esophagus, proventriculus, ventriculus), age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Paracuaria tridentata* (Linstow, 1877)
Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (3), 185-189
Rissa tridactyla (ventriculus): Agdenes area, Norway
- Paracuaria tridentata* (Linstow, 1877)
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 105-124
Larus argentatus
L. canus
L. crassirostris
L. schistisagus
Stercorarius parasiticus
(muscular stomach of all): all from coast of Sea of Okhotsk
- Paracuaria tridentata* (Linstow, 1877)
Bishop, C. A.; and Threlfall, W., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 25-35
Somateria mollissima: insular Newfoundland and/or southern Labrador
- Paracuaria tridentata*
Courtney, C. H.; and Forrester, D. J., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (esophagus, proventriculus): Florida; Louisiana
- Paracuaria tridentata* (Linstow, 1877) Barus, 1967
Keppner, E. J., 1973, *Tr. Am. Micr. Soc.*, v. 92 (2), 288-291
Larus californicus (proventriculus): city dump of Laramie, Wyoming
- Paracuaria tridentata* (Linstow, 1877)
Sergeeva, T. P., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 146-155
Larus genei: Azov Sea
L. ridibundus: Azov Sea; Tuva
L. argentatus: Azov Sea
L. minutus: Azov Sea
Sterna paradisaea: Yenisei
Larus ichthyætus: Tuva
- Paradujardinia* Travassos, 1933, *illus.*
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Toxocarinae
key
Syn.: *Dujardinia Gedoelst*, 1916, in part, nec *Quatrefages*, 1844
- Parafilaria bovicola*
Chauhan, P. P. S.; et al., 1976, *Indian J. Animal Sc.*, v. 46 (3), 152-153
buffalo (lacrimal duct)
- Parafilaria bovicola*, *illus.*
Nevill, E. M., 1975, *Onderstepoort J. Vet. Research*, v. 42 (1), 41-48
Parafilaria bovicola of cattle, potential fly vectors as determined by field collections and laboratory studies, seasonal variation in rates of infection in flies, only female flies were infected
Musca lusoria (head, abdomen) (nat. and exper.)
M. xanthomelas (head) (nat. and exper.)
Musca n. sp. (head, abdomen, thorax) (nat. and exper.)
all from Transvaal

- Parafilaria bovicola* (Tubangui 1934)
Viljoen, J. H., 1976, J. South African Vet. Ass., v. 47 (3), 161-169
Parafilaria bovicola, calves, prepatent period, clinical aspects, filaricidal effects of 9 compounds; levamisole hydrochloride and fenbendazole reasonably effective: vicinity of Onderstepoort
- Parafilaria multipapillosa*, illus.
Andersson, P.; Jalkanen, L.; and Nurmio, P., 1976, Finsk Vet.-Tidskr., v. 82 (2), 87-89
horse, case report: Finland, imported from Russia
- Parafilaroides decorus*
Sweeney, J. C.; and Gilmartin, W. G., 1974, J. Wildlife Dis., v. 10 (4), 370-376
survey, diseases in California sea lions, diagnosis, treatment
Zalophus californianus: southern California beaches
- Parafilaroides gymnuris* Railliet
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Phoca vitulina (lung): European waters
- Parafilaroides krascheninnikovi* Jurachno et A. Skriabin, 1971
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (lungs): Sakhalin Bay
- Paragordius varius* (Leidy), illus.
Ali-Khan, F. E. A.; and Ali-Khan, Z., 1977, J. Parasitol., v. 63 (1), 174-176
pseudoparasitism
human (stool): Quebec, Canada
- Paragordius varius*, illus.
Zapotosky, J. E., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 209-221
Paragordius varius, larval stage, fine structure of preseptum
- Paragordius varius* (Leidy, 1851), illus.
Zapotosky, J. E., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 103-111
Paragordius varius larvae, postseptum fine structure using electron microscopy
- Paragraphidium Freitas et Mendonca*, 1959
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Paraheligionella Durette-Desset*, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligionellidae, Heligionellinae
- Paraheligionina Durette-Desset*, 1971
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Paraheligionina* (Ortlepp, 1939)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligionellidae, Brevistriatinae
Syn.: Heligionobaylisia Mawson, 1961
- Paraheterotyphlum Johnston & Mawson*, 1948
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
"excluded from the key since . . . incompletely described"
- Paraheterotyphlum ophiophagos* n. sp., illus.
Schmidt, G. D.; and Kuntz, R. E., 1973, Am. Midland Naturalist, v. 89 (2), 481-484
Laticauda colubrina (small intestine): Tai-tung, Tai-tung Hsien, Taiwan
- Parahistiocephalus Belopolskaja*, 1953
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Ancyracanthopsis* Diesing, 1861
- Parahistiostrongylus Vigueras*, 1941
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Parahistiostrongylus viguerasi* (Lopez-Neyra, 1946), illus.
Sanchez-Acedo, C.; Otero, J.; and Albala-Perez, F., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 245-252
Rhinolophus ferrum equinum
Myotis myotis
all from Spain
- Paraiotonchium* gen. n.
Slobodianiuk, O. V., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 156-168
Sphaerulariidae, Iotonchiinae [p. 160 put in Allantonematinae]
tod: *P. autumnalis* (Nickle, 1967) comb. n.
- Paraiotonchium autumnalis* (Nickle, 1967) comb. n. (tod), illus.
Slobodianiuk, O. V., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 156-168
Syn.: *Heterotylenchus autumnalis* Nickle, 1967
Musca autumnalis (body cavity): Bakhchisaraisk region, Krymsk oblast
- Paraiotonchium nicholasi* sp. nov., illus.
Slobodianiuk, O. V., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 156-168
Syn.: *Heterotylenchus* sp. Nicholas and Hughes, 1970
Musca vetustissima (body cavity): Australia
- Paraleiuris Vaz & Pereira*, 1929
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"cannot be classified . . . probably a member of the Ascaropsinae"

- Paraleptus H. W. Wu, 1927, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Proleptinae
key
- Paraleptus Wu, 1927
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Physalopteridae, Physalopterinae
key
- Paraleptus australis Johnston and Mawson, 1943
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
[as syn. of] Neoleptus australis (Johnston and Mawson, 1943) comb. nov.
- Paralibyostrongylus Ortlepp, 1939
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Libyostrongylinae
- Parallintoschius [sic] Araujo, 1940
Durette-Desset, M.-C.; and Chabaud, A.-G., 1975, Ann. Parasitol., v. 50 (3), 303-337
as syn. of Allintoschius [sic] Chitwood, 1937
- Parallintoschius Araujo, 1940
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Allintoschius Chitwood, 1937
- Paramecistocirrus Roetti, 1941
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
- Paramermis (?) foveata sp. n., illus.
Rubtsov, I. A., 1976, Zool. Zhurnal, v. 55 (9), 1292-1298
Lestes sponsa: Volynsk obl., g. Novovolynsk
- Paramidostomum Freitas et Mendonca, 1950
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Paramonoviola gen. n.
Blinova, S. L.; and Vosilite, B. S., 1976, Zool. Zhurnal, v. 55 (1), 131-133
Diplogasteridae, Diplogasterinae
mt: P. rhagii sp. n.
- Paramonoviola rhagii sp. n. (mt), illus.
Blinova, S. L.; and Vosilite, B. S., 1976, Zool. Zhurnal, v. 55 (1), 131-133
development of egg and larvae
Rhagium inquisitor (fat body): Serebriano-borsk preserve, Moskovsk oblast
Ips sexdentatus (fat body) (exper.)
- Paranematospira Sprehn, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Heligmosomoides Hall, 1916
- Paranisakinea nov. trib.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinae
key; key to genera
includes: Paranisiakiopsis; Paranisakis
- Paranisiakiopsis
Gibson, D. I.; and Taylor, A. L., 1976, Parasitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Paranisiakiopsis Yamaguti, 1941
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Paranisakinea nov. trib.
key
- Paranisiakiopsis [sp.]
Lichtenfels, J. R.; et al., 1976, Tr. Am. Micr. Soc., v. 95 (2), 265-266 [Abstract]
anisakid larvae, resembles Paranisiakiopsis, from commercially important shellfish, description of 4th stage, nearly 100% hyperparasitized by haplosporidan
Spisula solidissima
Busycon canaliculata
Lunatia heros
all from coastal waters from New Jersey to North Carolina
- Paranisakis
Gibson, D. I.; and Taylor, A. L., 1976, Parasitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Paranisakis Baylis, 1923
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Paranisakinea nov. trib.
key; synonymy
- Parapharyngodon maplestonei Chatterji 1933
Pinnell, J. L.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 337-340
Sphenomorphus emigrans: Komodo Island, Indonesia
- Pararhabdonema Kreis, 1945
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Libyostrongylinae
- Parascaris
Blanchard, J., 1971, Prat. Vet. Equine, v. 3 (1), 29-31
gastrointestinal nematodes in race horses, clinical trials with equigard, good immediate results especially with ascarids; rapid reinfection rates (2 months after therapy), suggests contaminated paddocks; recommendations therefore for program of routine treatment: France

- Parascaris Yorke & Maplestone, 1926
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridinae
key
- Parascaris
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Parascaris
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Parascaris equorum
Abdel-Rahman, M. S.; et al., 1972, Parasitol. Hungar., v. 5, 225-237
Strongylus spp., Trichonema spp., Parascaris equorum, Trichostrongylus axei in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Parascaris equorum
Ardans, A.; and Walters, G., 1975, Am. J. Vet. Res., v. 36 (11), 1589-1590
Parascaris equorum, strongyles, Quarter horses, treatment with 3 formulations of cambendazole (paste, pellets, suspension) vs. thiabendazole (suspension)
- Parascaris equorum
Bemrick, W. J.; O'Leary, T. P.; and Averbeck, G. A., 1977, Immunology, v. 32 (4), 567-572
Parascaris equorum, guinea pigs, homocytotropic anaphylactic response to antigen extracts from formalized and unformalized worms
- Parascaris equorum
Blaszko, W., 1972, Med. Wet., v. 28 (4), 244-245
Parascaris equorum, obstructing horse ileum, surgical removal
- Parascaris equorum, illus.
Clayton, H. M.; and Duncan, J. L., 1977, Research Vet. Sc., v. 23 (1), 109-114
Parascaris equorum, foals (exper.), clinical findings, haematological findings, parasitology, establishment of patency, pathology
- Parascaris equorum
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Parascaris equorum
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Research Vet. Sc., v. 14 (1), 134-136
Parascaris equorum, foals, critical trials with morantel tartrate
- Parascaris equorum (Goeze, 1782)
Derkmann, K.; and Hasslinger, M. A., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (5), 95-98
Parascaris equorum, naturally infected horses, panacur, good results with high dosage: Munich area
- Parascaris equorum
Drudge, J. H.; and Lyons, E. T., 1977, Am. J. Vet. Research, v. 38 (10), 1581-1586
internal parasites, horses, antiparasitic drugs, methods of evaluation, critical tests, controlled tests and clinical trials, review of methods and criteria
- Parascaris equorum
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Parascaris equorum
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, Prakt. Tierarzt, v. 55 (8), 417-422
nematodes of horses, Fenbendazol, good results
- Parascaris equorum
Furmaga, S.; Gundlach, J. L.; and Patyra, J., 1976, Med. Wet., v. 32 (12), 734-737
roundworms, horses, fenbendazol and cambendazol very effective
- Parascaris equorum
Greve, J. H.; and Paul, J. W., 1976, Vet. Med. and Small Animal Clin., v. 71 (12), 1737-1740, 1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Parascaris equorum
Hiepe, T., 1972, Schweiz. Arch. Tierh., v. 114 (12), 613-614
Parascaris equorum; Strongyloides westeri, sports horses of all ages, Morantel tartrate, good results
- Parascaris equorum (Goeze, 1782)
Ismailov, T., 1976, Dokl. Akad. Nauk UzSSR (9), 65-66
Parascaris equorum, effect of humidity and temperature on embryogenesis under field conditions in Uzbekistan; metamorphosis from April to October, anabiosis from November to March
- Parascaris equorum
Krvavica, S.; Francetic, D.; and Zivkovic, D., 1976, Vet. Arhiv, Zagreb, v. 46 (9-10), 231-239
nematodes, trematodes, cestodes, activity, distribution and cofactor dependence of malic enzymes, majority are located in mitochondria in all investigated parasites

- Parascaris equorum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Parascaris equorum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, J. Parasitol., v. 62 (3), 453-459
Parascaris equorum, foals (nat. and exper.), development and location of migrating larvae at specific time intervals after infection, evaluation of possible chemotherapeutic activity of several anthelmintics against migrating and early enteric-stage larvae
- Parascaris equorum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, Am. J. Vet. Research, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and Gasterophilus (inactive against latter)
- Parascaris equorum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Parascaris equorum
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 721-723
helminths and bots in horses, thiabendazole and trichlorfon sequentially administered via stomach tube, critical testing, drug efficacies, good results
- Parascaris equorum
McCall, J. P.; and McCullough, C., 1977, Southwest. Vet., v. 30 (2), 159-160
strongylids, Parascaris equorum, horses (feces), thiabendazole, piperazine phosphate, good results; failure to demonstrate acquired drug resistance
- Parascaris equorum
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Parascaris equorum
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Parascaris equorum
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Parascaris equorum
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Parascaris equorum, illus.
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Parascaris equorum (Goeze, 1782)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Parascaris equorum
Smith, H. J., 1976, Canad. J. Comp. Med., v. 40 (4), 327-333
strongyles, Oxyuris equi, Parascaris equorum, naturally-infected ponies, thiabendazole, results support regular repeated treatment
- Parascaris equorum
Stevenson, P.; and Jacobs, D. E., 1977, J. Helminth., v. 51 (2), 149-154
Toxocara canis, T. cati, Ascaris suum, Toxascaris leonina, Parascaris equorum, pigs (exper.), in vitro larval precipitate test and indirect fluorescent antibody test using T. canis larvae as antigen, indirect fluorescent antibody test using A. suum larvae as antigen, specificity
- Parascaris equorum
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- P[arascaris] equorum
Tolone, G.; et al., 1972, Pathol. et Microbiol., v. 38 (3), 192-199
role of mast cells and eosinophiles in tissue injury after injection of P[arascaris] equorum fluid into rat (exper.) peritoneal cavity
- Parascaris equorum, illus.
Trimble, J. J. III; and Thompson, S. A., 1976, Cell and Tissue Research, v. 172 (3), 357-363
Ascaris suum, Parascaris equorum, distribution of concanavalin A binding site on nematode intestinal epithelium
- Parascaris equorum
Wlodarczak, C., 1972, Med. Wet., v. 28 (4), 221-222
strongyloidosis, Parascaris equorum, horses, Pyrequan

- Parascaris equorum* race *univalens*
Moritz, K. B.; and Roth, G. E., 1976, *Nature* (5538), v. 259, 55-57
Ascaris lumbricoides, *Parascaris equorum*, complexity of germline and somatic DNA
- Parascarophis Campana-Rouget*, 1955, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Parasitaphelenchus sexdentati*
Vosilite, B. S., 1975, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Parasitorhabditis sexdentati*
Vosilite, B. S., 1975, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 25, 13-17
nematode infection of *Ips sexdentatus* in relation to host life cycle, generations and seasonal distribution: Lithuanian SSR
- Paraspidodera uncinata* (Rudolphi, 1819) *Travassos*, 1914, *illus.*
Ibanez Herrera, N., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 15-20
synonymy, redescription
Cavia porcellus (ciego): Peru
- Paraspidodera uncinata*
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 214
Meerschweinchen
- Paraspidodera uncinata* (Rudolphi, 1819) *Travassos*, 1914
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 39-42
morphometric data
Ctenomys maulinus maulinus
C. m. brunneus
(intestino grueso of all): all from Chile
- Paraspidodera uruguayana* Khalil y Vogelsang, 1931 [et auct.]
Ibanez Herrera, N., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 15-20
as syn. of *Paraspidodera uncinata* (Rudolphi, 1819) *Travassos*, 1914
- Paraspirura* Sandground, 1936, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruridae
key
- Parastrongyloides peramelis* Mackerras
Beveridge, I.; and Barker, I. K., 1976, *Austral. J. Zool.*, v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (intestine): Powelltown, Victoria
- Parastrongyloides winchesi* Morgan, 1928
Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
Sorex: Catalan Pyrenean Mountains
- Parastrongylus dujardini* (Drozdz & Doby, 1970)
Mas-Coma, S.; and Gallego, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (3-4), 261-281
Clethrionomys glareolus: Catalan Pyrenean Mountains
- Parathelandros armatus* (Walton, 1933) n. comb.
Specian, R. D.; and Ubelaker, J. E., 1974, *Tr. Am. Micr. Soc.*, v. 93 (3), July, 413-415
Syn.: *Pharyngodon armatus* Walton, 1933
- Parathelandros texanus* n. sp., *illus.*
Specian, R. D.; and Ubelaker, J. E., 1974, *Tr. Am. Micr. Soc.*, v. 93 (3), July, 413-415
"Walker & Matthias (1973) reported *Pharyngodon warneri* from *Urosaurus ornatus* in Arizona. Examination of these specimens, kindly loaned to us by Dr. Walker, indicates that they are conspecific with *Parathelandros texanus* n. sp. as described above."
Sceloporus merriami: west Texas
S. undulatus: west Texas
Cophosaurus texana: west Texas
Cnemidophorus tigris: west Texas
C. inornatus: west Texas
C. scalaris: west Texas
Urosaurus ornatus: west Texas; Arizona (large intestine of all)
- Paraastrostrongylus* Mawson, 1973
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Parelaphostrongylus andersoni*, *illus.*
Nettles, V. F.; and Prestwood, A. K., 1976, *Vet. Path.*, v. 13 (5), 381-393
Parelaphostrongylus andersoni, *Odocoileus virginianus* (exper.), gross and microscopic lesions, lungs, muscles, clinical signs, egg production, course of infection
- Parelaphostrongylus andersoni*
Prestwood, A. K.; and Nettles, V. F., 1977, *J. Parasitol.*, v. 63 (6), 974-978
Parelaphostrongylus andersoni, repeated low-level infection of *Odocoileus virginianus*, clinical, parasitologic, and pathologic findings, apparent production of active immunity, results suggest that wild deer become infected by isolated chance encounters with infected gastropods
- Parelaphostrongylus andersoni* Prestwood 1972
Prestwood, A. K.; Nettles, V. F.; and Kellogg, F. E., 1974, *J. Wildlife Dis.*, v. 10 (4), 404-409
survey, prevalence of *Parelaphostrongylus andersoni* and *P. tenuis* among white-tailed deer
Odocoileus virginianus: southeastern United States
- Parelaphostrongylus andersoni* Prestwood, 1972
Pursglove, S. R., jr., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 107-108
Odocoileus virginianus (musculature): Cumberland County, New Jersey

- Parelaphostrongylus tenuis* Pryadko and Boev, 1971, *illus.*
Carpenter, J. W.; Jordan, H. E.; and Ward, B. C., 1973, *J. Wildlife Dis.*, v. 9 (2), 148-153
Parelaphostrongylus tenuis, *Cervus canadensis* (meninges and parenchyma of brain), clinical signs of neurologic disturbances, histopathologic lesions in CNS apparently related to nematode damage: Oklahoma
- Parelaphostrongylus tenuis* (Dougherty)
Gilbert, F. F., 1973, *J. Wildlife Dis.*, v. 9 (2), 136-143
Parelaphostrongylus tenuis, prevalence in *Odocoileus virginianus* males vs. females, fawns vs. adult deer, areas of high vs. low deer density, localization within cranial cavity, implications for transmission: Maine
- Parelaphostrongylus tenuis*, *illus.*
Lankester, M. W.; Crichton, V. J.; and Timmermann, H. R., 1976, *Canad. J. Zool.*, v. 54 (5), 680-684
1st stage protostrongylid larvae in *Rangifer tarandus caribou* (feces) may be *Elaphostrongylus* sp., lack of pathogenic effects seems to rule out *Parelaphostrongylus tenuis*: northwestern Ontario and Manitoba
- Parelaphostrongylus tenuis* (Dougherty 1945)
Pryadko & Boev 1971
Prestwood, A. K.; Nettles, V. F.; and Kellogg, F. E., 1974, *J. Wildlife Dis.*, v. 10 (4), 404-409
survey, prevalence of *Parelaphostrongylus andersoni* and *P. tenuis* among white-tailed deer
Odocoileus virginianus: southeastern United States
- Parelaphostrongylus tenuis*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Odocoileus virginianus: Hardy County, West Virginia
- Parelaphostrongylus tenuis* (Dougherty, 1945)
Pursglove, S. R., jr., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 107-108
Odocoileus virginianus (brain): Cumberland County, New Jersey; Oklahoma
- Parelaphostrongylus tenuis*
Stackhouse, L. L., 1977, *J. Am. Vet. Med. Ass.*, v. 171 (9), 987-988
cerebral nematodiasis in *Alces alces*, histopathologic features in brain compatible with cerebrospinal nematodiasis resulting from infection by *Parelaphostrongylus tenuis*: New Hampshire
- Parhadjelia* Lent & Freitas, 1939
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Hadjelia* Seurat, 1916
- Parhadjelia neglecta*
George, R. R.; and Bolen, E. G., 1975, *J. Wildlife Dis.*, v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas
- Parlitomosa zakii*, tentative identification, *illus.*
Chalifoux, L. V.; et al., 1973, *Lab. Animal Sc.*, v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saimiri sciureus
Saguinus oedipus
S. tamarinus
Ateles geoffroyi
Aotus trivirgatus
all from New England Regional Primate Research Center
- Paronchocerca rousseloti*
Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Parostertagia* Schwartz et Alicata, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, *Graphidiinae*
- Paryseria* Johnston, 1938
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Stegophorus* Wehr, 1934
- Passalurus*
Neppert, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Passalurus ambiguus*
Kutzer, E.; and Frey, H., 1976, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 89 (24), 480-483
Lepus europaeus: Austria
- Passalurus ambiguus* Rudolphi, 1819
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 267-271
Oryctolagus cuniculus: Huesca, Alto Aragon
- Passalurus ambiguus* (Rudolphi 1819) Dujardin 1845, *illus.*
Romero Rodriguez, J.; Guevara Pozo, D.; and Lizcano Herrera, J., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 315-329
description, review of life cycle
Oryctolagus cuniculus domestica: Spain

- Passalurus nonanulatus* Skinker 1932, illus.
Romero Rodriguez, J.; Guevara Pozo, D.; and Lizcano Herrera, J., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 315-329
description, review of life cycle
Lepus granatensis: Spain
- Patricialina* Inglis, 1968
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Paucipectines*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Pterygodermatites*
key
- Pectinospirura* Wehr, 1933, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Pectinospirura argentata* Wehr, 1933
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
comparison of *Pectinospirura multidentata* and *P. argentata*, differential diagnosis
- Pectinospirura multidentata* Sobolev, 1943
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Heteroscelus incanus brevis: Keta lake
- Pectinospirura multidentata* Sobolev, 1943, illus.
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
comparison of *Pectinospirura multidentata* and *P. argentata*, differential diagnosis
Syn.: *P. sobolevi* Turemuratov 1965
Larus argentatus: Yenisei
- Pectinospirura sobolevi* Turemuratov 1965
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
as syn. of *Pectinospirura multidentata* Sobolev, 1943
- Pelecitus* n. sp., adult and microfilariae
Dissanaïke, A. S.; and Fernando, M. A., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 138 [Demonstration]
Gallus gallus spadiceus (blood and lung smears, among tendons at base of legs): Malaysia
- Pelecitus* sp.
Ramakrishnan, K.; and Aziz bin Ahmad, A., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 149 [Demonstration]
Wuchereria bancrofti, *Pelecitus* sp., membrane feeding technique for infecting mosquitoes with filarial parasites
- Pelecitus helix* (Linstow, 1899)
Aleksiev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Botaurus stellaris (leg joint): Rimsko-Korsakov islands
- Pelodera strongyloides*, illus.
Bergeland, M. E.; Todd, K. S., jr.; and Ohlen-dorf, L. F., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 230-231
Pelodera strongyloides, dermatitis in sheep: northern Illinois
- Pelodera* (*Rhabditis*) *strongyloides*
Farrington, D. O.; Lundvall, R. L.; and Greve, J. H., 1976, Vet. Med. and Small Animal Clin., v. 71 (9), 1199, 1202
Pelodera strongyloides dermatitis, horse, case history, thiabendazole, good results: Iowa
- Pelodera strongyloides*, illus.
Kipnis, R. M.; and Todd, K. S., jr., 1977, Feline Pract., v. 7 (2), 16-19
Pelodera strongyloides, cat (urine), cystitis, case history, possible case of pseudo-parasitism: northern Illinois; Green Bay, Wisconsin
- Pelodera strongyloides*
Stringfellow, F., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 4-10
Pelodera strongyloides, hydroxyl ion, attractant to males, introduction of therapeutic agents to block nematode attractant could significantly decrease reproductive potential
- Pelodera strongyloides*, illus.
Stringfellow, F., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 206-211
Pelodera strongyloides in culture, determination of presence and distribution of carbonic anhydrase in worms, addition of diamox to cultures resulted in inhibition of carbonic anhydrase, reduced quantity of ammonia nitrogen production, and decreased ability of worms to find each other
- Pentadentoptera Schachnasarova, 1949
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key
Syn.: *Physalopteriata* Sobolev, 1949
- Peramelistrongylus* Mawson, 1961
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Mackerrastrongylinae
- Peramelistrongylus skedastos* Mawson
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (stomach): Powelltown, Victoria

- Pereiraia* Cuocolo, 1943
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Physocephalus* Diesing, 1861
- Peritrachelius* Diesing, 1861
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakis* Dujardin, 1845
- Perostrongylus* pridhami
Barber, D. L.; and Lockard, L. L., 1973, Great Basin Nat., v. 33 (1), 53-60
Mustela vison (lungs): Gallan and Madison counties, Montana
- Perutilimermis* Nickle
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
"Diximermis peterseni and *Perutilimermis* culicis must be regarded as genera et species inquirendarum."
- Perutilimermis* culicis Nickle
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
"Diximermis peterseni and *Perutilimermis* culicis must be regarded as genera et species inquirendarum."
- Pesteria* Tadros, 1966
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Dracunculus* (Reichard, 1759)
- Petiellus* n. sub.gen.
Durette-Desset, M.-C.; and Chabaud, A.-G., 1975, Ann. Parasitol., v. 50 (2), 173-185
subgen. of *Nycteridostromylus*; tod: *N. (Petiellus) petersi* n. sp.
- Petrovinema*
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Petrovinema* poculatum (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Petroviprocta* Schachtachtinskaja, 1951
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Avioserpens* Wehr & Chitwood, 1934
- Petrowimeres* Chertkova, 1953
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
discussion of criteria for differentiating from other subgenera of *Tetrameres* includes: *T. fissispina* Diesing 1861; *T. australis* Johnston and Mawson 1941; *T. biziurae* Johnston and Mawson 1941; *T. crami* Swales 1933; *T. galericulatus* Oschmarin 1956; *T. mohtedai* Bhalerao and Rao 1944; *T. nettatus* Ali 1970; *T. pavonis* Chertkova 1953; *T. rijikovi* Chuan 1961; *T. somateriae* Ryjikov 1963; *T. striatus* Oschmarin 1956; *T. indiana* Ali 1970; *T. plectropteri* Thwaite 1926
- Petrowospirura* Matschulsky, 1952
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cylicospirura* Vevers, 1922
- Pharurus* Leuckart 1848
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
key; key to species; diagnosis, differentiation from *Torynurus*
Syn.: *Otophocaenurus*
- Pharurus* alatus (Leuckart 1848) Stiles and Hasall 1905
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
synonymy; redescription; key
Monodon monoceros: off Baffin Is., Canada
- Pharurus* convolutus (Kuhn) Dougherty 1943
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829) Baylis and Daubney 1925
- Pharurus* dalli (Yamaguti) Yamaguti 1962
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Torynurus dalli* (Yamaguti 1951) Delyamure 1972
- Pharurus* minor (Kuhn) Cobbold 1879
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis and Daubney 1925
- Pharurus* oserskaiaae (Skrjabin) Dougherty 1949
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden 1870) n. comb.

- Pharurus pallasii* (van Beneden 1870) n. comb.,
illus.
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
redescription; key
Syns.: *Strongylus pallasii* van Beneden 1870; *S. arcticus* Cobb 1888; *Pseudalius arcticus* (Cobb) von Linstow 1900; *Stenurus arcticus* (Cobb) Baylis and Daubney 1925; *Otophocaenurus oserskoi* Skrjabin 1942; *Stenurus pallasii* (van Beneden) Dougherty 1943; *Pharurus oserskaiaae* (Skrjabin) Dougherty 1949
Delphinapteras leucas: MacKenzie River Delta, Canada; New Brunswick, Canada; Churchill, Manitoba, Canada
- Pharyngodon* Diesing, 1861
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
key to species, includes: *Pharyngodon mudgi* sp. n.; *P. kartana* Johnston and Mawson, 1941; *P. spinicauda* Dujardin, 1845; *P. yucatanensis* Chitwood, 1938; *P. hindlei* Thapar, 1925; *P. geckinis* Liu and Wu, 1941; *P. tiliquae* Baylis, 1930; *P. inermicauda* Baylis, 1923; *P. neyrae* Calvente, 1948; *P. australis* Johnston and Mawson, 1944; *P. mamillatus* (Linstow, 1897); *P. papilliocauda* Hannum, 1942; *P. cesarpintoi* Pereira, 1935
- Pharyngodon* sp., illus.
Bilqees, F. M.; and Siddiqui, M. H., 1975, *Pakistan J. Scient. and Indust. Research*, v. 18 (6), 261-264
Gecko sp. (intestine): Karachi
- Pharyngodon* sp., illus.
Guimaraes, J. F., 1975, *Bol. Inst. Biol., Bahia*, v. 14 (1), 44-52
description
Tropidurus torquatus (intestino delgado): Ondina, Salvador, Bahia, Brasil
- Pharyngodon anolis* (Chitwood, 1934)
Acholonu, A. D., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 106-116
Anolis cristatellus
A. evermanni
Ameiva exsul
(intestines of all): all from Puerto Rico
- Pharyngodon armatus* Walton, 1933
Specian, R. D.; and Ubelaker, J. E., 1974, *Tr. Am. Micr. Soc.*, v. 93 (3), July, 413-415
as syn. of *Parathelandros armatus* (Walton, 1933) n. comb.
- Pharyngodon cnemidophori* Read and Armein, 1954
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
Cnemidophorus tigris: Javelina Creek bed, west Texas
- Pharyngodon giganticus*
Pearce, R. C.; and Tanner, W. W., 1973, *Great Basin Nat.*, v. 33 (1), 1-18
Sceloporus occidentalis
Sceloporus undulatus
(cecum of all): all from Great Basin and Upper Colorado Plateau, Utah
- Pharyngodon kirbii* sp. n., illus.
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
key
Cnemidophorus scalaris (large intestine): Stairway Mountain, Black Gap Wildlife Area, Brewster County, Texas
- Pharyngodon kuntzi* Gupta 1959
Pinnell, J. L.; and Schmidt, G. D., 1977, *J. Parasitol.*, v. 63 (2), 337-340
Gekko gekko
Sphenomorphus emigrans
all from Komodo Island, Indonesia
- Pharyngodon medinae* Calvente, 1948
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
as syn. of *Skrjabinodon medinae* (Calvente, 1948) n. comb.
- Pharyngodon mudgi* sp. n., illus.
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
key
Coleonyx brevis (large intestine): Black Gap Wildlife Management Area, Brewster County, Texas
- Pharyngodon paratectipenis* Chabaud and Golvan, 1957
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
as syn. of *Spauligodon paratectipenis* (Chabaud and Golvan, 1957) n. comb.
- Pharyngodon tectipenis sensu Calvente, 1948 nec P. tectipenis Gedoelst, 1919*
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
as syn. of *Spauligodon paratectipenis* (Chabaud and Golvan, 1957) n. comb.
- Pharyngodon travassosi* Pereira, 1935
Acholonu, A. D., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 106-116
Anolis cristatellus
A. evermanni
(small intestines of all): all from Puerto Rico
- Pharyngodon warneri* Harwood, 1932
Specian, R. D.; and Ubelaker, J. E., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 46-51
Cnemidophorus inornatus: west Texas
- Pharyngodon warneri*
Specian, R. D.; and Ubelaker, J. E., 1974, *Tr. Am. Micr. Soc.*, v. 93 (3), July, 413-415
"Walker & Matthias (1973) reported *Pharyngodon warneri* from *Urosaurus ornatus* in Arizona. Examination of these specimens, kindly loaned to us by Dr. Walker, indicates that they are conspecific with *Parathelandros texanus* n. sp. as described above."

- Pharyngostrongylus gallardi* Johnston & Mawson, 1942
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
as syn. of *Cyclostrongylus wallabiae* Johnston & Mawson, 1939
- Pharyngostrongylus parma*
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
as syn. of *Cyclostrongylus parma* (Johnston & Mawson 1939) [n. comb.]
- Pheromermis* n. gen.
Poinar, G. O., jr.; Lane, R. S.; and Thomas, G. M., 1976, Nematologica, v. 22 (3), 360-370
Mermithidae, tod: *P. pachysoma* n. comb.
- Pheromermis pachysoma* (von Linstow) n. gen., n. comb. (tod), illus.
Poinar, G. O., jr.; Lane, R. S.; and Thomas, G. M., 1976, Nematologica, v. 22 (3), 360-370
redescription
Gumaga griseolum (intestinal cells, muscles)
Tipula sp. (gut wall, fat body)
beetle (gut wall)
ephemerid (gut wall)
Vespula pennsylvanica (body cavity)
Culex pipiens (exper.)
Callibaetis pictus (exper.)
all from University of California Field Station, outside of Hopland, Mendocino County, California
- Philometra*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Philometra*
key
- Philometra* Costa, 1845, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key; synonymy
includes subgens.: *Ranjhinema*; *Alinema*;
Philometra
- Philometra* Costa, 1945
Molnar, K.; and Fernando, C. H., 1975, J. Helminth., v. 49 (2), 101-105
Syn.: *Thwaitia* Rasheed, 1963
- Philometra* sp.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Euthynnus affinis (ovaries): South China Sea
- Philometra* sp., illus.
Reichenbach-Klinke, H. H., 1975, Fisch u. Umwelt (1), 113-121
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Philometra abdominalis* Nybelin, 1928
Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmit. Lab., v. 16, 87-110
as syn. of *Thwaitia abdominalis* (Nybelin, 1928) Rasheed, 1963
- Philometra abdominalis*, illus.
Reichenbach-Klinke, H. H., 1975, Fisch u. Umwelt (1), 113-121
Nematoda in fresh water fish as food hygiene problems, possible controls, review
- Philometra carassii* (Ishii, 1934)
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Carassius auratus (between caudal fin rays): Sacramento-San Joaquin Delta, California
- Philometra cylindracea* Ward and Magath, 1916, illus.
Ashmead, R. R.; and Crites, J. L., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 143-145
description of male, redescription of female
Perca flavescens: Lake Erie, Put-in-Bay Twp., Ottawa Co., Ohio
- Philometra cylindracea* (Ward and Magath, 1917)
Cooper, C. L.; Ashmead, R. R.; and Crites, J. L., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 96
prevalence, comparison with previous years
Perca flavescens (body cavity): western Lake Erie
- Philometra cylindracea* (Ward and Magath, 1916), illus.
Molnar, K.; and Fernando, C. H., 1975, J. Helminth., v. 49 (1), 19-24
Philometra cylindracea, description of male, redescription of female, developmental cycle
Syn.: *Ichthyonema cylindraceum* Ward and Magath, 1916
Perca flavescens (under serosa of air bladder, body cavity) (nat. and exper.): Laurel Creek, Ontario, Canada
Cyclops vernalis (exper.)
- Philometra cylindracea* Ward & Magath, 1916
White, G. E., 1974, Tr. Am. Micr. Soc., v. 93 (2), Apr., 280-282
Catostomus commersoni: Kentucky River drainage system
- Philometra kobuleji* sp. n., illus.
Molnar, K.; and Fernando, C. H., 1975, J. Helminth., v. 49 (2), 101-105
Catostomus commersoni (under serosa of air bladder, body cavity): Laurel Creek, Waterloo, and Bronte Creek, near Milton, Ontario, Canada
- Philometra nodulosa*
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Micropterus salmoides (nasal cavities, mouth): Eagle Mountain Lake, Texas
- Philometra obturans* Prenant, 1886
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland

- Philometra obturans* (Prenant, 1886), illus.
Molnar, K., 1976, Acta Vet., Budapest, v. 26 (2), 183-188
Philometra obturans, male and female described, life cycle
Esox lucius (nat. and exper.) (beneath serosa of air bladder, beneath peritoneum, gill vessels): river Danube, nearby Paks, Hungary; river Tisza, nearby Szolnok, Hungary
Cyclops strenuus (exper.) (body cavity)
Acanthocyclops viridis (exper.) (body cavity)
Cyprinus carpio (exper.)
- Philometra rischta* Skrjabin, 1917
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland
- Philometra rischta* Skrjabin, 1917
Kakacheva-Avramova, D., 1972, Izvest. Tsentral. Khelmit. Lab., v. 15, 89-107
Vimba vimba melanops (mucous membrane of gill cover): River Tundzha
- Philometra sanguinea* (Rud., 1819)
Fagerholm, H.-P., 1976, Norwegian J. Zool., v. 24 (4), 466 [Abstract]
Finland
- Philometridae Baylis & Daubney, 1926
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Draconculicidea
key; key to subfamilies
includes: Philoneminae; Philometrinae; Phlyctainophorinae
- Philometridae [sp.]
Bussieras, J.; and Baudin-Laurencin, F., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (4), 13a-19a
Thunnus albacares (ovaires): tropical Atlantic
- Philometrinae (Baylis & Daubney, 1926, fam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometridae
key; key to genera
includes: Ichthyofilaria; Thwaitia; Rumi; Philometra; Spirophilometra; Nilonema; Buckleyella; Philometroides
- Philometroides Yamaguti, 1935, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key
Syn.: Pseudophilometroides Parukhin, 1966
- Philometroides huronensis n. sp., illus.
Uhazy, L. S., 1976, Canad. J. Zool., v. 54 (3), 369-376
Catostomus commersoni (pectoral, pelvic, dorsal, and anal fins; peritoneum around swim bladder): Southern Lake Huron, Ontario, Canada
- Philonema Kuitunen-Ekbaum, 1933, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philoneminae
? Syn.: Coregonema Bauer, 1946
- Philonema oncorhynchi
Boyce, N. P.; and Yamada, S. B., 1977, J. Fish. Research Bd. Canada, v. 34 (5), 706-709
Oncorhynchus nerka: outlet of Babine Lake, central British Columbia
- Philonema oncorhynchi Kuitunen-Ekbaum, 1933
Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, Fish. Bull., National Oceanic and Atmos. Admir., v. 71 (1), 267-277
helminths, incidence and intensity of infection in young and adult *Oncorhynchus nerka*, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Philoneminae (Skrjabin, Sobolev & Khromova, 1971)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometridae
key
includes: *Philonema*
- Philostrongylus Wolfgang, 1951
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Viannaia* Travassos, 1914
- Phlyctainophora Steiner, 1921, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Phlyctainophorinae
- Phlyctainophorinae (Roman, 1965, fam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometridae
key
includes: Phlyctainophora
- Phocanema Myers, 1959
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinea
key
- Phocanema (= Terranova)
Myers, B. J., 1976, Tr. Am. Micr. Soc., v. 95 (2), 137-142
human anisakiasis, historical review
- Phocanema sp.
Bier, J. W.; et al., 1976, Tr. Am. Micr. Soc., v. 95 (2), 264-265 [Abstract]
Anisakis sp. and *Phocanema* sp. in pigs (exper.) (stomach), gross and microscopic pathology

- Phocanema sp.
Bier, J. W.; Jackson, G. J.; and Gerding, T. A., 1976, Tr. Am. Micr. Soc., v. 95 (2), 265 [Abstract]
Anisakis sp., Phocanema sp., blood analysis of experimentally infected pigs showed mild eosinophilia and increased levels of amylase, lactic dehydrogenase, and bilirubin
- Phocanema-type larvae
Chitwood, M., 1975, Am. J. Trop. Med. and Hyg., v. 24 (4), 710-711
Phocanema-type larva coughed up by boy who 4 days earlier had eaten marinated raw fish, case report: California
- Phocanema sp.
Jackson, G. J.; Bier, J. W.; and Payne, W. L., 1976, Tr. Am. Micr. Soc., v. 95 (2), 264 [Abstract]
Anisakis sp. and Phocanema sp. in miniature pigs (exper.) (stomach), course of infection
- Phocanema sp. larva
Juels, C. W.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 942-944
Phocanema sp. larva coughed up from throat of man several days after he had eaten raw fish, case report, evidence of possible temporary tissue invasion because of mild transitory eosinophilia: California
- Phocanema sp., illus.
Lichtenfels, J. R.; and Brancato, F. P., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 691-693
Phocanema sp., fourth-stage larva, pulled from throat of Eskimo, case report, parasite morphology: Alaska
- Phocanema spp. larvae
van Thiel, P. H., 1976, Trop. and Geogr. Med., v. 28 (2), 75-85
human anisakiasis, current status, review (diagnosis, treatment, epidemiology, distribution, definitive and intermediate hosts)
- Phocanema decipiens, illus.
Goh, S. L.; and Davey, K. G., 1976, Canad. J. Zool., v. 54 (5), 752-771
Phocanema decipiens, nervous system, distribution and localization of acetylcholinesterase, presence in synapses of large dense-core vesicles and small lucent vesicles
- Phocanema decipiens, illus.
Goh, S. L.; and Davey, K. G., 1976, Internat. J. Parasitol., v. 6 (5), 403-411
Phocanema decipiens, nervous system, catecholaminergic structures, localization and distribution using formaldehyde-induced and glyoxylic acid fluorescence histochemical techniques
- Phocanema decipiens, illus.
Goh, S. L.; and Davey, K. G., 1976, Tissue and Cell, v. 8 (3), 421-435
Phocanema decipiens, nervous system, selective uptake of noradrenaline, dopa, and 5-hydroxytryptamine, light autoradiographic and ultrastructural study
- Phocanema decipiens
Margolis, L., 1977, J. Fish. Research Bd. Canada, v. 34 (7), 887-898
Phocanema decipiens in humans that eat raw, lightly marinated, or undercooked marine fishes, diagnosis, treatment, public health implications, review
- Phocanema decipiens (Krabbe, 1878)
van Thiel, P. H., 1976, Trop. and Geogr. Med., v. 28 (2), 75-85
human anisakiasis, current status, review (diagnosis, treatment, epidemiology, distribution, definitive and intermediate hosts)
- Phocascaris Hoest, 1932
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Contraceacinea
key
- Phocascaris
Soleim, O., 1976, Norwegian J. Zool., v. 24 (4), 464 [Abstract]
"It is concluded that the genus Thynnascaris should be maintained and that Phocascaris become a synonym of Contraceacum."
- Phocascaris cystophorae Berland, 1963
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (stomach): Sakhalin Bay
- Phocascaris cystophorae Berland, 1963
Popov, V. N., 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
age dynamics of infection
Histriophoca fasciata (stomach, intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Phocascaris phocae
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Phoca vitulina (stomach): European waters
- Pholidostrongylus Baer, 1959
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Trichocheenia Kou, 1958
- Physaloptera Rudolphi, 1819, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key
- Physaloptera
Isenbuegel, E., 1976, Prakt. Tierarzt, v. 57, Sondernummer, 21-27
carbon tetrachloride, unsuccessful
Igel

NEMATODA

8

- Physaloptera sp.
 Babero, B. B., 1973, Tr. Am. Micr. Soc.,
 v. 92 (2), 265-272
 "may be assignable to *P. massino*"
Ammospermophilus leucurus (stomach): Nevada
- Physaloptera sp., illus.
 Babero, B. B.; and Cattán, P. E., 1975, Bol.
 Chileno Parasitol., v. 30 (3-4), 68-76
Octodon degus (estomago): Quebrada de la
 Plata, Santiago, Chile
- Physaloptera sp.
 Cattán, P. E.; George-Nascimento, M.; and
 Rodríguez, J., 1976, Bol. Chileno Parasitol.,
 v. 31 (1-2), 16-20
 prevalence survey of helminths of *Octodon
 degus*, seasonal variations, age and sex of
 hosts: Chile
- Physaloptera sp.
 Courtney, C. H.; and Forrester, D. J., 1974,
 Proc. Helminth. Soc. Washington, v. 41 (1),
 89-93
 prevalence and intensity, age of host
Pelecanus occidentalis (small intestine):
 Florida
- Physaloptera sp.
 Croft, R. E.; and Kingston, N., 1975, J. Wild-
 life Dis., v. 11 (2), 229-233
Falco mexicanus: Wyoming
- Physaloptera sp.
 Davidson, W. R., 1976, Proc. Helminth. Soc.
 Washington, v. 43 (2), 211-217
 epizootiologic and pathologic study of endo-
 parasitism of selected populations of gray
 squirrels
Sciurus carolinensis (stomach): southeastern
 United States
- Physaloptera sp.
 Dyer, W. G.; and Brandon, R. A., 1973, Tr.
 Illinois Acad. Sc., v. 66 (1-2), 23-29
Plethodon dorsalis (stomach): Equality
 Cave, southwest of Equality, Saline County,
 Illinois
- Physaloptera spp.
 McCurdy, H. D.; and Guerrero, J., 1977, West.
 Med. and Small Animal Clin., v. 72 (11),
 1731-1733
 helminths, dogs, mebendazole powder, con-
 trolled critical studies, good results
 against all helminths except *Dipylidium
 caninum*: Kansas; New Jersey; Texas
- Physaloptera sp.
 Martin, D. R., 1976, Proc. Helminth. Soc.
 Washington, v. 43 (1), 85-86
Tadarida brasiliensis: Texas; Louisiana
- Physaloptera sp. 1
 Mushkambayeva, M. G., 1973, Ekol. Nasekom.
 Turkmen. (Tashiev), 20-35
Trigonoseelus gigas
T. punctipletus
 all from Turkmenia
- Physaloptera sp. 2
 Mushkambayeva, M. G., 1973, Ekol. Nasekom.
 Turkmen. (Tashiev), 20-35
Aesmia servillei Schatzmayr
Trigonoseelus gigas
T. punctipletus
Pisterotarsa gigantea subsp. *zoubkoffi*
P. kessleri
 all from Turkmenia
- Physaloptera (Physaloptera) sp.
 Peters, W.; et al., 1973, Tr. Roy. Soc. Trop.
 Med. and Hyg., v. 67 (1), 3-4 [Demonstration]
Callioscirtus nigrovittatus (mucosa of stom-
 ach): Sabah
- Physaloptera sp.
 Singh, M.; and Cheong Chee Hock, 1971, South-
 east Asian J. Trop. Med. and Pub. Health, v. 2
 (4), 516-521
Rattus rattus argentiventer
R. r. rumpia
R. bowersi
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
 all from Malaysia
- Physaloptera sp.
 Thornton, J. E.; Bell, R. R.; and Reardon,
 M. J., 1974, J. Wildlife Dis., v. 10 (3),
 232-236
Canis latrans: Nueces County, Texas
- Physaloptera sp.
 Yonders, P. C.; and Dixon, C. F., 1977, J.
 Alabama Acad. Sc., v. 48 (3), 55-56 [Abstract]
 seasonal incidence rates
Crotaphytus collaris collaris: Baxter,
 Boone, Carroll, Izard, Marion, and Newton
 counties, Arkansas
- Physaloptera accipiteri sp. nov., illus.
 Kumar, P.; and Gupta, S. P., 1977, Current Sc.,
 Bangalore, v. 46 (14), 490-491 [Letter]
Accipiter badius (gizzard): Lucknow
- Physaloptera alata alata
 Vaidova, S. M., 1975, Izvest. Akad. Nauk
 Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
 distribution of avian helminths in relation
 to habitat zones (high mountain, mountain
 forest, forest and scrub, lowlands):
 Azerbaidzhan
- Physaloptera alata var. *chevreuxi* Sourat, 1914,
 illus.
 Sanchez-Acedo, C.; and Vericad, J. R., 1974,
 Rev. Iber. Parasitol., v. 34 (3-4), 197-203
Buteo buteo: Aragon Pyrenees
- Physaloptera anadonta Schaldybin, 1960
 Babayev, Ia.; and Kolodenko, A. I., 1975, Iz-
 vest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk
 (4), 71-75
 [*Hemiechinus auritus*]
 [*H. hypomedas*]
 all from Turkmenistan

- Physaloptera hispida*
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (stomach): Florida
- Physaloptera maxillaris* Molin, illus.
Cawthorn, R. J.; and Anderson, R. C., 1976, Canad. J. Zool., v. 54 (3), 313-323
Physaloptera maxillaris, description of development and stages in *Mephitis mephitis* (exper.), attempted experimental infection of other definitive hosts produced no patent infections but worms were found in *Canis familiaris* and *Mustela furo* (*Felis catus*, *Procyon lotor*, *Rattus norvegicus*, *Mustela vison* all negative), experimental investigation of possible paratenic hosts (*Rana pipiens*, *Thamnophis sirtalis*, *Mus musculus*, *Gallus* sp.)
- Physaloptera maxillaris* Molin
Cawthorn, R. J.; and Anderson, R. C., 1976, Canad. J. Zool., v. 54 (4), 442-448
Physaloptera maxillaris larvae in *Acheta pennsylvanicus*, effects of temperature, age of host, and previous infection on parasite development; resulting infectivity to *Mephitis mephitis*
- Physaloptera maxillaris* Molin
Cawthorn, R. J.; and Anderson, R. C., 1976, Canad. J. Zool., v. 54 (4), 522-525
Physaloptera maxillaris in *Mephitis mephitis*, seasonal distribution of adults and third- and fourth-stage larvae, analysis of data relates various stages found to feeding behavior of skunk throughout year, third-stage larvae determined to be overwintering stage: Guelph, Ontario
- Physaloptera myotis* (Babos, 1954), larvae, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Myotis oxygnathus
M. myotis
Rhinolophus hipposideros
Miniopterus schreibersi
(stomach wall of all): all from Moldavia
- Physaloptera myotis* Babos, 1954
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Nyctalus noctula: Moldavia
- Physaloptera praeputialis* Von Linstow, 1919
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 267-271
Felis sylvestris (estomago): Huesca, Alto Aragon
- Physaloptera rara*
Barnstable, R. W.; and Dyer, W. G., 1974, Tr. Illinois State Acad. Sc., v. 67 (4), 451-460
Procyon lotor (stomach and small intestine): southern Illinois
- Physaloptera rara* or *Gnathostoma procyonis*
Bartsch, R. C.; and Ward, B. C., 1976, Vet. Path., v. 13 (4), 241-249
"may have been either *Gnathostoma procyonis* or *Physaloptera*"
raccoons (gastric mucosa): southeastern Florida
- Physaloptera rara* Hall & Wigdor 1918
Gilbertson, D. E., 1977, J. Parasitol., v. 63 (1), 162-163
Vulpes fulva (stomach): Dakota County, Minnesota
- Physaloptera retusa*
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus magister (cecum)
Sceloporus occidentalis (stomach)
Sceloporus undulatus (cecum)
all from Great Basin and Upper Colorado Plateau, Utah
- Physaloptera retusa*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Physaloptera* (*Physaloptera*) *sibirica* Petrow et Gorbunow, 1931, illus.
Quentin, J. C.; and Biocca, E., 1976, Ann. Parasitol., v. 51 (2), 255-259
description
Eliomys quercinus (tube digestif): Hautes-Alpes, sur le versant francais, dans les regions de Meyries et de St-Veran, et sur le versant italien dans le Parc national "del Gran Paradiso", province d'Aoste, val di Cogne, Lonson
- Physaloptera turgida*
Prestwood, A. K.; Nettles, V. F.; and Farrell, R. L., 1977, Am. J. Vet. Research, v. 38 (4), 529-532
Didelphis marsupialis: Georgia
- Physaloptera turgida*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Physaloptera turgida*
Young, P. L.; and Babero, B. B., 1975, Proc. Oklahoma Acad. Sc., v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)

- Physalopterata Sobolev, 1949
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of Pentadentoptera Schachnasarova, 1949
- Physalopterid larva
Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, J. Parasitol., v. 61 (3), 547-548
Grus canadensis pratensis (lower small intestine): Florida
- Physalopteridae (Railliet, 1893, subfam.) Leiper, 1908
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopteroidea
key to subfamilies
includes: Thubunaeinae; Proleptinae;
Physalopterinae
- Physalopteridae [sp.], probably Physaloptera sp., illus.
Nicolaidis, N. J.; et al., 1977, Pathology, v. 9 (2), 129-135
Physalopteridae [sp.], probably Physaloptera sp. causing small bowel infarction in 11-month-old infant, infection thought to have resulted from ingestion of insects on grass eaten while at play in an area contaminated by bandicoots (probable definitive host); pathology resulted when larvae attempted tissue migration for re-encystment in a foreign host, clinical report: Queensland, Australia
- Physalopterinae Railliet, 1893
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopteridae
key; key to genera
includes: Pseudabbreviata; Skrjabinoptera; Abbreviata; Pseudophysaloptera; Physaloptera; Turgida; Pentadentoptera
- Physalopterinae
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
Physalopteridae
key to genera from fishes
- Physalopteroidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Physalopteroidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
includes: Physalopteridae
- Physalopteroides Wu & Liu, 1940, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thubunaeinae
key
- Physalopteroides geckonis n. sp., illus.
Bilqees, F. M.; and Siddiqui, M. H., 1975, Pakistan J. Scient. and Indust. Research, v. 18 (6), 261-264
Gecko sp. (intestine): Karachi
- Physocephaloides Maplestone, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Physocephalus Diesing, 1861
- Physocephalus Diesing, 1861, illus.
1 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascaropsinae
key; synonymy
- Physocephalus quadrialatus Kirschenblat, 1949, illus.
Meszaros, F., 1977, Acta Zool. Acad. Scient. Hungar., v. 23 (1-2), 133-138
description
Cricetus cricetus (stomach): Hungary
- Physocephalus sexualatus Molin, 1915, illus.
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972, Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera beetles, possible importance in biological control
Blaps polychresta (haemocoelae): Cairo, Egypt
Scarabaeus sacer: Abu-Rawash, Cairo, Egypt
- Physocephalus sexualatus (Molin, 1860) larva
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Charadrius hiaticula: Keta lake
- Physocephalus sexualatus
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
Sus scrofa domesticus x Sus scrofa cristatus (stomach): Aransas National Wildlife Refuge, southern Texas
- Physocephalus sexualatus
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (stomach): Georgia
- Physocephalus sexualatus
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (stomach): Florida
- Physocephalus sexualatus
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile

- Physocephalus sexalatus*
Peterson, P. M.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1778-1780
Ascarops strongylina, *Physocephalus sexalatus*, *Hyostrongylus rubidus*, natural incidence, comparison with earlier surveys hogs (stomachs): Georgia
- Physocephalus sexalatus* (Molin, 1860) larvae, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93
description, geographic distribution
Syn.: *Agamospirura*, nec *Skarbilovitsch*, 1946, Henry et Sisoff, 1912, n. syn.
Eptesicus serotinus
Vespertilio murinus
Pipistrellus pipistrellus
P. nathusii
P. kuhli
Myotis daubentoni
M. mystacinus
Barbastella barbastella
Nyctalus noctula
N. leisleri
all from Moldavia
- Physocephalus sexalatus* (Molin, 1860)
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Rhinolophus hipposideros
Myotis oxygnathus
M. dasycneme
M. daubentoni
M. bechsteini
M. nattereri
M. mystacinus
Plecotus auritus
Barbastella barbastella
Nyctalus leisleri
N. noctula
Eptesicus serotinus
all from Moldavia
- Physocephalus sexalatus*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (26), 123-128
[pig, wild]: Primorskii krai
- Physocephalus sexalatus*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Aphodius fimetarius
A. lungens
Geotrupes impressus
Copris lunaris
Gymnopleurus aciculatus
Onthophagus amyntas
all from Uzbekistan
- Physocephalus sexalatus*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago): *Planta Faenadora de Carnes Socoagro, Valdivia, Chile*
- Physocephalus sexalatus dromedarii* Muschkambarowa, 1967
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Scarabaeus sacer
Copris lunaris
Adesmia biseriata
Cerambycidae
all from Uzbekistan
- Piayussunema* [sic] Kohn, Gomes & Motta, 1968
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Ancyracanthus* Diesing, 1858
- Pintonema* Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Moennigia* Travassos, 1935
- Pintonema*
Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
as syn. of *Moennigia*
- Piscillania* [sic] Yeh, 1960
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Oncophora* Diesing, 1851
- Pithecostrongylus* Lubimov, 1930
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Placentonema* Gubanov, 1951, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Crassicaudinae
key
- Placoconus lotoris* (Schwartz, 1925) Webster, 1956
Balasingam, E., 1964, Med. J. Malaya, v. 19 (1), 46-47
growth and morphology of *Placoconus lotoris*
- Placoconus lotoris*
Barnstable, R. W.; and Dyer, W. G., 1974, Tr. Illinois State Acad. Sc., v. 67 (4), 451-460
Procyon lotor (small intestine): southern Illinois
synonymy
- Plectostrongylus* sp. (undescribed)
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (lung parenchyma): Powelltown, Victoria

- Plicatolabia Mozgovoi*, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Heterocheilus* Diesing, 1839
- Pneumospirura* Wu & Hu, 1938, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Pneumospiruridae
key
- Pneumospirura*
Pence, D. B.; and Stone, J. E., 1977, J. Parasitol., v. 63 (6), 979-991
systematics of family Pneumospiruridae
- Pneumospirura*
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
as syn. of *Metathelazia*
- Pneumospirura bassaris* sp. n., illus.
Pence, D. B.; and Stone, J. E., 1977, J. Parasitol., v. 63 (6), 979-991
pathology
Bassariscus astutus (primary and secondary bronchioles of lungs): Davis Mountains, 9.5 km north of Fort Davis, Jeff Davis Co., Texas
- Pneumospirura rodentium* n. sp., illus.
Wertheim, G.; and Giladi, M., 1977, Ann. Parasitol., v. 52 (6), 643-646
Gerbillus dasyurus
Meriones crassus
(lung parenchyma of all): all from Israel [and/or] Southern Sinai
- Pneumospirura rodentium*
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
as syn. of *Metathelazia rodentium* (Wertheim and Giladi, 1977) n. comb.
- Pneumospiruridae Wu & Hu, 1938
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelazioidea
key; key to genera
includes: *Pneumospirura*; *Metathelazia*; *Vogeloides*
- Pneumospiruridae
Pence, D. B.; and Stone, J. E., 1977, J. Parasitol., v. 63 (6), 979-991
systematics of family Pneumospiruridae
- Pneumospiruridae
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
Thelazioidea
cephalic structures, scanning electron microscopy, results used in revision of family
includes: *Vogeloides*; *Metathelazia*
- Pneumostrongylus calcaratus* Monnig, 1932, illus.
Anderson, I. G. H., 1976, J. South African Vet. Med. Ass., v. 47 (1), 23-27
morphology of 1st, 2nd, and 3rd stage larvae, exper. transmission to impala unsuccessful
Urocyclus flavescens (nat. and exper.):
Nyala game ranch
impala: Nyala game ranch
- Pneumostrongylus calcaratus* Monnig
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (lungs): Kenya
- Pneumostrongylus cornigerus* Ortlepp, 1962
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damaliscus dorcas dorcas: captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Pneumostrongylus tenuis*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (meningeal surface of brain): Kentucky
- Poekilostrongylus* Schmidt et Whittaker, 1975
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Polycaecum* Maplestone, 1930
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Multicaecum* Baylis, 1923
- Polydelphinae Mozgovoi & Shakhmatova, 1971
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Ophidascaridinea* (Hartwich, 1954, subfam.) Chabaud, 1965
- Polydelphis* Dujardin, 1845
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ophidascaridinea
key
- Polydelphis anoura*
McKenzie, R. A.; and Green, P. E., 1976, J. Wildlife Dis., v. 12 (3), 405-408
Morelia spilotes variegata (intestine)
- Polydelphis attenuata* (Duj., 1845)
Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, Dobuts. Zasshi, Tokyo, v. 84 (3), 258-261
seasonal differences of infection rate, worm burden
Vipera russelli: West Bengal villages

- Pontochoa* gen. nov.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 30-35
Rhabdochonidae, Rhabdochoninae
tod: *P. dorabi* sp. nov.
- Pontochoa* Mamaev, 1968
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Heptochona* Rasheed, 1965
- Pontochoa caudata* gen. et sp. nov., illus.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 30-35
Thunnus thynnus (intestine): South China Sea
- Pontochoa caudata* gen. et sp. nov. [nom. nud.]
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Thunnus thynnus (intestine): South China Sea
- Pontochoa dorabi* gen. et sp. nov. (tod), illus.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 30-35
Chirocentrus dorab (intestine, stomach):
South China Sea
- Popovastrongylus* n. gen.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
Trichonematidae
key to species, tod: *P. wallabiae* (Johnston & Mawson) [n. comb.]
Syn.: *Macropostrongylus* Yorke & Maplestone P.P.
- Popovastrongylus irma* n. sp., illus.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
key
Macropus irma (stomach): Perth, W. Australia
- Popovastrongylus pearsoni* (Johnston & Mawson) [n. comb.], illus.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
key, description, syn.: *Macropostrongylus pearsoni* Johnston & Mawson, 1940
Macropus eugenii: Kangaroo I., S. Australia
M. rufogriseus: Launceston, Tasmania
- Popovastrongylus wallabiae* (Johnston & Mawson) [n. comb.] (tod), illus.
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (2-4), 51-62
key, syn.: *Macropostrongylus wallabiae* Johnston & Mawson, 1939; *Gelanostrongylus wallabiae*: Popova, 1952
Macropus rufogriseus: Logan Village, Queensland; Launceston, Tasmania
- Porrocaecum*
Allison, V. F.; et al., 1973, Tr. Am. Micr. Soc., v. 92 (2), 291-297
Syn.: *Sulcascaris* Hartwich (1957)
- Porrocaecum* Railliet & Henry, 1912, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Toxocarinae
key; synonymy
- Porrocaecum* (Laymanicaecum) Mozgovoi, 1951, in part
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Porrocaecum* Railliet & Henry, 1912
- Porrocaecum*
Myers, B. J., 1976, Tr. Am. Micr. Soc., v. 95 (2), 137-142
human anisakiasis, historical review
- Porrocaecum* sp., illus.
Bakke, T. A.; and Barus, V., 1975, Norwegian J. Zool., v. 23 (3), 183-191
seasonal occurrence, age of host, morphological features
Larus canus (alimentary canal): Agdenes area, Norway
- Porrocaecum* sp.
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
nematodes of *Larus canus*, age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Porrocaecum* sp.
George, R. R.; and Bolen, E. G., 1975, J. Wildlife Dis., v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas
- Porrocaecum* sp. 1 larvae
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela putorius
Meles meles
(gastrointestinal tract of all): all from Karelia
- Porrocaecum* sp. 2
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes (intestine): Karelia
- Porrocaecum* (*Porrocaecum*) *accipiteri* sp. nov., illus.
Kumar, P.; and Gupta, S. P., 1977, Current Sc., Bangalore, v. 46 (14), 491-492 [Letter]
Accipiter badius (intestine): Lucknow
- Porrocaecum americanum*
Anderson, M. M.; and McDaniel, J. S., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
Blarina brevicauda: eastern North Carolina
- Porrocaecum angusticolle*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan

- Porrocaecum crassum (Deslongchamps, 1824)
Railliet et Henry, 1912
Kamburov, P.; and Vasilev, I., 1972, Izvest.
Sentral. Khelmit. Lab., v. 15, 109-133
Anas platyrhynchos (small intestine): Bulgaria
- Porrocaecum encapsulatum
Anderson, M. M.; and McDaniel, J. S., 1975, J.
Elisha Mitchell Scient. Soc., v. 91 (2), 73
Blarina brevicauda: eastern North Carolina
- Porrocaecum ensicaudatum (Zeder, 1800), illus.
Bakke, T. A.; and Barus, V., 1975, Norwegian
J. Zool., v. 23 (3), 183-191
seasonal occurrence, measurements, sex and
age of host
Larus canus (corneous membrane in gizzard,
alimentary canal): Agdenes area, Norway
- Porrocaecum ensicaudatum, larvae
Bakke, T. A.; and Barus, V., 1976, Norwegian
J. Zool., v. 24 (1), 7-31
nematodes of Larus canus, age and sex of
host, seasonal variations, distribution in
alimentary canal: Agdenes, Norway
- Porrocaecum ensicaudatum (Zeder, 1800)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 20, 35-45
Actitis hypoleucos
Calidris minuta
all from Keta lake
- Porrocaecum ensicaudatum
Cooper, C. L.; and Crites, J. L., 1974, J.
Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (intestine): South Bass
Island, Ohio
- Porrocaecum ensicaudatum
Cooper, C. L.; and Crites, J. L., 1974, J.
Wildlife Dis., v. 10 (4), 399-403
survey, helminths of red-winged blackbirds
including a check list of previous findings
Agelaius phoeniceus (intestine): South Bass
Island, Ohio
- Porrocaecum ensicaudatum (Zeder, 1800)
Cooper, C. L.; and Crites, J. L., 1974, Proc.
Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (intestine):
South Bass Island, Ottawa County, Ohio
- Porrocaecum ensicaudatum
Cooper, C. L.; and Crites, J. L., 1976, J.
Parasitol., v. 62 (1), 105-110
similarity index of helminth faunas of 7
passerine bird species, index of association
of 10 species of helminths identified as hav-
ing foci of infection, competition for inver-
tebrate food resources and aggregation into
mixed feeding flocks maximizes transmission:
South Bass Island, Ottawa County, Ohio
- Porrocaecum ensicaudatum Baylis, 1920 (larvae)
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint.
Lab., Akad. Nauk SSSR, v. 17, 95-103
Rattus norvegicus (body cavity): Karelia
- Porrocaecum ensicaudatum (Zeder, 1800)
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus: Azov Sea
L. ridibundus: Tuva
Sterna paradisea: Yenisei
Stercorarius longicaudatus: Yenisei
- Porrocaecum ensicaudatum
Vaidova, S. M., 1975, Izvest. Akad. Nauk
Azerbaijdzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation
to habitat zones (high mountain, mountain
forest, forest and scrub, lowlands):
Azerbaijdzhan
- Porrocaecum ensicaudatum (Zeder, 1800)
Young, P. L.; and Babero, B. B., 1975, Proc.
Oklahoma Acad. Sc., v. 55, 169-174
helminthic diseases, cockroaches may play
an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Porrocaecum heteroura (Creplin, 1829)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 20, 35-45
Pluvialis apricaria altifrons
Philomachus pugnax
Limosa limosa lapponica
all from lower Yenisei
- Porrocaecum heteroura (Creplin, 1929), illus.
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 20, 146-155
description
Sterna paradisea
Stercorarius longicaudatus
S. pomarinus
all from Yenisei
- Porrocaecum sulcatum (Rudolphi, 1819), illus.
Allison, V. F.; et al., 1973, Tr. Am. Micr.
Soc., v. 92 (2), 291-297
redescription
Chelone mydas (stomach): marine waters of
North Carolina
- Porrocaecum sulcatum (Rud. 1819), illus.
Boero, J. J.; and Led, J. E., 1974, Rev. Agron.
y Vet., v. 3 (1), 16-17
description
Thalassochelys caretta
(duodeno, intestino delgado): Jardin Zoo-
logico, La Plata, Argentina (captured near
Mar del Plata)
- Porrocaecum talpae (Schrank, 1788)
Mas-Coma, S.; and Gallego, J., 1975, Rev.
Iber. Parasitol., v. 35 (3-4), 261-281
Sorex araneus (cavidades abdominal y toraci-
ca): Catalan Pyrenean Mountains
- Poteriostomum
Colglazier, M. L.; Enzie, F. D.; and Kates, K.
C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, com-
parative efficacy of 4 benzimidazoles eval-
uated by critical test method

- Potterostomum**
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Potterostomum imparidentatum**
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Potterostomum imparidentatum**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Potterostomum imparidentatum**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Potterostomum ratzii**
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Potterostomum ratzii**
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
Cyathostominae in horses (large intestine), prevalence, relative abundance, site distribution, seasonal variation, epidemiological implications: south-west England
- Primasubulura distans**, illus.
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Prionoderma Rudolphi**, 1810, nec Cuvier, 1817
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Goezia Zeder*, 1800
- Pristionchus uniformis Fedorko et Stanuszek**
Bajan, C.; et al., 1976, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (3), 171-173
Pristionchus uniformis, influence on Colorado beetle, reduced number of eggs laid, protracted period of oviposition, reduced hatchability
- Probstmayria Ransom 1907**
File, S. K., 1976, J. Parasitol., v. 62 (2), 256-258
key to species, includes: *Probstmayria vivipara*; *P. simiae*; *P. gorillae*; *P. suis*; *P. gombensis*
- Probstmayria**
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Probstmayria**
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Probstmayria gombensis** sp. n., illus.
File, S. K., 1976, J. Parasitol., v. 62 (2), 256-258
key
Pan troglodytes schweinfurthii (intestine):
Gombe National Park, Tanzania
- Probstmayria gombensis** File
File, S. K.; McGrew, W. C.; and Tutin, C. E. G., 1976, J. Parasitol., v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces):
Gombe National Park, Tanzania
- Probstmayria vivipara**
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Probstmayria vivipara**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Probstmayria vivipara**
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency

- Probstmayria vivipara*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974,
Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Procammallanus Baylis* (1923)
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
key to species from fishes of Pakistan and
India, includes *P. aspliculus* Khera, 1955;
P. daccai Gupta, 1959; *P. heteropneustus*,
Ali, 1956; *P. planoratus* Kulkarni, 1935; *P.*
muelleri Agrawal, 1966; *P. mehrii* Agrawal
1930; *P. attui* Pande et al., 1963; *P. walla-*
gus Rehana and Bilquees, 1973; *P. bagarii*
Karve and Naik, 1951; *P. globoconchus* Ali,
1960; *P. gubernaculus* Khera, 1955; *P. vitta-*
tusi Sood, 1967; *P. alii* Kalyankar, 1971;
P. berdii (Khan and Yaseen, 1969); *P. vachai*
(Sinha and Sahay, 1965); *P. sparus*, sp. nov.;
P. dussumieri Bilquees et al. 1971; and *P.*
singhi Ali, 1956
Syn.: *Spirocammallanus* Olsen (1952)
- Procammallanus Baylis*, 1923, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Para-*
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Camallanidae
key; synonymy
- Procammallanus* (Baylis, 1923) Ali, 1956
Gupta, N. K.; and Duggal, C. L., 1973, *Riv.*
Parassitol., Roma, v. 34 (4), 295-304
key to the species of subgenus, includes:
P. (P.) glossobii (Pearse, 1933); *P. (P.)*
spiculogubernaculus Agarwal, 1958; *P. (P.)*
annulatus Yamaguti, 1934; *P. (P.) laevicon-*
chus (Wedl, 1862) Baylis, 1923; *P. (P.)*
brevis Kung, 1948; *P. (P.) bilaspurensis* n.
sp.; *P. (P.) heteropneustus* Ali, 1956; *P.*
(P.) clarius Ali, 1956; *P. (P.) attui* Pande,
Bhatia and Rai, 1962; *P. (P.) sphaeroconchus*
Tornquist, 1931; *P. (P.) sigami* Yamaguti,
1935; *P. (P.) lonis* Yamaguti, 1941; *P. (P.)*
gubernaculus Khera, 1955; *P. (P.) bagarii*
Karve and Naik, 1951; *P. (P.) vittatusi* Sood,
1967; *P. (P.) iheringi* Travassos et al.,
1928; *P. (P.) rarus* Travassos et al., 1928;
P. (P.) monotaxis Olsen, 1952; *P. (P.) mur-*
rayensis Johnston and Mawson, 1940; *P. (P.)*
mazabukae Yeh, 1957; *P. (P.) pereirai* Anner-
eaux, 1946; *P. (P.) amarali* Vaz and Pereira,
1934; *P. (P.) spiralis* Baylis, 1923; *P. (P.)*
fariasi Pereira, 1935; *P. (P.) hilarii* Vaz
et Pereira, 1934; *P. (P.) barroslimai* Perei-
ra, 1935; *P. (P.) viviparus* Ali, 1956;
P. (P.) ophicephalus Ali, 1960; *P. (P.) glo-*
boconchus Ali, 1960; *P. (P.) fulvidraconis*
Ali, 1935; *P. (P.) hyderabadensis* Ali, 1956;
P. (P.) singhi Ali, 1956
- Procammallanus bengalensis* new name for *P. (S[pi-*
rocammallanus?]) olsenii (Bashirullah, 1973) n.
comb., preoccupied
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
[as syn. of] *P. singhi* Ali, 1956
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus* (Procammallanus) *bilaspurensis* n.
sp., *illus.*
Gupta, N. K.; and Duggal, C. L., 1973, *Riv.*
Parassitol., Roma, v. 34 (4), 295-304
key
Mastacembelus armatus (intestine): Bilaspur
(H. P.), India
- Procammallanus daccai* Gupta, 1959
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
key
synonymy
- Procammallanus devendri* Sinha and Sahay, 1966
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
as syn. of *P. daccai* Gupta, 1959
- Procammallanus fasciatusi* Sood, 1967
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
as syn. of *P. gubernaculus* Khera, 1955
- Procammallanus* (P.) *globoconchus* Ali, 1960, *illus.*
Gupta, N. K.; and Duggal, C. L., 1973, *Riv.*
Parassitol., Roma, v. 34 (4), 295-304
description, key
Mystus seenghala (intestine): Jullundur
(PB.), India
- Procammallanus gomtii* Sood, 1967
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
as syn. of *P. gubernaculus* Khera, 1955
- Procammallanus gubernaculus* Khera, 1955
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
key
synonymy
- Procammallanus hindensis* [sic] Lal, 1965
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
as syn. of *P. daccai* Gupta, 1959
- Procammallanus* (S[pirocammallanus?]) *incarocae*
(Freitas and Ibanez, 1970) n. comb.
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus* (S[pirocammallanus?]) *istiblenni*
(Noble, 1966) n. comb.
Akram, M., 1975, *Biologia*, Lahore, v. 21 (2),
93-100
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus laeviconchus* (Wedl, 1861)
Khalil, L. F.; and Thurston, J. P., 1973,
Rev. Zool. et Botan. Africaines, v. 87 (2),
209-248
Synodontis victoriae: Entebbe, Lake Vic-
toria, Uganda
S. afro-fischeri: Entebbe, Lake Victoria,
Uganda
S. schall: Lake Albert, Uganda
(stomach of all)

- Procamallanus laeviconchus* (Weld, 1862)
Vassiliades, G., 1972, Bull. Inst. Fond. Afr.
rique Noire, s. A, v. 34 (3), 529-533
Clarias senegalensis: Sangalkam, Senegal
- Procamallanus magurii* Lal, 1965
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
as syn. of *P. daccii* Gupta, 1959
- Procamallanus* (S[*pirocammallanus*?]) *mazabukae*
(Yeh, 1957) n. comb.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procamallanus* (S[*pirocammallanus*?]) *olseni* (Bash-
irullah, 1973) n. comb., preoccupied, renamed
P. bengalensis new name
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
as syn. of *Procammallanus singhi* Ali, 1956
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus* (S[*pirocammallanus*?]) *olseni* (Cam-
pana-Rouget 1965) n. comb.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus* (S[*pirocammallanus*?]) *ompoci* (Ma-
jumdar and Datta, 1972) n. comb.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
as syn. of *P. gubernaculus* Khera, 1955
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus ottuei* sp. nov., illus.
Varma, S.; and Varma, S., 1971, Indian J.
Animal Research, v. 5 (1), 29-32
Heteropneustes fossilis (stomach): Ottu
reservoir, Sirsa (District Hissar), Haryana,
India
- Procammallanus singhi* Ali, 1956
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
key
synonymy
- Procammallanus sparus* sp. nov., illus.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
key
Sparus spinifer (intestine): Karachi Coast
- Procammallanus* (S[*pirocammallanus*?]) *timmi* (Bash-
irullah, 1973) n. comb.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
as syn. of *P. singhi* Ali, 1956
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procammallanus* (S[*pirocammallanus*?]) *vachai* (Sinha
and Sahay, 1965) n. comb.
Akram, M., 1975, Biologia, Lahore, v. 21 (2),
93-100
key
"The author supports Campana-Rouget (1961)
in suppressing the subgenera of the genus
Procammallanus . . ."
- Procyrnea* (Chabaud, 1958, subgen.), illus.
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Habroneematinae
key; synonymy
- Proleptinae* (Schulz, 1927)
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Physalopteridae
key; key to genera
includes: *Proleptus*; *Paraleptus*; *Helicone-*
ma; *Bulbocephalus*
- Proleptus* Dujardin, 1845, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 1-27
Proleptinae
key
- Proleptus* Dujardin, 1845
Specian, R. D.; Ubelaker, J. E.; and Dailey,
M. D., 1975, Proc. Helminth. Soc. Washington,
v. 42 (1), 14-21
Physalopteridae, *Physalopterinae*
synonymy, redescription, key, key to species,
includes: *P. obtusus* Dujardin, 1845; *P.*
problematicus Kries, 1940; *P. inflatus* (Lin-
stow, 1890); *P. acutus* Dujardin, 1845; *P.*
malayi Sandosham, 1954; *P. soridus* Lent et
Freitas, 1948; *P. trygonorrhonae* Johnston
and Mawson, 1943; *P. australis* Baylis, 1933;
P. africanus (Linstow, 1899)
- Proleptus acutus* Dujardin 1845
Dailey, M. D.; and Carvajal, J., 1976, J. Para-
sitol., v. 62 (6), 939-942
Rhinobatos planiceps: Juan Lopez Beach, An-
tofagasta, Chile
- Proleptus acutus* Dujardin, 1845, illus.
Specian, R. D.; Ubelaker, J. E.; and Dailey,
M. D., 1975, Proc. Helminth. Soc. Washington,
v. 42 (1), 14-21
synonymy, redescription, key
Platyrrhinoides triseriata
Heterodontus francisci
Mustelus henlei
M. californicus
all from Pacific waters off the southern
California coast
- Proleptus anabantis* Pearse, 1933
Specian, R. D.; Ubelaker, J. E.; and Dailey,
M. D., 1975, Proc. Helminth. Soc. Washington,
v. 42 (1), 14-21
species inquirenda
- Proleptus coronatus* (Beneden, 1858)
Specian, R. D.; Ubelaker, J. E.; and Dailey,
M. D., 1975, Proc. Helminth. Soc. Washington,
v. 42 (1), 14-21
species inquirenda

- Proleptus dogiyeli* Osmanov, 1940
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus acutus* Dujardin, 1845
- Proleptus elegans* (Orley, 1885)
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
species inquirenda
- Proleptus gordiodes* [sic] (Beneden, 1858)
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
species inquirenda
- Proleptus rajae* (Diesing, 1851)
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
species inquirenda
- Proleptus robustus* (Beneden, 1871) Seurat, 1916
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus acutus* Dujardin, 1845
- Proleptus tortus* (Linstow, 1906)
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
species inquirenda
- Proleptus urolophi* Johnston and Mawson, 1951
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Heliconema urolophi* (Johnston and Mawson, 1951) comb. nov.
- Prosthecosacter alatus* (Leuckart) Diesing 1851
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus alatus* (Leuckart 1848) Stiles and Hassall 1905
- Prosthecosacter convolutus* (Kuhn) Diesing 1851
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829) Baylis and Daubney 1925
- Prosthecosacter inflexus* (Rudolphi) Diesing 1851
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pseudalius inflexus* (Rudolphi 1808) Schneider 1866
- Prosthecosacter minor* (Kuhn) Diesing 1851
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis and Daubney 1925
- Prosungulonema* Roytman, 1963
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Johnstonmawsonia* Campana-Rouget, 1955
- Protospirura* Seurat, 1914, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruridae
key
Syn.: *Thaprospirura* Sood & Parshad, 1974
- Protospirura* sp.
O'Farrell, T. P., 1975, Am. Midland Naturalist, v. 93 (2), 377-387
Perognathus parvus
Peromyscus maniculatus
all from Arid Lands Ecology Reserve, Benton County, Washington
- Protospirura-Mastophorus* spp.
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus rattus argentiventer
R. r. rumpia
R. bowersi
R. canus
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
R. whiteheadi
all from Malaysia
- Protospirura chanchanensis* sp. n., illus.
Ibanez, N., 1966, Bol. Chileno Parasitol., v. 21 (2), 34-37
Rattus rattus norvegicus (estomago): Trujillo, Peru
- Protospirura muricola*, illus.
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Protospirura muris*, illus.
Taniguchi, M.; et al., 1977, Bull. Coll. Agric. and Vet. Med., Nihon Univ. (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo

- Protospirura numidica criceticola* Quentin, Kari-mi, and Rodriguez De Almeida, 1968
 Healey, M. C.; and Grundmann, A. W., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 59-63
Protospirura numidica criceticola, influence of intermediate hosts on infection pattern in *Peromyscus maniculatus sonoriensis*, yearly infections seasonal in desert habitats, low steady rate in mountains
Peromyscus maniculatus sonoriensis (nat. and exper.): Bonneville Basin, Utah
Eleodes tuberculata patruelis (exper.)
Melanoplus femur-rubrum (exper.)
M. atlantis (exper.)
Gryllus pennsylvanicus (exper.)
- Protostrongylid[ae]
 Maklakova, L. P., 1975, Trudy Gel'mint Lab., Akad. Nauk SSSR, v. 25, 102-106
 distribution of mollusc intermediate hosts in relation to habitat
- Protostrongylidae [sp.], illus.
 Lankester, M. W.; Crichton, V. J.; and Timmermann, H. R., 1976, Canad. J. Zool., v. 54 (5), 680-684
 1st stage protostrongylid larvae in *Rangifer tarandus caribou* (feces) may be *Elaphostrongylus* sp., lack of pathogenic effects seems to rule out *Parelaphostrongylus tenuis*: northwestern Ontario and Manitoba
- Protostrongylus
 Dzhabbarov, D. G., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (4), 90-92
 Protostrongylus, sheep, seasonal and age dynamics of infection: Lesser Caucasus
- Protostrongylus
 Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
 sheep helminths, effective control with Nil-vern and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Protostrongylus
 Melikov, Iu. F., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 80-82
 distribution in sheep of sheep farms of Kura-Araksinsk lowland, Azerbaidzhan
- Protostrongylus
 Nemeseri, L., 1976, Magy. Allat. Lapja, v. 98, v. 32 [i. e. 31] (7), 459-461
Dictyocaulus filaria, trichostrongylids, *Fasciola hepatica*, sheep, successful treatment with combination of diamphenetide and tetramisole; no effect on Protostrongylidae
- Protostrongylus
 Paul, I.; et al., 1975, Lucrar. Stiint. Inst. Agron. I. Ionescu Brad Iasi, II. Zootehn., Med. Vet., 71-72
 Protostrongylus, *Muellerius*, *Cystocaulus*, bronchopneumonia, adult sheep, morphopathology, result of association of several species
- Protostrongylus spp.
 Aller, B.; and Aller, J. M., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 337-346
 nematodes of sheep lungs, lower incidence of fungi in parasitized lungs: Leon
- Protostrongylus [sp.]
 Helle, O., 1976, Norwegian J. Zool., v. 24 (4), 463 [Abstract]
 goats (feces): Norway
- Protostrongylus sp. probably *stilesi*
 Hibler, C. P.; et al., 1974, J. Wildlife Dis., v. 10 (1), 39-41
Protostrongylus sp., prob. *stilesi*, evidence of transplacental route of infection in big-horn sheep, larvae recovered from tissues of fetuses and neonatal lambs: Pike's Peak, Colorado
- Protostrongylus spp.
 Rojo Vazquez, J., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 51-101
 Protostrongylinae, frequency in ovine lungs, primary nodes are infected by bacteria at a lower rate than non-parasitized lung areas: municipal slaughterhouse, Leon
- Protostrongylus sp., illus.
 Summaliev, P., 1973, Izvest. Tsentral. Khel-mint. Lab., v. 16, 161-167
Neostrongylus linearis, *Muellerius capillaris*, *Cystocaulus ocreatus*, *Protostrongylus* sp., differential diagnosis of infective larvae based on biometric studies
Helicella obvia
Zebrina detrita
 all from Bulgaria
- Protostrongylus *capensis*
 Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damalisca dorcas dorcas: captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Protostrongylus *commutatus*
 Kutzer, E.; and Frey, H., 1976, Berl. u. Munchen. Tierarztl. Wchnschr., v. 89 (24), 480-483
Lepus europaeus: Austria
- Protostrongylus *commutatus*
 Kutzer, E.; and Frey, H., 1976, Ztschr. Parasitenk., v. 50 (2), 213-214
Lepus europaeus
- Protostrongylus *kochi*
 Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
 nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Protostrongylus *kochi*
 Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
 nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nil-vern to nematode incidence on pastures, three-year study, overall decrease

- Protostrongylus macrootis*
Greiner, E. C.; Worley, D. E.; and O'Gara, B. W., 1974, *J. Wildlife Dis.*, v. 10 (1), 70-73
Antilocapra americana (lungs): Yellowstone National Park, Wyoming and Montana
- Protostrongylus macrootis*
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, *Canad. J. Zool.*, v. 54 (3), 307-312
helminths of Alces alces, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Protostrongylus rufescens* (Leuckart, 1865)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (trachea, bronchi): vicinity of Nowy Targ, Carpathian Mountains
- Protostrongylus rufescens*
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*, v. 99 (7), 129
Dictyocaulus filaria, Protostrongylus rufescens, and Cystocaulus ocreatus in sheep, fenbendazole effectively eliminated lung-worms
- Protostrongylus rufescens*
Ferreira, L. D. B. B., 1975, *Rev. Portug. Cien. Vet.* (433-434), v. 70, 25-39
broncho-pulmonary nematodes of sheep, life cycles, epizootiology, instructive review for veterinarians
- Protostrongylus rufescens*
Goda, Fawzy F. M., 1974, *Bull. Epizoot. Dis. Africa*, v. 22 (1), 75-78
sheep: Benghazi, Libya
- Protostrongylus rufescens*
Ramisz, A.; Urban, E.; and Piechocki, B., 1975, *Med. Wet.*, v. 31 (11), 677-679
Protostrongylidae of sheep, tetramisole effective against all species except Muellerius capillaris
- Protostrongylus rufescens*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Protostrongylus rupicaprae*
Polley, L.; and Hoerning, B., 1977, *Rev. Suisse Zool.*, v. 84 (3), 675-680
Rupicapra rupicapra (tracheas and larger bronchi): Switzerland
- Protostrongylus rushi*
Forrester, D. J.; and Littell, R. C., 1976, *J. Wildlife Dis.*, v. 12 (1), 48-51
Protostrongylus stilesi, P. rushi, influence of rainfall on infection levels
Ovis c. canadensis (lungs): western Montana
- Protostrongylus shiozawai*, illus.
Shiozawa, M.; et al., 1975, *Bull. Nippon Vet. and Zotech. Coll.* (24), 76-86
Protostrongylus shiozawai in the Japanese serow, histopathology of verminous pneumonia
- Protostrongylus skrjabini* (Boev, 1936), illus.
Azimov, D. A.; et al., 1976, *Dokl. Akad. Nauk UzSSR* (10), 65-67
Protostrongylus skrjabini, life cycle
Helicella candaharica (exper.)
Subzebrinus albiplicatus (exper.)
[Ovis aries] (exper.)
[Capra hircus] (exper.)
[Citellus] (exper.)
[Cavia porcellus] (exper.)
- P[rotostrongylus] skrjabini
Isakova, D. T.; et al., 1976, *Dokl. Akad. Nauk UzSSR* (11), 59-60
P[rotostrongylus] skrjabini, sheep, changes in serum proteins during 90 day course of infection
- Protostrongylus stilesi*
Forrester, D. J.; and Littell, R. C., 1976, *J. Wildlife Dis.*, v. 12 (1), 48-51
Protostrongylus stilesi, P. rushi, influence of rainfall on infection levels
Ovis c. canadensis (lungs): western Montana
- Protostrongylus tauricus*
Kutzer, E.; and Frey, H., 1976, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 89 (24), 480-483
Lepus europaeus: Austria
- Protostrongylus terminalis* (Passerini, 1884)
Kamensky, 1905
Mozgovoi, A. A.; et al., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 95-103
Lepus timidus (lungs): Karelia
- Protrellina aurifluus*
Leong, L.; and Paran, T. P., 1966, *Med. J. Malaya*, v. 20 (4), 349
Periplaneta americana: Singapore
- Proyseria Petter*, 1959, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key
- Psammomermis* sp., illus.
Klein, M. G.; et al., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 235-236
Psammomermis sp. in Popillia japonica, possible biological control: Brattleboro, Vermont; Wallingford, Cheshire and New Haven, Connecticut
- Pseudabbreviata Lichtenfels & Quigley*, 1968, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key

Pseudaliinae

- Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
key to genera
- Pseudalius* Dujardin 1845
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
key
- Pseudalius alatus* (Leuckart) von Linstow 1888
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Pharurus alatus* (Leuckart 1848)
Stiles and Hassall 1905
- Pseudalius arcticus* (Cobb) von Linstow 1900
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden 1870) n. comb.
- Pseudalius bicostatus* von Linstow 1906
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829)
Baylis and Daubney 1925
- Pseudalius convolutus* (Kuhn) Schneider 1866
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829)
Baylis and Daubney 1925
- Pseudalius filum* Dujardin 1845
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Pseudalius inflexus* (Rudolphi 1808) Schneider 1866
- Pseudalius inflexus* (Rudolphi 1808) Schneider 1866, *illus*
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
synonymy; redescription
Phocoena phocoena: Bay of Fundy, Canada;
North Sea off Netherlands
- Pseudalius minor* (Kuhn) Schneider 1866
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis and Daubney 1925
- Pseudalius tumidus* Schneider 1866
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
as syn. of *Halocercus invaginatus* (Quekett 1841) Dougherty 1943
- Pseudamidostomum* Boulenger, 1926
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae, Epomidiostomatinae
- Pseudancyracanthus* Skrjabin, 1923
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cystidicola* Fischer, 1798
- Pseudanisakis* Layman & Borovkova, 1926
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
synonymy, diagnosis, history
proposed type species: *P. tricupola* nom. nov.
- Pseudanisakis* Yamaguti, 1941
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
as syn. of *Pseudanisakis* Layman & Borovkova, 1926
- Pseudanisakis*
Gibson, D. I.; and Taylor, A. L., 1976, *Parasitology*, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Pseudanisakis* (Layman & Borovkova, 1926, subgen.) *Mozgovoi*, 1951
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Acanthocheilidae
key; synonymy
- Pseudanisakis* Yamaguti, 1941, *nec* Layman & Borovkova, 1926
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Metanisakis* *Mozgovoi*, 1951
- Pseudanisakis* *baylisi* nom. nov., *illus*.
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
description, syns.: *Eustoma rotundatum* (Rud.) of Punt (1941); *E. truncata* van Beneden, 1871, of Hartwich (1957) (in part)
Raja clavata
R. montagui
all from English Channel
- Pseudanisakis* *rajae* Yamaguti, 1941, *sensu nov.*
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
synonymy
- Pseudanisakis* *rotundata* (Rud.) of *Mozgovoi* (1950, 1953), of Williams (1965), of Williams & Richards (1968), and of Williams et al. (1970)
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
as syn. of *Pseudanisakis* *tricupola* nom. nov.

- Pseudanisakis tricupola* nom. nov., illus.
Gibson, D. I., 1973, *J. Nat. Hist.*, v. 7 (3), 319-340
proposed for type species, redescription, first description of larvae, measurements
Syns.: *Ascaris rotundata* Rud. of von Linstow (1880), of Jagerskiold (1894); *Anisakis* (*Pseudanisakis*) *rotundata* (Rud.) of Layman & Borovkova (1926); *Anacanthocheilus rotundatus* (Rud.) of Wulker (1930), of Baylis (1939); *Pseudanisakis rotundata* (Rud.) of Mosgovi (1950, 1953), of Williams (1965), of Williams & Richards (1968), and of Williams et al. (1970); *Eustoma truncata* van Beneden of Hartwich (1957) (in part); possibly *Eustoma rotundata* (Rud.) of Polyanski (1955), of Campana-Rouget (1955), of Kreis (1958, adults only), of Myers (1959), of Berland (1961), and of Threlfall (1969)
Raja radiata: Icelandic waters
R. batis: Icelandic waters
R. brachyrops: north of Falkland Islands
- Pseudaprocta decorata* Li, 1933
Jaron, W., 1969, *Acta Parasitol. Polon.*, v. 16 (1-19), 1968-1969, 137-152
helminth fauna of adult swallows just returning from migration compared with young birds; dynamics of infection, species composition of helminths, various stages of nesting season
Corvus corax
Pica caudata
Delichon urbica
(body cavity of all): all from Poland
- Pseudocystidicola* Layman, 1933
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Ascarophis* van Beneden, 1871
- Pseudofilaria giraffae* n. sp., illus.
Shoho, C.; and Sachs, R., 1975, *Tropenmed. und Parasitol.*, v. 26 (4), 489-493
Giraffa camelopardalis (subcutis and ligamentum nuchae): Serengeti National Park in northern Tanzania, East Africa
- Pseudogoezia* subgen.
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
key
- Pseudoheligmosomum* Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, Pudicinae
- Pseudometabronema* Bogdanova, 1963
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Salvelinema* Trofimenko, 1962
- Pseudonymus hydrophili* (Galeb, 1878) Stiles et Hassall, 1905
Hristovski, N. D., 1972, *Acta Parasitol. Iugoslavica*, v. 3 (2), 109-115
Hydrous piceus: Jugoslaviya (Belgrad; Zagreb)
Hydrous aterrimus: Jugoslaviya (Bitola)
- Pseudonymus hydrophili*
Hristovski, N. D., 1973, *Acta Parasitol. Iugoslavica*, v. 4 (2), 87-91
Hydrous piceus
Hydrous aterrimus
all from Macedonia, Yugoslavia
- Pseudonymus islamabadi*
Hristovski, N. D., 1973, *Acta Parasitol. Iugoslavica*, v. 4 (2), 87-91
Hydrous piceus
Hydrous aterrimus
all from Macedonia, Yugoslavia
- Pseudophilometroides* Parukhin, 1966
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Philometroides* Yamaguti, 1935
- Pseudophysaloptera* Baylis, 1934, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key
- Pseudophysaloptera kahmanni* n. sp., illus.
Mas-Coma, S.; and Gallego, J., 1977, *Ann. Parasitol.*, v. 52 (1), 19-24
pathologic lesions, mode of attachment to host
Eliomys quercinus ophiusae (estomac): Ile de Formentera (Balears, Espagne)
- Pseudoproleptus* Khera, 1953, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key; synonymy
- Pseudoproleptus* Khera, 1955
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (1), 14-21
Physalopteridae, Physalopterinae
key
- Pseudoproleptus africanus* n. sp., illus.
Khalil, L. F., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 795-807
Mormyrops sp. (intestine): Ibembo, Zaire
- Pseudorhabdochona* Liu & Wu, 1941
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Rhabdochona* Railliet, 1916
- Pseudorhabdochona* Liu et Wu, 1941
Rehana, R.; and Bilqees, F. M., 1976, *Agric. Pakistan*, v. 26 (4), 1975, 521-528
as syn. of *Rhabdochona* Railliet, 1916
- Pseudorictularia* Dollfus & Desportes, 1944
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
"seem to belong to the Physalopteroidea but, as yet, are not well enough known to be classified"

- Pseudorictularia* Dollfus & Desportes, 1945
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
"insufficiently known," "excluded from the key."
- Pseudostenurus* Yamaguti 1951
Arnold, P. W.; and Gaskin, D. E., 1975, *Canad. J. Zool.*, v. 53 (6), 713-735
key
- Pseudostenurus* sp.
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Phocoena phocoena: insular Newfoundland and its adjacent waters
- Pseudostertagia* Orloff, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, Libyostromylinae
- Pseudoterranova* Mozgovoi, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
"excluded from the key since . . . incompletely described"
- Psyllotylenchus*, n. gen.
Poinar, G. O., jr.; and Nelson, B. C., 1973, *J. Med. Entom.*, v. 10 (4), 349-354
Allantonematidae, tod: *P. viviparus* n. sp.
- Psyllotylenchus pawlowskyi* (Kurochkin, 1960)
n. comb.
Poinar, G. O., jr.; and Nelson, B. C., 1973, *J. Med. Entom.*, v. 10 (4), 349-354
Syn.: *Heterotylenchus pawlowskyi* Kurochkin, 1960
- Psyllotylenchus viviparus* n. sp. (tod), illus.
Poinar, G. O., jr.; and Nelson, B. C., 1973, *J. Med. Entom.*, v. 10 (4), 349-354
Monopsyllus wagneri: Mono County, Tom's Place, California
Catallagia sculleni rutherfordi: Plumas County, Lake Almanor, Main Campground, California
Catallagia sp.: Modoc County, Hackamore, California
Monopsyllus ciliatus protinus: Plumas county, Antelope Reservoir, Lone Rock Campground, California
Diamanus montanus: Plumas County, Lake Almanor, Main Campground and El Dorado County, Lake Tahoe, Sugar Pine Point State Park, California
- Pterygodermatites*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Pterygodermatites*
key
- Pterygodermatites* Wedl, 1861
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Rictulariidae
key
includes subgens.: Paucipectines; Neopaucipectines; *Pterygodermatites*; Mesopectines; Multipectines
- Pterygodermatites* (*Multipectines*) *affinis* (Jagerskiold, 1904), illus.
Quentin, J. C.; Seureau, C.; and Vernet, R., 1976, *Ann. Parasitol.*, v. 51 (1), 51-64
Pterygodermatites affinis, life cycle, development, morphology, cellular reaction in insect host
Tachyderma hispida (intra-epitheliale dans la paroi de l'intestin posterieur en arriere des tubes de Malpighi): Beni-Abbes, Algeria
chat (feces) (exper.)
Locusta migratoria (partie anterieure de l'ileon, au niveau de la vulve proctodeale, et dans l'epaisseur de la paroi du tube digestif) (exper.)
- Pterygodermatites dipodomis* (Tiner, 1948)
King, S. R.; and Babero, B. B., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 241-248
Dipodomys merriami
D. deserti
D. microps
(small intestine of all): all from Nevada
- Pterygodermatites hispanica* Quentin, 1973, illus.
Quentin, J. C.; and Seureau, C., [1975], *Ann. Parasitol.*, v. 49 (6), 1974, 701-719
Pterygodermatites hispanica, life cycle and development in *Apodemus sylvaticus* (exper.) and *Locusta migratoria* (exper.), morphology and organogenesis of larval stages, ontogenesis of cephalic structures, migration of larvae and cellular reactions in insect host
- Pterygodermatites hispanica*
Seureau, C.; and Quentin, J. C., 1977, *Ann. Parasitol.*, v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Pterygodermatites ondatrae*
Kinsella, J. M., 1974, *Am. Mus. Novitates* (2540), 1-12
Sigmodon hispidus (small intestine): Florida
- Pterygodermatites parkeri*
Davidson, W. R., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States

Pterygodermatites taterilli, illus.

Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
 comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction

Pterygodermatites (Mesopectines) *witenbergi* n. sp., illus.

Quentin, J. C.; and Wertheim, G., 1975, Ann. Parasitol., v. 50 (1), 63-85
Acomys cahirinus
Gerbillus dasyurus
G. pyramidum
G. gerbillus
Gerbillus sp.
Mus musculus
Sekeetamys calurus
 all from Israel

Pudica Travassos et Darriba, 1929 (type genus)
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Heligmonellidae, Pudicinae

Pudicinae (Skrjabin et Schikhobalova, 1952, tribu) Durette-Desset, 1971
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Heligmonellidae
 includes: *Pudica* (type genus); *Evandroia*; *Heligmostrongylus*; *Pseudoheligosomum*; *Sciurodendrium*

Pulchrostrongylus Travassos, 1935
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 as syn. of *Moennigia* Travassos, 1935

Pygarginema Kadenatsy, 1948, illus.
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 Ascaropsinae
 key

- Quasithelazia Maplestone, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Schistorophus Railliet, 1916
- Quentinstrongylus Durette-Desset, 1969
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Quentinstrongylus Durette-Desset, 1969
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Quimperia lanceolata Gendre, 1926
Vassiliades, G., 1972, Bull. Inst. Fond. Africaine Noire, s. A, v. 34 (3), 529-533
Ctenopoma kingsleyae: Sangalkam, Senegal guppy d'aquarium (exper.)
- Ranjhinema, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of Philometra
key
- Raphidascaridinae Hartwich, 1954
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakidae
key; key to tribes
includes: Lappetascaridinea; Aliasascaridinea; Raphidascaridinea; Paranisakinea nov. trib.
- Raphidascaridinea Chabaud, 1965
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinae
key; key to genera
includes: Raphidascaris; Raphidascaroides; Thynnascaris; Heterotyphlum
- Raphidascaris
Gibson, D. I.; and Taylor, A. L., 1976, Parasitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Raphidascaris Railliet & Henry, 1915, illus.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinea
key; synonymy
- Raphidascaris sp.
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Alosa sapidissima (coelom and mesentery): Sacramento-San Joaquin Delta, California
- Raphidascaris sp.
Sakaguchi, Y.; and Katamine, D., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 159-169
anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
Sillago japonica: sea near Nagasaki
Pneumatophorus tapeinocephalus: "
Decapterus maruadsi: East China Sea
Caranx equula: "
Trachurus japonicus: "
Lepidotrigla microptera: "
Trichiurus lepturus: "
Lutjanus sebae: South China Sea
Abalistes stellatus: "
Megalospis cordyla: "
Plectorhynchus pictus: "
- Raphidascaris acus (Block, 1779)
Bogoiavlenskii, Iu. K.; and Demchenko, A. N., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 281-285
Raphidascaris acus, structure of cuticle, hypodermis, and somatic musculature
- Raphidascaris acus Bloch, 1779
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Salmo trutta (alimentary tract)
Perca fluviatilis (intestine)
Esox lucius
all from Loch Leven, Scotland
- Raphidascaris acus (Bloch, 1779) Railliet et Henry, 1915
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 195-201
Lota l. lota (pyloric appendices, stomach, anterior portion of intestine)
Esox lucius
Anguilla anguilla
Perca fluviatilis
Silurus glanis
Leuciscus idus
all from Poland
- Raphidascaris acus (Bloch, 1779) Railliet et Henry, 1915
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
Silurus glanis (stomach, intestine): river Biebrza basin, Poland
- Raphidascaris acus (Bloch, 1779), illus.
Guimaraes, J. F., 1975, Bol. Inst. Biol., Bahia, v. 14 (1), 44-52
description
Lutjanus vivanus (estomago): Salvador, Bahia, Brasil
- Rhaphidascaris acus
Perłowska, R., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 27-32
Esox lucius
Rutilus rutilus
all from Zegrzynski Reservoir
- Rhaphidascaris acus (Bloch, 1779)
Puciłowska, A., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 33-46
helminths of fishes, dynamics of infection following formation of artificial body of water, seasonal distribution, brief description
Tinca tinca
Abramis brama
Leuciscus idus
all from Zegrzynski Reservoir

- Raphidascaroides Yamaguti, 1941
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Raphidascaridinea
key
Syn.: Ryjиковасcaris Mozgovoi, 1951
- Raphidascaroides blochii n. sp., illus.
Bilqees, F. M.; and Khanum, Z., 1974, Pakistan J. Zool., v. 6 (1-2), 151-155
Sphyrna blochii (intestine): Karachi coast
- Reesimermis Tsai and Grundmann, 1969
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
as syn. of Romanomermis Coman, 1961
- Reesimermis muspratti sp. nov., illus.
Obiamiwe, B. A.; and Macdonald, W. W., 1973, Ann. Trop. Med. and Parasitol., v. 67 (4), 439-444
life cycle
Aedes: Livingstone, Zambia
Culex: Livingstone, Zambia
Aedes aegypti ([?] exper.)
Aedes polynesiensis ([?] exper.)
Culex pipiens molestus ([?] exper.)
Culex pipiens fatigans ([?] exper.)
Anopheles stephensi ([?] exper.)
Anopheles albimanus ([?] exper.)
- Reesimermis muspratti Obiamiwe and Macdonald, 1973
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
as syn. of Octomyomermis muspratti (Obiamiwe and Macdonald, 1973) n. comb.
- Reesimermis nielsenii Tsai and Grundman, illus.
Chen, P. S., 1976, Bull. Inst. Zool., Acad. Sinica, v. 15 (1), 21-28
Reesimermis nielsenii as possible biological control agent for Culex pipiens fatigans: mass production; transstadial transmission; importance of water pH in limiting habitat range; field trials: Taiwan
- Reesimermis nielsenii Louisiana strain
Finngermis nielsenii, Louisiana strain
v. 37 v. J. R.; et al., 1977, Mosquito News, (1), 6-11
as syn. of Romanomermis culicivora
- Reesimermis nielsenii
Kerdipibul, V.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 150-151 [Demonstration of biological control of Mansonia uncinata vectors of human filariasis through nematode parasitism with Reesimermis nielsenii preliminary laboratory trials]
- Reesimermis nielsenii Tsai & Grundmann
Levy, R., Murphy, L. J., jr.; and Miller, T. W., jr., 1976, Mosquito News, v. 36 (4), 498-501
Reesimermis nielsenii, effects of pressure and nozzle impact of simulated aerial spray system of dissemination, no apparent loss of viability, infectivity, or development, tested against field-collected and laboratory-reared Culex pipiens quinquefasciatus larvae
- Reesimermis nielsenii, illus.
Nickle, W. R.; and Hoegger, Ch. H., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 173-177
Reesimermis nielsenii, scanning electron microscope study, infective stage, external anatomy of parts of male
- Reesimermis nielsenii Tsai & Grundmann
Petersen, J. J., 1973, J. Med. Entom., v. 10 (1), 75-79
Reesimermis nielsenii, mass production using Culex pipiens quinquefasciatus, effects of host density, parasite-host ratio, and amount of food fed to host on percentage of parasitism and female nemas produced
- Reesimermis nielsenii Tsai and Grundmann
Petersen, J. J.; and Willis, O. R., 1976, Mosquito News, v. 36 (3), 339-342
Reesimermis nielsenii, control of floodwater mosquitoes by release of nematode in appropriate habitat, collection of wild larval mosquitoes to measure success: Louisiana
Aedes atlanticus
A. tormentor
Psorophora columbiae
P. howardii
- Reesimermis nielsenii s. str.
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
as syn. of Romanomermis nielsenii (Tsai and Grundmann) n. comb.
- Reesimermis nielsenii auct. [pro parte]
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
as syn. of Romanomermis culicivora n. sp.
- Rhabdias sp.
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Rana ridibunda
R. esculenta
all from Samara river valley, Ukrainian SSR
- Rhabdias sp., illus.
Jacobson, E., 1976, J. Zoo Animal Med., v. 7 (2), 14-15
Rhabdias sp., levamisole hydrochloride, snakes
- Rhabdias sp.
Koller, R. L.; and Gaudin, A. J., 1977, Southwest Nat., v. 21 (4), 503-509
helminth recovery at 2 sites with diverse climates, statistical analysis indicates correlations between incidence and/or intensity of infection and host species, locality, and sex and size of host
Hyla regilla
Bufo boreas
all from Los Angeles County, California (Malibu Creek; Big Tujunga Wash)
- Rhabdias spp.
Pinnell, J. L.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 337-340
Gekko gekko: Komodo Island, Indonesia

- Rhabdias bufonis* (Schrank, 1788)
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Bufo bufo
Rana ridibunda
R. terrestris
R. esculenta
all from Samara river valley, Ukrainian SSR
- Rhabdias bufonis* (Schrank, 1788)
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
Rana temporaria: Macedonia
- Rhabdias bufonis* (Schrank, 1788), *illus.*
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Rana ridibunda
R. esculenta
R. agilis
Bufo bufo
Bombina bombina
(pluca of all): all from Yugoslavia
- Rhabdias bufonis* (Schrank, 1788)
Plasota, K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 47-60
helminths of frogs, comparison of aquatic and terrestrial hosts, relation of parasite fauna to environment, food supplies and food habits, host life cycle, temperature, rainfall, season, age and sex of host, competition between species of parasite, localization within host
Rana terrestris (lungs, intestine): Kampanos National Park, Poland
- Rhabdias fuscoverrosus*
Kurashvili, B. E., 1975, Izvest. Akad. Nauk Gruzinsk. SSR, s. Biol., v. 1 (4), 317-320
antagonistic and synergetic interrelationships between intestinal parasites
- Rhabdias microoris* Semenov, 1929
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (2), 82-84
Bombina bombina
Rana ridibunda
R. esculenta
all from Samara river valley, Ukrainian SSR
- Rhabdias sphaerocephala* Goodey, 1934
Caballero Deloya, J., 1974, An. Inst. Biol., Univ. Nac. Mexico, s. Zool., v. 45 (1), 45-50
Bufo horribilis (pulmones): Laguna de Catemaco, Veracruz, Mexico
- Rhabditida*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Secernentea
key
includes: Rhabditoidea
- Rhabditida* Chitwood, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Secernentea
includes: Rhabditina; Diplogasterata; Cephalobata
- Rhabditides*
Beaucournu, J. C.; and Deunff, J., [1976], Ann. Parasitol., v. 50 (6), 1975, 831-835
as hyperparasites of fleas, causing parasitic castration
siphonapteres: France
- Rhabditina* Chitwood, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Rhabditida
- Rhabditis axei* (Cobbold, 1884) Dougherty, 1955, *illus.*
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972, Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera beetles, possible importance in biological control
Blaps polychresta (externally on abdominal segments): Cairo, Egypt
- Rhabditis pellio*
Somers, J. A.; Shorey, H. H.; and Gaston, L. K., 1977, J. Chem. Ecol., v. 3 (4), 467-474
Rhabditis pellio, quantitative bioassay for female-produced attractant pheromone by measuring male migration response, age in relation to production and response, daily rhythms
- Rhabditis strongyloides*, *illus.*
Colby, E. D.; and McGrew, L. S., 1976, Vet. Med. and Small Animal Clin., v. 71 (10), 1426
Rhabditis strongyloides, dog, alopecia, case report
- Rhabditoidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Rhabditida
- Rhabditolaimus crassus* Korner, 1954, *illus.*
Ali, M.; Wahab, A.; and El-Kifel, A. H., 1972, Parasitol. Hungar., v. 5, 177-201
survey of nematode spp. invading Coleoptera beetles, possible importance in biological control
Scarabaeus sacer: Abu-Rawash, Cairo, Egypt
- Rhabditostomum* n. gen.
Chabaud, A. G.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 721-727
Oesophagostominae
tod: R. traguli (Maplestone, 1932) n. gen. n. comb.
- Rhabditostomum traguli* (Maplestone, 1932) n. gen. n. comb. (tod), *illus.*
Chabaud, A. G.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 721-727
redescription
Syn.: Oesophagostomoides traguli Maplestone, 1932
Tragulus javanicus (intestin grele): Selenkor, Jinjang, Bt. Legong Forest Reserve

- Rhabdochona Railliet, 1916, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochonidae
key; synonymy
includes subgens.: Rhabdochona; Filochona; Globochona
- Rhabdochona, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of Rhabdochona
key
- Rhabdochona Railliet, 1916
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
discussion of systematic position
- Rhabdochona Railliet, 1916
Rehana, R.; and Bilqees, F. M., 1976, *Agric. Pakistan*, v. 26 (4), 1975, 521-528
synonymy
- Rhabdochona sp.
Beacham, B. E.; and Haley, A. J., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 232-233
Morone americana (stomach): Chesapeake Bay
- Rhabdochona sp.
Combs, D. L.; Harley, J. P.; and Williams, J. C., 1977, *Tr. Kentucky Acad. Sc.*, v. 38 (3-4), 128-131
Moxostoma erythrurum (gut): Kentucky River
- Rhabdochona sp. 2
Kakacheva-Avramova, D., 1973, *Izvest. Tsentral. Khelmit. Lab.*, v. 16, 87-110
Barbus meriodionalis petenyi (intestine): Balkan Mountain river
- Rhabdochona sp. juv.
Kakacheva-Avramova, D., 1973, *Izvest. Tsentral. Khelmit. Lab.*, v. 16, 87-110
P[erca] fluviatilis (intestine): Balkan Mountain river
- Rhabdochona sp.
Lockard, L. L.; Parsons, R. R.; and Schaplow, B. M., 1975, *Great Basin Nat.*, v. 35 (4), 442-448
Salmo trutta (upper digestive tract).
relationship of incidence and intensity of nematode infection to age and sexual maturity of host, higher infection rate in sexually mature trout due to aggressive feeding behavior: streams in southern and western Montana
- Rhabdochona sp.
Mudry, D. R.; and Anderson, R. S., 1977, *J. Fish Biol.*, v. 11 (1), 21-33
Salmo clarki: Jasper National Park, Canada
Salvelinus fontinalis: Banff National Park, Canada
- Rhabdochona sp.
Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, *Fish. Bull., National Oceanic and Atmos. Admin.*, v. 71 (1), 267-277
helminths, incidence and intensity of infection in young and adult Oncorhynchus nerka, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Rhabdochona alii n. sp., *illus.*
Kalyankar, S. D., 1972, *Riv. Parassitol., Roma*, v. 33 (4), 281-288
Labeo rohita (intestine): Nanded, Maharashtra, India
- Rhabdochona baylisi Rai, 1969
Sahay, U.; and Narayan, S., 1971, *Indian J. Animal Research*, v. 5 (2), 51-54
as syn. of R. garuai Agrawal
- Rhabdochona cascadilla Wigdor
Cloutman, D. G., 1976, *Southwest Nat.*, v. 21 (1), 67-70
Campostoma anomalum pullum (gut): White River, Arkansas
- Rhabdochona cascadilla Wigdor
Lang, B. Z.; and Edson, S. A., 1976, *J. Parasitol.*, v. 62 (1), 93
Rhinichthys osculus: Turnbull National Wildlife Refuge, Spokane County, Washington
- Rhabdochona cascadilla Wigdor, 1918
Mudry, D. R.; and Anderson, R. S., 1977, *J. Fish Biol.*, v. 11 (1), 21-33
Salvelinus fontinalis: Banff National Park, Canada
- Rhabdochona cavasius Rehana & Bilqees, 1973
Rehana, R.; and Bilqees, F. M., 1976, *Agric. Pakistan*, v. 26 (4), 1975, 521-528
Mystus cavasius (intestine): Kalri Lake, Sind area
- Rhabdochona chanawanensis n. sp., *illus.*
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
Eutropiichthys vacha (intestine): Chanawan Fish Farm, Wazirabad, Pakistan
- Rhabdochona congolensis Campana-Rouget, 1961
Khalil, L. F.; and Thurston, J. P., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (2), 209-248
Haplochromis angustifrons
H. elegans
H. nigripinnus
H. squamipinnus
Haplochromis sp. (H. brassy of Greenwood in M.S.)
Haplochromis sp. (H. brassy brassy of Greenwood in M. S.)
(intestine of all): all from Lake George, Uganda
- Rhabdochona congolensis Campana-Rouget, 1961, *illus.*
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Chrysichthis nigrodigitatus: Cameroun
Synodontis notatus: Zaire
Haplochromis sp.: Uganda, Lake Victoria
Ctenopoma kingsleyae: Senegal
(intestine of all)
- Rhabdochona cotti (Gustafson, 1949)
Mudry, D. R.; and Anderson, R. S., 1977, *J. Fish Biol.*, v. 11 (1), 21-33
Cottus cognatus: Yoho National Park, Canada

- Rhabdochona decaturensis*
Grüniger, T. L.; Murphy, C. E.; and Britton, J. C., 1977, *Southwest. Nat.*, v. 22 (4), 525-535
Ictalurus punctatus
Aplodinotus grunniens
all from Eagle Mountain Lake, Texas
- Rhabdochona denudata* (Dujardin, 1845) Railliet, 1916
Ejsymont, L., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 203-216
description
Silurus glanis (anterior and middle parts of intestine): river Biebrza basin, Poland
- Rhabdochona* (*Rhabdochona*) *denudata* (Dujardin, 1845)
Kakacheva-Avramova, D., 1972, *Izvest. Tsentral. Khelmit. Lab.*, v. 15, 89-107
Leuciscus cephalus
Alburnus alburnus
(intestine of all): all from River Tundzha
- Rhabdochona denudata* (Dujardin, 1845)
Kakacheva-Avramova, D., 1973, *Izvest. Tsentral. Khelmit. Lab.*, v. 16, 87-110
L[euiscus] cephalus
V[imba] vimba tenella
Alb[urnus] alburnus
Alb[urnoides] bipunctatus
(intestine of all): all from Balkan Mountain river(s)
- Rhabdochona denudata*
Shakhmatova, V. I., 1966, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 17, 277-289
Lutra lutra: Karelia
- Rhabdochona filamentosa* (Bykovskaya-Pavlov, 1936)
Fagerholm, H.-P., 1976, *Norwegian J. Zool.*, v. 24 (4), 466 [Abstract]
Finland
- Rhabdochona garuui* Agrawal
Sahay, U.; and Narayan, S., 1971, *Indian J. Animal Research*, v. 5 (2), 51-54
Syn.: *R. baylisi* Rai, 1969
- Rhabdochona gendrei* Campana-Rouget, 1961, illus.
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Barbus camptacanthus (estomac, intestin): Cameroun, Olounou
- Rhabdochona hellichi* (Sramek, 1901)
Kakacheva-Avramova, D., 1973, *Izvest. Tsentral. Khelmit. Lab.*, v. 16, 87-110
Barbus barbus
B. meridionalis petenyi
(intestine of all): all from Balkan Mountain river(s)
- Rhabdochona labeonis* n. sp., illus.
Kalyankar, S. D., 1972, *Riv. Parassitol.*, Roma, v. 33 (4), 281-288
Labeo rohita (intestine): Nanded, Maharashtra, India
- Rhabdochona magna* Khan and Yaseen, 1969, illus.
Zaidi, D. A.; and Khan, D., 1975, *Pakistan J. Zool.*, v. 7 (1), 51-73
female redescribed
Rita rita (intestine): Panjnad Headworks, Pakistan
- Rhabdochona milleri* Choquette, 1951
Mudry, D. R.; and Anderson, R. S., 1977, *J. Fish Biol.*, v. 11 (1), 21-33
Salvelinus fontinalis: Yoho and Banff National Parks, Canada
Prosopium coulteri: Yoho National Park, Canada
Salvelinus malma: Yoho National Park, Canada
Salmo gairdneri: Yoho National Park, Canada
S. clarki: Waterton Lakes National Park, Canada
- Rhabdochona moravecii* sp. n., illus.
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Aphyosemion camerounensis (estomac, intestin): Cameroun, Olounou
- Rhabdochona sarana* Karve and Naik, 1951
Kalyankar, S. D., 1972, *Riv. Parassitol.*, Roma, v. 33 (4), 281-288
Labeo rohita (intestine): Aurangabad, India
- Rhabdochonidae* (Travassos, Artigas & Pereira, 1928, subfam.) Skrjabin, 1946
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Thelazioidea
key; key to genera
includes: *Rhabdochona*; *Johnstonmawsonia*; *Freitasia*; *Vasorhabdochona*; *Heptochona*; *Hepatinema*; *Trichospirura*
- Rhabdochonidae*
Chabaud, A. G.; and Krishnasamy, M., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 813-820
Trichospirura, should be placed in *Rhabdochonidae*, evolutionary position, host range, osmoexcretory apparatus, relations between *Rhabdochonidae* and *Cystidicolidae*
- Rhabdochonidae*
Puylaert, F. A., 1973, *Rev. Zool. et Botan. Africaines*, v. 87 (4), 647-665
Physalopteroidea
- Rhabdochonoides Janizewska*, 1955
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Rhabdochona* Railliet, 1916
- Rhaphidascaris*. See *Rhaphidascaris*.
- Rhigonema critesi* sp. n., illus.
Ramirez, J., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 131-134
Orthoporus typotypyge (hindgut): west of Juan Santamaria International Airport, Alajuela Province, Costa Rica
- Rictularia Froelich*, 1802
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Rictulariidae
key

- Rictularia Froelich, 1802
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93 comparative characteristics of species from bats, includes: *R. bovieri*; *R. plagiostoma*; *R. macdonaldi*; *R. lucifugus*; *R. nana*
- Rictularia sp. Froelich 1802
Bienek, G. K.; and Klikoff, L. G., 1974, Am. Midland Naturalist, v. 91 (1), 251-253
Dipodomys merriami vulcani: Dixie State Park, Washington Co., Utah
- Rictularia sp.
Kenney, M.; et al., 1975, Am. J. Trop. Med. and Hyg., v. 24 (4), 596-599
gravid female worm and eggs found in appendix at autopsy of elderly man: New York
- Rictularia sp.
Martin, D. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 85-86
Tadarida brasiliensis: Texas
- Rictularia sp.
O'Farrell, T. P., 1975, Am. Midland Naturalist, v. 93 (2), 377-387
Perognathus parvus
Peromyscus maniculatus
all from Arid Lands Ecology Reserve, Benton County, Washington
- Rictularia [sp.]
Saxena, A.; and Nama, H. S., 1977, Geobios, v. 4 (6), 243-244
Rattus rattus (colon): Jodhpur, India
- Rictularia sp.
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 39-42
Rattus norvegicus (intestino delgado): Chile
- Rictularia bovieri
Durette-Desset, M.-C.; and Chabaud, A.-G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Myotis mystacinus: grotte de Vallorbe, Vaud, Suisse; Col de Jaman, Vaud, Suisse
- Rictularia bovieri Blanchard, 1886, illus.
Skvortsov, V. G., 1971, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (7), 75-93 description, geographic distribution
Vespertilio murinus
Myotis myotis
M. oxygnathus
Miniopterus schreibersi
(small intestine of all): all from Moldavia
- Rictularia bovieri Blandchard, 1886
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155 ecological analysis of bat helminth fauna, geographic distribution
Myotis oxygnathus: Moldavia
- Rictularia lucifugus Douvres, 1956
Cain, G. D.; and Studier, E. H., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 113-114
Myotis lucifugus: New Mexico
- Rictularia proni, larva
Killick-Kendrick, R.; et al., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 22 [Demonstration]
Phlebotomus ariasi: Gard, southern France
- Rictularia proni Seurat 1915, illus.
Sanchez-Acedo, C.; and Vericad, J. R., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 197-203 measurements
Arvicola sapidus: Aragon Pyrenees
- Rictularia proni, illus.
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Rictularia ratti
Nama, H. S.; and Parihar, A., 1976, J. Helminth., v. 50 (2), 99-102
Rattus rattus rufescens (intestine): Jodhpur City area, India
- Rictularia shaldybini Skrjabin, Sobolev et Ivaschkin, 1967
Babaev, Ia.; and Kolodenco, A. I., 1975, Izvest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk (4), 71-75
[*Hemiechinus auritus*]: Turkmenistan
- Rictularia tani Hoespli, illus.
Kamiya, M., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 139-141
Rictularia tani, prevalence survey in rodents, male and female morphometric data and measurements, comparisons of present and previous studies
Rattus norvegicus (small intestines)
R. rattus " "
all from vicinity of Bangkok, Thailand
- Rictularia tani (Hoespli, 1929)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus r. jarak
R. cremoriventer
R. jalorensis
R. mulleri
R. rajah subsp.
R. sabanus
all from Malaysia
- Rictularia tani (Hoespli, 1929)
Wioreno, W., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 136-138
Rattus rattus diardi: Bogar, West Java, Indonesia

- Rictulariidae (Hall, 1915, subfam.) Railliet, 1916
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Rictularioidea
 key to genera
 includes: Rictularia; Pterygodermatites
- Rictulariidae gen. sp. I, larvae
 Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
 role of Tenebrionidae as intermediate hosts
 Blaps oblonga: Tadzhik SSR
- Rictulariidae gen. sp. II, larvae
 Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
 role of Tenebrionidae as intermediate hosts
 Penthyucus granulosis: Tadzhik SSR
- Rictulariidae gen. sp.
 Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
 Trigonoscelis gigas: Turkmenia
- Rictulariidae gen. sp. 1
 Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
 Trigonoscelis punctipleuris: Turkmenia
- Rictularioidea
 Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
 Spirurina
 key
- Rictularioidea
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Spirurina
 includes: Rictulariidae
 "The rictularioids, generally classified within the Spiruroidea or Thelazioidea are here given their own superfamily. . ."
- Rictularioides Hall, 1916
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 "may be a juvenile Pterygodermatites,"
 "excluded from the key."
- Rinadia Grigorian, 1951
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Trichostrongylidae, Ostertagiinae
- Rinadia mathevossiani (Ruchliadev, 1948) Andreeva, 1957
 Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
 Capreolus capreolus (rennet): southern Bulgaria
- Rinadia mathevossiani
 Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
 nematodes, seasonal dynamics in deer
- Rinadia mathevossiani
 Schweisgut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
 Rotwild: Nationalpark Bayerischer Wald
- Rodentocaulus ondatrae Schulz, Orloff, Kutas, 1933
 Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
 Ondatra zibethica (lungs): Karelia
- Romanomermis Coman, 1961
 Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
 resurrection, revised description, key, key to species
 Syns.: Reesimermis Tsai and Grundmann, 1969;
 Eurymermis in part: Rubtsov 1972
- Romanomermis sp. auct. (in relation to the North American fauna)
 Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
 as syn. of Romanomermis culicivorax n. sp.
- Romanomermis cazanica
 Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
 key
- Romanomermis culicivorax n. sp., illus.
 Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
 key
 Syns.: Reesimermis nielseni auct. [pro parte]; Romanomermis sp. auct. (in relation to the North American fauna)
 Culex pipiens quinquefasciatus
 Anopheles crucians
 Psorophora confinnis
- Romanomermis culicivorax
 Brown, B. J.; Platzer, E. G.; and Hughes, D. S., 1977, Mosquito News, v. 37 (4), 603-608
 Romanomermis culicivorax, field trials for reduction of mosquito larvae, infection percentage dependent on mosquito subfamily, application rate, and test site
 Anopheles franciscanus
 A. freeborni
 Culex tarsalis
 Culiseta inornata
 all from California
- Romanomermis culicivorax
 Finney, J. R.; et al., 1977, Mosquito News, v. 37 (1), 6-11
 Romanomermis culicivorax and growth regulator Altosid SE used separately and concurrently, effective in controlling pupal and larval populations of Aedes aegypti
 Syn.: Reesimermis nielseni, Louisiana strain
- Romanomermis culicivorax, illus.
 Hansen, E. L.; and Hansen, J. W., 1976, IRCS J. Med. Sc., v. 4 (11), 508
 Romanomermis culicivorax, experimental parasitism of Simulium damnosum vectors of Onchocerca volvulus, possible use as biological control agent and as laboratory model for developing similar techniques with other mermithids

- Romanomermis culicivora* Ross & Smith (= *Reesimermis nielsenii* Tsai & Grundmann, auct., partim.)
Levy, R.; and Miller, T. W., jr., 1977, Environment. Entom., v. 6 (3), 447-448
Romanomermis culicivora, effect of pesticides and growth regulators used in mosquito control operations on viability and infectivity
- Romanomermis culicivora* (*Reesimermis nielsenii*), illus.
Poinar, G. O., jr.; and Hess, R., 1977, Nature, London (5599), v. 266, 256-257 [Letter]
Romanomermis culicivora, parasitic juveniles contained virus-like particles in hypodermal cords, damage could lower efficiency of this nematode as biological control agent
- Romanomermis culicivora*, illus.
Poinar, G. O., jr.; and Hess, R., 1977, Exper. Parasitol., v. 42 (1), 27-33
Romanomermis culicivora, parasitic juveniles, morphological evidence of possible transport system for transcuticular uptake of nutrients
- Romanomermis hermaphrodita* n. sp., illus.
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
key, high incidence of intersexuality
Aedes (*Ochlerotatus*) *nigripes*: tundra pools on the coast of Hudson Bay, about 8 km east of Fort Churchill, Manitoba
- Romanomermis iyengari*
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
key
- Romanomermis kiktoreak* n. sp., illus.
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
key
Aedes (*Ochlerotatus*) *impiger*: tundra pools about 1 km north of Baker Lake, N.W.T., Canada
A. communis (exper.)
A. rempeli (exper.)
- Romanomermis nielsenii* (Tsai and Grundmann) n. comb.
Ross, J. F.; and Smith, S. M., 1976, Canad. J. Zool., v. 54 (7), 1084-1102
key
Syn.: *Reesimermis nielsenii* s. str.
- Rondonema Artigas*, 1926
Rao, V. J., 1973, Riv. Parassitol., Roma, v. 34 (3), 205-212
key to species, includes: *R. caballeri* Travassos et Kloss, 1960; *R. alatum* Kloss, 1965; *R. spirostreptus* Rao and Kumari, 1967; *R. spinifera* Rao, 1958; *R. sureshi* n. sp.; *R. rondoni* Artigas, 1926; *R. thapari* Farooqui, 1967
- Rondonema spinifera* Rao, 1958, illus.
Rao, V. J., 1973, Riv. Parassitol., Roma, v. 34 (3), 205-212
key, redescription
Spirostreptus sp. (intestine): Hyderabad (Andhra Pradesh), India
- Rondonema sureshi* n. sp., illus.
Rao, V. J., 1973, Riv. Parassitol., Roma, v. 34 (3), 205-212
key
Spirostreptus (intestine): Mananoor (Andhra Pradesh), India
- Roundworm
Sehgal, S. C.; Vinayak, V. K.; and Gupta, U., 1977, Indian J. Med. Research, v. 65 (4), 509-512
human helminthic ova in feces, diagnosis using the Kato thick smear technique more successful than commonly used techniques, recommended for epidemiologic surveys: Chandigarh, India
- Rumai Travassos, 1960
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key
- Ruschiella Freitas*, 1967
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Procyrnea* (Chabaud, 1958, subgen.)
- Rusguniella Seurat*, 1919, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key; synonymy
- Rusguniella* (*Rusgunoides*) Williams, 1929
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Rusguniella Seurat*, 1919
- Rusguniella wedli*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Ryjikovascaris Mozgovoi*, 1951
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Raphidascaroides Yamaguti*, 1941
- Ryzhikovistoma* subg. n.
Ali, M. M., 1970, Acta Parasitol. Polon., v. 17 (20-38), 237-246
subgen. of *Cyathostoma*
key, key to species
diagnosis, tod: *Cyathostoma* (*R.*) *coscorobae* Chapin, 1925 comb. n.

- Salvelinema Trofimenko*, 1962, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
Syn.: *Pseudometabronema Bogdanova*, 1963
- Sandnema n.* subgen.
Chabaud, A.-G.; and Bain, O., 1975, Ann. Parasitol., v. 51 (3), 365-397
subgen. of *Tetrapetalonema*; key
tod of subgen.: *Tetrapetalonema (Sandnema) digitata* (Chandler, 1929)
- Sanguinifilaria Yamaguti*, 1935
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Philometra Costa*, 1845
- Sarconema eurycerca*
Irwin, J. C., 1975, J. Wildlife Dis., v. 11 (1), 8-12
Sarconema eurycerca causing mortality in Olor columbianus, pathology: Lake St. Clair, Ontario
- Sarconema eurycerca*
Seegar, W. S.; et al., 1976, Science (4266), v. 194, 739-740
Sarconema eurycerca in *Cygnus columbianus columbianus* (blood), transmission to *C. olo* using larvae obtained from infected lice (*Trinoton anserinum*), results show mallophagan as natural cyclodevelopmental vector *Cygnus olor* (exper.) (blood)
C. columbianus columbianus (blood): North Carolina
Trinoton anserinum: North Carolina; Black Sea, U.S.S.R.
- Sarconema pseudolabiata nov. sp.*, illus.
Belogurov, O. I.; Daiia, G. G.; and Sonin, M. D., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 3-6
Syns.: *Loainae gen. sp.* Lubimow, 1926; *Filaria sp.* Serkova, 1948; *Ornithofilaria sp.* Borgarenko, 1960; *Aproctidae gen. sp.* 2 Oschmarin, 1963; *Aproctinae gen. sp.* Sonin et Borgarenko, 1965
Anas platyrhynchos
A. acuta
A. clypeata
A. crecca
A. penelope
Aix galericulata
(subcutaneous tissue in region of esophagus of all): all from SSSR
- Sarwaria Drozd*, 1965
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, *Ostertagiinae*
- Saurofilaria Caballero*, 1954
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
as syn. of *Macdonaldius Khanna*, 1933
- Schistogendra Chabaud & Rousselot*, 1956, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Schistorophinae
key
- Schistorophid larvae
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (esophagus, proventriculus): Florida
- Schistorophinae Travassos*, 1918
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariidae
key; key to genera
includes: *Schistogendra*; *Schistorophus*; *Sciadiocara*; *Viktorocara*; *Ancyracanthopsis*
- Schistorophus Railliet*, 1916, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Schistorophinae
key; synonymy
- Schistorophus sp.*
Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 76-79
degree of helminth infection in different age groups of *Cololabis saira*: region of Kuril'sk and Japan
- Schistorophus cornutus Sobolev*, 1943
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Xenus cinereus: lower Yenisei and Keta lake
- Schistorophus lii Daija*, Bondarenko et Gubanov, in litt. [nomen nudum]
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Numenius ph. phaeopus: Keta lake
- Schistorophus skrjabini Wassilkowa*, 1916
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus: Yenisei
- Schrankiana Strand* 1942
Dyer, W. G.; and Altig, R., 1977, Herpetologica, v. 33 (3), 293-296
synonymy
- Schrankiana schranki* (Travassos 1925) Strand 1942
Dyer, W. G.; and Altig, R., 1977, Herpetologica, v. 33 (3), 293-296
Leptodactylus pentadactylus (large intestine): Santa Cecilia, Napo Province, Ecuador
- Schranknema Travassos* 1949
Dyer, W. G.; and Altig, R., 1977, Herpetologica, v. 33 (3), 293-296
as syn. of *Schrankiana Strand* 1942

- Schulzia Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
- Schulzitriconema
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Schwenkiella icemi (Schwenk, 1926) Basir, 1956
Leong, L.; and Paran, T. P., 1966, Med. J. Malaya, v. 20 (4), 349
Periplaneta americana
Blatta orientalis
all from Singapore
- Sciadiocara Skrjabin, 1916, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Schistorophinae
key
- Sciadiocara chabaudi
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Sciadiocara chabaudi Schmidt and Kinsella, 1972
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans
Porphyryla martinica
(gizzard lining of all): all from Florida
- Sciadiocara umbellifera (Molin, 1869)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Xenus cinereus
Numenius ph. phaeopus
all from lower Yenisei [and/or] Keta lake
- Sciadiocara umbellifera
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Sciadiocara umbellifera Molin, 1860
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus genei: Azov Sea
- Sciurodendrium Durette-Desset, 1971
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Pudicinae
- Secernentea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Nematoda
key; key to orders
includes: Rhabditida; Strongylida; Oxyurida; Ascaridida; Spirurida
- Secernentea (=Phasmodia) Dougherty, 1958
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Nematoda
includes: Rhabditida; Strongylida; Ascaridida; Spirurida; Tylenchida
- Serpinema Yeh, 1960, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanidae
key
- Serratospiculum amaculatum
Croft, R. E.; and Kingston, N., 1975, J. Wildlife Dis., v. 11 (2), 229-233
Falco mexicanus: Wyoming
- Serratospiculum amaculata
Kocan, A. A.; and Gordon, L. R., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 908
Serratospiculum amaculata in Falco mexicanus (thoracic air sac), cause of death related to heavy infection and possibly to bronchial obstruction with edema, fluid, and blood after administration of anaesthetic
- Serratospiculum amaculata
Kocan, A. A.; and Locke, L. N., 1974, J. Wildlife Dis., v. 10 (1), 8-10
Haliaeetus leucocephalus: Wisconsin; Illinois
- Serticeps Railliet, 1916
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Viguiera Seurat, 1913
- Setaria
Neppert, J., 1974, Tropenmed. u. Parasitol., v. 25 (4), 454-463
cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Setaria
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, Am. J. Vet. Research, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Setaria sp. proche labiatopapillosa
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
Bubalus bubalis: Sofia; New Delhi; Mathura; Kathmandu
- Setaria sp., illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 589-599
Equus greyvi: proche du Soudan

- Setaria* sp.
Wilson, D. E.; and Hirst, S. M., 1977, *Wild-life Monogr.* (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Setaria bernardi*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii krai
- Setaria boulengeri*
Young, E.; et al., 1973, *Research J. National Parks Republic South Africa* (16), 77-81
Redunca fulvorufula (abdominal cavity): Mountain Zebra National Park
- Setaria bubali* [? n. comb.], illus.
Shoho, C., 1976, *Ann. Parasitol.*, v. 51 (5), 577-588
re-examination of specimens of Rudolphi 1819 of *Filaria bubali*
- Setaria cervi*
Ansari, J. A., 1977, *Indian J. Animal Sc.*, v. 47 (3), 115-119
Setaria cervi, *Bubalus bubalis*, incidence and seasonal variation of adult worms and microfilariae, periodic concentration of microfilariae: Aligarh district, Uttar Pradesh
- Setaria cervi*
Anwar, N.; et al., 1977, *Ztschr. Parasitenk.*, v. 51 (3), 275-283
Setaria cervi, enzymes of glycolysis and PEP-succinate pathway
- Setaria cervi*
Anwar, N.; Ansari, A. A.; and Ghatak, S., 1976, *Proc. Indian Nat. Sc. Acad., Part B, Biol. Sc.*, v. 41 (6), 1975, 550-558
Setaria cervi, hexose utilization and glycogen synthesis in vitro
- Setaria cervi*
Baqui, A.; and Ansari, J. A., 1975, *Indian J. Zool.*, v. 3 (1-2), 43-48
Setaria cervi, white rats (blood) (exper.), leucocytic response
- Setaria cervi*
Schweisgut, I., 1975, *Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75*, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Setaria cervi*, illus.
Sharma, B. N.; Singh, S. P.; and Sahai, B. N., 1977, *Pantnagar J. Research*, v. 2 (2), 188-189
Setaria digitata, *S. cervi*, blood of rabbits (exper.), differences in morphology of microfilariae
- Setaria cervi*
Singhal, K. C.; Madan, B. R.; and Saxena, P. N., 1977, *Indian J. Med. Research*, v. 66 (3), 517-521
Setaria cervi, effects of various chemicals on parasite nerve-muscle complexes and locomotion using worms with cuticular permeability barriers removed
- Setaria cervi* (Rudolphi, 1819), illus.
Subramanian, G.; and Srivastava, V. K., 1973, *Riv. Parassitol., Roma*, v. 34 (1), 59-62
description, synonymy
goat (peritoneal cavity)
- Setaria digitata*, illus.
Jemelka, E. D., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (5), 673-675
Setaria digitata, horse (eye), surgical removal, case report: El Kabayo Stables, Subic Bay Naval Base, the Philippines
- Setaria digitata*
Mohan, R. N., 1977, *Indian J. Animal Sc.*, v. 45 (11), 1975, 914-915
Setaria digitata, cattle and buffaloes, cells of peritoneal exudate, eosinophils and mesothelial cells
- Setaria digitata*, illus.
Sharma, B. N.; Singh, S. P.; and Sahai, B. N., 1977, *Pantnagar J. Research*, v. 2 (2), 188-189
Setaria digitata, *S. cervi*, blood of rabbits (exper.), differences in morphology of microfilariae
- Setaria digitata*
Shoho, C., 1976, *Ann. Parasitol.*, v. 51 (5), 577-588
boeufs domestiques: Ceylan; Madras; Kerala; Trichur; Orissa; Dacca; Hissar
Bubalus bubalis: Kathmandu; Trichur; Kerala
- Setaria digitata*, illus.
Yoshikawa, T.; Oyamada, T.; and Yoshikawa, M., 1976, *Japan. J. Vet. Sc.*, v. 38 (2), 105-115
Setaria digitata, cattle, accidental parasitic entry in bovine urinary bladder from abdominal cavity, pathology of granulomatous lesions, histological findings
- Setaria equina*
Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Setaria equina* Abildgaard
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool., London*, v. 174 (3), 397-406
Equus burchelli (wall of abdominal cavity): Kenya
- Setaria equina* (Abildgaard, 1789)
Young, P. L.; and Babero, B. B., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)

- Setaria equina dafaallai* n. sub. sp., illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 589-599
Equus caballus: Shambat, pres Khartoum Nord; Conakry, Guinee; Fort Lamy, Tchad; Fort-Archambault, R.C.A.
Equus hemionus: Fort Lamy
- Setaria equina equina* (Abildgaard, 1789) Raillet et Henry, 1911, illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 589-599
Equus caballus: hongrois, abattoir de Geneve; Transvaal; Venezuela; Macedonie; Bulgarie; Kazakhstan; Mogadiscio, Somalie; Honshu, Japon; Hokkaido, Japon
- Setaria equina theilerae* n. sub. sp., illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 589-599
Equus burchelli: Kirawira, Serengeti, Tanzanie; Athi River, Kenya; Selous Game Reserve, Tanzanie; Transvaal; Justicia, Pilgrimsrest; Kruger National Park
Equus burchelli boehmi: Diatoka, Congo
Equus zebra: Etosha National Park
- Setaria indica* (Dutt, 1963)
Subramanian, G.; and Srivastava, V. K., 1973, Riv. Parasitol., Roma, v. 34 (1), 59-62
as syn. of *Setaria cervi* (Rudolphi, 1819)
- Setaria labiato-papillosa*
Green, P. E.; and Trueman, K. F., 1971, Austral. Vet. J., v. 47 (12), 624 [Letter]
cattle, incidence: Queensland, Australia
- Setaria labiatopapillosa*
Mougey, Y.; and Bain, O., 1976, Ann. Parasitol., v. 51 (1), 95-110
5 filaria-vector sets, stochastic models for assessing relation between numbers of microfilariae ingested and numbers of microfilariae passing into hemocoel, limitation phenomenon
- Setaria labiatopapillosa*
Niel, G.; et al., 1972, Medecine et Malad. Infect., v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalonema vitae* and *Ascaris suum* as antigens
- Setaria labiatopapillosa* and/or *S. yehi*
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
helminths of *Alces alces*, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Setaria labiatopapillosa*
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
Syncerus caffer: Tchad
chevre: Senegal and/or Mauritanie
mouton: Senegal and/or Mauritanie
boeufs domestiques: Soudan; Somalie; Tchad; R.C.A.; Congo; Kenya; Tanzanie; Madagascar; Hissar; Karachi; Pakistan Nord; Turquie; Alma Ata; Espagne; France; Italie
- Setaria labiatopapillosa* (A. Alessandrini 1848), illus.
Shoho, C.; and Sachs, R., 1975, Tropenmed. und Parasitol., v. 26 (4), 489-493
Giraffa camelopardalis (peritoneal cavity): Serengeti National Park in northern Tanzania, East Africa
- Setaria labiato-papillosa*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Aedes caspius caspius: Uzbekistan
- Setaria leichungwingi* Chen, 1937 (stat. nov.), illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
description
Bubalus bubalis (cavite peritoneale): abattoirs de Bangkok; Nord Viet-Nam; Philippines; Kuala-Lumpur; Hong-Kong
- Setaria marshalli* pandei
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
Bubalus bubalis: Kerala; Trichur
- Setaria nelsoni* n. sp., illus.
Shoho, C., 1976, Ann. Parasitol., v. 51 (5), 577-588
Syncerus caffer (cavite peritoneale, tissus connectifs peritracheaux): Kianga, Ancoles, Ouganda; Rift Valley; Ancoles, Queen Elizabeth Park, Ouganda; Kirawira, Serengeti, Tanzanie; Acholi et Toro, Queen Elizabeth Park, Ouganda; Cameroun; R.C.A.; Tchad; Selous Game Reserve, Tanzanie
Kobus defassa: Lac Manyara
Adenota kob: Acholi, partie occidentale du Queen Elizabeth Park, Ouganda
- Setaria tundrae*, illus.
Rehbinder, C.; Christensson, D.; and Glatthard, V., 1975, Nord. Vet.-Med., v. 27 (10), 499-507
Onchocerca sp. in subperitoneal and subcutaneous granulomas and *Setaria tundrae* in encapsulations in peritoneum of reindeer, increasing incidence, association with liver lesions caused by Corynebacteria, found in forest herds but not mountain herds
- Setaria tundra capreoli* Chusaburo Choho, 1959
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
Capreolus capreolus (abdominal cavity): southern Bulgaria
- Setaria yehi*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (peritoneal cavity): Kentucky
- Setaria yehi* Desset, 1966
Phillips, J. H.; Harley, J. P.; and Rudersdorf, W. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 250
Dama dama (mesenteries): western Kentucky

- Setaria yehi*
Prestwood, A. K.; and Pursglove, S. R., 1977, J. Am. Vet. Med. Ass., v. 171 (9), 933-935
Odocoileus virginianus (abdominal and thoracic cavities), prevalence according to age and sex of host, distribution, pathogenicity: southeastern United States
- Setaria yehi* Desset, 1966
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (abdominal cavity): Cumberland County, New Jersey; Oklahoma
- Setaria yehi* and/or *S. labiatopapillosa*
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
helminths of *Alces alces*, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Setaria yehi*
Weinmann, C. J.; et al., 1973, J. Wildlife Dis., v. 9 (3), 213-220
Wehrdikmansia cervipedis, *Elaeophora schneideri*, *Setaria yehi*, prevalence of infection in various age classes of *Odocoileus hemionus columbianus*: northern California
- Seurattia* Skrjabin, 1916, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Seurattiinae
key
- Seurattia puffini* Yamaguti, 1941
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus: Yenisei
- Seurattiinae* Chitwood & Wehr, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariidae
key; key to genera
includes: *Ingliseria*; *Rusguniella*; *Aviculariella*; *Proyseria*; *Stegophorus*; *Seurattia*; *Streptocara*
- Seuratoidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Ascaridida
key
- Seuratum cadarachense* Desportes, 1947, illus.
Quentin, J. C.; and Seureau, C., 1975, Ztschr. Parasitenk., v. 47 (1), 55-68
Seuratum cadarachense, first larval stages, organogenesis, migration and cellular reactions in *Locusta migratoria* (exper.) (lumiere de l'intestin moyen), imperfect adaptation of nematode to intermediate host; comparison with other nematode life cycles, speculations on evolution
Eliomys quercinus (niveau de l'intestin): pied de Alpilles a Eygalieres (Bouches-du-Rhone)
- Seuratum cadarachense*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Seuratum cancellatum* Chitwood, 1938, illus.
Specian, R. D.; and Ubelaker, J. E., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 59-65
Seuratum cancellatum, redescription, cephalic morphology, external cuticular modifications, scanning electron microscopy
Anthrozous pallidus (small intestine)
Eptesicus fuscus (body cavity)
Eumops perotis " "
Myotis californicus (body cavity)
M. yumanesis (body cavity)
Plecotus townsendii (body cavity)
Tadarida brasiliensis (body cavity)
Pipistrellus hesperus (body cavity)
all from Black Gap Wildlife Management Area, Brewster County, Texas.
- Seuratum mucronatum*
Durette-Desset, M.-C.; and Chabaud, A.-G., 1975, Ann. Parasitol., v. 50 (3), 303-337
Plecotus auritus: Commugny, Vaud, Suisse
Tadarina teniotis: Col de Bretolet, Valais, Suisse
- Seurocyrnea* Strand, 1929
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cyrnea* Seurat, 1914
- Severianoia severiano* (Schwenk, 1926) Basir, 1956
Leong, L.; and Paran, T. P., 1966, Med. J. Malaya, v. 20 (4), 349
Periplaneta americana: Singapore
- Sexansocara* Sobolev & Sudarikov, 1939, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Sicarius* Li, 1934, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Habronematinae
key
- Sicarius caudatus* n. sp., illus.
Quentin, J. C.; and Wertheim, G., 1975, Ann. Parasitol., v. 50 (1), 63-85
Pycnonotus capensis: Jerusalem, Israel
- Sicarius dipterum*
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Pisterotarsa gigantea subsp. *zoubkoffi*: Turkmenia

- Simondsia Cobbold, 1864, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascaropsinae
key
- Sincosta Roe, 1929
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Heligmosomoides Hall, 1916
- Singhfilaria hayesi
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (connective tissue): Florida
- Sinicaspirura Skrjabin et al., 1963
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Gendrespirura Chabaud, 1958
- Skrjabillaninae (Shigin & Shigina, 1958, fam.)
Chabaud, 1965
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Anguillicolidae
key; key to genera
includes: Molnaria; Skrjabillanus
- Skrjabillanus Shigin & Shigina, 1958, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Skrjabillaninae
key
Syn.: Agrachanus Tichomirova in Skrjabin et al., 1971
- Skrjabinagia (Kassimov, 1942) Altaev, 1952
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
synonymy
- Skrjabinagia boeivi
Bryan, R. P.; Bainbridge, M. J.; and Kerr, J. D., 1976, Austral. J. Zool., v. 24 (3), 417-421
Bubalus bubalis (large and small intestine): Northern Territory, Australia
cattle (large and small intestine): Northern Territory, Australia
Bos taurus (exper.)
- Skrjabinagia kolchida (Popova, 1937) Andreeva, 1957
Drozd, J.; and Bylund, G., 1970, Acta Parasitol. Polon., v. 17 (20-38), 259-260
Alces alces (abomasa): Poland
- Skrjabinagia kolchida (Popova, 1937) Andreeva, 1956
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
synonymy
Capreolus capreolus (rennet, small intestine): southern Bulgaria
- Skrjabinagia kolchida
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Skrjabinagia kolchida
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Skrjabinagia odocoilei
Eve, J. H.; and Kellogg, F. E., 1977, J. Wildlife Management, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (Odocoileus virginianus) density: south-eastern United States
- Skrjabinagia odocoilei
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of Odocoileus virginianus and Ovis aries on common range, deer unlikely reservoir host for sheep parasites
Odocoileus virginianus: Hardy County, West Virginia
- Skrjabinagia odocoilei (Dikmans, 1931)
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Cumberland County, New Jersey; Oklahoma
- Skrjabinalius Delyamure, in Skrjabin 1942
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
key
- Skrjabinaria heteromorpha Kreis
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Phoca vitulina (heart): European waters
- Skrjabinaria spirocauda Leidy
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Phoca vitulina (heart): European waters
- Skrjabinema ovis, illus.
Hanuskova, Z.; and Tilc, K., 1974, Veterinarstvi, v. 24 (11), 516
Cameroon goats: Brnenske zoo
- Skrjabinema ovis (Skrjabin, 1915) Vereshchagin, 1926, illus.
Martinez Gomez, F.; Hernandez Rodriguez, S.; and Calero Carretero, R., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 331-336
description
Ovis aries (ciego)
cabra
all from Spain
- Skrjabinema ovis
Martinez Gomez, F.; Hernandez Rodriguez, S.; and Calero Carretero, R., 1973, Rev. Iber. Parasitol., v. 33 (4), 625-631
Capra hircus: Municipal Slaughterhouse, Cordoba, Spain

- Skrjabinema ovis*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Skrjabinema ovis*
Wilson, D. E.; and Hirst, S. M., 1977, Wild-life Monogr. (54), Suppl., 3-111
Hippotragus equinus: Percy Fyfe Nature Reserve, South Africa
- Skrjabinogylus nasicola* (Leuckart 1842)
Duncan, N., 1976, Mammal Rev., v. 6 (2), 63-74
Skrjabinogylus nasicola, theoretical aspects of transmission to stoats and weasels based on laboratory study of food habits under conditions of food abundance and food shortage; extent of skull damage in weasels
Mustela erminea
M. nivalis
all from Newbrough, Aberdeenshire
- Skrjabinogylus nasicola*, illus.
King, C. M., 1977, J. Zool., London, v. 182 (2), 225-249
Skrjabinogylus nasicola, incidence and extent of damage caused in *Mustela nivalis*, climate, age, sex and body size of host: Britain
- Skrjabinogylus nasicola* (Leuckart, 1845) Petrov, 1927
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Mustela erminea: Pechora river basin
- Skrjabinogylus nasicola* (Leuckart, 1842) Petrov, 1927
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Mustela lutreola
Mustela nivalis
Mustela putorius
Mustela erminea
(frontal and nasal cavities of all): all from Karelia
- Skrjabinogylus petrowi* Bageanov, 1936
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Gulo gulo
(frontal and nasal cavities of all): all from Karelia
- Skrjabinispirura* subgen. of *Oxyspirura*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Hamulofilaria* subgen. of *Oxyspirura*
- Skrjabinitectus* Majumdar & Banerjee, 1966
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
"too poorly described to be assigned to any family"
- Skrjabinobronema* Guschanskaya, 1950
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Ancyracanthopsis* Diesing, 1861
- Skrjabinocapillaria eubursata* Skarbilovitsch, 1946
Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
ecological analysis of bat helminth fauna, geographic distribution
Rhinolophus hipposideros
Myotis daubentonii
M. mystacinus
Plecotus auritus
Nyctalus leisleri
N. noctula
Eptesicus serotinus
all from Moldavia
- Skrjabinocara* Kurashvili, 1941
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Chordocephalus* [sic] Alegret, 1941
"We presume that *Chordocephalus* has priority but this is not certain."
- Skrjabinocerca* Shikhobalova, 1930, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Skrjabinocerca prima* Schikhobalova, 1930
Aleksiev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Halcyon pileata: Rimsko-Korsakov islands
- Skrjabinocerca prima* Schikhobalova, 1930
Tsimbaliuk, A. K.; et al., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 129-152
Orchestia ochotensis (body cavity)
Uria aalge (esophagus)
Lunda cirrhata "
Charadrius mongolus (esophagus)
Calidris alpina (esophagus)
C. maritima (esophagus)
Tringa incana "
Arenaria interpres (esophagus)
Phalaropus lobatus "
Numenius phaeopus "
Motacilla alba (esophagus)
Anthus gustavi "
Calcarius lapponicus (esophagus)
Plectrophenax nivalis "
Troglodytes troglodytes (esophagus)
Cuculus canorus (esophagus)
Rattus norvegicus "
all from Bering Island
- Skrjabinocercina*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Cylicospirura*
key

- Skrjabinochona Guschanskaja, 1931
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cyrnea* Seurat, 1914
- Skrjabinoclava Sobolev, 1943, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
Syn.: *Cordonema* Schmidt & Kuntz, 1971
- Skrjabinoclava brevispicula Bondarenko et Daija, in litt. [nomen nudum]
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Limosa limosa lapponica: lower Yenisei
- Skrjabinoclava decorata (Solonitzin, 1928)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Xenus cinereus: lower Yenisei and Keta lake
- Skrjabinoclava halcyoni Ryjnikov et Hohlova, 1964
Aleksiev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Halcyon pileata (intestine): Rimsko-Korsakov islands
- Skrjabinoclava horrida (Rudolphi, 1809)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Heteroscelus incanus brevipes
Charadrius hiaticula
Numenius ph. phaeopus
all from Keta lake
- Skrjabinoclava horridae (Rud., 1809)
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Stercorarius longicaudatus: Yenisei
- Skrjabinoclava soricus (Tiner, 1951)
Babaev, Ia.; and Kolodenco, A. I., 1975, Izvest. Akad. Nauk Turkmen. SSR; s. Biol. Nauk (4), 71-75
[*Crocidura suaveolens*]: Turkmenistan
- Skrjabinoclava thapari Teixeira de Freitas, 1953
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (proventriculus): Florida
- Skrjabinodon medinae (Calvente, 1948) n. comb.
Specian, R. D.; and Ubelaker, J. E., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 46-51
Syn.: *Pharyngodon medinae* Calvente, 1948
- Skrjabinofilaria Travassos, 1925
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
key
- Skrjabinofilaria philanderi (Foster, 1939)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
as syn. of *Skrjabinofilaria skrjabini* Travassos, 1925
- Skrjabinofilaria skrjabini
Bain, O., 1976, Bull. World Health Organ., v. 54 (4), 397-401
human filariasis, number of developing and infective larvae dependent upon number of microfilariae penetrating into haemocoel of vector, relationship based on proportionality, facilitation and limitation, application to disease control and treatment methods
- Skrjabinofilaria skrjabini Travassos, 1925, illus.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
Syn.: ? *Skrjabinofilaria philanderi* (Foster, 1939)
- Skrjabinofilaria skrjabini
Mougey, Y.; and Bain, O., 1976, Ann. Parasitol., v. 51 (1), 95-110
5 filaria-vector sets, stochastic models for assessing relation between numbers of microfilariae ingested and numbers of microfilariae passing into hemocoel, limitation phenomenon
- Skrjabinoptera Schulz, 1927, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Physalopterinae
key
- Skrjabinoptera sp. larvae
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of *Tenebrionidae* as intermediate hosts
Trigonoscelis gemmulata
Blaps fausti bactriana
all from Tadzhik SSR [and/or] Uzbek SSR
- Skrjabinoptera phrynosoma
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus magister
Sceloporus undulatus
(stomach of all): all from Great Basin and Upper Colorado Plateau, Utah
- Smetaleksenema* Schmidt & Kuntz, 1972
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Ancyracanthopsis* Diesing, 1861
- Soboleviatia* new gen. [nomen nudum]
Aleksiev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104

- Soboleviatia pileati Smetanina et Alexejev in lit. [nomen nudum]
 Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
 Halcyon pileata (muscular stomach): Rimsko-Korsakov islands
- Sobolevicephalus Parukhin, 1964
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
 as syn. of Hadjelia Seurat, 1916
- Sobolevingylus sp. Romanov, 1952
 Craig, R. E.; and Borecky, R. A., 1976, Canad. J. Zool., v. 54 (5), 806-807
 Martes pennanti (lungs): Ontario
- Sobolevingylus petrowi Romanov, 1952
 Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
 Martes martes
 Gulo gulo
 all from Karelia
- Soboliphyme sp.
 Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
 Mustela erminea: Pechora river basin
- Soboliphyme baturini (Petrov, 1930). illus.
 Bogoiavlenskii, Iu. K.; and Khatkevich, L. M., 1970, Parazitologiya, Leningrad, v. 4 (3), 223-230
 4 spp. of Diactophymata, fine structure of somatic musculature, distribution of DNA and RNA
- Spauligodon paratectipenis (Chabaud and Golvan, 1957) n. comb.
 Specian, R. D.; and Ubelaker, J. E., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 46-51
 Syns.: Pharyngodon paratectipenis Chabaud and Golvan, 1957; Pharyngodon tectipenis sensu Calvente, 1948 nec P. tectipenis Ge-doeIst, 1919
- Sphaerularia bombi Dufour
 Lundberg, H.; and Svensson, B. G., 1975, Norsk Entom. Tidsskr., v. 22 (2), 129-134
 Sphaerularia bombi, bumble-bee queens, correlation between parasitism and behavior (nest-seeking, hibernacula-seeking and foraging behavior)
 Bombus alpinus (abdomen)
 B. balteatus "
 B. hypnorum "
 B. jonellus "
 B. pratorum "
 B. lapponicus "
 B. pascuorum "
 B. lucorum "
 all from Abisko, northern Sweden
- Sphaerularia bombi (Dufour, 1837), illus.
 Madel, G.; and Scholtzseck, E., 1976, Ztschr. Parasitenk., v. 49 (1), 81-92
 Sphaerularia bombi, evaginated tubular reproductive tract, light and electron microscopy of tube cells, cytochemistry of fat body cells of parasite
 Bombus terrestris
 B. lapidarius
 B. pratorum
 all from Botanischer Garten, Bonn
- Spiculimermis uncus sp. n., illus.
 Gafurov, A. K., 1976, Dokl. Akad. Nauk Tadzhiksk. SSR, v. 19 (9), 58-62
 Diamesa sp. (posterior half of body):
 Varzobsk gorge, 1 km. from Takob settlement; Khodzha-Obigarm; 14 km. south of Varzob river
- Spiculocaulus austriacus (Gebauer, 1932) Dougherty and Goble, 1946
 Polley, L.; and Hoerning, B., 1977, Rev. Suisse Zool., v. 84 (3), 675-680
 Spiculocaulus austriacus, morphometry
 Rupicapra rupicapra (tracheas and larger bronchi): Switzerland
- Spiculopteragia Orloff, 1933
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Trichostrongylidae, Ostertagiinae
 synonymy
- Spiculopteragia [sp.]
 Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuaris, v. 3 (1), 3-7
 gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Spiculopteragia alcis
 Reh binder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
 reindeer (abomasum): Sweden
- Spiculopteragia (Ostertagia) asymmetrica (Ware, 1925), illus.
 Goffredo, G.; and Sobrero, R., 1972, Parassitologia, v. 14 (1), 143-148
 Dama dama (intestine): foresta Umbra (promontorio garganico, provincia di Foggia)
- Spiculopteragia asymmetrica (Ware, 1925) Orloff, 1933
 Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
 Capreolus capreolus (rennet): southern Bulgaria
- Spiculopteragia boehmi
 Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
 roe deer (digestive tract): Czechoslovakia

- Spiculopteragia boehmi*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Spiculopteragia boehmi*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Spiculopteragia boehmi*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Spiculopteragia boehmi* Gebauer, 1931
Rojo Vazquez, F. A.; and Cordero del Campillo, M., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 41-47
valid species
- Spiculopteragia dagestanica* (Altaev, 1953)
Andreeva, 1957
Drozd, J.; and Bylund, G., 1970, Acta Parasitol. Polon., v. 17 (20-38), 259-260
Alces alces (abomasa): Poland
- Spiculopteragia peruvianus*
Guerrero, C.; Rojas, M.; and Vargas, J., 1974, Rev. Invest. Pecuarias, v. 3 (1), 9-14
gastrointestinal nematodes, alpacas, activity of 1-tetramisole, significant body weight gain in treated animals
- Spiculopteragia peruvianus*
Vargas, J.; Guerrero, C.; and Rojas, M., 1972, Rev. Invest. Pecuarias, v. 1 (2), 137-144
levamisole, nematodes of alpacas, slight toxicity
- Spiculopteragia spiculoptera* (Guschanskaja, 1931) Orloff, 1933
Ianchev, I., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 205-220
synonymy
Capreolus capreolus (rennet): southern Bulgaria
- Spiculopteragia spiculoptera*
Rehbinder, C.; and Christensson, D., 1977, Nord. Vet.-Med., v. 29 (12), 556-557
reindeer (abomasum): Sweden
- Spiculopteragia spiculoptera* (Guschanskaia, 1931), illus.
Rojo Vazquez, F. A.; and Cordero del Campillo, M., 1977, An. Fac. Vet. Leon, Oviedo, v. 21 (21), 1975, 41-47
valid species, morphological characteristics of males and females, hosts and geographical distribution listed
Cervus elaphus: Portilla de la Reina (Leon), Iberian Peninsula
- Spiculopteragia spiculoptera*
Schweigsut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Spinicauda komodoensis* sp. n., illus.
Pinnell, J. L.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 337-340
Psammodynastes pulverulentus (intestine): Komodo Island, Indonesia
- Spinitectoides* Petter, 1969, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Spinitectus* Fourment, 1883, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Cystidicolidae
key
- Spinitectus* sp.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Auxis thazard (stomach): South China Sea
- Spinitectus* [sp.]
Von Essen, S. G., 1976, Proc. Nebraska Acad. Sc., 27
fish
Hexagenia sp.
all from Nebraska lakes
- Spinitectus agonostomi* Moravec et Barus, 1971, illus.
Petter, A. J.; Golvan, Y. J.; and Tchepprakoff, R., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 159-171
description
Agonostomus monticola (intestin): rivieres de la Basse Terre, Guadeloupe
- Spinitectus allaeri* Campana-Rouget, 1961
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Botan. Africaines, v. 87 (2), 209-248
Xenoclarias eupogon (stomach): Lake Victoria, Uganda
- Spinitectus caballeri* sp. nov., illus.
Datta, B. K.; and Majumdar, G., [1974], An. Inst. Biol. Univ. Nac. Auton. Mexico, s. Cien. Mar y Limnol., v. 43 (1), 1972, 85-92
Bagarius bagarius (stomach): Kansain river, Midnapore, West Bengal, India
- Spinitectus carolini*
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Micropterus salmoides
M. punctulatus
Lepomis gulosus
Pomoxis annularis
L. macrochirus
L. megalotis
all from Eagle Mountain Lake, Texas

- Spinitectus carolini*
Niederborn, J. Y., 1974, Tr. Missouri Acad. Sci., v. 7-8, 1973-1974, 160-163
Lepomis cynellus: Johnson County, Missouri
- Spinitectus carolini*
Rubertone, J. A.; and Hall, J. E., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 58-59
Ambloplites rupestris
Lepomis auritus
L. gibbosus
Lepomis sp.
Micropterus dolomieu
(intestine of all): all from Greenbrier River below Alderson, West Virginia
- Spinitectus gracilis* (Ward and Magath, 1916)
Baker, J. C.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 37-39
Ictalurus punctatus (intestines): island region of western Lake Erie
- Spinitectus gracilis*
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Ictalurus punctatus
Aplodinotus grunniens
(intestine of all): all from Eagle Mountain Lake, Texas
- Spinitectus gracilis* Ward and Magath, 1917, illus.
Keppner, E. J., 1975, Am. Midland Naturalist, v. 93 (2), 411-423
larvae compared with *S. micracanthus*
Hexagenia sp.: Racehorse Lake, Johnson Co., Missouri
- Spinitectus micracanthus* Christian, 1972, illus.
Keppner, E. J., 1975, Am. Midland Naturalist, v. 93 (2), 411-423
life cycle, development; larvae compared with *S. gracilis*
Lepomis macrochirus (intestine)(nat. and exper.): Lake Morel and a farm pond, Johnson Co., Missouri
Hexagenia sp. (abdominal muscles)(exper.)
- Spinitectus* (?) *mollis* sp. nov., illus.
Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
Euthynnus affinis (stomach)
Auxis thazard
all from South China Sea
- Spinitectus pandharinathi* n. sp., illus.
Kalyankar, S. D., 1973, Riv. Parassitol., Roma, v. 34 (1), 55-58
Ophiocephalus striatus (stomach): Aurangabad (Maharashtra, India)
- Spinitectus seenghalai* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Mystus seenghala (intestine): Taunsa Barrage, Pakistan
- Spinostromylus* Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylineae
- Spiralatus* Chabaud, Brygoo & Durette, 1963, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirocercinae
key
- Spirocamallanus* Olsen (1952)
Akram, M., 1975, Biologia, Lahore, v. 21 (2), 93-100
as syn. of *Procamallanus* Baylis (1923)
- Spirocamallanus* Olsen, 1952
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
review
- Spirocamallanus* Olsen, 1952, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Camallanidae
key
- Spirocamallanus* sp.
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Botan. Africaines, v. 87 (2), 209-248
Synodontis afro-fischeri (stomach): Lake Victoria, Uganda
- Spirocamallanus bagarii* (Karve y Naik, 1951) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus berdii* Khan and Yaseen, 1969, illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
female redescribed
Sillago sihama (intestine): Fish Harbour, Karachi, Pakistan
- Spirocamallanus chimusensis* sp. n., illus.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
Pygidium punctulatum (intestino): Chilete, Cajamarca, Peru
- Spirocamallanus crossorhombi* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Crossorhombus azureus (intestine): Fish Harbour, Karachi, Pakistan
- Spirocamallanus dessetae* n. sp., illus.
Petter, A. J.; Golvan, Y. J.; and Tchepprakoff, R., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 159-171
Agonostomus monticola (intestine): Grande Riviere a Goyaves (Domaine Duclos) and other rivers of Guadeloupe
- Spirocamallanus* (*Procamallanus*) *fulvidraconis* (Li, 1935)
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224

- Spirocamallanus globoconchus* (Ali, 1960) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus gubernaculus* (Khera, 1955) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus* (*Procamallanus*) *hyderabadensis* (Ali, 1956) n. comb.
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
- Spirocamallanus hyderabadensis* (Ali, 1956) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus inglisi* n. sp., illus.
Bashirullah, A. K. M.; and Hafizuddin, A. K. M., 1973, Riv. Parassitol., Roma, v. 34 (2), 114-119
Clupisoma murius (stomachs and intestines):
Dacca, Bangladesh
- Spirocamallanus notopteri* n. sp., illus.
Bashirullah, A. K. M.; and Hafizuddin, A. K. M., 1973, Riv. Parassitol., Roma, v. 34 (2), 114-119
Notopterus notopterus (stomachs and intestines): Dacca, Bangladesh
- Spirocamallanus olseni* n. sp., illus.
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
Channa (*Ophiocephalus*) *striatus* (intestine):
Dacca, Bangladesh
- Spirocamallanus ophicephalus* (Ali, 1960) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus pereirai*
Haaker, P. L., 1975, Fish Bull. (165), State Calif., Resources Agency, Dept. Fish and Game, 137-151
Paralichthys californicus (digestive tract):
Anaheim Bay
- Spirocamallanus pereirai*
Tasto, R. N., 1975, Fish Bull. (165), State Calif., Resources Agency, Dept. Fish and Game, 123-135
Leptocottus armatus (stomach, gastrointestinal tract): Anaheim Bay
- Spirocamallanus* (*Procamallanus*) *singhi* (Ali, 1956) n. comb.
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
- Spirocamallanus singhi* (Ali, 1956) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocamallanus spiralis* (Baylis, 1923)
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Botan. Africaines, v. 87 (2), 209-248
Synodontis victoriae (stomach): Lake Victoria, Uganda
- Spirocamallanus timmi* n. sp., illus.
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
Mystus cavasius
M. vittatus
Heteropneustes fossilis
(stomach of all): all from Sunamganj, Sylhet, Bangladesh
- Spirocamallanus* (*Procamallanus*) *viviparus* (Ali, 1956) n. comb.
Bashirullah, A. K. M., 1973, Am. Midland Naturalist, v. 90 (1), 221-224
- Spirocamallanus viviparus* (Ali, 1956) comb. n.
Teixeira de Freitas, J. F.; and Ibanez, N., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 146-148
- Spirocaudata* Sharma, 1971
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Viguiera* Seurat, 1913
- Spirocerca*
Rajan, A.; and Mohiyuddeen, S., 1975, Kerala J. Vet. Sc., v. 5 (2), 139-142
spirocercosis, pathology, dogs (thoracic oesophagus, aorta, stomach, thymus, lung, bronchial and mediastinal lymphnodes): Bangalore city
- Spirocerca* sp., illus.
Chattopadhyay, S. K.; and Sharma, R. M., 1972, Indian J. Animal Sc., v. 42 (9), 698-705
Spirocerca sp., lesions in aortas of sheep and goats, pathological studies
- Spirocerca arctica* Petrov, 1927
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Vulpes vulpes
Alopex lagopus
all from Pechora river basin
- Spirocerca lupi*
Bannor, T. T., 1976, Vet. Rec., v. 98 (15), 302
Spirocerca lupi, cause of fatal hemorrhage in alsatian dog, case history: Ghana
- Spirocerca lupi* (Rudolphi 1809)
Blancou, J.; and Albignac, R., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (2), 127-130
Spirocerca lupi, high mortality from aneurysms of thoracic aorta of lemurs
Lemur fulvus rufus
L. f. sanfordi
L. f. fulvus
L. f. collaris
L. macaco macaco
L. catta
all from zoological park

- Spirocerca lupi*
Brodey, R.S.; et al., 1977, *Vet. Parasitol.*, v. 3 (1), 49-59
Spirocerca lupi in native and pet dogs, prevalence, localization, clinicopathological and laboratory findings, epizootologic observations; also observed in *Canis mesomelas*: Kenya
- Spirocerca lupi*, *illus.*
Bwangamoi, O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (4), 363-370
recovery from dogs using Lindsey's method: case report, dog (oesophagus, aorta, stomach, lung, caecum): Uganda
- Spirocerca lupi* (Rudolphi, 1809) Chitwood, 1933, *illus.*
Chhabra, R. C.; and Singh, K. S., 1977, *Indian J. Animal Sc.*, v. 47 (1), 42-47
Spirocerca lupi, pathological changes in stomach, aorta and oesophagus of dogs (exper.)
- Spirocerca lupi* (Rudolphi, 1809), *illus.*
Chhabra, R. C.; and Singh, K. S., 1977, *Indian J. Animal Sc.*, v. 47 (4), 178-184
Spirocerca lupi, third-stage juveniles, morphology, life history, effect of temperature and humidity on development, viability, and infectivity
Onthophagus bonasus
O. dama
Onitis philemon
Catharsius pithecius
dog
(all exper.)
- Spirocerca lupi*
Corkish, J. D., 1977, *Trop. Animal Health and Prod.*, v. 9 (2), 81-84
prevalence in dogs (oesophagus, aorta, stomach, pleura, mediastinum, lung), post mortem examination: Accra, Ghana
- Spirocerca lupi*
Danks, B. C.; Adams, J. W. E.; and Roberts, H. M., 1977, *Rhodesian Vet. J.*, v. 7 (4), 82-85
dogs, pathology, clinical diagnosis, surgery, review
- Spirocerca lupi*, *illus.*
Ivoghli, B., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (8), 834
Spirocerca lupi, dog, fatal aortic aneurysm and rupture: Iran
- Spirocerca lupi*, *illus.*
Mayaudon T., H.; and Hoepf, A., 1972, *Rev. Med. Vet. y Parasitol.*, Maracay, v. 24 (1-8), 1971-1972, 57-60
Spirocerca lupi, atypical localization, causing rectal prolapse
Canis familiaris (rectum)
- Spirocerca lupi*
Mullin, S. W.; Stevens, S.; and Min, L. L., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (1), 90 [Demonstration]
dogs (oesophageal and aortic nodules): Kuala Lumpur, Malaysia
- Spirocerca lupi*
Nakasala-Situma, J., 1976, *Magy. Allat. Lapja*, v. 98, v. 31 (3), 185-187
Spirocerca lupi, case histories, symptoms, relationship to secondary pulmonary osteoarthropathy as a possible complication
dogs: Uganda
- Spirocerca lupi*, *illus.*
Ndiritu, C. G.; and Al-Sadi, H. I., 1976, *Mod. Vet. Pract.*, v. 57 (11), 924, 930, 934
Spirocerca lupi, dogs, pathogenesis and lesions, clinical aspects reviewed
- Spirocerca lupi*
Petrick, S. W., 1977, *J. South African Vet. Ass.*, v. 48 (2), 105-107
Spirocerca lupi, *Filaroides osleri*, ascarids, dogs, gastrointestinal fibroscope, useful diagnostic aid
- Spirocerca lupi*, *illus.*
Puliaevskaia, N. V., 1969, *Trudy Gel'mint. Lab.*, Akad. Nauk SSSR, v. 20, 119-122
Metastrongylus salmi and *Spirocerca lupi* females, morphology of genital tract
- Spirocerca lupi*, *illus.*
Radhakrishnan, C. V.; and Tirgari, M., 1976, *Indian Vet. J.*, v. 53 (8), 627-631
Spirocerca lupi, dogs, incidence survey, diagnosis, pathological findings: Iran
- Spirocerca lupi*
Retnasabapathy, A.; and Khoo Teik San, 1976, *Malaysian Vet. J.*, v. 6 (2), 69-71
Spirocerca lupi, dogs (thoracic oesophagus, aorta), incidence, disophenol, good results: Petaling Jaya
- Spirocerca lupi*, *illus.*
Singh, N. P.; and Tewari, A. N., 1976, *Indian J. Animal Sc.*, v. 46 (4), 211-214
Spirocerca lupi, pathoanatomical and histopathological study, dogs, oesophagus and aorta, fibrosarcoma in some cases
- Spirocerca lupi*
Thornton, J. E.; Bell, R. R.; and Reardon, M. J., 1974, *J. Wildlife Dis.*, v. 10 (3), 232-236
Canis latrans (wall of the thoracic esophagus): Nueces County, Texas
- Spirocerca lupi*, *illus.*
Wandera, J. G., 1976, *Vet. Rec.*, v. 99 (18), 348-351
Spirocerca lupi, dogs, incidence, pathological variations, oesophageal sarcomas, age of host, site of incidence, 11 year period: Kenya
- Spirocerca sanguinolenta*
Rep, B. H.; and Heinemann, D. W., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 104-110
dog (oesophageal wall): Surinam

- Spirocercella* Thwaite, 1928
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Leiuris* Leuckart, 1850
- Spirocercidae* (Chitwood & Wehr, 1932, subfam.)
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruroidea
key; key to subfamilies
includes: Spirocercinae; Ascaropsinae; Mastophorinae
- Spirocercinae* Chitwood & Wehr, 1932
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirocercidae
key; key to genera
includes: *Spiralatus*; *Cylicospirura*; *Spirocerca*; *Didelphonema*; *Cyathospirura*; *Vigispirura*
- Spironoura* sp., larval
Ernst, E. M.; and Ernst, C. H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 176-178
Chrysemys picta: Prince Georges County, Maryland
Terrapene carolina: Montgomery County, Maryland
- Spironoura armenica*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Emys orbicularis: Macedonia, Yugoslavia
- Spironoura petrei* Khalil, 1970
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Botan. Africaines, v. 87 (2), 209-248
Distichodus niloticus (intestine): Lake Albert, Uganda
- Spironoura tikasinghi* sp. n., illus.
Schoenecker, S. A.; Schmidt, G. D.; and Everard, C. O. R., 1977, J. Parasitol., v. 63 (2), 341-343
Geoemyda punctularia (cecum, ileum): Aripo Savannah, Ture Forest, Trinidad, West Indies
- Spirophilometra* Parukhin, 1971, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Philometrinae
key
- Spiroptera* Rudolphi, 1819
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus* Dujardin, 1845
- Spiroptera robusta* Linstow, 1903
Specian, R. D.; Ubelaker, J. E.; and Dailey, M. D., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 14-21
as syn. of *Proleptus acutus* Dujardin, 1845
- Spirostrongylus gallardi*: Mawson, 1955
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
as syn. of *Cyclostrongylus wallabiae* Johnston & Mawson, 1939
- Spirostrongylus kartana*
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
as syn. of *Cyclostrongylus kartana* (Mawson 1955) [n. comb.]
- Spirostrongylus parma*: Mawson, 1955
Mawson, P. M., 1977, Tr. Roy. Soc. South Australia, v. 101 (1), 19-20
as syn. of *Cyclostrongylus parma* (Johnston & Mawson 1939) [n. comb.]
- Spiroxyinae* Baylis & Lane, 1920
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatidae
key
includes: Spiroxys
- Spiroxys* Schneider, 1866, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spiroxyinae
- Spiroxys* sp.
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Lepomis megalotis
L. microlophus
(mesenteries of all): all from Eagle Mountain Lake, Texas
- Spiroxys* sp.
Hensley, G. H.; and Nahhas, F. M., 1975, Calif. Fish and Game, v. 61 (4), 201-208
Morone saxatilis (intestine, stomach, caeca, mesentery): Sacramento-San Joaquin Delta, California
- Spiroxys* sp. Schneider, 1866
Platt, T. R., 1977, Ohio J. Sc., v. 77 (2), 97-98
Emydoidea blandingii (stomach): Ottawa National Wildlife Refuge, Ottawa Co., Ohio
- Spiroxys contortus* Rudolphi, 1819
Kulakiv'ska, O. P., 1976, Vestnik. Zool., Akad. Nauk Ukrainsk. SSR, Inst. Zool. (4), 82-84
Umbra crameri (mucous membrane of intestine): Duna delta
- Spiroxys contortus* (Rudolphi, 1819)
Platt, T. R., 1977, Ohio J. Sc., v. 77 (2), 97-98
Chrysemys picta marginata
Emydoidea blandingii
(stomach of all): all from Ottawa National Wildlife Refuge, Ottawa Co., Ohio

- Spiroxys figueiredoi* Freitas & Dobbin Jr., 1962, illus.
Vicente, J. J., 1966, Atas Soc. Biol. Rio de Janeiro, v. 10 (1), 7-8
description
Kinosternon scorpioides scorpioides (estomago): Rio Bujaru, Estado do Para, Brasil
- Spirura* Blanchard, 1849
Babero, B. B., 1973, Tr. Am. Micr. Soc., v. 92 (2), 265-272
Spirurinae; key to North American species, includes: *Spirura michiganensis*; *S. leucurusi* n. sp.; *S. infundibuliformis*; *S. zapi*
- Spirura* Blanchard, 1849, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruridae
key; synonymy
- Spirura* Blanchard, 1849
Khalil, L. F., 1975, J. Helminth., v. 49 (2), 93-99
key to African species, includes: *S. hipposiderosi* n. sp.; *S. diplocyphos*; *S. nycterisi* n. sp.; *S. spinicaudata*; *S. congolense*; *S. dentata*; *S. rothschildi*; *S. rytipleurites seurati*; *S. portesiana*
- Spirura*
Quentin, J. Cl.; and Krishnasamy, M., [1976], Ann. Parasitol., v. 50 (6), 1975, 795-812
Spirura, evolution and distribution
- Spirura* (s.l.) sp. (Johnston & Mawson, 1952)
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
as syn. of *Stammerinema* sp.
- Spirura* sp.
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Trigonoscelis gemmulata
Blaps fausti bactriana
Cyphogenia gibba
Pachyscelis laevicollis
all from Tadzhik SSR [and/or] Uzbek SSR
- Spirura* sp. (Gafurov, 1968)
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Cyphostete komarovi: Turkmenia
- Spirura* sp. 1
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Adesmia servillei schatzmayri: Turkmenia
- Spirura aurangabadensis* (Ali et Lovekar, 1966), illus.
Quentin, J. Cl.; and Krishnasamy, M., [1976], Ann. Parasitol., v. 50 (6), 1975, 795-812
description
Nycticebus coucang
Tupaia glis
all from Ulu Gombak, Forest reserve, Gombak, Selangor, Malaisie occidentale
- Spirura hipposiderosi* n. sp., illus.
Khalil, L. F., 1975, J. Helminth., v. 49 (2), 93-99
key
Hipposideros ruber (stomach): Kisarawe, Tanzania
- Spirura leucurusi* n. sp., illus.
Babero, B. B., 1973, Tr. Am. Micr. Soc., v. 92 (2), 265-272
key
Ammospermophilus leucurus (stomach and small intestine): Clark County, Nevada
- Spirura malayensis* n. sp., illus.
Quentin, J. Cl.; and Krishnasamy, M., [1976], Ann. Parasitol., v. 50 (6), 1975, 795-812
Tupaia glis: W. Malaysia, Selangor Gombak, Ulu Gombak, Forest Reserve
T. minor: W. Malaysia, Selangor Gombak, Ulu Gombak, Forest Reserve
Nycticebus coucang: Kota Kinabalu, Sabah
Blatella germanica (exper.)
- Spirura nycterisi* n. sp., illus.
Khalil, L. F., 1975, J. Helminth., v. 49 (2), 93-99
key
Nycteris thebaica
Hipposideros ruber
(stomach of both): both from Kisarawe, Tanzania
- Spiruracerca* Erickson, 1938
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Spirura* Blanchard, 1849
- Spirurata* gen. sp.
Sharpilo, L. D., 1976, Vestnik Zool., Akad. Nauk Ukrain. SSR, Inst. Zool. (1), 62-67
rodents as reservoir hosts for game and domestic animal infestation with larval helminths
[*Sicista subtilis*]: Ukraine
- Spirurida*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Secernentea
key; key to suborders
includes: Camallanina; Spirurina
- Spirurida*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
includes: Camallanina; Spirurina
- Spirurida* Chitwood, 1933
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Secernentea
includes: Camallanina; Spirurina
- Spiruridae* Oerley, 1885
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spiruroidea
key; key to genera
includes: *Paraspirura*; *Spirura*; *Protospirura*

- Spiruriden-type, illus.
Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Spirurides
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Spirurids, larval
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Spirurina
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurida
key; key to superfamilies
includes: Gnathostomatoidea; Physalopteroid-
ea; Rictularioidea; Thelazioidea; Spiruroi-
dea; Habronematoidea; Acuarioidea; Filarioi-
dea; Aproctoidea; Diplotriaeoidea
- Spirurina
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurida
includes: Gnathostomatoidea; Physalopteroid-
ea; Rictularioidea; Thelazioidea; Spiruroi-
dea; Habronematoidea; Acuarioidea; Filarioi-
dea; Aproctoidea; Diplotriaeoidea
- Spirurina Railliet, 1914
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Spirurida
- Spiruroid larvae, illus.
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
brief description
Antechinus stuartii (stomach digests): Victoria
- Spiruroid larvae, probably Tetrameres grusi
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (proventriculus, under gizzard lining): Florida
- Spiruroidea
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Spiruroidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
key
- Spiruroidea
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Spirurida
key to families
includes: Gongylonematidae; Spiruridae;
Spirocercidae; Hartertiidae
- Spiruroidea [sp.]
Coggins, J. R., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
parasitic fauna, effect of host diet and habitat
Turdus migratorius: Kellogg Bird Sanctuary, Michigan
- Spiruroidea [sp.] ova
Faust, B. S.; and Pappas, P. W., 1977, J. Zoo Animal Med., v. 8 (1), 18-23
Haliaeetus leucocephalus
Pavo muticus
(feces of all): all from Columbus (Ohio)
Zoo
- Spiruroidea larva
Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
Sceloporus occidentalis: Great Basin and Upper Colorado Plateau, Utah
- Spiruroidea [sp.]
Tasto, R. N., 1975, Fish Bull. (165), State Calif., Resources Agency, Dept. Fish and Game, 123-135
Leptocottus armatus (mesentery): Anaheim Bay
- Splendidofilaria Skrjabin, 1923
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key to males of species
- Splendidofilaria sp.
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (heart): Florida
- Splendidofilaria sp.
Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Splendidofilaria spp., illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
- Splendidofilaria algonquinensis
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (heart, innominate artery): Franklin and Ottawa counties, Ohio
- Splendidofilaria algonquinensis (Anderson, 1955), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria boehmi Supperer, 1958, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key

- Splendidofilaria brevispiculum* Singh, 1949, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria californiensis* (Wehr and Her-
man, 1956), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria caperata* Hibler, 1964, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria columbensis* n. sp., illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
Columba fasciata fasciata (fascia of thighs):
Colorado, U.S.A.
- Splendidofilaria falconis* (Sonin, 1966), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria fallisensis* (Anderson, 1954), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria gedoelsti* Travassos, 1926, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria gretillati* Chabaud, Anderson, and Brygoo, 1959, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria hiblieri* n. sp., illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
Columba fasciata fasciata (fascia of thighs):
Colorado, U.S.A.
- Splendidofilaria kashmirensis* Amir and Ali, 1960, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria mavis* (Leiper, 1909), illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria papilloerca* (Lubimov, 1946) illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria passerina* Koch and Huizinga, 1971, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria pawlowskyi* Skrjabin, 1923, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria pectoralis* Gibson, 1967, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria picacardina* Hibler, 1964, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria rotundicephala* Oschmarin, 1950, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria singhi* Sultana, 1962, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria tuvensis* Spassky and Sonin, 1957, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria verrucosa* Oschmarin, 1950, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilaria wehri* Anderson, 1961, illus.
Olsen, O. W.; and Braun, C. E., [1977], Great Basin Nat., v. 36 (4), 1976, 445-457
key
- Splendidofilariinae* [sp.]
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Wash-
ington, v. 42 (2), 119-127
Meleagris gallopavo (upper esophagus and crop?): Florida
- Sprattia* n. gen.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
Onchocercinae
tod: *S. venacavincola* (Spratt et Varughese, 1975) n. comb.
- Sprattia capilliforme* (Baylis, 1934) n. comb.
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Sprattia venacavincola* (Spratt et Varughese, 1975) n. comb. (tod)
Chabaud, A.-G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Squamane* van Thiel, 1925
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Parabronema* Baylis, 1921

- Squamastromylos Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558 as syn. of Heligmostrongylus Travassos, 1917
- Srivastavanema new rank
Durette-Desset, M. C.; and Lim, B. L., 1975, Ann. Parasitol., v. 50 (1), 87-96 raised from subgeneric to generic status, Brevistriatinae, definition, tod: *S. longispicularis* (Singh, 1962) n. comb.
- Srivastavanema (Singh, 1962)
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Srivastavanema (Singh, 1962)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Srivastavanema bhagwansinghi n. sp., illus.
Durette-Desset, M. C.; and Lim, B. L., 1975, Ann. Parasitol., v. 50 (1), 87-96
Petaurista petaurista (intestin grele):
Malaisie (Pahang, Montane Forest, Kampong Rajah, Cameron Highland)
- Srivastavanema longispicularis (Singh, 1962) n. comb. (tod)
Durette-Desset, M. C.; and Lim, B. L., 1975, Ann. Parasitol., v. 50 (1), 87-96
- Srivastavanema musasabi (Yamaguti, 1941) n. comb.
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Syn.: Longistriata musasabi Yamaguti, 1941
- Srivastavanema yapi n. sp., illus.
Durette-Desset, M. C.; and Lim, B. L., 1975, Ann. Parasitol., v. 50 (1), 87-96
Aeromys tephromelas: Malaisie (Pahang, Gunung Benom, Foret Primaire; Selangor, Bukit Lagong, Forest Reserve, Batu Kuala Lumpur; Selangor, Forest Reserve, Bukit Rajah, Klang)
Hylopetes spadiceus: Malaisie (Selangor, Forest Reserve, Bukit Rajah, Klang)
Petaurista petaurista (intestin grele):
Malaisie (Selangor, Bukit Lagong Forest Reserve, Batu, Kuala Lumpur; Pahang, Cameron Highland)
- Srivastavanema yapi Durette-Desset et Lim-Boo-Liat, 1975
Durette-Desset, M. C.; and Krishnasamy, M., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 697-710
Aeromys tephromelas (intestin grele): Pahang, Gunang, Benom
- Stammerinema Osche, 1955, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
- Stammerinema sp.
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
Syn.: *Spirura* (s.l.) sp. (Johnston & Mawson, 1952)
Hydromys chrysogaster: South Australia
- Stammerinema suffodiata sp. n., illus.
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
pathology
Antechinus stuartii: Healesville, Powelltown, and Sherbrooke, Victoria
Rattus fuscipes: Victoria
R. lutreolus: Victoria (stomach of all)
- Stammerinema suffodiata Beveridge & Barker
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (stomach): Powelltown, Victoria
- Stefanskostrongylus soricis (Soltys, 1954), illus.
Mas-Coma, S., 1977, Ann. Parasitol., v. 52 (4), 447-456
Sorex minutus (bronches): valles de Ribas (Pyrenees-orientales espagnoles) et de Viladrau (Montseny), Espagne
description
- Stefanskostrongylus soricis (Soltys, 1954), illus.
Mas-Coma, S.; and Gallego, J., 1975, Rev. Iber. Parasitol., v. 35 (3-4), 261-281
Sorex minutus (bronquios): Catalan Pyrenean Mountains
- Stegophorus Wehr, 1934, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key
Syn.: *Paryseria* Johnston, 1938
- Stegophorus stellae-polaris (Parona, 1901)
Alekseev, V. M.; and Smetanina, Z. B., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 97-104
Ciceronia pusilla (muscular stomach): Rimsko-Korsakov islands
- Stegophorus stellaepolaris (Parona, 1901), illus.
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
description, nematodes of *Larus canus* (esophagus), age and sex of host, seasonal variations, distribution in alimentary canal:
Agdenes, Norway

- Stegophorus stellaepolaris* (Parona, 1901), *illus.*
illus.
 Bakke, T. A.; and Barus, V., 1976, Norwegian
 J. Zool., v. 24 (3), 185-189
 redescription
Hydrobates pelagicus (ventriculus): island
 of Rost, Norway
 Plautus alle: Taelavag, Sotra, Hordaland
- Stegophorus stellae-polaris* (Parona, 1901)
 Sergeeva, T. P., 1969, Trudy Gel'mint. Lab.,
 Akad. Nauk SSSR, v. 20, 146-155
Stercorarius longicaudatus
Larus argentatus
 all from Yenisei
- Stegophorus stercorarii* nov. sp., *illus.*
 Leonov, V. A.; Sergeeva, T. P.; and Tsimbaliuk,
 A. K., 1966, Trudy Gel'mint. Lab., Akad. Nauk
 SSSR, v. 17, 91-94
Stercorarius longicaudatus: Chukotka; lower
 Enisei
Fratercula cirrhata: Chukotka
Fratercula corniculata: Chukotka
Aethia cristatella: Chukotka
Fulmarus glacialis: Komandorskiye Islands
 (under cuticle of muscular stomach of all)
- Stegophorus stercorarii* Leonov, Sergeeva et
 Cimbaluk, 1966, *illus.*
 Belogurov, O. I.; Leonov, V. A.; and Zueva,
 L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana
 (Skriabin), 105-124
 description
Stercorarius longicaudatus (muscular stomach):
 coast of Sea of Okhotsk (Ol'sk region)
- Stegophorus stercorarii* Leonov, Sergeeva et
 Zimbaluk, 1966
 Sergeeva, T. P., 1969, Trudy Gel'mint. Lab.,
 Akad. Nauk SSSR, v. 20, 146-155
Stercorarius longicaudatus
S. pomarinus
 all from Yenisei
- Steinernema*
 Mracek, Z., 1977, J. Invert. Path., v. 30 (1),
 87-94
 diagnostic characters distinguishing *Stein-*
ernema from *Neoaplectana*
- Steinernema kraussei*, *illus.*
 Mracek, Z., 1977, J. Invert. Path., v. 30 (1),
 87-94
 description of first giant generation, of
 normal-size generation, of female, of male,
 of larvae, and of specimens grown in axenic
 culture
Cephaloia abietis (body cavity): vicinity
 of Cesky Rudolec, southern Bohemia, Czecho-
 slovakia
- Steinernematidae* Chitwood and Chitwood, 1937
 Khan, A.; Brooks, W. M.; and Hirschmann, H.,
 1976, J. Nematol., v. 8 (2), 159-168
 diagnosis emended
- Stellobronema* Guschanskaja, 1937
 Chabaud, A. G., 1975, CIH Keys Nematode Para-
 sites Vertebrates (Anderson, Chabaud, and
 Willmott) (3), 29-58
 as syn. of *Hadjelia* Seurat, 1916
- Stellocaronema* Gilbert, 1930
 Chabaud, A. G., 1975, CIH Keys Nematode Para-
 sites Vertebrates (Anderson, Chabaud, and
 Willmott) (3), 29-58
Histiocephalinae
 key
- Stellocaronema fausti* (Li, 1934) Adams et Gibson,
 1969
 Quentin, J. C.; and Wertheim, G., 1975, Ann.
 Parasitol., v. 50 (1), 63-85
 redescription
Glareola pratincola
Hoploterus spinosus
Philomachus pugnax
 all from Israel
- Stellocaronema indica* sp. n., *illus.*
 Ali, M. M., 1970, Acta Parasitol. Polon.,
 v. 17 (20-38), 357-363
Metopodius indicus (gizzard): Hyderabad,
 Andhra Pradesh, India
- Stellocaronema skrjabini* Gilbert, 1930
 Bondarenko, S. K., 1969, Trudy Gel'mint. Lab.,
 Akad. Nauk SSSR, v. 20, 35-45
Philomachus pugnax
Charadrius hiaticula
Calidris temminckii
 all from lower Yenisei [and/or] Keta lake
- Stenurus* Dujardin 1845
 Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
 J. Zool., v. 53 (6), 713-735
 key; key to species
- Stenurus alatus* (Leuckart) Yorke and Maplestone
 1926
 Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
 J. Zool., v. 53 (6), 713-735
 as syn. of *Pharurus alatus* (Leuckart 1848)
 Stiles and Hassall 1905
- Stenurus arcticus* (Cobb) Baylis and Daubney,
 1925
 Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
 J. Zool., v. 53 (6), 713-735
 as syn. of *Pharurus pallasii* (van Beneden
 1870) n. comb.
- Stenurus arctomarinus* Delyamure and Kleinenberg
 1958, *illus.*
 Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
 J. Zool., v. 53 (6), 713-735
 redescription; key
Delphinapteras leucas: MacKenzie River Del-
 ta, Canada; Churchill, Manitoba, Canada
- Stenurus globicephalae* Baylis and Daubney 1925,
illus.
 Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
 J. Zool., v. 53 (6), 713-735
 redescription; key
Globicephala melaena: off Newfoundland,
 Canada
Grampus griseus: off Newfoundland, Canada
Lagenorhynchus acutus: off Cape Cod, Massa-
 chusetts, U.S.A.
Globicephala macrorhyncha: off Castries,
 St. Lucia, Lesser Antilles

- Stenurus globicephalae* (Baylis & Daubney),
illus.
Cannon, L. R. G., 1977, Austral. J. Marine and
Freshwater Research, v. 28 (6), 717-722
Peponocephala electra (guttural pouches, ear
canals): Tugun Beach; Moreton Island
- Stenurus inflexus* Dujardin 1845
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis
and Daubney 1925
- Stenurus minor* (Kuhn 1829) Baylis and Daubney
1925, illus.
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
synonymy; redescription; key
Phocoena phocoena: Bay of Fundy, Canada;
off Newfoundland, Canada; North Sea off
Netherlands
- Stenurus ovatus*
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
key
- Stenurus pallasii* (van Beneden) Dougherty 1943
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden
1870) n. comb.
- Stenurus phocoenae* Dougherty 1943
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis
and Daubney 1925
- Stenurus vagans* (Eschricht 1841) Dougherty 1943
Arnold, P. W.; and Gaskin, D. E., 1975, Canad.
J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis
and Daubney 1925
- Stephanofilaria* [sp.]
Hiregoudar, L. S., 1974, Indian Vet. J., v. 51
(1), 72
Boselaphus tragocamelus (ears): Gir Forest,
Gujarat State
- Stephanofilaria* sp.
Hiregoudar, L. S., 1976, Indian Vet. J.,
v. 53 (3), 237
Boselaphus tragocamelus (ear): Gir forest,
Gujarat State, India
- Stephanofilaria* sp. Kono, 1965
Ueno, H.; and Chibana, T., 1977, National Inst.
Animal Health Quart., v. 17 (1), 16-26
as syn. of *Stephanofilaria okinawaensis* n.
sp.
- Stephanofilaria assamensis*
Ahmed, S.; and Ali, M. I., 1973, Vet. Med.
Rev. (2), 136-142
Stephanofilaria assamensis, cattle, Anti-
mosan-solution injected under sores followed
by local applications of gentian violet,
drug trials, good results: East Pakistan
- Stephanofilaria assamensis* Pande, 1936, illus.
Das, P. K.; Tripathy, S. B.; and Misra, S. K.,
1977, Indian J. Animal Sc., v. 45 (8), 1975,
543-545
Stephanofilaria assamensis, cattle, patho-
anatomy of skin
- Stephanofilaria assamensis*
Dutta, P. K.; and Hazarika, R. N., 1976,
Indian Vet. J., v. 53 (3), 221-224
Stephanofilaria assamensis, cattle, chemo-
therapeutic trials, comparative efficacy
of various drugs: Khanapara
- Stephanofilaria assamensis*, illus.
Patnaik, B., 1973, Ztschr. Tropenmed. u. Para-
sitol., v. 24 (4), 457-466
Stephanofilaria assamensis, life cycle com-
pleted by experimental reproduction of typi-
cal humpsores lesion on a calf using labora-
tory-raised *Musca conducens* as vectors,
parasite growth slow
Musca conducens (nat. and exper.): Orissa
Red-Sindhi calf (exper.)
- Stephanofilaria assamensis* (Pande, 1936)
Rahman, A.; and Khaleque, A., 1974, Vet. Med.
Rev. (4), 379-382
Stephanofilaria assamensis, cattle, Neguvon:
Bangladesh
- Stephanofilaria kaeli* Buckley, 1937, illus.
Fadzil, M., 1975, Kajian Vet., v. 7 (1), 1-7
Stephanofilaria kaeli microfilaria, develop-
ment in *Musca conducens* (exper.)
- Stephanofilaria kaeli*
Loke, Y. W.; and Ramachandran, C. P., 1966,
Med. J. Malaya, v. 20 (4), 348
histopathology of *Stephanofilaria kaeli*
lesions in cattle
- Stephanofilaria kaeli*
Mullin, S. W., 1971, Southeast Asian J. Trop.
Med. and Pub. Health, v. 2 (1), 84 [Demonstra-
tion]
Stephanofilaria kaeli, preliminary attempts
to establish life cycle show *Musca conducens*
as probable vector: Malaya
- Stephanofilaria kaeli*
Ramachandran, C. P.; Loke, Y. W.; and Nagen-
dram, C., 1966, Med. J. Malaya, v. 20 (4),
344-347
Stephanofilaria kaeli in cattle, intensity
of infection (size and condition of lesion
in relation to worm burden), location of
worms, morphology of adult worms, identifi-
cation of microfilariae: Malaya
- Stephanofilaria okinawaensis* n. sp., illus.
Ueno, H.; and Chibana, T., 1977, National Inst.
Animal Health Quart., v. 17 (1), 16-26
Syn.: *Stephanofilaria* sp. Kono, 1965
cows (teats, muzzle): Ishigaki Island, Nan-
sei (Southwestern) Islands, Japan
- Stephanofilaria okinawaensis*, illus.
Ueno, H.; Chibana, T.; and Yamashiro, E., 1977,
Vet. Parasitol., v. 3 (1), 41-48
Stephanofilaria okinawaensis, cattle, derma-
titis of teats, clinical and histopathologi-
cal observations, relationship to dermatitis
on muzzle: Nansei Islands, Okinawa Prefec-
ture, Japan

- Stephanofilaria stilesi* (Chitwood, 1934)
Azimov, D. A.; et al., 1976, Dokl. Akad. Nauk UzSSR (8), 53-54
Stephanofilaria stilesi, bovine, ecology, seasonal distribution, intermediate hosts and their daily activity in relation to temperature
Lyperosia irritans
L. titillans
Stomoxys calcitrans
all from southern Uzbekistan
- Stephanofilaria stilesi*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Lyperosia irritans
L. titillans
all from Uzbekistan
- Stephanofilaria zaheeri* Singh 1958
Agrawal, M. C.; and Dutt, S. C., 1976, Indian Vet. J., v. 53 (6), 475-476
Stephanofilaria zaheeri, buffaloes (ears), predominance of male adult worms: Jabalpur City abattoir
- Stephanofilaria zaheeri* Singh, 1958
Agrawal, M. C.; and Dutt, S. C., 1977, Indian Vet. J., v. 54 (1), 77-79
Stephanofilaria zaheeri, buffaloes, in vitro and in vivo drug trials, high efficacy with malathion: Jabalpur
- Stephanofilaria zaheeri* Singh, 1958
Agrawal, M. C.; and Dutt, S. C., 1977, Indian Vet. J., v. 54 (6), 497-498
Stephanofilaria zaheeri, buffaloes (ears), clinical and subclinical forms, prevalence, histopathology: Jabalpur
- Stephanofilaria zaheeri* Singh, 1958, illus.
Das, P. K.; Tripathy, S. B.; and Misra, S. K., 1977, Indian J. Animal Sc., v. 45 (12), 1975, 949-952
Stephanofilaria zaheeri, buffaloes (ear), histopathology, supona-20, fenitrothian, good results: Orissa
- Stephanofilariasis*
Muchlis, A.; and Soetijono, P., 1973, Vet. Med. Rev. (2), 134-135
stephanofilariasis, hoof myiasis, cattle, Asuntol ointment, good results: northern Sulawesi, Indonesia
- Stephanofilariasis*
Sen, S.; and Das, D. K., 1976, Indian J. Animal Health, v. 15 (1), 83
humpsores, cattle, Florocid treatment compared with phenothiazine-banocide ointment
- Stephanurus dentatus*
Coombs, D. W.; and Springer, M. D., 1974, J. Wildlife Dis., v. 10 (4), 436-441
pathological changes in infected wild pigs
Sus scrofa domesticus x *Sus scrofa cristatus* (liver, lungs, kidney pelvises, perirenal tissues): Aransas National Wildlife Refuge, southern Texas
- Stephanurus dentatus* Diesing, 1839
Dykova, I., 1977, Beitr. Trop. Landwirtschaft. u. Vet.-Med., v. 15 (4), 401-409
Stephanurus dentatus, pigs, relationship between incidence and zoohygienic conditions, pathology, economic importance, suggested control measures: Cuba
- Stephanurus dentatus*
Marti, O.G.; Fincher, G.T.; and Stewart, T.B., 1977, Vet. Parasitol., v. 3 (1), 89-93
Stephanurus dentatus, sows, cambendazole temporarily reduced hatchability but not production of eggs, not effective against adults, higher dosages should be evaluated
- Stephanurus dentatus*
Nowotny, F.; and Grestenberger, E., 1976, Wien. Tierarztl. Monatsschr., v. 63 (10), 311-312
imported pig (kidney): Wien
- Stephanurus dentatus*, illus.
Pandey, V. K.; Tripathy, S. B.; and Dey, P. C., 1977, Indian J. Animal Health, v. 16 (1), 91-92
Stephanurus dentatus, pigs (kidneys, bronchi), case history, pathology, post mortem examination, preventive measures and treatment with piperazine adipate and copper sulfate
- Stephanurus dentatus*, illus.
Sadana, J. R.; and Kalra, D. S., 1977, Indian J. Animal Sc., v. 44 (11), 1974, 865-868
Stephanurus dentatus, visceral larva migrans, pigs (lungs), pathology: abattoirs at Hissar and Delhi
- Stephanurus dentatus* (Diesing, 1839)
Sharma, R. K.; Singh, K.; and Sharma, P. K., 1976, Ztschr. Parasitenk., v. 49 (2), 187-192
Stephanurus dentatus, non-specific phosphomonoesterases, activity, distribution in various tissues
- Stephanurus dentatus*
Stewart, T. B.; et al., 1977, Am. J. Vet. Research, v. 38 (12), 2081-2083
Stephanurus dentatus, sows, levamisole HCl, subcutaneously and in corn meal feed, effect on hatchability and development of eggs and survival and development of ensuing larvae to 3rd stage
- Stephanurus dentatus*
Tromba, F. G.; and Romanowski, R. D., 1976, J. Parasitol., v. 62 (2), 250-255
Stephanurus dentatus, swine, evaluation as vaccines of 9 somatic antigens derived from excretory gland cells
- Sterliadochona* Skrjabin, 1946
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Cystidicoloides* Skinner, 1931

- Sterliadochona tenuissima* (Zeder, 1800) Spasskii and Roitman 1959
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Salmo trutta (stomach): Loch Leven, Scotland
- Stilestrongylus Freitas*, Lent et Almeida, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Nippostrongylinae
Syn.: *Mirandaia Travassos*, 1937
- Stilestrongylus manni* n. sp., illus.
Denke, M. A.; and Murua, R., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 127-131
Oryzomys longicaudatus (intestin grele): San Martin, Province de Valdivia, Chili
Akodon olivaceus: Province de Valdivia, Chili
A. longipilis: Province d'Orsono, Chili
- Stilestrongylus moreli* n. sp., illus.
Diaw, O. T., [1977], Bull. Mus. National Hist. Nat., Paris, 3. s. (405), 1976, Zool. (282), 1065-1089
Phyllotis boliviensis (intestin): Rio Lep-ture, Bolivie
- Stomachida Pereboom*, 1780
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Ascaris* L., 1758
- Stomachidae* (Johnston & Mawson, 1945, subfam.)
Hartwich, 1957
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakidae* (Railliet & Henry, 1912, subfam.) Skrjabin & Karokhin, 1945
- Stomachinae* Johnston & Mawson, 1945
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakinae* Railliet & Henry, 1912
- Stomachus* Goeze in Zeder, 1800, nom. nud.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Anisakis* Dujardin, 1845
- Streptocara* Railliet, Henry & Sisoff, 1912, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Seuratiinae
key
Syn.: *Yseria* Gedoelst, 1919
"Koriakinema Oshmarin, 1949 is too poorly known to be classified but it may be a synonym of *Streptocara*."
- Streptocara* sp.
Lancaster, M. B., 1973, Vet. Rec., v. 92 (10), 261-262 [Letter]
muscovy duckling (crop): England
- Streptocara crassicauda* (Creplin, 1829), illus.
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (3), 185-189
description
Rissa tridactyla (ventriculus): Agdenes area, Norway
- Streptocara crassicauda* (Creplin, 1892)
Bishop, C. A.; and Threlfall, W., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 25-35
Somateria mollissima (proventriculus, gizzard): insular Newfoundland and/or southern Labrador
- Streptocara crassicauda* (Creplin, 1829)
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Limosa limosa lapponica: lower Yenisei
- Streptocara crassicauda* (Creplin, 1829) Skrjabin, 1915
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anas platyrhynchos (under cuticle of muscular stomach): Bulgaria
- Streptocara crassicauda* (Creplin, 1829)
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus genei: Azov Sea
- Streptocara crassicauda* (Creplin, 1829)
Turner, B. C.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 157-169
parasites of *Anas crecca* and *A. discors*, incidence and intensity, age and sex of host
Anas discors (beneath the gizzard lining): eastern Canada
- Streptocara incognita* Gibson, 1968, illus.
Fox, J. G.; et al., 1974, J. Wildlife Dis., v. 10 (1), 66-69
Streptocara incognita in *Phoenicopterus chilensis* causing severe debilitating condition and eventual death, gross and histopathological findings: San Francisco Zoological Gardens
- Streptocara recta* (Linstow, 1879)
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skrjabin), 105-124
Stercorarius parasiticus (muscular stomach): coast of Sea of Okhotsk (Tuguro-Chumikansk region)
- Streptopharagus* Blanc, 1912, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascaropsinae
key

- Streptopharagus armatus*
McConnell, E. E.; et al., 1974, Onderstepoort J. Vet. Research, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (stomach, small intestine):
Kruger National Park, Transvaal
- Streptopharagus armatus*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Streptopharagus kutassi* (Schulz, 1927)
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Blaps fausti bactriana
B. deplanata reichardti
Trigonoscelis gemmulata
Pisterotarsa kessleri
P. gigantea
Cyphogenia gibba
Adesmia gebleri
Prosodes bactriana
Thriptera sp.
all from Tadzhik SSR
- Streptopharagus kutassi*
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Cyphostete komarovi
Adesmia servillei schatzmayri
Trigonoscelis gigas
T. punctipleuris
Pisterotarsa gigantea subsp. zoubkoffi
P. kessleri
Blaps fausti
all from Turkmenia
- Streptopharagus kutassi*, illus.
Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
- Streptopharagus lerouxi*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Strongyles
Bussieras, J., 1977, Rev. Med. Vet., Toulouse, v. 128 (3), 343-348
strongyles, ascarids, horses, pyrantel pamoate orally or by naso-oesophageal catheter, no bad effects
- Strongylacantha* Van Beneden, 1873
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Strongylacanthidae
- Strongylacanthidae (Yorke et Maplestone, 1926, sub. fam.), Chabaud, 1960
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongyloidea
includes: *Strongylacantha*
- [Strongylata] strongiliati
Panchin, O. G.; et al., 1975, Veterinariia, Kiev (40), 100-104
helminths and coccidia, sheep, seasonal incidence on pastures, measures for control: Kalanchats'k region, Kherson oblast
- Strongyles
Ardans, A.; and Walters, G., 1975, Am. J. Vet. Res., v. 36 (11), 1589-1590
Parascaris equorum, strongyles, Quarter horses, treatment with 3 formulations of cambendazole (paste, pellets, suspension) vs. thiabendazole (suspension)
- Strongyles
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
ruminants, survey, treatment, economic importance: Kaedi area (Mauritania)
- Strongyles
El-Moukdad, A.-R., 1977, Wien Tierarztl. Monatschr., v. 64 (3), 85-91
effect of disinfectants on eggs of *Ascaris suum*, *Toxascaris leonina*, small horse strongyles and coccidia oocysts
- Strongyles
Enigk, K.; et al., 1975, Zentralbl. Vet.-Med., Reihe B, v. 22 (8), 687-702
survival of resistant external stages of parasites during fermentation of liquid cattle manure at high temperatures
- Strongyles
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1976, Tierarztl. Umschau, v. 31 (8), 360-362
swine nematodes, mebendazole treatment
- Strongyles
Fagot, G.; Clery, P.; and Pascal, D., 1977, Prat. Vet. Equine, v. 9 (3), 169-170
ascarids, strongyles, *Oxyuris* in race horses, clinical trials with strongid-P most successful in eradicating strongyles: France
- Strongyles, small
Folz, S. D., 1977, Vet. Parasitol., v. 3 (4), 377-381
kinetics of natural expulsion of some equine parasites from nontreated horses during interim of critical test, concluded that natural elimination would not have biased efficacy of an anthelmintic or boticide
- Strongyles
Greve, J. H.; and Paul, J. W., 1976, Vet. Med. and Small Animal Clin., v. 71 (12), 1737-1740, 1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Strongyles
Krishna Iyer, P. P.; and Peter C. T., 1975, Kerala J. Vet. Sc., v. 5 (2), 121-123
gastrointestinal nematodes, goats, methyridine

- Strongyles**
Lange, E.; and Tomsen, K., 1976, Tierarztl. Umschau, v. 31 (10), 441-443
Strongyloides, strongyles, ascarides, horses, fenbendazole (aqueous suspension, granulate), field trials
- Strongyles**
McLeod, C. C., 1976, N. Zealand J. Exper. Agric., v. 4 (2), 215-218
thiabendazole, tetramisole, pre- and post-weaning anthelmintic drenching, live-weight gain, wool weight in Romney ewe lambs with strongyle infections better in treated than in untreated lambs, no significant differences between anthelmintics
- Strongyles**
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Strongyles**
Slocombe, J. O. D.; and Cote, J. F., 1977, Canad. Vet. J., v. 18 (8), 212-217
small strongyles, horses, cross resistance to cambendazole, thiabendazole, and mebendazole, susceptibility to non-benzimidazole compounds
- Strongyles**
Smith, H. J., 1976, Canad. J. Comp. Med., v. 40(4), 327-333
strongyles, Oxyuris equi, Parascaris equorum, naturally-infected ponies, thiabendazole, results support regular repeated treatment
- Strongyles**
Smith, H. J., 1976, Canad. J. Comp. Med., v. 40 (4), 334-340
strongyles, reinfection of thiabendazole treated ponies, development of patent infections, cyclic pattern of worm egg outputs
- Strongyles**
Tassi, P.; and Widenhorn, O., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 506-507
strongyles, sheep, seasonal variation in incidence: central-south and insular Italy
- Strongylida**
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Secernentea
key; key to superfamilies
includes: Diaphanocephaloidea; Ancylostomatoidea; Strongyloidea; Trichostrongyloidea; Metastrongyloidea
- Strongylida** Diesing, 1851
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Secernentea
- Strongylidae**
Cabaret, J., 1977, Rec. Med. Vet., v. 153 (6), 419-427
gastrointestinal strongyles (Trichostrongylidae, Strongylidae), inhibition of larval development (hypobiosis), weather conditions, epidemiological aspects, review
- Strongylidae**
Furmaga, S.; Gundlach, J. L.; and Patyra, J., 1976, Med. Wet., v. 32 (12), 734-737
roundworms, horses, fenbendazol and cambendazol very effective
- Strongylidae**
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Strongylidae**
Setasuban, P., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 390-394
Bathostomum sangeri in Elephas indicus, light and scanning electron microscopy, morphometric data, confirmation that parasite is probably a Strongylidae rather than Ancylostomidae: circus elephants in Brisbane, Australia (primary origin unknown)
- Strongyliden**
Loeliger, H. C., 1974, Prakt. Tierarzt, v. 55, Sondernummer, 6-9
rabbits, clinical signs, diagnosis, control, review
- Strongyliden**
Manz, D.; and Dingeldein, W., 1974, Prakt. Tierarzt, v. 55 (8), 422-425
nematodes of European and exotic herbivores, Banminth, good results
Equus quagga
- Strongyliden-type, illus.**
Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Strongylides, petits et gros**
Düwel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Strongylides, petits et grands**
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongylids**
Danielli, Y.; and Neuman, M., 1975, Refuah Vet., v. 32 (4), 94-95, 153-154
mixed parasites, cattle, good results following repeated chemotherapy: Birkat Ata
- Strongylids**
Lindberg, R., 1976, Svensk Vet.-Tidn., v. 28 (11), 509-514
horse strongylid eggs, longevity on grass plots when deposited early or late in grazing season or in winter, seasonal and climatic aspects, preliminary field study

- Strongylids**
 McCall, J. P.; and McCullough, C., 1977, Southwest. Vet., v. 30 (2), 159-160
 strongylids, *Parascaris equorum*, horses (feces), thiabendazole, piperazine phosphate, good results; failure to demonstrate acquired drug resistance
- Strongyloidea**
 Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
 Strongylida
 key
- Strongyloidea [sp.]**
 Coggins, J. R., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
 parasitic fauna, effect of host diet and habitat
Turdus migratorius: Kellogg Bird Sanctuary, Michigan
- Strongyloides**
 Basset, M.; et al., 1975, Medecine Trop., v. 35 (6), 475-476
 mixed *Strongyloides* infections apparent after treatment of *Necator americanus* and *Ancylostoma duodenale* with alcopar or didakene, case reports, statistics
- Strongyloides**
 Biagi, F.; Smyth, J.; and Gonzalez, C., 1974, Prensa Med. Mexicana, v. 39 (1-2), 51-53
 human intestinal helminths, successful clinical trials using mebendazole, drug well tolerated with minimal side effects: Mexico
- Strong[yloides]**
 Brunson, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
 lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Strongyloides**
 Chhabra, R. C.; Bali, H. S.; and Toor, L. S., 1976, J. Research, Punjab Agric. Univ., v. 13 (3), 308-311
 gastrointestinal strongyles in sheep, critical drug evaluation, thiabendazole (most effective), tetramisole (good results), morantel tartrate (fair results), clioxanide and methyridine (least effective): India
- Strongyloides**
 Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
 fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Strongyloides**
 Forstner, M. J.; Kopp, H.; and Wiesner, H., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (9), 180-183
 nematodes of ruminants, mebendazole, good results: Hellabrunn Zoo, Munich
- Strongyloides**
 Gheorghescu, P.; et al., 1976, Med. Interne, Bucarest, v. 14 (1), 31-38
 Giardia infections alone or in combination with *Strongyloides* or *Taenia solium*, absorption studies before and after treatment
- Strongyloides**
 Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
 nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Strongyloides**
 Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
 gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Strongyloides**
 Klein Mori, J., 1972, Rev. Med. Vet. y Parasitol., Maracay, v. 24 (1-8), 1971-1972, 207-226
 gastrointestinal nematodes, sheep, Neguvon, Ripercol, Thibenzoline, comparison, various management systems, all effective, Ripercol easiest to administer, Neguvon somewhat toxic
- Strongyloides**
 Krishna Iyer, P. P.; and Peter C. T., 1975, Kerala J. Vet. Sc., v. 5 (2), 121-123
 gastrointestinal nematodes, goats, methyridine
- Strongyloides**
 Lange, E.; and Tomsen, K., 1976, Tierarztl. Umschau, v. 31 (10), 441-443
 Strongyloides, strongyles, ascarides, horses, fenbendazole (aqueous suspension, granulate), field trials
- Strongyloides**
 Lodha, K. R.; Raisinghani, P. M.; and Karwar, R. S., 1977, Indian J. Animal Sc., v. 47 (10), 677-682
 helminths, camels, promintic and banminth. II effective, nilverm inconsistent in action, thiabendazole ineffective
- Strongyloides**
 McFarlane, H., 1976, Proc. Nutrition Soc., v. 35 (3), 263-272
 parasitic infestation, preschool children, malnutrition and impaired immune response, brief review comment: Nigeria
- Strongyloides**
 Manz, D.; and Dingeldein, W., 1974, Prakt. Tierarzt, v. 55 (8), 422-425
 nematodes of European and exotic herbivores, Banminth, good results
 Ovis musimon
Antidorcas marsupialis
Antilope cervicapra
- Strongyloides**
 Most, H., 1972, N. England J. Med., v. 287 (10), 495-498; (14), 698-702
 common parasitic infections of man encountered in the United States, recommendations for treatment, review

Strongyloides

Neppert, J.; and Warns, C.-M., 1974, Tropenmed. u. Parasitol., v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from Ascaris, hookworm, Onchocerca, Dirofilaria immitis, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination

Strongyloides

Raynaud, J.-P.; Sennelier, J.; and Irisarri, E., 1975, Folia Vet. Latina, v. 5 (3), 412-429
gastrointestinal helminths, swine, post natal infection of piglets in contact with infected mothers, comparison of various methods of husbandry and hygiene, studies during pregnancy and lactation, routine daily hygiene recommended

Strongyloides

Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, not effective

Strongyloides

Sewell, M. M. H., 1973, Vet. Rec., v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review

Strongyloides

Smeal, M. G.; et al., 1977, Austral. Vet. J., v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales

Strongyloides

Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction

Strongyloides

Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report

Strongyloides

Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date

Strongyloides

Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidial drug (Cozurone), good control of all except Strongyloides

Strongyloides

Zielinski, J., 1972, Med. Wet., v. 28 (9), 566-567
parasites, sheep, Nilverm, copper sulfate

Strongyloides sp., illus.

Brown, R. C.; and Girardeau, M. H. F., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 215-219
Strongyloides fuelleborni, Necator americanus, Ancylostoma duodenale, prevalence survey and study of possible transmammmary passage, presence of Strongyloides sp. larvae in milk of one nursing mother suggests that S. fuelleborni may be transmitted via milk in humans: Bulape, Zaire

Strongyloides sp.

Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (small intestine): Florida

Strongyloides sp.

Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand

Strongyloides sp.

Croll, N. A.; and Killick-Kendrick, R., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 688-689 [Letter]
Strongyloides sp. in human sputum, possible epidemiologic importance of spread by promiscuous expectoration: Bristol, United Kingdom

Strongyloides spp.

Dafalla, A. A.; Satti, M. H.; and Abdel Nur, O. M., 1977, J. Trop. Med. and Hyg., v. 80 (3), 63-64
cutaneous larva migrans, preliminary epidemiologic survey shows high prevalence in children especially during rainy seasons, etiology unknown but high incidence of Strongyloides spp. in soil samples, suggested control measures: Northern Kordofan, Sudan

Strongyloides spp.

El-Abdin, Y. Z.; et al., 1975, Egypt. J. Vet. Sc., v. 12 (1), 31-43
serum constituents and serum enzyme activities, normal and nematode infested Camelus dromedarius: Cairo abattoir

Strongyloides sp.

Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (duodenum, lower small intestine, ceca): Florida

- Strongyloides* sp.
Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, *J. Parasitol.*, v. 61 (3), 547-548
Grus canadensis pratensis (duodenum, lower small intestine, large intestine, ceca, cloaca): Florida
- Strongyloides* sp.
Grundmann, A. W.; and Lombardi, P. S., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 39-46
Ochotona princeps cinemomea: Tushar Mountains, Utah
- Strongyloides* sp.
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 119-127
Meleagris gallopavo (duodenum; lower small intestine; ceca): Florida
- Strongyloides* sp.
Kagei, N.; et al., 1976, *Bull. Inst. Pub. Health, Tokyo*, v. 25 (3), 140-144
Filaroides hirthei, *Strongyloides* sp., *Toxocara canis*, and *Trichuris vulpis* in imported dogs, inability to experimentally infect other dogs with *Strongyloides stercoralis* of human origin: Japan, imported from U.S.A.
- Strongyloides* sp., *illus.*
Kelly, A.; Little, M. D.; and Voge, M., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mintagezole, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides* new species
Kelly, A.; and Voge, M., 1973, *Papua N. Guinea Med. J.*, v. 16 (1), 59
Strongyloides n. sp. reported from humans (feces), preliminary morphologic characteristics of ova: Middle Fly River area, Kiunga, West District, Papua New Guinea
- Strongyloides* sp.
Kinsella, J. M., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 127-130
Aphelocoma c. coerulescens (small intestine): Florida
- Strongyloides* sp.
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, *Am. Midland Naturalist*, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans
Porphyryla martinica
(small intestine of all): all from Florida
- Strongyloides* spp.
McConnell, E. E.; et al., 1974, *Onderstepoort J. Vet. Research*, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (small intestine): Kruger National Park, Transvaal
- Strongyloides* spp., *illus.*
Page, F. T.; and Reeves, D. S., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (2), 256-259
Strongyloides spp., accelerated auto-infection in patient with terminal carcinomatosis, low-grade infection thought to have existed for 26 years before erupting, unusual symptoms with numerous larvae in sputum and feces and no evidence of adult worms even at autopsy, danger of infection spread through larvae-infected sputum: Bristol, United Kingdom
- Strongyloides* spp.
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, *Proc. 3. National Wild Turkey Symp.*, 27-32
Meleagris gallopavo silvestris: southeastern United States
- Strongyloides* sp.
Pursglove, S. R.; et al., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (9), 896-900
Strongyloides sp. in *Odocoileus virginianus*, pathology, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Strongyloides* [sp.]
Reed, D. E.; et al., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (9), 975-979
Odocoileus hemionus (rectal contents): South Dakota
- Strongyloides*-type, *illus.*
Schuetze, H. R., 1974, *Prakt. Tierarzt*, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Strongyloides* spp.
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongyloides* sp.
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, *Onderstepoort J. Vet. Research*, v. 42 (1), 29-31
Damaliscus dorcas dorcas: captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria
- Strongyloides avium* Cram, 1929
Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (1), 7-31
Syn.: *S. oswaldoi* Travassos, 1930
- Strongyloides avium* (Cram, 1929)
Fabiyl, J. P., 1972, *Bull. Epizoot. Dis. Africa*, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria

- Strongyloides avium* Cram, 1929
Fabiyl, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (caeca): Vom area, Benue Plateau State, Nigeria
- Strongyloides cebus*
Kelly, A.; Little, M. D.; and Voge, M., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mintezol, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides cebus, illus.*
King, N. W., jr., 1976, Scient. Publication (317). Pan Am. Health Organ., 169-198
- Strongyloides fuelleborni*
Brown, R. C.; and Girardeau, M. H. F., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 215-219
Strongyloides fuelleborni, *Necator americanus*, *Ancylostoma duodenale*, prevalence survey and study of possible transmammmary passage, presence of *Strongyloides* sp. larvae in milk of one nursing mother suggests that *S. fuelleborni* may be transmitted via milk in humans: Bulape, Zaire
- Strongyloides fuelleborni*
File, S. K.; McGrew, W. C.; and Tutin, C. E. G., 1976, J. Parasitol., v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces): Gombe National Park, Tanzania
- Strongyloides fuelleborni von Linstow, 1905, illus.*
Hira, P. R.; and Patel, B. G., 1977, Am. J. Trop. Med. and Hyg., v. 26 (4), 640-643
Strongyloides fuelleborni rhabditiform larvae and eggs cultured to free living adults obtained from human feces, discussion of differential diagnosis from *S. stercoralis*, prominent morphologic features, mode of human infections, survey of prevalence in predominantly urban and suburban areas of Zambia
- Strongyloides fuelleborni, illus.*
Kelly, A.; Little, M. D.; and Voge, M., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mintezol, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides fuelleborni, illus.*
Pampiglione, S.; and Ricciardi, M. L., 1972, Parasitologia, v. 14 (2-3), 329-338
Strongyloides fuelleborni, survey, geographic distribution, incidence in human feces, endemic in tropical forest regions, sporadic in savannah regions, slightly higher prevalence in children: West, Central and East Africa
- Strongyloides fuelleborni*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Strongyloides martis* Petrow, 1940
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes
Mustela lutreola
Mustela erminea
Lutra lutra
Mustela nivalis
Mustela putorius
(intestine of all): all from Karelia
- Strongyloides oswaldoi* Travassos, 1930
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
as syn. of *S. avium* Cram, 1929
- Strongyloides papillosus*
Bezubik, B., 1969, Acta Parasitol. Polon., v. 17 (1-19), 1-9
Strongyloides papillosus, rabbits, human gamma globulin beneficial as expressed by host weight, parasite egg production, and blood values; infection more severe in younger hosts
- Strongyloides papillosus*
Bezubik, B., 1969, Acta Parasitol. Polon., v. 17 (1-19), 11-16
Strongyloides papillosus, pregnant rabbits infected percutaneously, no infection in newborn offspring
- Strongyloides papillosus*
Bezubik, B.; and Borowik, M. M., 1969, Acta Parasitol. Polon., v. 17 (1-19), 17-24
Strongyloides papillosus, rats, natural resistance not broken down by cortisone
- Strongyloides papillosus* (Wedl, 1856)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (small intestine): vicinity of Nowy Targ, Carpathian Mountains
- S[trongyloides] *papillosus*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Strongyloides papillosus*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Strongyloides papillosus*
Bryan, R. P., 1976, Austral. Vet. J., v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide

- Strongyloides papillosus*
Bryan, R. P.; Bainbridge, M. J.; and Kerr, J. D., 1976, Austral. J. Zool., v. 24 (3), 417-421
Bubalus bubalis
cattle
(large and small intestine of all): all from Northern Territory, Australia
- Strongyloides papillosus*
Chomicz, L., 1969, Acta Parasitol. Polon., v. 17 (1-19), 47-54
Strongyloides papillosus, immature and mature rabbits, blood picture, age and sex resistance
- Strongyloides papillosus*
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs and heifers, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Strongyloides papillosus*
Colglazier, M. L.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with Trichuris spp. and Moniezia expansa
- Strongyloides papillosus*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Strongyloides papillosus*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 20 mg/kg moderate to high
- Strongyloides papillosus*
Forrester, D. J.; Taylor, W. J.; and Nair, K. P. C., 1974, J. Wildlife Dis., v. 10 (1), 11-17
strongyloidosis in captive Odocoileus virginianus fawns, high mortality, transmission strongly presumed to be intrauterine, effective control by removal from does after 2 days of age and thiabendazole treatment; possible implications for wild deer populations: Florida
- Strongyloides papillosus*
Frackowiak, B.; et al., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 21-25
Strongyloides papillosus, sheep, repeated infections, weekly or semiweekly intervals, serum transaminase activity during course of infection, possible relation to pathological damage by parasite
- Strongyloides papillosus*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Strongyloides papillosus*
Kelly, A.; Little, M. D.; and Voge, M., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 694-699
Strongyloides sp. (S. fuelleborni-like) infections in natives discovered during intestinal parasitic survey, mitezol, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides papillosus*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Strongyloides papillosus*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Strongyloides papillosus*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med., Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Strongyloides papillosus*
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Strongyloides papillosus*
Mantovani, A.; et al., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 569-570
role of *Strongyloides papillosus* larvae in penetration and spread of Sphaerophorus necrophorus in rabbits
- Strongyloides papillosus*
Mantovani, A.; et al., 1972, Parassitologia, v. 14 (1), 149-162
Strongyloides papillosus larvae transmit the bacteria Sphaerophorus necrophorus to rabbits and spread it to host lungs by migration
- Strongyloides papillosus*
Moncol, D. J.; and Grice, M. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 1-4
Strongyloides papillosus, goats, sheep (nat. or exper.), transmammmary passage

- Strongyloides papillosus*
Niec, R.; et al., 1976, *Gac. Vet.*, Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary ant-helmintic treatment that could prevent normal buildup of resistance
- Strongyloides papillosus*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Strongyloides papillosus* (Wedl, 1856)
Stankiewicz, M., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 39-46
Strongyloides papillosus larvae, sheep and rabbit strains, responses to light of various intensities, desiccation and temperature; pattern of migration from water by dense group of larvae; reaction to various chemicals; destruction by fungi; no differences between strains
- Strongyloides papillosus*
Stankiewicz, M., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 147-159
Strongyloides papillosus, sheep, repeated infections with sheep and rabbit strains, egg production, hematocrit, body weight; lower pathogenicity of rabbit-adapted strains, high immunity produced by both strains
- Strongyloides papillosus*
Stankiewicz, M., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 161-173
Strongyloides papillosus, sheep, single and multiple infections with sheep and rabbit strains, changes in leukocyte composition of peripheral blood
- Strongyloides papillosus*
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Strongyloides papillosus*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongyloides papillosus*, *illus.*
Triantaphyllou, A. C.; and Moncol., D. J., 1977, *J. Parasitol.*, v. 63 (6), 961-973
Strongyloides ransomi, *S. papillosus*, chromosomal complement, gametogenesis, mode of reproduction, sex determination, hybridization tests
- Strongyloides papillosus*
Troncy, P. M.; and Oumate, O., 1973, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 26 (2), 199-202
Neoscaris vitulorum, *Strongyloides papillosus*, Zebu calves, morantel tartrate: Tchad
- Strongyloides papillosus*
Troncy, P. M.; and Oumate, O., 1976, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 29 (3), 229-232
gastrointestinal parasites, *Camelus dromedarius*, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Strongyloides papillosus*
Volf, K.; and Volfova, M., 1974, *Veterinarstvi*, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Strongyloides papillosus*
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, *Vet. Glasnik*, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Strongyloides papillosus*
Williams, J. C.; and Knox, J. W., 1976, *Am. J. Vet. Research*, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Strongyloides papillosus*
Zaffagnini, F., 1972, *Parassitologia*, v. 14 (1), 209-214
Strongyloides papillosus females, calf strain, diploid parthenogenesis and diploid chromosome number $2n=4$; maturation of oocytes: ameiotic parthenogenesis in parasitic females and meiotic parthenogenesis in free-living females
- Strongyloides papillosus*
Zajicek, D.; and Kozdon, O., 1977, *Veterinarstvi*, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Strongyloides ransomi*
Chauhan, H. V. S.; Dwivedi, P.; and Kalra, D. S., 1974, *Haryana Vet.*, v. 13 (1), 5-21
protozoan and helminth parasites, transmitted through milk to newborn animals, review
- Strongyloides ransomi*
Dey-Hazra, A., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 198
helminths, pigs, mode of pathogenicity, review

- Strongyloides ransomi*
Dey-Hazra, A.; Enigk, K.; and Kolm, H. P., 1977, *Research Vet. Sc.*, v. 22 (3), 353-356
Strongyloides ransomi, piglets (exper.), moderate and heavy infections, intestinal absorption rates of palmitate and 2-aminoisobutyric acid, comparison with uninfected piglets
- Strongyloides ransomi*
Enigk, K.; and Dey-Hazra, A., 1975, *Vet. Parasitol.*, v. 1 (1), 69-75
Strongyloides ransomi-infected piglets, measurement of plasma and red cell loss into intestinal tract, leakage of plasma protein into gut is greater than can be accounted for in terms of whole blood loss
- Strongyloides ransomi*
Enigk, K.; and Dey-Hazra, A., 1976, *Prakt. Tierarzt*, v. 57 (4), 232-234
Strongyloides ransomi, prepatent *Ascaris suum*, experimental infection in swine, Camendazol in feed, effective treatment
- Strongyloides ransomi*
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1976, *Tierarztl. Umschau*, v. 31 (8), 360-362
swine nematodes, mebendazole treatment
- Strongyloides ransomi*
Enigk, K.; Dey-Hazra, A.; and Eduardo, S. L., 1976, *J. Comp. Path.*, v. 86 (2), 243-250
Strongyloides ransomi, pathogenesis in pigs, activity of disaccharidases and dipeptidases of intestinal mucosa during mild and severe infections
- Strongyloides ransomi*
Getler, K., 1972, *Med. Wet.*, v. 28 (8), 476-477
nematodes, pigs on industrial swill feeding farm, Atgard
- Strongyloides ransomi*
Hale, O. M.; Stewart, T. B.; and Johnson, J. C., jr., 1977, *Research Bull. (203) Agric. Exper. Stations Univ. Georgia*, 3-21
Strongyloides ransomi, *Ascaris suum*, naturally infected gilt and/or barrow crossbred pigs, superimposed *S. ransomi* infection, differences in performance with diets of varying levels of protein and vitamins not significant
- Strongyloides ransomi*
Johnson, J. C., jr.; Stewart, T. B.; and Hale, O. M., 1975, *J. Parasitol.*, v. 61 (3), 517-524
responses of pigs to natural infections of *Strongyloides ransomi* and *Ascaris suum* and to superimposed artificial infection with *Strongyloides ransomi*: effects of breed (Duroc, Hampshire, Duroc-Hampshire crossbred), level of *Strongyloides ransomi* infection, and season (spring, fall) on performance of growing-finishing pigs
- Strongyloides ransomi*
Kelly, A.; Little, M. D.; and Voge, M., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mntezol, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides ransomi*
Leland, S. E., jr., 1972, *J. Am. Vet. Med. Ass.*, v. 160 (1), 58-60
Strongyloides ransomi, pigs (exper.), hygromycin B, no protection against death or debilitating effects, no activity against migrating larvae or egg-laying process, adult worm burden reduced
- Strongyloides ransomi* Schwartz and Alicata, 1930, *illus.*
Moncol, D. J., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 86-92
Strongyloides ransomi, biology and morphology of transcolostr phase in pigs
- Strongyloides ransomi*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Strongyloides ransomi*, *illus.*
Romero Rodriguez, J., 1975, *Rev. Iber. Parasitol.*, v. 35 (1-2), 163-165
Sus scrofa domestica (base of tongue): Jabugo, Huelva (Spain)
- Strongyloides ransomi*
Stewart, T. B., 1976, *J. Parasitol.*, v. 62 (4), 650-651
Strongyloides ransomi in pigs, comparisons of exper. infections resulting from percutaneous vs. subcutaneous exposure, subcutaneously infected pigs had higher and earlier mean eggs per gram of feces and higher mean number of worms in close association with small intestinal mucosa and less % increase in body weight
- Strongyloides ransomi*
Stewart, T. B.; Stone, W. M.; and Marti, O. G., 1976, *Am. J. Vet. Research*, v. 37 (5), 541-544
Strongyloides ransomi, frequency of prenatal and transmammary infection, pigs of sequential litters from dams experimentally exposed as weanlings
- Strongyloides ransomi*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongyloides ransomi*, *illus.*
Triantaphyllou, A. C.; and Moncol, D. J., 1977, *J. Parasitol.*, v. 63 (6), 961-973
Strongyloides ransomi, *S. papillosus*, chromosomal complement, gametogenesis, mode of reproduction, sex determination, hybridization tests

- Strongyloides ratti*
Bailenger, J.; et al., [1977], Ann. Parasitol., v. 51 (6), 1976, 653-665
Strongyloides ratti, rats, inhibition of self cure by treatment with glucocorticosteroids or ACTH
- Strongyloides ratti*
Bailenger, J.; and Cabannes, A., 1976, Ann. Parasitol., v. 51 (5), 563-576
Strongyloides ratti, lactating rats, inhibition of self-cure, decrease in intensity of parasitemia, plasma corticosteroid levels
- Strongyloides ratti*
Bailenger, J.; and Faraggi, G., 1975, Ann. Parasitol., v. 50 (2), 187-197
Strongyloides ratti, rats (exper.), mechanism of hypocorticosteronemia
- Strongyloides ratti*
Bailenger, J.; and Faraggi, G., 1975, Ann. Parasitol., v. 50 (2), 199-208
Strongyloides ratti, rats (exper.), reserpine treatment inhibits self-cure reaction and causes hypercorticosteronemia, implications for role of hypocorticosteronemia normally associated with this parasite
- Strongyloides ratti*
Gavaghan, A. D.; and Nunn, A. J., 1972, Pharm. Acta Helvetiae, v. 47 (11-12), 719-726
Nippostrongylus brasiliensis, *Strongyloides ratti*, *Nematospiroides dubius*, synthesis of thiocyanine and hemithiocyanine derivatives of benzothiazole, significant anthelmintic activity in laboratory trials with mice
- Strongyloides ratti*
Kazacos, K. R., 1976, J. Parasitol., v. 62 (3), 493-494
Trichinella spiralis-immunized rats, increased resistance to *Strongyloides ratti*; lack of cross-reacting precipitating antibodies in in vitro tests
- Strongyloides ratti*, illus.
Kazacos, K. R.; and Thorson, R. E., 1975, J. Parasitol., v. 61 (3), 525-529
rats, immunization with *Nippostrongylus brasiliensis* or *Strongyloides ratti* protected against homologous and heterologous challenge; precipitates formed on infective larvae incubated in vitro in homologous or heterologous immune globulins
- Strongyloides ratti*
Mishra, G. S.; and Gonzalez, J. P., 1975, Arch. Inst. Pasteur Tunis, v. 52 (1-2), 71-87
Rattus norvegicus (intestin grele): Tunisia
- Strongyloides ratti*
Moqbel, R.; and Denham, D. A., 1977, J. Helminthol., v. 51 (4), 301-308
Strongyloides ratti, primary and secondary infections in small intestine of rats: course of infections; changes in size of worms; distribution and migration of adult worms in intestine
- Strongyloides ratti*
Roberts, T. M.; and Thorson, R. E., 1977, J. Parasitol., v. 63 (4), 764-766
Nippostrongylus brasiliensis, pairing between adults and other species of nematodes in vitro, comparison of intra and interspecific response patterns, results suggest that pairing is not related entirely to sexual attraction
- Strongyloides ratti* (Sandground, 1925)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus jalorensis: Malaysia
- Strongyloides ratti*, illus.
Taniguchi, M.; et al., 1977, Bull. Coll. Agric. and Vet. Med., Nihon Univ. (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo
- Strongyloides ratti*
Wilson, P. A. G., 1977, Parasitology, v. 75 (2), 233-239
Strongyloides ratti, rats (exper.), maternal worm burden when weaning is varied in relation to injection, effect of short-term stimulus (only 1 hr suckling) on maternal worm burden, working hypothesis to explain path-finding by migrating worms in lactating rats
- Strongyloides ratti*
Wilson, P. A. G.; Gentle, M.; and Scott, D. S., 1976, Parasitology, v. 73 (3), 399-406
Strongyloides ratti, dynamic determinants of route of larval migration in lactating rats; *Nippostrongylus brasiliensis*, *Strongyloides ratti*, control of exper. error in quantitative studies of milk transmission of skin-penetrating roundworms
- Strongyloides ratti*
Wilson, P. A. G.; Gentle, M.; and Scott, D. S., 1976, Parasitology, v. 72 (3), 355-360
rats, milk-borne infection, *Strongyloides ratti* (real and important vehicle for infection) vs. *Nippostrongylus brasiliensis* (possible but quantitatively insignificant)
- Strongyloides robustus*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Strongyloides sigmodontis*
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (small intestine): Florida

- Strongyloides stercoralis*
Adickman, M.; and Tuthill, T. M., 1976, Postgrad. Med., v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Strongyloides stercoralis*, *illus.*
Ali-Khan, Z.; and Seemayer, T. A., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (5-6), 473-476
Strongyloides stercoralis, fatal bowel infarction and septicemia in man with systemic strongyloidiasis, clinical case report, possible relationships between auto-infection and long-term steroid therapy: Montreal, Canada (immigrant from Hong Kong)
- Strongyloides stercoralis*, *illus.*
Bartholomew, C.; et al., 1977, Postgrad. Med. J., London (617), v. 53, 139-142
Strongyloides stercoralis, humans, case reports of sprue-like symptoms and pseudo-intestinal obstruction, pathologic findings, good results with thiabendazole therapy: Trinidad, West Indies
- Strongyloides stercoralis*
Best, J. C.; et al., 1976, Med. J. Australia, v. 1 (1-2), 14-20
assessment of growth-rate and growth-retardation in Australian Aboriginal children before and after treatment for common intestinal parasites
- Strongyloides stercoralis*
Bezjak, B., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 945-948
Strongyloides stercoralis, determination of immunoglobulins in duodenal contents and feces of infected humans, elevation of serum IgE levels
- Strongyloides stercoralis*
Biagi, F.; Lopez, R.; and Viso, J., 1975, Progr. Drug Research, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review
- Strongyloides stercoralis*
Biagi, F.; Smyth, J.; and Gonzalez, C., 1975, Prensa Med. Mexicana, v. 40 (5-6), 189-192
human intestinal parasites, clinical trials with mebendazole show it to be useful drug against many parasites and therefore recommended for mass therapy in low socioeconomic areas where multiple parasitism is likely to be present: Mexico
- Strongyloides stercoralis*
Bigorie, B.; et al., 1976, Nouv. Presse Med., v. 5 (13), 853-855
Strongyloides stercoralis in woman with resulting digestive tract complications and malabsorption syndrome, thiabendazole, clinical case report: France, immigrant from Guadeloupe
- Strongyloides stercoralis*
Botero R., D., 1975, Progr. Drug Research, v. 19, 28-43
human intestinal soil transmitted helminths, epidemiology and public health importance, pathology, extensive review
- Strongyloides stercoralis*, *illus.*
Buss, D. H., 1971, North Carolina Med. J., v. 32 (7), 269-274
Strongyloides stercoralis in woman, severe disseminated fatal infection associated with leukemia, and immunosuppressive therapy, clinical aspects: North Carolina
- Strongyloides stercoralis*
Bwangamoi, O., 1973, Bull. Epizoot. Dis. Africa, v. 21 (4), 363-370
Strongyloides stercoralis, *Dipylidium caninum*, *Taenia hydatigena*, recovery from dogs using Lindsey's method: Uganda
- Strongyloides stercoralis*
Cerva, L., 1976, Ceskoslov. Epidemiol., Mikrobiol., Immunol., v. 25 (6), 365-367
helminths, diagnostic staining method for eggs and larvae in fresh feces or those fixed by formalin
- Strongyloides stercoralis*, *illus.*
Cerva, L., 1976, Immun. u. Infekt., v. 4 (6), 279-282
intestinal helminths, diagnostic method for staining of eggs and larvae in smears of fresh and fixed stool samples
- Strongyloides stercoralis*
Cook, G. C., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (6), 419-436
role of *Giardia lamblia*, *Strongyloides stercoralis* and possibly other intestinal parasites in producing malabsorption in Africans
- Strongyloides stercoralis*
Copeman, R.; Pashen, D.; and Burger, G., 1975, Med. J. Australia, v. 1, suppl. 2, 8-13
association of common intestinal parasites to growth, nutrition and living situation of Aboriginal children: Cunnamulla, Western Queensland
- Strongyloides stercoralis*
Dancescu, P., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (2), 162-163
Strongyloides stercoralis in humans, observations on worm burden, clinical manifestations and infection duration as affected by host age: Romania
- Strongyloides stercoralis*
Doury, P.; et al., 1975, Nouv. Presse Med., v. 4 (11), 805 [Letter]
Strongyloides stercoralis in elderly woman resulting in symptoms of arthritis with multiple joint involvement, relief of symptoms after thiabendazole therapy: Paris

- Strongyloides stercoralis**
 Draghici, O., 1973, *Med. Int.*, Bucuresti, v. 25 (8), 945-953
 clinical manifestations of human *Strongyloides stercoralis*, eosinophilia changes before and after treatment with telmid or gentian violet, case reports
- Strongyloides stercoralis**
 Dwork, K. G.; Jaffe, J. R.; and Lieberman, H. D., 1975, *N. York State J. Med.*, v. 75 (8), 1230-1234
Strongyloides stercoralis, massive hyperinfection in humans with attenuated cell-mediated immunity, case report of mixed infection with toxoplasmosis, literature review of known cases, thiabendazole therapy helpful
- Strongyloides stercoralis**
 Fay, J. T.; and Disque, F. C., 1977, *Oral Surg.*, v. 43 (6), 898-901
Strongyloides stercoralis systemic infection in Vietnamese woman discovered as a result of routine biopsy and excision of impacted molar, increased eosinophilia led to medical work-up for possible parasitism, case history, successful treatment with di-thiazanine: Georgia
- Strongyloides stercoralis**
 de Figueiredo, N.; et al., 1968, *Bol. Chileno Parasitol.*, v. 23 (1-2), 57-59
 intestinal histopathology as ascertained by oral biopsies, common human intestinal parasites
- Strongyloides stercoralis**
 Forman, D. W.; et al., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 598-601
 parasitic pathogens discovered during etiologic survey of causes of diarrheal disease in U.S. Marines in Vietnam
- Strongyloides stercoralis** (Bavey, 1876), illus.
 Fossati, C., 1973, *Rev. Iber. Parasitol.*, v. 33 (1), 39-64
Strongyloides stercoralis, human, extensive literature review, two case histories with entero-pulmonary symptoms
- Strongyloides stercoralis**
 Galliard, H., 1974, *Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol.* (Vienna, Austria, Sept. 18-20, 1973), 295-304
 visceral and cutaneous larva migrans, review
- Strongyloides stercoralis**
 Garcia, F. T.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (5, part 1), 859-865
Strongyloides stercoralis, humans, results of study of intestinal function and morphology in strongyloidiasis show that associated malabsorption syndrome is secondary to concomitant malnutrition and not to the parasite per se
- Strongyloides stercoralis**
 Gherman, I.; et al., 1970, *Med. Int.*, Bucuresti, v. 22 (3), 335-344
 giardiasis, *Strongyloides stercoralis*, attempt to establish parasitic duodenitis as a clinical entity
- Strongyloides stercoralis**
 Gill, G. V.; Bell, D. R.; and Reid, H. A., 1977, *Brit. Med. J.* (6067), v. 1, 1007
Strongyloides stercoralis, long-standing infections in ex-Far East prisoners of war frequently presenting as rash with urticarial eruptions, good response to thiabendazole; similar therapy suggested for persons who have been in tropics and develop typical rash: Liverpool, England
- Strongyloides stercoralis**
 Goulart, E. G.; et al., 1977, *J. Helminth.*, v. 51 (2), 131-132
 ancylostomiasis, *Strongyloides stercoralis*, human, field trials of control by means of mass anthelmintic treatment combined with introduction of plants inhibitory to the free-living larval stages, reduction in prevalence: Ilha Do Governador, Rio de Janeiro, Brazil
- Strongyloides stercoralis**
 Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, *J. Infect. Dis.*, v. 131 (6), 755-758
 algorithms in the diagnosis and management of human *Strongyloides stercoralis*
- Strongyloides stercoralis**
 Hazel, J. R.; and Grant, A. K., 1969, *Med. J. Australia*, v. 2 (23), 1159-1160
Strongyloides stercoralis in Australian soldier who had served in South-East Asia, marked eosinophilia, malaise and skin rash as presenting symptoms, case report, cured with thiabendazole
- Strongyloides stercoralis**
 Hira, P. R.; and Patel, B. G., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (4), 640-643
Strongyloides fuelleborni rhabditiform larvae and eggs cultured to free living adults obtained from human feces, discussion of differential diagnosis from *S. stercoralis*, prominent morphologic features, mode of human infections, survey of prevalence in predominantly urban and suburban areas of Zambia
- Strongyloides stercoralis**
 Hsieh, H. C., 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 568 [Demonstration]
Ancylostoma duodenale, *Necator americanus* and *Strongyloides stercoralis*-related spp., differentiation of species using a modified test-tube filter-paper culture
- Strongyloides stercoralis**
 Islam, N.; and Chowdhury, N. A., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 81-84
 human intestinal parasites, comparative treatment trials using mebendazole and pyrantel pamoate, recommendations for use: Dacca, Bangladesh
- Strongyloides stercoralis**
 Jacquemin, J. L.; and Le Flohic, A. M., 1975, *Nouv. Presse. Med.*, v. 4 (31), 2271 [Letter]
Strongyloides stercoralis, possible role of public swimming pools in transmission of infection from man to man

- Strongyloides stercoralis*
Jakubowski, A.; Jakubowska, K.; and Brzechwa-Ajdukiewicz, A., 1975, *Polski Tygod. Lekar.*, v. 30 (28), 1175-1178
Strongyloides stercoralis, diagnostic radiologic findings in human infections
- Strongyloides stercoralis*
Jose, D. G.; and Welch, J. S., 1970, *Med. J. Australia*, v. 1 (8), 349-356
possible role of intestinal parasitism in growth-retarded, anemic and malnourished Australian Aboriginal children, comparison with normal Aboriginal children: Queensland
- Strongyloides stercoralis*
Junod, C.; 1972, *Medecine et Malad. Infect.*, v. 2 (2), 55-60
Strongyloides stercoralis, predominant intestinal parasite of natives of Antilles now residing in France, statistics of infection, means of detection, approved treatment with mintezol
- Strongyloides stercoralis*
Kagei, N.; et al., 1976, *Bull. Inst. Pub. Health, Tokyo*, v. 25 (3), 140-144
Filaroides hirthei, Strongyloides sp., Toxocara canis, and Trichuris vulpis in imported dogs, inability to experimentally infect other dogs with Strongyloides stercoralis of human origin: Japan, imported from U.S.A.
- Strongyloides stercoralis*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Strongyloides stercoralis*
Krubwa, F.; et al., 1974, *Medecine Trop.*, v. 34 (5), 679-687
trial administration of mebendazole to school children on quarterly basis for attempted eradication of intestinal nematodes, recommendations for use as mass control measure: Zaire
- Strongyloides stercoralis*
Layrisse, M.; and Vargas, A., 1975, *Progr. Food and Nutrition Sc.*, v. 1 (10), 645-667
human intestinal parasites, mechanisms by which parasites interfere with host nutrition (competition for nutrients, malabsorption, blood loss, excess nutrient utilization), extensive review
- Strongyloides stercoralis*
Levai, J.; Lengyel, A.; and Banki, G., 1972, *Med. Universalis*, v. 5 (1), 31-34
clinical trials using levamisole to treat human intestinal helminths, good results against Ascaris, Ancylostoma, Trichostrongylus, ineffective against Trichuris trichiura: Hungary
- Strongyloides stercoralis*
Matern, B.; et al., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 217-218
Strongyloides stercoralis, studies in anthropoid apes, evidence for autoinfection and prenatal infection: Zoologischen Gartens, Frankfurt
- Strongyloides stercoralis*
Mougeot, G.; et al., 1975, *Medecine et Malad. Infect.*, v. 5 (4), 208-212
analysis of 400 cases of strongyloidiasis diagnosed in patients in Paris hospital and geographic origin
- Strongyloides stercoralis*
Nauenberg, W.; Edelman, M. H.; and Spingarn, C. L., 1970, *Mount Sinai J. Med.*, v. 37 (5), 607-611
chronic Strongyloides stercoralis infection in humans, 96% cure rate in trials with thiabendazole: New York
- Strongyloides stercoralis*
O'Brien, W., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 69-77
Strongyloides stercoralis, intestinal malabsorption associated with high eosinophilia occurring in young soldiers probably resulting from parasitic infestation: Borneo
- Strongyloides stercoralis*
Otsuru, M., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Strongyloides stercoralis, illus.*
Owor, R.; and Wamukota, W. M., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (5-6), 1976, 497-499
Strongyloides stercoralis in man, fatal overwhelming strongyloidiasis associated with pyogenic meningitis and presence of larvae in meninges, probably depression of cellular immunity (unknown cause) contributed to fatal infection: Uganda
- Strongyloides stercoralis*
Poltera, A. A., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (1), 81-90
Strongyloides stercoralis, man, 5 fatal cases with autoinfection, case reports, histopathological findings: Uganda
- Strongyloides stercoralis, illus.*
Powles, A. C. P., 1973, *N. Zealand Med. J.* (490), v. 77, 169-171
Strongyloides stercoralis hyperinfection in previously healthy young man being treated with steroids to suppress brain edema caused by head injury, fatal illness, case report, thiabendazole therapy unsuccessful: New Zealand
- Strongyloides stercoralis*
Purtilo, D. T.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (2), 229-232
parasitized children with protein-calorie malnutrition, complete blood count, serum immunoglobulin concentration, significant relationship between intensity of parasitism and Ig levels: Brazil

- Strongyloides stercoralis*
Radermecker, M.; et al., 1974, Internat. Arch. Allergy and Applied Immunol., v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months
- Strongyloides stercoralis*
Rees, P. H.; and Marsden, P. D., 1970, Brit. J. Clin. Pract., v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment
- Strongyloides stercoralis*
Reid, B. D.; Reid, T. E.; and Ullstrop, G., 1971, Med. J. Zambia, v. 5 (2), 61-69
study of health of schoolchildren, parasitic survey and possible associations with nutritional status: Zambia
- Strongyloides stercoralis*
Rifaat, M.A.; et al., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (5), 722-723 [Letter]
Strongyloides stercoralis, 50-year old male farmer with nocturia and urinary incontinence cured by dithiazanine treatment, evidence supported diagnosis of infection of genitourinary system by *S. stercoralis*: Egypt
- Strongyloides stercoralis*
Scoggin, C. H.; and Call, N. B., 1977, Ann. Inst. Med., v. 87 (4), 456-458
disseminated *Strongyloides stercoralis*, infection in man who had undergone renal transplant, fatal illness unsuccessfully treated with thiabendazole, clinical case report: North Carolina
- Strongyloides stercoralis*
Seah, S. K. K., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (4), 534-542
intestinal parasites, persons living in non-endemic areas who acquired infections while travelling or who have immigrated from endemic areas, pyrantel pamoate successful for *Ascaris lumbricoides*, results with other parasites varied: Montreal, Canada
- Strongyloides stercoralis, illus.*
Smith, J. D.; Goette, D. K.; and Odom, R. B., 1976, Arch. Dermat., Chicago, v. 112 (8), 1161-1163
human systemic infection with *Strongyloides stercoralis* presenting as skin lesions of larva currens, clinical case report and management, discussion of autoinfection, thiabendazole therapy: California (lived earlier in southeastern United States)
- Strongyloides stercoralis, illus.*
Smith, S. B.; et al., 1977, J. Pediat., St. Louis, v. 91 (4), 607-609
Strongyloides stercoralis, fatal autoinfection by disseminated strongyloidiasis in young child (lungs, liver, peritoneum, small intestine, colon, respiratory diaphragm, heart, lymph nodes, skeletal muscle, perirenal and peripancreatic fat), presenting symptoms of acute abdominal distress and mixed intestinal parasites, clinical case report: Miami, Florida
- Strongyloides stercoralis*
Sohi, B. K., 1975, Pediat. Clin. India, v. 10 (3), 255
mixed strongyloidiasis and ancylostomiasis in young child, presenting symptom of acute abdomen, cure with mintezole: Pathankot, India
- Strongyloides stercoralis, illus.*
Soroczan, W., 1976, Ann. Univ. Mariae Curie-Skłodowska, Sect. C, Biol., v. 31, 281-289
Strongyloides stercoralis, morphology of developmental forms of parasitic and free-living generations, diagnosis determined by morphological features of filariform larvae only after 48 hours of culture
- Strongyloides stercoralis, illus.*
Sotolongo Guerra, F., 1972, Parassitologia, v. 14 (1), 29-33
Strongyloides stercoralis, human case, adult parasitic or strongyloid type males found, apparent variation on usual life cycle, more knowledge about life cycle needed
- Strongyloides stercoralis*
Viens, P.; and Beal, C. B., 1973, Medecine Afrique Noire, v. 20 (12), 1009-1011
Strongyloides stercoralis, human diagnosis and evaluation of thiabendazole treatment using duodenal intubation
- Strongyloides stercoralis*
Walker-Smith, J. A.; et al., 1969, Med. J. Australia, v. 2 (25), 1263-1265
Aboriginal infant with small bowel obstruction caused by *Strongyloides stercoralis*, case report, surgical and medical management: Sydney, Australia
- Strongyloides stercoralis, illus.*
Whitehead, R., 1973, Major Problems Path., v. 3, 105-110
human intestinal infection, diagnosis, pathological appearance of mucosal biopsy of gastrointestinal tract
- Strongyloides stercoralis*
Williams, D. M.; Krick, J. A.; and Remington, J. S., 1976, Am. Rev. Resp. Dis., v. 114 (3), 593-627
parasitic pulmonary infections in the compromised host, immunopathologic aspects, clinical manifestations, diagnosis and treatment, review

- Strongyloides stercoralis*
Wing, E. J.; and Remington, J. S., 1977, West. J. Med., San Francisco, v. 126 (1), 14-31
role of cell-mediated immunity (and enhancement by levamisole) in host defense against human parasitic diseases, extensive review
- Strongyloides stercoralis*
Zuidema, P. J., 1976, Nederl. Tijdschr. Geneesk., v. 120 (21), 901-906
human intestinal parasites, differential diagnosis of causes of diarrhea in hikers returning from visits to India: Netherlands
- Strongyloides stercoralis*, *illus.*
Zuidema, P. J., 1976, Nederl. Tijdschr. Geneesk., v. 120 (43), 1849-1854
human intestinal helminths, differential diagnosis and clinical management of parasitic infestations seen in immigrants from Surinam: Netherlands
- Strongyloides tumefaciens*, *illus.*
Malone, J. B.; et al., 1977, J. Am. Vet. Med. Ass., v. 171 (3), 278-280
cats, case report
- Strongyloides turkmenica* Kurtieva, 1953, *illus.*
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
description, nematodes of *Larus canus* (intestine), age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Strongyloides venezuelensis*
Kelly, A.; Little, M. D.; and Voge, M., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mintagez, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides westeri*
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, Prakt. Tierarzt, v. 55 (8), 417-422
nematodes of horses, Fenbendazol, good results
- Strongyloides westeri*
Hiepe, T., 1972, Schweiz. Arch. Tierh., v. 114 (12), 613-614
Parascaris equorum; *Strongyloides westeri*, sports horses of all ages, Morantel tartrate, good results
- Strongyloides westeri*
Kelly, A.; Little, M. D.; and Voge, M., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 694-699
Strongyloides sp. (*S. fuelleborni*-like) infections in natives discovered during intestinal parasitic survey, mintagez, morphological comparisons with other *Strongyloides* spp.: Fly River, Kiunga, Papua
- Strongyloides westeri*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 721-723
horses (feces)
- Strongyloides westeri*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 889-892
Strongyloides westeri, foals (exper.), prepatent period, efficacy of cambendazole and thiabendazole against 3rd stage larvae in milk of nursing mares during postpartum period, good results with cambendazole
- Strongyloides westeri*, *illus.*
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Strongyloides westeri*, *illus.*
Roedel, I. E.; and Lyons, E. T., 1977, Ann. Clin. and Lab. Sc., v. 7 (5), 405-410
Strongyloides westeri causing cutaneous larva migrans with severe allergic hypersensitivity in laboratory workers exposed to live *Strongyloides* larvae, clinical case reports, recommendations for control; similar lesions and pathologic changes produced in rabbit experimentally
- Strongyloides westeri*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongyloidiasis*
Bitoun, A.; et al., 1972, Nouv. Presse Med., v. 1 (29), 1935-1937
massive severe strongyloidiasis in young woman being treated with cortico-steroids, thiabendazole therapy, clinical case report: Paris, France (native of Gabon)
- Strongyloidiasis*
Burke, G. J., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (4), 402-405
10 patients with presumed parasitological disease, circulating absolute eosinophil levels over a 24 hour period, periodicity, steroid administration will not separate parasitic from other causes of eosinophilia
- Strongyloidiasis*
Cezner, M.; et al., 1970, Medicina, Rijeka, v. 7 (1), 77-81
epidemiology and clinical aspects of human strongyloidiasis in the Rijeka region of Yugoslavia
- Strongyloidiasis*
Coulaud, J. P.; et al., 1975, Nouv. Presse Med., v. 4 (16), 1208-1209 [Letter]
human massive fatal strongyloidiasis associated with immunosuppressive therapy after renal transplant, clinical review

- Strongyloidiasis**
 Coulaud, J. P.; Pasticier, A.; and Saint-Louvent, P., 1976, *Nouv. Presse. Med.*, v. 5 (44), 3013-3014 [Letter]
 human strongyloidiasis with accompanying acute epigastric distress, clinical aspects, thiabendazole
- Strongyloidiasis**
 Dale, D. C.; and Petersdorf, R. G., 1973, *Med. Clin. North Am.*, v. 57 (5), 1277-1287
 parasitic infections in the immuno-suppressed steroid-treated patient, review
- Strongyloidiasis**
 Diop, B.; and Bao, O., 1974, *Medecine Afrique Noire*, v. 21 (1), 31-40
 human intestinal helminths, clinical indications for treatment, suggested dosage, efficacy, tolerances, possible toxicities
- Strongyloidiasis**
 Marques, J. C.; and Ogawa, K. M., 1967, *Maternidade e Infancia*, v. 26 (4), 405-412
 infection in children, case reviews, clinical management, complications: Sao Paulo, Brasil
- Strongyloidiasis**
 Molet, B.; and Derr-Harf, C., 1975, *Medecine Trop.*, v. 35 (4), 336-337
 ancylostomiasis, strongyloidiasis, improved coproculture technique for diagnosis
- Strongyloidiasis**
 Pasticier, A., 1975, *Nouv. Presse. Med.*, v. 4 (37), 2659-2660
 single and mixed parasitic infections in humans, good results with pyrantel pamoate
- Strongyloidiasis**
 Poirot, J. L.; et al., 1976, *Medecine et Malad. Infect.*, v. 6 (1), 32-35
 survey of origination of patients infected with strongyloidiasis and ancylostomiasis in Paris hospital
- Strongyloidiasis**
 Tarabini Castellani, G., 1970, *Minerva Gastroenterol.*, v. 16 (1), 45-49
 human intestinal parasites, clinical trials using FI 6518 with and without thymol, acceptable drug for mass therapy: Somalia
- Strongyloidiasis**
 Thomas, J.; et al., 1969, *Marseille Med.*, v. 106 (12), 1033-1040
 strongyloidiasis, human, thiabendazole, good results: France
- Strongyloidiasis**
 Thomas, J.; et al., 1971, *Medecine Trop.*, v. 31 (2), 177-181
 bephenium hydroxynaphthoate, piperazine and thiabendazole in comparative clinical tests, institutionalized children with protein deficiency
- Strongyloidiasis**
 Wlodarczak, C., 1972, *Med. Wet.*, v. 28 (4), 221-222
 strongyloidosis, *Parascaris equorum*, horses, Pyrequan
- Strongyloidosis.** See **Strongyloidiasis.**
- Strongylosis**
 Buonaccorsi, A.; et al., 1972, *Atti Soc. Ital. Sc. Vet.*, v. 26, 312-315
 horses, chronic strongylosis, dysfunction of liver and adrenal cortex
- Strongylosis**
 Tassi, P.; and Widenhorn, O., 1972, *Parassitologia*, v. 14 (2-3), 381-398
 gastrointestinal strongylosis, survey by fecal examination, incidence, cattle, sheep, goats, horses, pigs, 1,788 farms, various regions: Italy
- Strongyluris chamaeleonis** (Baylis & Daubney 1922), illus.
 Sinha, A.; and Sahay, U., 1969, *Indian J. Helminth.*, v. 21 (1), 18-22
 description
 Calotes versicolor (intestine, stomach): Ranchi (Morhabadi Hill)
- Strongylus**
 Blanchard, J., 1971, *Prat. Vet. Equine*, v. 3 (1), 29-31
 gastrointestinal nematodes in race horses, clinical trials with equigard, good immediate results especially with ascarids; rapid reinfection rates (2 months after therapy), suggests contaminated paddocks; recommendations therefore for program of routine treatment: France
- Strongylus**
 Grellck, H.; Hoerchner, F.; and Woehrl, H., 1977, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 90 (19), 371-373
Fasciola hepatica-infected ponies, orally and intraperitoneally, rate of development of parasite, most liver flukes found in animals severely infected with *Strongylus* and *Trichonema*
- Strongylus**
 Neppert, J., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 454-463
 cross-reacting antigens among some filariae and other helminths, closed hexagonal immunodiffusion technique, implications for serodiagnosis of filariasis
- Strongylus**
 Reinecke, R. K.; and le Roux, D. J., 1972, *J. South African Vet. Ass.*, v. 43 (3), 287-294
 adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective

- Strongylus**
Shimura, K.; et al., 1975, Exper. Rep. Equine Health Lab. (12), 30-43
rabbits, sensitized with extract of equine *Strongylus*, same extract injected into ileoceocolic artery, pathological changes in mesentery and cecum; injection in ligatured loops of jejunum, ileum and vermiform appendix, erythrocytes, hemoglobin and serous exudate in lumina
- Strongylus [sp.]**
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
donkeys, survey, treatment, economic importance: Kaedi area (Mauritania)
- Strongylus spp.**
Ismail, E.; Tawfik, A. A.; and El-Ebrashi, N. M. A., 1977, Arzneimittel-Forsch., v. 27 (7), 1393-1394
6-hydroxy-4-methoxy-5-(p-methoxy-cinnamoyl)-benzofuran, in vitro broad spectrum anthelmintic activity against livestock helminths, promising results indicate need for future research
- Strongylus alatus Leuckart 1848**
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus alatus* (Leuckart 1848)
Stiles and Hassall 1905
- Strongylus (Pharurus) alatus (Leuckart) Diesing 1851**
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus alatus* (Leuckart 1848)
Stiles and Hassall 1905
- Strongylus arcticus Cobb 1888**
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden 1870) n. comb.
- Strongylus convolutus Kuhn 1829**
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829)
Baylis and Daubney 1925
- Strongylus edentatus**
Abdel-Rahman, M. S.; et al., 1972, Parasitol. Hungar., v. 5, 225-237
Strongylus spp., *Trichonema* spp., *Parascaris equorum*, *Trichostrongylus axei* in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Strongylus edentatus**
Chiejina, S. N.; and Mason, J. A., 1977, Vet. Rec., v. 100 (17), 360-361
horse: [Britain]
- Strongylus edentatus**
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Strongylus edentatus**
Drudge, J. H.; and Lyons, E. T., 1977, Am. J. Vet. Research, v. 38 (10), 1581-1586
internal parasites, horses, antiparasitic drugs, methods of evaluation, critical tests, controlled tests and clinical trials, review of methods and criteria
- Strongylus edentatus**
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Strongylus edentatus**
Duncan, J. L.; et al., 1977, Equine Vet. J., v. 9 (3), 146-149
immature strongyles, ponies, fenbendazole
- Strongylus edentatus**
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, Prakt. Tierarzt, v. 55 (8), 417-422
nematodes of horses, Fenbendazol, good results
- Strongylus edentatus**
Folz, S. D., 1977, Vet. Parasitol., v. 3 (4), 377-381
kinetics of natural expulsion of some equine parasites from nontreated horses during interim of critical test, concluded that natural elimination would not have biased efficacy of an anthelmintic or boticide
- S[trongylus] edentatus**
Grelck, H.; Hoerchner, F.; and Woehrl, H. E., 1977, Prakt. Tierarzt, v. 58 (4), 265-268
strongyles, horses, seasonal development of infective larvae on pastures, survival rate
- Strongylus edentatus**
Greve, J. H.; and Paul, J. W., 1976, Vet. Med. and Small Animal Clin., v. 71 (12), 1737-1740-1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Strongylus edentatus**
Lewis, L. M.; et al., 1977, Biol. Bull., v. 153 (2), 436 [Abstract]
Strongylus edentatus, ultrastructure of syncytial intestinal epithelium, possible contraction mechanisms
- Strongylus edentatus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis

- Strongylus edentatus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, *Am. J. Vet. Research*, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and *Gasterophilus* (inactive against latter)
- Strongylus edentatus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (6), 721-723
helminths and bots in horses, thiabendazole and trichlorfon sequentially administered via stomach tube, critical testing, drug efficacies, good results
- Strongylus edentatus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Strongylus edentatus**
Mirzayans, A., 1973, *Vet. Rec.*, v. 92 (10), 262 [Letter]
pony (intestinal tract): northern Iran
- Strongylus edentatus**
Nawalinski, T.; and Theodorides, V. J., 1976, *Am. J. Vet. Research*, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Strongylus edentatus**
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Equus caballus: Chile
- Strongylus edentatus**
Ogbourne, C. P., 1976, *J. Helminth.*, v. 50 (3), 203-214
horses (large intestine): south-west England
- Strongylus edentatus**
Ooms, L., 1975, *Vlaams Diergeneesk. Tijdschr.*, v. 44 (3), 95-118
strongyles, horses, epidemiology, cycles, pathogenesis, symptoms, control, diagnosis, immunity, review
- Strongylus edentatus**
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Strongylus edentatus, illus.**
Rizzoli-Stalder, C.; et al., 1976, *Schweiz. Arch. Tierh.*, v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Strongylus edentatus**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongylus equinus**
Abdel-Rahman, M. S.; et al., 1972, *Parasitol. Hungar.*, v. 5, 225-237
Strongylus spp., *Trichonema* spp., *Parascaris equorum*, *Trichostrongylus axei* in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Strongylus equinus**
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, *J. Parasitol.*, v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Strongylus equinus**
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, *Prakt. Tierarzt*, v. 55 (8), 417-422
nematodes of horses, Fenbendazol, good results
- S[trongylus] equinus**
Grelck, H.; Hoerchner, F.; and Woehrl, H. E., 1977, *Prakt. Tierarzt*, v. 58 (4), 265-268
strongyles, horses, seasonal development of infective larvae on pastures, survival rate
- Strongylus equinus**
Greve, J. H.; and Paul, J. W., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (12), 1737-1740, 1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Strongylus equinus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, *Am. J. Vet. Research*, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and *Gasterophilus* (inactive against latter)
- Strongylus equinus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (6), 721-723
helminths and bots in horses, thiabendazole and trichlorfon sequentially administered via stomach tube, critical testing, drug efficacies, good results
- Strongylus equinus**
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, *Am. J. Vet. Research*, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations

- Strongylus equinus*
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Strongylus equinus*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Strongylus equinus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Strongylus equinus*
Ooms, L., 1975, Vlaams Diergeneesk. Tijdschr., v. 44 (3), 95-118
strongyles, horses, epidemiology, cycles, pathogenesis, symptoms, control, diagnosis, immunity, review
- Strongylus equinus* Mueller
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Equus burchelli (intestines): Kenya
- Strongylus equinus*
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Strongylus equinus*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Strongylus inflexus* Rudolphi 1808
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pseudalius inflexus* (Rudolphi 1808) Schneider 1866
- Strongylus inflexus major* (Rudolphi) Raspail 1829
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pseudalius inflexus* (Rudolphi 1808) Schneider 1866
- Strongylus invaginatus* Quekett 1841
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Halocercus invaginatus* (Quekett 1841) Dougherty 1943
- Strongylus major* Raspail 1829
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pseudalius inflexus* (Rudolphi 1808) Schneider 1866
- Strongylus minor* Kuhn 1829
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Stenurus minor* (Kuhn 1829) Baylis and Daubney 1925
- Strongylus pallasii* van Beneden 1870
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Pharurus pallasii* (van Beneden 1870) n. comb.
- Strongylus vagans* Eschricht 1841
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of *Torynurus convolutus* (Kuhn 1829) Baylis and Daubney 1925
- Strongylus vulgaris*
Abdel-Rahman, M. S.; et al., 1972, Parasitol. Hungar., v. 5, 225-237
Strongylus spp., *Trichonema* spp., *Parascaris equorum*, *Trichostrongylus axei* in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Strongylus vulgaris*
Amborski, G. F.; Bello, T. R.; and Torbert, B. J., 1974, Am. J. Vet. Research, v. 35 (9), 1181-1188
Strongylus vulgaris, more adverse host reaction in parasite-free ponies than in ponies sensitized by previous natural infection, changes in serum glycoprotein patterns may be related to arterial damage associated with larval migrations
- Strongylus vulgaris*
Bello, T. R.; Amborski, G. F.; and Torbert, B. J., 1973, Am. J. Vet. Research, v. 34 (10), 1291-1297
methotrexate, immunosuppressive and toxic effects used alone or in combination with antihistamine compounds on parasite-free or *Strongylus vulgaris*-infected ponies
- Strongylus vulgaris*
Chiejina, S. N.; and Mason, J. A., 1977, Vet. Rec., v. 100 (17), 360-361
horse (anterior mesenteric artery and associated thrombus): [Britain]
- Strongylus vulgaris*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Strongylus vulgaris*
Drudge, J. H.; and Lyons, E. T., 1977, Am. J. Vet. Research, v. 38 (10), 1581-1586
internal parasites, horses, antiparasitic drugs, methods of evaluation, critical tests, controlled tests and clinical trials, review of methods and criteria

- Strongylus vulgaris*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Strongylus vulgaris*
Duncan, J. L.; et al., 1977, Equine Vet. J., v. 9 (3), 146-149
immature strongyles, ponies, fenbendazole
- Strongylus vulgaris*
Duncan, J. L.; and Pirie, H. M., 1975, Research Vet. Sc., v. 18 (1), 82-93
Strongylus vulgaris, single experimental infections of worm-free pony foals, clinical signs, pathology (intestinal and arterial lesions), and clinical pathology (haematology, serum proteins)
- Strongylus vulgaris*
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, Prakt. Tierarzt, v. 55 (8), 417-422
nematodes of horses, Fenbendazol, good results
- Strongylus vulgaris*
Fatzer, R., 1976, Schweiz. Arch. Tierh., v. 118 (11), 499-502
cerebral nematodiasis (probably Strongylus vulgaris), donkeys, possible cause of haemorrhagic, purulent and necrotizing meningoencephalitis
- Strongylus vulgaris*
Folz, S. D., 1977, Vet. Parasitol., v. 3 (4), 377-381
kinetics of natural expulsion of some equine parasites from nontreated horses during interim of critical test, concluded that natural elimination would not have biased efficacy of an anthelmintic or boticide
- Strongylus vulgaris*
Greatorex, J. C., 1977, Vet. Rec., v. 101 (10), 184-197
Strongylus vulgaris, horses, diagnosis and treatment of "verminous aneurysm" formation, dextran as antithrombotic agent, successful recovery
- S[trongylus] vulgaris
Grelck, H.; Hoerchner, F.; and Woehrl, H. E., 1977, Prakt. Tierarzt, v. 58 (4), 265-268
strongyles, horses, seasonal development of infective larvae on pastures, survival rate
- Strongylus vulgaris*
Greve, J. H.; and Paul, J. W., 1976, Vet. Med. and Small Animal Clin., v. 71 (12), 1737-1740, 1742
nematodes, horses, enteric-coated microencapsulated trichlorfon, critical and field evaluations, drug efficacy
- Strongylus vulgaris, illus.*
Little, P. B., 1972, J. Am. Vet. Med. Ass., v. 160 (10), 1407-1413
Strongylus vulgaris, horses, cerebrospinal nematodiasis, clinical syndromes, pathology, case reports
- Strongylus vulgaris*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Strongylus vulgaris*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, Am. J. Vet. Research, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and Gasterophilus (inactive against latter)
- Strongylus vulgaris*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 721-723
helminths and bots in horses, thiabendazole and trichlorfon sequentially administered via stomach tube, critical testing, drug efficacies, good results
- Strongylus vulgaris*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Strongylus vulgaris*
McCraw, B. M.; and Slocombe, J. O. D., 1976, Canad. Vet. J., v. 17 (6), 150-156
horse, development, pathogenesis, clinical signs, diagnosis, epidemiology, control and treatment, brief review
- Strongylus vulgaris*
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxibendazole
- Strongylus vulgaris*
Ober, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Equus caballus: Chile
- Strongylus vulgaris*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Strongylus vulgaris*
Ooms, L., 1975, Vlaams Diergeneesk. Tijdschr., v. 44 (3), 95-118
strongyles, horses, epidemiology, cycles, pathogenesis, symptoms, control, diagnosis, immunity, review

- Strongylus vulgaris*
Ooms, L.; et al., 1976, Vlaams Diergeneesk. Tijdschr., v. 45 (9-10), 290-299
helminthiasis in horses with particular reference to verminous aneurysms (*Strongylus vulgaris*), diagnosis, value of electrophoresis of serum proteins in addition to clinical and coprological examination
- Strongylus vulgaris*
Orr, J. P., 1972, Vet. Rec., v. 90 (20), 571
Ascaris equorum in Thoroughbred foal (peritoneal cavity, stomach), probable cause of fatal perforated duodenal ulcer; accompanied by aneurysm at root of anterior mesenteric artery containing near-adult *Strongylus vulgaris*, case report
- Strongylus vulgaris*
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Strongylus vulgaris*
Patton, S.; and Drudge, J. H., 1977, Am. J. Vet. Research, v. 38 (12), 2059-2066
Strongylus vulgaris, ponies given repeated small doses of infective larvae, acquired resistance against challenge exposure; clinical and hematologic responses, corticosteroid and/or antibiotic therapy did not alter immune response
- Strongylus vulgaris*
Perce, R. B.; and Shideler, R. K., 1976, Vet. Med. and Small Animal Clin., v. 71 (8), 1103-1107
Strongylus vulgaris, horse, septic thrombosis, case report, pathology
- Strongylus vulgaris* Looss
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Equus burchelli (intestines): Kenya
- Strongylus vulgaris, illus.*
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Strongylus vulgaris*
Slocombe, J. O. D.; et al., 1977, Canad. J. Comp. Med., v. 41 (2), 137-145
Strongylus vulgaris, ponies, value of arteriography for revealing vascular lesions
- Strongylus vulgaris*
Sutoh, M.; et al., 1976, National Inst. Animal Health Quart., v. 16 (2), 59-64
Strongylus vulgaris, pathology in foals: Japan
- Strongylus vulgaris, illus.*
Sutoh, M.; et al., 1976, Exper. Rep. Equine Health Lab. (13), 60-78
Strongylus vulgaris, foals, occurrence, clinical signs, pathologic changes in mesenteric artery and related blood vessels caused by larval migration: Hidaka district, Hokkaido
- Strongylus vulgaris*
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Struthiofilaria* gen. n.
Noda, R.; and Nagata, S., 1976, Bull. Univ. Osaka Prefect., s. B, Agric. and Biol., v. 28, 1-4
Onchocercidae, Onchocercinae; tod: *S. megalocephala* sp. n.
- Struthiofilaria megalocephala* sp. n. (tod), *illus.*
Noda, R.; and Nagata, S., 1976, Bull. Univ. Osaka Prefect., s. B, Agric. and Biol., v. 28, 1-4
Struthio camelus (body cavity): Misaki Park Zoo, Osaka Prefecture, Japan (imported from Republic of South Africa)
- Stunkardionema* Arnold, 1941
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of *Heligmostrongylus* Travassos, 1917
- Subulura* sp. I
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Dila bucharica
Trigonoscelis gemmulata
Blaps fausti bactriana
Pseudeuthripta tadzhiikistana
Pachyscelis laevicollis
Adesmia gebleri
all from Tadzhiik SSR [and/or] Uzbek SSR
- Subulura* sp. II
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Prosodes bactriana
Pelorocnemis punctata
Pachyscelis banghaasi
Adesmia gebleri
Blaps fausti bactriana
Dissonomus sp.
all from Tadzhiik SSR
- Subulura* sp.
Gupta, N. K.; and Dutt, K., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 185-188
description
Loris tardigradus (intestine): Madras (India)

- Subulura* spp.
Radhakrishnan, C. V.; and Ebrahimina, A., 1975, J. Vet. Fac. Univ. Tehran, v. 30 (4), 1-4
chickens (ceca): Darab, Fars Province, Iran
- Subulura allodapa*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Subulura andersoni* Cobbold, 1876
Peters, W.; et al., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (1), 3-4 [Demonstration]
Calliosciurus nigrovittatus (ileocaecal region): Sabah
- Subulura andersoni* (Cobbold, 1876)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus bowersi
R. cremoriventer
all from Malaysia
- Subulura baylisi* Lopez-Neyra, 1946
Sanchez-Acedo, C.; and Vericad, J. R., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 197-203
Garrulus glandarius: Aragon Pyrenees
- Subulura distans* (Rudolphi, 1809) Railliet et Henry, 1913
Gupta, N. K.; and Dutt, K., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 185-188
measurements
Loris tardigradus (intestine): India
- Subulura elongata*, *illus.*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Subulura hindi* Mirza, 1936
Gupta, N. K.; and Dutt, K., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 185-188
measurements
Sciurus palmarum (intestine): Chandigarh
- Subulura minetti* Bhalerao, 1941
Bali, H. S.; and Kalra, I. S., 1975, J. Research, Punjab Agric. Univ., v. 12 (3), 313-316
fowl, domestic
fowl, desi
all from Punjab State, India
- Subulura nevadense* n. sp., *illus.*
Babero, B. B., 1973, Tr. Am. Micr. Soc., v. 92 (2), 265-272
[figs. for *S. nevadense* mislabelled as *S. leucurus*]
Spermophilus tereticaudus
Ammospermophilus leucurus
(caecum of all): all from Clark, Lincoln, and/ or Nye counties, Nevada
- Subulura samanamudi* sp. n., *illus.*
Ibanez H., N., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 137-139
Crotapha sulcirrostris (intestino): Moche, Peru
- Subulura suctoria* (Molin, 1860)
Fabiyl, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Subulura suctoria* Molin, 1860
Fabiyl, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 235-238
Numida meleagris galeata (caeca): Vom area, Benue Plateau State, Nigeria
- Subulura suctoria*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Adesmia biseriata
A. gracilentata
A. septemcostata
Blaps oblonga
all from Uzbekistan
- Subulura suctoria*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Subulura williaminglisi*
Seureau, C.; and Quentin, J. C., 1977, Ann. Parasitol., v. 52 (4), 457-470
comparison of larval migration of 17 subulurid and spirurid nematodes in *Locusta migratoria* (exper.), course and duration of migration, histopathologic consequences, brief discussion of relation to phylogeny of nematodes and host hemocytic defense reaction
- Subuluroidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Ascaridida
key
- Sulcascaris* Hartwich (1957)
Allison, V. F.; et al., 1973, Tr. Am. Micr. Soc., v. 92 (2), 291-297
as syn. of *Porrocaecum*
- Sulcascaris*
Gibson, D. I.; and Taylor, A. L., 1976, Parasitology, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function

- Sulcascaris* Hartwich, 1957
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinea
key
- Sulcascaris*
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (4), 379-387
genus upheld and redefined
- Sulcascaris sulcata*, illus.
Sprent, J. F. A., 1977, J. Helminthol., v. 51 (4), 379-387
redescription, life history
Chelonia mydas
Caretta caretta (stomach) (nat. and exper.): Eastern Mediterranean; off coast of Queensland
Amusium balloti (adductor muscle): Queensland coastal waters
Chlamys sp. (adductor muscle): Queensland coastal waters
- Sulphuretylenchus pugionifer* sp. n., illus.
Slankis, A., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 156-157
Hylastes cunicularius (body cavity): Shchelkovsk forest, Moskovsk oblast
- Suncinema Durette-Desset*, 1973
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmosomidae, Heligmosominae
- Suncinema witenbergi* n. sp., illus.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Crocidura russula (intestin grele): Masmiya junction, Israel
- Sychnotylenchidae* Paramonov, 1967
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Tylenchoidea
diagnosis, revised, keys to subfam. and gen. includes: *Sychnotylenchinae* subfam. n.; *Neoditylenchinae* subfam. n.
- Sychnotylenchinae* subfam. n.
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Sychnotylenchidae Paramonov, 1967 [lapsus as *Sylenchotylenchinae*], key, type gen.: *Sychnotylenchus* Ruehm, 1956
- Sychnotylenchus* Ruehm, 1956 (type gen.)
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
Sychnotylenchidae, *Sychnotylenchinae* subfam. n.
key
- Sylenchotylenchinae* subfam. n. [lapsus for *Sychnotylenchinae*]
Kakuliia, G. A.; and Devdariani, Ts. G., 1975, Soobshch. Akad. Nauk Gruzinsk. SSR, v. 78 (3), 713-716
- Syncuaria* Gilbert, 1927, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
Syn.: *Decorataria* Sobolev, 1949
- Syncuaria* sp.
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Syngamus*
Valenza, F., 1974, Atti Soc. Ital. Sc. Vet., v. 28, 704-705
Syngamus, pheasants, tracheal granulomas, pathology, description of lesions
- Syngamus* sp.
Faust, B. S.; and Pappas, P. W., 1977, J. Zoo Animal Med., v. 8 (1), 18-23
Aix galericulata (feces): Columbus (Ohio) Zoo
- Syngamus*-type, illus.
Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
helminths of pet birds, diagnosis of eggs in fecal examination
- Syngamus anterogonimus* Ryjikov, 1949
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Sterna hirundo (trachea): coast of Sea of Okhotsk (Tuguro-Chumikansk and Ol'sk regions)
- Syngamus gibbocollis* Ryjikov, 1949
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Gallinago gallinago: Keta lake
- Syngamus palustris* Ryjikov, 1949
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Philomachus pugnax: lower Yenisei and Keta lake
- Syngamus trachea* (Montagu, 1811)
Bakke, T. A., 1973, Norwegian J. Zool., v. 21 (4), 299-303
Syngamus trachea, geographic distribution, epizootiology (climatic conditions, infection of *Sterna paradisaea* occurs before northward migration)
Pica pica: Oslo, Norway
Pavo cristatus: Jaeren, Norway
Passer domesticus: Halden, Norway
Turdus pilaris: Oslo/Baerum, Norway
Sterna paradisaea (trachea): Agdenes, Norway
- Syngamus trachea*
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (trachea): Florida

- Syngamus trachea*
 Connan, R. M.; and Wise, D. R., 1977, Vet. Rec., v. 101 (2), 34-35
Syngamus trachea in pheasants and turkeys (both exper.), dichloroxylenol (no effect), tetramisole (moderate efficacy against three- and four-day-old larvae, poor results against seven-day-old and older larvae)
- Syngamus trachea*
 Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (trachea): South Bass Island, Ohio
- Syngamus trachea*
 Cooper, C. L.; and Crites, J. L., 1974, J. Wildlife Dis., v. 10 (4), 399-403
 survey, helminths of red-winged blackbirds including a check list of previous findings
Agelaius phoeniceus (trachea): South Bass Island, Ohio
- Syngamus trachea* (Montagu, 1811)
 Cooper, C. L.; and Crites, J. L., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (trachea): South Bass Island, Ottawa County, Ohio
- Syngamus trachea*
 Cooper, C. L.; and Crites, J. L., 1976, J. Parasitol., v. 62 (1), 105-110
 similarity index of helminth faunas of 7 passerine bird species, index of association of 10 species of helminths identified as having foci of infection, competition for invertebrate food resources and aggregation into mixed feeding flocks maximizes transmission: South Bass Island, Ottawa County, Ohio
- Syngamus trachea*
 Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (trachea): Ottawa county, Ohio
- Syngamus trachea* (Montagu, 1811)
 Fabiyi, J. P., 1972, Bull. Epizoot. Dis. Africa, v. 20 (3), 229-234
 survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Syngamus trachea*
 Feteanu, A.; et al., 1973, Isotopes and Radiation Parasitol. III, 101-111
Syngamus trachea, chicks, pheasants, immunization with irradiated larval antigen, fluorescent antibody technique for detection of serum antibodies
- Syngamus trachea* (Montagu, 1811)
 Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (trachea): Florida
- Syngamus trachea*
 Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, J. Parasitol., v. 61 (3), 547-548
Grus canadensis pratensis (trachea): Florida
- Syngamus trachea*
 Frank, C., 1976, Ang. Parasitol., v. 17 (2), 99-100
 treatment with Chinosol, mortality in young storks
Ciconia ciconia (lungs, bronchi): Burgenland
- Syngamus trachea*
 Packer, D. E., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 15 [Demonstration]
Syngamus trachea, chicks (exper.), relationships between size of challenge infection, worm burden and egg production
- Syngamus trachea*
 Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix: CSSR
- Syngamus trachea*
 Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Syngamus trachea*
 Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
 fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Syngamus trachea*
 Winward, L. D., 1976, Exper. Parasitol., v. 40 (1), 74-76
Syngamus trachea, larvae injected into embryonic chicken eggs survived entire period of time necessary for embryos to develop and hatch, some migrated to trachea and completed life cycle in chickens hatched from these eggs
- Syngamus trachea*
 Winward, L. D.; and Russell, B. R., 1976, Exper. Parasitol., v. 40 (1), 77-79
Syngamus trachea, turkeys, infections produced by parenteral inoculations (larvae injected into veins, skin, peritoneum, musculature, and trachea)
- Syngamus trachea*
 Wissler, K.; and Halvorsen, O., 1975, J. Wildlife Dis., v. 11 (2), 245-247
Lagopus lagopus: Karlsoy Island and Sennaland, northern Norway

- Synhimantus* Railliet, Henry & Sisoff, 1912
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Acuariinae
key
includes subgens.: *Synhimantus*; *Dispharynx*
- Synhimantus*, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of *Synhimantus*
key
- Synhimantus* sp.
Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (esophagus): Florida
- Synhimantus* sp.
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (under gizzard lining): Florida
- Synhimantus* sp.
Forrester, D. J.; Bush, A. O.; and Williams, L. E., jr., 1975, J. Parasitol., v. 61 (3), 547-548
Grus canadensis pratensis (under gizzard lining): Florida
- Synhimantus* sp.
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (gizzard lining): Florida
- Synhimantus* sp.
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans (proventriculus): Florida
- Synhimantus australiensis* (Johnston and Mawson, 1952) Yamaguti, 1961, *illus.*
Beveridge, I.; and Barker, I. K., 1975, J. Helminth., v. 49 (4), 211-227
redescription
Syn.: *Cosmocephalus australiensis* Johnston and Mawson, 1952
Antechinus stuartii (stomach, lumen):
Powelltown, Sherbrooke, Dartmouth, and Mt. Sabine, Victoria
- Synhimantus australiensis* (Johnson & Mawson)
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, *Antechinus stuartii*, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (stomach): Powelltown, Victoria
- Synhimantus cramae* n. sp., *illus.*
Sharma, R. K., 1973, Riv. Parassitol., Roma, v. 34 (2), 111-114
Anastomus oscitans (gizzard, under the horny layer): Bareilly, U.P., India
- Synhimantus invaginatus*
Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (esophagus): Louisiana
- Synhimantus laticeps*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Synhimantus longevaginatus* (Molin 1860), *illus.*
Boero, J. J.; Led, J. E.; and Brandetti, E., 1972, Analecta Vet., v. 4 (1), 17-34
Euxenura maguari (esofago): Argentine Republic
- Syphacia*
Quentin, J. C., 1977, Ann. Parasitol., v. 52 (5), 559-567
Syphacia of gerbillids, murids, and microtids, hypothesis on biogeographic origin
- Syphacia*
Sharp, J. W.; and Wescott, R. B., 1976, Lab. Animal Sc., v. 26 (2, pt. I), 222-223
Aspiculuris, *Syphacia*, mice, mebendazole, good results
- Syphacia* sp.
Mozgovoï, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Microtus agrestis
Clethrionomys glareolus
Rattus norvegicus
(intestine of all): all from Karelia
- Syphacia* sp.
Nama, H. S.; and Parihar, A., 1976, J. Helminth., v. 50 (2), 99-102
Rattus rattus rufescens (intestine): Jodhpur City area, India
- Syphacia* sp.
O'Farrell, T. P., 1975, Am. Midland Naturalist, v. 93 (2), 377-387
Perognathus parvus: Arid Lands Ecology Reserve, Benton County, Washington
- Syphacia* [sp.]
Saxena, A.; and Nama, H. S., 1977, Geobios, v. 4 (6), 243-244
Rattus rattus (colon): Jodhpur, India
- Syphacia* sp.
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Mustela nivalis: Karelia

- Syphacia* sp.
Sood, M. L.; and Parshad, V. R., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 189-196
infections in *Millardia melitana*, survey of seasonal distribution, possible correlations between host diet and sex and incidence of infection
- Syphacia* (*Syphatineria*) *callosciuri* n. sp., illus.
Quentin, J. C., 1977, Ann. Parasitol., v. 52 (5), 559-567
phylogenetic relation to *Syphacia muris*
Callosciurus caniceps: Kuala Krai W. Malaysia
- Syphacia citelli* Tiner & Rausch, 1950
Babero, B. B., 1973, Tr. Am. Micr. Soc., v. 92 (2), 265-272
Spermophilus tereticaudus
Ammospermophilus leucurus
all from Nevada
- Syphacia mesocriceti*
Shava, F. H. M.; and Lewis, J. W., 1977, Parasitology, v. 75 (2), xxv-xxvi [Abstract]
Syphacia stroma, *S. obvelata*, *S. mesocriceti*, differences in general body surface, lip regions, eggs, and mamelons, scanning electron microscopy
- Syphacia montana*
Merkusheva, I. V., 1975, Vestsi Akad. Navuk BSSR, s. Biial. Navuk (6), 82-86
helminths of rodents as model for quantitative indices in analysis of faunistic and ecological studies
- Syphacia muris*, illus.
van der Gulden, W. J. I.; and van Aspert-van Erp, A. J. M., 1976, Exper. Parasitol., v. 39 (1), 40-44
Syphacia muris, egg hatching: effects of 22°C, 37°C, and cysteine on larval motility within closed egg and on subsequent hatching; effects of temperature, cysteine, and trypsin on permeability of water through eggshell; effect of water on opening of operculum
- Syphacia muris*
van der Gulden, W. J. I.; and van Aspert-van Erp, A. J. M., 1976, Exper. Parasitol., v. 39 (1), 45-50
Syphacia muris, effect of external stimuli on egg hatching (enzymes of intestinal tract, temperature, pH, pCO₂, redox potential), results indicate hatching mechanism of oxyurids identical to that of various nematodes which hatch in intestinal tract but dependent on environment to appreciably lesser extent
- Syphacia muris*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Ratte
- Syphacia muris*
Quentin, J. C., 1977, Ann. Parasitol., v. 52 (5), 559-567
phylogenetic relation to *Syphacia callosciuri* n. sp.
- Syphacia muris* (Yamaguti, 1935)
Singh, M.; and Cheong Chee Hock, 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (4), 516-521
Rattus rattus argentiventer
R. sabanus
all from Malaysia
- Syphacia muris* (Yamaguti, 1935) Yamaguti 1941
Singhvi, A.; and Johnson, S., 1977, J. Parasitol., v. 63 (5), 858-860
Aspiculuris ratti, *Syphacia muris*, female to male ratio of nematodes in concurrent infections in *Rattus rattus*, no significant correlation with worm burden, possible explanations
- Syphacia muris*
Strasser, H.; and Tiefenbach, B., 1976, Deutsche Tierarztl. Wchnschr., v. 83 (5), 224-226
Syphacia muris, rat breeding colony, fenbendazole
- Syphacia muris*, illus.
Taniguchi, M.; et al., 1977, Bull. Coll. Agric. and Vet. Med., Nihon Univ. (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo
- Syphacia obvelata*
Behnke, J. M., 1975, J. Helminth., v. 49 (2), 85-90
Aspiculuris tetraptera and *Syphacia obvelata*, survey of levels of infection in wild *Mus musculus*, prevalence of infection of *A. tetraptera* greater in male than in female mice: London Zoo
- Syphacia obvelata*
Berenguer Puvia, F. J.; and Gallego Berenguer, J., 1973, Rev. Iber. Parasitol., v. 33 (1), 81-106
Aspiculuris tetraptera, *Syphacia obvelata*, *Nippostrongylus brasiliensis*, mice, piperazine and phenothiazine compared with 4 phenothiazine-piperazine derivatives; piperazine: good activity, phenothiazine: low activity against *N. brasiliensis*, low toxicity of both; derivatives: more toxic, no anthelmintic activity
- Syphacia obvelata*
Berenguer Puvia, F. J.; and Gallego Berenguer, J., 1973, Rev. Iber. Parasitol., v. 33 (4), 573-598
Aspiculuris tetraptera, *Syphacia obvelata*, natural infections in mice, treatment with phenothiazine, piperazine hydrate, piperazine anhydrate; at various doses, calculation of elimination index and statistical analysis
- Syphacia obvelata*
Farell-Sala, A.; Berenguer-Puvia, F. J.; and Gallego-Berenguer, J., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 331-353
Aspiculuris tetraptera, *Syphacia obvelata*, mice, piperazine treatment, comparison of methods of measuring drug activity (deparasitization index and elimination index); relationships of dosage and activity, statistical analysis

- Syphacia obvelata*
Howes, H. L., jr., 1972, Proc. Soc. Exper. Biol. and Med., v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that *T. muris*-mouse infection could be test model for antiwhipworm studies
- Syphacia obvelata*
Lewis, J. W.; and Shava, F. H. M., 1977, Parasitology, v. 75 (2), iv [Abstract]
Syphacia obvelata, *Nematospiroides dubius*, differences in periodicity of egg deposition can be correlated with differences in transmission of infective stages to definitive host
- Syphacia obvelata*
McNair, D. M.; and Timmons, E. H., 1977, Lab. Animal Sc., v. 27 (1), 38-42
Syphacia obvelata and *Aspiculuris tetraptera*, effects on exploratory behavior of inbred mouse strain (exper.)
- Syphacia obvelata*
Martin, O. C., 1975, Philippine Agric., v. 59 (3-4), 114-118
brief description
Mus musculus: Bureau of Research and Laboratories, Alabang, Rizal
- Syphacia obvelata*
Merkusheva, I. V., 1975, Vestsi Akad. Navuk BSSR, s. Biial. Navuk (6), 82-86
helminths of rodents as model for quantitative indices in analysis of faunistic and ecological studies
- Syphacia obvelata*
Mishra, G. S.; and Gonzalez, J. P., 1975, Arch. Inst. Pasteur Tunis, v. 52 (1-2), 71-87
Rattus norvegicus (caecum): Tunis, Tunisia
- Syphacia obvelata*, illus.
Munn, E. A., 1977, Tissue and Cell, v. 9 (1), 23-34
Haemonchus contortus, structure of intestinal cells, helical polymeric extracellular protein associated with luminal surface for which name contortin is proposed, *Ostertagia circumcincta* also contained contortin-like material but *Nippostrongylus brasiliensis* and *Syphacia obvelata* contained material associated with outer surface of microvilli which was quite distinct from contortin
- Syphacia obvelata*
Owen, D., 1976, Lab. Animals, v. 10 (3), 271-278
Rattus norvegicus
Mus musculus
all from Carshalton
- Syphacia obvelata*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Maus
- Syphacia obvelata*
Shava, F. H. M.; and Lewis, J. W., 1977, Parasitology, v. 75 (2), xxv-xxvi [Abstract]
Syphacia stroma, *S. obvelata*, *S. mesocriceti*, differences in general body surface, lip regions, eggs, and mamelons, scanning electron microscopy
- Syphacia obvelata*
Singhal, K. C., 1976, Indian J. Exper. Biol., v. 14 (3), 345-347
berberine hydrochloride, in vivo activity against *Syphacia obvelata*, *Nippostrongylus brasiliensis*, and *Hymenolepis nana*, mice; elimination of *S. obvelata* only, drug considered equipotent to piperazine citrate
- Syphacia obvelata*
Taffs, L. F., 1975, J. Helminth., v. 49 (3), 173-177
continuous feed medication with thiabendazole for removal of *Hymenolepis nana*, *Syphacia obvelata*, and *Aspiculuris tetraptera* in naturally infected laboratory mice, unexplained deaths among inbred strain C3H/Hef Nimir mice
- Syphacia obvelata*
Taffs, L. F., 1976, Vet. Rec., v. 99 (8), 143-144
Hymenolepis nana, *Syphacia obvelata*, *Aspiculuris tetraptera*, mice, efficacy of thiabendazole given in diet
- Syphacia obvelata*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Syphacia obvelata*
Wescott, R. B.; Malczewski, A.; and Van Hoosier, G. L., 1976, Lab. Animal Sc., v. 26 (5), 742-745
filter top caging effective method for preventing pinworm infection in pathogen-free mice being introduced into laboratory colony where *Aspiculuris tetraptera* and *Syphacia obvelata* were enzootic
- Syphacia obvelata*, illus.
Wright, K. A., 1976, Organ. Nematodes (Croll), 71-105
cephalic anatomy of nematodes with astomatous and stomatous buccal capsules, integration of cephalic sense organs into the nematode head, definitions of "lips", "buccal capsule", and "stoma"
- Syphacia obvelata* (Rud., 1802)
Young, P. L.; and Babero, B. B., 1975, Proc. Oklahoma Acad. Sc., v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)

- Syphacia petrusewiczii* Bernard, 1966, illus.
 Sharpilo, L. D., 1975, Vestnik Zool., Akad.
 Nauk Ukrainsk. SSR, Inst. Zool. (4), 79-81
 description of male
Microtus subterraneus
Clethrionomys glareolus
 all from Zhitomirsk, Zakarpatsk and Kievsk
 oblasts, Ukrainian SSR
- Syphacia petrusewiczi* Bernard, 1966
 Wiger, R.; Lien, L.; and Tenora, F., 1976,
 Norwegian J. Zool., v. 24 (2), 133-135
Clethrionomys glareolus
Microtus agrestis
 (caecum of all): all from Kviteseid, Norway
- Syphacia petrusewiczi rauschi*, illus.
 Quentin, J. C.; and Gran, M. C., 1977, Ann.
 Parasitol., v. 52 (2), 231-234
 description of male
Clethrionomys rutilus: Alaska
- Syphacia sigmodoni*
 Kinsella, J. M., 1974, Am. Mus. Novitates
 (2540), 1-12
Sigmodon hispidus (cecum and large intes-
 tine): Florida
- Syphacia stroma*
 Shava, F. H. M.; and Lewis, J. W., 1977, Para-
 sitology, v. 75 (2), xxv-xxvi [Abstract]
Syphacia stroma, *S. obvelata*, *S. mesocri-
 ceti*, differences in general body surface,
 lip regions, eggs, and mamelons, scanning
 electron microscopy
- Syphacia stroma* (v Linstow, 1884)
 Wiger, R.; Lien, L.; and Tenora, F., 1976,
 Norwegian J. Zool., v. 24 (2), 133-135
Apodemus sylvaticus (small intestine):
 Oslo, Norway
- Syphacia thompsoni*
 Davidson, W. R., 1976, Proc. Helminth. Soc.
 Washington, v. 43 (2), 211-217
 epizootiologic and pathologic study of endo-
 parasites of selected populations of gray
 squirrels
Sciurus carolinensis (cecum, large intes-
 tine): southeastern United States
- Sypharista kamegaii* Quentin, 1971, illus.
 Kamiya, H.; Hasegawa, H.; and Chiba, K., 1976,
 Japan. J. Vet. Research, v. 24 (3-4), 99-100
 description, case of pseudoparasitism
Martes melampus (large intestine): Nakakubi-
 ki district, Niigata Prefecture

- Tachygonetria sp.
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Testudo graeca
Testudo haermani
all from Macedonia, Yugoslavia
- Tachygonetria dentata nearctica n. s. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
Gopherus sp., "vraisemblablement" G. agasizii: desert de Sonora, Mexique
Gopherus polyphemus: sud de Lake Placid, comte de Highlands, Floride (colon of all)
- Tachygonetria macrolaimus tetrapapillata (Caballero, 1944, espece) [n. rank], illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (389), Zool. (271), 731-768
description
Gopherus sp., "vraisemblablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraisemblablement" G. agasizii: desert de Sonora, Mexique
Gopherus polyphemus: sud de Lake Placid, comte de Highlands, Floride (colon of all)
- Tachygonetria robusta Drasche, 1883, illus.
Hanuskova, Z.; and Tilc, K., 1975, Acta Vet. Brno, v. 44 (4), 407-412
nematodes, incidence in Agrionemys horsfieldi (intestine) with regard to unsuitable conditions and food: Czechoslovakia, imported from USSR
- Tanqua Blanchard, 1904, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Gnathostomatinae
key
- Tanqua gigantea (Blanchard, 1904)
Majumder, S. S.; Mukherjee, O. P.; and Ghosh, P., 1975, Dobuts. Zasshi, Tokyo, v. 84 (3), 258-261
seasonal differences of infection rate, worm burden
Naja hannah: West Bengal villages
- Tanqua tiara (Linstow, 1879), illus.
Kan, S. P., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (3), 351-358
Tanqua tiara, ultrastructure of intestinal epithelium, presence of polymorphic inclusion bodies possibly associated with parasite's mode of feeding
Varanus salvator (digestive tract): Singapore
- Taphozoa Ali & Lovekar, 1967
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of Spirura Blanchard, 1849
- Tarsubulura perarmata, illus.
Quentin, J. C.; Krishnasamy, M.; and Tcheparkoff, R., 1977, Ann. Parasitol., v. 52 (2), 159-170
Tarsubulura perarmata, life cycle, organogenesis and morphology of larval stages, ontogenesis of cephalic structures from third stage to adult
Tarsius bancanus: region de Kuala Lumpur, Malaisie
Tupaia glis: region de Kuala Lumpur, Malaisie
Tupaia minor: region de Kuala Lumpur, Malaisie
Valanga (exper.)
Oxya (exper.)
Blatella germanica (exper.)
- Tejeraia Diaz-Ungria, 1963, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascaropsinae
key
- Teladorsagia Andreeva et Satubaldin, 1953
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Ostertagiinae
synonymy
- Teladorsagia
Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Teladorsagia
Fudalewicz-Niemczyk, W.; et al., 1976, Acta Zootech., Bratislava (32), 5-19
gastrointestinal helminths, mountain sheep, nilverm and zanil, favorable influence on body weight and wool production of treated animals, no influence of treatment on fertility: Poland
- Teladorsagia circumcincta (Stadelman, 1894)
Drozd, 1965
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Teladorsagia circumcincta
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lams, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Teladorsagia davtiani, illus.
Daskalov, P., 1974, Izvest. Tsentral. Khel-mint. Lab., v. 17, 59-72
Ostertagia circumcincta, O. trifurcata, Teladorsagia davtiani, males crossed with different morphological types of female Ostertagia, no reproductive barriers between them, proposed that they be considered the Ostertagia circumcincta complex

- Teladorsagia davtiani*
Downey, N. E., 1977, *Vet. Rec.*, v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Teladorsagia davtiani* Andreeva et Satubaldin, 1954; *sensu* Becklund et Walker, 1971
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus: southern Bulgaria
- Teladorsagia davtiani*
Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Teladorsagia davtiani* Andreeva y Satubaldin, 1954, *illus.*
Oberger, C.; and Valenzuela, G., 1976, *Bol. Chileno Parasitol.*, v. 31 (3-4), 85-86
morphometric data, differentiation from *Ostertagia trifurcata* by examination of genital cone
Ovis aries (abomasum): Provincia de Valdivia, Chile
- Teladorsagia pinnata* (Daubney, 1933) Drozd, 1965
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Teladorsagia trifurcata* (Ransom, 1907) Drozd, 1965
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Teladorsagia trifurcata*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Tenorastrongylus Durette-Desset*, 1970
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmonellidae, Nippostrongylinae
- Tenorastrongylus josephi* n. sp., *illus.*
Wertheim, G.; and Durette-Desset, M. C., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 735-762
Mus musculus (intestin grele): Nahar Rubin, Dan, Bet Alpha, Caesarea, and Yad Kennedy (Jerusalem), Israel
- Ternidens deminutus*
McConnell, E. E.; et al., 1974, *Onderstepoort J. Vet. Research*, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (small intestine): Kruger National Park, Transvaal
- Ternidens deminutus*
Rogers, S.; and Goldsmid, J. M., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (4), 503-504
Ternidens deminutus, man, indirect fluorescent antibody test evaluated for possible diagnostic use, some cross reactions with *Necator americanus*, promising epidemiologic tool
- Terranova Leiper & Atkinson*, 1914
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Anisakinea
key; synonymy
- Terranova* (*Sauronema*) *Mozgovoi*, 1951
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
as syn. of *Terranova Leiper & Atkinson*, 1914
- Terranova*
Otsuru, M., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Terranova* Type I
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 227-232
incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
Apogonichthys poecilopterus
Euthynnus alletteratus
Kishinoella tonggol
Plectropomus maculatus
Pranesus ogilbyi
Scomberomorus commerson
S. nipponius
all from south-eastern Queensland
- Terranova* Type II
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 227-232
incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
Caranx emburyi
C. leptolepis
Carangoides fulvoguttatus
Carcharinus nasuta
Eulamia spallanzani
Elagatis bipinnulatus
Epinephalus tauvina
Euthynnus alletteratus
Johnius antarctica
J. australis
Lutjanus amabilis
Nemipterus aurifilum
Parapercis nebulosus
Parastromateus niger
Platycephalus indicus
Plectropomus maculatus
Pranesus ogilbyi
Pseudorhombus arsius
Saurida undosquamis
Sciaena dussumieri
Scoliodon jordani
Scomberomorus commerson
S. nipponius
Trichiurus savala
Upeneus tragula
all from south-eastern Queensland

- Terranova sp. (Type I), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Terranova sp. (Type II), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Terranova sp.
Popov, V. N., 1976, *Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR* (145), year 19, (1), 49-53
Histriophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Terranova sp. (Type B), *illus.*
Sakaguchi, Y.; and Katamine, D., 1971, *Nettai Igaku (Trop. Med.)*, v. 13 (4), 159-169
anisakid larvae in marine fishes, prevalence survey, morphometric comparisons
Zeus japonicus: East China Sea
Nemipterus virgatus: South China Sea
Lutjanus sebae: "
Saurida tumbil: "
Carcarrhinus menisorrah: "
Abalistes stellatus: "
Lutjanus basmira: "
Pristipomoides sieboldi: "
Epinephelus septemfasciatus: "
Clidoderma asperrimum: "
Plectorhynchus pictus: "
Caranx equula: "
Lethrinus haematopterus: "
- Terranova sp., *illus.*
Widera, L., 1976, *Med. Wet.*, v. 32 (8), 498-500
Terranova sp., cod fish, histopathology of lesions in meat
- Terranova azarasi (Yamaguti et Arima, 1942)
Deliamure, S. L.; and Popov, V. N., 1975, *Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR* (142), year 18, (10), 7-10
Erignathus barbatus nauticus (stomach): Sakhalin Bay
- Terranova decipiens
Ackman, R. G., 1976, *J. Fish. Research Bd. Canada*, v. 33 (12), 2819-2821
Terranova decipiens, volatile ketones and alcohols from axenic culture medium, from worms from culture and from fish, study techniques
- Terranova decipiens (Krabbe, 1878)
Baeva, O. M., 1968, *Gel'mint. Zhivot. Tikhogo Okeana (Skriabin)*, 80-88
helminth distribution among age groups of Pleurogrammus azonus (musculature): Peter the Great Bay, Sea of Japan
- Terranova decipiens
Bonner, W. N., 1972, *Oceanogr. and Marine Biol. Ann. Rev.*, v. 10, 461-507
role of seals as definitive hosts of Terranova decipiens, larvae of which are found in cod and other commercially prepared seafood
Halichoerus grypus
Phoca vitulina
(stomach of all): all from European waters
- Terranova decipiens (Krabbe, 1873)
Deliamure, S. L.; and Popov, V. N., 1975, *Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR* (142), year 18, (10), 7-10
Erignathus barbatus nauticus (stomach): Sakhalin Bay
- Terranova decipiens (Krabbe, 1878)
Korotaeva, V. D., 1968, *Gel'mint. Zhivot. Tikhogo Okeana (Skriabin)*, 89-96
Enophrys diceraus
Icelus spiniger
Myoxocephalus jaok
M. brandti
(muscles of all): all from Sea of Japan
- Terranova decipiens
McClelland, G., 1976, *Tr. Am. Micr. Soc.*, v. 95 (2), 265 [Abstract]
Terranova decipiens in Phoca vitulina and Halichoerus grypus (both exper.) (stomach of both), course of infection and pathology
- Terranova scoliodontis
Gibson, D. I.; and Taylor, A. L., 1976, *Parasitology*, v. 73 (2), v [Abstract]
Ascaridoidea, excretory system, comment upon taxonomic significance and function
- Tetanonema Steiner, 1937
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)* (3), 1-27
incertae sedis
- Tetrabothriostrogylus Mawson, 1960
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Amidostomatidae, Mackerrastrongylineae
- Tetrabothriostrogylus mackerrasae Mawson
Beveridge, I.; and Barker, I. K., 1976, *Austral. J. Zool.*, v. 24 (2), 265-272
helminths and arthropods, Antechinus stuartii, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (intestine): Powelltown, Victoria
- Tetragomphius sp., *illus.*
Ashizawa, H.; et al., 1976, *Bull. Fac. Agric. Univ. Miyazaki*, v. 23 (2), 371-381
Tetragomphius sp. in Meles meles anakuma, pathologic changes in pancreatic duct: Miyazaki Prefecture; Oita Prefecture
- Tetragomphius sp., *illus.*
Ashizawa, H.; et al., 1976, *Bull. Fac. Agric. Univ. Miyazaki*, v. 23 (2), 383-393
Tetragomphius sp. in Martes melampus melampus (pancreatic duct), mixed infection with small flukes (probably Concinnum ten), pathological changes: Miyazaki Prefecture

- Tetrameres Creplin, 1846
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Tetramerinae
key
includes subgens.: Tetrameres; Microtetrameres
- Tetrameres, *illus.*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
subgen. of Tetrameres
key
- Tetrameres (Gynaecophila)
Pence, D. B.; Mollhagen, T.; and Forrester, D. J., 1975, J. Parasitol., v. 61 (3), 494-498
"Pending examination of *T. (G.) gynaecophila* (the subgenus type) and the results of studies under way ... it is deemed necessary at present to view the subgeneric arrangement of the genus *Tetrameres* with suspicion. ... a subgenus that was originally perceived as lacking cuticular spines cannot continue to be regarded as characteristically having them."
- Tetrameres
Pence, D. B.; Mollhagen, T.; and Forrester, D. J., 1975, J. Parasitol., v. 61 (3), 494-498
subgenus of *Tetrameres*
subgeneric status discussed, probable revision indicated
- Tetrameres, subgen.
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
"Herewith placed in the subgenus *Tetrameres* are *Tetrameres cordoniferens* Rasheed 1960; *T. cygni* Ryjnikov and Kozlov 1960; and *T. pahnicopterus* [sic] Ali 1970."
- Tetrameres sp.
Ellis, C. J.; and Calderwood, G., 1977, Proc. Iowa Acad. Sc., v. 84 (1), 30-31
Porzana carolina
Seiurus aurocapillus
(proventriculi of all): all from Iowa
- Tetrameres sp.
Keppner, E. J., 1973, Tr. Am. Micr. Soc., v. 92 (2), 288-291
Larus californicus (proventriculus): city dump of Laramie, Wyoming
- Tetrameres sp.
Sergeeva, T. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 146-155
Larus argentatus
Sterna paradisea
Stercorarius longicaudatus
all from Yenisei
- Tetrameres americana
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (proventriculus): Ottawa county, Ohio
- Tetrameres (Petrowimeres) australis* Johnston and Mawson 1941
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Petrowimeres) biziurae* Johnston and Mawson 1941
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Tetrameres) cardinalis* sp. n., *illus.*
Quentin, J. C.; and Barre, N., 1976, Ann. Parasitol., v. 51 (1), 65-81
Tetrameres cardinalis n. sp., life cycle, development
Richmondia cardinalis (nat. and exper.)
(paroi du proventricule): Mexique
Locusta migratoria (exper.) (tissu adipeux)
- Tetrameres coloradensis* Schmidt, 1962
Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 343
"It remains possible that the type specimen of *T. dubia* is aberrant, lacking body spines, and postanal papillae normal for the species, but until this can be proved it seems justified to retain *T. coloradensis* as a valid species."
- Tetrameres (Tetrameres) cordoniferens* Rasheed 1960
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres crami* Swales, 1933
Buck, O. D.; Cooper, C. L.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 233-234
Larus argentatus: Bass Island region of Lake Erie
- Tetrameres (Petrowimeres) crami* Swales 1933
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Tetrameres) cygni* Ryjnikov and Kozlov 1960
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres dubia* Travassos, 1917
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Pluvialis apricaria altifrons
Gallinago media
Philomachus pugnax
Gallinago gallinago
Heteroscelus incanus brevipes
Tringa glareola
Calidris minuta
all from lower Yenisei [and/or] Keta lake

- Tetrameres dubia* Travassos, 1917
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63 (2), 343
"the description by Travassos is accurate, including measurements. It remains possible that the type specimen of *T. dubia* is aberrant, lacking body spines, and postanal papillae normal for the species, but until this can be proved it seems justified to retain *T. coloradensis* as a valid species."
- Tetrameres fissispina* (Diesing, 1861)
Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (1), 7-31
nematodes of *Larus canus* (proventriculus), age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Tetrameres fissispina* (Diesing, 1861)
Fabiyyi, J. P., 1972, *Bull. Epizoot. Dis. Africa*, v. 20 (3), 229-234
survey of helminths of chickens, comparison of techniques of management (native extensive, deep-litter (intensive) and semi-intensive systems) on worm burden; suggested preventive measures and treatment with piperazine: Vom area, Benue-Plateau State, Nigeria
- Tetrameres fissispina* Diesing, 1861
Fabiyyi, J. P., 1972, *Bull. Epizoot. Dis. Africa*, v. 20 (3), 235-238
Numida meleagris galeata (proventriculus): Vom area, Benue Plateau State, Nigeria
- Tetrameres fissispina* Diesing, 1861
Kamburov, P.; and Vasilev, I., 1972, *Izvest. Tsentral. Khelmint. Lab.*, v. 15, 109-133
Anser anser
Anas platyrhynchos
A. clypeata
A. acuta
A. crecca
A. querquedula
Aythya nyroca
(crop of all): all from Bulgaria
- Tetrameres fissispina*, illus.
Orlandi, M.; and Colombani, B., 1975, *Ann. Fac. Med. Vet. Pisa*, v. 27, 1974, 113-128
histopathology
Anas querquedula (proventriculus): Tombolo (Pisa)
- Tetrameres* (Petrowimeres) *fissispina* Diesing 1861
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829
- Tetrameres* (Petrowimeres) *galericulatus* Oschmarin 1956
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829
- Tetrameres globosa* (Linstow, 1879)
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, *Am. Midland Naturalist*, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans
Porphyryla martinica
(proventriculus of all): all from Florida
- Tetrameres grusi*
Forrester, D. J.; et al., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 55-59
Grus canadensis tabida (proventriculus): Florida
- Tetrameres* (T.) *gubanovi* Shigin, 1957, illus.
Aleksiev, V. M.; and Smetanina, Z. B., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 97-104
redescription
Podiceps griseigena (proventriculus): Rimsko-Korsakov islands
- Tetrameres* (Gynaecophila) *gynaecophila*
Pence, D. B.; Mollhagen, T.; and Forrester, D. J., 1975, *J. Parasitol.*, v. 61 (3), 494-498
"Pending examination of *T. (G.) gynaecophila* (the subgenus type) and the results of studies under way ... it is deemed necessary at present to view the subgeneric arrangement of the genus *Tetrameres* with suspicion. ... a subgenus that was originally perceived as lacking cuticular spines cannot continue to be regarded as characteristically having them."
- Tetrameres* (Petrowimeres) *indiana* Ali 1970
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829
- Tetrameres inerme*
Courtney, C. H.; and Forrester, D. J., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (proventriculus): Florida
- Tetrameres* (Petrowimeres) *mohtedai* Bhalerao and Rao 1944
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829
- Tetrameres* (Petrowimeres) *nettatus* Ali 1970
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829
- Tetrameres paraaraliensis* Oschmarin, 1956
Bondarenko, S. K., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 35-45
Eudromias morinellus: Keta lake
- Tetrameres* (Petrowimeres) *pavonis* Chertkova 1953
Pence, D. B.; Mollhagen, T.; and Prestwood, A. K., 1975, *J. Parasitol.*, v. 61 (5), 825-829

- Tetrameres (Tetrameres) phaenicopterus Ali 1970
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Petrowimeres) plectropteri Thwaite
1926
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Petrowimeres) rijikovi Chuan 1961
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres ryjikovi Khuan Shen-i, 1961
Turner, B. C.; and Threlfall, W., 1975, Proc.
Helminth. Soc. Washington, v. 42 (2), 157-169
parasites of *Anas crecca* and *A. discors*,
incidence and intensity, age and sex of host
Anas crecca
A. discors
(mucous glands of proventriculus of all):
all from eastern Canada
- Tetrameres (T.) skrjabini Panowa, 1926
Alekseev, V. M.; and Smetanina, Z. B., 1968,
Gel'mint. Zhivot. Tikhogo Okeana (Skriabin),
97-104
Larus crassirostris: Rimsko-Korsakov islands
- Tetrameres skrjabini
Belogurov, O. I.; Leonov, V. A.; and Zueva,
L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana
(Skriabin), 105-124
Larus argentatus
L. crassirostris
Sterna hirundo
Stercorarius longicaudus
S. parasiticus
(proventriculus of all): all from coast of
Sea of Okhotsk
- Tetrameres somateriae (Ryzhikov, 1963)
Bishop, C. A.; and Threlfall, W., 1974, Proc.
Helminth. Soc. Washington, v. 41 (1), 25-35
Somateria mollissima: insular Newfoundland
and/or southern Labrador
- Tetrameres (Petrowimeres) somateriae Ryjikov
1963
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Petrowimeres) striatus Oschmarin
1956
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
- Tetrameres (Gynaecophila) strigiphila sp. n.,
illus.
Pence, D. B.; Mollhagen, T.; and Forrester, D.
J., 1975, J. Parasitol., v. 61 (3), 494-498
Strix varia georgica (encysted in a fibrous
capsule in the tunica muscularis of proven-
tricularis): near Ocala, Florida
- Tetrameres (Tetrameres) tinamicola sp. n., illus.
Pence, D. B.; Mollhagen, T.; and Prestwood, A.
K., 1975, J. Parasitol., v. 61 (5), 825-829
Eudromia elegans elegans
E. elegans albida
(proventriculus of all): all from Argentina
- Tetrameres williamsi
Bush, A. O.; and Forrester, D. J., 1976, Proc.
Helminth. Soc. Washington, v. 43 (1), 17-23
Eudocimus albus (proventriculus): Florida
- Tetrameridae Travassos, 1914
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Habronematoidea
key; key to subfams.
includes: Tetramerinae; Geopetitiinae;
Crassicaudinae
- Tetramerinae (Travassos, 1914, fam.)
Chabaud, A. G., 1975, CIH Keys Nematode Para-
sites Vertebrates (Anderson, Chabaud, and
Willmott) (3), 29-58
Tetrameridae
key; key to genera
includes: Microhadjelia; Tetrameres
- Tetrapetalonema Faust, 1935
Chabaud, A. G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
key
includes subgenera: Sandnema n. subgen.;
Tetrapetalonema; Esslingeria n. subgen.
- Tetrapetalonema [? n. rank]
Chabaud, A. G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
subgen. of Tetrapetalonema
key
tod: T. (T.) marmosetae Faust, 1935
- Tetrapetalonema (Tetrapetalonema) atelensis
McCoy, 1936
Chabaud, A. G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Tetrapetalonema atelensis, illus.
Chalifoux, L. V.; et al., 1973, Lab. Animal
Sc., v. 23 (2), 211-220
differentiation of 11 types of circulating
microfilariae in blood smears from 7 spp. of
New World monkeys based on differences in
histochemical localization of acid phosphatase
Ateles geoffroyi: New England Regional Pri-
mate Research Center
- Tetrapetalonema (Tetrapetalonema) barbascalensis
(Esslinger et Gardiner, 1974), illus.
Chabaud, A. G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397
- Tetrapetalonema (Sandnema) digitata (Chandler,
1929) (tod of subgen.)
Chabaud, A. G.; and Bain, O., 1976, Ann. Para-
sitol., v. 51 (3), 365-397

- Tetrapetalonema (*Tetrapetalonema*) *dunni* Mullin et Orihel, 1972, *illus.*
Chabaud, A.-G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Tetrapetalonema*) *interstitium* (Price, 1962)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Esslingeria*) *leopoldi* (Berghe, Chardome et Peel, 1964)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Tetrapetalonema*) *llewellyni* (Price, 1962)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Tetrapetalonema*) *marmosetae* Faust, 1935 (tod of subgen.)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema *marmosetae*, *illus.*
Chalifoux, L. V.; et al., 1973, *Lab. Animal Sc.*, v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saimiri sciureus
Saguinus oedipus
S. tamarinus
all from New England Regional Primate Research Center
- Tetrapetalonema *marmosetae*, *illus.*
Dreizen, S.; Smith, W. N.; and Levy, B. M., 1970, *Oral Surg.*, v. 30 (4), 527-532
Tetrapetalonema *marmosetae*, histopathologic study of infection in Saguinus oedipus (blood vessels and connective tissues of lips, tongue, periodontium, dental pulp, salivary glands) kept as research colony by University of Texas Dental Science Institute
- Tetrapetalonema (*Tetrapetalonema*) *nicollei* (?) (Mazza, 1930)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Tetrapetalonema*) *obtusa* (McCoy, 1936)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema *panamensis*, tentative identification, *illus.*
Chalifoux, L. V.; et al., 1973, *Lab. Animal Sc.*, v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saimiri sciureus
Cebus albifrons
C. apella
Saguinus tamarinus
S. oedipus
Aotus trivirgatus
all from New England Regional Primate Research Center
- Tetrapetalonema *papionis*, *illus.*
McConnell, E. E.; et al., 1974, *Onderstepoort J. Vet. Research*, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (skin and skeletal muscle fascia): Kruger National Park, Transvaal
- Tetrapetalonema (*Tetrapetalonema*) *parvum* McCoy, 1936, *illus.*
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Esslingeria*) *perstans* (Manson, 1891), *illus.* (tod of subgen.)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Esslingeria*) *rodhaini* (Peel et Chardome, 1947)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Esslingeria*) *streptocerca* (Macfie et Corson, 1922)
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Sandnema*) *sunci* Sandground, 1933
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema (*Tetrapetalonema*) *tamarinae* Dunn et Lambrecht, 1963
Chabaud, A. G.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (3), 365-397
- Tetrapetalonema *tamarinae*, *illus.*
Chalifoux, L. V.; et al., 1973, *Lab. Animal Sc.*, v. 23 (2), 211-220
differentiation of 11 types of circulating microfilariae in blood smears from 7 spp. of New World monkeys based on differences in histochemical localization of acid phosphatase
Saguinus tamarinus: New England Regional Primate Research Center

- Tetrapetalonema* (Esslingeria) vanhoofi (Peel et Chardome, 1946)
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Tetrapetalonema* (Tetrapetalonema) zakii (Nagaty, 1935) Sandground, 1958
Chabaud, A. G.; and Bain, O., 1976, Ann. Parasitol., v. 51 (3), 365-397
- Toxicospirura* M. B. Chitwood & Cordero del Campillo, 1966, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
Ascarosinae
key
- Thalassonema* Ward, 1933
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
Marineremithidae fam. n.
- Thaparina macrocephala* n. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. 5. (389), Zool. (271), 751-768
Gopherus sp., "vraissementablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraissementablement" G. agassizii: desert de Sonora, Mexique (colon of all)
- Thaparina microcephala* n. sp., illus.
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. 5. (389), Zool. (271), 751-768
Gopherus sp., "vraissementablement" G. flavo-marginatus: desert de Coahuila, Mexique
Gopherus sp., "vraissementablement" G. agassizii: desert de Sonora, Mexique (colon of all)
- Thaprosipirura* Sood & Parshad, 1974
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Protospirura* Seurat, 1914
- Thaprosipirura meltadi* Sood et Parshad 1974
Sood, M. L.; and Parshad, V. R., 1975, Riv. Parasitol., Roma, v. 36 (2-3), 189-196
infections in *Milgardia meltada*, survey of seasonal distribution, possible correlations between host diet and sex and incidence of infection
- Thelastoma* sp.
Hornick, W. M., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (5), 385 [Demonstration]
The lastoma sp., bacterial infection of cuticle of pinworms inhabiting hindgut of laboratory reared *Periplaneta americana*, bacterial preference for *Thelastoma* sp. over *Hammerschmidtella dieisingi* possibly related to structure or cuticle
- Thelastoma pachyulii* (Parona, 1896) Travassos, 1925
Hristovski, N. D., 1972, Acta Parasitol. Iugoslavica, v. 5 (2), 109-115
Julidae gen. et sp.: Jugoslavia (Skopje; Split); Grcija (Lerin)
- Thelastoma pachyulii*
Hristovski, N. D., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 87-91
Julus sp.: Macedonia, Yugoslavia
- Thelastoma singaporensis* Leong, 1965
Leong, L.; and Paran, T. P., 1966, Med. J. Malaya, v. 20 (6), 349
Periplaneta americana
Blattella orientalis
all from Singapore
- Thelastomatidae* [sp.]
Carter, J. B., 1976, J. Applied Ecol., v. 13 (1), 103-122
Tipula paludosa
Tipula luna
T. unca
all from north-east England
- Thelastomoides*
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. 5. (389), Zool. (271), 751-768
as syn. of *Alaeuris*
- Thelastomoides longicollis* Walton, 1927
Petter, A. J.; and Douglass, J. F., 1976, Bull. Mus. National Hist. Nat., Paris, 3. 5. (389), Zool. (271), 751-768
as syn. of *Alaeuris longicollis* (Walton, 1927)
- Thelazia* Bosc, 1819
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelaziinae
key
Syn.: *Annulofilaria* G. N. Hsu, 1957
includes subgen.: *Thelaziella*; *Thelazia*
- Thelazia*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Thelazia*
key
- Thelazia* sp.
Frechette, J. L.; Marcoux, M.; and St.-Pierre H., 1976, Canad. Vet. J., v. 17 (4), 114
Thelazia lacrymalis in horses operated on for cataracts; *Thelazia* sp., keratoconjunctivitis in dairy cows; Quebec
- Thelazia* sp.
Khamboonruang, C.; and Saeng-Udom, C., 1971, Southeast Asian J. Trop. Med. and Pub. Health v. 2 (4), 588 [Demonstration]
Thelazia sp., Thai boy presenting with symptoms of conjunctivitis, worm discovered in purulent discharge from eye: Chiangmai Province, Thailand
- Thelazia* spp. larvae, one resembling *T. lacrymalis*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, J. Parasitol., v. 62 (5), 877-880
Musca autumnalis: Kentucky

- Thelazia* [sp.]
Miyamoto, K.; Shinonaga, S.; and Kano, R., 1975, Eisei Dobutsu (Japan. J. San. Zool.), v. 26 (4), 203-206
Musca hervei: Kobayashi City, Miyazaki Prefecture
- Thelazia gulosa*, illus.
Arbuckle, J. B. R.; and Khalil, L. F., 1976, Vet. Rec., v. 99 (19), 376-377
cattle (eyes): abattoir, Surrey, Great Britain
- Thelazia gulosa*
Arora, G. S.; et al., 1977, Indian J. Animal Sc., v. 45 (12), 1975, 953-957
Thelazia skrjabini, *T. rhodesii*, and *T. gulosa*, buffaloes (eyeballs), incidence and intensity of infection, seasonal fluctuations: [India]
- Thelazia gulosa* Railliet and Henry 1910, illus.
Lyons, E. T.; and Drudge, J. H., 1975, J. Parasitol., v. 61 (6), 1119-1122
Thelazia gulosa, *T. skrjabini*, single and mixed infections, cattle (eyes), measurements: near Georgetown, Kentucky
- Thelazia gulosa*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Musca sp.: Uzbekistan
- Thelazia lacrymalis* (Gurll, 1831)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (conjunctival sac): Kruger National Park
- Thelazia lacrymalis*
Frechette, J. L.; Marcoux, M.; and St.-Pierre, H., 1976, Canad. Vet. J., v. 17 (4), 114
Thelazia lacrymalis in horses operated on for cataracts; *Thelazia* sp., keratoconjunctivitis in dairy cows: Quebec
- Thelazia lacrymalis* (Gurll 1831) Railliet and Henry 1910, illus.
Lyons, E. T.; and Drudge, J. H., 1975, J. Parasitol., v. 61 (6), 1122-1124
Thelazia lacrymalis, horses (surface of eyeball, under upper and lower eyelids, under conjunctiva, under nictitating membrane, from lacrymal gland and its excretory ducts), measurements: near Lexington, Kentucky
- Thelazia lacrymalis*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, J. Parasitol., v. 62 (6), 877-880
horses (nat. and exper.): Kentucky
Musca autumnalis (exper.)
- Thelazia rhodesii*
Arora, G. S.; et al., 1977, Indian J. Animal Sc., v. 45 (12), 1975, 953-957
Thelazia skrjabini, *T. rhodesii*, and *T. gulosa*, buffaloes (eyeballs), incidence and intensity of infection, seasonal fluctuations: [India]
- Thelazia rhodesii* (Desmarest, 1828)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (conjunctival sac): Kruger National Park
- Thelazia rhodesii*
Ecimovic, T. J., 1973, Bull. Epizoot. Dis. Africa, v. 21 (2), 129-131
Thelazia rhodesii, incidence, cattle, conjunctivitis parasitaria: Mbeya region of Tanzania
- Thelazia rhodesii*
Michalski, L., 1976, Med. Wet., v. 32 (7), 417-419
Thelazia rhodesii, cattle, levamisole injected into conjunctival sac highly efficient, tetramisole applied orally less effective
- Thelazia rhodesii*
Miyamoto, K.; Shinonaga, S.; and Kano, R., 1975, Eisei Dobutsu (Japan. J. San. Zool.), v. 26 (4), 203-206
cows (eyes): Kobayashi City, Miyazaki Prefecture
- Thelazia rhodesii*
Shinonaga, S.; et al., 1974, J. Med. Entom., v. 11 (5), 595-598
Thelazia rhodesii, *T. skrjabini*, seasonal fluctuation in cattle closely correlated with numbers of *Musca hervei* infesting cattle and natural infection rate of *Thelazia* spp. in *M. hervei* cattle (eyes)
Musca hervei (wall of body cavity, malpighian tubules, fat body tissues)
all from Chiba city, Japan
- Thelazia rhodesii* (Demarest, 1828)
Somasundaram, M.; and Rajamanickam, C., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (3), 453 [Demonstration]
Thelazia rhodesii, eye worm infection in cattle, clinical aspects, treatment with boric acid solution and antibiotic: Peninsular Malaysia
- Thelazia rhodesii*
Sultanov, M. A.; and Kabilov, T., 1976, Dokl. Akad. Nauk UzSSR (11), 57-58
Musca domestica
M. larvipara
all from Uzbekistan
- Thelazia rhodesii*
Vassiliades, G.; et al., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (3), 315-317
Thelazia rhodesii, bovine, treatment with morantel tartrate, levamisole, tetramisole and levamisole chlorhydrate, all effective; lugol, ineffective: Casamance (Senegal)
- Thelazia skrjabini*
Arbuckle, J. B. R.; and Khalil, L. F., 1976, Vet. Rec., v. 99 (19), 376-377
cattle (eyes): abattoir, Surrey, Great Britain

- Thelazia skrjabini*
Arora, G. S.; et al., 1977, Indian J. Animal Sc., v. 45 (12), 1975, 953-957
Thelazia skrjabini, *T. rhodesii*, and *T. gulosa*, buffaloes (eyeballs), incidence and intensity of infection, seasonal fluctuations: [India]
- Thelazia skrjabini*
Chauhan, P. P. S.; et al., 1976, Indian J. Animal Sc., v. 46 (3), 152-153
buffalo (vitreous humour, lacrimal duct of eye)
- Thelazia skrjabini* Ershov 1928, illus.
Lyons, E. T.; and Drudge, J. H., 1975, J. Parasitol., v. 61 (6), 1119-1122
Thelazia gulosa, *T. skrjabini*, single and mixed infections, cattle (eyes), measurements: near Georgetown, Kentucky
- Thelazia skrjabini*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, J. Parasitol., v. 62 (6), 877-880
horses (nat. and exper.): Kentucky
Musca autumnalis: Kentucky
- Thelazia skrjabini*
Miyamoto, K.; Shinonaga, S.; and Kano, R., 1975, Eisei Dobutsu (Japan. J. San. Zool.), v. 26 (4), 203-206
cows (eyes): Kobayashi City, Miyazaki Prefecture
- Thelazia skrjabini*
Shinonaga, S.; et al., 1974, J. Med. Entom., v. 11 (5), 595-598
Thelazia rhodesii, *T. skrjabini*, seasonal fluctuation in cattle closely correlated with numbers of *Musca hervei* infesting cattle and natural infection rate of *Thelazia* spp. in *M. hervei* cattle (eyes)
Musca hervei (wall of body cavity, malpighian tubules, fat body tissues)
all from Chiba city, Japan
- Thelaziella*, illus.
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
subgen. of *Thelazia*
key
- Thelaziidae* Skrjabin, 1915
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelazioidae
key; key to subfamilies
includes: *Oxyspirurinae*; *Thelaziinae*
- Thelaziinae* (Skrjabin, 1915, fam.) Baylis & Daubney, 1926
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Thelaziidae
key; key to genera
includes: *Hempelia*; *Ceratospira*; *Thelazia*
- Thelazioidae*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (1), 6-17
Spirurina
key
- Thelazioidae*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
Spirurina
key to families
includes: *Pneumospiruridae*; *Thelaziidae*; *Rhabdochoniidae*
- Thominx aerophilus* (Creplin, 1839)
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Vulpes vulpes: Pechora river basin
- Thominx aerophilus* (Creplin, 1839) Skrjabin et Schikhobalova, 1954
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Martes martes (trachea, bronchi): Karelia
- Thominx anatis* (Schrank, 1790) Skrjabin et Schikhobalova, 1954
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anser erythropus
Anas platyrhynchos
A. penelope
A. querquedula
Aythya ferina
A. nyroca
Netta rufina
Anas crecca
(caecum of all): all from Bulgaria
- Thominx collaris*
Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix: CSSR
- Thominx contorta* (Creplin, 1839)
Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
synonymy, nematodes of *Larus canus* (esophagus, proventriculus, ventriculus, intestine), age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Thominx contorta* (Creplin, 1899) Travassos, 1915
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmint. Lab., v. 15, 109-133
Anas platyrhynchos
A. clypeata
A. acuta
A. crecca
A. querquedula
Aythya nyroca
(esophagus of all): all from Bulgaria
- Thominx limicolae* Gubanov et Mamaev, 1959
Bondarenko, S. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 35-45
Xenus cinereus: Keta lake

- Thominx neopulchra (Babos, 1954) Skrjabin et Schikhobalova, 1954
 Skvortsov, V. G., 1973, Parazity Zhivot. i Rasten., Akad. Nauk Moldavsk. SSR (9), 92-155
 ecological analysis of bat helminth fauna, geographic distribution
 Myotis oxygnathus
 M. dasycneme
 M. daubentoni
 M. bechsteini
 M. mystacinus
 all from Moldavia
- Thominx perforans (Kotlan et Orosz, 1931) Skrjabin et Schikhobalova, 1954
 Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmit. Lab., v. 15, 109-133
 Anas acuta
 A. platyrhynchos
 (esophagus of all): all from Bulgaria
- Thominx railletii
 Bakke, T. A.; and Barus, V., 1976, Norwegian J. Zool., v. 24 (1), 7-31
 as syn. of Thominx contorta (Creplin, 1839)
- Thominx skrjabini Lubimova, 1947, illus.
 Daiia, G. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 49-53
 redescription
 Anas acuta
 A. penelope
 (caecum of all): all from Yakut ASSR
- Thominx wavilovoi
 Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
 Tetrao urogallus: CSSR
- Thubunaea Seurat, 1914, illus.
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Thubunaeinae
 key
- Thubunaea iguanae
 Pearce, R. C.; and Tanner, W. W., 1973, Great Basin Nat., v. 33 (1), 1-18
 Sceloporus magister (stomach): Great Basin and Upper Colorado Plateau, Utah
- Thubunaeinae (Sobolev, 1949, tribe)
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Physalopteridae
 key; key to genera
 includes: Thubunaea; Physalopteroides
- Thwaitia Rasheed, 1963, illus.
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Philometrinae
 key
- Thwaitia Rasheed, 1963
 Molnar, K.; and Fernando, C. H., 1975, J. Helminth., v. 49 (2), 101-105
 as syn. of Philometra Costa, 1945
- Thwaitia abdominalis (Nybelin, 1928) Rasheed, 1963
 Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmit. Lab., v. 16, 87-110
 Syn.: Philometra abdominalis Nybelin, 1928
 G[obio] gobio (body cavity): Balkan Mountain river(s)
- Thylaconema Chandler (1929)
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 "valid but the head structure is not well known, especially in en face view. Perhaps it is related to Ceratospira."
- Thynnascaris Dollfus, 1933
 Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
 Raphidascaridinea
 key; synonymy
- Thynnascaris Dollfus, 1933
 Kalyankar, S. D., 1972, Riv. Parassitol., Roma, v. 33 (3), 203-208
 valid genus
- Thynnascaris
 Soleim, O., 1976, Norwegian J. Zool., v. 24 (4), 464 [Abstract]
 "It is concluded that the genus Thynnascaris should be maintained and that Phocascaris become a synonym of Contracaecum."
- Thynnascaris Type I
 Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 227-232
 incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
 Amentum devisi
 Cynoglossus bilineatus
 Gerres ovatus
 Mylio australis
 Paraplotosus albilabris
 Platycephalus indicus
 Polynemus sheridani
 Saurida undosquamis
 Sphaeroides hamiltoni
 Synaptura orientalis
 Tachysurus australis
 Tragulichthys jaculiferus
 Triacanthus biaculeatus
 all from south-eastern Queensland
- Thynnascaris Type II
 Cannon, L. R. G., 1977, Internat. J. Parasitol., v. 7 (3), 227-232
 incidence, intensity, host diet, habitat; ecological relationships of larval ascaridoids from marine fishes
 Choerodon venustus
 Plectorhynchus chrysotaenia
 all from south-eastern Queensland

- Thynnascaris* Type III
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 227-232
incidence, intensity, host diet, habitat;
ecological relationships of larval ascari-
doids from marine fishes
Nemipterus aurifilum
Parapercis nebulosus
Parastromateus niger
Pseudorhombus arsius
Sciaena dussumieri
all from south-eastern Queensland
- Thynnascaris* Type IV
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 227-232
incidence, intensity, host diet, habitat;
ecological relationships of larval ascari-
doids from marine fishes
Johnius australis
Lutjanus amabilis
Nemipterus aurifilum
Pranesus ogilbyi
Pseudorhombus arsius
Sciaena dussumieri
Scoliodon jordani
all from south-eastern Queensland
- Thynnascaris* sp. (Type I), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Thynnascaris* sp. (Type II), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Thynnascaris* sp. (Type III), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Thynnascaris* sp. (Type IV), *illus.*
Cannon, L. R. G., 1977, *Internat. J. Parasitol.*, v. 7 (3), 233-243
description, key
- Thynnascaris* [sp.] (type MB of Norris and Overstreet, 1976)
Ebert, D. J.; and Norris, D. E., 1976, *J. Mississippi Acad. Sc.*, Suppl., v. 21, 70
[Abstract]
Thynnascaris (type MB of Norris and Overstreet, 1976) larvae in *Trichiurus lepturus* (mesentery), migration and localization in white mice (exper.)
- Thynnascaris* sp.
Munson, D. A., 1974, *J. Wildlife Dis.*, v. 10 (3), 256-262
Liparis atlanticus (mesenteries): Rye, New Hampshire
- Thynnascaris adunca* (Rudolphi, 1802), *illus.*
Bakke, T. A.; and Barus, V., 1975, *Norwegian J. Zool.*, v. 23 (3), 183-191
measurements
Larus canus (alimentary canal): Agdenes area, Norway
- Thynnascaris adunca*
Bakke, T. A.; and Barus, V., 1976, *Norwegian J. Zool.*, v. 24 (1), 7-31
nematodes of *Larus canus*, age and sex of host, seasonal variations, distribution in alimentary canal: Agdenes, Norway
- Thynnascaris aduncum*
McVicar, A. H., 1977, *J. Helminth.*, v. 51 (1), 11-21
intestinal helminths of *Raja naevus*, incidence, intensity, pattern of infection with host age and sex, geographical differences in composition of parasite burden
Raja naevus (stomach, spiral intestine): Loch Ewe; off Aberdeen; off Plymouth
- Thynnascaris adunca* (Rud. 1802)
Soleim, Ø., 1976, *Norwegian J. Zool.*, v. 24 (4), 319-323
Thynnascaris adunca from *Gadus morhua*, comparison of 2 populations, relative age of parasites indicates that cod in warmer Norwegian coastal waters is subject to loss of parasites and re-infection more often than colder Barents Sea cod
Syn.: *Contraceacum aduncum* (Rud.)
Gadus morhua: Barents Sea; Bergen fish-market, Norway
- Thynnascaris adunca* (Rud. 1802)
Soleim, O., 1976, *Norwegian J. Zool.*, v. 24 (4), 464 [Abstract]
study of morphology using scanning electron and light microscopy
- Thynnascaris inquires* (Linton, 1901) Rasheed, 1965, *illus.*
Kalyankar, S. D., 1972, *Riv. Parassitol.*, Roma, v. 33 (3), 203-208
description of larval forms
Elacate niger (stomach)
sea-crabs (gills)
all from Ratnagiri (Maharashtra: India)
- Thynnascaris serrani* n. sp., *illus.*
Kalyankar, S. D., 1972, *Riv. Parassitol.*, Roma, v. 33 (3), 203-208
Serranus ferio (intestine): Malvan, Maharashtra, India
- Torquatella* Yorke & Maplestone, 1926 (preoccupied)
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Torquatoides* (Williams, 1929, subgen.) Inglis, 1965
- Torquatella* York and Maplestone, 1926
Pence, D. B.; and Casto, S., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 24-28
"Torquatoides is used as the only available replacement for *Torquatella* York and Maplestone, 1926."
- Torquatoides* (Williams, 1929, subgen.) Inglis, 1965, *illus.*
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
Histiocephalinae
key; synonymy

- Torquatoides Williams, 1929
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
synonymy
"only available replacement for Torquatella York and Maplestone, 1926."
- Torquatoides balanocephala (Gendré, 1922) [n. comb.]
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
- Torquatoides conocephala Molin (1860) [n. comb.]
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
- Torquatoides crotophaga Williams, 1929 [n. comb.], illus.
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
redescription
Geococcyx californianus
Crotophaga sulcirostris
(under koilon of gizzard of all): all from Millett, LaSalle County, Texas
- Torquatoides torquata (Gendré, 1922) [n. comb.]
Pence, D. B.; and Casto, S., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 24-28
- Torrestrongylus Vigueras, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Torynurus Baylis and Daubney 1925
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
key; key to species; diagnosis, differentiation from Pharurus
- Torynurus alatus (Leuckart) Delyamure 1952
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of Pharurus alatus (Leuckart 1848)
Stiles and Hassall 1905
- Torynurus bicostatus (von Linstow) Schmidt-Ries 1939
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
as syn. of Torynurus convolutus (Kuhn 1829)
Baylis and Daubney 1925
- Torynurus convolutus (Kuhn 1829) Baylis and Daubney 1925, illus.
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
synonymy; redescription; key
Phocoena phocoena: Bay of Fundy, Canada; Vancouver Is., British Columbia; Canada; North Sea off Netherlands; NE Atlantic
- Torynurus dalli (Yamaguti 1951) Delyamure 1972
Arnold, P. W.; and Gaskin, D. E., 1975, Canad. J. Zool., v. 53 (6), 713-735
synonymy; redescription; key
Phocoenoides dalli: off Long Beach, California
- Toxascaris Leiper, 1907
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascaridinae
key
- Toxascaris spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Toxascaris leonina
El-Moukdad, A. R., 1977, Wien Tierarztl. Monatschr., v. 64 (3), 85-91
effect of disinfectants on eggs of Ascaris suum, Toxascaris leonina, small horse strongyles and coccidia oocysts
- Toxascaris leonina (Linstow 1902)
Gilbertson, D. E., 1977, J. Parasitol., v. 63 (1), 162-163
Vulpes fulva (intestine): Dakota County, Minnesota
- Toxascaris leonina
Girdwood, R. W. A.; et al., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 284
[Demonstration]
Toxocara canis, Toxascaris leonina, incidence in dogs and in soil samples from public places: Glasgow, Scotland
- Toxascaris leonina
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Toxascaris leonina
Guterbock, W. M.; and Levine, N. D., 1977, J. Am. Vet. Med. Ass., v. 170 (12), 1411-1413
cats (feces): east central Illinois
- Toxascaris leonina
Hass, D. K.; and Collins, J. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 135-137
helminths, dogs, comparative efficacy of vinclofos, ticarbodine, mebendazole
- Toxascaris leonina (Linstow, 1902)
Hinaidy, H. K., 1976, Zentralbl. Vet.-Med., Reihe B, v. 23 (1), 66-73
Vulpes vulpes: Osterreich
- Toxascaris leonina
Hogarth-Scott, R. S., 1967, Internat. Arch. Allergy and Applied Immunol., v. 32 (2), 201-207
Toxocara canis, T. cati, Toxascaris leonina, Ascaris suum, rabbits (exper.), presence of reagin-like antibodies demonstrable by homologous passive cutaneous anaphylaxis, responsible allergens were common to all 4 nematode species

- Toxascaris leonina*
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, *Vet. Rec.*, v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects
- Toxascaris leonina* Linstow, 1902
Kozlov, D. P., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 71-78
Canis familiaris
Vulpes vulpes
Alopex lagopus
all from Pechora river basin
- Toxascaris leonina*
McCurdy, H. D.; and Guerrero, J., 1977, *Vet. Med. and Small Animal Clin.*, v. 72 (11), 1731-1733
helminths, dogs, mebendazole powder, controlled critical studies, good results against all helminths except *Dipylidium caninum*: Kansas; New Jersey; Texas
- Toxascaris leonina*
Pegg, E. J., 1977, *Brit. Vet. J.*, v. 133 (4), 427-431
Toxocara canis and other parasite ova, horticultural flame-gun for control on concrete-floored kennel runs
- Toxascaris leonina* Linstow, 1902
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, *Rev. Iber. Parasitol.*, v. 33 (2-3), 267-271
Felis sylvestris: Huesca, Alto Aragon
- Toxascaris leonina*
Ray, D. K.; Negi, S. K.; and Srivastava, P. S., 1975, *Indian J. Animal Research*, v. 9 (2), 75-78
wild cat: Tarai area, Uttar Pradesh
- Toxascaris leonina*
Read, M. A.; and Thompson, R. C. A., 1976, *J. Helminth.*, v. 50 (2), 95-96
Toxocara canis, *Toxascaris leonina*, prevalence of ova in dog faeces deposited on streets, potential human health hazard: Leeds, England
- Toxascaris leonina*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, *Am. J. Vet. Research*, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections
- Toxascaris leonina* (Linstow, 1902)
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Canis familiaris
Felis catus
all from insular Newfoundland
- Toxascaris leonina*
Stevenson, P.; and Jacobs, D. E., 1977, *J. Helminth.*, v. 51 (2), 149-154
Toxocara canis, *T. cati*, *Ascaris suum*, *Toxascaris leonina*, *Parascaris equorum*, pigs (exper.), in vitro larval precipitate test and indirect fluorescent antibody test using *T. canis* larvae as antigen, indirect fluorescent antibody test using *A. suum* larvae as antigen, specificity
- Toxascaris leonina*
Tharaldsen, J., 1973, *Norwegian J. Zool.*, v. 21 (4), 327-328 [Abstract]
dogs (feces): quarantine station, Oslo, Norway
- Toxascaris leonina*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Toxascaris leonina*
Turner, T.; and Pegg, E., 1977, *Vet. Rec.*, v. 100 (14), 284-285
survey of patent nematode infestations in dogs (faeces): north-west suburban London
- Toxascaris leonina*
Williams, B. M., 1976, *Brit. Vet. J.*, v. 132 (3), 309-312
Vulpes vulpes (intestine): southwest Wales
- Toxascaris leonina*
Yang, J.; and Scholten, T., 1977, *Am. J. Clin. Path.*, v. 67 (3), 300-304
diagnosis of human intestinal parasites, fecal examination technique using Junod's fixative for concentration and permanent staining procedures, comparison with results using formalin-ether procedure
- Toxascaris leonina* Linstow, 1902
Young, P. L.; and Babero, B. B., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Toxocara* Stiles, 1905, *illus.*
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Toxocarinae
key; synonymy
- Toxocara*, *illus.*
Karel, I.; et al., 1977, *Ophthalmologica*, Basel, v. 174 (1), 14-19
Toxocara larva migrans, woman with granulomatous ocular lesion and active larva in the pupillary area, intolerance to mintezol therapy, case report: Czechoslovakia

- Toxocara**
Lifshitz G., A.; Butron P., L.; and Ariza A., A., 1976, *Prensa Med. Mexicana*, v. 41 (9-10), 323-327
Toxocara larval granulomatosis in man associated with immunoblastic lymphadenopathy, clinical case report, possible relationships: Mexico
- Toxocara**
Shields, J. A.; Lerner, H. A.; and Felberg, N. T., 1977, *Am. J. Ophth.*, Chicago, v. 84 (3), 319-322
probable Toxocara endophthalmitis in child with white fundus mass in right eye, differential diagnosis from retinoblastoma on basis of increased eosinophilia and normal lactate dehydrogenase levels in aqueous aspiration, conservative treatment resulted in resolution of eye mass: Philadelphia, Pennsylvania
- Toxocara**
Smythe, R. H., 1977, *Vet. Rec.*, v. 100 (12), 251-252 [Letter]
Toxocara, dogs, resulting excrement following worming may provide means for human infection
- Toxocara spp.**
Borg, O. A.; and Woodruff, A. W., 1976, *Brit. Med. J.* (6036), v. 2, 621-622
technique for recovery of Toxocara spp. ova from soil in order to assess environmental contamination
- Toxocara [sp.]**
Kipnis, R. M.; and Todd, K. S., jr., 1977, *Feline Pract.*, v. 7 (2), 16-19
cat: northern Illinois; Green Bay, Wisconsin
- Toxocara sp.**
Klein, J. B.; and Bradley, R. E., sr., 1976, *Vet. Med. and Small Animal Clin.*, v. 71 (5), 598-599
dogs, cats, sansalid, critical testing, good results
- Toxocara larva**
Lemmingson, W., 1972, *Mod. Problems Ophth.*, v. 10, 312-318
Toxocara larval infestation of eye with resulting secondary retinal detachment, clinical and surgical aspects
- Toxocara sp.**
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1975, *Indian Vet. J.*, v. 52 (6), 451-456
Ancylostoma caninum, dogs (nat. and exper.), nitroxylnil subcutaneously, drug efficacy, good results; nitroxylnil not effective against Taenia sp., Dipylidium sp., Toxocara sp.
- Toxocara sp.**
Rep, B. H.; and Heinemann, D. W., 1976, *Trop. and Geogr. Med.*, v. 28 (2), 104-110
dog: Surinam
- Toxocara spp.**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Toxocara canis**
Adickman, M.; and Tuthill, T. M., 1976, *Postgrad. Med.*, v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Toxocara canis**
Arnaud, J. P., 1976, *Medecine Infant.*, v. 83 (1), 47-54
Toxocara cati, T. canis, clinical aspects of infection in children, clinical forms, diagnosis, prophylaxis, treatment
- Toxocara canis, illus.**
Bwangamoi, O., 1973, *Bull. Epizoot. Dis. Africa*, v. 21 (4), 363-370
dog (stomach to rectum including caecum): Uganda
- Toxocara canis (Werner, 1782)**
Chari, S. S.; and Subramanian, G., 1972, *Indian J. Animal Sc.*, v. 42 (11), 957-960
Toxocara canis, histopathological and histochemical changes in orally infected mice
- Toxocara canis**
Charleston, W. A. G., 1977, *N. Zealand Vet. J.*, v. 25 (7), 171-172
Toxocara canis, T. cati, potential importance of infection in dogs and cats to public health, review: New Zealand
- Toxocara canis**
Chauhan, H. V. S.; Dwivedi, P.; and Kalra, D. S., 1974, *Haryana Vet.*, v. 13 (1), 5-21
protozoan and helminth parasites, transmitted through milk to newborn animals, review
- Toxocara canis, illus.**
Church, E. M.; Wyand, D. S.; and Lein, D. H., 1975, *Am. J. Vet. Research*, v. 36 (3), 331-335
cerebrospinal nematodiasis, experimentally induced in *Oryctolagus cuniculus* with *Ascaris columnaris*, A. suum, or Toxocara canis, naturally occurring in *Sylvilagus floridanus* and *O. cuniculus*, clinical signs, gross and microscopic changes, duration of infection and parasite morphology and distribution in CNS, potential of rabbits as intermediate or paratenic hosts for ascarids of carnivorous origins

- Toxocara canis**
Collins, R. F.; and Ivey, M. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 455-459
skin test responses in guinea pigs infected with small numbers of *Toxocara canis* or *Ascaris suum* and challenged intradermally with several adult and larval somatic antigenic preparations
- Toxocara canis**
Collins, R. F.; and Ivey, M. H., 1975, Am. J. Trop. Med. and Hyg., v. 24 (3), 460-464
passive cutaneous anaphylaxis responses of sensitized guinea pigs to various antigens of adult and larval stages of *Toxocara canis* or *Ascaris suum*; homologous reactions; *Ascaris* larval antigen reacted with *Toxocara* antiserum
- Toxocara canis**
Congdon, L. L.; and Ames, E. R., 1973, Am. J. Vet. Research, v. 34 (3), 417-418
Toxocara canis, prepatent infections, dogs, thiabendazole effective
- Toxocara canis**
Coriglione, G.; Corso, P.; and Gorgone, G., 1969, Minerva Oftal., v. 11 (3), 99-103
larva migrans of *Toxocara canis*, probable cause of macular chorioretinal granuloma in youth, case report, diagnostic problems: Italy
- Toxocara canis**
Cypess, R. H.; et al., 1977, J. Infect. Dis., v. 135 (4), 633-640
visceral larva migrans, human, serum precipitating antibodies specific for larval antigens of *Toxocara canis* as determined by double diffusion in agar, enzyme-linked immunosorbent assay was more sensitive and revealed high titers of antibodies to *Toxocara* larvae in all patients with VLM
- Toxocara canis**
Cypess, R. H.; and Glickman, L. T., 1976, Mod. Vet. Pract., v. 57 (6), 462-464
prevalence of antibody to *Toxocara canis*, human and dogs, enzyme linked immunosorbent assay
- Toxocara canis**
Dafalla, A. A., 1975, Tr. Roy. Soc. Trop. Med. and Hyg., v. 69 (1), 146-147
Toxocara canis, *T. cati* in humans, immunodiagnosis using the capillary-tube precipitin test, cross reaction with *Ascaris* could be eliminated by absorption with *Ascaris* antigen
- Toxocara canis**
Davies, P.; and Nicholas, W. L., 1977, Austral. Vet. J., v. 53 (5), 247-248 [Letter]
dogs (feces): Goodradigbee Shire, New South Wales
- [Toxocara] canis**
Deiana, S.; and Arru, E., 1972, Parassitologia, v. 14 (2-3), 269-273
presence in dogs being treated with Mansonil for test of cestode control
- Toxocara canis**
Dobson, C.; and Welch, J. S., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 223-228
survey for antibodies against *Dirofilaria immitis*, *Toxocara canis*, *Ascaris suum*, *Angiostrongylus cantonensis*, *A. mackerrasae*, in patients with eosinophilia using fluorescent antibody test and passive reversed Arthus test in guinea pigs; *D. immitis* implicated as etiologic agent of human eosinophilic meningitis: Australia
- Toxocara canis**
Enayat, M. S.; and Pezeshki, M., 1977, J. Helminth., v. 51 (2), 143-148
Toxocara canis, guinea pigs (exper.), comparison of counterimmunoelectrophoresis with indirect haemagglutination test for detection of antibodies, possible use of these techniques for immunodiagnosis of human visceral larva migrans
- Toxocara canis**
Engel, H.; et al., 1972, Medicina Alemana, v. 13 (6), 826-837
case report of acute meningomyelitis in young man apparently associated with visceral larva migrans of *Toxocara canis*: Spain
- Toxocara canis**
Fernando, S. T.; and Soulsby, E. J. L., 1974, J. Comp. Path., v. 84 (4), 569-576
Toxocara canis, immunoglobulin classes of antibodies in infected *Macaca sinica*
- Toxocara canis**
Forsdahl, A.; and Brunborg, I., 1971, Tidsskr. Norske Laegefor., v. 91 (13), 958-960
Toxocara canis in dogs, life cycle, implications for serious infections in man and possible associations with epilepsy: Norway
- Toxocara canis, illus.**
Fosella, P. V.; et al., 1974, Atti Soc. Ital. Sc. Vet., v. 28, 858-862
Toxocara canis, infected white mice, electrocardiographic changes related to microascariid pneumonia and localization of larva in myocardium, verified histologically
- Toxocara canis**
Galliard, H., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 295-304
visceral and cutaneous larva migrans, review
- Toxocara canis**
Genchi, C., 1976, Arch. Vet. Ital., v. 27 (3-4), 98-99
dog intestinal helminth ova, incidence in soil samples in public parks, potential public health problem: Milan
- Toxocara canis**
Ghadirian, E.; et al., 1976, Canad. J. Pub. Health, v. 67 (6), 495-498
human toxocariasis, prevalence survey of *Toxocara* spp. and other helminth ova in dogs and soil from city parks, larvae survival over winter months results in continuing contamination of soil and increasing public health problem: Montreal

- Toxocara canis*
Gietko, M.; and Zapart, W., 1975, *Pediatr. Polska*, v. 50 (1), 61-68
visceral larva migrans in young children with severe eosinophilia, diagnosed by intradermal skin test antigens of *Ascaris lumbricoides* and *Toxocara canis*
- Toxocara canis*
Girdwood, R. W. A.; et al., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (4), 284 [Demonstration]
Toxocara canis, *Toxascaris leonina*, incidence in dogs and in soil samples from public places: Glasgow, Scotland
- Toxocara canis*
Glickman, L. T.; and Cypess, R. H., 1977, *Am. J. Pub. Health*, v. 67 (12), 1193-1195
Toxocara canis, epidemiologic survey using the enzyme-linked immunosorbent assay to measure antibodies to *Toxocara* in employees of an animal hospital; results showed that there was no statistical association with either job exposure to dogs or with dog ownership: New York
- Toxocara canis*
Guildal, J. A.; and Clausen, B., 1973, *Norwegian J. Zool.*, v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Toxocara canis*
Hass, D. K.; and Collins, J. A., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 135-137
helminths, dogs, comparative efficacy of vincofos, ticarbodine, mebendazole
- Toxocara canis, illus.*
Hayden, D. W.; and Van Kruiningen, H. J., 1973, *J. Am. Vet. Med. Ass.*, v. 162 (5), 379-384
dogs, eosinophilic gastroenteritis occurring simultaneously with visceral larva migrans and manifesting as chronic diarrhea, clinicopathology and histopathology, case reports, nematode larvae found in lesions of 3 of 5 cases, identified as *Toxocara canis* in 1
- Toxocara canis*
Hayden, D. W.; and Van Kruiningen, H. J., 1975, *Am. J. Vet. Res.*, v. 36 (11), 1605-1614
Toxocara canis, dogs (exper.), eosinophilic gastroenteritis, hematologic findings, serum proteins (β -globulin content as potential diagnostic tool), precipitating humoral antibodies, intradermal test, histopathology, comparison with naturally occurring disease
- Toxocara canis, illus.*
Ho, C. C.; and Chen, K. Y., 1976, *J. Chinese Soc. Vet. Sc.*, v. 2 (2), 80-83
puppy, cavity of tunica vaginalis, gastrointestinal tract, fatal infection apparently of prenatal origin, case history
- Toxocara canis*
Hogarth-Scott, R. S., 1967, *Internat. Arch. Allergy and Applied Immunol.*, v. 32 (2), 201-207
Toxocara canis, *T. cati*, *Toxascaris leonina*, *Ascaris suum*, rabbits (exper.), presence of reagin-like antibodies demonstrable by homologous passive cutaneous anaphylaxis, responsible allergens were common to all 4 nematode species
- Toxocara canis*
Hogarth-Scott, R. S.; and Feery, B. J., 1976, *Austral. J. Exper. Biol. and Med. Sc.*, v. 54 (4), 317-327
existence of cross-reacting antigens between *Toxocara canis* and *Ascaris* spp. and probably between *T. canis* and other nematodes confirmed by in vitro and in vivo tests, such cross-reactions compromise usefulness of skin tests in diagnosis
- Toxocara canis*
Holt, P. E., 1976, *Vet. Rec.*, v. 98 (19), 383
Toxocara canis, puppies, incidence of infection, piperazine citrate: Oldham, Lancashire
- Toxocara canis, illus.*
Huisman, H., 1977, *Ophthalmologica*, Basel, v. 174 (1), 10-13
2 case reports of human *Toxocara canis* solitary granuloma ocular infection, one case previously misdiagnosed as toxoplasmosis, treatment with minzolium and/or photocoagulation with steroid therapy
- Toxocara canis*
Huntley, C. C., 1976, *N. England J. Med.*, v. 294 (23), 1295 [Letter]
Toxocara canis in children, speculation on role of parasite in stimulating IgE antibody levels with resulting allergy to antigen and in turn occurrence of asthma
- Toxocara canis*
Huntley, C. C.; et al., 1976, *Pediatrics*, *Am. Acad. Pediat.*, v. 57 (6), 875-883
Toxocara canis and other helminthiasis affecting humans, suspected but not proved relationship between helminth parasitism of mother and ABO hemolytic disease in the infant, comparison study of populations in Puerto Rico and North Carolina
- Toxocara canis*
Jacobs, D. E.; et al., 1977, *Brit. Med. J.*, (6052), v. 1, 51
Toxocara canis, results of dog breeding kennel survey suggest that infection "not readily acquired by kennel staff maintaining a reasonable standard of personal hygiene"

- Toxocara canis*, *illus.*
Kagei, N.; et al., 1976, Bull. Inst. Pub. Health, Tokyo, v. 25 (3), 140-144
Filaroides hirthei, Strongyloides sp.,
Toxocara canis, and *Trichuris vulpis* in imported dogs, inability to experimentally infect other dogs with *Strongyloides stercoralis* of human origin: Japan, imported from U.S.A.
- Toxocara canis*
Kays, S. G.; and Oaks, J. A., 1976, Am. J. Trop. Med. and Hyg., v. 25 (4), 575-580
effect of inoculum size and length of infection on distribution of *Toxocara canis* larvae in mouse, analysis of variance of larval recovery from various organs, crowding effect observed in heavy infections manifested as altered dispersion rates
- Toxocara canis*
Kessler, G. F.; Frick, O. L.; and Gold, W. M., 1974, Internat. Arch. Allergy and Applied Immunol., v. 47 (3), 313-328
Toxocara canis or *Ascaris suum*-sensitive dogs, experimental asthma, immunologic and physiologic characterization of role of reaginic antibodies
- Toxocara canis*
Khalil, H. M.; et al., 1971, Tr. Roy. Soc. Trop. Med. and Hyg., v. 65 (5), 599-601
toxocariasis in young children, significantly higher incidence of parasitic infection in children suffering from chronic poliomyelitis than in normal children: Egypt
- Toxocara canis*
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, Vet. Rec., v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects
- Toxocara canis* (Werner, 1782) Stiles, 1905
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Vulpes vulpes: Pechora river basin
- Toxocara canis*
Krakowka, S., 1977, J. Am. Vet. Med. Ass., 1977, v. 171 (8), 750-753
transplacentally acquired parasitic diseases of dogs, diagnostic features, review
- Toxocara canis*, *illus.*
Lamina, J., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 305-311
Toxocara canis, human visceral larva migrans, attempted diagnosis using micro-precipitation with living larvae and Ouchterlony gel-precipitin test using *Toxocara* and some other helminth antigens, some cross-reactions
- Toxocara canis*
Lee, K. T.; Min, H. K.; and Soh, C. T., 1976, J. Parasitol., v. 62 (3), 460-465
Toxocara canis, migration and distribution in nonpregnant mice and mice infected at various intervals before and after pregnancy, results suggest that transplacental migration is influenced by developmental stages of placenta
- Toxocara canis*
Loeffler, K., 1974, Prakt. Tierarzt, v. 55, Sondernummer, 68-72
parasites, possible transmission from small domestic animals to man, brief review
- Toxocara canis*
Loret, P.; and Pascalon, J. P., 1972, Medecine Trop., v. 32 (5), 630-635
Toxocara canis infection in 19-month old child characterized by eosinophilia and hepatosplenomegalia of year's duration, case report, mintagez
- Toxocara canis*
McCurdy, H. D.; and Guerrero, J., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1731-1733
helminths, dogs, mebendazole powder, controlled critical studies, good results against all helminths except *Dipylidium caninum*: Kansas; New Jersey; Texas
- Toxocara canis*
Neal, W. P., 1971, Maternal and Child Health, v. 3 (12), 38-39
persistent eosinophilia, diagnostic in suspected human parasitic infestations
- Toxocara canis*
Olson, L. J., 1976, Internat. J. Parasitol., v. 6 (3), 247-251
Toxocara canis larvae in mouse eye, distribution within various eye tissues and effect of previous infection on numbers and distribution, onset and development of hemorrhagic and white cell lesions in anterior eye following challenge of immunized and control mice
- Toxocara canis*, *illus.*
Pavri, K. M.; et al., 1975, Tr. Roy. Soc. Trop. Med. and Hgy., v. 69 (1), 99-110
Toxocara canis in albino mice (exper.), marked synergistic effect in mixed viral infections, possible role of visceral larval migrans in creating "acute encephalopathy syndrome" in presence of simultaneous viral infections
- Toxocara canis*
Pegg, E. J., 1977, Brit. Vet. J., v. 133 (4), 427-431
Toxocara canis and other parasite ova, horticultural flame-gun for control on concrete-floored kennel runs
- Toxocara canis* Werner, 1782
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 267-271
Felis sylvestris: Huesca, Alto Aragon

- Toxocara canis*
Ray, D. K.; Negi, S. K.; and Srivastava, P. S., 1975, Indian J. Animal Research, v. 9 (2), 75-78
jackal: Tarai area, Uttar Pradesh
- Toxocara canis*
Read, M. A.; and Thompson, R. C. A., 1976, J. Helminth., v. 50 (2), 95-96
Toxocara canis, *Toxascaris leonina*, prevalence of ova in dog faeces deposited on streets, potential human health hazard: Leeds, England
- Toxocara canis*
Roberson, E. L.; and Ager, A. L., 1976, Am. J. Vet. Research, v. 37 (12), 1479-1482
cestodes, nematodes, dogs, natural infections, uredofos highly effective, no toxicosis
- Toxocara canis*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, Am. J. Vet. Research, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections
- Toxocara canis*
Robinson, M.; Hooke, F. G.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (2), 104-108
pyrantel pamoate, critical trials against roundworms and hookworms in naturally and experimentally infected dogs, drug efficacy compared with piperazine citrate: Australia
- Toxocara canis*
Robinson, M.; Hooke, F.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (3), 173-176
Trichuris vulpis, *Ancylostoma caninum*, *Toxocara canis*, dogs, oxantel pamoate separately and combined with pyrantel pamoate, drug efficacy, good results: Australia
- Toxocara canis*
Ronald, N. C.; and Bell, R. R., 1976, South-west. Vet., v. 29 (3), 217-218
dogs, critical evaluation of butamisole hydrochloride, highly effective against *Ancylostoma caninum* and *Trichuris vulpis*, not effective against *Toxocara canis*; *Dipylidium caninum* unaffected
- Toxocara canis*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Toxocara canis*
Ruitenbergh, E. J.; et al., 1976, Nederl. Tijdschr. Geneesk., v. 120 (15), 645-649
Toxocara canis, survey of 253 children for complement-fixing antibodies against *Toxocara* shows low incidence; eosinophilia attributed to presence of *Enterobius vermicularis*: Netherlands
- Toxocara canis*
Ruitenbergh, E. J.; and Buys, J., 1976, Vet. Parasitol., v. 1 (3), 231-237
Toxocara canis, puppies (exper.), serodiagnosis with immunofluorescence test
- Toxocara canis*
Ruitenbergh, E. J.; and van Knapen, F., 1977, J. Infect. Dis., v. 136, suppl., S267-S273
application of the enzyme-linked immunosorbent assay to the detection of human and animal helminthic and protozoal infections, advantages of assay for seroepidemiology, discussion of performance of assay in tubes and microplates
- Toxocara canis*
de Savigny, D. H., 1975, J. Parasitol., v. 61 (4), 781-782
Toxocara canis, larvae, in vitro maintenance, simple method of production of excretory-secretory antigen for use in serodiagnostic tests for visceral larva migrans
- Toxocara canis*
de Savigny, D. H.; and Tizard, I. R., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 501-507
Toxocara larva migrans, larval excretions and secretions from in vitro cultures used as antigen in passive hemagglutination and fluorescent antibody tests to diagnose visceral larva migrans in man and laboratory animals (exper.), preliminary evaluation for serodiagnostic purposes, no cross reactions with *Ascaris suum* infections
- Toxocara canis*
Schantz, P. M.; and Prezioso, U., 1976, Am. J. Vet. Research, v. 37 (5), 619-620
immature *Echinococcus granulosus*, dogs, efficacy of divided doses of fospirate (70-94%); also active against *Ancylostoma caninum* and *Toxocara canis*
- Toxocara canis* (Werner, 1782)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Canis familiaris: insular Newfoundland
- Toxocara canis*
Stevenson, P.; and Jacobs, D. E., 1976, Parasitology, v. 73 (2), 1-ii [Abstract]
pigs, milkspot lesion of liver, evidence from serological surveys supports view of *Ascaris suum* as major factor in aetiology and gives little indication that *Toxocara* spp. play any significant role: N. Wales; East Anglia

- Toxocara canis**
Stevenson, P.; and Jacobs, D. E., 1977, *J. Helminth.*, v. 51 (2), 149-154
Toxocara canis, T. cati, Ascaris suum, Toxascaris leonina, Parascaris equorum, pigs (exper.), in vitro larval precipitate test and indirect fluorescent antibody test using T. canis larvae as antigen, indirect fluorescent antibody test using A. suum larvae as antigen, specificity
- Toxocara canis**
Stoye, M., 1976, *Deutsche Tierarztl. Wchnschr.*, v. 83 (3), 107-108
Toxocara canis, beagle dogs infected at conception or parturition, degree of prenatal and lactogenic infection respectively in offspring
- Toxocara canis**
Stromberg, B. E.; and Soulsby, E. J. L., 1977, *Vet. Parasitol.*, v. 3 (2), 169-175
Ascaris suum, guinea pigs, heterologous resistance induced by Toxocara canis and Ancylostoma caninum but not by Haemonchus contortus, Caenorhabditis briggsae, or Turbatrix aceti
- Toxocara canis**
Terziiski, A., 1972, *Izvest. Tsentral. Khel-mint. Lab.*, v. 15, 199-207
Ascaris suum, A. lumbricoides, Toxocara canis, migration in white mice, comparison
- Toxocara canis**
Tharaldsen, J., 1973, *Norwegian J. Zool.*, v. 21 (4), 327-328 [Abstract]
dogs (feces): quarantine station, Oslo, Norway
- Toxocara canis**
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Toxocara canis**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Toxocara canis**
Tomimura, T.; Yokota, M.; and Takiguchi, H., 1976, *Japan. J. Vet. Sci.*, v. 38 (6), 533-548
Toxocara canis in Macaca spp., visceral larva migrans, clinical, hematological, biochemical, and gross pathological observations, occurrence of severe neurological disorders
Macaca fascicularis
M. nemestrina
M. cyclopis
M. fuscata
(all exper.)
- Toxocara canis**
Tongson, M. S.; and Dayrit, A. M., 1975, *Philippine J. Vet. Med.*, v. 14 (2), 53-64
Toxocara canis larvae, dl-tetramisole hydrochloride, white rats
- Toxocara canis**
Tribouley-Duret, J.; et al., 1976, *Compt. Rend. Soc. Biol., Paris*, v. 170 (2), 349-352
Toxocara canis, mice, rabbits, detection of antibodies using antigen prepared from adult worm rather than larva, precipitation, complement fixation, hemagglutination, results show such antigen should be suitable for diagnosis of visceral larva migrans
- Toxocara canis**
Turner, T.; and Pegg, E., 1977, *Vet. Rec.*, v. 100 (14), 284-285
survey of patent nematode infestations in dogs (faeces): north-west suburban London
- Toxocara canis, illus.**
Ubelaker, J. E.; and Allison, V. F., 1975, *J. Parasitol.*, v. 61 (5), 802-807
eggs, fine external morphology, scanning electron microscopy
- Toxocara canis**
Warren, K. S.; and Mahmoud, A. A. F., 1977, *J. Infect. Dis.*, v. 135 (5), 868-872
human ascariasis and toxocariasis, algorithms in diagnosis and clinical management
- Toxocara canis**
Welch, J. S.; and Dobson, C., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (6), 466-472
comparative fluorescent antibody test survey of Aborigines and Caucasians for presence of antibodies to Dirofilaria immitis and correlations with canine filariasis; cross-reactions to Toxocara canis observed only in presence of eosinophilia: Queensland, Australia
- Toxocara canis**
Wendler, H., 1972, *Munchen. Med. Wchnschr.*, v. 114 (39), 1634-1640
visceral larva migrans syndrome in humans, clinical aspects, diagnosis, prophylaxis, case reports, review
- Toxocara canis**
Williams, B. M., 1976, *Brit. Vet. J.*, v. 132 (3), 309-312
Vulpes vulpes (intestine): southwest Wales
- Toxocara canis**
Wiseman, R. A.; Woodruff, A. W.; and Pettitt, L. E., 1971, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 65 (5), 591-598
Toxocara canis, effects of diethylcarbamazine and thiabendazole on survival of larvae in mice (exper.); humans treated with diethylcarbamazine showed decreases in antibody levels soon after therapy started

- Toxocara canis*
Wong, H. S. W.; Embil, J. A.; and Ozere, R. L., 1976, *Exper. Parasitol.*, v. 40 (3), 421-426
Ascaris suum, *Toxocara canis*, guinea pigs sensitized with egg extract antigens, dermal reactivity, macrophage migration inhibition test, and lymphocyte transformation using homologous and heterologous antigens
- Toxocara canis*
Yang, J.; and Scholten, T., 1977, *Am. J. Clin. Path.*, v. 67 (3), 300-304
diagnosis of human intestinal parasites, fecal examination technique using Junod's fixative for concentration and permanent staining procedures, comparison with results using formalin-ether procedure
- Toxocara canis*
Yeoh, T. S.; et al., 1975, *Internat. Arch. Allergy and Applied Immunol.*, v. 49 (3), 371-380
characteristics of anaphylactic histamine release in vitro from peritoneal cells of rats infected with *Toxocara canis*, both disodium cromoglycate and levamisole produced dose-related inhibition of histamine release
- Toxocara canis*
Young, P. L.; and Babero, B. B., 1975, *Proc. Oklahoma Acad. Sc.*, v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
rats (lungs)
(all exper.)
- Toxocara canis*, *illus.*
Zyngier, F. R., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (2), 225-228
Toxocara canis, mice (exper.), histopathology of liver, lung, brain, and muscle
- Toxocara canis*
Zyngier, F. R., 1976, *N. England J. Med.*, v. 295 (26), 1483
Toxocara canis-infected mice, inclusion bodies in neutrophils observed on bronchial lumen, assumed to be Charcot-Leyden crystals
- Toxocara canis*
Zyngier, F. R.; and Brockbank, A., 1974, *Ann. Trop. Med. and Parasitol.*, v. 68 (2), 229-233
Toxocara canis, mice (exper.), pathology of lung involvement, electron microscopy
- Toxocara canis*, *illus.*
Zyngier, F. R.; and Santa-Rosa, G., 1976, *Ann. Trop. Med. and Parasitol.*, v. 70 (4), 445-448
Toxocara canis, mice, histopathology of multiple infection, influence of antihistamines and corticosteroids
- Toxocara cati*
Adickman, M.; and Tuthill, T. M., 1976, *Postgrad. Med.*, v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Toxocara cati*
Arnaud, J. P., 1976, *Medecine Infant.*, v. 83 (1), 47-54
Toxocara cati, *T. canis*, clinical aspects of infection in children, clinical forms, diagnosis, prophylaxis, treatment
- Toxocara cati*
Charleston, W. A. G., 1977, *N. Zealand Vet. J.*, v. 25 (7), 171-172
Toxocara canis, *T. cati*, potential importance of infection in dogs and cats to public health, review: New Zealand
- Toxocara cati*
Dafalla, A. A., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (1), 146-147
Toxocara canis, *T. cati* in humans, immunodiagnosis using the capillary-tube precipitin test, cross reaction with *Ascaris* could be eliminated by absorption with *Ascaris* antigen
- Toxocara cati*
Ghadirian, E.; et al., 1976, *Canad. J. Pub. Health*, v. 67 (6), 495-498
human toxocarasis, prevalence survey of *Toxocara* spp. and other helminth ova in dogs and soil from city parks, larvae survival over winter months results in continuing contamination of soil and increasing public health problem: Montreal
- Toxocara cati*
Gorgas, M., 1976, *Zool. Garten N. F.*, v. 46 (1-2), 82-85
Toxocara leonina, *T. cati*, tigers, cerebral disorders caused by somatically migrating larvae, treatment with neguvon and thiabendazole
- Toxocara cati*
Gregory, G. G.; and Munday, B. L., 1976, *Austral. Vet. J.*, v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island
- Toxocara cati*
Hogarth-Scott, R. S., 1967, *Internat. Arch. Allergy and Applied Immunol.*, v. 32 (2), 201-207
Toxocara canis, *T. cati*, *Toxascaris leonina*, *Ascaris suum*, rabbits (exper.), presence of reagin-like antibodies demonstrable by homologous passive cutaneous anaphylaxis, responsible allergens were common to all 4 nematode species
- Toxocara cati* (Schrank, 1788) Brumpt, 1927
Mirzayans, A., 1973, *Vet. Rec.*, v. 92 (10), 262 [Letter]
Syn.: *T. mystax* (Zeder, 1800)
pony (small intestine): northern Iran

- Toxocara cati*
Neal, W. P., 1971, Maternal and Child Health, v. 3 (12), 38-39
persistent eosinophilia, diagnostic in suspected human parasitic infestations
- Toxocara cati*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Toxocara cati*
de Savigny, D. H.; and Tizard, I. R., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 501-507
Toxocara larva migrans, larval excretions and secretions from in vitro cultures used as antigen in passive hemagglutination and fluorescent antibody tests to diagnose visceral larva migrans in man and laboratory animals (exper.), preliminary evaluation for serodiagnostic purposes, no cross reactions with *Ascaris suum* infections
- Toxocara cati*
Stevenson, P.; and Jacobs, D. E., 1976, Parasitology, v. 73 (2), i-ii [Abstract]
pigs, milkspot lesion of liver, evidence from serological surveys supports view of *Ascaris suum* as major factor in aetiology and gives little indication that *Toxocara* spp. play any significant role: N. Wales; East Anglia
- Toxocara cati*
Stevenson, P.; and Jacobs, D. E., 1977, J. Helminth., v. 51 (2), 149-154
Toxocara canis, *T. cati*, *Ascaris suum*, *Toxascaris leonina*, *Parascaris equorum*, pigs (exper.), in vitro larval precipitate test and indirect fluorescent antibody test using *T. canis* larvae as antigen, indirect fluorescent antibody test using *A. suum* larvae as antigen, specificity
- Toxocara cati*
Uhlikova, M.; and Huebner, J., 1974, J. Protozool., v. 21 (3), 458-459 [Abstract]
Toxoplasma gondii, *Toxocara cati*, concomitant incidence of infection in 8 children
- Toxocara cati*
Warren, K. S.; and Mahmoud, A. A. F., 1977, J. Infect. Dis., v. 135 (5), 868-872
human ascariasis and toxocariasis, algorithms in diagnosis and clinical management
- Toxocara cati* Schrank, 1788
Young, P. L.; and Babero, B. B., 1975, Proc. Oklahoma Acad. Sc., v. 55, 169-174
helminthic diseases, cockroaches may play an important role in transmission
Periplaneta americana
Blattella germanica
Blaberus giganteus
Parcoblatta sp.
(all exper.)
- Toxocara felis* Goeze, 1782
Ramon Vericad, J.; and Sanchez Acedo, C., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 267-271
• *Felis sylvestris*: Huesca, Alto Aragon
- Toxocara leonina*
Ghadirian, E.; et al., 1976, Canad. J. Pub. Health, v. 67 (6), 495-498
human toxocariasis, prevalence survey of *Toxocara* spp. and other helminth ova in dogs and soil from city parks, larvae survival over winter months results in continuing contamination of soil and increasing public health problem: Montreal
- Toxocara leonina*
Gorgas, M., 1976, Zool. Garten N. F., v. 46 (1-2), 82-85
Toxocara leonina, *T. cati*, tigers, cerebral disorders caused by somatically migrating larvae, treatment with neguvon and thiabendazole
- Toxocara mystax*
Guterbock, W. M.; and Levine, N. D., 1977, J. Am. Vet. Med. Ass., v. 170 (12), 1411-1413
cats (feces): east central Illinois
- Toxocara mystax*
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, Vet. Rec., v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects
- Toxocara mystax* (Zeder, 1800) Stiles, 1907
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Felis catus dom.: Pechora river basin
- Toxocara mystax*
Loeffler, K., 1974, Prakt. Tierarzt, v. 55, Sondernummer, 68-72
parasites, possible transmission from small domestic animals to man, brief review
- Toxocara mystax* (Zeder, 1800)
Mirzayans, A., 1973, Vet. Rec., v. 92 (10), 262 [Letter]
as syn. of *T. cati* (Schrank, 1788) Brumpt, 1927
- Toxocara mystax* (Zeder, 1800)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Felis catus: insular Newfoundland
- Toxocara mystax*
Torres, P.; and Barriga, O. O., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 79-85
Ascaris suum, *A. lumbricoides*, *Toxocara mystax*, *Ascaridia galli*, comparative antigenic analysis by gel double diffusion and immunoelectrophoresis
- Toxocara mystax*, illus.
Ubelaker, J. E.; and Allison, V. F., 1975, J. Parasitol., v. 61 (5), 802-807
eggs, fine external morphology, scanning electron microscopy

- Toxocara* (*Neoascaris*) *vitulorum*
Mia, S.; et al., 1975, *Trop. Animal Health and Prod.*, v. 7 (3), 153-156
Toxocara vitulorum, infection of buffalo calves via colostrum
- Toxocariasis**
Daffalla, A. A.; and Ali Omer, M. I., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 544-545
probable toxocariasis in 7-year old boy with high eosinophilia, fever and abdominal distention; positive toxocaral skin test and fluorescent antibody test, marked improvement with diethylcarbamazine therapy: Gezira area, Sudan
- Toxocariasis**
Pattanayak, P. C., 1974, *Indian J. Animal Research*, v. 8 (2), 56-58
Ancylostoma sp., toxocariasis, pups, clinical trial with tetramisole, effective safe medication
- Toxocariasis**
Patterson, R.; et al., 1975, *Pediatrics*, *Am. Acad. Pediat.*, v. 56 (3), 417-420
differentiation of visceral larva migrans caused by toxocariasis from that caused by ascariasis using radioimmunoassay to demonstrate serum immunoglobulins
- Toxocariasis**
Woodruff, A. W., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (6), 755-769
toxocariasis as a public health problem, review
- Toxocarinae** (Hartwich, 1954, fam.) Osche, 1958
Hartwich, G., 1974, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (2), pp. 1-15
Ascarididae
key; key to genera
includes: *Toxocara*; *Porrocaecum*; *Paradujardinia*
- Travassosius Khalil**, 1922
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongylidae, *Trichostrongylinae*
- Travassosius rufus** Kh.
Volokh, A. M.; and Samarskii, S. L., 1977, *Vestnik Zool.*, *Akad. Nauk Ukrainsk. SSR, Inst. Zool.* (3), 89-90
Stichorchis subtriquetus, *Travassosius rufus*, incidence in relation to sex and age of host [Castor fiber]: Middle Dnieper area
- Travassospirura Moennig**, 1958
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Spirura* Blanchard, 1849
- Travassostrongylus Orloff**, 1933
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, *Viannaiinae*
Syn.: *Camerostongylus* Wolfgang, 1951
- Travassostrongylus callis** (Travassos, 1914)
Diaw, O. T., 1976, *Ann. Parasitol.*, v. 51 (3), 355-363
trichostrongyloid nematode fauna of *Didelphis marsupialis* compared to that of *Metachirops opossum*, localization within intestine
Didelphis marsupialis (intestin): Guyane francaise
- Travassostrongylus callis** (Travassos, 1914), *illus.*
Diaw, O. T., [1977], *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (405), 1976, *Zool.* (282), 1065-1089
redescription
Didelphis marsupialis (intestin): Guyane francaise
- Travassostrongylus orloffii** Travassos, 1935
Diaw, O. T., 1976, *Ann. Parasitol.*, v. 51 (3), 355-363
trichostrongyloid nematode fauna of *Didelphis marsupialis* compared to that of *Metachirops opossum*, localization within intestine
Didelphis marsupialis (intestin): Guyane francaise
- Travassostrongylus orloffii** Travassos, 1935, *illus.*
Diaw, O. T., [1977], *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (405), 1976, *Zool.* (282), 1065-1089
redescription
Didelphis marsupialis (intestin): Guyane francaise
- Travassostrongylus paraquintus** n. sp., *illus.*
Durette-Desset, M. C., 1974, *Ann. Parasitol.*, v. 49 (5), 555-566
4 spp. of *Trichostrongyloidea*, localization in intestine, larval and adult synophes compared, implications for taxonomy and evolution
Metachirops opossum (intestin): Guyane Francaise
- Travassostrongylus tourei** Diaw, 1976 [nomen nuda]
Diaw, O. T., 1976, *Ann. Parasitol.*, v. 51 (3), 355-363
trichostrongyloid nematode fauna of *Didelphis marsupialis* compared to that of *Metachirops opossum*, localization within intestine
Didelphis marsupialis (intestin): Guyane francaise
- Travassostrongylus tourei** n. sp., *illus.*
Diaw, O. T., [1977], *Bull. Mus. National Hist. Nat.*, Paris, 3. s. (405), 1976, *Zool.* (282), 1065-1089
Didelphis marsupialis (partie anterieure de l'intestin): Guyane francaise
- Trichinella**
Berman, H. A.; and Weinstein, L., 1970, *Med. J. Australia*, v. 2 (13), 583-584
serum survey of 109 persons for presence of *Trichinella* antibodies, negative findings: Western Samoa; Fiji

- Trichinella**
 Ershov, V. S.; et al., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 343-348
 anaphylactic shock in guinea pigs after sensitization with free-living or plant-parasitic nematodes and challenge with various helminth antigens indicates antigenic components in common; intradermal tests using antigen from free-living nematode in cases of ascariasis, trichinellosis, and cysticercosis; possible use of free-living nematode to immunize against dictyocaulosis and ascariasis
- Trichinella**
 Kassur, B.; Januszkiewicz, J.; and Poznanska, H., 1970, Mater. Med. Pol. (4), v. 2 (2-3), 32-43
 experimental and human trichinosis, changes in enzyme activity in serum and muscle tissue, possible relationships to pathologic processes and diagnosis
- Trichinella**
 Most, H., 1972, N. England J. Med., v. 287 (10), 495-498; (14), 698-702
 common parasitic infections of man encountered in the United States, recommendations for treatment, review
- Trichinella, illus.**
 Thomsen, D. U., 1976, Medlemsbl. Danske Dyr-laegeforen., v. 59 (11), 481-490
 Trichinella, improved diagnostic technique for pig meat post mortem using digestion in Colworth Stomacher 3500; 50% better results than trichinoscopic method, time reduced from six hours to one
- Trichinella, illus.**
 Timonov, E. V., 1970, Parazitologiya, Leningrad, v. 4 (3), 237-240
 Trichinella, morphogenesis, luminescent microscopy with acridine orange and rhodamine C
- Trichinella [sp.]**
 Hoerning, B., 1977, Schweiz. Arch. Tierh., v. 119 (8), 337-339
 Vulpes vulpes
 Meles meles
 Martes foina
 M. martes
 all from Schweiz
- Trichinella [sp.]**, probably *T. nativa*, Britov and Boev, 1972, illus.
 Nelson, G. S.; et al., 1975, J. Helminth., v. 49 (4), 301-303
 xeroradiographic visualization of trichinae in polar bear muscle
- Trichinella [sp.]**, probably *T. nativa* Britov and Boev, 1972
 Nelson, G. S.; et al., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 10 [Demonstration]
 Trichinella [sp.] cysts, visualization in polar bear diaphragm muscle using xeroradiography: London Zoological Society Garden (Imported from Moscow Zoo)
- Trichinella larvae**
 Ridley, D. S.; and Hedge, E. C., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (6), 522-525
 microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Trichinella nativa**
 Orlov, I. V.; Britov, V. A.; and Boev, S. N., 1976, Vestnik Sel'skokhoz. Nauki (243) (12), 61-68
 Trichinella spp., experimental hybridization between species shows very limited crossing, reproductive isolation; useful technique for species diagnosis
- Trichinella nelsoni, illus.**
 Mutafova, T.; and Komandarev, S., 1976, Ztschr. Parasitenk., v. 48 (3-4), 247-250
 Trichinella nelsoni, laboratory strain, karyotype of males and females
- Trichinella nelsoni**
 Orlov, I. V.; Britov, V. A.; and Boev, S. N., 1976, Vestnik Sel'skokhoz. Nauki (243) (12), 61-68
 Trichinella spp., experimental hybridization between species shows very limited crossing, reproductive isolation; useful technique for species diagnosis
- Trichinella pseudospiralis**
 Orlov, I. V.; Britov, V. A.; and Boev, S. N., 1976, Vestnik Sel'skokhoz. Nauki (243) (12), 61-68
 Trichinella spp., experimental hybridization between species shows very limited crossing, reproductive isolation; useful technique for species diagnosis
- Trichinella pseudospiralis Garkavi, 1972**
 Tomasovicova, O., 1975, Biologia, Bratislava, s. B, Zool., v. 30 (11), 821-826
 experimental infection of chickens, *Gallus gallus dom[esticus]*, infection of mice with *Trichinella pseudospiralis* from chickens, viability and infectivity maintained; experimental infection of chickens with *T. spiralis* only of brief duration; possibility that wild and domestic birds actively transfer and disseminate *T. pseudospiralis* but act only as passive hosts of intestinal stage *T. spiralis*
- Trichinella spiralis**
 Ahmad, R.; and Harpur, R. P., 1977, Canad. Fed. Biol. Soc., Programme and Proc. 20. Ann. Meet., v. 20, 72 [Abstract]
 Trichinella spiralis, course of infection in rats, exercise had no effect on number of larvae recovered, physically fit rats had higher blood pressures and lower haematocrit values

- Trichinella spiralis*
Aita, J. F.; and Kramer, M. D., 1973, Maryland State Med. J., v. 22 (12), 40-46
case reports and clinical review of human trichinosis, mintezol: Maryland
- Trichinella spiralis*
Alvarez, V.; et al., 1970, Bol. Chileno Parasitol., v. 25 (1-2), 83-86
Trichinella spiralis, review of surveys for possible reservoirs in Chile, positive findings in dogs, cats and rats but no infections found in wild mammals or whales
- Trichinella spiralis*
Ambia Medina, J.; and Quiroz Romero, H., 1976, Veterinaria, Mexico, v. 7 (1), 17-19
dogs, prevalence (intercostal, masseter, and diaphragmatic muscles): urban zones of Mexico city
- Trichinella spiralis*
Andrews, J. S.; Hill, C. H.; and Henson, L. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 81-84
Trichinella spiralis, pigs, trichina-cyst antigen, intradermal diagnosis, results unreliable
- Trichinella spiralis*, illus.
Auger, P.; et al., 1976, Canad. Med. Ass. J., v. 114 (6), 522-525
clinical discussion of 4 cases of *Trichinella spiralis* infection in Montreal family, positive diagnosis by muscle biopsy, immunoserological and pathologic aspects reviewed: Canada
- Trichinella spiralis*
Barnett, J. B.; and Justus, D. E., 1975, Infect. and Immun., v. 11 (6), 1342-1351
Trichinella spiralis, mice, no direct relationship between mast cell degranulation, anaphylaxis, and production of homocytotropic antibodies
- Trichinella spiralis*
Barrett-Connor, E.; et al., 1976, J. Infect. Dis., v. 133 (4), 473-477
Trichinella spiralis, outbreak in campers after eating roasted wild pig, diagnosis by eosinophilia and sero-immunologic studies; diagnostic test comparisons, skin-test antigen inconclusive: California (infected in Hawaii)
- Trichinella spiralis*
Barriga, O. O., 1968, Bol. Chileno Parasitol., v. 23 (1-2), 33-37
Trichinella spiralis, resistance of chicks to experimental infection (intestinal phase)
- Trichinella spiralis*
Barriga, O. O., 1977, J. Clin. Microbiol., v. 6 (3), 274-279
Trichinella spiralis, different antigenic fractions, reactivity and specificity (tested for cross-reactions against *Ascaris suum*) in cutaneous (immediate and delayed) and serological (bentonite agglutination, hemagglutination, hemagglutination inhibition) tests, implications for clinical diagnosis of trichinellosis
- Trichinella spiralis*
Behnke, J. M.; et al., 1976, Parasitology, v. 73 (2), xv [Abstract]
Trichinella spiralis expulsion from mice, effect on concurrent helminth infections (*Hymenolepis diminuta*, *H. microstoma*, *Aspicularis tetraptera*)
- Trichinella spiralis*
Behnke, J. M.; Bland, P. W.; and Wakelin, D., 1977, Parasitology, v. 75 (1), 79-88
rejection phase of *Trichinella spiralis* infection in mice had marked negative effect on growth and survival of *Hymenolepis diminuta*; this effect was not mediated by direct cross-immunity nor was it a direct consequence of inter-specific competition
- Trichinella spiralis*
Behnke, J. M.; Wakelin, D.; and Wilson, M. M., 1977, Parasitology, v. 75 (2), xxxiv-xxxv [Abstract]
interactions between intestinal phase of *Trichinella spiralis* and *Nematospiroides dubius*
- Trichinella spiralis*
Bessonov, A. S., 1977, [Trichinosis], 110 pp., illus.
handbook on epizootiology, epidemiology, clinical aspects, pathology, diagnosis, prophylaxis, therapy, review
- Trichinella spiralis*
Blackwood, L. L.; and Molinari, J. A., 1976, Abst. Ann. Meet. Am. Soc. Microbiol., 82
Trichinella spiralis, mice, delayed hypersensitivity to heterologous antigens, immune capabilities dependent upon phase of parasitic life cycle
- Trichinella spiralis*
de Boni, U.; Lenczner, M. M.; and Scott, J. W., 1977, Canad. Med. Ass. J., v. 117 (5), 472
Trichinella spiralis cyst discovered near subcutaneous border of intercostal muscle during autopsy and histopathologic examination of Egyptian mummy
- Trichinella spiralis*
Brown, A. R.; and Crandall, C. A., 1976, J. Immunol., v. 116 (4), 1105-1109
mice, *Ascaris suum*-induced phosphorylcholine-binding component identified as IgM antibody having idiotypic determinants in common with PC-binding IgA myeloma TEPC 15, response not duplicated by immunization with dead *Ascaris* larvae or by infection with *Heligmosomoides polygyrus* or *Trichinella spiralis*
- Trichinella spiralis*
Bruce, R. G.; Rose, M.; and Parrott, D. M. V., 1976, Parasitology, v. 73 (2), xvii-xviii [Abstract]
Trichinella spiralis, mice, lymphoblasts, enhanced migration to and localization in small intestinal tissue at 2 and 4 days after infection but not at 6-10 days, primary effectors of cell-mediated response in gut

- Trichinella spiralis*
Bruce, R. G.; and Wakelin, D., 1977, *Parasitology*, v. 74 (2), 163-173
Trichinella spiralis, *Trichuris muris*, concurrent infection in mice, interactive expulsive response considered an example of indirect cross-immunity with no element of antigenic similarity, involvement of cell-mediated inflammatory response strongly suggested
- Trichinella spiralis*, *illus.*
Brzosko, W. J.; and Gancarz, Z., 1970, *Med. Dosw. i Mikrobiol.*, v. 22 (1), 91-94
Trichinella spiralis, electron microscopy of larval cuticular antigenic structure
- Trichinella spiralis*
Buerger, H. J., 1976, *Zentralbl. Vet.-Med.*, Reihe B, v. 23 (8), 678-697
Trichinella spiralis, rats, mechanism of immune elimination, dose of infection and sex of rats affected time of onset of worm expulsion; number of female trichinellae decreased earlier than number of male worms
- Trichinella spiralis*
Buerger, H. J., 1976, *Zentralbl. Vet.-Med.*, Reihe B, v. 23 (9), 705-732
Trichinella spiralis, rats immunized with sensitized cells from spleen, lymph nodes, or thymus eliminated worm burdens earlier than normal cell controls
- Trichinella spiralis*
Buerger, H. J., 1976, *Zentralbl. Vet.-Med.*, Reihe B, v. 23 (10), 793-800
Trichinella spiralis, rats, passively immunized during plateau phase with isologous or allogous immune globulin from infected rats, worm burdens and body lengths not changed, PCA titers significantly reduced, IHA antibodies elevated
- Trichinella spiralis*
Buerger, H. J., 1977, *Zentralbl. Vet.-Med.*, Reihe B, v. 24 (1), 1-24
Trichinella spiralis, interference with the cellular immunological system of rats, antithymocyte serum and an antiserum retarded immune elimination of *T. spiralis* from intestine, more pronounced effects in neonatally thymectomized rats; homologous immune serum unable to restore the retarding effect of neonatal thymectomy and/or antithymocyte serum treatment; results suggest that T cells are involved in immune elimination
- Trichinella spiralis*
Bura, M. W. T.; and Willett, W. C., 1977, *East African Med. J.*, v. 54 (4), 185-193
Trichinella spiralis, extensive epidemiologic survey of trichinosis outbreak (11 persons of whom 2 died of infection) in Iraqw native tribe, source of infection probably a warthog killed and shared by 4 families, potential increasing public health problem if domestic pigs come in contact with carcasses of infected wild pigs: Tanzania
- Trichinella spiralis*
Bussieras, J., 1976, *Rec. Med. Vet.*, v. 152 (4), 229-234
Trichinella spiralis, epidemiology, role of wild and domestic mammals, review, suggestions for controlling transmission to man: France
- Trichinella spiralis*
Campbell, W. C.; and Blair, L. S., 1975, *J. Parasitol.*, v. 61 (6), 1116-1117
Trichinella spiralis, failure to confirm reported lethal effect of cytotoxic drugs on encapsulated larvae in mice
- Trichinella spiralis*
Campbell, W. C.; Malanga, C. M.; and Conroy, J. A., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (2), 163
Trichinella spiralis infected mice, no significant protection against subsequent *Trypanosoma cruzi* infection
- Trichinella spiralis*
Capron, A.; et al., 1977, *Ann. Immunol.*, v. 128C (1-2), 541-556
impairment of immune response in parasitic infections characterized by high prevalence of autoantibodies and by immunosuppression, review discussing malaria, trypanosomiasis, trichinosis, and schistosomiasis, with some original material on the last
- Trichinella spiralis*
Carney, I. F., 1976, *Internat. Arch. Allergy and Applied Immunol.*, v. 50 (3), 322-328
Trichinella spiralis, guinea pigs, IgE-mediated anaphylactic bronchoconstriction, severity reduced by disodium cromoglycate treatment
- Trichinella spiralis*
Castro, G. A.; et al., 1974, *Proc. Soc. Exper. Biol. and Med.*, v. 146 (3), 703-706
Trichinella spiralis, *Hymenolepis diminuta*, rats (exper.) in which all nutrients were derived from parenteral or exocrino-enteric circulation rather than by ingesting food orally; *H. diminuta* failed to develop and *T. spiralis* showed differences from normal population size thus suggesting the importance of food in the host intestine in regulating development of tissue and lumen-dwelling parasites
- Trichinella spiralis*
Castro, G. A.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (6), 848-853
intestinal parasites, rats, serum and antral gastrin levels, *Trichinella spiralis* associated with inflammatory changes in small bowel mucosa and with significant increase in serum gastrin, neither changes in hormone level nor inflammation induced by *Hymenolepis diminuta*, findings suggest that pathologic changes caused by enteric parasites may be due to changes in functions that are regulated by gastrointestinal hormones

- Trichinella spiralis*
Castro, G. A.; et al., 1976, Gastroenterology, v. 71 (4), 620-625
Trichinella spiralis, increased propulsive activity in parasitized rats with associated inflammatory changes and a significant reduction in disaccharidase levels in gut mucosa
- Trichinella spiralis*
Castro, G. A.; et al., 1976, J. Parasitol., v. 62 (3), 353-359
course of infection with Trichinella spiralis and Hymenolepis diminuta when a parasitized, enterally fed rat is switched to total parenteral nutrition
- Trichinella spiralis*
Castro, G. A.; Post, C. A.; and Roy, S. A., 1977, J. Parasitol., v. 63 (4), 713-719
Trichinella spiralis-immunized rats, challenge infection does not elicit changes in intestinal motility in contrast to a primary infection of equal size which enhances intestinal transit
- Trichinella spiralis*
Castro, G. A.; Roy, S. A.; and Schanbacher, L. M., 1975, J. Parasitol., v. 61 (6), 1053-1060
Trichinella spiralis, untreated worms or worms exposed to phytohemagglutinin or immune serum, in vitro effects of lamina propria cells from small intestine of immunized rats, deleterious effect of disrupted (but not intact) cells on juveniles and adults (but not larvae), vermucidal component not linked to peroxidase-H₂O₂-halide system
- Trichinella spiralis*
Chang, G. N.; and Wang, W. Y., 1976, Taiwan J. Vet. Med. and Animal Husb. (28), 30-34
survey in rats and slaughtered pigs, no positive case seen: southern Taiwan
- Trichinella spiralis*
Chimyshkyan, K. L.; et al., 1976, Biomedicine, v. 25 (5), 176-180
inhibition of transplantation immunity and ability of lymphoid cells to induce graft-versus-host reactions during certain phases of Trichinella spiralis infections
- Trichinella spiralis*
Cironeanu, I., 1975, Rev. Crest. Animalelor, v. 25 (2), 81-82
Sarcocystis miescheriana, Trichinella spiralis, differential diagnosis of cysts
- Trichinella spiralis*
Clausen, B.; and Henriksen, S. A., 1976, Nord. Vet.-Med., v. 28 (4-5), 265-270
Vulpes vulpes: Denmark
- Trichinella spiralis*
Clinard, E. H., 1975, Am. J. Vet. Research, v. 36 (5), 615-618
Trichinella spiralis, swine (exper.), soluble-antigen fluorescent antibody test evaluated at intervals from 7 days to 1 year, possibly adequate as mass screening test for surveillance and control of trichinosis in swine
- Trichinella spiralis*
Colella, G., 1975, Vet. Ital., v. 26 (9-12), 371-377
distribution of cases in man, wolf, dog and fox
dogs: province of Matera, Italy
- Trichinella spiralis*
Colley, D. G., 1976, Cellular Immunol., v. 24 (2), 328-335
spleen or lymph node cells from Schistosoma mansoni; infected mice respond to challenge with soluble egg antigenic preparation by elaboration of eosinophil stimulation promoter, culture conditions, antigen requirements, production kinetics, and immunologic specificity of this lymphokine, ability to stimulate eosinophil migration from eosinophil-rich peritoneal exudates from either S. mansoni- or Trichinella spiralis-infected mice
- Trichinella spiralis*
Corba, J.; and Spaldonova, R., 1974, Biologia, Bratislava, s. B, Zool. (1), v. 29 (2), 167-173
Trichinella spiralis, mice, immunosuppressive substances given at intestinal phase cause significant increase of muscle trichinellae, but only slight increase when given at migratory phase; host immunity mechanism more effective at intestinal phase and its inhibition causes longer stay in intestine, higher reproduction and more larvae in muscle phase
- Trichinella spiralis*
Crandall, R. B., 1975, J. Parasitol., v. 61 (3), 566-567
Trichinella spiralis, C57Bl/6J mice, decreased resistance with age, prior infection prevented increased susceptibility of aged mice
- Trichinella spiralis*
Cremers, H. J. W. M.; Jansen, J.; and Swierstra, D., 1975, Tijdschr. Diergeneesk., v. 100 (22), 1209-1211
Rattus sp.: Netherlands
- Trichinella spiralis*
Crum, E. D.; Despommier, D. D.; and McGregor, D. D., 1977, Immunology, v. 33 (6), 787-795
Trichinella spiralis, rats, immunization by series of methyridine-terminated oral infections with larvae, thoracic duct lymphocytes from immunized animals can protect normal rats against challenge, protective cells belong to 2 different populations, immune serum and lymph fail to transfer resistance
- Trichinella spiralis*
Curlphey, J. E., 1971, Oral Surg., v. 31 (1), 19-24
Trichinella spiralis occurring as mass on right mandible in man who had no other symptoms or clinical signs, complete recovery after surgical removal of mass: England

- Trichinella spiralis*, illus.
Cypess, R. H.; et al., 1977, J. Med. Primatol., v. 6 (1), 23-32
Trichinella spiralis in *Macaca mulatta* (exper.), parasite distribution throughout muscular system, clinical and pathological changes, monkeys develop trichinosis which is clinically, pathologically and morphologically similar to human trichinosis
- Trichinella spiralis*
Czerpak, R., 1970, Acta Parasitol. Polon., v. 17 (20-38), 285-292
Trichinella spiralis larvae, mice given gold-thioglucose and vitamin A, oxygen uptake by diaphragm muscles, influence of host sex and age
- Trichinella spiralis*, illus.
Davis, M. J.; et al., 1976, Neurology, v. 26 (1), 37-40
Trichinella spiralis in man, severe muscle involvement (4,046 larvae per gram of muscle), clinical case report, history of frequent consumption of pickled pigs' feet, recovery after thiabendazole given concomitantly with prednisone: New York City
- Trichinella spiralis*, illus.
Despommier, D., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, Pt. 2), 68-75
Trichinella spiralis, immunity, correlation of biological activities of various stages of infection with host protective mechanisms active against these worm stages, workshop report
- Trichinella spiralis*
Despommier, D. D.; et al., 1977, Immunology, v. 33 (6), 797-805
Trichinella spiralis, influence of immune lymphocytes on life cycle in normal and in irradiated rats, localization of labelled lymphocytes in tissues, evidence that immunity is directed against enteral stage of parasite and is expressed in rapid expulsion of worms from small intestine, immune T cells probably have 'helper' function in promoting formation of protective B cells
- Trichinella spiralis*
Despommier, D. D.; Campbell, W. C.; and Blair, L. S., 1977, Parasitology, v. 74 (1), 109-119
Trichinella spiralis, correlation of in vitro adult worm fecundity with recoverable muscle larvae in immunized and non-immunized rats; in vitro fecundity of individual adult female worms recovered from non-immunized mice; effects of a high vs. a low dose of antigen on adult counts, adult fecundity, and number of recoverable muscle larvae in mice
- Trichinella spiralis*, illus.
Despommier, D. D.; and Mueller, M., 1976, J. Parasitol., v. 62 (5), 775-785
Trichinella spiralis, stichosome ultrastructure, stichocytes contain two major types of granules which contain antigens identical to those previously described from excretion-secretion products of mature muscle larvae
- Trichinella spiralis*, illus.
Dukic, Z.; et al., 1976, Vet. Glasnik, v. 30 (7), 629-635
Trichinella spiralis, distribution, routes of infection in wild and domestic swine and man, rats and foxes as possible vectors: Serbia
- Trichinella spiralis*
Enigk, K.; Feder, H.; and Dey-Hazra, A., 1976, Tropenmed. u. Parasitol., v. 27 (1), 57-69
mineral contents of blood and chemical composition of muscles of *Trichinella spiralis*-infected pigs, comparison with normal controls
- Trichinella spiralis*
Farris, K. N.; and Harley, J. P., 1977, Exper. Parasitol., v. 41 (1), 17-30
Trichinella spiralis-infected mice, alteration of gastrocnemius muscle kinetics, correlation with known morphological, biochemical, and histochemical changes
- Trichinella spiralis*
Faubert, G. M., 1976, Immunology, v. 30 (4), 485-489
Trichinella spiralis, depression of plaque-forming cells to sheep red blood cells by new-born larvae in vivo (mice) and in vitro, transitory phenomenon
- Trichinella spiralis*
Faubert, G., 1977, Ann. Immunol., v. 128C (1-2), 281-282
Trichinella spiralis, mice, temporary immunodepression may be related to migrating phase only
- Trichinella spiralis*
Faubert, G. M., 1977, Exper. Parasitol., v. 43 (2), 336-341
Trichinella spiralis in Swiss mice, expulsion rate during primary and challenge infections, numbers of encysted muscle larvae also needed as assay for immunity, response of plaque-forming cells to sheep red blood cells in challenge infections used to determine timing of immunosuppression
- Trichinella spiralis*
Faubert, G. M.; and Tanner, C. E., 1975, Immunology, v. 28 (6), 1041-1050
Trichinella spiralis, leucoagglutinating and leucotoxic activity of serum of infected mice and of saline extracts of larvae, capacity of infected mouse sera to prolong skin allografts
- Trichinella spiralis*
Feizulliaev, N. A.; Litvinov, V. P.; and Litvinov, V. F., 1977, Dokl. Akad. Nauk Azerbaidzhan. SSR, v. 33 (2), 61-62
Trichinella spiralis, prevalence in predatory mammals, wild boars sold for meat as potential source of human infection
[Meles meles]
[Canis lupus]
[Vulpes]
[Canis aureus]
[Felis chaus]
all from Kyzyl-Agach reservation

- Trichinella spiralis*
Feldman, A.; Rosenkrantz, H. S.; and Despomier, D., 1975, *J. Parasitol.*, v. 61 (3), 570-571
Trichinella spiralis, mature muscle larva, guanine-cytosine content of DNA, determined from buoyant density and thermal-helix coil transition measurements
- Trichinella spiralis*
Fernando, S. S. E.; and Denham, D. A., 1976, *J. Parasitol.*, v. 62 (6), 874-876
Trichinella spiralis, effect of mebendazole and fenbendazole on different life cycle stages, mice
- Trichinella spiralis*, *illus.*
Fontan, R.; Beauchamp, F.; and Beaver, P. C., 1975, *Bull. Soc. Path. Exot.*, v. 68 (6), 557-566
case reports of infections in humans (muscles): Laos
- Trichinella spiralis*, *illus.*
Fossati, C., 1972, *Rev. Iber. Parasitol.*, v. 32 (3-4), 181-198
Trichinella spiralis, human, extensive literature review including symptomatology, case history with infection of diaphragm presenting with respiratory symptoms
- Trichinella spiralis*, *illus.*
Gaertner, L.; et al., 1976, *Arch. Exper. Vet.-Med.*, v. 30 (2), 227-238
Trichinella spiralis, rabbits, guinea pigs, problems with use of immunofluorescence for diagnosis
- Trichinella spiralis*
Gancarz, Z., 1974, *Med. Dosw. i Mikrobiol.*, v. 26 (1-2), 157-167
Trichinella spiralis, pigs, rabbits (exper.), bentonite flocculation most successful for mass human diagnosis, comparative laboratory trials
- Trichinella spiralis*, *illus.*
Gardiner, C. H., 1976, *J. Parasitol.*, v. 62 (6), 865-870
Trichinella spiralis in mice, exact niche of adult worms during intestinal phase (epithelial layer of mucosa of small intestine, at base of villi and in glandular crypts), reproductive behavior of intestinal worms
- Trichinella spiralis*
Gentilini, M.; et al., 1976, *Nouv. Presse Med.*, v. 5 (11), 720 [Letter]
Trichinella spiralis in humans, value of indirect immunofluorescence in diagnosis during two recent epidemics in France
- Trichinella spiralis*
Gomez, V.; et al., 1974, *Rev. Iber. Parasitol.*, v. 34 (3-4), 317-322
Trichinella spiralis, human, C-reactive proteins in serum associated with non-specific reactions to haemagglutination tests
- Trichinella spiralis*
Gonzalez, H.; and Plaza, J., 1968, *Bol. Chileno Parasitol.*, v. 23 (3-4), 138-141
Trichinella spiralis, *Taenia solium*, prevalence as discovered during meat inspection at 107 abattoirs from 1963 to 1965: Chile
- Trichinella spiralis*
Gonzalez F., H.; and Plaza S., J., 1976, *Bol. Chileno Parasitol.*, v. 31 (1-2), 29-32
trichinosis, *cysticercosis*, *echinococcosis*, *fascioliasis*, survey of reasons for condemnations of swine slaughtered from 1959-1973, economic importance: Santiago, Chile
- Trichinella spiralis*
Grove, D. I.; Hamburger, J.; and Warren, K. S., 1977, *J. Infect. Dis.*, v. 136 (4), 562-570
Trichinella spiralis, kinetics of infection in mice (exper.), immunologic responses, resistance to reinfections, pathology
- Trichinella spiralis*
Grove, D. I.; Mahmoud, A. A. F.; and Warren, K. S., 1977, *J. Exper. Med.*, v. 145 (3), 755-759
Trichinella spiralis-infected mice depleted of eosinophils, no effect on spontaneous expulsion of adult worms from small intestines but numbers of larvae in muscles almost doubled, concluded that eosinophils contribute to resistance to systemic phase of trichinosis
- Trichinella spiralis*, *illus.*
Grove, D. I.; and Warren, K. S., 1976, *Ann. Trop. Med. and Parasitol.*, v. 70 (4), 449-453
Trichinella spiralis, mice, niridazole suppresses cell-mediated reactions but leaves humoral antibody formation relatively intact
- Trichinella spiralis*
Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, *J. Infect. Dis.*, v. 132 (4), 485-488
algorithms in the diagnosis and management of human *Trichinella spiralis*
- Trichinella spiralis*, *illus.*
Gudiol, F.; et al., 1974, *Med. Clin., Barcelona*, v. 63 (3), 125-137
human *Trichinella spiralis*, comprehensive resume of finding observed in 22 human infections (predominance of females, inverse relationship to alcoholic consumption, eosinophilia, changes in blood protein levels, metabolic activity of parasitized muscles, treatment with thiabendazole)
- Trichinella spiralis*
Guevara Pozo, D.; and Pinero Venegas, F., 1972, *Rev. Iber. Parasitol.*, v. 32 (3-4), 167-180
Trichinella spiralis, experimental infection of white mice with standard dose, comparison of three methods of measuring infection (direct count, artificial digestion, homogenization), statistical analysis, artificial digestion least accurate technique

- Trichinella spiralis*
Gustowska, L.; Gabryel, P.; and Zeromski, J., 1970, *Patol. Polska*, v. 21 (4), 663-669
human *Trichinella spiralis*, high specificity of immunofluorescence test demonstrated in diagnostic comparisons with muscle biopsy, biopsy recommended after obtaining positive results with immunofluorescence in order to assess degree of muscle invasion and damage
- Trichinella spiralis*
Henriksen, S. A.; and Clausen, B., 1977, *Nord. Vet.-Med.*, v. 29 (12), 543-545
Sus scrofa (muscles), rate of infection: game park in Northwest Jutland, Denmark
- Trichinella spiralis*
Horwitz, M. A.; and Hughes, J. M., 1976, *J. Infect. Dis.*, v. 134 (3), 306-312
Trichinella spiralis, *Toxoplasma gondii*, Anisakidae, implicated in human food borne diseases in United States, compiled for 1974 by Center for Disease Control, Atlanta
- Trichinella spiralis*
Howes, H. L., jr., 1972, *Proc. Soc. Exper. Biol. and Med.*, v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that *T. muris*-mouse infection could be test model for antiwhipworm studies
- Trichinella spiralis*
Hughes, J. M.; Merson, M. H.; and Pollard, R. A., jr., 1975, *J. Infect. Dis.*, v. 132 (2), 224-228
summary of data from food-borne disease outbreaks in the United States reported to the Center for Disease Control in Atlanta, Ga., in 1973, includes cases of *Trichinella spiralis*
- Trichinella spiralis*, illus.
Hughes, J. T., 1974, *Major Problems Path.*, v. 4, 122-131
general review of human trichinosis, clinical findings, epidemiology, pathology
- Trichinella spiralis*
Hughes, W. L.; and Harley, J.P., 1977, *Exper. Parasitol.*, v. 42 (2), 363-373
Trichinella spiralis, first-stage migratory larvae, taxes, response to chemicals associated with energy source for muscular contraction, response to electrical stimuli associated with membrane potentials and to KCl gradient
- Trichinella spiralis*
Jackson, G., 1977, *Brit. Vet. J.*, v. 133 (3), 318-319
Trichinella spiralis larvae, recommendation that pig muscle samples for diagnosis by peptic digestion should not be frozen
- Trichinella spiralis*, illus.
Jacobson, E. S.; and Jacobson, H. G., 1977, *Am. J. Clin. Path.*, v. 68 (6), 791-794
Trichinella spiralis in an immunosuppressed human (leukemia patient), case report, clinical presentation and autopsy findings
- Trichinella spiralis*
James, E. R., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (4), 433 [Demonstration]
Trichinella spiralis, miniaturization of technique for diagnosis using standard indirect fluorescent antibody test
- Trichinella spiralis*
James, E. R.; and Denham, D. A., 1975, *J. Helminth.*, v. 49 (1), 43-47
mice immunized to intestinal stage of *Trichinella spiralis* by drug-abbreviated infections, significant reduction in muscle larvae which encysted following normal complete challenge infection, no significant protection against challenge with parenteral stages, stage-specificity of immune response
- Trichinella spiralis*.
James, E. R.; Moloney, A.; and Denham, D. A., 1977, *J. Parasitol.*, v. 63 (4), 720-723
Trichinella spiralis, mice, immunogenicity of parenteral phase confirmed, resistance stimulated by this phase does not affect the intestinal phase
- Trichinella spiralis*
Jones, J. F.; Crandall, C. A.; and Crandall, R. B., 1976, *Cellular Immunol.*, v. 27 (1), 102-110
Trichinella spiralis, mice, T-dependent suppression of primary antibody response to sheep erythrocytes
- Trichinella spiralis*
Kazacos, K. R., 1975, *Vet. Parasitol.*, v. 1 (2), 165-174
Trichinella spiralis-immunized rats, increased resistance to *Nippostrongylus brasiliensis*, heterologous and homologous tests of immune precipitate formation on infective larvae
- Trichinella spiralis*
Kazacos, K. R., 1976, *J. Parasitol.*, v. 62 (3), 493-494
Trichinella spiralis-immunized rats, increased resistance to *Strongyloides ratti*; lack of cross-reacting precipitating antibodies in in vitro test
- Trichinella spiralis*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 1-12
meat and offal-borne anthroponotic helminthiases in Australia
- Trichinella spiralis*
Kennedy, M. W., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (4), 285 [Demonstration]
Trichinella spiralis in NIH mice, kinetics of establishment and rejection of enteral phase of primary infection
- Trichinella spiralis*
Khamboonruang, C.; and Nateewatana, N., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (1), 74-78
Trichinella spiralis, outbreak of trichinosis in 27 native villagers who had eaten raw pork as part of native dish, epidemiologic survey: Mae Sruay District, northern Thailand

- Trichinella spiralis*
Kim, C. W.; Fragola, A. C.; and Rega, R. J., 1977, *J. Parasitol.*, v. 63 (6), 1133-1135
Trichinella spiralis, low dose of antigen in combination with Freund's complete adjuvant is effective in inducing and transferring delayed hypersensitivity in the guinea pig as manifested by skin test reactions, typical histopathology and absence of circulating antibody
- Trichinella spiralis*
Kingsbury, W. D.; et al., 1976, *J. Med. Chem.*, v. 19 (6), 839-840
laboratory trials comparing parbendazole, benzimidazole and 1- and 2-substituted indazoles, *Trichinella spiralis*-infected mice and nematode-infected sheep
- Trichinella spiralis*
van Knapen, F.; Framstad, K.; and Ruitenbergh, E. J., 1976, *J. Parasitol.*, v. 62 (2), 332-333
Trichinella spiralis, reliability of enzyme-linked immunosorbent assay as control method for detection of infections in naturally infected slaughter pigs, compared with direct methods of diagnosis (trichinoscopy; digestion method) and other serological tests (immunofluorescence; counter-electrophoresis; Ouchterlony agar gel diffusion)
- Trichinella spiralis*
van Knapen, F.; Framstad, K.; and Ruitenbergh, E. J., 1976, *Tijdschr. Diergeneesk.*, v. 101 (17), 952-956
Trichinella spiralis, slaughter pigs, detection by enzyme-linked immunosorbent assay, more sensitive than other diagnostic methods
- Trichinella spiralis*
Komandarev, S.; et al., 1977, *Dokl. Bolgar. Akad. Nauk*, v. 30 (4), 567-569
Trichinella spiralis, rats (exper.), increase in peritoneal macrophages creates protection in subsequent infection by *Erysipelothrix rhusiopathiae*
- Trichinella spiralis*
Komandarev, S.; and Dragneva, N., 1973, *Izvest. Tsentral. Khelmin. Lab.*, v. 16, 127-132
Trichinella spiralis, guinea pigs, infection and superinfection, leucocytolysis in vitro
- Trichinella spiralis*
Komandarev, S.; and Mikhov, L., 1972, *Izvest. Tsentral. Khelmin. Lab.*, v. 15, 135-142
Trichinella spiralis, mice, experiments do not give evidence of intrauterine transmission of immunological tolerance or immunity
- Trichinella spiralis*
Komandarev, S.; and Mikhov, L., 1973, *Izvest. Tsentral. Khelmin. Lab.*, v. 16, 111-118
Trichinella spiralis-immunized mice, immunity to homologous and heterologous (*Ascaris suum*) challenge
- Trichinella spiralis*
Komandarev, S.; and Mikhov, L., 1974, *Izvest. Tsentral. Khelmin. Lab.*, v. 17, 81-87
Trichinella spiralis, water-soluble antigens, fractionation and characterization
- Trichinella spiralis*
Komandarev, S.; Poliakova-Krusteva, O.; and Dragneva, N., 1973, *Izvest. Tsentral. Khelmin. Lab.*, v. 16, 119-126
Trichinella spiralis, guinea pigs, infection and superinfection, plasmocyte reaction
- Trichinella spiralis*
Kozek, W. J., 1976, *Rev. Biol. Trop.*, v. 24 (1), 175-183
Trichinella spiralis, quantitative data on reproductive potential of male and female worms in mice, occurrence of multiple inseminations; presence or absence of larvae in diaphragm of a mouse can be used as an absolute criterion to confirm or rule out infection
- Trichinella spiralis* (Owen, 1835)
Kozlov, D. P., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 71-78
Vulpes vulpes
Felis catus dom.
Gulo gulo
all from Pechora river basin
- Trichinella spiralis*, illus.
Kramer, M. D.; and Aita, J. F., 1972, *Neurology*, v. 22 (5), 485-491
Trichinella spiralis in human with central nervous system involvement and unilateral rectus paresis, case report, history of pork chop consumption, thiabendazole in conjunction with prednisone: Baltimore
- Trichinella spiralis*
Larsh, J. E., jr.; and Weatherly, N. F., 1975, *Advances Parasitol.*, v. 13, 183-222
principles of delayed (cellular) hypersensitivity, cell-mediated immunity against parasitic worms, extensive review
- Trichinella spiralis*
Leenstra, F.; Elgersma, A.; and Ruitenbergh, E. J., 1977, *Trop. and Geogr. Med.*, v. 29 (2), 206 [Abstract]
Trichinella spiralis, *Hymenolepis diminuta*, infected congenitally athymic mice and their thymus-bearing heterozygous littermates (exper.), immunologic reactions shown to be dependent on immune status of host whereas non-specific histopathologic changes were thymus-independent
- Trichinella spiralis*
Levin, D. M.; et al., 1976, *Infect. and Immun.*, v. 13 (1), 27-30
Trichinella spiralis, rat model, temporal development of antigen-reactive cells in Peyer's patches and other lymphoid tissues, cellular reactivity (lymphocyte blastogenesis) not evident in Peyer's patches during earliest stages of infection
- Trichinella spiralis*
Ljungström, I., 1976, *Pathophysiol. Parasit. Infect.*, 247-253
Trichinella spiralis, mice, delay in allograft rejection

- Trichinella spiralis*
Ljungstroem, I.; and Huldt, G., 1977, Acta Path. et Microbiol. Scand., v. 85C (2), 131-141
Trichinella spiralis, mice, humoral and cellular immune responses to unrelated antigens at different stages of infection, humoral response depressed during short period of infection but depression of cell mediated response is more severe and longer lasting
- Trichinella spiralis*
Ljungström, I.; and Ruitenber, E. J., 1976, Clin. and Exper. Immunol., v. 24 (1), 146-156
Trichinella spiralis, thymectomized lethally irradiated CBA mice, immunohistological and serological response to oral infection, data support thymus dependency of host response against *Trichinella*
- Trichinella spiralis*, illus.
Londono, I., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 64-71
Trichinella spiralis, antigenic differences between larvae and adults demonstrated by development of stage-specific precipitin antibodies
- Trichinella spiralis*, illus.
Londono, I., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 72-78
Trichinella spiralis, association of antilarval and anti-adult precipitin antibodies with specific immunoglobulins, rats
- Trichinella spiralis*
Love, R. J.; and Ogilvie, B. M., 1977, Exper. Parasitol., v. 41 (1), 124-132
Nippostrongylus brasiliensis, *Trichinella spiralis*, rats, increased localization of thoracic duct lymphoblasts in small intestine, not correlated with presence of antigen but related to factors associated with inflammation, no increase in blast localization in lactating vs. nonlactating rats but increased localization in lactating mammary gland
- Trichinella spiralis*
Love, R. J.; Ogilvie, B. M.; and McLaren, D. J., 1976, Immunology, v. 30 (1), 7-15
Trichinella spiralis, duration of infections in rats (young, adult, and lactating), rapidity of expulsion from previously infected rats, transfer of immunity with antiserum and lymph node cells, ultrastructural signs of antibody damage to worms, results suggest that mechanism of immune expulsion requires both antibody and cells, comparison with *Nippostrongylus brasiliensis*
- Trichinella spiralis*
Lubiniecki, A. S.; and Cypess, R. H., 1975, Tropenmed. u. Parasitol., v. 26 (3), 329-333
Trichinella spiralis in mice (exper.), effects of helminth infections on tumor development and variations in host immune reactions
- Trichinella spiralis*
Lubiniecki, A. S.; Cypess, R. H.; and Lucas, J. P., 1974, Tropenmed. u. Parasitol., v. 25 (3), 345-349
Trichinella spiralis-infected mice, altered distribution of intravenously injected sheep erythrocytes (reduced uptake by spleen) but no significant depression of resultant plaque-forming cells or circulating HA antibody
- Trichinella spiralis*
Luengo, J.; and Barriga, O. O., 1966, Bol. Chileno Parasitol., v. 21 (1), 2-7
Trichinella spiralis in *Rattus norvegicus* (exper.), high single dose of thiabendazole effective on enteral infection but showed little effect on migrating larvae and no effect on encysted parasites, increased weight gain in treated rats
- Trichinella spiralis*
Matossian, R. M.; Rebeiz, J.; and Stephan, E., 1975, Ann. Trop. Med. and Parasitol., v. 69 (3), 387-392
Trichinella spiralis, human, early diagnosis of acute trichinosis can be made by indirect fluorescent antibody test using freeze-dried larval antigen, human serum, and anti-human IgG, IgM, and IgA fluorescein conjugates
- Trichinella spiralis*
Matossian, R. M.; Salti, I.; and Stephan, E., 1977, J. Helminth., v. 51 (1), 1-4
Trichinella spiralis, human, time course development of serum immunoglobulin levels
- Trichinella spiralis*
Meerovitch, E.; and Ackerman, S. J., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (5), 417 [Letter]
pre-existing *Trichinella spiralis* infection in rats (exper.) greatly reduced by level of *Trypanosoma lewisi* parasitemia, possible cell-mediated immune reaction
- Trichinella spiralis*
Meerovitch, E.; and Bomford, R., 1977, Ann. Trop. Med. and Parasitol., v. 71 (2), 245-247
Trichinella spiralis, potentiator of macrophage-mediated immunity
- Trichinella spiralis*
Menard, E.; et al., 1973, Bol. Chileno Parasitol., v. 28 (3-4), 73-77
Trichinella spiralis outbreak in 13 of 19 persons of a religious group, had eaten pork not raised and slaughtered under sanitary conditions: Chile
- Trichinella spiralis*
Mikhail, E. G.; and Milad, M., 1975, Med. J. Cairo Univ., v. 43 (1), 65-71
exper. *Trichinella spiralis* myocarditis, rats, pathologic findings, occurs during migration phase of infection during first two weeks of disease, generally no permanent damage

- Trichinella spiralis*
Molinari, J. A.; Cypess, R. H.; and Appel, B. N., 1975, Internat. Arch. Allergy and Applied Immunol., v. 48 (6), 776-783
Trichinella spiralis- and/or BCG-infected mice, histopathologic changes in thymus, possible relation to functional alterations of immune system
- Trichinella spiralis*
Molinari, J. A.; and Ebersole, J. L., 1976, Abst. Ann. Meet. Am. Soc. Microbiol., 84
Trichinella spiralis, mice, nematode induced potentiation of delayed hypersensitivity, induces stimulation of host anti-neoplastic activity
- Trichinella spiralis*
Molinari, J. A.; and Ebersole, J. L., 1977, Internat. Arch. Allergy and Applied Immunol., v. 55 (1-6), 444-448
Trichinella spiralis, mice, antineoplastic effects of long-term infection on B-16 melanoma, apparently related to potentiation of cellular immune response
- Trichinella spiralis*
Molinari, J. A.; and Ebersole, J. L., 1977, Internat. Arch. Allergy and Applied Immunol., v. 55 (1-6), 449-457
Trichinella spiralis-induced immunopotential of delayed-type hypersensitivity reactions to BCG, in vitro responses of spleen cells from infected mice
- Trichinella spiralis*
Motyka, S., 1976, Bull. Acad. Vet. France, v. 49 (1), 95-99
Trichinella spiralis, human outbreak probably caused by eating imported horse meat: region of Paris, France
- Trichinella spiralis*
Munoz, J. J.; and Cole, R. L., 1977, Infect. and Immun., v. 15 (1), 84-90
Trichinella spiralis-infected mice, relative unresponsiveness to passive cutaneous anaphylaxis induced with hen egg albumin and its corresponding antibodies, believed to be due to increase in production of IgE which competitively blocks mast cell sites for other IgE molecules
- Trichinella spiralis*
Muzquiz Moracho, J. L., 1976, Veterinaria, Barcelona, v. 41 (10-12), 161-163
Trichinella spiralis, epidemiological study, not found in survey of 100 dogs, Oct. 1974-May, 1975: Zaragoza (Espana)
- Trichinella spiralis*
Neal, W. P., 1971, Maternal and Child Health, v. 3 (12), 38-39
persistent eosinophilia, diagnostic in suspected human parasitic infestations
- Trichinella spiralis*
Ngwenya, B. Z., 1976, Cellular Immunol., v. 24 (1), 116-122
Trichinella spiralis, effect of lactation on cell-mediated immunity, cell transfer studies with lactating and non-lactating mice, lactogenic hormones apparently suppressed expression of adoptive immunity
- Trichinella spiralis*
Ngwenya, B. Z., 1976, J. Parasitol., v. 62 (6), 871-873
Trichinella spiralis, suppression of rejection in mice treated with ovine prolactin
- Trichinella spiralis*
Ngwenya, B. Z.; 1977, Internat. J. Parasitol., v. 7 (1), 41-45
Trichinella spiralis, effect of lactation on worm expulsion in (1) lactating, (2) induced agalactic post-parturient, (3) previously sensitized, and (4) mice sensitized during lactation, results indicate lactation suppressed either expression or induction of immunity
- Trichinella spiralis*
Nitsche, W., 1973, Munchen. Med. Wchnschr., v. 115 (5), 141-146
Trichinella spiralis in humans, dangers of being infected during journeys to areas without meat inspection laws, control through prophylaxis: Germany
- Trichinella spiralis*
Norman, L.; and Kagan, I. G., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 58-64
Trichinella spiralis, rabbit and human serum, evaluation of crude and fractionated antigens and comparison of effectiveness of serologic tests for diagnosis
- Trichinella spiralis*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Sus scrofa: Chile
- Trichinella spiralis*
Orlandi, V., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 482-484
Trichinella spiralis, incidence in foxes and badgers: Ascoli Piceno province
- Trichinella spiralis*
Orlov, I. V.; Britov, V. A.; and Boev, S. N., 1976, Vestnik Sel'skokhoz. Nauki (243) (12), 61-68
Trichinella spp., experimental hybridization between species shows very limited crossing, reproductive isolation; useful technique for species diagnosis
- Trichinella spiralis*
Ossola, A.; et al., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 123-127
Trichinella spiralis outbreak in 31 persons, epidemiologic survey, meat from 3 pigs implicated as source: Mercedes, Chile

- Trichinella spiralis*
 Ottesen, E. A.; Smith, T. K.; and Kirkpatrick, C. H., 1975, Internat. Arch. Allergy and Applied Immunol., v. 49 (3), 396-410
Trichinella spiralis, mice, chronic infection, cellular immune responsiveness, sequential development of antigen-reactive cells in various lymphoid cell populations, antibody responses (haemagglutination titers, homocytotropic antibody)
- Trichinella spiralis*
 Panitz, E., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 257-259
 adult *Trichinella spiralis*, distribution in intestine of *Meriones unguiculatus*, immunosuppressive effect of betamethasone
- Trichinella spiralis*, illus.
 Perez, C.; and Luengo, J., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 163
Trichinella spiralis, nine larvae demonstrated in single cyst of pig muscle
- Trichinella spiralis*
 Perrudet-Badoux, A.; and Binaghi, R. A., 1977, Ann. Immunol., v. 128C (1-2), 243-244
Trichinella spiralis, antibody-mediated adherence of rat peritoneal exudate cells to larvae
- Trichinella spiralis*
 Perrudet-Badoux, A.; and Binaghi, R. A., 1977, Compt. Rend. Acad. Sc., Paris, v. 284, s. D (6), 497-498
Trichinella spiralis, antibody-mediated adherence of peritoneal exudate cells to larvae
- Trichinella spiralis*
 Perrudet-Badoux, A.; and Binaghi, R. A., 1977, Immunology, v. 33 (6), 881-885
Trichinella spiralis, rats, IgE antibodies not transferred from mother to young during lactation nor during pregnancy although they are sometimes secreted in the milk
- Trichinella spiralis*
 Perrudet-Badoux, A.; Binaghi, R. A.; and Bous-sac-Aron, Y., 1976, Immunochemistry, v. 13 (5), 443-445
Trichinella spiralis-infected rats, quantitative study of production of different classes of immunoglobulins
- Trichinella spiralis*
 Perrudet-Badoux, A.; Binaghi, R. A.; and Bous-sac-Aron, Y., 1977, Ann. Immunol., v. 128C (4-5), 965-966 [Abstract]
Trichinella spiralis, rats, enhanced antibody response to unrelated antigen
- Trichinella spiralis*
 Pinero Venegas, F.; and Guevara Pozo, D., 1973, Rev. Iber. Parasitol., v. 33 (1), 3-10
Trichinella spiralis, differences in susceptibility of two strains of laboratory mice measured by survival time of mice and degree of muscular infestation
- Trichinella spiralis* (Owen, 1835)
 Podhajecky, K.; Spaldonova, R.; and Tomasovicova, O., 1974, Biologia, Bratislava, s. B, Zool. (1), v. 29 (2), 163-166
Trichinella spiralis in mice given intravenous calcium, development in and duration of intestinal phase unaffected, moderate retardation of encapsulation process in muscle phase, small increase in worm reproductive capacity
- Trichinella spiralis*
 Podhajecky, K.; and Tomasovicova, O., 1976, Biologia, Bratislava, s. B, Zool. (3), v. 31 (8), 595-599
Trichinella spiralis, reproductive potential in relation to size of infecting dose and to season of year, mice
- Trichinella spiralis*
 Poirot, J. L.; et al., 1976, Ann. Parasitol., v. 51 (4), 447-452
Trichinella spiralis, mice, effect of cyclophosphamide treatment on parasite sex ratio and survival, transplantation of 18-day old worms from treated vs. untreated into treated vs. untreated mice
- Trichinella spiralis*
 Portaro, J. K.; et al., 1977, J. Parasitol., v. 63 (1), 172-174
 differential response of *Brugia pahangi*-sensitized splenocytes to antigens from *Brugia pahangi*, *Dirofilaria immitis*, and *Trichinella spiralis*, possible diagnostic use
- Trichinella spiralis*
 Potter, M. E.; et al., 1976, Am. J. Pub. Health, v. 66 (12), 1194-1196
Trichinella spiralis, outbreak of trichinosis in 23 of 50 persons who had eaten home-made summer sausage; economic importance of this avoidable infection and need for nationwide program of control of trichinosis in swine: Illinois
- Trichinella spiralis*
 Przyjałkowski, Z.; et al., 1976, Bull. Acad. Polon. Sc., Cl. II., s. Sc. Biol., v. 24 (12), 751-755
Trichinella spiralis, antibody response of germfree, gnotobiotic, and conventional mice compared
- Trichinella spiralis*, illus.
 Przyjałkowski, Z.; and Warton, A., 1977, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (11), 1976, 675-678
Trichinella spiralis in bacteria-free or conventional mice, changes in ultrastructure of epithelium of small intestines
- Trichinella spiralis*
 Puccini, V., 1975, Vet. Ital., v. 26 (9-12), 378-392
Trichinella spiralis, dogs, incidence, epidemiology: province of Foggia, Italy

- Trichinella spiralis*
Pullen, M. M.; Seymour, M. R.; and Zimmermann, W. J., 1977, J. Am. Vet. Med. Ass., v. 171 (11), 1171-1172
prevalence in diaphragmatic samples of sows slaughtered at abattoir: Kentucky (marketed from Kentucky, Ohio and Indiana)
- Trichinella spiralis*
Ramisz, A.; and Komorowski, A., 1975, Polskie Arch. Wet., v. 17 (4), 623-631
Trichinella spiralis, muscle phase in mice, fenchlorphos and bromophos, inhibition of host cholinesterase activity in motor end plates; increased activity of cholinergic system as main factor in pathogenesis
- Trichinella spiralis*, illus.
Ramisz, A.; and Szankowska, Z., 1976, Folia Histochem. et Cytochem., v. 14 (4), 321-326
Trichinella spiralis, mice, administration of paraoxon under the protection of toxobidin (an acetylcholinesterase reactivator), effect on cholinesterases in host skeletal muscles
- Trichinella spiralis*
Rees, P. H.; and Marsden, P. D., 1970, Brit. J. Clin. Pract., v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment
- Trichinella spiralis*
Rivera-Ortiz, C.-I.; and Nussenzweig, R., 1976, Exper. Parasitol., v. 39 (1), 7-17
Trichinella spiralis, differential ability of several inbred mouse strains of different histocompatibility locus specificities to produce reagin and IgG, antibodies in response to infection, relationship between production of anaphylactic antibodies and larval and adult recoveries, stage of life cycle which induces antibody formation
- Trichinella spiralis*
Roberts, T. M.; and Thorson, R. E., 1977, J. Parasitol., v. 63 (4), 764-766
Nippostrongylus brasiliensis, pairing between adults and other species of nematodes in vitro, comparison of intra and interspecific response patterns, results suggest that pairing is not related entirely to sexual attraction
- Trichinella spiralis*
Rodriguez, M.; Gomez, V.; and Gonzalez, J., 1974, Rev. Iber. Parasitol., v. 34 (3-4), 253-256
survey of hospital outpatients by indirect immunofluorescence test, three positive results out of 499 samples, absence of eosinophilia or other symptoms: Provincia de Granada
- Trichinella spiralis*
Rose, M. L.; Parrott, D. M. V.; and Bruce, R. G., 1976, Immunology, v. 31 (5), 723-730
Trichinella spiralis, syngeneic mice, migration of mesenteric lymphoblasts and mesenteric T lymphoblasts at various times after infection, enhanced accumulation in small intestine at days 2 and 4
- Trichinella spiralis*
Ruitenbergh, E. J., 1973, Norwegian J. Zool., v. 21 (4), 326 [Abstract]
Trichinella spiralis in rats (exper.), immune response directed towards intestinal phase; *Corynebacterium parvum* sensitization prolonged expulsion
- Trichinella spiralis*
Ruitenbergh, E. J., 1977, Trop. and Geogr. Med., v. 29 (3), 319 [Abstract]
Trichinella spiralis-infected nude mice, failure of infections to induce gut mast cell response; both gut and blood eosinophils increased during infection, the phenomenon being T-cell dependent
- Trichinella spiralis*
Ruitenbergh, E. J.; et al., 1976, Tijdschr. Diergeneesk., v. 101 (2), 57-70
Trichinella spiralis, conventionally raised pigs with experimental infections at various doses, enzyme-linked immunosorbent assay more sensitive than immunofluorescence test, further ways to improve reliability
- Trichinella spiralis*
Ruitenbergh, E. J.; et al., 1977, Immunology, v. 33 (4), 581-587
Trichinella spiralis, comparison of infection in congenitally athymic (nude) mice and their heterozygous thymus-bearing littermates: expulsion of adult worms; yield of muscle larvae; production of specific antibodies; number of pyroninophilic cells, intra-epithelial lymphocytes, and eosinophils in small intestine; blood eosinophilia; data support thymus dependence of worm expulsion, plasma cell and antibody production, and tissue and blood eosinophilia
- Trichinella spiralis*
Ruitenbergh, E. J.; and Elgersma, A., 1976, Nature (5583), v. 264, 258-260
Trichinella spiralis, congenitally athymic mice, absence of intestinal mast cell response, results indicate T-cell dependence of host protection
- Trichinella spiralis*
Ruitenbergh, E. J.; and van Knapen, F., 1977, J. Infect. Dis., v. 136, suppl., S267-S273
application of the enzyme-linked immunosorbent assay to the detection of human and animal helminthic and protozoal infections, advantages of assay for seroepidemiology, discussion of performance of assay in tubes and microplates

- Trichinella spiralis*
Ruitenber, E. J.; and van Knapen, F., 1977, *Vet. Parasitol.*, v. 3 (4), 317-326
Trichinella spiralis, pigs, enzyme-linked immunosorbent assay as diagnostic method, comparison with conventional digestion method
- Trichinella spiralis*
Ruitenber, E. J.; Leenstra, F.; and Elgersma, A., 1977, *Brit. J. Exper. Path.*, v. 58 (3), 311-314
Trichinella spiralis, athymic mice, intestinal pathology; immunological reaction dependent upon host immune status, non-specific histopathological changes thymus-independent
- Trichinella spiralis*
Ruitenber, E. J.; and Steerenberg, P. A., 1976, *J. Parasitol.*, v. 62 (1), 164-166
Trichinella spiralis, rats, immunization with newborn larvae, challenge with newborn larvae, significant decrease in yield of muscle larvae in immunized challenged rats compared with challenged control rats
- Trichinella spiralis*
Ruitenber, E. J.; Teppema, J. S.; and Steerenberg, P. A., 1974, *Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol.* (Vienna, Austria, Sept. 18-20, 1973), 319-326
Trichinella spiralis intestinal phase, no evidence found for immunity-induced changes in enzyme histochemical staining pattern of adult worms, detection of antigen or of antigen-antibody complexes using bridge immuno-peroxidase anti-peroxidase technique, immunoglobulins found around cuticle of adult worms even in ATS-treated animals
- Trichinella spiralis*
Sadun, E. H.; Williams, J. S.; and Gore, R. W., 1973, *Isotopes and Radiation Parasitol.* III, 73-90
Schistosoma mansoni, *S. haematobium*, *Trichinella spiralis*, development of radioactive antigen microprecipitin assay (RAMP), comparison with soluble antigen fluorescent antibody and passive cutaneous anaphylaxis tests, results indicate RAMP measures antibody primarily of I_gE class
- Trichinella spiralis*
Sagua, H.; et al., 1972, *Bol. Chileno Parasitol.*, v. 27 (1-2), 58-60
eosinophilia, intradermal, ring precipitation and bentonite flocculation tests, comparison of results in persons at time of *Trichinella spiralis* infection and 8 1/2 years after infection
- Trichinella spiralis*
Saowakontha, S., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (1), 37-40
Trichinella spiralis, absorptive functions of intestines of infected rats on low and high protein diets were not impaired either in early or late intestinal phases of infections as tested by D-xylose absorption
- Trichinella spiralis*
Saowakontha, S., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (4), 586-592
Trichinella spiralis, rats (exper.), effects of different levels of protein in diet on numbers of intestinal worms and muscular larvae and on intensity of infections in hosts of varying weights and ages
- Trichinella spiralis*
Saowakontha, S., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (1), 79-81
Trichinella spiralis, no differences between number of worms in immunized rats (exper.) fed low or high protein diets, significantly lower number of worms in immunized rats compared with rats not immunized
- Trichinella spiralis*
Sapunar, J.; and Szekely, R., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 31-36
Trichinella spiralis, analysis of clinical findings in 76 persons with trichinosis, symptoms, pathology, findings possibly useful as prognosis index
- Trichinella spiralis*
Saunders, G. C.; et al., 1977, *J. Infect. Dis.*, v. 136, suppl., S258-S266
Trichinella spiralis, application of the indirect enzyme-labeled antibody microtest for the detection of trichinosis in swine, experimental trials testing the feasibility for use in meat packinghouses
- Trichinella spiralis*
Saunders, G. C.; and Clinard, E. H., 1976, *J. Clin. Microbiol.*, v. 3 (6), 604-608
Trichinella spiralis, swine, rapid micro-method for mass screening for antibodies using indirect enzyme-labeled antibody test, high number of false positives
- Trichinella spiralis*
Schantz, P. M.; Juranek, D. D.; and Schultz, M. G., 1977, *J. Infect. Dis.*, v. 136 (5), 712-716
Trichinella spiralis, humans, increased incidence of infections reported in 1975, statistics of epidemiologic study: United States
- Trichinella spiralis*
Schenone, H.; et al., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 2-10
Trichinella spiralis, mild outbreak affecting 36 persons, exact etiology unknown but hogs feeding on garbage dump which also had abundant *T. spiralis*-infected rat population believed to be source: Antofagasta, Chile
- Trichinella spiralis, illus.*
Schenone, H.; et al., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 32-37
heavy concurrent fatal infection of *Trichinella spiralis* and *Taenia solium* in 17-year-old boy with history of eating raw pork, others in family less heavily infected, clinical case report: Santiago, Chile

- Trichinella spiralis*
Schenone, H.; et al., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 118-121
Trichinella spiralis, human, skin test and bentonite flocculation test, surveys show progressive decline in infection: Santiago, Chile
- Trichinella spiralis*
Schenone, H.; et al., 1972, Bol. Chileno Parasitol., v. 27 (1-2), 33-36
trichinosis reported in 49 persons who had eaten insufficiently cooked pork, clinical findings, serologic diagnosis: Temuco, Chile
- Trichinella spiralis*
Schenone, H.; et al., 1972, Bol. Chileno Parasitol., v. 27 (3-4), 103-107
Trichinella spiralis, humans, prevalence survey, evidence of increasing infection: Chile
- Trichinella spiralis*
Schenone, H.; et al., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 107-108
heterogenetic relationship between *Trichinella spiralis* and typhoid and paratyphoid antigenic fractions
- Trichinella spiralis*
Schenone, H.; and Reyes, H., 1968, Bol. Chileno Parasitol., v. 23 (1-2), 62-65
statistics of survey of 41,799 autopsies for evidence of previously undiagnosed echinococcosis, cysticercosis, and *Trichinella spiralis*: Chile
- Trichinella spiralis*
Schenone, H.; Reyes, H.; and Rojas, A., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 152-153
Trichinella spiralis, 1000 autopsy surveys indicate decreasing incidence in humans probably resulting from improved meat inspection: Santiago, Chile
- Trichinella spiralis*
Schmitt, N.; et al., 1976, Canad. J. Pub. Health, v. 67 (1), 21-24
incidence survey of infection in various wild mammal species as possible reservoir for human trichinosis
Ursus arctos
U. americanus
Canis latrans
Martes americana
Lynx canadensis
Mephitis mephitis
Felis concolor
Lynx rufus
Mustela frenata
Gulo luscus
Sorex vagrans
Peromyscus maniculatus
Spermophilus columbianus
Tamiasciurus hudsonicus
dog
all from Kootenay areas of British Columbia
- Trichinella spiralis* (Owen, 1835)
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Mustela vison
Martes martes
[Vulpes vulpes]
[Canis lupus]
all from Karelia
- Trichinella spiralis*
Singal, M.; Schantz, P. M.; and Werner, S. B., 1976, Am. J. Trop. Med. and Hyg., v. 25 (5), 675-681
Trichinella spiralis, report of outbreak in passengers of pleasure cruise, ground beef contaminated by pork implicated as transmission vehicle: luxury liner en route to Alaska
- Trichinella spiralis*
Sluifers, J. F.; and Ruitenbergh, E. J., 1974, Nederl. Tijdschr. Geneesk., v. 118 (7), 230-233
Trichinella spiralis, incidence survey shows evidence of larvae in fattening pigs: Netherlands
- Trichinella spiralis*
Smith, H. J., 1976, J. Wildlife Dis., v. 12 (2), 256-257
investigation of North Atlantic whales, negative results
- Trichinella spiralis*
Spaldonova, R.; et al., 1973, Izvest. Tsentral. Khelmint. Lab., v. 16, 197-203
Trichinella spiralis, white mice, effect of aloxan diabetes on dynamics of intestinal trichinosis and intensity of muscular infection, hyperglycaemia may inhibit allergic reaction of host to infection
- Trichinella spiralis*
Spaldonova, R.; and Corba, J., 1977, Biologia, Bratislava, s. B, Zool. (4), v. 32 (11), 855-860
Trichinella spiralis, intestinal stages, mice, efficacy of cambendazole, decreases as worms mature, females more susceptible
- Trichinella spiralis*
Spry, C. J. F., 1972, Immunology, v. 22 (4), 663-675
Trichinella spiralis, rats, origin, recirculation kinetics, and distribution of large lymphocytes from thoracic duct, no definite conclusions on mechanism by which large lymphocytes in rats with trichinosis stimulate eosinopoiesis
- Trichinella spiralis*
Stankiewicz, M.; and Jeska, E. L., 1973, Immunology, v. 26 (1), 827-834
Trichinella spiralis, cell adherence reactions to infective larvae, importance of heat labile and heat stable substances in peritoneal exudate fluid

- Trichinella spiralis*
Steele, J. H.; and Arambulo, P. V. III, 1975, Internat. J. Zoonoses, v. 2 (2), 55-75
Trichinella spiralis, distribution of trichinosis throughout the world, extensive sylvatic reservoirs, review
- Trichinella spiralis*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (26), 123-128
[pig, wild]: Primorskii krai
- Trichinella spiralis*
Sturrock, R. F.; et al., 1977, Parasitology, v. 75 (1), 89-100
Papio anubis, eosinophilia following oral infection with *Trichinella spiralis*, eosinophilia following intravenous administration of *Trichinella spiralis* and the effect on subsequent exposure to *Schistosoma mansoni*, latter appears to be suitable method of experimental induction of non-specific eosinophilia to further investigate possible immune mechanisms to *Schistosoma mansoni* in the baboon
- Trichinella spiralis*
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcvi
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Trichinella spiralis*
Tomasovicova, O., 1975, Biologia, Bratislava, s. B, Zool., v. 30 (11), 821-826
experimental infection of chickens, *Gallus gallus dom[esticus]*, infection of mice with *Trichinella pseudospiralis* from chickens, viability and infectivity maintained; experimental infection of chickens with *T. spiralis* only of brief duration; possibility that wild and domestic birds actively transfer and disseminate *T. pseudospiralis* but act only as passive hosts of intestinal stage *T. spiralis*
- Trichinella spiralis* (Owen, 1835)
Tomasovicova, O.; and Spaldonova, R., 1974, Biologia, Bratislava, s. B, Zool. (1), v. 29 (2), 159-162
Trichinella spiralis, intestinal phase in mice having experimental diabetes, longer worms than in normal mice but development not affected
- Trichinella spiralis*
Toshkov, A.; et al., 1977, Dokl. Bolgar. Akad. Nauk, v. 30 (6), 895-896
Trichinella spiralis-infected rats followed by infection with *Erysipelothrix rhusiopathiae*, effect of ACTH on defence mechanisms is counteracted by *T. spiralis* (inhibition of non-specific protective factors)
- Trichinella spiralis*
Wakelin, D.; and Lloyd, M., 1976, Parasitology, v. 72 (2), 173-182
Trichinella spiralis, young and older NIH strain mice, dynamics of establishment and expulsion of primary and challenge infections, parameters of immunity must be established for each host strain
- Trichinella spiralis*
Wakelin, D.; and Lloyd, M., 1976, Parasitology, v. 72 (3), 307-315
Trichinella spiralis, mice given mesenteric lymph node cells or serum or both from infected donors, acceleration of worm expulsion
- Trichinella spiralis*
Wakelin, D.; and Wilson, M. M., 1977, Parasitology, v. 74 (3), 215-224
Trichinella spiralis, mice, transfer of immunity with mesenteric lymph node cells: time of appearance of effective cells in donors; expression of immunity in recipients (worm expulsion and impaired worm reproduction may represent independent aspects of immune response)
- Trichinella spiralis*
Wakelin, D.; and Wilson, M. M., 1977, Parasitology, v. 74 (3), 225-234
Trichinella spiralis, mice, inhibition of worm expulsion by host irradiation, attempts at reconstitution of immune response gave evidence for involvement of bone marrow-derived cell population in immune expulsion
- Trichinella spiralis*
Wakelin, D.; and Wilson, M. M., 1977, Parasitology, v. 75 (2), xiv [Abstract]
Trichinella spiralis, expulsion from mice appears to be dependent on cooperation of immune mesenteric lymph node cells and a bone marrow cell component
- Trichinella spiralis*
Walls, R. S.; and Beeson, P. B., 1972. Clin. and Exper. Immunol., v. 12 (1), 111-119
Trichinella spiralis, rats, findings suggest that eosinophilia characteristic of macro-parasitic infestations is related to character of local inflammatory reaction excited by parasites in organs which harbor them
- Trichinella spiralis*
Warren, K. S.; et al., 1976, J. Infect. Dis., v. 134 (3), 277-280
Trichinella spiralis, human, murine, eosinophil stimulation promoter test, aid in diagnosis, specificity established by lack of cross-reactions with *Schistosoma mansoni*
- Trichinella spiralis*
Williams, L. P., jr.; Nelson, C. B.; and Zymet, C. L., 1969, Minnesota Med., v. 52 (7), 1153-1158
Trichinella spiralis outbreak involving 37 persons, infection apparently from consumption of summer sausage all had obtained from local meatpacking source: Minnesota; Iowa

- Trichinella spiralis*
Wing, E. J.; et al., 1977, Nature, London (5621), v. 268, 642-644
macrophages activated by infection with *Toxoplasma gondii* or *Trichinella spiralis*, ability to inhibit tumor cell DNA synthesis and to inhibit intracellular multiplication of *Toxoplasma gondii*, results show that macrophages characterized as activated by one criterion may not satisfy other criteria of activation and that differences in functional capacity depend on the method used to activate the macrophages
- Trichinella spiralis*
Young, E.; and Whyte, I. J., 1975, J. South African Vet. Ass., v. 46 (3), 233-234
Trichinella spiralis, wildlife, epizootiology, review; first case in *Viverra civetta*: Kruger National Park, South Africa
- Trichinella spiralis*
Zander, B.; and Hoerchner, F., 1976, Ztschr. Parasitenk., v. 50 (2), 178
Trichinella spiralis, guinea pigs, serum titration before and after mebendazole treatment
- Trichinella spiralis*
Zapart, W.; Adonajło, A.; and Gancarz, Z., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 9-19
Trichinella spiralis, sera from infected humans, evaluation of fractionated antigens, complement fixation and ring precipitation tests
- Trichinella spiralis*, illus.
Zeromski, J.; and Jazbor, A., 1969, Acta Parasitol. Polon., v. 17 (1-19), 119-125
Trichinella spiralis, rats, localization of antigens by direct and indirect immunofluorescence, methods compared, antigens found only within developing parasite, probably immunological mechanisms not active in production of clinical symptoms
- Trichinella spiralis*, illus.
Zeromski, J.; and Jazbor, A., 1969, Acta Parasitol. Polon., v. 17 (1-19), 127-130
Trichinella spiralis, rats, immunofluorescent fixation of heterologous complement by larval antigens; antilarval antibodies apparently present in host serum very early
- Trichinella spiralis*
Zimmermann, W. J., 1977, Am. J. Epidemiol., v. 106 (2), 167-171
Trichinella spiralis in *Ursus americanus*, age prevalence rates, intensity of infection, public health hazard: California; Idaho; Wisconsin
- Trichinosis, illus.
Andy, J. J.; et al., 1977, Am. J. Med., v. 63 (5), 824-829
trichinosis, acute fatal infection in woman who at autopsy was found to have extensive ventricular endocardial damage with superimposed thrombosis, was known to have eaten raw meat frequently, clinical and morphologic report, review of other autopsies for other trichinosis-associated heart involvements
- Trichinosis
Artemenko, Iu. G., 1976, Veterinariia, Kiev (43), 84-86
trichinosis, comparative infectivity of domestic (from domestic swine) and wild (from Siberian bears) strains to experimental swine, polar foxes, foxes, mink and cats
- Trichinosis
Bessonov, A. S., 1974, Proc. 6. Internat. Conf. World Ass. Adv. Vet. Parasitol. (Vienna, Austria, Sept. 18-20, 1973), 179-186
perspectives on eradication of several helmintho-zoonotic diseases in the USSR
- Trichinosis
Buendia, E.; et al., 1974, Nouv. Presse Med., v. 3 (36), 2334 [Letter]
human trichinosis, alterations of blood proteins in presence of infections
- Trichinosis
Burgos, H., 1973, Bol. Chileno Parasitol., v. 28 (1-2), 37-38
echinococcosis, cysticercosis, fascioliasis and trichinosis prevalence in livestock slaughtered in abattoirs: Bio-Bio Province, Chile
- Trichinosis, illus.
Dickel, H., 1977, Prakt. Tierarzt, v. 58 (2), 92, 94, 96
trichinosis, hunting dog probably infected by wild swine, injection of neguvon, good results; incidence in wild and domestic swine: Germany
- Trichinosis
Ferencz, A.; et al., 1972, Orvosi Hetilap, v. 113 (52), 3194-3196, 3199
human trichinosis, electrocardiographic changes during infection
- Trichinosis, illus.
Gabryel, P.; and Gustowska, L., 1970, Patol. Polska, v. 21 (1), 55-67
human trichinosis, morphologic and histochemical analysis of pathologic changes in diagnostic muscle biopsy material
- Trichinosis
Hoerning, B., 1977, Schweiz. Arch. Tierh., v. 119 (8), 337-339
raw horse meat as a probable source of human trichinosis: Italy and France
- Trichinosis
Kim, C. W., 1975, Progr. Clin. Path., v. 6, 267-288
extensive review of techniques used to diagnose human parasitic diseases
- Trichinosis
Klimowicz, J.; Gancarz, Z.; and Wyrzykowski, J., 1975, Przegl. Lek., v. 32 (12), 876-878
human trichinosis, use of immunofluorescence and passive hemagglutination tests for diagnosis and for epidemiologic surveys

- Trichinosis**
Kozakiewicz, B., 1972, *Med. Wet.*, v. 28 (9), 563-565
modified trichinoscope for parasitological examination
- Trichinosis**
Lupi, D.; et al., 1975, *Med. Arh.*, v. 29 (6), 605-608
clinical aspects of small epidemic of human trichinosis in rural village, diagnostic measures, successful treatment with mintage: Lipovica, Yugoslavia
- Trichinosis**
Mantovani, A.; et al., 1976, *Bull. Acad. Vet. France*, v. 49 (2), 213-222
human trichinosis, source of infections appears to be contaminated horsemeat imported from Poland or Yugoslavia, case report: Bagnolo in Piano (Reggio Emilia), Italy
- Trichinosis**
Puelma, E.; et al., 1970, *Bol. Chileno Parasitol.*, v. 25 (3-4), 140-142
epidemiologic survey using immunologic methods to ascertain incidence of echinococcosis, cysticercosis, trichinosis, fascioliasis and trypanosomiasis in mining town of Sewell, Chile
- Trichinosis**
Schenone, H.; et al., 1970, *Bol. Chileno Parasitol.*, v. 25 (1-2), 46-51
epidemiologic survey shows increasing incidence of human trichinosis, possible association with increased pork consumption and lack of satisfactory meat inspection: Santiago Province, Chile
- Trichinosis**
Smith, H. J.; Anzengruber, A.; and DuPlessis, D. M., 1976, *Canad. Vet. J.*, v. 17 (3), 72-75
incidence of trichinosis in swine, field and laboratory investigations from 1968-1975: abattoirs in Atlantic provinces, Canada
- Trichinosis**
Snobl, A.; and Kuklik, K., 1976, *Veterinarstvi*, v. 26 (6), 272-273
trichinosis, examination of swine meat under large-scale conditions by digestive method
- Trichinosis**
Szarfman, A.; et al., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (1), 19-24
evaluation of specificity of EVI factor observed in *Trypanosoma cruzi* cases using indirect fluorescent antibody test on sera from patients with other parasitic diseases, immunofluorescence of anti-skeletal muscle antibody from leishmaniasis
- Trichinosis**
Szekely, R., 1972, *Bol. Chileno Parasitol.*, v. 27 (3-4), 108-114
application of intradermal skin tests to diagnose human parasites, advantages and limitations, review of techniques
- Trichinosis**
Vliegen, J.; Froescher, W.; and Gullotta, F., 1976, *Nervenarzt*, v. 47 (7), 439-443
human trichinosis, amyotrophic lateral sclerosis and other neurologic complications resulting from parasite infection, case reports: Germany
- Trichinosis**
Walther, G.; Fasl, H.; and Kauer, E., 1971, *Med. Welt*, v. 22 (25), 1041-1044
trichinosis, human epidemic, case report of fatal illness resulting from associated myocarditis: Germany
- Trichinosis**
Wojcik, A. R.; and Grzywinski, L., 1975, *Medycyna Wet.*, v. 31 (10), 597-598
incidence in swine, economic losses: Torun slaughterhouse
- Trichinosis**
Zambeli, M., 1975, *Med. Arh.*, v. 29 (6), 627-629
report of small epidemic of trichinosis in Yugoslavia
- Trichobaylisia Travassos, 1937**
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
as syn. of *Heligmonina Baylis*, 1928
- Trichocephalata Skryabin & Schul'ts, 1928**
Maggenti, A. R., 1976, *Organ. Nematodes (Croll)*, 1-10
Trichocephalida
- Trichocephales**
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Trichocephaliasis. See Trichuriasis.**
- Trichocephalida Skryabin & Schul'ts, 1938**
Maggenti, A. R., 1976, *Organ. Nematodes (Croll)*, 1-10
Enoplia
includes: Trichocephalata; Diotophymatina
- Trichocephalus**
Bulucea, D., 1974, *Med. Int.*, Bucuresti, v. 26 (2), 249-256
ascariasis and Trichocephalus as causes of massive eosinophilia
- Trichocephalus**
Bulucea, D.; and Rarinca, A., 1972, *Microbiol., Parazitol., Epidemiol.*, Bucuresti, v. 17 (2), 169-174
mixed ascariasis and Trichocephalus infection in woman complicated by pleuro-pulmonary staphylococcal pathology, probable parasitic allergic reactions involved in etiology, case report

- Trichocephalus*
Diop, B.; and Bao, O., 1974, *Medecine Afrique Noire*, v. 21 (1), 31-40
human intestinal helminths, clinical indications for treatment, suggested dosage, efficacy, tolerances, possible toxicities
- Trichocephalus*
Grigorescu-Sido, P.; et al., 1972, *Pediatria, Bucuresti*, v. 21 (2), 115-124
human gastrointestinal parasitic diseases, possible relationships with lactase deficiency in children
- Trichocephalus*
Thomas, J.; et al., 1971, *Medecine Trop.*, v. 31 (2), 177-181
bephenium hydroxynaphthoate, piperazine and thiabendazole in comparative clinical tests, institutionalized children with protein deficiency
- Trichocephalus*
Tsolov, B.; and Tsanov, G., 1975, *Vet. Sbirka*, v. 73 (9), 20-22
strongyloids of sheep, economics of tetramisole treatment, body weight, milk production, not effective against *Trichocephalus*
- Trichocephalus*
Vignola, G.; Mori, P. G.; and Cottafava, F., 1969, *Minerva Pediat.*, v. 21 (33), 1526-1530
mixed giardiasis and *Trichocephalus* infections in young child resulting in severe malabsorption syndrome, pathology of intestinal mucosa
- Trichocephalus* [sp.]
Volf, K.; and Volfova, M., 1974, *Veterinarstvi*, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Trichocephalus capreoli*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Trichocephalus capreoli*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichocephalus capreoli*
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Trichocephalus dispar*
Hlavon, J., 1968, *Ceskoslov. Pediat.*, v. 23 (3), 261-262
Trichocephalus dispar in 2-year-old boy with presenting symptoms of malabsorption syndrome, diagnostic difficulties, treatment with telmid: Czechoslovakia
- Trichocephalus dispar*
Rudobielska, M.; et al., 1971, *Pediat. Polska*, v. 46 (9), 1103-1110
significant reduction in serum vitamin B 12 levels in children infected with *Lamblia intestinalis*, no variation seen in children infected with *Trichocephalus dispar* when compared with normal controls
- Trichocephalus muris* Schrank, 1788
Mozgovoi, A. A.; et al., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 95-103
Microtus agrestis
Clethrionomys glareolus
Clethrionomys sp.
all from Karelia
- Trichocephalus muris*
Shikhobalova, N. P.; and Korsak-Paruzhinskaia, L. S., 1975, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 25, 204-214
Ascaridia galli, *Ganguleterakis spumosa*, *Trichocephalus muris*, ionizing radiation of eggs, postradiation effect of high and low temperatures on radiosensitivity as evidenced by development; radiation as possible control of helminths
- Trichocephalus muris*
Shikhobalova, N. P.; and Paruzhinskaia, L. S., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 191-194
Trichocephalus muris, change in radiosensitivity of eggs as a result of irradiation in successive generations
- Trichocephalus nutria*
Sadykhov, I. A., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk* (1), 74-78
influence of ecological factors (age and sex of host, wild or caged animals, season of year) on parasitism
[*Myocastor coypus*]: Azerbaidzhan
- Trichocephalus ovis* Abildgaard, 1795
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (large intestine, cecum): vicinity of Nowy Targ, Carpathian Mountains
- Trichocephalus ovis*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny

- Trichocephalus ovis*
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Trichocephalus ovis*
Dyk, V.; and Chroust, K., 1975, *Veterinarstvi*, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Trichocephalus ovis* Abilgaard, 1795
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
Capreolus capreolus (large intestine): southern Bulgaria
- Trichocephalus ovis*
Kozdon, O.; and Zajicek, D., 1976, *Vet. Med.*, Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field conditions, seasonal distribution as modified by dehelminthization, possible management strategies for effective timing of dehelminthization: Western Bohemia
- Trichocephalus ovis*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Trichocephalus ovis*
Zajicek, D.; and Kozdon, O., 1977, *Veterinarstvi*, v. 27 (6), 257-258
nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Trichocephalus skrjabini* (Baskakow, 1924)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (large intestine, cecum): vicinity of Nowy Targ, Carpathian Mountains
- Trichocephalus skrjabini*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- [*Trichocephalus suis*] *Trikhotosefalus suis*
Bonev, B.; et al., 1975, *Vet. Sbirka*, v. 73 (11), 14-15
pigs, [*Ascaris suum*], [*Trichocephalus suis*], [*Oesophagostomum dentatum*], hygromycin B in feed, completely controls ascariasis
- Trichocephalus suis*
Eimontas, Z. I.; and Medzevicius, A. K., 1974, *Lietuvos TSR Moks. Akad. Darbai*, s. C (68) (4), 65-76
Trichocephalus suis, pigs, immunization with gamma globulin from immune and normal pigs, antibody dynamics, investigation by precipitation and agglutination
- Trichocephalus suis*
Getler, K., 1972, *Med. Wet.*, v. 28 (8), 476-477
nematodes, pigs on industrial swill feeding farm, Atgard
- Trichocephalus suis*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, *Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (26), 123-128
[pig, wild]: Primorskii krai
- Trichocephalus suis*
Tarczynski, S.; Romaniuk, K.; and Szlagiewicz-Czosnek, M., 1972, *Med. Wet.*, v. 28 (4), 217-218
intestinal nematodes, swine, Suiverm
- Trichocephalus suis*
Ziomko, I.; and Paciejewski, S., 1976, *Med. Wet.*, v. 32 (12), 750-752
Trichocephalus suis, pigs, subclinical infection caused decrease in weight gains
- Trichocephalus trichiura*
Asmera, J.; et al., 1975, *Ceskoslov. Pediat.*, v. 30 (4), 171-172
human intestinal helminths, mebendazole used as successful wide-spectrum anthelmintic in clinical trials: Czechoslovakia
- Trichocephalus trichiura*, *illus.*
Cerva, L., 1976, *Immun. u. Infekt.*, v. 4 (6), 279-282
intestinal helminths, diagnostic method for staining of eggs and larvae in smears of fresh and fixed stool samples
- Trichocephalus trichiuris*
Obtulowicz, K., 1972, *Polski Tygod. Lekar.*, v. 27 (8), 296-298
human intestinal parasites, possible role in etiology of asthma and other allergic conditions
- Trichocephalus trichiurus*
Spaldonova, R.; and Virgala, J., 1974, *Ceskoslov. Pediat.*, v. 29 (6), 345-347
Trichocephalus trichiurus in children, clinical trials using mebendazole and recommendations for drug usage: Czechoslovakia
- Trichocephalus vulpis*
Kozakiewicz, B., 1976, *Med. Wet.*, v. 32 (5), 293-295
Trichocephalus vulpis, dogs, mebendazole very effective, tetramisole ineffective

- Trichocheenia Kou, 1958
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Molineinae
synonymy
- Trichohelix Ortlepp, 1922
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Tricholeiperia Travassos, 1935
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Tricholeiperia sp.
Martin, D. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 85-86
Tadarida brasiliensis: Texas
- Tricholinstowia Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligionellidae, Heligionellinae
synonymy
- Trichonema
Blanchard, J., 1971, Prat. Vet. Equine, v. 3 (1), 29-31
gastrointestinal nematodes in race horses, clinical trials with equigard, good immediate results especially with ascarids; rapid reinfection rates (2 months after therapy), suggests contaminated paddocks; recommendations therefore for program of routine treatment: France
- Trichonema
Grelck, H.; Hoerchner, F.; and Woehrl, H., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (19), 371-373
Fasciola hepatica-infected ponies, orally and intraperitoneally, rate of development of parasite, most liver flukes found in animals severely infected with Strongylus and Trichonema
- Trichonema
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Trichonema spp.
Abdel-Rahman, M. S.; et al., 1972, Parasitol. Hungar., v. 5, 225-237
Strongylus spp., Trichonema spp., Parascaris equorum, Trichostrongylus axei in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Trichonema [sp.]
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
donkeys, survey, treatment, economic importance: Kaedi area (Mauritania)
- Trichonema spp.
Duncan, J. L.; et al., 1977, Equine Vet. J., v. 9 (3), 146-149
immature strongyles, ponies, fenbendazole
- Trichonema spp.
Grelck, H.; Hoerchner, F.; and Woehrl, H. E., 1977, Prakt. Tierarzt, v. 58 (4), 265-268
strongyles, horses, seasonal development of infective larvae on pastures, survival rate
- Trichonema spp.
Jeggio, M. H.; and Sewell, M. M. H., 1977, Vet. Rec., v. 101 (10), 187
Trichonema spp., horse, mebendazole and pyrantel embonate failed to prevent onset of disease, successfully treated with fenbendazole
- Trichonema sp.
Mirzayans, A., 1973, Vet. Rec., v. 92 (10), 262 [Letter]
pony (intestinal tract): northern Iran
- Trichonema spp.
Ooms, L., 1975, Vlaams Diergeneesk. Tijdschr., v. 44 (3), 95-118
strongyles, horses, epidemiology, cycles, pathogenesis, symptoms, control, diagnosis, immunity, review
- Trichonema spp., illus.
Rizzoli-Stalder, C.; et al., 1976, Schweiz. Arch. Tierh., v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Trichonema [sp.], larvae
Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 77-81
Equus zebra zebra (intestinal wall): Mountain Zebra National Park
- Trichonema catinatum
Chiejina, S. N.; and Mason, J. A., 1977, Vet. Rec., v. 100 (17), 360-361
Trichonema longibursatum, T. catinatum, horses, maturation of inhibited 3rd stage larvae in intestinal mucosa causing diarrhoea in spring: [Britain]
- Trichonema catinatum (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Trichonema longibursatum
Chiejina, S. N.; and Mason, J. A., 1977, Vet. Rec., v. 100 (17), 360-361
Trichonema longibursatum, T. catinatum, horses, maturation of inhibited 3rd stage larvae in intestinal mucosa causing diarrhoea in spring: [Britain]

- Trichoneminae
Furmaga, S.; Gundlach, J. L.; and Patyra, J., 1976, *Med. Wet.*, v. 32 (12), 734-737
roundworms, horses, fenbendazol and cambendazol very effective
- Trichoskrjabinia Travassos, 1937
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Molineidae, Molineinae
- Trichoskrjabinia secundus sp. n., illus.
Pinnell, J. L.; and Schmidt, G. D., 1977, *J. Parasitol.*, v. 63 (2), 337-340
Sphenomorphus striolatus (small intestine): Nununay Vau, Komodo Island, Indonesia
- Trichosomoides crassicauda
Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
Rattus norvegicus (vessie): Tunis, Tunisia
- Trichosomoides crassicauda
Owen, D., 1976, *Lab. Animals*, v. 10 (3), 271-278
Rattus norvegicus: Carshalton
- Trichosomoides crassicauda
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 214
Ratte
- Trichosomoides crassicauda (Bellingham, 1840)
Singh, M.; and Cheong Chee Hock, 1971, *South-east Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 516-521
Rattus sabanus: Malaysia
- Trichosomoides crassicauda, illus.
Taniguchi, M.; et al., 1977, *Bull. Coll. Agric. and Vet. Med., Nihon Univ.* (34), 202-217
Rattus norvegicus
R. rattus
all from Setagaya-ku area, Tokyo
- Trichosomoides crassicauda (Bellingham, 1840)
Railliet, 1895
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, *BoI. Chileno Parasitol.*, v. 31 (1-2), 39-42
Rattus norvegicus (vejiga urinaria): Chile
- Trichospirura Smith & Chitwood, 1967
Chabaud, A. G., 1975, *CIH Keys Nematode Parasites Vertebrates* (Anderson, Chabaud, and Willmott) (3), 1-27
Rhabdochoniidae
key
- Trichospirura
Chabaud, A. G.; and Krishnasamy, M., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 813-820
Trichospirura, should be placed in Rhabdochoniidae, evolutionary position, host range, osmoexcretory apparatus, relations between Rhabdochoniidae and Cystidicolidae
- Trichospirura sp., illus.
Chabaud, A. G.; and Krishnasamy, M., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 813-820
description
Myotis mystacinus (intestin): Bukit Lagong-Kepong
- Trichospirura leptostoma, illus.
King, N. W., jr., 1976, *Scient. Publication* (317). *Pan Am. Health Organ.*, 169-198
- Trichospirura willmottae n. sp., illus.
Chabaud, A. G.; and Krishnasamy, M., [1976], *Ann. Parasitol.*, v. 50 (6), 1975, 813-820
Tupaia glis (canaux de glandes salivaires): Uly Gombak (proximite de Kuala-Lumpur)
- [Trichostrongyles] trichostrongilly
Nurtazin, A. T., 1975, *Vestnik Sel'skokhoz. Nauki Kazakhstana* (4), 84-86
sheep, trichostrongyles, anthelmintic efficacy of banminth, good results
- Trichostrongyles
Varshney, T. R.; and Singh, Y. P., 1976, *Indian Vet. J.*, v. 53 (9), 672-676
Haemonchus contortus, trichostrongyles, Trichuris, lambs, critical trial, efficacy and economic value, various anthelmintics: Central Sheep and Wool Research Station, Pashulok Rishikesh (Dehradun)
- Trichostrongylidae
Cabaret, J., 1977, *Rec. Med. Vet.*, v. 153 (6), 419-427
gastrointestinal strongyles (Trichostrongylidae, Strongylidae), inhibition of larval development (hypobiosis), weather conditions, epidemiological aspects, review
- Trichostrongylidae (Leiper, 1908, sub. fam.),
Leiper, 1912
Durette-Desset, M. C.; and Chabaud, A. G., 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Trichostrongyloidea
includes: Libyostrongylinae; Cooperiinae; Graphidiinae; Ostertagiinae; Trichostrongylinae; Haemonchinae
- Trichostrongylidae [sp.]
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, *Canad. J. Zool.*, v. 54 (3), 307-312
helminths of Alces alces, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Trichostrongyliden
Huetz, J.; and Lamina, J., 1975, *Bayer. Landwirtsch. Jahrb.*, v. 52 (7), 819-836
cattle, development of parasite, diagnosis, treatment, economic importance, review

- Trichostrongyliden**
 Manz, D.; and Dingeldein, W., 1974, Prakt. Tierarzt, v. 55 (8), 422-425
 nematodes of European and exotic herbivores,
 Banminth, good results
 Dama dama mesopotamica
 Cervus elaphus
 Ovis musimon
 Nyala angasi
 Taurotragus oryx derbianus
 Antidorcas marsupialis
 Antilope cervicapra
- Trichostrongyliden-type, illus.**
 Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
 helminths of pet birds, diagnosis of eggs in fecal examination
- Trichostrongylids**
 Bergstrom, R. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 61-63
 Marshallagia marshalli, other trichostrongylids, incidence and intensity in Ovis aries under 3 types of management (farm flocks, fenced range, and seasonally herded range) and in wild Antilocapra americana; evidence for transmission of M. marshalli from antelope to sheep where range is shared: Wyoming
- Trichostrongylid ova**
 Hiregoudar, L. S., 1976, Indian Vet. J., v. 53 (3), 237
 Gazella gazella (feces): Gir forest, Gujarat State, India
- Trichostrongylids**
 Jansen, J., 1977, Trop. and Geogr. Med., v. 29 (3), 316-317 [Abstract]
 trichostrongylosis, sheep, questionable correlation between breed susceptibility to infections and hemoglobin types of breeds
- Trichostrongylid infections**
 Kelly, J. D., 1974, Internat. J. Zoonoses, v. 1 (1), 13-24
 anthroponozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Trichostrongylids**
 Kirsch, R., 1976, Ztschr. Parasitenk., v. 50 (2), 223
 trichostrongylid eggs, morphological changes after fenbendazole treatment of infected lambs
- Trichostrongylids**
 Nemeseri, L., 1976, Magy. Allat. Lapja, v. 98, v. 32 [i. e. 31] (7), 459-461
 Dictyocaulus filaria, trichostrongylids, Fasciola hepatica, sheep, successful treatment with combination of diamphenetide and tetramisole; no effect on Protostrongylidae
- Trichostrongylinae Leiper, 1912**
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 Trichostrongylidae
 includes: Trichostrongylus (type genus); Graphidioides; Travassosius
- Trichostrongyloidea**
 Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
 Strongylida
 key
- Trichostrongyloidea**
 Durette-Desset, M. C.; and Chabaud, A. G., 1975, Ann. Parasitol., v. 50 (3), 303-337
 Trichostrongyloidea of Chiroptera, evolutionary relationships
- Trichostrongyloidea**
 Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
 classification, hypotheses on evolution includes: Strongylacanthidae; Amidostomatidae; Molineidae; Helogmosomidae; Heligmonellidae; Trichostrongylidae; Ollulanidae; Dictyocaulidae
- Trichostrongyloidea**
 Durette-Desset, M. C.; Chabaud, A. G.; and Cassone, J., 1977, Bull. Mus. National Hist. Nat., Paris, 3. s. (428), Zool. (298), 133-158
 Trichostrongyloidea from Brazilian Xenarthra not related to those from American marsupials, seem to have evolved from reptilian or amphibian forms
- Trichostrongyloidea [sp.] ova**
 Faust, B. S.; and Pappas, P. W., 1977, J. Zoo Animal Med., v. 8 (1), 18-23
 Cygnus atratus (feces): Columbus (Ohio) Zoo
- Trichostrongyloidiasis**
 Hollo, F., 1971, Parasitol. Hungar., v. 4, 104-114
 trichostrongyloidiasis in sheep causing gastro-enteritis, comparative trials of various anthelmintics showed tetramisole to be treatment of choice
- Trichostrongylosis**
 Jansen, J., 1977, Trop. and Geogr. Med., v. 29 (3), 316-317 [Abstract]
 trichostrongylosis, sheep, questionable correlation between breed susceptibility to infections and hemoglobin types of breeds
- Trichostrongylosis**
 Tarazona, J. M., 1975, An. Inst. Nac. Invest. Agrar., s. Hig. y San. Animal (2), 11-17
 trichostrongylidosis, ovine, seasonal incidence, 1971-1975: Spain
- Trichostrongylus**
 Baines, D. M.; and Colegrave, A. J., 1977, Vet. Rec., v. 100 (11), 217-219
 gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales
- Trichostrongylus**
 Bliss, D. H.; and Todd, A. C., 1977, Vet. Med. and Small Animal Clin., v. 72 (10), 1612-1617
 milk production in dairy cows exposed to mixed trichostrongylid larvae, results indicate that greatest milk loss occurs during the first 90 days of lactation, relationship between exposure time and stage of lactation

- Trichostrongylus**
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Trich[ost]rongylus**
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Trichostrongylus**
Bueno, L.; Dorchies, P.; and Ruckebusch, Y., 1976, Compt. Rend. Soc. Biol., Paris, v. 169 (6), 1975, 1627-1632
Trichostrongylus, lambs (exper.), disturbances in gastrointestinal motility preceding diarrhea, electromyographic analysis, effect of thiabendazole treatment
- Trichostrongylus**
Chhabra, R. C.; Bali, H. S.; and Toor, L. S., 1976, J. Research, Punjab Agric. Univ., v. 13 (3), 308-311
gastrointestinal strongyles in sheep, critical drug evaluation, thiabendazole (most effective), tetramisole (good results), morantel tartrate (fair results), clioquinide and methyridine (least effective): India
- Trichostrongylus**
Chowaniec, W.; et al., 1975, Med. Wet., v. 31 (12), 741-743
Fasciola hepatica, Haemonchus, Trichostrongylus, Nematodirus, cattle, Nilzan, Zanil, field trials, good results
- Trichostrongylus**
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Vet. Rec., v. 92 (20), 551-554
control of clinical infections of gastrointestinal nematodes and lungworms in calves using morantel/diethylcarbazine solution, field trials, good results as measured by growth response and clinical symptoms; routine treatment economically sound under conditions of heavy infection
- Trichostrongylus**
Crowley, J. W., jr.; et al., 1977, Am. J. Vet. Research, v. 38 (5), 689-692
lungworms, gastrointestinal parasites, cattle, 3 controlled critical trials, highly effective
- Trichostrongylus**
Dewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Trichostrongylus** Looss, 1905 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Trichostrongylidae, Trichostrongylinae
synonymy
- Trichostrongylus**
Fudalewicz-Niemczyk, W.; et al., 1975, Med. Wet., v. 31 (11), 666-668
sheep helminths, effective control with Nilverm and Zanil, increased weight gains and shearing yields: Hanczowa, Gorlice district
- Trichostrongylus**
Fudalewicz-Niemczyk, W.; et al., 1976, Acta Zootech., Bratislava (32), 5-19
gastrointestinal helminths, mountain sheep, nilverm and zanil, favorable influence on body weight and wool production of treated animals, no influence of treatment on fertility: Poland
- Trichostrongylus**
Grzywinski, L.; et al., 1975, Medycyna Wet., v. 31 (9), 524-526
cattle, Nilverm by injection for control
- Trichostrongylus**
Guarino, C.; and Rivellini, P., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 487-490
nematode larvae in grass samples from various types of pasture, degree of infestation: province of Avellino
- Trichostrongylus**
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Trichostrongylus**
Hausfater, G.; and Watson, D. F., 1976, Nature, London (5570), v. 262, 688-689
emission of parasite ova (primarily Trichuris and Trichostrongylus) by Papio cynocephalus in relation to host social and reproductive condition, high-ranking adult males had higher egg emission than more subordinate individuals, sexually cycling females had higher emissions than anoestrous females
- Trichostrongylus**
Henriksen, Sv. Aa.; et al., 1976, Vet. Parasitol., v. 2 (3), 259-272
gastro-intestinal nematodes, young calves during first grazing season, infection levels, blood findings, body weight gains, comparison of animals grazing same pasture entire season with those moved in early July and between levamisole-treated and untreated animals: Denmark
- Trichostrongylus**
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1974, Indian J. Animal Research, v. 8 (2), 75-78
Haemonchus contortus, other nematodes, experimentally or naturally infected sheep, nitroxylin highly effective, critical testing; in vitro testing against H. contortus

- Trichostrongylus**
Qadir, A. N. M. A., 1976, Indian Vet. J., v. 53 (11), 855-858
gastrointestinal nematodes, goats and calves, urea for pasture control of free-living stages
- Trichostrongylus**
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, not effective
- Trichostrongylus**
Sewell, M. M. H., 1973, Vet. Rec., v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Trichostrongylus**
Tharaldsen, J., 1976, Acta Vet. Scand., v. 17, Suppl. 61, 1-21
trichostrongylid infections, calves, survival of larvae on pasture, occurrence of larvae not influenced by artificial irrigation; treatment with thiabendazole did not effectively control infection due to overwintering larvae, neither improved weight gain nor reduced egg production: Norway
- Trichostrongylus**
Theodorides, V. J.; et al., 1973, Brit. Vet. J., v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Trichostrongylus**
Theodorides, V. J.; et al., 1976, Experientia, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Trichostrongylus**
Todd, A. C.; et al., 1976, Am. J. Vet. Research, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Trichostrongylus**
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Trichostrongylus**
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, Vet. Glasnik, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbazine effective against Dictyocaulus filaria and most gastrointestinal helminths except Strongyloides papillosus, Trichuris ovis, and Moniezia sp.
- Trichostrongylus**
Wallnoefer, E., 1977, Wien. Tierarztl. Monatsschr., v. 64 (4), 129-131
sheep parasites, Mebenvet, good results when treatment was repeated after 14 days: Austria
- Trichostrongylus**
Whitlock, J. H.; and Georgi, J. R., 1976, Parasitology, v. 72 (3), 207-224
biological controls in mixed trichostrongylid infections (predominantly Haemonchus contortus cayugensis) in sheep, different ecosystems (barn vs. pasture) and different treatment groups, course of infections (erythrocyte loss, fecal egg counts, hematocrit values), "Anaphylactoid 'self-cure' did not occur in this experiment but something like premunition certainly did."
- Trichostrongylus**
Zeakes, S. J.; et al., 1976, Am. J. Vet. Research, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos crumbles and naftalofos boluses
- Trichostrongylus**
Zielinski, J., 1972, Med. Wet., v. 28 (9), 566-567
parasites, sheep, Nilverm, copper sulfate
- Trichostrongylus spp.**
Arfaa, F.; et al., 1977, Am. J. Trop. Med. and Hyg., v. 26 (2), 230-233
Ascaris, Trichostrongylus spp., Ancylostoma duodenale, evaluation of different methods of control of soil transmitted helminths (sanitation measures, mass-therapy, combined mass-therapy and sanitation) in villagers of Khuzestan, southwest Iran
- Trichostrongylus sp.**
Bali, M. K.; and Singh, R. P., 1976, Indian J. Animal Research, v. 10 (2), 111-112
Haemonchus contortus, Oesophagostomum sp., Trichostrongylus sp., Trichuris sp., sheep, goats, morantel tartrate, good results against all parasites except for Trichuris sp.
- Trichostrongylus [sp.]**
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
ruminants, donkeys, survey, treatment, economic importance: Kaedi area (Mauritania)
- Trichostrongylus sp.**
Canale, A.; et al., 1972, Atti Soc. Ital. Sc. Vet., v. 26, 306-310
Fasciola hepatica, calves, light experimental infection alone or in combination with gastrointestinal nematodes, digestive function not impaired
- Trichostrongylus spp.**
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs and heifers, efficacy of fenbendazole, thiabendazole and tetramisole compared

- Trichostrongylus spp.
Colglazier, M. L.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with Trichuris spp. and Moniezia expansa
- Trichostrongylus sp.
Cornwell, R. L.; Jones, R. M.; and Pott, J. M., 1973, Brit. Vet. J., v. 129 (6), 526-532
cattle, morantel tartrate, good results against Cooperia sp., Ostertagia sp., and Trichostrongylus sp., increased weight gain in treated cattle, field trials: United Kingdom
- Trichostrongylus spp.
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Trichostrongylus spp.
Düewel, D.; et al., 1974, Prakt. Tierarzt, v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Trichostrongylus spp.
El-Abdin, Y. Z.; et al., 1975, Egypt. J. Vet. Sc., v. 12 (1), 31-43
serum constituents and serum enzyme activities, normal and nematode infested Camelus dromedarius: Cairo abattoir
- Trichostrongylus spp.
Farahmandian, I.; et al., 1977, Chemotherapy, v. 23 (2), 98-105
human intestinal Ascaris, hookworm and Trichostrongylus spp., statistics of comparative trials using pyrantel, levamisole, mebendazole, thiabendazole and bephenium: Iran
- [Trichostrongylus] sp. "Trikostrongilus"
Georgieva, D.; Vladimirova, A.; and Monov, M., 1975, Vet. Sbirka, v. 73 (11), 18, 20
nematodes of lambs, comparative tests of tetramisole, group and individual applications
- Trichostrongylus sp.
Gonzalez, H.; and Plaza, J., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 134-137
gastrointestinal nematodes of sheep, comparative therapeutic trials using banminth, phenothiazine, and thiabendazole
- Trichostrongylus spp.
Ismail, E.; Tawfik, A. A.; and El-Ebrashi, N. M. A., 1977, Arzneimittel-Forsch., v. 27 (7), 1393-1394
6-hydroxy-4-methoxy-5-(p-methoxy-cinnamoyl)-benzofuran, in vitro broad spectrum anthelmintic activity against livestock helminths, promising results indicate need for future research
- Trichostrongylus sp.
Jacobson, H. A.; and Kirkpatrick, R. L., 1974, J. Wildlife Dis., v. 10 (4), 384-391
comparison of selected physiological measurements in untreated parasitized cottontail rabbits and those treated with 1-tetramisole hydrochloride and 2,2-dichlorovinyl, dimethyl phosphate: Montgomery County, Virginia
- Trichostrongylus [sp.]
Leguia, G.; and Bendezu, P., 1974, Rev. Invest. Pecuarias, v. 3 (1), 3-7
gastrointestinal nematodes, variation in fecal egg counts, 2 year period, pregnant Lama pacos: Central Sierra of Peru (Dept. Pasco)
- Trichostrongylus spp.
Leimbacher, F.; Nicolas, J. A.; and Delahaye, J., 1976, Rev. Med. Vet., Toulouse, v. 127 (6), 941-958
oxfendazole, comparison with tetramisole, gastrointestinal strongylosis, lambs
- Trichostrongylus spp.
Levine, N. D.; et al., 1975, Am. J. Vet. Research, v. 36 (10), 1459-1464
lambs grazing with their ewes under 2 pasture rotation systems, lambs under rotation had more nematodes and gained less weight than nonrotated control lambs, rotation is not recommended to control nematode parasitism of sheep in Illinois
- Trichostrongylus sp. 4th stage
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Trichostrongylus sp.
Mutafova, T., 1972, Izvest. Tsentral. Khelminth. Lab., v. 15, 143-150
Haemonchus contortus, Ostertagia sp., Trichostrongylus sp., sheep, seasonal changes in egg-production, spring rise observed irrespective of age or sex of host
- Trichostrongylus spp.
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Trichostrongylus sp.
Qadir, A. N. M. A., 1976, Indian Vet. J., v. 53 (6), 448-450
Haemonchus sp., Trichostrongylus sp., Oesophagostomum sp., larvicidal action of 4 chemical compounds on infective nematode larvae in experimental outdoor plots; urea most effective

- Trichostrongylus sp., *illus.*
Sapunar, J.; and Bertossi, E., 1966, Bol. Chileno Parasitol., v. 21 (3), 94-95
mixed Trichostrongylus, Trichuris trichiura, and Giardia lamblia infection in young man diagnosed by fecal examination, case report: Chile
- Trichostrongylus sp.
Schillhorn van Veen, T.; and Brinckman, W. L., 1975, Samaru Agric. Newsletter, v. 17 (2), 70-74
lambs, regular drenching with thiabendazole at regular intervals during rainy season, better weight gain, cost/benefit; possible influence of resistance and breed of sheep
- Trichostrongylus spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Trichostrongylus sp.
Tongson, M. S.; and Montenegro, M. M., 1975, Philippine J. Vet. Med., v. 13 (1-2), 170-182
purified microfine phenothiazine + lead arsenate, anthelmintic efficiency, good results, goats: Philippines
- Trichostrongylus spp.
Torres, P.; Figueroa, L.; and Navarrete, N., 1972, Bol. Chileno Parasitol., v. 27 (1-2), 52-55
Trichostrongylus spp. survey in humans, morphological differentiation for diagnosis: Chile
- Trichostrongylus spp.
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, Camelus dromedarius, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Trichostrongylus sp.
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago, intestino delgado): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Trichostrongylus sp.
Vassiliades, G.; and Toure, S. M., 1975, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 28 (4), 481-489
digestive strongylosis, sheep, morantel tartrate, with or without anticoccidian drug (Cozurone), good control of all except Strongyloides
- Trichostrongylus spp.
Vlassoff, A., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 281-284
trichostrongyle larvae on pasture, seasonal incidence, residual pasture infestation more important than ewes as source of infection for lambs in spring, autumn infections acquired from eggs passed by lambs themselves: New Zealand
- Trichostrongylus [sp.]
Volf, K.; and Volfova, M., 1974, Veterinarstvi, v. 24 (3), 125-126
jeleni zvere
srnci zvere
all from Trebic District
- Trichostrongylus sp.
Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
Psammomys obesus: Israel
- Trichostrongylus sp.
Wikerhauser, T.; et al., 1974, Acta Parasitol. Jugoslavica, v. 5 (2), 79-81
trichostrongylids, cattle, fenbendazole compared with thiabendazole, good results from both
- Trichostrongylus affinis
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (cecum, large intestine): Georgia; Alabama; Mississippi
- Trichostrongylus affinis
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (cecum): Florida
- Trichostrongylus affinis
Knight, R. A., 1977, J. Parasitol., v. 63 (5), 957-958
Trichostrongylus affinis, Nematodirus spathiger, rabbits (exper.), effect of dexamethasone (higher egg counts but no difference in numbers of worms that developed, did not prevent worm expulsion)
- Trichostrongylus affinis
Yonders, P. C., 1975, J. Alabama Acad. Sc., v. 46 (3-4), 95 [Abstract]
Trichostrongylus affinis infected domestic rabbits, ⁵⁵Fe incorporation, compared with non-infected rabbits, results indicate that the infection is altering the iron metabolism of the host
- Trichostrongylus askivali
Eve, J. H.; and Kellogg, F. E., 1977, J. Wildlife Management, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (Odocoileus virginianus) density: south-eastern United States

- Trichostrongylus askivali Dunn, 1964
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Oklahoma
- Trichostrongylus axei
Abdel-Rahman, M. S.; et al., 1972, Parasitol. Hungar., v. 5, 225-237
Strongylus spp., Trichonema spp., Parascaris equorum, Trichostrongylus axei in horses, field trials testing efficacy of various anthelmintics; phenothiazine, banminth and thiabendazole most effective: Egypt
- Trichostrongylus axei
van Adrichem, P. W. M.; and Shaw, J. C., 1977, J. Animal Sc., v. 45 (3), 417-422
gastrointestinal nematodes, monozygous twin cattle, comparison of treated and untreated pairs infected naturally on pasture, growth performance, results indicate that the reduced growth may be long-lasting
- Trichostrongylus axei
van Adrichem, P. W. M.; and Shaw, J. C., 1977, J. Animal Sc., v. 45 (3), 423-429
gastrointestinal nematodes, effects on growth performance and milk production in cambendazole-treated vs. non-treated monozygous twin cattle naturally infected on pasture during the first lactation period
- Trichostrongylus axei
Anderson, N., 1977, Research Vet. Sc., v. 23 (3), 298-302
Ostertagia ostertagi, Trichostrongylus axei, cattle, efficacy of levamisole, thiabendazole, and fenbendazole
- Trichostrongylus axei
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, Trichostrongylus, and Nematodirus, oxfendazole highly effective against adult stages in sheep
- Trichostrongylus axei (Cobbold, 1879)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of Syncerus caffer, some pathological findings, age of host
Syncerus caffer (small intestine): Kruger National Park
- Trichostrongylus axei
Benz, G. W.; and Ernst, J. V., 1977, Am. J. Vet. Research, v. 38 (9), 1425-1426
gastrointestinal nematodes, calves (exper.), albendazole significantly reduced infestations
- Trichostrongylus axei (Cobbold, 1879)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Trichostrongylus axei
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Trichostrongylus axei
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- T[richostrongylus] axei
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Trichostrongylus axei
Bryan, R. P., 1976, Austral. Vet. J., v. 52 (9), 403-408
nematodes, paramphistomes, young beef cattle, growth rates, levamisole, niclosamide
- Trichostrongylus axei
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Trichostrongylus axei
Ciordia, H.; et al., 1977, Am. J. Vet. Research, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Trichostrongylus axei
Coles, G. C.; and Simpkin, K. G., 1977, Research Vet. Sc., v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant Haemonchus contortus and Trichostrongylus colubriformis to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Trichostrongylus axei
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method

- Trichostrongylus axei*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Trichostrongylus axei*
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (abomaso): Sardegna
- Trichostrongylus axei*
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxbendazole, drug efficacy
- Trichostrongylus axei*
Curr, C., 1977, Austral. Vet. J., v. 53 (9), 425-428
nematodes, calves, levamisole, efficiency of pour-on formulation, drug trials, good results
- Trichostrongylus axei*
Dorn, H.; and Federmann, M., 1976, Vet.-Med. Nachr. (1), 5-17
gastrointestinal nematodes in cattle (nat. and exper.), citarin-L spot-on, application on skin, good results
- Trichostrongylus axei*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Trichostrongylus axei* (Cobbold, 1879) Railliet et Henry, 1909
Drozd, J.; and Bylund, G., 1970, Acta Parasitol. Polon., v. 17 (20-38), 259-260
Alces alces (abomasa): Poland
- Trichostrongylus axei*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Trichostrongylus axei*
Duncan, J. L.; et al., 1976, Vet. Rec., v. 98 (17), 342
Ostertagia ostertagi (inhibited 4th stage larvae), *Trichostrongylus axei*, *Cooperia* spp., fenbendazole, good results, compared with levamisole
- Trichostrongylus axei*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Trichostrongylus axei*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon
Capreolus capreolus
(digestive tract of all): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichostrongylus axei*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitology, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon
Capreolus capreolus
all from Czechoslovakia
- Trichostrongylus axei*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Trichostrongylus axei*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Trichostrongylus axei*
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1974, Prakt. Tierarzt, v. 55 (8), 417-422
nematodes of horses, fenbendazol, good results
- Trichostrongylus axei*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichostrongylus axei*
Eve, J. H.; and Kellogg, F. E., 1977, J. Wildlife Management, v. 41 (2), 169-177
technique for using intensity of abomasal parasite infections as an index to deer (*Odocoileus virginianus*) density: southeastern United States
- Trichostrongylus axei*
Foix, J., 1977, Rev. Med. Vet., Toulouse, v. 128 (8-9), 1111-1119
Dicrocoelium dendriticum, sheep, cambendazole, good results against flukes as well as gastrointestinal strongyles
- Trichostrongylus axei*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high

- Trichostrongylus axei*
Ghadirian, E.; and Arfaa, F., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus axei*
Goldberg, A., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 109-110
control of helminth parasitism, infectiousness of pastures, rested or grazed by resistant cattle
- Trichostrongylus axei*
Gonzalez, H.; and Plaza, J., 1968, Bol. Chileno Parasitol., v. 23 (3-4), 134-137
gastrointestinal nematodes of sheep, comparative therapeutic trials using banminth, phenothiazine, and thiabendazole
- Trichostrongylus axei*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Trichostrongylus axei*
Herlich, H., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.), oxbendazole, efficacy against adult and larval stages
- Trichostrongylus axei*
Herlich, H., 1976, Vet. Parasitol., v. 2 (4), 377-383
Trichostrongylus axei, *T. colubriformis*, rabbits, efficacy of 8 anthelmintics at dose rates known to be effective in ruminants, evaluation of this system as preliminary screen for testing potential anthelmintics
- Trichostrongylus axei*
Herlich, H., 1977, Am. J. Vet. Research, v. 38 (8), 1247-1248
efficacy of albendazole against gastrointestinal nematodes and *Fasciola hepatica* in cattle (exper.); comparison of critical vs. controlled tests
- Trichostrongylus axei*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Trichostrongylus axei* (Cobbold, 1879) Railliet et Henry, 1909
Ianchev, I., 1973, Izvest. Tsentral. Khelminth. Lab., v. 16, 205-220
Capreolus capreolus (rennet, small intestine): southern Bulgaria
- Trichostrongylus axei*
Ingolfsson, A.; and Gislason, G., 1975, Islen. Landbunadarranns., v. 7 (1-2), 3-7
cattle, nautgripa (abomasum): southwestern Iceland, slaughterhouse in Reykjavik
- Trichostrongylus axei*
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired *Dictyocaulus filaria* infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Trichostrongylus axei*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Trichostrongylus axei*
Kerboeuf, D.; and Leimbacher, F., 1977, Rec. Med. Vet., v. 153 (1), 19-25
Ostertagia circumcincta, *Trichostrongylus axei*, *Haemonchus contortus*, lambs, use of serum pepsinogen measurements to assess average worm burden in a herd
- Trichostrongylus axei*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Trichostrongylus axei*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Trichostrongylus axei*
Le Jambre, L. F.; and Royal, W. M., 1976, Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (fecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except *Nematodirus* resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales
- Trichostrongylus axei*
Lukovich, R.; et al., 1977, Gac. Vet., Buenos Aires (318), v. 39, 91-95
helminths, cattle, levamisole, results from injectable and dermal application similar

- Trichostrongylus axei*
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Trichostrongylus axei*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1976, Am. J. Vet. Research, v. 37 (6), 701-702
horses, thiabendazole (paste formulation), efficacy determined by critical testing method against large nematodes and *Gasterophilus* (inactive against latter)
- Trichostrongylus axei*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (6), 721-723
horses (feces)
- Trichostrongylus axei*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Trichostrongylus axei*
McBain, D. G.; et al., 1977, Vet. Rec., v. 101 (14), 285-286
helminths, calves, fenbendazole in feed blocks
- Trichostrongylus axei*
McKenna, P. B., 1976, N. Zealand J. Exper. Agric., v. 4 (2), 235-237
post-mortem recovery of *Haemonchus contortus*, *Ostertagia* spp., *Trichostrongylus axei*, sheep, peptic digestion of ovine abomasum unlikely to be of diagnostic value for field-submitted specimens
- Trichostrongylus axei*
Martinez Gomez, F.; and Hernandez Rodriguez, S., 1973, Rev. Iber. Parasitol., v. 33 (1), 11-20
Ovis aries (abomasum): Cordoba, Spain
- Trichostrongylus axei*
Martinez Gomez, F.; Hernandez Rodriguez, S.; and Calero Carretero, R., 1973, Rev. Iber. Parasitol., v. 33 (4), 625-631
Capra hircus: Municipal Slaughterhouse, Cordoba, Spain
- Trichostrongylus axei*
de Matos, P. F.; and Costa, J. O., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 173-180
gastrointestinal helminths, horses, levamisole, haloxon, crufomate, anthelmintic efficiency
- Trichostrongylus axei*
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with oxfendazole
- Trichostrongylus axei*
Niec, R.; et al., 1976, Gac. Vet., Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary anthelmintic treatment that could prevent normal buildup of resistance
- Trichostrongylus axei*
Nowosad, B., 1975, Zeszyty Nauk. Akad. Rolnicz. Krakow. (98), Zootech. (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Trichostrongylus axei*
Oberger, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus
Ovis aries
Sus scrofa
Equus caballus
all from Chile
- Trichostrongylus axei*
Panitz, E., 1977, J. Helminth., v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Trichostrongylus axei* Cobbold
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella thomsonii (gut)
Alcelaphus buselaphus cokei (digestive tract)
all from Kenya
- Trichostrongylus axei*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, J. Wildlife Dis., v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Trichostrongylus axei*
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Trichostrongylus axei* (Cobbold, 1879)
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (abomasum): Oklahoma

- Trichostrongylus axei*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Trichostrongylus axei*
Raynaud, J. P.; and Bouchet, A., 1976, *Ann. Recherches Vet.*, v. 7 (3), 253-280
bovine ostertagiosis, analysis of types and syndromes, total worm counts, post mortem examinations, survey of 74 cattle: France
- Trichostrongylus axei*
Reid, J. F. S., 1976, *Vet. Rec.*, v. 98 (25), 496-499
gastrointestinal nematodes, coccidiosis, diarrhea of sheep, age and seasonal factors: Britain
- [*Trichostrongylus*] *axei*
Reinecke, R. K., 1972, *Onderstepoort J. Vet. Research*, v. 39 (3), 153-178
gastrointestinal nematodes of cattle, use of modified nonparametric method to evaluate anthelmintic efficacy of levamisole and mebendazole against various parasite stages, detailed description of each step of procedure
- Trichostrongylus axei*, *illus.*
Rizzoli-Stalder, C.; et al., 1976, *Schweiz. Arch. Tierh.*, v. 118 (9), 367-375
gastrointestinal parasites, horses, influence of pasturing and deworming on infestation, two test groups, higher infestation in group receiving regular anthelmintic treatment probably due to high density of animals on pasture
- Trichostrongylus axei*
Ronald, N. C.; Bell, R. R.; and Craig, T. M., 1977, *J. Am. Vet. Med. Ass.*, v. 170 (3), 317-319
gastrointestinal nematodes, calves, levamisole phosphate, effective at one-half recommended dosage
- Trichostrongylus axei*
Rose, J. H., 1971, *Symposia Brit. Soc. Parasitol.*, v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Trichostrongylus axei*
Rothwell, T. L. W.; et al., 1976, *Vet. Parasitol.*, v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Trichostrongylus axei*
Schweisgut, I., 1975, *Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75*, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Trichostrongylus axei*
Searson, J. E.; and Doughty, F. R., 1977, *Austral. Vet. J.*, v. 53 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult *Ostertagia ostertagi* than larval forms): southern New South Wales
- Trichostrongylus axei*
Smeal, M. G.; et al., 1977, *Austral. Vet. J.*, v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Trichostrongylus axei*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, *Austral. J. Agric. Research*, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Trichostrongylus axei*
Stewart, T. B.; Ciordia, H.; and Utley, P. R., 1975, *Am. J. Vet. Research*, v. 36 (6), 785-787
feedlot cattle with subclinical parasitism (heifer calves, yearling heifers, yearling steers), treatment with levamisole HCl or morantel tartrate or not treated, correlation with worm populations, worm egg counts, weight gains, and feed conversion efficiencies, possible economic advantage of treatment
- Trichostrongylus axei*, *illus.*
Stringfellow, F., 1977, *Proc. Helminth. Soc. Washington*, v. 44 (1), 76-81
monospecific and dual species infections of *Ostertagia ostertagi* and *Trichostrongylus axei*, calves, histochemical studies of abomasal tissue
- Trichostrongylus axei*
Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Trichostrongylus axei*
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Trichostrongylus axei*
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole

- Trichostrongylus axei*
Thornton, J. E.; et al., 1973, *J. Wildlife Dis.*, v. 9 (2), 160-162
Antelope cervicapra (abomasum, small intestine): Texas
- Trichostrongylus axei*
Valenzuela, G.; et al., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago, intestino delgado): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Trichostrongylus axei*
Williams, J. C.; and Knox, J. W., 1976, *Am. J. Vet. Research*, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Trichostrongylus axei*
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, *Am. J. Vet. Research*, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Trichostrongylus calcaratus*
Davidson, W. R., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (cecum, large intestine): southeastern United States
- Trichostrongylus calcaratus*
Pursglove, S. R.; et al., 1976, *J. Am. Vet. Med. Ass.*, v. 169 (9), 896-900
intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Trichostrongylus capricola* Ransom, 1907
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (abomasum): vicinity of Nowy Targ, Carpathian Mountains
- Trichostrongylus capricola*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Trichostrongylus capricola*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichostrongylus capricola*
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Trichostrongylus capricola*
Ghadirian, E.; and Arfaa, F., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus capricola* Ransom, 1907
Ianchev, I., 1973, *Izvest. Tsentral. Khelmint. Lab.*, v. 16, 205-220
synonymy
Capreolus capreolus (rennet, small intestine): southern Bulgaria
- Trichostrongylus capricola*
Ingolfsson, A.; and Gislason, G., 1975, *Islen. Landbunadarranns.*, v. 7 (1-2), 3-7
cattle, nautgripa (small intestine): southwestern Iceland, slaughterhouse in Reykjavik
- Trichostrongylus capricola* or *T. vitrinus*
Ingolfsson, A.; and Gislason, G., 1975, *Islen. Landbunadarranns.*, v. 7 (1-2), 3-7
cattle, nautgripa (small intestine): southwestern Iceland, slaughterhouse in Reykjavik
- Trichostrongylus capricola*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Trichostrongylus capricola*
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Trichostrongylus capricola*
Ramajo Martin, V.; and Simon Vicente, F., 1975, *Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C.*, v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Trichostrongylus colubriformis*
Adams, D. B.; and Cripps, A. W., 1977, *Austral. J. Exper. Biol. and Med. Sc.*, v. 55 (5), 509-522
Trichostrongylus colubriformis-infected sheep, cellular changes in intestinal lymph, diminished traffic of lymphocytes in intestinal lymph and reduced numbers of mitogen and nematode antigen-reactive lymphocytes in both blood and intestinal lymph during early stages of infection are closely related to slow development of protective immunity

- Trichostrongylus colubriformis*
Adams, D. B.; and Rothwell, T. L. W., 1977, *Exper. Parasitol.*, v. 42 (1), 121-128
Trichostrongylus colubriformis, guinea pigs, passive transfer of immunity using mesenteric lymph node cells, influence of various factors (immunization schedule for cell donors; size of cell dose transferred; size of challenge dose; age of both cell donors and recipients), rate of worm rejection from recipients
- Trichostrongylus colubriformis*
Anderson, P. J. S.; and Marais, F. S., 1972, *J. South African Vet. Ass.*, v. 43 (3), 271-285
nematodes of sheep and goats, morantel tartrate, efficiency trials
- Trichostrongylus colubriformis*
Askenase, P. W., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (6, Pt. 2), 96-103
immune inflammatory responses to parasites, interconnections between immediate and delayed hypersensitivities, role of basophils, mast cells, and vasoactive amines (*Trichostrongylus colubriformis*; ticks; *Schistosoma mansoni*), workshop report
- Trichostrongylus colubriformis*
Baker, N. F.; and Fisk, R. A., 1977, *Am. J. Vet. Research*, v. 38 (9), 1315-1316
Ostertagia, *Trichostrongylus*, and *Nematodirus*, oxfendazole highly effective against adult stages in sheep
- Trichostrongylus colubriformis*
Bergstrom, R. C.; Kinnison, J. L.; and Werner, B. A., 1977, *Am. J. Vet. Research*, v. 38 (6), 887-888
Trichostrongylus colubriformis, *Eimeria ninakohlyakimovae*, changes in weight gains, feed conversion efficiency, and wool fiber diameter in lambs maintained on 2 different diets (good ration; marginal diet)
- Trichostrongylus colubriformis*
Bergstrom, R. C.; Maki, L. R.; and Werner, B. A., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (2), 171-174
trichostrongylid eggs in cattle or sheep feces, dung beetles (*Aphodius* spp.; *Canthon practicola*) as possible biological control agents, laboratory studies showed decreased eggs in feces when beetles were present
- Trichostrongylus colubriformis* (Giles, 1892)
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Trichostrongylus colubriformis*
Boag, B.; and Thomas, R. J., 1973, *Research Vet. Sc.*, v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Trichostrongylus colubriformis*
Boag, B.; and Thomas, R. J., 1977, *Research Vet. Sc.*, v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Trichostrongylus colubriformis*
Campbell, W. C.; and Thomson, B. M., 1973, *Austral. Vet. J.*, v. 49 (2), 110-111
ensheathed and exsheathed nematode larvae, survival rates after liquid nitrogen freezing, cryoprotective effect of exsheathment; exsheathed larvae of *Trichostrongylus colubriformis* proved uninfective even if they had not been frozen
- Trichostrongylus colubriformis*
Chalmers, K., 1977, *N. Zealand Vet. J.*, v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Trichostrongylus colubriformis*
Ciordia, H.; et al., 1977, *Am. J. Vet. Research*, v. 38 (9), 1335-1339
gastrointestinal parasitism of cattle on fescue pastures fertilized with broiler litter vs. NH_4NO_3 , prevalence, yearly and seasonal variation; parasite burden lower in calves raised on broiler litter-fertilized pastures (where available forage was greater), no significant differences in adult cows nor in calf weight gains
- Trichostrongylus colubriformis*
Coles, G. C.; and Simpkin, K. G., 1977, *Research Vet. Sc.*, v. 22 (3), 386-387
resistance of normal nematode eggs and eggs of benzimidazole-resistant *Haemonchus contortus* and *Trichostrongylus colubriformis* to ovicidal activity of benzimidazoles, observed that eggs from benzimidazole-resistant nematodes are resistant to benzimidazoles, may be useful as simple screen for detecting resistance
- Trichostrongylus colubriformis*
Coop, R. L.; Sykes, A. R.; and Angus, K. W., 1976, *Research Vet. Sc.*, v. 21 (3), 253-258
Trichostrongylus colubriformis, lambs, sub-clinical infection, reduced growth rate and food intake, hypophosphataemia, hypoalbuminaemia, hyperglobulinaemia; possible use of plasma constituents in diagnosis
- Trichostrongylus colubriformis*
Cornwell, R. L., 1975, *Research Vet. Sc.*, v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Trichostrongylus colubriformis*
Corticelli, B.; and Lai, M., 1972, *Parassitologia*, v. 14 (1), 95-96
Ovis musimon (tenue): Sardegna

- Trichostrongylus colubriformis*
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxbendazole, drug efficacy
- Trichostrongylus colubriformis*
Dineen, J. K.; et al., 1977, Internat. J. Parasitol., v. 7 (3), 211-215
Trichostrongylus colubriformis-vaccinated sheep, high level of protection against single-species homologous challenge, lowered level of protection against single-species challenge with *T. vitrinus*, no protection against single-species challenge with *Nematodirus spathiger*, high level of protection against all 3 species to simultaneous challenge with all 3 species, latter suggests that terminal effectors of resistance are immunologically non-specific
- Trichostrongylus colubriformis*
Downey, N. E., 1977, Vet. Rec., v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Trichostrongylus colubriformis*
Düewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Trichostrongylus colubriformis*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichostrongylus colubriformis*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Trichostrongylus colubriformis*
Dyk, V.; and Chroust, K., 1975, Veterinarstvi, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Trichostrongylus colubriformis*
Eichler, D. A., 1973, Brit. Vet. J., v. 129 (6), 533-543
nematodes, sheep (nat. and exper.), calves (exper.), thiophanate, drug efficacy, useful as a broad spectrum anthelmintic
- Trichostrongylus colubriformis*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichostrongylus colubriformis*
Folz, S. D.; Rector, D. L.; and Geng, S., 1976, J. Parasitol., v. 62 (2), 281-285
gastrointestinal nematodes and cestodes, lambs, p-toluoyl chloride phenylhydrazone, efficacy at dose levels of 20, 30, 40, and 50 mg/kg moderate to high
- Trichostrongylus colubriformis*
Gemési, G., 1971, Parasitol. Hungar., v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Trichostrongylus colubriformis*
Ghadirian, E.; and Arfaa, F., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus colubriformis*
Gibson, T. E.; and Parfitt, J. W., 1972, Research Vet. Sc., v. 13 (6), 529-535
Trichostrongylus colubriformis, lambs, increased ability to develop resistance with increasing age, importance of grazing management designed to reduce hazard of infection for young animals
- Trichostrongylus colubriformis*
Gregg, P.; Dineen, J. K.; and Griffiths, D. A., 1976, Vet. Parasitol., v. 2 (4), 363-375
Trichostrongylus colubriformis in guinea pigs and sheep, effects of broad range of gamma-radiation dose levels on viability, infectivity, and pattern of infection
- Trichostrongylus colubriformis*
Grieve, R. B.; Bergstrom, R. C.; and Belden, E. L., 1977, Vet. Parasitol., v. 3 (4), 343-348
dual infection of lambs with *Trichostrongylus colubriformis* and chlamydiae
- Trichostrongylus colubriformis*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Trichostrongylus colubriformis*
Hennessy, D. R.; Prichard, R. K.; and Griffiths, D. A., 1977, Exper. and Molecular Path., v. 27 (2), 143-151
Trichostrongylus colubriformis-infected sheep, thyroxine and thyroxine-binding proteins

- Trichostrongylus colubriformis*
Herlich, H., 1975, Proc. Helminth. Soc. Wash-
ington, v. 42 (2), 135-137
gastrointestinal nematodes, cattle (exper.),
oxibendazole, efficacy against adult and
larval stages
- Trichostrongylus colubriformis*
Herlich, H., 1976, Vet. Parasitol., v. 2 (4),
377-383
Trichostrongylus axei, *T. colubriformis*,
rabbits, efficacy of 8 anthelmintics at dose
rates known to be effective in ruminants,
evaluation of this system as preliminary
screen for testing potential anthelmintics
- Trichostrongylus colubriformis*
Herlich, H., 1977, Am. J. Vet. Research, v. 38
(8), 1247-1248
efficacy of albendazole against gastroin-
testinal nematodes and *Fasciola hepatica* in
cattle (exper.); comparison of critical vs.
controlled tests
- Trichostrongylus colubriformis*
Hogarth-Scott, R. S.; et al., 1976, Research
Vet. Sc., v. 21 (2), 232-237
Haemonchus contortus, *Trichostrongylus colu-*
briformis, thiabendazole-resistant strains,
efficacy of fenbendazole, sheep, implica-
tions for management of helminth disease in
grazing animals with respect to benzimidazole-resistance
- Trichostrongylus colubriformis*
Horak, I. G.; Honer, M. R.; and Schroeder,
J., 1976, J. South African Vet. Ass., v. 47
(4), 247-251
helminths and *Oestrus ovis*, merino sheep,
treated at four-weekly intervals or strate-
gically, live mass gains, wool production
and fecal worm egg counts, compared with
untreated controls: Eastern Transvaal High-
veld
- Trichostrongylus colubriformis*
Horton, G. M. J.; et al., 1977, J. South
African Vet. Ass., v. 48 (2), 99-103
Trichostrongylus colubriformis, lambs
(exper.), effect of host diet low in sele-
nium as compared to diets supplemented with
vitamin E and selenium, pathology, haema-
tological changes, serum enzyme changes
- Trichostrongylus colubriformis*
Hsu, C. K.; and Levine, N. D., 1977, Am. J.
Vet. Research, v. 38 (8), 1115-1119
Haemonchus contortus, *Trichostrongylus colu-*
briformis, development of infective larvae
under cyclic vs. constant conditions of
temperature and humidity, degree-day concept
appears to be applicable
- Trichostrongylus colubriformis*
Huxtable, C. R.; and Rothwell, T. L. W., 1975,
Austral. J. Exper. Biol. and Med. Sc., v. 53
(6), 437-445
Trichostrongylus colubriformis, guinea pig
small intestinal mucosa, basophils and
eosinophils at site of infection, ultrastruc-
tural changes as compared with bone marrow
and buffy coat cells
- Trichostrongylus colubriformis* (Giles, 1892)
Ianchev, I., 1973, Izvest. Tsentral. Khelmint.
Lab., v. 16, 205-220
Capreolus capreolus (rennet, small intes-
tine): southern Bulgaria
- Trichostrongylus colubriformis*
Kelly, J. D.; et al., 1975, Research Vet.
Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole
against naturally acquired *Dictyocaulus*
filaria infection associated with concu-
rent infection of gastro-intestinal nema-
todes in sheep
- Trichostrongylus colubriformis*
Kelly, J. D.; et al., 1977, Research Vet. Sc.,
v. 22 (2), 161-168
Haemonchus contortus, *Trichostrongylus colu-*
briformis, resistant or susceptible to thia-
bendazole, sheep, effect of route of adminis-
tration (oral, intra-ruminal or intra-abo-
masal) on efficacy of benzimidazole anthel-
mintics
- Trichostrongylus colubriformis*
Kennedy, T. J.; and Todd, A. C., 1975, Am. J.
Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy
of fenbendazole at dose levels of 3.5, 5.0,
and 7.5 mg/kg of body weight
- Trichostrongylus colubriformis*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A.,
1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect
of breed and birth date on parasite acqui-
sition: Clay Center, Nebraska
- Trichostrongylus colubriformis*
Kozdon, O.; and Zajicek, D., 1976, Vet. Med.,
Praha, v. 49, v. 21 (11), 693-702
nematodes, sheep under natural field con-
ditions, seasonal distribution as modified
by dehelminthization, possible management
strategies for effective timing of dehel-
minthization: Western Bohemia
- Trichostrongylus colubriformis*
Larsh, J. E., jr.; and Weatherly, N. F., 1975,
Advances Parasitol., v. 13, 183-222
principles of delayed (cellular) hypersensi-
tivity, cell-mediated immunity against para-
sitic worms, extensive review
- Trichostrongylus colubriformis*
Le Jambre, L. F.; and Royal, W. M., 1976,
Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally
infected Angora goats and Merino sheep
grazing intraspecifically or in mixed ex-
perimental paddocks (fecal egg counts showed
no significant within-host differences);
increased resistance of sheep to all worms
except *Nematodirus* resulted in a signifi-
cantly lower worm burden for sheep: North-
ern Tablelands of New South Wales

- Trichostrongylus colubriformis*
Levai, J.; Lengyel, A.; and Banki, G., 1972, *Med. Universalis*, v. 5 (1), 31-34
clinical trials using levamisole to treat human intestinal helminths, good results against *Ascaris*, *Ancylostoma*, *Trichostrongylus*, ineffective against *Trichuris trichiura*: Hungary
- Trichostrongylus colubriformis*
Lyons, E. T.; et al., 1975, *Am. J. Vet. Research*, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Trichostrongylus colubriformis*
McConnell, E. E.; et al., 1974, *Onderstepoort J. Vet. Research*, v. 41 (3), 97-168
pathological and parasitological survey of 100 free-ranging chacma baboons
Papio ursinus (small intestine): Kruger National Park, Transvaal
- Trichostrongylus colubriformis*
Niec, R.; et al., 1976, *Gac. Vet.*, Buenos Aires (315), v. 38, 457-466
gastrointestinal nematodes, sheep, effect of thiabendazole drenches on buildup of host resistance; might be advisable to accept moderate degree of parasitism in sheep up to 9-10 months of age, avoid unnecessary antihelminthic treatment that could prevent normal buildup of resistance
- Trichostrongylus colubriformis*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Roln. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Trichostrongylus colubriformis*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Ovis aries
Sus scrofa
all from Chile
- Trichostrongylus colubriformis*
Okon, E. D.; and Emyeenih, U. K., 1975, *Bull. Office Internat. Epizoot.*, v. 83 (111-112), 1139-1144
Haemonchus contortus, *Trichostrongylus colubriformis*, lambs grazed on contaminated pastures, control by pasture rotation
- Trichostrongylus colubriformis* (Giles)
Pestter, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Gazella thomsonii (gut)
G. granti (abomasum, small intestine)
Alcelaphus buselaphus cokei (digestive tract)
Connochaetes taurinus (small intestine)
all from Kenya
- Trichostrongylus colubriformis*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Trichostrongylus colubriformis*
Prosl, H., 1976, *Ztschr. Parasitenk.*, v. 50 (2), 214
Rhesus affe
- Trichostrongylus colubriformis*
Ramajo Martin, V.; and Simon Vicente, F., 1975, *Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C.*, v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Trichostrongylus colubriformis*
Reid, J. F. S., 1976, *Vet. Rec.*, v. 98 (25), 496-499
gastrointestinal nematodes, coccidiosis, diarrhoea of sheep, age and seasonal factors: Britain
- Trichostrongylus colubriformis*
Reveron, A. E.; et al., 1971, *Proc. Nutrition Soc.*, v. 30 (3), 83A-84A
Trichostrongylus colubriformis, lambs (exper.), effects of nematode parasitism on calcium and phosphorus retention, results show that infestation adversely affected bone mineralization with marked decrease in absorption and retention of phosphorus but little effect on calcium metabolism
- Trichostrongylus colubriformis*
Rose, J. H., 1971, *Symposia Brit. Soc. Parasitol.*, v. 9, 109-121
gastrointestinal nematodes and lungworms of farm animals, isolation and maintenance in vivo, extensive review
- Trichostrongylus colubriformis*
Rose, J. H., 1973, *Research Vet. Sc.*, v. 14 (3), 326-333
Ostertagia circumcincta, *O. ostertagi*, *Hyostromyulus rubidus*, culture from infective larva to adult worm in WAE medium, other species of gastrointestinal nematodes underwent limited development in this medium or a modification thereof
- Trichostrongylus colubriformis*
Roseby, F. B., 1977, *Austral. J. Agric. Research*, v. 28 (1), 155-164
Trichostrongylus colubriformis, sheep, effects on characteristics of digesta flow and fermentation, derangements of nitrogen metabolism
- Trichostrongylus colubriformis*
Roseby, F. B., 1977, *Austral. J. Agric. Research*, v. 28 (4), 713-719
Trichostrongylus colubriformis, sheep, glucose metabolism, rate of glucose synthesis higher in infested sheep compared with controls on same feed intake

- Trichostrongylus colubriformis*
Rothwell, T. L. W.; et al., 1976, *Vet. Parasitol.*, v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Trichostrongylus colubriformis*
Rothwell, T. L. W.; and Griffiths, D. A., 1977, *J. Parasitol.*, v. 63 (4), 761-762
Trichostrongylus colubriformis, kinetics of expulsion from previously uninfected, reinfected, and vaccinated guinea pigs compared
- Trichostrongylus colubriformis*
Rothwell, T. L. W.; and Huxtable, C. R., 1976, *Austral. J. Exper. Biol. and Med. Sc.*, v. 54 (4), 329-335
Trichostrongylus colubriformis, guinea pigs, basophil leucocytes in cutaneous hypersensitivity reactions
- Trichostrongylus colubriformis*
Rothwell, T. L. W.; Love R. J.; and Goodrich, B. S., 1977, *Internat. Arch. Allergy and Applied Immunol.*, v. 53 (1), 93-95
Trichostrongylus colubriformis, guinea pigs, prostaglandins not inducing immune expulsion of worms from intestine; prostaglandin synthetase inhibitors not inhibiting expulsion by immune guinea pigs
- Trichostrongylus colubriformis*
Sharpe, M. J.; and Lee, D. L., 1977, *Parasitology*, v. 75 (2), xvi [Abstract]
Trichostrongylus colubriformis, *Nematospiroides dubius*, effect of levamisole on adenylate energy charge and on levels of acetylcholinesterase
- Trichostrongylus colubriformis*, *illus.*
Sobrero, R.; and Goffredo, G., 1972, *Parassitologia*, v. 14 (1), 193-198
Trichostrongylus colubriformis longispiculatus [nov. var.], distinction from ovine *T. colubriformis* by mathematical analysis of measurements
Camelus dromedarius
- Trichostrongylus colubriformis*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, *Austral. J. Agric. Research*, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Trichostrongylus colubriformis*
Sykes, A. R.; and Coop, R. L., 1976, *J. Agric. Sc.*, v. 86 (3), 507-515
Trichostrongylus colubriformis larvae, lambs (exper.), food intake and utilization after parasitic damage to small intestine, body weight changes, reduced host mineral metabolism
- Trichostrongylus colubriformis*
Sykes, A. R.; and Coop, R. L., 1976, *Proc. Nutrition Soc.*, v. 35 (1), 13A-14A [Abstract]
Trichostrongylus colubriformis, *Ostertagia circumcincta*, sheep (exper.), effects of extensive chronic parasitism on food intake and utilization by growing lambs, deposition of nitrogen and fat severely impaired and food intake reduced by 9%
- Trichostrongylus colubriformis*
Symons, L. E. A.; and Jones, W. O., 1975, *Austral. J. Agric. Research*, v. 26 (6), 1063-1072
Trichostrongylus colubriformis, sheep, radioisotope-labelled leucine for measurement of protein synthesis in muscles, liver and wool follicles, anorexia as major cause of protein metabolism changes
- Trichostrongylus colubriformis*
Theodorides, V. J.; et al., 1973, *Brit. Vet. J.*, v. 129 (6), xcvi-xcviii
oxibendazole, outstanding efficacy against gastrointestinal parasites in domestic and laboratory animals (nat. and exper.), well tolerated with no effects on host reproduction
- Trichostrongylus colubriformis*
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Trichostrongylus colubriformis*
Theodorides, V. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Trichostrongylus colubriformis*
Thornton, J. E.; et al., 1973, *J. Wildlife Dis.*, v. 9 (2), 160-162
Antilope cervicapra (small intestine): Texas
- Trichostrongylus colubriformis*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Trichostrongylus colubriformis*
Valenzuela, G.; et al., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago, intestino delgado): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Trichostrongylus colubriformis*
Vilimszky, Z.; and Szigetine Toth, N., 1971, *Parasitol. Hungar.*, v. 4, 97-101
Trichostrongylus colubriformis, increased incidence in human infections revealed large endemic focus, people living in close contact with ruminants, successfully treated with thiabendazole: Borsod-Abaúj-Zemplén county, Hungary

- Trichostrongylus colubriformis* (Giles, 1892)
 Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
 nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karroo
- Trichostrongylus colubriformis* (Giles, 1892), *illus.*
 Wertheim, G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 735-762
 redescription
Hystrix indicus (intestin grele): pres de Bet Alfa, Israel
- Trichostrongylus colubriformis*
 Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
 failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Trichostrongylus colubriformis*
 Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
 gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Trichostrongylus colubriformis*
 Wilson, D. E.; and Hirst, S. M., 1977, Wild-life Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Trichostrongylus colubriformis*
 Zajicek, D.; and Kozdon, O., 1977, Veterinarstvi, v. 27 (6), 257-258
 nematodes, sheep, relation of dehelminthization with pyrantel HCl, helmantac and nilverm to nematode incidence on pastures, three-year study, overall decrease
- Trichostrongylus colubriformis longispiculatus* [nov. var.], *illus.*
 Sobrero, R.; and Goffredo, G., 1972, Parassitologia, v. 14 (1), 193-198
Trichostrongylus colubriformis longispiculatus [nov. var.], distinction from ovine *T. colubriformis* by mathematical analysis of measurements
Camelus dromedarius
- Trichostrongylus dosteri*
 Eve, J. H.; and Kellogg, F. E., 1977, J. Wild-life Management, v. 41 (2), 169-177
 technique for using intensity of abomasal parasite infections as an index to deer (*Odocoileus virginianus*) density: south-eastern United States
- Trichostrongylus falculatus* Ransom, 1911
 Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
 nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karroo
- Trichostrongylus falculatus*
 Young, E.; et al., 1973, Research J. National Parks Republic South Africa (16), 195-198
Antidorcas marsupialis (duodenum): Mountain Zebra National Park near Cradock, Cape Province
- Trichostrongylus lerouxi* Biocca, Chabaud, and Ghadirian, 1974
 Ghadirian, E., 1977, Am. J. Trop. Med. and Hyg., v. 26 (6, part 1), 1212-1213
 distribution and prevalence in man (stool): Iran
- Trichostrongylus longispicularis*
 Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, Bol. Chileno Parasitol., v. 29 (3-4), 99-102
Bos taurus: Chile
- Trichostrongylus longispicularis*
 Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
 intestinal nematodes of *Odocoileus virginianus*, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: south-eastern United States
- Trichostrongylus longispicularis* (Gordon, 1933), *illus.*
 Tongson, M. S.; Arambulo, P. V. III; and Trovela, V., 1975, Philippine J. Vet. Med., v. 13 (1-2), 163-169
Trichostrongylus longispicularis, morphological features, body measurements
 Brahman cattle (duodenum): Philippines
- Trichostrongylus longispicularis*
 Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
 meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
 cerdos (intestino delgado): Planta Faenadora de Carnes Socogro, Valdivia, Chile
- Trichostrongylus longispicularis* (Gordon, 1933), *illus.*
 Valenzuela, G.; Oberg, C.; and Loyola, R., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 45-46
 morphology, description of male porcinos: Valdivia, Chile
- Trichostrongylus medius*
 Pav, J.; and Zajicek, D., 1974, Veterinarstvi, v. 24 (11), 517-520
Lyrus tetrix: CSSR

- Trichostrongylus minor*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (1), 65-77
roe deer (digestive tract): Czechoslovakia
- Trichostrongylus minor*
Dyk, V.; and Chroust, K., 1974, Acta Vet. Brno, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Capreolus capreolus (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichostrongylus minor*
Dyk, V.; and Chroust, K., 1975, Vet. Parasitol., v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Capreolus capreolus: Czechoslovakia
- Trichostrongylus orientalis*
Ghadirian, E.; and Arfaa, F., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus pergracilis* (Cobbold 1873)
Railliet and Henry 1909
Radhakrishnan, C. V.; and Ebrahimina, A., 1975, J. Vet. Fac. Univ. Tehran, v. 30 (4), 1-4
as syn. of *Trichostrongylus tenuis*
- Trichostrongylus probolurus*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichostrongylus probolurus*
Ghadirian, E.; and Arfaa, F., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus probolurus*
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella granti (small intestine)
Alcelaphus buselaphus cokei (digestive tract)
Gazella thomsonii (gut)
all from Kenya
- Trichostrongylus probolurus*
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antilope cervicapra (small intestine): Texas
- Trichostrongylus retortaeformis*
Ford, G. E., 1971, Immunology, v. 21 (6), 1073-1078
Trichostrongylus retortaeformis, heterogeneity of allergens and of homocytotropic antibody responses to them in rabbits
- Trichostrongylus retortaeformis*
Ford, G. E., 1971, Vet. Rec. (4883), v. 89 (26), 692 [Letter]
Trichostrongylus retortaeformis sheathed third-stage larvae, rabbits, infectivity increased for at least 2 to 4 weeks after disposition of larvae-containing feces, indicates maturation requirement
- Trichostrongylus retortaeformis*
Haupt, W.; and Hartung, J., 1977, Monatsh. Vet.-Med., v. 32 (9), 339-341
Lepus europaeus (Dunndarm): Leipzig area
- Trichostrongylus retortaeformis*
Kutzer, E.; and Frey, H., 1976, Berl. u. Münch. Tierärztl. Wchschr., v. 89 (24), 480-483
Lepus europaeus: Austria
- Trichostrongylus retortaeformis*
Kutzer, E.; and Frey, H., 1976, Ztschr. Parasitenk., v. 50 (2), 213-214
Lepus europaeus
- Trichostrongylus retortaeformis* (Zeder, 1800)
Looss, 1915
Mozgovoi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Lepus timidus (small intestine): Karelia
- Trichostrongylus retortaeformis*
Wilson, P. A. G., 1976, J. Zool., London, v. 179 (1), 135-141
Trichostrongylus retortaeformis growth patterns, moulting cycle, population growth profile
- Trichostrongylus rugatus*
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and *Oestrus ovis*, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Trichostrongylus rugatus*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Trichostrongylus rugatus* Monnig, 1925
Verster, A.; Imes, G. D., jr.; and Smit, J. P. J., 1975, Onderstepoort J. Vet. Research, v. 42 (1), 29-31
Damaliscus dorcas dorcas (small intestine): captured at Bontebok National Park, Swellendam and transferred to the National Zoological Gardens, Pretoria

- Trichostrongylus rugatus* Monnig, 1925
Viljoen, J. H., 1969, Onderstepoort J. Vet. Research, v. 36 (2), 233-263
nematodes of sheep, epizootiology: seasonal incidence and worm burden in relation to temperature and rainfall at three sites, availability of live infective larvae on pasture, drenching recommendations: the Karroo
- Trichostrongylus sigmodontis*
Kinsella, J. M., 1974, Am. Mus. Novitates (2540), 1-12
Sigmodon hispidus (small intestine): Florida
- Trichostrongylus skrjabini*
Ghadirian, E.; and Arfaa, F., 1975, Am. J. Trop. Med. and Hyg., v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus tenuis* (Mehlis, 1846) Railliet & Henry, 1909, illus.
Alcaino, H.; and Gorman, T., 1969, Bol. Chileno Parasitol., v. 24 (3-4), 157-159
Gallus gallus domesticus (ciego): Santiago, Chile
- Trichostrongylus tenuis*
Cervenka, J.; Zajicek, D.; and Nydl, J., 1975, Veterinarstvi, v. 25 (6), 263-264
helminths, geese, Mebendazole
- Trichostrongylus tenuis* (Mehlis, 1846)
Forrester, D. J.; et al., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 55-59
Grus canadensis tabida (duodenum): Florida
- Trichostrongylus tenuis*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (ceca): Florida
- Trichostrongylus tenuis* Mehlis, 1846
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelminth. Lab., v. 15, 109-133
Anser anser
A. albifrons
Anas penelope
A. querquedula
(caecum of all): all from Bulgaria
- Trichostrongylus tenuis*
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, Proc. 3. National Wild Turkey Symp., 27-32
Meleagris gallopavo silvestris: south-eastern United States
- Trichostrongylus tenuis*
Radhakrishnan, C. V.; and Ebrahimina, A., 1975, J. Vet. Fac. Univ. Tehran, v. 30 (4), 1-4
brief description, syn.: *Trichostrongylus pergracilis* (Cobbold 1873) Railliet and Henry, 1909
chickens (ceca): Darab, Fars Province, Iran
- Trichostrongylus triramosus* Schulz, 1931
Mozgovi, A. A.; et al., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 95-103
Lepus timidus (small and large intestine): Karelia
- Trichostrongylus vitrinus*
Baker, N. F.; and Fisk, R. A., 1977, Am. J. Vet. Research, v. 38 (9), 1315-1316
Ostertagia, *Trichostrongylus*, and *Nematodirus*, oxfendazole highly effective against adult stages in sheep
- Trichostrongylus vitrinus* Looss, 1905
Bezubik, B.; Stankiewicz, M.; and Baginska, G., 1969, Acta Parasitol. Polon., v. 17 (1-19), 25-37
brief description
sheep (abomasum, small intestine): vicinity of Nowy Targ, Carpathian Mountains
- Trichostrongylus vitrinus*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Trichostrongylus vitrinus*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Trichostrongylus vitrinus*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Trichostrongylus vitrinus*
Cornwell, R. L., 1975, Research Vet. Sc., v. 18 (1), 1-5
yearly pattern of infection with gastrointestinal nematodes in young fattening lambs at pasture, degree of infection and incidence of different genera: United Kingdom
- Trichostrongylus vitrinus*
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon (tenue): Sardegna
- Trichostrongylus vitrinus*
Dineen, J. K.; et al., 1977, Internat. J. Parasitol., v. 7 (3), 211-215
Trichostrongylus colubriformis-vaccinated sheep, high level of protection against single-species homologous challenge, lowered level of protection against single-species challenge with *T. vitrinus*, no protection against single-species challenge with *Nematodirus spathiger*, high level of protection against all 3 species to simultaneous challenge with all 3 species, latter suggests that terminal effectors of resistance are immunologically non-specific

- Trichostrongylus vitrinus*
Downey, N. E., 1977, *Vet. Rec.*, v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Trichostrongylus vitrinus*
Dyk, V.; and Chroust, K., 1974, *Acta Vet. Brno*, v. 43 (2), 123-131
helminths and coccidians of *Ovis ammon musimon* and *Capreolus capreolus*, intensity variation with age of host, lack of evidence for parasite exchange between mouflons and roe deer
Ovis ammon musimon (digestive tract): School Forest Enterprise, University of Agriculture Brno, Krtiny
- Trichostrongylus vitrinus*
Dyk, V.; and Chroust, K., 1975, *Vet. Parasitol.*, v. 1 (2), 145-150
coccidia and helminths in mouflon and roe deer, incidence and intensity, possible cross transmission, implications for game management
Ovis ammon musimon: Czechoslovakia
- Trichostrongylus vitrinus*
Dyk, V.; and Chroust, K., 1975, *Veterinarstvi*, v. 25 (7), 315-317
helminths, incidence by age of host, problem in mouflon husbandry: Brno oblast
- Trichostrongylus vitrinus*
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*, v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichostrongylus vitrinus*
Ghadirian, E.; and Arfaa, F., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (6, pt. 1), 935-941
Trichostrongylus spp., man and domestic animals, present status in Iran, means of transmission
- Trichostrongylus vitrinus* Looss, 1905
Ianchev, I., 1973, *Izvest. Tsentral. Khelmit. Lab.*, v. 16, 205-220
Capreolus capreolus (rennet, small intestine): southern Bulgaria
- Trichostrongylus vitrinus* or *T. capricola*
Ingolfsson, A.; and Gislaason, G., 1975, *Islan. Landbunadarranns.*, v. 7 (1-2), 3-7
cattle, nautgripa (small intestine): southwestern Iceland, slaughterhouse in Reykjavik
- Trichostrongylus vitrinus*
Kistner, T. P.; and Wyse, D., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Trichostrongylus vitrinus*
Musila, V., 1976, *Veterinarstvi*, v. 26 (6), 264
helminths of fallow deer, incidence:
Zehusice enclosure
- Trichostrongylus vitrinus*
Novy, H., 1976, *Veterinarstvi*, v. 26 (6), 263
helminths of white deer, incidence:
Zehusice enclosure
- Trichostrongylus vitrinus*
Nowosad, B., 1975, *Zeszyty Nauk. Akad. Rolnicz. Krakow.* (98), *Zootech.* (15), 219-251
lambs, experimental infection with various doses and combinations of gastrointestinal helminths, lowered yield of various cuts at slaughter
- Trichostrongylus vitrinus*
Oberger, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus
Ovis aries
all from Chile
- Trichostrongylus vitrinus*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Trichostrongylus vitrinus*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Trichostrongylus vitrinus*
Ramajo Martin, V.; and Simon Vicente, F., 1975, *Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C.*, v. 1, 137-163
Trichostrongylidae, sheep, development on pastures, climatic factors: Salamanca
- Trichostrongylus vitrinus*
Reid, J. F. S., 1976, *Vet. Rec.*, v. 98 (25), 496-499
gastrointestinal nematodes, coccidiosis, diarrhoea of sheep, age and seasonal factors: Britain
- Trichostrongylus vitrinus*
Rothwell, T. L. W.; et al., 1976, *Vet. Parasitol.*, v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of *Strongylida*, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific

- Trichostrongylus vitrinus*
Scnweisgut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Trichostrongylus vitrinus*
Southcott, W. H.; Major, G. W.; and Barger, I. A., 1976, Austral. J. Agric. Research, v. 27 (2), 277-286
sheep nematodes, seasonal pasture contamination, availability to infect grazing sheep, overwintering: Armidale, New South Wales
- Trichostrongylus vitrinus*
Valenzuela, G.; et al., 1977, Bol. Chileno Parasitol., v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (estomago, intestino delgado): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Trichostrongylus vitrinus* (Loos, 1905), illus.
Valenzuela, G.; Oberg, C.; and Loyola, R., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 21-22
morphometric data of adult males
porcinos (intestino delgado): Valdivia, Chile
- Trichosyringida* Goodey, 1963
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Enoplia
- Trichotravassosia* Lent et Freitas, 1938
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Trichuriasis*
Bekhli, A. F.; Braude, M. B.; and Bolotina, L. A., 1976, Khimiko-Farm. Zhurnal, v. 10 (6), 29-31
synthesis method, new anthelmintic for trichuriasis (bemosate), no trials against parasite
- Trichuriasis*
Biagi, F., 1975, Progr. Drug Research, v. 19 23-27
human hookworm and trichuriasis successfully treated with bitoscanate in clinical trials; mixed results when used to treat ascariasis in comparison trials with dithiazanine and piperazine
- Trichuriasis*
de Carneri, I.; de Dominici, M.; and Carozzi, A., 1973, Riv. Parasitol., Roma, v. 34 (3), 213-218
Enterobius vermicularis, school children, incidence, no correlation with age or sex or infection with trichuriasis, some correlation with socio-economic status: Pavia Province, Italy
- Trichuriasis*
Egorov, V. I., 1968, Botan. i Zool. Issled. Dal'nem Vostoke, v. 2, 250-255
ascariasis, trichuriasis, oesophagostomiasis, swine, single and mixed infections in various combinations, control studies, best system used dehelminthization 4 times a year
- Trichuriasis*
Egorov, V. I., 1968, Botan. i Zool. Issled. Dal'nem Vostoke, v. 2, 256-260
ascariasis, trichuriasis, swine, epizootiology, seasonal distribution of depositing eggs and larvae in barns and pastures: Primor'e
- Trichuriasis*
Nitzulescu, V.; et al., 1970, Pediatria, Bucuresti, v. 19 (4), 373-375
trichuriasis in humans, clinical trials using bemarsal, good results
- Trichuriasis*
Pawlowski, Z.; and Chodera, L., 1975, Polski Tygod. Lekar., v. 30 (24), 1035-1037
ascariasis, trichuriasis, ancylostomiasis, humans, clinical trials show mebendazole to be effective drug for all 3 infections: Poznan, Poland
- Trichuriasis*
Radermecker, M.; et al., 1974, Internat. Arch. Allergy and Applied Immunol., v. 47 (2), 285-295
various human helminthic or protozoal infections, serum IgE concentration, IgE level often raised in parasitosis with prominent tissue phases and remains normal with helminths restricted to lumen of digestive tract, IgE level tends to increase significantly and rapidly following specific treatment and then to decrease slowly and return to normal in a few months
- Trichuriasis*
Shah, P. M.; Junnarkar, A. R.; and Khare, R. D., 1975, Progr. Drug. Research, v. 19, 136-146
impact of periodic deworming on nutritional status of preschool children, evaluation of including periodic deworming in National Nutrition Programmes of India
- Trichuriasis*
Trifonov, T.; and Velkov, D., 1975, Vet. Sbirka, v. 73 (11), 16-17
ascariasis, trichuriasis, oesophagostomiasis, pigs, dehelminthization with suiverm or hygomix in feed, economically effective
- Trichuriasis*
Vakil, B. J.; and Dalal, N. J., 1975, Progr. Drug Research, v. 19, 166-175
comparative efficacy of newer anthelmintics in treating various human intestinal helminths, review
- Trichuris*
Baines, D. M.; and Colegrave, A. J., 1977, Vet. Rec., v. 100 (11), 217-219
gastrointestinal helminths, sheep, thiophanate, productivity and tolerance trials, compared with thiabendazole and tetramisole: England and Wales

- Trichuris
Biagi, F.; Smyth, J.; and Gonzalez, C., 1974, Prensa Med. Mexicana, v. 39 (1-2), 51-53
human intestinal helminths, successful clinical trials using mebendazole, drug well tolerated with minimal side effects: Mexico
- Trichuris
Boag, B.; and Thomas, R. J., 1975, Research Vet. Sc., v. 19 (3), 293-295
sheep nematodes, population dynamics, field studies, level of larval mortality may vary from year to year with prevailing climatic conditions, 'spring rise' in ewes is major source of pasture contamination causing wave of lamb infections in late August and September
- Trichuris
Bruch, K.; and Haas, J., 1976, Ann. Trop. Med. and Parasitol., v. 70 (2), 205-211
Ascaris, hookworm, Trichuris, human, single dose fenbendazole compared with pyrantel: Liberia
- T[rich]uris
Brunsdon, R. V., 1976, N. Zealand J. Exper. Agric., v. 4 (3), 275-279
lambs, effectiveness of single thiabendazole drench at weaning in controlling build-up of trichostrongyle worm burdens, relative importance of various sources of pasture contamination (overwintered larvae; larvae deposited by ewes and lambs in pre-weaning period; larvae deposited by lambs at weaning)
- Trichuris
Crowley, J. W.; et al., 1976, Am. J. Vet. Research, v. 37 (11), 1285-1286
nematodes, cattle, oxbendazole, drug efficacy
- Trichuris
Danielli, Y.; and Neuman, M., 1975, Refuah Vet., v. 32 (4), 94-95, 153-154
mixed parasites, cattle, good results following repeated chemotherapy: Birkat Ata
- Trichuris
Dey-Hazra, A., 1976, Ztschr. Parasitenk., v. 50 (2), 198
helminths, pigs, mode of pathogenicity, review
- Trichuris
Duewel, D., 1977, Cahiers Bleus Vet. (26), 201-215
fenbendazole, efficacy against nematodes in various animals, useful as broad spectrum anthelmintic, mechanism of action, pharmacokinetics, metabolism, toxicology
- Trichuris
Forstner, M. J.; Kopp, H.; and Wiesner, H., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (9), 180-183
nematodes of ruminants, mebendazole, good results: Hellabrunn Zoo, Munich
- Trichuris
Gilman, R. H.; Davis, C.; and Fitzgerald, F., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (4), 313-316
Trichuris infection and amoebic dysentery in Orang Asli aborigine children, comparison of two diseases, results support hypothesis that heavy Trichuris infection itself is responsible for a symptom complex: Malaysia
- Trichuris
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (1), 9-15
nematode parasitism, calves (Holstein x Zebu), female to male ratio of worms, higher number of females: State of Minas Gerais, Brazil
- Trichuris
Hausfater, G.; and Watson, D. F., 1976, Nature, London (5570), v. 262, 688-689
emission of parasite ova (primarily Trichuris and Trichostrongylus) by Papio cynocephalus in relation to host social and reproductive condition, high-ranking adult males had higher egg emission than more subordinate individuals, sexually cycling females had higher emissions than anoestrous females
- Trichuris
Kennedy, T. J.; and Todd, A. C., 1975, Am. J. Vet. Research, v. 36 (10), 1465-1467
gastrointestinal parasites, lambs, efficacy of fenbendazole at dose levels of 3.5, 5.0, and 7.5 mg/kg of body weight
- Trichuris
Krishna Iyer, P. P.; and Peter C. T., 1975, Kerala J. Vet. Sc., v. 5 (2), 121-123
gastrointestinal nematodes, goats, methyridine
- Trichuris
McFarlane, H., 1976, Proc. Nutrition Soc., v. 35 (3), 263-272
parasitic infestation, preschool children, malnutrition and impaired immune response, brief review comment: Nigeria
- Trichuris
Makkar, M. S.; Joshi, H. C.; and Gupta, I., 1974, Indian J. Animal Research, v. 8 (2), 75-78
Haemonchus contortus, other nematodes, experimentally or naturally infected sheep, nitroxylnil highly effective, critical testing; in vitro testing against H. contortus
- Trichuris
Most, H., 1972, N. England J. Med., v. 287 (10), 495-498; (14), 698-702
common parasitic infections of man encountered in the United States, recommendations for treatment, review

- Trichuris**
Neppert, J.; and Warns, C. M., 1974, *Tropenmed. u. Parasitol.*, v. 25 (4), 492-497
sera from Liberians with various helminthic infections, cross reactions with antigens from *Ascaris*, hookworm, *Onchocerca*, *Dirofilaria immitis*, closed hexagon immunodiffusion, complement fixation reaction, indirect haemagglutination
- Trichuris**
Sewell, M. M. H., 1973, *Vet. Rec.*, v. 94 (14), 371-372 [Letter]
anthelmintic treatment of ewes around lambing time to lessen gastrointestinal nematode worm burden in their lambs, variable results, review
- Trichuris**
Sewell, M. M. H.; and Urquhart, H. R., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (4), 287 [Demonstration]
technique for quantitative recovery of ascarid and *Trichuris* eggs from 100g samples of soil
- Trichuris**
Tantengco, V. O.; et al., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 524-533
Ascaris, *Trichuris*, hookworm infections apparently not contributory cause of nutritional anemia in schoolchildren: Philippine Islands
- Trichuris**
Theodorides, V. J.; et al., 1976, *Experientia*, v. 32 (6), 702-703
anthelmintic activity of albendazole against liver flukes, tapeworms, lung and gastrointestinal roundworms, brief preliminary report
- Trichuris**
Todd, A. C.; et al., 1976, *Am. J. Vet. Research*, v. 37 (4), 439-441
nematodes, calves (exper.), mixed infections, controlled evaluation of fenbendazole treatment
- Trichuris**
Turner, K. J.; Baldo, B. A.; and Anderson, H. R., 1975, *Internat. Arch. Allergy and Applied Immunol.*, v. 48 (6), 784-799
humans, serum IgE levels, no significant correlation with faecal egg counts to hookworm, *Ascaris lumbricoides*, and *Trichuris*, incidence of IgE antibodies to *Ascaris lumbricoides* not correlated with incidence of asthma but significantly elevated in patients with chronic obstructive lung disease, hypersensitivity to *Ascaris* apparently not factor of importance in etiology of asthma in this area: Highland area of Papua-New Guinea
- Trichuris**
Varshney, T. R.; and Singh, Y. P., 1976, *Indian Vet. J.*, v. 53 (9), 672-676
Haemonchus contortus, trichostrongyles, *Trichuris*, lambs, critical trial, efficacy and economic value, various anthelmintics: Central Sheep and Wool Research Station, Pashulok Rishikesh (Dehradun)
- Trichuris**
Zeakes, S. J.; et al., 1976, *Am. J. Vet. Research*, v. 37 (6), 709-710
cattle nematodes, efficacy of coumaphos crumbles and naftalofos boluses
- Trichuris sp.**
Bali, M. K.; and Singh, R. P., 1976, *Indian J. Animal Research*, v. 10 (2), 111-112
Haemonchus contortus, *Oesophagostomum* sp., *Trichostrongylus* sp., *Trichuris* sp., sheep, goats, morantel tartrate, good results against all parasites except for *Trichuris* sp.
- Trichuris sp.**
Cabrera, B. D., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 50-55
Rattus rattus (feces): Leyte, Philippines
- Trichuris spp.**
Colglazier, M. L.; et al., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (2), 145-150
gastrointestinal helminths, sheep, pasture trials, levamisole and thiabendazole, good to fair control except with *Trichuris* spp. and *Moniezia expansa*
- Trichuris spp.**
Downey, N. E., 1976, *Vet. Rec.*, v. 99 (14), 267-270
nematodes, calves (natural infections), oxfendazole compared with levamisole (oxfendazole showed higher efficacy than levamisole against *Ostertagia* spp., similar efficacy against other species), both drugs increased calves' weight gains
- Trichuris spp.**
Downey, N. E., 1977, *Vet. Rec.*, v. 101 (13), 260-263
gastrointestinal nematodes, sheep, controlled trial of oxfendazole before and after lambing, reduced egg output in ewes, high efficacy against nematodes in lambs, compared with levamisole
- Trichuris sp.**
File, S. K.; McGrew, W. C.; and Tutin, C. E. G., 1976, *J. Parasitol.*, v. 62 (2), 259-261
Pan troglodytes schweinfurthii (feces): Gombe National Park, Tanzania
- Trichuris sp.**
Graves, I. L.; Adams, W. H.; and Pyakural, S., 1975, *Am. J. Vet. Research*, v. 36 (6), 843-846
Babesia bigemina, *Bos grunniens* moved from high to low altitude and challenged with influenza A viruses, hemolytic anemia, possible explanations, death due to *Fasciola hepatica* and *F. gigantica*, incidental finding of *Bunostomum* sp., *Trichuris* sp., *Neoascaris vitulorum*, *Dictyocaulus* sp., coccidia, some reasons for poor survival of yaks at low altitude: Nepal
- Trichuris sp.**
Grundmann, A. W.; and Lombardi, P. S., 1976, *Proc. Helminth. Soc. Washington*, v. 43 (1), 39-46
Ochotona princeps uinta: Uinta Mountains, Utah

- Trichuris [sp.]
Hass, D. K.; and Chitwood, M. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 255
dog (feces): west central Ohio
- Trichuris spp., *illus.*
Henriksen, S. A., 1977, Medlemsbl. Danske Dyrlaegeforen., v. 60 (11), 482-485
morphology, life cycle, epidemiology, pathology, diagnosis, control, brief review
- Trichuris sp.
Hiregoudar, L. S., 1976, Indian Vet. J., v. 53 (3), 237
Cervus unicolor (feces)
Gazella gazella "
all from Gir forest, Gujarat State, India
- Trichuris spp.
Horak, I. G.; Honer, M. R.; and Schroeder, J., 1976, J. South African Vet. Ass., v. 47 (4), 247-251
helminths and Oestrus ovis, merino sheep, treated at four-weekly intervals or strategically, live mass gains, wool production and fecal worm egg counts, compared with untreated controls: Eastern Transvaal Highveld
- Trichuris sp.
Lyons, E. T.; et al., 1975, Am. J. Vet. Research, v. 36 (6), 777-780
calves, natural infections of gastrointestinal parasites and lungworms, controlled test of activity of levamisole administered via drinking water, subcutaneous injection, or alfalfa pellet premix
- Trichuris [sp.]
Manz, D.; and Dingeldein, W., 1974, Prakt. Tierarzt, v. 55 (8), 422-425
nematodes of European and exotic herbivores, Banminth, good results
Dama dama mesopotamica
Taurotragus oryx derbianus
- Trichuris sp.
Mirza, M. Y.; and al-Rawas, A. Y., 1975, J. Protozool., v. 22 (1), 23-24
Tatera indica (feces): Baghdad district, Iraq
- Trichuris sp.
Pester, F. R. N.; and Laurence, B. R., 1974, J. Zool., London, v. 174 (3), 397-406
Gazella granti (large intestine): Kenya
- Trichuris sp.
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 203-204
nematodes, seasonal dynamics in deer
- Trichuris sp.
Pursglove, S. R., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 107-108
Odocoileus virginianus (cecum): Oklahoma
- Trichuris sp.
Pursglove, S. R.; et al., 1976, J. Am. Vet. Med. Ass., v. 169 (9), 896-900
intestinal nematodes of Odocoileus virginianus, geographic distribution; deer insignificant in epizootiology of intestinal nematodes of domestic livestock: southeastern United States
- Trichuris sp.
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
helminths of Alces alces, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Trichuris sp.
Schneider, C. R.; et al., 1975, Ann. Trop. Med. and Parasitol., v. 69 (2), 227-232
Bubalus bubalis: Khong Island, Laos
- Trichuris [sp.] larva
Schweisgut, I., 1975, Untersuchungen über den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Trichuris spp.
Searson, J. E.; and Doughty, F. R., 1977, Austral. Vet. J., v. 53 (9), 456-457 [Letter]
nematodes, cattle, fenbendazole, good results (higher efficiency against adult Ostertagia ostertagi than larval forms): southern New South Wales
- Trichuris sp.
Seese, F. M., 1973, Am. Midland Naturalist, v. 89 (2), 257-265
key
Ochotona p. princeps (caecum): St. Joe Baldy Mountain, Benewah County, Idaho
- Trichuris spp.
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (10), 1207-1209
oxibendazole, cattle, drench and premix
- Trichuris spp.
Theodorides, V. J.; et al., 1976, Am. J. Vet. Research, v. 37 (12), 1517-1518
gastrointestinal nematodes, calves, albendazole
- Trichuris sp.
Thornton, J. E.; et al., 1973, J. Wildlife Dis., v. 9 (2), 160-162
Antelope cervicapra (large intestine): Texas
- Trichuris spp.
Tiefenbach, B., 1977, Cahiers Bleus Vet. (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date

- Trichuris* sp.
Torres, P.; Lopetegui, O.; and Gallardo, M., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 39-42
morphometric data
Ctenomys maulinus maulinus
C. m. brunneus
(intestino grueso of all): all from Chile
- Trichuris* sp.
Williams, J. C.; and Knox, J. W., 1976, Am. J. Vet. Research, v. 37 (4), 453-464
failure of stocker cattle to achieve projected weight gains at high stocking rates on Coastal bermudagrass pastures even with supplemental feeding and anthelmintic control of parasitism
- Trichuris* sp.
Williams, J. C.; Sheehan, D.; and Fuselier, R. H., 1977, Am. J. Vet. Research, v. 38 (12), 2037-2038
gastrointestinal nematodes, tapeworms, cattle, efficacy of albendazole (oral drench)
- Trichuris* sp.
Wilson, D. E.; and Hirst, S. M., 1977, Wildlife Monogr. (54), Suppl., 3-111
Hippotragus niger: Percy Fyfe Nature Reserve, South Africa
- Trichuris bradleyi* sp. n., illus.
Babero, B. B.; Cattán, P. E.; and Cabello, C., 1975, J. Parasitol., v. 61 (6), 1061-1063
Octodon degus (cecum, large intestine): Quebrada de la Plata, 30 km SW from Santiago, Chile
- Trichuris bradleyi*
Babero, B. B.; and Cattán, P. E., 1975, Bol. Chileno Parasitol., v. 30 (3-4), 68-76
Octodon degus: Quebrada de la Plata, Santiago, Chile
- Trichuris bradleyi* (Babero y col., 1975)
Cattán, P. E.; George-Nascimento, M.; and Rodríguez, J., 1976, Bol. Chileno Parasitol., v. 31 (1-2), 16-20
prevalence survey of helminths of *Octodon degus*, seasonal variations, age and sex of hosts: Chile
- Trichuris chiliensis* n. sp., illus.
Babero, B. B.; Cattán, P. E.; and Cabello, C., 1976, Tr. Am. Micr. Soc., v. 95 (2), 232-235
Akodon longipilis (cecum and large intestines): Codillera de Nahuelbuta, Malleco, west of Angol, Chile
- Trichuris dipodomis* Read
Bienek, G. K.; and Klikoff, L. G., 1974, Am. Midland Naturalist, v. 91 (1), 251-253
Dipodomys microps
- Trichuris discolor*
Georgi, J. R.; Whitlock, R. H.; and Flinton, J. H., 1972, Cornell Vet., v. 62 (1), 58-60
Trichuris discolor, heifer (feces, colon), fatal infection, case report
- Trichuris discolor*
Guimaraes, M. P.; et al., 1976, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 28 (2), 217-219
sheep, pastured with cattle: Patos de Minas, Minas Gerais, Brasil
- Trichuris globulosa* (Linstow, 1901)
Basson, P. A.; et al., 1970, Onderstepoort J. Vet. Research, v. 37 (1), 11-28
parasitic and other diseases of *Syncerus caffer*, some pathological findings, age of host
Syncerus caffer (large intestine): Kruger National Park
- Trichuris globulosa* (v. Linstow, 1901) Ransom, 1911
Jain, P. C.; and Kamalapur, S. K., 1971, Indian J. Animal Research, v. 5 (1), 43-44
sheep
goats
all from Madhya Pradesh
- Trichuris globulosa*
Lodha, K. R.; Raisinghani, P. M.; and Karwar, R. S., 1977, Indian J. Animal Sc., v. 47 (10), 677-682
helminths, camels, promintic and banminth II effective, nilverm inconsistent in action, thiabendazole ineffective
- Trichuris globulosa* (von Linstow, 1901) Ransom, 1911, illus.
Martinez Gomez, F.; Hernandez Rodriguez, S.; and Calero Carretero, R., 1973, Rev. Iber. Parasitol., v. 33 (2-3), 331-336
description
Ovis aries: Spain
- Trichuris globulosa*
Schweisgut, I., 1975, Untersuchungen uber den Endoparasitenbefall des Rotwildes im Nationalpark Bayerischer Wald in den Jagdjahren 1973/74 und 1974/75, 70 pp.
Rotwild: Nationalpark Bayerischer Wald
- Trichuris globulosa*
Troncy, P. M.; and Oumate, O., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (2), 189-198
Strongylidae of zebu, morantel tartrate, efficacy, toxicity: Tchad
- Trichuris globulosa*
Troncy, P. M.; and Oumate, O., 1976, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 29 (3), 229-232
gastrointestinal parasites, *Camelus dromedarius*, morantel tartrate, drug efficacy; good results against Strongylidae: Tchad
- Trichuris leporis*
Jacobson, H. A.; and Kirkpatrick, R. L., 1974, J. Wildlife Dis., v. 10 (4), 384-391
comparison of selected physiological measurements in untreated parasitized cottontail rabbits and those treated with 1-tetramisole hydrochloride and 2,2-dichlorovinyl, dimethyl phosphate: Montgomery County, Virginia

- Trichuris leporis*
Kutzer, E.; and Frey, H., 1976, Berl. u. Münch. Tierärztl. Wchnschr., v. 89 (24), 480-483
Lepus europaeus: Austria
- Trichuris leporis*
Kutzer, E.; and Frey, H., 1976, Ztschr. Parasitenk., v. 50 (2), 213-214
Lepus europaeus
- Trichuris muris*
Bruce, R. G.; and Wakelin, D., 1977, Parasitology, v. 74 (2), 163-173
Trichinella spiralis, *Trichuris muris*, concurrent infection in mice, interactive expulsive response considered an example of indirect cross-immunity with no element of antigenic similarity, involvement of cell-mediated inflammatory response strongly suggested
- Trichuris muris*
Howes, H. L., jr., 1972, Proc. Soc. Exper. Biol. and Med., v. 139 (2), 394-398
Trichuris muris and other helminths, dogs, mice (both exper.), CP-14,445 hydrochloride and pamoate compared with activity of known anthelmintics; dosage response data indicate that *T. muris*-mouse infection could be test model for antiwhipworm studies
- Trichuris muris*
Jenkins, S. N., 1976, Parasitology, v. 73 (2), xiv [Abstract]
Trichuris muris, immunization with whole male and stichocyte antigen preparations and with 'exo' antigen obtained by incubation of adult worms, analysis of functional antigens by immunodiffusion and physicochemical treatments
- Trichuris muris*
Jenkins, S. N., 1977, Parasitology, v. 75 (2), xiv [Abstract]
Trichuris muris, mice, cell transfer studies highlight complexity of interaction of humoral and cellular immune response
- Trichuris muris*
Jenkins, S. N.; and Behnke, J. M., 1977, Parasitology, v. 75 (1), 71-78
Trichuris muris, mice, primary immune expulsion markedly delayed by concurrent infection with *Nematospiroides dubius*, possible relevance in pathogenesis of concurrent tropical diseases
- Trichuris muris*
Jenkins, S. N.; and Behnke, J. M., 1977, Parasitology, v. 75 (2), xxxiv [Abstract]
Trichuris muris, delay of primary expulsion in mice concurrently infected with *Nematospiroides dubius*
- Trichuris muris*
Jenkins, S. N.; and Wakelin, D., 1977, Parasitology, v. 74 (2), 153-161
Trichuris muris, mice, vaccination with whole male worm extract, stichosome extract, and short-term incubation fluid in attempt to localize protective antigens and investigate them physico-chemically, concluded that one of protective immunogens is protein which can be associated with precipitin line and originates in stichosome
- Trichuris muris*
Jenkins, T.; Erasmus, D. A.; and Davies, T. W., 1977, Exper. Parasitol., v. 41 (2), 464-471
Trichuris suis, *T. muris*, intestinal inclusions, analysis of elemental composition using X-ray analysis in transmission electron microscope and cryo-ultramicrotomy
- Trichuris muris*
Kagei, N.; and Kihata, M., 1976, Bull. Inst. Pub. Health, Tokyo, v. 25 (2), 73-81
Trichuris muris, albino mice (exper.), cortisone-acetate, sharp reduction in worm numbers between 15 and 25 days after infection in untreated controls but not in treated mice
- Trichuris muris*
Mishra, G. S.; and Gonzalez, J. P., 1975, Arch. Inst. Pasteur Tunis, v. 52 (1-2), 71-87
Rattus norvegicus (caecum): Tunisia
- Trichuris muris*
Owen, D., 1976, Lab. Animals, v. 10 (3), 271-278
Rattus norvegicus: Carshalton
- Trichuris muris*
Phillips, R. S.; and Wakelin, D., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (4), 276 [Demonstration]
Trichuris muris-infected mice, suppression of immunity by concurrent infection with *Babesia* spp., severely depressed agglutinating antibody response to sheep red blood cells
- Trichuris muris*
Phillips, R. S.; and Wakelin, D., 1976, Exper. Parasitol., v. 39 (1), 95-100
mice concurrently infected with *Babesia* and *Trichuris muris*, marked immunodepression, normal immune expulsion of nematode delayed; *Babesia* infections had little effect on expulsion of challenge infections of *T. muris* from previously immunized mice; *Babesia* infections exerted profound immunodepressive effect on agglutinating antibody response to sheep red blood cells
- Trichuris muris*
Wakelin, D., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (4), 277 [Demonstration]
Trichuris muris in mice, possible genetic control of immune responses
- Trichuris muris*
Wakelin, D.; and Selby, G. R., 1974, Immunology, v. 26 (1), 1-10
Trichuris muris, mice, induction of immunological tolerance by treatment with cortisone

- Trichuris muris*
Wakelin, D.; and Selby, G. R., 1976, Parasitology, v. 72 (1), 41-50
Trichuris muris, immune expulsion from resistant mice, suppression by irradiation, attempts to restore by transfer of mesenteric lymph node cells, bone marrow, or immune serum, results confirm involvement of both antibody-mediated and lymphoid cell-mediated phases in immune expulsion
- Trichuris ovis*
Boag, B.; and Thomas, R. J., 1973, Research Vet. Sc., v. 14 (1), 11-20
gastrointestinal nematode parasites of sheep, effectiveness of 3 control measures applied at strategic points in lamb infection pattern (anthelmintic treatment of ewes at lambing, of lambs at weaning, and moving lambs to clean pasture at weaning--tested singly and in combination)
- Trichuris ovis*
Boag, B.; and Thomas, R. J., 1977, Research Vet. Sc., v. 22 (1), 62-67
gastro-intestinal nematodes, sheep, epidemiology, post mortem worm counts, faecal egg counts and pasture larval counts, seasonal number of generations and succession of species
- Trichuris ovis*
Chalmers, K., 1977, N. Zealand Vet. J., v. 25 (10), 266-269
gastrointestinal nematodes and cestodes, sheep, oxfendazole, drug efficacy, good results: New Zealand
- Trichuris ovis*
Chroust, K.; and Dyk, V., 1975, Deutsche Tierarztl. Wchnschr., v. 82 (12), 487-491
gastrointestinal nematodes of lambs, efficacy of fenbendazole, thiabendazole and tetramisole compared
- Trichuris ovis*
Corticelli, B.; and Lai, M., 1972, Parassitologia, v. 14 (1), 95-96
Ovis musimon: Sardegna
- Trichuris ovis*
Düewel, D.; et al., 1974, Prakt. Tierarzt, v. 55 (8), 425-427
sheep stomach and intestinal nematodes, controlled tests of Fenbendazol, good results
- Trichuris ovis*
Eslami, A. H.; and Anwar, M., 1976, Vet. Rec., v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichuris ovis* (Abildgaard, 1795), illus.
Goffredo, G.; and Sobrero, R., 1972, Parassitologia, v. 14 (1), 143-148
Dama dama (intestine): foresta Umbra (promontorio garganico, provincia di Foggia)
- Trichuris ovis*
Gupta, O. P.; et al., 1976, Indian J. Exper. Biol., v. 14 (3), 356-357
in vitro anthelmintic activity of embelin disalts, Paramphistomum cervi, Oesophagostomum columbianum, *Trichuris ovis*, Dipylidium caninum, good results
- Trichuris ovis*
Heuer, D. E.; et al., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 141-143
Odocoileus virginianus (cecum): Kentucky
- Trichuris ovis*
Kelly, J. D.; et al., 1975, Research Vet. Sc., v. 19 (1), 105-107
anthelmintic efficacy of fenbendazole against naturally acquired Dictyocaulus filaria infection associated with concurrent infection of gastro-intestinal nematodes in sheep
- Trichuris ovis*
Kistner, T. P.; and Wyse, D., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 93-97
nematodes of sheep, injectable levamisole, effective control of abomasal and small intestinal parasites with no evidence of skin damage or gross lesions at injection sites
- Trichuris ovis*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, Am. J. Vet. Research, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Trichuris ovis*
Le Jambre, L. F.; and Royal, W. M., 1976, Austral. Vet. J., v. 52 (4), 181-183
nematode worm burdens compared in naturally infected Angora goats and Merino sheep grazing intraspecifically or in mixed experimental paddocks (fecal egg counts showed no significant within-host differences); increased resistance of sheep to all worms except Nematodirus resulted in a significantly lower worm burden for sheep: Northern Tablelands of New South Wales
- Trichuris ovis*
Misra, S. C., 1972, Indian J. Animal Research, v. 6 (2), 95-96
parasitic gastro-enteritis, goats, epidemiology, seasonal incidence: Orissa
- Trichuris ovis*
Musila, V., 1976, Veterinarstvi, v. 26 (6), 264
helminths of fallow deer, incidence: Zehusice enclosure
- Trichuris ovis*
Novy, H., 1976, Veterinarstvi, v. 26 (6), 263
helminths of white deer, incidence: Zehusice enclosure

- Trichuris ovis*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Bos taurus
Ovis aries
all from Chile
- Trichuris ovis*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Trichuris ovis*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Trichuris ovis*
Randall, R. W.; and Gibbs, H. C., 1977, *Am. J. Vet. Research*, v. 38 (10), 1665-1668
gastrointestinal nematodes, dairy cattle, occurrence, degree of parasitism, and seasonal fluctuations: Maine
- Trichuris ovis*
Smeal, M. G.; et al., 1977, *Austral. Vet. J.*, v. 53 (12), 566-573
nematodes, cattle, occurrence, seasonal distribution, poor relationship between faecal egg counts and worm burdens: North Coast and Tableland regions of New South Wales
- Trichuris ovis* (Abilgaard, 1795)
Smith, F. R.; and Threlfall, W., 1973, *Am. Midland Naturalist*, v. 90 (1), 215-218
Ovis aries: insular Newfoundland
- Trichuris ovis*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Trichuris ovis*
Townsend, R. B.; et al., 1977, *Research Vet. Sc.*, v. 23 (3), 385-386
Moniezia expansa, *Trichuris ovis*, sheep, fenbendazole highly efficient
- Trichuris ovis*
Vujic, B.; Pop-Cenic, S.; and Blagojevic, R., 1976, *Vet. Glasnik*, v. 30 (1), 11-17
sheep, morantel tartarate + diethylcarbamazine effective against *Dictyocaulus filaria* and most gastrointestinal helminths except *Strongyloides papillosus*, *Trichuris ovis*, and *Moniezia* sp.
- Trichuris skrjabini*
Eslami, A. H.; and Anwar, M., 1976, *Vet. Rec.*, v. 99 (11), 214-215
gastrointestinal nematodes, sheep, fenbendazole, satisfactory results: Iran
- Trichuris skrjabini*
Knight, R. A.; Vegors, H. H.; and Glimp, H. A., 1973, *Am. J. Vet. Research*, v. 34 (3), 323-327
gastrointestinal nematodes, lambs, effect of breed and birth date on parasite acquisition: Clay Center, Nebraska
- Trichuris skrjabini*
Prestwood, A. K.; Pursglove, S. R.; and Hayes, F. A., 1976, *J. Wildlife Dis.*, v. 12 (3), 380-385
survey of parasites of *Odocoileus virginianus* and *Ovis aries* on common range, deer unlikely reservoir host for sheep parasites
Ovis aries: Hardy County, West Virginia
- Trichuris spiricollis* Solomon
Pester, F. R. N.; and Laurence, B. R., 1974, *J. Zool.*, London, v. 174 (3), 397-406
Gazella thomsonii (large intestine, caecum)
Alcelaphus buselaphus cokei (digestive tract)
all from Kenya
- Trichuris suis, illus.*
Ashizawa, H.; et al., 1975, *Bull. Fac. Agric. Miyazaki Univ.*, v. 22 (2), 211-220
Trichuris suis, pathology of swine cecum and colon: slaughterhouses in Kagoshima and Miyazaki Prefectures
- Trichuris suis*
Baines, D. M.; Dalton, S. E.; and Eichler, D. A., 1976, *Vet. Rec.*, v. 99 (7), 119-122
swine nematodes, field and exper. studies, thiophanate alone or with piperazine, compared with thiabendazole alone or with pica-dex
- Trichuris suis, illus.*
Batte, E. G.; et al., 1977, *Am. J. Vet. Research*, v. 38 (7), 1075-1079
Trichuris suis, swine, clinical signs, scanning electron microscopy of intestinal mucosa, biochemical changes
- Trichuris suis*
Batte, E. G.; McLamb, R. D.; and Vestal, T. J., 1976, *Pathophysiol. Parasit. Infect.*, 69-73
Trichuris suis, pigs, clinical signs, hypoalbuminaemia, increase in serum alpha, beta, and gamma globulin, depressed serum calcium levels, no correlated changes in zinc levels, serum creatinine, SGPT, blood sugars, hemoglobin, bilirubin, or BUN
- Trichuris suis, illus.*
Beer, R. J.; and Rutter, J. M., 1972, *Research Vet. Sc.*, v. 13 (6), 593-595
Trichuris suis, weaned pigs (exper.), syndrome resembling swine dysentery, demonstration of spirochaetal invasion of colonic mucosa, possible significance of association of nematode and bacteria

- Trichuris suis*
Burden, D. J.; and Hammet, N. C., 1976, *Vet. Parasitol.*, v. 2 (3), 307-311
Trichuris suis, comparison of infectivity of ova embryonated by 4 different methods, found that differences in method of culture profoundly affected ability of fully developed eggs to hatch and parasites to become established in pigs, ova of highest infectivity produced after culture in moist vermiculite
- Trichuris suis*
Corwin, R. M., 1977, *Am. J. Vet. Research*, v. 38 (4), 465-467
mixed nematode infections, pigs, oxfendazole, critical evaluation: Missouri
- Trichuris suis*
Enigk, K.; et al., 1975, *Zentralbl. Vet.-Med., Reihe B*, v. 22 (8), 687-702
survival of resistant external stages of parasites during fermentation of liquid cattle manure at high temperatures
- Trichuris suis*
Enigk, K.; Dey-Hazra, A.; and Batke, J., 1976, *Tierarztl. Umschau*, v. 31 (8), 360-362
swine nematodes, mebendazole treatment
- Trichuris suis*
Hall, G. A.; Rutter, J. M.; and Beer, R. J. S., 1976, *J. Comp. Path.*, v. 86 (2), 285-292
Trichuris suis, sequential development of large intestinal lesions in piglets (conventionally reared vs. specific-pathogen-free vs. gnotobiotic) studied histologically, synergistic effect of T. suis and bacterial flora in disease process
- Trichuris suis*
Henriksen, S. A., 1974, *Medlemsbl. Danske Dyrlaegeforen.*, v. 57 (24), 981-987
morphology, epidemiology, pathology, diagnosis, control, treatment, incidence, review: Denmark
- Trichuris suis*
Hubert, J., 1977, *Rec. Med. Vet.*, v. 153 (12), 923-929
Ascaris suum (exper.), Trichuris suis (exper.), Oesophagostomum spp., pigs, comparison of coproscopical method of count after dilution in dense solution and flotation in Mac Master Slides and count after sedimentation and flotation in dense solution in Mac Master Slides; various densities of solutions compared with both techniques
- Trichuris suis*
Jacobs, D. E.; Lean, I. J.; and Oakley, G. A., 1977, *Vet. Rec.*, v. 100 (3), 49
Trichuris suis adults, pigs, levamisole high activity by subcutaneous injection, poor results in feed
- Trichuris suis, illus.*
Jenkins, T.; Erasmus, D. A.; and Davies, T. W., 1977, *Exper. Parasitol.*, v. 41 (2), 464-471
Trichuris suis, T. muris, intestinal inclusions, analysis of elemental composition using X-ray analysis in transmission electron microscope and cryo-ultramicrotomy
- Trichuris suis*
Oberg, C.; Diaz, L.; and Valenzuela, G., 1974, *Bol. Chileno Parasitol.*, v. 29 (3-4), 99-102
Sus scrofa: Chile
- Trichuris suis*
Panitz, E., 1977, *J. Helminth.*, v. 51 (1), 23-30
ethyl-6-ethoxybenzothiazole-2-carbamate, evaluation of anthelmintic activity in ponies, swine, lambs, and chickens
- Trichuris suis*
Tiefenbach, B., 1977, *Cahiers Bleus Vet.* (26), 216-230
fenbendazole (available in 5 forms), efficacy against nematodes in various animals, well tolerated with no apparent effects on fertility or fetus, extensive summary of results to date
- Trichuris suis*
Valenzuela, G.; et al., 1977, *Bol. Chileno Parasitol.*, v. 32 (1-2), 23-26
meat inspection survey at local abattoir for evidence and frequency of intestinal parasites
cerdos (intestino grueso): Planta Faenadora de Carnes Socoagro, Valdivia, Chile
- Trichuris sylvilagi*
Haupt, W.; and Hartung, J., 1977, *Monatsh. Vet.-Med.*, v. 32 (9), 339-341
Lepus europaeus: Leipzig area
- Trichuris trichiura*
Alcasid, M. L. S.; et al., 1973, *N. York State J. Med.*, v. 73 (13), 1786-1788
human intestinal parasites, no significant correlation with bronchial asthma, results of study with control and asthmatic groups: New York
- Trichuris trichiura*
Alvarez Ch., R.; et al., 1976, *Rev. Invest. Salud Pub., Mexico*, v. 36 (2), 113-116
Trichuris trichiura, human double-blind clinical trials with diphetarsone
- Trichuris trichiura*
Alvarez Chacon, R.; Rodriguez Rodriguez, M.; and Cob Sosa, C. E., 1974, *Bol. Med. Hosp. Inf.*, v. 31 (6), 1125-1147
massive Trichuris trichiura infections, children, statistical survey, comparative clinical trial with oral and rectal thiabendazole and rectal hexylresorcinol
- Trichuris trichiura*
Anderson, R. I.; and Buck, A. A., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 447-456
complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad
- Trichuris trichiura*
Arnau, J. P.; and Danis, M., 1976, *Medecine Infant.*, v. 83 (1), 9-22
helminthiasis of digestive tract in children, differential diagnosis, current treatment methods

- Trichuris trichiura*
Bailenger, J.; et al., 1977, *Pharmacien Biol.* (109), v. 11, 267-277
human intestinal parasites, fecal examination using floatation or diphasic concentration, principles involved in both diagnostic procedures, changes in results when mercury or sodium merthiolate is added during procedure, special application of Janeckso and Urbanyi reaction
- Trichuris trichiura*
Best, J. C.; et al., 1976, *Med. J. Australia*, v. 1 (1-2), 14-20
assessment of growth-rate and growth-retardation in Australian Aboriginal children before and after treatment for common intestinal parasites
- Trichuris trichiura*
Bevanger, L., 1974, *Tidsskr. Norske Laegefor.*, v. 94 (10), 651-652
single and mixed intestinal parasitic infections in adoptive children from Asiatic areas, need for control measures: Norway
- Trichuris trichiura*
Biagi, F.; Lopez, R.; and Viso, J., 1975, *Progr. Drug Research*, v. 19, 10-22
human intestinal parasites, analysis of signs and symptoms related to infections, extensive review
- Trichuris trichiura*
Biagi, F.; Smyth, J.; and Gonzalez, C., 1975, *Prensa Med. Mexicana*, v. 40 (5-6), 189-192
human intestinal parasites, clinical trials with mebendazole show it to be useful drug against many parasites and therefore recommended for mass therapy in low socioeconomic areas where multiple parasitism is likely to be present: Mexico
- Trichuris trichiura*
Borlone Rojas, S., 1975, *Pediatrics*, Santiago, v. 18 (1), 28-30
severe massive infection in 6-year-old child, clinical case report and review of medical aspects of infections in Chile
- Trichuris trichiura*
Botero R., D., 1975, *Progr. Drug Research*, v. 19, 28-43
human intestinal soil transmitted helminths, epidemiology and public health importance, pathology, extensive review
- Trichuris trichiura*
Bull, F.; Oyarce, R.; and Stehr, I., 1967, *Bol. Chileno Parasitol.*, v. 22 (1), 10-15
prevalence and epidemiologic survey of human intestinal parasites in slum areas of Concepcion Province, Chile
- Trichuris trichiura*
Burden, D. J.; et al., 1976, *J. Hyg., Cambridge*, v. 77 (3), 377-382
Trichuris trichiura in humans, soil fumigants applied to contaminated soil of no value in killing parasite ova, eggs on untreated soil had only 20% viability after 18 months; difetarone equally unsuccessful in treatment of children thought to be source of contaminated soil: Great Britain
- Trichuris trichiura*
Camargo, P. A. M.; et al., 1974, *Pediat. Prat., S. Paulo*, v. 45 (10-12), 189-194
human intestinal helminths, comparative treatment trials using mebendazole and pyrantel pamoate, efficacy of drugs checked using the McMaster technique to determine reduction in egg production
- Trichuris trichiura*
Cervoni, W. A.; and Oliver-Gonzalez, J., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 589-591
pyrantel pamoate effective against human infection of *Ascaris lumbricoides* and *Necator americanus* but not *Trichuris trichiura* in clinical evaluations: Puerto Rico
- Trichuris trichiura*
Chanco, P. P., jr.; Goquiolay-Arellano, R.; and Atienza, M. R., 1974, *Philippine J. Pediat.*, v. 23 (2), 43-53
Trichuris trichiura, children with complication of rectal prolapse, successful therapy with mebendazole, clinical trials
- Trichuris trichiura*
Chodera, L.; et al., 1973, *Polski Tygod. Lekar.*, v. 28 (9), 338-340
Necator americanus, *Trichuris trichiura*, evaluation of bitoscanate as therapy; poor results in trichuriasis and frequent side effects in hookworm
- Trichuris trichiura*
Forman, D. W.; et al., 1971, *Am. J. Trop. Med. and Hyg.*, v. 20 (4), 598-601
parasitic pathogens discovered during etiologic survey of causes of diarrheal disease in U.S. Marines in Vietnam
- Trichuris trichiura*
Garcia, E. G., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (6), 914-915
Trichuris trichiura, human, oxantel, effective and safe single dose chemotherapeutic agent
- Trichuris trichiura*
Garrocho-Sandoval, C.; and Torres-Ruvalcaba, A., 1977, *Am. J. Clin. Path.*, v. 67 (6), 603-605
human intestinal parasites, comparison study of two different methods for collecting fecal samples for diagnostic purposes
- Trichuris trichiura*
Gemessi, G., 1971, *Parasitol. Hungar.*, v. 4, 47-63
human intestinal parasites, comparison of standard methods of egg count in fecal specimens with that of the new and improved Telemann method
- Trichuris trichiura*
de Groot, J. F. M.; et al., 1976, *Nederl. Tijdschr. Geneesk.*, v. 120 (51), 2263-2267
Trichuris trichiura, *Ascaris lumbricoides* and *Entamoeba histolytica*, fecal survey of nursing staff and inmates of hospital for mentally retarded for presence of intestinal parasites: Groot Schuylenburg, Apeldoorn, Netherlands

- Trichuris trichiura*
Hays, B. D., 1977, *J. Environ. Health*, v. 39 (6), 424-426
transmission of protozoan cysts and metazoan eggs from land application of sewage effluents and sludges, brief literature review, parasite survey from selected Pittsburgh area sludges, control measures
- Trichuris trichiura*
Herrero Alonso, C., 1969, *Med. Trop.*, Madrid, v. 45 (1-2), 11-20
thiabendazole drug of choice in treating human infections of *Ancylostoma duodenale*, *Trichuris trichiura* and *Ascaris lumbricoides*, clinical trials
- T[*richuris*] *trichiura*
Islam, N.; and Chowdhury, N. A., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (1), 81-84
human intestinal parasites, comparative treatment trials using mebendazole and pyrantel pamoate, recommendations for use: Dacca, Bangladesh
- Trichuris trichiura*
Iwanczuk, I.; 1969, *Acta Parasitol. Polon.*, v. 17 (1-19), 139-145
human parasite incidence in water and surfaces of swimming pools; change of incidence in children using swimming pool for 6 week period: Poland
- Trichuris trichiura*
Iwanowski, H.; and Herman, T., 1977, *Pediat. Polska*, v. 52 (6), 649-654
Ascaris lumbricoides in children, Soprunova urine test used successfully to diagnose ascariasis and differentiate from infections of *Trichuris trichiura*, *Enterobius vermicularis* and *Lamblia intestinalis*
- Trichuris trichiura*
Lyngkaran, N.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (4), 568-572
clinical trials using mebendazole to treat severe *Trichuris trichiura* infections in children; proctoscopy established as best method to assess worm burden and severity of involvement; mebendazole safe and effective if given over longer than currently recommended period, eggs post-treatment morphologically altered and no longer viable: Malaysia
- Trichuris trichiura*
Jones, H. I., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (4), 523-529
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, statistical epidemiologic survey of human intestinal parasites with discussion of socio-cultural and environmental factors influencing spread of infections: Venilale District of East Timor
- Trichuris trichiura*
Jones, H. I., 1976, *Papua N. Guinea Med. J.*, v. 19 (3), 165-172
Ascaris lumbricoides, *Necator americanus*, *Trichuris trichiura*, epidemiologic survey of human helminths in natives of Kar Kar Island, *Ascaris* infections highest and heaviest in children and principal cause of high eosinophilia in the communities: Madang Province, Papua New Guinea
- Trichuris trichiura*
Jose, D. G.; and Welch, J. S., 1970, *Med. J. Australia*, v. 1 (8), 349-356
possible role of intestinal parasitism in growth-retarded, anemic and malnourished Australian Aboriginal children, comparison with normal Aboriginal children: Queensland
- Trichuris trichiura*
Kagei, N., 1975, *Bull. Inst. Pub. Health, Tokyo*, v. 24 (3), 169-175
comparison of Kato thick smear and Tween 80 citric acid ether sedimentation methods for diagnosis of helminth ova
- Trichuris trichiura*
Kelly, J. D., 1974, *Internat. J. Zoonoses*, v. 1 (1), 13-24
anthropozoonotic helminthiases associated with domesticated and domiciliated vertebrates, developmental phases in man: Australia; New Zealand
- Trichuris trichiura*
Khan, S. M.; and Khairul Anuar, A., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 260-264
Ascaris lumbricoides, *Trichuris trichiura*, hookworms, prevalence survey of soil-transmitted intestinal helminths in patients admitted to the District Hospital in Balik Pulau, Penang, Malaysia
- Trichuris trichiura*
Krubwa, F.; et al., 1974, *Medecine Trop.*, v. 34 (5), 679-687
trial administration of mebendazole to school children on quarterly basis for attempted eradication of intestinal nematodes, recommendations for use as mass control measure: Zaire
- Trichuris trichiura*
Langer, A.; and Hung, C. T., 1973, *Obst. and Gynec.*, v. 42 (4), 564-567
human hookworm anemia associated with pregnancy, clinical management with tetrachlorethylene or with tetrachlorethylene and thiabendazole in *Trichuris trichiura*-associated infections
- Trichuris trichiura*
Lara Aguilera, R.; and Bernal, R. M., 1974, *Bol. Med. Hosp. Inf.*, v. 31 (4), 779-784
Balantidium coli, dysentery in young children, frequent association with *Trichuris trichiura* infections in endemic areas, case reports, metronidazole: Mexico

- Trichuris trichiura
Layrisse, M.; and Vargas, A., 1975, Progr. Food and Nutrition Sc., v. 1 (10), 645-667
human intestinal parasites, mechanisms by which parasites interfere with host nutrition (competition for nutrients, malabsorption, blood loss, excess nutrient utilization), extensive review
- Trichuris trichiura, illus.
Lee, E.-L.; et al., 1976, Am. J. Trop. Med. and Hyg., v. 25 (4), 563-567
evaluation of oxantel pamoate for treatment of severe clinical Trichuris trichiura infection in children, description of clinical trials and correlated treatment of frequently occurring mixed parasitic infections, oxantel suggested as safe and effective anthelmintic: Malaysia
- Trichuris trichiura
Lengyel, A.; Levai, J.; and Rovo, J. T., 1974, Med. Universalis, v. 7 (5), 263-265
Trichuris trichiura, mebendazole effective in short and long term treatment, clinical trials
- Trichuris trichiura
Levai, J.; Lengyel, A.; and Banki, G., 1972, Med. Universalis, v. 5 (1), 31-34
clinical trials using levamisole to treat human intestinal helminths, good results against Ascaris, Ancylostoma, Trichostrongylus, ineffective against Trichuris trichiura: Hungary
- Trichuris trichiura
Lie, K. J.; Kwo, E. H.; and Ow-Yang, C. K., 1971, Southeast Asian J. Trop. Med. and Pub. Health, v. 2 (2), 196-200
Ascaris lumbricoides, Trichuris trichiura, Necator americanus, fecal survey for evidence of soil-transmitted helminths in infants and children living near Kuala Lumpur, Malaysia
- Trichuris trichiura
McKay, D. A.; Hiap, C. Y.; and Virik, H. K., 1971, Med. J. Malaya, v. 26 (1), 20-24
treatment trials in children with diarrhea from severe Trichuris trichiura infection or mixed infections with Entamoeba histolytica, comparison of thiabendazole alone or in combination with hexylresorcinol as single retention enema: Kuala Lumpur
- Trichuris trichiura
O'Holohan, D. R.; and Hugce-Matthews, J., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 576-579
Trichuris trichiura, humans, successful clinical trials testing diphetarsones; 8 persons excreting E[ntamoeba] histolytica cysts at beginning of trials were cyst free after treatment
- Trichuris trichiura
Orren, A.; and Dowdle, E. B., 1975, Internat. Arch. Allergy and Applied Immunol., v. 49 (6), 814-830
serum IgE concentrations and immediate skin hypersensitivity to common allergens analyzed with respect to ethnic group (whites, Cape Coloreds, Africans), sex, allergic status, and evidence of intestinal helminthic infestation: Western Cape Province, South Africa
- Trichuris trichiura
Otsuru, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Trichuris trichiura
Pawlowski, Z.; and Chodera, L., 1972, Polski Tygod. Lekar., v. 27 (41), 1605-1607
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, humans, levo-tetramisole clinical trials successful except in trichuriasis
- T[richuris] trichiura
Penna, R.; and Grassi, L., 1972, Parassitologia, v. 14 (2-3), 339-341
T[richuris] trichiura, A[scaris] lumbricoides, H[ymenolepis] nana, survey of prevalence of geohelminthiasis in school children, helminthiasis related to socio-economic and hygienic conditions, not to scholastic achievement, prevalence higher in plain country than in mountain country: Province of Alesandria (Italy)
- Trichuris trichiura
Prosl, H., 1976, Ztschr. Parasitenk., v. 50 (2), 214
Rhesusaffe
- Trichuris trichiura
Purtilo, D. T.; et al., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 229-232
parasitized children with protein-calorie malnutrition, complete blood count, serum immunoglobulin concentration, significant relationship between intensity of parasitism and Ig levels: Brazil
- Trichuris trichiura
Quevedo, W.; et al., 1974, Pediatria, Bogota, v. 14 (4), 235-243
human Trichuris trichiura, difetarsones drug of choice in treatment: Colombia
- Trichuris trichiura
Rees, P. H.; and Marsden, P. D., 1970, Brit. J. Clin. Pract., v. 24 (1), 3-11
important intestinal parasites diagnosed in Britain, emphasis on clinical aspects, laboratory diagnosis and current treatment
- Trichuris trichiura
Reyes, H.; Olea, M.; and Hernandez, R., 1972, Bol. Chileno Parasitol., v. 27 (3-4), 115-116
survey of food handlers for intestinal parasites shows 50.5% infection rate: Santiago, Chile

- Trichuris trichiura**
Ripert, C.; and Avouac-Borzee, F., 1974, *Medecine Afrique Noire*, v. 21 (10), 713-715
Ascaris lumbricoides, Necator americanus, Trichuris trichiura, human, epidemiology: Mirebalais, Haiti
- Trichuris trichiura**
Robinson, H. M.; and Samorodin, C. S., 1976, *Arch. Dermat.*, Chicago, v. 112 (12), 1757-1760
Trichuris trichiura, woman, thiabendazole-induced toxic epidermal necrolysis syndrome: Baltimore, Maryland
- Trichuris trichiura**
Sagua, H.; et al., 1973, *Bol. Chileno Parasitol.*, v. 28 (3-4), 58-60
comparison of phenol-alcohol-formalin sedimentation and polyvinyl alcohol fixative tests in diagnosis of human intestinal helminths and protozoa
- Trichuris trichiura**
Samuel, M. R., 1975, *Progr. Drug Research*, v. 19, 96-107
human intestinal helminths, review of clinical experiences world wide comparing the efficacy and tolerance of bitoscanate with that of bephenium hydroxynaphthoate and tetrachlorethylene; found to be most useful against hookworm with results against other helminths still inconclusive
- Trichuris trichiura**
Sandars, D. F.; and Bianchi, G. N., 1970, *Med. J. Australia*, v. 1 (6), 261-266
Entamoeba histolytica epidemic in institutionalized mental defectives, high associated incidence of Trichuris trichiura suggests it as factor contributing to Entamoeba's pathogenicity; comparative study of effectiveness of intestopan and humatin in treatment of amoebiasis: Queensland, Australia
- Trichuris trichiura**
Sapunar, J.; and Bertossi, E., 1966, *Bol. Chileno Parasitol.*, v. 21 (3), 94-95
mixed Trichostrongylus, Trichuris trichiura, and Giardia lamblia infection in young man diagnosed by fecal examination, case report: Chile
- Trichuris trichiura**
Schenone, H.; et al., 1976, *Bol. Chileno Parasitol.*, v. 31 (3-4), 75-78
Enterobius vermicularis, Ascaris lumbricoides, and Trichuris trichiura in children, clinical trials of oral mebendazole highly successful, recommended for individual or mass therapy
- Trichuris trichiura**
Schenone, H.; Galdames, M.; and Cabello, C., 1975, *Bol. Chileno Parasitol.*, v. 30 (3-4), 89-90
intestinal parasites, young girls, combined therapy with mebendazole and thiabendazole
- Trichuris trichiura**
Scragg, J. N.; and Proctor, E. M., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (2), 198-203
Trichuris trichiura, mebendazole treatment of children with severe symptomatic infections, satisfactory results with cure in 74% and reduced egg counts in remaining children, possible use for mass therapy: Durban, South Africa
- Trichuris trichiura**
Seah, S. K. K., 1973, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 4 (4), 534-542
intestinal parasites, persons living in non-endemic areas who acquired infections while travelling or who have immigrated from endemic areas, pyrantel pamoate successful for Ascaris lumbricoides, results with other parasites varied: Montreal, Canada
- Trichuris trichiura**
Seah, S. K. K., 1976, *Canad. Med. Ass. J.*, v. 115 (8), 777-779
mebendazole suggested as drug of choice for Trichuris trichiura and mixed nematode infections after extensive clinical trials: Canada
- Trichuris trichiura**
Sehgal, S. C.; Vinayak, V. K.; and Gupta, U., 1977, *Indian J. Med. Research*, v. 65 (4), 509-512
human helminthic ova in feces, diagnosis using the Kato thick smear technique more successful than commonly used techniques, recommended for epidemiologic surveys: Chandigarh, India
- Trichuris trichiura**
Signorello, G., 1973, *Minerva Med.*, v. 64 (52), 2736-2740
Ascaris lumbricoides in children resulting in cutaneous allergic reactions, skin test diagnosis, frequent mixed infections with Trichuris trichiura also diagnosed
- Trichuris trichiura**
Singson, C. N.; Banzon, T. C.; and Cross, J. H., 1975, *Am. J. Trop. Med. and Hyg.*, v. 24 (6, pt. 1), 932-934
Capillaria philippinensis, human clinical trials using mebendazole for intestinal capillariasis, additionally effective against Ascaris lumbricoides, Trichuris trichiura and hookworm: Philippines
- Trichuris trichiura**
Stuart, J. E.; and Welch, J. S., 1973, *Med. J. Australia*, v. 2 (22), 1017-1019
treatment trials with Aboriginal children infected with Trichuris trichiura using thiabendazole in combination with viprynum, no significant difference between treated and nontreated groups: Queensland, Australia

- Trichuris trichiura*
Tantengco, V. O.; Marzan, A. M.; and de Castro, C. R., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 580-586
Ascaris lumbricoides, *Trichuris trichiura*, establishment of blood parameters in infected and control children, little significant differences except lowered albumin and elevated globulin levels in presence of infections: Philippine Islands
- Trichuris trichiura*
Tarabini Castellani, G., 1970, Minerva Gastroenterol., v. 16 (1), 45-49
human intestinal parasites, clinical trials using FI 6518 with and without thymol, acceptable drug for mass therapy: Somalia
- Trichuris trichiura*, *illus.*
Vanhaelen-Lindhout, E.; and Smit, A. M., 1971, Trop. and Geogr. Med., v. 23 (4), 381-384
abnormal *Trichuris trichiura* eggs in feces of children during and immediately after drug therapy with thiabendazole: Netherlands (previous residents of Surinam)
- Trichuris trichiura*
Vinayak, V. K.; and Sehgal, S. C., 1976, Indian J. Med. Research, v. 64 (9), 1347-1350
human helminthic and protozoan parasites, comparison of nigrosin-methylene blue diagnostic test with formol-ether method and direct examination
- Trichuris trichiura*
Warren, K. S.; and Mahmoud, A. A. F., 1976, J. Infect. Dis., v. 133 (2), 240-243
Trichuris trichiura, epidemiologic, clinical and diagnostic review of human trichuriasis
- Trichuris trichiura*
Zaman, V.; and Sabapathy, N. N., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (1), 103-105
Trichuris trichiura in humans, successful clinical trials testing CP-14,445, drug highly efficacious and without toxic side effects: West Malaysia
- Trichuris trichiura*, *illus.*
Zuidema, P. J., 1976, Nederl. Tijdschr. Geneesk., v. 120 (43), 1849-1854
human intestinal helminths, differential diagnosis and clinical management of parasitic infestations seen in immigrants from Surinam: Netherlands
- Trichuris vulpis*
Davies, P.; and Nicholas, W. L., 1977, Austral. Vet. J., v. 53 (5), 247-248 [Letter]
dogs (feces): Goodradigbee Shire, New South Wales
- Trichuris vulpis*
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Trichuris vulpis*
Hass, D. K.; and Collins, J. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 135-137
helminths, dogs, comparative efficacy of vincofos, ticarbondine, mebendazole
- Trichuris vulpis*, *illus.*
Kagei, N.; et al., 1976, Bull. Inst. Pub. Health, Tokyo, v. 25 (3), 140-144
Filaroides hirthei, *Strongyloides* sp., *Toxocara canis*, and *Trichuris vulpis* in imported dogs, inability to experimentally infect other dogs with *Strongyloides stercoralis* of human origin: Japan, imported from U.S.A.
- Trichuris vulpis*
Klein, J. B.; and Bradley, R. E., sr., 1976, Vet. Med. and Small Animal Clin., v. 71 (5), 598-599
dogs, sansalid, critical testing, fair results
- Trichuris vulpis*
McCurdy, H. D.; and Guerrero, J., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1731-1733
helminths, dogs, mebendazole powder, controlled critical studies, good results against all helminths except *Dipylidium caninum*: Kansas; New Jersey; Texas
- Trichuris vulpis*
Olsen, J. L.; et al., 1977, J. Am. Vet. Med. Ass., v. 171 (6), 542-544
Trichuris vulpis, dogs (nat. and exper.), dichlorvos administered orally, single and repeated doses; additional doses may be required to remove mature worms
- Trichuris vulpis*
Pegg, E. J., 1977, Brit. Vet. J., v. 133 (4), 427-431
Toxocara canis and other parasite ova, horticultural flame-gun for control on concrete-floored kennel runs
- Trichuris vulpis*
Rep, B. H.; and Heinemann, D. W., 1976, Trop. and Geogr. Med., v. 28 (2), 104-110
dog: Surinam
- Trichuris vulpis*
Roberson, E. L.; and Ager, A. L., 1976, Am. J. Vet. Research, v. 37 (12), 1479-1482
cestodes, nematodes, dogs, natural infections, uredofos highly effective, no toxicosis
- Trichuris vulpis*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, Am. J. Vet. Research, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections

- Trichuris vulpis*
Robinson, M.; Hooke, F.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (3), 173-176
Trichuris vulpis, *Ancylostoma caninum*, *Toxocara canis*, dogs, oxaⁿtel pamoate separately and combined with pyrantel pamoate, drug efficacy, good results: Australia
- Trichuris vulpis*
Ronald, N. C.; and Bell, R. R., 1976, South-west. Vet., v. 29 (3), 217-218
dogs, critical evaluation of butamisol^e hydrochloride, highly effective against *Ancylostoma caninum* and *Trichuris vulpis*, not effective against *Toxocara canis*; *Di^lpylidium caninum* unaffected
- Trichuris vulpis*
Rothwell, T. L. W.; et al., 1976, Vet. Parasitol., v. 1 (3), 221-230
14 common gastrointestinal nematodes, incidence and specificity of anti-acetylcholinesterase antibodies in infected hosts, results show that anti-AChE antibody production occurs in infections with some but not all genera of Strongylida, that not all infected hosts produce detectable antibody, and that the enzyme appears to be genus but not species specific
- Trichuris vulpis* (Froelich, 1789)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Canis familiaris: insular Newfoundland
- Trichuris vulpis*
Tharaldsen, J., 1973, Norwegian J. Zool., v. 21 (4), 327-328 [Abstract]
dogs
cats
(feces of all): all from quarantine station, Oslo, Norway
- Trichuris vulpis*
Todd, K. S.; and Yates, R. L., 1976, Am. J. Vet. Research, v. 37 (11), 1329-1330
Ancylostoma caninum, *Trichuris vulpis*, dogs (exper.), diuredosan, drug efficacy, good results
- Trichuris vulpis*
Turner, T.; and Pegg, E., 1977, Vet. Rec., v. 100 (14), 284-285
survey of patent nematode infestations in dogs (faeces): north-west suburban London
- Trichuris vulpis*
Yang, J.; and Scholten, T., 1977, Am. J. Clin. Path., v. 67 (3), 300-304
diagnosis of human intestinal parasites, fecal examination technique using Jumod's fixative for concentration and permanent staining procedures, comparison with results using formalin-ether procedure
- Trichuroidea*
Chabaud, A. G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott)(1), 6-17
Enoplida
key
- Trichuroidea* [sp.]
Coggins, J. R., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
parasitic fauna, effect of host diet and habitat
Turdus migratorius
Quiscalus quiscula
all from Kellogg Bird Sanctuary, Michigan
- Tricocefali*
Genchi, C., 1976, Arch. Vet. Ital., v. 27 (3-4), 98-99
dog intestinal helminth ova, incidence in soil samples in public parks, potential public health problem: Milan
- Tricocefali*
Genchi, C.; and Locatelli, A., 1974, Atti Soc. Ital. Sc. Vet., v. 28, 862-863
dog helminth eggs, contaminated samples from public parks, potential source of infection for humans and domestic animals: Milan
- Trifurcata* (Schulz, 1926)
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Molineidae, Anoplostrongylinae
- Trilobostrongylus bioccai* Anderson, 1963
Craig, R. E.; and Borecky, R. A., 1976, Canad. J. Zool., v. 54 (5), 806-807
Martes pennanti (lungs): Ontario
- Triodontophorus*
Colglazier, M. L.; Enzie, F. D.; and Kates, K. C., 1977, J. Parasitol., v. 63 (4), 724-727
gastrointestinal parasites of ponies, comparative efficacy of 4 benzimidazoles evaluated by critical test method
- Triodontophorus*
Reinecke, R. K.; and le Roux, D. J., 1972, J. South African Vet. Ass., v. 43 (3), 287-294
adult nematodes, critical tests on donkeys and modified critical tests on horses using mebendazole, highly effective
- Triodontophorus* [sp.]
Cabaret, J., 1976, Rev. Elevage et Med. Vet. Pays Trop. v. 29 (3), 221-226
donkeys, survey, treatment, economic importance: Kaedi area (Mauritania)
- Triodontophorus* spp.
Grelck, H.; Hoerchner, F.; and Woehrl, H. E., 1977, Prakt. Tierarzt, v. 58 (4), 265-268
strongyles, horses, seasonal development of infective larvae on pastures, survival rate
- Triodontophorus* spp.
Nawalinski, T.; and Theodorides, V. J., 1976, Am. J. Vet. Research, v. 37 (4), 469-471
gastrointestinal parasites, ponies, critical tests with o^xibendazole

- Triodontophorus brevicauda*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Triodontophorus brevicauda*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Triodontophorus brevicauda*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Triodontophorus brevicauda*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Triodontophorus minor*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Triodontophorus minor* (Looss, 1900)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Equus caballus: insular Newfoundland
- Triodontophorus serratus*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Triodontophorus serratus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Triodontophorus serratus*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Triodontophorus serratus*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Triodontophorus tenuicollis*
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1975, Am. J. Vet. Research, v. 36 (4), Part 1, 435-439
cambendazole, 3 formulations (suspension, paste, pellet), efficacy against major internal parasites of horses determined by critical testing method
- Triodontophorus tenuicollis*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 128-135
internal parasites of naturally infected horses, critical tests of levamisole alone or mixed with piperazine or trichlorfon, via stomach tube or in feed, varying rates of effectiveness, no toxicosis
- Triodontophorus tenuicollis*
Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1977, Am. J. Vet. Research, v. 38 (12), 2049-2053
internal parasites, horses, critical tests with oxfendazole, powder and pellet formulations
- Triodontophorus tenuicollis*
Ogbourne, C. P., 1976, J. Helminth., v. 50 (3), 203-214
horses (large intestine): south-west England
- Tripyloidina DeConinck*, 1965
Maggenti, A. R., 1976, Organ. Nematodes (Croll), 1-10
Araeolaimida
- Trispiculascaris Skrjabin*, 1916, gen. dub.
Hartwich, G., 1974, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (2), pp. 1-15
? as syn. of *Dujardinascaris Baylis*, 1927
- Trophomera* gen. n.
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
Marimermithidae fam. n., tod: *T. iturupiensis* sp. n.
- Trophomera iturupiensis* sp. n. (tod), illus.
Rubtsov, I. A.; and Platonova, T. A., 1974, Zool. Zhurnal, v. 53 (10), 1445-1458
host unknown: Kuril ridge, near Iturup Island, Dobroe Nachalo Gulf
- Tropisurus crami*
George, R. R.; and Bolen, E. G., 1975, J. Wildlife Dis., v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas

- Tropisurus* (*Petrowimeres*) *indiana* sp. n., illus.
 Ali, M. M., 1970, *Acta Parasitol. Polon.*,
 v. 17 (20-38), 315-327
Casarca ferruginea (proventriculus): Hyderabad,
 Andhra Pradesh, India
- Tropisurus* (*Petrowimeres*) *nettatus* sp. n., illus.
 Ali, M. M., 1970, *Acta Parasitol. Polon.*,
 v. 17 (20-38), 315-327
Netta rufina (proventriculus): Hyderabad,
 Andhra Pradesh, India
- Tropisurus* (*Petrowimeres*) *pavonis* (Tchertkova,
 1954), illus.
 Ali, M. M., 1970, *Acta Parasitol. Polon.*,
 v. 17 (20-38), 315-327
 redescription
Pavo cristatus (proventriculus): Hyderabad,
 Andhra Pradesh, India
- Tropisurus* (*Petrowimeres*) *spinosa* (Maplestone,
 1931) Rasheed, 1960, illus.
 Ali, M. M., 1970, *Acta Parasitol. Polon.*,
 v. 17 (20-38), 315-327
 redescription
Nyroca ferina (proventriculus): Hyderabad,
 Andhra Pradesh, India
- Trypanoxyuris* sp.
 Grundmann, A. W.; and Lombardi, P. S., 1976,
Proc. Helminth. Soc. Washington, v. 43 (1),
 39-46
Ochotona princeps cinnemomea: Tushar Moun-
 tains, Utah
- Tupaiostromylus* Dunn, 1963
 Durette-Desset, M. C.; and Chabaud, A. G.,
 1977, *Ann. Parasitol.*, v. 52 (5), 539-558
 Molineidae, Anoplostrongylineae
- Turgida* Schulz, 1927
 Chabaud, A. G., 1975, *CIH Keys Nematode Para-
 sites Vertebrates* (Anderson, Chabaud, and
 Willmott) (3), 1-27
 Physalopterinae
 key
- Tylenchid nematodes
 Ashford, R. W., 1974, *J. Med. Entom.*, v. 11
 (5), 605-616
Sergentomyia affinis (hemocoel): Ethiopia
- Tylenchida Thorne, 1949
 Maggenti, A. R., 1976, *Organ. Nematodes*
 (Croll), 1-10
 Secernentea
- Tylenchides
 Beaucournu, J. C.; and Deunff, J., [1976],
Ann. Parasitol., v. 50 (6), 1975, 831-835
 as hyperparasites of fleas, causing para-
 sitic castration
 siphonapteres: France
- Tylenchoidea, illus.
 Wright, K. A., 1976, *Organ. Nematodes* (Croll),
 71-105
- Typhlophorus* von Linstow, 1906
 Hartwich, G., 1974, *CIH Keys Nematode Para-
 sites Vertebrates* (Anderson, Chabaud, and
 Willmott) (2), pp. 1-15
 Multicaecinea
 key

- Uncinaria**
Biagi, F.; and Medina, J. A., 1974, Prensa Med. Mexicana, v. 39 (9-10), 440-443
human intestinal parasites, pyrantel pamoate, metabolism and pharmaceutical action
- Uncinaria**
Botto, C.; and Mane-Garzon, F., 1975, Rev. Biol. Uruguay, v. 3 (2), 127-141
influence of geographical factors in speciation
- Uncinaria**
Hernandez Chinas, C.; Biagi Finisola, F.; and Salmon Rodriguez, L., 1975, Bol. Med. Hosp. Inf., v. 32 (4), 621-627
uncinariasis, children, epidemiologic and statistical survey: central Mexico
- Uncinaria**
Ockens, N., 1977, Medlemsbl. Danske Dyrlægeforen., v. 60 (4), 147-148
Ancylostomum, Uncinaria, dogs, symptoms, good results with banminth
- Uncinaria [sp.]**
Ashford, R. W., 1977, Ann. Trop. Med. and Parasitol., v. 71 (1), 29-34
Vulpes vulpes (feces): Wales; Exmoor
- Uncinaria [sp.]**
Hass, D. K.; and Chitwood, M. B., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 255
dog (feces): west central Ohio
- Uncinaria hamiltoni platensis n. ssp., illus.**
Botto, C.; and Mane-Garzon, F., 1975, Rev. Biol. Uruguay, v. 3 (2), 127-141
Otaria flavescens (intestino delgado): Isla Rasa (Islas de Torres, costa oceanica del Uruguay)
- Uncinaria lotoris Schwartz, 1925**
Barnstable, R. W.; and Dyer, W. G., 1974, Tr. Illinois State Acad. Sc., v. 67 (4), 451-460
as syn. of Placoconus lotoris (Schwartz, 1925) Webster, 1956
- Uncinaria lucasi**
Chauhan, H. V. S.; Dwivedi, P.; and Kalra, D. S., 1974, Haryana Vet., v. 13 (1), 5-21
protozoan and helminth parasites, transmitted through milk to newborn animals, review
- Uncinaria (Megadeirides) olseni n. sp., illus.**
Chabaud, A. G.; and Durette-Desset, M. C., [1976], Ann. Parasitol., v. 50 (6), 1975, 789-793
Tupaia glis (intestin grele): Sepilok, region de Sandakan, Sabah (Borneo)
- Uncinaria stenocephala (Railliet, 1884)**
Alcaino, H., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 110-113
Uncinaria stenocephala, efficacy of tetramizole, dogs (exper.), comparative trials with thiabendazole which was not effective
- Uncinaria stenocephala, illus.**
Alcaino, H.; and Huerta, J., 1970, Bol. Chileno Parasitol., v. 25 (3-4), 136-137
Uncinaria stenocephala discovered in intestines of 2 dogs (perros) at necropsy, morphological descriptions: Chile
- Uncinaria stenocephala**
Davies, P.; and Nicholas, W. L., 1977, Austral. Vet. J., v. 53 (5), 247-248 [Letter]
dogs (feces): Goodradigbee Shire, New South Wales
- U[ncinaria] stenocephala**
Deiana, S.; and Arru, E., 1972, Parassitologia, v. 14 (2-3), 269-273
presence in dogs being treated with Mansoniil for test of cestode control
- Uncinaria stenocephala (Railliet 1884)**
Gilbertson, D. E., 1977, J. Parasitol., v. 63 (1), 162-163
Vulpes fulva (intestine): Dakota County, Minnesota
- Uncinaria stenocephala**
Gregory, G. G.; and Munday, B. L., 1976, Austral. Vet. J., v. 52 (7), 317-320
feral cats: Tasmanian Midlands and King Island
- Uncinaria stenocephala**
Guildal, J. A.; and Clausen, B., 1973, Norwegian J. Zool., v. 21 (4), 329-330 [Abstract]
Vulpes vulpes: Denmark
- Uncinaria stenocephala**
Hass, D. K.; and Collins, J. A., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 135-137
helminths, dogs, comparative efficacy of vincofos, ticarbondine, mebendazole
- Uncinaria stenocephala**
Jacobs, D. E., 1976, Research Vet. Sc., v. 21 (2), 238-239
Uncinaria stenocephala, seasonal fluctuations in numbers of third stage larvae on grass runs of greyhound kennel: Britain
- Uncinaria stenocephala**
Kingsbury, P. A.; Rees, T. A.; and Piercy, D. W. T., 1977, Vet. Rec., v. 101 (24), 477-479
nematodes, dogs, cats (both nat. and exper.), haloxon, efficacy and safety trials, good results with no significant side effects

- Uncinaria stenocephala* (Railliet, 1884) Railliet, 1885
Kozlov, D. P., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 71-78
Canis familiaris
Vulpes vulpes
Alopex lagopus
all from Pechora river basin
- Uncinaria stenocephala*
McCurdy, H. D.; and Guerrero, J., 1977, Vet. Med. and Small Animal Clin., v. 72 (11), 1731-1733
helminths, dogs, mebendazole powder, controlled critical studies, good results against all helminths except *Dipylidium caninum*: Kansas; New Jersey; Texas
- Uncinaria stenocephala*
Pegg, E. J., 1977, Brit. Vet. J., v. 133 (4), 427-431
Toxocara canis and other parasite ova, horticultural flame-gun for control on concrete-floored kennel runs
- Uncinaria stenocephala*
Roberson, E. L.; and Ager, A. L., 1976, Am. J. Vet. Research, v. 37 (12), 1479-1482
cestodes, nematodes, dogs, natural infections, uredofos highly effective, no toxicosis
- Uncinaria stenocephala*
Roberson, E. L.; Anderson, W. I.; and Hass, D. K., 1977, Am. J. Vet. Research, v. 38 (5), 597-600
intestinal nematodes, dogs, dichlorvos-mediated dry dog feed, fast vs. slow release rate, various doses; no drug-related complications from *Dirofilaria immitis* infections
- Uncinaria stenocephala*
Robinson, M.; Hooke, F. G.; and Iverson, K. E., 1976, Austral. Vet. Practitioner, v. 6 (2), 104-108
pyrantel pamoate, critical trials against roundworms and hookworms in naturally and experimentally infected dogs, drug efficacy compared with piperazine citrate: Australia
- Uncinaria stenocephala* (Railliet, 1884)
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Meles meles: Karelia
- Uncinaria stenocephala* (Railliet, 1884)
Smith, F. R.; and Threlfall, W., 1973, Am. Midland Naturalist, v. 90 (1), 215-218
Canis familiaris: insular Newfoundland
- Uncinaria stenocephala*
Strel'chik, V. A.; Shnaidmiller, A. P.; and Gapon, N. M., 1976, Sborn. Nauch. Rabot. SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (26), 123-128
[pig, wild]: Primorskii krai
- Uncinaria stenocephala*
Tharaldsen, J., 1973, Norwegian J. Zool., v. 21 (4), 327-328 [Abstract]
dogs (feces): quarantine station, Oslo, Norway
- Uncinaria stenocephala*
Turner, T.; and Pegg, E., 1977, Vet. Rec., v. 100 (14), 284-285
survey of patent nematode infestations in dogs (faeces): north-west suburban London
- Uncinaria stenocephala*
Williams, B. M., 1976, Brit. Vet. J., v. 132 (3), 309-312
Vulpes vulpes (intestine): southwest Wales
- Uncinariasis*
Biagi, F.; Alvarez, R.; and Delgado, R., 1972, Prensa Med. Mexicana, v. 37 (5-6), 221-223
human intestinal helminths, clinical trials with pyrantel pamoate, good results

- Varestrongylus [sp.]
Tewari, H. C.; Dhar, D. N.; and Singh, K. S.,
1973, *Isotopes and Radiation Parasitol.* III,
43-50
sheep: Kashmir, India
- Vasorhabdochona Martin & Zam, 1967
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 1-27
Rhabdochoniidae
key
- Vaznema Freitas & Lent, 1947
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
as syn. of *Odontospirura* Wehr, 1933
- Vexillata (Travassos, 1937)
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, Viannaiinae
- Viannaia Travassos, 1914 (type genus)
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, Viannaiinae
Syn.: *Philostrongylus* Wolfgang, 1951
- Viannaia metachirops n. sp., illus.
Durette-Desset, M. C., 1974, *Ann. Parasitol.*,
v. 49 (5), 555-566
4 spp. of *Trichostrongyloidea*, localization
in intestine, larval and adult synophes
compared, implications for taxonomy and
evolution
Metachirops opossum (intestin): Guyane
Francaise
- Viannaia viannai Travassos, 1914
Diaw, O. T., 1976, *Ann. Parasitol.*, v. 51 (3),
355-363
trichostrongyloid nematode fauna of *Didelphis
marsupialis* compared to that of *Metachirops
opossum*, localization within intestine
Didelphis marsupialis (intestin): Guyane
francaise
- Viannaia viannai Travassos, 1914
Diaw, O. T., [1977], *Bull. Mus. National Hist.
Nat.*, Paris, 3. s. (405), 1976, *Zool.* (282),
1065-1089
Didelphis marsupialis
- Viannaia viannai Travassos, 1914, illus.
Durette-Desset, M. C., 1974, *Ann. Parasitol.*,
v. 49 (5), 555-566
4 spp. of *Trichostrongyloidea*, localization
in intestine, larval and adult synophes
compared, implications for taxonomy and
evolution
Metachirops opossum (intestin): Guyane
Francaise
- Viannaiinae Neveu-Lemaire, 1934
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae
includes: *Viannaia* (type genus); *Heligmo-
skrjabinia*; *Hoineffia*; *Oswaldonema*; *Travas-
sostrongylus*; *Vexillata*; *Viannella*
- Viannella* Travassos, 1919
Durette-Desset, M. C.; and Chabaud, A. G.,
1977, *Ann. Parasitol.*, v. 52 (5), 539-558
Heligmosomidae, Viannaiinae
synonymy
- Viannella* sp.
Sood, M. L.; and Parshad, V. R., 1975, *Riv.
Parassitol.*, Roma, v. 36 (2-3), 189-196
infections in *Millardia meltada*, survey of
seasonal distribution, possible correla-
tions between host diet and sex and inci-
dence of infection
- Victorocara*. See *Viktorocara*.
- Vigisospirura* Petrow & Potekhina, 1953, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
Spirocercinae
key
"Chlamydopecta Chandler, 1954 may be a
synonym of *Vigisospirura*."
- Vigisospirura* potekhini (Petrow et Potekhina,
1953)
Gafurov, A. K., 1969, *Trudy Gel'mint. Lab.,
Akad. Nauk SSSR*, v. 20, 46-54
role of *Tenebrionidae* as intermediate hosts
Blaps fausti bactriana
Trigonoscelis gemmulata
T. ceromatica
Pisterotarsa kiritschenkoi
Pachyscelis banghaasi
P. laevicollis
Cyphogenia gibba
Stalagmoptera incostata
all from Tadjik SSR [and/or] Uzbek SSR
- Viguiera* Seurat, 1913, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
Histocephalinae
key; synonymy
- Viktorocara* Guschanskaja, 1950, illus.
Chabaud, A. G., 1975, *CIH Keys Nematode Para-
sites Vertebrates* (Anderson, Chabaud, and
Willmott) (3), 29-58
Schistorophinae
key
- Viktorocara* sp.
Bush, A. O.; and Forrester, D. J., 1976, *Proc.
Helminth. Soc. Washington*, v. 43 (1), 17-23
Eudocimus albus (gizzard lining): Florida
- Victorocara halcyoni* Ryjnikov et Hohlova, 1964
Aleksseev, V. M.; and Smetanina, Z. B., 1968,
Gel'mint. Zhivot. Tikhogo Okeana (Skriabin),
97-104
Halcyon pileata (esophagus): Rimsko-Korsakov
islands
- Victorocara schejkini* Guschanskaja, 1950
Bondarenko, S. K., 1969, *Trudy Gel'mint. Lab.,
Akad. Nauk SSSR*, v. 20, 35-45
Limosa limosa lapponica
Charadrius hiaticula
all from lower Yenisei [and/or] Keta lake

- Vogeloides Orlov, Davtian & Lubimov in Skrjabin, 1933, illus.
 Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
 Pneumospiruridae
 key
 Syn.: Osleroides Orlov, Davtian & Lubimov, 1933
- Vogeloides
 Pence, D. B.; and Stone, J. E., 1977, J. Parasitol., v. 63 (6), 979-991
 systematics of family Pneumospiruridae
- Vogeloides
 Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
 Thelazioidea, Pneumospiruridae
 cephalic structures, scanning electron microscopy
- Vogeloides chabaudi (Singh and Pande, 1956 [i.e., 1966]), n. comb.
 Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
- Vogeloides congolense (Vuylsteke, 1956), n. comb.
 Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
- Vogeloides felis (Vogel 1928) Davtian 1933, illus.
 Pence, D. B.; and Stone, J. E., 1977, J. Parasitol., v. 63 (6), 979-991
 synonymy
 redescription, pathology
 Felis rufus (bronchi, primary and secondary bronchioles of lung): Benjamin, Knox Co., Texas; Adams Ranch, King Co., Texas; Pitchfork and Beggs Ranches, Dickens Co., Texas
- Vogeloides oesophageus (Gerichter, 1948), Dougherty, 1952, illus.
 Wertheim, G.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (6), 647-657
 cephalic structures, scanning electron microscopy
 [Syn.]: Metathelazia oesophageus

- Waltonella Schacher, 1974 (= Waltonia Schacher et Crans, 1973, sous-genre)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
Onchocercidae; Waltonellinae n. subfam.
- Waltonella americana (Walton, 1929)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella brachyoptera (Wehr et Causey, 1939)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella confusa (Schmidt et Kuntz, 1969)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella convoluta (Molin, 1858)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella dolichoptera (Wehr et Causey, 1939)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella duboisi (Gedoelst, 1916)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella flexicauda (Schacher et Crans, 1973)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella flexicauda
Terwedow, H. A., jr.; and Craig, G. B., jr., 1977, Exper. Parasitol., v. 41 (2), 272-282
Waltonella flexicauda, development controlled by genetic factor in Aedes aegypti, this factor for susceptibility did not control development of Brugia pahangi or Dirofilaria immitis
- Waltonella flexicauda (Schacher and Crans, 1973) = Foleyella flexicauda Schacher and Crans (1973)
Terwedow, H. A., jr.; and Craig, G. B., jr., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 86-91
Waltonella flexicauda, development in culicine mosquitoes, variation in mosquito susceptibility, selection for susceptibility, defining refractoriness
Aedes aegypti (exper.)
A. annandalei (exper.)
A. atropalpus (exper.)
A. polynesiensis (exper.)
Culex pipiens quinquefasciatus (exper.)
Rana catesbeiana: Morris County, New Jersey
- Waltonella guyanensis n. sp., illus.
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
Bufo marinus (mesentere dorsal): Maripasoula, Guyane francaise
- Waltonella ranae (Walton, 1929)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella scalaris (Travassos, 1929)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella striatus (Ochoterena et Caballero, 1932)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonella vellardi (Travassos, 1929)
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
- Waltonellinae n. subfam.
Bain, O.; and Prod'hon, J., [1975], Ann. Parasitol., v. 49 (6), 1974, 721-739
Onchocercidae; includes: Waltonella; Ochoterena; Madochotera
- Warrenius Hall, 1916
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
as syn. of Citellinema Hall, 1916
- Wehrdikmansia sp.
Low, W. A., 1976, Canad. Field-Naturalist, v. 90 (2), 189-191
Rangifer tarandus caribou (over calcaneum subdermal): Tweedsmuir Provincial Park, British Columbia
- Wehrdikmansia cervipedis (Wehr et Dikmans, 1935)
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
Wehrdikmansia cervipedis returned to Onchocerca
- Wehrdikmansia cervipedis sensu Tarczynski, 1954
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
as syn. of Onchocerca tubingensis n. sp.
- Wehrdikmansia cervipedis
Samuel, W. M.; Barrett, M. W.; and Lynch, G. M., 1976, Canad. J. Zool., v. 54 (3), 307-312
helminths of Alces alces, 3 study areas, differences in parasite prevalence due to fauna and ecology of habitat and age of host: Alberta, Canada
- Wehrdikmansia cervipedis
Weinmann, C. J.; et al., 1973, J. Wildlife Dis., v. 9 (3), 213-220
Wehrdikmansia cervipedis, Elaeophora schneideri, Setaria yehi, prevalence of infection in various age classes of Odocoileus hemionus columbianus: northern California
- Wehrdikmansia rugosicauda Bohm et Supperer, 1953
Bain, O.; and Schulz-Key, H., 1974, Tropenmed. u. Parasitol., v. 25 (4), 437-449
"affinites avec les Dipetalonema sensu largo"

- Wehrdikmansia rugosicauda Bohm und Supperer 1953
Schulz-Key, H., 1975, Tropenmed. und Parasitol., v. 26 (4), 494-498
development and distribution of microfilariae (near nose, ears, skin of legs) and adults (subcutaneously in connective tissue of back) in Capreolus capreolus; Ixodes ricinus probable vector: Southern Germany
- Wehrdikmansia rugosicauda
Schulz-Key, H.; Bain, O.; and Wenk, P., 1976, Tropenmed. u. Parasitol., v. 27 (2), 229-232
distribution pattern of Onchocerca garmsi microfilariae in the skin of Cervus elaphus and comparisons with distribution of adults and microfilariae of Onchocerca tubingensis, O. cervicalis, O. tarsicola and Wehrdikmansia rugosicauda: Sueddeutschland
- Wellcomia sp.
O'Farrell, T. P., 1975, Am. Midland Naturalist, v. 93 (2), 377-387
Peromyscus maniculatus: Arid Lands Ecology Reserve, Benton County, Washington
- Whipworms
Chakraborty, B.; Ray, N. M.; and Sikdar, S., 1976, Indian J. Animal Health, v. 15 (2), 117-119
Ananas sativus, in vitro tests against whipworms and flatworms, fair anthelmintic activity but showed less potency than piperazine
- Whipworm
Oomi, H.; Watanabe, T.; and Yoshida, Y., 1976, J. Agric. Sc., Tokyo Nogyo Daigaku, v. 20 (3-4), 206-212
hookworm, whipworm, dogs, subcutaneous injection, methyridine + disophenol, good results
- Woodwardstrongylus Wahid, 1964
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Mackerrastrongylinae
- Woodwardstrongylus Wahid, 1964
Mawson, P. M., 1976, Tr. Roy. Soc. South Australia, v. 100 (3), 121-123
Woodwardstrongylus transferred to family Amidostomatidae
Syn.: Cristiceps Mawson, 1971
- Woodwardstrongylus obendorfi n. sp., illus.
Mawson, P. M., 1976, Tr. Roy. Soc. South Australia, v. 100 (3), 121-123
Macropus parryi
M. robustus
M. rufogriseus
(oesophagus of all): all from Warwick, Queensland
- Woolleya Mawson, 1973
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Amidostomatidae, Amidostomatinae
- Woolleya hickmani Mawson
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, Antechinus stuartii, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (intestine): Powelltown, Victoria
- Woolleya monodelphis Mawson
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, Antechinus stuartii, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (intestine): Powelltown, Victoria
- Woolleya sprengi Mawson
Beveridge, I.; and Barker, I. K., 1976, Austral. J. Zool., v. 24 (2), 265-272
helminths and arthropods, Antechinus stuartii, seasonal and sex-related variations in numbers of helminths, parasites unlikely directly involved in seasonal mortality of male host; ectoparasites may contribute to anemia in hosts
A. stuartii (intestine): Powelltown, Victoria
- Wuchereria bancrofti
Abdalla, R. E., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (1), 53-55
Wuchereria bancrofti and Onchocerca volvulus as cause of human filariasis with hydrocele and elephantiasis, newly reported cases in previously non-endemic areas, need for differentiation from Loa loa: Sudan
- Wuchereria bancrofti, illus.
Adickman, M.; and Tuthill, T. M., 1976, Postgrad. Med., v. 60 (3), 143-148
pulmonary symptoms and eosinophilia associated with human parasitic infections, diagnostic and clinical review, need for increased awareness in travelers to endemic areas, immigrants and military personnel
- Wuchereria bancrofti
Aikat, T. K.; and Das, M., 1977, Indian J. Med. Research, v. 65 (1), 58-64
Wuchereria bancrofti, modified statistical method for analysis of periodicity of microfilaria using harmonic wave equation
- Wuchereria bancrofti
Ambroise-Thomas, P.; and Kien Truong, T., 1974, Ann. Trop. Med. and Parasitol., v. 68 (4), 435-452
filariasis, human, diagnosis, indirect fluorescent antibody test on sections of adult filariae (Dipetalonema viteae, Dirofilaria immitis, Wuchereria bancrofti, Loa loa, Onchocerca volvulus), possible application to epidemiological surveys and post-therapeutic surveillance

- Wuchereria bancrofti*
Ambroise-Thomas, P.; Prod'hon, J.; and Venard, P., 1975, Bull. Soc. Path. Exot., v. 68 (6), 579-588
Wuchereria bancrofti, humans, epidemiologic prevalence survey comparing findings from microfilaremia determinations and those from indirect immunofluorescence studies using *Dipetalonema viteae* antigens: south-east coast of Madagascar
- Wuchereria bancrofti*
Anderson, R. I.; and Buck, A. A., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 447-456
complement levels in residents of rural village in relation to wide variety of clinical, laboratory, and epidemiological factors including parasitic diseases: Ouli Bangala, Republic of Chad
- Wuchereria bancrofti*, *illus.*
Aoki, Y., 1971, Nettai Igaku (Trop. Med.), v. 13 (3), 134-140
Wuchereria bancrofti, *Brugia malayi*, *B. pahangi*, demonstration of exsheathing of microfilariae on thick blood film or on agar plate, effects of temperature on exsheathment
- Wuchereria bancrofti*
Aoki, Y., 1971, Nettai Igaku (Trop. Med.), v. 13 (4), 170-179
Wuchereria bancrofti, *Brugia pahangi*, exsheathing of microfilariae on thick blood film or on agar plate, effects of temperature, salinity, and pH
- Wuchereria bancrofti*
Bain, O., 1976, Bull. World Health Organ., v. 54 (4), 397-401
human filariasis, number of developing and infective larvae dependent upon number of microfilariae penetrating into haemocoel of vector, relationship based on proportionality, facilitation and limitation, application to disease control and treatment methods
- Wuchereria bancrofti*
Barry, C.; Ahmed, A.; and Khan, A. Q., 1971, Am. J. Trop. Med. and Hyg., v. 20 (4), 592-597
statistics of prevalence and distribution survey of human *Wuchereria bancrofti* filariasis in East Pakistan, possible sociologic aspects affecting disease transmission
- Wuchereria bancrofti*
Brinkmann, U. K., 1976, Tropenmed. u. Parasitol., v. 27 (1), 50-56
Wuchereria bancrofti, humans, effect of periodicity on microfilaremia prevalence surveys with assessment of variations in density if surveying in sub-peak hours: Liberia
- Wuchereria bancrofti*
Brinkmann, U. K., 1977, Tropenmed. u. Parasitol., v. 28 (1), 71-76
Wuchereria bancrofti, epidemiologic survey comparing 5 areas demonstrated that bancroftian filariasis is major health problem in rural coastal Liberia
- Wuchereria bancrofti*
Bryan, J. H.; and Southgate, B. A., 1976, Tr. Roy. Soc. Trop. Med. and Hyg., v. 70 (1), 39-48
Wuchereria bancrofti, assessment of diethylcarbamazine administration campaign for eradication of human infection and its effects on local vector *Aedes polynesiensis*, recommendations for continued surveillance and control: Western Samoa
- Wuchereria bancrofti*, *illus.*
Chandrasoma, P. T.; Mendis, K. N.; and Kumararatne, D. S., 1977, Am. J. Trop. Med. and Hyg., v. 26 (3), 570-571
Wuchereria bancrofti microfilariae removed from breast nodule of woman suffering from tropical pulmonary eosinophilia: Colombo, Sri Lanka
- Wuchereria bancrofti*, *illus.*
Charters, A. D., 1970, Med. J. Australia, v. 2 (20), 919-922
Wuchereria bancrofti filariasis in immigrants living in Western Australia; case reports including woman who experienced chyluria with each of 7 pregnancies; public health importance of increasing immigration from endemic areas
- [*Wuchereria bancrofti*] bancroftian microfilariae
Chen, C. C.; and Fan, P. C., 1977, Southeast Asian J. Trop. Med. and Pub. Health, v. 8 (1), 53-55
bancroftian filariasis, residual microfilariae in blood of diethylcarbamazine-treated human carriers, ability to develop into infective larvae after being ingested by vector mosquitoes, possible source of disease transmission in endemic areas after suspension of control measures
- Wuchereria bancrofti*
Cheong, W. H.; and Omar, A. H. B., 1965, Med. J. Malaya, v. 20 (1), 74-75
Anopheles maculatus as new vector of *Wuchereria bancrofti* in Malaysia
- Wuchereria bancrofti* ?
Cheong, W. H.; and Tang, K. Y., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 138-139 [Demonstration]
probable *Wuchereria bancrofti* involvement of genitalia in man, excision of growth in left epididymis revealed apparent female adult filarial worm, no microfilariae found, case report: Kuala Lumpur, Malaysia
- Wuchereria bancrofti*
Chernin, E., 1977, Am. J. Trop. Med. and Hyg., v. 26 (5, part 2), 1065-1070
human filariasis, historical review of Patrick Manson's investigations on transmission
- Wuchereria bancrofti*
Chlebowski, H. O.; and Zielke, E., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (2), 181-182 [Letter]
Wuchereria bancrofti, *Onchocerca volvulus* in humans, diagnosis of microfilariuria using a modified membrane filter technique; application also to filtration of eggs from urine of persons suspected to have schistosomal infections: Liberia, West Africa

- Wuchereria bancrofti*
Chow, C. Y., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 404-418
Wuchereria bancrofti, *Brugia malayi*, up-to-date review of confirmed and suspected vectors in WHO Western Pacific Region, available data of natural and experimental infection and infective rates of most vector mosquitoes, remarks on ecology and control
- Wuchereria bancrofti*
Crans, W. J., 1973, J. Med. Entom., v. 10 (2), 189-193
Wuchereria bancrofti, laboratory strains of *Anopheles gambiae* and *Culex pipiens fatigans*, comparison of vector potential, *C. pipiens fatigans* far superior; differences may be due to strain rather than natural susceptibility
- Wuchereria bancrofti*
Dennis, D. T.; McConnell, E.; and White, G. B., 1976, Am. J. Trop. Med. and Hyg., v. 25 (2), 257-262
Wuchereria bancrofti, human, diagnosis by Nucleopore filter more sensitive than conventional methods (thick film, counting chamber), increasing volume of blood filtered from 1 to 5 ml increased numbers of carriers identified, filtering 5 ml of day blood is as sensitive as filtering 5 ml of night blood: southwestern Ethiopia
- Wuchereria bancrofti*
Desowitz, R. S.; and Southgate, B. A., 1973, Southeast Asian J. Trop. Med. and Pub. Health, v. 4 (2), 179-183
Wuchereria bancrofti, comparative efficacy of membrane filtration, counting-chamber and stained thick blood-film techniques in identifying persistent low density microfilaremia in natives following therapeutic course of diethylcarbamazine given during mass drug campaign: Fiji and Western Samoa
- Wuchereria bancrofti*
Desowitz, R. S.; and Una, S. R., 1976, J. Helminth., v. 50 (1), 53-57
Dirofilaria immitis-infected cats and dogs, *Wuchereria bancrofti*-infected humans, diagnosis, counterimmunoelectrophoresis using *D. immitis* adult and microfilarial antigens
- Wuchereria bancrofti*
Diesfeld, H. J.; Dutta, S. N.; and Braun-Munzinger, R., 1973, Ztschr. Tropenmed. u. Parasitol., v. 24 (4), 439-446
Wuchereria bancrofti-endemic area, survey of 225 Indians, microfilaremia, fluorescent antibody titer, clinical manifestations, eosinophilia, immunoglobulin levels: Dhanbad/Asansol, India
- W[*Wuchereria*] *bancrofti*
Diesfeld, H. J.; and Kirsten, C., 1975, Tropenmed. und Parasitol., v. 26 (4), 499-502
Dipetalonema vitae, localization of antigen-antibody reactions in male and female using immunofluorescence and serum from human filariasis patients; possible implications for diagnosis of human infection
- Wuchereria bancrofti*, illus.
Dissanaike, A. S., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (1), 139-141 [Demonstration]
larval development to fourth stage in pulmonary vessels of splenectomised hamster (exper.) but growth retarded because of unsuitability of host
- Wuchereria bancrofti*
Dondero, T. J., jr.; et al., 1976, Am. J. Trop. Med. and Hyg., v. 25 (1), 64-73
Wuchereria bancrofti, human, type and frequency of clinical manifestations in an area of high intensity year round exposure, genitalia of males most commonly affected but in general little noticeable serious filarial disease: Howrah, near Calcutta, India
- Wuchereria bancrofti*
Dondero, T. J., jr.; and Sivanandam, S., 1973, Med. J. Malaysia, v. 27 (4), 306-309
no evidence of human filariasis infection in "spot" blood survey of 4 localities, not public health problem in Kelantan River area of Malaysia
- [*Wuchereria bancrofti*] bancroftian filariasis
Dutta, S. N., 1977, Tr. Roy. Soc. Trop. Med. and Hyg., v. 71 (2), 180 [Letter]
Culex fatigans (bancroftian filariasis vector mosquito) found breeding in underground pits of coalmine which is in endemic filariasis area of India
- Wuchereria bancrofti*
Dutta, S. N.; Diesfeld, H. J.; and Kirsten, C., 1976, Tropenmed. u. Parasitol., v. 27 (4), 479-482
immunofluorescent antibody test using *Dipetalonema* viteae as antigen applied to mothers' blood and to the umbilical cord blood of their newborns, results show that in a *Wuchereria bancrofti* endemic area maternal antibodies against filariae are passed via placenta to newborn, findings not demonstrable after 6th month of life: India
- Wuchereria bancrofti*
Fan, P. C.; et al., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 211-222
Wuchereria bancrofti, humans, extensive epidemiologic survey of civilian and military populations of island to establish infection rates and possible means of control of transmission of filariasis from civilians to military troops garrisoned on Kinmen (Quemoy) Islands
- [*Wuchereria*] *bancrofti* "bancroftian filariasis"
Fan, P. C.; et al., 1975, Ann. Trop. Med. and Parasitol., v. 69 (4), 515-516
bancroftian filariasis, control by use of common salt medicated with diethylcarbamazine, reduced prevalence in humans and in *Culex p. fatigans*: Liehyu District (Little Kinmen), Kinmen (Quemoy) Islands, Republic of China

- Wuchereria bancrofti*
Fouques, M.; et al., 1972, *Medecine Afrique Noire*, v. 19 (2), 83-87
Wuchereria bancrofti, surgical treatment of elephantiasis and lymphatic system pathology resulting from infection
- Wuchereria bancrofti*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, *Medecine et Malad. Infect.*, v. 3 (8-9), 351-353
diagnostic review of human filariasis
- Wuchereria bancrofti*, *illus.*
Gonzaga dos Santos, L.; Santos, D. S.; and Azevedo, R., 1976, *Ann. Trop. Med. and Parasitol.*, v. 70 (2), 219-225
Wuchereria bancrofti, human, diagnosis, immunofluorescence using microfilariae treated with papain as antigen
- Wuchereria bancrofti*
Govindwar, S. L.; Ghirnikar, S. N.; and Hari-nath, B. C., 1976, *Indian J. Med. Research*, v. 64 (11), 1607-1610
Wuchereria bancrofti, microfilaria carriers, chronic filarial patients and normal controls, biochemical study of blood samples comparing parameters of proteins, lipids, electrolytes, enzymes
- Wuchereria bancrofti*
Grove, D. I.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (2), 220-229
antigen abstract prepared from subperiodic *Brugia malayi* compared with *Dirofilaria immitis* antigen in diagnosis of human filari-iasis, concluded that antigens from micro-filariae, adult worms, and 3rd-stage larvae of *B. malayi* are more sensitive than *D. immitis* antigens and do not have a signifi-cantly higher number of false positive re-actions: Philippines
- Wuchereria bancrofti*
Grove, D. I.; Warren, K. S.; and Mahmoud, A. A. F., 1975, *J. Infect. Dis.*, v. 132 (3), 340-352
algorithms in the diagnosis and management of human forms of filariases
- Wuchereria bancrofti*
Guptavani, P.; and Harinasuta, C., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 578 [Demonstration]
Brugia malayi, *Wuchereria bancrofti*, spon-taneous disappearance of microfilariae from 3 human carriers after they were transferred from an endemic area to a non-endemic area in Bangkok
- Wuchereria bancrofti*
Harinasuta, C.; Sucharit, S.; and Vutikes, S., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (1), 102-103 [Demonstration]
Wuchereria bancrofti, mosquito survey for possible vectors
Anopheles stephensi
A. minimus
A. vagus
A. subpictus malayensis
A. maculatus
A. philippinensis
Culex pallens
Anopheles hyr[canus] sinensis
Aedes togoi
A. aegypti
Culex molestus
(all exper.)
- Wuchereria bancrofti*
Hawking, F., 1974, *Progr. Drug Research*, v. 18, 173-190
Wuchereria bancrofti, *Brugia malayi*, review of present status of human infections in India with emphasis on public health issues and possible control measures
- Wuchereria bancrofti*
Hembree, S. C., 1974, *Mil. Med.*, v. 139 (4), 309-312
Wuchereria bancrofti, distribution and preva-lence in US Army Special Forces Camps: Republic of Vietnam
- Wuchereria bancrofti*, *illus.*
Hira, P. R., 1975, *Ann. Trop. Med. and Para-sitol.*, v. 69 (4), 521-522
human, first autochthonous case reported from Zambia; characteristics in relation to report of possible new microfilaria from Western India
- Wuchereria bancrofti*
Ismail, M. M.; and Nagaratnam, N., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (3), 405-409
filariasis as possible cause of arthritis, clinical features and laboratory findings in 33 cases, age distribution, diethylcarba-mazine treatment gave good results: Sri Lanka
- Wuchereria bancrofti*
Iwamoto, I., 1971, *Nettai Igaku (Trop. Med.)*, v. 13 (1), 1-6
Wuchereria bancrofti microfilariae in car-rier dogs, small amounts of diethylcarba-mazine in blood cause temporary appearance of larvae and disturbance of 24-hour micro-filarial rhythm
- Wuchereria bancrofti*
Iwamoto, I., 1972, *Nettai Igaku (Trop. Med.)*, v. 14 (3), 124-137
Dirofilaria immitis, *Wuchereria bancrofti*, in vitro survival in various media, effects of temperature; *D. immitis* survival in dogs (exper.), periodicity, migration and local-ization

- W[uchereria] bancrofti, illus.
Jones, T. C.; Mott, K.; and Pedrosa, L. C., 1975, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 69 (2), 243-246
technique for isolating and concentrating microfilariae from peripheral blood using gradient centrifugation
- Wuchereria bancrofti
Katamine, D., 1969, *Nettai Igaku (Trop. Med.)*, v. 11 (1), 1-10
Wuchereria bancrofti in humans, skin test diagnosis using purified antigen (FPT) prepared from *Dirofilaria immitis*, useful for tool in mass diagnostic survey
- Wuchereria bancrofti
Katamine, D.; et al., 1967, *Nettai Igaku (Trop. Med.)*, v. 9 (3), 143-157
Acanthocheilonema perstans, *Wuchereria bancrofti*, prevalence survey in area residents: Tanzania
- Wuchereria bancrofti
Katiyar, J. C.; et al., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (2), 169-170 [Letter]
human filariasis, persons undergoing diethyl-carbamazine therapy for *Wuchereria bancrofti*, interference with skin test reactions when *W. bancrofti* used as antigen, review of possible mechanisms
- Wuchereria bancrofti
Krafsur, E. S.; and Garrett-Jones, C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (2), 155-160
Wuchereria bancrofti, statistical survey of factors affecting survival and infectivity of *Wuchereria*-infected *Anopheles funestus* mosquito vectors living in natural environment: north-eastern Tanzania
- Wuchereria bancrofti
Kuhlow, F.; and Zielke, E., 1976, *Tropenmed. u. Parasitol.*, v. 27 (1), 93-100
night blood survey of 82 localities to establish distribution and prevalence of *Wuchereria bancrofti* in various parts of Liberia, humans
- Wuchereria bancrofti
Kurihara, T.; Ikemoto, T.; and Eshita, Y., 1977, *Mosquito News*, v. 37 (4), 770-772
Wuchereria bancrofti, survey of mosquito vectors
Culex quinquefasciatus: Bicol Region, southern Luzon, Philippines
- Wuchereria bancrofti
Lambrecht, F. L., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (2), 222-232
Wuchereria bancrofti, statistics of epidemiologic survey of periodic-nocturnal infections present in island natives, *Culex pipiens fatigans* established as the main and possibly only vector: Seychelles Islands
- Wuchereria bancrofti
Lambrecht, F. L.; and Fernando, J. B., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (1), 76-79
Wuchereria bancrofti, age-grading of *Culex pipiens fatigans* vector mosquitoes, assessment in relationship to vectorial capacities
- Wuchereria bancrofti
Lambrecht, F. L.; and Van Someren, E. C., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (4), 483-485
Wuchereria bancrofti, *Culex pipiens fatigans* vector mosquitoes found to be breeding on Chagos Archipelago, Indian Ocean
- Wuchereria bancrofti
Lysenko, A. J.; Pavlova, E. A.; and Glazunova, Z. I., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (5-6), 1976, 536 [Letter]
Wuchereria bancrofti in man who had lived only in Dominican Republic until travel to Moscow
- Wuchereria bancrofti
Maasch, H. J., 1973, *Ztschr. Tropenmed. u. Parasitol.*, v. 24 (4), 419-434
Wuchereria bancrofti, human, quantitative aspects of vector transmission in urban, suburban, and rural areas of Liberia
Anopheles gambiae (nat. and exper.): Liberia
A. melas (nat. and exper.): Liberia
A. funestus: Liberia
A. hancocki: Liberia
Culex pipiens fatigans (exper.)
- [Wuchereria bancrofti] bancroftian filariasis
Mahfudin, H.; et al., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 173-178
human bancroftian filariasis, statistics of blood and clinical survey of Kepu District, relationship of habitat to infection rate: Central Jakarta, Indonesia
- Wuchereria bancrofti
de Meillon, B., 1977, *Proc. 15. Internat. Cong. Entom.* (Washington, D.C., Aug. 19-27, 1976), 498-504
Wuchereria bancrofti, changing patterns of transmission, vector control, importance of new diagnostic techniques, review
- Wuchereria bancrofti
Miller, M. J.; Ratard, R. C.; and McNeeley, D. F., 1976, *J. Parasitol.*, v. 62 (5), 845-847
Wuchereria bancrofti, human, nocturnal microfilarial periodicity; presence of *Mansonia ozzardi* also reported: Haiti
- Wuchereria bancrofti
Moreau, J. P.; Barbier, D.; and Radanielina, R., 1975, *Nouv. Presse Med.*, v. 4 (15), 1136 [Letter]
Wuchereria bancrofti in humans, preliminary clinical trials using 1-tetramisole, good results: Madagascar


- Wuchereria bancrofti*
Moreau, J. P.; Radanielina, R.; and Barbier, P., 1975, *Medecine Trop.*, v. 35 (6), 451-455
levamisole in treatment of *Wuchereria bancrofti*, possible use in mass treatment
- Wuchereria bancrofti*
Mougey, Y.; and Bain, O., 1976, *Ann. Parasitol.*, v. 51 (1), 95-110
5 filaria-vector sets, stochastic models for assessing relation between numbers of microfilariae ingested and numbers of microfilariae passing into hemocoel, limitation phenomenon
- Wuchereria bancrofti*
Muller, R. L.; and Denham, D. A., 1974, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 68 (1), 8-9 [Demonstration]
filarial spp. larvae, field technique for recovery and preservation of infective larvae from their insect vectors, application to studies of transmission dynamics
- Wuchereria bancrofti*
Nelson, G. S.; and Davies, J. B., 1976, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (1), 16-17 [Demonstration]
Mansonella ozzardi, humans, epidemiologic survey, concentration of microfilariae in superficial capillaries, mixed infections with *Wuchereria bancrofti* differentiated using stained filters: Trinidad
- Wuchereri[a] bancrofti*
Niel, G.; et al., 1972, *Medecine et Malad. Infect.*, v. 2 (5), 193-202
filariasis, human, diagnosis by double-diffusion and immunoelectrophoresis, examination of possible use of *Setaria labiatopapillosa* as antigen, comparison with *Dipetalone-ma vitae* and *Ascaris suum* as antigens
- Wuchereria bancrofti*
Obiamiwe, B. A., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (4), 487-490
sub-periodic *Brugia malayi*, periodic *Wuchereria bancrofti*, influence of the sb gene in *Culex pipiens* vectors on susceptibility to parasite development
- Wuchereria bancrofti*, ill.
Omar, M. S., 1977, *Tropenmed. u. Parasitol.*, v. 28 (1), 100-108
Wuchereria bancrofti, *Brugia malayi*, *B. pahangi*, *Dirofilaria immitis*, distribution of acid phosphatase activity in larval stages in the mosquito, presence or absence of enzymic activity in the excretory cell complex and amphids of developing larvae useful as adjunctive diagnostic method
- Wuchereria bancrofti*
Omori, N.; et al., 1967, *Nettai Igaku (Trop. Med.)*, v. 9 (2), 97-110
Wuchereria bancrofti, mosquito vector control, usefulness of larvicide compared with insecticide affecting adult mosquitoes
- Wuchereria bancrofti*
Omori, N.; and Wada, Y., 1968, *Nettai Igaku (Trop. Med.)*, v. 10 (3), 154-160
Wuchereria bancrofti in humans, factors affecting transmission by mosquitoes
- Wuchereria bancrofti*
Otsuru, M., 1974, *Internat. Med. Found. Japan. Reporting series* (4), 49-64
human nematode infections, extensive review on epidemiology, treatment and control measures: Japan
- Wuchereria bancrofti*
Ottesen, E. A.; Weller, P. F.; and Heck, L., 1977, *Immunology*, v. 33 (3), 413-421
Wuchereria bancrofti, human, antigen-specific cellular immune unresponsiveness, unchanged 2 weeks after diethylcarbamazine treatment, this immunologic deficit may be of fundamental importance in pathogenesis of filarial disease: Mauke, Cook Islands
- Wuchereria bancrofti*
Partono, F.; and Idris, K. N., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (2), 158-164
factors that influence loss of microfilariae from stained thick blood films, results of laboratory experiments
- Wuchereria bancrofti*
Partono, F.; and Sri Oemijati, 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 516-518
Wuchereria bancrofti, susceptibility of *Culex pipiens fatigans* to infection, laboratory studies, genetic studies with three generations of mosquitoes, evidence of modifier genes
- Wuchereria bancrofti*
Picq, J. J.; and Roux, J., 1973, *Medecine Trop.*, v. 33 (5), 451-461
Onchocerca volvulus, human, microfilaruria in relation to age and sex of host, other filarial diseases, geographic region, cutaneous microfilarial densities, albuminuria during suramin treatment, eggs of *Schistosoma haematobium* in urine, and diethylcarbamazine chemotherapy
- [*Wuchereria*] *bancrofti*
Pinon, J. M.; and Gentilini, M., 1973, *Nouv. Presse Med.*, v. 2 (19), 1283-1287
human filariasis, application of cellular immunologic tests (rosette formation, macrophage migration) in diagnosis and comparison with serologic tests (fluorescent antibody, passive hemagglutination, gel diffusion)
- Wuchereria bancrofti*
Ponnampalam, J. T., 1971, *Med. J. Malaya*, v. 26 (1), 62-64
review of human chronic endemic filariasis in West Malaysia, case reports of varying presenting symptoms

- Wuchereria bancrofti*, *illus.*
Pradhan, S.; et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (1), 199-200
Wuchereria bancrofti, human, microfilariae detected in bone marrow smears in seven cases of anemia (probably incidental and unrelated) in persons with asymptomatic filariasis
- Wuchereria bancrofti*
Raccurt, C.; and Hodges, W., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (5), 452-453 [Letter]
Wuchereria bancrofti, epidemiologic and vector survey in Haiti
- Wuchereria bancrofti*
Rajagopalan, P. K.; Kazmi, S. J.; and Mani, T. R., 1977, *Indian J. Med. Research*, v. 66 (2), 200-215
Wuchereria bancrofti, statistical epidemiologic survey on microfilaremia (seasonal fluctuation of infection in *Culex pipiens fatigans* vectors, localization in vectors, human infection by age and sex, microfilarial periodicity, percentage of night biting, experimental laboratory infections): Pondicherry, India
- Wuchereria bancrofti*
Rakai, I. M.; et al., 1974, *J. Med. Entom.*, v. 11 (5), 588-594
diel biting times for various *Aedes* and *Culex* spp., vector potential of *A. polynesiensis* and *A. pseudoscutellaris* for *Wuchereria bancrofti* transmission: Fiji
- Wuchereria bancrofti*
Ramachandran, C. P.; et al., 1970, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 1 (4), 505-515
epidemiologic and entomologic survey, no vectors implicated: Trengganu State, West Malaysia
- Wuchereria bancrofti*
Ramachandran, C. P.; Sandosham, A. A.; and Sivanandam, S., 1966, *Med. J. Malaya*, v. 20 (4), 333
partial larval development of *Wuchereria bancrofti* in exper. cats
- W[uchereria] *bancrofti*
Ramakrishnan, K.; and Aziz bin Ahmad, A., 1974, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 5 (1), 149 [Demonstration]
W[uchereria] *bancrofti*, *Pelecitus* sp., membrane feeding technique for infecting mosquitoes with filarial parasites
- Wuchereria bancrofti*
Richet, P., 1973, *Medecine Afrique Noire*, v. 20 (11), 899-920
guidelines for mass therapy in human filarial infections, drugs in current use, review
- Wuchereria bancrofti*
Ridley, D. S.; and Hedge, E. C., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (6), 522-525
microfilariae of various spp., immunofluorescent reactions involving sheath, cuticle, and cytoplasm, relevance to immuno-evasive mechanisms: (1) microfilariae failed to adsorb non-specific immunoglobulins in contrast to other helminth larvae and non-blood protozoa; (2) sheath of *Wuchereria bancrofti* and *Loa loa* adsorbed specific A and B blood group antigens; (3) low titer reaction between microfilarial cytoplasm (*L. loa* and *W. bancrofti*) and host serum
- Wuchereria bancrofti*
Rodhain, F.; and Dodin, A., 1971, *Medecine et Malad. Infect.*, v. 1 (4), 185-188
Wuchereria bancrofti, *Loa loa*, variations in human antistreptolysin O titers before and after treatment for filariasis, possible antigenic immune reaction between filariae and *Streptococcus*
- Wuchereria bancrofti*
Rodhain, F.; and Rodhain-Rebourg, F., 1974, *Medecine et Malad. Infect.*, v. 4 (3), 133-138
Wuchereria bancrofti, review of geographic distribution surveys of human lymphatic filariasis in the African continent south of the Sahara
- Wuchereria bancrofti*
Rodhain, F.; and Rodhain-Rebourg, F., 1974, *Medecine et Malad. Infect.*, v. 4 (12), 621-630
geographic distribution of human filariasis in Central America and Mediterranean areas, review of literature
- Wuchereria bancrofti*
Rodhain, F.; and Rodhain-Rebourg, F., 1976, *Medecine et Malad. Infect.*, v. 6 (3), 108-114
Wuchereria bancrofti, *Brugia malayi*, review of geographic distribution of human filariasis on the Asian continent
- Wuchereria bancrofti*
Rosen, L., et al., 1976, *Am. J. Trop. Med. and Hyg.*, v. 25 (6), 906-913
unsuccessful field trial of competitive displacement of *Aedes polynesiensis* (principal vector of nonperiodic filariasis caused by *Wuchereria bancrofti*) by *Aedes albopictus* (refractory to development of human filariae): Taiaro, remote Pacific coral atoll
- Wuchereria bancrofti*
Salfield, S., 1975, *Med. J. Australia*, v. 1 (9), 264-267
clinical report of 19 cases of human arthritis attributed to filarial *Wuchereria bancrofti*, good response to treatment with diethylcarbamazine: Papua New Guinea
- Wuchereria bancrofti*
Sandosham, A. A.; and Sivanandam, S., 1968, *Med. J. Malaya*, v. 22 (3), 236
adult *Wuchereria bancrofti* removed from vesicle on human scrotum: Malaysia

- Wuchereria bancrofti
Sant, M. V.; Gatlwar, W. N.; and Menon, T.U.K., 1974, Progr. Drug Research, v. 18, 269-275
Wuchereria bancrofti, comparative epidemiologic survey of 4 Indian villages with varying prevalences of infection; evidence of serum iodine insufficiency in infected persons living in sea coast areas normally high in iodine suggests that bancroftian filarial worms do utilize host serum iodine for their metabolism
- Wuchereria bancrofti
Sasa, M., 1974, Internat. Med. Found. Japan. Reporting series (4), 3-48
human filariasis in the Americas, extensive review, epidemiology, geographic distribution, mosquito vectors, control measures, literature review
- Wuchereria bancrofti
Sasa, M.; et al., 1976, Southeast Asian J. Trop. Med. and Pub. Health, v. 7 (3), 370-376
Brugia malayi, Wuchereria bancrofti, field study to determine pattern of microfilarial periodicity conducted in locality where Brugia malayi was first described, both parasites of nocturnal periodic form: Bireuen, Indonesia
- Wuchereria bancrofti
Sasa, M.; and Tanaka, H., 1972, Southeast Asian J. Trop. Med. and Pub. Health, v. 3 (4), 518-536
Wuchereria bancrofti, Brugia malayi, humans, technique for statistical analysis of microfilarial periodicity survey data
- Wuchereria bancrofti
Scheiber, P.; Braun-Munzinger, R. A.; and Southgate, B. A., 1976, Tropenmed. u. Parasitol., v. 27 (2), 224-228
epidemiologic survey in endemic area of nocturnally periodic Wuchereria bancrofti to compare use of conventional blood films and the membrane filtration concentration technique in detecting infections: Togo
- Wuchereria bancrofti
Singh, S. N.; Rao, V. G.; and Rao, B. K., 1976, J. Helminth., v. 50 (2), 96-97
Wuchereria bancrofti, first report of Anopheles fluviatilis as exper. vector host, low level of infection in A. stephensi (exper.)
- [Wuchereria] bancrofti
Southgate, B. A., 1974, Tr. Roy. Soc. Trop. Med. and Hyg., v. 68 (3), 177-186
human bancroftian filariasis, simultaneous trials using 4 known field techniques to diagnose microfilaremia in order to obtain comparative epidemiologic profiles: Fiji
- [Wuchereria] bancrofti
Sroczyński, J., 1977, Polski Tygod. Lekar., v. 32 (16), 589-591
Necator americanus, [Wuchereria] bancrofti, Schistosoma mansoni, studies on hospitalized Africans to assess variations in blood picture during infections showed anemia in hookworm to be mainly iron deficiency while schistosomiasis caused protein deficiency, eosinophilia of peripheral origin rather than correlated with changes in bone marrow
- Wuchereria bancrofti
Stuerchler, D.; and Degremont, A., 1976, Schweiz. Med. Wchnschr., v. 106 (20), 682-688
extensive diagnostic and clinical review of filarial parasites frequently encountered by travelers to endemic tropical areas: Switzerland
- [Wuchereria] bancrofti
Subra, R.; and Hebrara, G., 1975, Tropenmed. u. Parasitol., v. 26 (1), 48-59
[Wuchereria] bancrofti, ecology of vector Culex pipiens fatigans larvae in an area of high filariasis endemicity: Mayotte, Archipel des Comores
- Wuchereria bancrofti
Sucharit, S.; et al., 1975, Southeast Asian J. Trop. Med. and Pub. Health, v. 6 (4), 549-554
Wuchereria bancrofti, Brugia spp., attempted differentiation using morphological characteristics
- Wuchereria bancrofti
Sullivan, T. J. III; Carnahan, V. A.; and Cutting, R. T., 1970, Mil. Med., v. 135 (9), 797-798
Wuchereria bancrofti, soldier returned from Vietnam, filariasis without microfilaremia, diagnostic difficulties, need for awareness in returning military personnel
- Wuchereria bancrofti
Suswillo, R. R.; et al., 1977, J. Helminth., v. 51 (2), 132-134
slight development in Meriones unguiculatus (exper.) (abdominal cavity, heart, lungs, para-aortic lymphatics)
- Wuchereria bancrofti
Suzuki, T.; and Sone, F., 1975, Nettare Igaku (Trop. Med.), v. 16 (3), 147-156
Wuchereria bancrofti, survey of Aedes polynesiensis and A. samoanus vector mosquitoes pre- and post- mass diethylcarbamazine therapy to evaluate effectiveness of disease control efforts: Western Samoa
- Wuchereria bancrofti
Takahashi, J.; and Sato, K., 1976, Japan. J. Exper. Med., v. 46 (1), 7-13
fractionation and purification of Dirofilaria immitis antigens by column chromatography and disc electrophoresis, evaluation for use in diagnosis of human Wuchereria bancrofti by hemagglutination test

- Wuchereria bancrofti*, *illus.*
Terwedow, H. A., jr.; and Huff, R. L., 1976, *J. Parasitol.*, v. 62 (1), 172-174
Wuchereria bancrofti microfilariae, localization and pattern of acid phosphatase activity, possible taxonomic tool in differentiating microfilariae of different species but strain and technique variation must be taken into account
- Wuchereria bancrofti*
Thomas, V., 1974, *Med. J. Malaysia*, v. 29 (1), 73-77
feeding experiments with *Culex pipiens fatigans* to determine strain resistance and infectivity to rural strain of *Wuchereria bancrofti*: Malaysia
- Wuchereria bancrofti*
Thomas, V.; and Ramachandran, C. P., 1970, *Med. J. Malaya*, v. 24 (3), 196-199
laboratory trials with *Culex pipiens fatigans* to replace indigenous parasite-susceptible strains with parasite-resistant strains as possible biological control measure against *Wuchereria bancrofti*
- Wuchereria bancrofti*
Uemura, S., 1967, *Nettai Igaku (Trop. Med.)*, v. 9 (1), 24-38
Wuchereria bancrofti, *Anopheles sinensis* as potential vector, laboratory studies: Okinawa main island
- Wuchereria bancrofti*
Webber, R. H., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (3), 430-434
Wuchereria bancrofti, area survey for possible disease control as additional result of Malaria Eradication Programme since both infections transmitted by same mosquito vector: Choiseul and Shortland of Western Solomon Islands
- Wuchereria bancrofti*
Webber, R. H., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 70 (5-6), 1976, 537-538 [Letter]
human microfilaria, comparison of use of counting chamber and measured blood films for epidemiologic estimations, survey of *Wuchereria bancrofti*-endemic area in the Solomon Islands
- Wuchereria bancrofti*
Webber, R. H., 1977, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 71 (5), 396-400
control measures carried out by the Malaria Eradication Programme against *Anopheles* vectors of human malaria also resulted in the natural decline in the prevalence of human *Wuchereria bancrofti* infections since both diseases are transmitted by the same vector in the Solomon Islands
- Wuchereria bancrofti*
Weinstock, H.; et al., 1977, *Am. J. Trop. Med. and Hyg.*, v. 26 (6, part 1), 1148-1152
Wuchereria bancrofti, epidemiological survey, microfilaremia rates, distribution by age, sex, and race, public health importance, larvae found in *Culex pipiens fatigans*: Puerto Limon, Costa Rica
- Wuchereria bancrofti*
Wenceslao, J. M.; Oban, E.; and Cabrera, B. D., 1972, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 3 (4), 552-561
epidemiologic survey established widespread endemicity of human filariasis in Eastern Samar province, survey for vector mosquitoes *Aedes poecilus*
Culex fatigans
all from Republic of the Philippines
- Wuchereria bancrofti*
Wijers, D. J. B., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (3), 313-331
bancroftian filariasis, humans, extensive prevalence survey of adult males for microfilaremia and pathologic evidence of infection: Coast Province, Kenya
- Wuchereria bancrofti*
Wijers, D. J. B., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (4), 451-463
human bancroftian filariasis, dynamics of disease distribution and transmission as related to climate and vector breeding sites; microfilaremia rates as related to evidence of hydroceles and elephantiasis: Kenya
- Wuchereria bancrofti*
Wijers, D. J. B.; and Kiilu, G., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (3), 347-359
Wuchereria bancrofti, humans, extensive entomological survey comparing 2 geographically different sites for prevalence of infected mosquito vectors, correlations with seasons and socio-economic status of natives *Culex pipiens fatigans*
Anopheles gambiae
A. funestus
all from Coastal Province, Kenya
- [*Wuchereria*] *bancrofti*
Wijers, D. J. B.; and Kinyanjui, H., 1977, *Ann. Trop. Med. and Parasitol.*, v. 71 (3), 333-345
bancroftian filariasis, extensive prevalence survey comparing natives of a small sea shore village and those from a sparsely populated rural area for evidence of microfilaremia and visible pathology: Coast Province, Kenya
- Wuchereria bancrofti*
Woodruff, A. W., 1971, *Brit. J. Clin. Pract.*, v. 25 (12), 529-535
achievements in tropical medicine during past 25 years, control, prophylaxis, treatment
- Wuchereria bancrofti*
Yong, W. K., 1973, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 67 (3), 338-344
Wuchereria bancrofti, human, immunodiagnosis by indirect fluorescent antibody test using actual causal agent as antigen (micro-fragments of microfilariae and/or infective stage larvae)

- Wuchereria bancrofti*
Young, A. E.; and Kinmonth, J. B., 1976, Proc. Roy. Soc. Med., v. 69 (9), 708-709
Wuchereria bancrofti, case report of human chronic filariasis with scrotal edema, enlarged megalymphatics and observation of live adult worms in lymphatics of spermatic cord: London (Indian native)
- Wuchereria bancrofti*
Zaman, V.; and Lal, M., 1973, Tr. Roy. Soc. Trop. Med. and Hyg., v. 67 (4), 610 [Letter]
Wuchereria bancrofti, human, treatment with levamisole, severe reaction in one of two patients probably similar to pattern seen with diethylcarbamazine
- Wuchereria bancrofti*
Zerner, J.; and Barber, H. R. K., 1977, N. York State J. Med., v. 77 (3), 414-417
Wuchereria bancrofti in young Oriental woman, development of chyluria immediately post-partum with microfilaria discovered in blood and urine, clinical case report: New York City (immigrated 6 months earlier from Hong Kong)
- Wuchereria bancrofti*
Zielke, E., 1975, Tropenmed. u. Parasitol., v. 26 (3), 345-347
Wuchereria bancrofti, migration patterns of third stage larvae in *Anopheles gambiae* vectors
- Wuchereria bancrofti*
Zielke, E., 1976, Tropenmed. u. Parasitol., v. 27 (2), 160-164
feeding experiments with *Anopheles* and *Culex* vectors of human *Wuchereria bancrofti*, comparison of infection rates and loss of worm burden when feeding on blood chamber, guinea pig or sugar water solution
- Wuchereria bancrofti*
Zielke, E.; and Kuhl, F., 1977, Tropenmed. u. Parasitol., v. 28 (1), 68-70
Wuchereria bancrofti, variations in susceptibility of *Culex pipiens fatigans* vector mosquitoes to infection based on genetic factors
- Wuchereria bancrofti pacifica*
Gentilini, M.; Pinon, J. M.; and Danis, M., 1973, Medecine et Malad. Infect., v. 3 (8-9), 351-353
diagnostic review of human filariasis
- Wuchereria bancrofti* var. *pacifica*
Outin-Fabre, D.; Moreau, J. P.; and Stanghellini, A., 1972, Medecine Afrique Noire, v. 19 (2), 89-92
Wuchereria bancrofti, filariasis in endemic area of Polynesia, review of campaign for mass control
- Xericola Durette-Desset*, 1974
Durette-Desset, M. C., 1976, Bull. Mus. National Hist. Nat., Paris, 3. s. (388), Zool. (270), 711-720
Brevistriatinae
key; evolution of morphological characters, distribution of species among hosts and geographical regions, good correlation
- Xericola Durette-Desset*, 1974
Durette-Desset, M. C.; and Chabaud, A. G., 1977, Ann. Parasitol., v. 52 (5), 539-558
Heligmonellidae, Brevistriatinae
- Yorkeispirura* subgen. of *Oxyspirura*
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Oxyspirura* subgen. of *Oxyspirura*
- Yseria* Gedoelst, 1919
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 29-58
as syn. of *Streptocara* Railliet, Henry & Sisoff, 1912
- ZyLANEMA* [sic] Yeh, 1960
Agrawal, N., 1976, Indian J. Zoot., v. 15 (1), 1974, 11-13
valid genus
- Zeylanema* Yeh, 1960
Chabaud, A. G., 1975, CIH Keys Nematode Parasites Vertebrates (Anderson, Chabaud, and Willmott) (3), 1-27
as syn. of *Camallanus* Railliet & Henry, 1915
- Zeylanema barragi* n. sp., illus.
Zaidi, D. A.; and Khan, D., 1975, Pakistan J. Zool., v. 7 (1), 51-73
Mastacembelus armatus (intestine): Taunsa Barrage, Pakistan
- Zeylanema mastacembeli* Sahay et Sinha, 1966
Sinha, A.; and Sahay, U., 1971, Indian J. Animal Research, v. 5 (2), 67-72
valid species
Syn.: *Camallanus mastacembeli* Agrawal, 1967
- ZyLANEMA* [sic] *varanasiensis* n. sp., illus.
Agrawal, N., 1976, Indian J. Zoot., v. 15 (1), 1974, 11-13
Rita rita (intestine): river Ganges, Varanasi, India
- ZyLANEMA*. See *Zeylanema*.



- Acanthocephala**
 Gabrisch, K., 1976, Prakt. Tierarzt, v. 57, Sondernummer, 37-40
 parasites of reptiles, diagnosis, treatment, brief review
- Acanthocephala**
 Lindquist, W. D., 1970, Dis. Swine (Dunne), 3. ed., 708-744
 swine, pathology, diagnosis, control, text-book
- Acanthocephala**
 Lindquist, W. D., 1975, Dis. Swine (Dunne), 4. ed., 780-815
 helminths of swine, emphasis on nematodes, morphology, pathology, life cycle, diagnosis, treatment and control, review
- Acanthocephala**
 Ryzhikov, K. M., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 124-135
 helminths of birds in Russia, number of species in each class of helminths, comparison with numbers worldwide, review of literature
- Acanthocephal[a]**
 Schuetze, H. R., 1974, Prakt. Tierarzt, v. 55 (8), 429-432
 helminths of pet birds, diagnosis of eggs in fecal examination
- Acanthocephala**
 Weinmann, C. J., 1970, Immun. Parasitic Animals (Jackson, Herman and Singer), v. 2, 1021-1059
 cestodes, immunology, review with brief summary on acanthocephalans
- Acanthocephala sp.**
 Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
 Aplodinotus grunniens (intestine): Eagle Mountain Lake, Texas
- Acanthocephalus sp.**
 Combs, D. L.; Harley, J. P.; and Williams, J. C., 1977, Tr. Kentucky Acad. Sc., v. 38 (3-4), 128-131
 Minytrema melanops (gut): Kentucky River
 Moxostoma erythrurum (gut): Kentucky River
- Acanthocephalus anguillae (Mueller, 1780)**
 Dabrowska, Z., 1970, Acta Parasitol. Polon., v. 17 (20-38), 189-193
 Cyprinus carpio
 Tinca tinca
 Barbus barbus
 Abramis brama x Blicca bjoerkna
 Leuciscus idus
 Esox lucius
 Anguilla anguilla
 Lota lota
 Perca fluviatilis
 (intestine of all): all from Vistula River near Warsaw
- Acanthocephalus anguillae (Mueller, 1780)**
 Luehe, 1911
 Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 195-201
 Lota l. lota (stomach)
 Abramis brama
 Blicca bjoerkna
 Leuciscus idus
 L. cephalus
 Rutilus rutilus
 Anguilla anguilla
 Perca fluviatilis
 Silurus glanis
 all from Poland
- Acanthocephalus anguillae (Mueller, 1780)**
 Luehe, 1911
 Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
 Silurus glanis (stomach): river Biebrza basin, Poland
- Acanthocephalus anguillae (Mueller, 1780)**
 Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmit. Lab., v. 16, 87-110
 L[euiscus] cephalus (intestine): Balkan Mountain river
- Acanthocephalus anguillae**
 Øien, K., 1976, Norwegian J. Zool., v. 24 (4), 466-467 [Abstract]
 Rutilus rutilus
 Leuciscus idus
 (intestine of all): all from Lake Oyeren
- Acanthocephalus anguillae**
 Perłowska, R., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 27-32
 Esox lucius
 Tinca tinca
 Abramis brama
 Leuciscus idus
 Rutilus rutilus
 all from Zegrzynski Reservoir
- Acanthocephalus anguillae (Mueller, 1780)**
 Puciłowska, A., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 33-46
 helminths of fishes, dynamics of infection following formation of artificial body of water, seasonal distribution, brief description
 Perca fluviatilis: Zegrzynski Reservoir
- Acanthocephalus anguillae (Mueller, 1780)**
 Willemse, J. J., 1968, Bull. Zool. Mus. Univ. Amsterdam, v. 1 (8), 83-87
 Anguilla anguilla: IJsselmeer
- Acanthocephalus clavula (Dujardin, 1845)**
 Andrews, C.; and Rojanapaibul, A., 1976, Parasitology, v. 73 (2), xi-xii [Abstract]
 Acanthocephalus clavula, seasonal changes in incidence, intensity, and maturation, effect of host species, temperature thought to be a factor
 Perca fluviatilis
 Coregonus lavaratus
 Rutilus rutilus
 Cottus gobio
 Anguilla anguilla
 all from Llyn Tegid, North Wales

- Acanthocephalus clavula* (Dujardin, 1845)
Rojanapaibul, A., 1976, *Parasitology*, v. 73 (2), xi [Abstract]
Acanthocephalus clavula, life cycle and development
Asellus meridianus (exper.)
bullhead (exper.)
- Acanthocephalus dirus* (Van Cleave, 1931)
Amin, O. M., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 81-88
distribution, structural observations, effects of host size (age) on worm burden and site of infection
Catostomus commersoni (small and large intestine): southeastern Wisconsin
- Acanthocephalus falcatus* (Froelich, 1789)
Hristovski, N. D., 1975, *Acta Parasitol. Yugoslavica*, v. 6 (1), 3-5
Rana graeca: Bitola district, Macedonia, Yugoslavia
- Acanthocephalus galaxii* n. sp., illus.
Hine, P. M., 1977, *J. Roy. Soc. N. Zealand*, v. 7 (1), 51-57
Acanthocephalus galaxii n. sp., larval development
Paracalliope fluviatilis (exper.)
Galaxias maculatus (intestine): Waimeha Stream, Waikanae, North Island, New Zealand
G. argenteus (intestine): Waimeha Stream, Waikanae, North Island, New Zealand
Retropinna retropinna (intestine): Waimeha Stream, Waikanae, North Island, New Zealand
Gobiomorphus cotidianus (intestine): Waimeha Stream, Waikanae, North Island, New Zealand
Anguilla australis (intestine): Waimeha Stream, Waikanae, North Island, New Zealand; Lake Ellesmere, south of Christchurch, New Zealand
A. dieffenbachii (intestine): Waimeha Stream, Waikanae, North Island, New Zealand; Makara Stream near Wellington, New Zealand
- Acanthocephalus jacksoni* Bullock, 1962
Muzzall, P. M.; and Rabalais, F. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (1), 31-34
Acanthocephalus jacksoni from fish, seasonal periodicity, effect of water temperature, definitive host spawning period, vegetation, and intermediate host
Lirceus lineatus (nat. and exper.)
Etheostoma blennoides
Lepomis cyanellus
L. macrochirus
Ambloplites rupestris
Notropis spilopterus
N. hudsonius
N. umbratilis
N. crysocephalus
Carassius auratus
Cyprinus carpio
Campostoma anomalum
Hypentelium nigricans
Lepomis gibbosus
Catostomus commersoni
Semotilus atromaculatus
all from Jackson Cutoff, Wood County, Ohio
- Acanthocephalus jacksoni* Bullock, 1962
Muzzall, P. M.; and Rabalais, F. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (1), 35-38
Acanthocephalus jacksoni cystacanths, sex, position, and orientation in *Lirceus lineatus*, host pigmentation, sex, size, and gut content, possible sterility in host females; precocious development and marked sexual dimorphism in cystacanths: Jackson Cutoff
- Acanthocephalus jacksoni* Bullock, 1962
Muzzall, P. M.; and Rabalais, F. C., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 116-118
Acanthocephalus jacksoni-infected *Lirceus lineatus*, altered behavioral responses and increased conspicuousness of parasitized isopods increases likelihood of transmission of cystacanth to definitive host
- Acanthocephalus jacksoni* Bullock, 1962
White, G. E., 1974, *Tr. Am. Micr. Soc.*, v. 93 (2), Apr., 280-282
Catostomus commersoni: Kentucky River drainage system
- Acanthocephalus jacksoni*
White, G. E.; and Harley, J. P., 1973, *Tr. Kentucky Acad. Sc.*, v. 34 (3, 4), 53-54
Catostomus commersoni: Lake Wilgreen, Madison County, Kentucky
- Acanthocephalus lucii* (Mueller, 1776)
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 189-193
Cyprinus carpio
Barbus barbus
Leuciscus idus
Esox lucius
Lota lota
Perca fluviatilis
(intestine of all): all from Vistula River near Warsaw
- Acanthocephalus lucii* (Mueller, 1776) Luehe, 1911
Ejsymont, L., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 195-201
Lota l. lota (pyloric appendices)
Abramis brama
Carassius carassius
Leuciscus idus
Rutilus rutilus
Scardinius erythrophthalmus
Tinca tinca
Esox lucius
Acerina cernua
Perca fluviatilis
Anguilla anguilla
Silurus glanis
all from Poland
- Acanthocephalus lucii* (Mueller, 1776) Luehe, 1911
Ejsymont, L., 1970, *Acta Parasitol. Polon.*, v. 17 (20-38), 203-216
Silurus glanis (intestines): river Biebrza basin, Poland

- Acanthocephalus lucii* (Mueller, 1787)
Kulakiv's'ka, O. P., 1976, Vestnik. Zool., Akad. Nauk Ukrain'sk. SSR, Inst. Zool. (4), 82-84
 Umbra crameri (intestine): Duna delta
- Acanthocephalus lucii* (Muller 1776)
Lee, R. L. G., 1977, Lond. Naturalist (1976) (56), 57-70
 Gymnocephalus cernua
 Perca fluviatilis
 (gut of all): all from Serpentine lake, Hyde Park and Kensington Gardens, central London
- Acanthocephalus lucii*
Perłowska, R., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 27-32
 Esox lucius
 Perca fluviatilis
 Tinca tinca
 all from Zegrzynski Reservoir
- Acanthocephalus lucii* (Mueller, 1776)
Pucilowska, A., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 33-46
 helminths of fishes, dynamics of infection following formation of artificial body of water, seasonal distribution, brief description
 Leuciscus idus: Zegrzynski Reservoir
- Acanthocephalus lucii* (Mueller, 1777)
Willemse, J. J., 1968, Bull. Zool. Mus. Univ. Amsterdam, v. 1 (8), 83-87
 Esox lucius: Vinkeveen
 Perca fluviatilis: Amsterdam (Geuzenveld); Amsterdam (Slotermeer); Velsen; IJsselmeer
- Acanthocephalus ranae* (Schrank, 1788)
Antsyshkina, L. M.; et al., 1976, Vestnik Zool., Akad. Nauk Ukrain'sk. SSR, Inst. Zool. (2), 82-84
 Rana ridibunda
 Bombina bombina
 Rana terrestris
 R. esculenta
 all from Samara river valley, Ukrainian SSR
- Acanthocephalus ranae* Luehe
Bozhkov, D., 1974, Izvest. Tsentral. Khelmint. Lab., v. 17, 25-31
 8 helminth species in *Rana ridibunda* fed to *Natrix natrix* or *N. tessellata*, found that *Diplodiscus subclavatus*, *Opisthioglyphe ranae*, *Cephalogonimus retusus*, and *Cosmocerca ornata* can pass alive from body of ingested frog to intestine of *Natrix natrix*, and *D. subclavatus* to *N. tessellata*
- Acanthocephalus ranae* (Schrank 1788) Luehe 1911
Hristovski, N. D.; and Lees, E., 1973, Acta Parasitol. Iugoslavica, v. 4 (2), 93-97
 Rana temporaria: Macedonia
- Acanthocephalus ranae* (Schrank, 1788), illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
 Rana ridibunda
 R. esculenta
 R. temporaria
 R. agilis
 all from Yugoslavia
- Acanthocephalus ranae* (Schrank, 1788)
Plasota, K., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 47-60
 helminths of frogs, comparison of aquatic and terrestrial hosts, relation of parasite fauna to environment, food supplies and food habits, host life cycle, temperature, rainfall, season, age and sex of host, competition between species of parasite, localization within host
 Rana esculenta
 R. terrestris
 all from Kampinos National Park, Poland
- Acanthocephalus ranae* (Schrank 1788, Luhe 1911), illus.
Rozman, M., 1971, Acta Parasitol. Iugoslavica, v. 2 (2), 67-77
 description
 synonymy
 Rana esculenta (Tanko crijevo): environs of Novi Sad, Yugoslavia
- Acanthocephalus serendibensis* Cruzs & Mills, 1970, illus.
Cruzs, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
 redescription
 Cnemaspis kandianus kandianus (small intestine): Haputale Estate (1418 m), Ceylon
 Ceratophora stoddarti (small intestine): Hakgala Strict Natural Reserve, Ceylon
- Acanthocephalus srilankensis* sp. nov., illus.
Cruzs, H.; and Ching, C. C., [1976], Ann. Parasitol., v. 50 (5), 1975, 531-537
 Rhacophorus cruciger eques
 R. microtypanum
 (small intestine of all): all from Ceylon (Hakgala Strict Natural Reserve; Horton Plains (2195 m))
- Acanthocephalus tahlequahensis* sp. n., illus.
Oetinger, D. F.; and Buckner, R. L., 1976, J. Parasitol., v. 62 (2), 237-241
 Etheostoma punctulatum (hindgut)
 E. spectabile (hindgut)
 Nocomis asper (hindgut)
 Notropis pilsbryi (hindgut)
 Micropterus dolomieu (mesentery)
 all from Black Fox Creek, northeast Tahlequah, Oklahoma
- Acanthogyrus* (*Acanthosentis*) *tilapiae* (Baylis, 1948)
Khalil, L. F.; and Thurston, J. P., 1973, Rev. Zool. et Botan. Africae, v. 87 (2), 209-248
 Tilapia nilotica: Lake George, Uganda
 Haplochromis squampinnus: Lake George, Uganda
 Tilapia esculenta: Lake Victoria, Uganda
 Haplochromis sp.: Lake Victoria, Uganda (intestine of all)
- Acanthosentis* sp., illus.
Anantaraman, S.; and Ravindranath, M. H., 1976, Ztschr. Parasitenk., v. 48 (3-4), 227-238
 Acanthosentis sp. (identified in footnote as *A. oligospinus*), egg envelopes of acanthor, layers, histochemistry, permeability, phase-contrast microscopy
 Mystus gulio (intestine)

- Acanthosentis acanthuri* Cable et Quick 1954, illus.
Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (5), 865-867
redescription
Acanthurus coeruleus: Speyside, Tobago, W.I.
A. chirurgus
- Acanthosentis oligospinus*
Anantaraman, S.; and Ravindranath, M. H., 1976, *Ztschr. Parasitenk.*, v. 48 (3-4), 227-238
Acanthosentis sp. (identified in footnote as *A. oligospinus*), egg envelopes of acanthor, layers, histochemistry, permeability, phase-contrast microscopy
Mystus gulio (intestine)
- Acanthosentis tilapiae*, illus.
Marchand, B.; and Mattei, X., 1976, *J. Ultrastructure Research*, v. 55 (3), 391-399
Acanthosentis tilapiae, spermatozoon, variation in number of central fibers in flagellum
- Andracantha* gen. n.
Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
Polymorphidae
tod: *A. gravida* (Alegret 1941) comb. n.
- Andracantha gravida* (Alegret 1941) comb. n. (tod), illus.
Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
redescription
[Syn.]: *Corynosoma gravida* Alegret 1941
Phalacrocorax auritus floridanus: Florida
Pelecanus occidentalis carolinensis: Florida; Louisiana
- Andracantha mergi* (Lundstrom, 1941) comb. n., illus.
Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
redescription
[Syn.]: *Corynosoma mergi* Lundstrom 1941
Mergus serrator (small intestine): Alaska
Gavia stellata (small intestine): Alaska
Aythya affinis (small intestine): New York
Nycticorax nycticorax (small intestine): New Hampshire
- Andracantha phalacrocoracis* (Yamaguti, 1939) comb. n., illus.
Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
redescription
[Syn.]: *Corynosoma phalacrocoracis* Yamaguti 1939
Phalacrocorax pelagicus (small and large intestine): Shikoku Islands, Japan; St. Lawrence Island, Nunivak Island, Alaska
- Apororhynchus* sp., illus.
Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 245-259
description
Motacilla flava (large intestine): Chukotsk
- Arhythmorhynchus* Luehe, 1911
Khokhlova, I. G., 1975, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 25, 195-203
revision
key to species, includes: *Arhythmorhynchus roseum* (Molin, 1858); *A. eroliae* (Yamaguti, 1939); *A. plicatus* (Linstow, 1883); *A. distinctus* Baer, 1956; *A. comptus* Van Cleave, Raush, 1950; *A. rubicundus* (Molin, 1858); *A. pumilirostris* Van Cleave, 1916; *A. jeffreyi* Schmidt, 1963; *A. invaginabilis* (Linstow, 1902); *A. frassoni* (Molin, 1858); *A. johnstoni* Golvan, 1960; *A. trichocephalus* (Kaiser, 1893); *A. limosae* Edmonds, 1971; *A. tringi* Gubanov, 1952; *A. uncinatus* (Kaiser, 1893); *A. brevis* Van Cleave, 1916; *A. capellae* (Yamaguti, 1935); *A. teres* Van Cleave, 1920; *A. tigrinum* Moghe, Das, 1953; *A. siluricola* Dollfus, 1929; *A. frontospinosus* (Tubangui, 1935)
- Arhythmorhynchus anser* Florescu, 1942
Khokhlova, I. G., 1975, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 25, 195-203
as syn. of *Arhythmorhynchus invaginabilis* (Linstow, 1902)
- Arhythmorhynchus comptus* Van Cleave et Rausch, 1950, illus.
Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 245-259
description
Calidris alpina
C. minuta
C. temmincki
C. melanotos
Eurynorhynchus pygmaeus
Limnodromus griseus
Phalaropus lobatus
P. fulicarius
(small intestine of all): all from Chukotsk
- Arhythmorhynchus frassoni* (Molin, 1858)
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, *Gel'mint. Zhivot. Tikhogo Okeana* (Skriabin), 105-124
Larus argentatus
L. canus
L. crassirostris
L. ridibundus
Sterna hirundo
(small intestine of all): all from coast of Sea of Okhotsk
- Arhythmorhynchus frassoni* (Molin, 1858) Luhe, 1911
Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 260-276
Terekia cinerea (small intestine): Siberia
- Arhythmorhynchus invaginabilis* (Linstow, 1902) Luhe, 1912
Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 260-276
Terekia cinerea (small intestine): Siberia
- Arhythmorhynchus invaginabilis* (Linstow, 1902)
Khokhlova, I. G., 1975, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 25, 195-203
synonymy, key

- Arhythmorhynchus longicollis* (Villot, 1875) of Golvan, 1956; Belopolskaia, 1959
Khokhlova, I. G., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 195-203
as syn. of *Arhythmorhynchus invaginabilis* (Linstow, 1902)
- Arhythmorhynchus macrourus* (Bremser, 1821) nom. nud.
Khokhlova, I. G., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 195-203
- Arhythmorhynchus plicatus*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Arhythmorhynchus sachalinensis* Krotov et Petrotchenko, 1958
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Arenaria interpres
Calidris alpina
C. minuta
Phalaropus fulicarius
Plectrophenax nivalis
(small intestine of all): all from Chukotsk
- Arhythmorhynchus sachalinensis* Krotov, Petrotchenko, 1958
Khokhlova, I. G., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 195-203
as syn. of *Arhythmorhynchus teres* Van Cleave, 1920
- Arhythmorhynchus teres* Van Cleave, 1920, illus.
Khokhlova, I. G., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 195-203
key
description, syn.: *Arhythmorhynchus sachalinensis* Krotov, Petrotchenko, 1958
Larus ridibundus: Kamchatka
- Bolbosoma balaenae*
Dailey, M. D.; and Perrin, W. F., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (2), 455-471
Stenella graffmani
S. cf. *S. longirostris*
all from eastern tropical Pacific
- Bolbosoma caenoforme* (Heitz, 1920)
Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 80-88
helminth distribution among age groups of *Pleurogrammus azonus*: Peter the Great Bay, Sea of Japan
- Bolbosoma caenoforme* Heitz, 1920
Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (1), 267-277
helminths, incidence and intensity of infection in young and adult *Oncorhynchus nerka*, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Bolbosoma nipponicum* Yamaguti, 1939
Popov, V. N., 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
Histiophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Bolbosoma vasculosum* (Rudolphi, 1819)
Bussieras, J.; and Baudin-Laurencin, F., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (4), 13a-19a
+tuna (appareil digestif): region nord de l'Atlantique tropical oriental
- Bolbosoma vasculosum*
Dailey, M. D.; and Perrin, W. F., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (2), 455-471
Stenella graffmani
S. cf. *S. longirostris*
all from eastern tropical Pacific
- Centrorhynchus* sp. A, illus.
Acholonu, A. D., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 106-116
description
Anolis cristatellus (body cavity): Cabo Rojo, Puerto Rico
- Centrorhynchus* sp. B, illus.
Acholonu, A. D., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 106-116
description
Anolis cristatellus (body cavity): Cabo Rojo, Puerto Rico
- Centrorhynchus* sp.
Kocan, A. A.; and Locke, L. N., 1974, J. Wildlife Dis., v. 10 (1), 8-10
Haliaeetus leucocephalus: Maine; Florida; Iowa, New Jersey
- Centrorhynchus aluconis* (Mueller, 1780) Luehe, 1911, illus.
Milka, R., 1976, Veterinaria, Sarajevo, v. 25 (3), 449-476
Rana ridibunda
R. esculenta
all from Yugoslavia
- Centrorhynchus conspectus*
Anderson, M. M.; and McDaniel, J. S., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
Blarina brevicauda: eastern North Carolina
- Centrorhynchus corvi*, illus.
Parshad, V. R.; and Guraya, S. S., 1977, Ann. Biol. Animale Biochem. Biophys., v. 17 (6), 953-959
Centrorhynchus corvi, lipids, correlative biochemical and histochemical studies
- Centrorhynchus corvi*, illus.
Parshad, V. R.; and Guraya, S. S., 1977, Parasitology, v. 74 (3), 243-253
Centrorhynchus corvi, ovarian balls, morphology and histochemistry

- Centrorhynchus falconis* (Das, 1950), illus. Rengaraju, V.; and Das, E. N., 1976, *Acta Histochem.*, v. 57 (2), 263-269
Centrorhynchus falconis, histochemistry
- Centrorhynchus golvani* n. sp. [nom. nud.] Anantaraman, S., 1963, *J. Marine Biol. Ass. India*, v. 5 (1), 137-139
Hieraetus pennatus: Madras Coast
- Centrorhynchus lancea* Vaidova, S. M., 1975, *Izvest. Akad. Nauk Azerbaidzhan. SSR*, s. *Biol. Nauk* (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Centrorhynchus lanceoides* Petrotchenko, 1949 Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 260-276
Charadrius apricarius
C. morinellus
C. hiaticola
(small intestine of all): all from Siberia
- Centrorhynchus migrans* n. sp., illus. Zuberi, H. B.; and Farooq, M., 1974, *Pakistan J. Zool.*, v. 6 (1-2), 147-150
Milvus migrans (intestine): Karachi, Pakistan
- Centrorhynchus milvus* Ward, 1956, illus. Marchand, B.; and Mattei, X., 1976, *Compt. Rend. Soc. Biol., Paris*, v. 170 (1), 237-240
Centrorhynchus milvus, spermatozoon, ultrastructure
- Corynosoma* sp., illus. Boero, J. J.; Led, J. E.; and Brandetti, E., 1972, *Analecta Vet.*, v. 4 (1), 17-34
Spheniscus magellanicus (intestino, estomago): Argentine Republic
- Corynosoma* sp. Courtney, C. H.; and Forrester, D. J., 1974, *Proc. Helminth. Soc. Washington*, v. 41 (1), 89-93
prevalence and intensity, age of host
Pelecanus occidentalis (small intestine): Florida
- Corynosoma* sp. George, R. R.; and Bolen, E. G., 1975, *J. Wildlife Dis.*, v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas
- Corynosoma constrictum* Van Cleave, 1918 Turner, B. C.; and Threlfall, W., 1975, *Proc. Helminth. Soc. Washington*, v. 42 (2), 157-169
parasites of *Anas crecca* and *A. discors*, incidence and intensity, age and sex of host
Anas crecca
A. discors
(small intestine of all): all from eastern Canada
- Corynosoma enhydridis* Hennessy, S. L.; and Morejohn, G. V., 1977, *Calif. Fish and Game*, v. 63 (4), 268-272
seasonal incidence
Enhydra lutris: off coastal California
- Corynosoma gravaida* Alegret 1941 Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
[as syn. of] *Andracantha gravaida* (Alegret 1941) comb. n.
- Corynosoma mergi* Lundstrom 1941 Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
[as syn. of] *Andracantha mergi* (Lundstrom, 1941) comb. n.
- Corynosoma peposacae* George, R. R.; and Bolen, E. G., 1975, *J. Wildlife Dis.*, v. 11 (1), 17-22
endoparasites of *Dendrocygna autumnalis*, prevalence higher in juveniles, pathology: Nueces County, southern Texas
- Corynosoma phalacrocoracis* Yamaguti, 1939 Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 245-259
Gavia stellata
G. immer
Phalacrocorax pelagicus
Somateria spectabilis
(small intestine of all): all from Chukotsk
- Corynosoma phalacrocoracis* Yamaguti 1939 Schmidt, G. D., 1975, *J. Parasitol.*, v. 61 (4), 615-620
[as syn. of] *Andracantha phalacrocoracis* (Yamaguti, 1939) comb. n.
- Corynosoma semerme* Forss Bonner, W. N., 1972, *Oceanogr. and Marine Biol. Ann. Rev.*, v. 10, 461-507
Halichoerus grypus
Phoca vitulina
(gut of all): all from European waters
- Corynosoma semerme* (Frossell, 1904) Luhe, 1905 Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 17, 245-259
Gavia stellata
G. immer
(small intestine of all): all from Chukotsk
- Corynosoma semerme* (Frossell, 1904) Luhe, 1905 Kozlov, D. P., 1969, *Trudy Gel'mint. Lab., Akad. Nauk SSSR*, v. 20, 71-78
Alopex lagopus: Pechora river basin
- Corynosoma semerme* (Frossell, 1904) Popov, V. N., 1976, *Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR* (145), year 19, (1), 49-53
age dynamics of infection
Histiophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island

- Corynosoma strumosum* (Rudolphi, 1802)
Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 80-88
helminth distribution among age groups of *Pleurogrammus azonus* (intestine): Peter the Great Bay, Sea of Japan
- Corynosoma strumosum* (Rudolphi, 1802)
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus crassirostris (small intestine): coast of Sea of Okhotsk (Ol'sk region)
L. argentatus: coast of Sea of Okhotsk
- Corynosoma strumosum* Rudolphi
Bonner, W. N., 1972, Oceanogr. and Marine Biol. Ann. Rev., v. 10, 461-507
Halichoerus grypus
Phoca vitulina
(gut of all): all from European waters
- Corynosoma strumosum* (Rudolphi, 1802)
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (intestine): Sakhalin Bay
- Corynosoma strumosum* (Rudolphi, 1802) Luhe, 1904
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Phalaropus lobatus
Larus hyperboreus
Gavia stellata
G. immer
Aythya marila
Clangula hyemalis
Somateria mollissima
S. fischeri
S. stelleri
S. spectabilis
(small intestine of all): all from Chukotsk
- Corynosoma strumosum* (Rudolphi, 1802)
Korotaeva, V. D., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 89-96
Icelus spiniger
Hemilepidotus gilberti
all from Sea of Japan
- Corynosoma strumosum* (Rudolphi, 1902)
Popov, V. N., 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
age dynamics of infection
Histiophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Corynosoma strumosum* (Rudolphi, 1802) Luhe, 1904
Shakhmatova, V. I., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 277-289
Mustela vison: Karelia
- Corynosoma strumosum*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Corynosoma validum* Van Cleave, 1953
Deliamure, S. L.; and Popov, V. N., 1975, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (142), year 18, (10), 7-10
Erignathus barbatus nauticus (intestine): Sakhalin Bay
- Corynosoma validum* Van Cleave, 1953
Popov, V. N., 1976, Biol. Nauk., Min. Vyssh. i Sredn. Spetsial. Obrazovan. SSSR (145), year 19, (1), 49-53
age dynamics of infection
Histiophoca fasciata (intestine): northern shore of Okhotsk Sea from Lisiansk peninsula to Iamsk island
- Disteganius* Lehman 1953, nom. nud.
Schmidt, G. D., 1977, J. Parasitol., v. 63 (1), 112-116
as syn. of *Mediorhynchus* Van Cleave 1916
- Dollfusentis chandleri*
Overstreet, R. M.; and Howse, H. D., 1977, Ann. N. York Acad. Sc., v. 298, 427-462
helminths and protozoans of estuarine fishes, incidence and intensity; possible relationships with water pollutants
Micropogon undulatus: estuaries of Mississippi
- Echinopardalis lamasi* Freitas et Costa
Schmidt, G. D., 1977, J. Parasitol., v. 63 (3), 508-510
as syn. of *Oncicola lamasi* (Freitas et Costa 1964) comb. n.
- Echinorhynchidae* [sp.]
Beacham, B. E.; and Haley, A. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 232-233
Morone americana (intestine): Hooper's Island, Chesapeake Bay
- Echinorhynchus* sp.
Vooren, C. M.; and Tracey, D., 1976, N. Zealand J. Marine and Freshwater Research, v. 10 (3), 499-509
incidence, intensity
Cheilodactylus macropterus (body cavity): New Zealand
- Echinorhynchus borealis* Linstow, 1901 emend.
Grabda-Kazubska et Ejsymont, 1969
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 195-201
Lota l. lota (pyloric appendices, stomach, intestine): Poland
- Echinorhynchus borealis* Linstow, 1901 emend.
Grabda-Kazubska et Ejsymont, 1969
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
Silurus glanis (stomach, middle and posterior portions of intestine): river Biebrza basin, Poland

- Echinorhynchus borealis* Linstow, 1901, *illus.*
Grabda-Kazubaska, B.; and Ejsymont, L., 1969,
Acta Parasitol. Polon., v. 17 (1-19), 65-87
synonymy, morphology, morphological variation
Silurus glanis: northern Poland; Ladoga
lake
Lota lota: northern Poland; Volga river;
Amur river
Thymallus thymallus: northern Poland
T. arcticus: Taimyr lake
Esox lucius: northern Poland
Perca fluviatilis: Yenisei river
Pallasea quadrispinosa: Pertozero lake
Coregonus ussuriensis: Amgun river
- Echinorhynchus boschadis* Schrank, 1788
Pavlov, A. V., 1966, *Trudy Gel'mint. Lab.*,
Akad. Nauk SSSR, v. 17, 104-127
as syn. of *Polymorphus minutus* (Goeze, 1782)
- Echinorhynchus cinctulus* Porta, 1905
Grabda-Kazubaska, B.; and Ejsymont, L., 1969,
Acta Parasitol. Polon., v. 17 (1-19), 65-87
as syn. of *Echinorhynchus borealis* Linstow,
1901
- Echinorhynchus clavula* Duj. *nec* Ham.
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*,
v. 17 (20-38), 189-193
Cyprinus carpio
Lota lota
(intestine of all): all from Vistula River
near Warsaw
- Echinorhynchus clavula* Dujardin, 1845 sensu
Luhe, 1911
Grabda-Kazubaska, B.; and Ejsymont, L., 1969,
Acta Parasitol. Polon., v. 17 (1-19), 65-87
as syn. of *Echinorhynchus borealis* Linstow,
1901
- Echinorhynchus clavula* Dujardin, 1845
Willemsse, J. J., 1968, *Bull. Zool. Mus. Univ.*
Amsterdam, v. 1 (8), 83-87
Lota lota: Vinkeveen
- Echinorhynchus gadi* Mueller, 1776
Baeva, O. M., 1968, *Gel'mint. Zhivot. Tikhogo*
Okeana (Skriabin), 80-88
helminth distribution among age groups of
Pleurogrammus azonus: Peter the Great Bay,
Sea of Japan
- Echinorhynchus gadi* Mueller, 1776
Korotaeva, V. D., 1968, *Gel'mint. Zhivot.*
Tikhogo Okeana (Skriabin), 89-96
Enophrys diceraus
Icelus spiniger
Myoxocephalus jaok
M. brandti
all from Sea of Japan
- Echinorhynchus gadi*
McVicar, A. H. 1977, *J. Helminth.*, v. 51 (1),
11-21
intestinal helminths of *Raja naevus*, inci-
dence, intensity, pattern of infection with
host age and sex, geographical differences
in composition of parasite burden
Raja naevus (stomach): off Aberdeen
- Echinorhynchus gadi*
Moeller, H., 1976, *J. Marine Biol. Ass. United*
Kingdom, v. 56 (3), 781-785
intestinal helminths, elimination from host
held in captivity, high rate of elimination
of helminths unattached or slightly at-
tached to host, lower elimination rate of
helminths attached to host
Gadus morhua
Zoarces viviparus
Myoxocephalus scorpius
Platichthys flesus
(intestine of all): all from Kiel Fjord
(western Baltic Sea)
- Echinorhynchus gadi*, *illus.*
Munson, D. A., 1974, *J. Wildlife Dis.*, v. 10
(3), 256-262
Liparis atlanticus (intestine): Rye, New
Hampshire
- Echinorhynchus gadi* (Mueller, 1776)
Pennell, D. A.; Becker, C. D.; and Scofield,
N. R., 1973, *Fish. Bull.*, National Oceanic
and Atmos. Admin., v. 71 (1), 267-277
helminths, incidence and intensity of
infection in young and adult *Oncorhynchus*
nerka, life cycle review: Kvichak River
system, Bristol Bay, Alaska
- Echinorhynchus globulosus*
Samuel, N.; Nickol, B. B.; and Mayes, M. A.,
1976, *Am. Midland Naturalist*, v. 96 (2), 391-
406
species inquirendum
- Echinorhynchus hexagrammi* Bajewa, 1965
Baeva, O. M., 1968, *Gel'mint. Zhivot. Tikhogo*
Okeana (Skriabin), 80-88
helminth distribution among age groups of
Pleurogrammus azonus (intestine): Peter the
Great Bay, Sea of Japan
- Echinorhynchus nudus* (Harada, 1938)
Mamaev, I. L., 1968, *Gel'mint. Zhivot. Tikhogo*
Okeana (Skriabin), 5-27
Thunnus thynnus
Euthynnus affinis
Auxis thazard
(intestine of all): all from South China Sea
- Echinorhynchus salmonis* Mueller, 1780
Dabrowska, Z., 1970, *Acta Parasitol. Polon.*,
v. 17 (20-38), 189-193
Vimba vimba
Esox lucius
Perca fluviatilis
(intestine of all): all from Vistula River
near Warsaw
- Echinorhynchus tanagrae* Rud. 1819
Schmidt, G. D.; and Kuntz, R. E., 1977, *J.*
Parasitol., v. 63 (3), 500-507
"unidentifiable"
- Echinorhynchus truttae* Schrank, 1788
Campbell, A. D., 1974, *Proc. Roy. Soc. Edinb.*,
sect. B, *Biol.*, v. 74, 347-364
Salmo trutta (intestine)
Perca fluviatilis
Esox lucius (intestinal wall)
Gasterosteus aculeatus
all from Loch Leven, Scotland

- Echinorhynchus truttae Schrank, 1788
Ejsymont, L., 1970, Acta Parasitol. Polon.,
v. 17 (20-38), 195-201
Lota l. lota
Salmo trutta
S. trutta m. fario
Thymallus thymallus
Esox lucius
Anguilla anguilla
all from Poland
- Echinorhynchus truttae
Schuetze, H. R.; and Ankel, W. E., 1976,
Ztschr. Parasitenk., v. 50 (2), 197-198
Echinorhynchus truttae-infected Gammarus
pulex fossarum, population dynamics in
stream: Oberhessen
- Echinorhynchus truttae Schrank, 1788
Willemse, J. J., 1968, Bull. Zool. Mus. Univ.
Amsterdam, v. 1 (8), 83-87
Lota lota: Vinkeveen
- Echinorhynchus variabilis Diesing 1851 nec 1856
Kritscher, E., 1976, Ann. Naturh. Mus. Wien,
v. 80, 443-449
as syn. of Octospinifer variabilis (Diesing
1851) nov. comb.
- Empodius alecturae Johnston et Edmonds
Schmidt, G. D.; and Kuntz, R. E., 1977, J.
Parasitol., v. 63 (3), 500-507
as syn. of Mediorhynchus alecturae (Johnston
et Edmonds 1947) Byrd et Kellogg 1971
- Empodius otidis Miescher
Schmidt, G. D.; and Kuntz, R. E., 1977, J.
Parasitol., v. 63 (3), 500-507
as syn. of Mediorhynchus otidis (Miescher
1841) Byrd et Kellogg 1871
- Empodius segmentatus Marval, 1902
Fabiyyi, J. P., 1972, Bull. Epizoot. Dis.
Africa, v. 20 (3), 235-238
Numida meleagris galeata (intestine):
Vom area, Benue Plateau State, Nigeria
- Empodius turnixena T.
Schmidt, G. D.; and Kuntz, R. E., 1977, J.
Parasitol., v. 63 (3), 500-507
as syn. of Mediorhynchus turnixena (Tubangui
1933) Byrd et Kellogg 1971
- Filicollis anatis (Schrank, 1788)
de Jong, N., 1976, Netherlands J. Zool., v. 26
(2), 306-318
intestinal helminths of Anas platyrhynchos,
survey, influence of host migration on para-
site prevalence, exact site in intestine
Anas platyrhynchos (jejunum, ileum): the
Naardermeer, The Netherlands
- Filicollis anatis Schrank, 1788
Kamburov, P.; and Vasilev, I., 1972, Izvest.
Tsentral. Khel'mint. Lab., v. 15, 109-133
Anas platyrhynchos (small intestine): Bul-
garia
- Filicollis anatis (Schrank, 1788) Luhe, 1911
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 17, 245-259
Limnodromus griseus
Anas acuta
A. clypeata
A. penelope
Aythya marila
Melanitta americana
Clangula hyemalis
(small intestine of all): all from Chukotsk
- Filicollis anatis (Schrank, 1788) Luhe, 1911
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 17, 260-276
Philomachus pugnax
Aythya fuligula
A. marila
Melanitta nigra
(small intestine of all): all from Siberia
- Filicollis anatis (Schrank, 1788)
Pavlov, A. V., 1966, Trudy Gel'mint. Lab.,
Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated
list: Russia
Fulica atra: Volga delta
- Golvanacanthinae nov. subfam.
Paggi, L.; and Orecchia, P., 1972, Parassitol-
ogia, v. 14 (1), 175-181
Palaeacanthocephala, Echinorhynchoidea,
Rhadinorhynchidae
key, includes: Golvanacanthus nov. gen.,
type gen.
- Golvanacanthus nov. gen. (type gen. of subfam.)
Paggi, L.; and Orecchia, P., 1972, Parassitol-
ogia, v. 14 (1), 175-181
Rhadinorhynchidae, Golvanacanthinae nov.
subfam.
tod: G. blennii nov. sp.
- Golvanacanthus blennii nov. sp. (tod), illus.
Paggi, L.; and Orecchia, P., 1972, Parassitol-
ogia, v. 14 (1), 175-181
Blennius pavo (contenuto intestinale): Golfo
di Gaeta (provincia di Latina)
- Gorgorhynchinae
Paggi, L.; and Orecchia, P., 1972, Parassitol-
ogia, v. 14 (1), 175-181
Rhadinorhynchidae, key
- Quadrigyrus. See Quadrigyrus.
- Illiosentis furcatus var. africana Golvan,
1956, illus.
Marchand, B.; and Mattei, X., 1976, J. Ultra-
structure Research, v. 54 (3), 347-358
Illiosentis furcatus var. africana, sperma-
togenesis, ultrastructure
Albula vulpes (tube digestif): Dakar

- Leptorhynchoides aphredoderi* sp. n., illus.
Buckner, R. L.; and Buckner, S. C., 1976, J. Parasitol., v. 62 (6), 955-958
Aphredoderus sayanus (pyloric ceca): Tributary of Tchefuncte River, 5 miles west of Folsom, St. Tammany Parish, Louisiana; Tributary of Bogue Chitto River, 4 miles east of Enon, Washington Parish, Louisiana
- Leptorhynchoides thecatus*
Esch, G. W.; et al., 1976, Tr. Am. Fish. Soc., v. 105 (3), 486-490
Pomphorhynchus bulbocolli, *Leptorhynchoides thecatus*, helminth recruitment, bluegills, modified live-box technique, tethered and untethered fish compared, parasite spatial distribution
Lepomis macrochirus (intestine, caeca): Gull Lake, Kalamazoo County, Michigan
- Leptorhynchoides thecatus*
Esch, G. W.; Johnson, W. C.; and Coggins, J. R., 1975, Proc. Oklahoma Acad. Sc., v. 55, 122-127
Proteocephalus ambloplitis population dynamics, smallmouth bass (*Micropterus dolomieu*), lake temperature profile and infection rates, host hormones as possible stimulus for parenteric plerocercoid migration; suggested absence of competitive interaction between *P. ambloplitis* and *Leptorhynchoides thecatus*, densities of acanthocephalans and tapeworms and number of pyloric ceca present suggested potential space available for attachment not fully exploited: Gull Lake, Kalamazoo County, Michigan
- Leptorhynchoides thecatus* (Linton)
Lang, B. Z.; and Edson, S. A., 1976, J. Parasitol., v. 62 (1), 93
Rhinichthys osculus: Turnbull National Wildlife Refuge, Spokane County, Washington
- Leptorhynchoides thecatus* (Linton, 1891) Kostylev, 1924
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
Lepomis cyanellus (intestine; mesentery or liver)
L. gibbosus (intestine; mesentery or liver)
L. macrochirus (intestine)
Micropterus salmoides (intestine; mesentery or liver)
Pomoxis nigromaculatus (intestine)
Esox lucius (intestine)
E. vermiculatus (intestine)
Morone chrysops (intestine)
Perca flavescens (intestine)
all from Nebraska
- Leptorhynchoides thecatus*, illus.
Uznanski, R. L.; and Nickol, B. B., 1976, J. Parasitol., v. 62 (4), 569-573
Leptorhynchoides thecatus eggs, external fibrillar band, structure, function (to increase intimacy of association between eggs and filamentous algae, eggs associated with algae are more likely to produce infections in intermediate host (*Hyalella azteca*) than eggs not so associated)
- Lueheia inscripta* (Westrumb, 1821) Travassos, 1919, illus.
Acholonu, A. D., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 106-116
description, life cycle
Anolis cristatellus (intestine, body cavity)
Periplaneta americana (abdomen)
all from Puerto Rico
- Macracanthorhynchus catulinus* Kostylew, 1927
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
Pachyscelis banghaasi
Trigonoscelis gemmulata
Stalagmoptera incostata
Adesmia gebleri
Dissonomus sp.
all from Tadzhikskaja SSR [and/or] Uzbekskaja SSR
- Macracanthorhynchus catulinus*
Mushkambarova, M. G., 1973, Ekol. Nasekom. Turkmen. (Tashliev), 20-35
Cyphostete komarovi
Adesmia servillei schatzmayri
Trigonoscelis gigas
T. punctipleuris
Cyphogenia limbata
Pisterotarsa gigantea subsp. zoubkoffi
P. kessleri
all from Turkmenia
- Macracanthorhynchus hirudinaceus*
Denbo, J. R.; and Miller, D. M., 1975, Tr. Illinois State Acad. Sc., v. 68 (1), 73-82
Macracanthorhynchus hirudinaceus, various dilutions of sea water, osmolarity and ionic composition of pseudocolloemic fluid, sodium, calcium and potassium ions; weak forms of ionic and osmotic regulatory mechanisms
- Macracanthorhynchus hirudinaceus*, illus.
Dunagan, T. T.; and Miller, D. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 199-208
Macracanthorhynchus hirudinaceus, muscular anatomy of the praesoma, light and scanning electron microscopy
- Macracanthorhynchus hirudinaceus*
Hightower, K.; Miller, D. M.; and Dunagan, T. T., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 71-79
Macracanthorhynchus hirudinaceus, physiology of circular and longitudinal body wall muscles, activation system
- Macracanthorhynchus hirudinaceus*, illus.
Kliks, M.; Tantachamrun, T.; and Chaiyaporn, V., 1974, Southeast Asian J. Trop. Med. and Pub. Health, v. 5 (2), 303-309
Macracanthorhynchus hirudinaceus, human, fertilized female worm removed from ulcerous area of intestinal wall, infection probably resulted from ingestion of intermediate host beetle as food, clinical case report, morphology of recovered worm: Thailand
- Macracanthorhynchus hirudinaceus*, illus.
Miller, D. M.; and Dunagan, T. T., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 99-106
Macracanthorhynchus hirudinaceus, morphology of lacunar system of body wall

- Macracanthorhynchus hirudinaceus*
Saxon, D. J.; and Dunagan, T. T., 1976, *Comp. Biochem. and Physiol.*, v. 55 (3B), 377-380
Macracanthorhynchus hirudinaceus vs. *Neoechinorhynchus* spp., pentose phosphate pathway enzymes, specific activities at different incubation temperatures, relationship to body temperature of homothermic vs. poikilothermic hosts
- Macracanthorhynchus ingens*
Barnstable, R. W.; and Dyer, W. G., 1974, *Tr. Illinois State Acad. Sc.*, v. 67 (4), 451-460
Procyon lotor (small intestine): southern Illinois
- Macracanthorhynchus ingens* (Linstow, 1879)
Meyer, 1933
Shakhmatova, V. I., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 277-289
Mustela lutreola (intestine): Karelia
- Mediorhynchus* Van Cleave 1916
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63 (1), 112-116
synonymy
Mediorhynchus, particularly *M. robustus*, description of praesomal musculature, implications for taxonomy
- Mediorhynchus* Van Cleave 1916
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
revision, proposed system of describing hook arrangement; key to species, includes: *M. conirostris* Ward 1966; *M. centurorum* Nickol 1969; *M. numidae* (Baer 1925) Meyer 1933; *M. otidis* (Miescher 1841) Byrd et Kellogg 1871; *M. taeniatis* (Linstow 1901) Dollfus 1936; *M. petrochenkoi* Gvosdev et Soboleva 1966; *M. micracanthus* (Rud. 1819) Meyer 1933; *M. grandis* Van Cleave 1916; *M. alecturae* (Johnston et Edmonds 1947) Byrd et Kellogg 1971; *M. giganteus* Meyer 1931; *M. mirabilis* (Marvel 1905) Travassos 1924; *M. gallinarum* (Bhalerao 1937) Van Cleave 1947; *M. meiringi* Bisseru 1960; *M. empodius* (Skrjabin 1913) Meyer 1933; *M. turnixena* (Tubangui 1933) Byrd et Kellogg 1971; *M. murtensis* Lunstrom 1942; *M. tenuis* Meyer 1931; *M. leptis* Ward 1966; *M. kuntzi* Ward 1960; *M. sipocotensis* Tubangui 1935; *M. corcoracis* Johnston et Edmonds 1950; *M. papillosus* Van Cleave 1916; *M. rodensis* Cosin 1971; *M. edmondsi* sp. n.; *M. oswaldocruzi* Travassos 1923; *M. emberizae* (Rud. 1819) Travassos 1924; *M. robustus* Van Cleave 1935; *M. orientalis* Belopolskaya 1953; *M. wardi* Schmidt et Canaris 1967
- Mediorhynchus* sp.?
Bisseru, B.; and Lim, K. C., 1971, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 2 (3), 412 [Demonstration]
Corvus splendens protegatus
- Mediorhynchus* spp.
Prestwood, A. K.; Kellogg, F. E.; and Doster, G. L., 1975, *Proc. 3. National Wild Turkey Symp.*, 27-32
Meleagris gallopavo silvestris: southeastern United States
- Mediorhynchus alecturae* (Johnston et Edmonds 1947) Byrd et Kellogg 1971
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
Syn.: *Empodius alecturae* Johnston et Edmonds
key
- Mediorhynchus armenicus* Petrochenko, 1958, *illus.*
Khokhlova, I. G., 1966, *Trudy Gel'mint. Lab.*, *Akad. Nauk SSSR*, v. 17, 245-259
description
Stercorarius longicaudatus
Motacilla flava
Turdus minimus
(small intestine of all): all from Chukotsk
- Mediorhynchus armenicus* Petrochenko 1958
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
as syn. of *Mediorhynchus micracanthus* (Rud. 1819) Meyer 1933
- Mediorhynchus bakeri* Byrd et Kellogg 1971
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
as syn. of *Mediorhynchus papillosus* Van Cleave 1916
- Mediorhynchus bullocki* Gupta et Jain 1973
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
as syn. of *Mediorhynchus orientalis* Belopolskaya 1953
- Mediorhynchus cambellensis* Soota, Srivastava et Ghosh 1969
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
"unrecognizable"
- Mediorhynchus centurorum* Nickol 1969, *illus.*
Nickol, B. B., 1977, *J. Parasitol.*, v. 63 (1), 104-111
Mediorhynchus centurorum, life cycle, host specificity
Parcoblatta pensylvanica (nat. and exper.): 18 mi south of Baton Rouge, Louisiana
Centurus carolinus (exper.)
Melanerpes erythrocephalus (exper.)
Colaptes auratus (exper.)
Dendrocopos villosus (exper.)
- Mediorhynchus colini* Webster 1948
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
as syn. of *Mediorhynchus papillosus* Van Cleave 1916
- Mediorhynchus edmondsi* sp. n., *illus.*
Schmidt, G. D.; and Kuntz, R. E., 1977, *J. Parasitol.*, v. 63 (3), 500-507
key
Chloropsis palawanensis (small intestine): Terabanon Concepcion, Palawan, Republic of the Philippines

- Mediorhynchus gallinarum* (Bhalerao 1937) Van Cleave 1947
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
key
Syn.: *Mediorhynchus selengensis* Harris 1973
Gallus gallus: Terabanon Concepcion, Palawan Island
- Mediorhynchus garruli* Yamaguti 1939
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
as syn. of *Mediorhynchus robustus* Van Cleave 1935
- Mediorhynchus grandis* Van Cleave, 1916
Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Mediorhynchus grandis* Van Cleave 1916 of Kelly and Finnie, 1972
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"probably a misidentification"
- Mediorhynchus lagodekhiensis* Kuraschvili 1955
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"species probably is valid but the hook sizes are not known"
- Mediorhynchus micracanthus* (?) Marval, 1905, illus.
Jaron, W., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 137-152
description, helminth fauna of adult swallows just returning from migration compared with young birds; dynamics of infection, species composition of helminths, various stages of nesting season
Delichon urbica (rectum): Poland
- Mediorhynchus micracanthus* (Rud. 1819) Meyer 1933
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
key
Syn.: *Mediorhynchus armenicus* Petrochenko 1958
- Mediorhynchus orientalis* Belopolskaya 1953, illus.
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
redescription, key
Syn.: *Mediorhynchus bullocki* Gupta et Jain 1973
Charadrius dubius curonicus
Acridotheres tristis
Alcippe brunnea brunnea
A. morrisonia morrisonia
Garrulax canorus taewanus
Pomatorhinus ruficollis musicus
Zosterops palpebrosa batanis
Cettia diphone cantans
Lioichla steeri steeri
Emberiza spodocephala
Pycnonotus zeylanicus
Charadrius dominicus fluvus: Hawaii (small intestine of all)
- Mediorhynchus oswaldocruzi*
Schmidt, G. D., 1977, J. Parasitol., v. 63 (2), 343
examination of type male and female showed that the number of spines and hooks on the proboscis were accurate according to Travassos' original description
- Mediorhynchus otidis* (Miescher 1841) Byrd et Kellogg 1871
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
Syn.: *Empodius otidis* Miescher
key
- Mediorhynchus papillosum*
Hon, L. T.; Forrester, D. J.; and Williams, L. E., jr., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 119-127
Meleagris gallopavo (lower small intestine): Florida
- Mediorhynchus papillosus* Van Cleave, 1916, illus.
Ivashkin, V. M.; and Shmytova, G. I., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 62-63
Mediorhynchus papillosus, life cycle, brief description
Tenthyria taurica
Pimelia subglobosa
(body cavity of all): all from Chernomorsk region, Krymsk oblast
- Mediorhynchus papillosus* Van Cleave, 1916
Kayton, R. J.; and Schmidt, G. D., 1975, J. Helminth., v. 49 (2), 115-119
Petrochelidon pyrrhonota: Colorado
- Mediorhynchus papillosus* Van Cleave, 1916, illus.
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
description
Anthus cervina (small intestine): Siberia
- Mediorhynchus papillosus* Van Cleave, 1916
Pavlov, A. V., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated list: Russia
- Mediorhynchus papillosus* Van Cleave 1916, illus.
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
synonymy, redescription, key
Alauda gulgula wattersi
A. arvensis pescadorei
Erithacus calliope calliope
Dicrurus macrocercus harterti
(small intestine of all)
- Mediorhynchus passeris* Das 1951
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"may be valid but discrepancies in the description do not permit a definite conclusion"
- Mediorhynchus pauciuncinatus* Dollfus 1959
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"too poorly known to be identified with certainty"

- Mediorhynchus pinto* Travassos 1923
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"too poorly known to be included in the key"
- Mediorhynchus robustus* Van Cleave, 1916
Kinsella, J. M., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 127-130
Aphelocoma c. coerulescens (small intestine): Florida
- Mediorhynchus robustus* Van Cleave 1916
Schmidt, G. D., 1977, J. Parasitol., v. 63 (1), 112-116
Mediorhynchus, particularly *M. robustus*, description of praesomal musculature, implications for taxonomy
- Mediorhynchus robustus* Van Cleave 1935
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
key
Syn.: *Mediorhynchus garruli* Yamaguti 1939
"Baker and Hamon (1968) reported *M. sipocotensis* Tubangui 1935. . . it is probable that the specimens were misidentified. Most likely they are *M. robustus*"
- Mediorhynchus selengensis* Harris 1973
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
as syn. of *Mediorhynchus gallinarum* (Bhalerao 1937) Van Cleave 1947
- Mediorhynchus sharmai* Gupta et Lata 1967
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"is undoubtedly *Centrorhynchus*"
- Mediorhynchus sipocotensis* Tubangui 1935 of
Baker and Hamon, 1968
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"it is probable that the specimens were misidentified. Most likely they are *M. robustus*"
- Mediorhynchus turnixena* (Tubangui 1933) Byrd et Kellogg 1971
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
key
Syn.: *Empodius turnixena* T.
Turnix suscitator fasciata
- Mediorhynchus vaginatus* (Diesing 1851) Meyer 1933
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"unrecognizable"
- Mediorhynchus zosteropis* (Porta 1913) Meyer 1933
Schmidt, G. D.; and Kuntz, R. E., 1977, J. Parasitol., v. 63 (3), 500-507
"unidentifiable"
- Metechinorhynchus baeri* (Kostylew, 1928)
Grigorian, Dzh. A.; Minasian, A. K.; and Vartanian, L. K., 1976, Biol. Zhurnal Armenii, v. 29 (1), 102-105
Barbus goktschaicus: lake Sevan, Armenia
- Metechinorhynchus salmonis* Muller, 1784
Mudry, D. R.; and McCart, P. J., 1976, J. Fish. Research Bd. Canada, v. 33 (2), 271-275
Salvelinus alpinus (intestine): Alaska
- Metechinorhynchus salmonis* (Müller, 1780)
Skriabina, E. S., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 169-182
Acipenser baeri: Yenisei and Lena Rivers
- Metechinorhynchus truttae* (Schrank, 1788), illus.
Rydlo, M., 1975, Fisch u. Umwelt (1), 105-112
Acanthocephala, importance in fish culture, life cycles, descriptions, pathogenicity, review: Middle Europe
- Moniliformis* sp., possibly *M. moniliformis*
Goldsmid, J. M.; Smith, M. E.; and Fleming, F., 1974, Ann. Trop. Med. and Parasitol., v. 68 (3), 363-364
Moniliformis sp., human (12-month old child), successful treatment with mebendazole: Rhodesia
- Moniliformis* sp.
Nama, H. S.; and Parihar, A., 1976, J. Helminth., v. 50 (2), 99-102
Rattus rattus rufescens (intestine): Jodhpur City area, India
- Moniliformis clarki* (Ward 1917) Chandler 1921, illus.
Buckner, S. C.; and Nickol, B. B., 1975, J. Parasitol., v. 61 (6), 991-995
comparison of *Moniliformis clarki* and *M. moniliformis* reflects distinctness of species, definitive and intermediate host specificity, laboratory life cycles, failure to hybridize
Spermophilus tridecemlineatus (nat. and exper.): near Lincoln, Nebraska
Ceuthophilus fusiformis (nat. and exper.): near Lincoln, Nebraska
C. pallidus: near Lincoln, Nebraska
Mesocricetus auratus (exper.)
Rattus norvegicus (exper.)
- Moniliformis clarki* (Ward 1917) Chandler 1921
Buckner, S. C.; and Nickol, B. B., 1975, J. Parasitol., v. 61 (6), 996-998
comparison of *Moniliformis moniliformis* and *M. clarki*, inter- and intraspecific morphological variation in usual definitive hosts and laboratory hosts
- Moniliformis clarki*
Davidson, W. R., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 211-217
epizootiologic and pathologic study of endoparasites of selected populations of gray squirrels
Sciurus carolinensis (small intestine): southeastern United States
- Moniliformis dubius*
Abele, L. G.; and Gilchrist, S., 1977, Science (4298), v. 197, 81-83
Moniliformis dubius, report of homosexual rape, interpretation of this behavior and other aspects of biology of acanthocephalans in context of parental investment and sexual selection

- Moniliformis dubius*
Asaolu, S. O., 1976, *Parasitology*, v. 73 (2), xxviii [Abstract]
Moniliformis dubius, ovarian ball development
- Moniliformis dubius*, *illus.*
Atkinson, K. H.; and Byram, J. E., 1976, *J. Morphol.*, v. 148 (4), 391-426
Moniliformis dubius, morphology and development of ovarian balls, oogenesis, rat (exper.)
- Moniliformis dubius*
Crompton, D. W. T., 1976, *Parasitology*, v. 73 (2), xxviii [Abstract]
Moniliformis dubius, unfertilized and fertilized females, ovarian balls, estimates of numbers and sizes during course of infection in male rats
- Moniliformis dubius*, *illus.*
Crompton, D. W. T.; Arnold, S.; and Walters, D. E., 1976, *Parasitology*, v. 73 (1), 65-72
Moniliformis dubius, unfertilized and fertilized females, average numbers and sizes of ovarian balls during course of infection in rats
- Moniliformis dubius*
Crompton, D. W. T.; and Nesheim, M. C., 1977, *Parasitology*, v. 75 (2), xxi-xxii [Abstract]
Moniliformis dubius, rats, effect of host dietary starch on course of infection
- Moniliformis dubius*
Nesheim, M. C.; et al., 1977, *Proc. Roy. Soc., London*, s. B (1128), v. 197, 363-383
Moniliformis dubius, course of infection, growth, and reproduction in rats fed on diets of various compositions
- Moniliformis dubius*
Starling, J. A., 1975, *Tr. Am. Micr. Soc.*, v. 94 (4), 508-523
Hymenolepis diminuta and *Moniliformis dubius*, tegumental hexose transport, compared to glucose transport of other tapeworms and mucosal brush border of the vertebrate intestine, correlation between mechanisms of membrane transport and biochemical environment of absorptive surfaces
- Moniliformis dubius* (Meyer 1933)
Starling, J. A.; and Fisher, F. M., jr., 1975, *J. Parasitol.*, v. 61 (6), 977-990
Moniliformis dubius, females, kinetics and specificity of hexose absorption
- Moniliformis dubius* (Meyer, 1933)
Wirreno, W., 1975, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 6 (1), 136-138
Rattus rattus diardi (intestines): Bogar, West Java, Indonesia
- Moniliformis moniliformis* (Bremser, 1811) Travassos 1915
Babaev, Ia.; and Kolodenko, A. I., 1975, *Izvest. Akad. Nauk Turkmen. SSR*, s. Biol. Nauk (4), 71-75
[*Hemiechinus auritus*]
[*H. hypomelas*]
all from Turkmenistan
- Moniliformis moniliformis* (Bremser 1811) Travassos 1915, *illus.*
Buckner, S. C.; and Nickol, B. B., 1975, *J. Parasitol.*, v. 61 (6), 991-995
comparison of *Moniliformis clarki* and *M. moniliformis* reflects distinctness of species, definitive and intermediate host specificity, laboratory life cycles, failure to hybridize
Rattus norvegicus
Mesocricetus auratus
Periplaneta americana
(all exper.)
- Moniliformis moniliformis* (Bremser 1811) Travassos 1915
Buckner, S. C.; and Nickol, B. B., 1975, *J. Parasitol.*, v. 61 (6), 996-998
comparison of *Moniliformis moniliformis* and *M. clarki*, inter- and intraspecific morphological variation in usual definitive hosts and laboratory hosts
- Moniliformis moniliformis*, *illus.*
Dunagan, T. T.; and Miller, D. M., 1976, *J. Parasitol.*, v. 62 (3), 442-450
Moniliformis moniliformis, nerves originating from cerebral ganglion
- Moniliformis moniliformis*
Dunagan, T. T.; and Miller, D. M., 1977, *J. Morphol.*, v. 52 (2), 171-175
Moniliformis moniliformis, description of new ganglion identified as bursal ganglion
- Moniliformis moniliformis*
Khairul Anuar, A.; and Paran, T. P., 1976, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 7 (3), 415-416
Periplaneta americana, probable active intermediate host with ecological association with field rats: Penang, Malaysia
- Moniliformis moniliformis* (= *M. dubius*)
Mishra, G. S.; and Gonzalez, J. P., 1975, *Arch. Inst. Pasteur Tunis*, v. 52 (1-2), 71-87
Rattus norvegicus (intestin grele, gros intestin): Tunis, Tunisia
- Moniliformis moniliformis* (Bremser 1811) Travassos 1915, *illus.*
al-Rawas, A. Y.; et al., 1977, *J. Parasitol.*, v. 63 (2), 396-397
human child (feces), first finding in Iraq
- Moniliformis moniliformis* (Bremser, 1811) Sultanov, N. A.; Kabilov, T.; and Davlatov, N., 1974, *Uzbek. Biol. Zhurnal* (2), 55-57
incidence and intensity of infection in intermediate hosts, larval measurements in 4 different hosts
Scarabaeus sacer
Geotrupes impressus
Onthophagus amyntas
O. koshantschikoff
Adesmia biseriata
A. gebleri
A. gracilentata
A. septemcostata
Blaps deplanata
B. ferganica
B. oblonga

- Moniliformis moniliformis. -- Continued.
Sultanov, N. A.; Kabilov, T.; and Davlatov, N., 1974, Uzbek. Biol. Zhurnal (2), 55-57
- Blaps sp.
Prosodes pygmaea
P. nitida
Phodhomala fausti
Tentyria ballionis
Zophosus scabriuscula
all from Fergansk valley, Uzbekistan
- Moniliformis moniliformis
Tobias, R. C.; and Schmidt, G. D., 1977, J. Parasitol., v. 63 (3), 588-589
Moniliformis moniliformis juveniles, in vitro cultivation, partial growth achieved
- Moniliformis m. moniliformis (Bremser, 1811)
Gafurov, A. K., 1969, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 20, 46-54
role of Tenebrionidae as intermediate hosts
Prosodes bififormis
P. vincens
Blaps deplanata reichardti
all from Tadzhik SSR
- Neoechinorhynchus sp.
Amin, O. M., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 43-46
Lepomis cyanellus (intestine): southeastern Wisconsin
- Neoechinorhynchus sp.
Niederborn, J. Y., 1974, Tr. Missouri Acad. Sci., v. 7-8, 1973-1974, 160-163
Lepomis cynellus: Johnson County, Missouri
- Neoechinorhynchus spp.
Saxon, D. J.; and Dunagan, T. T., 1976, Comp. Biochem. and Physiol., v. 55 (3B), 377-380
Macracanthorhynchus hirudinaceus vs. Neoechinorhynchus spp., pentose phosphate pathway enzymes, specific activities at different incubation temperatures, relationship to body temperature of homothermic vs. poikilothermic hosts
- Neoechinorhynchus cristatus Lynch, 1936
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
Catostomus commersoni (intestine): Nebraska
- Neoechinorhynchus cylindratus (Van Cleave, 1913)
Cooper, C. L.; Ashmead, R. R.; and Crites, J. L., 1977, Proc. Helminth. Soc. Washington, v. 44 (1), 96
prevalence, comparison with previous years
Perca flavescens (intestine): western Lake Erie
- Neoechinorhynchus cylindratus
Esch, G. W.; Johnson, W. C.; and Coggins, J. R., 1975, Proc. Oklahoma Acad. Sc., v. 55, 122-127
Micropterus dolomieu (intestine): Gull Lake, Kalamazoo County, Michigan
- Neoechinorhynchus cylindratus
Eure, H., 1976, Parasitology, v. 73 (3), 355-370
Neoechinorhynchus cylindratus in Micropterus salmoides, seasonal periodicity (attempt to determine effects of water temperature, seasonally related host feeding habits, availability of infected intermediate hosts, host's sex and age, host location within reservoir), attempted analyses of parasite recruitment rate, maturation cycle, and sex ratio: heated reservoir (Par Pond), Energy Research and Development Administration's Savannah River Plant, Aiken, South Carolina
- Neoechinorhynchus cylindratus
Gruninger, T. L.; Murphy, C. E.; Britton, J. C., 1977, Southwest. Nat., v. 22 (4), 525-535
Micropterus salmoides
M. punctulatus
Lepomis gulosus
L. macrochirus
(intestine of all): all from Eagle Mountain Lake, Texas
- Neoechinorhynchus cylindratus
Rubertone, J. A.; and Hall, J. E., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 58-59
Micropterus dolomieu (intestine): Greenbrier River below Alderson, West Virginia
- Neoechinorhynchus cylindratus (Van Cleave, 1913)
Van Cleave, 1919
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
Ambloplites rupestris (intestine)
Lepomis cyanellus (intestine)
L. gibbosus (intestine; mesentery or liver)
Micropterus dolomieu (intestine)
M. salmoides (intestine)
Pomoxis annularis (intestine)
Esox lucius (intestine)
E. vermiculatus (intestine)
Ictalurus melas (intestine; mesentery or liver)
I. punctatus (intestine)
Morone americana (intestine)
M. chrysops (intestine)
Perca flavescens (intestine)
Stizostedion vitreum (intestine)
all from Nebraska
- Neoechinorhynchus cylindratum (Van Cleave, 1913)
Williams, E. H., jr., 1975, Tr. Am. Microsc. Soc., v. 94 (3), 340-346
Moxostoma sp. (intestine): Miller Creek, north of Valley, Alabama, Lee County
- Neoechinorhynchus prolixoides
Combs, D. L.; Harley, J. P.; and Williams, J. C., 1977, Tr. Kentucky Acad. Sc., v. 38 (3-4), 128-131
Moxostoma erythrurum (gut): Kentucky River
- Neoechinorhynchus prolixus Van Cleave and Timmons, 1952
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
Carpiodes carpio
C. cyprinus
(intestine of all): all from Nebraska

- Neoechinorhynchus rutili* (Mueller, 1790)
Baker, J. C.; and Crites, J. L., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 37-39
Ictalurus punctatus (intestines): island region of western Lake Erie
- Neoechinorhynchus rutili* (Muller, 1780)
Campbell, A. D., 1974, Proc. Roy. Soc. Edinb., sect. B, Biol., v. 74, 347-364
Perca fluviatilis
Esox lucius
all from Loch Leven, Scotland
- Neoechinorhynchus rutili* (Mueller, 1780)
Dabrowska, Z., 1970, Acta Parasitol. Polon., v. 17 (20-38), 189-193
Leuciscus cephalus
L. idus
Aspius aspius
Perca fluviatilis
(intestine of all): all from Vistula River near Warsaw
- Neoechinorhynchus rutili* (Mueller, 1780)
Dickinson, A. B.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (2), 111-116
helminths of *Fundulus heteroclitus*, seasonal variations, preferred site of attachment, host size and sex
Fundulus heteroclitus: Newfoundland
- Neoechinorhynchus rutili* (Mueller, 1780)
Dickinson, A. B.; and Threlfall, W., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 86-87
Pungitius pungitius (intestine): insular Newfoundland
- Neoechinorhynchus rutili* (Mueller, 1780)
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 195-201
Lota l. lota (pyloric appendices, stomach, anterior portion of intestine)
Aspius aspius
Blicca bjoerkna
Gobio gobio
Rutilus rutilus
Scardinius erythrophthalmus
Esox lucius
Silurus glanis
all from Poland
- Neoechinorhynchus rutili* (Mueller, 1780) Hamann, 1892
Ejsymont, L., 1970, Acta Parasitol. Polon., v. 17 (20-38), 203-216
Silurus glanis (anterior portion of intestine): river Biebrza basin, Poland
- Neoechinorhynchus rutili* (Mueller, 1780) Hamann, 1892, illus.
Khatkevich, L. M., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 181-185
Neoechinorhynchus rutili, histology of excretory ducts of female reproductive system
- Neoechinorhynchus rutili* (Mueller, 1780)
Mudry, D. R.; and Anderson, R. S., 1977, J. Fish Biol., v. 11 (1), 21-33
Catostomus catostomus: Waterton Lakes National Park, Canada
- Neoechinorhynchus rutili*
Niederkorn, J. Y., 1974, Tr. Missouri Acad. Sci., v. 7-8, 1973-1974, 160-163
Lepomis cynellus: Johnson County, Missouri
- Neoechinorhynchus rutili*
Øien, K., 1976, Norwegian J. Zool., v. 24 (4), 466-467 [Abstract]
Rutilus rutilus
Leuciscus idus
(intestine of all): all from Lake Oyeren
- Neoechinorhynchus rutili* (Mueller, 1780)
Pennell, D. A.; Becker, C. D.; and Scofield, N. R., 1973, Fish. Bull., National Oceanic and Atmos. Admin., v. 71 (1), 267-277
helminths, incidence and intensity of infection in young and adult *Oncorhynchus nerka*, life cycle review: Kvichak River system, Bristol Bay, Alaska
- Neoechinorhynchus rutili* Mueller, 1780
Puczyłowska, A., 1969, Acta Parasitol. Polon., v. 16 (1-19), 1968-1969, 33-46
helminths of fishes, dynamics of infection following formation of artificial body of water, seasonal distribution, brief description
Rutilus rutilus: Zegrzynski Reservoir
- Neoechinorhynchus rutili* (Mueller, 1780), illus.
Rydlo, M., 1975, Fisch u. Umwelt (1), 105-112
Acanthocephala, importance in fish culture, life cycles, descriptions, pathogenicity, review: Middle Europe
- Neoechinorhynchus rutili* (Mueller, 1780)
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
Hybognathus hankinsoni
Notropis dorsalis
Pimephales promelas
Rhinichthys cataractae
Semotilus atromaculatus
(intestine of all): all from Nebraska
- Neoechinorhynchus rutili* Muller, 1780
Skriabina, E. S., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 169-182
Acipenser baeri: Lena River
- Neoechinorhynchus rutili* (Mueller, 1780)
Willemsse, J. J., 1968, Bull. Zool. Mus. Univ. Amsterdam, v. 1 (8), 83-87
Rutilus rutilus: Amsterdam
Scardinius erythrophthalmus: Amsterdam (Nieuwe Meer)
Pygosteus pungitius: Amsterdam (Slotermeer)
Abramis brama: IJsselmeer
- Nephridiiorhynchus major* (Bremser, 1811) Meyer, 1931
Babaev, Ia.; and Kolodenko, A. I., 1975, Izvest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk (4), 71-75
[*Hemiechinus auritus*]
[*H. hypomelas*]
all from Turkmenistan

- Octospinifer macilentus* Van Cleave, 1919
Samuel, N.; Nickol, B. B.; and Mayes, M. A.,
1976, *Am. Midland Naturalist*, v. 96 (2), 391-
406
Catostomus commersoni (intestine): Nebraska
- Octospinifer macilentus* Van Cleave, 1922
White, G. E., 1974, *Tr. Am. Micr. Soc.*, v. 93
(2), Apr., 280-282
Catostomus commersoni: Kentucky River drain-
age system
- Octospinifer variabilis* (Diesing 1851) nov.
comb., *illus.*
Kritscher, E., 1976, *Ann. Naturh. Mus. Wien*,
v. 80, 443-449
redescription, syn.: *Echinorhynchus varia-*
bilis Diesing 1851 nec 1856
Plecostomus commersonii: Sao Leopoldo (Rio
Grande do Sul), Rio do Sinos
- Oligacanthorhynchus microcephala* (Rud. 1819)
Schmidt 1972
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Didelphis albiventris: Chaco Boreal, Para-
guay
- Oligacanthorhynchus tortuosa* (Leidy 1850)
Schmidt 1972
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Euphractus sexcinctus: Chaco Boreal, Para-
guay
- Oncicola lamasi* (Freitas et Costa 1964) comb. n.
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Syn.: *Echinopardalis lamasi* Freitas et Cos-
ta
- Oncicola luehei* (Travassos 1917) Schmidt 1972
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Nasua nasua: Chaco Boreal, Paraguay
- Oncicola martini* sp. n., *illus.*
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Felis geoffroyi (small intestine): Estancia
Juan de Zalazar, Chaco Boreal, Paraguay
- Oncicola oncicola* (Ihering 1902) Travassos 1916
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Felis goeffroyi: Chaco Boreal, Paraguay
- Oncicola paracampanulata* Machado, 1963
Schmidt, G. D., 1977, *J. Parasitol.*, v. 63
(3), 508-510
Felis yagouaroundi: Chaco Boreal, Paraguay
- Oncicola venezuelensis* n. sp., *illus.*
Marteau, M., 1977, *Ann. Parasitol.*, v. 52 (1),
25-33
Felis pardalis (intestin): Venezuela
- Paracanthocephalus tenuirostris* Achmerov et
Doubrowskaja-Achmerova, 1941
Kakacheva-Avramova, D., 1973, *Izvest. Tsentral.*
Khelmit. Lab., v. 16, 87-110
Barbus meriodionalis petenyi
L[eu]ciscus cephalus
(intestine of all): all from Balkan Mountain
river(s)
- Paulisentis missouriensis* Keppner, 1974
Samuel, N.; Nickol, B. B.; and Mayes, M. A.,
1976, *Am. Midland Naturalist*, v. 96 (2), 391-
406
Semotilus atromaculatus (intestine):
Nebraska
- Pilum* gen. n.
Williams, E. H., jr., 1976, *J. Parasitol.*,
v. 62 (1), 102-104
Echinorhynchidae
tod: *P. pilum* sp. n.
- Pilum pilum* sp. n. (tod), *illus.*
Williams, E. H., jr., 1976, *J. Parasitol.*,
v. 62 (1), 102-104
Lepomis gulosus: unnamed tributary of Chat-
tahoochee River, near Huguley, Alabama
(Appalachicola River Drainage); Beaver
Swamp Creek, near Shorter, Alabama and Up-
hatee Creek, NE of Tuskegee, Alabama (Mo-
bile Bay Drainage)
L. auritus: unnamed tributary of Chatta-
hoochee River, near Huguley, Alabama (Appa-
lachicola River Drainage)
L. macrochirus: unnamed tributary of Chatta-
hoochee River, near Huguley, Alabama (Appa-
lachicola River Drainage); Beaver Swamp
Creek, near Shorter, Alabama (Mobile Bay
Drainage)
L. punctatus: unnamed tributary of Chatta-
hoochee River, near Huguley, Alabama (Appa-
lachicola River Drainage); Beaver Swamp
Creek, near Shorter, Alabama (Mobile Bay
Drainage)
Esox americanus: unnamed tributary of Chat-
tahoochee River, near Huguley, Alabama
(Appalachicola River Drainage)
Ictalurus natalis: unnamed tributary of
Chattahoochee River, near Huguley, Alabama
(Appalachicola River Drainage); Beaver
Swamp Creek, near Shorter, Alabama (Mobile
Bay Drainage); High Pine Creek, W of Roa-
noke, Alabama (Mobile Bay Drainage)
I. nebulosus: unnamed tributary of Chatta-
hoochee River, near Huguley, Alabama (Appa-
lachicola River Drainage)
Aphredoderus sayanus: Small Creek, NE of
Ogechee River, near Savannah, Georgia (At-
lantic Coast Drainage)
- Plagiorhynchus formosus*
Cooper, C. L.; and Crites, J. L., 1974, *J.*
Wildlife Dis., v. 10 (4), 397-398
Turdus migratorius (intestine): South Bass
Island, Ohio
- Plagiorhynchus formosus*
Cooper, C. L.; and Crites, J. L., 1974, *J.*
Wildlife Dis., v. 10 (4), 399-403
survey, helminths of red-winged blackbirds
including a check list of previous findings
Agelaius phoeniceus (intestine): South Bass
Island, Ohio

- Plagiorhynchus formosus* Van Cleave, 1918
Cooper, C. L.; and Crites, J. L., 1974, Proc. Helminth. Soc. Washington, v. 41 (2), 233-237
Quiscalus quiscula versicolor (intestine):
South Bass Island, Ottawa County, Ohio
- Plagiorhynchus formosus*
Cooper, C. L.; and Crites, J. L., 1976, J. Parasitol., v. 62 (1), 105-110
similarity index of helminth faunas of 7 passerine bird species, index of association of 10 species of helminths identified as having foci of infection, competition for invertebrate food resources and aggregation into mixed feeding flocks maximizes transmission:
South Bass Island, Ottawa County, Ohio
- Plagiorhynchus formosus*
Cooper, C. L.; Troutman, E. L.; and Crites, J. L., 1973, Ohio J. Sc., v. 73 (6), 376-380
Molothrus a. ater (intestine): Ottawa county, Ohio
- Polymorphidae Meyer 1931
Schmidt, G. D., 1975, J. Parasitol., v. 61 (4), 615-620
problems in taxonomy discussed, with reference to trunk spines
- Polymorphus altmani*
Hennessy, S. L.; and Morejohn, G. V., 1977, Calif. Fish and Game, v. 63 (4), 268-272
Enhydra lutris: off coastal California
- Polymorphus boschadis* (Schrank, 1788) Raillet, 1919
de Jong, N., 1976, Netherlands J. Zool., v. 26 (2), 306-318
intestinal helminths of *Anas platyrhynchos*, survey, influence of host migration on parasite prevalence, exact site in intestine
Syn.: *P. minutus* (Zeder, 1800)
Anas platyrhynchos (jejunum, ileum): the Naardermeer, The Netherlands
- Polymorphus botulus* (Van Cleave, 1916)
Bishop, C. A.; and Threlfall, W., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 25-35
Somateria mollissima (small intestine):
insular Newfoundland and/or southern Labrador
- Polymorphus contortus* (Bremser, 1821) Travassos, 1926
Ahern, W. B.; and Schmidt, G. D., 1976, Parasitology, v. 73 (3), 381-398
Recurvirostra americana (large intestine):
Colorado
- Polymorphus diploinflatus* Lundstrom, 1942, illus.
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
description
Aythya marila
Melanitta americana
M. deglandi
Clangula hyemalis
(small intestine of all): all from Chukotsk
- Polymorphus diploinflatus* Lundstrom, 1942
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
Aythya fuligula
Melanitta nigra
M. fusca
(small intestine of all): all from Siberia
- Polymorphus gavii* Hohlova, 1965
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Gavia arctica
G. immer
(small intestine of all): all from Chukotsk
- Polymorphus kenti* Van Cleave, 1947
Hennessy, S. L.; and Morejohn, G. V., 1977, Calif. Fish and Game, v. 63 (4), 268-272
Enhydra lutris: off coastal California
- Polymorphus magnus* Skrjabin, 1913
Kamburov, P.; and Vasilev, I., 1972, Izvest. Tsentral. Khelmin. Lab., v. 15, 109-133
Anas platyrhynchos
A. acuta
A. crecca
A. querquedula
(small intestine of all): all from Bulgaria
- Polymorphus magnus* Skrjabin, 1913
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Grus canadensis
Larus argentatus
Sterna hirundo
Stercorarius longicaudatus
S. parasiticus
Gavia stellata
Anas acuta
A. clypeata
A. crecca
A. formosa
A. penelope
Aythya marila
Melanitta americana
M. deglandi
Clangula hyemalis
(small intestine of all): all from Chukotsk
- Polymorphus magnus* Skrjabin, 1913, illus
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
description
Charadrius hiaticola (small intestine)
Philomachus pugnax (small intestine)
Phalaropus lobatus (small intestine)
Chlidonias leucoptera (small intestine)
Sterna paradisea (small intestine)
Cygnus bewicki (small intestine)
Anser fabalis (small intestine)
Anas acuta (small intestine)
A. crecca (small intestine)
A. penelope (small intestine)
Aythya fuligula (small intestine)
A. marila (small intestine)
Melanitta nigra (small intestine)
M. fusca (small intestine)
Clangula hyemalis (small intestine)
Mergus albellus (small intestine)
Gammarus locusta (body cavity)
all from Siberia

- Polymorphus magnus* Skrjabin, 1913
Pavlov, A. V., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated list: Russia
- Polymorphus major* Lundstrom, 1942
Hennessy, S. L.; and Morejohn, G. V., 1977, Calif. Fish and Game, v. 63 (4), 268-272
Enhydra lutris: off coastal California
- Polymorphus marilis* Van Cleave, 1939
Pavlov, A. V., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated list: Russia
- Polymorphus mathevossianae* Petrochenko, 1949
KinSELLA, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Porphyryla martinica (small intestine): Florida
- Polymorphus minutus* (Goeze, 1782)
Belogurov, O. I.; Leonov, V. A.; and Zueva, L. S., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 105-124
Larus argentatus
Uria lomvia
Cephus carbo
(small intestine of all): all from coast of Sea of Okhotsk
- Polymorphus minutus*
Eley, T. J., jr., 1976, Calif. Fish and Game, v. 62 (2), 156-157
Fulica americana (small intestine): lower Colorado River
- Polymorphus minutus* (Zeder, 1800)
de Jong, N., 1976, Netherlands J. Zool., v. 26 (2), 306-318
as syn. of *P. boschadis* (Schrank, 1788)
Raillet, 1919
- Polymorphus minutus* (Goeze, 1872) Luhe 1911
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Aythya marila
Melanitta americana
M. deglandi
(small and large intestine of all): all from Chukotsk
- Polymorphus minutus* (Goeze, 1782) Luhe, 1911
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
Aythya fuligula
A. marila
Melanitta fusca
Clangula hyemalis
Mergus albellus
(small & large intestine of all): all from Siberia
- Polymorphus minutus* (Goeze, 1782)
Pavlov, A. V., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated list: Russia
Syn.: *Echinorhynchus boschadis* Schrank, 1788
- Polymorphus minutus*
Spencer, L. T., 1974, Am. Midland Naturalist, v. 91 (2), 505-509
Polymorphus minutus, seasonal fluctuation and effect on reproduction in *Gammarus lacustris*: Sheep Lakes, Rocky Mountain National Park, Colorado
- Polymorphus minutus*
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Polymorphus paradoxus*, illus.
Holmes, J. C.; and Bethel, W. M., 1972, Zool. J. Linn. Soc., London, v. 51, Suppl. 1, 123-149
parasite modification of intermediate host behavior in order to increase vulnerability of intermediate host to predation by definitive host; reversal of phototaxis by cystacanths of *Polymorphus paradoxus* produces altered evasive response in *Gammarus lacustris* vectors which in turn increases vulnerability of infected vectors to ingestion by mallard duck hosts
- Polymorphus phippsi*, illus.
Ivanova, G. V.; and Makhanbetov, Sh., 1975, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 25, 33-37
Polymorphus phippsi males, innervation of genital system
- Polymorphus phippsi* Kostylew, 1922
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Calidrus alpina
Rissa tridactyla
Sterna paradisea
Gavia stellata
G. arctica
Fratercula corniculata
Clangula hyemalis
Somateria mollissima
S. fischeri
S. stelleri
S. spectabilis
(small intestine of all): all from Chukotsk
- Polymorphus pupa* (Linstow, 1905) Kostylew, 1922
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Somateria mollissima
S. spectabilis
(small intestine of all): all from Chukotsk
- Polymorphus strumosoides* Lundstrom, 1942
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Aythya marila
Melanitta americana
Clangula hyemalis
(small intestine of all): all from Chukotsk

- Polymorphus strumosoides* Lundstrom, 1942, illus. Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
description
Aythya fuligula
Melanitta fusca
Clangula hyemalis
Mergus albellus
(small & large intestine of all): all from Siberia
- Polymorphus trochus*
Eley, T. J., jr., 1976, Calif. Fish and Game, v. 62 (2), 156-157
Fulica americana (intestines): lower Colorado River
- Polymorphus trochus* Van Cleave, 1945
Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 245-259
Phalaropus lobatus
Larus canus
Anas acuta
A. crecca
A. penelope
Melanitta americana
Clangula hyemalis
(small intestine of all): all from Chukotsk
- Polymorphus trochus* Van Cleave, 1945, illus. Khokhlova, I. G., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 260-276
description
Calidris temmincki (large & small intestine)
Philomachus pugnax (large & small intestine)
Cygnus bewicki (large & small intestine)
Anas acuta (large & small intestine)
A. crecca (large & small intestine)
A. penelope (large & small intestine)
Melanitta fusca (large & small intestine)
Clangula hyemalis (large & small intestine)
Gammarus locusta (body cavity)
Gmelinoides fasciata (body cavity)
Euligammarus viridis (body cavity)
all from Siberia
- Polymorphus trochus* Van Cleave, 1945
Kinsella, J. M.; Hon, L. T.; and Reed, P. B., jr., 1973, Am. Midland Naturalist, v. 89 (2), 467-473
comparison of helminth fauna of common and purple gallinules
Gallinula chloropus cachinnans (small intestine): Florida
- Pomphorhynchidae [sp.]
Beacham, B. E.; and Haley, A. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 232-233
Morone americana (intestine): Hooper's Island, Chesapeake Bay
- Pomphorhynchus* sp. (acanthella)
Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmit. Lab., v. 16, 87-110
C[obitis] taenia (intestine): Balkan Mountain river(s)
- Pomphorhynchus bosniacus* Kiskaroly et Cankovic, 1967
Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmit. Lab., v. 16, 87-110
V[imba] vimba tenella (intestine): Balkan Mountain river
- Pomphorhynchus bulbocolli* (Linkins, 1919) Van Cleave, 1919
Amin, O. M., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 43-46
Catostomus commersoni (intestine): southeastern Wisconsin
- Pomphorhynchus bulbocolli*
Esch, G. W.; et al., 1976, Tr. Am. Fish. Soc., v. 105 (3), 486-490
Pomphorhynchus bulbocolli, *Leptorhynchoides thecatus*, helminth recruitment, bluegills, modified live-box technique, tethered and untethered fish compared, parasite spatial distribution
Lepomis macrochirus (intestine): Gull Lake, Kalamazoo County, Michigan
- Pomphorhynchus bulbocolli*
Harley, J. P., 1977, Tr. Kentucky Acad. Sci., v. 38 (3-4), 136-138
Pomoxis annularis (intestine): Lake Wilgreen, Madison County, Kentucky
- Pomphorhynchus bulbocolli* Linkins in Van Cleave, 1919
Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
redescription, comparison with *P. rocci*
Carpiodes carpio (intestine)
Catostomus commersoni (intestine)
Lepomis cyanelus (intestine; mesentery or liver)
L. gibbosus (intestine; mesentery or liver)
Cyprinus carpio (intestine; mesentery or liver)
Hybopsis gracilis (intestine)
Rhinichthys cataractae (intestine)
Semotilus atromaculatus (intestine)
Esox vermiculatus (intestine)
Ictalurus melas (intestine)
I. punctatus (intestine)
all from Nebraska
- Pomphorhynchus laevis* (Muller, 1776)
Grigorian, Dzh. A.; Minasian, A. K.; and Vartanian, L. K., 1976, Biol. Zhurnal Armenii, v. 29 (1), 102-105
Barbus goktschaicus: lake Sevan, Armenia
- Pomphorhynchus laevis*, Mueller, 1776
Kakacheva-Avramova, D., 1972, Izvest. Tsentral. Khelmit. Lab., v. 15, 89-107
Leuciscus cephalus
Rhodeus sericeus amarus
Gobio gobio
Tinca tinca
Alburnus alburnus
Stizostedion lucioperca
Cyprinus carpio
Barbus tauricus cyclolepis
all from River Tundzha

- Pomphorhynchus laevis (Mueller, 1776) Monticelli, 1905
Kakacheva-Avramova, D., 1973, Izvest. Tsentral. Khelmin. Lab., v. 16, 87-110
Barbus meridionalis petenyi
B. barbus
Alb[urnus] alburnus
L[euiscus] cephalus
P[erca] fluviatilis
Ph[oxinus] phoxinus
Salmo trutta morpha fario
G[obio] gobio
(intestine of all): all from Balkan Mountain river(s)
- Pomphorhynchus laevis
Kennedy, C. R.; Broughton, P. F.; and Hine, P. M., 1976, Parasitology, v. 72 (2), 195-206
Pomphorhynchus laevis, distribution along alimentary tract of 7 fish species, factors affecting site preference
- Pomphorhynchus laevis
Kennedy, C. R.; and Rumpus, A., 1977, J. Fish. Biol., v. 10 (1), 35-42
Pomphorhynchus laevis in Gammarus pulex and Leuciscus leuciscus, incidence and intensity of infection, long-term changes in P. laevis population size: River Avon, Hampshire
- Pomphorhynchus laevis, illus.
Leake, L. D., 1975, Comparative histology. An introduction to the microscopic structure of animals, 738 pp.
parasites, comparative histology, textbook
- Pomphorhynchus laevis
Marshall, J. P., 1976, Parasitology, v. 73 (2), xxix [Abstract]
Pomphorhynchus laevis, envelope surrounding larvae in Gammarus pulex, may protect from haemocytic encapsulation
- Pomphorhynchus laevis (Mueller, 1776), illus.
Rydlo, M., 1975, Fisch u. Umwelt (1), 105-112
Acanthocephala, importance in fish culture, life cycles, descriptions, pathogenicity, review: Middle Europe
- Pomphorhynchus laevis (Mueller, 1776)
Willemsse, J. J., 1968, Bull. Zool. Mus. Univ. Amsterdam, v. 1 (8), 83-87
Osmerus eperlanus: IJsselmeer
Zoarces viviparus: Molengat (Texel)
- Prosthenorchis elegans, illus.
King, N. W., jr., 1976, Scient. Publication (317). Pan Am. Health Organ., 169-198
- Prosthorhynchus sp.
Coggins, J. R., 1975, J. Elisha Mitchell Scient. Soc., v. 91 (2), 73
parasitic fauna, effect of host diet and habitat
Turdus migratorius
Quiscalus quiscula
Agelaius phoeniceus
all from Kellogg Bird Sanctuary, Michigan
- Prosthorhynchus formosum (Van Cleave, 1918)
Andrews, S. E.; and Threlfall, W., 1975, Proc. Helminth. Soc. Washington, v. 42 (1), 24-28
Corvus brachyrhynchos (mid- and posterior section of small intestine): insular Newfoundland
- Prosthorhynchus reticulatus (Westrumb, 1821)
Pavlov, A. V., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 104-127
helminth fauna of Ralliformes, annotated list: Russia
- Prosthorhynchus transversus
Vaidova, S. M., 1975, Izvest. Akad. Nauk Azerbaidzhan. SSR, s. Biol. Nauk (3), 74-79
distribution of avian helminths in relation to habitat zones (high mountain, mountain forest, forest and scrub, lowlands): Azerbaidzhan
- Pseudoechinorhynchus clavula (Dujardin, 1845)
Skriabina, E. S., 1966, Trudy Gel'mint. Lab., Akad. Nauk SSSR, v. 17, 169-182
Acpenser baeri: Yenisei River
- Quadrigyrus cholodkowskyi Kostylew, 1928
Grigorian, Dzh. A.; Minasian, A. K.; and Vartanian, L. K., 1976, Biol. Zhurnal Armenii, v. 29 (1), 102-105
Barbus goktschaicus: lake Sevan, Armenia
- Rhadinorhynchidae Travassos, 1923
Paggi, L.; and Orecchia, P., 1972, Parassitologia, v. 14 (1), 175-181
key to subfamilies, includes: Golvanacanthinae nov. subfam.; Serrasentinae; Rhadinorhynchinae; Gorgorhynchinae
- Rhadinorhynchidae [sp.]
Beacham, B. E.; and Haley, A. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (2), 232-233
Morone americana (intestine): Hooper's Island, Chesapeake Bay
- Rhadinorhynchinae
Paggi, L.; and Orecchia, P., 1972, Parassitologia, v. 14 (1), 175-181
Rhadinorhynchidae, key
- Rhadinorhynchus cadenati Golvan et Houin, 1964
Bussieras, J.; and Baudin-Laurencin, F., 1973, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 26 (4), 13a-19a
+tuna (appareil digestif): region nord de l'Atlantique tropical oriental
- Rhadinorhynchus decapteri sp. n., illus.
Parukhin, A. M.; and Kovalenko, L. M., 1976, Zool. Zhurnal, v. 55 (1), 137-138
Decapterus pinnulatus (intestine): Pacific Ocean, region of Hawaii

- Rhadinorhynchus trachuri Harada, 1935
 Baeva, O. M., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 76-79
 degree of helminth infection in different age groups of Cololabis saira: region of Kuril'sk and Japan
- Serrasentinae
 Paggi, L.; and Orecchia, P., 1972, Parassitologia, v. 14 (1), 175-181
 Rhadinorhynchidae
 key
- Serrasentis socialis (Leidy, 1851)
 Mamaev, I. L., 1968, Gel'mint. Zhivot. Tikhogo Okeana (Skriabin), 5-27
 Euthynnus affinis (stomach): South China Sea
- Serrasentis socialis Van Cleave, 1924, illus. Marchand, B.; and Mattei, X., 1977, J. Ultrastructure Research, v. 59 (3), 263-271
 Serrasentis socialis, course of spermatogenesis, migration of centriolar derivative
 Rachycentron canadum (tube digestif)
- Southwellina dimorpha Schmidt, 1973
 Bush, A. O.; and Forrester, D. J., 1976, Proc. Helminth. Soc. Washington, v. 43 (1), 17-23
 Eudocimus albus (small intestine): Florida
- Southwellina hispida
 Courtney, C. H.; and Forrester, D. J., 1974, Proc. Helminth. Soc. Washington, v. 41 (1), 89-93
 prevalence and intensity, age of host
 Pelecanus occidentalis (small and large intestine, ceca): Florida; Louisiana
- Southwellina hispida (Van Cleave 1925) Witenburg 1932
 Miller, R. L.; Olson, A. C., jr.; and Miller, L. W., 1973, Calif. Fish and Game, v. 59 (3), 196-206
 Micropterus salmoides (mesentery near intestine): southern California reservoirs
- Tanaorhamphus longirostris (Van Cleave, 1913) Ward, 1917
 Samuel, N.; Nickol, B. B.; and Mayes, M. A., 1976, Am. Midland Naturalist, v. 96 (2), 391-406
 Carpiodes carpio
 Dorosoma cepedianum
 Morone chrysops
 Ictalurus punctatus
 (intestine of all): all from Nebraska

INDEX-CATALOGUE OF MEDICAL
AND VETERINARY ZOOLOGY
SUPPLEMENT 22, PART 4

NAME

OUT

IN

