

**EFFECTS OF AGING AND EXERCISE TRAINING ON THE MECHANISMS
OF ANGIOTENSIN II-INDUCED VASOCONSTRICTION IN RAT SKELETAL
MUSCLE ARTERIOLES**

A Dissertation

by

YOONJUNG PARK

Submitted to the Office of Graduate Studies of
Texas A&M University
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

December 2006

Major Subject: Kinesiology

**EFFECTS OF AGING AND EXERCISE TRAINING ON THE MECHANISMS
OF ANGIOTENSIN II-INDUCED VASOCONSTRICTION IN RAT SKELETAL
MUSCLE ARTERIOLES**

A Dissertation

by

YOONJUNG PARK

Submitted to the Office of Graduate Studies of
Texas A&M University
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

Approved by:

Chair of Committee, Robert B. Armstrong
Committee Members, Michael D. Delp
Christopher R. Woodman
James D. Fluckey
Emily Wilson
Head of Department, Robert B. Armstrong

December 2006

Major Subject: Kinesiology

ABSTRACT

Effects of Aging and Exercise Training on the Mechanisms of Angiotensin II-Induced Vasoconstriction in Rat Skeletal Muscle Arterioles. (December 2006)

Yoonjung Park, B.A., Seoul National University;

M.S., Seoul National University;

M.A., The University of Texas at Austin

Chair of Advisory Committee: Dr. Robert B. Armstrong

Aging is associated with increases in regional and systemic vascular resistance and impaired ability to increase blood flow to active muscles during exercise. Aging enhances vasoconstrictor responsiveness in both humans and animals, and an increase in Angiotensin II-induced vasoconstriction is one possible mechanism for old age-associated increase in muscle vascular resistance. The purpose of this study was to determine 1) whether aging alters Ang II-induced vasoconstriction, 2) whether exercise training attenuates the age-associated alteration in Ang II-mediated vasoconstriction, and 3) the mechanism(s) through which aging and exercise training alter Ang II-induced vasoconstriction in rat skeletal muscle arterioles. Male Fischer 344 rats were assigned to 4 groups: Young sedentary (YS; 4 months), old sedentary (OS; 24 months), young trained (YT) and old trained (OT). Exercise-trained groups performed treadmill exercises for 60 min/day at 15 m/min, on a 15° incline for 5 days/week for 10-12 weeks. First-order (1A) arterioles were isolated from soleus and gastrocnemius muscles for *in*

in vitro experimentation. Intraluminal diameter changes were determined in response to the cumulative addition of Ang II (3×10^{-11} - 3×10^{-5} M). Ang II dose responses were then determined following the removal of endothelium and treatment with *N*^G-nitro-L-arginine methyl ester (L-NAME, 10^{-5} M), a nitric oxide synthase (NOS) inhibitor. Ang II-induced vasoconstriction was augmented in the aged skeletal muscle arterioles, both in soleus and gastrocnemius muscles, and age-associated increases in Ang II-induced vasoconstriction were abolished with the removal of endothelium and with L-NAME. Exercise training ameliorated the age-induced increase in Ang II-vasoconstriction, and this alteration was eliminated by the removal of endothelium and with NOS inhibition. These findings suggest that aging enhances Ang II-induced vasoconstrictor responses in the arterioles from both soleus, high oxidative, and white portion of gastrocnemius, low oxidative glycolytic muscles, and this age-associated change occurs through an endothelium-dependent NOS signaling pathway. These results also demonstrated that exercise training can ameliorate the age-associated increase in Ang II vasoconstriction in the arterioles from both high oxidative and low oxidative glycolytic muscles through an endothelium-mediated NOS mechanism.

ACKNOWLEDGEMENTS

My deepest gratitude goes to my advisor, Dr. Michael Delp, who is my mentor and role model as a scientist. His scientific and personal advice and encouragement have carried me through difficult times, and his tremendous efforts to guide me and insightful advice have made all my research possible. Having him as my advisor is a very significant turning point in my science career and in my life.

I would like to express my gratitude to all of my committee members. This dissertation would not have been possible without them. First, Dr. Robert Armstrong has served as the chair of my committee and given me great advice on how to be a good scientist. His advice is always warm, like the advice my father gives me. Second, Dr. Christopher Woodman has given insightful comments on my research and served me not only as just one of the committee members but also as an advisor. Third, Dr. James Fluckey has given me scientific insight on my dissertation and encouraged me with discerning comments. Lastly, Dr. Emily Wilson has taught me the biochemistry of nitric oxide synthase and shown me her support and her beautiful smile.

I would like to express sincere gratitude to all of the Delp lab members for their collaboration. Especially, I want to give very special thanks to Dr. Judy Muller-Delp for teaching me all the techniques necessary to accomplish my dissertation. I am appreciative of Dr. Rhonda Prisby for her collaboration and discussions, Dr. Brad Behnke for his nice comments, and Patrick Dougherty for helping train the rats.

I am deeply grateful to my parents and family, who trust me and have provided endless support. Especially, I would like to give special thanks to my wife, Jiyoun Paik, and son, John Hayoung Park, for their love and considerate support to complete this work.

Above all, I greatly thank my God for giving me all opportunities and precious people in my life.

TABLE OF CONTENTS

	Page
ABSTRACT	iii
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS	vii
LIST OF FIGURES	viii
LIST OF TABLES	x
CHAPTER	
I INTRODUCTION	1
1.1 Aging, exercise training, cardiovascular system, and control of muscle blood flow	1
1.2 Aging, vasoconstrictor mechanisms of blood flow and exercise training	6
1.3 Ang II	8
1.4 Aging, exercise, and Ang II	12
1.5 Purpose and hypotheses	13
II EFFECTS OF AGING AND EXERCISE TRAINING ON THE MECHANISMS OF ANGIOTENSIN II-INDUCED VASOCONSTRICTION IN RAT SKELETAL MUSCLE ARTERIOLES	16
2.1 Introduction	16
2.2 Methods	19
2.3 Results	23
2.4 Discussion	38
III SUMMARY AND CONCLUSION	48
REFERENCES	50
VITA	61

LIST OF FIGURES

		Page
Figure 2.1	Comparison of vasoconstrictor response to the cumulative addition of Angiotensin II between young sedentary (YS) and old sedentary (OS) in soleus (A) and gastrocnemius (B) muscle arterioles	28
Figure 2.2	Comparison of vasoconstrictor response to the single dose of Angiotensin II [10^{-8} M] between young sedentary (YS) and old sedentary (OS) in soleus (A) and gastrocnemius (B) muscle arterioles	29
Figure 2.3	Effect of removal of endothelium on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS), endothelium removed young sedentary (YS-E), old sedentary (OS) and endothelium removed old sedentary (OS-E) in soleus (A) and gastrocnemius (B) muscle arterioles.....	30
Figure 2.4	Effect of L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS) young sedentary with L-NAME (YS+L-NAME), old sedentary (OS) and old sedentary with L-NAME (OS+L-NAME) in soleus (A) and gastrocnemius (B) muscle arterioles.	32
Figure 2.5	Effect of exercise training on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS), young exercise trained (YT), old sedentary (OS) and old exercise trained (OT) in soleus (A) and gastrocnemius (B) muscle arterioles.....	33
Figure 2.6	Effect of removal of endothelium on vasoconstrictor response to the cumulative addition of Angiotensin II among old sedentary (OS), old exercise trained (OT), endothelium removed old sedentary (OS-E), and endothelium removed old exercise trained (OT-E) in old soleus (A) and gastrocnemius (B) muscle arterioles	34

	Page
Figure 2.7	Effect of L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II among old sedentary (OS), old exercise trained (OT), old sedentary with L-NAME (OS+L-NAME), and old exercise trained with L-NAME (OT+L-NAME) in old soleus (A) and gastrocnemius (B) muscle arterioles35
Figure 2.8	Comparison of effect of removal of endothelium and L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II in young and old sedentary soleus (A) and gastrocnemius (B) muscle arterioles36
Figure 2.9	Comparison of effect of removal of endothelium and L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II in young and old exercise trained soleus (A) and gastrocnemius (B) muscle arterioles37

LIST OF TABLES

	Page
Table 2.1	Animal characteristics of YS, OS, YT, and OT groups.25
Table 2.2	Characteristics of 1A arterioles from soleus and the superficial portion of gastrocnemius muscles from YS, OS, YT, and OT groups.26
Table 2.3	Vascular sensitivity (EC_{50}) to Ang II in arterioles from soleus and the superficial portion of gastrocnemius muscles.27

CHAPTER I

INTRODUCTION

1.1 Aging, exercise training, cardiovascular system, and control of muscle blood flow

By the year 2020, the average life expectancy will be 82.0 years for women and 74.2 years for men, and by the year 2040, it will increase to 83.1 years for women and 75.0 years for men (94). The number of the population in the United States over 65 years old will rise to 52 million by the year 2020 and to 68 million by the year 2040. This rapid growth of the elderly will have an enormous impact on future health care costs (94). A characteristic of this population is the high prevalence of cardiovascular diseases, which negatively affect both life expectancy and quality of life, and are associated with an impaired ability for adaptation to environmental change (91). Epidemiological studies have reported that advancing age, genetic factors, diabetes, lipid levels, and sedentary lifestyle are all risk factors for cardiovascular disease, such as coronary artery disease, hypertension, congestive heart failure, and stroke (39, 57). For example, advancing aging is highly correlated with the prevalence of hypertension, incidence of atherothrombotic stroke, and incidence of coronary heart disease (57).

This dissertation follows the style and format of *American Journal of Physiology-Heart and Circulatory Physiology*.

The old age-related impairment of the cardiovascular system is associated with various detrimental changes of cardiovascular structure and function. In terms of cardiovascular structure, aging results in increased vascular intimal thickness, vascular stiffness, left ventricular wall thickness and left atrial size (37, 55, 57, 76). Functional changes of the cardiovascular system with aging include decreased end diastolic filling pressure at rest (68), although systolic function is not significantly altered with aging, which preserves ejection fraction and stroke volume (90). Also, cardiac responsiveness to β -adrenergic stimuli is decreased with aging, confirming that increases in heart rate and myocardial contractility induced by catecholamines or exercise are diminished in elderly people and peak cardiac output at maximal exercise is consequently reduced with aging (32, 56).

One of the key interventions that is able to prevent and reduce the risk of cardiovascular disease is long term regular physical activity. The benefits of chronic physical activity have been reported both in healthy subjects and old patients (12). However, aging itself limits elderly individuals to continue a physically active life style due to a significantly reduced exercise capacity and maximal aerobic capacity (41, 78). This decline in exercise capacity is in part due to an attenuated ability to increase blood flow to working muscle during exercise, as well as a diminished ability to elevate cardiac output during exercise (63). It has been reported that skeletal muscle perfusion during muscle stimulation and exercise is lower in old animals (50, 75) and humans (5, 60, 64, 82, 84, 86, 106), although skeletal muscle blood flow is not altered with aging during rest (18, 41, 111). Irion et al. (50) reported that blood flow is significantly lower

in aged rats undergoing intermittent tetanic contractions compared to younger animals. Likewise, Musch et al. (75) reported that muscle blood flow in conscious exercising rats was reduced with advanced age to highly oxidative muscle, but was increased with aging to low oxidative muscle although blood flow to total hindlimb muscles was not different between young and old rats during exercise. Human data also indicate that leg muscle blood flow is lower with advanced age during submaximal exercise in sedentary men (5, 82) and women (84) and endurance-trained men (86). For example, Proctor et al. (86) demonstrated during whole body exercise that leg blood flow and vascular conductance were lower in aged individuals compared to younger counterparts. Several potential problems with the experimental approach of using whole body exercise to determine whether local vascular factors limit skeletal muscle perfusion with aging is that the lower old age-associated exercise hyperemia could result from limited cardiac output (32, 56) or a smaller muscle mass (35) among elderly subjects. To overcome these potential limitations, Lawrenson et al. (60) measured muscle blood flow and vascular resistance during small muscle mass knee extensor exercise which would not elicit maximal cardiac output and found that knee extensor blood flow was lower and vascular resistance higher across a range of work rates in aged men relative to young subjects. These findings indicate that local factors, such as decreased responsiveness to vasodilator stimuli and/or increased responsiveness to vasoconstrictor stimuli could underlie the old age-associated reduction in muscle blood flow during exercise (60).

One of the potential local mechanisms that could mediate an impaired ability to increase skeletal muscle blood flow with aging is a diminished endothelium-dependent

vasodilator function. Furchgott and Zawadzki (34) firstly found the important role of the endothelium in the vasodilation of the vascular system caused by the endothelium-dependent vasodilator ACh. They demonstrated the role of endothelial cells to release a factor that causes vasodilation in the vessel, termed endothelium-derived relaxing factor (EDRF), which modulates the vascular response (34). This factor has been identified as nitric oxide (NO) (81). NO synthesis is catalyzed by the constitutive enzyme, nitric oxide synthase (NOS), from the terminal guanidine of L-arginine, which is calcium (Ca^{++})/calmodulin-dependent (80). Since NO is rapidly diffusible, once it is synthesized, it can easily move into the smooth muscle cell through the cell membrane where it plays an important role in regulating the constriction of smooth muscle in the vascular system (33).

Although the effect of aging on endothelium-dependent vasoreactivity varies with species and vascular beds (48, 112), aging-induced impairment of endothelial function in various vascular beds has been well documented (16, 21, 22, 30, 36, 46, 71, 98, 118). Desouza et al. (22) reported that vascular responsiveness to the endothelium-dependent vasodilator, ACh, is decreased, but vasodilatation to an endothelium-independent vasodilator, nitroprusside, is unchanged in the forearm of aged humans. Woodman et al. (118) reported similar finding with isolated feed artery from the rat skeletal muscles. They found that endothelium-dependent dilation to ACh was lower in old soleus muscle feed arteries, whereas endothelium-independent dilation to sodium nitroprusside (SNP) was not different in skeletal muscle feed arteries from young and old rats. Moreover, our laboratory has shown similar results in the arterioles isolated from

skeletal muscle in aged rats. Specifically, it was reported that endothelium-dependent vasodilator responses to ACh are diminished with aging in first-order (1A) arterioles from the soleus muscle, a highly oxidative muscle, but not from the superficial portion of gastrocnemius muscle, a low-oxidative glycolytic muscle (71, 98). Furthermore, age-related differences in ACh-induced vasodilatation of soleus muscle arterioles between young and old rats are abolished with N^G -nitro-L-arginine methyl ester (L-NAME), a NOS antagonist (71, 98). These findings indicate that aging impairs endothelium-dependent and NO-mediated vasodilatation in 1A soleus muscle arterioles.

Long-term aerobic exercise training has been reported to ameliorate the age-associated dysfunction of both central and peripheral cardiovascular function (78). For example, 16 weeks of endurance training results in a significant increase in leg blood flow and vasodilator capacity in both men and women (64). Also, cross-sectional studies report that physically active individuals and aged endurance athletes have greater endothelium-dependent vasodilation through the NOS mechanism (22, 106). Moreover, several animal studies have supported the finding that exercise training results in enhanced endothelium-dependent vasodilation in conduit arteries (19, 20) and skeletal muscle arterioles (71, 98). Spier et al. (98) reported that the exercise training-induced reversal of the age-related endothelial dysfunction in soleus muscle arterioles is mediated through the NO signaling pathway.

1.2 Aging, vasoconstrictor mechanisms of blood flow and exercise training

In addition to impaired endothelium-dependent vasodilator function with aging, the age-associated reduction in skeletal muscle blood flow capacity could be due to an enhancement of resistance vessel vasoconstrictor responsiveness. Three major vasoconstrictor mechanisms have been described: noradrenergic-, endothelin-1 (ET-1)- and Ang II- mediated mechanisms. Since these are potentially important factors that control muscle blood flow at rest and during exercise, understanding the mechanism of effects of aging and exercise training on these factors is important.

Aging causes progressive increases in the sympathetic vasoconstrictor outflow to skeletal muscle in resting humans, and it is evidenced by elevations in muscle sympathetic nerve activity (MSNA) and basal norepinephrine (NE) spillover rates (97). However, age-associated responsiveness or sensitivity to α -adrenergic receptors is controversial. For example, no changes in age-associated responsiveness or sensitivity to α -adrenergic stimulation (26, 96), age-related attenuated vasoconstrictor responsiveness to sympathetic stimulation (14, 104), or elevated adrenergic sensitivity of the leg vasculature in older men (25, 54) have been reported. Although some studies have reported that leg blood flow is not changed with aging in isolated contracting muscles (51), in exercising rats (75), and in humans performing leg exercise (85), studies have supported an age-associated reduction in the ability to increase muscle blood flow during exercise as previously mentioned (50, 52, 60, 62, 86). Reductions in basal limb blood flow and vascular conductance with aging have been reported to be related to enhanced sympathetic α -adrenergic vasoconstriction (25). Koch et al. (54) reported greater

reduction in leg vascular conductance to sympathetic stimulation during exercise in older men whereas increase in sympathetic outflow to local cold stimulation during the cycle ergometer exercise was not different. Also, impaired functional sympatholysis in the forearm vascular beds during rhythmic handgrip exercise was found in older men indicating the impaired ability to blunt sympathetic α -adrenergic vasoconstriction with advancing age causing the reduction in blood flow to exercising muscle (24).

Although the mechanisms through which aging and exercise training alter endothelium-dependent vasodilation have been previously investigated (21, 71, 98, 106), the effects of aging and exercise training on the modulation of vasoconstrictor responses in the peripheral resistance vasculature have not been clearly delineated. Alterations in resistance vessel sensitivity to vasoconstrictors may contribute to the age-related reduction in exercise tolerance and skeletal muscle blood flow capacity. The mechanisms through which aging and exercise training affect noradrenergic- and ET-1-mediated vasoconstriction have been recently investigated in our laboratory (27, 28). Donato et al. (27) found that aging is associated with an augmented α -adrenergic vasoconstriction in soleus muscle arterioles and exercise training attenuated this augmentation in old rats. The aging- and training-associated alterations in α -adrenergic vasoconstriction are mediated through an endothelium-dependent mechanism, since the aging and exercise training effects are abolished with the removal of the endothelium.

Donato et al. (28) also reported that aging is associated with an enhancement of ET-1 sensitivity in 1A arterioles from the white portion of the gastrocnemius muscle, but not in 1A arterioles from the soleus muscle. This enhanced vasoconstrictor response to

ET-1 in gastrocnemius muscle arterioles from old rats is due to augmented vasoconstrictor response mediated through ET_A receptors on the smooth muscle, but not through ET_B receptors. Thus, aging has been shown to enhance vasoconstrictor responses of skeletal muscle arterioles through α -adrenergic and ET receptors. However, despite α -adrenergic and ET receptors both being present on vascular endothelial and smooth muscle cells, the aging and exercise training effects of α -receptor-mediated vasoconstriction occurred through the endothelial cells, whereas the aging effect mediated through the ET_A-receptor occurred through the smooth muscle cells. No study has been reported that investigated the mechanisms for the effects of aging and training on Ang II-induced vasoconstriction, which is one of the possible vasoconstriction mechanisms in the skeletal muscle arterioles.

1.3 Ang II

Ang II is the main biologically active peptide of the renin-angiotensin system (RAS), which plays a major physiological role in regulation of the cardiovascular system. Disorders of the RAS are associated with the pathophysiology of renal diseases, hypertension, and chronic heart failure (61). Ang II is generated by the angiotensin-converting enzyme (ACE), which is an ectoenzyme that catalyzes the extracellular conversion of Ang I to Ang II. Also, another enzyme, renin, is involved in regulation of the Ang II synthesis in the RAS. Renin is released from the juxtaglomerular cells of the kidney into the circulation where it converts angiotensinogen to Ang I. Ang II also plays an important physiological role in the regulation of blood pressure, plasma volume, and sympathetic nervous activity (7); it is in this capacity that Ang II, as a potent

vasoconstrictor, can serve to regulate blood flow through alteration in vascular tone and conductance, and ultimately serve to regulate mean arterial blood pressure (102, 105).

The vascular effects of Ang II are mediated through direct action of Ang II on Ang II receptors. Final cardiovascular responses to Ang II are determined by the result of combined action of Ang II receptors. Two major subtypes of Ang II receptors have been mainly defined on the basis of their different pharmacological and biochemical properties: Ang II type 1 receptor (AT₁R) and Ang II type 2 receptor (AT₂R) (53, 69, 74).

The locations of AT₁R and AT₂R vary depending on species of animal and tissue type. AT₁R are located in the cardiovascular, renal, endocrine, and nervous systems in humans (3). In the vasculature, AT₁R are primarily concentrated on smooth muscle cells with relatively low levels in the adventitia (2, 121) and endothelial cells (88). In contrast to AT₁R, AT₂R are highly present in fetal tissues, and its expression is rapidly decreased after birth, but, in adults, AT₂R expression is detectable in various tissues including the vasculature (1). AT₂R are expressed both in endothelial and smooth muscle cells in rat mesenteric arteries and skeletal muscle arterioles (65, 77).

AT₁Rs are involved in most of the well-known physiological effects of Ang II, and this subtype is the major effector mechanism of Ang II-mediated vascular functions (92). Activation of AT₁R results in stimulation of vasoconstriction, vascular cell hypertrophy and hyperplasia, and sodium retention (7). Also, physiological effects of this receptor have also been reported to include stimulation of reactive oxygen species (ROS) (89) and induction of inflammatory (72), thrombotic (113), and fibrotic (107) processes. In terms of vasoconstriction, binding of Ang II to AT₁R stimulates G protein-

coupled activation of phospholipase C (PLC) and results in phosphatidylinositol (PIP₃) hydrolysis and formation of inositol triphosphate (IP₃) and diacylglycerol (DAG). IP₃ activates the movement of Ca²⁺ from sarcoplasmic reticulum (SR), and DAG leads protein kinase C (PKC) to activate the sodium/hydrogen (Na⁺/H⁺) exchanger (110). Consequently, it results in increased intracellular free Ca²⁺ concentration and actin-myosin interaction in vascular smooth muscle cells and, subsequently, vascular smooth muscle contraction. Thus the overall signaling pathway through AT₁R binding results in vasoconstriction in the vasculature (110).

AT₂R has a less well-defined role in the cardiovascular system, but there is growing evidence that the AT₂R plays an important role in cardiovascular physiology. Overall, AT₂R appears to counter-regulate the excitatory effect of AT₁R, including vasodilation, antigrowth, antihypertrophic effects, and depressor regulation of blood pressure (11, 47, 66). There are evidences showing AT₂R has a significant role to control vascular tone by mediating vasodilation and counterbalancing the AT₁R-mediated vasoconstriction of Ang II. AT₂R stimulates endothelium-dependent vasodilation by release of NO (11, 61), which counteracts with the direct smooth muscle contraction that is mainly mediated by the AT₁R. Moreover, Ichiki et al. (49) found that AT₂R knock-out mice have higher blood pressure compared to the wild-type control mice, and Munzenmaier et al. (73) reported that pharmacological blockade of AT₂R augments the pressor effect of Ang II in the rat. AT₂R-mediated signaling works through a G protein coupling mechanism to mediate cardiovascular actions (47). Binding to AT₂R stimulates several cascades, including activation of protein phosphatases and

protein dephosphorylation, stimulation of phospholipase A (PLA₂) and release of arachidonic acid (AA), sphingolipid-derived ceramide, and increase in cyclic guanosine 3',5'-monophosphate (cGMP) level through a NO mechanism, which cause the vasodilation (47).

It has been well-documented that activation of AT₁R mediates vasoconstriction of conduit coronary arteries (103), but several studies relating to pharmacological properties suggest that AT₂R may mediate vasodilation of large conduit vessels (95, 117) and the coronary microcirculation (119) through the endothelium. In addition, studies have reported that, in endothelial cells, AT₁R plays a role of vasodilation through an eNOS mechanism (8), whereas activation of AT₂R on the smooth muscle cells exert a functional vasodilation via neuronal NOS (nNOS) and a soluble guanylate cyclase (sGC) pathway (15). Since Ang II exerts direct effects on both endothelial and smooth muscle cells in the vasculature, Ang II-induced vasoconstriction is determined by the net interactions between smooth muscle and endothelial cells. In order to explain the mechanisms of Ang II-induced vasoconstriction to control blood flow, Ang II-induced vasodilation through the endothelium is also critical (40), although the majority of studies in the literature have focused on the Ang II-induced vasoconstriction role through smooth muscle. Moreover, Gruetter et al. (38) reported that Ang II-induced vasoconstriction is attenuated by the release of a substance from the endothelium and enhanced by removal of endothelium. Specifically, Ang II activates NO production by vascular endothelial cells through AT₁R activation (10) and/or through AT₂R activation (93, 116). Therefore, identifying the interaction between the smooth muscle and

endothelial cell response is important for understanding the mechanism(s) of effects of aging and exercise training on Ang II-induced vasoconstriction in resistance arterioles.

Ang II has been shown to play a role in the vasomotor activity in various microvascular beds (44), but few studies investigating the direct effects of Ang II on the microcirculation have been performed. The degree of Ang II-induced vasoconstriction gradually diminishes with increasing Ang II concentration above a certain point of concentration. Zhang et al. (119) recently reported that the more potent vasoconstriction occurs at a lower concentration of Ang II (1nmol/L), where there is less vasoconstriction at higher concentrations (10nmol/L) in coronary arterioles. This is consistent with the findings in the perfused heart showing that the magnitude of coronary vasodilation is increased with increasing Ang II concentrations (83). This lower constrictor response at higher concentrations of Ang II may depend on the interaction of the smooth muscle cell and endothelial cell response to Ang II.

1.4 Aging, exercise, and Ang II

Limited information is available regarding changes in Ang II vasoreactivity with aging in the microcirculation. Although aging is associated with lower plasma renin activity in normal individuals (115), evidence suggests no aging effect on plasma Ang II concentration (29). Moreover, few studies have investigated possible changes of Ang II receptor density with aging, and what has been reported provides no clear pattern. Daubert et al. (13) reported that the density of Ang II receptor binding sites is lower in old mouse brains, whereas Heymes et al. (45) found that mRNA levels of both AT₁R and AT₂R subtypes are markedly up-regulated in the myocardium of aged rats. More

significantly, the effect of aging on Ang II-induced vasoconstriction in skeletal muscle arterioles has not been shown, although understanding the effect of aging on Ang II-induced vasoconstriction in skeletal muscle arterioles may be potentially important since Ang II-induced vasoconstriction may be one of the important contributing factors to the age-associated reduction in blood flow to skeletal muscles during exercise.

During dynamic exercise, the plasma Ang II levels are increased in an intensity-dependent manner (101, 108), and as a result, Ang II may be more involved in determining the blood flow response during exercise. Ang II is a potent vasoconstrictor in the cardiovascular system during exercise (102, 105) in that it increases the pressor response and may be involved in redirecting blood flow from the splanchnic and renal regions toward active muscles (102). However, no study has been reported regarding the effect of long-term exercise training on Ang II-induced changes in blood flow and vascular conductance. Moreover, the effect of exercise training on Ang II-induced vasoconstriction in the skeletal muscle resistance vessels also has not been investigated.

Overall, the role of Ang II has been investigated in the control of blood flow and vascular conductance during acute exercise. However, studies on the effect of aging and endurance exercise training on Ang II-mediated vascular responses are lacking.

1.5 Purpose and hypotheses

The central hypothesis for this study is that aging diminishes vascular conductance due to a shift of interaction between vascular smooth muscle cells and endothelial cells, and that exercise training serves to attenuate this effect. One possible mechanism for old age-related reductions in vascular conductance is an enhanced

vasoconstrictor responsiveness to Ang II, and that exercise training may serve to diminish Ang II-mediated vasoconstriction.

The overall purpose of this dissertation research is to determine whether aging alters Ang II vasoreactivity in skeletal muscle resistance arterioles and whether exercise training can ameliorate the putative old age-associated alteration in Ang II-induced vasoconstriction. A secondary purpose is to elucidate the mechanism(s) of the aging and exercise training effects on Ang II-induced vasoconstrictor response of skeletal muscle resistance arterioles from old rats. Therefore, six hypotheses were tested:

- 1) There will be higher Ang II-induced vasoconstriction of arterioles from soleus and gastrocnemius muscles in old rats compared to that in young animals.
- 2) Removal of the endothelium will abolish the age-associated difference of Ang II-mediated vasoconstriction in arterioles from soleus and gastrocnemius muscles.
- 3) Inhibition of NOS activity with L-NAME will abolish the age-associated difference in Ang II-mediated vasoconstriction in soleus and gastrocnemius muscle arterioles.
- 4) Exercise training will decrease Ang II-mediated vasoconstriction in aged arterioles from soleus and gastrocnemius muscles to levels at or near that in young sedentary animals.
- 5) Removal of endothelium will abolish the exercise training-associated reduction in Ang II-mediated vasoconstriction in aged arterioles from soleus and gastrocnemius muscles.

- 6) Inhibition of NOS activity with L-NAME will abolish the exercise training-associated reduction of Ang II-mediated vasoconstriction in aged arterioles from both soleus and gastrocnemius muscles.

CHAPTER II

EFFECTS OF AGING AND EXERCISE TRAINING ON THE MECHANISMS OF ANGIOTENSIN II-INDUCED VASOCONSTRICTION IN RAT SKELETAL MUSCLE ARTERIOLES

2.1 Introduction

It is well documented that aging has a detrimental effect on the cardiovascular system and that aging is an independent risk factor for cardiovascular diseases, such as atherosclerosis, hypertension, and coronary artery disease (31). In addition to increased risk of cardiovascular disease with aging, aerobic exercise capacity declines with advancing age. This decline in exercise capacity is partly due to an attenuated ability to increase blood flow to working muscle during exercise, through both declines in maximal cardiac output (63) and skeletal muscle blood flow during exercise (60).

One of the potential local mechanisms that could mediate an impaired ability to increase skeletal muscle blood flow with aging is an impaired endothelium-dependent and nitric oxide (NO)-mediated vasodilator function (16, 21, 22, 30, 36, 46, 71, 98, 118). Long-term aerobic exercise training has been also reported to ameliorate the age-associated dysfunction through the endothelium-dependent vasodilation and the NOS signaling pathway mechanism in humans and animals (22, 98, 106).

Although the mechanisms through which aging and exercise training alter endothelium-dependent vasodilation have been previously investigated, the effects of

aging and exercise training on the modulation of vasoconstrictor responses in the peripheral resistance vasculature have not been clearly delineated. Recently, however, Donato et al. (27) reported that aging is associated with an augmented α -adrenergic vasoconstriction in soleus muscle arterioles and exercise training attenuates this augmentation in old rats through an endothelium-dependent mechanism. Donato et al. (28) also found that aging is associated with an enhancement of ET-1 sensitivity in 1A arterioles from the white portion of the gastrocnemius muscle through augmented vasoconstrictor response mediated through ET_A receptors on the smooth muscle and not through the endothelium.

Ang II is the main biologically active peptide of the renin-angiotensin system (RAS), which exerts both hemodynamic and renal effects. Ang II also plays an important physiological role in the regulation of blood pressure, plasma volume, and sympathetic nervous activity in the cardiovascular system (7). In regard to the regulation of blood pressure, Ang II is a potent substance capable of constricting arterioles. Two subtypes of Ang II receptors have been mainly defined on the basis of their different pharmacological and biochemical properties. Ang II type 1 receptor (AT₁R), which is involved in most of the well-known physiological effects of Ang II and exerts potent vasoconstriction in the blood vessels, and Ang II type 2 receptor (AT₂R), which has a less well-defined role but appears capable of counterbalancing some of the effects of AT₁R stimulation (53, 69, 74). However, studies have reported that, in endothelial cells, AT₁R also plays a role of vasodilation through an eNOS mechanism (8,

87), whereas activation of AT₂R in the smooth muscle cell still exert a functional vasodilation via nNOS and sGC pathway (15).

Since Ang II exerts direct effects on both endothelial and smooth muscle cells in the vasculature, Ang II-induced vasoconstriction is determined by the interaction of vasoconstrictor and vasodilator influences mediated by smooth muscle and endothelial cells. Therefore, understanding the role of the smooth muscle and endothelial cells is important to elucidate the mechanism(s) of the effect of aging and exercise training on Ang II-induced vasoconstriction in the resistance arteriole.

Limited information is available regarding changes in Ang II vasoreactivity with aging in the microcirculation. Moreover, no direct studies have been conducted regarding the effects of aging or exercise training on Ang II-induced vasoconstriction in skeletal muscle arterioles. Therefore, the purpose of this study is to determine whether and through what mechanism(s) aging and exercise training affect Ang II-mediated vasoconstriction in rat skeletal muscle resistance arterioles. Based on results showing age enhances α -adrenoceptor-mediated vasoconstriction via diminished endothelium dependent NOS mechanism (27), we hypothesized that aging would enhance Ang II-induced vasoconstriction through an impaired endothelium-dependent NOS signaling pathway, and that exercise training would attenuate the increased Ang II-mediated vasoconstriction through the endothelium NOS mechanism.

2.2 Methods

2.2.1 Animals

Male Fischer 344 young (3-6 months) and old (22-24 months) rats were obtained from the National Institutes for Aging (NIA/Harlan) and housed in a temperature-controlled ($23\pm 2^{\circ}\text{C}$) room with a 12:12 light-dark cycle. Water and rat chow were provided ad libitum. All animal procedures were approved by the Texas A&M University Laboratory Animal Care Committee and complied with the guidelines of the National Research Council *Guide for the Care and Use of Laboratory Animals*.

2.2.2 Exercise Training

Training consisted of the following 10-12 weeks running program, executed on a motor-driven treadmill. During habituation, the rats walked on the treadmill at 10 m/min (0° incline) and then speed was increased to 15 m/min, 5 min/day for 3 days. After habituation on the treadmill, young and old rats were assigned to one of four groups, young sedentary (YS), young exercise-trained (YT), old sedentary (OS), and old exercise-trained (OT). Exercise-trained rats performed treadmill running at 15 m/min on a 15° incline, 60 min/day, 5 days a week for 10 to 12 weeks as previously described (20, 98). A minimum of 48 hours was allowed between the execution of experiments and the final bout of exercise.

2.2.3 Microvessel Preparation

The rats were anesthetized with pentobarbital sodium (60 mg/kg ip) and the gastrocnemius-plantaris-soleus muscle group from the hindlimb was carefully dissected free and placed in cold (4°C) physiological saline solution (PSS) that contained 145.0 mM NaCl, 4.7 mM KCl, 2.0 mM CaCl₂, 1.17 mM MgSO₄, 1.2 mM NaH₂PO₄, 5.0 mM glucose, 2.0 mM pyruvate, 0.02 mM EDTA, 3.0 mM MOPS buffer, and 1 g/100 ml BSA at pH 7.4. With a dissecting microscope (Olympus SVH10), 1A arterioles from the soleus muscle and the white portion of the gastrocnemius muscle were isolated and removed from the surrounding muscle tissue as previously described (67, 70). The arterioles were transferred to a Lucite chamber that contained PSS equilibrated with room temperature. Each end of the arteriole was cannulated with a micropipette and secured with nylon suture. After cannulation of the arterioles, the microvessel chamber was transferred to the stage of an inverted microscope (Olympus IX70) equipped with a video camera (Panasonic BP310), video caliper (Microcirculation Research Institute), and data acquisition system (MacLab/Macintosh) for on-line recording of intraluminal diameter. The arterioles were initially pressurized to 70 cmH₂O with two independent hydrostatic pressure reservoirs. Leaks were detected by pressurizing the vessel and then closing the valves to the reservoirs and verifying that intraluminal pressure remained constant. The arterioles that did not hold pressure were discarded. The arterioles that were free from leaks were warmed to 37°C and allowed to develop initial spontaneous tone during a 30- to 60-min equilibration period.

2.3.4 Experimental Design

Protocol I. Concentration-response relations to the cumulative addition of Ang II [3×10^{-11} to 3×10^{-5} M] were determined in arterioles from the soleus and gastrocnemius muscles from YS, OS, YT, and OT groups. Diameter was recorded for 3 minutes following each addition of Ang II. The arterioles were allowed to develop at least 15% spontaneous tone prior to addition of Ang II.

Protocol II. To determine whether alterations induced by aging and exercise training from protocol I were mediated through the vascular endothelium, the endothelium was denuded from the gastrocnemius and soleus muscle arterioles from the YS, YT, OS, and OT by passing 5 ml of air through the lumen of the vessel. In order to insure full removal of the endothelium, the arterioles were exposed to ACh [3×10^{-5} M]. Vessels that exhibited vasodilation of more than 5% were excluded from further study. Following the ACh test, the vessels were washed several times with PSS and allowed to develop spontaneous tone prior to the Ang II dose response. The diameters of denuded 1A arterioles from gastrocnemius and soleus muscles were measured in response to increasing concentrations of Ang II [3×10^{-11} to 3×10^{-5} M].

Protocol III. Since results from protocol II indicated that the effects of aging and exercise training were endothelium-dependent, another series of studies was performed to determine whether the alteration of the endothelium by aging and exercise training was mediated through the NOS signaling pathway. After the arterioles were allowed to develop spontaneous tone, they were incubated for 20 minutes with L-NAME [10^{-5} M] and the Ang II dose response [3×10^{-11} to 3×10^{-5} M] was performed.

2.2.5 Muscle Citrate Synthase Activity

Sections of the soleus and white gastrocnemius muscles from each animal were stored at -80°C for determination of citrate synthase activity (100), a measure of muscle oxidative capacity, to determine the efficacy of the training regimen. Likewise, the heart was removed to determine whether exercise training elevated heart-to-body mass ratio, an indicator of an exercise-trained state.

2.2.6 Data Analysis

Actual diameter was measured in response to Ang II and was expressed as a percentage of constrictor response according to the following formula:

$$\text{Vasoconstriction (\% Maximal Response)} = [(D_b - D_s)/(D_b) \times 100]$$

where D_b is the initial baseline diameter recorded immediately before the addition of the Ang II and D_s is the steady-state diameter measured after each dose of Ang II. Dose response curves were analyzed by two-way ANOVA with repeated measure on one factor (Ang II dose). Pairwise comparisons between specific levels were made through post-analysis (LSD). A one-way ANOVA was performed to determine significance of differences among groups in citrate synthase activity, body weight, and muscle weight. All values were presented as mean \pm SEM. Significant differences were indicated by $P \leq 0.05$.

2.3 Results

2.3.1 Animal Characteristics

The animals' age at the time of study was approximately 6 months (range, 4-7 month) for the young rats and 25 months (range, 24-25 month). Body mass was greater in old sedentary than in young sedentary rats (YS: 342 ± 6 g; OS: 431 ± 4 g), and exercise training reduced body mass in old rats (OS: 431 ± 4 g; OT: 396 ± 6 g), but not in young rats (YS: 342 ± 6 g; YT: 331 ± 7 g) (Table 2.1). Although only soleus muscle mass was increased with age, both soleus and gastrocnemius muscle mass-to-body mass ratio were decreased with aging and exercise training in young and old rats. Also, only soleus muscle mass-to-body mass ratios was increased with exercise training in young and old rats (Table 2.1).

Heart mass and left ventricle mass-to-body mass ratio were higher in the exercise trained groups and citrate synthase activity was higher in soleus muscle from both young and old trained rats (Soleus; YS 20.0 ± 0.6 , OS: 17.5 ± 0.9 , YT: 25.9 ± 0.9 , OT: 23.4 ± 1.3) indicating the efficacy of the exercise training regimen ($P < 0.05$). However, citrate synthase activity in the white portion of gastrocnemius was not altered by exercise training in either young or old rats (Table 2.1).

2.3.2 Isolated Vessel Characteristics

Maximal intraluminal diameters of soleus muscle arterioles were not different among groups, but maximal intraluminal diameters of gastrocnemius muscle arterioles were increased with age. Exercise training tended to increase maximal diameter of gastrocnemius muscle arterioles in old rats ($P = 0.057$) (YS: 151 ± 4 , OS: 172 ± 5 , YT:

156 ± 7, OT: 188 ± 7) (Table 2.2). Initial spontaneous tone developed was not different among groups in arterioles from soleus and gastrocnemius muscles.

2.3.3 Ang II Vasoconstrictor Studies

Vascular sensitivity (EC_{50}). There were no aging and exercise training effect on vascular sensitivity (EC_{50}) in the rat skeletal muscle arterioles (Table 2.3).

Effect of aging. Aging enhanced Ang II-mediated vasoconstrictor response in the 1A arterioles from both soleus (Figure 2.1 A) and gastrocnemius (Figure 2.1 B) muscles.

Vasoconstrictor response to single dose of Ang II. Ang II-induced vasoconstriction with a single dose of Ang II at the concentration of 1×10^{-8} M also resulted in a higher vasoconstrictor response in aged arterioles from soleus (A) and gastrocnemius (B) muscles (Figure 2.2). This difference was similar to the concentration with the cumulative addition of Ang II (Figure 2.1). These results indicate that the difference in Ang II-induced vasoconstriction between young and old skeletal muscle arterioles is concentration-dependent, not time-dependent.

Effect of removal of endothelium. The biphasic responses of Ang II-mediated vasoconstriction, which is a lower vasoconstrictor response at higher concentration of Ang II (higher than 3×10^{-7} M), were abolished with the removal of the endothelium in both young and old rats (Figure 2.3). More importantly, the removal of the endothelium resulted in an elimination of the age-associated difference in Ang II-mediated vasoconstriction of soleus (Figure 2.3 A) and gastrocnemius (Figure 2.3 B) muscle arterioles from young and old rats.

Table 2.1 Animal characteristics of YS, OS, YT, and OT groups.

	Sedentary		Exercise-Trained	
	Young (YS)	Old (OS)	Young (YT)	Old (OT)
N	29	29	23	20
Body Mass (g)	342 ± 6	431 ± 4 [*]	331 ± 7	396 ± 6 [‡]
Soleus Muscle Mass (mg)	150 ± 4	170 ± 4 [*]	158 ± 5	174 ± 6
Gastrocnemius Muscle Mass (mg)	1,756 ± 33	1,804 ± 37	1,713 ± 30	1,702 ± 47
Soleus Muscle Mass/Body Mass Ratio (mg/kg)	441 ± 10	395 ± 10 [*]	480 ± 13 [†]	438 ± 12 [‡]
Gastrocnemius Muscle Mass/Body Mass Ratio (mg/kg)	5,160 ± 98	4,182 ± 85 [*]	5,208 ± 119	4,295 ± 94
Heart Mass (mg)	909 ± 33	1,168 ± 37 [*]	949 ± 36	1,176 ± 37
Heart Mass/Body Mass Ratio (mg/kg)	2,657 ± 124	2,709 ± 94	2,867 ± 90 [†]	2,969 ± 117 [‡]
LV Mass/Body Mass Ratio (mg/kg)	1,822 ± 71	1,887 ± 35	2,252 ± 59 [†]	2,215 ± 66 [‡]
Soleus muscle citrate synthase activity (μmol/min/g wet Wt)	20.0 ± 0.6	17.5 ± 0.9 [*]	25.9 ± 0.9 [†]	23.4 ± 1.3 [‡]
White portion of gastrocnemius muscle citrate synthase activity (μmol/min/g wet Wt)	12.0 ± 0.3	12.1 ± 0.6	12.1 ± 0.6	13.0 ± 1.0

Wt is weight; LV is left ventricle. * indicates significant difference between young sedentary and old sedentary, † indicates significant difference between young sedentary and young trained, ‡ indicates difference between old sedentary and old trained, $P < 0.05$. Values are means ± SEM

Table 2.2 Characteristics of 1A arterioles from soleus and the superficial portion of gastrocnemius muscles from YS, OS, YT, and OT groups.

	Sedentary		Exercise-Trained	
	Young (YS)	Old (OS)	Young (YT)	Old (OT)
N	36	32	21	15
Soleus Muscle Arteriole Lumen Diameter (μm)	120 \pm 3	125 \pm 4	112 \pm 4	118 \pm 4
Gastrocnemius Muscle Arteriole Lumen Diameter (μm)	151 \pm 4	172 \pm 5*	156 \pm 7	188 \pm 7
Soleus Muscle Arteriole Spontaneous Tone (%)	51 \pm 3	45 \pm 3	50 \pm 4	40 \pm 4
Gastrocnemius Muscle Arteriole Spontaneous Tone (%)	45 \pm 3	40 \pm 3	50. \pm 4	47 \pm 4

* indicates significant difference between young sedentary and old sedentary, † indicates significant difference between young sedentary and young trained, ‡ indicates difference between old sedentary and old trained, $P < 0.05$. Values are means \pm SEM

Table 2.3 Vascular sensitivity (EC_{50}) to Ang II in arterioles from soleus and the superficial portion of gastrocnemius muscles.

		Soleus Muscle Arterioles EC_{50} (M)	Gastrocnemius Muscle Arterioles EC_{50} (M)
Sedentary	YS	$2.3 E^{-10} \pm 0.6 E^{-10}$	$1.5 E^{-10} \pm 0.3 E^{-10}$
	Young (YS) YS-E	$2.4 E^{-9} \pm 1.0 E^{-9}$	$1.7 E^{-8} \pm 0.3 E^{-8}^{\#}$
	YS+L-NAME	$4.0 E^{-9} \pm 2.2 E^{-9}$	$1.9 E^{-8} \pm 0.9 E^{-8}$
	OS	$3.0 E^{-10} \pm 1.5 E^{-10}$	$1.6 E^{-10} \pm 0.5 E^{-10}$
	Old (OS) OS-E	$8.6 E^{-9} \pm 6.6 E^{-9}^{\#}$	$7.4 E^{-8} \pm 3.4 E^{-8}^{\#}$
	OS+L-NAME	$4.0 E^{-9} \pm 2.2 E^{-9}$	$2.6 E^{-8} \pm 1.1 E^{-8}^{\Psi}$
Exercise-Trained	YT	$1.2 E^{-10} \pm 0.4 E^{-10}$	$1.3 E^{-10} \pm 0.6 E^{-10}$
	Young (YT) YT-E	$1.0 E^{-8} \pm 0.4 E^{-8}$	$1.3 E^{-8} \pm 0.7 E^{-8}$
	YT+L-NAME	$1.6 E^{-8} \pm 0.6 E^{-8}^{\S}$	$7.0 E^{-9} \pm 3.2 E^{-9}$
	OT	$1.7 E^{-10} \pm 0.6 E^{-10}$	$1.5 E^{-10} \pm 0.5 E^{-10}$
	Old (OT) OT-E	$1.8 E^{-8} \pm 1.0 E^{-8}^{\#}$	$6.4 E^{-8} \pm 1.4 E^{-8}^{\#}$
	OT+L-NAME	$6.7 E^{-9} \pm 3.4 E^{-9}$	$1.4 E^{-9} \pm 0.5 E^{-9}^{\Psi}$

-E is endothelium removed. # indicates significant effect of removal of endothelium, § indicates significant effect of treatment of L-NAME, Ψ indicates significant different between endothelium removed and L-NAME, $P < 0.05$. Values are means \pm SEM




Figure 2.1 Comparison of vasoconstrictor response to the cumulative addition of Angiotensin II between young sedentary (YS) and old sedentary (OS) in soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.2 Comparison of vasoconstrictor response to the single dose of Angiotensin II [10⁻⁸ M] between young sedentary (YS) and old sedentary (OS) in soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM. * indicates significantly different between YS and OS. (P<0.05)




Figure 2.3 Effect of removal of endothelium on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS), endothelium removed young sedentary (YS-E), old sedentary (OS) and endothelium removed old sedentary (OS-E) in soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.

Effect of NOS inhibition. Treatment with L-NAME resulted in a linear increase in the vasoconstrictor response to Ang II rather than the biphasic response occurring at the higher concentrations of Ang II in soleus and gastrocnemius muscle arterioles. (Figure 2.4). More importantly, age-related difference in Ang II-mediated vasoconstriction of soleus (Figure 2.4 A) and gastrocnemius (Figure 2.4 B) muscle arterioles were abolished by the removal of the endothelium.

Effect of exercise training. Exercise training decreased Ang II-mediated vasoconstrictor responses in arterioles from both soleus ($P < 0.05$) (Figure 2.5 A) and gastrocnemius ($P < 0.05$) (Figure 2.5 B) muscles of old rats, but training had no effect on arterioles from young rats from the soleus ($P = 0.607$) and gastrocnemius ($P = 0.965$) muscles.

Effect of removal of endothelium in old exercise trained arterioles. Following removal of vascular endothelium, the above-mentioned exercise training-mediated alterations in Ang II-induced vasoconstriction of soleus (Figure 2.6 A) and gastrocnemius (Figure 2.6 B) muscle arterioles from old rats were abolished.

Effect of NOS inhibition in old exercise trained arterioles. In the presence of L-NAME, the training-related decrease in Ang II-mediated vasoconstriction of old rat soleus (Figure 2.7 A) and gastrocnemius (Figure 2.7 B) muscle arterioles were likewise abolished.




Figure 2.4 Effect of L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS) young sedentary with L-NAME (YS+L-NAME), old sedentary (OS) and old sedentary with L-NAME (OS+L-NAME) in soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.5 Effect of exercise training on vasoconstrictor response to the cumulative addition of Angiotensin II among young sedentary (YS), young exercise trained (YT), old sedentary (OS) and old exercise trained (OT) in soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.6 Effect of removal of endothelium on vasoconstrictor response to the cumulative addition of Angiotensin II among old sedentary (OS), old exercise trained (OT), endothelium removed old sedentary (OS-E), and endothelium removed old exercise trained (OT-E) in old soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.7 Effect of L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II among old sedentary (OS), old exercise trained (OT), old sedentary with L-NAME (OS+L-NAME), and old exercise trained with L-NAME (OT+L-NAME) in old soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.8 Comparison of effect of removal of endothelium and L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II in young and old sedentary soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.




Figure 2.9 Comparison of effect of removal of endothelium and L-NAME on vasoconstrictor response to the cumulative addition of Angiotensin II in young and old exercise trained soleus (A) and gastrocnemius (B) muscle arterioles. Values are means \pm SEM.

Comparison between removal of endothelium and L-NAME. Ang II-mediated vasoconstriction appeared to be identical between arterioles with the endothelium removed and those treated with L-NAME from young and old soleus muscle (Figure 2.8). In both young and old exercise trained arterioles from soleus muscle, the effects of removal of the endothelium on Ang II-mediated vasoconstriction were also not different from treatment with of L-NAME (Figure 2.9 A). However, in arterioles from exercise trained gastrocnemius muscle, Ang II-mediated vasoconstriction in L-NAME treated arterioles was higher compared to endothelium removed arterioles in old trained rats; responses in young trained rat gastrocnemius muscle arterioles were not different (Figure 2.8 B).

2.4 Discussion

The purpose of this study was to determine 1) whether aging alters Ang II-induced vasoconstriction of skeletal muscle arterioles, 2) whether exercise training modulates Ang II-induced vasoconstriction in young and old animals, and 3) mechanisms of putative aging- and exercise training-induced differences in Ang II-induced vasoconstriction in skeletal muscle arterioles. The present study provides several unique findings. First, aging enhances Ang II-induced vasoconstrictor responses in arterioles from both the highly oxidative soleus muscle and low-oxidative superficial portion of gastrocnemius muscle. Second, the age-associated enhancement of Ang II-induced vasoconstriction occurs through an endothelium-dependent NOS signaling mechanism. And third, exercise training ameliorates the age-associated increase in Ang

II-mediated vasoconstriction in both soleus and gastrocnemius muscles, and this too occurs through an endothelium-dependent NOS signaling pathway.

2.4.1 Determination of Ang II-induced vasoconstriction

Unlike other vasoconstrictor responses in the vasculature, Ang II-induced vasoconstriction is a biphasic response, where the most potent vasoconstriction occurs at the lower concentrations of Ang II (119). Our results (Figure 2.1) are consistent with those findings showing lower vasoconstrictor responsiveness with the highest concentrations of Ang II. Many studies have reported that activation of AT₁R results in vasoconstriction through increased Ca⁺⁺ availability in the smooth muscle cell and binding to AT₂R causes vasodilation through increased NO synthesis (7, 11, 42, 65, 110). Thus, it appears that the net Ang II-induced vasoconstriction is the result of an interaction between vasoconstrictor influence via AT₁R and vasodilator influence through AT₂R. However, this view may be somewhat misleading for gaining understanding of the mechanisms of Ang II-induced vasoconstriction because several studies have reported the presence of vascular endothelial cell AT₁R which, when stimulated, result in vasodilation (8, 87). What is clear is that the endothelium plays a very important role in modulating Ang II-induced contraction of blood vessels through an endothelium dependent vasodilator mechanism (8, 38, 40, 109, 120). Therefore, Ang II-induced vasoconstriction may be determined by the interaction of vasoconstrictor signals mediated through AT₁R and the vasodilator influence mediated through endothelial cells AT₁R and AT₂R. Although AT₂R are also located on smooth muscle cells in various vascular beds which mediate vasodilation (15), vasoconstriction

mediated through AT₁R is the predominant smooth muscle cell response. Our findings in skeletal muscle arterioles support the idea that the endothelium plays an important role in determining the biphasic Ang II-induced vasoconstrictor response (Figure 2.3).

2.4.2 Age-associated enhancement in Ang II-induced vasoconstriction

In humans, it has been shown that aging results in lower leg blood flow at rest and during exercise (5, 23, 60, 82, 84, 86). Although an impairment of endothelium-dependent vasodilator function has been reported as a mechanism for this age-associated decrease in muscle blood flow in human (21, 71, 98, 118) and rats (21, 71, 98, 118), the role of vasoconstrictor mechanisms to explain the reduced blood flow capacity to the skeletal muscle with advancing aging has not been as thoroughly explored. Recently, our laboratory found that old age-related reductions in the ability to increase muscle blood flow during exercise may be due to alterations in vasoconstrictor responsiveness. Donato et al. (27, 28) found that an augmented α -adrenergic vasoconstriction and ET 1-mediated vasoconstriction may play an important role in determining age-associated reduction in skeletal muscle blood flow. One other possible vasoconstrictor mechanism may be an enhanced Ang II-mediated vasoconstriction, since it is well established that Ang II acts as a potent vasoconstrictor (102). Studies concerning the effect of aging on Ang II-induced vasoconstriction are limited and equivocal. Mixed findings have been reported regarding the effect of aging on Ang II-related vascular responses; decreased vascular responsiveness to Ang II in aorta in rats with advancing age (9) and no effect of aging on constrictor response to Ang II in rat mesenteric resistance arteries (58). The discrepancy in these results may be due to differences in vessels studied. The present study

demonstrates that there is greater Ang II-induced vasoconstriction in arterioles from skeletal muscles of old rats (Figure 2.1).

Musch et al. (75) reported that although total hindlimb muscle blood flow is not altered with advanced age during submaximal treadmill exercise in rats, blood flow to specific muscles or muscle parts are altered based on the oxidative potential of the muscle. For example, perfusion of highly oxidative muscles (e.g., the soleus muscle) is lower during exercise in old rats, whereas blood flow to low oxidative muscles (e.g., the white portion of gastrocnemius muscle) is greater. Results from the present study showing enhanced Ang II-mediated vasoconstriction with old age in soleus muscle arterioles are consistent with the lower exercise hyperemia in the soleus muscle of aged rats. However, data from gastrocnemius muscle arterioles showing enhanced Ang II-mediated vasoconstriction appear to be at odds with the observation by Musch et al. (75) of a greater muscle perfusion during exercise in this low oxidative muscle. Two factors could serve to explain this apparent discrepancy. First, several previous studies (6, 70, 71, 98) as well as the present study (Table 2.2) report that the luminal diameter of feed arteries and arterioles are enlarged in the white portion of gastrocnemius muscle from old rats. The increased size of the resistance vasculature in low oxidative muscles such as the white gastrocnemius muscle could contribute to the old age-related elevation in muscle blood flow during exercise despite a greater Ang II-induced vasoconstriction. A second potential factor contributing to the higher perfusion of low oxidative muscle is a change in the muscle recruitment pattern during exercise with aging. During moderate to high intensity treadmill running (20 m/min), the intensity of exercise used by Musch et

al. (75) to measure muscle blood flow in young and old rats, the muscles recruited in the young animals would predominantly be the high oxidative muscles, such as the soleus muscle, with little recruitment of low oxidative muscle (17, 59). However, perfusion of the highly oxidative hindlimb muscles in the old rats is lower than that in the young animals (75). The compromised delivery of oxygen to high oxidative muscle would likely cause premature fatigue of these muscles and necessitate the recruitment of other low oxidative muscles in order to maintain the exercise intensity imposed on the animals. Thus, the higher blood flow to low oxidative muscles in the older rats may more closely reflect a change in the motor unit recruitment pattern with age and consequently greater muscle activity and metabolism in the low oxidative muscle rather than a change in the vasomotor properties of the arterioles.

2.4.3 Mechanisms for age-related increase in Ang II-induced vasoconstriction

The present study indicates that the mechanisms for the age-associated alteration in Ang II-induced vasoconstriction is an endothelium-dependent mechanism, since aging differences were abolished with the removal of the endothelial cell layer (Figure 2.3). Previous studies have shown that aging-induced alterations in NE-mediated vasoconstrictor responses occur through an endothelium-dependent mechanism (27), whereas aging-induced enhancement of ET 1-mediated vasoconstriction is a smooth muscle cell-dependent mechanism through the ET_A receptors (28). Results from the present study, as well as these of NE-mediated vasoconstriction (27), are consistent with previous studies reporting that endothelial vasodilator function declines with age (21, 71, 118). Moreover, previous studies demonstrating Ang II-induced vasoconstriction has an

endothelium-dependent component supports the finding of this study. For example, Gruetter et al. (38) reported that malfunction of the endothelium significantly increases Ang II-induced contraction in rat aorta and bovine coronary artery, and Haberl et al. (40) suggested that Ang II produces a vasodilator factor from the endothelium in the rat cerebral arterioles. Therefore, dysfunction of endothelial cell signaling with aging can be an important mechanism to determine an age-associated enhancement of Ang II-mediated vasoconstriction.

As previously described, endothelium-dependent vasodilation is an important factor in determining Ang II-induced vasoconstrictor responsiveness. Three pathways responsible for endothelium-dependent vasodilation are currently known: NO, which is released by action of NOS, prostacyclin (PGI₂), which is formed through cyclooxygenase (COX), and endothelium-derived hyperpolarizing factor (EDHF). The present results indicate that the NOS signaling pathway plays the major role in determining the age-associated alteration in Ang II-induced vasoconstriction, since this age-associated difference is eliminated with treatment of L-NAME in the arterioles from soleus (Figure 2.4 A) and gastrocnemius (Figure 2.4 B) muscles. Moreover, as illustrated in Figure 2.8, the effect of NOS blockade and effect of the removal of endothelium on Ang II-induced vasoconstrictions are comparable in arterioles from soleus (Figure 2.8 A) and gastrocnemius (Figure 2.8 B) muscles. These results suggest that NO is the major mechanism responsible for the endothelium-dependent vasodilator component of Ang II-induced vasoconstriction in rat skeletal muscle arterioles. Although some studies have reported that Ang-II induced vasodilation through the endothelium is dependent on the

COX mechanism in rat cerebral arterioles (40) or, neither NOS nor COX (4), the preponderance of evidence demonstrates that Ang II-induced constriction is modulated by the release of endothelium-derived NO through endothelial AT₁R in rat carotid artery (8) and cultured endothelial cell from rat aorta (87), and endothelial AT₂R (65) in rat mesenteric resistance arteries and in rat aorta (116). Pueyo et al. (87) found that NO production in the endothelial cell is augmented by Ang II infusion and this increased NO production is abolished by losartan, an AT₁R-specific antagonist. They also found that cyclic GMP levels are increased with Ang II infusion, but it is reduced with losartan. These data suggest that Ang II can elicit endothelium-dependent vasodilation by Ang II-induced NO release and it may modulate the direct vasoconstrictor effects of Ang II on smooth muscle cells through increased cGMP production. Moreover, Ang II infusion with NG-monomethyl-L-arginine (L-NMMA), another NOS antagonist, results in less NO production and the treatment of L-NAME with Ang II decreases blood flow and vascular conductance (105).

2.4.4 The effect of exercise training on Ang II-induced vasoconstriction

Although several studies have been performed to investigate the effects of chronic exercise training on vasoconstrictor responsiveness to NE and ET-1 (20, 27, 99), no direct study elucidating the effect of endurance training on Ang II-induced vasoconstriction has been reported. It is believed that this present study is the first to examine the long-term exercise training effect on vasoconstrictor responsiveness to Ang II in rat skeletal muscle arterioles in young and old rats. Our results indicate that exercise training reduces Ang II-mediated vasoconstriction in arterioles from soleus and

gastrocnemius muscles of old rats, but has no effect on arterioles from young rats (Figure 2.5). This present study is consistent with the study of Donato et al. (27) showing that the effect of exercise training on the NE-mediated vasomotor response occurs only in aged skeletal arterioles, whereas previous studies also reported that long-term exercise training reduces NE-mediated vasoconstrictor responsiveness in abdominal aortas from young animals as well (20, 99). This discrepancy related to the effects of exercise training on NE-mediated vasoconstriction in young animals may be due to different adaptations to exercise in different vascular beds or differences in the type of artery studied, i.e., conduit arteries vs. resistance arterioles. It has shown that endurance training can increase leg blood flow in aged humans (22, 64, 106) and that the mechanism for this exercise training-induced increase in blood flow in the elderly is through endothelium-dependent vasodilation (19, 20, 22, 98, 106). Our results suggest that endurance exercise training can also ameliorate an age-associated enhancement of Ang II- induced vasoconstriction through an endothelium-dependent pathway, and that this may be one contributing mechanism to enhance blood flow in trained elderly individuals.

2.4.5 The mechanism for exercise training-associated reduction in Ang II-induced vasoconstriction

In addition, the present results suggest that the mechanism of this reduced Ang II-induced vasoconstriction in exercise trained arterioles from old skeletal muscle is the endothelium-dependent NOS signaling pathway (Figure 2.6 & Figure 2.7). It has been shown that activation of endothelial AT₁R and AT₂R results in the release of NO (8, 65,

87, 116) and PGI₂ (40), which promote vasodilation in the blood vessel. The present study indicates that NO is fully responsible for the exercise training-associated reduction in Ang II-induced vasoconstriction in the rat skeletal muscle arterioles (Figure 2.9). Although no direct studies were performed to investigate the possible PGI₂ and EDHF mechanisms on Ang II-induced vasoconstrictor responses, the notion that training primarily affects the endothelial NOS signaling pathway is supported by the finding of Spier et al. (98), who found that exercise training enhances the endothelium-mediated vasodilation via the NOS pathway in rat skeletal muscle arterioles from old rats.

One of the mechanisms to improve vasodilator function by exercise training is increased eNOS mRNA and protein expression in the vascular endothelium through a vascular shear stress stimulus (98). Exposure of the endothelium to exercise-induced increases in Ang II concentration is another possible mechanism to increase eNOS expression in the endothelium. The plasma concentration of Ang II is increased during dynamic exercise, and there is a reported doubling during exercise at 80% of maximal heart rate (114). Also, it has been reported that Ang II increases eNOS mRNA and protein expression (43, 79). Hennington et al. (43) found that acute Ang II-infusion (8 ng/kg/min) for 110 minutes resulted in increased renal eNOS mRNA, but no change in renal eNOS protein concentration, whereas chronic Ang II-infusion (5 ng/kg/min) for 10 days increased renal eNOS protein content, but no change in renal eNOS mRNA (43). Olson et al. (79) also found that in the bovine pulmonary artery endothelium that eNOS mRNA expression was increased 2.4-fold with 4 hours of 1 μ M Ang II infusion and eNOS protein content was elevated 3.3-fold with Ang II infusion for 8 hours. These data

suggest that chronic elevations in plasma Ang II concentration with dynamic exercise training could serve as a stimulus to increase levels of eNOS mRNA and protein in the endothelium. Consequently, there could be a greater vasodilator component to Ang II vasomotor responses in arterioles from exercise trained animals, resulting in a diminished net vasoconstrictor response.

2.4.6 Conclusion and significance

In conclusion, Ang II-induced vasoconstriction is determined by the net effects of a potent smooth muscle cell vasoconstrictor response and less potent endothelium-dependent vasodilator influence via a NOS signaling pathway. With aging, Ang II-mediated vasoconstriction is enhanced due to an age-associated dysfunction of the endothelium-dependent NOS vasodilator mechanism. Exercise training can ameliorate this age-associated enhancement of Ang II-induced vasoconstriction through an enhancement in the endothelium-dependent NOS signaling pathway.

Although the majority of studies in the literature have focused on the role of Ang II in the regulation of renal and splanchnic blood flow, Ang II is also important for the regulation of skeletal muscle blood flow during exercise. Since aging can elevate the Ang II-induced vasoconstrictor responsiveness of skeletal muscle arterioles and exercise training can ameliorate this enhanced vasoconstriction to Ang II, these data in the present study suggest that the effects of aging and exercise training on skeletal muscles vasomotor responsiveness to Ang II may play a role in old age-associated reductions in skeletal muscle blood flow and the training-induced restoration of muscle perfusion during exercise.

CHAPTER III

SUMMARY AND CONCLUSION

The overall purpose of this dissertation research was to determine whether aging alters Ang II vasoreactivity of skeletal muscle resistance arterioles and whether exercise training can ameliorate possible alterations in Ang II-induced vasoconstriction. A secondary purpose was to elucidate the mechanism(s) of possible aging and exercise training effects on Ang II-induced vasoconstrictor responses of skeletal muscle resistance arterioles from old rats. We tested six hypotheses and found that:

- 1) there was higher Ang II-induced vasoconstriction in arterioles from soleus and gastrocnemius muscles of old vs. young rats (Figure 2.1),
- 2) removal of the endothelium abolished the age-associated difference of Ang II-mediated vasoconstriction in soleus and gastrocnemius muscle arterioles (Figure 2.3),
- 3) inhibition of NOS activity with L-NAME abolished the age-associated difference in Ang II-mediated vasoconstriction in soleus and gastrocnemius muscle arterioles (Figure 2.4),
- 4) exercise training decreased Ang II-mediated vasoconstriction in aged arterioles from soleus and gastrocnemius muscles (Figure 2.5),

- 5) removal of the endothelium eliminated the exercise training-associated reduction in Ang II-mediated vasoconstriction in arterioles from soleus and gastrocnemius muscles of old rats (Figure 2.6), and
- 6) inhibition of NOS activity with L-NAME eliminated the exercise training-associated reduction of Ang II-mediated vasoconstriction in arterioles from both soleus and gastrocnemius muscles of aged rats (Figure 2.7).

In conclusion, aging results in enhanced Ang II-mediated vasoconstriction in the arterioles from the rat skeletal muscle due to age-associated dysfunction of endothelium-dependent NOS signaling pathway. However, exercise training could ameliorate this age-associated increase in Ang II-induced vasoconstriction in the arterioles from the rat skeletal muscle through an enhanced endothelium-dependent NOS signaling mechanism.

REFERENCES

1. **Aguilera G, Kapur S, Feuillan P, Sunar-Akbasak B, and Bathia AJ.** Developmental changes in angiotensin II receptor subtypes and AT1 receptor mRNA in rat kidney. *Kidney Int* 46: 973-979, 1994.
2. **Allen AM, Zhuo J, and Mendelsohn FAO.** Localization and function of angiotensin AT1 receptors. *Am J Hypertens* 13: 31S-38S, 2000.
3. **Allen AM, Zhuo JL, and Mendelsohn FAO.** Localization of angiotensin AT1 and AT2 receptors. *J Am Soc Nephrol* 10: S23-S29, 1999.
4. **Arima S, Endo Y, Yaoita H, Omata K, Ogawa S, Tsunoda K, Abe M, Takeuchi K, Abe K, and Ito S.** Possible role of P-450 metabolite of arachidonic acid in vasodilator mechanism of angiotensin II type 2 receptor in the isolated microperfused rabbit afferent arteriole. *J Clin Invest* 100: 2816-2823, 1997.
5. **Beere PA, Russell SD, Morey MC, Kitzman DW, and Higginbotham MB.** Aerobic exercise training can reverse age-related peripheral circulatory changes in healthy older men. *Circulation* 100: 1085-1094, 1999.
6. **Behnke B, Prisby RD, Lesniewski LA, Donato A J, Olin H, and Delp MD.** Influence of aging and physical activity on vascular morphology in rat skeletal muscle. *J Physiol (in Press)*, 2006.
7. **Berry C, Touyz R, Dominiczak AF, Webb RC, and Johns DG.** Angiotensin receptors: signaling, vascular pathophysiology, and interactions with ceramide. *Am J Physiol Heart Circ Physiol* 281(6): H2337-2365, 2001
8. **Boulanger CM, Caputo L, and Levy B.** Endothelial AT1-mediated release of nitric oxide decreases angiotensin II contractions in rat carotid artery. *Hypertension* 26: 752 -757, 1995.
9. **Cai G, Gurdal H, Seaholtz TM, and Johnson MD.** Age-related changes in angiotensin II-stimulated vascular contraction and inositol phosphate accumulation in Fischer 344 rats. *Mech Ageing Dev* 76: 125-133, 1994.
10. **Caputo L, Benessiano J, Boulanger CM, and Lévy BI.** Angiotensin II increases cGMP content via endothelial angiotensin II AT1 subtype receptor in the rat carotid artery. *Arterioscler Thromb Vasc Biol* 15: 1646-1651, 1995.
11. **Carey RM, Wang ZQ, and Siragy HM.** Update: role of the angiotensin type-2 (AT(2)) receptor in blood pressure regulation. *Curr Hypertens Rep* 2: 198-201, 2000.

12. **Cherubini A, Donahue JL, Lowenthal DT, Mecocci P, Meuleman JR, Maggio D, and Senin U.** The effect of aging and exercise on cardiovascular performance in healthy elderly. *Am J Geriat Cardiol* 7: 25-32, 1998.
13. **Daubert DL, Meadows GG, Wang JH, Sanchez PJ, and Speth RC.** Changes in angiotensin II receptors in dopamine-rich regions of the mouse brain with age and ethanol consumption. *Brain Res* 816: 8-16, 1999.
14. **Davy KP, Seals DR, and Tanaka H.** Augmented cardiopulmonary and integrative sympathetic baroreflexes but attenuated peripheral vasoconstriction with age. *Hypertension* 32: 298-304, 1998.
15. **de Godoy MA, de Oliveira AM, and Rattan S.** Angiotensin II-induced relaxation of anococcygeus smooth muscle via desensitization of AT1 receptor, and activation of AT2 receptor associated with nitric-oxide synthase pathway. *J Pharmacol Exp Ther* 311: 394-401, 2004.
16. **Delp MD, Brown M, Laughlin MH, and Hasser EM.** Rat aortic vasoreactivity is altered by old age and hindlimb unloading. *J Appl Physiol* 78: 2079-2086, 1995.
17. **Delp MD, and Duan C.** Composition and size of type I, IIA, IID/X, and IIB fibers and citrate synthase activity of rat muscle. *J Appl Physiol* 80: 261-270, 1996.
18. **Delp MD, Evans MV, and Duan C.** Effects of aging on cardiac output, regional blood flow, and body composition in Fischer-344 rats. *J Appl Physiol* 85(5): 1813-1822, 1998.
19. **Delp MD, and Laughlin MH.** Time course of enhanced endothelium-mediated dilation in aorta of trained rats. *Med Sci Sports Exerc* 29(11): 1454-1461, 1997.
20. **Delp MD, McAllister RM, and Laughlin MH.** Exercise training alters endothelium-dependent vasoreactivity of rat abdominal aorta. *J Appl Physiol* 75: 1354-1363, 1993.
21. **DeSouza CA, Clevenger CM, Greiner JJ, Smith DT, Hoetzer GL, Shapiro LF, and Stauffer BL.** Evidence for agonist-specific endothelial vasodilator dysfunction with ageing in healthy humans. *J Physiol (Lond)* 542: 255-262, 2002.
22. **DeSouza CA, Shapiro LF, Clevenger CM, Dinunno FA, Monahan KD, Tanaka H, and Seals DR.** Regular aerobic exercise prevents and restores age-related declines in endothelium-dependent vasodilation in healthy men. *Circulation* 102: 1351-1357, 2000.

23. **Dinenno FA, Jones PP, Seals DR, and Tanaka H.** Limb blood flow and vascular conductance are reduced with age in healthy humans : relation to elevations in sympathetic nerve activity and declines in oxygen demand. *Circulation* 100: 164-170, 1999.
24. **Dinenno FA, Masuki S, and Joyner MJ.** Impaired modulation of sympathetic alpha-adrenergic vasoconstriction in contracting forearm muscle of ageing men. *J Physiol* 567: 311-321, 2005
25. **Dinenno FA, Tanaka H, Stauffer BL, and Seals DR.** Reductions in basal limb blood flow and vascular conductance with human ageing: role for augmented - adrenergic vasoconstriction. *J Physiol (Lond)* 536: 977-983, 2001.
26. **Docherty JR.** Cardiovascular responses to ageing: a review. *Pharmacol Rev* 42: 103-125, 1990.
27. **Donato AJ, Lesniewski LA, and Delp MD.** The effects of aging and exercise training on adrenergic vasoreactivity in skeletal muscle arterioles. (*In Preparation*), 2006.
28. **Donato AJ, Lesniewski LA, and Delp MD.** The effects of aging and exercise training on endothelin-1 vasoconstrictor responses in rat skeletal muscle arterioles. *Cardiovasc Res* 66: 393-401, 2005
29. **Duggan J, Kilfeather S, O'Brien E, O'Malley K, and Nussberger J.** Effects of aging and hypertension on plasma angiotensin II and platelet angiotensin II receptor density. *Am J Hypertens* 5(10): 687-693, 1992.
30. **Egashira K, Inou T, Hirooka Y, Kai H, Sugimachi M, Suzuki S, Kuga T, Urabe Y, and Takeshita A.** Effects of age on endothelium-dependent vasodilation of resistance coronary artery by acetylcholine in humans. *Circulation* 88: 77-81, 1993.
31. **Ferrari AU, Radaelli A, and Centola. M.** Aging and the cardiovascular system. *J Appl Physiol* 95: 2591 - 2597, 2003.
32. **Fleg JL, O'Connor FC, Gerstenblith G, Becker LC, Clulow J, Schulman SP, and Lakatta EG.** Impact of age on the cardiovascular response to dynamic upright exercise in healthy men and women. *J Appl Physiol* 78: 890-900, 1995.
33. **Furchgott RF.** The role of endothelium in the response of vascular smooth muscle to drugs. *Annu Rev Pharmacol Toxicol* 24: 175-197, 1984.

34. **Furchgott RF, and Zawadzki JV.** The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature (Lond)* 288: 373-376, 1980.
35. **Gallagher D, Visser M, De Meersman RE, Sepulveda D, Baumgartner RN, Pierson RN, Harris T, and Heymsfield SB.** Appendicular skeletal muscle mass: effects of age, gender and ethnicity. *J Appl Physiol* 83: 229-239, 1997.
36. **Gerhard M, Roddy MA, Creager SJ, and Creager MA.** Aging progressively impairs endothelium-dependent vasodilation in forearm resistance vessels of humans. *Hypertension* 27: 849-853, 1996.
37. **Gerstenblith G, Frederiksen J, Yin FC, Fortuin NJ, Lakatta EG, and Weisfeldt ML.** Echocardiographic assessment of a normal adult ageing population. *Circulation* 56: 273-278, 1977.
38. **Gruetter CA, Ryan ET, Lemke SM, Fox MK, and Fox DD.** Endothelium-dependent modulation of angiotensin II-induced contraction in blood vessels. *Eur J Pharmacol* 146: 85-95, 1988.
39. **Grundy SM, Pasternak R, Greenland P, Smith S, and Fuster V.** Assessment of cardiovascular risk by use of multiple-risk-factor assessment equations: a statement for healthcare professionals from the American Heart Association and the American College of Cardiology. *Circulation* 100: 1481-1492, 1999.
40. **Haberl RL, Anneser F, Villringer A, and Einhaupl KM.** Angiotensin II induces endothelium-dependent vasodilation of rat cerebral arterioles. *Am J Physiol Heart Circ Physiol* 258: H1840-H1846, 1990.
41. **Haidet GC, and Parsons D.** Reduced exercise capacity in senescent beagles: an evaluation of the periphery. *Am J Physiol* 260: H173-182, 1991
42. **Hannan RE, and Widdop RE.** Vascular angiotensin II actions mediated by angiotensin II type 2 receptors. *Curr Hypertens Rep* 6: 117-123, 2004.
43. **Hennington BS, Zhang H, Miller MT, Grange rJP, and Reckelhoff JF.** Angiotensin II stimulates synthesis of endothelial nitric oxide synthase. *Hypertension* 31: 283-288, 1998
44. **Henrion D, Kubis N, and Levy BI.** Physiological and pathophysiological functions of the AT(2) subtype receptor of angiotensin II: from large arteries to the microcirculation. *Hypertension* 38(5): 1150-1157, 2001.

45. **Heymes C, Silvestre JS, Cortes CL, Chevalier B, Marotte F, Levy B, Swynghedauw B, and Samuel JL.** Cardiac senescence is associated with enhanced expression of angiotensin II receptor subtypes. *Endocrinology* 139: 2579-2587, 1998.
46. **Hongo K, Nakagomi T, Kassell NF, Sasaki T, Lehman M, Vollmer DG, Tsukahara T, Ogawa H, and Torner J.** Effects of aging and hypertension on endothelium-dependent vascular relaxation in rat carotid artery. *Stroke* 19: 892-897, 1988.
47. **Horiuchi M, Akishita M, and Dzau VJ.** Recent progress in angiotensin II type 2 receptor research in the cardiovascular system. *Hypertension* 33: 613-621, 1999.
48. **Hynes MR, and Duckles SP.** Effect of increasing age on the endothelium-mediated relaxation of rat blood vessels in vitro. *J Pharmacol Exp Therap* 241: 387-392, 1987.
49. **Ichiki T, Labosky PA, Shiota C, Okuyama S, Imagawa Y, Fugo A, Niimura F, Ichikawa I, Hogan BL, and Inagami T.** Effects on blood pressure and exploratory behavior of mice lacking angiotensin II type-2 receptor. *Nature* 377: 748-750, 1995.
50. **Irion GL, Vasthare US, and Tuma RF.** Age-related change in skeletal muscle blood flow in the rat. *J Gerontol* 42(6): 660-665, 1987.
51. **Irion GL, Vasthare US, and Tuma RF.** Preservation of skeletal muscle hyperemic response to contraction with aging in female rats. *Exp Gerontol* 23: 183-188, 1988.
52. **Jorfeldt L, and Wahren J.** Leg blood flow during exercise in man. *Clin Sci* 41: 459-473, 1971.
53. **Kambayashi Y, Bardhan S, Takahashi K, Tsuzuki S, Inui H, Hamakubo T, and Inagami T.** Molecular cloning of a novel angiotensin II receptor isoform involved in phosphotyrosine phosphatase inhibition. *J Biol Chem* 268: 24543-24546, 1993.
54. **Koch DW, Leuenberger UA, and Proctor DN.** Augmented leg vasoconstriction in dynamically exercising older men during acute sympathetic stimulation. *J Physiol* 551: 337-444, 2003.
55. **Lakatta EG.** Age-associated cardiovascular changes in health: impact on cardiovascular disease in older persons. *Heart Fail Rev* 7: 29-49, 2002.

56. **Lakatta EG.** Deficient neuroendocrine regulation of the cardiovascular system with advancing age in healthy humans. *Circulation* 87(2): 631-636, 1993.
57. **Lakatta EG, and Levy D.** Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part I: aging arteries: a "set up" for vascular disease. *Circulation* 107(1): 139-146, 2003.
58. **Lang MG, Noll G, and Luscher TF.** Effect of aging and hypertension on contractility of resistance arteries: modulation by endothelial factors. *Am J Physiol* 269: H837-H844, 1995.
59. **Laughlin MH, and Armstrong RB.** Muscular blood flow distribution patterns as a function of running speed in rats. *Am J Physiol* 243: H296-306, 1982.
60. **Lawrenson L, Poole JG, Kim J, Brown C, Patel P, and Richardson RS.** Vascular and metabolic response to isolated small muscle mass exercise: effect of age. *Am J Physiol Heart Circ Physiol* 285(3): H1023-1031, 2003.
61. **Levy BI.** Can angiotensin II type 2 receptors have deleterious effects in cardiovascular disease? Implications for therapeutic blockade of the renin-angiotensin system. *Circulation* 109: 8-13, 2004.
62. **Magnusson G, Kaijser L, Isberg B, and Saltin B.** Cardiovascular responses during one- and two-legged exercise in middle-aged men. *Acta Physiol Scand* 150: 353-362, 1994.
63. **Marin J.** Age-related changes in vascular responses: a review. *Mech Ageing Dev* 79: 71-114, 1995.
64. **Martin WHI, Kohrt WM, Malley MT, Korte E, and Stoltz S.** Exercise training enhances leg vasodilatory capacity of 65-yr-old men and women. *J Appl Physiol* 69: 1804-1809, 1990.
65. **Matrougui K, Loufrani L, Heymes C, Levy BI, and Henrion D.** Activation of AT2 receptors by endogenous angiotensin II is involved in flow-induced dilation in rat resistance arteries. *Hypertension* 34: 659-665, 1999.
66. **Matsubara H.** Pathophysiological role of angiotensin II type 2 receptor in cardiovascular and renal diseases. *Circ Res* 83: 1182-1191, 1998.
67. **McCurdy MR, Colleran PN, Muller-Delp J, and Delp MD.** Physiology of a microgravity environment: Selected contribution: effects of fiber composition and hindlimb unloading on the vasodilator properties of skeletal muscle arterioles. *J Appl Physiol* 89: 398-405, 2000.

68. **Miller TR, Grossman SJ, Schechtman KB, Biello DR, Ludbrook PA, and Ehsani AA.** Left ventricular diastolic filling and its association with age. *Am J Cardiol* 58: 531-535, 1986.
69. **Mukoyama M, Nakajima M, Horiuchi M, Sasamura H, Pratt RE, and Dzau VJ.** Expression cloning of type-2 angiotensin II receptor reveals a unique class of 7-transmembrane receptors. *J Biol Chem* 268: 24539-24542, 1993.
70. **Muller-Delp J, Spier SA, Ramsey MW, Lesniewski LA, Papadopoulos A, Humphrey JD, and Delp MD.** Effects of aging on vasoconstrictor and mechanical properties of rat skeletal muscle arterioles. *Am J Physiol Heart Circ Physiol* 282: H1843-1854, 2002.
71. **Muller-Delp JM, Spier SA, Ramsey MW, and Delp MD.** Aging impairs endothelium-dependent vasodilation in rat skeletal muscle arterioles. *Am J Physiol Heart Circ Physiol* 283: H1662-1672, 2002.
72. **Muller DN, Dechend R, Mervaala EMA, Park JK, Schmidt F, Fiebeler A, Theuer J, Breu V, Ganten D, Haller H, and Luft FC.** NF-B inhibition ameliorates angiotensin II-induced inflammatory damage in rats. *Hypertension* 35: 193-201, 2000.
73. **Munzenmaier DH, and Greene AS.** Opposing actions of angiotensin II on microvascular growth and arterial blood pressure. *Hypertension* 27: 760-765, 1996.
74. **Murphy TJ, Alexander RW, Griendling KK, Runge MS, and Bernstein KE.** Isolation of a cDNA encoding the vascular type-1 angiotensin II receptor. *Nature* 351: 233-236, 1991.
75. **Musch TI, Eklund KE, Hageman KS, and Poole DC.** Altered regional blood flow responses to submaximal exercise in older rats. *J Appl Physiol* 96(1): 81-88, 2004.
76. **Nagai J, Metter EJ, Earley CJ, Kemper MK, Becker LC, Lakatta EG, and Fleg JL.** Increased carotid artery intimal-medial thickness in asymptomatic older subjects with exercise-induced myocardial ischemia. *Circulation* 98: 1504-1509, 1998.
77. **Nora EH, Munzenmaier DH, Hansen-Smith FM, Lombard JH, and Greene AS.** Localization of the ANG II type 2 receptor in the microcirculation of skeletal muscle. *Am J Physiol Heart Circ Physiol* 275: H1395-H1403, 1998.

78. **Ogawa T, Spina RJ, Martin WH, Kohrt WM, Schechtman KB, Holloszy JO, and Ehsani AA.** Effects of aging, sex, and physical training on cardiovascular responses to exercise. *Circulation* 86: 494-503, 1992.
79. **Olson SC, Dowds TA, Pino PA, Barry MT, and Burke-Wolin T.** ANG II stimulates endothelial nitric oxide synthase expression in bovine pulmonary artery endothelium. *Am J Physiol* 273: L315-L321, 1997.
80. **Palmer RM, and Moncada S.** A novel citrulline-forming enzyme implicated in the formation of nitric oxide by vascular endothelial cells. *Biochem Biophys Res Commun* 158: 348-352, 1989.
81. **Palmer RMJ, Ferrige AG, and Moncada S.** Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 327: 524-526, 1987.
82. **Poole JG, Lawrenson L, Kim J, Brown C, and Richardson RS.** Vascular and metabolic response to cycle exercise in sedentary humans: effect of age. *Am J Physiol Heart Circ Physiol* 284: H1251-H1259, 2003.
83. **Porsti I, Hecker M, Bassenge E, and Busse R.** Dual action of angiotensin II on coronary resistance in the isolated perfused rabbit heart. *Naunyn Schmiedebergs Arch Pharmacol* 348: 650-658, 1993.
84. **Proctor DN, Koch DW, Newcomer SC, Le KU, and Leuenberger UA.** Impaired leg vasodilation during dynamic exercise in healthy older women. *J Appl Physiol* 95: 1963-1970, 2003.
85. **Proctor DN, Newcomer SC, Koch DW, Le KU, MacLean DA, and Leuenberger UA.** Leg blood flow during submaximal cycle ergometry is not reduced in healthy older normally active men. *J Appl Physiol* 94: 1859-1869, 2003.
86. **Proctor DN, Shen PH, Dietz NM, Eickhoff TJ, Lawler LA, Ebersold EJ, Loeffler DL, and Joyner MJ.** Reduced leg blood flow during dynamic exercise in older endurance-trained men. *J Appl Physiol* 85(1): 68-75, 1998.
87. **Pueyo ME, Arnal JF, Rami J, and Michel JB.** Angiotensin II stimulates the production of NO and peroxynitrite in endothelial cells. *Am J Physiol* 274: C214-C220, 1998.
88. **Pueyo ME, Diaye NN, and Michel JB.** Angiotensin II-elicited signal transduction via AT1 receptors in endothelial cells. *Br J Pharmacol* 118: 79-84, 1996.

89. **Rajagopalan S, Kurz S, Munzel T, Tarpey M, Freeman BA, Griending KK, and Harrison DG.** Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation: contribution to alterations of vasomotor tone. *J Clin Invest* 97: 1916-1923, 1996.
90. **Rodeheffer RJ, Gerstenblith G, Becker LC, Fleg JL, L WM, and Lakatta EG.** Exercise cardiac output is maintained with advancing age in healthy human subjects: cardiac dilatation and increased stroke volume compensate for diminished heart rate. *Circulation* 69: 203-213, 1984.
91. **Roth GS.** Hormone/neurotransmitter action during aging: the calcium hypothesis of impaired signal transduction. *Rev Biol Res Aging* 4: 243-252, 1990.
92. **Sadoshima J.** Versatility of the angiotensin II type I receptor. *Circ Res* 82: 1352-1355, 1998.
93. **Scheure DA, and Perrone MH.** Angiotensin II type 2 receptors mediated depressor phase of biphasic pressure response to angiotensin II *Am J Physiol* 33: R917-R923, 1993.
94. **Schneider EL, and Guralnick JM.** The aging of America: impact on health care costs. *JAMA* 263: 2335-2340, 1990.
95. **Schuijt MP, Basdew M, van Veghel R, de Vries R, Saxena PR, Schoemaker RG, and Danser AH.** AT2 receptor-mediated vasodilation in the heart: effect of myocardial infarction. *Am J Physiol Heart Circ Physiol* 281: H2590-H2596, 2001.
96. **Scott PJW, and Reid JL.** The effect of age on the responses of human isolated arteries to noradrenaline. *Br J Clin Pharmacol* 13: 237-239, 1982.
97. **Seals DR, and Esler MD.** Human ageing and the sympathoadrenal system. *J Physiol* 528: 407-417, 2000.
98. **Spier SA, Delp MD, Meininger CJ, Donato AJ, Ramsey MW, and Muller-Delp JM.** Effects of ageing and exercise training on endothelium-dependent vasodilatation and structure of rat skeletal muscle arterioles *J Physiol* 556: 947-958, 2004.
99. **Spier SA, Laughlin MH, and Delp MD.** Effects of acute and chronic exercise on vasoconstrictor responsiveness of rat abdominal aorta. *J Appl Physiol* 87: 1752-1757, 1999.
100. **Srere PA.** Citrate synthase. *Methods Enzymology* 13: 3-5, 1969.

101. **Staessen J, Fagard R, Hespel P, Lijnen P, Vanhees L, and Amery A.** Plasma renin system during exercise in normal men. *J Appl Physiol* 63: 188-194, 1987.
102. **Stebbins CL, and Symons JD.** Role of angiotensin II in hemodynamic responses to dynamic exercise in miniswine. *J Appl Physiol* 78: 185-190, 1995.
103. **Sudhir K, MacGregor JS, Gupta M, Barbant SD, Redberg R, Yock PG, and Chatterjee K.** Effect of selective angiotensin II receptor antagonism and angiotensin converting enzyme inhibition on the coronary vasculature in vivo: intravascular two-dimensional and Doppler ultrasound studies. *Circulation* 87: 931-938, 1993.
104. **Sugiyama Y, Matsukawa T, Shamsuzzaman ASM, Okada H, Watanabe T, and Mano T.** Delayed and diminished pressor response to muscle sympathetic nerve activity in the elderly. *J Appl Physiol* 80: 869-875, 1996.
105. **Symons JD, Musch TI, Hageman KS, and Stebbins CL.** Regional blood flow responses to acute ANGII infusion: effects of nitric oxide synthase inhibition. *J Cardiovasc Pharmacol* 34: 116-123, 1999.
106. **Taddei S, Galetta F, Viridis A, Ghiadoni L, Salvetti G, Franzoni F, Giusti C, and Salvetti A.** Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation* 101: 2896-2901, 2000.
107. **Takahashi T, Taniguchi T, Konish iH, Kikkawa U, Ishikawa Y, and Yokoyama M.** Activation of Akt protein kinase B after stimulation with angiotensin II in vascular smooth muscle cells. *Am J Physiol Heart Circ Physiol* 276: H1927-H1934, 1999.
108. **Tidgren B, Hjendahl P, Theodorsson E, and Nussberger J.** Renal neurohormonal and vascular responses to dynamic exercise in humans. *J Appl Physiol* 70: 2279-2286, 1991.
109. **Toda N.** Endothelium-dependent relaxation induced by angiotensin II and histamine in isolated arteries of dog. *Br J Pharmacol* 81: 301-307, 1984.
110. **Touyz RM, and Schiffrin EL.** Signal transduction mechanisms mediating the physiological and pathophysiological actions of angiotensin II in vascular smooth muscle cells. *Pharmacol Rev* 52: 639-672, 2000.
111. **Tuma RF, Irion GL, Vasthare US, and Heinel LA.** Age-related changes in regional blood flow in the rat. *Am J Physiol* 249(3 Pt 2): H485-491, 1985.
112. **Vanhoutte PM.** Aging and vascular responsiveness. *J Cardiovasc Pharmacol* 12 (supply. 8): S11-S19, 1988.

113. **Vaughan DE, Lazos SA, and Tong K.** Angiotensin II regulates the expression of plasminogen-activator inhibitor-1 in cultured endothelial cells: a potential link between the renin-angiotensin system and thrombosis. *J Clin Invest* 95: 995-1001, 1995.
114. **Warren JH, Lewis W, Wraa CE, and Stebbins CL.** Central and peripheral effects of angiotensin II on the cardiovascular response to exercise. *J Cardiovasc Pharmacol* 38(5): 693-705, 2001.
115. **Weidmann P, De Myttenaere-Bursztein S, Maxwell MH, and de Lima J.** Effect on aging on plasma renin and aldosterone in normal man. *Kidney Int* 8(5): 325-333, 1975.
116. **Wiemer G, Schölkens BA, Busse R, Wagner A, Heitsch H, and Linz W.** The functional role of angiotensin II subtype AT₂-receptors in endothelial cells and isolated ischemic rat hearts. *Pharm Pharmacol Lett* 3: 24-27, 1993.
117. **Wiemer G, Scholkens BA, Wagner A, Heitsch H, and Linz W.** The possible role of angiotensin II subtype AT₂ receptors in endothelial cells and isolated ischemic rat hearts. *J Hypertens* 11: S234-S235, 1993.
118. **Woodman CR, Price EM, and Laughlin MH.** Aging induces muscle-specific impairment of endothelium-dependent dilation in skeletal muscle feed arteries. *J Appl Physiol* 93: 1685-1690, 2002.
119. **Zhang C, Hein TW, Wang W, and Kuo L.** Divergent roles of angiotensin II AT₁ and AT₂ receptors in modulating coronary microvascular function. *Circ Res* 92: 322-329, 2003.
120. **Zhang JJ, Van Meel CA, Pfaffendorf M, Zhang J, and Van Zwiten PA.** Endothelium-dependent, nitric oxide-mediated inhibition of angiotensin II-induced contraction in rabbit aorta. *Eur J Pharmacol* 262: 247-253, 1994.
121. **Zhuo J, Moeller I, Jenkins T, Chai SY, Allen AM, Mitsuru O, and Mendelsohn AO.** Mapping tissue angiotensin-converting enzyme and angiotensin AT₁, AT₂ and AT₄ receptors. *J Hypertens* 16: 2027-2037, 1998.

VITA

Yoonjung Park

ADDRESS (Office)

Department of Veterinary Physiology & Pharmacology
 College of Veterinary Medicine and Biomedical Sciences
 Texas A&M University
 College Station, TX 77843-4466
 (979) 845-0976

EDUCATION

<u>INSTITUTION</u>	<u>DEGREE</u>	<u>YEAR</u>	<u>FIELD OF STUDY</u>
Texas A&M University	Ph.D.	2003-2006	Exercise Physiology
The University of Texas at Austin	M.A.	2000-2003	Exercise Physiology
Seoul National University	M.S.	1996-1998	Exercise Physiology
Seoul National University	B.A.	1992-1996	Physical Education

PROFESSIONAL EXPERIENCE

2003-2006	Graduate Research/Teaching Assistant, Department of Health and Kinesiology, Texas A&M University, College Station, TX
2001-2003	Graduate Teaching Assistant, Department of Kinesiology & Biological Science, The University of Texas, Austin, TX
1998-2000	Exercise Specialist, Sports and Health Medical Center, Asan Medical Center, Seoul, Korea
1997-1998	Graduate Research Assistant, Department of Physical Education Seoul National University, Seoul, Korea

BIBLIOGRAPHY

1. Delp MD, Spier SA, **Park Y**, Wu G, Muller-Delp JM. Aging diminishes endothelium-dependent vasodilation and tetrahydrobiopterin content in rat skeletal muscle arterioles. (*In Preparation*), 2006
2. Starnes JW, Taylor RP, **Park Y**. Exercise improves postischemic function in aging hearts. *Am J Physiol Heart Circ Physiol* 285(1):H347-H351, 2003
3. **Park YJ**, Lee HJ, Kim YK, Jin YS. The effects of exercise training on MVO₂ and QTc interval in obese women. *Korean J Sports Medicine* 17(1):188-196, 1999