

FORCED TWO LAYER BETA-PLANE QUASI-GEOSTROPHIC FLOW

A Dissertation

by

CONSTANTIN ONICA

Submitted to the Office of Graduate Studies of  
Texas A&M University  
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

December 2005

Major Subject: Mathematics

FORCED TWO LAYER BETA-PLANE QUASI-GEOSTROPHIC FLOW

A Dissertation

by

CONSTANTIN ONICA

Submitted to the Office of Graduate Studies of  
Texas A&M University  
in partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY

Approved by:

Chair of Committee,	Ciprian Foias
Committee Members,	R. Lee Panetta
	Carl Percy
	Emil Straube
Head of Department,	Al Boggess

December 2005

Major Subject: Mathematics

## ABSTRACT

Forced Two Layer Beta-Plane Quasi-Geostrophic Flow. (December 2005)

Constantin Onica, B.S., Al. I. Cuza University, Iasi, Romania;

M.S., Al. I. Cuza University, Iasi, Romania

Chair of Advisory Committee: Dr. Ciprian Foias

We consider a model of quasigeostrophic turbulence that has proven useful in theoretical studies of large scale heat transport and coherent structure formation in planetary atmospheres and oceans. The model consists of a coupled pair of hyperbolic PDE's with a forcing which represents domain-scale thermal energy source. Although the use to which the model is typically put involves gathering information from very long numerical integrations, little of a rigorous nature is known about long-time properties of solutions to the equations. In the first part of my dissertation we define a notion of weak solution, and show using Galerkin methods the long-time existence and uniqueness of such solutions. In the second part we prove that the unique weak solution found in the first part produces, via the inverse Fourier transform, a classical solution for the system. Moreover, we prove that this solution is analytic in space and positive time.

## ACKNOWLEDGMENTS

First, I would like to thank my advisor Dr. Ciprian Foias who guided me during the last 4 years.

I also thank Dr. Carl Percy for his help and support since I came to Texas A&M University.

I am thankful to Dr. Lee Panetta for the joint work done on the model for my dissertation.

Also, I would like to thank Dr. Emil Straube for serving on my Ph.D. committee.

## TABLE OF CONTENTS

CHAPTER		Page
I	INTRODUCTION . . . . .	1
II	PRELIMINARIES . . . . .	5
	A. The model system . . . . .	5
	B. The wave-vector formulation . . . . .	10
III	WEAK SOLUTIONS . . . . .	13
	A. Galerkin approximations . . . . .	13
	B. Existence of weak solutions . . . . .	19
	C. Uniqueness of weak solutions . . . . .	28
IV	REGULARITY . . . . .	35
	A. Estimates for the Galerkin approximations . . . . .	35
	B. Time and space analyticity . . . . .	43
V	CONCLUSION . . . . .	47
	REFERENCES . . . . .	48
	VITA . . . . .	51

## CHAPTER I

## INTRODUCTION

Among several challenging aspects of weather prediction, one recognized very early was the large range of time and space scales involved if attempts are based on fundamental equations of continuum mechanics. “Weather” here refers to motions of relatively low frequency when compared with sound or gravity waves. Pioneering attempts [3], [4] with the first computers to predict extra-tropical weather patterns on spatial scales of order 1000 km used a series of observationally motivated approximations to derive a system of equations which “filtered” out relatively high frequency motions, thereby substantially reducing the range of timescales and easing the computational burden to the point where the goal of a useful forecast came within reach. The assumptions and approximations, now collectively called quasigeostrophic theory, placed special emphasis on observations that the evolution of the horizontal velocity and pressure gradient fields appeared to nearly preserve a “geostrophic” balance between Coriolis and pressure gradients forces, on large space scales and time scales exceeding a day. While computational technology now allows forecasts using equations derived under less restrictive assumptions, and the theory is now but one of a class based on geophysically relevant “balances” (see [15], [16]), quasigeostrophic theory and its numerical models remain of interest to meteorologists and oceanographers because they capture a number of physically important features while possessing a structure amenable to mathematical analysis and extensive numerical experimentation.

This dissertation concerns a simple quasigeostrophic model used by the author

---

The journal model is *IEEE Transactions on Automatic Control*.

of [13] to study a problem of pattern formation believed to be important in climate studies. The same model has been used for other purposes ([8], [12], [9], [11]). It is a coupled pair of 2D vorticity equations, in which the coupling term has the physical interpretation of a temperature field and is of central importance to its use. The system is forced by stimulation of a geophysically important instability present in the system. Numerical integrations indicate that the instability is typically arrested by nonlinearity, and all variables of interest come *eventually* to fluctuate irregularly about a *suitably defined* average value. Different variables take differing amounts of integration time to reveal this behavior; if this occurs for all variables of interest, the system is judged to be at “statistically steady state.” Statistically steady states are not always observed: for some choices of model parameters the system energy grows without bound and integrations must be stopped because of exponential overflow. No analysis has been done that explains this experience.

The model is typically used when many long-time numerical integrations of geophysical turbulence are required for purposes related to climate studies, purposes for which use of a climate model would be unnecessarily (and often prohibitively) demanding of computational time. Reliance on the model has been based on the convincing representation it gives of certain observed phenomena. Data from long numerical integrations are subjected to various averaging procedures to extract information about statistically steady states; these averages constitute the “climate” of the model, and sensitivity of these averages to parametric changes in the model is of interest to theories of climate behavior. No analytical guidance exists for the proper construction, or interpretation, of these averages.

*Our primary motivation in undertaking this study is to put on a firm mathematical ground the calculations in [13].* We expect that this analytic study will clarify the theoretical difficulties referred to in the preceding paragraph. Also, as the reader will

see in the next chapter, the model system sits in an interesting position between 2D and 3D Navier Stokes, so the problem may have some independent interest. The most closely related analytical work appears to be that of [1], which establishes finite-time existence and uniqueness for the quasigeostrophic model proposed by [2], with estimates of that finite time based on the size of initial data and the size of the forcing. (We mention recent work on a less closely related equation in the next chapter.)

The plan of the dissertation is as follows. In Section A of Chapter II we present the model in physical space variables, place it in context with recent related work, give some discussion of the forcing, and motivate an energy norm chosen for the subsequent analysis. In Section B of Chapter II we reformulate the model in wave-vector space, define relevant function spaces and norms, and present our notion of a weak solution. Chapter III follows an approach presented in [7] for study of the Navier-Stokes equations. In Section A of Chapter III we define a sequence of approximating Galerkin systems. Each system is a finite set of ODE's with quadratic nonlinearity, constructed by truncating the full wave-vector system at a wavenumber  $N$ . The long time existence of a classical solution (called there an  $N$ -solution) for each such system follows from the theory of ordinary differential equations. Key steps involve obtaining bound on energy injection by the forcing and certain algebraic observations that are analogues of integration-by-parts arguments. Section B of Chapter III then establishes (Theorem B.1) the existence of a weak solution by first verifying equicontinuity and uniform boundedness of the family of  $N$ -solutions, for a fixed wavenumber and time interval  $[0, T]$  of integer length  $T$ . Applications of the Arzela-Ascoli Theorem, diagonalizing over wavenumbers and  $T$ , produces a limit which is then shown to be a weak solution. Section C of Chapter III demonstrates the uniqueness of the weak solution. In each of these sections the main effort is to control the non-linear term: key steps in the proof of Theorem B.1 involve combinations of Holder's and



Ladyzhenskaya's inequalities with a Gronwall argument. In Chapter IV we show that our unique weak solution is in fact a classical solution. In addition we will prove that the mentioned solution is time and space analytic. Meantime, L. Panetta, E. Titi and M. Ziane have announced in [14] existence and uniqueness results (as well as a dissipativity property) for the strong solutions of our system under a more restrictive condition on the dissipative terms of the system.

## CHAPTER II

## PRELIMINARIES

## A. The model system

In this section we employ non-dimensionalizations that we do not discuss. Details can be found in [13], [15], [16]. Common to all versions of quasigeostrophic theory is the assumption that the horizontal velocity field has a streamfunction

$$\vec{u} = \nabla^\perp \psi, \quad (2.1)$$

(a non-dimensional form of geostrophic balance), together with an evolution equation for a quantity  $Q$

$$\frac{\partial Q}{\partial t} + \frac{\partial \psi}{\partial x_1} \frac{\partial Q}{\partial x_2} - \frac{\partial \psi}{\partial x_2} \frac{\partial Q}{\partial x_1} = F[\psi] + D[\psi]. \quad (2.2)$$

Here  $\nabla^\perp \psi = (-\frac{\partial \psi}{\partial x_2}, \frac{\partial \psi}{\partial x_1})$ ,  $(x_1, x_2)$  are horizontal coordinates,  $F$  and  $D$  are forcing and dissipation terms, and  $Q$  is related to  $\psi$  by a linear differential operator  $L$  in space variables

$$Q = L[\psi]. \quad (2.3)$$

Different choices for  $L$  give different versions of the theory: the general form is

$$L[\psi] = \beta x_2 + \Delta \psi + a(x_3) \frac{\partial}{\partial x_3} \left( b(x_3) \frac{\partial}{\partial x_3} \psi \right). \quad (2.4)$$

Here  $\Delta \equiv \frac{\partial}{\partial x_1^2} + \frac{\partial}{\partial x_2^2}$ ,  $\beta \geq 0$  is a constant and  $a(x_3)$ ,  $b(x_3)$  are functions related to a reference state density structure which is not explained by the theory. In this form  $Q$  is called the continuously stratified version of potential vorticity; in numerical models the vertical dependence is expressed in terms of fluid layers or modes, with appropriate treatments of the vertical derivatives. Thorough discussions from different points of view are given by [15] and [16].

The quantity

$$\tau \equiv \frac{\partial \psi}{\partial x_3} \quad (2.5)$$

appearing in (2.4) plays an important role in the theory: it is a representation of temperature (or buoyancy), and in view of (2.1), its horizontal gradient is related to vertical shear:

$$\frac{\partial \vec{u}}{\partial x_3} = \nabla^\perp \tau. \quad (2.6)$$

The presence of non-zero  $\tau$  also allows a form of vorticity generation not present in 2D flow. Versions of the theory that assume  $\tau \equiv 0$ , are called barotropic, and ones that do not are called baroclinic. (Note that barotropic versions with  $\beta = 0$  are simply 2D incompressible Navier Stokes equations.) For baroclinic versions, an equation for evolution of temperature on the boundary is included. Recent interest has in fact focused on the model that emerges when  $Q$  is assumed constant within the interior of the domain, and the evolution equation (2) is replaced by one governing boundary temperature field: this model, with  $L[\psi] = -(-\Delta)^{-1/2}\psi$  is called “surface geostrophic theory” and presents an interesting connection with the 3D Euler and Navier Stokes equations ([10], [6],[5]).

The model we study here uses the same vertical discretization of (2.2), (2.4) used in the early forecast attempts [4], but with the periodic boundary conditions motivated by [2] and with a special form of forcing that we describe briefly. Details are in [13], [8]. The model is defined in terms of a pair of streamfunctions  $(\psi_1, \psi_2)$ . In the physical interpretation, the flow given by  $\psi_1$  is at a greater altitude ( $x_3$  value) than that given by  $\psi_2$ . The analogue of the temperature variable (2.5) is

$$\hat{\psi} = \frac{\psi_1 - \psi_2}{2} \quad (2.7)$$

and there is a relation corresponding naturally to (2.6) between horizontal derivatives

of  $\hat{\psi}$  and vertical velocity differences. It is assumed that the flow takes place in the presence of an imposed, horizontally uniform temperature gradient, with a strength sufficient to excite an exponential instability at a number of scales. This gradient, like the reference stratification, cannot be altered by the flow's evolution. It is a stronger physical assumption than a simple imposition of a temperature drop across the domain. What actually appears in the equations is the vertical velocity difference related to the temperature gradient, which we denote in this section by  $2\hat{U}$ . The equations are

$$\frac{\partial q_1}{\partial t} + \frac{\partial \psi_1}{\partial x_1} \frac{\partial q_1}{\partial x_2} - \frac{\partial \psi_1}{\partial x_2} \frac{\partial q_1}{\partial x_1} = - \left[ 2\hat{U} \frac{\partial q_1}{\partial x_1} + (\beta + \hat{U}) \frac{\partial \psi_1}{\partial x_1} \right] - \nu(-\Delta)^p q_1 \quad (2.8)$$

$$\frac{\partial q_2}{\partial t} + \frac{\partial \psi_2}{\partial x_1} \frac{\partial q_2}{\partial x_2} - \frac{\partial \psi_2}{\partial x_2} \frac{\partial q_2}{\partial x_1} = - \left[ (\beta - \hat{U}) \frac{\partial \psi_2}{\partial x_1} \right] - \nu(-\Delta)^p q_2 - \kappa_M \Delta \psi_2. \quad (2.9)$$

Here the  $q_i$  are related to the  $\psi_i$  by

$$q_1 = \Delta \psi_1 - \hat{\psi} \quad (2.10)$$

$$q_2 = \Delta \psi_2 + \hat{\psi} \quad (2.11)$$

Solutions  $(\psi_1(x_1, x_2, t), \psi_2(x_1, x_2, t))$  to these equations are sought which are *periodic* on the domain  $\Omega \equiv [0, 2\pi\hat{L}]^2$ , where  $\hat{L}$  is a nondimensional real number. It is also assumed in [13] that such solutions have vanishing horizontal average. (Note: the velocity difference  $2\hat{U}$  is actually used to non-dimensionalize the equations in [13], [8], and so should be replaced by the value 1/2. We keep it, in this section alone, to mark terms related to the forcing and to show below how the imposed temperature gradient enters in the energy equation.)

The linear term involving  $\beta$  is a representation in this planar geometry of an effect of sphericity in planetary scale flow ([15], [16]); non-zero  $\beta$  is crucial to the formation of jets and introduces long timescales in the solutions [13]. (Getting estimates regarding

this effect is one of our aims.) The term involving  $\kappa_M$  is a parameterization of a boundary layer effect called Ekman pumping ([15], [16]). In the terms involving  $\nu$ , choices of  $p > 1$  are not as directly based on physical principles, and have more to do with expectations regarding energy and enstrophy cascades, and most often are made for computational convenience: they are designed to produce dissipative terms, and to concentrate the dissipation processes in simulations at the smallest small spatial scales included in the calculation. The hope is that this does not affect in any important way non-linear interactions at larger scales. When  $p > 1$  the value of  $\nu$  has only phenomenological justification. (We note that in [13] the high order Laplacian operator is not applied to the  $q_i$ , but instead to the  $\psi_i$ . The analysis we present for the equations here differs inessentially from what would be needed in that case. We choose this form of the equations because it the one being used in currently ongoing numerical studies, and it also agrees with [12], [9], and [11].)

A useful view of the roles of the terms on the right-hand sides of (2.8,2.9) comes from deriving the energy equation for the model. To do this, each layer equation is multiplied by its streamfunction, the equations are integrated horizontally, and the results are added. Using the notation (in this section alone)

$$\langle F \rangle = \int_{\Omega} F(x_1, x_2, t) dx_1 dx_2 \quad (2.12)$$

what results after several integrations by parts and uses of periodicity is

$$\frac{\partial E}{\partial t} = 2\hat{U} \langle \frac{\partial \tilde{\psi}}{\partial x_1} \hat{\psi} \rangle - \kappa_M \langle |\nabla \psi_2|^2 \rangle - \nu P \quad (2.13)$$

where  $\tilde{\psi} \equiv \frac{\psi_1 + \psi_2}{2}$  and the total energy  $E$  is defined by

$$E = \frac{\langle |\nabla \psi_1|^2 + |\nabla \psi_2|^2 \rangle}{2} + \langle \hat{\psi}^2 \rangle \quad (2.14)$$

is the sum of terms representing the kinetic energies in each layer and the model's

form of potential energy. The term  $P$  is positive definite:

$$P = \begin{cases} \langle (\Delta^{m+1}\psi_1)^2 + (\Delta^{m+1}\psi_2)^2 + 2|\nabla(\Delta^m\hat{\psi})|^2 \rangle & \text{if } p = 2m + 1 \\ \langle |\nabla(\Delta^m\psi_1)|^2 + |\nabla(\Delta^m\psi_2)|^2 + 2(\Delta^m\hat{\psi})^2 \rangle & \text{if } p = 2m \end{cases} \quad (2.15)$$

The only term not clearly sign-definite is that involving  $\hat{U}$  and is the energy source term for the model. It corresponds to the net flux of heat down the mean temperature gradient represented by the imposed vertical shear  $\hat{U}$ . This is as in models of thermal convection, where the energy generation for turbulent motions may also be related to the net down-gradient heat flux.

Notice that formal use of Cauchy-Schwartz and Poincare inequalities (recall the assumption of zero horizontal average for the  $\psi_i$ ) gives the crude estimate

$$\begin{aligned} \langle \frac{\partial \tilde{\psi}}{\partial x_1} \hat{\psi} \rangle &\leq \left( \langle |\nabla \tilde{\psi}|^2 \rangle \right)^{1/2} \left( \langle |\hat{\psi}|^2 \rangle \right)^{1/2} \\ &\leq \hat{L} \left( \langle |\nabla \tilde{\psi}| \rangle \right)^{1/2} \left( \langle |\nabla \hat{\psi}| \rangle \right)^{1/2} \\ &= \frac{\hat{L}}{2} \frac{\langle |\nabla \psi_1|^2 + |\nabla \psi_2|^2 \rangle}{2} \leq \frac{\hat{L}}{2} E. \end{aligned}$$

So from the energy equation (2.13) we get

$$\frac{\partial E}{\partial t} + \nu P + \kappa_M \langle |\nabla \psi_2|^2 \rangle \leq \hat{U} \hat{L} E. \quad (2.16)$$

An analogue of this argument will be used in Section A of Chapter III. Notice that no mention of the parameter  $\beta$  occurs in this estimate of the domain-integrated energy. (It does, however, appear in the equation for enstrophy equation). Nevertheless, experience with the model has indicated that the presence of the term  $\beta$  fundamentally affects the manner in which energy transfers within the domain occur, and the timescales present in numerical solutions.

We now drop further mention of the constant  $\hat{U}$ , using instead its value  $1/2$ .

### B. The wave-vector formulation

Let  $\widehat{L} > 0$ ,  $\Omega$  be the square  $[0, 2\pi\widehat{L}]^2 \subset \mathbb{R}^2$  and  $\alpha$  be an arbitrary nonnegative real number. We consider the equations (2.8)-(2.11) in  $\Omega$  with periodic boundary conditions:

$$\frac{\partial q_1}{\partial t} + \left( \frac{\partial \psi_1}{\partial x_1} \frac{\partial q_1}{\partial x_2} - \frac{\partial \psi_1}{\partial x_2} \frac{\partial q_1}{\partial x_1} \right) = -\frac{\partial q_1}{\partial x_1} - \left( \beta + \frac{1}{2} \right) \frac{\partial \psi_1}{\partial x_1} - \nu(-\Delta)^{1+\alpha} q_1 \quad (2.17)$$

$$\frac{\partial q_2}{\partial t} + \left( \frac{\partial \psi_2}{\partial x_1} \frac{\partial q_2}{\partial x_2} - \frac{\partial \psi_2}{\partial x_2} \frac{\partial q_2}{\partial x_1} \right) = -\kappa_M \Delta \psi_2 - \left( \beta - \frac{1}{2} \right) \frac{\partial \psi_2}{\partial x_1} - \nu(-\Delta)^{1+\alpha} q_2, \quad (2.18)$$

where

$$q_1 = \Delta \psi_1 - \frac{\psi_1 - \psi_2}{2} \text{ and } q_2 = \Delta \psi_2 + \frac{\psi_1 - \psi_2}{2}. \quad (2.19)$$

If  $\varphi$  is a  $2\pi\widehat{L}$ -periodic complex-valued scalar or vector function which is integrable over  $\Omega$ , we define its Fourier coefficients by

$$\varphi(\mathbf{k}) = \frac{1}{(2\pi\widehat{L})^2} \int_{\Omega} e^{-\frac{i}{L}\mathbf{k}\cdot\mathbf{x}} \varphi(\mathbf{x}) d\mathbf{x}, \quad \mathbf{k} \in \mathbb{Z}^2.$$

Its Fourier series will then be

$$\sum_{\mathbf{k} \in \mathbb{Z}^2} \varphi(\mathbf{k}) e^{\frac{i}{L}\mathbf{k}\cdot\mathbf{x}}.$$

Moreover, if  $\varphi = \varphi(\mathbf{x}, t) : \mathbb{R}^2 \times [0, T]$  (or  $[0, \infty)$ )  $\longrightarrow \mathbb{C}^d$ ,  $d \in \mathbb{N}$ , is  $2\pi\widehat{L}$ -periodic in the plane variable, we denote by  $\{\varphi(\mathbf{k}, t)\}_{\mathbf{k} \in \mathbb{Z}^2}$  the Fourier coefficients of  $\varphi(\cdot, t)$ .

By formally replacing in (2.17)-(2.19)  $\psi_j(\mathbf{x}, t)$  with  $\sum_{\mathbf{k} \in \mathbb{Z}^2} \psi_j(\mathbf{k}, t) e^{\frac{i}{L}\mathbf{k}\cdot\mathbf{x}}$  and  $q_j(\mathbf{x}, t)$  with  $\sum_{\mathbf{k} \in \mathbb{Z}^2} q_j(\mathbf{k}, t) e^{\frac{i}{L}\mathbf{k}\cdot\mathbf{x}}$ ,  $j = 1, 2$ , and identifying the corresponding Fourier coefficients we obtain the following equations:

$$\begin{aligned} & \frac{d}{dt} q_1(\mathbf{k}, t) + \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_1(\mathbf{h}, t) q_1(\mathbf{l}, t) \\ &= -\frac{i}{\widehat{L}} k_1 q_1(\mathbf{k}, t) - \left( \beta + \frac{1}{2} \right) \frac{i}{\widehat{L}} k_1 \psi_1(\mathbf{k}, t) - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} q_1(\mathbf{k}, t), \end{aligned} \quad (2.20)$$

$$\begin{aligned}
& \frac{d}{dt}q_2(\mathbf{k}, t) + \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2)\psi_2(\mathbf{h}, t)q_2(\mathbf{l}, t) \\
&= \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \psi_2(\mathbf{k}, t) - \left(\beta - \frac{1}{2}\right) \frac{i}{\widehat{L}} k_1 \psi_2(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} q_2(\mathbf{k}, t), \quad (2.21)
\end{aligned}$$

with

$$q_1(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \psi_1(\mathbf{k}, t) - \frac{\psi_1(\mathbf{k}, t) - \psi_2(\mathbf{k}, t)}{2}, \quad (2.22)$$

$$q_2(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \psi_2(\mathbf{k}, t) + \frac{\psi_1(\mathbf{k}, t) - \psi_2(\mathbf{k}, t)}{2}, \quad (2.23)$$

for every  $\mathbf{k} \in \mathbb{Z}^2$ . Since  $\psi_j(\mathbf{x}, t), j = 1, 2$ , are real-valued functions we have that

$$\psi_j(-\mathbf{k}, t) = \overline{\psi_j(\mathbf{k}, t)}, \mathbf{k} \in \mathbb{Z}^2, j = 1, 2, \quad (2.24)$$

where for a complex number  $z$  we denote by  $\bar{z}$  the complex conjugate of  $z$ . The equations (2.20)-(2.24) are called the wave-vectors formulation of the equations (2.17)-(2.19) for plane  $2\pi\widehat{L}$ -periodic solutions. Let

$$\begin{aligned}
\mathcal{K} := \{ \vec{\psi} = (\{\psi_1(\mathbf{k})\}_{\mathbf{k} \in \mathbb{Z}^2}, \{\psi_2(\mathbf{k})\}_{\mathbf{k} \in \mathbb{Z}^2}) : \psi_j(\mathbf{k}) \in \mathbb{C}, \psi_j(-\mathbf{k}) = \overline{\psi_j(\mathbf{k})}, j = 1, 2, \\
\mathbf{k} \in \mathbb{Z}^2, \psi_1(\mathbf{0}) + \psi_2(\mathbf{0}) = 0 \} \quad (2.25)
\end{aligned}$$

and

$$H := \left\{ \vec{\psi} \in \mathcal{K} : |\vec{\psi}|^2 := \sum_{\mathbf{k} \in \mathbb{Z}^2} E(\vec{\psi})(\mathbf{k}) < \infty \right\}, \quad (2.26)$$

where

$$E(\vec{\psi})(\mathbf{k}) := \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|\psi_1(\mathbf{k})|^2 + |\psi_2(\mathbf{k})|^2) + \frac{|\psi_1(\mathbf{k}) - \psi_2(\mathbf{k})|^2}{2}.$$

The space  $\mathcal{K}$  with the metric

$$d(\vec{\psi}, \vec{\varphi}) := \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \sum_{j=1}^2 \frac{|\psi_j(\mathbf{k}) - \varphi_j(\mathbf{k})|}{1 + |\psi_j(\mathbf{k}) - \varphi_j(\mathbf{k})|} \right) 2^{-|\mathbf{k}|^2}, \quad (2.27)$$

is a Frechet space, and  $H$  with the norm (as above) given by the scalar prod-



uct  $\langle \vec{\psi}, \vec{\varphi} \rangle := \sum_{\mathbf{k} \in \mathbb{Z}^2} \left[ \frac{|\mathbf{k}|^2}{\tilde{L}^2} \left( \psi_1(\mathbf{k}) \overline{\varphi_1(\mathbf{k})} + \psi_2(\mathbf{k}) \overline{\varphi_2(\mathbf{k})} \right) + \frac{(\psi_1(\mathbf{k}) - \psi_2(\mathbf{k}))(\overline{\varphi_1(\mathbf{k})} - \overline{\varphi_2(\mathbf{k})})}{2} \right]$  is a Hilbert space. For each  $\gamma > 0$  define

$$V_\gamma := \left\{ \vec{\psi} \in H : |\vec{\psi}|_\gamma^2 := \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \frac{|\mathbf{k}|}{\tilde{L}} \right)^{2\gamma} E(\vec{\psi})(\mathbf{k}) < \infty \right\}. \quad (2.28)$$

We denote by  $C([0, \infty), \mathcal{K})$  the space of all  $\mathcal{K}$ -valued continuous functions on  $[0, \infty)$ , where the continuity is with respect to the metric defined by (2.27). We also define the spaces  $L_{loc}^\infty([0, \infty), H)$  and  $L_{loc}^2([0, \infty), V_\gamma)$  by the following:

$$L_{loc}^\infty([0, \infty), H) = \left\{ \vec{\psi} : [0, \infty) \longrightarrow H : \text{ess-sup}_{0 \leq t \leq T} |\vec{\psi}(t)| < \infty, \text{ for every } T \in [0, \infty) \right\}$$

and

$$L_{loc}^2([0, \infty), V_\gamma) = \left\{ \vec{\psi} : [0, \infty) \longrightarrow V_\gamma : \int_0^T |\vec{\psi}(t)|_\gamma^2 dt < \infty, \text{ for every } T \in [0, \infty) \right\}.$$

Now we are ready to give the definition of a weak solution for (2.20)-(2.24) with initial data  $\vec{\psi}^0 \in H$ .

**Definition B.1.** Let  $\vec{\psi}^0 \in H$ . A  $H$ -valued function  $\vec{\psi}$  is called weak solution for the equations (2.20)-(2.24) with initial data  $\vec{\psi}^0$  if it has the following properties:

- 1)  $\vec{\psi} \in C([0, \infty), \mathcal{K}) \cap L_{loc}^\infty([0, \infty), H) \cap L_{loc}^2([0, \infty), V_{1+\alpha})$ ,
- 2)  $q_1(\mathbf{k}, t) = q_1(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\tilde{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) + \frac{i}{\tilde{L}} k_1 q_1(\mathbf{k}, \tau) + (\beta + \frac{1}{2}) \frac{i}{\tilde{L}} k_1 \psi_1(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\tilde{L}} \right)^{2(1+\alpha)} q_1(\mathbf{k}, \tau) \right\} d\tau$ ,  
 $q_2(\mathbf{k}, t) = q_2(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\tilde{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) - \kappa_M \frac{|\mathbf{k}|^2}{\tilde{L}^2} \psi_2(\mathbf{k}, \tau) + (\beta - \frac{1}{2}) \frac{i}{\tilde{L}} k_1 \psi_2(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\tilde{L}} \right)^{2(1+\alpha)} q_2(\mathbf{k}, \tau) \right\} d\tau$ ,  $\forall t \in [0, \infty)$ ,  $\forall \mathbf{k} \in \mathbb{Z}^2$ , where  $q_1(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\tilde{L}^2} \psi_1(\mathbf{k}, t) - \frac{\psi_1(\mathbf{k}, t) - \psi_2(\mathbf{k}, t)}{2}$ ,  $q_2(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\tilde{L}^2} \psi_2(\mathbf{k}, t) + \frac{\psi_1(\mathbf{k}, t) - \psi_2(\mathbf{k}, t)}{2}$ ,  $\forall \mathbf{k} \in \mathbb{Z}^2$ ,  
and

- 3)  $\psi_j(\mathbf{k}, 0) = \psi_j^0(\mathbf{k})$ ,  $j = 1, 2$ ,  $\forall \mathbf{k} \in \mathbb{Z}^2$ .

## CHAPTER III

## WEAK SOLUTIONS

## A. Galerkin approximations

In order to prove the existence of a weak solution for the equations (2.20)-(2.24) we will use the Galerkin approximations technique. Notice that

$$\begin{pmatrix} q_1(\mathbf{k}) \\ q_2(\mathbf{k}) \end{pmatrix} = \begin{pmatrix} -\left(\frac{|\mathbf{k}|^2}{L^2} + \frac{1}{2}\right) & \frac{1}{2} \\ \frac{1}{2} & -\left(\frac{|\mathbf{k}|^2}{L^2} + \frac{1}{2}\right) \end{pmatrix} \begin{pmatrix} \psi_1(\mathbf{k}) \\ \psi_2(\mathbf{k}) \end{pmatrix}.$$

Denote

$$A_{\mathbf{k}} = \begin{pmatrix} -\left(\frac{|\mathbf{k}|^2}{L^2} + \frac{1}{2}\right) & \frac{1}{2} \\ \frac{1}{2} & -\left(\frac{|\mathbf{k}|^2}{L^2} + \frac{1}{2}\right) \end{pmatrix}$$

and note that  $A_{\mathbf{k}}$  is invertible for every  $\mathbf{k} \neq \mathbf{0}$ . For every  $\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}$ , the equations (2.20) and (2.21) become

$$\begin{aligned} \frac{d}{dt} \begin{pmatrix} \psi_1(\mathbf{k}, t) \\ \psi_2(\mathbf{k}, t) \end{pmatrix} &= A_{\mathbf{k}}^{-1} \begin{pmatrix} -\frac{1}{L^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \psi_1(\mathbf{h}, t) q_1(\mathbf{l}, t) \\ -\frac{1}{L^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \psi_2(\mathbf{h}, t) q_2(\mathbf{l}, t) \end{pmatrix} + \\ A_{\mathbf{k}}^{-1} &\begin{pmatrix} -\frac{i}{L} k_1 q_1(\mathbf{k}, t) - (\beta + \frac{1}{2}) \frac{i}{L} k_1 \psi_1(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{L}\right)^{2(1+\alpha)} q_1(\mathbf{k}, t) \\ \kappa_M \frac{|\mathbf{k}|^2}{L^2} \psi_2(\mathbf{k}, t) - (\beta - \frac{1}{2}) \frac{i}{L} k_1 \psi_2(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{L}\right)^{2(1+\alpha)} q_2(\mathbf{k}, t) \end{pmatrix}. \end{aligned} \quad (3.1)$$

For  $N \in \mathbb{N}$  fixed we consider the system:

$$\begin{aligned} \frac{d}{dt} \begin{pmatrix} \varphi_1(\mathbf{k}, t) \\ \varphi_2(\mathbf{k}, t) \end{pmatrix} &= A_{\mathbf{k}}^{-1} \begin{pmatrix} -\frac{1}{L^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, t) r_1(\mathbf{l}, t) \\ -\frac{1}{L^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_2(\mathbf{h}, t) r_2(\mathbf{l}, t) \end{pmatrix} + \\ A_{\mathbf{k}}^{-1} &\begin{pmatrix} -\frac{i}{L} k_1 r_1(\mathbf{k}, t) - (\beta + \frac{1}{2}) \frac{i}{L} k_1 \varphi_1(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{L}\right)^{2(1+\alpha)} r_1(\mathbf{k}, t) \\ \kappa_M \frac{|\mathbf{k}|^2}{L^2} \varphi_2(\mathbf{k}, t) - (\beta - \frac{1}{2}) \frac{i}{L} k_1 \varphi_2(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{L}\right)^{2(1+\alpha)} r_2(\mathbf{k}, t) \end{pmatrix}, \end{aligned} \quad (3.2)$$

for  $\mathbf{k} \neq \mathbf{0}$ ,  $|\mathbf{k}| \leq N$ , and

$$\frac{d}{dt} \begin{pmatrix} \varphi_1(\mathbf{0}, t) \\ \varphi_2(\mathbf{0}, t) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (3.3)$$

where

$$r_1(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \varphi_1(\mathbf{k}, t) - \frac{\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)}{2}, \quad (3.4)$$

$$r_2(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \varphi_2(\mathbf{k}, t) + \frac{\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)}{2}. \quad (3.5)$$

We will be referring to equations (3.2) and (3.3) together with (3.4) and (3.5) as the  $N$ -system.

**Definition A.1.** Let  $\mathbb{Z}_N^2 = \{\mathbf{k} \in \mathbb{Z}^2 \mid |\mathbf{k}| \leq N\}$ . A  $N$ -solution is a family of functions  $\{(\varphi_1(\mathbf{k}, \cdot), \varphi_2(\mathbf{k}, \cdot))\}_{\mathbf{k} \in \mathbb{Z}_N^2}$  satisfying the  $N$ -system.

**Lemma A.1.** Let  $N \in \mathbb{N}$  and  $\overrightarrow{\psi}^0 \in H$ . Then

(a) there exist  $t_0 > 0$  and  $\{(\varphi_1(\mathbf{k}, \cdot), \varphi_2(\mathbf{k}, \cdot))\}_{\mathbf{k} \in \mathbb{Z}_N^2}$  such that

(i)  $\varphi_j(\mathbf{k}, \cdot) \in C^\infty([0, t_0]; \mathbb{C})$ ,

(ii)  $\{(\varphi_1(\mathbf{k}, \cdot), \varphi_2(\mathbf{k}, \cdot))\}_{\mathbf{k} \in \mathbb{Z}_N^2}$  is a  $N$ -solution with  $\varphi_j(\mathbf{k}, 0) = \psi_j^0(\mathbf{k})$ ,  $\forall |\mathbf{k}| \leq N$ ,  
 $j = 1, 2$ , and

(iii)  $\overline{\varphi_j(\mathbf{k}, t)} = \varphi_j(-\mathbf{k}, t)$ ,  $\forall |\mathbf{k}| \leq N, j = 1, 2$ ,

(b) for every  $T \in (0, \infty)$  with the property that the above solution exists on  $[0, T)$  there exists  $M > 0$  such that

$$|\varphi_j(\mathbf{k}, t)| \leq M, \forall t \in [0, T), \forall |\mathbf{k}| \leq N, j = 1, 2. \quad (3.6)$$

Moreover, the  $N$ -solution  $\{(\varphi_1(\mathbf{k}, \cdot), \varphi_2(\mathbf{k}, \cdot))\}_{\mathbf{k} \in \mathbb{Z}_N^2}$  with initial data  $\overrightarrow{\psi}^0$  is unique in the interval of existence.

*Proof.* Part (a) follows immediately from the classical theory of systems of ordinary differential equations and the fact that  $\{\overline{\varphi_j(\mathbf{k}, t)}\}_{|\mathbf{k}| \leq N, j=1,2}$  and  $\{\varphi_j(-\mathbf{k}, t)\}_{|\mathbf{k}| \leq N, j=1,2}$  are solutions for the same system of ODEs with the same initial data (since  $\overline{\psi_j^0(\mathbf{k})} = \psi_j^0(-\mathbf{k})$ , for every  $\mathbf{k} \in \mathbb{Z}^2$ ). For (b) we start by noticing that from (3.3) we have that

$$\varphi_j(\mathbf{0}, t) = \psi_j^0(\mathbf{0}), \forall t \in [0, T], j = 1, 2. \quad (3.7)$$

Using the equations (3.4) and (3.5) we also get

$$\begin{aligned} & \operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left( \left( \frac{d}{dt} r_1(\mathbf{k}, t) \right) \overline{\varphi_1(\mathbf{k}, t)} + \left( \frac{d}{dt} r_2(\mathbf{k}, t) \right) \overline{\varphi_2(\mathbf{k}, t)} \right) = \\ & \operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left\{ -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \left( \frac{d}{dt} \varphi_1(\mathbf{k}, t) \right) \overline{\varphi_1(\mathbf{k}, t)} - \frac{|\mathbf{k}|^2}{\widehat{L}^2} \left( \frac{d}{dt} \varphi_2(\mathbf{k}, t) \right) \overline{\varphi_2(\mathbf{k}, t)} - \right. \\ & \left. \left( \frac{d}{dt} \left( \frac{\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)}{2} \right) \right) \overline{\varphi_1(\mathbf{k}, t)} + \left( \frac{d}{dt} \left( \frac{\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)}{2} \right) \right) \overline{\varphi_2(\mathbf{k}, t)} \right\} = \\ & -\frac{1}{2} \frac{d}{dt} \sum_{|\mathbf{k}| \leq N} \left\{ \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|\varphi_1(\mathbf{k}, t)|^2 + |\varphi_2(\mathbf{k}, t)|^2) + \frac{|\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)|^2}{2} \right\}. \quad (3.8) \end{aligned}$$

We will extend a  $N$ -solution in a natural way to a function  $\vec{\psi}_N$  such that for every  $t$  in the interval of existence of our  $N$ -solution we have  $\vec{\psi}_N(t) \in \mathcal{K}$ , namely:

$$\psi_N(\mathbf{k}, t) = \varphi(\mathbf{k}, t), \text{ if } |\mathbf{k}| \leq N \text{ and } \psi_N(\mathbf{k}, t) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \text{ if } |\mathbf{k}| > N. \quad (3.9)$$

For  $\vec{\psi}_N$  we then obtain from (3.8) that

$$\frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 = -\operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left( \frac{d}{dt} r_1(\mathbf{k}, t) \right) \overline{\varphi_1(\mathbf{k}, t)} + \left( \frac{d}{dt} r_2(\mathbf{k}, t) \right) \overline{\varphi_2(\mathbf{k}, t)},$$

and using (3.2) we get

$$\frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 = -\operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left\{ -\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, t) r_1(\mathbf{l}, t) \right\}$$

$$\begin{aligned}
& -\frac{i}{\widehat{L}}k_1r_1(\mathbf{k}, t) - (\beta + \frac{1}{2})\frac{i}{\widehat{L}}k_1\varphi_1(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} r_1(\mathbf{k}, t)\overline{\varphi_1(\mathbf{k}, t)} \\
& + \left(-\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2)\varphi_2(\mathbf{h}, t)r_2(\mathbf{l}, t) + \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2}\varphi_2(\mathbf{k}, t) \right. \\
& \left. - (\beta - \frac{1}{2})\frac{i}{\widehat{L}}k_1\varphi_2(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} r_2(\mathbf{k}, t)\overline{\varphi_2(\mathbf{k}, t)}\right),
\end{aligned}$$

which implies that

$$\begin{aligned}
\frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 &= \frac{1}{\widehat{L}^2} \operatorname{Re} \sum_{|\mathbf{k}| \leq N} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2) (\varphi_1(\mathbf{h}, t)r_1(\mathbf{l}, t)\overline{\varphi_1(\mathbf{k}, t)} \\
& + \varphi_2(\mathbf{h}, t)r_2(\mathbf{l}, t)\overline{\varphi_2(\mathbf{k}, t)}) + \operatorname{Re} \left( \frac{i}{\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1r_1(\mathbf{k}, t)\overline{\varphi_1(\mathbf{k}, t)} \right) \\
& - \kappa_M \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\varphi_2(\mathbf{k}, t)|^2 \\
& + \nu \operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} (r_1(\mathbf{k}, t)\overline{\varphi_1(\mathbf{k}, t)} + r_2(\mathbf{k}, t)\overline{\varphi_2(\mathbf{k}, t)}). \tag{3.10}
\end{aligned}$$

Using (iii) from part (a) of Lemma A.1 we deduce that

$$\begin{aligned}
S_1 &: = \sum_{|\mathbf{k}| \leq N} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2) \varphi_1(\mathbf{h}, t)r_1(\mathbf{l}, t)\overline{\varphi_1(\mathbf{k}, t)} \\
&= \sum_{\mathbf{h}+\mathbf{k}=\mathbf{0}, |\mathbf{h}|, |\mathbf{l}|, |\mathbf{k}| \leq N} (h_2l_1 - h_1l_2) \varphi_1(\mathbf{h}, t)r_1(\mathbf{l}, t)\varphi_1(\mathbf{k}, t),
\end{aligned}$$

and after we interchange  $\mathbf{h}$  with  $\mathbf{k}$  we obtain

$$\begin{aligned}
S_1 &= \sum_{\mathbf{h}+\mathbf{k}=\mathbf{0}, |\mathbf{h}|, |\mathbf{l}|, |\mathbf{k}| \leq N} (k_2l_1 - k_1l_2) \varphi_1(\mathbf{k}, t)r_1(\mathbf{l}, t)\varphi_1(\mathbf{h}, t) \\
&= \sum_{\mathbf{h}+\mathbf{k}=\mathbf{0}, |\mathbf{h}|, |\mathbf{l}|, |\mathbf{k}| \leq N} ((-h_2 - l_2)l_1 - (-h_1 - l_1)l_2) \varphi_1(\mathbf{k}, t)r_1(\mathbf{l}, t)\varphi_1(\mathbf{h}, t) \\
&= \sum_{\mathbf{h}+\mathbf{k}=\mathbf{0}, |\mathbf{h}|, |\mathbf{l}|, |\mathbf{k}| \leq N} (-h_2l_1 + h_1l_2) \varphi_1(\mathbf{k}, t)r_1(\mathbf{l}, t)\varphi_1(\mathbf{h}, t) = -S_1.
\end{aligned}$$

Therefore,  $S_1 = 0$ . Similarly,

$$S_2 := \sum_{|\mathbf{k}| \leq N} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_2(\mathbf{h}, t) r_2(\mathbf{l}, t) \overline{\varphi_2(\mathbf{k}, t)} = 0.$$

Thus, (3.10) becomes

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 + \kappa_M \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\varphi_2(\mathbf{k}, t)|^2 &= \operatorname{Re} \left( \frac{i}{\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1 r_1(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right) \\ + \nu \operatorname{Re} \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} &(r_1(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} + r_2(\mathbf{k}, t) \overline{\varphi_2(\mathbf{k}, t)}). \end{aligned} \quad (3.11)$$

From (3.4) and (3.5) we easily get that

$$r_1(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} + r_2(\mathbf{k}, t) \overline{\varphi_2(\mathbf{k}, t)} = -E(\vec{\varphi})(\mathbf{k}) \quad (3.12)$$

and

$$\begin{aligned} S_3 &: = \operatorname{Re} \left( \frac{i}{\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1 r_1(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right) \\ &= \operatorname{Re} \frac{i}{\widehat{L}} \left( - \sum_{|\mathbf{k}| \leq N} k_1 \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + \frac{1}{2} \right) |\varphi_1(\mathbf{k}, t)|^2 + \frac{1}{2} \sum_{|\mathbf{k}| \leq N} k_1 \varphi_2(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right) \\ &= \operatorname{Re} \left( \frac{i}{2\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1 \varphi_2(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right). \end{aligned} \quad (3.13)$$

Using (3.12) and (3.13), (3.11) becomes

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 + \kappa_M \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\varphi_2(\mathbf{k}, t)|^2 &= \operatorname{Re} \left( \frac{i}{2\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1 \varphi_2(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right) \\ - \nu \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} &E(\vec{\psi}_N)(\mathbf{k}). \end{aligned} \quad (3.14)$$

Next we notice that

$$\left| \frac{i}{2\widehat{L}} \sum_{|\mathbf{k}| \leq N} k_1 \varphi_2(\mathbf{k}, t) \overline{\varphi_1(\mathbf{k}, t)} \right| \leq \frac{1}{2\widehat{L}} \sum_{|\mathbf{k}| \leq N} |\mathbf{k}| |\varphi_2(\mathbf{k}, t)| |\varphi_1(\mathbf{k}, t)|$$

$$\begin{aligned}
&\leq \frac{\widehat{L}}{2} \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} |\varphi_1(\mathbf{k}, t)| \right) \left( \frac{|\mathbf{k}|}{\widehat{L}} |\varphi_2(\mathbf{k}, t)| \right) \\
&\leq \frac{\widehat{L}}{4} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|\varphi_1(\mathbf{k}, t)|^2 + |\varphi_2(\mathbf{k}, t)|^2) \leq \frac{\widehat{L}}{4} |\vec{\psi}_N(t)|^2.
\end{aligned} \tag{3.15}$$

From (3.14) and (3.15) we obtain

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 + \kappa_M \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\varphi_2(\mathbf{k}, t)|^2 + \nu \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{\psi}_N)(\mathbf{k}) \\
&\leq \frac{\widehat{L}}{4} |\vec{\psi}_N(t)|^2, \quad \forall t \in [0, T].
\end{aligned} \tag{3.16}$$

Therefore,  $\frac{1}{2} \frac{d}{dt} |\vec{\psi}_N(t)|^2 \leq \frac{\widehat{L}}{4} |\vec{\psi}_N(t)|^2$ ,  $\forall t \in [0, T)$  which implies that  $|\vec{\psi}_N(t)|^2 \leq e^{\frac{\widehat{L}}{2}t} |\vec{\psi}^0|^2 \leq e^{\frac{\widehat{L}}{2}T} |\vec{\psi}^0|^2$ ,  $\forall t \in [0, T)$ . From here and relation (3.7) we easily get that  $\exists M > 0$  such that  $|\varphi_j(\mathbf{k}, t)| \leq M$ ,  $\forall t \in [0, T), \forall |\mathbf{k}| \leq N, j = 1, 2$ .  $\square$

From Lemma A.1 (b) and the classical theory of ODE's it follows immediately that

**Corollary A.1.** *For given  $\vec{\psi}^0 \in H$  there exists a unique  $N$ -solution with initial data  $\vec{\psi}^0$  defined on  $[0, \infty)$ .*

**Corollary A.2.** *The function  $\vec{\psi}_N$  defined by (36) belongs to  $C([0, \infty), \mathcal{K})$ .*

*Proof.* Recall that  $\overline{\varphi_j(\mathbf{k}, t)} = \varphi_j(-\mathbf{k}, t)$ ,  $\forall |\mathbf{k}| \leq N, j = 1, 2$ , and notice also that  $\varphi_1(\mathbf{0}, t) + \varphi_2(\mathbf{0}, t) = \psi_1^0(\mathbf{0}) + \psi_2^0(\mathbf{0}) = 0$ ,  $\forall t \in [0, \infty)$ . Therefore,

$$\vec{\psi}_N(t) \in \mathcal{K}, \quad \forall t \in [0, \infty).$$

Since  $\varphi_j(\mathbf{k}, \cdot)$  is continuous on  $[0, \infty)$ ,  $\forall |\mathbf{k}| \leq N, j = 1, 2$ , we see that  $\vec{\psi}_N(\cdot) \in C([0, \infty), \mathcal{K})$ .  $\square$

## B. Existence of weak solutions

Applying the process from Section A of this chapter for every  $N \in \mathbb{N}$  we get the sequence  $\{\vec{\psi}_N(\cdot)\}_{N \in \mathbb{N}} \subset C([0, \infty), \mathcal{K})$ . On the space  $C([0, \infty), \mathcal{K})$  we define the metric

$$\text{dist}(\vec{\psi}(\cdot), \vec{\varphi}(\cdot)) = \sum_{T=1,2,\dots} \frac{1}{2^T} \frac{\sup\{d(\vec{\psi}(t), \vec{\varphi}(t)) : 0 \leq t \leq T\}}{1 + \sup\{d(\vec{\psi}(t), \vec{\varphi}(t)) : 0 \leq t \leq T\}}.$$

**Remark B.1.** *The convergence  $\text{dist}(\vec{\varphi}_m(\cdot), \vec{\varphi}(\cdot)) \rightarrow 0$  as  $m \rightarrow \infty$  is equivalent to, for every  $\mathbf{k} \in \mathbb{Z}^2$  and  $t_0 \in [0, \infty)$ ,  $\varphi_{m,j}(\mathbf{k}, t) \rightarrow \varphi_j(\mathbf{k}, t)$  uniformly on  $[0, t_0]$ ,  $j = 1, 2$ .*

The proof of the existence of weak solutions for (2.20)-(2.24) with initial data  $\vec{\psi}^0 \in H$  will be split in two parts. First we prove that there exists a subsequence  $\{\vec{\psi}_{N_p}(\cdot)\}_{p \in \mathbb{N}}$  of  $\{\vec{\psi}_N(\cdot)\}_{N \in \mathbb{N}}$  converging to some  $\vec{\psi}(\cdot)$  in  $C([0, \infty), \mathcal{K})$ . After that we will show that the limit  $\vec{\psi}(\cdot)$  is our desired weak solution. The first part is covered by the following lemma.

**Lemma B.1.** *There exist a subsequence  $\{\vec{\psi}_{N_p}(\cdot)\}_{p \in \mathbb{N}}$  of  $\{\vec{\psi}_N(\cdot)\}_{N \in \mathbb{N}}$  and a function  $\vec{\psi}(\cdot) \in C([0, \infty), \mathcal{K})$  such that  $\lim_{p \rightarrow \infty} \text{dist}(\vec{\psi}_{N_p}, \vec{\psi}) = 0$ .*

*Proof.* Let  $T, N \in \mathbb{N}$  be fixed. Using (3.2) we can write

$$\begin{aligned} \frac{d}{dt} (r_1(\mathbf{k}, t) + r_2(\mathbf{k}, t)) &= -\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) (\varphi_1(\mathbf{h}, t) r_1(\mathbf{l}, t) \\ &+ \varphi_2(\mathbf{h}, t) r_2(\mathbf{l}, t)) - \frac{i}{\widehat{L}} k_1 r_1(\mathbf{k}, t) - \left(\beta + \frac{1}{2}\right) \frac{i}{\widehat{L}} k_1 \varphi_1(\mathbf{k}, t) - \left(\beta - \frac{1}{2}\right) \frac{i}{\widehat{L}} k_1 \varphi_2(\mathbf{k}, t) \\ &+ \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \varphi_2(\mathbf{k}, t) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} (r_1(\mathbf{k}, t) + r_2(\mathbf{k}, t)). \end{aligned} \quad (3.17)$$

Next we add (3.4) with (3.5) and we divide by  $-\frac{|\mathbf{k}|^2}{\widehat{L}^2}$ . With the use of (3.17) we get

$$\frac{d}{dt} (\varphi_1(\mathbf{k}, t) + \varphi_2(\mathbf{k}, t)) = \frac{1}{|\mathbf{k}|^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) (\varphi_1(\mathbf{h}, t) r_1(\mathbf{l}, t)$$



$$\begin{aligned}
& +\varphi_2(\mathbf{h}, t)r_2(\mathbf{l}, t) + \frac{i\widehat{L}k_1}{|\mathbf{k}|^2}r_1(\mathbf{k}, t) + \left(\beta + \frac{1}{2}\right)\frac{i\widehat{L}k_1}{|\mathbf{k}|^2}\varphi_1(\mathbf{k}, t) \\
& + \left(\beta - \frac{1}{2}\right)\frac{i\widehat{L}k_1}{|\mathbf{k}|^2}\varphi_2(\mathbf{k}, t) - \kappa_M\varphi_2(\mathbf{k}, t) - \nu\left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)}(\varphi_1(\mathbf{k}, t) + \varphi_2(\mathbf{k}, t)). \quad (3.18)
\end{aligned}$$

Now define  $\widetilde{\varphi}(\mathbf{k}, t) := \varphi_1(\mathbf{k}, t) + \varphi_2(\mathbf{k}, t)$  and  $\widehat{\varphi}(\mathbf{k}, t) := \varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)$ . For  $s, t \in [0, T]$ ,  $s < t$ , from (3.18) we obtain

$$\begin{aligned}
|\widetilde{\varphi}(\mathbf{k}, t) - \widetilde{\varphi}(\mathbf{k}, s)| & \leq \frac{1}{|\mathbf{k}|^2} \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} \int_s^t (h_2l_1 - h_1l_2)(\varphi_1(\mathbf{h}, \tau)r_1(\mathbf{l}, \tau) + \right. \\
& \varphi_2(\mathbf{h}, \tau)r_2(\mathbf{l}, \tau))d\tau \left. + \frac{\widehat{L}}{|\mathbf{k}|} \int_s^t |r_1(\mathbf{k}, \tau)|d\tau + \left(\beta + \frac{1}{2}\right)\frac{\widehat{L}}{|\mathbf{k}|} \int_s^t |\varphi_1(\mathbf{k}, \tau)|d\tau + \right. \\
& \left. \left(\beta - \frac{1}{2}\right)\frac{\widehat{L}}{|\mathbf{k}|} \int_s^t |\varphi_2(\mathbf{k}, \tau)|d\tau + \kappa_M \int_s^t |\varphi_2(\mathbf{k}, \tau)|d\tau + \right. \\
& \left. \nu\left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} \int_s^t (|\varphi_1(\mathbf{k}, \tau)| + |\varphi_2(\mathbf{k}, \tau)|)d\tau. \quad (3.19)
\end{aligned}$$

Recall that we proved that

$$\begin{aligned}
|\vec{\psi}_N(t)|^2 & = \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|\varphi_1(\mathbf{k}, t)|^2 + |\varphi_2(\mathbf{k}, t)|^2) + \frac{|\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)|^2}{2} \right) \\
& \leq e^{\frac{LT}{2}} |\vec{\psi}^0|^2, \quad (3.20)
\end{aligned}$$

for every  $t \in [0, T]$ . Therefore, for  $\mathbf{k} \neq \mathbf{0}$ , we have

$$|\varphi_j(\mathbf{k}, t)| \leq \frac{\widehat{L}}{|\mathbf{k}|} e^{\frac{LT}{4}} |\vec{\psi}^0|, \quad \forall t \in [0, T], \quad j = 1, 2.$$

Using this we see that all the integrals from the right hand side of (3.19) except the first one are bounded by  $c_1(t - s)$  where  $c_1$  is some positive constant which doesn't depend on  $N$ . From (3.4) we obtain

$$\sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2l_1 - h_1l_2)\varphi_1(\mathbf{h}, \tau)r_1(\mathbf{l}, \tau) =$$

$$\begin{aligned}
& - \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \frac{|\mathbf{l}|^2}{\widehat{L}^2} \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) \\
& - \frac{1}{2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) (\varphi_1(\mathbf{l}, \tau) - \varphi_2(\mathbf{l}, \tau)). \tag{3.21}
\end{aligned}$$

We notice that

$$\begin{aligned}
S_4 & : = \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \frac{|\mathbf{l}|^2}{\widehat{L}^2} \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) \\
& = \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \frac{\mathbf{l} \cdot (\mathbf{k} - \mathbf{h})}{\widehat{L}^2} \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) \\
& = \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) (\mathbf{l} \cdot \mathbf{k}) \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) - \\
& \quad \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) (\mathbf{l} \cdot \mathbf{h}) \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau). \tag{3.22}
\end{aligned}$$

By interchanging  $\mathbf{h}$  with  $\mathbf{l}$  in the last sum of (3.22) we get that the indicated sum is 0, and, therefore

$$\begin{aligned}
S_4 & = \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) (\mathbf{l} \cdot \mathbf{k}) \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) \\
& = \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2(k_1 - h_1) - h_1(k_2 - h_2)) (\mathbf{l} \cdot \mathbf{k}) \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau) \\
& = \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 k_1 - h_1 k_2) (\mathbf{l} \cdot \mathbf{k}) \varphi_1(\mathbf{h}, \tau) \varphi_1(\mathbf{l}, \tau). \tag{3.23}
\end{aligned}$$

From (3.21) and (3.23) we deduce that

$$\begin{aligned}
& \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) \right| \leq \\
& \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} |\mathbf{h}| |\mathbf{k}|^2 |\mathbf{l}| |\varphi_1(\mathbf{h}, \tau)| |\varphi_1(\mathbf{l}, \tau)| \\
& + \frac{\widehat{L}^2}{2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} \frac{|\mathbf{h}|}{\widehat{L}} |\varphi_1(\mathbf{h}, \tau)| \frac{|\mathbf{l}|}{\widehat{L}} |\varphi_2(\mathbf{l}, \tau)|,
\end{aligned}$$

and using Cauchy-Schwartz inequality we get that the left term in the above inequality is less or equal then

$$\begin{aligned}
& |\mathbf{k}|^2 \left( \sum_{|\mathbf{h}| \leq N} \frac{|\mathbf{h}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{h}, \tau)|^2 \right)^{\frac{1}{2}} \left( \sum_{|\mathbf{l}| \leq N} \frac{|\mathbf{l}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{l}, \tau)|^2 \right)^{\frac{1}{2}} \\
& + \frac{\widehat{L}^2}{2} \left( \sum_{|\mathbf{h}| \leq N} \frac{|\mathbf{h}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{h}, \tau)|^2 \right)^{\frac{1}{2}} \left( \sum_{|\mathbf{l}| \leq N} \frac{|\mathbf{l}|^2}{\widehat{L}^2} |\varphi_2(\mathbf{l}, \tau)|^2 \right)^{\frac{1}{2}} \leq \\
& \left( |\mathbf{k}|^2 + \frac{\widehat{L}^2}{2} \right) |\vec{\psi}_N(\tau)|^2 \leq \left( |\mathbf{k}|^2 + \frac{\widehat{L}^2}{2} \right) e^{\frac{LT}{2}} |\vec{\psi}^0|^2, \forall \tau \in [0, T]. \tag{3.24}
\end{aligned}$$

Similarly,

$$\left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}, |\mathbf{h}|, |\mathbf{l}| \leq N} (h_2 l_1 - h_1 l_2) \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau) \right| \leq c_2, \forall \tau \in [0, T], \tag{3.25}$$

where  $c_2$  is a positive constant which doesn't depend on  $N$ . Thus, there exists  $\tilde{c} > 0$  which doesn't depend on  $N$  such that

$$|\tilde{\varphi}(\mathbf{k}, t) - \tilde{\varphi}(\mathbf{k}, s)| \leq \tilde{c}(t - s), \forall t, s \in [0, T], s < t. \tag{3.26}$$

Using the same idea we can easily get that  $\widehat{c} > 0$  such that

$$|\widehat{\varphi}(\mathbf{k}, t) - \widehat{\varphi}(\mathbf{k}, s)| \leq \widehat{c}(t - s), \forall t, s \in [0, T], s < t. \tag{3.27}$$

From (3.26) and (3.27) we obtain that there exists  $c > 0$  which doesn't depend on  $N$  such that

$$|\psi_{N,j}(\mathbf{k}, t) - \psi_{N,j}(\mathbf{k}, s)| \leq c(t - s), \forall t, s \in [0, T], j = 1, 2. \tag{3.28}$$

Notice that we can choose  $c$  such that the following is also true

$$|\psi_{N,j}(\mathbf{k}, 0)| = |\psi_j^0(\mathbf{k})| \leq c |\vec{\psi}^0|. \tag{3.29}$$

The relations (3.28) and (3.29) allow us to apply Arzela-Ascoli Theorem for the se-

quence  $\{\psi_{N,j}(\mathbf{k}, \cdot)\}_{N \in \mathbb{N}}$ . We get that for  $T = 1$  and a fixed  $\mathbf{k} \in \mathbb{Z}^2$  there exist a subsequence  $\{\psi_{N_h,j}(\mathbf{k}, \cdot)\}_{h \in \mathbb{N}}$  of  $\{\psi_{N,j}(\mathbf{k}, \cdot)\}_{N \in \mathbb{N}}$  and a function  $\psi_{1,j}(\mathbf{k}, \cdot) \in C([0, 1], \mathbb{C})$  such that  $\{\psi_{N_h,j}(\mathbf{k}, \cdot)\}_{h \in \mathbb{N}}$  converges to  $\psi_{1,j}(\mathbf{k}, \cdot)$  uniformly on  $[0, 1]$ . By applying Cantor's diagonal method for  $\mathbf{k} \in \mathbb{Z}^2$  (written as a sequence) we prove the existence of a subsequence of  $\{\vec{\psi}_N(\cdot)\}_{N \in \mathbb{N}}$  which converges to a function  $\vec{\psi}_1(\cdot)$  in  $C([0, 1], \mathcal{K})$ . For this subsequence we repeat the above argument with  $T = 2$  to get another subsequence which converges to a function  $\vec{\psi}_2(\cdot)$  in  $C([0, 2], \mathcal{K})$ . We continue with  $T = 3, 4, \dots$ , and we apply Cantor's diagonal method to obtain that there exist a subsequence  $\{\vec{\psi}_{N_p}(\cdot)\}_{p \in \mathbb{N}}$  of  $\{\vec{\psi}_N(\cdot)\}_{N \in \mathbb{N}}$  and  $\vec{\psi}(\cdot) \in C([0, \infty), \mathcal{K})$  such that  $\{\vec{\psi}_{N_p}(\cdot)\}_{p \in \mathbb{N}}$  converges to  $\vec{\psi}(\cdot)$  in  $C([0, \infty), \mathcal{K})$ .  $\square$

Now we are ready to prove the main theorem of this section.

**Theorem B.1.** *The function  $\vec{\psi}$  provided by Lemma B.1 is a weak solution for (2.20)-(2.24) with initial data  $\vec{\psi}^0$ .*

*Proof.* Since  $\{\vec{\psi}_{N_p}\}_{p \in \mathbb{N}}$  converges to  $\vec{\psi}$  in  $C([0, \infty), \mathcal{K})$  we get that for every  $T \in [0, \infty)$

$$\psi_{N_p,j}(\mathbf{k}, t) \rightarrow \psi_j(\mathbf{k}, t) \text{ uniformly on } [0, T], j = 1, 2. \quad (3.30)$$

If  $\vec{\theta} \in \mathcal{K}$  and  $M \in \mathbb{N}$  define  $P_M \vec{\theta} \in \mathcal{K}$  by

$$(P_M \vec{\theta})(\mathbf{k}) = \theta(\mathbf{k}) \text{ if } |\mathbf{k}| \leq M, \text{ and } (P_M \vec{\theta})(\mathbf{k}) = \mathbf{0}, \text{ if } |\mathbf{k}| > M.$$

Then, if  $M \in \mathbb{N}$  and  $N_p \geq M$  we have

$$|P_M \vec{\psi}_{N_p}(t)| \leq |\vec{\psi}_{N_p}(t)| \leq e^{\frac{LT}{4}} |\vec{\psi}^0|, \forall t \in [0, T].$$

Letting  $p \rightarrow \infty$  and using (3.30) we obtain

$$|P_M \vec{\psi}(t)| \leq e^{\frac{LT}{4}} |\vec{\psi}^0|, \forall t \in [0, T], \forall M \in \mathbb{N},$$

and by letting  $M \rightarrow \infty$  we get

$$|\vec{\psi}(t)| \leq e^{\frac{LT}{4}} |\vec{\psi}^0|, \forall t \in [0, T], \quad (3.31)$$

which shows that  $\vec{\psi} \in L_{loc}^\infty([0, \infty), H)$ . By integrating (3.16) we deduce that

$$\begin{aligned} \nu \int_0^T |\vec{\psi}_N(t)|_{1+\alpha}^2 dt &= \nu \int_0^T \sum_{|\mathbf{k}| \leq N} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{\psi}_N)(\mathbf{k}) dt \\ &\leq \frac{1}{2} |\vec{\psi}^0|^2 + \frac{\widehat{L}}{4} \int_0^T |\vec{\psi}_N(t)|^2 dt \leq \left( \frac{1}{2} + \frac{\widehat{L}T}{4} e^{\frac{LT}{2}} \right) |\vec{\psi}^0|^2. \end{aligned} \quad (3.32)$$

If  $M \in \mathbb{N}$  and  $N_p \geq M$  we have

$$\begin{aligned} &\nu \int_0^T \sum_{|\mathbf{k}| \leq M} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{\psi}_{N_p})(\mathbf{k}) dt \\ &\leq \nu \int_0^T \sum_{|\mathbf{k}| \leq N_p} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{\psi}_{N_p})(\mathbf{k}) dt \\ &\leq \left( \frac{1}{2} + \frac{\widehat{L}T}{4} e^{\frac{LT}{2}} \right) |\vec{\psi}^0|^2. \end{aligned}$$

Therefore, by using (3.30), if  $p \rightarrow \infty$  we obtain

$$\nu \int_0^T \sum_{|\mathbf{k}| \leq M} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{\psi}_{N_p})(\mathbf{k}) dt \leq \left( \frac{1}{2} + \frac{\widehat{L}T}{4} e^{\frac{LT}{2}} \right) |\vec{\psi}^0|^2, \forall M \in \mathbb{N}.$$

Now we apply Beppo-Levi Theorem to get that

$$\int_0^T |\vec{\psi}(t)|_{1+\alpha}^2 dt \leq \frac{1}{\nu} \left( \frac{1}{2} + \frac{\widehat{L}T}{4} e^{\frac{LT}{2}} \right) |\vec{\psi}^0|^2, \forall T \in [0, \infty), \quad (3.33)$$

which proves that  $\vec{\psi} \in L_{loc}^2([0, \infty), V_{1+\alpha})$ . Thus  $\vec{\psi}$  satisfies the condition 1) from Definition B.1 of Chapter II. From (3.2) and (3.3) we easily get that

$$\begin{aligned} q_{N_p,1}(\mathbf{k}, t) &= q_{N_p,1}(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_{N_p,1}(\mathbf{h}, \tau) q_{N_p,1}(\mathbf{l}, \tau) \right. \\ &\quad \left. + \frac{i}{\widehat{L}} k_1 q_{N_p,1}(\mathbf{k}, \tau) + \left( \beta + \frac{1}{2} \right) \frac{i}{\widehat{L}} k_1 \psi_{N_p,1}(\mathbf{k}, \tau) \right\} d\tau \end{aligned}$$

$$+\nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} q_{N_p,1}(\mathbf{k}, \tau) \} d\tau, \forall N_p \geq |\mathbf{k}|. \quad (3.34)$$

Using (3.30) it is clear that all the terms under the integral except the first one converge to the corresponding ones for  $\vec{\psi}$ . We need to show that

$$\delta := \int_0^t \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) q_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau)) d\tau \rightarrow 0 \quad (3.35)$$

as  $p \rightarrow \infty$ . For this we have

$$\begin{aligned} & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) q_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau)) \\ = & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{|\mathbf{l}|^2}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ & + \frac{1}{2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ & + \frac{1}{2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,2}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_2(\mathbf{l}, \tau)). \end{aligned}$$

The first sum on the right hand side is equal to

$$\begin{aligned} & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{\mathbf{l} \cdot (\mathbf{k} - \mathbf{h})}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ = & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{\mathbf{l} \cdot \mathbf{k}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ & - \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{\mathbf{l} \cdot \mathbf{h}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \end{aligned}$$

and since the last sum is zero we get that

$$\begin{aligned} & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{|\mathbf{l}|^2}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ = & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \frac{\mathbf{l} \cdot \mathbf{k}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \\ = & \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2(k_1 - h_1) - h_1(k_2 - h_2)) \frac{\mathbf{l} \cdot \mathbf{k}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \end{aligned}$$

$$= \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2k_1 - h_1k_2) \frac{\mathbf{l} \cdot \mathbf{k}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)),$$

It follows that

$$\begin{aligned} & \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) q_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau)) \right| \\ & \leq \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2k_1 - h_1k_2) \frac{\mathbf{l} \cdot \mathbf{k}}{\widehat{L}^2} (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \right| \\ & \quad + \frac{1}{2} \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_1(\mathbf{l}, \tau)) \right| \\ & \quad + \frac{1}{2} \left| \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2l_1 - h_1l_2) (\psi_{N_p,1}(\mathbf{h}, \tau) \psi_{N_p,2}(\mathbf{l}, \tau) - \psi_1(\mathbf{h}, \tau) \psi_2(\mathbf{l}, \tau)) \right| \\ & \leq \left( |\mathbf{k}|^2 + \frac{\widehat{L}^2}{2} \right) \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} \left( \frac{|\mathbf{h}|}{\widehat{L}} |\psi_{N_p,1}(\mathbf{h}, \tau) - \psi_1(\mathbf{h}, \tau)| \frac{|\mathbf{l}|}{\widehat{L}} |\psi_{N_p,1}(\mathbf{l}, \tau)| + \right. \\ & \quad \left. \frac{|\mathbf{h}|}{\widehat{L}} |\psi_1(\mathbf{h}, \tau)| \frac{|\mathbf{l}|}{\widehat{L}} |\psi_{N_p,1}(\mathbf{l}, \tau) - \psi_1(\mathbf{l}, \tau)| + \right. \\ & \quad \left. \frac{\widehat{L}^2}{2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} \left( \frac{|\mathbf{h}|}{\widehat{L}} |\psi_{N_p,1}(\mathbf{h}, \tau) - \psi_1(\mathbf{h}, \tau)| \frac{|\mathbf{l}|}{\widehat{L}} |\psi_{N_p,2}(\mathbf{l}, \tau)| + \right. \right. \\ & \quad \left. \left. \frac{|\mathbf{h}|}{\widehat{L}} |\psi_1(\mathbf{h}, \tau)| \frac{|\mathbf{l}|}{\widehat{L}} |\psi_{N_p,2}(\mathbf{l}, \tau) - \psi_2(\mathbf{l}, \tau)| \right) \right. \\ & \leq (|\mathbf{k}|^2 + \widehat{L}^2) (|\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau)| |\vec{\psi}_{N_p}(\tau)| + |\vec{\psi}(\tau)| |\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau)|). \end{aligned}$$

Using the last estimate and Holder's inequality we get that

$$\delta \leq 2(|\mathbf{k}|^2 + \widehat{L}^2) \left( \int_0^t |\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau)|^2 d\tau \right)^{1/2} \sqrt{te^{\frac{Lt}{4}}} |\vec{\psi}^0|.$$

Now we can see that in order to prove (3.35) it suffices to show that

$$\int_0^t |\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau)|^2 d\tau \rightarrow 0 \text{ as } p \rightarrow \infty.$$

Since  $\psi_{N_p,j}(\mathbf{k}, \tau) \rightarrow \psi_j(\mathbf{k}, \tau)$  uniformly for  $\tau \in [0, t]$ , for each fixed  $\mathbf{k} \in \mathbb{Z}^2$ , we have for each  $M = 1, 2, \dots$

$$\begin{aligned}
\lambda & : = \limsup_{p \rightarrow \infty} \int_0^t |\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau)|^2 d\tau \\
& = \limsup_{p \rightarrow \infty} \int_0^t |(I - P_M)(\vec{\psi}_{N_p}(\tau) - \vec{\psi}(\tau))|^2 d\tau \\
& \leq \limsup_{p \rightarrow \infty} \left[ 2 \int_0^t |(I - P_M)\vec{\psi}_{N_p}(\tau)|^2 d\tau \right] + 2 \int_0^t |(I - P_M)\vec{\psi}(\tau)|^2 d\tau. \quad (3.36)
\end{aligned}$$

We also have

$$\begin{aligned}
|(I - P_M)\vec{\psi}_{N_p}(\tau)|^2 & = \sum_{|\mathbf{k}| > M} E(\vec{\psi}_{N_p}(\tau))(\mathbf{k}) \leq \frac{\widehat{L}}{M} \sum_{|\mathbf{k}| > M} \frac{|\mathbf{k}|}{\widehat{L}} E(\vec{\psi}_{N_p}(\tau))(\mathbf{k}) \\
& \leq \frac{\widehat{L}}{M} |\vec{\psi}_{N_p}(\tau)|_{1+\alpha}^2. \quad (3.37)
\end{aligned}$$

From (3.32) and (3.37) we get

$$\begin{aligned}
\int_0^t |(I - P_M)\vec{\psi}_{N_p}(\tau)|^2 d\tau & \leq \frac{\widehat{L}}{M} \int_0^t |\vec{\psi}_{N_p}(\tau)|_{1+\alpha}^2 d\tau \\
& \leq \frac{\widehat{L}}{\nu M} \left( \frac{1}{2} + \frac{\widehat{L}t}{4} e^{\frac{Lt}{2}} \right) |\vec{\psi}^0|^2. \quad (3.38)
\end{aligned}$$

Applying Lebesgue's dominated convergence theorem we have that

$$\lim_{M \rightarrow \infty} \int_0^t |(I - P_M)\vec{\psi}(\tau)|^2 d\tau = 0. \quad (3.39)$$

Using (3.38) and (3.39) we let  $M \rightarrow \infty$  in (3.36) and we obtain that  $\lambda = 0$ . Next we let  $p \rightarrow \infty$  in (3.34) to get the first equation of 2) in Definition B.1 of Chapter II. In a similar fashion we deduce the second equation of 2) in Definition B.1 of Chapter II. It is easy to see that  $\psi_j(\mathbf{k}, 0) = \psi_j^0(\mathbf{k})$ ,  $\forall \mathbf{k} \in \mathbb{Z}^2$ ,  $j = 1, 2$ , and the proof that  $\vec{\psi}$  is a weak solution for (2.20)-(2.24) with initial data  $\vec{\psi}^0$  is complete.  $\square$



### C. Uniqueness of weak solutions

In this section we prove that  $\overrightarrow{\psi}$  (the weak solution found in the previous section) is the unique weak solution for (2.20)-(2.24). For this we need a few preliminary results.

**Lemma C.1.** *Let  $\varphi_0, \psi_0 \in \mathbb{R}^d$ ,  $f, g \in L^2([0, T]; \mathbb{R}^d)$  and let*

$$\varphi(t) = \varphi_0 + \int_0^t f(\tau) d\tau, \quad \psi(t) = \psi_0 + \int_0^t g(\tau) d\tau, \quad \forall 0 \leq t \leq T. \quad (3.40)$$

*Then*

$$\varphi(t) \cdot \psi(t) = \varphi_0 \cdot \psi_0 + \int_0^t (f(\tau) \cdot \psi(\tau) + \varphi(\tau) \cdot g(\tau)) d\tau, \quad \forall 0 \leq t \leq T. \quad (3.41)$$

*Proof.* If  $f, g \in C([0, T]; \mathbb{R}^d)$  then from (3.40) we get that  $\varphi'(t) = f(t), \psi'(t) = g(t)$  and (3.41) is easily obtained by integrating

$$\frac{d}{dt}(\varphi \cdot \psi) = \frac{d\varphi}{dt} \cdot \psi + \varphi \cdot \frac{d\psi}{dt}. \quad (3.42)$$

The proof is complete by noticing that  $C([0, T]; \mathbb{R}^d)$  is dense in  $L^2([0, T]; \mathbb{R}^d)$ .  $\square$

**Corollary C.1.** *Let  $\varphi_0 \in \mathbb{C}$ ,  $f \in L^2([0, T]; \mathbb{C})$  and let*

$$\varphi(t) = \varphi_0 + \int_0^t f(\tau) d\tau, \quad \forall 0 \leq t \leq T.$$

*Then*

$$|\varphi(t)|^2 = |\varphi_0|^2 + 2\operatorname{Re} \int_0^t f(\tau) \overline{\varphi(\tau)} d\tau, \quad \forall 0 \leq t \leq T.$$

The next result that we will use in the proof of uniqueness is the following variant of Ladyzhenskaya's inequality. With  $\Omega = [0, 2\pi\widehat{L}]^2 \subset \mathbb{R}^2$  there exists  $c_L > 0$  such that

$$\|u\|_{L^4(\Omega)} \leq c_L \|u\|_{L^2(\Omega)}^{1/2} \|\nabla u\|_{(L^2(\Omega))^2}^{1/2}, \quad (3.43)$$

for every  $u$  in

$$H_{per}^1(\Omega) = \{v \in L^2(\Omega) : \sum_{\mathbf{k} \in \mathbb{Z}^2} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |v(\mathbf{k})|^2 < \infty\} \quad (3.44)$$

with average zero (i.e.,  $\int_{\Omega} u(x) dx = 0$ ). (Recall that in (3.44),  $\{v(\mathbf{k})\}_{\mathbf{k} \in \mathbb{Z}^2}$  are the Fourier coefficients of  $v$ .)

Next we prove the main theorem of this section.

**Theorem C.1.** *For every given initial data in  $H$  the equations (2.20)-(2.24) have a unique weak solution.*

*Proof.* Suppose that  $\vec{\varphi}$  is another weak solution for (2.20)-(2.24) with initial data  $\vec{\psi}^0$ . Let  $\vec{w} = \vec{\psi} - \vec{\varphi}$  and  $\vec{y} = \vec{q} - \vec{r}$ , where

$$r_j(\mathbf{k}, t) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \varphi_j(\mathbf{k}, t) + (-1)^j \frac{\varphi_1(\mathbf{k}, t) - \varphi_2(\mathbf{k}, t)}{2}, j = 1, 2.$$

Since  $\vec{\psi}$  and  $\vec{\varphi}$  are weak solutions we have that

$$\begin{aligned} q_1(\mathbf{k}, t) = & q_1(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) + \frac{i}{\widehat{L}} k_1 q_1(\mathbf{k}, \tau) + \right. \\ & \left. (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \psi_1(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} q_1(\mathbf{k}, \tau) \right\} d\tau \end{aligned} \quad (3.45)$$

and

$$\begin{aligned} r_1(\mathbf{k}, t) = & r_1(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) + \frac{i}{\widehat{L}} k_1 r_1(\mathbf{k}, \tau) \right. \\ & \left. + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \varphi_1(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} r_1(\mathbf{k}, \tau) \right\} d\tau. \end{aligned} \quad (3.46)$$

By subtracting (3.46) from (3.45) we get that

$$\begin{aligned} y_1(\mathbf{k}, t) = & y_1(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau)) \right. \\ & \left. + \frac{i}{\widehat{L}} k_1 y_1(\mathbf{k}, \tau) + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_1(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} y_1(\mathbf{k}, \tau) \right\} d\tau. \end{aligned} \quad (3.47)$$

Similarly,

$$q_2(\mathbf{k}, t) = q_2(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) - \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \psi_2(\mathbf{k}, \tau) \right. \\ \left. + (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \psi_2(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} q_2(\mathbf{k}, \tau) \right\} d\tau,$$

$$r_2(\mathbf{k}, t) = r_2(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau) \right. \\ \left. - \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \varphi_2(\mathbf{k}, \tau) + (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \varphi_2(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} r_2(\mathbf{k}, \tau) \right\} d\tau,$$

and

$$y_2(\mathbf{k}, t) = y_2(\mathbf{k}, 0) - \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) - \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) \right. \\ \left. - \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} w_2(\mathbf{k}, \tau) + (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_2(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} y_2(\mathbf{k}, \tau) \right\} d\tau. \quad (3.48)$$

Next we define  $\widetilde{w} = w_1 + w_2$  and  $\widehat{w} = w_1 - w_2$ . An easy calculation gives us that

$$y_1(\mathbf{k}) + y_2(\mathbf{k}) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \widetilde{w}(\mathbf{k}) \text{ and } y_1(\mathbf{k}) - y_2(\mathbf{k}) = -\left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) \widehat{w}(\mathbf{k}).$$

Adding (3.47) and (3.48) we obtain that

$$\frac{|\mathbf{k}|^2}{\widehat{L}^2} \widetilde{w}(\mathbf{k}, t) = \frac{|\mathbf{k}|^2}{\widehat{L}^2} \widetilde{w}(\mathbf{k}, 0) + \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) \right. \\ \left. - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) + \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) - \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) + \frac{i}{\widehat{L}} k_1 y_1(\mathbf{k}, \tau) \right. \\ \left. + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_1(\mathbf{k}, \tau) - \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} w_2(\mathbf{k}, \tau) + (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_2(\mathbf{k}, \tau) \right. \\ \left. + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} \left( -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \widetilde{w}(\mathbf{k}, \tau) \right) \right\} d\tau. \quad (3.49)$$

Applying Corollary C.1 with  $\varphi(t) = \frac{|\mathbf{k}|}{\widehat{L}} \widetilde{w}(\mathbf{k}, t)$ , for every  $\mathbf{k} \neq \mathbf{0}$  we obtain

$$\begin{aligned}
\frac{|\mathbf{k}|^2}{\widehat{L}^2} |\widetilde{w}(\mathbf{k}, t)|^2 &= \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\widetilde{w}(\mathbf{k}, 0)|^2 + 2\operatorname{Re} \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \right. \\
&(\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) + \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) - \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) \overline{\widetilde{w}(\mathbf{k}, \tau)} \\
&+ \frac{i}{\widehat{L}} k_1 y_1(\mathbf{k}, \tau) \overline{\widetilde{w}(\mathbf{k}, \tau)} + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_1(\mathbf{k}, \tau) \overline{\widetilde{w}(\mathbf{k}, \tau)} - \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} w_2(\mathbf{k}, \tau) \overline{\widetilde{w}(\mathbf{k}, \tau)} \\
&\left. + (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_2(\mathbf{k}, \tau) \overline{\widetilde{w}(\mathbf{k}, \tau)} - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(2+\alpha)} |\widetilde{w}(\mathbf{k}, \tau)|^2 \right\} d\tau. \quad (3.50)
\end{aligned}$$

Subtracting (3.48) from (3.47) we get

$$\begin{aligned}
\left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) \widehat{w}(\mathbf{k}, t) &= \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) \widehat{w}(\mathbf{k}, 0) + \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \right. \\
&(\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) - \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) + \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) \\
&+ \frac{i}{\widehat{L}} k_1 y_1(\mathbf{k}, \tau) + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_1(\mathbf{k}, \tau) + \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} w_2(\mathbf{k}, \tau) \\
&\left. - (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_2(\mathbf{k}, \tau) + \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} \left( - \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) \widehat{w}(\mathbf{k}, \tau) \right) \right\} d\tau.
\end{aligned}$$

Applying again Corollary C.1 we deduce that

$$\begin{aligned}
\left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) |\widehat{w}(\mathbf{k}, t)|^2 &= \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) |\widehat{w}(\mathbf{k}, 0)|^2 + 2\operatorname{Re} \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \right. \\
&(\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau) - \psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) + \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) \overline{\widehat{w}(\mathbf{k}, \tau)} \\
&+ \frac{i}{\widehat{L}} k_1 y_1(\mathbf{k}, \tau) \overline{\widehat{w}(\mathbf{k}, \tau)} + (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_1(\mathbf{k}, \tau) \overline{\widehat{w}(\mathbf{k}, \tau)} + \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} w_2(\mathbf{k}, \tau) \overline{\widehat{w}(\mathbf{k}, \tau)} \\
&\left. - (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 w_2(\mathbf{k}, \tau) \overline{\widehat{w}(\mathbf{k}, \tau)} - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) |\widehat{w}(\mathbf{k}, \tau)|^2 \right\} d\tau. \quad (3.51)
\end{aligned}$$

Recall that

$$|\vec{w}|^2 = \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|w_1(\mathbf{k})|^2 + |w_2(\mathbf{k})|^2) + \frac{|w_1(\mathbf{k}) - w_2(\mathbf{k})|^2}{2} \right)$$

$$= \frac{1}{2} \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\widetilde{w}(\mathbf{k})|^2 + \left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + 1 \right) |\widehat{w}(\mathbf{k})|^2 \right).$$

Using this and the relations (3.50) and (3.51), after summing over  $\mathbf{k} \in \mathbb{Z}^2$  we obtain

$$\begin{aligned} |\vec{w}(t)|^2 &= |\vec{w}(0)|^2 + 2\operatorname{Re} \int_0^t \left\{ \frac{1}{\widehat{L}^2} \sum_{\mathbf{k} \in \mathbb{Z}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_1(\mathbf{h}, \tau) q_1(\mathbf{l}, \tau) \right. \\ &\quad \left. - \varphi_1(\mathbf{h}, \tau) r_1(\mathbf{l}, \tau)) \overline{w_1(\mathbf{k}, \tau)} + \frac{1}{\widehat{L}^2} \sum_{\mathbf{k} \in \mathbb{Z}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\psi_2(\mathbf{h}, \tau) q_2(\mathbf{l}, \tau) \right. \\ &\quad \left. - \varphi_2(\mathbf{h}, \tau) r_2(\mathbf{l}, \tau)) \overline{w_2(\mathbf{k}, \tau)} + \frac{i}{\widehat{L}} \sum_{\mathbf{k} \in \mathbb{Z}^2} k_1 y_1(\mathbf{k}, \tau) \overline{w_1(\mathbf{k}, \tau)} - \kappa_M \sum_{\mathbf{k} \in \mathbb{Z}^2} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |w_2(\mathbf{k}, \tau)|^2 \right. \\ &\quad \left. - \nu \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{w})(\mathbf{k}) \right. \end{aligned}$$

Therefore,

$$\begin{aligned} |\vec{w}(t)|^2 + \kappa_M \int_0^t \sum_{\mathbf{k} \in \mathbb{Z}^2} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |w_2(\mathbf{k}, \tau)|^2 d\tau + \nu \int_0^t \sum_{\mathbf{k} \in \mathbb{Z}^2} \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} E(\vec{w})(\mathbf{k}) d\tau = \\ 2\operatorname{Re} \frac{i}{\widehat{L}} \int_0^t \sum_{\mathbf{k} \in \mathbb{Z}^2} k_1 y_1(\mathbf{k}, \tau) \overline{w_1(\mathbf{k}, \tau)} d\tau + 2\operatorname{Re} \frac{1}{\widehat{L}^2} \sum_{j=1}^2 \int_0^t \sum_{\mathbf{k} \in \mathbb{Z}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \\ (w_j(\mathbf{h}, \tau) q_j(\mathbf{l}, \tau) \overline{w_j(\mathbf{k}, \tau)} + \varphi_j(\mathbf{h}, \tau) y_j(\mathbf{l}, \tau) \overline{w_j(\mathbf{k}, \tau)}) d\tau. \end{aligned} \quad (3.52)$$

Using the same steps as when we proved that  $S_1 = 0$  we can show that

$$\sum_{\mathbf{k} \in \mathbb{Z}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) w_j(\mathbf{h}, \tau) q_j(\mathbf{l}, \tau) \overline{w_j(\mathbf{k}, \tau)} = 0, \quad j = 1, 2.$$

Also we have

$$\begin{aligned} \sum_{\mathbf{k} \in \mathbb{Z}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) y_1(\mathbf{l}, \tau) \overline{w_1(\mathbf{k}, \tau)} = \\ \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}=\mathbf{0}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) w_1(\mathbf{k}, \tau) \left( -\frac{|\mathbf{l}|^2}{\widehat{L}^2} w_1(\mathbf{l}, \tau) - \frac{w_1(\mathbf{l}, \tau) - w_2(\mathbf{l}, \tau)}{2} \right), \end{aligned}$$

and

$$\begin{aligned}
& \left| \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} (h_2 l_1 - h_1 l_2) \varphi_1(\mathbf{h}, \tau) w_1(\mathbf{k}, \tau) \frac{|\mathbf{l}|^2}{\widehat{L}^2} w_1(\mathbf{l}, \tau) \right| \\
&= \left| \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} (h_2 k_1 - h_1 k_2) \varphi_1(\mathbf{h}, \tau) w_1(\mathbf{k}, \tau) \frac{\mathbf{h} \cdot \mathbf{l}}{\widehat{L}^2} w_1(\mathbf{l}, \tau) \right| \\
&\leq \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} \left( \frac{|\mathbf{h}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{h}, \tau)| \right) \left( \frac{|\mathbf{k}|}{\widehat{L}} |w_1(\mathbf{k}, \tau)| \right) \left( \frac{|\mathbf{l}|}{\widehat{L}} |w_1(\mathbf{l}, \tau)| \right).
\end{aligned}$$

Next we define the auxiliary functions

$$f(x) = \sum_{\mathbf{k} \in \mathbb{Z}^2} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{k}, \tau)| e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}} \quad \text{and} \quad g(x) = \sum_{\mathbf{k} \in \mathbb{Z}^2} \frac{|\mathbf{k}|}{\widehat{L}} |w_1(\mathbf{k}, \tau)| e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}}.$$

Then,

$$\begin{aligned}
S & : = \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} \left( \frac{|\mathbf{h}|^2}{\widehat{L}^2} |\varphi_1(\mathbf{h}, \tau)| \right) \left( \frac{|\mathbf{k}|}{\widehat{L}} |w_1(\mathbf{k}, \tau)| \right) \left( \frac{|\mathbf{l}|}{\widehat{L}} |w_1(\mathbf{l}, \tau)| \right) \\
&= \frac{1}{(2\pi \widehat{L})^2} \int_{\Omega} f(x) g^2(x) dx.
\end{aligned}$$

Applying Holder's and Ladyzhenskaya's inequalities we obtain that

$$\begin{aligned}
|S| &\leq \frac{1}{(2\pi \widehat{L})^2} \|f\|_{L^2(\Omega)} \|g\|_{L^4(\Omega)}^2 \\
&\leq \frac{1}{(2\pi \widehat{L})^2} c_L^2 \|f\|_{L^2(\Omega)} \|g\|_{L^2(\Omega)} \|\nabla g\|_{(L^2(\Omega))^2} \\
&\leq c |\vec{\varphi}(\tau)|_1 |\vec{w}(\tau)| |\vec{w}(\tau)|_1, \quad \forall \tau \in [0, T].
\end{aligned}$$

Using the above estimate and similar estimates for the other terms, from (3.52) we get

$$\begin{aligned}
|\vec{w}(t)|^2 + \nu \int_0^t |\vec{w}(\tau)|_{1+\alpha}^2 d\tau &\leq |\vec{w}(0)|^2 + c \int_0^t |\vec{w}(\tau)|^2 d\tau \\
&\quad + c \int_0^t |\vec{\varphi}(\tau)|_1 |\vec{w}(\tau)| |\vec{w}(\tau)|_1 d\tau, \quad (3.53)
\end{aligned}$$

for every  $t \in [0, T]$ , where  $c > 0$  depends on  $T$ . In the last integral of (3.53) we use

the inequality  $2ab \leq a^2 + b^2$  to get

$$\begin{aligned} |\vec{w}(t)|^2 + \nu \int_0^t |\vec{w}(\tau)|_{1+\alpha}^2 d\tau &\leq |\vec{w}(0)|^2 + c \int_0^t |\vec{w}(\tau)|^2 d\tau + \nu \int_0^t |\vec{w}(\tau)|_1^2 d\tau \\ &\quad + \tilde{c} \int_0^t |\vec{\varphi}(\tau)|_1^2 |\vec{w}(\tau)|^2 d\tau, \end{aligned}$$

which implies that

$$|\vec{w}(t)|^2 \leq |\vec{w}(0)|^2 + \hat{c} \int_0^t |\vec{\varphi}(\tau)|_1^2 |\vec{w}(\tau)|^2 d\tau.$$

Using Lemma C.2 (below) we deduce that  $|\vec{w}(t)|^2 \leq |\vec{w}(0)|^2 e^{\hat{c} \int_0^t |\vec{\varphi}(\tau)|_1^2 d\tau}$ ,  $\forall t \in [0, T]$ . But  $\vec{w}(0) = \mathbf{0}$ , and thus,  $\vec{w}(t) = \mathbf{0}$ ,  $\forall t \in [0, T]$ . Since  $T$  was arbitrary we conclude that  $\vec{\psi}(t) = \vec{\varphi}(t)$ ,  $\forall t \in [0, \infty)$ , and the proof is complete.  $\square$

The lemma below is a generalization of Gronwall's inequality. The proof is elementary and it is omitted.

**Lemma C.2.** *Let  $f_0 \geq 0$  and  $f \in L^\infty([0, T], \mathbb{R})$ ,  $g \in L^1([0, T], \mathbb{R})$  be nonnegative functions such that*

$$f(t) \leq f_0 + \int_0^t g(\tau) f(\tau) d\tau, \forall t \in [0, T].$$

Then

$$f(t) \leq f_0 e^{\int_0^t g(\tau) d\tau}, \forall t \in [0, T].$$

**Remark C.1.** *Since every limit point in  $C([0, \infty), \mathcal{K})$  of  $\{\vec{\psi}_N\}_{N \in \mathbb{N}}$  is a weak solution for (2.20)-(2.24) we easily get as a consequence of uniqueness that  $\{\vec{\psi}_N\}_{N \in \mathbb{N}}$  converges to  $\vec{\psi}$  in  $C([0, \infty), \mathcal{K})$ .*

## CHAPTER IV

## REGULARITY

## A. Estimates for the Galerkin approximations

If  $\vec{\varphi} \in \mathcal{K}$ ,  $s \in \mathbb{R}$  and  $\zeta \in \mathbb{C}$  we denote by  $A^s \vec{\varphi}$  and  $e^{\zeta A^s} \vec{\varphi}$  the following

$$\begin{aligned} A^s \vec{\varphi} &= \left( \left\{ \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2s} \varphi_1(\mathbf{k}) \right\}_{\mathbf{k} \in \mathbb{Z}^2}, \left\{ \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2s} \varphi_2(\mathbf{k}) \right\}_{\mathbf{k} \in \mathbb{Z}^2} \right) \\ e^{\zeta A^s} \vec{\varphi} &= \left( \left\{ e^{\zeta \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2s}} \varphi_1(\mathbf{k}) \right\}_{\mathbf{k} \in \mathbb{Z}^2}, \left\{ e^{\zeta \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2s}} \varphi_2(\mathbf{k}) \right\}_{\mathbf{k} \in \mathbb{Z}^2} \right). \end{aligned}$$

We define the space

$$H_{\mathbb{C}} = \{ \vec{\psi} + i \vec{\varphi} : \vec{\psi}, \vec{\varphi} \in H \}$$

with the scalar product

$$(\vec{\psi} + i \vec{\varphi}, \vec{u} + i \vec{v})_{\mathbb{C}} = (\vec{\psi}, \vec{u}) + (\vec{\varphi}, \vec{v}) + i[(\vec{\varphi}, \vec{u}) - (\vec{\psi}, \vec{v})].$$

Similarly, we complexify the space  $V_1$  to get the space  $V_{1,\mathbb{C}}$  with the corresponding scalar product  $(\cdot, \cdot)_{1,\mathbb{C}}$ . For every  $N \in \mathbb{N}$  we consider the following Galerkin system:

$$\begin{aligned} \frac{d}{d\zeta} \begin{pmatrix} \Psi_1(\mathbf{k}, \zeta) \\ \Psi_2(\mathbf{k}, \zeta) \end{pmatrix} &= A_{\mathbf{k}}^{-1} \begin{pmatrix} -\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_1(\mathbf{h}, \zeta) Q_1(\mathbf{l}, \zeta) \\ -\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_2(\mathbf{h}, \zeta) Q_2(\mathbf{l}, \zeta) \end{pmatrix} + \\ A_{\mathbf{k}}^{-1} \begin{pmatrix} -\frac{i}{\widehat{L}} k_1 Q_1(\mathbf{k}, \zeta) - (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \Psi_1(\mathbf{k}, \zeta) - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} Q_1(\mathbf{k}, \zeta) \\ \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \Psi_2(\mathbf{k}, \zeta) - (\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \Psi_2(\mathbf{k}, \zeta) - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} Q_2(\mathbf{k}, \zeta) \end{pmatrix} \end{aligned} \quad (4.1)$$

for  $\mathbf{k} \neq \mathbf{0}$ ,  $|\mathbf{k}| \leq N$ , and

$$\frac{d}{d\zeta} \begin{pmatrix} \Psi_1(\mathbf{0}, \zeta) \\ \Psi_2(\mathbf{0}, \zeta) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad (4.2)$$

where



$$Q_1(\mathbf{k}, \zeta) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \Psi_1(\mathbf{k}, \zeta) - \frac{\Psi_1(\mathbf{k}, \zeta) - \Psi_2(\mathbf{k}, \zeta)}{2}, \quad (4.3)$$

$$Q_2(\mathbf{k}, \zeta) = -\frac{|\mathbf{k}|^2}{\widehat{L}^2} \Psi_2(\mathbf{k}, \zeta) + \frac{\Psi_1(\mathbf{k}, \zeta) - \Psi_2(\mathbf{k}, \zeta)}{2}. \quad (4.4)$$

The sums in (4.1) are taken only for  $|\mathbf{h}|, |\mathbf{l}| \leq N$ . For  $\vec{\psi}^0 \in H$  we consider the initial condition

$$\Psi_j(\mathbf{k}, 0) = \psi_j^0(\mathbf{k}), |\mathbf{k}| \leq N, j = 1, 2. \quad (4.5)$$

The system (4.1) together with (4.5) admits a unique analytic solution  $\vec{\Psi}^{(N)}(\zeta)$  for  $\zeta$  in a complex neighborhood of the origin. The solution  $\vec{\Psi}^{(N)}(\zeta)$ , for  $\zeta$  real, coincides with the usual Galerkin approximation. From (4.1) we have that

$$\begin{aligned} & \frac{d}{d\zeta} Q_1(\mathbf{k}, \zeta) + \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_1(\mathbf{h}, \zeta) Q_1(\mathbf{l}, \zeta) \\ &= -\frac{i}{\widehat{L}} k_1 Q_1(\mathbf{k}, \zeta) - \left(\beta + \frac{1}{2}\right) \frac{i}{\widehat{L}} k_1 \Psi_1(\mathbf{k}, \zeta) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} Q_1(\mathbf{k}, \zeta), \end{aligned} \quad (4.6)$$

and

$$\begin{aligned} & \frac{d}{d\zeta} Q_2(\mathbf{k}, \zeta) + \frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_2(\mathbf{h}, \zeta) Q_2(\mathbf{l}, \zeta) \\ &= \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \Psi_2(\mathbf{k}, \zeta) - \left(\beta - \frac{1}{2}\right) \frac{i}{\widehat{L}} k_1 \Psi_2(\mathbf{k}, \zeta) - \nu \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(1+\alpha)} Q_2(\mathbf{k}, \zeta). \end{aligned} \quad (4.7)$$

Let  $a \geq 0$ ,  $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$  and take  $\zeta$  of the form  $\zeta = \sigma e^{i\theta}$  for  $\sigma > 0$ . We want to evaluate  $\frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1, \mathbb{C}}^2$ . For this we first notice that

$$\frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1, \mathbb{C}}^2 = \frac{1}{2} \frac{d}{d\sigma} \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k})$$

where

$$E(\vec{\Psi}(\zeta))(\mathbf{k}) := \frac{|\mathbf{k}|^2}{\widehat{L}^2} (|\Psi_1(\mathbf{k}, \zeta)|^2 + |\Psi_2(\mathbf{k}, \zeta)|^2) + \frac{|\Psi_1(\mathbf{k}, \zeta) - \Psi_2(\mathbf{k}, \zeta)|^2}{2}.$$

Therefore,

$$\begin{aligned} \frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 &= \sum_{|\mathbf{k}| \leq N} a \cos \theta \frac{|\mathbf{k}|^3}{\widehat{L}^3} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) + \\ &\frac{1}{2} \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} \frac{d}{d\sigma} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}). \end{aligned}$$

The derivative with respect to  $\sigma$  of  $E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k})$  is equal to

$$-2\operatorname{Re}e^{i\theta} \left[ \left( \frac{d}{d\zeta} Q_1(\mathbf{k}, \zeta) \right)_{\zeta=\sigma e^{i\theta}} \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} + \left( \frac{d}{d\zeta} Q_2(\mathbf{k}, \zeta) \right)_{\zeta=\sigma e^{i\theta}} \overline{\Psi_2(\mathbf{k}, \sigma e^{i\theta})} \right]$$

and using (4.6) and (4.7) we get

$$\begin{aligned} \frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 &= a \cos \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^3}{\widehat{L}^3} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) - \\ \operatorname{Re}\{e^{i\theta} \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} [ &(-\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_1(\mathbf{h}, \sigma e^{i\theta}) Q_1(\mathbf{l}, \sigma e^{i\theta}) - \\ \frac{i}{\widehat{L}} k_1 Q_1(\mathbf{k}, \sigma e^{i\theta}) - (\beta + \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \Psi_1(\mathbf{k}, \sigma e^{i\theta}) - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} &Q_1(\mathbf{k}, \sigma e^{i\theta}) \\ \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} + (-\frac{1}{\widehat{L}^2} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_2(\mathbf{h}, \sigma e^{i\theta}) Q_2(\mathbf{l}, \sigma e^{i\theta}) + \kappa_M \frac{|\mathbf{k}|^2}{\widehat{L}^2} \Psi_2(\mathbf{k}, \sigma e^{i\theta}) & \\ -(\beta - \frac{1}{2}) \frac{i}{\widehat{L}} k_1 \Psi_2(\mathbf{k}, \sigma e^{i\theta}) - \nu \left( \frac{|\mathbf{k}|}{\widehat{L}} \right)^{2(1+\alpha)} Q_2(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_2(\mathbf{k}, \sigma e^{i\theta})}] \}. & \end{aligned}$$

Thus

$$\begin{aligned} \frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 &= a \cos \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^3}{\widehat{L}^3} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) + \\ \operatorname{Re}\{e^{i\theta} \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^4} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) &(\Psi_1(\mathbf{h}, \sigma e^{i\theta}) Q_1(\mathbf{l}, \sigma e^{i\theta}) \\ \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} + \Psi_2(\mathbf{h}, \sigma e^{i\theta}) Q_2(\mathbf{l}, \sigma e^{i\theta}) \overline{\Psi_2(\mathbf{k}, \sigma e^{i\theta})}) \} &+ \\ \operatorname{Re}\{ie^{i\theta} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} \frac{k_1}{\widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} Q_1(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} \} &+ \end{aligned}$$

$$\begin{aligned}
& (\beta + \frac{1}{2}) \operatorname{Re}\{ie^{i\theta} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2 k_1}{\widehat{L}^2 \widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_1(\mathbf{k}, \sigma e^{i\theta})|^2\} - \\
& \quad \kappa_M \operatorname{Re}\{e^{i\theta} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^4}{\widehat{L}^4} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_2(\mathbf{k}, \sigma e^{i\theta})|^2\} + \\
& (\beta - \frac{1}{2}) \operatorname{Re}\{ie^{i\theta} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2 k_1}{\widehat{L}^2 \widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_2(\mathbf{k}, \sigma e^{i\theta})|^2\} + \\
& \nu \operatorname{Re}\{e^{i\theta} \sum_{|\mathbf{k}| \leq N} \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(2+\alpha)} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} (Q_1(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} + \\
& \quad Q_2(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_2(\mathbf{k}, \sigma e^{i\theta})})\}.
\end{aligned}$$

The last equality is equivalent with a new one which has the left-hand side equal to

$$\begin{aligned}
& \frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1, \mathbb{C}}^2 + \kappa_M \cos \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^4}{\widehat{L}^4} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_2(\mathbf{k}, \sigma e^{i\theta})|^2 + \\
& \quad \nu \cos \theta \sum_{|\mathbf{k}| \leq N} \left(\frac{|\mathbf{k}|}{\widehat{L}}\right)^{2(2+\alpha)} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k})
\end{aligned}$$

and the right-hand side equal to

$$\begin{aligned}
& a \cos \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^3}{\widehat{L}^3} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) - \\
& (\beta + \frac{1}{2}) \sin \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2 k_1}{\widehat{L}^2 \widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_1(\mathbf{k}, \sigma e^{i\theta})|^2 - \\
& (\beta - \frac{1}{2}) \sin \theta \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2 k_1}{\widehat{L}^2 \widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} |\Psi_2(\mathbf{k}, \sigma e^{i\theta})|^2 + \\
& \operatorname{Re}\{ie^{i\theta} \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2 k_1}{\widehat{L}^2 \widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} Q_1(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})}\} + \\
& \operatorname{Re}\{e^{i\theta} \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^4} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) (\Psi_1(\mathbf{h}, \sigma e^{i\theta}) Q_1(\mathbf{l}, \sigma e^{i\theta}) \\
& \quad \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} + \Psi_2(\mathbf{h}, \sigma e^{i\theta}) Q_2(\mathbf{l}, \sigma e^{i\theta}) \overline{\Psi_2(\mathbf{k}, \sigma e^{i\theta})})\}.
\end{aligned}$$

Next step is to estimate the terms from the right-hand side of the equality.

For this we have

$$\begin{aligned}
& \left| \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} \frac{k_1}{\widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} Q_1(\mathbf{k}, \sigma e^{i\theta}) \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} \right| \\
&= \left| \sum_{|\mathbf{k}| \leq N} \frac{|\mathbf{k}|^2}{\widehat{L}^2} \frac{k_1}{\widehat{L}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \left[ -\left( \frac{|\mathbf{k}|^2}{\widehat{L}^2} + \frac{1}{2} \right) |\Psi_1(\mathbf{k}, \sigma e^{i\theta})|^2 + \right. \right. \\
&\quad \left. \left. \frac{1}{2} \Psi_2(\mathbf{k}, \sigma e^{i\theta}) \cdot \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} \right] \right| \\
&\leq c_1 |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}
\end{aligned}$$

and

$$\begin{aligned}
&= \left| \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^4} \sum_{\mathbf{h}+\mathbf{l}=\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_1(\mathbf{h}, \sigma e^{i\theta}) Q_1(\mathbf{l}, \sigma e^{i\theta}) \overline{\Psi_1(\mathbf{k}, \sigma e^{i\theta})} \right| \\
&= \left| \sum_{|\mathbf{k}| \leq N} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^4} \sum_{\mathbf{h}+\mathbf{l}=-\mathbf{k}} (h_2 l_1 - h_1 l_2) \Psi_1(\mathbf{h}, \sigma e^{i\theta}) Q_1(\mathbf{l}, \sigma e^{i\theta}) \overline{\Psi_1(-\mathbf{k}, \sigma e^{i\theta})} \right| \\
&\leq \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} e^{2a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} \frac{|\mathbf{h}|}{\widehat{L}} \frac{|\mathbf{l}|}{\widehat{L}} |\Psi_1(\mathbf{h}, \sigma e^{i\theta})| |\Psi_1(-\mathbf{k}, \sigma e^{i\theta})| \left[ \frac{|\mathbf{l}|^2}{\widehat{L}^2} |\Psi_1(\mathbf{l}, \sigma e^{i\theta})| \right. \\
&\quad \left. + \frac{1}{2} (|\Psi_1(\mathbf{l}, \sigma e^{i\theta})| + |\Psi_2(\mathbf{l}, \sigma e^{i\theta})|) \right]. \tag{4.8}
\end{aligned}$$

Let  $S$  be the sum

$$\begin{aligned}
S &: = \sum_{\mathbf{h}+\mathbf{l}+\mathbf{k}=\mathbf{0}} \left( e^{a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\Psi_1(-\mathbf{k}, \sigma e^{i\theta})| \right) \left( e^{a\sigma \cos \theta \frac{|\mathbf{h}|}{L}} \frac{|\mathbf{h}|}{\widehat{L}} |\Psi_1(\mathbf{h}, \sigma e^{i\theta})| \right) \\
&\quad \left( e^{a\sigma \cos \theta \frac{|\mathbf{l}|}{L}} \frac{|\mathbf{l}|^3}{\widehat{L}^3} |\Psi_1(\mathbf{l}, \sigma e^{i\theta})| \right)
\end{aligned}$$

and define the auxiliary functions

$$\begin{aligned}
u(\mathbf{x}) &= \sum_{|\mathbf{k}| \leq N} e^{a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\Psi_1(-\mathbf{k}, \sigma e^{i\theta})| e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}}, \\
v(\mathbf{x}) &= \sum_{|\mathbf{k}| \leq N} e^{a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|}{\widehat{L}} |\Psi_1(\mathbf{k}, \sigma e^{i\theta})| e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}}, \text{ and} \\
w(\mathbf{x}) &= \sum_{|\mathbf{k}| \leq N} e^{a\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^3}{\widehat{L}^3} |\Psi_1(\mathbf{k}, \sigma e^{i\theta})| e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}}.
\end{aligned}$$

Then, using Holder's inequality, we obtain that

$$S = \frac{1}{(2\pi\widehat{L})^2} \int_{\Omega} u(\mathbf{x})v(\mathbf{x})w(\mathbf{x})d\mathbf{x} \leq \frac{1}{(2\pi\widehat{L})^2} \|u\|_{L^4(\Omega)} \|v\|_{L^4(\Omega)} \|w\|_{L^2(\Omega)}.$$

Applying Ladyzhenskaya's inequality we get that

$$\begin{aligned} S &\leq \frac{c_L^2}{(2\pi\widehat{L})^2} \|u\|_{L^2(\Omega)}^{1/2} \|\nabla u\|_{L^2(\Omega)}^{1/2} \|v\|_{L^2(\Omega)}^{1/2} \|\nabla v\|_{L^2(\Omega)}^{1/2} \|w\|_{L^2(\Omega)} \\ &\leq \widetilde{c}_2 |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^{3/2} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}^{3/2}. \end{aligned} \quad (4.9)$$

Estimating the other terms in a similar way we deduce that

$$\begin{aligned} &\frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \nu \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2+\alpha,\mathbb{C}}^2 \leq \\ &(a \cos \theta + |\sin \theta|) c_1 |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}} + \\ &\quad \widehat{L}(\beta + \frac{1}{2}) |\sin \theta| |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \\ &\quad c_2 \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^{3/2} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}^{3/2}. \end{aligned} \quad (4.10)$$

By using the inequalities  $xy \leq \frac{1}{2}(x^2 + y^2)$  and  $xy \leq \frac{x^4}{4} + \frac{3y^{4/3}}{4}$ , (4.10) implies that

$$\begin{aligned} &\frac{1}{2} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \nu \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2+\alpha,\mathbb{C}}^2 \leq \\ &\frac{(a \cos \theta + |\sin \theta|)^2 c_1^2}{\nu \cos \theta} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \frac{\nu \cos \theta}{4} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}^2 + \\ &\quad \widehat{L}(\beta + \frac{1}{2}) |\sin \theta| |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \frac{\nu \cos \theta}{4} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}^2 + \\ &\quad \frac{27c_2^4}{4\nu^3} \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^6. \end{aligned}$$

Therefore,

$$\begin{aligned} &\frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \nu \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{2+\alpha,\mathbb{C}}^2 \leq \\ &2 \left[ \frac{(a \cos \theta + |\sin \theta|)^2 c_1^2}{\nu \cos \theta} + \widehat{L}(\beta + \frac{1}{2}) |\sin \theta| \right] |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 + \\ &\quad \frac{27c_2^4}{2\nu^3} \cos \theta |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^6, \end{aligned}$$

which gives us that

$$\begin{aligned} \frac{d}{d\sigma} |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 &\leq \gamma_1^2 |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2 \\ &\quad + \gamma_2^2 |e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^6, \end{aligned} \quad (4.11)$$

where

$$\gamma_1 = \sqrt{2} \left[ \frac{(a \cos \theta + |\sin \theta|)^2 c_1^2}{\nu \cos \theta} + \widehat{L}(\beta + \frac{1}{2}) |\sin \theta| \right]^{1/2}, \quad (4.12)$$

$$\gamma_2 = \left( \frac{27c_2^4}{2\nu^3} \cos \theta \right)^{1/2}. \quad (4.13)$$

If we denote  $|e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}}^2$  by  $g(\sigma)$  (4.11) becomes

$$\frac{d}{d\sigma} g(\sigma) \leq \gamma_1^2 g(\sigma) + \gamma_2^2 g^3(\sigma).$$

We have that

$$\begin{aligned} \frac{d}{d\sigma} (\gamma_1 + \gamma_2 g(\sigma)) &= \gamma_2 \frac{d}{d\sigma} g(\sigma) \leq (\gamma_1^2 + \gamma_2^2 g^2(\sigma)) \gamma_2 g(\sigma) \\ &\leq (\gamma_1 + \gamma_2 g(\sigma))^2 \gamma_2 g(\sigma) \leq (\gamma_1 + \gamma_2 g(\sigma))^3. \end{aligned}$$

The last inequality implies that

$$\gamma_1 + \gamma_2 g(\sigma) \leq \sqrt{2}(\gamma_1 + \gamma_2 g(0)) \text{ for } 0 \leq \sigma \leq \frac{1}{4}(\gamma_1 + \gamma_2 g(0))^{-2}.$$

This proves the following proposition.

**Proposition A.1.** *There exists  $\Gamma > 0$  independent of  $N$  such that*

$$|e^{a\sigma e^{i\theta} A^{1/2}} \overrightarrow{\Psi}^{(N)}(\sigma e^{i\theta})|_{1,\mathbb{C}} \leq \Gamma, \quad \forall N \in \mathbb{N} \quad (4.14)$$

for  $|\theta| < \frac{\pi}{2}$  and  $0 \leq \sigma \leq \frac{1}{4}(\gamma_1 + \gamma_2 |\overrightarrow{\psi}^0|_1^2)^{-2}$ .

In the next section we will use this proposition with  $a = 1$ . However we will also need the following result (in the proof of which we will take  $a = 0$ ).

**Proposition A.2.** *If  $\vec{\psi}$  is the unique weak solution for our system with initial data  $\vec{\psi}^0 \in H$  then  $\vec{\psi}(t) \in V_1, \forall t > 0$ .*

*Proof.* In (4.9) we have in fact that

$$S \leq \tilde{c}_2 |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{\mathbb{C}}^{1/2} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}} |e^{a\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{2,\mathbb{C}}^{3/2}. \quad (4.15)$$

If we take  $a = 0$  and  $\theta = 0$  in (4.10) and use (4.15) instead of (4.9) we get that

$$\frac{1}{2} \frac{d}{dt} |\vec{\psi}^{(N)}(t)|_1^2 + \nu |\vec{\psi}^{(N)}(t)|_{2+\alpha}^2 \leq c_2 |\vec{\psi}^{(N)}(t)|_1 |\vec{\psi}^{(N)}(t)|^{1/2} |\vec{\psi}^{(N)}(t)|_2^{3/2}, \forall t \geq 0.$$

As before, this implies that

$$\frac{1}{2} \frac{d}{dt} |\vec{\psi}^{(N)}(t)|_1^2 \leq \frac{c_3}{\nu^3} |\vec{\psi}^{(N)}(t)|_1^4 |\vec{\psi}^{(N)}(t)|^2, \forall t \geq 0.$$

From here we obtain that

$$|\vec{\psi}^{(N)}(t)|_1^2 \leq |\vec{\psi}^{(N)}(t_0)|_1^2 e^{\frac{c_3}{\nu^3} \int_{t_0}^t |\vec{\psi}^{(N)}(\tau)|_1^2 |\vec{\psi}^{(N)}(\tau)|^2 d\tau}, \forall t \geq t_0.$$

For a fixed  $T \in (0, \infty)$  we get by using (3.20) and (3.32) that

$$\int_{t_0}^t |\vec{\psi}^{(N)}(\tau)|_1^2 |\vec{\psi}^{(N)}(\tau)|^2 d\tau \leq b_0(T), \forall 0 \leq t_0 \leq t \leq T,$$

with  $b_0(T)$  independent of  $N$ . Therefore,

$$|\vec{\psi}^{(N)}(t)|_1^2 \leq b_1(T) |\vec{\psi}^{(N)}(t_0)|_1^2, \forall 0 \leq t_0 \leq t \leq T,$$

where  $b_1(T)$  is independent of  $N$ . Integrating in  $t_0$  between 0 and  $t$  we get

$$t |\vec{\psi}^{(N)}(t)|_1^2 \leq b_1(T) \int_0^t |\vec{\psi}^{(N)}(t_0)|_1^2 dt_0 \leq b_2(T), \forall t \in [0, T],$$

where  $b_2(T)$  doesn't depend on  $N$ . For  $M \in \mathbb{N}$  and  $M < N$  we have

$$|P_M \vec{\psi}^{(N)}(t)|_1^2 \leq |\vec{\psi}^{(N)}(t)|_1^2 \leq \frac{1}{t} b_2(T), \forall t \in (0, T].$$

Letting  $N \rightarrow \infty$  we get that

$$|P_M \vec{\psi}(t)|_1^2 \leq \frac{1}{t} b_2(T), \forall t \in (0, T].$$

This implies, by letting  $M \rightarrow \infty$ , that

$$|\vec{\psi}(t)|_1^2 \leq \frac{1}{t} b_2(T), \forall t \in (0, T].$$

Since  $T$  was arbitrary we conclude that  $\vec{\psi}(t) \in V_1, \forall t > 0$ . □

## B. Time and space analyticity

We can now state the first main result of this chapter.

**Theorem B.1.** *If  $\vec{\psi}^0 \in V_1$  then the unique weak solution extends to an analytic function  $\vec{\Psi}$  from  $\mathcal{D} = \{\zeta = \sigma e^{i\theta} : |\theta| < \frac{\pi}{4}, 0 < \sigma < \sigma_0\}$  into  $V_{1,\mathbb{C}}$  satisfying*

$$|e^{\sigma e^{i\theta} A^{1/2}} \vec{\Psi}(\sigma e^{i\theta})|_{1,\mathbb{C}} \left( = \left( \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{2\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) \right)^{1/2} \right) \leq \Gamma \quad (4.16)$$

in  $\mathcal{D}$ , where

$$\sigma_0 = \frac{1}{4} \left( \left[ 4\sqrt{2}c_1^2\nu^{-1} + \widehat{L}(\beta + \frac{1}{2})\sqrt{2} \right]^{1/2} + \frac{3\sqrt{3}c_2^2}{\sqrt{2}\nu^{3/2}} |\vec{\psi}^0|_1^2 \right)^{-2}.$$

*Proof.* We use Proposition A.1 with  $a = 1$  and  $|\theta| < \frac{\pi}{4}$ . Then using also (4.12) and (4.13) we get that

$$|e^{\sigma e^{i\theta} A^{1/2}} \vec{\Psi}^{(N)}(\sigma e^{i\theta})|_{1,\mathbb{C}} \leq \Gamma, \forall N \in \mathbb{N} \text{ in } \mathcal{D}.$$

Thus (by virtue of the classical Vitali theorem for operator-valued analytic functions) there exist a subsequence  $\{\vec{\Psi}^{(N_p)}\}_{p \in \mathbb{N}}$  and an analytic function  $\Theta$  defined for  $\zeta$  in  $\mathcal{D}$  such that

$$e^{\zeta A^{1/2}} \vec{\Psi}^{(N_p)}(\zeta) \rightarrow \Theta(\zeta) \text{ weakly in } V_{1,\mathbb{C}}$$



for all  $\zeta \in \mathcal{D}$ . Let  $M \in \mathbb{N}$  be fixed. We have

$$|P_M e^{\zeta A^{1/2}} \overrightarrow{\Psi}^{(N_p)}(\zeta) - P_M \Theta(\zeta)|_{1, \mathbb{C}} \rightarrow 0.$$

But from the convergence of  $\overrightarrow{\psi}^{(N)}(\cdot)$  to  $\overrightarrow{\psi}(\cdot)$  (see Remark C.1 of Chapter III) we also have

$$|e^{tA^{1/2}} P_M \overrightarrow{\psi}^{(N_p)}(t) - e^{tA^{1/2}} P_M \overrightarrow{\psi}(t)|_{1, \mathbb{C}} \rightarrow 0, \forall t \in [0, \sigma_0].$$

Therefore,  $e^{tA^{1/2}} P_M \overrightarrow{\psi}(t) = P_M \Theta(t)$ , *a.e.* on  $[0, \sigma_0]$  for every  $M \in \mathbb{N}$ . This easily implies that  $\overrightarrow{\psi}(t) = e^{-tA^{1/2}} \Theta(t)$ ,  $\forall t \in [0, \sigma_0]$ . The analytic function  $\overrightarrow{\Psi}(\zeta) = e^{-\zeta A^{1/2}} \Theta(\zeta)$  extends  $\overrightarrow{\psi}$  and satisfies (4.16). This completes the proof.  $\square$

The property (4.16) of the solution implies that it is an analytic function in the time and space variables. Indeed we have the following.

**Lemma B.1.** *Let*

$$\Phi(\zeta, \mathbf{x}) = \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}} \Phi_{\mathbf{k}}(\zeta), (\zeta, \mathbf{x}) \in \mathcal{D} \times \mathbb{R}^2,$$

*satisfy the following conditions*

- (i)  $\Phi_{\mathbf{k}}(\zeta)$  ( $\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}$ ) are  $\mathbb{C}$ -valued analytic functions in  $\mathcal{D}$ , and
- (ii)  $\sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{Re\zeta \frac{|\mathbf{k}|}{L}} |\Phi_{\mathbf{k}}(\zeta)| < \infty$  in  $\mathcal{D}$ .

*Then*

$$\sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{z}} \Phi_{\mathbf{k}}(\zeta)$$

*is absolutely convergent on*

$$\Xi = \{(\zeta, \mathbf{z}) : \zeta \in \mathcal{D}, |Im\mathbf{z}| < Re\zeta\}$$

*and the sum  $\Phi_{ex}(\zeta, \mathbf{z})$  is analytic in  $\Xi$  and extends  $\Phi(\zeta, \mathbf{x})$  to  $\Xi$ .*

*Proof.* We have

$$\begin{aligned}
\sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} |e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{z}} \Phi_{\mathbf{k}}(\zeta)| &\leq \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{-\frac{1}{L} \mathbf{k} \cdot \text{Imz}} |\Phi_{\mathbf{k}}(\zeta)| \\
&\leq \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\text{Re}\zeta \frac{|\mathbf{k}|}{L}} |\Phi_{\mathbf{k}}(\zeta)| e^{-\frac{1}{L} \mathbf{k} \cdot \text{Imz} - \text{Re}\zeta \frac{|\mathbf{k}|}{L}} \\
&\leq \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\text{Re}\zeta \frac{|\mathbf{k}|}{L}} |\Phi_{\mathbf{k}}(\zeta)| e^{-(\text{Re}\zeta - |\text{Imz}|) \frac{|\mathbf{k}|}{L}} \\
&\leq \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\text{Re}\zeta \frac{|\mathbf{k}|}{L}} |\Phi_{\mathbf{k}}(\zeta)| < \infty
\end{aligned}$$

if  $|\text{Imz}| \leq \text{Re}\zeta$ . Thus  $\Phi_{ex}$  is well defined in  $\Xi$  and extends  $\Phi$ . By noting that the series is also uniform convergent for  $|\text{Imz}| \leq (1 - \varepsilon)\text{Re}\zeta$  for any  $0 < \varepsilon < 1$ , it is easy to infer that  $\Phi_{ex}$  is also analytic in  $\Xi$ .  $\square$

We can now pass to the second main result of this chapter.

**Theorem B.2.** *Let  $\vec{\psi}^0 \in H$  and  $\vec{\psi}$  be the unique weak solution with initial data  $\vec{\psi}^0$ . Then the functions*

$$\psi_j(t, \mathbf{x}) = \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\frac{i}{L} \mathbf{k} \cdot \mathbf{x}} \psi_{j, \mathbf{k}}(t), j = 1, 2$$

*can be extended to analytic functions on some open neighborhood of  $(0, \infty) \times \mathbb{R}^2$  in  $\mathbb{C} \times \mathbb{C}^2$  and also satisfy our system in the classical sense.*

*Proof.* The theory of weak solutions that we developed so far allows us to consider any  $t_0 > 0$  as initial time. So it is clear that it suffices to assume that  $\vec{\psi}^0 \in V_1$  and prove first that  $\psi_j, j = 1, 2$  can be extended to analytic functions on  $\Xi$ . For this we need to show that the extensions for  $\vec{\psi}_j, j = 1, 2$  given by Theorem B.1 satisfy Lemma B.1. Indeed this is true because

$$\sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} e^{\text{Re}\zeta \frac{|\mathbf{k}|}{L}} |\psi_{j, \mathbf{k}}(\zeta)| \leq \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{\mathbf{0}\}} \frac{\widehat{L}^2}{|\mathbf{k}|^2} e^{\text{Re}\zeta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} |\psi_{j, \mathbf{k}}(\zeta)|$$

$$\begin{aligned}
&\leq \left( \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{0\}} \frac{\widehat{L}^4}{|\mathbf{k}|^4} \right)^{1/2} \left( \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{0\}} e^{2\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^4}{\widehat{L}^4} |\psi_{j,\mathbf{k}}(\zeta)|^2 \right)^{1/2} \\
&\leq c \left( \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{0\}} e^{2\sigma \cos \theta \frac{|\mathbf{k}|}{L}} \frac{|\mathbf{k}|^2}{\widehat{L}^2} E(\vec{\Psi}(\sigma e^{i\theta}))(\mathbf{k}) \right)^{1/2} \leq c\Gamma.
\end{aligned}$$

Thus  $\psi_j, j = 1, 2$  are analytic as functions of  $(\zeta, \mathbf{z})$  for  $(\zeta, \mathbf{z})$  in a neighborhood in  $\mathbb{C}^3$  of  $(0, \infty) \times \mathbb{R}^2$ . In particular  $\psi_j \in C^\infty((0, \infty) \times \mathbb{R}^2)$ . Therefore, the way we got the wave-vector formulation of our system automatically shows that  $\psi_j, j = 1, 2$  satisfy the system in the classical sense.  $\square$

**Remark B.1.** (i)  $\psi_j(t, \cdot) \rightarrow \psi_j(0, \cdot)$  as  $t \searrow 0$  in  $H^1(\Omega)$ .

(ii) If  $\psi_j(0, \cdot) \in H^2(\Omega), j = 1, 2$  then  $\vec{\psi}(0) \in V_1$ . This implies that  $\vec{\psi}(t) \rightarrow \vec{\psi}(0)$  as  $t \searrow 0$  in  $V_1$  which gives us that  $\psi_j(t, \cdot) \rightarrow \psi_j(0, \cdot)$  as  $t \searrow 0$  in  $H^2(\Omega)$ . By the classical Sobolev inequalities one obtains that  $\psi_j(t, \mathbf{x}) \rightarrow \psi_j(0, \mathbf{x})$  as  $t \searrow 0$  uniformly in  $\mathbf{x}$ .

## CHAPTER V

## CONCLUSION

This dissertation is the beginning of a research project. Our primary motivation in undertaking this study was to put on a firm mathematical ground the behavior of our system. Also the model system sits in an interesting position between 2D and 3D Navier-Stokes, so the problem may have some independent interest. In the first part of our study we defined a notion of weak solution, and showed using Galerkin methods the long-time existence and uniqueness of such solutions. In the second part we showed that our unique weak solution is in fact a classical solution. In addition we proved that the mentioned solution is time and space analytic. Next we plan on working on the convergence of a numerical scheme related to our model, estimating the errors in the same time. From physical point of view the time average of solutions are very important. Because of that one of our future plans is to estimate the energy norm of the time average of solution in terms of fluctuations.

## REFERENCES

- [1] A.F. Bennett and P.E. Kloeden, “The periodic quasigeostrophic equations: existence and uniqueness of strong solutions,” *Proc. Roy. Soc. Edin.*, vol. 93A, pp. 185-203, 1982.
- [2] J. Charney, “Geostrophic turbulence,” *J. Atmos. Sci.*, vol. 28, pp. 1087-1095, 1971.
- [3] J. Charney, R. Fjortoft, and J. von Neumann, “Numerical integration of the barotropic vorticity equation,” *Tellus*, vol. 2, pp. 237-254, 1950.
- [4] J. Charney and N. A. Phillips, “Numerical integration of the quasi-geostrophic equations for barotropic and simple baroclinic flows,” *J. Met.*, vol. 10, pp. 71-91, 1953.
- [5] P. Constantin, D. Cordoba, and J. Wu, “On the critical dissipative quasi-geostrophic equation,” *Indiana Univ. Math. J.*, vol. 50, pp. 97-107, 2001.
- [6] P. Constantin, A.J. Majda, and E. Tabak, “Formation of strong fronts in the 2d quasi-geostrophic thermal active scalar,” *Nonlinearity*, vol. 7, pp. 1495-1533, 1994.
- [7] C. Foias, “Lecture notes on the Navier-Stokes equations”, class notes, 2002-2004.
- [8] D.B. Haidvogel and I.M. Held, “Homogeneous quasi-geostrophic turbulence driven by a uniform temperature gradient,” *J. Atmos. Sci.*, vol. 42, pp. 2644-2660, 1980.

- [9] I.M. Held and V.D. Larichev, "A scaling theory for horizontally homogeneous baroclinically unstable flow on a beta plane," *J. Atmos. Sci.*, vol. 53, pp. 946-952, 1996.
- [10] I.M. Held, R.T. Pierrehumbert, S.T. Garner, and K.L. Swanson, "Surface quasi-geostrophic dynamics," *J. Fluid Mech.*, vol 282, pp. 1-20, 1995.
- [11] G. Lapeyre and I.M. Held, "Diffusivity, kinetic energy dissipation, and closure for the poleward eddy heat flux," *J. Atmos. Sci.*, vol. 60, pp. 2907-2916, 2003.
- [12] V.D. Larichev and I.M. Held, "Eddy amplitudes and fluxes in a homogeneous model of fully developed baroclinic turbulence," *J. Phys. Ocean.*, vol. 25, pp. 2285-2297, 1995.
- [13] R.L. Panetta, "Zonal jets in wide baroclinically unstable regions: Persistence and scale selection," *J. Atmos. Sci.*, vol. 50, pp. 2073-2106, 1994.
- [14] R.L. Panetta, E. Titi and M. Ziane, "Nonlinear stability of a quasi-geostrophic model of zonal jets," paper presented at the 996<sup>th</sup> AMS Meeting, Los Angeles, April 3-4, 2004.
- [15] J. Pedlosky, *Geophysical Fluid Dynamics*, 2nd ed. New York: Springer-Verlag, , 1987.
- [16] R. Salmon, *Lectures on Geophysical Fluid Dynamics*, Oxford, U.K.: Oxford University Press, 1998.

Supplemental sources consulted

P. Constantin and C. Foias, *Navier-Stokes Equations*, Chicago: University of Chicago Press, 1988.

C. Foias and R. Temam, “Gevrey class regularity for the solutions of the Navier-Stokes equations,” *J. Funct. Anal.*, vol. 87, pp. 359-369, 1989.

C. Onica and R.L. Panetta, “Forced two layer beta-plane quasi-geostrophic flow, part I: long-time existence and uniqueness of weak solutions,” submitted, 2005.

C. Onica and R.L. Panetta, “Forced two layer beta-plane quasi-geostrophic flow, part II: time and space analyticity,” submitted, 2005.

## VITA

**Name:** Constantin Onica

**Date of birth:** May 17, 1973

**Education:**

M.S., Al. I. Cuza University, Iasi, Romania, 1998

B.S., Al. I. Cuza University, Iasi, Romania, 1997

**Address:** Department of Mathematics, Texas A&M University, College Station, TX  
77843-3368

**e-mail:** onica@math.tamu.edu

**Membership:** Member of the American Mathematical Society.