

RUMINANT METHANOGENIC ACTIVITY IN THE UNITED STATES

BEEF CATTLE INDUSTRY

A Thesis

by

SARAH B. KLOPATEK

Submitted to the Office of Graduate and Professional Studies of
Texas A&M University
in partial fulfillment of the requirements for the degree of

MASTER OF SCIENCE

Chair of Committee,	Tryon A. Wickersham
Co-Chair,	Luis O. Tedeschi
Committee Members,	Jason E. Sawyer
	Todd R. Callaway
Head of Department,	H. Russell Cross

August 2016

Major Subject: Animal Science

Copyright 2016 Sarah B. Klopatek

ABSTRACT

In the U.S., gastrointestinal fermentation from cattle is estimated to account for approximately 25% of total anthropogenic related methane (CH₄) emissions. In addition, 2-8% of gross energy consumed by cattle is lost in the form of CH₄, representing an energetic cost to the animal. Thereby, in order to decrease greenhouse gas emissions (GHG) and improve the efficiency of cattle production additional research on gastrointestinal CH₄ emissions from cattle is needed. In ruminants carbohydrate (CHO) catabolism and nitrogen (N) utilization have a tremendous impact on ruminal methanogenesis. However, the impact of purified carbohydrates in the presence of a variety of N sources on rates of CH₄ and VFA production remains unknown. In order to determine these rates for use in predictive models of the ruminal fermentation, we formulated a fractional rate equation to fit the rate of CH₄ production and measured the concentration of CH₄ and VFA and using purified CHO with a variety of N sources in two *in vitro* mixed ruminal microorganism fermentation studies. In both studies, a CHO treatment × incubation time (IT) effect was observed for both VFA and CH₄ ($P < 0.01$). There was also a N × IT interaction for CH₄ production at 24 h in Study 2, where nitrogen free and NH₃ treatments produced greater concentrations of CH₄ than treatments with amino acids ($P < 0.01$). A nonlinear equation for the conversion of carbohydrates to CH₄ was able to fit starch treatments in Study 1 and glucose treatments in Study 2. Overall, this study demonstrated different fermentation patterns among all CHO and N sources and was the first step in determining rates for *in vitro* CH₄ production. Although cattle contribute with high amounts of anthropogenic GHG, they

are not the only methanogenic producing food source in the U.S. Rice and wild ruminants (e.g. bison, elk, deer) are also methanogenic producing food sources. The objective of this final study was to compare the efficiency of beef and milk production to pre-settlement wild ruminants and rice production on a kilogram of CH₄ emitted to kilogram human-edible protein production basis. Bison had the highest ratio of 13.93 kg CH₄: Protein, followed by elk (12.50) deer (6.66) and beef (2.47). Overall, wild ruminants emitted 296 to 564 percent more CH₄ per kilogram of human-edible protein produced than current beef cattle production systems. Rice yielded the second lowest CH₄ to human-edible protein ratio (0.83), followed by dairy cattle milk production (0.50). We believe, this analysis provides insight on the efficiency of methanogenic food sources that may aid in the development of regulatory guidelines of CH₄ production.

DEDICATION

I dedicate this thesis to my mother and father. Without their continual love and support both as parents and as professors, this thesis would not have been possible. Thank you

Ma'ma and Daddy, I love you both.

ACKNOWLEDGEMENTS

I would like to thank my undergraduate professors in the Animal Science department at the University of Arizona. Your time, dedication, and continual support gave me the foundation needed to succeed at Texas A&M.

Thank you to my professors at Texas A&M University. Your energy and teaching gave me the time and knowledge needed to find my place in a tumultuous and confusing world. I don't know where I would be without your guidance.

To my "boxing coach" Daryn, you helped me stand as an individual and instilled within me the courage to follow my dreams. Madam Curie remarked, "Nothing in life is to be feared, it is only to be understood. Now is the time to understand more, so that we may fear less." Thank you for taking away the fear, replacing it with understanding, and showing me how wonderful and exciting life is. You will always be in my heart.

Suzanne, thank you for being my other mom, my best friend, and for always providing me with encouragement. You once told me I would never influence the world by being like it. Thank you for teaching me how to be an individual and to stand on my own. I will make you proud Ma'ma Rogers.

Mom and Dad, if not for your love, nonstop encouragement, and scientific expertise I would not have been able to follow my passion in agricultural sustainability. I will always value you as parents, mentors, and scientists.

TABLE OF CONTENTS

	Page
ABSTRACT.....	ii
DEDICATION.....	iv
ACKNOWLEDGEMENTS.....	v
TABLE OF CONTENTS.....	vi
LIST OF FIGURES.....	viii
LIST OF TABLES.....	ix
CHAPTER I INTRODUCTION.....	1
CHAPTER II REVIEW OF THE LITERATURE.....	4
Introduction.....	4
The ruminant.....	5
Carbohydrate degradation.....	7
Methanogenesis.....	10
Methane mitigation strategies and impactors.....	12
U.S. cattle gastrointestinal CH ₄ production.....	19
Conclusion.....	24
CHAPTER III METHANE AND VFA PRODUCTION RATES BY IN VITRO MIXED RUMINAL MICROORGANISM FERMENTATIONS OF PURIFIED CARBOHYDRATES AND A VARIETY OF NITROGEN SOURCES.....	26
Introduction.....	26
Materials and methods.....	27
Results.....	31
Discussion.....	39
Conclusion.....	43

CHAPTER IV COMPARING U.S. METHANE GENERATING PROTEIN SYSTEMS.....	44
Introduction.....	44
Materials and methods.....	46
Results and discussion.....	52
Conclusion.....	60
CHAPTER V SUMMARY.....	61
REFERENCES.....	64

LIST OF FIGURES

FIGURE		Page
1.1	The fermentation of cellobiose by <i>R. albus</i> grown as a pure culture.....	11
1.2	The fermentation of cellobiose by <i>R. albus</i> grown in a co-culture with a methanogen.....	11
2.1	pH of fermentations for each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1.....	32
2.2	Methane production of fermentations of each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1.....	33
2.3	Volatile fatty acid (VFA) production of fermentations of each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1.....	34
2.4	pH of fermentations for each carbohydrate source (1600 mg/L) in mixed ruminal microorganism fermentation in Study 2.....	36
2.5	Methane production of fermentations of each carbohydrate source (1600 mg/L) in mixed ruminal microorganism fermentation in Study 2.....	37
2.6	Methane production of fermentations of each nitrogen source (900 mg N/L) in mixed ruminal microorganism fermentation in Study 2.....	37
2.7	Volatile fatty acid (VFA) production of fermentations of each carbohydrate source (1600 mg/ L) in mixed ruminal microorganism fermentation in Study 2.....	38
2.8	Volatile fatty acid (VFA) Production of fermentations of each nitrogen source (900 mg N/L) in Study 2.....	39

LIST OF TABLES

TABLE		Page
2.1	Fractional Rates of CH ₄ production among CHO Sources.....	35
3.1	Historic ruminant gastrointestinal CH ₄ emissions.....	54
3.2	Pre-settlement wild ruminant and current beef cattle gastrointestinal emissions of CH ₄ and production of human- edible protein.....	56
3.3	Dairy cattle gastrointestinal CH ₄ emissions and human-edible protein ratio.....	58
3.4	Rice CH ₄ emissions and human-edible protein.....	59
3.5	Methane emissions to human-edible protein yield.....	59

CHAPTER I

INTRODUCTION

Ruminants are able to degrade cellulose due to the symbiotic relationship they have with the native anaerobic microorganisms that inhabit their rumen. However, cellulose utilization under anaerobic conditions requires the regeneration of reducing equivalents, often by interspecies hydrogen transfer resulting in CH₄ production. Methane represents an energetic loss from 2 to 8% of gross energy to the animal, as well as a loss of profit to the producer (Dong et al., 2006). In 2014, the EPA listed beef cattle as the leading cause of agriculturally-related CH₄ emissions in the United States, accounting for approximately 38% of total agricultural CH₄ emissions (U.S. EPA, 2014). Furthermore, CH₄ has a 12-year atmospheric lifetime and global warming potential 25 times greater than that of CO₂ (U. S. EPA, 2016), thus strategies to mitigate enteric CH₄ production are of critical interest worldwide to environmentalists, as well as to cattle producers.

Many strategies have been evaluated to reduce CH₄ including modifying dietary composition, inclusion of ionophores, organic acids, and plant compounds, defaunation and vaccinations. These abatement strategies directly or indirectly target ruminal methanogen populations, resulting in varying degrees of efficacy. Although strides have been made in the reduction of methanogenesis, there is a continual need for ruminal CH₄ mitigating research.

Ruminal CH₄ production models can also be utilized as an instrument to help reduce emissions at the animal, farm, regional and global scale (McAllister et al., 2011). These models are not only vital from an agricultural systems perspective, but from both policy-making and regulatory perspectives. Although, predictive modeling has become an essential tool in animal nutrition, its accuracy is difficult to assess and at times is less precise than desired (McAllister et al., 2011). It is well known that carbohydrate catabolism and nitrogen (N) utilization have a tremendous impact on ruminal methanogenesis as well as ruminant nutrition. However, the impact of purified carbohydrates in the presence of a variety of N sources on rates of CH₄ and VFA production remains unknown. In order to clarify these rates for use in predictive models of ruminal fermentation, it is important to formulate fractional rate equations to fit the rate of CH₄ production. Ultimately, by constructing nonlinear rate equations for the conversion of carbohydrates to CH₄, the predictive accuracy of current CH₄ models can be improved upon.

Although cattle production provides the population with an abundant and complete source of protein (National Academy of Sciences, 2005; U.S. Department of Health and Human Services, 2016), some believe because cattle contribute to anthropogenic GHG that they are a non-sustainable food source (Pollan, 2002; Bittman, 2008). However, one factor that has been remiss from the dialog is that beef cattle have not always been the dominant source of agricultural CH₄ emissions nor are they the only current methanogenic food source.

Bison, elk and deer, like cattle, are ruminants and it has been theorized that prior to European settlement, wild ruminants produced sizable amounts of CH₄ (Subak, 1994; Hristov, 2012). In addition to wild ruminants the grain rice, one of the most heavily consumed food sources in the world, is methanogenic, constituting one of the largest sources of agricultural CH₄ emissions (van Groenigen et al., 2013; IPCC, 2007). Therefore, in order to understand and address the impending issues of food sustainability, the efficiencies of pre-settlement and current methanogenic agricultural food sources should be compared on an CH₄ output and human-edible protein basis. Ultimately, by performing innovative comparison studies, we can provide insight on the efficiency of methanogenic food sources and one day may aid in the development of environmental and governmental food sustainability guidelines.

CHAPTER II

LITERATURE REVIEW

Introduction

Ruminants are an evolutionary marvel because their symbiotic relationship with anaerobic microbes in their reticulorumen allows for the conversion of cellulose and other structural carbohydrates into meat, milk, and fiber. However, this ability to utilize forages comes at a cost. Methane (CH₄) is an inevitable end product of all ruminal fermentation and is a sink for reducing equivalents. Therefore, a certain amount of methanogenesis is inevitable from ruminal fermentation.

Methane is a potent greenhouse gas (GHG), with a global warming potential 23 times greater than that of carbon dioxide (U.S. EPA, 2016). In 2012, the Environmental Protection Agency (EPA) listed gastrointestinal fermentation from beef and dairy cattle as one of the leading cause of agriculturally related CH₄ emissions in the United States, accounting for approximately 25% of total agricultural CH₄ emissions (U.S. EPA, 2014). Additionally, CH₄ production is a cost to animal production, representing a loss of 2 to 8% of gross energy (Gerber et al., 2013). Therefore, it is of critical interest to cattle producers as well as environmentalists that CH₄ emissions from gastrointestinal fermentation be reduced.

Many strategies including diet modification, ionophores, and feed additives, have been used to reduce methanogenesis, with varying degrees of success (Van Nevel and Demeyer, 1977; Beauchemin and McGinn., 2006; Ellis et al. 2007). Over the last

quarter century, gastrointestinal CH₄ production from the entire beef cattle industry has increased by 0.6% while beef production has increased by 14%, meaning the U.S. produces more beef with fewer emissions. However, the continuing improvement of CH₄ mitigation will be dictated by the availability and quantity of feed, physiological state of the animals, governmental regulatory agencies, public perception, and the development of new technologies.

The ruminant

Food animals can be categorized as either non-ruminants or ruminants. Non-ruminants, such as pigs and horses, are postgastric fermenters and have simple, single chamber stomachs. Ruminants, such as cattle and sheep, are pregastric fermenters and have a complex, four chambered “stomach”s. Ruminant digestion differs from non-ruminant digestion as the majority of microbial fermentation occurs in the rumen, prior to passage through the gastric stomach (abomasum) opposed to non-ruminants where the majority of fermentation occurs in the large intestine or cecum. In cattle, the rumen is a blind pouch that is an anaerobic fermentation chamber with a volume of 100 to 180 L that comprises approximately 50% of the gastrointestinal tract volume. The Rumen is populated by a wide variety of microorganisms including protozoa, bacteria, and fungi that synergistically degrade and ferment feedstuffs. Remarkably, the rumen is one of the world’s richest microbial habitats in the world. In a droplet of rumen fluid there are more bacteria present than there have been people that have ever lived on earth (Russell

and Hespell, 1981; Prescott et al., 2005). Thus far, over 200 species of bacteria and 20 species of protozoa have been cultured in rumen fluid (Russell and Hespell, 1981).

The Ruminant microbial fermentation end-products include volatile fatty acids (VFA), microbial crude protein, NH_4 , CO_2 , and CH_4 . Volatile fatty acids are absorbed across the epithelium and serve as the primary energy source for ruminants. The most common VFA are acetate, propionate, and butyrate and their relative ratios vary according to dietary composition. Branched-chain VFA, include isovalerate, valerate, and isobutyrate, are also produced. Although BC VFA comprise only a small percentage of the total VFA production, they are imperative to ruminal syntrophic processes (Pitt et al., 1996).

Production of VFA decreases ruminal pH; to counteract this, ruminants attempt to maintain ruminal pH homeostasis by producing large quantities of saliva that contains bicarbonate, a buffer. Domestic cattle secrete between 100 to 150 liters of saliva per day, depending on diet and size of the animal (Bowen, 2009). If ruminal pH falls below 5.5, severe health problems can occur (Garrett et al., 1999).

Cattle typically meet the majority of their protein requirements by degrading and digesting ruminal microbes that pass out of the reticulorumen. This ability to utilize microbial crude protein derived from fermentation of forage enables cattle to effectively fill a unique environmental niche. Although, microbial protein is also synthesized in the large intestine, most herbivores cannot use this as a protein source because amino acids, dipeptides and tripeptides, the end products of protein digestion, cannot be absorbed from the large intestine.

Carbohydrate degradation

Cellulose is the most abundant polymer on land comprising 20-40% of plant matter on a DM basis (Leschine, 1995). Cellulose is a β -1,4 linked glucose homopolymer with glucose molecules oriented linearly, allowing polymerized molecules to stack (Leschine, 1995). Thus far, no vertebrate animals have been identified with the ability to produce cellulose-degrading enzymes, and only specific varieties of microbes, possess the enzymatic ability to hydrolyze β 1,4 glycosidic bonds (Watanabe and Tokuda, 2000).

Microorganisms have evolved several strategies to degrade cellulose. One strategy, is the release of cellulose degrading enzymes into the ruminal fluid. The other dominant strategy is direct adhesion. Carbohydrate-binding modules (CBM) adhere their enzymes directly to cellulose to create a maximum interface between the catalytic domain and the substrate before the enzyme diffuses from the cellulose particle (Wilson, 2011). In addition to CBM, Cellulosomes are multi-enzyme complexes that use scaffoldin subunits or cellulose-binding domains, to adhere carbohydrates to the microbial cell wall (Ding et al., 2008). Overall, adhesion strategies are advantageous to the microorganisms because they allow for the protection of their enzymatic resources by directly attaching to plant cell walls and closely capturing degradation products.

Pure cellulose is a biological rarity. In nature, cellulose is predominantly associated with hemicellulose, lignin, pectin, and proteins in a plant (Van Soest, 1994; Leschine, 1995). The degree to which cellulose is bound to other structural components affects nutrient availability and digestibility. Lignin, an organic polymer important to

plant structural integrity, is an anti-nutritional factor that slows down fiber degradation by ruminal microbes. As plants mature, lignin is deposited and cross-linked with cellulose within the cell-wall. Thus, as the plant matures, the ability of ruminal microorganisms to degrade cellulose decreases due to the increased lignification (Jung and Allen, 1995; Moore and Jung, 2001).

Both physical and chemical factors that alter the rumen environment can influence rate of cellulose degradation. One of the principal chemical factors impacting cellulose degradation is pH. Bacteria responsible for cellulose degradation, such as *Fibrobacter succinogenes*, *Ruminococcus albus*, and *Ruminococcus flavefaciens* are pH sensitive. If the pH declines below the optimum fibrolytic enzyme pH, 6.2 (Greve et al., 1984; Matte and Forsberg, 1992), fiber digestion is inhibited (Russell and Dombrowski, 1980; Hoover, 1986; Grant and Mertens, 1992).

Cellulose molecules can be oriented in tightly packed crystalline constructions, or more loosely arranged in amorphous structures. The percent of crystallinity within native cellulose ranges from 60-90% (Leschine, 1995). Although crystallinity may affect cellulose digestibility, it has been observed that surface area or cellulose may have a greater impact on digestibility (Weimer et al., 1990). Particles with larger surface areas have a higher rate of microbial colonization than smaller surface areas and thus have a higher rate of cellulose degradation (Weimer et al., 1990).

Plants primarily store energy in the form of starch (Van Soest, 1994). Starch is predominantly composed of amylopectin and amylose, also both glucose polysaccharides. Amylose is a linear α -1,4 glucopyranosid consisting of several hundred

chains (Van Soest, 1994). Amylopectin, is a much larger polymer than amylose and consists of both α -1,4-linkages and α -1,6 linkages (Van Soest, 1973). Incorporation of α -1,6 linkages causes branching in starch and increases accessibility to microbes, opposed to amylose which is structurally condensed. Unlike β -1,4 bonds, α -1,4 and α -1,6 bonds can be hydrolyzed by microbial or vertebrate enzymes. Factors that affect the rate of starch digestion include the proportion of amylose to amylopectin, and degree of crystallinity (Theurer, 1986). Starch, like cellulose, can be divided into amorphous or crystalline regions. Crystalline starch's compact structure makes it nearly impermeable to microbial attack, therefore as the percentage of crystalline structure increases, starch digestibility decreases (Sveinbjörnsson, 2006).

Processing methods such as steam flaking and grinding increase the rate and extent of starch degradation by ruminal microbes (Theurer, 1986). Amyolytic starch degradation from starch from cereal grains by both ruminal microbial and pancreatic enzyme sources are increased by processing methods. It is thought that the improvement in starch digestion by processing techniques is the primary reason for enhanced feed conversion of cattle fed high grain diets (Theurer, 1986).

As starch enters the rumen, starch-fermenting bacteria, such as *Prevotella ruminicola*, *Selenomonas ruminatium*, and *Streptococcus bovis*, begin to ferment starch into lactate and acetate (Ørskov, 1986; Russell and Rychlik, 2001). Because lactate has a low pKa (3.9), it dissociates into lactate and Hydronium ions which causes a decrease in pH. When the ruminal pH decreases below 6.0 many species of bacteria grow more slowly, including fibrolytic bacteria, but many amyolytic bacteria are not pH sensitive

until pH falls below 5.5 (Reece et al., 2015). In addition, starch fermenting bacteria have a generation time significantly faster than fiber fermenting bacteria (Hungate, 1966). Accordingly, inclusion of starch in the diet alters the rumen environment in favor of starch fermenting microorganisms, inhibiting fiber digestion (Piwonka and Firkins, 1996).

Starch is added to the diet to increase energetic efficiency by increasing the propionate to acetate ratio. Propionate is gluconeogenic and provides more moles of ATP per mole of propionate than acetate (Russell and Rychlik, 2001). The rapid fermentation of starch can also cause ruminal health issues. If the VFA production rate is greater than the rate of absorption, ruminal pH begins to fall. If pH falls below 5.5 then ruminal acidosis, either acute or subacute ruminal acidosis, can occur (Blood and Radostits, 1989; Plaizier et al., 2008). Clinical signs of acidosis depend on the severity of the pH depression as well the time the pH remains depressed. Signs include poor body condition, decreased rumen motility, laminitis, abscesses, and even result in the mortality of the animal (Blood and Radostits, 1989; Underwood, 1992).

Methanogenesis

Ruminal methanogens are autotrophic archaea that live in symbiosis with the ruminal consortium of protozoa, fungi and bacteria (Van Soest, 1982). Two classes of methanogenic archaea predominate the rumen, *Methanobacteriales* and *Methanomicrobiales* (St-Pierre et al., 2015); thus far, microbiologists, have isolated seven species from rumen fluid (Janssen and Kirs, 2008).

Ruminal methanogens produce CH₄ as an end product of ruminal fermentation. Although, the production of CH₄ represents an energetic loss to the animal, the presence of methanogens in the rumen are important for microbial energetic efficiency. Aerobic organisms produce ATP via oxidation phosphorylation, but because the rumen is anaerobic, microbes cannot utilize oxygen as the terminal electron acceptor (Murray, 2012). Coupled redox reactions are utilized for energy generation within the rumen, utilizing carbon, instead of oxygen, as the final electron acceptor (Russell, 2002). Through the process of interspecies hydrogen transfer, CO₂ and hydrogen protons are converted to CH₄ by methanogens (Scheifinger et al., 1975; Latham and Wolin, 1977). Methanogens are a major means of reducing equivalent disposal. By keeping the partial pressure of H₂ low via hydrogenase, methanogens allow for ruminal bacteria to more efficiently metabolize energy.




Figure 1.1: The Fermentation of cellobiose by *R. albus* grown as a pure culture. (Taken from Russell, 2002).




Figure 1.2: The Fermentation of cellobiose by *R. albus* grown in a co-culture with a methanogen. (Taken from Russell, 2002).

In a pure culture of *R. albus* (Fig. 1.1) hydrogenases are unable to oxidize NADH, thereby alcohol production becomes the alternative method for reducing equivalent disposal reducing the total ATP yield. In contrast when grown in a co-culture of *R. albus* and a methanogen (Fig. 1.2), the methanogen is able to use hydrogens produced by the reductions of G-6-P to Pyruvate and Acetyl CoA to acetate. Ultimately, this allows for the regeneration of reducing equivalents, enabling the production of two acetate and two ATP. Accordingly, methanogens are crucial for microbial energetic efficiency and the complete elimination of methanogens from the rumen would be detrimental to both microbes as well as the host animal.

Methane mitigation strategies and impactors

Despite the intrinsic necessity for a certain level of CH₄ production in the rumen, minimizing CH₄ production is a goal of all producers. Gastrointestinal CH₄ production can be inhibited by directly inhibiting methanogens or indirectly by decreasing substrate availability to the methanogens. Numerous strategies have been developed and implemented to reduce gastrointestinal CH₄ production with varying degrees of efficacy.

Dietary impacts

One of the most utilized strategies to reduce methanogenic activity is diet modification. Dietary factors that influence rate of passage, rate of fermentation, rumination or pH, alter H⁺ concentrations affecting microbiota present, including methanogens. Specific dietary factors include DMI, metabolizable energy intake, digestibility, and the percent of forage in the diet (Ellis et al., 2007).

Dry matter intake may account for as much as 52 to 64% of the variation in daily CH₄ production (Boadi et al., 2002; Hammond et al., 2009). In general, as DMI increases total CH₄ production increases (Blaxter, 1967; McAllister, et al. 1996). However, when intake is increased from maintenance, the amount of energy lost as CH₄ per unit of feed intake decreases (Blaxter, 1967). Studies have indicated that when forage DM intake increases by 30% kg^{-0.75}, Y_m (percent of gross energy lost in the form of CH₄) values decreased from 7% to 6.5% (McAllister et al., 1996).

In addition to feed intake, diet digestibility also has a substantial impact on CH₄ production (Hegarty et al., 2010). Factors that affect the digestibility of the feedstuff include, but are not limited to, plant type, plant maturity, physical processing methods, lignification, neutral detergent fiber content, and acid detergent fiber content. As plants mature, the fraction of insoluble fiber increases, which causes a decrease in plant digestibility. Thus, ruminal CH₄ production generally increases concurrently with the maturity of the forage fed (Tyrrell and Moe, 1992; Jung and Allen, 1995; Boadi et al., 2002). Digestibility of feedstuffs can be improved by physical processing methods (e.g. chopping, pelleting, and grinding) which can lower GHG emissions. Studies have indicated that CH₄ production is lower for finely ground and pelleted forages than chopped and/or long stem forages (Hironaka et al. 1996).

Forage type also has an effect on ruminal methanogenesis (e.g., legume forages generally yield lower CH₄ production from ruminal fermentation than do grass forages) (Varga et al. 1985). Varga et al. (1985) suggested that because the legume diet had a higher proportion of soluble carbohydrates compared to the grass diet that the grass diet

was favorable to ruminal methanogenesis. Strategies that have been recommended to decrease emissions without negatively affecting intake include replacing grass silage with either corn or legume silages, and incorporate highly digestible fibers to the diet, such as beet pulp (Kujawa, 1994; Hristov et al., 2013).

Increasing the proportion of concentrate to the diet significantly effects methanogenic activity (Moe and Tyrrell, 1979; Kebreab et al., 2008). Johnson and Johnson (1995) reported that animals fed at maintenance lost on average 6-7% Y_m on 100 percent forage diets compared to 2-3% Y_m for animals fed high concentrate diets (up to 90 percent). This decline in CH_4 production caused by the inclusion of concentrate feedstuffs happens because of several factors. Adding concentrate to cattle diet increases the propionate to acetate ratio by providing substrate to amylolytic bacteria (van Kessel and Russell, 1996). When propionate is produced the hydrogen availability for methanogens is decreased, ultimately decreasing CH_4 production (van Kessel and Russell, 1996). Additionally, the decreased pH and increased rate of fermentation caused by the inclusion of dietary starch, may decrease methanogen activity by inhibiting ciliate protozoa (Demeyer, 1975; McAllister et al., 1996). Methanogenic archaea are metabolically and synergistically associated with ciliate protozoa (Finlay et al., 1994). Due to protozoa's intricate relationship with ruminal methanogens, it has been estimated that ciliate protozoa are responsible for 9 to 37% of the CH_4 production within the rumen (Newbold et al., 1995). Thereby, if protozoa are reduced methanogenesis is impacted.

Forage generally has a slower rate of digestion compared to starch and prolonging the ruminal residence time of feed particles increases the amount of CH₄ produced per unit of feed digested (McAllister, 1996). Overall, incorporating concentrates into cattle diets decreases methanogenic activity and is a crucial strategy for providing sustainable beef that limits GHG emissions.

Ionophores

Monensin is a naturally occurring polyether ionophore produced from *Streptomyces cinnamonensis*. The benefits of ionophores to ruminants occur because they alter the end products of the ruminal fermentation through secondary effects of a disruption of the ion gradients maintained by Gram-positive bacteria, killing the gram-positive bacteria (Russell and Strobel, 1989). Ionophores were first utilized to help combat coccidiosis in the poultry industry. When poultry litter was fed to cattle, it led to the discovery that ionophores inhibited methanogenesis. Early research determined that monensin resulted in improvements in feed utilization, increasing feed to gain ratios up to 17% (Raun et al., 1976). *In vitro* research determined that monensin did not have a direct effect on methanogens, but rather inhibited of formate degradation decreasing hydrogen availability for methanogens (Van Nevel and Demeyer, 1977). Today, monensin is the predominant ionophore used in the feedlot cattle industry increasing feed efficiency by 6% and 15% for feedlot and grass-fed animals, respectively (Russell, 2002).

Chlorinated Hydrocarbons

Chlorinated hydrocarbons inhibit CH₄ production both *in vitro* and *in vivo* (Leng, 2014). One of the most successful chlorinated hydrocarbons tested *in vivo* is bromochloromethane (BCM). When BCM was added to ruminal fluid *in vitro*, it reduced CH₄ production by 94% (Martinez-Fernandez et al., 2015). In an *in vivo* study, when BCM was included in Brahman cattle diets, the Y_m values were significantly decreased as BCM inclusion levels increased (Tomkins and Hunter, 2004).

The chlorinated hydrocarbon, 9, 10-anthraquinone (AQ) has also been shown to decrease CH₄ production (Garcia-Lopez et al., 1996; St-Pierre et al., 2015) by directly affecting methanogenic bacteria (Kung et al., 2003). Inclusion of AQ in ovine diets decreased CH₄ emissions without compromising digestibility or animal health (Kung et al., 2003). However, long-term studies are needed to determine the longevity and persistence of efficacy of AQ as a CH₄ abatement strategy for cattle.

3-Nitrooxypropanol

The compound 3-Nitrooxypropanol (NOP) is a structural analog to methyl-coenzyme M (MCR), a cofactor involved in the final reduction stages of methanogenesis (Haisan et al., 2014). NOP inhibits MCR, subsequently causing a decrease in methanogenesis (Van Nevel and Demeyer, 1995). During *in vivo* trials, NOP was shown to reduce CH₄ emissions in lactating dairy cows without decreasing DMI or milk production (Haisan et al., 2014). However in a contrasting study, although a reduction of CH₄ was observed the addition of NOP also reduced diet digestibility (Reynolds et al.,

2014). Overall, additional research is needed to determine NOP dose, delivery method, as well as the longevity of efficacy (Reynolds et al., 2014).

Vaccinations

One of the most recent CH₄ mitigating strategies being investigated is an anti-methanogenic vaccination. In one vaccine trial, sheep were assigned one of two different treatments, either a 3-methanogen cocktail or a 7-methanogen cocktail (Wright et al., 2004). It was thought that the vaccine would use the ruminant's immune system to produce antibodies against methanogens and directly inhibit CH₄ production. The 3-methanogen cocktail reduced CH₄ production per kg dry matter intake by 7.7%, but the seven-methanogen cocktail had no effect on methanogenic activity. In a later trial, a vaccine based on five-methanogen strains was developed and administered to sheep at 0, 28, and 103 days (Williams et al., 2009). Although the vaccine targeted 52% of the known methanogen population, total CH₄ output actually increased by 18%. Currently, it does not appear that vaccination is a viable CH₄ abatement strategy.

Fats

Unsaturated fats, which are often included in the ruminant diet, have inhibitory effects on ruminal CH₄ emission. Unsaturated fatty acids may be used as hydrogen acceptors as an alternative to the reduction of carbon dioxide. In addition, fatty acids have the ability to directly inhibit methanogens by binding to the methanogen cell membrane interrupting CH₄ transport (Dohme et al., 2001). The addition of fat is also thought to decrease CH₄ by decreasing protozoa (Newbold et al., 2015).

Refined soy oil, linseed oil, coconut oil and sunflower oil have all been shown to reduce gastrointestinal CH₄ (Jordan et al., 2006; Beauchemin and McGinn., 2006; McGinn et al., 2004). *In vitro* studies demonstrate that the addition of fat, at 5.3%, reduced total CH₄ production by as much as 20% (Dohme et al., 2001). Overall, the addition of fats to the diet has been considered a more favorable means of CH₄ mitigation for it is more natural than chemical additives (Toprak, 2015). However, if dietary fat is fed in excess, feed intake and fiber digestibility decreases, which significantly impacts animal performance and lowers feed efficiency (McGinn et al., 2004).

Dicarboxylic organic acids

Dicarboxylic Organic acids such as malate, aspartate, and fumarate have been evaluated as dietary additives to improve animal performance and decrease methanogenesis (Newbold and Rode, 2006). It was hypothesized that organic acids would increase the propionate to acetate ratio, thereby decreasing H₂ availability for methanogens. However, the effectiveness of organic acids as a cattle CH₄ mitigating strategy are highly variable between experimentations. When the organic acids DL malate and fumarate were added to ruminal *in vitro* fermentations there was little effect on CH₄ concentrations (Callaway et al, 1997). Interestingly, when DL malate was fermented in the presence of monensin, CH₄ production decreased as inclusion levels of DL malate increased (Callaway et al., 1997). In an *in vivo* study when 100 g/kg fumarate was supplemented to growing lambs, depending on type of encapsulation of the acid, CH₄ emissions were reduced by 62% to 76% (Williams et al., 2009).

Contrastingly, in other *in vivo* trials when fumarate was fed to growing beef cattle it was not found to significantly reduce CH₄ emissions (McGinn, 2004; Beauchemin et al., 2010). Ultimately, additional studies need to be performed to determine the effectiveness of dicarboxylic organic acids usage in large scale cattle operations.

Defaunation

The anti-methanogenic effects of many feed additives and dietary treatments have been directly or indirectly associated with decreasing protozoa activity (Ffoulkes and Leng, 1988; Morgavi et al., 2012; Newbold et al., 2015). Elimination of protozoa (defaunation) has been suggested as a way to mitigate CH₄ emissions strategy (Newbold et al., 1995; Boadi et al., 2004; Hristov and Jouany, 2005), and the decrease in CH₄ production in the absence of protozoa has been observed both *in vitro* and *in vivo*. Studies have reported a range in CH₄ reduction from 5 to 20% (Hegarty, 1999; Martin et al., 2010). Although short term trials have demonstrated efficacy in reducing methanogen populations no long term defaunation trials have demonstrated efficacy.

U.S. cattle gastrointestinal CH₄ production

The beef cattle production chain can be divided between sectors including cow-calf, stocker, and feedlot. Life cycle analysis have determined that It was determined that the cow-calf system accounted for about 80% of total GHG emissions and the feedlot system only 20% (Beauchemin et al., 2010). Of the cow-calf sector approximately 84% of gastrointestinal CH₄ was from mature cows. A review by Broucek (2014) determined that CH₄ emissions from beef cattle ranged from 161 to 323

g of CH₄ per day. For mature beef cows, emissions range from 240 g to 396 g of CH₄ per day (Broucek, 2014). It has been suggested that to reduce emissions, additional CH₄ mitigation research should be devoted to the cow-calf sector.

U.S. EPA ruminant CH₄ estimates

There are two dominant factors when estimating livestock production of CH₄, average daily feed intake as gross energy (GE; MJ/d) and CH₄ conversion rates (Y_m ; Crosson et al., 2011). Animal intake and digestibility are the factors impacting of Y_m , while DMI is the largest contributing factor for total CH₄ production (Hristov et al. 2013). Dry matter intake for cattle varies depending on diet, age, sex, region and stage of production. International Panel of Climate Change (IPCC) assessments have estimated the Y_m of feedlot cattle 3.0 ± 1.0 while cattle consuming temperate-climate grasslands were estimated at 6.5 ± 1.0 (IPCC 2006).

Using IPCC's calculation and Tier 2 methodology (characterization of diets and animal growth curves for each category) the U.S. EPA estimated that for the year 2012 gastrointestinal CH₄ emissions for the U.S. cattle industry (both dairy and beef) were $6.71 \text{ CH}_4 \text{ Tg yr}^{-1}$. For beef cattle 4.79 Tg yr^{-1} , approximately 170 g of CH₄ per head per day. In total, gastrointestinal CH₄ emissions from livestock in the U.S. constituted 25% of total U.S. anthropogenic GHG emissions.

Although this contribution may seem high, it is consistent with other estimates. The Emissions Database for Global Atmospheric Research (EDGAR) estimated that total U.S. cattle gastrointestinal CH₄ emissions for 2005 were 6.45 Tg yr^{-1} , which was within 5% of EPA's assessment. Hristov et al. (2014) using a "bottom-up" approach,

similar to the U.S. EPA, estimated that total gastrointestinal CH₄ emissions from the total beef and dairy cattle herd to be were 6.24 Tg yr⁻¹ which was comparable to current, U.S. EPA estimates.

However, other reports have suggested that the U.S. EPA underestimated ruminant GHG emissions. Using a “top-down” approach, Miller et al. (2013) concluded that the current U.S. EPA assessment substantially underestimated ruminant CH₄ emissions. In this study, U.S. anthropogenic CH₄ sources were estimated from atmospheric CH₄ observations, extensive spatial datasets, and a high-resolution atmospheric transport model. According to Miller’s estimates, U.S. cattle CH₄ emissions from gastrointestinal fermentation were approximately 12.7 ± 5.0 Tg yr⁻¹, which was double the U.S. EPA assessments. However, this assessment was unable to be substantiated by animal scientists (Hristov et al., 2014).

Comparing emissions: Grass-fed vs. grain-fed beef

In the past 10 years, the demand for grass-fed beef has grown annually at a rate of 25-30% (Windrock International, 2012), in part due to the belief that grass-fed is more environmental friendly than conventionally raised beef (Walsh, 2008). Despite the recent increase in demand for grass-fed beef a life cycle analysis has demonstrated that grain-fed beef has a lower environmental footprint than grass-fed beef systems (Crosson et al., 2011; Peters et al, 2011; Capper, 2012).

Capper (2012) determined that if the U.S. resorted to consuming only grass-fed animals to meet the current beef demands, the number of cattle would need to increase by 64.6 million animals and this would require more than 200,000 square miles be

devoted to beef rearing, equivalent to 75% of the state of Texas (Capper, 2012). To produce a billion kg of conventional beef, production systems require 56.3% fewer animals, 24.8% less water, 55.3% less land, and require 71.4% of the fossil fuels compared to grass-fed systems (Capper, 2012). To switch to an all grass-fed beef from a GHG standpoint CH₄ emissions would increase by 0.35 Tg yr⁻¹ (Capper, 2012). Overall, the combination of technologies and management strategies used in conventional cattle production greatly reduce resource use and GHG emissions per unit of beef produced as compared to grass-fed operations (Capper, 2012).

Trends in the industry

Over the last several decades the beef industry has made tremendous improvements in animal efficiency. In 1977 it took 25% more animals to produce the same amount of beef than in 2007 (Capper, 2011). In addition, the time to produce beef, from calving to harvest, has been dramatically reduced. Today cattle on average are harvested at 485 days compared to 30 years ago when cattle were harvested at approximately 609 days (Capper, 2011). Ultimately, with improved genetics, nutrition, and management, cattle are grown more efficiently and each animal produces greater quantities of edible beef (Capper, 2011). Today's beef production requires 81.4% of feedstuffs, 67% of the land resources and 87.9% of water as compared to systems from the late 1970's (Capper, 2011).

According to the U.S. EPA from 1990 to 2012, emissions from beef cattle gastrointestinal fermentation increased by 0.6 percent (US EPA, 2014). Yet, beef cattle populations actually declined by 5 percent while beef production increased 14 percent

(U.S. EPA, 2014). This indicates that although emission factors per head are increasing, emission factors per unit of product are going down and the U.S. is able to produce more beef with fewer emissions.

GHG emissions for protein sources-rice and wild ruminants

The world population is expected to grow to 9.5 billion people by 2050 (FAO, 2009). In order to meet the nutritional needs of a growing population, world-wide food production must increase by 100% and protein resources need to increase by 70% (FAO, 2011). However, in order to counter the effects of climate change, GHGs need to be immediately reduced, rendering food production a conundrum. Although there is no food production “silver bullet”, life cycle analyses can be performed to help determine which food sources produce the greatest amounts of specific nutrients at the lowest GHG impact.

While beef production produces a significant amount of CH₄, it is not the only methanogenic food source. Rice, one of the most widely consumed human foodstuffs in the world, produces large quantities of CH₄ (Yan et al., 2009; Neue, 2014). Specifically, global rice production accounts for 31-112 Tg CH₄ yr⁻¹ compared to the estimate of worldwide total ruminant production of 76 to 92 Tg CH₄ yr⁻¹ (IPCC, 2007). In the United States rice is the third largest agricultural CH₄ source, producing more than 0.35 Tg CH₄ yr⁻¹ (U.S. EPA, 2014).

Historically and currently, cattle are not the only large ruminants in the United States. Wild ruminants, such as bison, elk, and deer produce large quantities of CH₄ on a per head basis (Crutzen et al., 1986; Galbraith et al., 1998; Kelliher and Clark, 2010).

It has been estimated that wild ruminant produce 9% of GE as CH₄ (Crutzen et al., 1986), which is significantly higher than either grass-fed or grain fed cattle, 6.5% and 3% respectively (Dong et al., 2006). Currently, wild ruminants contribute a low proportion of U.S. greenhouse gases, approximately 0.28 Tg yr⁻¹ (Hristov, 2012). However, prior to European settlement (pre-settlement) approximately 30-75 million bison inhabited the United States (McHugh, 1972; Dary, 1989; Isenberg, 2001). Pre-settlement CH₄ emissions for wild ruminants for the contiguous U.S. has been estimated from 2.9 to 7.3 Tg yr⁻¹, depending on assumed bison population size (30-75 million). Assuming a population of 50 million, bison would have historically produced 86 percent of the present-day CH₄ emissions from domestic ruminants in the U.S. (Hristov, 2012). Overall, this points out that there has always been CH₄ production from a large ruminant population in North America and brings into question whether or not today's cattle production should be condemned for their associated GHG emissions.

Conclusion

Cattle production, ruminant nutrition, GHG production, and food security and sustainability are complex and inextricably linked. With continual improvements and improvements in technology and beef cattle management, beef production in the U.S. has become more efficient. Currently, the U.S. is able to produce more beef, with fewer animals and resources, without increasing herd gastrointestinal CH₄ emissions (Capper, 2011; U.S. EPA, 2014). Although cattle production requires a significant investment of resources, it still plays an important role in the United States food supply, and

production of beef and dairy when viewed as part of the larger world-wide food supply is a sustainable source of high quality protein for a growing world population.

CHAPTER III

METHANE AND VFA PRODUCTION RATES BY *IN VITRO* MIXED RUMINAL MICROORGANISMS FERMENTATIONS OF PURIFIED CARBOHYDRATES AND A VARIETY OF N SOURCES

Introduction

Ruminants are able to degrade cellulose due to the symbiotic relationship they have with the native anaerobic microorganisms that inhabit their rumen (pre-gastric fermentation chamber). However, cellulose utilization under anaerobic conditions requires the regeneration of reducing equivalents, often by interspecies hydrogen transfer resulting in CH₄ production. Methane represents an energetic loss from 2 to 8% of gross energy to the animal, as well as a loss of profit to the producer (Dong et al., 2006). In 2012, the U.S. EPA described gastrointestinal fermentation from livestock, particularly beef cattle, as one of the leading causes of agriculturally-related CH₄ emissions in the United States, accounting for approximately 25% of total anthropogenic CH₄ emissions (U.S. EPA, 2014). Furthermore, CH₄ has a 12-year atmospheric lifetime and global warming potential 25 times greater than that of CO₂ (U.S. EPA, 2016), thus strategies to mitigate gastrointestinal CH₄ production are of critical interest worldwide, to environmentalists, as well as to cattle producers.

Ruminal CH₄ production models can be utilized as an instrument to help reduce emissions at the animal, farm, regional and global scale (McAllister et al., 2011). These models are not only vital from an agricultural systems perspective, but from both policy-

making and regulatory perspectives. Although, predictive modeling has become an essential tool in animal nutrition, its accuracy is difficult to assess and at times is less precise than desired (McAllister et al., 2011). Carbohydrate degradation kinetics have been extensively researched (Groot et al., 1996) and are incorporated into the Cornell Net Carbohydrate and Protein System (CNCPS) model (Russell et al., 1992; Sniffen et al., 1992); however, research has not established a definitive link between the kinetics of carbohydrate degradation and the rate and extent of CH₄ production. Therefore the objective of this study was to 1) determine the concentrations of VFA and CH₄ from mixed ruminal microorganism degradation of purified carbohydrate substrates fermented in the presence of a variety of nutrient conditions and 2) formulate a fractional rate for the conversion of carbohydrates into CH₄.

Materials and methods

Study 1: Methane and VFA production rates by in vitro mixed ruminal microorganism fermentations of purified carbohydrates

All animals were maintained in accordance with a protocol approved by the Southern Plains Agricultural Research Center Animal Care and Use Committee. One Holstein steer (BW 550 kg) and one Jersey cow (BW 360 kg) were provided *ad libitum* access to water, minerals, and Bermudagrass pasture at all times. Ruminal contents were collected by hand from at least five locations (at random) from the ventral sac of the rumen at 0700. Immediately after removal from the rumen, contents were squeezed through a fine mesh nylon strainer (Reaves and Co., Durham, NC) and pooled together,

filling a 1000 ml flask (500 ml per animal). Ruminal fluid was then transported to the laboratory and incubated at 39 °C for 45 min, to allow gas production to buoy feed particles to the top of the flask and protozoa to sediment to the bottom. The middle layer of mixed ruminal fluid was combined (33% vol/vol) with an anoxic basal medium containing (per liter): 292 mg K₂HPO₄, 202 mg KH₂PO₄, 436 mg NH₄SO₄, 480 mg NaCl, 100 mg MgSO₄·7H₂O, 64 mg CaCl₂·H₂O, 4,000 mg Na₂CO₃, 600 mg cysteine hydrochloride (Cotta and Russell, 1982). The resultant suspensions were anaerobically transferred to 18 × 150 mm Balch tubes (Bellco Glass, Vineland, NJ; 10 ml per tube) that contained 0 or 0.5 grams of carbohydrate substrate (amorphous cellulose, corn starch, or glucose), that were flushed under a CO₂ gas phase. Amorphous cellulose was cordially provided by Dr. P. Wiemer (USDA-ARS Dairy Forage Research Center). The amorphous structure of the cellulose was determined by x-ray crystallography (Isogai and Atalla, 1991). Tubes in triplicate for each of 8 time points (n = 24 tubes/substrate) were then sealed using butyl rubber stoppers with aluminum crimps and incubated for 24 h at 39 °C under a CO₂ headspace. Samples were collected from tubes (n=3 at each time point) after 0, 1, 2, 4, 8, 12, 18, and 24 h of incubation. Prior to liquid sample removal after 4, 12, and 24 h of incubation, a headspace gas sample (1.0 ml) was removed from each tube via gastight syringe and analyzed for CH₄ using a Gow Mac thermal conductivity series 550 gas chromatograph (Gow Mac Instrument, Bridgewater, NJ) equipped with a Carbosieve S 8100 column (Supelco, Inc., Bellefonte, PA). The gas flow (N₂) was 20 ml/min, and the column temperature was 125 °C and the detector temperature was 150 °C. Immediately upon opening, pH for each fermentation tube was

measured using an pH meter (Orion 2 Star) and ruminal fluid samples (5 ml) were collected and centrifuged ($10,000 \times g$, 10 min, 24 °C) to remove particulate matter, and the resultant supernatant was frozen at -20 °C until further analysis. Volatile fatty acids were determined using an Agilent 7890A gas chromatograph equipped with an FID detector, and an Agilent 7693 autosampler. Agilent DB-FAPP capillary columns were used and the inlet temperature was 230 °C, the oven temperature ranged from 40 °C to 200 °C, and the detector was heated to 300 °C.

Study 1 was conducted as a completely randomized design with a 6×3 factorial treatment structure. Factors included incubation time (IT) and carbohydrate (CHO) as fixed effects, and triplicate was treated as a random effect. Carbohydrates included corn starch, purified glucose and amorphous cellulose. Data were analyzed using the mixed models procedure of SAS[®] 9.4. Model effects included IT and CHO and all 2-way interactions. Means within time were separated using Tukey's HSD (honest significant difference) using the PDMIX800 macro of SAS. The exponential equation, $[CH_4]=A-B \cdot \exp^{-k_f \cdot \text{time}}$, was formulated to fit CH₄ concentration using PROC NLIN of SAS. K_f was representative of the fractional rate, A was the asymptote, and B was the slope.

Study 2: Methane and VFA production rates by in vitro mixed ruminal microorganism fermentations of purified carbohydrates and a variety of N sources.

All animals were maintained in accordance with a protocol approved by the Texas A&M University Animal Care and Use Committee. Two ruminally cannulated Angus cross steers (BW 350kg) were provided *ad libitum* access to water and minerals at all times. Rumen fluid was collected from cattle fed 0.9 kg/d of dried distiller's grains

with *ad libitum* access to pasture and Bermudagrass hay at the time of ruminal fluid collection. Ruminal contents were collected by hand from at least five locations (at random) in the ventral sac of the rumen at 0700. Ruminal contents were squeezed through 8 layers of cheesecloth and transported to the laboratory and prepared as described in Study 1.

Rumen fluid containing mixed ruminal bacteria was anaerobically transferred (33% v/v) to nitrogen-free anoxic media as described above (Chen and Russell, 1991). Mixed ruminal microorganism media was subdivided into four aliquots, based upon added nitrogen source. Each aliquot contained 0 mg/L added nitrogen (NF; negative control), or 900mg/ L of either: ammonia (NH₃), casamino acids and trypticase (AA), or an equimolar mixture of NH₃ and the casamino acids and trypticase mixture (AA+NH₃). Each ruminal fluid aliquot containing these nitrogen sources were anaerobically transferred into Balch fermentation tubes that contained one of the three carbohydrate sources used previously (amorphous cellulose, glucose, starch) to achieve final concentrations of 1600 mg/L. Fermentations of each N source with no added carbohydrate were used to estimate the contribution of the residual carbohydrate in the ruminal fluid and these negative controls were subtracted from production values.

Triplicate tubes (n = 3) for each time point (n = 36 tubes/substrate) were then sealed using butyl rubber stoppers with aluminum crimps and incubated for 24 h at 39 °C under a CO₂ atmosphere. Headspace gas samples and pH were measured at times 4, 12, and 24 h, samples were frozen for later analysis. Immediately upon opening, pH for each fermentation tube was measured using an pH meter (Orion 2 Star) and ruminal

fluid samples (5 ml) were collected and centrifuged ($10,000 \times g$, 10 min, 24 °C) to remove particulate matter, and the resultant supernatant was frozen at -20 °C until further analysis. Procedures for gas and VFA measurements were as described above.

Study 2 was implemented as a randomized complete block design with a 3×4 factorial treatment structure, and day was treated as a blocking factor. Data were analyzed using PROC MIXED of SAS. Model effects included the fixed effects of IT, N and CHO source, and all 2- and 3-way interactions. Means within time were separated using Tukey's HSD in the PDMIX800 macro of SAS. Estimation of exponential production of CH₄ production was determined using a nonlinear regression. The exponential equation, $[CH_4]=A-B*\exp^{-k_f*time}$, was formulated to fit CH₄ concentration using PROC NLIN of SAS. K_f was representative of the fractional rate, A was the asymptote, and B was the slope. Equations for both studies were divided by CHO treatment.

Results

Study 1

There was a CHO treatment effect and an IT \times CHO treatment interaction on pH ($P < 0.01$; Figure 2.1). During fermentation there were no differences in pH between substrates until 2 h, when glucose presented with lower pH, followed by starch and cellulose ($P < 0.05$). The pH declined for all CHOs over the 24 hour period ($P < 0.05$). By 24 h glucose and starch fermentations reached their nadir at pH's of 4.6 and 4.9, respectively; whereas cellulose fermentations was 5.6 ($P < 0.05$). During the

fermentation the pH did fall below 5.5 suggesting that fibrolytic bacteria may have been inhibited (Russell and Dombrowski, 1980; Hoover, 1986).




Figure 2.1: pH of fermentations for each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1

There was both a CHO treatment effect, and an IT \times CHO treatment interaction for CH₄ production ($P < 0.01$; Figure 2.2). Methane production differed between substrates at all time points except at 1 h. By 2 h, glucose was fermented the most rapidly and produced significantly higher concentrations of CH₄ than both starch and cellulose ($P < 0.05$). Fermentations containing glucose continued to produce greater concentrations of CH₄ through 8 h ($P < 0.05$). Starch fermentation CH₄ concentrations at 12 h were 29% higher than glucose and 52% greater than cellulose ($P < 0.05$). By 18 h, starch containing fermentations contained the greatest concentration of CH₄, followed

by cellulose, then glucose ($P < 0.05$). At 24 h, cellulose concentrations were similar to starch ($P > 0.05$) and 49% greater than glucose ($P < 0.05$).




Figure 2.2: Methane production of fermentations of each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1

Volatile fatty acid (VFA) production demonstrated a CHO treatment effect and an IT \times CHO treatment interaction ($P < 0.01$; Figure 2.3). From 2 h through 4 h fermentations of glucose produced more VFA than those of starch or cellulose ($P < 0.05$). However, at 8 h starch fermentations contained the highest concentrations of VFA; concentrations were 32% greater than glucose ($P < 0.05$) and 96% higher than cellulose ($P < 0.05$). At 12 h VFA concentrations produced from fermentations of cellulose increased from 1.27 mM to 12.36 mM. Although VFA production from cellulose fermentation increased, starch and glucose fermentations contained higher concentrations than cellulose fermentations, (24% and 66% greater, respectively [$P <$

0.05]). By 18 h glucose and cellulose fermentations VFA concentrations were similar ($P > 0.05$), and starch concentrations were 31% greater than the other CHO fermentations ($P < 0.05$). By 24 h, VFA concentrations in starch fermentations were 25% greater than those in cellulose fermentations ($P < 0.05$) with VFA concentrations from glucose fermentation being intermediate between starch and cellulose ($P > 0.05$ to each). There was no effect of CHO source on the acetate to propionate (A:P) ratio ($P > 0.05$) at any time point.

The nonlinear regression equation, $A-B \cdot \exp^{-kf \cdot \text{time}}$, criteria was met for glucose CH_4 production (Table 2.1). A k_f value for CH_4 concentrations was valued at 0.31 (Table 2.1). Cellulose CH_4 production was unable to converge. Starch CH_4 production was able to converge, but the k_f value of zero was in the confidence interval, suggesting a zero rate.




Figure 2.3: Volatile fatty acid (VFA) production of fermentations of each carbohydrate source (2500 mg/L) in mixed ruminal microorganism fermentation in Study 1

Table 2.1: Fractional rates of CH₄ production among CHO sources

CHO	Variable	Estimate	Approx. STD Error	Approximate 95% Confidence Limits	
Study 1					
Glucose	a	3.31	0.17	2.95	3.68
	b	4.18	0.54	3.03	5.32
	kf	0.319	0.084	0.14	0.5
Study 2					
Starch	a	5.52	0.296	4.92	6.12
	b	10.59	1.33	7.89	13.3
	kf	0.18	0.038	0.11	0.26

Study 2

Glucose pH decreased the most rapidly of the CHO sources, from 6.3 to 5.6 within the first four hours ($P < 0.05$; Figure 2.4). The pH of both cellulose- and starch-containing fermentations did not decrease until 12 h; but by 24 h, glucose fermentation pH increased to 5.8, while cellulose- and starch-containing fermentation pH decreased to 5.94 and 5.76, respectively ($P < 0.05$). At no time did the pH fall below 5.5, suggesting that fibrolytic bacteria were not completely inhibited in these fermentations (Russell and Dombrowski, 1980; Hoover, 1986). In this study there was no effect of N source on pH ($P > 0.05$).

There was a CHO treatment effect and an IT \times CHO treatment interaction in CH₄ production ($P < 0.01$ Figure 2.5). During initial fermentation at 4 h, glucose was fermented the most rapidly and resulted in CH₄ concentration approximately 61% greater than both starch- and cellulose-containing fermentations ($P < 0.05$). At 12h of

fermentation, starch fermentations contained the concentrations of CH₄ that were 43% higher ($P < 0.05$) than glucose- and cellulose-containing fermentations which had similar CH₄ concentrations ($P > 0.05$).

This pattern continued at 24 h and starch fermentation produced approximately 30% more CH₄ than did cellulose or glucose fermentations ($P < 0.05$). There was an effect of N source on CH₄, but this only became significant at 24h ($P < 0.05$; Figure 2.6), at which time NH₃ and NF containing fermentations produced 24% more CH₄ than did fermentations containing amino acids ($P < 0.05$).




Figure 2.4: pH of fermentations for each carbohydrate source (1600 mg/L) in mixed ruminal microorganism fermentation in Study 2




Figure 2.5: Methane production of fermentations of each carbohydrate source (1600 mg/L) in mixed ruminal microorganism fermentation in Study 2




Figure 2.6: Methane production of fermentations of each nitrogen source (900 mg N/L) in mixed ruminal microorganism fermentation in Study 2

Volatile fatty acid concentrations were affected by N source, CHO source, and their subsequent two and three-way interactions with IT (to be consistent) ($P < 0.01$; Figure 2.7 and 2.8). At 4 h, glucose fermentations contained greater proportions of VFA than either cellulose or starch fermentations, (56% and 73% greater, respectively [$P < 0.05$]). By 12 h, both glucose and starch fermentations contained higher VFA concentrations than did cellulose fermentations ($P < 0.05$). At 24 h, glucose fermentations contained marginally higher VFA concentrations than starch fermentations ($P = 0.05$), and significantly greater concentrations than from cellulose fermentations ($P < 0.05$). There were no distinguishable differences between N sources on VFA production at 4 h or 12 h ($P > 0.05$; Figure 2.8). However at 24 h, NH_3 treatments produced on average 40% greater VFA concentrations than all other N treatments ($P < 0.05$).




Figure 2.7: Volatile fatty acid (VFA) production of fermentations of each carbohydrate source (1600 mg/ L) in mixed ruminal microorganism fermentation in Study 2




Figure 2.8: Volatile fatty acid (VFA) Production of fermentations of each nitrogen source (900 mg N/L) in Study 2

There was an IT \times CHO treatment interaction on A:P ratio ($P < 0.01$). At 24 h, cellulose had the highest A:P ratio, followed by starch and glucose ($P < 0.05$). There was no effect of N treatment on A:P ratio. In Study 2, CH₄ production rates from starch fermentation converged to fit the nonlinear rate equation, $A-B \cdot \exp^{-kf \cdot \text{time}}$, with a kf value of 0.18. Neither cellulose nor glucose CH₄ productions converged with the equation (Table 2.1).

Discussion

In both studies, there were effects of CHO source on CH₄ concentrations at different times. As expected, glucose was the most readily available carbohydrate.

In Study 1 and Study 2, glucose fermentations produced the highest concentrations of CH₄ within the first four hours, followed by starch and cellulose fermentations. This result was expected because complex carbohydrates, such as starch and cellulose, take longer to be fermented than do more simple carbohydrates, such as glucose (Beuvink and Spoelstra, 1992; Cone and van Gelder, 1999). Although, glucose initially (< 4 h) had 50% greater CH₄ concentrations than cellulose or starch, in both studies CH₄ concentrations during this early period were minute compared to later time points.

In Study 1, at 24 h cellulose fermentations had greater CH₄ concentrations than starch ($P > 0.05$) or glucose ($P < 0.05$; Figure 2.2). Contrastingly, during Study 2, starch-containing fermentations produced significantly greater concentrations of CH₄ than from cellulose at 24 h (Figure 2.5). One hypothesis for the different outcomes, may have been due to the varying diets in each trial. In the first study, animals were fed a diet lacking concentrate, compared to Study 2 which incorporated 0.9 kg of DDG. Rumen fluid collected from animals in Study 2 initially had lower pH than did rumen fluid collected during Study 1 (6.25 compared to 6.5). The incorporation of DDG may have caused the depression in ruminal pH, selecting for more starch fermenting bacteria in the ruminal microbiome rather than fibrolytic bacteria.

Fermentations lacking amino acids resulted in a higher level of CH₄ production compared with fermentations containing amino acids (Figure 2.6). Previous studies have indicated that the inclusion of protein on *in vitro* fermentations decreased total CH₄ production (Cone and van Gelder, 1999). The increase in methanogenic activity in fermentations without amino acids may be similar to the phenomenon of energy spilling.

Bacteria spill energy in non-growth functions if their growth medium is nitrogen-limited and the energy source is in 'excess' (Russell, 2007). It has been concluded that many bacteria have this reaction in order to dissipate excess ATP when the catabolic rate is faster than the anabolic rate (Russell and Cook, 1995). As a result when energy spilling occurs large amounts of H₂ are dissipated that subsequently can be used by methanogens, which can lead to an increase in CH₄ production.

Ammonia inclusion resulted in the greatest VFA production amongst all N sources (Figure 2.8). This result could also be linked to an energy spilling effect. Without a source of amino acids bacteria were unable to use their N source rapidly enough along with the carbon skeletons derived from CHO fermentation for microbial crude protein synthesis, and as a result diverted their carbon to VFA. Treatments lacking any N source produced VFA concentrations similar to those of treatments that contained amino acids. This may be due to the fact that although there was no additional nitrogen added to the treatments, residual nitrogen was included in the rumen fluid and bacterial protein from cell lysis and death would be available over time. This would be plausible because the ration was not nitrogen limiting and DDG are included as a high quality protein source (Belyea et al., 2004).

As expected both experiments had a CHO × IT interaction on VFA production. However, results differed between experiments. In Study 1 and 2, glucose fermentation produced the greatest concentrations of VFA compared to both starch and cellulose treatments at 4 h ($P < 0.05$). This result was expected for unlike polymers (starch and glucose), glucose can immediately be converted to VFA and fermentation of glucose

does not generally demonstrate a lag time before fermentation (Beuvink and Spoelstra, 1992). In Study 1, at 24 h starch produced marginally higher concentrations of VFA than glucose. Yet in Study 2, glucose fermentation produced greater VFA concentrations as compared to starch.

In Study 2, at 24 h glucose fermentations contained the lowest A:P ratio ($P < 0.05$) as well as the lowest CH_4 concentrations compared with starch or cellulose fermentations ($P < 0.05$). An increase in propionate concentration decreases in hydrogen availability for methanogens, which can lead to a reduction in CH_4 in the rumen (Russell and Rychlik, 2001). Surprisingly, starch and cellulose fermentations produced similar A:P ratios, which is atypical of these very different substrates. Starch fermentations are known to have a lower A:P ratio than cellulose fermentations (Armentano and Young, 1983), so it is unknown why in this experiment starch and cellulose had similar A:P ratios, but it may be related to the amorphous nature of the cellulose used in this study.

Rates were unable to be determined for all of the CHO treatments in both Study 1 and 2. Although a formula converged for starch in Study 1 and cellulose in Study 2; the k_f value of zero was in the confidence interval, indicating a linear relationship for fermentations. However, carbohydrate degradation is known to be an exponential relationship (Russell et al., 1992; Sniffen et al., 1992; Russell, 2009), making it very unlikely that the conversion of amorphous cellulose to CH_4 is a truly linear relationship. Lengthening the fermentation time of cellulose beyond 24 h could exhibit a nonlinear production rate for CH_4 . In addition to more accurately fit fractional rate equations to

CH₄ production, additional measurements would need to be taken over the fermentation period amongst all CHO treatments.

Conclusion

This is one of the first studies to determine the effects of a purified CHO and variety of N sources on *in vitro* ruminal fermentation. Results indicated that both CHO and N sources had a profound impact on CH₄ and VFA production over time.

Collectively, the results gathered in these studies are the first steps necessary to build nonlinear rate equations for the conversion of carbohydrates to CH₄ and one day may be used to improve the predictive accuracy of CH₄ models.

CHAPTER IV

COMPARING METHANE GENERATING PROTEIN SYSTEMS

Introduction

The recent discussion over food sustainability and livestock production comes at a time when the awareness of the effects of global climate change has never been greater (Tedeschi et al., 2015). According to the U.S. EPA (2014); “Methane emissions from enteric (gastrointestinal) fermentation represent 25.0 percent of anthropogenic activities.” Methane, a potent greenhouse gas (GHG) with a 12-year atmospheric lifetime and global warming potential 25 times higher than that of carbon dioxide (EPA, 2016), has significant environmental ramifications (Stephenson, 2009). Mitigating climate change must include greenhouse gasses (GHG) reduction. Yet, over the next 25 years, the predicted exponential human population growth necessitating protein production to increase by 70% (IPCC, 2006). Although cattle production provides the population with an abundant and complete source of protein (National Academy of Sciences, 2005; U.S. Department of Health and Human Services, 2016), some believe because cattle contribute to anthropogenic GHG that they are a non-sustainable food source (Pollan, 2002; Bittman, 2008). This conflicting issue involving environmental vs. human protein needs has been referred to as “wicked problem” (Peterson, 2011). However, one factor that has been remiss from the dialog is that beef cattle have not always been the dominant source of agricultural CH₄ emissions nor is it the only source that should be examined. Therefore, in an effort to determine the efficiency of current

animal protein producing systems both on nutritional and environmental bases life cycle assessments (LCA) have been performed.

In a recent LCA, it was determined that beef production was the least efficient protein source and produced five times more GHG than the average of the other livestock (egg, swine, dairy, and poultry) protein producing categories (Eshel et al., 2014). In another LCA it was determined that the increase in dairy consumption suggested in 2010 USDA dietary guideline recommendations, was not a sustainable decision because of dairy cows associated GHG emissions (Heller and Keoleian, 2015). Subsequently, cattle production have been thought to be a non-sustainable food source (Pollan, 2002; Bittman, 2008).

Although these assessments addressed cattle substantial production of agricultural CH₄, over the course of United States' history, there has been a multitude of agricultural food sources, other than cattle, that have substantially contributed to CH₄ emissions that were not taken into account. For example, Bison, elk and deer, like cattle, are ruminants and it has been theorized that prior to European settlement (pre-settlement) produced sizable amounts of CH₄ (Subak, 1994; Hristov, 2012). In addition to wild ruminants the grain rice, one of the most heavily consumed food sources in the world, is methanogenic, constituting one of the largest sources of agricultural CH₄ emissions (van Groenigen et al., 2013; IPCC, 2007). In order to understand and address the impending issues of food sustainability, the efficiencies of pre-settlement and current methanogenic agricultural food sources should be compared on an CH₄ output and human-edible protein basis. To our knowledge, such comparisons have not been

performed. Thus, the objective of this study was to compare current cattle producing food sources (beef and milk) to pre-settlement populations of bison, elk and deer, and current rice production on a CH₄ emitted to human-edible protein output basis.

Materials and methods

Literature sources, both historic and present day, were used to determine historic ruminant CH₄ emission and human-edible protein production. The following equation were used to calculate CH₄ emissions and human-edible protein for wild ruminants:

$$\begin{aligned} \text{Human-edible Protein Produced} &= \text{Animal Live Weight (kg)} \times \text{percent of hot carcass} \times \text{percent of human-edible meat per hot carcass} \times \text{protein content (g/kg of product)} \times \text{population size} \times \text{sustainable harvest rate} \end{aligned} \quad [1]$$

$$\begin{aligned} \text{Annual Herd CH}_4 \text{ Production} &= \text{CH}_4 \text{ yield per unit of DM ((g of CH}_4\text{)/(kg of DMI))} \times \text{DMI ((kg per head)/day)} \times 365 \text{ days} \times \text{population size} \end{aligned} \quad [2]$$

$$\begin{aligned} \text{Methane Emitted to Human-edible Protein Produced} &= \text{Herd CH}_4 \text{ Emissions} / \text{Annual Herd Human-edible Protein Produced} \end{aligned} \quad [3]$$

Assumptions that had to be made for this species included intake, weight, and population size. It was postulated that pre-settlement, prior to the settlement of Europeans, bison populations could have exceeded 75 million, although many sources

have suggested that the maximum carrying capacity for bison could not exceed 30 million head, this estimate was used for this analysis (Dary, 1989; Isenberg, 2001). Bison intake was based on Kelliher and Clark (2010), calculated dry matter intake (DMI) based on IPCC Tier 2 methodology, estimated that bison consumed 2.2% of body weight as DM. Previous assessments of bison mass for females and males ranged from 318-554 and 544 to 977 kg, respectively (Meagher, 1986). Assuming a 1:1 male to female sex ratio the bison weight on average 600 kg.

The annual sustainable harvest rate for bison populations was determined to be 15% of the herd, or 4.5 million animals (Frost, 2015). This value represented the largest population that could be harvested without decreasing population growth the subsequent year. Dressing percentage used for bison was 57% (NBA, 2016). Approximately, 70% of the hot carcass was boneless lean meat, which equated to 227 kg of human-edible meat (Hawley, 1986; Koch et al., 1995b). Average protein yield per carcass, assuming grass fed, averaged 202 g per kg of meat (USDA, 2015b). Total herd protein was calculated using Equation [1].

The amount of CH₄ produced per kg of DMI per day for this model was based on Hristov (2012) calculations, who determined that bison produced 21 CH₄ per kg of DMI consumed. Hristov (2012) calculated this value off of the average of two previous estimates: 1) Galbraith et al. (1998) empirically based CH₄ losses of 20 g of CH₄ per kg of DMI per day, and 2) Kelliher and Clark (2010) who calculated that animals produced 21.4 g of CH₄ per kg of DMI per day. Dry matter intake was based off of Kelliher and

Clark (2010), who concluded using IPCC tier 2 methodology, that bison consumed 2.2% body weight as DM. Total CH₄ production was calculated using Equation [2].

Elk

Historic elk populations was valued at 10 million in the contiguous U.S. pre-settlement (Rocky Mountain Elk Foundation [RMEF], 2016). Body weight for elk averaged 227 kg for cows and 318 kg bulls (RMEF, 2016). Elk body weight, assuming a 1:1 sex ratio averaged 272 kg (RMEF, 2016). Field dressed weight (without hide, head, or feet) was determined to be 58% (Field, 2003b). The percent of consumable hot carcass was 58%, yielding 92 kg of human-edible meat per animal (Field, 2003b). Elk's sustainable harvest was based on 15% of population per year, or 1,500,000 animals (Frost, 2015). Protein yield per kg of human-edible meat averaged 213 g (USDA, 2015b). Total herd protein was calculated using Equation [1].

Based on Galbraith et al. (1998) and Hristov (2012) calculations, elk produced 16 g of CH₄ per kg of DMI per day. For the purpose of this analysis the amount of DM consumed relative to body weight (kg) was based on the Small Ruminant NRC (2007), valued at 2.3%. Total CH₄ production was calculated using Equation [2]. Methane to human-edible protein ratio was calculated using Equation [3].

Deer

Although a variety of deer species and sub-species currently exist in the United States, historically, there were two dominant species, White Tailed and Mule deer. Pre-settlement populations for White Tailed and Mule deer were approximately 30 million and 13 million, respectively (Peek, 2003). The size of Mule deer ranged from 60 kg to

128 kg (Bauer and Bauer, 1995). White Tailed deer averaged 50 to 136 kg (Fulbright and Ortega-S, 2006). Average weight between species, assuming a 1:1 female to male sex ratio, averaged 58 kg (Smith, 1991; Ferguson, 2005). The field dressed body weight (without skin, feet, or head) and dressing percentage among deer species averaged 34 kg and 59% (Field, 2003a). Approximately, 57% of the hot carcass was consumable, yielding 18 kg of human-edible meat (Field, 2003a). Protein yield per kg of human-edible meat averaged 223 g (USDA ARS, 2015). Sustainable harvest rate's for deer were higher than that of elk or bison, averaged 22.5% or 9.6 million head (Guynn, 1985; Frost, 2015). Total herd protein was calculated using Equation [1].

Methane emissions for both species of deer were based on Galbraith's (1998) and Hristov (2012) studies that calculated 10 g CH₄ were emitter per kg of DMI per day. Average DMI per animal was assumed to be 3.2% of total body weight, based on the NRC (2007). Total CH₄ production was calculated using Equation [2]. Methane to human-edible protein ratio was calculated using Equation [3].

Beef cattle

In 2012, total beef cattle populations in the U.S. totaled 77 million head (USDA National Agricultural Statistical Service [USDA NASS], 2012). This figure included, all beef cows, beef replacement heifers, bulls, beef heifers and steers. From this population a total of 43%, approximately 38 million animals, were harvested (USDA NASS, 2012). Federally inspected live weights and hot carcass weights across all beef cattle harvested for the year 2012 averaged 593 kg and 359 kg, respectively (Bertramsen, 2015). Percent of hot carcass consumable was 74.4%, based on yield 3 grade carcass

with 0.64 cm fat trim, yielding 251 kg of human-edible (Griffin et al., 1992). Total edible protein per kg of human-edible meat was determined to be 214 g (United States Department of Agriculture [USDA], 2015b). Using EPA's estimations for gastrointestinal CH₄ emissions for 2012, total herd emissions were valued at 4,789 CH₄ Gg yr⁻¹, approximately 170 CH₄ g per head per day (U.S. EPA, 2014). Total herd protein was calculated using Equation [1]. Annual herd CH₄ production was then divided by annual herd human-edible protein produced to formulate a CH₄ emitted to human-edible protein ratio.

Dairy cattle

In 2012, there were 13.7 million head of dairy cattle in the United States including cows and replacement heifers (USDA NASS, 2012). In the same year the total herd produced 9.10×10^{10} kg of milk (USDA NASS, 2012). For a kg of milk, there were 36.6 g of protein (USDA Nutrition, 2015). Methane production for dairy cattle was based on the EPA Inventory of U.S. Greenhouse Gas Emission and Sinks: 1990-2012, which utilized Tier 2 IPCC methodology and CH₄ values generated by the COWPOLL model (Kebreab et al., 2008) and other models. Total herd CH₄ for the year 2012 was estimated to be 1,668 Gg yr⁻¹ or averaging 388 CH₄ g d⁻¹ per head (U.S. EPA, 2014). It was assumed that 100% of milk produced was consumable. Annual herd CH₄ production was then divided by total human-edible protein produced to formulate the CH₄ to human-edible protein ratio.

Rice production

We chose to compare rice production to ruminant production for rice is one of the world's most common crops and it is known to produce CH₄. Predominate rice growing regions in the United States include Arkansas, California, and Louisiana (USDA, 2014). Long grain is grown almost exclusively in the south and accounts for more than 70 percent of U.S. production (USDA, 2012). Medium grain, grown both in California and the South, account for more than one-fourth of total U.S. production and forms most of California's rice crop. Arkansas accounts for most of the southern medium-grain production. Short grain accounts for 1-2 percent of total U.S. rice production and is grown almost exclusively in California. Among U.S. rice species harvested protein content averaged 68 g per kg of product (USDA, 2015b). Methane emissions from rice production was based on EPA Inventory of U.S. Greenhouse Gas Emission and Sinks: 1990-2012, valued at 351 CH₄ Gg yr⁻¹ (U.S. EPA, 2014). In 2012, 2.6 million acres were harvested yielding 10.1×10³ kg (USDA NASS, 2012). Approximately, 68% of the crop was consumable and consumable and retail ready (Wilson, 2015). Protein yield was determined by amount of kg of rice produced × percent consumable × g protein produced per kg of rice. Methane to human-edible protein ratio for rice was calculated by CH₄ emissions for the year 2012 divided by annual human-edible protein produced.

Results and discussion

Before European settlement the United States, plains and forests were dominated by wild ruminants including bison (*Bison Bison*), elk (*Cervus canadensis*), and deer (*Odocoileus virginianus* and *Odocoileus hemionus*). Like cattle, wild ruminants produce CH₄ as an end product of microbial fermentation. It has been speculated that during pre-settlement times, 30 to 75 million bison, along with millions of deer and elk, inhabited the U.S. (McHugh, 1972; Dary, 1989; Isenberg, 2001) and at this time were responsible for producing substantial amounts of CH₄ (Subak, 1994).

Determining CH₄ emissions from pre-settlement animals is a difficult task because neither daily animal emissions nor populations could be measured at this time. As such, many assumptions had to be made in this comparison. Factors determining emissions include forage type, DMI, and the size of the animal, which can be difficult to measure. Ruminants are classified as grazers and browsers, which makes it problematic to decipher DMI and type of forage being consumed. Crutzen et al. (1986) theorized that because wild ruminants lived entirely on roughage and herbs at maintenance levels, ruminants would produce on average 9 percent of gross energy lost in the form of CH₄ (Y_m). However, this value is difficult to confirm. Therefore, in order to empirically record wild ruminant emissions, Galbraith et al. (1998) built respiratory chambers for yearling female bison (195.7 kg ± 7.52), elk (151.3 kg ± 4.1), and deer (34.4 kg ± 1.45) (Galbraith et al., 1998a). Using the chambers, Y_m values for bison, elk and deer measured 6.6, 5.2, and 3 percent, respectively. Interestingly, these values were significantly lower than Crutzen's (1986) previous assessments. Animals utilized in

Galbraith's (1998) trials were fed a diet of alfalfa pellets and no long stem forages. Feeding pellets rather than long stem hay, reduces rumen retention time, which can decrease total CH₄ production (Galbraith et al., 1998b). This may be why Galbraith's Y_m values were lower than Cruzen's (1986) estimates. To date, these are the only empirically collected CH₄ data for wild ruminants and were used as the source for elk and deer emissions and one of two sources for bison emissions in this model.

Bison

According to the National Bison Association (2015), average dressing percentages of bison was valued at 57 percent. Dressing percentages in the literature evaluated the dressed carcasses higher than the National Bison Association, closer to 60 percent (Hawley, 1986; Koch et al., 1995a; López-Campos et al., 2013). However, these estimates were biased on farmed raised bison consuming a diet that consisted of both forage and grain. For beef cattle fed 100 percent forages diet, animals yielded lower dressing percentages than those that were finished on grain (Leheska et al., 2008). Therefore, it can be reasonably assumed that free range bison dressing percentages would be lower than grain fed animals. As such, the dressing percentage of 57, based on the estimate provided by National Bison Association, was used in this analysis. Using the USDA Nutrition Database (2015), each animal produced 4.8×10^4 g of protein (Table 3.2). Overall, this study determined that bison weighing 600 kg, consuming 2.2 percent DMI of BW, producing 21 CH₄ g per kg DMI produced 277 CH₄ g per head per day (Table 3.1). For 30 million bison, producing 277 CH₄ g per head per day a total of 3,035 Gg CH₄ yr⁻¹ was emitted (Table 3.2). On a CH₄ to protein biases,

bison emitted 13.93 kg of CH₄ for every kg of human-edible protein produced (Table 3.2).

Table 3.1: Historic Ruminant Gastrointestinal CH₄ Emissions

Items	Bison	Elk	Deer
Body Weight, kg	600 ¹	272 ²	57.6 ³
DMI of BW, %	2.2 ⁴	2.3 ⁵	3.1 ⁵
DMI, kg/head/d ⁶	13.2	6.26	1.79
CH ₄ Emissions, g/kg of DM/d	21	16	10
CH ₄ Emissions, g/head/d	277.2	100.1	17.9

1 Meagher (1986)

2 Peek (2003)

3 Fulbright and Ortega (2006)

4 Kelliher and Clark (2010)

5 NRC (2007)

6 Galbraith et al. (1998)

Elk

For historic elk populations (10 million), assuming a skinned carcass percentage of 58 percent (Fields et al., 2003b), elk produced 19.5 kg of protein per animal or annually 2.9×10^7 kg of protein for the herd (Table 3.2). Elk weighing 272 kg, consuming 2.3 percent of body weight as DM, emitting 16 g CH₄ per kg of DMI, produced 100.1 g CH₄ d⁻¹ (Table 3.1). For a population of 10 million elk, 365 Gg of CH₄ yr⁻¹ was emitted (Table 3.2). For every kilogram of human-edible protein produced, 12.50 kg of CH₄ would be emitted, which was only slightly lower than bison's ratio (Table 3.2).

Deer

Although there were two species of deer used for this analysis, CH₄ emissions for both populations were based on Galbraith's et al. (1998) study using White Tailed deer

at measured 10 g of CH₄ per kg of DMI. Historic deer populations, assuming a field dressing percentage of 59 percent, produced 4.04 kg of protein per animal. The herd annually produced 4.2×10⁷ kg of protein (Table 3.2). For deer weighing 58 kg, consuming 3.1 percent of their BW as DM, producing 10 g of CH₄ per kg of DMI, deer emitted 17.86 CH₄ d⁻¹ (Table 3.1). For a population of 43 million deer, emission rate was valued at 280 Gg of CH₄ yr⁻¹ (Table 3.2). Thereby, 6.66 kg of CH₄ was emitted for every kg of human-edible protein produced (Table 3.2).

Beef Cattle

Beef cattle CH₄ emissions were based on the EPA Inventory of U.S Greenhouse Gas Emissions and Sinks for 2012 (U.S. EPA, 2014). The U.S. EPA assessment used IPCC's Tier 2 methodology incorporating a total of 177 input variables for cattle emissions. Based on this assessment, gastrointestinal fermentation from beef cattle for the year 2012 valued 4,789 Gg CH₄ yr⁻¹, averaging 170 g d⁻¹ per head (U.S. EPA, 2014). The EPA assessment included all beef cows, beef replacement heifers, bulls, and all feedlot animals. The average hot carcass weight of all federal inspected cattle for 2012 averaged 359 kg (Bertramsen, 2015). Federally inspected cattle encompassed 98.5 percent of cattle slaughtered in the United States (Bertramsen, 2015). It was assumed that the proportion of consumable meat from the hot carcass was 74.4 percent, based on previous studies by Griffin et al. (1992) that broke down sub-primal yields of beef carcasses (Griffin et al., 1992). Using USDA's Nutritional Database the raw beef

Table 3.2: Pre-settlement wild ruminant and current beef cattle gastrointestinal emissions of CH₄ and production of human-edible protein

Protein Source	Bison	Elk	Deer	Beef Cattle
Herd Size ¹ , in millions	30	10	43	77
Daily Animal CH ₄ ^{1,2} , g/d/hd	277.2	100.1	17.86	170.4
Daily Herd CH ₄ , kg/d	8,316,000	1,000,960	767,808	13,113,100
Annual Herd CH ₄ , Gg/yr	3,035	365	280	4,790
Sustainable Harvest ^{3,4}	0.15	0.15	0.23	0.44
Head Harvested ⁵	4,500,000	1,500,000	9,675,000	33,880,000
Live Weights ^{1,5} , kg	600	272	58	593
Hot Carcass Weight ⁵ , kg	342	190	34.2	359
Percent of Hot Carcass Consumable	0.70	0.58	0.57	0.74
Human Edible Product, kg	239	91.5	18.1	267
Human Edible Protein ⁷ , g/kg	202	213	223	214
Protein Yield per Head, kg	48.4	19.5	4.35	57.1
Herd Protein Yield, Gg	218	29.2	42.1	2000
CH ₄ per Human-edible Meat, kg/kg	2.82	0.40	0.33	0.23
CH ₄ per Human-edible Protein, kg/kg	13.9	12.5	6.66	2.47

¹ See Table 3.1

² U.S. EPA (2014)

³ Frost (2015)

⁴ Guynn (1985)

⁵ Bertramsen (2015)

⁶ USDA (2015a)

composite of trimmed retail cuts, separable lean and fat, trimmed to 1/8 in. (0.32 cm) fat, all grades, averaging 214 g of protein per kg of beef, approximately 57.1 kg protein per harvested beef animal (Table 3.2). This equated to 2.47 kg CH₄ emitted for every kg of human-edible protein produced (Table 3.2, Table 3.5).

Historic Wild Ruminant Herd Emissions

We showed in this model that wild ruminates produced 77 percent of CH₄ emissions as compared to beef cattle emissions. However, this calculation was derived from Galbraith's empirically based data that could have underestimated bison emissions.

Using Crutzen's (1986) Y_m value of 9 percent, and Galbraith's et al. (1998) calculations of bison daily energy intake (kj), bison would produce 370 g of kg $CH_4 d^{-1}$. Using this value, wild ruminants produced approximately 4,047 Gg of $CH_4 yr^{-1}$, equating to 98% of beef cattle gastrointestinal CH_4 emissions for the year 2012.

Dairy

Heller and Keoleian (2015) determined that the increase in dairy consumption recommended by 2010 USDA dietary guidelines would contribute significantly to increased GHG emissions and would not be an ideal protein alternative to meat. Furthermore, this statement was repeated in the USDA Scientific Report of the 2015 Nutritional Guidelines Recommendations (USDA, 2015a). However, of all the food sources evaluated in our model, milk produced the fewest emissions per kg of human edible protein produced. For 13.7 million dairy cattle emitting 1,668 Gg $CH_4 yr^{-1}$, producing 91,000 Gg of milk yr^{-1} , with a protein content of 36.6 g per kg of milk, the CH_4 to human-edible protein ratio was 0.50 (Table 3.3). Overall, dairy produced 20.2% of beefs CH_4 emissions per kilogram of protein produced (Table 3.5). Our model shows the U.S. dairy production is extremely efficient at producing a food source.

Rice

We compared ruminants to rice production, not to condemn one food producing system over another, but merely to contrast it with CH_4 production for beef and milk protein productions. Rice, one of the most widely consumed human feedstuffs in the world, is methanogenic, constituting a predominant source of anthropogenic CH_4 (Agnihotri et al., 1999; Yan et al., 2009; Neue, 2014). It is expected that total rice

consumption will continue to increase concurrently with the rate of population growth (Mohanty, 2012).

Table 3.3: Dairy cattle gastrointestinal CH₄ emissions and human consumable protein ratio

Herd Size ¹ , millions	13.7
Daily Head CH ₄ , g/d/hd	334
Daily Herd CH ₄ , kg/d	4,600,000
Annual Herd CH ₄ ² , Gg/yr	1,668
Annual Herd Milk Production ¹ , Gg/yr	91,000
Annual Milk Production per Hd, kg/yr	9,873.6
Percent Consumable, %	1.0
Consumable Product, kg	91,000
Consumable Protein ³ , g/kg	36.6
Protein Yield per Head, kg	243
Herd Protein Yield, Gg	3,330
CH ₄ per Milk, kg/kg	0.02
CH ₄ per Human-edible Protein, kg/kg	0.50

¹ USDA NASS (2012)

² U.S. EPA (2014)

³ USDA (2015a)

Therefore, it can be expected that rice production and its associated CH₄ emissions will increase over the course of this century. Currently, worldwide rice production accounts for 31-112 Tg of CH₄ yr⁻¹, which is similar to total ruminant production that accounts for 76 to 92 Tg CH₄ yr⁻¹ (IPCC, 2007). Most of the world's rice, and all rice in the United States, is grown on flooded fields (Borris, 2006). The flooding of field's induces the aerobic decomposition of organic material that depletes soil oxygen creating anaerobic growing conditions. Methanogens present in the soil produce CH₄ through the anaerobic decomposition of soil organic matter (Holzapfel-Pschorn et al., 1985).

In 2012, 351 Gg of CH₄ was emitted from 2.68 ×10⁵ acres of rice patties across the U.S. (Table 3.4; U.S. EPA, 2012; USDA NASS, 2016). When rice was compared to beef production (2.47 kg CH₄ per kg of human- edible protein) rice produced 66% fewer CH₄ emissions on a human-edible protein basis (Table 3.4, Table 3.5). When rice was compared to dairy, the dairy CH₄ ratio (0.50 kg CH₄ per kg of human-edible protein) lower than rice’s ratio (Table 3.5).

Table 3.4: Rice CH₄ emissions and human-edible protein

Acre harvested ¹ , acre	268,000
Yield per Acre ¹ , kg/yr	3,400
Total Produced, Gg/yr	9,110
CH ₄ per Hectare, g/d/ha	359
Daily CH ₄ , kg/d	962,000
Annual Total CH ₄ ² , Gg/yr	351
Percent of Crop Consumable ³ , %	0.68
Human Edible Product, kg	6,195
Human Edible Protein ⁴ , g/kg	68.0
Protein Yield per Acre, kg	157
Harvest Protein Yield, kg	421
CH ₄ Emissions per Human-edible Rice, kg/kg	0.06
CH ₄ Emissions per Human-edible Protein, kg/kg	0.83

¹USDA NASS (2012)

²U.S. EPA (2014)

³Wilson (2015)

⁴USDA (2015a)

Table 3.5: CH₄ emissions to human-edible protein yield

Protein Source	CH ₄ to Protein (kg/kg)	Relative to Beef, %
Bison	13.93	564
Elk	12.50	506
Deer	6.66	296
Beef	2.47	100
Rice	0.83	33.6
Milk	0.50	20.2

Conclusion

Currently, the U.S. has a population of 320 million people that is steadily increasing. In order to meet the protein demands of the country's growing population, both in terms of quantity and quality, food production must be produced as efficiently as possible. Additionally, in order to minimize the impact on the environment we need to generate food that minimizes negative environmental feedback. Thus, it is imperative that we evaluate and compare all current methanogenic food producing systems along with historic methanogenic food sources in order to make informed agricultural decisions that are advantageous for both human and environmental health. This model was the first of its kind to compare pre-settlement wild ruminants, and current rice, beef, and milk production on a CH₄ to human-edible protein basis. The implications of this model demonstrate that, when compared to other methanogenic producing food sources, beef may not be as an inefficient food source speculated previously. Ultimately, this model provides insight on the efficiency of methanogenic food sources and one day may aid in the development of environmental and governmental food sustainability guidelines.

CHAPTER V

SUMMARY

Approximately 2-8% of energy consumed by cattle is converted into CH₄ (Yan et al., 2009) a significant energy loss from the animal. Many strategies have been evaluated to reduce CH₄ including modifying dietary composition, inclusion of ionophores, organic acids, and plant compounds, defaunation and vaccinations. Abatement strategies directly or indirectly target ruminal methanogen populations, resulting in varying degrees of efficacy. Methane production not only comes at a cost to the animal, it also contributes to anthropogenic GHG. Specifically, according to the U.S. EPA (2014) gastrointestinal fermentation accounts for 38% of total agriculturally related CH₄ emissions in the U.S. Furthermore, CH₄ has a 12-year atmospheric lifetime and global warming potential 25 times greater than that of CO₂ (U.S. EPA, 2016), thus strategies to mitigate gastrointestinal CH₄ production are of critical interest to environmentalists, as well as to cattle producers.

In ruminants carbohydrate (CHO) catabolism and nitrogen (N) utilization have a tremendous impact on ruminal methanogenesis. However, the impact of purified carbohydrates in the presence of a variety of N sources on rates of CH₄ and VFA production remains unknown. In order to determine these rates for use in predictive models of the ruminal fermentation, we formulated a fractional rate equation to fit the rate of CH₄ production and measured the concentration of CH₄ and VFA and using purified CHO with a variety of N sources in two *in vitro* mixed ruminal microorganism

fermentation studies. Results indicated that both CHO and N sources had a profound impact on CH₄ and VFA production over time. Collectively, the results gathered in these studies are the first steps necessary to build nonlinear rate equations for the conversion of carbohydrates to CH₄ and one day may be used to improve the predictive accuracy of CH₄ models.

The objective of the final study was to compare the efficiency of beef and milk production to pre-settlement wild ruminants and rice production on a kilogram of CH₄ emitted to kilogram human-edible protein production basis. Currently, the U.S. has a population of 320 million people that is steadily increasing. In order to meet the protein demands of the country's growing population, both in terms of quantity and quality, food production must be produced as efficiently as possible. Additionally, in order to minimize the impact on the environment we need to generate food that minimizes negative environmental feedback. Thus, it is imperative that we evaluate and compare all current methanogenic food producing systems along with historic methanogenic food sources in order to make informed agricultural decisions that are advantageous for both human and environmental health. Overall, this model was the first of its kind to compare pre-settlement wild ruminants, and current rice, beef, and milk production on a CH₄ to human-edible protein basis. Bison had the highest ratio of 13.93 kg CH₄: Protein, followed by elk (12.50) deer (6.66) and beef (2.47). Wild ruminants emitted 296 to 564 percent more CH₄ per kilogram of human-edible protein produced than current beef cattle production systems. Rice yielded the second lowest CH₄ to human-edible protein ratio (0.83), followed by dairy cattle milk production (0.50). Ultimately,

this model provides insight on the efficiency of methanogenic food sources and one day may aid in the development of environmental and governmental food sustainability guidelines.

In conclusion, cattle production, ruminant nutrition, GHG production, and food security and sustainability are complex and inextricably linked. With continual improvements in technology and beef cattle management, the U.S. able to produce more beef without increasing herd gastrointestinal CH₄ emissions (Capper, 2011; U.S. EPA, 2014). Although beef require a great amount of recourses, they still play an intricate role in the United States food system. For it is a balance between the myriad of protein production systems that will ensure U.S livestock's economic, social and environmentally sustainability.

REFERENCES

- Agnihotri, S., K. Kulshreshtha, and S. N. Singh. 1999. Mitigation strategy to contain methane emission from rice-fields. *Environmental Monitoring and Assessment*. 58: 95-104.
- Armentano, L. E., and J. W. Young. 1983. Production and metabolism of volatile fatty acids, glucose and CO₂ in steers and the effects of monensin on volatile fatty acid kinetics. *The Journal of Nutrition*. 113: 1265-1277.
- Bauer, E. A., and P. Bauer. 1995. *Mule deer : behavior, ecology, conservation*. Voyageur Press. Stillwater, MN.
- Beauchemin, K.A. and McGinn S.M. 2006. Methane emissions from beef cattle: effects of fumaric acid, essential oil, and canola oil. *Journal of Animal Science*. 84: 1489–1496.
- Beauchemin, K.A., H. H. Janzen, S. M. Little, T. A. McAllister, and S. M. McGinn. 2010. Life cycle assessment of greenhouse gas emissions from beef production in western Canada: A case study. *Agriculture Systems*. 103:371–379.
- Belyea, R. L., K. D. Rausch, and M. E. Tumbleson. 2004. Composition of corn and distillers dried grains with solubles from dry grind ethanol processing. *Bioresource Technology*. 94: 293-298.
- Bertramsen, S. 2015. NASS Cattle Slaughter Specialist. Personal Communication. April 6, 2015.

- Beuvink, J. M. W., and S. F. Spoelstra. 1992. Interactions between substrate, fermentation end-products, buffering systems and gas production upon fermentation of different carbohydrates by mixed rumen microorganisms in vitro. *Applied Microbiology and Biotechnology*. 37: 505-509.
- Bittman, M. 2008. Rethinking the Meat-Guzzler. Accessed March 28, 2016. <http://www.nytimes.com/2008/01/27/weekinreview/27bittman.html>
- Blaxter, K. L. 1967. *The Energy Metabolism of Ruminants*. Hutchinson and Co., London, UK.
- Blood, D. C., and O. M. Radostits. 1989. *Veterinary Medicine*. 7th ed. Balliere and Tindall, London, England.
- Boadi, D. A., K. M. Wittenberg, and A. D. Kennedy. 2002. Validation of the sulphur hexafluoride (SF₆) tracer gas technique for measurement of methane and carbon dioxide production by cattle. *Canadian Journal of Animal Science*. 82: 125-131.
- Boadi, D. A., C. Benchaar, J. Chiquette, and D. Massé. 2004. Mitigation strategies to reduce enteric methane emissions from dairy cows: update review. *Canadian Journal of Animal Science*. 84: 319–335.
- Borris, H. 2006. Rice Profile. Agricultural Issue Center. University of California, Davis, CA. Accessed March 5, 2016. <http://www.agmrc.org/commodities-products/grains-oilseeds/rice-profile/>.
- Bowen, R. 2009. *Digestive Physiology of Ruminants*. Accessed March, 28 2016.

<http://arbl.cvmbs.colostate.edu/hbooks/pathphys/digestion/herbivores/rumination.html>

- Broucek, J. 2014. Production of methane emissions from ruminant husbandry: a review. *Journal Environmental Protection*. 5: 1482-1493.
- Callaway, T. R., A. M. Carneiro De Melo, and J. B. Russell. 1997. The effect of nisin and monensin on ruminal fermentations in vitro. *Current Microbiology*. 35: 90-96.
- Capper, J. L. 2011. The environmental impact of beef production in the United States: 1977 compared with 2007. *Journal of Animal Science*. 89: 4249-4261.
- Capper, J. L. 2012. Is the grass always greener? Comparing the environmental impact of conventional, natural and grass-fed beef production systems. *Animals*. 2.2: 127-143.
- Chen, G. J., and J. B. Russell. 1991. Effect of monensin and a protonophore on protein degradation, peptide accumulation, and deamination by mixed ruminal microorganisms in vitro. *Journal of Animal Science*. 69: 2196-2203.
- Cone, J. W., and A. H. van Gelder. 1999. Influence of protein fermentation on gas production profiles. *Animal Feed Science and Technology*. 76: 251-264.
- Cotta, M. A., and J. B. Russell. 1982. Effect of peptides and amino acids on efficiency of rumen bacterial protein synthesis in continuous culture. *Journal of Dairy Science*. 65: 226-234.
- Council for Agricultural Science and Technology (CAST). 1999. *Animal Agriculture and Global Food Supply*. Task Force Report. No. 135.

CAST, Ames, Iowa. Accessed April 15, 2014. <http://www.cast-science.org/>.

Crosson, P., L. Shalloo, D. O'Brien, G. J. Lanigan, P. A. Foley, T. M. Boland, and D. A. Kenny. 2011. A review of whole farm systems models of greenhouse gas emissions from beef and dairy cattle production systems. *Animal Feed Science and Technology*. 166–167: 29–4.

Crutzen, P. J., I. Aselmann, and W. Seiler. 1986. Methane production by domestic animals, wild ruminants, other herbivorous fauna, and humans. *Tellus*. 388: 271-284.

Dary, D. 1989. *The buffalo book : the full saga of the American animal*. Swallow Press/Ohio University Press, Chicaco, IL.

Demeyer, D. I. 1975. Methanogenesis: an integrated part of carbohydrate fermentation and its control. In: M. a. A. C. Wamer (ed.) *Digestion and metabolism in the ruminant*. The University of New England Publishing Unit, Armidale, Australia. p 366-382.

Ding, S.-Y., Q. Xu, M. Crowley, Y. Zeng, M. Nimlos, R. Lamed, E. A. Bayer, and M. E. Himmel. 2008. A biophysical perspective on the cellulosome: new opportunities for biomass conversion. *Current Opinion in Biotechnology*. 19: 218-227.

Dohme, F., A. Machmuller, A. Wasserfallen, and M. Kreuzer. 2001. Ruminant methanogenesis as influenced by individual fatty acids supplemented to complete ruminant diets. *Letters to Applied Microbiology*. 32: 47-51.

- Dong H., Mangino J., McAllister T. A., Hatfield J. L., Johnson D. E., Lassey K. R.,
Aparecida de Lima M. & Romanovskaya A. 2006. Emissions from livestock and
manure management, Chapter 10. Guidelines for National Greenhouse Gas
Inventories. 4: 10.1–10.87.
- Ellis, J. L., Kebreab, E., Odongo, N. E., McBride, B. W., Okine, E. K., & France, J.
2007. Prediction of methane production from dairy and beef cattle. *Journal of
Dairy Science*. 90: 3456-3466.
- Eshel, G., A. Shepon, T. Makov, and R. Milo. 2014. Land, irrigation water, greenhouse
gas, and reactive nitrogen burdens of meat, eggs, and dairy production in the
United States. *Proceedings of the National Academy of Sciences*. 111: 11996-
12001.
- FAO. 2011. *The State of Food Insecurity in the World 2011*. Accessed April 15, 2016.
<http://www.fao.org/docrep/014/i2330e/i2330e00.htm>.
- Ferguson, K. I. 2006. Mule deer (*Odocoileus hemionus*). *Natural Resources
Conservation Service, Washington, D.C. Wildlife Habitat Council, Silver Spring
Maryland, MD*. Accessed March 15, 2016.
<http://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=18506.wb>
a
- Ffoulkes, D., and R. A. Leng. 1988. Dynamics of protozoa in the rumen of cattle.
The British Journal of Nutrition. 59: 429-436.
- Field, R. A., F.C. Smith, W.G. Hepworth, and W.J. Means. 2003a. *The Mule Deer
Carcass*. University of Wyoming Agricultural Experiment Station. B-589R.

- Field, R. A., W.G. Hepworth, W.J. Means. 2003b. The Elk Carcass. University of Wyoming Agricultural Experiment Station. B-594R.
- Finlay, B.J., Esteban, G., Clarke, K.J., Williams, A.G., Embley, T.M. and Hirt, R.P. 1994. Some rumen ciliates have endosymbiotic methanogens. *FEMS Microbiology Letters*. 117: 157–162.
- Frost, G. 2015. Wyoming Parks and Wildlife Biologist. Personal Communication. August 20, 2015.
- Fulbright, T. E., and J. A. Ortega-S. 2006. White-tailed deer habitat : ecology and management on rangelands. Texas A&M University Press, College Station, TX.
- Galbraith, J. K., G. W. Mathison, R. J. Hudson, T. A. McAllister, and K.-J. Cheng 1998. Intake, digestibility, methane and heat production in bison, wapiti and white-tailed deer. *Canadian Journal of Animal Science*. 78: 681–691.
- Garcia-Lopez, P. M., L. Kung, Jr., and J. M. Odom. 1996. In vitro inhibition of microbial methane production by 9,10-anthraquinone. *Journal of Animal Science* 74: 2276-2284.
- Garrett, E. F., M. N. Pereira, K. V. Nordlund, L. E. Armentano, W. J. Goodger, and G. R. Oetzel. 1999. Diagnostic methods for the detection of subacute ruminal acidosis in dairy cows. *Journal of Dairy Science* 82: 1170-1178.
- Gerber, P. J., B. Henderson, and H. P. S. Makkar. 2013. Mitigation of greenhouse gas emissions in livestock production : a review of technical options for non-CO₂ emissions. Rome: Food and Agriculture Organization of the United Nations.

- Grant, R. J., and D. R. Mertens. 1992. Influence of buffer pH and raw corn starch addition on in vitro fiber digestion kinetics. *Journal of Dairy Science*. 75: 2762-2768.
- Greve, L. C., J. M. Labavitch, and R. E. Hungate. 1984. alpha-L-arabinofuranosidase from *Ruminococcus albus* 8: purification and possible role in hydrolysis of alfalfa cell wall. *Applied and Environmental Microbiology*. 47: 1135-1140.
- Griffin, D. B., J. W. Savell, J. B. Morgan, R. P. Garrett, and H. R. Cross. 1992. Estimates of subprimal yields from beef carcasses as affected by USDA grades, subcutaneous fat trim level, and carcass sex class and type. *Journal of Animal Science*. 70: 2411-2430.
- Groot, J. C. J., J. W. Cone, B. A. Williams, F. M. A. Debersaques, and E. A. Lantinga. 1996. Multiphasic analysis of gas production kinetics for in vitro fermentation of ruminant feeds. *Animal Feed Science and Technology*. 64: 77-89.
- Guynn, D. C., Jr. 1985. Deer harvest-what percent should be harvested In: White, L. D.; Guynn, D. E.; Troxel, T. R., eds. *Proceedings of the International Ranchers' Roundup*. Texas Agricultural Extension Service, Laredo, TX. 29: 438-443.
- Haisan, J. Sun, L. L. Guan, K. A. Beauchemin, A. Iwaasa, S. Duval, D. R. Barreda, and M. Oba. 2014. The effects of feeding 3-nitrooxypropanol on methane emissions and productivity of Holstein cows in mid lactation. *Journal of Dairy Science*. 97: 3110-3119.
- Hammond, K. J., S. Muetzel, G. C. Waghorn, C. S. Pinares-Patino, J. L. Burke, and S. O. Hoskin. 2009. Exploring variation in methane emissions from sheep and cattle

- fed pasture determined by either SF₆ marker dilution or direct calorimetry. Proceedings of the New Zealand Society of Animal Production. 69:174–178.
- Hawley, A. W. L. 1986. Carcass characteristics of bison (*Bison bison*) steers. Canadian Journal of Animal Science. 66.1: 293-295.
- Hegarty, R. S. 1999. Reducing rumen methane emissions through elimination of rumen protozoa. Australian Journal of Agricultural Research. 50: 1321-1328.
- Hegarty, R.S., Alcock, D., Robinson, D.L., Goopy, J.P. and Vercoe, P.E. 2010. Nutritional and flock management options to reduce methane output and methane per unit product from sheep enterprises. Anim. Prod. Sci. 50: 1026–1033.
- Heller, M. C., and G. A. Keoleian. 2015. Greenhouse gas emission estimates of U.S. dietary choices and food loss. Journal of Industrial Ecology. 19: 391-401.
- Hironaka, R., G.W. Mathison, B. K. Kerrigan, and I. Vlach. 1996. The effect of pelleting of alfalfa hay on methane production and digestibility by steers. Science of the Total Environment. 180: 221–227.
- Holzappel-Pschorn, A., R. Conrad, and W. Seiler. 1985. Production, oxidation and emission of methane in rice paddies. FEMS Microbiology Letters. 31: 343.
- Hoover, W. H. 1986. Chemical factors involved in ruminal fiber digestion. Journal of Dairy Science. 69: 2755-2766.
- Hristov, A. N. 2012. Historic, pre-European settlement, and present-day contribution of wild ruminants to enteric methane emissions in the United States. Journal of Animal Science. 90: 1371-1375.

- Hristov, A.N. and J. P. Jouany. 2005. Factors affecting the efficiency of nitrogen utilization in the rumen. In E. Pfeffer & A.N. Hristov, eds. Nitrogen and Phosphorus Nutrition of Cattle: Reducing the Environmental Impact of Cattle Operations, Wallingford, U.K., CAB International. p. 117–166.
- Hristov, A.N., J. Oh, C. Lee, R. Meinen, F. Montes, T. Ott, J. Firkins, A. Rotz, C. Dell, A. Adesogan, W. Yang, J. Tricarico, E. Kebreab, G. Waghorn, J. Dijkstra, S. Oosting. 2013. Mitigation of greenhouse gas emissions in livestock production – A review of technical options for non-CO₂ emissions. In: Gerber, P. J.; Benjamin Henderson, B., Makkar, H. P. S. (Eds). FAO Animal Production and Health Paper No. 177. FAO, Rome, Italy.
- Hristov A.N., K. A. Johnson, and E. Kebreab. 2014. Livestock methane emissions in the United States. Proceedings of the National Academy of Sciences USA. 111: E1320.
- Hungate, R. E. 1966. The Rumen and its Microbes. Academic Press, Ithica, New York.
- IPCC (Intergovernmental Panel on Climate Change). 2007. Climate Change 2007: Working Group I: The Physical Science Basis, 7.4.1 Methane. Accessed March 28, 2016. https://www.ipcc.ch/publications_and_data/ar4/wg1/en/ch7s7-4-1.html Accessed
- Isenberg, A. 2001. The Destruction of the Bison: An Environmental History, 1750-1920. Cambridge University Press, Cambridge, Mass.

- Isogai, A., and R. H. Atalla. 1991. Amorphous celluloses stable in aqueous media: Regeneration from SO₂-amine solvent systems. *Journal of Polymer Science Part A: Polymer Chemistry*. 29: 113-119.
- Janssen, P. H., and M. Kirs. 2008. Structure of the archaeal community of the rumen. *Applied and Environmental Microbiology*. 74: 3619-3625.
- Johnson, K., M. Huyler, H. Westberg, B. Lamb, and P. Zimmerman. 1994. Measurement of methane emissions from ruminant livestock using a sulfur hexafluoride tracer technique. *Environmental Science & Technology*. 28: 359-362.
- Johnson, K. A., and D. E. Johnson. 1995. Methane emissions from cattle. *Journal of Animal Science*. 73: 2483-2492.
- Jordan, E. Lovett, D. K., Monahan, F. J., Callan, J., Flynn, B., and O'Mara, F. P. 2006. Effect of refined coconut oil or copra meal on methane output and on intake and performance of beef heifers. *Journal of Animal Science*. 84: 162-170.
- Jung, H. G., and M. S. Allen. 1995. Characteristics of plant cell walls affecting intake and digestibility of forages by ruminants. *Journal of Animal Science*. 73: 2774-2790.
- Kebreab, E., D. Pape, T. Wirth, K. A. Johnson, and S. L. Archibeque. 2008. Model for estimating enteric methane emissions from United States dairy and feedlot cattle. *Journal of Animal Science*. 86: 2738-2748.
- Kelliher, F. M., and H. Clark. 2010. Methane emissions from bison—An historic herd estimate for the North American Great Plains. *Agricultural and Forest Meteorology*. 150: 473-477.

- Koch, R. M., H. G. Jung, J. D. Crouse, V. H. Varel, and L. V. Cundiff. 1995a. Growth, digestive capability, carcass, and meat characteristics of *Bison bison*, *Bos taurus*, and *Bos x Bison*. *Journal of Animal Science*. 73: 1271-1281.
- Kujawa, M. A. 1994. Energy partitioning in steers fed cottonseed hulls and beet pulp. Ph.D. Dissertation. Colorado State Univ, Fort Collins.
- Kung, L., K. A. Smith, A. M. Smagala, K. M. Endres, C. A. Bessett, N. K. Ranjit, and J. Yaissle. 2003. Effects of 9,10 anthraquinone on ruminal fermentation, total-tract digestion, and blood metabolite concentrations in sheep. *Journal of Animal Science*. 81: 323-328.
- steLatham, M. J., and M. J. Wolin. 1977. Fermentation of cellulose by *Ruminococcus flavefaciens* in the presence and absence of *Methanobacterium ruminantium*. *Applied and Environmental Microbiology*. 34: 297-301.
- Leheska, J. M., L. D. Thompson, J. C. Howe, E. Hentges, J. Boyce, J. C. Brooks, B. Shriver, L. Hoover, and M. F. Miller. 2008. Effects of conventional and grass-feeding systems on the nutrient composition of beef. *Journal of Animal Science*. 86: 3575-3585.
- Leng, R. A. 2014. Interactions between microbial consortia in biofilms: a paradigm shift in rumen microbial ecology and enteric methane mitigation. *Animal Production Science*. 54: 519-543.
- Leschine, S. B. 1995. Cellulose degradation in anaerobic environments. *Annual Review of Microbiology*. 49: 399-426.

- López-Campos, Ó., J. L. Aalhus, J. Galbraith, I. L. Larsen, M. Juarez, B. Uttaro¹, and W. M. Robertson. 2013. The relation of carcass physiological maturity to meat quality in the Canadian bison grading system. *Canadian Journal of Animal Science*. 94: 55-62.
- Martin, C., D. P. Morgavi, and M. Doreau. 2010. Methane mitigation in ruminants: from microbe to the farm scale. *Animal*. 4: 351-365.
- Martínez-Fernández, G., L. A. Abecia, I. Martín-García, E. Ramos-Morales, S. E. Denman, C. J. Newbold, E. Molina-Alcaide, and D. R. Yáñez-Ruiz. 2015. Response of the rumen archaeal and bacterial populations to anti-methanogenic organosulphur compounds in continuous-culture fermenters. *FEMS Microbiology Ecology*. 91:079-084.
- Matte, A., and C. W. Forsberg. 1992. Purification, characterization, and mode of action of endoxylanases 1 and 2 from *Fibrobacter succinogenes* S85. *Applied and Environmental Microbiology*. 58: 157-168.
- McAllister, T. A., K. J. Cheng, E. K. Okine, and G. W. Mathison. 1996. Dietary, environmental and microbiological aspects of methane production in ruminants. *Canadian Journal of Animal Science*. 76: 231-243.
- McAllister, T. A., K. A. Beauchemin, S. M. McGinn, and X. Hao. 2011. Greenhouse gases in animal agriculture--Finding a balance between food production and emissions. *Animal Feed Science and Technology*. 166-167: 1-6.
- McHugh, T. 1972. *The Time of the Buffalo*. Alfred A. Knopf, New York, NY.

- McGinn, S. M., K. A. Beauchemin, T. Coates, and D. Colombatto. 2004. Methane emissions from beef cattle: Effects of monensin, sunflower oil, enzymes, yeast, and fumaric acid. *Journal of Animal Science*. 82: 3346-3356.
- Meagher, M. 1986. *Bison bison*. American Society of Mammalogists. 266:1-8
- Miller S.M., W.C. Steven Michalak, E.A. Kort, A. E. Andrews, S. C. Biraud, E. J. Dlugokencky, J. Eluszkiewicz, M. L. Fischer, G. Janssens-Maenhout, B. R. Miller, J. B. Miller, S. A. Montzka, T. Nehrkorn, and C. Sweeneyet. 2013. Anthropogenic emissions of methane in the United States. *Proceedings in the National Academy of Sciences USA*. 110: 20018–20022.
- Moe, P. W., and H. F. Tyrrell. 1979. Methane production in dairy cows. *Journal of Dairy Science*. 62: 1583-1586.
- Mohanty, S. 2012. Trends in global rice production. *Rice Today*. 12:44-45.
- Moore, K. J., and H. J. G. Jung. 2001. Lignin and fiber digestion. *Journal of Range Management*. 54: 420-430.
- Morgavi, D. P., C. Martin, J. P. Jouany, and M. J. Ranilla. 2012. Rumen protozoa and methanogenesis: not a simple cause-effect relationship. *The British Journal of Nutrition*. 107: 388-397.
- Murray, R. K. 2012. *Harper's Illustrated Biochemistry*. 29th ed. New York, NY, McGraw-Hill Medical.
- National Academy of Sciences (NAS). 2005. Institute of Medicine. Food and Nutrition Board. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. National

Academy Press. Washington, DC.

- Neue, H. 2014. Methane emission from rice fields: Wetland rice fields may make a major contribution to global warming. *Bioscience*. 43: 466-473.
- Newbold, C. J., and L. M. Rode. 2006. Dietary additives to control methanogenesis in the rumen. *International Congress Series 1293*: 138-147.
- Newbold, C. J., B. Lassalas, and J. P. Jouany. 1995. The importance of methanogens associated with ciliate protozoa in ruminal methane production in vitro. *Letters to Applied Microbiology*. 21: 230-234.
- Newbold, C. J., G. de la Fuente, A. Belanche, E. Ramos-Morales, and N. R. McEwan. 2015. The Role of Ciliate Protozoa in the Rumen. *Frontiers in Microbiology*. 6: 1-14.
- NRC. 2007. *Nutrient Requirements of Small Ruminants : Sheep, Goats, Cervids, and New World camelids*. National Academies Press, Washington, D.C.
- Ørskov, E. R. 1986. Starch digestion and utilization in ruminants. *Journal of Animal Science*. 63: 1624-1633.
- Peek, J. M. 2003. *Wapiti (Cervus elaphus): Wild mammals of North America : biology, management, and conservation*. Second Edition. Johns Hopkins University Press, Baltimore, Mass.
- Peters, G.M., H. V. Rowley, S. Wiedemann, R. Tucker, M. D. Short, and M. Schulz. 2010. Red meat production in Australia: Life cycle assessment and comparison with overseas studies. *Environmental Science and Technology*. 44: 1327–1332.

- Peterson, H. C. 2011. An epistemology for agribusiness: peers, methods and engagement in the agri-food-bio system. *International Food and Agribusiness Management Review*. 14: 11–26.
- Pitesky, M. E., K. R. Stackhouse, and F. M. Mitloehner. 2009. Chapter 1: Clearing the Air. *Livestock's Contribution to Climate Change. Advances in Agronomy*. 103: 1-40.
- Pitt, R. E., J. S. van Kessel, D. G. Fox, A. N. Pell, M. C. Barry, and P. J. Van Soest. 1996. Prediction of ruminal volatile fatty acids and pH within the net carbohydrate and protein system. *Journal of Animal Science*. 74: 226-244.
- Piwonka, E. J., and J. L. Firkins. 1996. Effect of glucose fermentation on fiber digestion by ruminal microorganisms in vitro. *Journal of Dairy Science* 79: 2196-2206.
- Plaizier, J. C., D. O. Krause, G. N. Gozho, and B. W. McBride. 2008. Subacute ruminal acidosis in dairy cows: The physiological causes, incidence and consequences. *The Veterinary Journal*. 176: 21-31.
- Pollan, M. 2002. Power Steer *The New York Times: Magazine*. March 31, 2012. *The New York Times*, New York, NY. Accessed March 15, 2016. <http://www.nytimes.com/2002/03/31/magazine/powersteer.html> pagewanted=all.
- Prescott, L. M., J. P. Harley, and D. A. Klein. 2005. *Microbiology. Sixth Edition*. McGraw-Hill, New York, NY.
- Raun, A. P., C. O. Cooley, E. L. Potter E. L., R. P Rathmaeher, and L. F. Richardson. 1976. Effect of monensin on feed efficiency of feedlot cattle.

- Journal of Animal Science. 43:670.
- Reece, W. O., H. H. Erickson, J. P. Goff, and E. E. Uemura. 2015. Dukes' Physiology of Domestic Animals. Thirteenth edition. Wiley Blackwell, Ames, IA.
- Reynolds, C. K., D.J. Humphries, P. Kirton, M. Kindermann, S. Duval, and W. Steinberg. 2014. Effects of 3-nitrooxypropanol on methane emission, digestion, and energy and nitrogen balance of lactating dairy cows. Journal of Dairy Science. 97: 3777-3789.
- Ripple, W. J., P. Smith, H. Haberl, S.A. Montzka, C. McAlpine, and D. H. Boucher. 2014. Ruminants, climate change and climate policy. Nature Climate Change. 4: 2-5.
- Rockey Mountain Elk Foundation (RMEF). 2016. Elk Facts. Rockey Mountain Elk Foundataion. Accessed March 5, 2015. <http://www.rmef.org/ElkFacts.aspx>
- Russell, J. B. 2002. Rumen Microbiology and Its Role in Ruminant Nutrition. University Press, Ithaca, NY.
- Russell, J. B. 2007. The energy spilling reactions of bacteria and other organisms. Journal of Molecular Microbiology and Biotechnology. 13: 1-11.
- Russell, J. B., and D. B. Dombrowski. 1980. Effect of pH on the efficiency of growth by pure cultures of rumen bacteria in continuous culture. Applied and Environmental Microbiology. 39: 604-610.
- Russell, J. B., and R. Hespell. 1981. Microbial Rumen Fermentation. Journal of Dairy Science. 64: 1153-1169.

- Russell J. B., and H. J. Strobel. 1989. Effect of ionophores on ruminal fermentation. *Applied and Environmental Microbiology*. 55.1:1-6.
- Russell, J. B., and G. M. Cook. 1995. Energetics of bacterial growth: balance of anabolic and catabolic reactions. *Microbiological Reviews*. 59: 48-62.
- Russell, J. B., and J. L. Rychlik. 2001. *Science*. 292: 1119-1122.
- Russell, J. B., J. D. O'Connor, D. G. Fox, P. J. Van Soest, and C. J. Sniffen. 1992. A net carbohydrate and protein system for evaluating cattle diets: I. Ruminal fermentation. *Journal of Animal Science*. 70: 3551-3561.
- Scheifinger, C. C., B. Linehan, and M. J. Wolin. 1975. H₂ production by *Selenomonas ruminantium* in the absence and presence of methanogenic bacteria. *Applied Microbiology*. 29: 480-483.
- Smith, W. P. 1991. *Odocoileus virginianus*. *The American Society of Mammologists*. 388: 1-13.
- Sniffen, C. J., J. D. O'Connor, P. J. Van Soest, D. G. Fox, and J. B. Russell. 1992. A net carbohydrate and protein system for evaluating cattle diets: II. Carbohydrate and protein availability. *Journal of Animal Science*. 70: 3562-3577.
- St-Pierre, B., L. M. Cersosimo, S. L. Ishaq, and A. D. G. Wright. 2015. Toward the identification of methanogenic archaeal groups as targets of methane mitigation in livestock animals. *Frontiers in Microbiology*. 6: 776.
- Stephenson, J. B. 2009. Climate change science : high quality greenhouse gas emissions data are a cornerstone of programs to address climate change : testimony before the Subcommittee on Energy and Environment, Committee on Science and

Technology, House of Representatives. U.S. Govt. Accountability Office,
Washington, D.C.

Subak, S. 1994. Methane from the House of Tudor and the Ming Dynasty:

Anthropogenic emissions in the sixteenth century. *Chemosphere*. 29: 843-854.

Sveinbjörnsson, J. 2006. Substrate levels, carbohydrate degradation rate and their effects on ruminal end-product formation. Doctoral Thesis. Swedish University of Agricultural Sciences, Sweden.

Tedeschi, L. O., J. P. Muir, D. G. Riley, and D. G. Fox. 2015. The role of ruminant animals in sustainable livestock intensification programs. *International Journal of Sustainable Development & World Ecology*. 22: 452-465.

Theurer, C. B. 1986. Grain processing effects on starch utilization by ruminants. *Journal of Animal Science*. 63: 1649-1662.

Tomkins, N. W., and R. A. Hunter. 2004. Methane reduction in beef cattle using a novel antimethanogen. *Animal Production in Australia*. 25: 329.

Toprak, N. N. 2015. Do fats reduce methane emission by ruminants? - a review. *Animal Science Papers & Reports*. 33: 305-321.

Tyrrell, H.F., and P.W Moe. 1972. Net energy value for lactation of a high and low concentrate ration containing corn silage. *Journal of Dairy Science*. 55: 1106–1112.

Underwood, W. J. 1992. Rumen lactic acidosis: Part 1. Epidemiology and pathophysiology. *Continual Education in Veterinary Practices*. 14:1127–1133.

- USDA. 2012. U.S. Rice Production: Background. US Department of Agriculture, Washington, D.C. Accessed June 15, 2015
<http://www.ers.usda.gov/topics/crops/rice/background.aspx>
- USDA. 2015a. Crop Production Summary: ISSN: 1057-7823. United States Department of Agriculture, Washington, D.C. Accessed October 15, 2015.
<http://www.usda.gov/nass/PUBS/TODAYRPT/cropan15.pdf>.
- USDA. 2015b. 2015 Dietary Guidelines Advisory Committee Report. United States Department of Agriculture, Washington D.C. Accessed March 15, 2016.
<http://health.gov/dietaryguidelines/2015-scientific-report/PDFs/Scientific-Report-of-the-2015-Dietary-Guidelines-Advisory-Committee.pdf>.
- USDA-ARS. 2015. United States Department of Agriculture Agricultural Research Service National Nutrient Database for Standard Reference Release 28. United States Department of Agriculture, Washington, D.C. Accessed Jul. 10, 2015.
<https://ndb.nal.usda.gov/ndb/search>.
- USDA NASS. 2012. United States Department of Agriculture National Agricultural Statistics, Washington, D.C. Accessed on June 11, 2015.
<http://www.nass.usda.gov/>.
- U.S. Department of Health and Human Services (USDHHS) and U.S. Department of Agriculture. 2015-2020 Dietary Guidelines for Americans. Eighth Edition. December 2015. Accessed on 1 May 2016.
<http://health.gov/dietaryguidelines/2015/guidelines/>.

- U.S. EPA. 2014. Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990-2012. U.S. Environmental Protection Agency, Washington, DC. Accessed July 15, 2016. <https://www3.epa.gov/climatechange/Downloads/ghgemissions/US-GHG-Inventory-2014-Main-Text.pdf>
- U.S. EPA. 2016. Methane Emissions: Overview of Greenhouse Gases. U.S. Environmental Protection Agency, Washington, D.C. Accessed July 10, 2015. <https://www3.epa.gov/climatechange/ghgemissions/gases/ch4.html>
- van Groenigen, K. J., C. van Kessel, and B. A. Hungate. 2013. Increased greenhouse-gas intensity of rice production under future atmospheric conditions. *Nature Climate Change*. 3: 288-291.
- van Kessel, J. A. S., and J. B. Russell. 1996. The effect of pH on ruminal methanogenesis. *FEMS Microbiology Ecology*. 20: 205-210.
- Van Nevel, C. J., and D. I. Demeyer. 1977. Effect of monensin on rumen metabolism in vitro. *Applied and Environmental Microbiology*. 34: 251-257.
- Van Soest, P. J. 1973. The uniformity and nutritive availability of cellulose. *Federation Proceedings*. 32: 1804-1808.
- Van Soest, P. J. 1994. *Nutritional Ecology of the Ruminant*. Second Edition. Comstock Publishing, Ithaca, NY.
- Varga, G.A., H. F. Tyrell, D. R. Waldo, G. B. Huntington, and B. P. Glenn. 1985. Effect of alfalfa orchardgrass silages on energy and nitrogen utilization for growth by Hostein steers. *10th International Symposium on Energy and Protein Metabolism and Nutrition*. 32:86.

- Watanabe, H., and G. Tokuda. 2000. Animal cellulases. *Cellular and Molecular Life Sciences*. 58: 1167-1178.
- Weimer, P. J., J. M. Lopez-Guisa, and A. D. French. 1990. Effect of cellulose fine structure on kinetics of its digestion by mixed ruminal microorganisms in vitro. *Applied and Environmental Microbiology*. 56: 2421-2429.
- Williams, Y. J., S. Popovski, S. M. Rea, L. C. Skillman, A. F. Toovey, K. S. Northwood, and A. D. G. Wright. 2009. A vaccine against rumen methanogens can alter the composition of archaeal populations. *Applied and Environmental Microbiology*. 75: 1860-1866.
- Wilson, C. E. 2015. University of Arkansas Rice Research and Extension Center Director. Personal Communication. June 15, 2015.
- Wilson, D. B. 2011. Microbial diversity of cellulose hydrolysis. *Current Opinion in Microbiology*. 14: 259-263.
- Wright, A. D., G. P. Kennedy, C. J. O'Neill, A. F. Toovey, S. Popovski, S. M. Rea, C. L. Pimm, and L. Klein. 2004. Reducing methane emissions in sheep by immunization against rumen methanogens. *Vaccine*. 22: 3976-3985.
- Windrock International. 2012. Expanding grass based animal agriculture in the midwest: the pasture project final report. Windrock International, Arlington, VA. Accessed May 1, 2016. <http://www.houseworthrealty.com/Ranch/Pasture%20Project%20Final%20Report%20Phase%20I%20for%20WEB%20SITE.pdf>

Yan, X., H. Akiyama, K. Yagi, and H. Akimoto. 2009. Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 Intergovernmental Panel on Climate Change Guidelines. *Global Biogeochemical Cycles*. 23: 1-15.