


FERTILIZER RECOMMENDATIONS for the Gulf Coast Prairie



Adapted from Texas Agricultural Experiment Station Bulletin 431, by W. T. Carter.

FERTILIZER RECOMMENDATIONS

for the Gulf Coast Prairie

M. K. Thornton, Extension Agricultural Chemist
Bluefford Hancock, Extension Horticulturist
J. C. Smith, Superintendent, Substation No. 3
Texas A. and M. College System

For best results with fertilizers, other factors should be favorable, such as a well-prepared seed bed, good stand, absence of disease, adequate moisture and good cultural practices. Good cropping systems with legumes in rotation aid in a favorable response of crops to fertilizers.

High-analysis fertilizers usually are cheaper. Low-analysis fertilizers cost less per bag, but the cost per acre is greater for the same amount of nutrients. The grades, 5-10-5 and 10-20-10, have the same ratio (1-2-1) of nutrients, but 10-20-10 has twice as much fertilizing value as 5-10-5. It requires only one-half as much 10-20-10 per acre to supply the same amount of plant nutrients.

Fertilizers containing phosphorus should be drilled or plowed into the land. Phosphorus does not move freely into the soil. Liquid fertilizers may be used instead of solid fertilizers. Liquid fertilizers usually are much more expensive per unit of nutrients.

Anhydrous ammonia is a liquefied gas and is an excellent source of nitrogen. It is the cheapest nitrogen fertilizer when used at moderate to high rates on large acreages. When anhydrous ammonia is used as a fertilizer, the opening made by the applicator should be covered immediately to prevent loss of ammonia. Likewise, the soil should be in good tilth.

FERTILIZER RECOMMENDATIONS:

The following recommendations are expressed in pounds of nutrients per acre and do not represent fertilizer grades. The nutrients must be obtained from materials or fertilizer mixtures sold on the market.

For example, a recommendation calling for 30-60-30, which is a 1-2-1 ratio can be obtained by applying 600 pounds of 5-10-5 or 250 pounds of 12-24-12 or 300 pounds of 10-20-10. Again, if a recommendation calls for 15-60-0, this may be obtained by applying about 400 pounds of a 4-16-0 or 125 pounds of 11-48-0.

Row Crops: Fertilizer usually is applied at the time of planting or just before. Fertilizers are more efficiently used by most crops when applied in a band two to three inches to the side and two to three inches below the seed.

If equipment for applying fertilizers in bands while planting or cultivating is not available, apply the fertilizer in the water furrow and bed on it.

If large quantities of nitrogen fertilizer are to be applied part of the nitrogen should be drilled into the soil with the phosphorus and potash and the remainder applied 35 to 45 days later as a side or topdressing.

Small Grains (except rice): Fertilizers for small grains should be applied in a band midway between the seed rows through a separate drill spout. Fertilizers containing nitrogen and potash should not be allowed to touch the seed.

Phosphorus, potash and part of the nitrogen should be applied at or before seeding. The rest of the nitrogen should be applied in the spring before plants begin to joint.

Pastures: For establishing improved pastures, phosphorus, potash and part of the nitrogen should be drilled or plowed in. The remainder should be applied in 40 to 60 days as a topdressing. For maintenance, topdress as needed.

Fruit Trees: Fertilizers for fruit trees should be applied over the area covered by the spread of the limbs and cultivated into the soil.

Recommendations for fertilizers in this circular are those found best by experiments, tests and practical experience in the field. They range from the calcareous (limy) river valley clays to the sands of the uplands. If your farm contains both clays and loams, there will be two recommendations for your land.

When crops follow legumes turned under, the amount of fertilizer to be applied at planting may be reduced. Side or topdress with the amount of fertilizer suggested.

The letters NR mean that the crop is not recommended for this class of soils.

THE FERTILIZER RECOMMENDATIONS CONTAINED HEREIN ARE GENERAL IN SCOPE. SOILS VARY IN FIELDS AND AREA SO MUCH THAT SOIL TESTS SHOULD BE MADE IN ORDER TO RECOMMEND MORE DEFINITE AND ECONOMICAL FERTILIZER APPLICATIONS.

Gulf Coast Prairie

POUNDS OF NUTRIENTS TO BE APPLIED PER ACRE AT OR BEFORE PLANTING

	Calcareous alluvial clays and clay loams	Calcareous alluvial loams and sandy loams	Non-calcareous alluvial loams and sandy loams	Black clay and clay loams	Black loams and sandy loams	Gray sands and loamy sands	Remarks
FIELD CROPS							
Alfalfa	0-40-0	15-60-0	20-80-40	NR	NR	NR	0-60-0 annually for maintenance. For acid clays, 2 tons lime, for acid loams and sands, 1 ton lime
Corn Grain sorghum	40-40-0	40-40-0	30-60-60	30-60-0	30-60-60	30-60-90	Sidedress with 60-0-0 in 35 days
Sweet sorghum Sudan	40-40-0	40-40-0	30-60-60	30-60-0	30-60-30	30-60-60	Topdress with 30-0-0 each cutting or each time grazed down when soil is moist
Cotton	40-40-0	40-40-0	30-60-30	60-60-0	60-60-30	60-60-60	
Flax	30-30-0	30-30-0	30-60-30	30-30-0	30-30-30	30-60-60	Also topdress in January with 30-0-0
Legumes	0-40-0	0-40-0	0-60-30	0-60-0	0-60-30	15-60-60	
PASTURES							
Grasses and Legumes	30-60-0	30-60-0	30-60-30	30-60-0	30-60-30	30-60-60	On acid soils, 2 tons lime on clays, 1 ton on loams and sandy loams
PASTURES							
Grasses	30-30-0	30-30-0	30-60-30	30-60-0	30-60-30	30-60-60	On acid soils 1 ton lime. Topdress with 30-0-0 each time grazed or cut when soil is moist
Peanuts	NR	NR	NR	20-40-0	20-40-20	20-40-40	
Rice	NR	NR	80-40-0	80-40-0	40-40-20	40-40-20	If fields are weedy the fertilizer should be applied 30-45 days after planting
TRUCK CROPS							
Lettuce, cabbage, Mustard, Collards	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	Sidedress with 60-0-0 when heads begin to form. For mustard, 10 days from emergence
Carrots, beets, Turnips	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	Sidedress with 60-0-0 when plants are in full leaf
Sweet potatoes	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	
Irish potatoes	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	Sidedress with 60-0-0 in 40-50 days
Tomatoes, peppers Eggplant	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	Sidedress with 30-0-0
Cantaloupes, Squash, Cucumbers	30-30-0	30-60-0	30-60-30	30-60-0	30-60-30	30-60-60	Sidedress with 30-0-0 when vines start to bloom
Figs	30-30-0	30-60-0	30-60-30	30-60-0	30-60-30	30-60-60	Sidedress in May or June with 30-0-0

Figs	30-30-0	30-60-0	30-60-30	30-60-0	30-60-30	30-60-60	Sidedress in May or June with 30-0-0
Watermelons	20-40-0	20-80-0	20-80-40	20-80-0	20-80-40	20-80-60	Sidedress with 30-0-0 when vines start to bloom
Onions	40-40-0	40-80-0	40-80-40	40-80-0	40-80-40	40-80-80	
General garden	40-80-0	40-80-40	40-40-40	40-80-0	40-80-40	40-80-80	Sidedress in 40 days with 40-0-0
Pecans	Pounds of Fertilizer per bearing tree per inch diameter						Apply in Feb. or early March—Young trees 1/3 to 1/2 quantity for bearing trees
	1 lb. 33-0-0 or 1 1/2 lb. 20-0-0	1 lb. 33-0-0 or 1 1/2 lb. 20-0-0	6 lb. 10-10-5	6 lb. 10-10-5	6 lb. 10-10-10		