

for the *High Plains*



TEXAS AGRICULTURAL EXTENSION SERVICE
G. G. Gibson, Director, College Station, Texas

FERTILIZER RECOMMENDATIONS

for the High Plains

M. K. Thornton, Extension Agricultural Chemist
Donald L. Jones, Superintendent, Substation No. 8
Texas A. & M. College System

For best results with fertilizers, other factors should be favorable, such as a well-prepared seed bed, good stand, absence of disease, adequate moisture and good cultural practices. Good cropping systems with legumes in rotation aid in a favorable response of crops to fertilizers. Where soil conditions are very favorable even higher rates of fertilization than those shown may be economically advantageous.

High-analysis fertilizers usually are cheaper. Low-analysis fertilizers cost less per bag, but the cost per acre is greater for the same amount of nutrients. The grades, 5-10-5 and 10-20-10, have the same ratio (1-2-1) of nutrients, but 10-20-10 has twice as much fertilizing value as 5-10-5. It requires only one-half as much 10-20-10 per acre to supply the same amount of plant nutrients.

Fertilizers containing phosphorus should be drilled or plowed into the land. Phosphorus does not move freely into the soil. Liquid fertilizers may be used instead of solid fertilizers. Liquid fertilizers usually are much more expensive per unit of nutrients.

Anhydrous ammonia is a liquefied gas and is an excellent source of nitrogen. It is the cheapest nitrogen fertilizer when used at moderate to high rates on large acreages. When anhydrous ammonia is used as a fertilizer, the opening made by the applicator should be covered immediately to prevent loss of ammonia. Likewise, the soil should be in good tilth.

FERTILIZER RECOMMENDATIONS:

The following recommendations are expressed in pounds of nutrients per acre and do not represent fertilizer grades. The nutrients must be obtained from materials or fertilizer mixtures sold on the market.

For example, a recommendation calling for 30-60-30, which is a 1-2-1 ratio can be obtained by applying 600 pounds of 5-10-5 or 250 pounds of 12-24-12 or 300 pounds of 10-20-10. Again, if a recommendation calls for 15-60-0, this may be obtained by applying about 400 pounds of a 4-16-0 or 125 pounds of 11-48-0.

Row Crops: Fertilizer usually is applied at the time of planting or just before. Fertilizers are more efficiently used by most crops when applied in a band two to three inches to the side and two to three inches below the seed.

If equipment for applying fertilizers in bands while planting or cultivating is not available, apply the fertilizer in the water furrow and bed on it.

If large quantities of nitrogen fertilizer are to be applied part of the nitrogen should be drilled into the soil with the phosphorus and potash and the remainder applied 35 to 45 days later as a side or topdressing.

Small Grains: Fertilizers for small grains should be applied in a band midway between the seed rows through a separate drill spout. Fertilizers containing nitrogen and potash should not be allowed to touch the seed.

Phosphorus, potash and part of the nitrogen should be applied at or before seeding. The rest of the nitrogen should be applied in the spring before plants begin to joint.

Pastures: For establishing improved pastures, phosphorus, potash and part of the nitrogen should be drilled or plowed in. The remainder of the nitrogen should be applied broadcast in 40 to 60 days as a topdressing. For maintenance, topdress as needed.

Fruit Trees: Fertilizer for fruit trees may be applied over the entire area covered by the orchard when the trees are mature. In nonbearing orchards, the fertilizer should be applied over the area covered by the spread of the limbs. Keep fertilizer 1 foot away from tree trunks. Cultivate fertilizer applications into the soil.

Recommendations for fertilizers in this circular are those found best by experiments, tests and practical experience in the field. They range from the calcareous (limy) river valley clays to the sands of the uplands. If your farm contains both clays and loams, there will be two recommendations for your land.

When crops follow legumes turned under, the amount of fertilizer to be applied at planting may be reduced. Side or topdress with the amount of fertilizer suggested.

The letters NR mean that the crop is not recommended for this class of soils.

THE FERTILIZER RECOMMENDATIONS CONTAINED HEREIN ARE GENERAL IN SCOPE. SOILS VARY IN FIELDS AND AREA SO MUCH THAT SOIL TESTS SHOULD BE MADE IN ORDER TO RECOMMEND MORE DEFINITE AND ECONOMICAL FERTILIZER APPLICATIONS.

High Plains

(Irrigated Land)

POUNDS OF NUTRIENTS TO BE APPLIED PER ACRE AT OR BEFORE PLANTING

	Clays and clay loams	Loams and sandy loams	Sands and loamy sands	Remarks
FIELD CROPS				
Alfalfa & Biennial Sweetclovers	0-60-0	0-60-0	30-60-30	For maintenance 0-60-0 annually
Corn Grain sorghum	60-0-0	60-30-0	60-60-30	
Sorghum for hay Sudan	30-0-0	30-30-0	30-60-30	Sidedress with 30-0-0 after each cutting
Cotton, sesame	0-0-0*	0-0-0*	30-60-30	
Annual legumes	0-30-0	0-40-0	15-60-30	
Oats, wheat, and other small grain	0-0-0	30-30-0	30-60-30	Topdress with 40-0-0 in February or early March if the soil moisture is adequate
Pastures (cultivated) grasses only Grasses and legumes	0-0-0	0-60-0	30-60-30	Topdress with 30-0-0 each time grazed down or cut
Sugar beets, stock beets	30-0-0	30-60-0	30-60-30	Sidedress with 60-0-0
TRUCK CROPS				
Beans, peas, black-eye, etc.	0-40-0	20-60-0	20-60-30	
Cabbage, lettuce, mustard, etc.	30-30-0	40-80-0	40-80-40	Sidedress with 60-0-0 when plants begin to bud
Cantaloupes, squash, cucumbers	30-30-0	40-80-0	40-80-40	Sidedress with 30-0-0 when vines begin to bloom
Carrots, beets, turnips	0-30-0	30-60-0	30-60-30	
Irish potatoes	30-0-0	30-60-30	30-60-30	Sidedress with 40-0-0 in 40 days
Onions	40-40-0	40-80-0	40-80-40	
Sweetpotatoes	NR	30-60-30	30-60-60	
Tomatoes, peppers	40-40-0	50-100-0	50-100-50	Sidedress with 30-0-0 in 40 days
Watermelons	20-40-0	30-60-0	30-60-30	Sidedress with 30-0-0 when vines begin to bloom

Fertilizers not recommended on dry land.

*Twenty years of experimental results have failed to show any advantage for use of fertilizers on cotton. Frequently visual response will be obtained but this rarely if ever results in economically increased yields.