



DESTRUCTIVE MITES IN THE GARDEN AND HOME LANDSCAPE

Philip J. Hamman and Garland McIlveen*

Many mite species are associated with home lawns, vegetable gardens and ornamental plants. Because mites are very small they usually are not noticed until present in large numbers. This publication provides the homeowner with information needed to recognize and control damaging mite infestations.

Adult mites typically have eight legs and no antennae or wings; thus, they are in the same class as ticks, spiders, scorpions and "harvestmen" or "daddy long-legs." Mites differ from insects in the following ways:

Adult Mites	Insects
8 legs	6 legs
no antennae	2 antennae
no wings	usually 4 wings

Most mite species feed not on plants but on organic matter in the soil, on other mites or on small insects. However, several mite species do attack house and garden plants. The most common of these are spider mites and eriophyid mites.

Spider mites are oval-shaped, about 1/25-inch in diameter and, when viewed through a magnifying glass, appear to be sparsely covered with long hairs. Many members of this family are gregarious, and some produce large quantities of fine webbing.

Eriophyid mites, commonly called gall mites, rust mites, or bud mites, are much smaller than spider mites (about 1/25-inch long) and produce no webbing. These cigar-shaped mites are difficult to see,

even with the aid of a magnifying glass. Mites from both families feed by inserting their mouthparts into a plant and sucking out plant juices.

Spider Mites (Family Tetranychidae)

Spider mites usually begin colonies on the undersides of leaves. Early plant damage appears as "stippling" or "sandblasting" at the site of the colony. Spider mites develop through egg, larval and two nymphal stages before molting to adults. The eggs are spherical in shape and about one-quarter to one-third as large as the adult female mite. Newly deposited eggs are translucent and vary from cream color to bright orange or red, depending on the mite species. A six-legged larva hatches from the egg. Two nymphal stages follow the larval stage and each nymph has eight legs, as does the adult mite.

Spider mites can reproduce asexually. In some spider mites species, only females are known. In other species, unmated adult females produce only male young, though after mating mostly female young are produced. Males are about one-half to three-fourths as large as adult females and are arrowhead-shaped. The following are the most common spider mites in Texas.

Two-Spotted Spider Mite (*Tetranychus urticae* Koch)

Description and Life History. The two-spotted spider mite feeds on a wide variety of plants and is one of the most common mites attacking home

*Extension urban entomologists, The Texas A&M University System.




Figure 1. A spider mite of the family Tetranychidae.

gardens and ornamental plants. The adult female is pale yellow or pale to dark green with conspicuous dark spots on each side. These darkened areas are caused by food showing through the transparent body wall. The eggs are spherical, clear and colorless when first laid, becoming opaque and finally ivory just before hatching. The newly hatched larva is colorless at first, though it becomes greenish after feeding. Both nymphal stages are green to pale yellow. Two-spotted spider mites usually overwinter as adult females on the host plant or in nearby debris. These overwintering females are pale orange to straw-colored and active, though they do not produce eggs during the winter. In mild climates or in greenhouses, adult mites may remain active and reproduce year-round.

Two-spotted spider mites usually begin colonies on the undersides of plant leaves. Webbing is visible to the naked eye, and all stages of mite development occur in and beneath the webbing. Many female mites may be present in each colony, and several colonies may be present on each infested leaf. Small colonies usually begin near a leaf vein and with favorable weather spread rapidly to nearby leaves and plants.

The rate of population increase highly depends on the weather. Dry hot weather (80 to 90 degrees F.) is most favorable to mite reproduction. A single female mite produces from 2 to 20 eggs per day, for a total of about 150 to 200 eggs during her 2- to 3-week lifetime. Development from egg to adult requires 5 to 15 days.

During dry, hot weather, feeding damage to the plant intensifies. As the host plant becomes seriously injured and damaged leaves turn yellow or brown, mites begin to migrate to uninfested plants. During heavy migrations, thousands of mites cluster on leaf tips at the tops of infested plants. Migrating mites occasionally form long chains of individuals hanging from the plant. Individual mites drop from the chains and walk or are carried by wind to other plants.

Damage. Mites usually feed on lower surfaces, though occasionally they infest upper leaf surfaces.




Figure 2. A rust mite of the family Eriophyidae.

Light to moderate infestations on lower surfaces cause stippling on upper leaf surfaces. As feeding continues and the mite population increases, leaves become lighter in color as a result of the mites removing the green chlorophyll from leaf tissue. Heavily damaged leaves turn tan or bronze, become desiccated and usually die.

Carmine Spider Mite (*Tetranychus cinnabarinus* Boisduval)

Description and Life History. The carmine spider mite is dark red to carmine in color with a dark area on each side. Eggs of this mite are light orange to red. The biology of the carmine spider mite is essentially the same as the two-spotted spider mite, and these species attack the same host plants. Carmine mites are quite common on outdoor plants and in greenhouses, and can multiply explosively.

Damage. The carmine spider mite usually feeds on the lower surface of the leaf as does the two-spotted spider mite. Light infestations cause stippling which becomes progressively worse with higher populations. Webbing is usually abundant, and high populations may web over entire leaves.

Pecan Spider Mite (*Eotetranychus hicoriae* McGregor)

Description and Life History. The pecan spider mite is a straw-colored to pale green mite with several to many dark spots along the back and sides. This species feeds on pecan, chestnut, hickory and oak trees throughout the eastern United States. It initiates small colonies on both sides of leaves, though it occurs principally on the undersides. Bright yellow females hibernate in protected places on tree bark or in limb crotches.

A second species of spider mite, *Oligonychus viridis* (Banks), also attacks pecan trees in southeastern portions of the United States. This mite prefers the upper leaf surfaces but occurs on both sides of infested leaves.

Damage. Pecan spider mites are not considered major pests because natural enemies normally control them. However, occasional outbreaks occur, especially after trees are treated with pesticides to control other pecan pests. These mites cause dark brown or liver-colored blotches and moderate to heavy webbing on the leaflets. Heavy infestations in the early fall may cause premature leaf fall that starts at the lower branches. Leaves remaining on the tree appear dry or scorched. Excessive premature foliage loss may prevent trees from blooming properly the following spring.

Banks Grass Mite (*Oligonychus pratensis* Banks)

Description and Life History. The Banks grass mite attacks grassy plants such as garden corn, Johnson-grass and Bermudagrass lawns. The adult female is pale to deep green with numerous dark spots on the back and sides.

Mite infestations usually develop first on the undersides of leaves along the main veins. Heavy populations cause the lower portions of grasses to become matted with webbing. High temperatures are most favorable to population buildups of this mite. Adult females hibernate in plant debris at the bases of host plants.

Damage. This mite usually causes serious damage only in areas of meager rainfall or infrequent irrigation. Thus, lawns or gardens frequently watered are not normally damaged. Light infestations on Bermudagrass lawns cause stippled leaves and light to moderate webbing at the bases of the plants. Usually, lawn infestations need not be controlled except in late summer when temperatures are high.

Early damage to corn appears at stippling on the mature leaves, followed by a red coloration at the sites of heavy mite feeding. With dry, hot weather, high populations develop very quickly and can kill corn at the tassel stage of development. Such plants look as though they died from lack of moisture and are usually heavily enveloped in sheets of webbing produced by the many thousands of mites.

Brown Wheat Mite (*Petrobia latens* Muller)

Description and Life History. The brown wheat mite sometimes attacks onions, carrots, lettuce, melons and strawberries, although these mites prefer to feed on grass-type plants. High populations will migrate and may damage gardens and ornamental plants. Only female mites of this species occur. The adult female is dark brown or greenish brown with long, light yellow legs. The front part of the legs is much longer than the body. These mites produce no webbing and may drop to the ground or run rapidly when disturbed.

Brown wheat mites do not deposit eggs on the foliage but on soil particles in the vicinity of host plants. They produce two types of eggs, depending

on environmental conditions. During dry, hot summer weather or during the fall when host plants are in poor condition, the female produces a white egg which does not hatch for many months, and then only after exposure to moisture. Subsequent population development depends on a lack of heavy rainfall. During dry spring or fall weather, the female produces a cherry-red egg, and populations increase rapidly. The brown wheat mite occasionally deposits its eggs inside dwellings. To discourage this maintain grass-free band 18 to 24 inches wide around buildings.

Damage. The brown wheat mite is primarily a dry-weather pest. Damage is quite similar to that caused by drought, even where there appears to be sufficient moisture. Infested plants's leaves have a fine mottling. Such plants appear bronze or yellow when observed at a distance. In gardens, infestations usually start in a localized area and then spread rapidly. During early stages of infestation, plant foliage turns yellow and later becomes dry and brittle.

Clover Mite (*Bryobia praetiosa* Koch)

Description and Life History. The clover mite feeds on a wide variety of plants including clovers, lawn grasses and ornamental flowers. Populations usually are highest in spring, late summer or early fall. The mite overwinters in the egg stage away from the host plant, though some active stages may survive the winter in warmer climates. Larvae hatch in spring when temperatures are 65 to 75 degrees F. Newly hatched individuals seek host plants on which to feed, leaving the host plants for each molt to the next developmental stage. The dark-red females behave similarly, leaving the host plant to lay their bright red eggs and returning to feed. When conditions become unfavorable for the mites, such as when the lawn is mowed or hot summer or cold winter weather approaches, these mites seek shelter on trees or in dwellings, where they remain until conditions improve or until they die. Typically, only the eggs survive unfavorable conditions.

When disturbed, these mites drop from the plant, curl their legs, and remain motionless.

Damage. Clover mites produce no webbing, and injury to plants first appears as a winding, narrow trail of tiny bleached spots on infested leaves. More extensive damage causes the foliage to turn yellow or brown, and leaves eventually wilt and die. To discourage mites from entering dwellings for molting or egg laying, maintain a grass-free band 18 to 24 inches wide around such buildings.

For more information on this mite, refer to Extension publication L-2040 *Clover Mites*.

Texas Citrus Mite (*Eutetranychus banksi* McGregor)

Description and Life History. The Texas citrus mite is tan to brownish green with dark brown to greenish spots on sides. This mite occurs year-round on Texas

citrus, with smallest populations occurring from February to April and largest numbers during May and June. Low relative humidity, temperatures of about 80 degrees F. and the absence of rainfall favor buildups of this mite. Low to moderate populations usually occur in the south quadrant of the tree. Texas citrus mites appear on upper leaf surfaces and produce no webbing.

Damage. Injury to citrus leaves includes stippling, light colored spots and a greyish or silvery appearance. Under conditions of drought or low relative humidity with winds, the combined effects of mites and weather may cause heavy leaf and fruit drop and, occasionally, twig die-back.

Eriophyid Mites (Family Eriophyidae)

Eriophyid mites are very small, four-legged mites which cause many kinds of plant damage. Many of these mites inject salivary chemicals into plants during feeding, which causes either leaf discoloration or leaf shape distortion. Each eriophyid mite species is generally quite specific as to the host plant it will attack.

The Bermudagrass or couchgrass mite, *Eriophyes cynodontiensis* (Sayed), infests Bermudagrass in climates where atmospheric humidity is low. The mites cause thickened shoots to develop where they then congregate, living under the broadened leaf bases. Grass infested by Bermudagrass mites is tightly bunched and stunted.

The tomato rust mite, *Aculops lycopersici* (Masse), attacks many solanaceous plants including tomato, pepper, potato, eggplant, tobacco and petunia. Plant injury begins at the base of the plant and spreads upward over the leaves and stems. Damage to tomatoes consists of bronzed stems which may crack longitudinally. Leaves turn brown, become paper-like in appearance (though they do not wilt) and may drop if injury is severe. Potatoes do not show stem damage, but the leaves of potatoes dry severely and plants may die from heavy infestations. Eggplant usually supports large populations of the mite, with the damage consisting of distorted or crinkled leaves. The most favorable growing conditions for the tomato rust mite is a temperature of 80 degrees F. and 30 percent relative humidity. Developmental time from egg to adult may be only 6 or 7 days. Adult female mites may produce up to 50 or more eggs during their lifetimes.

The citrus rust mite, *Phyllocoptruta oleivora* (Ashmead), is an occasional pest of citrus in Texas. This eriophyid mite feeds on both upper and lower leaf surfaces and on fruit; however, they prefer green fruit and the undersides of leaves. Warm humid con-

ditions favor development of the mite. At average summer temperatures of 91 degrees F., development from egg to adult requires about 7 to 10 days. During the winter months or at an average 72 degrees F., development requires 16 days or more. Each female mite lays about 20 eggs during her 10- to 20-day lifetime. The female lays these eggs on fruit and on leaf surfaces. Heavy populations of this mite cause bronzing of leaves and green twigs, and loss of vitality. Damage to fruit is only superficial, though deterioration of such fruit may be more rapid than normal.

MITE CONTROL

The most important aspect of mite control is early detection. Make routine checks for infestations, especially in ornamental or vegetable gardens. Check both upper and lower surfaces of several leaves from several plants to discern webbing or adult mites. Even low populations can cause damage easily seen by the watchful gardener. Quickly check yellowing or other unusual leaf appearances.

Chemical Control. Chemical controls for mites in the home vegetable garden are suggested in Table 1, and a list of miticides that can be used in the home landscape is given in Table 2. Mite control is often difficult because mites resist the miticide used. The homeowner must be sure to achieve complete spray coverage of infested plants. Many of the miticides are contact materials, and the spray solution must come in contact with the mites to achieve satisfactory control. Repeat applications (two to four) of a miticide are generally required for satisfactory control. Read all label instructions carefully before using a miticide material. Store all unused materials in a locked area to avoid accidental poisoning of small children or pets.

Non-Chemical Control. Plant-feeding mites, particularly spider mites, are generally highly susceptible to destruction by water. When feasible, a fine, hard stream of water (through an ordinary garden nozzle) usually kills most mites present. Hold nozzle 4 to 8 inches from the infested leaf surface and move across the leaf to get full coverage. Direct water stream at the leaf surface where most mites are present and wet leaf completely. Destruction of the webbing by water washes most mites from the plant and causes others to become helplessly entangled. In either case, those mites wetted by water are killed and cause no further damage. However, eggs are unaffected by this procedure and a few mites are frequently missed in the initial washing. Thus, washings should be repeated two or three times at 3- to 5-day intervals to kill newly hatched mites and escaped adults.

The information given herein is for educational purposes only. Reference to commercial products or trade names is made with the understanding that no discrimination is intended and no endorsement by the Cooperative Extension Service is implied.

Table 1. Suggested Miticides for the Home Vegetable Garden

Pesticides

Vegetables	Diazinon	Kelthane® ¹ (dicofol)	Malathion	Pyrethrins	Sulfur ²
Beans	x	x	x	x	x
Beets					x
Cantaloupes	x	x	x	x	
Carrots	x				x
Corn	x			x	x
Cucumbers	x	x	x	x	
Eggplant		x	x	x	x
Lettuce	x		x	x	x
Mustard Greens			x		x
Okra					x
Peas	x		x	x	x
Peppers	x	x	x	x	x
Potatoes	x		x	x	x
Squash	x	x	x	x	
Tomatoes	x	x	x	x	x
Watermelons	x	x	x	x	

¹Does not harm beneficial insects.

²Do not apply to cucumbers, squash or melons unless plants are dry or injury may result.

Always read and follow carefully the instructions on the container label. Apply miticides only to the crop specified.

This information is provided for informational purposes only. It is not intended to be used as a substitute for professional advice. Always read and follow the instructions on the container label.

Table 2. Suggested Miticides for the Home Landscape

Plants	Pesticides	acephate (Orthene®)	carbophenothion (Trithion®)	chlorbenzilate	chlorpyrifos (Dursban®)	diazinon	dicofof (Kelthane®)	disulfoton (Di-Syston®)	ethion	malathion	oils, dormant and summer	oxydemeton-methyl (Metasystox-R®)	pyrethrins	sulfur
Bermudagrass						x	x		x					x
House plants	x				x		x						x	
Roses	x				x	x	x	x		x	x	x	x	x
Woody shrubs	x				x	x	x		x		x	x		x
Citrus			x	x			x		x		x			
Marigolds					x		x							x
Pecans						x	x			x	x			x
Spring flowering annuals	x				x	x	x			x		x	x	
Summer flowering annuals	x				x	x	x			x		x	x	

Follow label directions carefully for rates of application.

The original printing was prepared by Frank E. Gilstrap, Associate Professor, Department of Entomology, Texas Agricultural Experiment Station and Clifford E. Hoelscher, Extension entomologist, The Texas A&M University System.

Educational programs conducted by the Texas Agricultural Extension Service serve people of all ages regardless of socioeconomic level, race, color, sex, religion, handicap or national origin.

Cooperative Extension Work in Agriculture and Home Economics, The Texas A&M University System and the United States Department of Agriculture cooperating. Distributed in furtherance of the Acts of Congress of May 8, 1914, as amended, and June 30, 1914.
20M—8-83, Revision