



US006680195B1

(12) **United States Patent**
Patti et al.

(10) **Patent No.:** **US 6,680,195 B1**
(45) **Date of Patent:** **Jan. 20, 2004**

(54) **EXTRACELLULAR MATRIX-BINDING PROTEINS FROM STAPHYLOCOCCUS AUREUS**

(75) Inventors: **Joseph M. Patti**, Cumming, GA (US); **Timothy J. Foster**, Dublin (IE); **Elisabet Josefsson**, Gothenburg (SE); **Deidre Ni Eidhin**, Dublin (IE); **Magnus A. O. Hook**, Houston, TX (US); **Samuel E. Perkins**, Houston, TX (US)

(73) Assignees: **Inhibitex, Inc.**, Norcross, GA (US); **Bioresearch Ireland**, Dublin (IE); **The Texas A&M University System**, College Station, TX (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/200,650**

(22) Filed: **Nov. 25, 1998**

Related U.S. Application Data

(60) Provisional application No. 60/098,427, filed on Aug. 31, 1998, and provisional application No. 60/066,815, filed on Nov. 26, 1997.

(51) **Int. Cl.**⁷ **C07H 21/04**; C12N 15/00

(52) **U.S. Cl.** **435/320.1**; 536/23.7

(58) **Field of Search** 435/320.1, 325, 435/252.3, 254.11, 257.2, 6; 534/23.7, 24.32; 424/185.1, 243.1; 530/350

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,008,341 A * 12/1999 Foster et al.

FOREIGN PATENT DOCUMENTS

EP 0 786 519 A2 7/1997
WO 97/30070 8/1997

OTHER PUBLICATIONS

Josefsson et al., "Three new members of the serine-aspartate repeat protein multigene family of *Staphylococcus aureus*", *Microbiology* (1998), vol. 144, pp. 3387-3395.

McCrea et al., "B-47. A Family of Putative Adherence Proteins Related to the Clumping Factor of *Staphylococcus aureus*", May 17, 1998.

Boden and Flock, *Infect. Immun.* 57:2358-2363, 1989.

Boden and Flock, *Mol. Microbiol.* 12:599-606, 1994.
Cheung et al., *J. Clin. Invest.* 87:2236-2245, 1991.
Cheung et al., *Infect. Immun.* 63:1914-1920, 1995.
Dickinson et al., *Infect. Immun.* 63:3143-3150, 1995.
Hermann et al., *J. Infect. Dis.* 167:312-322, 1993.
Homonylo-McGavin et al., *Infect. Immun.* 61:2479-2485, 1993.
Jones et al., *Nature* 321:522-525, 1986.
McDevitt et al., *Mol. Microbiol.* 11:237-248, 1994.
McDevitt et al., *Mol. Microbiol.* 16:895-907, 1995.
McDevitt et al., *Eur. J. Biochem.* 247:416-424, 1997.
Moreillon et al., *Infect. Immun.* 63:4738-4743, 1995.
Patti et al., *Ann. Rev. Microbiol.* 48:585-617, 1994.
O'Connell et al., *J. Biol. Chem.*, 273:6821-6829, 1998.
Tempest et al., *Biotechnology* 9:266-273, 1991.
Vaudaux et al., *J. Infect. Dis.* 160:865-875, 1989.
Vaudaux et al., *Infect Immun.* 63:585-590, 1995.
Wolz et al., *Infect. Immun.* 64:3142-3147, 1996.

* cited by examiner

Primary Examiner—Patricia A. Duffy


(74) *Attorney, Agent, or Firm*—Larson & Taylor PLC

(57) **ABSTRACT**

Isolated extracellular matrix-binding proteins, designated ClfB, SdrC, SdrD and SdrE, and their corresponding amino acid and nucleic acid sequences and motifs are described. The proteins, peptides, fragments thereof or antigenic portions thereof are useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection and as scientific research tools. Further, antibodies or antibody fragments to the proteins, peptides, fragments thereof or antigenic portions thereof are also useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection. In particular, the proteins or antibodies thereof may be administered to wounds or used to coat biomaterials to act as blocking agents to prevent or inhibit the binding of *S. aureus* to wounds or biomaterials. ClfB is a cell-wall associated protein having a predicted molecular weight of approximately 88 kDa and an apparent molecular weight of approximately 124 kDa, which binds both soluble and immobilized fibrinogen. ClfB binds both the alpha and beta chains of fibrinogen and acts as a clumping factor. SdrC, SdrD and SdrE are cell-wall associated proteins that exhibit cation-dependent ligand binding to the extracellular matrix. It has been discovered that in the A region of SdrC, SdrD, SdrE, ClfA and ClfB, there is a highly conserved amino acid sequence that can be used to derive a consensus motif of TYTFTDYVD (SEQ ID NO: 16).

6 Claims, 34 Drawing Sheets

ClfA



ClfA





FIG. 1

FIG. 2



Probe used to
screen mutants

Fragment modified in mutants





FIG. 3



1 kb

FIG. 4

S>
TAGAAATGAAATGGAGTAATATTTTGAAAAAAGAATTGATTATTTGTGCAATAAGCAGAATAAGTATTCGATTAGACGTTTTA
CAGT
N G V I F L K K R I D Y L S N K Q N K Y S I R R F
T V 22

A>
AGGTACCACATCAGTAATAGTAGGGCAACTATACTATTTGGGATAGGCAATCATCAAGCACAAGCTTCAGAACAATCGAACGATA
CAAC
G T T S V I V G A T I L F G I G N H Q A Q A S E Q S N D
T T 52
GCAATCTTCGAAAAATAATGCAAGTGCAGATTCGAAAAAACAATATGATAGAAACACCTCAATTAATAACACGGCTAATGATA
CATC
Q S S K N N A S A D S E K N N M I E T P Q L N T T A N D
T S 82
TGATATTAGTGCAAACACAAACAGTGCAGATTCGAAAAAACAATATGATAGAAACACCTCAATTAATAACACGGCTAATGATA
AGCC
D I S A N T N S A N V D S T T K P M S T Q T S N T T T T
E P 112
AGCTTCAACAAATGAAACACCTCAACCGACGGCAATTAATAATCAAGCAACTGCTGCAAAAATGCAAGATCAAACTGTTCTCAAG
AAGG
A S T N E T P Q P T A I K N Q A T A A K M Q D Q T V P Q
E G 142
AAATTCCTCAAGTAGATAATAAAACAACGAATGATGCTAATAGCATAGCAACAACAGTGAGCTTAAAAATTCTCAACATTAGATT
TACC
N S Q V D N K T T N D A N S I A T N S E L K N S Q T L D
L P 172
ACAATCATACCACAAACGATTTCCAATGCGCAAGGAAGTAGTAAACCAAGTGTAGAACGAGAGCTGTACGTAGTTTAGCTGTTG
CTGA
Q S S P Q T I S N A Q G T S K P S V R T R A V R S L A V
A E 202
ACCGGTAGTAAATGCTGCTGATGCTAAAGGTACAAATGTAATGATAAAGTTACGGCAAGTAATTTCAAGTTAGAAAAGACTACAT
TTGA
P V V N A A D A K G T N V N D K V T A S N F K L E K T T
F D 232
CCCTAATCAAAGTGGTAACACATTTATGGCGGCAATTTTACAGTGACAGATAAAGTGAATCAGGGGATATTTTACAGCGAAGT
TACC
P N Q S G N T F M A A N F T V T D K V K S G D Y F T A K
L P 262
AGATAGTTAACTGGTAATGGAGACGTGGATTATTCCTAATTCAAATAATACGATGCCAATTGCAGACATTAAAAGTACGAATGGCG
ATGT
D S L T G N G D V D Y S N S N N T M P I A D I K S T N G
D V 292
TGTAGCTAAAGCAACATATGATATCTTGACTAAGACGTATACATTTGTCTTTACAGATTATGTAATAATAAAGAAAAATATTAACG

FIG. 5A

GACA

V A K A T Y D I L T K T Y T F V F T D Y V N N K E N I N
G Q 322ATTTTCATTACCTTTATTTACAGACCGAGCAAAGGCACCTAAATCAGGAACATATGATGCCAATATTAATATTGCCGGATGAAATGT
TTAAF S L P L F T D R A K A P K S G T Y D A N I N I A D E M
F N 352TAATAAAATTACTTATAACTATAGTTCGCCAATTGCAGGAATTGATAAACCAATGGCGCAACATTTCTCTCAAATTATTGGTG
TAGAN K I T Y N Y S S P I A G I D K P N G A N I S S Q I I G
V D 382TACAGCTTCAGGTCAAACACATACAAGCAAACAGTATTTGTTAACCTAAGCAACGAGTTTLAGGTAATACGTGGGTGTATATTA
AAGGT A S G Q N T Y K Q T V F V N P K Q R V L G N T W V Y I
K G 412CTACCAAGATAAAATCGAAGAAAGTAGCGGTAAGTAAGTGTACAGATACAAAACGAGAATTTTGAAGTGAATGATACATCTA
AATTY Q D K I E E S S G K V S A T D T K L R I F E V N D T S
K L 442ATCAGATAGCTACTATGCAGATCCAAATGACTCTAACCTTAAAGAAGTAACAGACCAATTTAAAAATAGAATCTATTATGACCATC
CAAAS D S Y Y A D P N D S N L K E V T D Q F K N R I Y Y E H
P N 472TGTAGTAGTATTAATTTGGTGATATTACTAAAACATATGTAGTATTAGTAGAAGGCATTACGACAATACAGGTAAGAACTTAA
AAACV A S I K F G D I T K T Y V V L V E G H Y D N T G K N L
K T 502TCAGGTTATTCAAGAAAATGTTGATCCTGTAACAAATAGAGACTACAGTATTTTCGGTTGGAATAATGAGAATGTTGTACGTTATC
GTGGQ V I Q E N V D P V T N R D Y S I F G W N N E N V V R Y
G G 532

P>

TGGAAGTGCTGATGGTGATTCAGCAGTAAATCCGAAAGACCCAACTCCAGGGCCCGGTTGACCCAGAACCAAGTCCAGACCCAG
AACCG S A D G D S A V N P K D P T P G P P V D P E P S P D P
E P 562

R>

AGAACCAACGCCAGATCCAGAACCAAGTCCAGACCCAGAACC GAACCAAGCCAGACCCGGATCCGGATTCGGATTCAGACAGTG
ACTCE P T P D P E P S F D P E P E P S P D P D P D S D S D S
D S 592AGGCTCAGACAGCGACTCAGGTTGAGATAGCGACTCAGAATCAGATAGCGATTCGGATTCAGACAGTGATTCAGATTCAGACAGCG
ACTC**FIG. 5B**

G S D S D S G S D S D S E S D S D S D S D S D S D S D S
 D S 622
 AGAATCAGATAGCGATTGATTCAGAATCAGATAGCGACTCAGATTGATTCAGATAGCGATTGATTCAGATAGCGATTGATTCAGATAGCG
 ATTC
 E S D S D S E S D S D S D S D S D S D S D S D S D S D S D S
 D S 652
 GGATTGACAGCTGATTCAGATTGACAGCGACTCAGAATCAGATAGCGACTCAGAATCAGATAGTGAGTCAGATTGACAGCTG
 ACTC
 D S D S D S D S D S D S E S D S D S E S D S E S D S D S
 D S 682
 GGACTCAGACAGTATTCAGACTCAGATAGCGATTGACTCAGATAGCGATTGATTCAGACAGCGACTCAGATTGACAGCG
 ACTC
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S
 D S 712
 AGACTCAGATAGCGACTCAGACTCAGACAGCGACTCAGATTGATTCAGATAGCGATTGACTCAGACAGCGACTCAGACTCAGACAGCG
 ACTC
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S
 D S 742
 AGACTCAGATAGCGACTCAGATTGATTCAGATAGCGATTGACTCAGACAGCGACTCAGATTGATTCAGATAGCGATTGACTCAGACAGCG
 ATTC
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S
 D S 772
 AGATTGACAGCGACTCAGACTCGGATAGCGATTGATTCAGATAGCGATTGATTCAGACAGTATTCAGATTGACAGCG
 ACTC
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S
 D S 802
 AGACTCGGATAGCGACTCAGACTCAGACAGCGATTGACTCAGATAGCGACTCAGACTCGGATAGCGACTCGGATTGATTCAGATAGCG
 ACTC
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S
 D S 832

W>

AGACTCAGATAGTGACTCCGATTCAAGAGTTACACCACCAATAATGAACAGAAAGCACCATCAATCCTAAAGGTGAAGTAAACC
 ATTC
 D S D S D S D S R V T P P N N E Q K A P S N P K G E V N
 H S 862

M>

TAATAAGGTATCAAAACAACACAAAAGTATGCTTTACCAGAAACAGGAGATAAGAGCGAAAACACAAATGCAACTTTATTTGGTG
 CAAT
 N K V S K Q H K T D A L P E T G D K S E N T N A T L F G
 A M 892
 GATGGCATTATTAGGATCATTACTATTGTTTAGAAAACCGAAGCAAGATCATAAAGAAAAAGCGTAAATACTTTTTAGCCGAAT
 ACAT
 M A L L G S L L L F R K R K Q D H K E K A

FIG. 5C

913

TTGTATTCGGTTTTTTTGTTGAAAATGATTTTAAAGTGAATTGATTAAGCGTAAAATGTTGATAAAGTAGAATTAGAAAGGGTCA
TGAC

GTATGGCTTATATTCATTAAACTATCATTACCAACAATTGGTATGCATCAAATTTGACAGTCATTTTACCGGAAGAACGAGAA
TTC

FIG. 5D

clfB
clfA

DVDYNSNNTMPIADIKSTNGDVVAKATYDILTKTYTFVFTDYVNNKENINGQFSLPLFT 329
-VTSTAKVPPIMAGDQVLANGVIDSDG-----NVIYTFDYVNTKDDVKATLTMPAYI 339
* . . . * ** ***** * . . . * .

FIG. 6

S>

ATGAATAATAAAAAGACAGCAACAAATAGAAAAGGCATGATACCAAATCGATTAACAATTTTCGATAAGAAAGTATTCTGTAGG
TACTM N N K K T A T N R K G M I P N R L N K F S I R K Y S V G
T 30

A>

GCTTCAATTTTAGTAGGGACAACATTGATTTTGGGTTAAGTGGTCATGAAGCTAAAGCGGCAGAACATACGAATGGAGAATTAAA
TCAAA S I L V G T T L I F G L S G H E A K A A E H T N G E L N
Q 60TCAAAAAATGAAACGACAGCCCAAGTGAAGAATAAAACAACATAAAAAAGTTGATAGTCGCAACTAAAAGACAATACGCCAACTGC
AACTS K N E T T A P S E N K T T K K V D S R Q L K D N T Q T A
T 90GCAGATCAGCCTAAAGTGACAATGAGTGATAGTGCAACAGTTAAAGAACTAGTAGTAACATGCAATCACCACAAAACGCTACAGC
TAATA D Q P K V T M S D S A T V K E T S S N M Q S P Q N A T A
N 120CAATCTACTACAAAAGTAGCAATGTAACAACAAATGATAAATCATCAACTACATATAAGTAATGAAACTGATAAAAGTAATTTAAC
ACAAQ S T T K T S N V I T N D K S S T T Y S N E T D K S N L T
Q 150GCAAAAGATGTTTCAACTACACCTAAAACAACGACTATTAACCAAGAACTTTAAATCGCATGCCAGTGAATACTGTTGCAGCTCC
ACAAA K D V S T T P K T T T I K P R T L N R M A V N T V A A P
Q 180CAAGGAACAAATGTTAATGATAAAGTACATTTTCAAATATTGACATTGCGATTGATAAAGGACATGTTAATCAGACTACTGGTAA
AACTQ G T N V N D K V H F S N I D I A I D K G H V N Q T T G K
T 210GAATTTTGGGCAACTTCAAGTATGTTTAAAATAAAAGCAAATTACACAATCGATGATTCTGTTAAAGAGGGCGATACATTAC
TTTTE F W A T S S D V L K L K A N Y T I D D S V K E G D T F T
F 240AAATATGGTCAATATTTCCGTCAGGATCAGTAAGATTACCTTCACAACTCAAATTTATATAATGCCCAAGGTAATATTATTGC
AAAAK Y G Q Y F R P G S V R L P S Q T Q N L Y N A Q G N I I A
K 270GGTATTTATGATAGTACAACAAACACAACATATACTTTTACGAACTATGTAGATCAATATACAAATGTTAGAGGTAGCTTTGA
ACAAG I Y D S T T N T T T Y T F T N Y V D Q Y T N V R G S F E
Q 300

GTTGCATTTGCGAAACGTAAAAATGCAACAACGATAAAACAGCTTATAAAATGGAAGTAACTTTAGGTAATGATACATATAGCGA

FIG. 7A

AGAA
V A F A K R K N A T T D K T A Y K M E V T L G N D T Y S E
E 330
ATCATTGTCGATTATGGTAATAAAAAAGCACACCCTTATTTCAAGTACAACTATATTAACAATGAAGATTATCGCGTAATAT
GACT
I I V D Y G N K K A Q P L I S S T N Y I N N E D L S R N M
T 360
GCATATGTAATCAACCTAAAAATACATATACTAAACAAACGTTTGTACTAATTTAACTGGATATAAATTTAATCCAATGCAAA
AAAC
A Y V N Q P K N T Y T K Q T F V T N L T G Y K F N P N A K
N 390
TTCAAAATTTACGAAGTGACAGATCAAAATCAATTTGTGGATAGTTTACCCTGATACTTCAAACTTAAAGATGTTACTGATCA
ATTC
F K I Y E V T D Q N Q F V D S F T P D T S K L K D V T D Q
F 420
GATGTTATTATAGTAATGATAATAAACAGCTACAGTCGATTTAATGAAAGGCCAAACAAGCAGCAATAACAATACATCATTCA
ACAA
D V I Y S N D N K T A T V D L M K G Q T S S N K Q Y I I Q
Q 450
GTTGCTTATCCAGATAATAGTTCAACAGATAATGGAAAAATTGATTATACTTTAGACACTGACAAAATAAATATAGTTGGTCAAA
TAGT
V A Y P D N S S T D N G K I D Y T L D T D K T K Y S W S N
S 480

B1>

TATTCAAATGTGAATGGCTCATCAACTGCTAATGGCGACAAAAGAAATATAATCTAGGTACTATGTATGGGAAGATACAAATAA
AGAT
Y S N V N G S S T A N G D Q K K Y N L G D Y V W E D T N K
D 510
GCTAAACAAGATGCCAATGAAAAGGGATTAAGGTGTTTATGTCATTCTTAAAGATAGTAACGGTAAAGAATTAGATCGTACGAC
AACA
G K Q D A N E K G I K G V Y V I L K D S N G K E L D R T T
T 540
GATGAAAATGGTAAATATCAGTTCAGTTCAGTGGTTAAGCAATGGAACCTATAGTGTAGAGTTTTCAACACCAGCCGGTTATACCCGAC
AACT
D E N G K Y Q F T G L S N G T Y S V E F S T P A G Y T P T
T 570
GCCAATGTAGGTACAGATGATGCTGTAGATTCTGATGGACTAACTACAACAGGTGTCATTAAAGACCGCTGACAACATGACATTAGA
TAGT
A N V G T D D A V D S D G L T T T G V I K D A D N M T L D
S 600

B2>

GGATTCTACAAACACCAAATATAGTTAGGTGATTATGTTGGTACGACAGTAATAAGATGGTAAACGAGATTCGACTGAAA
AGGA

FIG. 7B

G F Y K T P K Y S L G D Y V W Y D S N K D G K R D S T E K
G 630
ATTAAAGGTCTTAAAGTTACTTTGCAAAACGAAAAAGCGAAGTAATTGGTACAACGAAACAGATGAAAATGGTAAATACCGCTT
TGAT
I K G V K V T L Q N E K G E V I G T T E T D E N G K Y R F
D 660
AATTTAGATAGTGGTAAATACAAAGTTATCTTTGAAAAACCTGCTGGCTTAACTCAAACAGGTACAATACAACGAAAGATGATAA
AGAT
N L D S G K Y K V I F E K P A G L T Q T G T N T T E D D K
D 690

R>

GCCGATGGTGGCGAAGTTGATGTAACAATTACGGATCATGATGATTTCACTTGATAATGGCTACTACGAAGAAGAAACATCAGA
TAGC
A D G G E V D V T I T D H D D F T L D N G Y Y E E E T S D
S 720
GACTCAGATTCTGACAGCGATTGACTCAGACTCAGATAGCGACTCAGATTGATAGCGACTCAGATTGACAGCGATTGACAGCGCA
CTCA
D S D
S 750
GACTCAGATAGCGATTGACTCAGACTCAGACTCAGACAGCAATTCAGACTCGGATAGCGACTCAGACTCAGATAGCGA
CTCA
D S D S D S D S D S D S D S D S D S N S D S D S D S D S D S D S D S D S D S D S D S D S D S D S D
S 780
GATTCGGATAGCGACTCAGACTCAGATAGCGATTGACTCAGATTGATAGCGATTGCGACTCAGACAGTGATTGACTCAGACTCAGA
TAGC
D S D
S 810
GACTCAGATTCTGACAGCGATTGACTCAGACTCAGACTCAGACAGTGATTGACTCAGACTCAGACAGCGACTCAGATTGACA
TAGC
D S D
S 840
GACTCAGACTCAGATAGCGACTCAGACTCAGATAGCGACTCAGACTCGGATAGCGATTGACTCAGACTCAGACAGCGACTCAGATTGACA
TAGC
D S D
S 870

W>

GATTCGGACTCAGACAACGACTCAGATTGATAGCGATTGACTCAGATTGATAGCGATTGCGAGTAAACATACTCCGGCTAAACCAATGAGTAC
GGTT
D S D S D N D S D S D S D S D S D S D A G K H T P A K P M S T
V 900

M>

AAAGATCAGCATAAAACAGCTAAAGCATTACCAGAAACAGGTAGTAAAATAATAATTCAAATAATGGCACATTATTCGGTGGATT
ATTC

FIG. 7C

K D Q H K T A K A L P E T G S E N N N S N N G T L F G G L
F 930
GCGGCATTAGGATCATTATTGTCATTCGGTCGTCGTAACAAAATAAA
A A L G S L L S F G R R K K Q N K
947

FIG. 7D

S>
ATGCTAAACAGAGAAAAATAAACGGCAATAACAAGGAAAGGCATGGTATCCAATCGATTAAATAAATTTTCGATTAGAAAGTACAC
AGTG
M L N R E N K T A I T R K G M V S N R L N K F S I R K Y T
V 30

A>
GGAACAGCATCAATTTTAGTAGGTACAACATTAATTTTGGTCTGGGAACCAAGAAGCAAAGGCTGCAGAAAGTACTAATAAAGA
ATTG
G T A S I L V G T T L I F G L G N Q E A K A A E S T N K E
L 60
AACGAAGCGACAACCTTCAGCAAGTGATAATCAATCGAGTGATAAAGTTGATATGCAGCAACTAAATCAAGAAGACAATACTAAAA
TGAT
N E A T T S A S D N Q S S D K V D M Q Q L N Q E D N T K N
D 90
AATCAAAAAGAAATGGTATCATCTCAAGCTAATGAAACGACTTCAAATGGGAATAAATTAATAGAAAAAGAAAGTGTACAATCTAG
CACT
N Q K E M V S S Q G N E T T S N G N K L I E K E S V Q S T
T 120
GGAAATAAAGTTGAAGTTCAACTGCCAAATCAGATGAGCAAGCTTCACCAAAATCTACGAATGAAGATTTAAACACTAAACAAAC
TATA
G N K V E V S T A K S D E Q A S P K S T N E D L N T K Q T
I 150
AGTAATCAAGAAGCGTTACAACCTGATTTGCAAGAGAATAAATCAGTGGTAAATGTTCAACCAACTAATGAGGAAAAACAAAAGGT
AGAT
S N Q E A L Q P D L Q E N K S V V N V Q P T N E E N K K V
D 180
GCCAAAACCTGAATCACTACATTAATGTTAAAAGTGATGCTATCAAGAGTAATGATGAACTCTTGTTGATAACAATAGTAATC
AAAT
A K T E S T T L N V K S D A I K S N D E T L V D N N S N S
N 210
AATGAAAATAATGCAGATATCATTTTCCAAAAAGTACAGCACCTAAACGTTTGAATACAAGAATGCGTATAGCAGCAGTACAGCC
ATCA
N E N N A D I I L P K S T A P K R L N T R M R I A A V Q P
S 240
TCAACAGAGGCTAAAAATGTTAATGATTTAATCACATCAAATACAACATTAACCTGCTGTTGATGCAGATAAAAACAATAAAATCGT
ACCA
S T E A K N V N D L I T S N T T L T V V D A D K N N K I V
P 270
GCCCAAGATTATTATCATTAATAACACAAATTACAGTTGATGACAAAGTTAAATCAGGTGATTATTTACAATTAATACTCAGA
TACA
A Q D Y L S L K S Q I T V D D K V K S G D Y F T I K Y S D
T 300
GTACAAGTATATGGATTGAATCCGGAAGATATTAATAATTTGGTATATTAAGATCCAAATAATGGTGAACAATTCGGACTGC

FIG. 8A

AAAA
V Q V Y G L N P E D I K N I G D I K D P N N G E T I A T A
K 330
CATGATACTGCAAAATAATTTAATTACATATACATTTACAGATTATGTTGATCGATTTAATCTGTACAAATGGGAATTAATTATTC
AATT
H D T A N N L I T Y T F T D Y V D R F N S V Q M G I N Y S
I 360
TATATGGATGCTGATACAATTCCTGTTAGTAAAAACGATGTTGAGTTAATGTTACGATAGGTAATACTACAACAAAAACAACCTGC
TAAC
Y M D A D T I P V S K N D V E F N V T I G N T T T K T T A
N 390
ATTCAATATCCAGATTATGTTGTAATGAGAAAAATCAATTGGATCAGCGTTCCTGAAACAGTTTCACATGTTGGAATAAAGA
AAAT
I Q Y P D Y V V N E K N S I G S A F T E T V S H V G N K E
N 420
CCAGGGTACTATAAACAAACGATTTATGTAATCCATCGGAAAATCTTTAACAAATGCCAACTAAAAGTTCAAGCTTACCACCTC
AAGT
P G Y Y K Q T I Y V N P S E N S L T N A K L K V Q A Y H S
S 450
TATCCTAATAATATCGGGCAAATAAATAAGATGTAACAGATATAAAAAATATCAAGTTCCTAAAGGTTATACATTAATAAAGG
ATAC
Y P N N I G Q I N K D V T D I K I Y Q V P K G Y T L N K G
Y 480
GATGTGAATACTAAAGAGCTTACAGATGTAACAAATCAATACTTGCAGAAAATTACATATGGCGACAACAATAGCGCTGTTATTGA
TTTT
D V N T K E L T D V T N Q Y L Q K I T Y G D N N S A V I D
F 510
GGAAATGCAGATTCTGCTTATGTTGTAATGGTTAATACAAAATCCAATATACAAATAGCGAAAGCCCAACTTGTCAAATGGC
TACT
G N A D S A Y V V M V N T K F Q Y T N S E S P T L V Q M A
T 540

B1>

TTATCTTCAACAGGTAATAAATCCGTTTCTACTGGCAATGCTTTAGGATTTACTAATAACCAAAGTGGCGAGCTGGTCAAGAAGT
ATAT
L S S T G N K S V S T G N A L G F T N N Q S G G A G Q E V
Y 570
AAAATTGGTAACTACGTATGGGAAGATACTAATAAAAACGGTGTCAAGAATTAGGAGAAAAAGCGTTGGCAATGTAAGTGTAACT
TGTA
K I G N Y V W E D T N K N G V Q E L G E K G V G N V T V T
V 600
TTTGATAATAATACAAATACAAAAGTAGGAGAAGCAGTTACTAAGAAGATGGGTCACTTGTTCCTAACTTACCTAATGGAGA
TTAC
F D N N T N T K V G E A V T K E D G S Y L I P N L P N G D

FIG. 8B

Y 630

CGTGTAGAATTTTCAAACCTACCAAAGGTTATGAAGTAACCCCTTCAAACAAGGTAATAACGAAGAATTAGATTCAAACCGCTT
ATCTR V E F S N L P K C Y E V T P S K Q G N N E E L D S N G L
S 660

B2>

TCAGTTATTACAGTTAATGGCAAAGATAACTTATCTGCAGACTTAGGTATTTACAAACCTAAATACAACCTTAGGTGACTATGTCTG
GGAAS V I T V N G K D N L S A D L G I Y K P K Y N L G D Y V W
E 690GATACAAATAAAAATGGTATCCAAGACCAAGATGAAAAAGGTATATCTGGCGTAACGGTAACATTAAGATGAAAACGGTAACGT
GTTAD T N K N G I Q D Q D E K G I S G V T V T L K D E N G N V
L 720AAAACAGTTACAACAGACGCTGATGGCAAATATAAATTTACTGATTTAGATAATGGTAATTATAAAGTTGAATTTACTACACCAGA
AGGCK T V T T D A D G K Y K F T D L D N G N Y K V E F T T P E
G 750TATACACCGACTACAGTAACATCTGGTACCGACATTGAAAAAGACTCTAATGGTTAACAACAACAGGTGTTAATGGTCTGA
TAACY T P T T V T S G S D I E K D S N G L T T T G V I N G A D
N 780

B3>

ATGACATTAGATAGTGGATTCTACAAAACACCAAAATATAATTTAGGTAATTATGTATGGGAAGATACAAATAAAGATGGTAAGCA
GGATM T L D S G F Y K T P K Y N L G N Y V W E D T N K D G K Q
D 810TCAACTGAAAAAGGTATTTTCAGGCGTAACAGTTACATTGAAAAATGAAAACGGTCAAGTTTTACAAACAATAAACAGATAAAGA
TGGTS T E K G I S G V T V T L K N E N G E V L Q T T K T D K D
G 840AAATATCAATTTACTGGATTAGAAAATGCAACTTATAAAGTTGAATTCGAAACCCATCAGGTTACACCAACACAAGTAGGTTTC
AGGAK Y Q F T G L E N G T Y K V E F E T P S G Y T P T Q V G S
G 870ACTGATGAAGGTATAGATTCAAATGGTACATCAACAACAGGTGTCATTAAAGATAAAGATAACGATACTATTGACTCTGGTTTCTA
CAAAT D E G I D S N G T S T T G V I K D K D N D T I D S G F Y
K 900

B4>

CCGACTTACAACCTAGGTGACTATGTATGGGAAGATACAAATAAAAACGGTGTTCAGATAAAGATGAAAAGGGCATTTCAGGTGT
AACAP T Y N L G D Y V W E D T N K N G V Q D K D E K G I S G V**FIG. 8C**

T 930
 GTTACGTTAAAAGATGAAAACGACAAAGTTTTAAAAACAGTTACAACAGATGAAAATGGTAAATATCAATTCACCTGATTTAAACAA
 TGGA
 V T L K D E N D K V L K T V T T D E N G K Y Q F T D L N N
 G 960
 ACTTATAAAGTTGAATTCGAGACACCATCAGGTTATACACCAACTTCAGTAACCTCTGGAAATGATACTGAAAAAGATTCTAATGG
 TTTA
 T Y K V E F E T P S G Y T P T S V T S G N D T E K D S N G
 L 990

B5>

ACAACAACAGGTGTCATTAAGATGCAGATAACATGACATTAGACAGTGGTTTTCTATAAAACACCAAATATAGTTAGGTGATTA
 TGTT
 T T T G V I K D A D N M T L D S G F Y K T P K Y S L G D Y
 V 1020
 TGTTACGACAGTAATAAGACGGCAAACAAGATTCAACTGAAAAAGGTATCAAAGATGTTAAAGTTACTTTATTAATGAAAAAGG
 CGAA
 W Y D S N K D G K O D S T E K G I K D V K V T L L N E K G
 E 1050
 GTAATTGGAACAACATAAACAGATGAAAATGGTAAACTGCTTTGATAATTTAGATACCGGTAATACAAAGTTATTTTTGAAAA
 GCCT
 V I G T T K T D E N G K Y C F D N L D S G K Y K V I F E K
 P 1080
 GCTGGCTTAACACAAAACAGGTACAATACAACCTGAAGATGATAAAGATGCAGATGGTGGCGAAGTTGACGTAACAATTACGGATCA
 TGAT
 A G L T Q T G T N T T E D D K D A D G G E V D V T I T D H
 D 1110
 R>
 GATTCACACTTGATAATGGCTACTACGAAGAAGAAACATCAGATAGCGACTCAGATTCGGACAGCGACTCAGATTCAGACAGAGA
 CTCA
 D F T L D N G Y Y E E E T S D S D S D S D S D S D S D R D
 S 1140
 GACTCAGATAGTGATTCAGACTCGGATAGCGATTCAGATTCAGACAGCGATTCAGATTCAGATAGCGATTCAGATTCAGACAGAGA
 CTCA
 D S D S D S D S D S D S D S D S D S D S D S D S D S D R D
 S 1170
 GATAGTGATTCAGACTCAGATAGCGACTCAGATTCAGACAGCGACTCAGATTCAGACAGCGACTCAGACTCAGATAGTGATTCAGA
 CTCA
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S D
 S 1200
 GATAGCGACTCAGATTCGGATAGCGACTCAGATTCAGACAGCGACTCAGACTCGGATAGTGATTCAGACTCAGATAGCGACTCAGA
 CTCA
 D S D S D S D S D S D S D S D S D S D S D S D S D S D S D
 S 1230

FIG. 8D

W>
GATAGCGATTCAGATTCAGATAGCGACTCAGACTCAGACAGCGATTCAGACTCAGACAGCGACTCAGACTCAGATGCAGGTAAGCA
CACA
D S D S D S D S D S D S D S D S D S D S D S D S D A G K H
T 1260
CCTGTAAACCAATGAGTACTACTAAAGACCATCACAATAAAGCAAAAGCATTACCAGAAACAGGTAATGAAAATAGCGGCTCAAA
TAAC
P V K P M S T T K D H H N K A K A L P E T G N E N S G S N
N 1290
M>
GCAACGTTATTTGGCGGATTATTCGCAGCATTAGGATCATTATTGTTATTCGGTCGTCGTAACAAAACAAAATAAA
A T L F G G L F A A L G S L L L F G R R K K Q N K
1315

FIG. 8E

S>
ATGATTAACAGGGATAATAAAAAGGCAATAACAAAAAGGGTATGATTTCAAATCGCTTAAACAAATTTTCGATTAGAAAAGTATAC
TGTA
M I N R D N K K A I T K K G M I S N R L N K F S I R K Y T
V 30

A>
GGAAGTCATCGATTTTAGTAGGTACGACATTGATTTTTGGTCTAGGGAACCAAGAAGCTAAAGCTGCTGAAAACACTAGTACAGA
AAAT
G T A S I L V G T T L I F G L G N Q E A K A A E N T S T E
N 60
GCAAAACAAGATGATGCAACGACTAGTGATAATAAAGAAGTAGTGTGCGAAACTGAAAATAATTCGACAACAGAAAATAATTC AAC
AAAT
A K Q D D A T T S D N K E V V S E T E N N S T T E N N S T
N 90
CCAATTAAGAAAGAAACAATACTGATTCACAACCAGAAGCTAAAAAGAATCAACTTCATCAAGTACTCAAAAACAGCAAAATAA
CGTT
P I K K E T N T D S Q P E A K K E S T S S S T Q K Q Q N N
V 120
ACAGCTACAACGAAACTAAGCCTCAAAACATTGAAAAAGAAAATGTTAAACCTTCAACTGATAAAACTGCGACAGAAGATACATC
TGTT
T A T T E T K P Q N I E K E N V K P S T D K T A T E D T S
V 150
ATTTTAGAAGAGAAGAAAGCACC AAATAATACAAATAACGATGTAAC TACAAAACCATCTACAAGTGAACCATCTACAAGTGAAT
TCAA
I L E E K K A P N N T N N D V T T K P S T S E P S T S E I
Q 180
ACAAAACCAACTACACCTCAAGAATCTACAAATATTGAAAATTCACAACCGCAACCAACGCCTTCAAAGTAGACAATCAAGTTAC
AGAT
T K P T T P Q E S T N I E N S Q P Q P T P S K V D N Q V T
D 210
GCAACTAATCCAAAAGAACCAGTAAATGTGTCAAAAGAAGAACTTAAAATAATCCTGAGAAATTAAGAATTGGTTAGAAATGA
TAGC
A T N P K E P V N V S K E E L K N N P E K L K E L V R N D
S 240
AATACAGATCATTCAACTAAACCAGTTGCTACAGCTCCAACAAGTGTGCACCAAAACGTGTAACGCAAAAATGCGCTTTGCAGT
TGCA
N T D H S T K P V A T A P T S V A P K R V N A K M R F A V
A 270
CAACCAGCAGCAGTTGCTTCAAACAATGTAATGATTTAATTAAGTACGGAAGCAACCAATCAAACTTGGCGATGGTAAAGATAA
TGTC
Q P A A V A S N N V N D L I K V T K Q T I K V G D G K D N
V 300
GCAGCAGCGCATGACGGTAAAGATATTGAATATGATACAGAGTTTACAATTGACAATAAAGTCaAAAAAGCGGATACAATGACGAT

FIG. 9A

TAAT
A A A H D G K D I E Y D T E F T I D N K V K K G D T M T I
N 330
TATGATAAGAATGTAATTCCTTCGGATTTAACAGATAAAAATGATCCTATCGATATTACTGATCCATCAGGAGAGGTCATTGCTAA
AGGA
Y D K N V I P S D L T D K N D P I D I T D P S G E V I A K
G 360
ACATTTGATAAAGCAACTAAGCAAATCACATATACATTTACAGACTATGTAGATAAATATGAAGATATAAAAATCACGCTTAACTCT
ATAT
T F D K A T K Q I T Y T E T D Y V D K Y E D I K S R L T L
Y 390
TCGTATATTGATAAAAAACAGTTCCAAATGAGACAAGTTGAATTTAACATTTGCTACAGCAGGTAAGAAACAAGCCAAAATGT
CACT
S Y I D K K T V P N E T S L N L T F A T A G K E T S Q N V
T 420
GTTGATTATCAAGATCCAATGGTCCATGGTGATTCAACATTCATCTATCTTTACAAAATTAGATGAAGATAAGCAAATATTGA
ACAA
V D Y Q D P M V H G D S N I Q S I F T K L D E D K Q T I E
Q 450
CAAATTTATGTTAACCCATTGAAAAATCAGCAACCAACTAAAGTTGATATAGCTGGTAGTCAAGTAGATGATTATGGAAATAT
TAAA
Q I Y V N P L K K S A T N T K V D I A G S Q V D D Y G N I
K 480
CTAGGAAATGGTAGCACCATTATTGACCAAAATACAGAAATAAAGTTTATAAAGTTAACTCTGATCAACAATTGCCTCAAAGTAA
TAGA
L G N G S T I I D Q N T E I K V Y K V N S D Q Q L P Q S N
R 510
ATCTATGATTTTAGTCAATACGAAGATGTAACAAGTCAATTTGATAATAAAAAATCATTAGTAATAATGTAGCAACATGGATT
TGGT
I Y D F S Q Y E D V T S Q F D N K K S F S N N V A T L D F
G 540
GATATTAATTCAGCCTATATTATCAAAGTTGTTAGTAAATATACACCTACATCAGATGGCGAACTAGATATTGCCAAGGTACTAG
TATG
D I N S A Y I I K V V S K Y T P T S D G E L D I A Q G T S
M 570
AGAACAACCTGATAAATATGGTTATTATAATTATGCAGGATATTCAAACCTCATCGTAACTTCTAATGACACTGGCGGTGGCGACGG
TACT
R T T D K Y G Y Y N Y A G Y S N F I V T S N D T G G G D G
T 600

B1>

GTAAACCTGAAGAAAAGTTATACAAAATGGTGACTATGTATGGGAAGACGTTGATAAAGACGGTCTTCAAGGTACAGATTCAAA
AGAA
V K P E E K L Y K I G D Y V W E D V D K D G V Q G T D S K

FIG. 9B

E 630
AAACCAATGCCAAACGTTTTAGTTACATTAACCTACCCGGACGGTACTACAAAATCAGTAAGAACAGATGCTAATGGTCATTATGA
ATTC
K P M A N V L V T L T Y P D G T T K S V R T D A N G H Y E
F 660
GGTGGTTTGAAAGACGGAGAACTTATACAGTTAAATTCGAAACGCCAACTGGATATCTTCCAACAAAAGTAAATGGAACAACCTGA
TGGT
G G L K D G E T Y T V K F E T P T G Y L P T K V N G T T D
G 690
B2>
GAAAAAGACTCAAATGGTAGTTCGGTTACTGTAAAATTAATGGTAAAGATGATATGTCCTTAGATACTGGTTTTTACAAGAACC
TAAA
E K D S N G S S V T V K I N G K D D M S L D T G F Y K E P
K 720
TACAACCTAGGTGACTATGTATGGGAAGATACTAATAAAGATGGTATCCAAGATGCAAAATGAGCCAGGAATCAAAGATGTTAAGGT
TACA
Y N L G D Y V W E D T N K D G I Q D A N E P G I K D V K V
T 750
TTAAAAGATAGTACTGGAAAAGTTATTGGTACAACACTACTGATGCCTCGGGTAAATATAAATTTACAGATTTAGATAATGGTAA
CTAT
L K D S T G K V I G T T T T C A S G K Y K F T D L D N G N
Y 780
ACAGTAGAATTTGAAACACCAGCAGGTTACACGCCAACGGTTAAAAATACTACAGCTGATGATAAAGATTCTAATGGTTTAAACAAC
AACA
T V E F E T P A G Y T P T V K N T T A D D K D S N G L T T
T 810
B3>
GGTGTCAATAAAGATGCAGATAATATGACATTAGACAGGGGTTTCTATAAAAACACCAAAATACAGTTAGGTGATTATGTTGGTA
CGAC
G V I K D A D N M T L D R G F Y K T P K Y S L G D Y V W Y
D 840
AGTAATAAAGACGGCAAACAAGATTCAACTGAAAAAGGTATCAAAGATGTGACAGTTACATTGCAAAACGAAAAAGCCGAAGTAAT
TGGA
S N K D G K Q D S T E K G I K D V T V T L Q N E K G E V I
G 870
ACAACATAAAGACAGATGAAAATGGTAAATATCGTTTCGATAATTTAGATAGCGGTAAATACAAAGTTATTTTTGAAAAGCCTGCTGG
CTTA
T T K T D E N G K Y R F D N L D S G K Y K V I F E K P A G
L 900
ACACAAACAGTTACAAATACAACCTGAAGATGATAAAGATGCAGATGGTGGCGAAGTTGACGTAACAATTACGGATCATGATGATTT
CACA
T Q T V T N T T E D D K D A D G G E V D V T I T D H D D F
T 930

FIG. 9C

R>

CTTGATAACGGATACTCGAAGAAGATACATCAGACAGCGATTTCAGACTCAGATAGTACTCAGACAGCGACTCAGACTCAGACAG
CGAC

L D N G Y F E E D T S D S D S D S D S D S D S D S D S D S
D 960

TCAGACTCAGACAGTGTTCAGATTTCAGACAGCGACTCAGATTTCAGATAGCGACTCAGATTTCGGACAGCGATTTCAGACTCAGATAG
CGAC

S D S D S D S D S D S D S D S D S D S D S D S D S D S D S
D 990

TCAGATTTCAGATAGCGATTTCAGACTCAGACAGCGACTCAGATTTCAGATAGCGATTTCGGACTCAGACAGCGATTTCAGACTCAGATAG
CGAC

S D S D S D S D S D S D S D S D S D S D S D S D S D S D S
D 1020

TCAGACTCAGACAGCGACTCAGATTTCAGATAGCGATTTCGGACTCAGATAGCGACTCAGATTTCAGACAGCGATTTCAGACTCAGATAG
CGAC

S D S D S D S D S D S D S D S D S D S D S D S D S D S D S
D 1050

TCAGATTTCAGACAGCGATTTCAGACTCAGATAGCGACTCAGACTCAGACAGTGTTCAGATTTCAGACAGCGACTCAGACTCAGATAG
CGAC

S D S D S D S D S D S D S D S D S D S D S D S D S D S D S
D 1080

W>

TCAGATTTCAGACAGCGACTCAGACTCAGATAGCGACTCAGACTCAGACAGTGTTCAGACAGCGATTTCAGACTTCGGATGCAGGAAA
ACAT

S D S D S D S D S D S D S D S D S D S D S D S D S D A G K
H 1110

ACACCTGTAAACCAATGAGTACTACTAAAGACCATCACAATAAAGCAAAGCATTACCAGAACAGGTAGTAAAATAACGGCTC
AAAT

T P V K P M S T T K D H H N K A K A L P E T G S E N N G S
N 1140

M>

AACGCAACGTTATTTGGTGGATTATTTGCAGCATTAGGTTTATTGTTATTTCGGTCGTCGCAAAAAACAAAAAAA

N A T L F G G L F A A L G S L L L F G R R K K Q N K
1166

FIG. 9D




FIG. 10

$\%$	ClfB	SdrC	SdrD	SdrE
ClfA	27	20	24	25
ClfB		24	25	23
SdrC			24	23
SdrD				30

FIG. 11

1.			
SdrD	282	TVDDKVKSGDYFTIK	296
SdrE	316	TIDNKVKKGDTMTIN	330
ClfB	246	TVTDKVKSGDYFTAK	269
SdrC	227	TIDDSVKEGDTFFEK	241
ClfA	264	SVPNSAVKGDTEFKIT	278
		:: ... ** : .	
2.			
SdrD	324	ETIATAKHDTANNLITYTFTDYVD	347
SdrE	355	EVIAGTDFDKATKQITYTFTDYVD	378
ClfB	291	DVVAKATYDILTKITYTFVTDYVN	314
SdrC	266	NIIAKGIYDSTTNTTTYTFTNYVD	289
ClfA	302	QVLANGVIDSEGN-VIYFTDYVN	324
		: *.. * : : **:**:	
3.			
SdrD	463	TDIKIYQVPK	472
SdrE	492	TEIKVYKVNS	501
ClfB	429	TKLRIFEVND	438
SdrC	391	--FKIYEVD	398
ClfA	431	TSIKVYKVDN	440
		: : : * .	
4.			
SdrD	489	DVTNQ	493
SdrE	519	DVTSQ	523
ClfB	457	EVTDQ	461
SdrC	415	DVTDQ	419
ClfA	457	DVTNS	461
		: **..	
5.			
SdrCB1	496	KYNLGDYVWEDTNKDGKQ--DANEKGIKGVYVILKDSNGK-ELDRTTDENGKYQFTGLS	552
SdrCB2	607	KYSLGDYVWYDSNRDGR--DSTEKGIKGVKVTLQNEKGE-VIGTTE'DENGGYRFDNLD	663
SdrDB1	569	VYKIGNYVWEDTNKNGVQ--ELGEGVGNVTVTFDNTNTKVGEAVTKEDGSYLIPNLP	626
SdrDB2	681	KYNLGDYVWEDTNKNGIQ--DQDEKGISGVTVTLKDENG- <u>VLKTVTTDADGKYKFTDL</u>	737
SdrDB3	792	KYNLGNVWEDTNKDGKQ--DSTEKGISGVTVTLKNENGE- <u>VLQTTKTDKDGKYQFTGLE</u>	848
SdrDB4	902	TYNLGDYVWEDTNKNGVQ--DKDEKGISGVTVTLKDENDK- <u>VLKTVTTDENGKYQFTDLN</u>	958
SdrDB5	1013	KYSLGDYVWYDSNRDGR--DSTEKGIKDVVTLQNEKGE-VIGTTKTDENGGYCFDNLD	1069
SdrEB1	607	LYKIGDYVWEDVDKDGVOGTDSKEKPMANVLVTLTYPDG--TTKSVRTDANGHYEFGGLK	664
SdrEB2	720	KYNLGDYVWEDTNKNGIQ--DANEKGIKDVVTLKDSGK-VIGTTTTDASGKYFTDL	776
SdrEB3	830	KYSLGDYVWYDSNRDGR--DSTEKGIKDVVTLQNEKGE-VIGTTKTDENGGYRFDNLD	886
		*.:** * :*: : * : * : * : * : * : *	

FIG. 12A

SdrCB1	553	NG-TYSVEFST-PAGYTPPTANVGTDDAVDSGLTTTGVIKDADNMTLDSGFYKTP-	606
SdrCB2	664	SG-KYKVIFEK-PAGLTQTGTNTTEDD-KDADGGEVDVTITDHDDFTLDNGYEEET	717
SdrDB1	627	NG-DYRVEFSNLPKGYEVTPSKQGNNEELDSNGLSSVITVNGKDNLSADLGIYKP--	680
SdrDB2	738	NG-NYKVEFTT-PEGYTPPTVTSKSDIEKDSNGLTTTGVIKADNMTLDSGFYKTP-	791
SdrDB3	849	NG-TYKVEFET-PSGYTPPTQVGSCTDEGIDSNSTTTGVIKDKDNDTIDSGFYKTP--	901
SdrDB4	959	NG-TYKVEFET-PSGYTPPTSVTSKSDIEKDSNGLTTTGVIKADNMTLDSGFYKTP-	1012
SdrDB5	1070	SG-KYKVIFEK-PAGLTQTGTNTTEDD-KDADGGEVDVTITDHDDFTLDNGYEEET	1123
SdrEB1	665	DGETYTVKFET-PTGYLPTKVNGTIDGKDSNGSSVTVKINGKDDMSLDTGFYKEP-	719
SdrEB2	777	NG-NYTVFEFET-PAGYTPPTVKNNTADD-KDSNGLTTTGVIKADNMTLDRGFYKTP-	829
SdrEB3	887	SG-KYKVIFEK-PAGLTQTVTNTTEDD-KDADGGEVDVTITDHDDFTLDNGYFEEDT	940

. * * * * . * * * * : * : * * : .. * : : * * * :

FIG. 12B




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19

Staphylococcus aureus Consensus and Variable Motif

Protein	Motif									
ClfA	I	Y	T	F	T	D	Y	V	N	
ClfB	T	F	V	F	T	D	Y	V	N	
SdrC	T	Y	T	F	T	D	Y	V	D	
SdrD	T	Y	T	F	T	D	Y	V	D	
SdrE	T	Y	T	F	T	D	Y	V	D	
Consensus	T	Y	T	F	T	D	Y	V	D	
Variable Motif	T/I	Y/F	T/V	F	T	D/N	Y	V	D/N	

FIG. 20




FIG. 21




FIG. 22A

FIG. 22B




FIG. 23

1

EXTRACELLULAR MATRIX-BINDING PROTEINS FROM *STAPHYLOCOCCUS AUREUS*

This application claims priority to U.S. provisional appli- 5
cation numbers 60/098,427, filed Aug. 31, 1998 and 60/066,
815, filed Nov. 26, 1997. The U.S. Government has rights in
this invention arising out of National Institutes of Health
grant number AI20624.

FIELD OF THE INVENTION

The present invention is in the fields of microbiology and 10
molecular biology. The invention includes the isolation and
use of extracellular matrix-binding proteins and genes that
express the proteins from *Staphylococcus aureus* to inhibit,
prevent and diagnose *S. aureus* infection.

BACKGROUND OF THE INVENTION

In hospitalized patients *Staphylococcus aureus* is a major 15
cause of infections associated with indwelling medical
devices, such as catheters and prostheses, and related infec-
tions of surgical wounds. A significant increase in *Staphy-
lococcus aureus* isolates that exhibit resistance to most
known antibiotics has been observed in hospitals throughout 20
the world. The recent emergence of resistance to
vancomycin, the last remaining antibiotic for treating
methicillin-resistant *Staphylococcus aureus* (MRSA)
infections, has emphasized the need for alternative prophyl-
actic or vaccine strategies to reduce the risk of nosocomial 25
S. aureus infections.

Initial localized infections of wounds or indwelling medi- 30
cal devices can lead to serious invasive infections such as
septicemia, osteomyelitis, and endocarditis. In infections
associated with medical devices, plastic and metal surfaces
become coated with host plasma and extracellular matrix 35
proteins such as fibrinogen and fibronectin shortly after
implantation. The ability of *S. aureus* to adhere to these
proteins is of crucial importance for initiating infection.
Vascular grafts, intravenous catheters, artificial heart valves, 40
and cardiac assist devices are thrombogenic and prone to
bacterial colonization. *S. aureus* is the most damaging
pathogen to cause such infections.

Fibrin is the major component of blood clots, and 45
fibrinogen/fibrin is one of the major plasma proteins depos-
ited on implanted biomaterials. Considerable evidence exists
to suggest that bacterial adherence to fibrinogen/fibrin is
important in the initiation of device-related infection. For
example, as shown by Vaudaux et al., *S. aureus* adheres to
in vitro plastic that has been coated with fibrinogen in a
dose-dependent manner (*J. Infect. Dis.* 160:865–875
(1989)). In addition, in a model that mimics a blood clot or
damage to a heart valve, Herrmann et al. demonstrated that 50
S. aureus binds avidly via a fibrinogen bridge to platelets
adhering to surfaces (*J. Infect. Dis.* 167: 312–322 (1993)). *S.
aureus* can adhere directly to fibrinogen in blood clots
formed in vitro, and can adhere to cultured endothelial cells
via fibrinogen deposited from plasma acting as a bridge
(Moreillon et al., *Infect. Immun.* 63:4738–4743 (1995);
Cheung et al., *J. Clin. Invest.* 87:2236–2245 (1991)). As 60
shown by Vaudaux et al. and Moreillon et al., mutants
defective in the fibrinogen-binding protein clumping factor
(ClfA) exhibit reduced adherence to fibrinogen in vitro, to
explanted catheters, to blood clots, and to damaged heart
valves in the rat model for endocarditis (Vaudaux et al., 65
Infect. Immun. 63:585–590 (1995); Moreillon et al., *Infect.
Immun.* 63: 4738–4743 (1995)).

2

An adhesin for fibrinogen, often referred to as “clumping 5
factor,” is located on the surface of *S. aureus* cells. The
interaction between the clumping factor on bacteria and
fibrinogen in solution results in the instantaneous clumping
of bacterial cells. The binding site on fibrinogen is located in
the C-terminus of the gamma chain of the dimeric fibrinogen
glycoprotein. The affinity is very high and clumping occurs
in low concentrations of fibrinogen. Scientists have recently
shown that clumping factor also promotes adherence to solid
phase fibrinogen, to blood clots, and to damaged heart
valves (McDevitt et al., *Mol. Microbiol.* 11: 237–248
(1994); Vaudaux et al., *Infect. Immun.* 63:585–590 (1995);
Moreillon et al., *Infect. Immun.* 63: 4738–4743 (1995)).

The gene for a clumping factor protein, designated ClfA, 15
has been cloned, sequenced and analyzed in detail at the
molecular level (McDevitt et al., *Mol. Microbiol.* 11:
237–248 (1994); McDevitt et al., *Mol. Microbiol.*
16:895–907 (1995)). The predicted protein is composed of
933 amino acids. A signal sequence of 39 residues occurs at
the N-terminus followed by a 520 residue region (region A),
which contains the fibrinogen binding domain. A 308 resi-
due region (region R), composed of 154 repeats of the
dipeptide serine-aspartate, follows. The R region sequence is
encoded by the 18 basepair repeat GAYTCNGAYT 20
CNGAYAGY (SEQ ID NO: 9) in which Y equals pyrim-
idines and N equals any base. The C-terminus of ClfA has
features present in many surface proteins of Gram-positive
bacteria such as an LPDTG (SEQ ID NO: 10) motif, which
is responsible for anchoring the protein to the cell wall, a
membrane anchor, and positive charged residues at the
extreme C-terminus.

The platelet integrin alpha IIB β 3 recognizes the 25
C-terminus of the gamma chain of fibrinogen. This is a
crucial event in the initiation of blood clotting during
coagulation. ClfA and alpha IIB β 3 appear to recognize
precisely the same sites on fibrinogen gamma chain because
ClfA can block platelet aggregation, and a peptide corre-
sponding to the C-terminus of the gamma chain (198–411)
can block both the integrin and ClfA interacting with
fibrinogen (McDevitt et al., *Eur. J. Biochem.* 247:416–424
(1997)). The fibrinogen binding site of alpha IIB β 3 is close
to, or overlaps, a Ca²⁺ binding determinant referred to as an
“EF hand”. ClfA region A carries several EF hand-like
motifs. A concentration of Ca²⁺ in the range of 3–5 mM
blocks these ClfA-fibrinogen interactions and changes the
secondary structure of the ClfA protein. Mutations affecting
the ClfA EF hand reduce or prevent interactions with
fibrinogen. Ca²⁺ and the fibrinogen gamma chain seem to
bind to the same, or to overlapping, sites in ClfA region A.

The alpha chain of the leucocyte integrin, alpha M β 2, has 30
an insertion of 200 amino acids (A or I domain) which is
responsible for ligand binding activities. A novel metal
ion-dependent adhesion site (MIDAS) motif in the I domain
is required for ligand binding. Among the ligands recog-
nized is fibrinogen. The binding site on fibrinogen is in the
gamma chain (residues 190–202). It was recently reported
that *Candida albicans* has a surface protein, alpha Int1p,
having properties reminiscent of eukaryotic integrins. The
surface protein has amino acid sequence homology with the
I domain of alpha M β 2, including the MIDAS motif.
Furthermore, alpha Int1p binds to fibrinogen.

ClfA region A also exhibits some degree of sequence 35
homology with alpha Int1p. Examination of the ClfA region
A sequence has revealed a potential MIDAS motif. Mutations
in supposed cation coordinating residues in the
DXSXS (SEQ ID NO: 13) portion of the MIDAS motif in
ClfA results in a significant reduction in fibrinogen binding.

A peptide corresponding to the gamma-chain binding site for alpha M β 2 (190–202) has been shown by O'Connell et al. to inhibit ClfA-fibrinogen interactions (O'Connell et al., *J. Biol. Chem.*, in press). Thus it appears that ClfA can bind to the gamma-chain of fibrinogen at two separate sites. The ligand binding sites on the ClfA are similar to those employed by eukaryotic integrins and involve divalent cation binding EF-hand and MIDAS motifs.

Scientists have recently shown that *S. aureus* expresses proteins other than ClfA that may bind fibrinogen (Boden and Flock, *Mol. Microbiol.* 12:599–606 (1994)). One of these proteins is probably the same as the broad spectrum ligand-binding protein reported by Homonylo-McGavin et al., *Infect. Immun.* 61:2479–2485 (1993). Another is coagulase, as reported by Boden and Flock, *Infect. Immun.* 57:2358–2363 (1989), a predominantly extracellular protein that activates the plasma clotting activity of prothrombin. Coagulase binds prothrombin at its N-terminus and also interacts with soluble fibrinogen at its C-terminus. Cheung et al., *Infect. Immun.* 63:1914–1920 (1995) have described a variant of coagulase that binds fibrinogen. There is some evidence that coagulase can contribute, in a minor way, to the ability of *S. aureus* cells to bind fibrinogen. As shown by Wolz et al., *Infect. Immun.* 64:3142–3147 (1996), in an agr regulatory mutant, where coagulase is expressed at a high level, coagulase appears to contribute to the binding of soluble fibrinogen to bacterial cells. Also, as shown by Dickinson et al., *Infect. Immun.* 63:3143–3150 (1995), coagulase contributes in a minor way to the attachment of *S. aureus* to plasma-coated surfaces under flow. However, it is clear that clumping factor ClfA is the major surface-located fibrinogen-binding protein responsible for bacterial attachment to immobilized fibrinogen/fibrin.

The identification and isolation of additional *S. aureus* extracellular matrix binding proteins would be useful for the development of therapies, diagnosis, prevention strategies and research tools for *S. aureus* infection.

Accordingly it is an object of the present invention to provide isolated cell-wall associated extracellular matrix-binding proteins of *S. aureus* and active fragments thereof.

It is a further object of the invention to provide methods for preventing, diagnosing, treating or monitoring the progress of therapy for bacterial infections caused by *S. aureus*.

It is a further object of the present invention to provide isolated *S. aureus* surface proteins that are related in amino acid sequence to ClfA and are able to promote adhesion to the extracellular matrix or host cells.

It is another object of the present invention to generate antisera and antibodies to cell-wall associated extracellular matrix-binding proteins of *S. aureus*, or active fragments thereof.

It is a further object of the present invention to provide *S. aureus* vaccines, including a DNA vaccine.

It is a further object of the present invention to provide improved materials and methods for detecting and differentiating *S. aureus* organisms in clinical and laboratory settings.

It is a further object of the invention to provide nucleic acid probes and primers specific for *S. aureus*.

It is a further object of the invention to provide isolated extracellular matrix-binding proteins or peptides of *S. aureus*.

SUMMARY OF THE INVENTION

Isolated extracellular matrix-binding proteins, designated ClfB, SdrC, SdrD and SdrE, and their corresponding amino

acid and nucleic acid sequences and motifs are described. The proteins, peptides, fragments thereof or antigenic portions thereof are useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection and as scientific research tools. Further, antibodies or antibody fragments to the proteins, peptides, fragments thereof or antigenic portions thereof are also useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection. The proteins, peptides, peptide fragments, antibodies, or antibody fragments can be administered in an effective amount to a patient in need thereof in any appropriate manner, preferably intravenously or otherwise by injection, to impart active or passive immunity. In an alternative embodiment, the proteins or antibodies thereof can be administered to wounds or used to coat biomaterials to act as blocking agents to prevent or inhibit the binding of *S. aureus* to wounds or biomaterials.

Specifically, extracellular matrix-binding proteins from *S. aureus* designated as ClfB, SdrC, SdrD, and SdrE are provided.

ClfB is a fibrinogen binding protein. The nucleic acid and amino acid sequences of ClfB are provided in FIG. 5. The amino acid sequence of ClfB is SEQ ID NO:1, and the nucleic acid sequence of ClfB is SEQ ID NO:2.

SdrC has been discovered to bind to several extracellular matrix proteins of the host, including for example, bone sialoprotein (BSP), decorin, plasmin, fibrinogen and vitronectin. The amino acid and nucleic acid sequences of SdrC are SEQ ID NOS:3 and 4 respectively and are provided in FIG. 7.

Another of the discovered proteins, SdrD, binds at least vitronectin. The amino acid and nucleic acid sequences of SdrD are SEQ ID NOS:5 and 6 respectively and are provided in FIG. 8.

SdrE binds to extracellular matrix proteins, for example, bone sialoprotein (BSP). The amino acid and nucleic acid sequences of SdrE are SEQ ID NOS:7 and 8 respectively and are provided in FIG. 9.

ClfB has a predicted molecular weight of approximately 88 kDa and an apparent molecular weight of approximately 124 kDa. ClfB is a cell-wall associated protein and binds both soluble and immobilized fibrinogen. In addition, ClfB binds both the alpha and beta chains of fibrinogen and acts as a clumping factor. SdrC, SdrD and SdrE are cell-wall associated proteins that exhibit cation-dependent ligand binding of extracellular matrix proteins such as decorin, plasmin, fibrinogen, vitronectin and BSP.

It has been discovered that in the A region of SdrC, SdrD, SdrE, ClfA, and ClfB, there is a highly conserved amino acid sequence that can be used to derive a consensus TYTFTDYVD (SEQ ID NO: 18) motif (See FIG. 20). The motif can be used in vaccines to impart broad spectrum immunity against bacterial infections. The motif can also be used as an antigen in the production of monoclonal or polyclonal antibodies to impart broad spectrum passive immunity. In an alternative embodiment, any combination of the variable sequence motif (T/I) (Y/F) (TN) (F) (T) (D/N) (Y) (V) (D/N) (SEQ ID NO: 23) can be used as an immunogen or antigen, or in the preparation of antibodies.

The ClfB, SdrC, SdrD and SdrE proteins or the consensus or variable motifs thereof are useful as scientific research tools to identify *S. aureus* binding sites on the extracellular matrix. They are further useful as research tools to promote an understanding of the mechanisms of bacterial pathology and the development of antibacterial therapies.

The ClfB, SdrC, SdrD and SdrE nucleic acid sequences or selected fragments thereof, including the sequences encod-

ing the consensus or variable motifs, are useful as nucleic acid probes for the identification of other *S. aureus* extracellular matrix-binding proteins. Alternatively, the amino acid sequences of the proteins, or selected fragments thereof, can be used as probes to identify the corresponding nucleic acid sequences.

The ClfB, SdrC, SdrD and SdrE nucleic acid sequences or the sequences encoding the consensus or variable motifs are further useful as polynucleotides which comprise contiguous nucleic acid sequences capable of being expressed. The nucleic acid sequences may be inserted into a vector and placed in a microorganism for the production of recombinant ClfB, SdrC, SdrD and SdrE proteins or the variable or consensus amino acid motifs. This allows for the production of the gene product upon introduction of said polynucleotide into eukaryotic tissues *in vivo*. The encoded gene product preferably either acts as an immunostimulant or as an antigen capable of generating an immune response. Thus, the nucleic acid sequences in this embodiment encode an MSCRAMM (Microbial Surface Components Recognising Adhesive Matrix Molecules) immunogenic epitope, and optionally a cytokine or a T-cell costimulatory element, such as a member of the B7 family of proteins.

There are several advantages of immunization with a gene rather than its gene product. The first is the relative simplicity with which native or nearly native antigen can be presented to the immune system. A second advantage of DNA immunization is the potential for the immunogen to enter the MHC class I pathway and evoke a cytotoxic T cell response. Cell-mediated immunity is important in controlling infection. Since DNA immunization can evoke both humoral and cell-mediated immune responses, its greatest advantage may be that it provides a relatively simple method to survey a large number of *S. aureus* genes for their vaccine potential.

Antibodies immunoreactive with ClfB, SdrC, SdrD and SdrE proteins, or their active fragments, including with the consensus or variable amino acid motifs, are provided herein. Vaccines or other pharmaceutical compositions containing the proteins or amino acid motifs are additionally provided herein.

Antibodies and antisera to the consensus TYTFTDYVD (SEQ ID NO: 18) sequence epitope or the variable (T/I) (Y/F) (TtV) (F) (T) (D/N) (Y) (V) (D/N) (SEQ ID NO: 23) sequence, specifically TYTFTNYVD (SEQ ID NO: 19) in SdrC, TYTFTDYVD (SEQ ID NO: 18) in SdrD and SdrE, TFFVFTDYVN (SEQ ID NO: 20) in ClfB or IYTFTDYVN (SEQ ID NO: 21) in ClfA are provided herein. Vaccines or other pharmaceutical compositions containing the epitopes are also provided herein.

In addition, diagnostic kits containing nucleic acid molecules, the proteins, antibodies or antisera to ClfB, SdrC, SdrD, SdrE or their active fragments, including the consensus or variable amino acid motifs and the appropriate reagents for reaction with a sample are also provided.

In one embodiment of the invention, the diagnostic kit is used to identify patients or animals that have levels of antibodies to ClfB ClfB, SdrC, SdrD, or SdrE that are above a population norm. The plasma of the patients or animals can be obtained, processed, and administered to a host in need of passive immunity to *S. aureus* infection.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a schematic representation comparing features of unprocessed ClfA and ClfB proteins. S indicates the signal sequence. A indicates the conserved region (region A). P

indicates the proline-rich region (repeats are indicated by gray boxes). R indicates the SD repeat region (region R). W indicates the wall-spanning region. M indicates the membrane spanning and anchoring regions. EF hand I of ClfA and its partial homologue on ClfB are indicated by black vertical bars. The MIDAS motifs are indicated by hatched (DXSXS) (SEQ ID NO: 13) and narrow vertical lines (downstream T and D residues) connected by dashed lines.

FIG. 2 is a schematic representation of general plasmid and probe constructions for sequencing *clfB*. A repeat-carrying EcoRI fragment was cloned from phage clone A1-1 into pGEM 7Z (f)+ to give pA1-1E (top), and subsequently reduced by deletion of an XbaI fragment to give pA1-1EX, which contains the entire *clfB* gene. A SmaI fragment containing *clfB* and 500 bp of upstream DNA was cloned into pCU1 for overexpression and complementation work (pA1-1EA). The HpaI probe used to screen mutants, and the hybridizing BamHI fragment are also indicated.

FIG. 3 is a schematic representation showing construction of a cassette for allele replacement. *clfB* was interrupted by blunt-end cloning the Tc determinant from pT181 into the HpaI site in the middle of the gene in pA1-1EX. pTS2 was then cloned into the SmaI site of the cassette to enable temperature sensitive propagation in *S. aureus*.

FIG. 4 is a schematic representation of a physical map of the *sdrC sdrD sdrE* locus in *S. aureus* strain Newman. The extents of the plasmid clones are delineated. A6-2 is a LambdaGEM@-12 clone. pEJ1, pEJ2 and pEJ3 are A6-2 fragments subcloned in the pGEM 7Z (f)+ (pEJ1 and pEJ2) and the pBluescript KS+ vector (pEJ3). pC1 is a HindIII fragment directly cloned from strain Newman in the pBluescript KS+ vector. Arrows indicate the direction of transcription of *sdrC*, *sdrD* and *sdrE*.

FIG. 5 is the nucleic acid sequence of *clfB* and flanking DNA, and amino acid translation of the ORF. The likely start codon is double underlined, and the principal regions indicated using the abbreviations of FIG. 1. Two salient features of region A, the DYSNS (SEQ ID NO: 11) of the putative MIDAS motif, and the sequence FTDYVN (SEQ ID NO: 12), the longest region of identity with ClfB, are underlined. Vertical bars indicate the repeats in the proline-rich region. An inverted repeat specifying a possible transcription termination signal is underlined.

FIG. 6 is an amino acid sequence alignment of part of region A of the ClfB and ClfA proteins in the region of strongest similarity. EF hand I of ClfA is underlined. Identical residues are denoted by an asterisk; conservative substitutions are denoted by a period. The DXSXS (SEQ ID NO: 13) portion of the MIDAS motif of ClfB is double underlined.

FIG. 7 is the nucleic acid sequence and amino acid translation of the *sdrC* gene. The consensus TYTFTDYVD (SEQ ID NO: 18) motif, expressed in SdrC as TYTFTNYVD (SEQ ID NO: 19), the EF hands in the B repeats, and the LPXTG (SEQ ID NO: 14) motif are underlined. Major regions, such as the signal sequence (S), region A (A), B repeats (B) region R (R), the wall-spanning domain (W), and the membrane-anchoring domain (M), are indicated.

FIG. 8 is the nucleic acid sequence and amino acid translation of the *sdrD* gene. The consensus TYTFTDYVD (SEQ ID NO: 18) motif, the EF hands in the B repeats, and the LPXTG (SEQ ID NO: 14) motif are underlined. Major regions, such as the signal sequence (S), region A (A), B repeats (B) region R (R), the wall-spanning domain (W), and the membrane-anchoring domain (M), are indicated.

FIG. 9 is the nucleic acid sequence and amino acid translation of the *sdrE* gene. The consensus TYTFTDYVD

(SEQ ID NO: 18) motif, the EF hands in the B repeats, and the LPXTG (SEQ ID NO: 14) motif are underlined. Major regions, such as the signal sequence (S), region A (A), B repeats (B) region R (R), the wall-spanning domain (W), and the membrane-anchoring domain (M), are indicated.

FIG. 10 is a schematic diagram of the region R-containing proteins. Numerals over the proteins denote numbers of amino acids in the regions, numerals under the proteins denote the location on the amino acid sequence of the motifs counted from the beginning of the signal peptide. Abbreviations: S: Signal peptide; A: Region A; B: B repeat; R: Region R; W.M: Wall and membrane spanning regions.

FIG. 11 is a chart showing similarities between A regions ClfA, ClfB, SdrC, SdrD and SdrE. Each sequence was aligned in pairwise combinations and the percent identical residues given.

FIG. 12 indicates Clustal™ multiple sequence alignments of areas of similarity of the A and B regions of the region R containing genes of strain Newman. An asterisk denotes identity of amino acids, and a colon represents increasing similarity of polarity and hydrophobicity/hydrophilicity of side chains of amino acids. Alignments 1–4 show areas from region A. Alignments 1, 3 and 4 show the common motifs. Alignment 2 shows homology in the vicinity of the ClfA EF-hand (underlined), with the consensus TYTFTDYVD (SEQ ID NO: 18) sequence conserved in all five genes. Alignment 5 shows the B repeats of proteins SdrD, SdrD and SdrE with possible EF hands underlined.

FIG. 13 is a time-course graph of ClfB expression in *S. aureus* Newman versus time, monitored by Western blotting. Shake flask cultures were sampled at specific time intervals. A standard number of cells was used to prepare lysates.

FIG. 14 is a graph of absorbance versus concentration of ClfA/ClfB comparing the binding of increasing concentrations of biotinylated recombinant region A from ClfA and ClfB to fibrinogen coated plates. Binding to BSA-coated plates is shown as a control. The closed square symbol represents fibrinogen-ClfA; the closed circle symbol represents fibrinogen-ClfB; the open square symbol represents BSA-ClfA; the open circle symbol represents BSA-ClfB.

FIG. 15 is a graph of cells bound versus fibrinogen concentration showing adherence of *S. aureus* Newman and mutants to fibrinogen immobilized on ELISA plates. Increasing amounts of fibrinogen were used to coat the plates, and a fixed concentration of cells from exponential phase cultures were added. The square symbol represents wild-type; the diamond symbol represents clfA; the circle symbol represents clfB; the triangle symbol represents clfA-clfB; the x symbol represents clfAclfB,clfB*.

FIG. 16 is a graph of cells bound versus fibrinogen concentration showing adherence of *S. aureus* Newman and mutants to fibrinogen immobilized on ELISA plates. Increasing amounts of fibrinogen were used to coat the plates, and a fixed concentration of cells from stationary phase cultures added. The square symbol represents wild-type; the diamond symbol represents clfA; the circle symbol represents clfB; the triangle symbol represents clfAclfB; the x symbol represents clfAclfB,clfB*.

FIG. 17 is a graph of cells bound versus IgG concentration showing effects of preincubation with anti-ClfB IgG on adherence of *S. aureus* Newman and mutants to immobilized fibrinogen. The square symbol represents wild-type; the diamond symbol represents clfA; the circle symbol represents clfB; the x symbol represents clfAclfB,clfB*.

FIG. 18 is a bar graph showing adherence of *S. aureus* Newman and mutants to explanted hemodialysis tubing.

Cells from two hour shake-flask cultures were used. The graph provides the means and SEM of three experiments.

FIG. 19 is a bar graph showing adherence of *S. aureus* Newman and mutants to fibrinogen immobilized on PMMA (polymethylmethacrylate) coverslips. Cells from two hour shake-flask cultures were used. The graph provides the means and SEM of three experiments.

FIG. 20 is a table which shows the highly conserved amino acid sequences in the A region of ClfA, ClfB, SdrC, SdrD and SdrE, which are used to provide consensus and variable motifs.

FIG. 21 is a graph of absorbance versus concentration of anti-TYTFTDYVD (SEQ ID NO: 18) antibodies, demonstrating the binding of increasing concentrations of (o the antibodies to ClfA, ClfB or BSA coated plates. BSA-coated plates are used as a control, and no significant binding is observed. The closed square symbol represents antibody bound to ClfB; the open diamond symbol represents antibody bound to ClfA; the open circle symbol represents BSA.

FIG. 22 is a Western Blot which illustrates the differing specificities of ClfA and ClfB in the binding of human fibrinogen. The Western Blot was created by the separation of human fibrinogen, and later, the incubation of the nitrocellulose membrane with the A region of either biotinylated ClfA or ClfB. Biotinylated ClfA region A binds the γ chain of fibrinogen, as is seen in lane A2. Biotinylated ClfB region A binds to both the α and β chains of fibrinogen, as seen in lane B2.

FIG. 23 is a bar graph showing adherence of recombinant SdrC region A (SdrCA) to ten different extracellular matrix proteins, BSA, BSP, two forms of collagen, decorin, fibrinogen, fibronectin, laminin, plasmin and vitronectin. The extracellular matrix proteins were immobilized on microtiter wells. Absorbance tests revealed reactivity of SdrCA with fibrinogen, BSP, decorin, plasmin and vitronectin.

DETAILED DESCRIPTION OF THE INVENTION

Isolated extracellular matrix-binding proteins, designated ClfB, SdrC, SdrD and SdrE, and their corresponding amino acid and nucleic acid sequences and motifs are described. The proteins, peptides, fragments thereof or antigenic portions thereof are useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection and as scientific research tools. Further, antibody or antibody fragments to the proteins, peptides, fragments thereof or antigenic portions thereof are also useful for the prevention, inhibition, treatment and diagnosis of *S. aureus* infection. In particular, the proteins or antibodies, or active fragments thereof may be administered as vaccines to induce either passive or cellular immunity.

ClfB binds to at least fibrinogen.

SdrC has been discovered to bind to extracellular matrix proteins of the host, including for example, BSP, decorin, plasmin, vitronectin and fibrinogen. SdrD binds to at least vitronectin. SdrE binds to extracellular matrix proteins, for example, bone sialoprotein (BSP).

The amino acid sequence of ClfB is SEQ ID NO:1. The nucleic acid sequence encoding ClfB is SEQ ID NO:2. The nucleic acid and amino acid sequences of ClfB are also provided in FIG. 5. The amino acid and nucleic acid sequences of SdrC are SEQ ID NOS:3 and 4 respectively and are provided in FIG. 7. The amino acid and nucleic acid sequences of SdrD are SEQ ID NOS:5 and 6 respectively

and are provided in FIG. 8. The amino acid and nucleic acid sequences of SdrE are SEQ ID NOS:7 and 8 respectively and are provided in FIG. 9. The term "isolated" is defined herein as free from at least some of the components with which it naturally occurs. In a preferred embodiment, an isolated component is at least 90% pure, and more preferably 95%.

ClfB has a predicted molecular weight of approximately 88 kDa and an apparent molecular weight of approximately 124 kDa. ClfB is a cell-wall associated protein and binds both soluble and immobilized fibrinogen. In addition, ClfB binds both the alpha and beta chains of fibrinogen and acts as a clumping factor. Despite the low level of identity between ClfA and ClfB, both proteins bind fibrinogen (on different chains) by a mechanism that is susceptible to inhibition by divalent cations, despite not sharing obvious metal binding motifs. The ClfB protein has been demonstrated to be a virulence factor in experimental endocarditis.

The SdrC, SdrD and SdrE proteins are related in primary sequence and structural organization to the ClfA and ClfB proteins and are localized on the cell surface. The SdrC, SdrD and SdrE proteins are cell wall-associated proteins, having a signal sequence at the N-terminus and an LPXTG (SEQ ID NO: 14) motif, hydrophobic domain and positively charged residues at the C-terminus. Each also has an SD repeat containing region R of sufficient length to allow, along with the B motifs, efficient expression of the ligand binding domain region A on the cell surface. With the A region of the SdrC, SdrD and SdrE proteins located on the cell surface, the proteins can interact with proteins in plasma, the extracellular matrix or with molecules on the surface of host cells. The Sdr proteins share some limited amino acid sequence similarity with ClfA and ClfB. Like ClfA and ClfB, SdrC, SdrD and SdrE also exhibit cation-dependent ligand binding of extracellular matrix proteins.

It was surprising to learn that the disclosed extracellular matrix-binding proteins share a unique dipeptide repeat region (region R) including predominately aspartate and serine residues. It had been reported by McDevitt et al., *Mol. Microbiol.* 11: 237–248 (1994); McDevitt et al., *Mol. Microbiol.* 16:895–907 (1995) that ClfA also has this R repeat region. He reported that there were genes in *S. epidermidis* that hybridized to the gene encoding the R domain containing protein. However, McDevitt et al did not know the function of the R region and had not discovered that other cell surface proteins from *S. aureus*, *S. hemolyticus*, *S. lugdenensis*, *S. schleriferi* share this unusual motif. Therefore, in one aspect of this invention, a method is provided for the identification of genes and encoding proteins from *S. aureus* (other than ClfA), *S. hemolyticus*, *S. lugdenensis*, *S. schleriferi* useful for the prevention, treatment, and diagnosis of bacterial infection that includes using the R repeat region as an identifying probe.

The DS repeat is encoded by 18 nucleotide repeats with the consensus (where Y equals pyrimidines and N equals any base) GAYTCNGAYT CNGAYACY (SEQ ID NO: 9), with TCN as the first and second serine codons and AGY as the third serine codon. The R region is near the C-terminus of the proteins and typically contains between 40 and 300 DS residues, or more particularly, greater than 40, 60, 80, 100, 125, 150, 200 or 250 repeating units, of which greater than 90, 95 or even 98% of the amino acids are D or S. The R region DS repeat varies in length between proteins, and while the R region itself does not bind extracellular matrix proteins, the R region enables the presentation of the binding regions of the protein on the cell surface of *S. aureus*. Thus, probes to the consensus DNA encoding the DS repeat (see

above) can be used to identify other genes encoding different binding proteins essential to the attachment of *S. aureus* to host tissues. Antibodies to an R region can be used to discover such additional binding proteins as well.

The sdr genes are closely linked and tandemly arrayed. The Sdr proteins have both organizational and sequence similarity to ClfA and ClfB. At the N-terminus secretory signal sequences precede A regions which are approximately 500 residues in length. The A regions of the Sdr and Clf proteins exhibit only 20–30% residue identity when aligned with any other member of the family.

It has been discovered that in the A region of SdrC, SdrD, SdrE, ClfA, and ClfB, there is highly conserved amino acid sequence that can be used to derive a consensus TYTFT-DYVD (SEQ ID NO: 18) motif. The motif exhibits slight variation between the different proteins. This variation, along with the consensus sequence of the motif is depicted in FIG. 20. In the Clf-Sdr proteins, this motif is highly conserved. The motif can be used in vaccines to impart broad spectrum cellular immunity to bacterial infections, and also can be used as an antigen in the production of monoclonal or polyclonal antibodies. Such an antibody can be used to impart broad spectrum passive immunity.

The Sdr proteins differ from ClfA and ClfB by having two to five additional 110–113 residue repeated sequences (B-motifs) located between region A and the R-region. Each B-motif contains a consensus Ca²⁺-binding EF-hand loop normally found in eukaryotic proteins. The structural integrity of a recombinant protein comprising the five B-repeats of SdrD was shown by bisANS fluorescence analysis to be Ca²⁺-dependent, suggesting that the EF-hands are functional. When Ca²⁺ was removed the structure collapsed to an unfolded conformation. The original structure was restored by addition of Ca²⁺. The C-terminal R-domains of the Sdr proteins contain 132–170 SD residues. These are followed by conserved wall-anchoring regions characteristic of many surface proteins of Gram positive bacteria. The sdr locus was present in all 31 *S. aureus* strains from human and bovine sources tested by Southern hybridization, although in a few strains it contained two rather than three genes.

In the Sdr and Clf proteins this B motif is highly conserved while a degenerate version occurs in fibronectin binding MSCRAMMS, as well as the collagen binding protein Cna. The B motifs, in conjunction with the R regions, are necessary for displaying the ligand-binding domain at some distance from the cell surface.

The repeated B motifs are one common denominator of the sub-group of SD repeat proteins described herein. These motifs are found in different numbers in the three Sdr proteins from strain Newman. There are clear distinctions between the individual B motifs. The most conserved units are those located adjacent to the R regions (SdrC B2, SdrD B5 and SdrE B3). They differ from the rest at several sites, especially in the C-terminal half. A noteworthy structural detail is that adjacent B repeats are always separated by a proline residue present in the C-terminal region, but a proline never occurs between the last B repeats and the R region. Instead this linker is characterized by a short acidic stretch. These differences are evidence that the end units have a different structural or functional role compared to the other B motifs. The N-terminal B motifs of SdrD and SdrE have drifted apart from the others, and there are numerous amino acid alterations, including small insertions and deletions whereas the remaining internal B motifs are more highly conserved. Note that each of the three Sdr proteins has at least one B motif of each kind.

The C-terminal R-domains of the Sdr proteins contain 132–170 SD residues. These are followed by conserved wall-anchoring regions characteristic of many surface proteins of Gram positive bacteria.

ClfB, SdrC, SdrD and SdrE subdomains are shown in FIG. 10 and, depending on the protein, include subdomains A and B1–B5.

The terms ClfB protein, SdrC protein, SdrD protein and SdrE protein are defined herein to include ClfB, SdrC, SdrD and SdrE subdomains, and active or antigenic fragments of ClfB, SdrC, SdrD and SdrE proteins, such as consensus or variable sequence amino acid motifs. Active fragments of ClfB, SdrC, SdrD, SdrE and consensus or variable sequence amino acid motifs peptides or proteins are defined herein as peptides or polypeptides capable of blocking the binding of *S. aureus* to extracellular matrix proteins. Antigenic fragments of ClfB, SdrC, SdrD, SdrE proteins or the consensus or variable amino acid motifs are defined herein as peptides or polypeptides capable of producing an immunological response.

Nucleic Acid Sequences

The nucleic acid sequences encoding ClfB, SdrC, SdrD, SdrE and the consensus or variable sequence amino acid motifs are useful for the production of recombinant extracellular matrix-binding proteins. They are further useful as nucleic acid probes for the detection of *S. aureus*-binding proteins in a sample or specimen with high sensitivity and specificity. The probes can be used to detect the presence of *S. aureus* in the sample, diagnose infection with the disease, quantify the amount of *S. aureus* in the sample, or monitor the progress of therapies used to treat the infection. The nucleic acid and amino acid sequences are also useful as laboratory research tools to study the organism and the disease, thus furthering the development of therapies and treatments for the disease.

It will be understood by those skilled in the art that ClfB, SdrC, SdrD, SdrE and the consensus or variable sequence amino acid motifs are also encoded by sequences substantially similar to the nucleic acid sequences provided in the sequence listing. By “substantially similar” is meant a DNA sequence which, by virtue of the degeneracy of the genetic code, is not identical with that shown in any of SEQ ID NOS:2, 4, 6, and 8, but which still encodes the same amino acid sequence; or a DNA sequence which encodes a different amino acid sequence but retains the activities of the proteins, either because one amino acid is replaced with another similar amino acid, or because the change (whether it be substitution, deletion or insertion) does not affect the active site of the protein. In the latter case, the sequence has substantial homology to the disclosed sequence if it encodes a protein with at least 70% 80%, 90%, 95% or even 98% of the same amino acids.

Also provided herein are sequences of nucleic acid molecules that selectively hybridize with nucleic acid molecules encoding the extracellular matrix-binding proteins from *S. aureus* described herein or complementary sequences thereof. By “selective” or “selectively” is meant a sequence which does not hybridize with other nucleic acids to prevent adequate detection of ClfB, SdrC, SdrD, SdrE or the consensus or variable sequence amino acid motifs. Therefore, in the design of hybridizing nucleic acids, selectivity will depend upon the other components present in a sample. The hybridizing nucleic acid should have at least 70% complementarity with the segment of the nucleic acid to which it hybridizes. As used herein to describe nucleic acids, the term

“selectively hybridizes” excludes the occasional randomly hybridizing nucleic acids, and thus, has the same meaning as “specifically hybridizing”. The selectively hybridizing nucleic acids of the invention can have at least 70%, 80%, 85%, 90%, 95%, 97%, 98%, and 99% complementarity with the segment of the sequence to which it hybridizes.

The invention contemplates sequences, probes and primers which selectively hybridize to the encoding DNA or the complementary, or opposite, strand of DNA as those specifically provided herein. Specific hybridization with nucleic acid can occur with minor modifications or substitutions in the nucleic acid, so long as functional species-specific hybridization capability is maintained. By “probe” is meant nucleic acid sequences that can be used as probes or primers for selective hybridization with complementary nucleic acid sequences for their detection or amplification, which probes can vary in length from about 5 to 100 nucleotides, or preferably from about 10 to 50 nucleotides, or most preferably about 18–24 nucleotides. Therefore, the terms “probe” or “probes” as used herein are defined to include “primers”. Isolated nucleic acids are provided herein that selectively hybridize with the species-specific nucleic acids under stringent conditions and should have at least 5 nucleotides complementary to the sequence of interest as described by Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.

If used as primers, the composition preferably includes at least two nucleic acid molecules which hybridize to different regions of the target molecule so as to amplify a desired region. Depending on the length of the probe or primer, the target region can range between 70% complementary bases and full complementarity and still hybridize under stringent conditions. For example, for the purpose of diagnosing the presence of the *S. aureus*, the degree of complementarity between the hybridizing nucleic acid (probe or primer) and the sequence to which it hybridizes (e.g., *S. aureus* DNA from a sample) is at least enough to distinguish hybridization with a nucleic acid from other bacteria.

The nucleic acid sequences encoding ClfB, SdrC, SdrD, SdrE active fragments thereof or consensus or variable sequence amino acid motifs can be inserted into a vector, such as a plasmid, and recombinantly expressed in a living organism to produce recombinant ClfB, SdrC, SdrD and SdrE proteins or fragments thereof, such as consensus or variable sequence amino acid motifs. For example, DNA molecules producing recombinant ClfB, SdrC, and both SdrD and SdrE were deposited in plasmids pA1-1EX, pC1 and lambda phage A6-2, respectively, at the NCIMB under the Accession Nos. 40903, 40902 and 40904, respectively on Oct. 13, 1997.

Methods for the Detection and Identification of *S. aureus*

Methods of using the nucleic acids described herein to detect and identify the presence of *S. aureus* are provided. The methods are useful for diagnosing *S. aureus* infections and disease such as upper respiratory tract infections (such as otitis media, bacterial tracheitis, acute epiglottitis, thyropharyngitis), lower respiratory infections (such as emphysema, lung abscess), cardiac (such as infective endocarditis), gastrointestinal (such as secretory diarrhea, splenic abscess, retroperitoneal abscess), central nervous system (such as cerebral abscess), ocular (such as blepharitis, conjunctivitis, keratitis, endophthalmitis, pre-

septal and orbital cellulitis, dacryocystitis), kidney and urinary tract (such as epididymitis, intrarenal and perinephric abscess, toxic shock syndrome), skin (such as impetigo, folliculitis, cutaneous abscesses, cellulitis, wound infection, bacterial myositis, bone and joint (such as septic arthritis, osteomyelitis).

The method involves the steps of obtaining a sample suspected of containing *S. aureus*. The sample may be taken from an individual, such as a wound, blood, saliva, tissues, bone, muscle, cartilage, or skin. The cells can then be lysed, and the DNA extracted, precipitated and amplified. Detection of *S. aureus* DNA is achieved by hybridizing the amplified DNA with a *S. aureus* probe that selectively hybridizes with the DNA as described above. Detection of hybridization is indicative of the presence of *S. aureus*.

Preferably, detection of nucleic acid (e.g. probes or primers) hybridization can be facilitated by the use of detectable moieties. For example, the probes can be labeled with biotin and used in a streptavidin-coated microtiter plate assay. Other detectable moieties include radioactive labeling, enzyme labeling, and fluorescent labeling, for example.

DNA may be detected directly or may be amplified enzymatically using polymerase chain reaction (PCR) or other amplification techniques prior to analysis. RNA or cDNA can be similarly detected. Increased or decreased expression of ClfB, SdrC, SdrD, SdrE and consensus or variable sequence amino acid motifs can be measured using any of the methods well known in the art for the quantitation of nucleic acid molecules, such as, amplification, PCR, RT-PCR, RNase protection, Northern blotting, and other hybridization methods.

Diagnostic assays which test for the presence of the ClfB or SdrC, SdrD or SdrE proteins, peptides, motifs, fragments thereof or antibodies to any of these may also be used to detect the presence of an infection. Assay techniques for determining protein or antibody levels in a sample are well known to those skilled in the art and include methods such as radioimmunoassay, Western blot analysis and ELISA (Enzyme-Linked Immunosorbant Assay) assays.

Amino Acid Sequences

It will be understood by those skilled in the art that minor amino acid substitutions or deletions may be present in functional ClfB, SdrC, SdrD, SdrE and consensus or variable sequence amino acid motifs, peptides, proteins, or fragments thereof. The amino acid sequences set forth herein and substantially similar amino acid sequences can be used to produce synthetic ClfB, SdrC, SdrD, SdrE and consensus or variable sequence amino acid motifs, peptides, proteins or active fragments thereof. Active ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs, peptide or protein fragments are defined herein as ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs, portions or peptides that bind to extracellular matrix proteins or compete with or prevent *S. aureus* organisms from binding to extracellular matrix proteins such as decorin, plasmin, fibrinogen, vitronectin or bone sialoprotein.

When used in conjunction with amino acid sequences, the term "substantially similar" means an amino acid sequence which is not identical to SEQ ID NOS:1, 3, 5, or 7, but which produces a protein having the same functionality and retaining the activities of ClfB, SdrC, SdrD, SdrE and consensus or variable sequence amino acid motifs, either because one amino acid is replaced with another similar amino acid, or because the change (whether it be substitution, deletion or

insertion) does not affect the active site of the protein or peptide. Two amino acid sequences are "substantially homologous" when at least about 70%, (preferably at least about 80%, and most preferably at least about 90% or 95%) of the amino acids match over the defined length of the sequences.

Extracellular Matrix-Binding Protein Antibodies

The isolated, recombinant or synthetic ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs, or peptides or active fragments thereof or fusion proteins thereof, are useful as scientific research tools to identify *S. aureus* binding sites on the extracellular matrix. This will promote an understanding of the mechanisms of bacterial pathology and the development of antibacterial therapies. Furthermore, the isolated, recombinant or synthetic protein, or antigenic portions thereof (including epitope-bearing fragments), or fusion proteins thereof can be administered to humans or animals as immunogens or antigens. It can be administered alone or in combination with an adjuvant, for the production of antisera reactive with ClfB, SdrC, SdrD, SdrE or motifs or peptides thereof. In addition, the peptides or proteins can be used to screen antisera for hyperimmune patients from whom can be derived antibodies having a very high affinity for the proteins.

Antibodies isolated from the antisera are useful for the specific detection of *S. aureus* or *S. aureus* extracellular matrix-binding proteins or as research tools. The term "antibodies" as used herein includes monoclonal antibodies, polyclonal, chimeric, single chain, bispecific, simianized, and humanized antibodies as well as Fab fragments, including the products of an Fab immunoglobulin expression library.

Monoclonal antibodies are generated by methods well known to those skilled in the art. The preferred method is a modified version of the method of Kearney, et al., *J. Immunol.* 123:1548-1558 (1979), which is incorporated by reference herein. Briefly, animals such as mice or rabbits are inoculated with the immunogen in adjuvant, and spleen cells are harvested and mixed with a myeloma cell line, such as P3X63Ag8,653. The cells are induced to fuse by the addition of polyethylene glycol. Hybridomas are chemically selected by plating the cells in a selection medium containing hypoxanthine, aminopterin and thymidine (HAT). Hybridomas producing the preferred antibodies are cloned, expanded and stored frozen for future production.

Techniques for the production of single chain antibodies are known to those skilled in the art and described in U.S. Pat. No. 4,946,778 and can be used to produce single chain antibodies to the proteins described herein. Phage display technology may be used to select antibody genes having binding activities for ClfB, SdrC, SdrD, SdrE, and consensus or variable sequence amino acid motifs, or antigenic portions thereof, from PCR-amplified v genes of lymphocytes from humans screened for having antibodies to ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs or naive libraries. Bispecific antibodies have two antigen binding domains wherein each domain is directed against a different epitope.

The antibody may be labeled directly with a detectable label for identification and quantitation of *S. aureus*. Labels for use in immunoassays are generally known to those skilled in the art and include enzymes, radioisotopes, and fluorescent, luminescent and chromogenic substances including colored particles such as colloidal gold and latex beads. Suitable immunoassays include enzyme-linked immunosorbent assays (ELISA).

Alternatively, the antibody may be labeled indirectly by reaction with labeled substances that have an affinity for immunoglobulin, such as protein A or G or second antibodies. The antibody may be conjugated with a second substance and detected with a labeled third substance having an affinity for the second substance conjugated to the antibody. For example, the antibody may be conjugated to biotin and the antibody-biotin conjugate detected using labeled avidin or streptavidin. Similarly, the antibody may be conjugated to a hapten and the antibody-hapten conjugate detected using labeled anti-hapten antibody. These and other methods of labeling antibodies and assay conjugates are well known to those skilled in the art.

Antibodies to the disclosed proteins may also be used in production facilities or laboratories to isolate additional quantities of the protein, such as by affinity chromatography.

The proteins, or antigenic portions thereof, are useful in the diagnosis of *S. aureus* bacterial infections and in the development of anti-*S. aureus* vaccines for active or passive immunization. When administered to a wound or used to coat polymeric biomaterials in vitro and in vivo, both the proteins and antibodies thereof are useful as blocking agents to prevent or inhibit the initial binding of *S. aureus* to the wound site or biomaterials. Preferably, the antibody is modified so that it is less immunogenic in the patient to whom it is administered. For example, if the patient is a human, the antibody may be "humanized" by transplanting the complementarity determining regions of the hybridoma-derived antibody into a human monoclonal antibody as described by Jones et al., *Nature* 321:522-525 (1986) or Tempest et al. *Biotechnology* 9:266-273 (1991).

Medical devices or polymeric biomaterials to be coated with the antibodies, proteins and active fragments described herein include, but are not limited to, staples, sutures, replacement heart valves, cardiac assist devices, hard and soft contact lenses, intraocular lens implants (anterior chamber, posterior chamber or phakic), other implants such as corneal inlays, kerato-prostheses, vascular stents, epikeratophalia devices, glaucoma shunts, retinal staples, scleral buckles, dental prostheses, thyroplastic devices, laryngoplastic devices, vascular grafts, soft and hard tissue prostheses including, but not limited to, pumps, electrical devices including stimulators and recorders, auditory prostheses, pacemakers, artificial larynx, dental implants, mammary implants, penile implants, cranio/facial tendons, artificial joints, tendons, ligaments, menisci, and disks, artificial bones, artificial organs including artificial pancreas, artificial hearts, artificial limbs, and heart valves; stents, wires, guide wires, intravenous and central venous catheters, laser and balloon angioplasty devices, vascular and heart devices (tubes, catheters, balloons), ventricular assists, blood dialysis components, blood oxygenators, urethral/ureteral/urinary devices (Foley catheters, stents, tubes and balloons), airway catheters (endotracheal and tracheostomy tubes and cuffs), enteral feeding tubes (including nasogastric, intragastric and jejunal tubes), wound drainage tubes, tubes used to drain the body cavities such as the pleural, peritoneal, cranial, and pericardial cavities, blood bags, test tubes, blood collection tubes, vacutainers, syringes, needles, pipettes, pipette tips, and blood tubing.

It will be understood by those skilled in the art that the term "coated" or "coating", as used herein, means to apply the protein, antibody, or active fragment to a surface of the device, preferably an outer surface that would be exposed to *S. aureus* infection. The surface of the device need not be entirely covered by the protein, antibody or active fragment.

Immunological and Pharmaceutical Compositions

Immunological compositions, including vaccine, and other pharmaceutical compositions containing the ClfB,

SdrC, SdrD, SdrE or consensus or variable sequence amino acid motif, peptides or proteins are included within the scope of the present invention. One or more of the ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motif, peptides, proteins, or active or antigenic fragments thereof, or fusion proteins thereof can be formulated and packaged, alone or in combination with other antigens, using methods and materials known to those skilled in the art for vaccines. The immunological response may be used therapeutically or prophylactically and may provide antibody immunity or cellular immunity such as that produced by T lymphocytes such as cytotoxic T lymphocytes or CD4⁺ T lymphocytes.

The immunological compositions, such as vaccines, and other pharmaceutical compositions can be used alone or in combination with other blocking agents to protect against human and animal infections caused by *S. aureus*. In particular, the compositions can be used to protect humans against endocarditis or to protect humans or ruminants against mastitis caused by *S. aureus* infections. The vaccine can also be used to protect canine and equine animals against similar *S. aureus* infections.

To enhance immunogenicity, the proteins may be conjugated to a carrier molecule. Suitable immunogenic carriers include proteins, polypeptides or peptides such as albumin, hemocyanin, thyroglobulin and derivatives thereof, particularly bovine serum albumin (BSA) and keyhole limpet hemocyanin (KLH), polysaccharides, carbohydrates, polymers, and solid phases. Other protein derived or non-protein derived substances are known to those skilled in the art. An immunogenic carrier typically has a molecular weight of at least 1,000 daltons, preferably greater than 10,000 daltons. Carrier molecules often contain a reactive group to facilitate covalent conjugation to the hapten. The carboxylic acid group or amine group of amino acids or the sugar groups of glycoproteins are often used in this manner. Carriers lacking such groups can often be reacted with an appropriate chemical to produce them. Preferably, an immune response is produced when the immunogen is injected into animals such as mice, rabbits, rats, goats, sheep, guinea pigs, chickens, and other animals, most preferably mice and rabbits. Alternatively, a multiple antigenic peptide comprising multiple copies of the protein or polypeptide, or an antigenically or immunologically equivalent polypeptide may be sufficiently antigenic to improve immunogenicity without the use of a carrier.

The ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motif, peptide, protein or proteins may be administered with an adjuvant in an amount effective to enhance the immunogenic response against the conjugate. At this time, the only adjuvant widely used in humans has been alum (aluminum phosphate or aluminum hydroxide). Saponin and its purified component Quil A, Freund's complete adjuvant and other adjuvants used in research and veterinary applications have toxicities which limit their potential use in human vaccines. However, chemically defined preparations such as muramyl dipeptide, monophosphoryl lipid A, phospholipid conjugates such as those described by Goodman-Snitkoff et al. *J. Immunol.* 147:410-415 (1991) and incorporated by reference herein, encapsulation of the conjugate within a proteoliposome as described by Miller et al., *J. Exp. Med.* 176:1739-1744 (1992) and incorporated by reference herein, and encapsulation of the protein in lipid vesicles such as Novasome™ lipid vesicles (Micro Vesicular Systems, Inc., Nashua, N.H.) may also be useful.

The term "vaccine" as used herein includes DNA vaccines in which the nucleic acid molecule encoding ClfB, SdrC,

SdrD, SdrE and consensus or variable sequence amino acid motifs, or nucleic acid molecules which are not identical to the disclosed sequences, but which are substantially homologous thereto and encode peptides or proteins which have the same functionality and activities, or antigenic portions thereof in a pharmaceutical composition is administered to a patient. For genetic immunization, suitable delivery methods known to those skilled in the art include direct injection of plasmid DNA into muscles (Wolff et al., *Hum. Mol. Genet.* 1:363 (1992)), delivery of DNA complexed with specific protein carriers (Wu et al., *J. Biol. Chem.* 264:16985 (1989), coprecipitation of DNA with calcium phosphate (Benvenisty and Reshef, *Proc. Natl. Acad. Sci.* 83:9551 (1986)), encapsulation of DNA in liposomes (Kaneda et al., *Science* 243:375 (1989)), particle bombardment (Tang et al., *Nature* 356:152 (1992) and Eisenbraun et al., *DNA Cell Biol.* 12:791 (1993)), and in vivo infection using cloned retroviral vectors (Seeger et al., *Proc. Natl. Acad. Sci.* 81:5849, 1984).

Methods of Administration and Dose of Pharmaceutical Compositions

Pharmaceutical compositions containing the ClfB, SdrC, SdrD or SdrE proteins, nucleic acid molecules, antibodies, or fragments thereof may be formulated in combination with a pharmaceutical carrier such as saline, dextrose, water, glycerol, ethanol, other therapeutic compounds, and combinations thereof. The formulation should be appropriate for the mode of administration. The compositions are useful for interfering with, modulating, or inhibiting *S. aureus* host cell binding interactions with the extracellular matrix.

Suitable methods of administration include, but are not limited to, topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal and intradermal administration.

For topical administration, the composition is formulated in the form of an ointment, cream, gel, lotion, drops (such as eye drops and ear drops), or solution (such as mouthwash). Wound or surgical dressings, sutures and aerosols may be impregnated with the composition. The composition may contain conventional additives, such as preservatives, solvents to promote penetration, and emollients. Topical formulations may also contain conventional carriers such as cream or ointment bases, ethanol, or oleyl alcohol.

In a preferred embodiment, a vaccine is packaged in a single dosage for immunization by parenteral (i.e., intramuscular, intradermal or subcutaneous) administration or nasopharyngeal (i.e., intranasal) administration. The vaccine is most preferably injected intramuscularly into the deltoid muscle. The vaccine is preferably combined with a pharmaceutically acceptable carrier to facilitate administration. The carrier is usually water or a buffered saline, with or without a preservative. The vaccine may be lyophilized for resuspension at the time of administration or in solution.

The carrier to which the protein may be conjugated may also be a polymeric delayed release system. Synthetic polymers are particularly useful in the formulation of a vaccine to effect the controlled release of antigens. For example, the polymerization of methyl methacrylate into spheres having diameters less than one micron has been reported by Kreuter, J., *MICROCAPSULES AND NANOPARTICLES IN MEDICINE AND PHARMACOLOGY*, M. Donbrow (Ed). CRC Press, p. 125-148.

Microencapsulation of the protein will also give a controlled release. A number of factors contribute to the selection of a particular polymer for microencapsulation. The

reproducibility of polymer synthesis and the microencapsulation process, the cost of the microencapsulation materials and process, the toxicological profile, the requirements for variable release kinetics and the physicochemical compatibility of the polymer and the antigens are all factors that must be considered. Examples of useful polymers are polycarbonates, polyesters, polyurethanes, polyorthoesters polyamides, poly (d,l-lactide-co-glycolide) (PLGA) and other biodegradable polymers. The use of PLGA for the controlled release of antigen is reviewed by Eldridge, J. H., et al. *CURRENT TOPICS IN MICROBIOLOGY AND IMMUNOLOGY*, 146:59-66 (1989).

One typical dose for human administration is from 0.01 mg/kg to 10 mg/kg. Based on this range, equivalent dosages for heavier body weights can be determined. The dose should be adjusted to suit the individual to whom the composition is administered and will vary with age, weight and metabolism of the individual. The vaccine may additionally contain stabilizers such as thimerosal (ethyl(2-mercaptobenzoate-S)mercury sodium salt) (Sigma Chemical Company, St. Louis, Mo.) or physiologically acceptable preservatives.

Protein-Label Conjugates

When labeled with a detectable biomolecule or chemical, the extracellular matrix-binding proteins described herein are useful for purposes such as in vivo and in vitro diagnostics and laboratory research. Various types of labels and methods of conjugating the labels to the proteins are well known to those skilled in the art. Several specific labels are set forth below. The labels are particularly useful when conjugated to a protein such as an antibody or receptor.

For example, the protein can be conjugated to a radiolabel such as, but not restricted to, ³²P, ³H, ¹⁴C, ³⁵S, ¹²⁵I, or ¹³¹I. Detection of a label can be by methods such as scintillation counting, gamma ray spectrometry or autoradiography.

Bioluminescent labels, such as derivatives of firefly luciferin, are also useful. The bioluminescent substance is covalently bound to the protein by conventional methods, and the labeled protein is detected when an enzyme, such as luciferase, catalyzes a reaction with ATP causing the bioluminescent molecule to emit photons of light.

Fluorogens may also be used to label proteins. Examples of fluorogens include fluorescein and derivatives, phycoerythrin, allo-phycoyanin, phycoyanin, rhodamine, and Texas Red. The fluorogens are generally detected by a fluorescence detector.

The protein can alternatively be labeled with a chromogen to provide an enzyme or affinity label. For example, the protein can be biotinylated so that it can be utilized in a biotin-avidin reaction, which may also be coupled to a label such as an enzyme or fluorogen. For example, the protein can be labeled with peroxidase, alkaline phosphatase or other enzymes giving a chromogenic or fluorogenic reaction upon addition of substrate. Additives such as 5-amino-2,3-dihydro-1,4-phthalazinedione (also known as Luminol) (Sigma Chemical Company, St. Louis, Mo.) and rate enhancers such as p-hydroxybiphenyl (also known as p-phenylphenol) (Sigma Chemical Company, St. Louis, Mo.) can be used to amplify enzymes such as horseradish peroxidase through a luminescent reaction; and luminogenic or fluorogenic dioxetane derivatives of enzyme substrates can also be used. Such labels can be detected using enzyme-linked immunoassays (ELISA) or by detecting a color change with the aid of a spectrophotometer. In addition, proteins may be labeled with colloidal gold for use

in immunoelectron microscopy in accordance with methods well known to those skilled in the art.

The location of a ligand in cells can be determined by labeling an antibody as described above and detecting the label in accordance with methods well known to those skilled in the art, such as immunofluorescence microscopy using procedures such as those described by Warren and Nelson, *Mol. Cell. Biol.* 7: 1326-1337 (1987).

Screening Methods

The ClfB, SdrC, SdrD and SdrE proteins, or fragments thereof, such as consensus or variable amino acid motifs, are useful in a method for screening materials to identify substances that inhibit *S. aureus* host cell binding interactions with the extracellular matrix. In accordance with the method for screening, the substance of interest is combined with one or more of the ClfB, SdrC, SdrD, or SdrE proteins, or fragments thereof, such as consensus or variable sequence amino acid motif peptides, and the degree of binding of the molecule to the extracellular matrix is measured or observed. If the presence of the substance results in the inhibition of binding, then the substance may be useful for inhibiting *S. aureus* in vivo or in vitro. The method could similarly be used to identify substances that promote *S. aureus* interactions with the extracellular matrix.

The method is particularly useful for identifying substances having bacteriostatic or bacteriocidal properties.

For example, to screen for *S. aureus* agonists or antagonists, a synthetic reaction mixture, a cellular compartment (such as a membrane, cell envelope or cell wall) containing one or more of the ClfB, SdrC, SdrD, SdrE proteins, or fragments thereof, such as consensus or variable sequence amino acid motifs, and a labeled substrate or ligand of the protein is incubated in the presence of a substance under investigation. The ability of the substance to agonize or antagonize the protein is shown by a decrease in the binding of the labeled ligand or decreased formation of substrate product. Substances that bind well and increase the rate of product formation from substrate are agonists. Detection of the rate or level of formation of product from substrate may be enhanced by use of a reporter system, such as a calorimetric labeled substrate converted to product, a reporter gene that is responsive to changes in ClfB, SdrC, SdrD, SdrE or consensus or variable amino acid sequence motifs' nucleic acid or protein activity, and binding assays known to those skilled in the art. Competitive inhibition assays can also be used.

Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs' nucleic acid molecules or proteins and thereby inhibit their activity or bind to a binding molecule (such as fibrinogen) to prevent the binding of the ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs' nucleic acid molecules or proteins to the binding molecule. For example, a compound that inhibits ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motifs' activity may be a small molecule that binds to and occupies the binding site of the ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motif peptide or protein, thereby preventing binding to cellular binding molecules. Examples of small molecules include, but are not limited to, small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules. Preferred antagonists include compounds related to and variants or derivatives of ClfB, SdrC, SdrD, SdrE or consensus or variable sequence amino acid motif peptides or proteins.

The nucleic acid molecules described herein may also be used to screen compounds for antibacterial activity.

Therapeutic Applications

In addition to the therapeutic compositions and methods described above, the ClfB, SdrC, SdrD, SdrE or consensus or variable amino acid motifs, peptides or proteins, nucleic acid molecules or antibodies are useful for interfering with the initial physical interaction between a pathogen and mammalian host responsible for infection, to mammalian extracellular matrix proteins on indwelling devices or to extracellular matrix proteins in wounds. they are further useful to block ClfB, SdrC, SdrD, SdrE, or active fragments thereof, including consensus or variable amino acid motifs, peptide or protein-mediated mammalian cell invasion. In addition, these molecules are useful to mediate tissue damage and to block the normal progression of pathogenesis in infections.

S. aureus Detection Kit

The invention further contemplates a kit containing one or more ClfB, SdrC, SdrD, SdrE proteins, peptides, or active fragments thereof, including consensus or variable amino acid motif- encoding nucleic acid probes. These probes can be used for the detection of *S. aureus* or *S. aureus* extracellular matrix-binding proteins in a sample. Such a kit can also contain the appropriate reagents for hybridizing the probe to the sample and detecting bound probe.

In an alternative embodiment, the kit contains one or more ClfB, SdrC, SdrD, or SdrE proteins, peptides or consensus or variable amino acid motif-specific antibodies, which can be used for the detection of *S. aureus* organisms or *S. aureus* extracellular matrix-binding proteins in a sample.

In yet another embodiment, the kit contains one or more ClfB, SdrC, SdrD or SdrE-proteins, or active fragments thereof, such as the consensus or variable sequence amino acid motifs, which can be used for the detection of *S. aureus* organisms or *S. aureus* extracellular matrix-binding antibodies in a sample.

The kits described herein may additionally contain equipment for safely obtaining the sample, a vessel for containing the reagents, a timing means, a buffer for diluting the sample, and a colorimeter, reflectometer, or standard against which a color change may be measured.

In a preferred embodiment, the reagents, including the protein or antibody, are lyophilized, most preferably in a single vessel. Addition of aqueous sample to the vessel results in solubilization of the lyophilized reagents, causing them to react. Most preferably, the reagents are sequentially lyophilized in a single container, in accordance with methods well known to those skilled in the art that minimize reaction by the reagents prior to addition of the sample.

EXAMPLES

The present invention is further illustrated by the following non-limiting examples, which are not to be construed in any way as imposing limitations upon the scope thereof. On the contrary, it is to be clearly understood that resort may be had to various other embodiments, modifications, and equivalents thereof which, after reading the description herein, may suggest themselves to those skilled in the art without departing from the spirit of the present invention.

Example 1

Gene Cloning, Sequencing and Expression

A fibrinogen-binding protein gene, designated clfB, was isolated, cloned and sequenced as follows:

The *E. coli* and *S. aureus* strains used for the cloning and sequencing of *clfB* are listed in Table 1, below. *Escherichia coli* was routinely grown on L-broth or agar. *S. aureus* was routinely grown on trypticase soy broth (Oxoid) or agar. The following antibiotics were incorporated into media where appropriate: ampicillin (Ap), 100 µg/ml; tetracycline (Tc), 2 µg/ml; chloramphenicol (Cm), 5 µg/ml; erythromycin (Em) 10 µg/ml.

TABLE 1

Bacterial strains used in the present study			
Bacterial strain	Genotype	Relevant properties/ Use in present study	Source/ reference
<i>E. coli</i>			
C600	F ⁻ , lacY1, leuB6, supE44, thi-1, thr-1, tonA21	Propagation of lambda recombinants	Appleyard, Genetics 39: 440-452 (1954)
DH5α	F ⁻ , φ80dlacZM15, deoR, endA1, gyrA96, hsdR17, (r _k ⁻ , m _k ⁺), (lacZYA-argF)U169, recA1, relA1, supE44, thi-1	Recombination deficient, host strain for plasmids and for DNA sequencing	Hanahan et al., J. Mol. Biol. 166:557-580 (1983)
JM101	supE, thi-1, (lac-proAB), [F ⁺ traD36, proAB, lac ⁺ ZM15]	Host strain for plasmid bank and for sequencing	Stratagene (La Jolla, CA)
LE392	F ⁻ , (r _k ⁻ , m _k ⁺), galK2, galT22, hsdR574, lacY1 or (lacIZY)6, metB1, supE44, supF58, tprR55	Propagation of lambda recombinants	(Promega Corp. (Madison, WI))
XL-1 Blue	[F ⁺ proAB, lac ⁺ ZM15, Tn10(tc ^r), endA1, gyrA96, hsdR17, lac, recA1, relA1, supE44, thi-1	Propagation of plasmids	Stratagene (La Jolla, CA)
<i>S. aureus</i>			
Newman		Strong adherence to fibrinogen	NCTC 8178; Duthie and Lorenz, J. Gen. Microbiol. 6: 95-107 (1952)
DU5876		<i>clfA2</i> ::Tn917, Em _r	McDevitt et al., Mol. Microbiol. 11: 237-248 (1994)
DU5943		<i>clfB</i> ::Tc _r , Tc _r	described herein
DU5944		<i>clfA</i> Δ <i>clfB</i> , Em _r , Tc _r	described herein
DU5874		<i>spa</i> ::Tc _r	Protein A-defective mutant of NewmanMcDevitt et al., Mol. Microbiol. 16:895-907 (1995)
Δ map			McDevitt, unpublished
8325-4		NCTC 8325 cured of prophages	Novick, Virology 33: 155-166 (1967)

TABLE 1-continued

Bacterial strains used in the present study			
Bacterial strain	Genotype	Relevant properties/ Use in present study	Source/ reference
ISP546		<i>agr</i> ::Tn551	8325-4 agrBrown and Pattee, Infect. Immun. 30: 36-42 (1980)
RN4220		Restriction deficient derivative of 8325-4	Kreiswirth et al., Nature 305: 709-712 (1983)
V8		Classic V8 protease producer, produces PV leukocidin	ATCC 27733
Cowan 1		Classic protein A producer, adheres well to fibrinogen and fibronectin	ATCC 12598
RN4282		TSST-1 producer	Kreiswirth et al., 1983 (as 3-14)
Phillips		Collagen binding strain	Patti et al., Infect. Immun. 62: 152-161 (1994)
V13		Septicaemia isolate	O'Reilly et al., Mol. Microbiol. 4: 1947-1955 (1990)
GH13		Methicillin resistant	Poston and Li Saw Hee, J. Med. Microbiol. 34: 193-201 (1991)
P1		Rabbit virulent strain	Sherertz et al., J. Infect. Dis. 167: 98-106 (1993)
M60		Bovine mastitis isolate	Anderson, Zentralbl Bakteriol Parasitenkd Infektionskr Hyg Abt. 1 Orig Reihe A 5(Suppl.): 783-790 (1976)

DNA Manipulation

Unless otherwise specified, DNA manipulations were done according to standard methods as described by Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY. New York, John Wiley and Sons (1987) and Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed. Cold Spring Harbour, N.Y., Cold Spring Harbour Laboratory Press (1989). Enzymes for DNA manipulation were obtained from New England Biolabs (Beverly, Mass.) and Promega (Madison, Wis.), and used as directed by the manufacturer. Genomic DNA from *S. aureus* Newman was prepared according to methods of Muller et al., Infect. Immun. 61:551-558 (1993). Smaller scale preparations were made by lysing cells in phosphate buffered saline (PBS) containing 12 µg/ml lysostaphin and 20 mM EDTA (ethylenediaminetetraacetic acid), followed by protease K treatment (500 µg/ml in 1% SDS) for 1 hour at 60° C., extraction with phenol and chloroform, and dialysis against 10 mM Tris HCL, pH 8.0, 1 mM EDTA. Plasmid

DNA was prepared from *S. aureus* according to the method of Vriesema et al., *Appl. Environ. Microbiol.* 62:3527–3529 (1996). *E. coli* plasmid DNA for use in polymerase chain reaction (PCR) and sequencing was routinely made by the modified alkaline lysis method of Feliciello and Chinali, *Anal. Biochem.* 212:394–401 (1993), and occasionally by large scale isolation and dye-buoyant density centrifugation. Screening of *E. coli* transformants for chimeric plasmids was routinely done by the rapid colony lysis-procedure of Le Gouill and Dery, *Nucl. Acids Res.* 19:6655 (1994).

Cloning of Repeat-containing Loci

A genomic library of *S. aureus* Newman was constructed in the LambdaGEM-12 replacement vector (obtained as prepared XhoI half-site arms from Promega Madison, Wis.) according to the manufacturer's instructions. Oligonucleotide probes specific for regions A and R of *S. aureus* Newman were made by polymerase chain amplification of these regions from the cloned gene on pCF14, as described by McDevitt and Foster, *Microbiology* 141:937–943 (1995), and random-primer labeled with [α - 32 P]dATP using the Promega Prime-a-Gene™ kit (Promega). The bank was screened by Southern blotting, using an overnight hybridization temperature of 65° C. Selected clones were single plaque purified twice, and plate-lysate stocks made for storage and for inoculation of liquid cultures for the large-scale preparation of phage for DNA isolation.

A 3.87-kb HindIII fragment containing homology to region R DNA was cloned from the genome of *S. aureus*. HindIII-cleaved genomic DNA in the range of 3–4 kb was excised from an agarose gel, purified, and ligated to the pBluescript cloning vector. Plasmids were transformed into *E. coli* JM101 and identification of a recombinant *E. coli* containing a region R DNA insert was identified by PCR screening. PCR products were generated using primers specific for region R DNA. Individual colonies within a pool producing a positive PCR reaction were then analyzed for their potential to generate a PCR product. One transformant, pC1, was identified and found to contain the 3.87-kb fragment with homology to region R.

DNA Sequencing

The DNA sequence of *clfB* was obtained from pA1-1EX, a plasmid containing a fragment subcloned from recombinant phage A1-1 into pGEM 7Z (f)+. Nested deletions were made using the Erase-a-Base™ Kit (Promega). The Flash Dye Primer Sequencing Kit (Genpak) was used for sequencing in a Model 373A sequencing system (Applied Biosystems, Foster City, Calif.). Confirmatory sequencing in the forward direction was carried out. Double stranded sequencing of *sdrD* and *sdrE* was done on the subclones pEJ1, pEJ2 and pEJ3, containing fragments subcloned from recombinant phage A6-2 in pGEM 7Z (f)+, by nested deletions and primer walking. Automated sequence analysis of *sdrC* and the 5' end of *sdrD* on plasmid clone pC1 was performed. Sequence analysis was performed on both strands by primer extension to known sequences.

Screening of *S. aureus* Strains for *clfB* Homologues

A probe specific for the region A-encoding portion of *clfB* was made by excising a 614 bp internal AccI fragment from pA1-1EX, purifying from an agarose gel using the GENECLEAN II™ kit (BIO 101 Inc., La Jolla, Calif.), and labeling with [α - 32 P]dATP as described in FIG. 2. A probe was similarly made to distal regions of the gene

(encoding region R, the wall and membrane-spanning regions, and about 100 bp of downstream DNA), using a 1.2 kb BamHI fragment from pA1-1EX. HindIII digests of genomic DNA from a panel of strains were Southern blotted and screened using these probes.

Expression of *clfB* Region A

Region A (encoding residues S45 to N542) of *clfB* was amplified from pA1-1EX by PCR using the following primers: (SEQ ID NO:15)

Forward: 5' CGAGGATCCTCAGGACAATCGAAC-GATACAACG 3' (SEQ ID NO: 16)

Reverse: 5' CGAGGTACCATTTACTGCTGAATCACC 3'.

Cleavage sites for BamHI and KpnI (underlined) were appended to the 5' ends of the respective primers to introduce these sites into the product and facilitate its cloning into expression vectors. The forward primer was subsequently found to include a single base mismatch (G, underlined), changing an E codon to a G codon. Reaction mixtures (50 μ l) contained 2 mM dNTPs, 1.5 mM MgCl₂, 1 ng pA1-1EX, 50 nM primers and 1.25 U Taq polymerase in standard Promega (Madison, Wis.) Taq reaction buffer. Amplification proceeded in a Perkin Elmer Cetus (Foster City, Calif.) thermocycler with an initial denaturation at 94° C. for 4 minutes, followed by 30 cycles with denaturation at 94° C. for 1 minute, annealing at 50° C., and extension at 72° C. for 1.5 minutes, with minimum heating and cooling between steps. The final extension was for 10 minutes. A single product was obtained, which was purified using the Wizard™ PCR purification kit (Promega). The product was initially cloned into the His-tag expression vector pQE30. However, because high-level expression was not obtained in this system, the product was recloned into an alternative vector, the GST fusion vector pGEX-KG, between the BamHI and HindIII sites. The recombinant protein was recovered from lysates by affinity chromatography on glutathione-sepharose (GST Gene Fusion System™, Pharmacia, Piscataway, N.J.) and from the glutathione-S-transferase fusion partner by thrombin cleavage.

Cloning of Repeat Carrying Loci

A library of *S. aureus* Newman genomic DNA was made using the replacement lambda vector LambdaGEM™-12. About 10 000 plaques were screened using the region R-specific probe. Of the 60 positive plaques retained, 26 were purified and counter-screened with a *clfA* region A-specific probe. One plaque hybridized with the latter, indicating that it contained the *clfA* gene; of the remaining, non-hybridizing plaques, three were selected at random, and the DNA isolated. The DNA was cut with several restriction enzymes and analyzed by Southern blotting using the region R probe. Clones A1-1 and A2-3 appeared to contain overlapping sequences. Restriction mapping and Southern blotting indicated that these clones contained a single region R homologue. Clone A6-2 was found to contain three region R homologues, since cleavage with EcoRV yielded three fragments hybridizing to the region R probe.

Clone A1-1 was chosen for more detailed study, as the hybridizing fragment was slightly longer than in clone A2-3. A 7.4 kb EcoRI fragment containing the repeat region was subcloned from lambda clone A1-1 into plasmid pGEM 7Z f(+) to generate plasmid pA1-1E. This insert was reduced to approximately 3 kb by excision of a 4.4 kb XbaI segment to form pA1-1EX as shown in FIGS. 2 and 3.

Clone A6-2 was restriction mapped and fragments subcloned into plasmid vectors for sequencing as shown in FIG.

4. Southern blotting with the region R probe and preliminary sequencing suggested that there were three tandemly arrayed genes carrying region R encoding sequences. On A6-2 there were two complete ORFs, sdrD and sdrE, and one incomplete ORF, sdrC.

The two complete ORFs were sequenced on fragments subcloned from lambdaA6-2 into plasmid vectors pGEM7Z f(+) (subclones pEJ1 and pEJ2) and pBluescript KS+ (subclone pEJ3). sdrC was cloned separately from *S. aureus* genomic DNA. A 3.87-kb HindIII fragment of strain Newman was cloned directly into plasmid pBluescript KS+, generating clone pCl (FIG. 4). This clone, containing a region R DNA insert, was identified by PCR screening. The sequence of sdrC and the 162 bp at the 5' end of sdrD were determined from pCl.

Plasmid pA1-1EX, carrying the clfB gene, was deposited at the National Collections of Industrial and Marine Bacteria on Oct. 13, 1997 under the Accession No. 40903. Plasmid pCl, carrying the gene for sdrC, was deposited at the National Collections of Industrial and Marine Bacteria under the Accession No. NCIMB 40902 on Oct. 13, 1997 and a recombinant lambda phage A6-2, carrying the sdrD and sdrE genes, was deposited at the NCIMB on Oct. 13, 1997 under the Accession No. NCIMB 40904. All deposits comply with the terms of the Budapest Treaty.

Features of ClfB

The translated open reading frame (ORF) contained within pA1-1EX is shown in FIG. 5. The ORF shows features reminiscent of secreted proteins of Gram positive cocci. Although the entire ORF is shown in FIG. 5, the start codon is unlikely to be the N codon. There is no ATG codon at the 5' end of the ORF. However, GTG and TTG are occasionally used as translational start codons in *S. aureus*, although methionine is the actual amino acid residue inserted, e.g., the fibronectin binding proteins (GTG), and protein A (TTG). The first TTG codon (L) may well be the initiation codon, as a possible ribosome binding site, GGAG, is suitably located upstream, starting at position-12. The N-terminal 44 amino acid residue region thus predicted has properties similar to signal sequences of secreted proteins of Gram positive cocci, i.e., an initial stretch of 19 mostly polar residues, with an overall positive charge, followed by 18 neutral residues with a high content of hydrophobic residues, and finally a short stretch of mainly polar residues with a good consensus cleavage site, AQA-S.

If the above prediction of the signal sequence is correct, region A of ClfB is 498 residues long, and shows 26.3% residue identity with the equivalent region in ClfA, or 44.4% homology when conservative substitutions are included. The most marked stretch of amino acid similarity between ClfA and ClfB occurs between residues 314-329 (ClfA) and 304-319 (ClfB), with 7 identical and 5 conserved residues. In ClfA, the stretch overlaps the C-terminal half of a putative Ca²⁺ binding loop, EF hand I, required for fibrinogen binding as shown in FIG. 6. The sequence DYSNS (SEQ ID NO:11), which obeys the consensus for the N-terminal moiety of a MIDAS motif, occurs a short distance upstream. Accordingly, the downstream sequence was inspected for D and T residues to complete the motif. D and T occur frequently throughout the protein, and T 339 is suitably located, 63 residues downstream. However, the consensus would require a D residue 14-23 residues downstream from the T, and in the present case, the nearest D residues are 9 or 28 residues away (D 348 and D 367).

At the C-terminal end of region A, a prominent proline-rich region occurs (21/42 residues are P; as shown in FIG.

5). There is a 14-residue repeat within this sequence. The DNA encoding the P-rich repeats is highly conserved. Of the three base substitutions, only one results in an amino acid replacement, a conservative substitution of S for T.

5 Region R is somewhat shorter in clfB than in clfA (272 residues instead of 308). The region R encoding sequence comprises the 18-bp consensus repeat observed in the equivalent part of clfA.

10 Following region R is a short stretch of predominantly hydrophilic residues, containing the distinctive LPETG (SEQ ID NO: 22) motif near its C-terminal end, presumably the cell-sorting signal. The C-terminal region of the predicted protein shows strong homology with the corresponding region in ClfA, with an initial stretch of mostly hydrophobic residues in a final stretch rich in positively charged residues, reminiscent of membrane spanning and anchoring domains, respectively. The general organization of ClfA and ClfB is compared in FIG. 1.

20 A putative transcription termination signal occurs 3' to clfB. No open reading frames occur within 260 bp 5' or 200 bp 3', suggesting that the gene is not part of an operon.

Features of SdrC, SdrD and SdrE

25 The DNA sequences and the translated amino acid sequences of sdrC, sdrD and sdrE are shown in FIGS. 7, 8 and 9. Each predicted protein has a putative signal sequence, an approximately 500 residue "region A" with limited homology to region A of ClfA (see FIG. 10), variable numbers of B repeats, an SD repeat containing region F, and LPXTG (SEQ ID NO: 14) cell wall sorting motif, a hydrophilic membrane anchor, and positively charged residues at the extreme C terminus.

30 The organization of the five region R containing proteins is shown in FIG. 10. The A regions of SdrC, SdrD and SdrE have limited sequence similarity to each other and to those of ClfA and ClfB as shown in FIG. 11. Alignments of those sequences more strongly conserved between all five proteins are shown in FIG. 12. The consensus motif TYTFTDYVD (SEQ ID NO: 18) overlaps the EF hand 1 motif of ClfA (alignment 2, FIG. 12). This region of ClfA has been shown to be of crucial significance in its ligand (fibrinogen) binding activity as described by O'Connell et al., *J. Biol. Chem.*, 273:6821-6829 (1998), and may also be of importance in the biological activity of the new proteins.

35 The three proteins SdrC, SdrD and SdrE form a separate subgroup of region R containing proteins: in addition to regions R and A they contain variable numbers of B repeats, located between region A and region R. The B repeats are 110-113 amino acids long and show considerable similarity (alignment 5, FIG. 12). The repeats SdrC B2, SdrD B5 and SdrE B3 adjacent to region R are 93-95% identical. There is a strongly conserved EF hand near the N-terminal end of each repeat.

clfB Homologues in Other *S. Aureus* Strains

40 Nine strains of *S. aureus* were screened for the clfB gene by Southern blotting. Genomic DNA was cut to completion with HindIII, and probed with an internal 0.6 kb AccI fragment of the region A coding sequence of clfB, shown in FIG. 2. The probe recognized a single HindIII fragment varying from 2 to 3 kb in length in all nine strains, indicating that each possesses a single clfB allele. A probe made from the region R and distal regions of clfB recognized an identical band in all strains, indicating that the clfB homologues in other strains also contain region R.

Expression of clfB

The portion of clfB encoding region A was amplified by PCR using primers incorporating suitable 5' restriction sites, and cloned into the *E. coli* expression vector pGEX-KG. A protein of 94.3 kDa was detected in lysates in induced bacteria. The GST-ClfB fusion protein was immobilized on a glutathione sepharose affinity column, cleaved with thrombin, and examined by SDS-PAGE. The predominant band was 42 kDa, whereas the calculated molecular weight of region A is 54 kDa. This protein was used to raise antibody in rabbits, to probe Western blots of cell lysates made from strain Newman grown under a variety of conditions, as described below. The antibody failed to detect any antigens in lysates made from plate cultures, statically grown broth cultures, or shake-flask cultures grown to stationary phase. A single 124-kDa band was detected in lysates made from exponential phase shake-flask cultures of strain Newman and derivatives. If it is assumed that processing removes the signal sequence and the C-terminal portion of the protein from the last G of the LPETG (SEQ ID NO:22), the predicted molecular weight of ClfB is 88.3 kDa. In a time-course of ClfB production by a shake-flask culture of strain Newman, the ClfB protein was most abundant in the early exponential phase and showed a sharp decline toward the end of exponential phase, after which levels became undetectable. The results of the time-course study are shown in FIG. 13.

Example 2

Production of Anti-ClfB Serum

Antibodies to recombinant region A were raised in two young New Zealand white rabbits (2 kg) showing no prior reaction with *E. coli* or *S. aureus* antigens in Western blots. Injections, given subcutaneously, contained 25 μ g of the antigen, diluted to 500 ml in phosphate buffered saline (PBS) emulsified with an equal volume of adjuvant. The initial injection contained Freund's complete adjuvant; the two to three subsequent injections, given at two-week intervals, contained Freund's incomplete adjuvant. When the response to the recombinant protein was judged adequate, the rabbits were bled, serum recovered, and total IgG purified by affinity chromatography on protein A sepharose (Sigma Chemical Co., St. Louis, Mo.).

SDS-PAGE and Western Blotting

Samples were analyzed by SDS-PAGE in 10 or 12% acrylamide gels. Isolated proteins and *E. coli* cell lysates were prepared for electrophoresis by boiling for five minutes in denaturation buffer. For *S. aureus*, cells were suspended to an OD₆₀₀ of 40 units in 100 mM PBS containing 10 mM EDTA. To each 500 μ l sample, 40 μ l protease inhibitors (Complete™ cocktail, Boehringer Mannheim, Indianapolis, Ind.), 5 μ l each of DNase and RNase (from 10 mg/ml stocks, Sigma Chemical Co.), and 60 μ l of a 2 mg/ml lysostaphin stock (Ambicin L™ recombinant lysostaphin, Applied Microbiology Inc., Tarrytown, N.Y.) were then added and the suspension incubated in a 37° C. water bath until it cleared. The samples were then processed as usual. Gels were stained with Coomassie blue or transferred to Nytran™ membrane by Western blotting in the Bio-Rad Semidry™ system (Bio-Rad Laboratories, Richmond, Calif.). For detection of native ClfB in *S. aureus*, blots were processed using the BM Chemiluminescence Detection System™ (POD) of Boehringer Mannheim, according to the manufacturer's instructions. Primary anti-ClfB antibody

was used at a 1/1000 dilution, for a two hour incubation at room temperature. Protein A conjugated with horse radish peroxidase (Sigma Chemical Co.) was used to detect bound antibody, diluted 1/2000 for a one hour incubation at room temperature. Blots requiring less sensitivity were treated in a similar way, except that 5% skim milk was used as a blocking agent, and the blots were developed using chloronaphthol and hydrogen peroxide.

To determine whether ClfB is cell wall-associated, whole cells from an exponential phase culture were treated with lysostaphin in buffer supplemented with 30% raffinose to stabilize the protoplasts. The protoplasts were harvested, and the protoplasts and supernatant analyzed separately by Western blotting. ClfB protein was detected only in the supernatant, indicating that all ClfB was cross-linked to the peptidoglycan, and could be released by lysostaphin without disruption of the protoplast.

ClfB expression was enhanced by growth in rich media, such as tryptone soy broth or brain heart infusion.

Several *S. aureus* strains known to contain clfB alleles were screened for ClfB production by Western blotting. Cultures were harvested in early exponential phase to maximize expression. Of the nine strains examined, 8325-4, RN4282, and V13 expressed immunoreactive antigens of similar size and intensity to that of Newman, whereas strains GH13 and P1 had very weak bands of this size. Strains P1, Cowan and M60 expressed smaller immunoreactive antigens which may be degradation products. Strains V8 and Phillips expressed no detectable ClfB protein. Strain RN4220, which was derived from 8325-4, expressed exceptionally high levels of ClfB.

Example 3

Immunoassay for ClfB Using Biotinylated Recombinant ClfB Region A

The DNA encoding region A of clfB (encoding residues S45 to N542) was amplified from genomic DNA of *S. aureus* Newman using the following primers: (SEQ ID NO:17)
Forward: 5' CGAAAGCTTGTCAGAACAATCGAAC-GATACAACG 3' (SEQ ID NO:16)
Reverse: 5' CGAGGATCCATTTACTGCTGAATCACC 3'

Cleavage sites for HindIII and BamHI (underlined) were appended to the 5' ends of the respective primers to facilitate cloning of the product into the His-tag expression vector pV4. Cloning employed *E. coli* JM101 as a host strain. The recombinant region A was purified by nickel affinity chromatography.

Enzyme Linked Immunosorbent Assay (ELISA)

Immulon 1™ plates (Dynatech™, Dynal, Inc., Great Neck, N.Y.) were coated overnight with 100 μ l of 10 μ g/ml human fibrinogen (Chromogenix). They were then blocked with 200 μ l of 5 mg/ml bovine serum albumin (BSA) for one hour. The plates were then incubated for three hours with 100 μ l biotinylated ClfB (His-tag recombinant region A) diluted to 0.1–10 μ g/ml. They were then given three five-minute washes with PBS containing 0.02% Tween 20 and 1 mg/ml BSA. The plates were then incubated for one hour with 100 μ l of a 1/10 000 dilution of streptavidin conjugated with alkaline phosphatase (Boehringer Mannheim, Indianapolis, Ind.), and washed as before. The plates were then developed for 30 minutes at 37° C. with 100 μ l of 1 M diethanolamine, pH 9.8, containing 1 mg/ml p-nitrophenyl phosphate (Sigma Chemical Co.). Plates coated with BSA only were used as negative controls. The absorbance was measured at 405 nm.

Western Affinity Blotting

A 20 μg quantity of human fibrinogen (Chromogenix) was subjected to SDS-PAGE on a 15% acrylamide gel for two hours. Proteins were transferred to nitrocellulose at 100 V for two hours. The membrane was blocked overnight in PBS containing 10% nonfat dry milk. The blot was then incubated with 2.5 $\mu\text{g}/\text{ml}$ biotinylated ClfB (His-tag recombinant region A) for one hour with shaking, the biotinylation being performed with EZ link-sulfo-NHS-LC-Biotin™ (Pierce, Rockford, Ill.). The blot was then given three five-minute washes in PBS containing 0.1% Tween 20. The blot was then incubated for one hour with avidin conjugated with horseradish peroxidase (Boehringer Mannheim) at a 1/200,000 dilution. The blot was then washed as before, and developed using the enhanced chemiluminescence system of Amersham (Little Chalfont, Bucks, UK). The band profile was compared with that obtained by subjecting fibrinogen to SDS-PAGE and Coomassie Blue staining.

In a Western affinity blot, in which biotinylated purified ClfB region A was used to probe blotted fibrinogen, a comparison with a lane of stained fibrinogen indicated that ClfB bound the alpha and beta-chains of fibrinogen. No bands were seen when ClfB was omitted. This experiment shows an important difference with ClfA, which is known to bind to the gamma-chain of fibrinogen.

Example 4

Mutagenesis of *clfB*

An insertion mutation in *claw* was created by introducing a fragment containing a Tc resistance marker into the middle of the gene on pA1-1EX as shown in FIG. 3. The 2.35-kb HindIII fragment from pT181 was filled in with Klenow enzyme, and blunt-end ligated into the HpaI site of pA1-1EX. Plasmid pTS2, with temperature sensitive replication and a Cm^r marker, was cloned into this construct at the SmaI site by cleaving with AvaI. This cloning step was carried out in *E. coli*, and transformants were selected on Ap and incubated at 30° C. to avoid selection of revertants to temperature independence. The plasmid was then purified and transformed into *S. aureus* RN4220 by electroporation and Tc^r transformants selected at 30° C. Five independent broth cultures grown at 30° C. were diluted 1/100 in fresh medium without antibiotics, and grown at 42° C. for six hours or 18 hours. The cultures were then diluted 1/100 and incubated at 42° C. for another time period. Six such dilutions and incubations were made, by which time Tc resistance had declined to approximately 1/1000 colony forming units (CFU). The cultures were then diluted to give approximately 100 CFU per plate on medium containing Tc, and incubated overnight at 37° C. Colonies which were Tc^r but Cm^s were presumed to have undergone a double cross-over event between the plasmid and host genome, leading to replacement of the wild-type gene with the mutated one, with subsequent loss of the plasmid. Five hundred colonies were screened per culture. Eleven presumptive mutants were isolated from four of the five cultures. Four representative mutants were selected and genomic DNA isolated. Mutant DU5944, deficient in both *clfA* and *clfB*, was constructed by transducing *clfA2::Tn917* from strain DU5876 into *clfB* mutant DU5943, selecting for Em^r.

To determine whether mutations known to affect exoprotein expression influenced *clfB*, strain 8325-4 and the agr mutant ISP546 were compared. No significant differences in the level or dynamics of ClfB expression were noted.

To determine the role of ClfB in bacteria-fibrinogen interactions, a *clfB* mutant of strain Newman was constructed by allele replacement as shown in FIG. 2. Genomic DNA of the mutant was digested with BamHI and subjected to Southern blotting with a labeled 1.3 kb HpaI fragment from plasmid pA1-1E containing the 5' half of *clfB* and about 150 bp of upstream sequence. A single band hybridized in each case, but as expected, the band was 2.35 kb longer in the mutant than in the wild-type. The mutation was initially isolated in RN4220 and then transduced into strain Newman, forming strain DU5943.

Overexpression of ClfB and Complementation of *clfB* Mutation

Overproduction of ClfB was enabled by subcloning a SmaI fragment containing the *clfB* gene and 500 bp of upstream DNA from pA1-1E into the high copy number shuttle plasmid pCU1. The construct was then transformed into strain RN4220 and transduced into strain Newman. Transductants were selected on Cm. Southern and Western blotting confirmed that the high copy number was maintained in strain Newman, and that ClfB was produced at higher than wild-type levels, indicating that the upstream DNA contained the promoter necessary for expression of the *clfB* gene. Transduction of the construct into *clfB* mutants restored ClfB synthesis to higher than wild-type levels. The construct was also transduced into *clfAclfB* double mutants for use in complementation studies.

To create a *clfAclfB* double mutant, a *clfA::Tn917* mutation was transferred by transduction from strain DU5876 into the *clfB::Tc^r* mutant DU5943, forming DU5944. The wild-type *clfB⁺* gene was cloned into shuttle plasmid pCU1 to give plasmid pA1-1EA, which was introduced into the *clfAclfB* mutant by transduction to test complementation. Western blotting with anti-ClfB serum showed that the ClfB protein was missing in mutant DU5943. It was expressed at a higher level than the wild-type in mutants carrying the complementing plasmid pA1-1EA, indicating overexpression of the protein due to gene dosage effect.

Example 5

ClfB Binding Assays

Clumping Assays

The role of ClfB in binding of *S. aureus* cells to soluble fibrinogen was investigated in clumping assays. Clumping assays were carried out in Sarstedt™ flat-bottomed multi-well test plates, using 50- μl volumes of human fibrinogen (Calbiochem Corp. (San Diego, Calif.) plasminogen free, >95% pure), diluted serially two-fold in PBS from a starting concentration of 1 mg/ml. *S. aureus* cultures were washed once in PBS, resuspended to a final OD₆₀₀ of 6, and 20 μl added to each well. Control wells contained PBS only. The plates were agitated briskly for five minutes and visually examined for clumping. The clumping titer was the lowest concentration of fibrinogen at which clumping occurred.

The results are set forth in Table 2, below. Results are the mean of concurrent duplicate assays.

TABLE 2

Strain	Clumping titer, $\mu\text{g/ml}$ fibrinogen	
	Exponential phase	Stationary phase
Wild-type	0.98	0.98
clfA	3.91	>1000.00
clfB	1.95	0.98
clfA clfB	>1000.00	>1000.00
clfA clfB (pA1-1EA; clfB ⁺)	2.93	250.00

The clumping titers of clfA and clfB single mutants were very similar to wild-type when exponential phase cultures were used. However, the double clfAclfB mutant failed to form clumps, even at the highest fibrinogen concentration. In contrast, the double mutant carrying the wild-type clfB gene on pA1-1EA formed clumps with almost the same avidity as the wild-type. These data show unambiguously that ClfB is a clumping factor.

The difference in clumping titer between the single mutants was much greater when stationary phase cultures were used, where only ClfA is present on cells. The wild-type strain and single clfB mutant had identical titers. The single clfA mutant failed to clump, and was thus indistinguishable from the double mutant. Interestingly, there was a slight restoration of clumping when the double mutant was complemented with the overexpressed clfB⁺ gene. This probably reflects over expression of the protein.

Plate Adherence Assays

To determine whether ClfB can promote bacterial attachment to immobilized fibrinogen, strains were tested for fibrinogen binding in a microtiter plate adherence assay. Binding of cells to fibrinogen immobilized on plates was measured by the assay of Wolz et al., *Infect. Immun.* 64:3142-3147 (1996). Fibrinogen was diluted in carbonate buffer (15 mM Na₂CO₃, 35 mM NaHCO₃, 3.2 μM Na₂N₃, pH 9.6) and 100 μl used to coat 96-well flat-bottomed ELISA plates (Immulon 4™, Dynatech) overnight at 4° C. Control wells contained carbonate buffer only. After washing in 150 mM NaCl, 0.05% Tween 20™ surfactant, the plates were blocked for one hour at 37° C. in 1% BSA, 0.05% Tween in PBS. After washing as before, 100 μl of a cell suspension (OD₆₀₀ of 0.4 in PBS) was added, and the plates incubated for two hours at 37° C. After gentle washing by hand, adherent cells were fixed by adding 100 μl of 25% aqueous formaldehyde, and incubating at room temperature for at least 30 minutes. The plates were then washed gently once more, stained with crystal violet, washed again, and the plates read by ELISA reader at 570 nm. To avoid inter-assay variation, experiments were designed so that a single plate provided a complete set of results.

The pattern of adherence strongly reflected that obtained in clumping assays (FIG. 15). Assays in which the concentration of cells was varied indicated that binding was approximately half the maximum value at a cell density of 0.4 OD (except for the double mutant), and this cell density was subsequently used routinely. Wild-type, clfA, clfB mutants and clfAclfB (pA1-1EA) showed a fibrinogen concentration-dependent increase in binding (FIG. 16). This

increase was less marked for the clfB mutant (expressing ClfA) than for the clfA mutant (expressing ClfB), suggesting that ClfB may be a less avid and/or abundant receptor. With stationary phase cells, the clfB mutant continued to behave like the wild-type strain, whereas the clfA mutant bound much less avidly. As with clumping, adherence was slightly higher with the complemented double mutant, presumably due to a gene dosage effect.

The clumping and adherence assays show that ClfB mediates binding both to soluble and immobilized fibrinogen, closely resembling the activity of ClfA.

The binding of increasing concentrations of biotinylated purified region A from ClfA and ClfB to solid phase fibrinogen was compared in a direct ELISA. The results are shown in FIG. 14. The adherence profiles of the two proteins were very similar, especially at the lower concentrations. At the highest concentration, binding of ClfA was approximately 50% greater than that of ClfB. Neither protein bound to BSA.

Effect of Anti-ClfB Antibody on Bacterial Adherence to Immobilized Fibrinogen

To study inhibition of fibrinogen binding by IgG, the cells used for the assay were preincubated with serial two-fold dilutions of purified IgG in PBS, starting with a concentration of 500 $\mu\text{g/ml}$. Preincubation was for two hours at 37° C. in Sarstedt™ multiwell test plates, and the cells were then transferred to ELISA plates coated with fibrinogen (2.5 $\mu\text{g/ml}$) and blocked as before. The rest of the assay was as before.

Cells from exponential phase cultures of wild-type and mutant Newman strains were preincubated with increasing concentrations of purified anti-ClfB IgG, and adherence to plastic surfaces coated with 2.5 $\mu\text{g/ml}$ fibrinogen examined. The results are shown in FIG. 17. Binding of the clfB mutant was not inhibited, and binding of wild-type cells was almost unaffected, even at the highest antibody concentration. However, binding of the clfA mutant showed an IgG concentration-dependent decrease, with an IC₅₀ of 16 $\mu\text{g/ml}$. The double mutant carrying clfB⁺ on a complementing plasmid was also inhibited by the antibody, although the IC₅₀ was higher (50 $\mu\text{g/ml}$), presumably because more ClfB was being expressed on the cell surface.

Effect of Divalent Cations on Bacterial Adherence to Immobilized Fibrinogen

The effect of metal ions on fibrinogen binding was studied in a similar manner, preincubating the cells with serial two-fold dilutions of MgCl₂, MnCl₂ or MgCl₂ in TBS (50 mM Tris HCl, pH 7.5, 150 mM NaCl), starting with a concentration of 50 mM. TBS was used instead of PBS, which causes precipitation of both calcium and manganese. Since the cells bound less well under these conditions, the starting cell concentration was doubled.

It is known that the interaction of ClfA and fibrinogen is inhibited by Ca²⁺ and Mn²⁺, but not Mg²⁺ ions. The effect of divalent cations on ClfB-promoted adherence to fibrinogen was thus tested. Preincubation of exponential phase cells of the wild-type strain and the clf mutants with CaCl₂ inhibited binding to fibrinogen. Those strains expressing ClfB alone showed greater sensitivity than the mutant expressing ClfA alone (clfB). The IC₅₀ for the wild-type strain and the clfB mutant were 17 and 14 mM, respectively, whereas for the clfA mutant and the clfB⁺ complemented double mutant the IC₅₀ was 1.05 and 0.60 mM, respectively. MnCl₂ also inhibited attachment of the wild-type strain and

mutants, with a stronger effect on strains expressing only clfB. The IC₅₀ for the wild-type and the clfB mutant was 3.3 and 6.4 mM, respectively, whereas for the clfA mutant and the double mutant carrying clfB+ on a complementing plasmid the IC₅₀ was 0.35 and 1.26 mM respectively. MgCl₂ had no effect on binding below 12.5 mM.

Thus, clfB promoted adherence of bacteria to immobilized fibrinogen is inhibited by Ca²⁺ and Mn²⁺ at similar concentrations to ClfA-promoted adherence. However, the mechanisms are likely to be different since ClfB does not contain a homologue of EF hand I implicated in Ca²⁺ promoted modulation of ClfA-fibrinogen interactions.

Platelet-fibrin Clot Adherence Assay

Adherence to platelet-fibrin clots was measured using a modification of an assay employed by Moreillon et al., *Infect. Immun.* 63: 4738–4743 (1995). Fresh canine blood was collected on 10% sodium citrate buffer (Sigma Chemical Co.), and centrifuged at 3000×g for 10 minutes at room temperature. The plasma fraction was removed and placed in a clean tube. Platelet-fibrin clots were made by mixing 0.5 ml volumes of plasma with 0.1 ml volumes of 0.2 mM CaCl₂ in 35 mm petri dishes. Thrombin (0.1 ml of 500 U/ml Sigma bovine thrombin) was then added, mixed in quickly, and the clots allowed to form. To measure bacterial adherence, 2 ml of PBS containing 5×10³ cfu/ml of bacteria (from a BHI-grown exponential phase culture) was added to each dish, and the dishes shaken for three minutes on an orbital shaker. The inoculum was drained off and the clots washed twice for five minutes each with 2 ml of PBS. The clots were then overlaid with 3 ml of molten TSA, incubated for 15 hours at 37° C., and the colonies counted. The bacterial suspension used as an inoculum was spread on TSA plates to obtain a total viable count, and the percentage of bound inoculum calculated. Results represent means of 6–10 plates per strain, and were analyzed statistically using the student's T test.

The clfB mutation reduced adherence when compared to the wild-type strain Newman, as did the clfA mutation which was previously shown by Moreillon et al. to have significantly reduced adherence in this model.

Assay for Adherence to Haemodialysis Tubing

In order to demonstrate that ClfB could serve as an adhesin for *S. aureus* in biomaterial-related infections, explanted human haemodialysis tubing was tested for promotion of bacterial adherence in vitro. The tubing was coated with a complex mixture of host plasma proteins including fibrinogen and fibronectin.

These experiments employed sections of haemodialysis tubing removed from patients 3 to 3.5 hours after implantation. Cultures were grown for two hours with shaking. Results, showing means with SEM of three experiments, are shown in FIG. 19.

Assay for Adherence to Fibrinogen-coated PMMA Coverslips

Adherence of *S. aureus* Newman and mutants to fibrinogen-coated polymethylmethacrylate (PMMA) coverslips was measured as described by Greene et al., *Mol. Microbiol.* 17:1143–1152 (1995), except that the coverslips were coated with pure fibrinogen (1 μg/ml). Cultures for the assay were grown for two hours with shaking. Results, showing the means and SEM of triplicate experiments, are shown in FIG. 18.

The pattern of adherence to the tubing segments resembled the pattern of binding seen for immobilized

fibrinogen in a parallel assay for adherence to fibrinogen immobilized on PMMA coverslips. The single clfA mutants had slightly lower levels of adherence compared to the wild-type whereas the double clfAclfB mutant was reduced to approximately 30% of wild-type level. Complementation of the single clfB mutant with the clfB gene on pA1-1EA restored binding to greater than wild-type levels, whereas complementation of the double mutant with the same plasmid restored binding only to the same level as the single clfA mutant.

Example 6

ClfB as a Virulence Factor in Experimental Endocarditis

Clumping factor A was shown to be a virulence factor promoting adherence to damaged heart valves in the rat model of experimental endocarditis of Moreillon et al., *Infect. Immun.* 63:4738–4743 (1995). Therefore, the role of ClfB in this infection was tested by comparing the infection rate of a clfB mutant and the mutant carrying the complementing clfB+ plasmid. Rats were infected intravenously at an ID₆₀ with 5×10³ cfu. 61% of the wild-type control animals' valves were infected (n=13), whereas only 30% of the clfB mutant infected animals were colonized (n=20). In contrast 77% (n=9) of the complemented mutant became infected. This clearly shows that ClfB is an adhesin and potential virulence factor in the endocarditis model.

Example 7

Generation of TYTFTDYVD (SEQ ID NO:18) Peptide Antibodies

The nanopptide, TYTFTDYVD (SEQ ID NO:18), was synthesized in multiple antigen peptide format (MAP; Research Genetics, Inc., Huntsville, Ala.). The peptide was conjugated to KLH according to manufacturers' directions (Pierce). Two female New Zealand White rabbits were immunized subcutaneously with the KLH-TYTFTDYVD (SEQ ID NO:18) conjugate emulsified with Freund's Complete Adjuvant. The rabbits were boosted 3 weeks later by subcutaneous injection of KLH-TYTFTDYVD (SEQ ID NO: 18) adjuvanted with Freund's Incomplete. A third boost was administered subcutaneously with KLH-TYTFTDYVD (SEQ ID NO:18) in PBS. The animals were analyzed for TYTFTDYVD (SEQ ID NO:18) specific antibodies 21 days after the final boost. For purification of antibodies, antisera was diluted 1:1 with Tris-HCl pH 8.0 and passed over a Protein A-Sepharose® column. After sequential washes with Tris-HCl pH 8.0, 0.5 M sodium chloride, the bound antibodies were eluted in 3.5 M MgCl₂, and dialyzed into PBS.

Immulon-2 microtiter plates (Dynex Technologies, Chantilly, Va.) were coated for 2 hr at room temperature with 1 μg ClfA, ClfB, or BSA. The protein coated plates were washed three times with PBS, 0.05% Tween 20 and then blocked with PBS, 1% BSA. The blocked plates were washed three times with PBS, 0.05% Tween 20. Fifty μl of the purified rabbit KLH-TYTFTDYVD (SEQ ID NO:18) antibodies were serially diluted in PBS and added to the microtiter plate and incubated at 25° C. on a rocker platform. The wells were washed three times with PBS, 0.05% Tween 20 and the secondary antibody was added to the wells and incubated for 1 hr at room temperature. The secondary antibody was alkaline phosphatase-conjugated goat anti-rabbit IgG (Bio-Rad), diluted 3000-fold in PBS. ELISA plates were developed for 1 hr at 37° C. with 1 mg/ml

p-nitrophenyl phosphate (Sigma) in 1 M diethanolamine, 0.5 mM MgCl₂, pH 9.8, and quantified at 405 nm on a Perkin Elmer HTS 7000 Bio-Assay reader.

The data is shown in FIG. 21. These data indicate that the anticonsensus sequence TYTFTDYVD (SEQ ID NO: 18) antibodies significantly bind to ClfA and ClfB proteins, but not the control protein, BSA.

Example 8

Passive Immunization With Rabbit ClfB IgG

The DNA encoding region A of clfB (encoding residues S45 to N542) was amplified from genomic DNA of *S. aureus* Newman using the following primers: (SEQ ID NO:17) Forward: 5' CGAAAGCTTGTTCAGAACAAATCGAAC-GATACAACG 3' (SEQ ID NO:16)

Reverse: 5' CGAGGATCCATTACTGCTGAATCACC 3'

Cleavage sites for HindIII and BamHI (underlined) were appended to the 5' ends of the respective primers to facilitate cloning of the product into the His-tag expression vector pV4. Cloning employed *E. coli* JM101 as a host strain. The recombinant region A was purified by nickel affinity chromatography. Antibodies were raised in rabbits with the purified recombinant A region according to standard procedures. Anti-ClfB A region IgG was purified by affinity chromatography on a Protein A sepharose column.

Twenty Swiss Webster mice (23–28 g) were used to determine if passive immunization with purified rabbit anti-ClfB A region IgG could prevent infection mediated by a methicillin resistant *S. aureus*.

Methicillin resistant *S. aureus* strain 601 was cultured on blood agar plates. A single colony was then inoculated into 10 mls of BHI broth and incubated at 37° C. overnight. The culture was diluted to a 1:100 dilution, placed into 10 ml of fresh BHI and grown to an optical density (O.D.) of 1.5–2.0. The culture was then centrifuged and washed in 1xPBS. The culture was re-suspended in 1xPBS containing 5% BSA and 10% dimethyl sulfoxide (DMSO) and kept frozen at –20° C. The bacterial solution was thawed, washed, diluted in PBS, and adjusted to the appropriate concentrations before dosing the mice.

The mice were divided into four treatment groups (5 mice per treatment group). Mice were assigned to treatment groups as follows:

Antibody/Bacteria	Dose CFU/mouse	No. of Mice
1 Normal rabbit IgG/ <i>S. aureus</i>	3.81 × 10 ⁷	5
2 Normal rabbit IgG/ <i>S. aureus</i>	7.62 × 10 ⁷	5
3 Rabbit anti-ClfB IgG/ <i>S. aureus</i>	3.81 × 10 ⁷	5
4 Rabbit anti-ClfB IgG/ <i>S. aureus</i>	7.62 × 10 ⁷	5

On day -1, ten mice were administered 10 mg rabbit anti-ClfB region A IgG and 10 mice were given 10 mg normal rabbit IgG. Both antibodies were given via intraperitoneal (i.p.) injection. On day 0, all mice were infected intravenously (i.v.) with either 3.81×10⁷ CFU *S. aureus* or 7.62×10⁷ CFU *S. aureus*.

Systemic infection was measured by evaluation of body weight loss. Body weight loss is one of the primary parameters that is evaluated when cases of illness and injury are being assessed in mice. The body weight of each animal was recorded on Day -1 and every other day thereafter, including terminal sacrifice. The animals were weighed to the nearest 0.1 gram.

Mice injected with normal rabbit IgG displayed a significantly larger weight loss at the end of the experiment

compared to mice passively immunized with rabbit anti-ClfB region A IgG (see table below). In addition, pathological evaluation of the mice at necropsy revealed a greater number of lesions and foci of infection in the kidneys from the mice receiving normal rabbit IgG compared to the kidneys from mice that were immunized with anti-ClfB region A IgG.

Day of Study	% Change in body weight (mean)			
	Normal IgG/ <i>S. aureus</i>	Anti-ClfB IgG/ <i>S. aureus</i>	Normal IgG/ <i>S. aureus</i>	Anti-ClfB IgG/ <i>S. aureus</i>
–1	3.81 × 10 ⁷	3.81 × 10 ⁷	7.62 × 10 ⁷	7.62 × 10 ⁷
1	0	0	0	0
3	2.9	3.6	3.9	5.8
5	10	5.1	8.5	8.2
5	8.3	1.5	8.0	6.6

Example 9

CUB Region A Binds α and β chains of Human Fibrinogen

Human fibrinogen (20 μg; Chromogenix) was separated by SDS-PAGE on a 15% acrylamide gel for 2 hours. Proteins were transferred to nitrocellulose at 100 V for 2h. The membranes were blocked overnight in PBS containing 10% non-fat dry milk and then incubated with 2.5 μg/ml biotinylated ClfB or ClfA region A protein for 1h with shaking. They were then given 3x5 mm washes with PBS containing 0.1% Tween 20 and incubated for 1 hr with avidin conjugated horseradish peroxidase (Boehringer Mannheim: 1:100,000 dilution). The filters were washed as before and developed using enhanced chemiluminescence (Amersham). The Western Blot (FIG. 22) illustrates the binding of biotinylated ClfA to the γ chain of fibrinogen and the binding of biotinylated ClfB to the α and β chains of fibrinogen.

Example 10

ClfB Region A Binds 75 kD and 50 kD Proteins From Human Rhabdomyosarcoma Cell Line

Human Rhabdomyosarcoma Cells were lysed with the SDS-PAGE running buffer and varying amounts (2–10 μd) of the protein lysate were separated by SDS-PAGE on a 15% acrylamide gel for 2h. Proteins were transferred to nitrocellulose at 100 V for 2h. The membranes were blocked overnight in PBS containing 10% non-fat dry milk and then incubated with 2.5 μg/ml biotinylated ClfB or ClfA region A protein for 1 hr with constant shaking. They were then given 3x5 min washes with PBS containing 0.1% Tween 20 and incubated for 1 hr with avidin conjugated horseradish peroxidase (Boehringer Mannheim; 1:100,000 dilution). The filters were washed as before and developed using enhanced chemiluminescence (Amersham). Two major bands were seen at 50 kD and 75 kD that reacted with the biotinylated ClfB region A protein.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1

<211> LENGTH: 918

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 1

```

Asn Gly Val Ile Phe Leu Lys Lys Arg Ile Asp Tyr Leu Ser Asn Lys
 1           5           10          15
Gln Asn Lys Tyr Ser Ile Arg Arg Phe Thr Val Gly Thr Thr Ser Val
          20          25          30
Ile Val Gly Ala Thr Ile Leu Phe Gly Ile Gly Asn His Gln Ala Gln
          35          40          45
Ala Ser Glu Gln Ser Asn Asp Thr Thr Gln Ser Ser Lys Asn Asn Ala
 50          55          60
Ser Ala Asp Ser Glu Lys Asn Asn Met Ile Glu Thr Pro Gln Leu Asn
 65          70          75          80
Thr Thr Ala Asn Asp Thr Ser Asp Ile Ser Ala Asn Thr Asn Ser Ala
          85          90          95
Asn Val Asp Ser Thr Thr Lys Pro Met Ser Thr Gln Thr Ser Asn Thr
          100         105         110
Thr Thr Thr Glu Pro Ala Ser Thr Asn Glu Thr Pro Gln Pro Thr Ala
          115         120         125
Ile Lys Asn Gln Ala Thr Ala Ala Lys Met Gln Asp Gln Thr Val Pro
          130         135         140
Gln Glu Gly Asn Ser Gln Val Asp Asn Lys Thr Thr Asn Asp Ala Asn
          145         150         155         160
Ser Ile Ala Thr Asn Ser Glu Leu Lys Asn Ser Gln Thr Leu Asp Leu
          165         170         175
Pro Gln Ser Ser Pro Gln Thr Ile Ser Asn Ala Gln Gly Thr Ser Lys
          180         185         190
Pro Ser Val Arg Thr Arg Ala Val Arg Ser Leu Ala Val Ala Glu Pro
          195         200         205
Val Val Asn Ala Ala Asp Ala Lys Gly Thr Asn Val Asn Asp Lys Val
          210         215         220
Thr Ala Ser Asn Phe Lys Leu Glu Lys Thr Thr Phe Asp Pro Asn Gln
          225         230         235         240
Ser Gly Asn Thr Phe Met Ala Ala Asn Phe Thr Val Thr Asp Lys Val
          245         250         255
Lys Ser Gly Asp Tyr Phe Thr Ala Lys Leu Pro Asp Ser Leu Thr Gly
          260         265         270
Asn Gly Asp Val Asp Tyr Ser Asn Ser Asn Asn Thr Met Pro Ile Ala
          275         280         285
Asp Ile Lys Ser Thr Asn Gly Asp Val Val Ala Lys Ala Thr Tyr Asp
          290         295         300
Ile Leu Thr Lys Thr Tyr Thr Phe Val Phe Thr Asp Tyr Val Asn Asn
          305         310         315         320
Lys Glu Asn Ile Asn Gly Gln Phe Ser Leu Pro Leu Phe Thr Asp Arg
          325         330         335
Ala Lys Ala Pro Lys Ser Gly Thr Tyr Asp Ala Asn Ile Asn Ile Ala
          340         345         350

```

-continued

Asp	Glu	Met	Phe	Asn	Asn	Lys	Ile	Thr	Tyr	Asn	Tyr	Ser	Ser	Pro	Ile
		355					360					365			
Ala	Gly	Ile	Asp	Lys	Pro	Asn	Gly	Ala	Asn	Ile	Ser	Ser	Gln	Ile	Ile
	370					375					380				
Gly	Val	Asp	Thr	Ala	Ser	Gly	Gln	Asn	Thr	Tyr	Lys	Gln	Thr	Val	Phe
385					390					395					400
Val	Asn	Pro	Lys	Gln	Arg	Val	Leu	Gly	Asn	Thr	Trp	Val	Tyr	Ile	Lys
				405					410					415	
Gly	Tyr	Gln	Asp	Lys	Ile	Glu	Glu	Ser	Ser	Gly	Lys	Val	Ser	Ala	Thr
			420					425					430		
Asp	Thr	Lys	Leu	Arg	Ile	Phe	Glu	Val	Asn	Asp	Thr	Ser	Lys	Leu	Ser
		435					440					445			
Asp	Ser	Tyr	Tyr	Ala	Asp	Pro	Asn	Asp	Ser	Asn	Leu	Lys	Glu	Val	Thr
	450					455					460				
Asp	Gln	Phe	Lys	Asn	Arg	Ile	Tyr	Tyr	Glu	His	Pro	Asn	Val	Ala	Ser
465					470					475					480
Ile	Lys	Phe	Gly	Asp	Ile	Thr	Lys	Thr	Tyr	Val	Val	Leu	Val	Glu	Gly
			485						490					495	
His	Tyr	Asp	Asn	Thr	Gly	Lys	Asn	Leu	Lys	Thr	Gln	Val	Ile	Gln	Glu
			500					505					510		
Asn	Val	Asp	Pro	Val	Thr	Asn	Arg	Asp	Tyr	Ser	Ile	Phe	Gly	Trp	Asn
		515					520					525			
Asn	Glu	Asn	Val	Val	Arg	Tyr	Gly	Gly	Gly	Ser	Ala	Asp	Gly	Asp	Ser
	530					535					540				
Ala	Val	Asn	Pro	Lys	Asp	Pro	Thr	Pro	Gly	Pro	Pro	Val	Asp	Pro	Glu
545					550					555					560
Pro	Ser	Pro	Asp	Pro	Glu	Pro	Glu	Pro	Thr	Pro	Asp	Pro	Glu	Pro	Ser
			565						570					575	
Pro	Asp	Pro	Glu	Pro	Glu	Pro	Ser	Pro	Asp	Pro	Asp	Pro	Asp	Ser	Asp
			580					585					590		
Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp
		595					600					605			
Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	610						615					620			
Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp
625					630					635					640
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			645						650					655	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp
			660					665					670		
Ser	Asp	Ser	Glu	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			675				680					685			
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	690					695				700					
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
705					710					715					720
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			725						730					735	
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			740					745					750		
Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
			755				760						765		

-continued

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 770 775 780

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 785 790 795 800

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 805 810 815

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 820 825 830

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Arg Val Thr
 835 840 845

Pro Pro Asn Asn Glu Gln Lys Ala Pro Ser Asn Pro Lys Gly Glu Val
 850 855 860

Asn His Ser Asn Lys Val Ser Lys Gln His Lys Thr Asp Ala Leu Pro
 865 870 875 880

Glu Thr Gly Asp Lys Ser Glu Asn Thr Asn Ala Thr Leu Phe Gly Ala
 885 890 895

Met Met Ala Leu Leu Gly Ser Leu Leu Leu Phe Arg Lys Arg Lys Gln
 900 905 910

Asp His Lys Glu Lys Ala
 915

<210> SEQ ID NO 2
 <211> LENGTH: 2969
 <212> TYPE: DNA
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 2

tagaaattga aatggagtaa tatttttgaa aaaaagaatt gattatttgt cgaataagca 60
 gaataagtat tcgattagac gttttacagt aggtaccaca tcagtaatag taggggcaac 120
 tatactatatt gggataggca atcatcaagc acaagcttca gaacaatcga acgatacaac 180
 gcaatcttcg aaaaataatg caatgacaga ttccgaaaaa aacaatatga tagaaacacc 240
 tcaattaaat acaacggcta atgatacatc tgatattagt gcaaacacaa acagtgcgaa 300
 tgtagatagc acaacaaaac caatgtctac acaaacgagc aataccacta caacagagcc 360
 agcttcaaca aatgaaacac ctcaaccgac ggcaattaa aatcaagcaa ctgctgcaaa 420
 aatgcaagat caaactgttc ctcaagaagg aaattctcaa gtagataata aaacaacgaa 480
 tgatgctaata agcatagcaa caaacagtga gcttaaaaaa tctcaaacat tagatttacc 540
 acaatcatca ccacaaacga tttccaatgc gcaaggaact agtaaaccaa gtggttagaac 600
 gagagctgta cgtagtttag ctgttgctga accggtagta aatgctgctg atgctaaaagg 660
 taaaaatgta aatgataaag ttacggcaag taatttcaag ttagaaaaga ctacatttga 720
 ccctaataca agtggttaaca catttatggc ggcaaathtt acagtgcagc ataaagttaa 780
 atcaggggat tattttacag cgaagttacc agatagttaa actggtaaat gagacgtgga 840
 ttatttctaat tcaataata cgatgccaat tgcagacatt aaaagtacga atggcgtatgt 900
 tgtagctaaa gcaacatatg atatcttgac taagacgtat acatttctct ttacagatta 960
 tgtaaataat aaagaaaata ttaacggaca attttcatta cctttattta cagaccgagc 1020
 aaaggcacct aaatcaggaa catatgatgc gaatattaat attgctgatg aaatgtttaa 1080
 taataaaaatt acttataact atagttcgcc aattgcagga attgataaac caaatggcgc 1140
 gaacatttct tctcaaatata ttgggtgtaga tacagcttca ggtcaaaaca catacaagca 1200
 aacagtatht gttaacccta agcaacagat tttaggtaat acgtgggtgt atattaagg 1260

-continued

```

ctaccaagat aaaatcgaag aaagtagcgg taaagtaagt gctacagata caaaactgag 1320
aatttttgaa gtgaatgata catctaaatt atcagatagc tactatgcag atccaaatga 1380
ctctaacctt aaagaagtaa cagaccaatt taaaaataga atctattatg agcatccaaa 1440
tgtagctagt attaaatttg gtgatattac taaaacatat gtagtattag tagaagggca 1500
ttacgacaat acaggtgaaga acttaaaaac tcagggttatt caagaaaatg ttgatcctgt 1560
aacaataga gactacagta ttttcggttg gaataatgag aatgttgtac gttatggtgg 1620
tggagtgct gatggtgatt cagcagtaaa tccgaaagac ccaactccag ggccgccggt 1680
tgaccagaa ccaagtccag acccagaacc agaaccaacg ccagatccag aaccaagtcc 1740
agaccagaa ccggaaccaa gcccagacc ggatccggat tcggattcag acagtgactc 1800
aggctcagac agcgactcag gttcagatag cgactcagaa tcagatagcg attcggattc 1860
agacagtgat tcagattcag acagcgactc agaatcagat agcgattcag aatcagatag 1920
cgactcagat tcagatagcg attcagattc agatagcgat tcagattcag atagcgattc 1980
ggattcagac agtgattcag attcagacag cgactcagaa tcagatagcg actcagaatc 2040
agatagtgag tcagattcag acagtgactc ggactcagac agtgattcag actcagatag 2100
cgattcagac tcagatagcg attcagattc agacagcgac tcagattcag acagcgactc 2160
agactcagat agcgactcag actcagacag cgactcagat tcagatagcg attcagactc 2220
agacagcgac tcagactcag acagcgactc agactcagat agcgactcag attcagatag 2280
cgattcagac tcagacagcg actcagattc agatagcgat tcggactcag acagcgattc 2340
agattcagac agcgactcag actcggatag gattcagat tcagatagcg attcggattc 2400
agacagtgat tcagattcag acagcgactc agactcggat agcgactcag actcagacag 2460
cgattcagac tcagatagcg actcagactc ggatagcgac tcggattcag atagcgactc 2520
agactcagat agtgactccg attcaagatg tacaccacca aataatgaac agaaagcacc 2580
atcaaatoct aaagtggaag taaaccattc taataaggta tcaaaacaac acaaaactga 2640
tgctttacca gaaacaggag ataagagcga aaacacaaat gcaactttat ttggtgcaat 2700
gatggcatta ttaggatcat tactattggt tagaaaacgc aagcaagatc ataaagaaaa 2760
agcgtaaata ctttttagg ccgaatacat ttgtattcgg ttttttggtt gaaaatgatt 2820
ttaaagtga ttgattaagc gtaaaatggt gataaagtag aattagaag gggtcatgac 2880
gtatggctta tatttcatta aactatcatt caccaacaat tggtatgcat caaaatttga 2940
cagtcatttt accggaagaa cgagaattc 2969

```

```

<210> SEQ ID NO 3
<211> LENGTH: 930
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 3

```

```

Met Asn Asn Lys Lys Thr Ala Thr Asn Arg Lys Gly Met Ile Pro Asn
  1             5             10             15
Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Ser Val Gly Thr Ala Ser
             20             25             30
Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Ser Gly His Glu Ala
             35             40             45
Lys Ala Ala Glu His Thr Asn Gly Glu Leu Asn Gln Ser Lys Asn Glu
             50             55             60

```

-continued

Thr	Thr	Ala	Pro	Ser	Glu	Asn	Lys	Thr	Thr	Lys	Lys	Val	Asp	Ser	Arg	65	70	75	80
Gln	Leu	Lys	Asp	Asn	Thr	Gln	Thr	Ala	Thr	Ala	Asp	Gln	Pro	Lys	Val	85	90	95	
Thr	Met	Ser	Asp	Ser	Ala	Thr	Val	Lys	Glu	Thr	Ser	Ser	Asn	Met	Gln	100	105	110	
Ser	Pro	Gln	Asn	Ala	Thr	Ala	Asn	Gln	Ser	Thr	Thr	Lys	Thr	Ser	Asn	115	120	125	
Val	Thr	Thr	Asn	Asp	Lys	Ser	Ser	Thr	Thr	Tyr	Ser	Asn	Glu	Thr	Asp	130	135	140	
Lys	Ser	Asn	Leu	Thr	Gln	Ala	Lys	Asp	Val	Ser	Thr	Thr	Pro	Lys	Thr	145	150	155	160
Thr	Thr	Ile	Lys	Pro	Arg	Thr	Leu	Asn	Arg	Met	Ala	Val	Asn	Thr	Val	165	170	175	
Ala	Ala	Pro	Gln	Gln	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	His	Phe	Ser	180	185	190	
Asn	Ile	Asp	Ile	Ala	Ile	Asp	Lys	Gly	His	Val	Asn	Gln	Thr	Thr	Gly	195	200	205	
Lys	Thr	Glu	Phe	Trp	Ala	Thr	Ser	Ser	Asp	Val	Leu	Lys	Leu	Lys	Ala	210	215	220	
Asn	Tyr	Thr	Ile	Asp	Asp	Ser	Val	Lys	Glu	Gly	Asp	Thr	Phe	Thr	Phe	225	230	235	240
Lys	Tyr	Gly	Gln	Tyr	Phe	Arg	Pro	Gly	Ser	Val	Arg	Leu	Pro	Ser	Gln	245	250	255	
Thr	Gln	Asn	Leu	Tyr	Asn	Ala	Gln	Gly	Asn	Ile	Ile	Ala	Lys	Gly	Ile	260	265	270	
Tyr	Asp	Ser	Thr	Thr	Asn	Thr	Thr	Thr	Tyr	Thr	Phe	Thr	Asn	Tyr	Val	275	280	285	
Asp	Gln	Tyr	Thr	Asn	Val	Arg	Gly	Ser	Phe	Glu	Gln	Val	Ala	Phe	Ala	290	295	300	
Lys	Arg	Lys	Asn	Ala	Thr	Thr	Asp	Lys	Thr	Ala	Tyr	Lys	Met	Glu	Val	305	310	315	320
Thr	Leu	Gly	Asn	Asp	Thr	Tyr	Ser	Glu	Glu	Ile	Ile	Val	Asp	Tyr	Gly	325	330	335	
Asn	Lys	Lys	Ala	Gln	Pro	Leu	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Asn	340	345	350	
Glu	Asp	Leu	Ser	Arg	Asn	Met	Thr	Ala	Tyr	Val	Asn	Gln	Pro	Lys	Asn	355	360	365	
Thr	Tyr	Thr	Lys	Gln	Thr	Phe	Val	Thr	Asn	Leu	Thr	Gly	Tyr	Lys	Phe	370	375	380	
Asn	Pro	Asn	Ala	Lys	Asn	Phe	Lys	Ile	Tyr	Glu	Val	Thr	Asp	Gln	Asn	385	390	395	400
Gln	Phe	Val	Asp	Ser	Phe	Thr	Pro	Asp	Thr	Ser	Lys	Leu	Lys	Asp	Val	405	410	415	
Thr	Asp	Gln	Phe	Asp	Val	Ile	Tyr	Ser	Asn	Asp	Asn	Lys	Thr	Ala	Thr	420	425	430	
Val	Asp	Leu	Met	Lys	Gly	Gln	Thr	Ser	Ser	Asn	Lys	Gln	Tyr	Ile	Ile	435	440	445	
Gln	Gln	Val	Ala	Tyr	Pro	Asp	Asn	Ser	Ser	Thr	Asp	Asn	Gly	Lys	Ile	450	455	460	
Asp	Tyr	Thr	Leu	Asp	Thr	Asp	Lys	Thr	Lys	Tyr	Ser	Trp	Ser	Asn	Ser	465	470	475	480

-continued

Tyr	Ser	Asn	Val	Asn	Gly	Ser	Ser	Thr	Ala	Asn	Gly	Asp	Gln	Lys	Lys
				485					490					495	
Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys
			500					505					510		
Gln	Asp	Ala	Asn	Glu	Lys	Gly	Ile	Lys	Gly	Val	Tyr	Val	Ile	Leu	Lys
		515					520					525			
Asp	Ser	Asn	Gly	Lys	Glu	Leu	Asp	Arg	Thr	Thr	Thr	Asp	Glu	Asn	Gly
	530					535						540			
Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Ser	Asn	Gly	Thr	Tyr	Ser	Val	Glu	Phe
545					550					555					560
Ser	Thr	Pro	Ala	Gly	Tyr	Thr	Pro	Thr	Thr	Ala	Asn	Val	Gly	Thr	Asp
				565					570					575	
Asp	Ala	Val	Asp	Ser	Asp	Gly	Leu	Thr	Thr	Thr	Gly	Val	Ile	Lys	Asp
			580					585					590		
Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr
		595					600					605			
Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser	Asn	Lys	Asp	Gly	Lys	Arg
	610					615					620				
Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	Gly	Val	Lys	Val	Thr	Leu	Gln	Asn
625					630					635					640
Glu	Lys	Gly	Glu	Val	Ile	Gly	Thr	Thr	Glu	Thr	Asp	Glu	Asn	Gly	Lys
			645						650					655	
Tyr	Arg	Phe	Asp	Asn	Leu	Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu
		660						665					670		
Lys	Pro	Ala	Gly	Leu	Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp
		675				680						685			
Lys	Asp	Ala	Asp	Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp
	690					695					700				
Asp	Phe	Thr	Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser
705					710					715					720
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			725					730						735	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			740					745						750	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asn	Ser
		755					760					765			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	770					775					780				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
785					790					795					800
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			805					810						815	
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			820					825					830		
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
		835					840					845			
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	850					855					860				
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Asn	Asp	Ser	Asp	Ser
865					870					875					880
Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Ala	Lys	Pro
			885						890						895

-continued

Met Ser Thr Val Lys Asp Gln His Lys Thr Ala Lys Ala Leu Pro Glu
 900 905 910
 Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn Gly Thr Leu Phe Gly Gly
 915 920 925
 Leu Phe
 930

<210> SEQ ID NO 4
 <211> LENGTH: 2841
 <212> TYPE: DNA
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 4

atgaataata aaaagacagc acaaataga aaaggcatga taccaaatcg attaaacaaa 60
 ttttcgataa gaaagtattc tgtaggtact gcttcaatth tagtagggac aacattgatt 120
 tttgggtaa gtggatcatga agctaaagcg gcagaacata cgaatggaga attaaatcaa 180
 tcaaaaaatg aaacgacagc cccaagttag aataaaacaa ctaaaaaagt tgatagtcgt 240
 caactaaaag acaatacagc aactgcaact gcagatcagc cttaaagtgc aatgagtgat 300
 agtgcaacag ttaaagaac tagtagtaac atgcaatcac cacaaaacgc tacagctaat 360
 caatctacta caaaaactag caatgtaaca acaaatgata aatcatcaac tacatatagt 420
 aatgaaactg ataaaagtaa tttaacacaa gcaaaagatg tttcaactac acctaaaaca 480
 acgactatta aaccaagaac tttaaatcgc atggcagtga atactgttg agctccacaa 540
 caaggaacaa atgttaataa taaagtacat tttcaaata ttgacattgc gattgataaa 600
 ggacatgtta atcagactac tggtaaaact gaattttggg caacttcaag tgatgtttta 660
 aaattaaaag caaattacac aatcgatgat tctgttaaag agggcgatc atttactttt 720
 aaatatggtc aatatttccg tccaggatca gtaagattac cttcacaaac tcaaaattta 780
 tataatgcc aaggaatat tattgcaaaa ggtatttatg atagtacaac aaacacaaca 840
 acatataact ttacgaacta tgtagatcaa tacaacaaat ttagaggtag ctttgaacaa 900
 gttgcatttg cgaacgtaa aaatgcaaca actgataaaa cagcttataa aatggaagta 960
 actttaggta atgatacata tagcgaagaa atcattgtcg attatggtaa taaaaagca 1020
 caaccgcta tttcaagtac aaactatatt aacaatgaag atttatcgcg taatatgact 1080
 gcatatgtaa atcaacctaa aaatacatat actaaacaaa cgtttgttac taatttaact 1140
 ggatataaat ttaatccaaa tgcaaaaaac ttcaaaatth acgaagtgc agatcaaaat 1200
 caattttggg atagtttcac cctgatact tcaaaactta aagatgttac tgatcaattc 1260
 gatgttatt atagtaataa taataaaaca gctacagtcg atttaatgaa aggcacaaaca 1320
 agcagcaata aacaatacat cattcaacaa gttgcttacc cagataatag ttcaacagat 1380
 aatggaaaaa ttgattatac tttagacact gacaaaacta aatatagttg gtcaaatagt 1440
 tattcaaatg tgaatggctc atcaactgct aatggcgacc aaaagaata taatctaggt 1500
 gactatgtat ggaagatc aaataaagat ggtaaacaaag atgccaatga aaaagggatt 1560
 aaaggtgttt atgtcattct taaagatagt aacggtaaag aattagatcg tacgacaaca 1620
 gatgaaaatg gtaaatatca gttcactggg ttaagcaatg gaacttatag tgtagagttt 1680
 tcaacaccag ccggttatac accgacaact gcaaatgtag gtacagatga tgctgtagat 1740
 tctgatggac taactacaac aggtgtcatt aaagacgctg acaacatgac attagatagt 1800
 ggattctaca aaacacaaa atatagttha ggtgattatg tttggtacga cagtaataaa 1860

-continued

```

gatggtaaac gagattcgac tgaaaaagga attaaaggtg ttaaagttac ttgcaaaac 1920
gaaaaaggcg aagtaattgg tacaactgaa acagatgaaa atggtaaata cgcctttgat 1980
aatttagata gtggtaaata caaagttatc ttgaaaaac ctgctggctt aactcaaaca 2040
ggtacaaata caactgaaga tgataaagat gccgatggtg gcgaagtga tgtaacaatt 2100
acggatcatg atgatttcac acttgataat ggctactacg aagaagaac atcagatagc 2160
gactcagatt ctgacagcga ttcagactca gatagcgact cagattcaga tagcgactca 2220
gattcagaca gcgattcaga cagcgactca gactcagata gcgattcaga ttcagacagc 2280
gactcagact cagacagcaa ttcagactcg gatagcgact cagactcaga tagcgactca 2340
gattcggata gcgactcaga ctcagatagc gattcagatt cagatagcga ttcggactca 2400
gacagtgatt cagattcaga ctcagatagc gactcagatt ctgacagcga ttcagactca 2460
gacagcgact cagactcaga cagtgattca gattcagaca gcgactcaga ttcagatagc 2520
gactcagact cagatagcga ctcagactca gatagcgact cagactcaga tagcgattca 2580
gattcagaca gcgactcaga ttcagatagc gattcggact cagacaacga ctcagattca 2640
gatagcgatt cagattcaga tgcaggtaaa catactccgg ctaaaccaat gagtacggtt 2700
aaagatcagc ataaaacagc taaagcatta ccagaaacag gtagtgaana taataattca 2760
aataatggca cattattcgg tggattatc gcggcattag gatcattatt gtcattcggg 2820
cgtcgtaaaa aacaaaataa a 2841

```

<210> SEQ ID NO 5

<211> LENGTH: 1315

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 5

```

Met Leu Asn Arg Glu Asn Lys Thr Ala Ile Thr Arg Lys Gly Met Val
 1             5             10            15
Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr
          20             25             30
Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln
          35             40             45
Glu Ala Lys Ala Ala Glu Ser Thr Asn Lys Glu Leu Asn Glu Ala Thr
          50             55             60
Thr Ser Ala Ser Asp Asn Gln Ser Ser Asp Lys Val Asp Met Gln Gln
          65             70             75             80
Leu Asn Gln Glu Asp Asn Thr Lys Asn Asp Asn Gln Lys Glu Met Val
          85             90             95
Ser Ser Gln Gly Asn Glu Thr Thr Ser Asn Gly Asn Lys Leu Ile Glu
          100            105            110
Lys Glu Ser Val Gln Ser Thr Thr Gly Asn Lys Val Glu Val Ser Thr
          115            120            125
Ala Lys Ser Asp Glu Gln Ala Ser Pro Lys Ser Thr Asn Glu Asp Leu
          130            135            140
Asn Thr Lys Gln Thr Ile Ser Asn Gln Glu Ala Leu Gln Pro Asp Leu
          145            150            155            160
Gln Glu Asn Lys Ser Val Val Asn Val Gln Pro Thr Asn Glu Glu Asn
          165            170            175
Lys Lys Val Asp Ala Lys Thr Glu Ser Thr Thr Leu Asn Val Lys Ser
          180            185            190

```

-continued

Asp	Ala	Ile	Lys	Ser	Asn	Asp	Glu	Thr	Leu	Val	Asp	Asn	Asn	Ser	Asn		
	195						200					205					
Ser	Asn	Asn	Glu	Asn	Asn	Ala	Asp	Ile	Ile	Leu	Pro	Lys	Ser	Thr	Ala		
	210					215					220						
Pro	Lys	Arg	Leu	Asn	Thr	Arg	Met	Arg	Ile	Ala	Ala	Val	Gln	Pro	Ser		
225					230					235				240			
Ser	Thr	Glu	Ala	Lys	Asn	Val	Asn	Asp	Leu	Ile	Thr	Ser	Asn	Thr	Thr		
				245					250					255			
Leu	Thr	Val	Val	Asp	Ala	Asp	Lys	Asn	Asn	Lys	Ile	Val	Pro	Ala	Gln		
			260					265					270				
Asp	Tyr	Leu	Ser	Leu	Lys	Ser	Gln	Ile	Thr	Val	Asp	Asp	Lys	Val	Lys		
	275						280					285					
Ser	Gly	Asp	Tyr	Phe	Thr	Ile	Lys	Tyr	Ser	Asp	Thr	Val	Gln	Val	Tyr		
	290					295					300						
Gly	Leu	Asn	Pro	Glu	Asp	Ile	Lys	Asn	Ile	Gly	Asp	Ile	Lys	Asp	Pro		
305					310					315					320		
Asn	Asn	Gly	Glu	Thr	Ile	Ala	Thr	Ala	Lys	His	Asp	Thr	Ala	Asn	Asn		
				325					330					335			
Leu	Ile	Thr	Tyr	Thr	Phe	Thr	Asp	Tyr	Val	Asp	Arg	Phe	Asn	Ser	Val		
			340					345					350				
Gln	Met	Gly	Ile	Asn	Tyr	Ser	Ile	Tyr	Met	Asp	Ala	Asp	Thr	Ile	Pro		
		355					360					365					
Val	Ser	Lys	Asn	Asp	Val	Glu	Phe	Asn	Val	Thr	Ile	Gly	Asn	Thr	Thr		
	370					375					380						
Thr	Lys	Thr	Thr	Ala	Asn	Ile	Gln	Tyr	Pro	Asp	Tyr	Val	Val	Asn	Glu		
385					390					395					400		
Lys	Asn	Ser	Ile	Gly	Ser	Ala	Phe	Thr	Glu	Thr	Val	Ser	His	Val	Gly		
			405						410					415			
Asn	Lys	Glu	Asn	Pro	Gly	Tyr	Tyr	Lys	Gln	Thr	Ile	Tyr	Val	Asn	Pro		
			420					425					430				
Ser	Glu	Asn	Ser	Leu	Thr	Asn	Ala	Lys	Leu	Lys	Val	Gln	Ala	Tyr	His		
	435						440					445					
Ser	Ser	Tyr	Pro	Asn	Asn	Ile	Gly	Gln	Ile	Asn	Lys	Asp	Val	Thr	Asp		
	450					455					460						
Ile	Lys	Ile	Tyr	Gln	Val	Pro	Lys	Gly	Tyr	Thr	Leu	Asn	Lys	Gly	Tyr		
465					470					475					480		
Asp	Val	Asn	Thr	Lys	Glu	Leu	Thr	Asp	Val	Thr	Asn	Gln	Tyr	Leu	Gln		
				485					490					495			
Lys	Ile	Thr	Tyr	Gly	Asp	Asn	Asn	Ser	Ala	Val	Ile	Asp	Phe	Gly	Asn		
			500					505					510				
Ala	Asp	Ser	Ala	Tyr	Val	Val	Met	Val	Asn	Thr	Lys	Phe	Gln	Tyr	Thr		
		515					520					525					
Asn	Ser	Glu	Ser	Pro	Thr	Leu	Val	Gln	Met	Ala	Thr	Leu	Ser	Ser	Thr		
						535					540						
Gly	Asn	Lys	Ser	Val	Ser	Thr	Gly	Asn	Ala	Leu	Gly	Phe	Thr	Asn	Asn		
545					550					555					560		
Gln	Ser	Gly	Gly	Ala	Gly	Gln	Glu	Val	Tyr	Lys	Ile	Gly	Asn	Tyr	Val		
				565					570					575			
Trp	Glu	Asp	Thr	Asn	Lys	Asn	Gly	Val	Gln	Glu	Leu	Gly	Glu	Lys	Gly		
			580					585					590				
Val	Gly	Asn	Val	Thr	Val	Thr	Val	Phe	Asp	Asn	Asn	Thr	Asn	Thr	Lys		
		595					600					605					

-continued

Val 610	Gly	Glu	Ala	Val	Thr	Lys 615	Glu	Asp	Gly	Ser	Tyr 620	Leu	Ile	Pro	Asn
Leu 625	Pro	Asn	Gly	Asp	Tyr 630	Arg	Val	Glu	Phe	Ser 635	Asn	Leu	Pro	Lys	Gly 640
Tyr	Glu	Val	Thr	Pro 645	Ser	Lys	Gln	Gly	Asn 650	Asn	Glu	Glu	Leu	Asp	Ser 655
Asn	Gly	Leu	Ser 660	Ser	Val	Ile	Thr	Val 665	Asn	Gly	Lys	Asp	Asn	Leu	Ser 670
Ala	Asp	Leu 675	Gly	Ile	Tyr	Lys	Pro 680	Lys	Tyr	Asn	Leu	Gly 685	Asp	Tyr	Val
Trp 690	Glu	Asp	Thr	Asn	Lys	Asn 695	Gly	Ile	Gln	Asp 700	Gln	Asp	Glu	Lys	Gly
Ile 705	Ser	Gly	Val	Thr 710	Val	Thr	Leu	Lys	Asp 715	Glu	Asn	Gly	Asn	Val	Leu 720
Lys	Thr	Val	Thr	Thr 725	Asp	Ala	Asp	Gly	Lys 730	Tyr	Lys	Phe	Thr	Asp	Leu 735
Asp	Asn	Gly	Asn 740	Tyr	Lys	Val	Glu	Phe 745	Thr	Thr	Pro	Glu	Gly 750	Tyr	Thr
Pro	Thr	Thr 755	Val	Thr	Ser	Gly	Ser 760	Asp	Ile	Glu	Lys	Asp 765	Ser	Asn	Gly
Leu 770	Thr	Thr	Thr	Gly 775	Val	Ile	Asn	Gly	Ala	Asp 780	Asn	Met	Thr	Leu	Asp
Ser 785	Gly	Phe	Tyr	Lys 790	Thr	Pro	Lys	Tyr	Asn 795	Leu	Gly	Asn	Tyr	Val	Trp 800
Glu	Asp	Thr	Asn	Lys 805	Asp	Gly	Lys	Gln	Asp 810	Ser	Thr	Glu	Lys	Gly	Ile 815
Ser	Gly	Val	Thr 820	Val	Thr	Leu	Lys	Asn 825	Glu	Asn	Gly	Glu	Val 830	Leu	Gln
Thr	Thr	Lys 835	Thr	Asp	Lys	Asp	Gly 840	Lys	Tyr	Gln	Phe	Thr 845	Gly	Leu	Glu
Asn 850	Gly	Thr	Tyr	Lys	Val	Glu 855	Phe	Glu	Thr	Pro	Ser 860	Gly	Tyr	Thr	Pro
Thr 865	Gln	Val	Gly	Ser	Gly 870	Thr	Asp	Glu	Gly	Ile 875	Asp	Ser	Asn	Gly	Thr 880
Ser	Thr	Thr	Gly	Val 885	Ile	Lys	Asp	Lys	Asp 890	Asn	Asp	Thr	Ile	Asp	Ser 895
Gly	Phe	Tyr	Lys 900	Pro	Thr	Tyr	Asn	Leu 905	Gly	Asp	Tyr	Val	Trp 910	Glu	Asp
Thr	Asn	Lys 915	Asn	Gly	Val	Gln	Asp 920	Lys	Asp	Glu	Lys	Gly 925	Ile	Ser	Gly
Val 930	Thr	Val	Thr	Leu	Lys 935	Asp	Glu	Asn	Asp	Lys	Val 940	Leu	Lys	Thr	Val
Thr 945	Thr	Asp	Glu	Asn	Gly 950	Lys	Tyr	Gln	Phe	Thr 955	Asp	Leu	Asn	Asn	Gly 960
Thr	Tyr	Lys	Val 965	Glu	Phe	Glu	Thr	Pro	Ser 970	Gly	Tyr	Thr	Pro	Thr	Ser 975
Val	Thr	Ser 980	Gly	Asn	Asp	Thr	Glu	Lys 985	Asp	Ser	Asn	Gly	Leu 990	Thr	Thr
Thr 995	Gly	Val	Ile	Lys	Asp	Ala	Asp 1000	Asn	Met	Thr	Leu	Asp 1005	Ser	Gly	Phe
Tyr 1010	Lys	Thr	Pro	Lys	Tyr	Ser	Leu 1015	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser

-continued

Asn Lys Asp Gly Lys Gln Asp Ser Thr Glu Lys Gly Ile Lys Asp Val
 1025 1030 1035 1040

Lys Val Thr Leu Leu Asn Glu Lys Gly Glu Val Ile Gly Thr Thr Lys
 1045 1050 1055

Thr Asp Glu Asn Gly Lys Tyr Cys Phe Asp Asn Leu Asp Ser Gly Lys
 1060 1065 1070

Tyr Lys Val Ile Phe Glu Lys Pro Ala Gly Leu Thr Gln Thr Gly Thr
 1075 1080 1085

Asn Thr Thr Glu Asp Asp Lys Asp Ala Asp Gly Gly Glu Val Asp Val
 1090 1095 1100

Thr Ile Thr Asp His Asp Asp Phe Thr Leu Asp Asn Gly Tyr Tyr Glu
 1105 1110 1115 1120

Glu Glu Thr Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1125 1130 1135

Asp Arg Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1140 1145 1150

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Arg
 1155 1160 1165

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1170 1175 1180

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1185 1190 1195 1200

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1205 1210 1215

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1220 1225 1230

Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser
 1235 1240 1245

Asp Ser Asp Ser Asp Ser Asp Ala Gly Lys His Thr Pro Val Lys Pro
 1250 1255 1260

Met Ser Thr Thr Lys Asp His His Asn Lys Ala Lys Ala Leu Pro Glu
 1265 1270 1275 1280

Thr Gly Asn Glu Asn Ser Gly Ser Asn Asn Ala Thr Leu Phe Gly Gly
 1285 1290 1295

Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys
 1300 1305 1310

Gln Asn Lys
 1315

<210> SEQ ID NO 6
 <211> LENGTH: 3945
 <212> TYPE: DNA
 <213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 6

atgctaaaca gagaaaataa aacggcaata acaaggaaag gcatggtatc caatcgatta	60
aataaat ttt cgattagaaa gtacacagtg ggaacagcat caat ttt tagt aggtacaaca	120
ttaatt ttttg gtctggggaa ccaagaagca aaggctgcag aaagtactaa taaagaattg	180
aacgaagcga caacttcagc aagtataat caatcgagtg ataaagt tga tatgcagcaa	240
ctaaatcaag aagacaatac taaaaatgat aatcaaaaag aaatggatc atctcaaggt	300
aatgaaacga cttcaaatgg gaataaatta atagaaaaag aaagtgtaca atctaccact	360
ggaataaag ttgaagtttc aactgcaaaa tcagatgagc aagcttcacc aaaatctacg	420

-continued

aatgaagatt	taaacactaa	acaaactata	agtaatcaag	aagcgttaca	acctgatttg	480
caagagaata	aatcagtggt	aaatgttcaa	ccaactaatg	aggaaaaaca	aaaggtagat	540
gccaaaactg	aatcaactac	atataaatgtt	aaaagtgatg	ctatcaagag	taatgatgaa	600
actcttggtg	ataacaatag	taattcaaat	aatgaaaata	atgcagatat	cattttgccca	660
aaaagtacag	cacctaaacg	tttgaatata	agaatgcgta	tagcagcagt	acagccatca	720
tcaacagagg	ctaaaaatgt	taattgattta	atcacatcaa	atacaacatt	aactgtcgtt	780
gatgcagata	aaaacaataa	aatcgtacca	gccaagatt	atttatoatt	aaaatcacia	840
attacagttg	atgacaaaagt	taaatcaggt	gattattttca	caattaaata	ctcagataca	900
gtacaagtat	atggattgaa	tccggaagat	attaaaaata	ttggtgatat	taaagatcca	960
aataatgggtg	aaacaattgc	gactgcaaaa	catgatactg	caaataattt	aattacatat	1020
acatttacag	attatggtga	tcgattttaat	tctgtacaaa	tggaataaa	ttattcaatt	1080
tatatggatg	ctgatacaat	tctctgttagt	aaaaacgatg	ttgagtttaa	tgttacgata	1140
ggtaatacta	caacaaaaac	aactgctaac	attcaatcag	cagattatgt	tgtaaatgag	1200
aaaaattcaa	ttggatcagc	gttcactgaa	acagtttcac	atggttgaaa	taaagaaaat	1260
ccagggact	ataacaaaac	gattttatgta	aatccatcgg	aaaattcttt	aacaaatgcc	1320
aaactaaaag	ttcaagotta	ccactcaagt	tatcctaata	atcctgggca	aataaataaa	1380
gatgtaacag	atataaaaaat	atatcaagtt	cctaaagggt	atacattaaa	taaaggatac	1440
gatgtgaata	ctaaagagct	tacagatgta	acaaatcaat	acttgcaaaa	aattacatat	1500
ggcgacaaca	atagcgotgt	tattgatttt	ggaaatgcag	attctgotta	tggtgtaatg	1560
gttaatacaa	aattccaata	tacaaatagc	gaaagcccaa	cacttggtca	aatggctact	1620
ttatcttcaa	caggtaataa	atccgtttct	actggcaatg	ctttaggatt	tactaataac	1680
caaagtggcg	gagctgggtca	agaagtatat	aaaattggta	actacgtatg	ggaagatact	1740
aataaaaaacg	gtgttcaaga	attaggagaa	aaaggcgttg	gcaatgtaac	tgtaactgta	1800
tttgataata	atacaaacac	aaaagtagga	gaagcagtta	ctaaagaaga	tggtgcatac	1860
ttgattccaa	acttacctaa	tggagattac	cgtgtagaat	tttcaactt	accaaaaggt	1920
tatgaagtaa	ccccttcaaa	acaaggtaat	aacgaagaat	tagattcaaa	cggcttatct	1980
tcagttatta	cagttaatgg	caaagataac	ttatctgcag	acttaggtat	ttacaaacct	2040
aaatacaact	taggtgacta	tgtctgggaa	gatacaata	aaaatgggat	ccaagaccaa	2100
gatgaaaaag	gtatatctgg	cgtaacggta	acattaaaag	atgaaaacg	taacgtgta	2160
aaaacagtta	caacagacgc	tgatggcaaa	tataaattta	ctgattttaga	taatggtaat	2220
tataaagttg	aatttactac	accagaaggc	tataaccga	ctacagtaac	atctggtagc	2280
gacattgaaa	aagactctaa	tggtttaaca	acaacaggtg	ttattaatgg	tgctgataac	2340
atgacattag	atagtggtat	ctacaaaaca	ccaaaatata	atttaggtaa	ttatgtatgg	2400
gaagatacaa	ataaagatgg	taagcaggat	tcaactgaaa	aaggatattc	aggcgtataca	2460
gttacattga	aaaatgaaaa	cgggtgaagtt	ttacaaaca	ctaaaacaga	taaagatggt	2520
aaatatcaat	ttactggatt	agaaaaatgga	acttataaag	ttgaattcga	aacaccatca	2580
ggttacacac	caacacaagt	aggttcagga	actgatgaag	gtatagattc	aatgggtaca	2640
tcaacaacag	gtgtcattaa	agataaagat	aacgatacta	ttgactctgg	tttctacaaa	2700
ccgacttaca	acttaggtga	ctatgtatgg	gaagatacaa	ataaaaacg	tgttcaagat	2760
aaagatgaaa	agggcatttc	agggtgaaca	gttacgttaa	aagatgaaa	cgacaaaagtt	2820

-continued

```

ttaaaaacag ttacaacaga tgaaaatggt aaatatcaat tcaactgattt aaacaatgga 2880
acttataaag ttgaattcga gacaccatca ggttatacac caacttcagt aacttctgga 2940
aatgatactg aaaaagattc taatggttta acaacaacag gtgtcattaa agatgcagat 3000
aacatgacat tagacagtgg tttctataaa acaccaaata atagtttagg tgattatggt 3060
tggtagcaca gtaataaaga cggcaaacaa gattcaactg aaaaaggat caaagatggt 3120
aaagttaact tattaaatga aaaaggcgaa gtaattggaa caactaaaac agatgaaaat 3180
ggtaaatact gctttgataa tttagatagc ggtaaataca aagttatgtt tgaaaagcct 3240
gctggcttaa cacaaacagg tacaataaca actgaagatg ataaagatgc agatggtggc 3300
gaagttgacg taacaattac ggatcatgat gatttcacac ttgataatgg ctactacgaa 3360
gaagaaacat cagatagcga ctcagattcg gacagcgact cagattcaga cagagactca 3420
gactcagata gtgattcaga ctcggatagc gattcagatt cagacagcga ttcagattca 3480
gatagcgatt cagattcaga cagagactca gatagtgatt cagactcaga tagcgactca 3540
gattcagaca gcgactcaga ttcagacagc gactcagact cagatagtga ttcagactca 3600
gatagcgact cagattogga tagcgactca gattcagaca gcgactcaga ctcggatagt 3660
gattcagact cagatagcga ctcagactca gatagcgatt cagattcaga tagcgactca 3720
gactcagaca gcgattcaga ctcagacagc gactcagact cagatgcagg taagcacaca 3780
cctgttaaac caatgagtac tactaaagac catcacaata aagcaaaagc attaccagaa 3840
acaggtaatg aaaatagcgg ctcaataaac gcaacgttat ttggcggatt attcgcagca 3900
ttaggatcat tattgttatt cggtcgtcgt aaaaaacaaa ataaa 3945

```

<210> SEQ ID NO 7

<211> LENGTH: 1166

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 7

```

Met Ile Asn Arg Asp Asn Lys Lys Ala Ile Thr Lys Lys Gly Met Ile
 1           5           10          15
Ser Asn Arg Leu Asn Lys Phe Ser Ile Arg Lys Tyr Thr Val Gly Thr
          20           25           30
Ala Ser Ile Leu Val Gly Thr Thr Leu Ile Phe Gly Leu Gly Asn Gln
          35           40           45
Glu Ala Lys Ala Ala Glu Asn Thr Ser Thr Glu Asn Ala Lys Gln Asp
          50           55           60
Asp Ala Thr Thr Ser Asp Asn Lys Glu Val Val Ser Glu Thr Glu Asn
          65           70           75           80
Asn Ser Thr Thr Glu Asn Asn Ser Thr Asn Pro Ile Lys Lys Glu Thr
          85           90           95
Asn Thr Asp Ser Gln Pro Glu Ala Lys Lys Glu Ser Thr Ser Ser Ser
          100          105          110
Thr Gln Lys Gln Gln Asn Asn Val Thr Ala Thr Thr Glu Thr Lys Pro
          115          120          125
Gln Asn Ile Glu Lys Glu Asn Val Lys Pro Ser Thr Asp Lys Thr Ala
          130          135          140
Thr Glu Asp Thr Ser Val Ile Leu Glu Glu Lys Lys Ala Pro Asn Asn
          145          150          155          160
Thr Asn Asn Asp Val Thr Thr Lys Pro Ser Thr Ser Glu Pro Ser Thr
          165          170          175

```

-continued

Ser Glu Ile Gln Thr Lys Pro Thr Thr Pro Gln Glu Ser Thr Asn Ile
 180 185 190

Glu Asn Ser Gln Pro Gln Pro Thr Pro Ser Lys Val Asp Asn Gln Val
 195 200 205

Thr Asp Ala Thr Asn Pro Lys Glu Pro Val Asn Val Ser Lys Glu Glu
 210 215 220

Leu Lys Asn Asn Pro Glu Lys Leu Lys Glu Leu Val Arg Asn Asp Ser
 225 230 235 240

Asn Thr Asp His Ser Thr Lys Pro Val Ala Thr Ala Pro Thr Ser Val
 245 250 255

Ala Pro Lys Arg Val Asn Ala Lys Met Arg Phe Ala Val Ala Gln Pro
 260 265 270

Ala Ala Val Ala Ser Asn Asn Val Asn Asp Leu Ile Lys Val Thr Lys
 275 280 285

Gln Thr Ile Lys Val Gly Asp Gly Lys Asp Asn Val Ala Ala Ala His
 290 295 300

Asp Gly Lys Asp Ile Glu Tyr Asp Thr Glu Phe Thr Ile Asp Asn Lys
 305 310 315 320

Val Lys Lys Gly Asp Thr Met Thr Ile Asn Tyr Asp Lys Asn Val Ile
 325 330 335

Pro Ser Asp Leu Thr Asp Lys Asn Asp Pro Ile Asp Ile Thr Asp Pro
 340 345 350

Ser Gly Glu Val Ile Ala Lys Gly Thr Phe Asp Lys Ala Thr Lys Gln
 355 360 365

Ile Thr Tyr Thr Phe Thr Asp Tyr Val Asp Lys Tyr Glu Asp Ile Lys
 370 375 380

Ser Arg Leu Thr Leu Tyr Ser Tyr Ile Asp Lys Lys Thr Val Pro Asn
 385 390 395 400

Glu Thr Ser Leu Asn Leu Thr Phe Ala Thr Ala Gly Lys Glu Thr Ser
 405 410 415

Gln Asn Val Thr Val Asp Tyr Gln Asp Pro Met Val His Gly Asp Ser
 420 425 430

Asn Ile Gln Ser Ile Phe Thr Lys Leu Asp Glu Asp Lys Gln Thr Ile
 435 440 445

Glu Gln Gln Ile Tyr Val Asn Pro Leu Lys Lys Ser Ala Thr Asn Thr
 450 455 460

Lys Val Asp Ile Ala Gly Ser Gln Val Asp Asp Tyr Gly Asn Ile Lys
 465 470 475 480

Leu Gly Asn Gly Ser Thr Ile Ile Asp Gln Asn Thr Glu Ile Lys Val
 485 490 495

Tyr Lys Val Asn Ser Asp Gln Gln Leu Pro Gln Ser Asn Arg Ile Tyr
 500 505 510

Asp Phe Ser Gln Tyr Glu Asp Val Thr Ser Gln Phe Asp Asn Lys Lys
 515 520 525

Ser Phe Ser Asn Asn Val Ala Thr Leu Asp Phe Gly Asp Ile Asn Ser
 530 535 540

Ala Tyr Ile Ile Lys Val Val Ser Lys Tyr Thr Pro Thr Ser Asp Gly
 545 550 555 560

Glu Leu Asp Ile Ala Gln Gly Thr Ser Met Arg Thr Thr Asp Lys Tyr
 565 570 575

Gly Tyr Tyr Asn Tyr Ala Gly Tyr Ser Asn Phe Ile Val Thr Ser Asn
 580 585 590

-continued

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1010 1015 1020

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1025 1030 1035 1040

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1045 1050 1055

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1060 1065 1070

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1075 1080 1085

Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp Ser Asp
 1090 1095 1100

Ser Asp Ala Gly Lys His Thr Pro Val Lys Pro Met Ser Thr Thr Lys
 1105 1110 1115 1120

Asp His His Asn Lys Ala Lys Ala Leu Pro Glu Thr Gly Ser Glu Asn
 1125 1130 1135

Asn Gly Ser Asn Asn Ala Thr Leu Phe Gly Gly Leu Phe Ala Ala Leu
 1140 1145 1150

Gly Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys Gln Asn Lys
 1155 1160 1165

<210> SEQ ID NO 8
 <211> LENGTH: 3498
 <212> TYPE: DNA
 <213> ORGANISM: Staphylococcus aureus
 <400> SEQUENCE: 8

atgattaaca gggataataa aaaggaaca acaaaaaagg gtatgatttc aaatcgctta 60
 aacaaatddd cgattagaaa gtatactgta ggaactgcat cgattttagt aggtacgaca 120
 ttgatttttg gtctagggaa ccaagaagct aaagctgctg aaaacactag tacagaaaat 180
 gcaaaacaag atgatgcaac gactagtgat aataaagaag tagtgtcggg aactgaaaat 240
 aattcgacaa cagaaaataa ttcaacaaat ccaattaaga aagaaacaaa tactgattca 300
 caaccagaag ctaaaaaaga atcaacttca tcaagtactc aaaaacagca aaataacggt 360
 acagctacaa ctgaaactaa gcctcaaaac attgaaaaag aaaatgttaa accttcaact 420
 gataaaaactg cgacagaaga tacatctggt attttagaag agaagaaagc accaaataat 480
 acaataaacg atgtaactac aaaaccatct acaagtgaac catctacaag tgaattcaa 540
 acaaaaccaa ctacacctca agaacttaca aatattgaaa attcacaacc gcaaccaacg 600
 ccttcaaaag tagacaatca agttacagat gcaactaatc caaaagaacc agtaaatgtg 660
 tcaaaagaag aacttaaaaa taatcctgag aaattaaaag aattggttag aatgatagc 720
 aatacagatc attcaactaa accagttgct acagctcaa caagtgttc accaaaacgt 780
 gtaaacgcaa aaatgcgctt tgcaagtgca caaccagcag cagttgcttc aaacaatgta 840
 aatgatttaa ttaaagtgac gaagcaaaca atcaaagttg gcgatggtaa agataatgtg 900
 gcagcagcgc atgacggtaa agatattgaa tatgatacag agtttacaat tgacaataaa 960
 gtcaaaaaag gcgatacaat gacgattaat tatgataaga atgtaattcc ttcggattta 1020
 acagataaaa atgatcctat cgatattact gatccatcag gagaggtcat tgctaaagga 1080
 acatttgata aagcaactaa gcaaatcaca tatacattta cagactatgt agataaatat 1140
 gaagatataa aatcacgctt aactctatat tcgtatattg ataaaaaac agttccaaat 1200
 gagacaagtt tgaatttaac atttgttaca gcaggtaaag aaacaagcca aatgtcact 1260

-continued

gttgattatc aagatccaat ggtccatggt gattcaaaca ttcaatctat ctttacaaaa	1320
ttagatgaag ataagcaaac tattgaacaa caaatattatg ttaaccatt gaaaaatca	1380
gcaaccaaca ctaaagttga tatagctggt agtcaagtag atgattatgg aatatataa	1440
ctaggaatg gtagcacat tattgaccaa aatacagaaa taaagttta taaagttaac	1500
tctgatcaac aattgcctca aagtaataga atctatgatt ttagtcaata cgaagatgta	1560
acaagtcaat ttgataataa aaatcattt agtaataatg tagcaacatt ggattttggt	1620
gatattaatt cagcctatat tatcaaagt gttagtaaat atacacctac atcagatggc	1680
gaactagata ttgccaagg tactagtagt agaacaactg ataaatatgg ttattataat	1740
tatgcaggat attcaaact catcgtaact tctaatgaca ctggcggtag cgacggtact	1800
gttaaacctg aagaaaagt atacaaaatt ggtgactatg tatgggaaga cgttgataaa	1860
gacggtgttc aaggtacaga ttcaaaagaa aaaccaatgg caaacgtttt agttacatta	1920
acttaccogg acggtactac aaatcagta agaacagatg ctaatggtca ttatgaattc	1980
ggtggtttga aagacggaga aacttataca gttaaattcg aaacgccaac tggatatctt	2040
ccaacaaaag taaatggaac aactgatggt gaaaagact caaatgtag ttcggttact	2100
gttaaaatta atggtaaaga tgatgtct ttagatactg gtttttaca agaacctaaa	2160
tacaacttag gtgactatgt atgggaagat actaataaag atggtatcca agatgcaaat	2220
gagccaggaa tcaaatggt taaggttaca ttaaaagata gtactggaaa agttattggt	2280
acaactacta ctgatgctc gggtaaatat aaatttacag atttagataa tggttaactat	2340
acagtagaat ttgaaacacc agcaggttac acgccaacgg ttaaaaaaac tacagctgat	2400
gataaagatt ctaatggtt aacaacaaca ggtgtcatta aagatgcaga taatatgaca	2460
ttagacaggg gtttctataa aacacaaaa tacagtttag gtgattatgt ttggtacgac	2520
agtaataaag acggcaaca agattcaact gaaaaggta tcaaatggt gacagttaca	2580
ttgcaaacg aaaaaggcga agtaattgga acaactaaa cagatgaaa tggtaaatat	2640
cgtttcgata atttagatag cggtaaatc aaagtattt ttgaaaagcc tgctggctta	2700
acacaaacag ttacaaatac aactgaagat gataaagatg cagatggtgg cgaagttgac	2760
gtaacaatta cggatcatga tgatttcaca cttgataacg gatacttca agaagataca	2820
tcagacagcg attcagactc agatagtgac tcagacagcg actcagactc agacagcgac	2880
tcagactcag acagtgattc agattcagac agcgactcag attcagatag cgactcagat	2940
tcggacagcg attcagactc agatagcgac tcagattcag atagcgattc agactcagac	3000
agcgactcag attcagatag cgattcggac tcagacagcg attcagactc agatagcgac	3060
tcagactcag acagcgactc agattcagat agcgattcgg actcagatag cgactcagat	3120
tcagacagcg attcagactc agatagcgac tcagattcag acagcgattc agactcagat	3180
agcgactcag actcagacag tgattcagat tcagacagcg actcagactc agatagcgac	3240
tcagattcag acagcgactc agactcagat agcgactcag actcagacag tgattcagac	3300
agcgattcag actcggatgc aggaaaacat acacctgta aaccaatgag tactactaaa	3360
gaccatcaca ataaagcaaa agcattacca gaaacaggta gtgaaaataa cggctcaaat	3420
aacgcaogt tatttggtag attatttgca gcattaggtt cattattggt attcggctcgt	3480
cgcaaaaaac aaaacaaa	3498

-continued

```

<210> SEQ ID NO 9
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Staphylococcus aureus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: n = (a or c or t or g)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (12)..(12)
<223> OTHER INFORMATION: n = (a or c or t or g)

```

```

<400> SEQUENCE: 9

```

```

gaytcngayt cngayagy

```

18

```

<210> SEQ ID NO 10
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 10

```

```

Leu Pro Asp Thr Gly
  1             5

```

```

<210> SEQ ID NO 11
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 11

```

```

Asp Tyr Ser Asn Ser
  1             5

```

```

<210> SEQ ID NO 12
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 12

```

```

Phe Thr Asp Tyr Val Asn
  1             5

```

```

<210> SEQ ID NO 13
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa = (Unspecified amino acid)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa = (Unspecified amino acid)

```

```

<400> SEQUENCE: 13

```

```

Asp Xaa Ser Xaa Ser
  1             5

```

```

<210> SEQ ID NO 14
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa = (Unspecified amino acid)

```

-continued

<400> SEQUENCE: 14

Leu Pro Xaa Thr Gly
 1 5

<210> SEQ ID NO 15

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 15

cgaggatcct caggacaatc gaacgatata acg

33

<210> SEQ ID NO 16

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 16

cgaggtacca tttactgctg aatcacc

27

<210> SEQ ID NO 17

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 17

cgaaagcttg tcagaacaat cgaacgatac aacg

34

<210> SEQ ID NO 18

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 18

Thr Tyr Thr Phe Thr Asp Tyr Val Asp
 1 5

<210> SEQ ID NO 19

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 19

Thr Tyr Thr Phe Thr Asn Tyr Val Asp
 1 5

<210> SEQ ID NO 20

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 20

Thr Phe Val Phe Thr Asp Tyr Val Asn
 1 5

<210> SEQ ID NO 21

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Staphylococcus aureus

<400> SEQUENCE: 21

Ile Tyr Thr Phe Thr Asp Tyr Val Asn
 1 5

-continued

```

<210> SEQ ID NO 22
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 22

```

```

Leu Pro Glu Thr Gly
 1             5

```

```

<210> SEQ ID NO 23
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Staphylococcus aureus

```

```

<400> SEQUENCE: 23

```

```

Thr Ile Tyr Phe Thr Val Phe Thr Asp Asn Tyr Val Asp Asn
 1             5             10

```

We claim:

1. An isolated nucleic acid molecule encoding an amino acid selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, and SEQ ID NO:7.

2. An isolated nucleic acid molecule having a sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, and SEQ ID NO:8.

3. A vector comprising the isolated nucleic acid of claim 1.

4. The vector of claim 3 that is capable of expressing the nucleic acid.

25 5. A vector comprising the isolated nucleic acid of claim 2.

6. The vector of claim 5 that is capable of expressing the nucleic acid.

* * * * *