



US007777096B2

(12) **United States Patent**
Mirkov et al.

(10) **Patent No.:** **US 7,777,096 B2**
(45) **Date of Patent:** ***Aug. 17, 2010**

(54) **ISOLATION OF PROTEINS INVOLVED IN POSTTRANSCRIPTIONAL GENE SILENCING AND METHODS OF USE**

(75) Inventors: **T. Erik Mirkov**, Harlingen, TX (US);
Ivan L. Ingelbrecht, Jabbeke (BE)

(73) Assignee: **The Texas A&M University System**,
College Station, TX (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 455 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **11/460,836**

(22) Filed: **Jul. 28, 2006**

(65) **Prior Publication Data**

US 2006/0272052 A1 Nov. 30, 2006

Related U.S. Application Data

(62) Division of application No. 11/252,080, filed on Oct. 17, 2005, now abandoned, which is a division of application No. 10/226,715, filed on Aug. 23, 2002, now Pat. No. 7,001,739.

(51) **Int. Cl.**
C12N 15/82 (2006.01)

(52) **U.S. Cl.** **800/278**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,939,541	A	8/1999	Vance et al.	536/24.1
6,040,496	A	3/2000	Law et al.	800/280
6,395,962	B1*	5/2002	Vance	800/278
7,001,739	B2	2/2006	Mirkov et al.	435/7.91
7,267,985	B2*	9/2007	Vance	435/468
2006/0269955	A1	11/2006	Mirkov et al.	435/6
2006/0272047	A1	11/2006	Mirkov et al.	800/278
2006/0275830	A1	12/2006	Mirkov et al.	435/7.1
2006/0286607	A1	12/2006	Mirkov et al.	435/7.1
2007/0067880	A1	3/2007	Mirkov et al.	800/320

FOREIGN PATENT DOCUMENTS

WO 03/018809 A2 8/2002

OTHER PUBLICATIONS

Chen et al., Arch. Virol., Jun. 2002, vol. 147, pp. 1237-1246.*
 Arencibia et al., Plant Cell Rep., 1995, vol. 14, pp. 305-309.*
 UniProt Acc. No. Q8QXL1, Jun. 1, 2002.*
 Anandalakshmi "A Calmodulin-Related Protein that Suppresses Posttranscriptional Gene Silencing in Plants" Science, American Association for the Advancement of Science, vol. 290, No. 5489 (pp. 142-144), Oct. 6, 2000.
 Anandalakshmi et al. "A viral suppressor of gene silencing in plants" Proc. Natl. Acad. Sci. USA vol. 95 (pp. 13079-13084), Oct. 1998.
 Atreya et al. "Mutational analysis of the helper component-proteinase gene of potyvirus: Effects of amino acid substitutions,

deletions, and gene replacement on virulence and aphid transmissibility" Proc. Natl. Acad. Sci. USA vol. 90 (pp. 11919-11923), Dec. 1993.

Brigneti et al. "Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*" The EMBO Journal, vol. 17, No. 22 (pp. 6739-6746), 1998.

Carrington et al. "Expression of potyviral polyproteins in transgenic plants reveals three proteolytic activities required for complete processing" The EMBO Journal, vol. 9, No. 5 (pp. 1347-1353), 1990.

Chuang et al. "Specific and Heritable Genetic Interference by Double-Stranded RNA in *Arabidopsis thaliana*" PNAS, vol. 97, No. 9 (pp. 4985-4990), Apr. 25, 2000.

Cogoni et al. "Conservation of transgene-induced post-transcriptional gene silencing in plants and fungi" Trends in Plant Science Reviews, vol. 2, No. 11 (pp. 438-443), Nov. 1997.

Cogoni et al. "Gene silencing in *Neurospora crassa* requires a protein homologous to RNA-dependent RNA polymerase" Nature, vol. 399 (pp. 166-169), May 13, 1999.

Cogoni et al. "Isolation of quelling-defective (*qde*) mutants impaired in posttranscriptional transgene-induced gene silencing in *Neurospora crassa*" Proc. Natl. Acad. Sci. USA vol. 94 (pp. 10233-10238), Sep. 1997.

Cogoni et al. "Posttranscriptional Gene Silencing in *Neurospora* by a RecQ DNA Helicase" Science, vol. 286 (pp. 2342-2344), Dec. 17, 1999.

Covey et al. "Plants combat infection by gene silencing" Nature, vol. 385 (pp. 781-782), Feb. 27, 1997.

Cronin et al. "Long-Distance Movement Factor: A Transport Function of the Potyvirus Helper Component Proteinase" The Plant Cell, vol. 7 (pp. 549-559), May 1995.

Dagkessamanskaia et al. FEMS Microbiol. Lett. vol. 200 (pp. 53-58), Jun. 2001.

Dalmay et al. "An RNA-Dependent RNA Polymerase Gene in *Arabidopsis* is Required for Posttranscriptional Gene Silencing Mediated by a Transgene but Not by a Virus" Cell, vol. 101 (pp. 543-553), May 26, 2000.

Dehio et al. "Identification of plant genetic loci involved in a post-transcriptional mechanism for meiotically reversible transgene silencing" Proc. Natl. Acad. Sci. USA, vol. 91 (pp. 5538-5542), Jun. 1994.

(Continued)

Primary Examiner—Ashwin Mehta
(74) Attorney, Agent, or Firm—King & Spalding L.L.P.

(57) **ABSTRACT**

The present disclosure includes methods of using viral and plant proteins to regulate silencing in plants such as monocots. Some embodiments include methods of increasing expression of transgenes in plants. Such methods may include selecting a protein involved in post-translational gene silencing (PTGS), for e.g., a protein active in PTGS or a suppressor of PTGS, in a plant and transforming the plant with a nucleic acid encoding the protein involved in PTGS wherein expression of the protein involved in PTGS increases expression of the transgene in the plant. One non-limiting example of such a protein is the Sorghum Mosaic Virus (SrMV) P1/HC-Pro. Methods of the disclosure may be used to regulate silencing of transgenes and to increase expression of transgenes in a variety of plants such as sugarcane, corn, sorghum, and/or rice.

OTHER PUBLICATIONS

- Depicker et al. "Post-transcriptional gene silencing in plants" *Current Opinion in Cell Biology*, vol. 9 (pp. 373-382), 1997.
- Elmayan et al. "*Arabidopsis* Mutants Impaired in Cosuppression" *The Plant Cell*, vol. 10 (pp. 1747-1757), Oct. 1998.
- Fields et al. "A novel genetic system to detect protein-protein interactions" *Nature*, vol. 340 (pp. 245-246), Jul. 20, 1989.
- Geering et al. "Genetic Diversity Among Banana Streak Virus Isolates from Australia" Database Embl AF215815 (2 pages), Jul. 18, 2000.
- Geering et al. "Genetic Diversity Among Banana Streak Virus Isolates from Australia" XP008017774, *Phytopathology*, vol. 90, No. 8 (pp. 921-927), Aug. 2000.
- Gindullis et al. "MAF1, A Novel Plant Protein Interacting with Matrix Attachment Region Binding Protein MFP1, Is Located at the Nuclear Envelope" *The Plant Cell*, vol. 11 (pp. 1755-1767), Sep. 1999.
- Guo et al. "Self-association and mapping of interaction domains of helper component-proteinase of potato A potyvirus" *Journal of General Virology*, vol. 80 (pp. 1127-1131), 1999.
- Hamilton et al. "A Species of Small Antisense RNA in Post-transcriptional Gene Silencing in Plants" *Science*, vol. 286 (pp. 950-952), Oct. 29, 1999.
- Ingelbrecht et al. "Posttranscriptional Gene Silencing in Transgenic Sugarcane Dissection of Homology-Dependent Virus Resistance in a Monocot That Has a Complex Polyploid Genome" *Plant Physiology*, vol. 119 (pp. 1187-1197), Apr. 1999.
- Ingelbrecht et al. "Posttranscriptional silencing of reporter transgenes in tobacco correlates with DNA methylation" *Proc. Natl. Acad. Sci. USA*, vol. 91 (pp. 10502-10506), Oct. 1994.
- Kasschau et al. "A Counterdefensive Strategy of Plant Viruses: Suppression of Posttranscriptional Gene Silencing" *Cell*, vol. 95 (pp. 461-470), Nov. 13, 1998.
- Kasschau et al. "Genome Amplification and Long-Distance Movement Functions Associated with the Central Domain of Tobacco Etch Potyvirus Helper Component-Proteinase" *Virology*, vol. 228, Article No. VY968368 (pp. 251-262), 1997.
- Kennerdell et al. "Use of dsRNA-Mediated Genetic Interference to Demonstrate that *frizzled* and *frizzled 2* Act in the Wingless Pathway" *Cell*, vol. 95 (pp. 1017-1026), Dec. 23, 1998.
- Ketting et al. "mut-7 of *C. elegans*, Required for Transposon Silencing and RNA Interference, Is a Homolog of Werner Syndrome Helicase and RNaseD" *Cell*, vol. 99 (pp. 133-141), Oct. 15, 1999.
- Kleiman et al. "Functional Interaction of BRCA1-Associated BARD1 with Polyadenylation Factor CstF-50" *Science*, vol. 285 (pp. 1576-1579), Sep. 3, 1999.
- Kohalmi et al. "Identification and characterization of protein interactions using the yeast 2-hybrid system" *Plant Molecular Biology Manuari*, vol. M1 (pp. 1-30), 1998.
- Kong "Complete Nucleotide Sequence and Analysis of the Putative Polyprotein of Maize Dwarf Mosaic Virus Genomic RNA (Bulgarian Isolate)" *Archives of Virology*, vol. 143, No. 9 (pp. 1791-1799), 1998.
- Kong "Complete Nucleotide Sequence and Analysis of the Putative Polyprotein of Maize Dwarf Mosaic Virus Genomic RNA (Bulgarian Isolate)" Database Embl AJ001691.1 (5 pages), May 6, 1998.
- Kumagai et al. "Cytoplasmic inhibition of carotenoid biosynthesis with virus-derived RNA" *Proc. Natl. Acad. Sci. USA*, vol. 92 (pp. 1679-1683), Feb. 1998.
- Kumpatla et al. "Genome intruder scanning and modulation systems and transgene silencing" *Trends in Plant Science*, vol. 3, No. 3 (pp. 97-104), Mar. 1998.
- Lindbo et al. "Induction of a Highly Specific Antiviral State in Transgenic Plants: Implications for Regulation of Gene Expression and Virus Resistance" *The Plant Cell*, vol. 5 (pp. 1749-1759), Dec. 1993.
- Mallory et al. "Suppression of RNA Silencing in Plants" *Phytopathology*, vol. 91, No. 6 Supplement (1 page), Jun. 2001.
- Matzke et al. "Epigenetic silencing of plant transgenes as a consequence of diverse cellular defense responses" *CMLS, Cell. Mol. Life Sci.*, vol. 54 (pp. 94-103), 1998.
- Matzke et al. "RNA-based Silencing Strategies in Plants" *Current Opinion in Genetics & Development*, vol. 11, No. 2 (pp. 221-227), Apr. 2001.
- Mirkov et al. "Monocot Host Proteins that Interact with a Viral Suppressor of Gene Silencing" *Phytopathology*, vol. 91, No. 6 Supplement (1 page), Jun. 2001.
- Misquitta et al. "Targeted disruption of gene function in *Drosophila* by RNA interference (RNA-i): A Role for *nautilus* in embryonic somatic muscle formation" *Proc. Natl. Acad. Sci. USA*, vol. 96 (pp. 1451-1456), Feb. 1999.
- Montgomery et al. "Double-stranded RNA as a mediator in sequence-specific genetic silencing and cosuppression" *TIG*, vol. 14, No. 7 (pp. 255-258), Jul. 98.
- Mourrain et al. "*Arabidopsis* SGS2 and SGS3 Genes Are Required for Posttranscriptional Gene Silencing and Natural Virus Resistance" *Cell*, vol. 101 (pp. 533-542), May 26, 2000.
- Napoli et al. "Introduction of a Chimeric Chalcone Synthase Gene into *Petunia* Results in Reversible Co-Suppression of Homologous Genes in trans" *The Plant Cell*, vol. 2 (pp. 279-289), Apr. 1990.
- Newberry et al. *Biochem.* vol. 38 (pp. 10678-10690), 1999.
- Palauqui et al. "Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions" *The EMBO Journal*, vol. 16, No. 15 (pp. 4738-4745), 1997.
- Pruss et al. "Plant Viral Synergism: The Potyviral Genome Encodes a Broad-Range Pathogenicity Enhancer That Transactivates Replication of Heterologous Viruses" *The Plant Cell*, vol. 9 (pp. 859-868), Jun. 1997.
- Ratcliff et al. "A Similarity Between Viral Defense and Gene Silencing in Plants" *Science*, vol. 276 (pp. 1558-1560), Jun. 6, 1997.
- Revers et al. "New Advances in Understanding the Molecular Biology of Plant/Potyvirus Interactions" *MPMI*, vol. 12, No. 5 (pp. 367-376), 1999.
- Ruiz et al. "Initiation and Maintenance of Virus-Induced Gene Silencing" *The Plant Cell*, vol. 10 (pp. 937-946), Jun. 1998.
- Sanchez Alvarado et al. "Double-stranded RNA specifically disrupts gene expression during planarian regeneration" *Proc. Natl. Acad. Sci. USA*, vol. 96 (pp. 5049-5054), Apr. 1999.
- Schiebel et al. "Isolation of an RNA-Directed RNA Polymerase-Specific cDNA Clone from Tomato" *The Plant Cell*, vol. 10. (pp. 2087-2101), Dec. 1998.
- Sehnke et al. "Consummating Signal Transduction: The Role of 14-3-3 Proteins in the Completion of Signal-Induced Transitions in Protein Activity" *The Plant Cell*, Supplement (pp. S339-S354), 2002.
- Sharp "RNAi and double-strand RNA" *Genes & Development*, vol. 13 (pp. 139-141), 1999.
- Shukla et al. "The Sugarcane Mosaic Virus Subgroup" *The Potyviridae*, CABI Chapter 11 (pp. 360-371), 1994.
- Smith et al. "Total silencing by intraspliced hairpin RNAs" *Nature*, vol. 407 (pp. 319-320), Sep. 21, 2000.
- Tabara et al. "The *rde-1* Gene, RNA Interference, and Transposon Silencing in *C. elegans*" vol. 99 (pp. 123-132), Oct. 15, 1999.
- Thombury et al. "Purification and Characterization of Potyvirus Helper Component" *Virology*, vol. 144 (pp. 260-267), Mar. 4, 1985.
- Urcuqui-Inchima et al. "Potyvirus Helper Component-Proteinase Self-Interaction in the Yeast Two-Hybrid System and Delineation of the Interaction Domain Involved" *Virology*, vol. 258 (pp. 95-99), 1999.
- van den Boogaart et al. "Can We Explain RNA-Mediated Virus Resistance by Homology-Dependent Gene Silencing?" *MPMI*, vol. 11, No. 7 (pp. 717-723), 1998.
- van der Krol et al. "Flavonoid Genes in *Petunia*: Addition of a Limited Number of Gene Copies May Lead to a Suppression of Gene Expression" *The Plant Cell*, vol. 2 (pp. 291-299), Apr. 1990.
- Vance et al. "RNA Silencing in Plants—Defense and Counter Defense" *Science*, vol. 292 (pp. 2277-2280), Jun. 22, 2001.
- Voinnet et al. "Suppression of gene silencing: A general strategy used by diverse DNA and RNA viruses of Plants" *PNAS*, vol. 96, No. 24 (pp. 14147-14152), Nov. 23, 1999.
- Walhout et al. "Protein Interaction Mapping in *C. elegans* Using Proteins Involved in Vulval Development" *Science*, vol. 287 (pp. 116-122), Jan. 7, 2000.

Waterhouse et al. "Virus resistance and gene silencing in plants can be induced by simultaneous expression of sense and antisense RNA" *Proc. Natl. Acad. Sci. USA*, vol. 95, (pp. 13959-13964), Nov. 1998.

Yang et al. "Sequence and Relationships of Sugarcane Mosaic and Sorghum Mosaic Virus Strains and Development of RT-PCR-Based RFLPs for Strain Discrimination" *Phytopathology*, vol. 87, No. 9 (pp. 932-939), May 12, 1997.

Schoonheim et al., "14-3-3 adaptor proteins are intermediates in ABA signal transduction during barley seed germination", *The Plant Journal* (2007), vol. 49, pp. 289-301, Sep. 25, 2005.

Stoica et al., "Interactions between the RNA Interference Effector Protein Ago1 and 14-3-3 Proteins", *Journal of Biological Chemistry* vol. 281 No. 49 pp. 37646-37651, Dec. 8, 2006.

Eiko Eurya Kuramae et al. "Identification of 14-3-3-like protein in sugarcane (*Saccharum officinarum*)" *Genetics and Molecular Biology*, 24 (1-4), pp. 43-48, 2001.

Yekti Asih Purwestri et al. "The 14-3-3 Protein GF14c Acts as a Negative Regulator of Flowering in Rice by Interacting with the Florigen Hd3a" *Plant Cell Physiol*, 50(3): pp. 429-438, 2009.

* cited by examiner

gaattcgccc	ttatttcagc	catggcagga	gcatggaaca	ctgtgactta	caagtggagg	60
ccgaatttgg	acaacgcaag	agatgttcga	aaagtgatgg	aacattttgc	agcaaagcac	120
caagtttatg	atgcaaagcg	tgacgagag	cataacagta	gaatccttcg	caggactttt	180
gtacaagaaa	ttgcaaaagc	acctgaagag	aagacttcat	acaaacctca	gggtgtgggtc	240
gaaaaacaag	ataacaatcc	aacgatccat	ctacactatg	ttagattcaa	aaacaaggag	300
aaaaagatat	taccggagat	cactccagga	tcggtcgcaa	agttaacgcg	acggatactt	360
gagctcagca	aaactacaaa	gcttgaagtt	gaactcattg	gtaagaaaag	acgcaaactg	420
actaaactag	caatcaaaaag	gcgcagaaac	agagagtatt	tgcatgtgga	aactagacac	480
gaaacaaaca	aatttaagcg	cgttgacatc	aacatagaac	gacactggtt	tccacttgtg	540
aagaagattt	caaagtgcta	tagtcacata	tcaccaagaa	tgtacaaaaa	catgagcaaa	600
ggcgacagtg	ggttaacatt	catccagaat	ggtgagttat	ttataatccg	aggaaaacga	660
gatggcgctc	tacttaatat	tatcaccaat	gaaactcgaa	ttaatgaaat	aacttatttt	720
agcgatgctc	aggcgaacga	cttctggcga	ggttacacag	atcatatggt	cgaaaatagg	780
ttaatttcta	caactcatac	agaacacata	cccacaataa	atntagagaa	gtgtggaaag	840
aggatggcat	tgtagagat	attgtttcac	tccacattca	aaattacgtg	taagcactgt	900
aacaatgacg	atcttgaact	atcggatgat	gagtttgag	aaagactata	taagaactta	960
atcagaattg	aagaaaagca	aaaagaatat	ttagctgaag	atcaaaagct	taagcgaatg	1020
atatactttc	tgaaggatag	atgcaatcca	aaatttgagc	atctaccatt	attatggcag	1080
gtcgtgtaaa	caattggaca	ttacactgat	aatcaagcaa	aacagatcct	tgaagttaat	1140
gaagcgctca	taaaagtga	cactctttct	gttgaagatg	cagtcaaacg	tagcgcacgc	1200
ttgctagaga	tttcaagatg	gtacaagaat	aggaaagaat	catcgaaaga	aggtacactt	1260
agtacattca	ggaataaaaat	ttcacctaaa	agtactatta	atacagcact	gatgtgtgat	1320
aatcagctcg	atacaaatgg	taacttccta	tggggaaaga	gagaatatca	tgccaagcga	1380
ttctttacaa	actattttga	agctgttgat	ccaaaagaca	cgtatgaaa	gcatgttact	1440
cggttcaatc	caaatgggtca	acgcaaacct	tcgattggaa	aactagtat	cccattagac	1500
ttccagaaga	ttcgtgaatc	atcttatagg	gttcaagttc	aaaaacaagc	aattagttaga	1560
gcgtgcttaa	gtaaaatcga	aaataattac	atataccctt	gctgttgtgt	aactacagaa	1620
tttggccaac	cggtttattc	agagatcatt	ccaccaacta	aagggtcatat	tactatttga	1680
aattcgaccg	acccaaaaat	tgtggatttg	cctaattccg	accaccaat	gatgtacata	1740
gcgaaagatg	gttattgtta	tttgaatata	tttttagctg	ctctgataaa	cgccaatgaa	1800
gattcagcaa	aagattacac	aaagtttttg	cgtgatgaac	taattgaaag	acttggaaag	1860
tggccaaaac	tcaaagacgt	ggcgacagca	tgttatgcat	tatcagtaat	gtttccagaa	1920
attaagaacg	cggagcttcc	acaaatacta	gtggaccacg	aacataaaac	catgcatgtg	1980
atagattcgt	acggatctct	cagtgttggc	ttccacatac	tcaaagcgaa	cacaatagga	2040
caattaatca	aaatgcaata	tgaatccatg	gaaagtgaaa	tgagagagta	tgtagtccgt	2100
tag						2103

(SEQ ID NO: 1)

Figure 1A

MAGAWNTVTVY	KWRPNLDNAR	DVRKVMHFHA	AKHQVYDAKR	AAEHNSRILR	RTFVQEIAKA	60
PEEKTSYKPQ	VWVEKQDNNP	TIHLHYVRFK	NKEKKILPEI	TPGSVAKLTR	RILELSKTTK	120
LEVELIGKKR	RKSTKLAIKR	RRNREYLHCE	TRHETNKFKR	VDINIERHWF	PLVKKISKCY	180
SHISPRMYKN	MSKGDSGLTF	IQNGELFIIR	GKRDGVLNS	ITNETRINEI	TYFSDAQAND	240
FWRGYTDHMV	ENRLISTTHT	EHIPTINLEK	CGKRMALLEI	LFHSTFKITC	KHCNDDLEL	300
SDDEFGERLY	KNLIRIEEQ	KEYLAEDQKL	KRMISFLKDR	CNPKFEHLPL	LWQVAETIGH	360
YTDNQAKQIL	EVNEALIKVN	TLSVEDAVKA	SASLLEISRW	YKNRKESKE	GTLSTFRNKI	420
SPKSTINTAL	MCDNQLDTNG	NFLWGKREYH	AKRFFFTNYFE	AVDPKDTYEK	HVTRFNPNGQ	480
RKLSIGKLV	PLDFQKIRE	FIGVQVQKQA	ISRACLSKIE	NNYIYPCCCV	TTEFGQPVYS	540
EIIPPTKGHI	TIGNSTDPKI	VDLPNSDPPM	MYIAKDGYCY	LNIFLAALIN	VNEDSAKDYT	600
KFLRDELIER	LGKWPCLKDV	ATACYALSVM	FPEIKNAELP	QILVDHEHKT	MHVIDSYGSL	660
SVGFHILKAN	TIGQLIKMQY	ESMESEMREY	VVG*			694

(SEQ ID NO: 2)

Figure 1B




Figure 2




Figure 3




Figure 4




Figure 5

```

ggccattatg gccggggaga aactgtatg aagaacatgg ccgcttctta tagtcaagac      60
ggcgagctcg atctcg tactgacgct cccgtcgatg ctagcgcgtc tcgttctgaa      120
cattcatctc agcttgctag ctctaactgg agatctgtga ttcaaaactc cccacctgat      180
ctcctatgcg gatgcggtag accggcaatt aggcgcacgg cagagactgc gaagaacaat      240
ggccgcatct ttgcacgctg tccggcgtgc aaaatatgga tttggcagga tctgctggac      300
agctatgtga atgctttgat aagctactgt cgtgatgcct ccattgattc ccttcagtca      360
gagcttgaat ctagccggtt ataatcttc gagaagcagg cacagatttc acgtttggag      420
aaacaattgg agacgctgca gccacttacc tcaaaatata ctgaacaatc tcgcagcatt      480
gctcaggcct ccattccatc atcacttttt ttcttgggaag cctgcagtct ccgacatcag      540
cggcggaggg tgaatgaaaa tcaaggaggg gctaatttca aaagaagcta ctggagcgat      600
taaagcttcg gttaaaaata agcaagaatc taacaccagc agcacaatc ccatgagctg      660
gctttttttt gggacaccct ttcatttttc atcaaaaaag ggggggcacc cccagtttcc      720
tccaaaaggc tcccctgtc cgacatcata ggtgatgtga ttacccaaaa acaggttgtc      780
ccgcttgctg actcgatgcc aaatttggat tcaatgctgc tcctgttgtt ttaacaatca      840
atcattttga ctaaaagcat tcccctaaa atgtttgta aatttattgt caaacttatt      900
accgcaaagt ccgttggcag gtaatcccc ccttttt      937

```

(SEQ ID NO: 3)

Figure 6A

```

GHYGRGEHCM KNMAASYSQD AQLDLVLPDA PVDASASRSE HSSQLASSNW RSVIQNSPPD      60
LLCGCRPAI RRTAETAKNN GRIFRTCPAC KIWIWQDLLD SYVNALISYC RDASIDSLQS      120
ELESSRLIS EKQAQISRLE KQLETLQPLI SKYTEQSRSI AQASIPSSLF FLEACSLRHQ      180
RRRVNENQGG ANFKRSYWS

```

(SEQ ID NO: 4)

Figure 6B


```
gaattcggcc attatggcgc gggcatgtgc agtgaatgcg ccaaggtcct gaggtaccaa      60
accactcggg gccccatctg caggcagcct gttgagcgtc tcctcgagat caaagtgagc      120
aacaatctg aagagcagca gcagacgcc caatcgcgc cgctcccagc cccagctctg      180
cagcaggaag aggtgtagcc gtgattaaag tcagttctga gacattatat ggaactagtt      240
tgcggccttc aggcctttcc ctaaaggttt gttctctcat ctgagcaacg gggaatgtaa      300
ccggtacttt acctttagcc tatgtaagct tctggcatcg catggetttg ccgacctctg      360
ctgtacctgc ttatctggag gtcggagacc aagatgccaa ggaaagtgtg taccgtatat      420
taaaaaaaaa aaaaaaaaaa aaaaaaaaaa
```

(SEQ ID NO: 5)

Figure 7A

```
EFGHYGRGMC SECAKVLRYQ TTRCPICRQP VERLLEIKVS NKSEEQQTP QSPPLPAPAL      60
QQEEV*P*LK SVLRHYMELV CGLQAFP*RF VLSSEQRGM* PVLYL*PM*A SGIAWLCRPL      120
LYLLIWRSET KMPRKVCTVY *KKKKKKKK
```

(SEQ ID NO: 6)

Figure 7B

gaattcggcc	attatggccg	ggaccaggaa	ctccaggaat	cagatgacaa	ctcgggggat	60
agagctcctg	aagtgaccat	gtccggtcag	tattctcaaa	agagtgatgt	ttacagcttt	120
gggtgctgca	tgcttgagct	actgactgga	cagaaagcat	ttgacagctc	tcgggcaagg	180
tcccagcaat	cactagtccg	gtgggcttca	ccgcagctgc	acgacatcga	ctcgttagat	240
cagatgggtg	atccaacctt	agaggggctg	taccatgcga	aatcactctc	tcggttcgca	300
gacgcaatcg	ctctctgtgt	ccagcctgaa	ccagaattca	ggccaccaat	gtcggaggtc	360
gtccagtcac	tggtcctctt	tgtgcagcga	gcaagcatgg	ggacagcact	aagcagcgag	420
tgggaattctt	gccagttcga	tgaatctggt	gatcacacgc	tctaggggaa	aatgatgtgt	480
atctcctaga	gagtcctgat	aggaaactata	gaaggctcac	aagtcataga	aacttgcagc	540
ttggcattgt	tgtgagttgt	gacggtgtga	catgtgccag	tgtcaggtga	aatgtgactt	600
ttacctatgc	cattttactg	agagtctgct	gcaacctgaa	gtaggggtga	aaagaaagtt	660
ccttctttaa	aaatatatat	ggttcattcg	gacgtgtata	tgaatatctt	ttgaagacaa	720
tcaactttct	gatttcgtct	ctgacgctg	tccaaaaatt	atcaggggag	atgtagcact	780
agtccctgca	cagaattagt	catctgtata	tctcagaaa	tccgaacct	atccagggaa	840
catcaacaga	ggacacgtcc	acatattcga	ac			872

(SEQ ID NO: 7)

Figure 8A

EFGHYGRDQE	LQESDDNSGY	RAPEVTMSGQ	YSQKSDVYSF	GVVMLELLTG	QKAFDSSRAR	60
SQQSLVRWAS	PQLHDIDSLD	QMVDPTLEGL	YHAKSLSRFA	DAIALCVQPE	PEFRPPMSEV	120
VQSLVRLVQR	ASMGTALSSE	WNSCQFDESG	DHTL*GKMMC	IS*RV*GTI	EGSQVIETCS	180
LALL*VVTV*	HVPVSGEM*L	LPMPFY*ESA	AT*SRGEKKV	PSLKIYMVHS	DVYMNIF*RQ	240
STF*FRL*SL	SKNYQGRGST	SPATELVICI	SSEIRTISRK	HQQRTRPHIR		290

(SEQ ID NO: 8)

Figure 8B

```

gaattcggcc attatggccg gggaccgcag ttccccgcac cacaccgttc cgccgcgcac 60
agaggccagc cccgcgccag gagtaagttt gttcttttta acaatatgtc gagggaggag 120
aatgtttaca tggccaagct ggctgagcag gccgaaaggt atgaggagat ggttgagtat 180
atggagaag tggctaagac tgtagatggt gaagagctca ctgtggagga gcgtaacctc 240
ctgtctgtcg catacaagaa tgtgattggg gctcgccgtg cttcatggcg cattgtctct 300
tccattgaac agaaggagga gtcccgtaa aacgaagagc atgtgaacct tatcaaggaa 360
taccgcgga agattgaggc tgaactgagc aacatctgtg atggcatcct gaaactgctt 420
gactcccacc tagtgccttc ctctactgct gctgaatcaa aggtcttcta cctcaagatg 480
aagggtgact atcacaggta tcttgccgaa ttttaagactg gtgctgagag gaaggaatct 540
gctgagagca caatggtagc ctacaaggct gctcaggaca ttgctctggc tgagctggca 600
cctacacatc cgataaggct tgggcttgc ctttaacttct cagtgttcta ttatgagatt 660
ctgaactccc cagacaaaagc ttgcaacctt gcaaagcagg cgtttgatga agctatctct 720
gagttagaca cccttgggga ggagtcatc aaagatagca ctctgatcat gcagctcctg 780
agggacaact tgaccctttg gacctctgac ctcacggagg atggtgctga tgagggcaaa 840
gaagcctcaa aaggtgatgc tggcgagggg cagtaatctt cggagagggc atggttgttc 900
agcctggttt tagatgctct atgctgtcga agctgtgccg tgccattatt gtagcagatt 960
tcctctcccc ctcacttcat ttgcctcata ttagtaggct ggtagtggtc gaattagttc 1020
ccattgcttt gtggttcagc tagttggcac taggtccgtg tggactggta ttgttcccct 1080
ggatttgaca agcatgtcct gtggtcgctc tagcgtttta ttgagctttg aagcctcgat 1140
t 1141

```

(SEQ ID NO: 9)

Figure 9A

```

EFGHYGRGPQ FPRPHRSAAH RGQPRARSKF VLFNMSREE NVYMAKLAEQ AERYEEMVEY 60
MEKVAKTVDV EELTVEERNL LSVAYKNVIG ARRASWRIVS SIEQKEESRK NEEHVNLIKE 120
YRGKIEAELS NICDGILKLL DSHLVPSSTA AESKVFYLMK KGDYHRYLAE FKTGAERKES 180
AESTMVAYKA AQDIALAELA PTHPIRLGLA LNFSVFYIEI LNSPKACNL AKQAFDEAIS 240
ELDTLGEESY KDSTLIMQLL RDNLTWTSR LTEDGADEGK EASKGDAGEG Q*SSERACCS 300
SLVLDALCCR SCAVPLL*QI SSPPHFICLI LVGW*WSN*F PLLCVAASWH *VRVDWYCSP 360
GFDKHLWLSL *RFIEL*SLD 380

```

(SEQ ID NO: 10)

Figure 9B

```

gaattcggcc attatggccg ggctgttcaa gctggaccgt tatatggat gggacacat 60
ggatcttcca ccacaattgc ttatggcggg gcatacttgc catattcttc ctcaactgga 120
caatcgagca ataatacatca agagcatgga ttctctgagc ggccagggca gcctgagtgt 180
caatatttta tgaggactgg aggttgcaaa tttggaacta tgtgtaaata taaccatcct 240
cgagattgga gcactcctaa gtccaactac atgttcagtc atctctgcct tccacttctg 300
ccgggtgctc agccttgtgc gtactatgca caaaatggat attgcagata tggagtgtgca 360
tgcaaataatg atcacccaat gggtagacta ggctacagtt catctgcttt acccctatct 420
gacatgccaa ttgctcccta ccctatcggc ttctctgttg ccacgttggc tccatcttca 480
tcttccccag aatatatttc aaccaaagat ccatcaatca accaagtagc atcaccagtg 540
cagcaccoga acatgttggg acaatcttgc caaaaggggt tcccttcggg atccattatg 600
cgaactcaac ttctacaagt gtccgcagtt caagcctggg gggcgctgat tttctgactg 660
ggggatgatc cttaacacaa atttctatac ttgaacagtt tgaagccttc aaggaataaa 720
aactggggcc ttgaaaaacc gggagggggt cttcccaaat aaaactgtgg tcaaacactca 780
tcttgaattg gtttctatt caaacggaag aggttttaga gtcacattg 829

```

(SEQ ID NO: 11)

Figure 10A

```

EFGHYGRAVQ AGPLYGMGHH GSSTTIAYGG AYL PYSSSTG QSSNNHQEHG FPERPGQPEC 60
QYFMRTGGCK FGTMCKYNHP RDWSTPKSNY MFSHLCLPLR PGAQPCAYYA QNGYCRYGVA 120
CKYDHPMGTL GYSSSALPLS DMPIAPYPIG FSVATLAPSS SSPEYISTKD PSINQVASPV 180
QHPNMLEQSC QKGFPPGSIM RTQLLQVSAV QAWGALIF*L GDDP*HKFLY LNSLKPSRNK 240
NWGLEKPGGV LPK*NCGQHS S*IGFLFKRK RFRSHI 276

```

(SEQ ID NO: 12)

Figure 10B




Figure 11A




Figure 11B




Figure 12




Figure 13




Figure 14

Figure 15





Figure 16

MWM His-S/14-3-3 His-S BSA HEWL RNase A



93

50

36

29

21





Figure 17




Figure 18




Figure 19




Figure 20




Figure 21

ISOLATION OF PROTEINS INVOLVED IN POSTTRANSCRIPTIONAL GENE SILENCING AND METHODS OF USE

CLAIM TO PRIOR APPLICATIONS

The present application is a divisional application of U.S. application Ser. No. 11/252,080 filed on Oct. 17, 2005 and published as US 2006 0090217, now abandoned, which is a divisional application of U.S. application Ser. No. 10/226,715 filed on Aug. 23, 2002, published as US 2003 0099984 on May 29, 2003 and issued as U.S. Pat. No. 7,001,739 on Feb. 21, 2006, both incorporated by reference herein.

TECHNICAL FIELD OF THE INVENTION

The present invention relates generally to sugarcane protein isolation and more particularly to isolation and characterization of proteins that are involved in posttranscriptional gene silencing. The invention also includes sugarcane and sorghum mosaic virus proteins involved in silencing and the cDNAs which encode them. Finally the invention includes use of these cDNAs and proteins to regulate silencing.

BACKGROUND OF THE INVENTION

In the course of evolution organisms have developed a spectrum of defense mechanisms that target alien, parasitic elements. The most specialized defense system is perhaps the vertebrate immune system, which provides effective protection against a wide range of infectious microbes. In the vertebrate immune system, it is essentially peptides that are recognized as non-self and eliminated. Transgenic plant studies have revealed the existence of another, more ancestral level of defense response. Homology-dependent gene silencing can be viewed as a novel, innate host defense system that is capable of recognizing foreign nucleic acids as non-self and inactivating or removing them from the cell. First recognized in plants and fungi, homology-dependent gene silencing mechanisms have now been shown to operate in a wide range of eukaryotic organisms. In plants, this type of gene silencing may occur at the level of DNA, by inhibition of transcription, or at the level of RNA, by enhanced RNA turnover. Both viral RNA and transgenes are subject to these host surveillance systems, which are only poorly understood at the molecular level. At the core of both silencing events resides a molecular mechanism that is able to recognize nucleic acid sequence homology.

Transgene-induced gene silencing in plants was originally described as the coordinated suppression of transgenes that share sequence similarity (Depicker and Van Montagu, 1997). This phenomenon is most often induced when multiple copies of a transgene are present at a single locus. Silencing not only affects all genes in that locus, i.e. in cis, but also acts in trans, and additionally down-regulates the expression of other, unlinked transgene(s). Silencing can also affect the expression of endogenous genes, provided they have sequence similarity to the silencing transgene, a phenomenon referred to as cosuppression (Napoli et al., 1990; Van der Krol et al., 1990).

In plants, cases of transgene-induced gene silencing belong to two different mechanistic classes: those that occur at the level of transcription and those that are due to enhanced RNA turnover. Transcriptional gene silencing (TGS) requires sequence identity in the promoter region and is associated with methylation and inactivation of the promoter sequences of the affected genes (Kumpatla et al., 1998). In posttranscrip-

tional gene silencing (PTGS), the (trans)genes remain actively transcribed but the steady-state RNA levels are highly reduced due to sequence-specific RNA degradation. Some instances of PTGS are associated with DNA methylation located in the transcribed portion of the genes (Ingelbrecht et al., 1994; 1999).

The expression level, number and configuration of the integrated transgenes as well as developmental and environmental factors can all influence the occurrence of transgene-induced gene silencing. Importantly, transgene-induced gene silencing in plants is reversible, and in the absence of the silencer locus, expression of endogenous genes or other transgenes can be restored to normal. The changes in gene expression are therefore not due to irreversible changes in DNA but rather are epigenetic.

PTGS behaves as a non-clonal event and, in agreement with this, it has been shown that a sequence-specific signal is involved in the systemic spread of PTGS (Palauqui et al., 1997; Voynet and Baulcombe, 1997). These experiments allow differentiation of separate initiation and maintenance phases in PTGS and further suggest that a molecular system amplifies the silencing signal during the course of long-distance movement of PTGS. Mutants that enhance (Dehio and Schell, 1994) or suppress (Elmayan et al., 1998; Mourrain et al., 2000; Dalmay et al., 2000) PTGS have been isolated in *Arabidopsis thaliana* but only two of the corresponding genes have been cloned (Mourrain et al., 2000; Dalmay et al., 2000). One of these has no significant similarity with any known or putative protein (Mourrain et al., 2000) and the other is similar to a RNA-dependent RNA polymerase (RdRp; Mourrain et al., 2000; Dalmay et al., 2000). Establishment of PTGS in plants requires separately identifiable initiation, spread, and maintenance phases, but the proteins involved in these pathways have not been characterized.

Plant virus studies have greatly contributed to the current understanding of gene silencing in general and PTGS in particular. Applying the concept of pathogen-derived resistance, viral genes were introduced into plants and resulted in virus resistant phenotypes. Many resistance phenotypes do not require the expression of a functional protein but are mediated at the level of RNA. It is now an established fact that a mechanism similar to PTGS is the underlying molecular mechanism in most of these cases (van den Boogaart et al., 1998).

Posttranscriptional silencing of an endogenous plant gene or transgene can be triggered by replication of a recombinant virus that carries sequences homologous to these genes (Kumagai et al., 1995; Ruiz et al., 1998). This process involves sequence-specific RNA turnover, similar to PTGS induced by transgenes, hence the term virus-induced gene silencing. Moreover, natural virus infection of non-transgenic plants can induce a resistance mechanism that is strain-specific and targeted against RNA, similar to RNA-mediated resistance induced by (silenced) transgenes (Ratcliff et al., 1997; Covey et al., 1997). Transgene- and virus-induced gene silencing are collectively described as homology-dependent gene silencing because these mechanisms all target homologous nucleic acid sequences. It was proposed that homology-dependent gene silencing acts as a natural plant defense mechanism against invading DNA or RNA elements (Matzke and Matzke, 1998).

The demonstration that plant viral proteins can suppress PTGS provides direct evidence that PTGS functions as a host defense response in plants (Anandalakshmi et al., 1998; Brigneti et al., 1998; Kasschau and Carrington, 1998). At least 5 different proteins encoded by unrelated DNA and RNA viruses of plants have now been shown to act as sup-

pressors of PTGS in *Nicotiana benthamiana*. Importantly, the suppression phenotypes induced by these viral proteins are distinct indicating that separate steps of the host PTGS defense system are targeted. For example, the potyviral helper-component proteinase (HC-Pro) can reverse the effects of PTGS in tissues that were previously silenced, whereas the 2b protein of Cucumber mosaic virus only affects initiation of PTGS (Voinnet et al., 1999). Although potyviral HC-Pro by itself is sufficient to suppress transgene-induced silencing, it appears that the potyviral P1 protein can enhance its ability to suppress virus-induced gene silencing (Anandalakshmi et al., 1998; V. Vance). The discovery of viral suppressors of silencing phenomena is unique to plants. So far, no animal or fungal viruses have been shown to suppress PTGS in these organisms.

It has been proposed that 'aberrant' RNA molecules trigger PTGS in plants (Lindbo et al., 1993). The exact nature of this aberrant RNA is unknown but it could be double-stranded RNA (dsRNA) (Waterhouse et al., 1998), prematurely terminated transcripts, levels of RNA that exceed a certain threshold, or some other unusual characteristic. These RNA molecules would serve as templates for an RNA-dependent RNA polymerase (RdRp) and lead to the production of short complementary RNAs (cRNA). These cRNAs would then anneal with homologous mRNAs or viral RNAs and the resulting double-stranded RNA would be degraded by double strand-specific RNases. This model accounts for the sequence-specific RNA turnover and several aspects of it are supported by experimental data. For example, an RdRp that is induced during viral infection has been cloned in tomato (Schiebel et al., 1998) and small cRNAs have recently been identified in transgenic plants that display PTGS (Hamilton and Baulcombe, 1999). The identification of a double strand-specific RNase in *Caenorhabditis elegans* and a RdRp-like protein in *Neurospora crassa*, and recently in *Arabidopsis*, as essential components of PTGS-like mechanisms in these organisms (see below) provides further support for this hypothesis.

RNA-mediated genetic interference (RNAi) in *C. elegans* is a process that closely resembles PTGS in plants: both act at the posttranscriptional level and result in sequence-specific RNA turnover (Tabara et al., 1998; Montgomery and Fire, 1998). The trigger for RNAi in *C. elegans* is well characterized and consists of dsRNA (Sharp, 1999). RNA-specific silencing can be induced by locally injecting homologous dsRNA molecules in a few cells. Silencing then spreads from the site of injection into neighboring cells and tissues and is even transmitted to the F1 progeny. The ability of silencing to move both in space and over time strongly suggests that amplification of the silencing signal is taking place, similar to PTGS in plants.

Recently, several genes have been identified in *C. elegans* that are required for this interference process. The MUT-7 gene encodes a homolog of RNaseD, which is a double strand-specific RNase (Ketting et al., 1999). The RDE-1 gene belongs to a family of genes that are conserved from plants to vertebrates and several members of this family are required for gene silencing mechanisms in animal systems (Tabara et al., 1999). Interestingly, mutations in both these genes reactivate mobilization of endogenous transposons, suggesting that one function of RNAi is transposon silencing. Sequence-specific inhibition of gene function by dsRNA has also been demonstrated in trypanosomes, *Drosophila* and *planaria* and has been used in these organisms as a method to determine gene functions (Kennerdell and Carthew, 1998; Misquitta and Patterson, 1999; Sanchez Alvarado and Newmark, 1999).

Transgene-induced PTGS is termed 'quelling' in the fungus *N. crassa* (Cogoni and Macino, 1997a). Quelling-defective (qde) mutants of *N. crassa*, in which transgene-induced gene silencing is impaired, have been isolated and could be classified in three qde complementation groups (Cogoni and Macino, 1997b). Two QDE genes that belong to two different complementation groups, have recently been cloned. The QDE-1 gene encodes a protein that contains an RdRp-motif (Cogoni and Macino, 1999a) and QDE-3 belongs to the RecQ DNA helicase family (Cogoni and Macino, 1999b).

As summarized above, there has been substantial progress in the general understanding of PTGS in plants and its importance as part of a general defense system is now fully appreciated. However, all of the biochemical pathways of PTGS and the enzymes that are involved have not yet been elucidated in plants. Insight into these mechanisms may come from analyzing mutants that are defective in PTGS. This approach has already been used with success in *Neurospora* and *C. elegans* and is currently also being followed for *Arabidopsis*. While this strategy is relatively straightforward and will surely result in the identification of genes that play a central role in this process, there are also limitations. For example, gene redundancies and possibly lethal, loss-of-function phenotypes might prevent identification of certain genes. There are also practical problems in generating and screening a sufficiently large number of mutants which limit this approach to model plants such as *Arabidopsis*.

An alternative or complementary approach involves directly identifying the host factors that mediate PTGS. The identification of viral proteins as suppressors of PTGS provides the necessary tools to pursue this strategy.

Identification and characterization of proteins that interact with a viral suppressor of PTGS will have an impact on understanding fundamentals of virus-host plant interactions, particularly on the mechanisms that plants employ to combat viral infection and on the counterdefensive strategies that viruses use to suppress or evade these responses. To date, viral suppression of PTGS is a process unique to plants. However, because PTGS is a defense mechanism that is conserved among various eukaryotic kingdoms, the identified protein interactions might also shed light on the molecular mechanisms of silencing phenomena in other organisms.

In addition to significance for basic (plant) molecular virology, establishing the biochemical pathways of host defense responses will facilitate the development of improved virus control strategies in plants. PTGS-based approaches for virus control are already in use but the lack of a solid understanding of the phenomenon necessitates a more empirical approach and has an uncertain outcome. Also, such approaches are currently limited because of their narrow range. Possible and realistic improvements involve enhanced and more predictable triggering and broadening the scope of the PTGS defense system.

Use of the method of the present invention will also contribute to plant genetic engineering in general. It is now clear that transgenes in plants (and other organisms) can be perceived as intrusive elements and consequently are inactivated. Developing procedures that allow stable and predictable transgene expression is one of the challenges of genetic engineering. The monocot crop plants provide the most important source of food worldwide and offer great potential for improvement through genetic transformation, not only for traits related to food production but also as recombinant expression systems for high value products.

Finally, gene silencing can be used as a way to produce 'knock-out' phenotypes in reverse genetic studies. This has already been successfully applied in animal systems and its

potential has been demonstrated in plants. With an increasing number of genes being discovered in sugarcane, many of which have no known function, it can be expected that these approaches will become even more important in the future.

Thus, the yeast two-hybrid method of the present invention has been used to unravel the pathway(s) of PTGS and plant defense responses and novel, key proteins involved in this process have been identified. In doing so, a cDNA library from silenced plant tissues rather than non-silence plant tissues has been used. These proteins and genes can be applied towards regulating PTGS of transgenes, endogenous plant genes, and viral genes. Specific applications of the present invention include but are not limited to, improved strategies for engineered virus resistance, increased expression of transgenes by inhibiting silencing, and modulation of silencing of native genes to obtain desirable traits or in functional genomic studies.

SUMMARY OF THE INVENTION

The present invention includes a method of isolating nucleic acid encoding a plant polypeptide active in PTGS. As used throughout the application, plant may mean a mature plant, an embryonic callus, or other stages of plant development. It may also mean a portion of a plant, a plant tissue, or a plant cell.

The first step of a method of the above method involves selecting a bait nucleic acid which encodes a bait protein active in PTGS in plants or suppressive of PTGS in plants. After bait selection, a cDNA prey library may be prepared from a plant that actively exhibits PTGS at the time of library generation. If an entire plant exhibiting PTGS is not available, tissues in which PTGS is exhibited may be selected.

After the bait and prey are selected, a yeast two-hybrid assay may be conducted with the bait and prey nucleic acids. Prey cDNA that yields a true positive yeast two-hybrid assay result encodes a polypeptide active in PTGS in the plant. True positive status may be verified using methods known in the art, such as null controls, reversal of bait and prey, and in vitro and in planta studies of interactions. Such in vitro assay may include farwestern blot assays and pull down assays. They may be performed under plant physiological conditions to eliminate false negatives and false positives. In planta studies may be performed in an embryonic callus or other plant tissue.

In an exemplary embodiment of the above method of the present invention, the bait nucleic acid comprises a sequence selected from SEQ.ID.NO. 1, SEQ.ID.NO. 3, SEQ.ID.NO. 5, SEQ.ID.NO. 7, SEQ.ID.NO. 9, or SEQ.ID.NO. 11. The entire nucleic acids of these sequences may be used or only portions thereof. Additionally, substituted nucleic acids and nucleic acids with similar identities may also be used.

In another exemplary embodiment the plant is a monocot, particularly sugarcane and more particularly *Saccharum* hybrid cultivar CP72-1210.

The present invention also includes several SrMV and sugarcane novel proteins and nucleic acids. Novel nucleic acids are provided in SEQ.ID.NOS. 1,3,5,7,9 and 11. Novel amino acid sequences are provided in SEQ.ID.NOS. 2,4,6,8,10 and 12. It will be apparent to one skilled in the art that portions of these nucleic acids and proteins or polypeptides may be used in various applications. Additionally, it will be apparent to one skilled in the art that nucleic acids and proteins or polypeptides with high similarity, particularly in regions related to functional domains, may also be substituted. The present invention includes such variations up to the point of disclosures already in the prior art. Each of these proteins is

involved in PTGS either as an activator or suppressor. Some proteins may fail to function alone and may rather require or assist another protein for their PTGS-related functions.

The present invention additionally includes transgenic plants including any of the above novel nucleic acids, proteins or polypeptides. In particular, the invention includes a transgenic plant in which PTGS is suppressed that includes the nucleic acid of SEQ.ID.NO. 1 or the protein of SEQ.ID.NO. 2. It also includes a transgenic plant in which PTGS is enhanced that includes the nucleic acid of SEQ.ID.NO. 3 or the protein of SEQ.ID.NO. 4.

The invention additionally includes a method of increasing viral resistance in a plant in which a protein suppressive of PTGS in the plant is selected and the plant is transformed with a nucleic acid encoding the PTGS suppressive protein.

Another method of the invention involves a method of increasing expression of a transgene in a plant by selecting a protein active in PTGS in the plant and transforming the plant with a nucleic acid encoding the protein active in PTGS.

Yet another method of the present invention involves suppressing expression of a native gene in a plant by preparing a vector including a nucleic acid with a sequence of the coding portion of the gene wherein the nucleic acid, upon transcription, products an mRNA molecule double stranded in the region corresponding to the coding portion of the gene. A plant is then transformed with the vector.

BRIEF DESCRIPTION OF THE DRAWINGS

For a more complete understanding of the present invention and the advantages thereof, reference is now made to the following description taken in conjunction with the accompanying drawings, wherein like reference numbers represent like parts, and which:

FIG. 1A provides the cDNA (SEQ.ID.NO. 1) and FIG. 1B provides the amino acid (SEQ.ID.NO. 2) sequences for Sorghum Mosaic Virus (SrMV) P1/HC-Pro according to an embodiment of the present invention;

FIG. 2 is a Northern analysis of SrMV (coat protein) CP mRNA in which the lane labeled "S" includes mRNA derived from a plant posttranscriptionally silenced for SrMV CP; lanes labeled "Retransformed S" include mRNA from samples of the silenced plant which were retransformed with either nptII alone or with nptII and SrMV P1/HC-Pro; the lane including mRNA from a plant that is not silenced is labeled "NS"; and the empty lane is labeled "b";

FIG. 3 is a 9-galactosidase filter lift assay on L40 yeast transformed with various DNA binding and activation domain constructs and grown on media that does not select for interactions (-leu +zeo) or media that does select for transformation and interaction (-leu, -his +zeo); where the region labeled "A" represents yeast transformed with SrMV HC-Pro bait+SrMV HC-Pro prey; the region labeled "B" represents yeast transformed with SrMV HC-Pro bait+empty prey; the region labeled "C" represents yeast transformed with lamin bait+SrMV HC-Pro prey; and the region labeled "D" represents yeast transformed with empty bait+SrMV HC-Pro prey; although the figure is presented in black and white, growth in the regions marked A on both plants appears blue white, on the -leu +zeo plate, growth in regions B-D appears red;

FIG. 4 is a gel analysis of the cDNA library cloned in pIVING1154; where Lane 1 contains a λ PstI size marker; Lane 2 contains a lower band of 0.8 kb which is the CP coding region amplified from the plasmid cDNA library and a top band of 6.6-kb which is the plasmid vector; and Lane 3

contains a SfiI digest of plasmid library showing the 6.6-kb vector and inserts visible as a smear with size range between 0.5 and 2.0 kb;

FIG. 5 is a β -galactosidase filter lift assay on L40 yeast transformed with various DNA binding and activation domain constructs and grown on media that does not select for interactions (-leu +zeo) or media that does select for transformation and interaction (-leu, -his +zeo); where the region labeled "A" represents yeast transformed with SrMV HC-Pro bait+SrMV HC-Pro prey; the region labeled "B" represents yeast transformed with empty bait+RNase H-like protein prey; the region labeled "C" represents yeast transformed with lamin bait+RNase H-like protein prey; and the region labeled "D" represents yeast transformed with SrMV HC-Pro bait+RNase H-like protein prey; although the figure is not presented in color, the regions marked A and D on both plates appear blue, while the regions marked B and C on the -leu +zeo plate appear red;

FIG. 6A is the cDNA sequence of the RNase H-like protein (SEQ.ID.NO. 3) and FIG. 6B is the encoded amino acid sequence (SEQ.ID.NO. 4), according to an embodiment of the present invention;

FIG. 7A is the cDNA sequence of the sugarcane RING zinc finger protein that interacts with SrMV HC-Pro (SEQ.ID.NO. 5); FIG. 7B is the encoded amino acid sequence (SEQ.ID.NO. 6) according to an embodiment of the present invention;

FIG. 8A is the cDNA sequence of the LRR (leucine-rich repeat) transmembrane protein kinase that interacts with SrMV HC-Pro (SEQ.ID.NO. 7); FIG. 8B is the encoded amino acid sequence (SEQ.ID.NO. 8) according to an embodiment of the present invention;

FIG. 9A is the cDNA sequence of a nucleic acid encoding the sugarcane 14-3-3 protein that interacts with the RNase H-like protein (SEQ.ID.NO. 9); FIG. 9B is the encoded amino acid (SEQ.ID.NO. 10), according to an embodiment of the invention;

FIG. 10A is the cDNA sequence of the sugarcane RING zinc finger protein that interacts with RNase H-like protein (SEQ.ID.NO. 11) and FIG. 10B provides the encoded amino acid sequence (SEQ.ID.NO. 12) according to an embodiment of the present invention;

FIG. 11A shows a 12% PA, CB stained gel; FIG. 11B shows a Large S-AP probed blot; in parts of FIG. 10, the molecular weight lane is indicated as "MW", lane 1 contains the *E. coli* expression product of Bugbuster™ (Novagen, Madison, Wis., affiliate of Merck KGaA, Darmstadt, Germany) Insoluble pET30 with no insert; lane 2 contains the *E. coli* expression product of Bugbuster™ Insoluble pET30 with an SrMV HC-Pro insert; lane 3 contains one preparation of Ni column purified *E. coli* expression product of Bugbuster™ Insoluble pET30 with an SrMV HC-Pro insert; and lane 4 contains a second preparation of Ni column purified *E. coli* expression product of Bugbuster™ Insoluble pET30 with an SrMV HC-Pro insert;

FIG. 12 shows a two hour exposure of a 15% SDS PAGE gel containing RNase H-like protein (lane labeled "RNase H") labeled with ³⁵S Cysteine according to the present invention; control lanes are provided in which no DNA was used in the preparation procedure (lane labeled "No DNA") and in which Luciferase DNA was used (lane labeled "Luciferase"); molecular weight markers are indicated;

FIG. 13 shows a 32 hour exposure of a farwestern blot probed under in vitro plant cell physiological conditions with a ³⁵S labeled RNase H-like protein transcription and translation (TNT) product; the lane labeled "His-S/HC-Pro" contains Ni column purified, His-S tagged SrMV HC-Pro protein

produced in *E. coli*; the lane labeled "His-S" contains the HIS-S tag only; the lane labeled "BSA" contains untagged bovine serum albumen; the lane labeled "HEWL" contains untagged hen egg white lysozyme and the lane labeled "RNase A" contains untagged RNase A;

FIG. 14 shows a 6 hour exposure of a farwestern blot probed under in vitro plant cell physiological conditions with a ³⁵S labeled RNase H-like protein transcription and translation (TNT) product; the lane labeled "His-S/HC-Pro" contains Ni column purified, His-S tagged SrMV HC-Pro protein produced in *E. coli*; the lane labeled "His-S/RNase H" contains Ni column purified, His-S tagged RNase H-like protein produced in *E. coli*; the lane labeled "His-S" contains the product in *E. coli* of the His-S tag only; the unlabeled lanes contain plant extracts which are from left to right, from: healthy sugarcane plant, SrMV infected sugarcane plant, sugarcane plant transgenic for SrMV P1/HC-Pro, sugarcane plant transgenic for delta N 12 SrMV HC-Pro.; and the lane labeled MWM contains molecular weight markers, which were used to produce the size indicators on the side of the blot;

FIG. 15 shows a 32 hour exposure of a 15% SDS PAGE gel on which Ni column pull down products of a TNT ³⁵S labeled protein/His-S tagged protein physiological incubation were run; the top lane marker indicates the source of DNA used in a TNT procedure in to obtain ³⁵S labeled protein, where "RNase" indicates the use of RNase H-like protein cDNA, "No DNA" indicates a control in which no DNA template was provided, and "Luciferase" indicates a control in which Luciferase DNA was provided; the bottom lane marker indicates the His-S tagged product produced in *E. coli*, where "His-S/HC-Pro" or "HS/HC-Pro" indicates Ni column purified His-S tagged SrMV HC-Pro protein, "His-S" indicates the His-S tag only, and "BSA" indicates untagged bovine serum albumen;

FIG. 16 shows a 3 hour exposure of a farwestern blot probed under in vitro plant cell physiological conditions with a ³⁵S labeled RNase H-like protein transcription and translation (TNT) product; the lane labeled "His-S/14-3-3" contains Ni column purified, His-S tagged sugarcane 14-3-3 protein produced in *E. coli*; the lane labeled "His-S" contains the His-S tag only; the lane labeled "BSA" contains untagged bovine serum albumen; the lane labeled "HEWL" contains untagged hen egg white lysozyme; and the lane labeled "RNase A" contains untagged RNase A.

FIG. 17A depicts a DNA construct containing sense and antisense sequences in opposite directions and separated by an intron which, following transcription to mRNA, form a double strand due to sense/antisense sequence complementation shown in FIG. 17B;

FIG. 18 depicts the pMCG161 plasmid expression vector, according to one embodiment of the present invention;

FIG. 19 depicts the results of transient expression of constructs inserted in the pMGC161 plasmid in embryonic sugarcane calli where the callus 1 is transformed with GUS under control of a Ubi promoter; callus 2 is transformed with GUS under control of a Ubi promoter and DNA that produces double stranded GUS mRNA under control of a 35Sint promoter; callus 3 is transformed with the same constructs as callus 2 and additionally with DNA that produces double stranded RNase H-like protein mRNA under control of the 35Sint promoter; callus 4 is transformed with the same constructs as callus 2 and additionally with SrMV P1/HC-Pro under the control of the Ubi promoter; callus 5 is transformed with the same constructs as callus 2 and additionally with SrMV HC-Pro-delta N 12 under control of the Ubi promoter for ease of plasmid construction to include a start ATG, the

first N-terminal amino acids of the full length SrMV HC-Pro are deleted via deletion of 36 cDNA base pairs); and callus 6 is transformed with the same constructs as callus 2 and additionally with DNA that produces double stranded GFP under control of the 35Sint promoter; although the figure is not provided in color, all darker regions in the calli of the figure are blue while lighter regions are pinkish tan, as will be apparent to one skilled in the art;

FIG. 20 is a graphical representation of three experiments such as that of FIG. 19; vertical bars represent the average of these three independent experiments while vertical line represent standard errors; treatment 1 represents transformation with GUS under control of a Ubi promoter; treatment 2 represents transformation with GUS under control of a Ubi promoter and DNA that produces double stranded GUS mRNA under control of a 35Sint promoter; treatment 3 represents transformation with the same constructs as treatment 2 and additionally with DNA that produces double stranded RNase H-like protein mRNA under control of the 35Sint promoter; treatment 4 represents transformation with the same constructs as treatment 2 and additionally with RNase H-like protein under control of the 35Sint promoter; treatment 5 represents transformation with the same constructs as treatment 2 and additionally with SrMV P1 and HC-Pro under the control of the Ubi promoter; treatment 6 represents transformation with the same constructs as callus 2 and additionally with SrMV HC-Pro-delta N 12 under control of the Ubi promoter; and treatment 7 represents transformation with the same constructs as callus 2 and additionally with DNA that produces double stranded GFP mRNA under control of the 35Sint promoter;

FIG. 21 depicts 4 embryonic calli transiently transformed with constructs including, in callus 1, GUS under control of the 35Sint promoter; in callus 2, GUS under control of the 35Sint promoter and DNA that produces double stranded GUS mRNA also under control of the 35Sint promoter; in callus 3, the same constructs as callus 2 and additionally RNase H-like protein under control of the 35Sint promoter; in callus 4, the same constructs as callus 2 and additionally DNA that produces RNase H-like protein mRNA under control of the 35Sint promoter; although the figures is not presented in color, the darker colored calli, calli 1 and 4 are blue while the lighter colored calli, calli 2 and 3 are pinkish tan as will be apparent to one skilled in the art.

DETAILED DESCRIPTION OF EXAMPLE EMBODIMENTS OF THE INVENTION

The present invention includes a novel method for determination of plant proteins active in PTGS or suppressive of PTGS. The invention may be used in both monocot and dicot plants, including sugarcane.

The methods of the present invention may be used, inter alia: (i) to identify, isolate and characterize cellular proteins that interact with these PTGS suppressive viral proteins or plant proteins involved in PTGS using a yeast two-hybrid system; and (ii) to evaluate these plant/viral protein interactions in vitro under physiological conditions and in vivo in either transient studies or in transgenic plants, such as sugarcane, expressing these viral proteins. Because many viral proteins such as P1 and HC-Pro are multifunctional proteins involved in various aspects of plant/potyvirus interactions, the methods may also be used to assess the role of interacting host proteins in gene silencing. The methods may be carried out using a combination of molecular genetics, immunological studies, transient antisense suppression studies, and plant transformation.

The overall method includes using a yeast two-hybrid system to search for plant proteins that interact with viral suppressors of PTGS to identify proteins involved in PTGS. These proteins or proteins identified as having a role in PTGS through other methods may then serve as bait to locate other proteins involved in PTGS. Because proteins that interact with other proteins involved in PTGS may be either suppressive of or active in PTGS, the yeast two-hybrid assays may identify both types of proteins. Whether the proteins involved in PTGS are active in or suppressive of PTGS may be determined through a variety of methods, including identification of motifs with particular functions, comparison with known proteins, and in planta studies.

In the present invention the bait protein may be either active in PTGS or may function as a suppressor. The prey used is derived from an expression library of the plant of interest. In an exemplary embodiment of the invention, the prey library is derived from a plant in which PTGS is occurring. This facilitates identification of proteins involved in PTGS because such proteins are actively being produced in the silenced plant and may therefore be better represented in the mRNA pool of a silenced plant than in a non-silenced plant.

After identification of an interacting prey, the bait and prey portions of the two-hybrid screen of the present invention may be reversed to help identify false positives. Comparison with controls designed to look for activation of the reporter system absent either the bait or prey may also be used to identify false positives.

Such yeast two-hybrid screens may then be followed by assays such as farwestern blots or pull down assays to further determine whether identified proteins are false positives and also to characterize their function. In planta physiological conditions may be used in such assays.

In planta studies may also be conducted using transiently or permanently transformed embryonic calli transformed with either a test protein or a DNA encoding a double stranded mRNA of the test protein (which induces PTGS of that protein) in order to further evaluate the suppressive or effective role of the test protein in PTGS.

Proteins identified in one or more of the above screens may also be further characterized by identifying putative functional domains and also be searching for overall cDNA and amino acid sequence similarity with other proteins.

Often proteins and cDNAs identified using the above methodologies may be novel and patentable. In the present invention, five such novel proteins and cDNAs have been identified: the sugarcane 14-3-3 protein and cDNA, the RNase H-like protein and cDNA, the sugarcane LRR transmembrane protein kinase and cDNA, and the two sugarcane RING zinc finger proteins and cDNAs. Additionally, the SrMV P1/HC-Pro protein and cDNA used in some embodiments of the method of the present invention are also novel.

The above cDNAs as well as other identified using the methodologies of the present invention may also be used to construct transgenic plants, plant cells and plant tissues (collectively "plant entities") in which PTGS is either enhanced or suppressed. The methodologies used in the assay methods to generate embryonic calli and other methods known to the art may be used to construct these transgenic plant entities. Transformation may be transient or permanent, depending upon the intended use of the transgenic plant entity.

Permanently transformed plant entities with increased PTGS may be virus resistant. PTGS may be suppressed in other plants to allow increased expression of a transgene for any of a variety of reasons, including improvement of plant health, adaptation to certain growing conditions, producing of a novel nutrient or vaccine, and production of a protein later

purified for medical or industrial uses. Finally, the PTGS-regulatory methods of the present invention may be used to induce PTGS of a particular gene, for instance by introducing a construct encoding dsRNA for a portion of the gene, thereby offering a novel method of producing knock-out plants.

The following examples are provided only to illustrate certain aspects of the invention and are not intended to embody the total scope of the invention or any aspect thereof. Variations of the exemplary embodiments of the invention below will be apparent to one skilled in the art and are intended to be included within the scope of the invention.

EXAMPLES

Example 1

Nucleotide Sequence of Sorghum Mosaic Virus

SrMV is a member of the genus Potyvirus and can cause mosaic disease and yield loss in poaceous plants such as sugarcane and sorghum (Shukla et al., 1994). A 2.0-kb region located at the 3' end of the SrMV strain H genomic RNA which encompasses the 3' untranslated region, the complete open reading frame for the coat protein and part of the NIb ORF has been previously sequenced (Yang and Mirkov, 1997). The remaining part of the SrMV genomic RNA has been sequenced in the present invention from overlapping RT-PCR products. From the combined sequences, a 9,581 bp consensus was derived that contains a continuous ORF encoding a 3079 amino acid putative polyprotein with ten gene products typical for members of the genus Potyvirus (Ingelbrecht et al., in preparation). The SrMV polyprotein consensus sequence was compared to that of Tobacco etch virus (TEV-HAT), Maize dwarf mosaic virus (MDMV-Bu), Plum pox virus (PPV-D) and Pea seedborne mosaic virus (PSbMV-D) in a multiple sequence alignment using Clustal X. Based on this alignment, putative proteolytic cleavage sites were positioned in the SrMV polyprotein. (See FIG. 1.) A novel nucleic acid (SEQ.ID.NO. 1) encoding a novel SrMV P1/HC-Pro protein (SEQ.ID.NO. 2) was developed.

Example 2

SrMV P1/HC-Pro Reverses PTGS

The potyviral HC-Pro is a multifunctional protein and harbors at least 3 functional domains. The N-terminal part contains a highly conserved KITC motif and is involved in aphid transmission (Thornbury et al., 1985; Atreya and Pirone, 1991). The central part is required for long distance movement and virus amplification (Cronin et al., 1995; Kasschau et al., 1997) while the C-terminal part represents a papain-like proteinase involved in polyprotein processing (Carrington et al., 1990). In addition, potyviral HC-Pro is the determinant of potyviral synergistic interactions with other viruses (Pruss et al., 1997).

Potyviral HC-Pro can reverse the effects of PTGS in tissues that were previously silenced. This suppression phenotype has been observed for the potyviral HC-Pro protein of three different potyviruses, TEV, PVY and PSbMV, suggesting that this particular function of potyviral HC-Pro may be conserved among different potyviruses (Voynet et al., 1999). From this combined with the presence of conserved amino-acid sequence motifs in the SrMV HC-Pro, it was postulated that the SrMV HC-Pro might also act as a suppressor of PTGS in sugarcane. However, the suppressor ability of SrMV HC-Pro could not be assumed based on sequence similarity alone

and experiments were required to determine if suppressor activity was exhibited in sugarcane and the extent of such suppression.

Reversal of PTGS by SrMV P1/HC-Pro was demonstrated by retransforming a plant which is posttranscriptionally silenced for the CP transgene (Ingelbrecht et al., 1999, plant #16). The silenced plant #16 had originally been generated by selection on bialaphos. Embryogenic callus generated from plant #16 was therefore retransformed with either nptII alone or with nptII and SrMV P1/HC-Pro, and transgenic plants were selected on geneticin. As shown in the Northern blot analysis of the CP gene in FIG. 2, in 5 of the 6 plants retransformed with nptII and SrMV P1/HC-Pro, PTGS has been suppressed and the steady state levels of the CP mRNA are equivalent to the levels seen in a plant that is not silenced (in FIG. 2, NS=plant #463 of Ingelbrecht et al., 1999). No such reversal was seen in plants retransformed with nptII alone. Southern and Northern analysis of nptII and SrMV P1/HC-Pro on these plants revealed that the one plant retransformed with nptII and SrMV P1/HC-Pro in which PTGS was not reversed was only transgenic for nptII (data not shown).

Example 3

The Yeast Two-hybrid System

The yeast two-hybrid system has become a powerful tool for the study of protein-protein interactions *in vivo*. It can also be used to search an expression library for proteins that interact with a target protein. This latter application is being increasingly used with success in both animal (Kleiman and Manley, 1999) and plant systems (Kohalmi et al., 1998; Gindullis et al., 1999). Currently, high throughput two-hybrid procedures are being developed to catalogue protein-protein interactions on a genome-wide scale (Walhout et al., 2000).

Before embarking on a screen, it is useful to study the behavior of the target protein, termed bait, in the yeast cells. For example, one should demonstrate that the bait protein by itself does not activate transcription of the marker genes that are used in the selection and screening procedures. False positives can still be encountered following this precaution, but the great majority of these can be eliminated in subsequent steps. Finally, protein-protein interactions identified in the yeast cells may be verified via independent experimental procedures, preferably in the original biological system or a physiologically equivalent environment.

A yeast two-hybrid screening procedure typically consists of the following steps:

1) Subclone the coding region of the target protein into a LexA or GAL4 binding domain vector to create a bait. In an example of the present method this bait is the SrMV HC-Pro sequence cloned into pHYBLexZeo to generate pIVING1281 or SrMV P1 cloned into pAS2 to generate pIVING1088-6.

2) Transform the bait into the yeast reporter strain to verify that the expressed protein does not activate the reporter gene(s) by itself and is not toxic to the yeast cells. In an example of the present method the yeast strain L40 is transformed with either pIVING1168, pIVING1281, or pIVING1088-6 alone, and also co-transformed with either a pGAD424 derivative (pIVING1154) that lacks an insert, empty pHYBLexZeo, or pHYBLexZeo containing a non-specific bait (lamin). These sets of transformed yeast strains function as negative controls for the screening procedures as known in the art.

3) Subclone a cDNA library of sequences into a GAL4 activator domain vector to create a library of prey. In an example of the present system, the cDNA library is derived

13

from a transgenic sugarcane plant that displays PTGS (Ingelbrecht et al., 1999, plant #16) and is cloned into a modified pGAD424 derivative (pIVING1154), which constitutes the prey.

4) Transform the yeast reporter strain carrying the bait plasmid with the cDNA/prey library.

5) Screen yeast bait and prey co-transformed colonies for expression of a reporter gene that is fused to a GAL4 activated promoter. In an example of the present system, the yeast strain L40 contains reporter constructs for expression of a β -galactosidase gene, as well as a reporter construct for the HIS gene. Cells in which bait and prey fusion proteins interact in the yeast nucleus will grow on minimal media in the absence of histidine and will produce β -galactosidase enzyme activity at levels elevated from the negative control cells.

6) Verify positive interactions of co-transformants and eliminate false positives by reestablishing and retesting yeast strains, testing yeast strains which only contain the previously identified prey construct and testing the prey construct with empty bait or bait that contains a non-specific protein (such as lamin). If the prey construct activates transcription of a reporter gene that is under regulation of a GAL4-inducible promoter under any of the above conditions, the prey may be considered a false positive.

This method, as used in the examples of the present system, may be used to isolate host proteins from sugarcane and other plants that are involved in PTGS using the HC-Pro and P1 proteins from SrMV or other proteins involved in PTGS as bait. Experiments using a library-scale yeast two-hybrid screen with SrMV HC-Pro as bait have identified 441 interacting proteins. At least 12 of these (including a protein with high similarity to a viral RNase H referred to herein as the "RNase H-like protein") have been confirmed to be true interactors. Additional proteins involved in PTGS, such as a novel sugarcane 14-3-3 protein, have also been identified using the above method based on their ability to interact with the RNase H-like protein used as bait.

Example 4

Self-interaction with HC-Pro or P1 in a Yeast Two-Hybrid System

The yeast two-hybrid method of this invention may be used to identify proteins from sugarcane that interact with HC-Pro or P1 of SrMV. The two-hybrid procedure is based on the reconstruction of a functional transcriptional activator such as GAL4 or LexA, whose DNA binding domain (DBD) and transcription activation domain (AD) are expressed on two different vectors (Fields and Song, 1989). In the present example, the bait protein, i.e. the SrMV HC-Pro, was fused to the DNA-binding domain and the cDNA library was constructed in the activation domain vector, which produces the prey. These vectors were introduced into the yeast strain L40, which has an endogenous β -galactosidase (*lacZ*) reporter gene and the nutritional marker HIS3 for selection.

A 1.4-kb RT-PCR fragment encompassing the complete open reading frame of HC-Pro was amplified from SrMV virion particles and subcloned in pCR4-TOPO by T/A cloning (Invitrogen, Carlsbad, Calif.), yielding pIVING1148-1. The complete HC-Pro ORF was cloned as a fusion protein into the two-hybrid vectors pHYBLexZeo (bait) and pGAD424 (prey), yielding the plasmids pIVING1281 and pIVING1168, respectively. The P1 ORF was subcloned as a 0.78-kb fragment in pAS2 yielding pIVING1088-6. The vector-insert junctions were confirmed by sequencing for all constructs.

14

Recent studies have shown that potyviral HC-Pro proteins can interact with themselves in a yeast two-hybrid system (Urcuqui-Inchima et al., 1999; Guo et al., 1999), in agreement with an earlier proposal that the potyviral HC-Pro is biologically active as a homodimer (Thornbury et al., 1985). The ability of the SrMV HC-Pro for self-interaction was tested in the present example. It was also verified that the SrMV HC-Pro bait construct does not activate transcription of the *lacZ* gene, either by itself or in combination with empty bait or prey vectors. These latter experiments were also conducted for the P1 bait construct pIVING1088-6. As shown in FIG. 3, the SrMV HC-Pro does interact with itself (A) in the two-hybrid system demonstrating the functionality of this construct. Secondly, no *lacZ* expression can be observed in combination with either empty prey (B) or empty bait (D) vectors or a bait vector containing a non-specific protein (lamin; C). Therefore, SrMV HC-Pro does not activate transcription by itself and can be used as bait to screen for interacting proteins. Similar results were obtained for the SrMV P1 bait construct (data not shown).

Example 5

Development of a cDNA Library from Silenced Sugarcane

The development of SrMV-resistant sugarcane plants via transformation with an untranslatable form of the capsid protein sequence has been previously described and it has been demonstrated that the underlying resistance mechanism is related to PTGS (Ingelbrecht et al., 1999). As described in Ingelbrecht et al., 1999, plant # 16 is a recovery plant that is immune to infection with SrMV after recovery from the initial infection and is posttranscriptionally silenced for the CP transgene. Although the CP transgenes are actively transcribed, the CP steady-state mRNA level is below the detection limit on an RNA gel blot (See Example 2 and FIG. 2).

Using poly(A)+RNA from silenced leaf tissue of this plant a cDNA fusion library was constructed in the prey vector pIVING 1154 using the SMART™ PCR cDNA library construction kit (offered by Clontech, Palo Alto Calif., a part of Beckton Dickinson, Franklin Lakes, N.J.). To create the vector pIVING1154, pGAD424 was modified to allow directional cloning of the cDNA into asymmetric SfiI sites. The library has an estimated 1.66×10^6 primary clones. As shown in FIG. 4, the inserts range between 0.5 to 2.0 kb with an average size of about 0.8 kb. Also, a 0.8-kb fragment containing the CP coding sequence could be readily amplified from the plasmid cDNA library indicating that low abundance mRNAs are represented.

Example 6

Library Scale Two-hybrid Screen with HC-Pro Bait

The yeast strain L40 containing the SrMV HC-Pro bait construct was transformed with the prey cDNA library representing the equivalent of approximately 1.6×10^6 primary *E. coli* clones (see Example 4). About 77 million primary yeast transformants were plated on selective media (-leucine, -histidine +zeocin) and 776 yeast colonies were recovered by the fourth day of growth. Of these, 441 showed a range of light to heavy blue staining in the standard β -galactosidase colony-lift filter assay. To date, all of these double transformants have been colony purified, and all of the prey plasmids have been isolated. A RNase H-like protein, a RING zinc finger protein and LRR transmembrane protein kinase have been identified

15

among the true positives. Some results of this screen and additional screens with SrMV P1 or RNase H like-protein as the bait are depicted in Table 1.

TABLE 1

Bait (DBD)	Prey (AD)	Interaction
SrMV HC-Pro	SrMV HC-Pro	++
SrMV P1	SrMV P1	-/+
SrMV HC-Pro	SrMV P1	-
SrMV P1	SrMV HC-Pro	-
SrMV HC-Pro	RNase H-like prot.	+++
RNase H-like prot.	SrMV HC-Pro	+++
SrMV P1	RNase H-like prot.	-
RNase H-like prot.	RNase H-like prot.	+++

A -/+plus designates that very light blue staining was observed only after a 24 hour incubation. +++ designates that strong blue staining was seen in less than 30 minutes, and ++ designates strong blue staining in less than 3 hours. The RNase H-like protein self interaction may indicate that the protein functions as a homodimer.

Example 7

RNase H-like Protein

One of the prey plasmids that was recovered during the experiments of Example 5 was retransformed into the SrMV HC-Pro bait yeast strain to reconfirm activation of the reporter genes. The plasmid encodes a protein with approximately 40% identity to a viral RNase H. This protein has been recovered several times using the methods of these examples. As shown in FIG. 5, the interaction with HC-Pro is very strong (D) and there is no interaction in combination with an empty bait plasmid or with a lamin bait plasmid (B and C, respectively). Furthermore, the RNase H-like gene is in the correct reading frame with respect to the fusion protein, so this is a true interactor. The cDNA sequence (SEQ.ID.NO. 3) for a nucleic acid encoding the RNase H-like protein and its amino acid sequence (SEQ.ID.NO. 4) are provided in FIG. 6.

Example 8

RING Zinc Finger Protein that Interacts with HC-Pro

Another protein identified as a true positive in HC-Pro interaction screens is a RING zinc finger protein. A partial cDNA sequence of a nucleic acid encoding this protein is provided in FIG. 7A (SEQ.ID.NO. 5). The encoded amino acid sequence is included as FIG. 7B (SEQ.ID.NO. 6). The nucleic acid sequence is sufficient to identify nucleic acids encoding the protein, but likely lacks approximately 300 bp at the 5' end in the figure. However, because a protein encoded by the possibly truncated nucleic acid sequence was identified in a two-hybrid screen, a nucleic acid with the sequence provided must encode a sufficient portion of the protein to allow interaction with SrMV HC-Pro. With the information disclosed in this invention, it is possible for a person skilled in the art to isolate the full length cDNA and protein sequence.

Example 9

LRR Transmembrane Protein Kinase

Another protein identified as a true positive in SrMV HC-Pro yeast two-hybrid screens is an LRR transmembrane pro-

16

tein kinase. A partial cDNA sequence of a nucleic acid encoding this protein is provided in FIG. 8A (SEQ.ID.NO. 7). The encoded amino acid sequence is included in FIG. 8B (SEQ.ID.NO. 8). The sequence is sufficient to identify nucleic acids encoding the protein, but likely lacks approximately 1 kb at the 5' end in the figure. However, because a protein encoded by the possibly truncated nucleic acid sequence was identified in a two-hybrid screen, a nucleic acid with the sequence provided must encode a sufficient portion of the protein to allow interaction with SrMV HC-Pro. With the information disclosed in this invention, it is possible for a person skilled in the art to isolate the full length cDNA and protein sequence.

Example 10

Library Scale Two-hybrid Screen with RNase H-like Protein as Bait

A library scale two-hybrid screen similar to that of Example 5 was conducted using the same prey library (described in Example 4) with the RNase H-like protein as bait. Using the RNase H-like protein as bait, at least two true positives for prey/host proteins were identified and further characterized. The results of some of these assays as well as assays with SrMV HC-Pro and SrMV P1 as bait are depicted in Table 2.

TABLE 2

Bait (DBD)	Prey (AD)	Interaction
SrMV HC-Pro	SrMV HC-Pro	++
RNase H-like prot.	SrMV HC-Pro	+++
SrMV P1	RNase H-like prot.	-
RNase H-like prot.	RNase H-like prot.	+++
RNase H-like prot.	Sugarcane 14-3-3	+++
SrMV HC-Pro	Sugarcane 14-3-3	-
SrMV P1	Sugarcane 14-3-3	-

A -/+ designates that very light blue staining was observed only after a 24 hour incubation. +++ designates that strong blue staining was seen in less than 30 minutes, and ++ designates strong blue staining in less than 3 hours.

Example 11

Sugarcane 14-3-3 Protein

One protein identified through its ability to interact with the RNase H-like protein is a sugarcane 14-3-3 protein. The sequence of a nucleic acid encoding the sugarcane 14-3-3 protein is provided in FIG. 9A (SEQ. ID. NO. 9). The encoded amino acid sequence is provided in FIG. 9B (SEQ. ID. NO. 10). The role of another 14-3-3 protein in signal transduction in the dicot *Arabidopsis* has recently been described in Sehnke et al., 2002. A similar role in monocots is suggested by the sequence disclosed in this example. Specifically, 14-3-3 proteins exhibit a phosphorylation-dependent association with proteins in dicots. The RNase H-like protein of the present invention contains a serine with a 99.1% chance of being phosphorylated. This serine is also located in a good consensus 14-3-3 protein client-binding site. More specifically, the putative phosphorylation/binding site is VIQNpSP-PDL (wherein "pS" designates phosphoserine) (SEQ. ID. NO. 13) beginning at amino acid 48 of the protein in FIG. 6B. Binding of sugarcane 14-3-3 and the RNase H-like protein in order to achieve a functional dimer (or as part of a functional

17

complex containing other proteins) is also suggested by the absence of DNase or RNase activity of either protein in isolation. As shown in Table 2, 14-3-3 does not interact noticeably with SrMV HC-Pro or SrMV P1.

Example 12

RING Zinc Finger Protein that Interacts with RNase H-like Protein

Another protein identified as a true positive in RNase H-like protein interaction screens is a RING zinc finger protein. This is not the same RING zinc finger protein identified as interacting with SrMV HC-Pro. A partial cDNA sequence of a nucleic acid encoding this protein is provided in FIG. 10A (SEQ.ID.NO. 11). The amino acid sequence encoded by the nucleic acid is provided in FIG. 10B(SEQ.ID.NO. 12)

Example 13

DNA Sequence Analyses of the Prey Genes Identified in the Two-hybrid Screen

Prey sequences that have been verified by the above screening procedures may be DNA sequenced utilizing standard fluorescence-based thermocycle sequencing methods, and restriction maps created. This sequence information may be utilized to search the genetic databases with BLASTx and BLASTp to determine sequence similarity with known genes or proteins. In cases where similarity is clear, one may verify that the sequence under investigation is inserted in the appropriate reading frame in the prey vector. If the sequence is not in the appropriate reading frame, the prey can be considered a false positive.

Example 14

Characterization of the Prey Nucleic Acids Identified in the Two-Hybrid Screen for the Ability of Their Corresponding Proteins to Interact With SrMV HC-Pro Protein or Other Bait Protein in Vitro Under Physiological Conditions

Although prey and bait combinations identified in the two-hybrid screen of the present invention may represent proteins which interact in yeast cells, several caveats of the assay may produce results which are not indicative of interactions that occur in planta. The yeast two-hybrid system assay requires that the bait and prey GAL4 fusion proteins both be imported into the yeast nucleus, which biochemically is a much more reducing environment than the yeast cytoplasm. This may lead to different patterns of protein folding than might otherwise occur for proteins whose operating environment is a more oxidative cytoplasm. Additionally, protein fragments placed in an unnatural position in a yeast two-hybrid assay (e.g. an N term region in the C term of the fusion protein) may fail to fold or function properly because of positional effects.

Given that in vivo studies often require specialized reagents in the present method, an initial in vitro screening procedure in which potential protein interactions are verified under physiological conditions may be used to save time and expense associated with plant transformations in the case of false positives. However, it does remain possible that in vitro conditions may lack certain requirements for interaction found only in living cells and thus generate false negatives. Therefore, in some instances it may be desirable to proceed

18

with in planta studies without or despite results in vitro. Such an approach is within the scope of the present invention.

If in vitro studies are preformed, host/prey proteins identified in the two-hybrid screen may be further tested for their ability to interact with SrMV P1 and SrMV HC-Pro or other proteins involved in PTGS in vitro under physiological conditions (i.e. pH 6.8 and 0.1 M ionic strength) as either polyhistidine, S-tag, or glutathione-s-transferase (GST) fusion-proteins. A series of vectors from Novagen, Inc. (Madison, Wis., affiliate of Merck KgaA, Darmstadt, Germany) including pET15b, pET29a,b,c and pET30a,b,c, allow the construction of N-terminal or C-terminal polyhistidine fusion proteins, which in the case of pET29 and pET30 derivatives may also contain S-tags. Pharmacia Corp. (Peapack, N.J.) also produces pGEX2, pGEX3 and pGEX4 derivatives in which GST fusion proteins may be produced. These vectors allow the specific purification of "tagged" proteins that have either polyhistidine tags (using Ni⁺-chelated resin), S-tags (using S-agarose), or GST-tags (using reduced glutathione-conjugated resins). Bait and prey inserts identified from the two-hybrid screen may be subcloned into one of these vectors. Other vectors encoding tag regions may also be used.

One approach is to construct two vector types, if possible, for the bait as well as for each prey that has been identified and to produce GST- and polyhistidine tagged proteins in *E. coli*. These tagged proteins may be purified with kits available from Pharmacia, Novagen, or other sources, or using methods known to the art and appropriate for the selected tag. His-S tagged SrMV HC-Pro produced in the pET30 vector purified with a Ni column is shown in FIG. 11 in lanes 3 and 4. As FIG. 11 indicates, tagged protein is produced only when an insert is present in the vector and may be largely purified using a Ni column.

Radiolabelled SrMV HC-Pro, SrMV P1, RNase H-like protein or other bait proteins of interest may be produced by translation of their corresponding transcripts in rabbit reticulocyte lysate. Using the TNS method, transcripts may be generated in a T7 in vitro transcription system or other system known to the art. Subsequent translation in the presence of S-35-labeled methionine or cysteine allows radiolabeling of the bait protein. FIG. 12 shows that RNase H-like protein produced using the above methodology is, in fact, radiolabeled. A small amount of this translated lysate may then be incubated with the tagged prey protein bound to its corresponding resin, and retention of radiolabeled bait protein may be assayed with SDS-PAGE gels and autoradiograms. Positive controls can include tagged bait protein, and negative controls can include the tags themselves, as well as the resins without any other proteins. These procedures or various modifications thereof readily determined by one skilled in the art allow one to determine whether prospective prey proteins are capable of interacting with either one or both viral proteins in vitro under physiological conditions.

Example 15

Farwestern Blots and Other Assays Indicate Interaction of Various Proteins

FIG. 13 shows a farwestern blot of His-S tagged SrMV HC-Pro probed with ³⁵S labeled RNase H-like protein. Both proteins were produced according to the methods of the above examples. A comparison of binding of labeled RNase H-like protein to the His-S tagged SrMV HC-Pro to the various controls indicates that the interaction is specific between the two proteins and also that it occurs in conditions approximating plant physiological conditions.

19

FIG. 14 shows a farwestern blot of His-S tagged SrMV HC-Pro and RNase H-like protein and a plant extracts probed with ³⁵S labeled RNase H-like protein. A comparison of the lanes indicates that the RNase H interacts with itself and SrMV HCPro and with other sugarcane proteins—from left to right—healthy sugarcane plant, SrMV infected sugarcane plant, sugarcane plant transgenic for SrMV P1/HC-Pro, sugarcane plant transgenic for SrMV delta N 12 HC-Pro. An extra band in the three plants expressing SrMV HC-Pro that is slightly smaller than the tagged SrMV HC-Pro may be seen in FIG. 14.

FIG. 15 shows the results of a Ni column pull down of His-S tagged protein and its binding partner. The proteins attached to the Ni column were removed and run on an SDS PAGE gel. The results indicate that His-S tagged SrMV HC-Pro and not control proteins pulled down labeled RNase H-like protein. Label controls were not pulled down. This indicates that the SrMV HC-Pro/RNase H-like protein interaction is specific and viable under physiological conditions.

FIG. 16 shows a farwestern blot of His-S tagged sugarcane 14-3-3 probed with ³⁵S labeled RNase H-like protein. A comparison of binding of labeled RNase H-like protein to the His-S tagged sugarcane 14-3-3 to the various controls indicates that the interaction is specific between the two proteins and also that it occurs in conditions approximating physiological conditions.

Example 16

HC-Pro Deletion Mutant and Interaction with RNase H-like Protein

In order to better determine the region of SrMV HC-Pro responsible for interaction with the RNase H-like protein, a series of deletion mutants were created and their interaction was tested in yeast two-hybrid and farwestern blot assays as described above. The results of these experiments are summarized in Table 3 and indicate that the C terminal portion, including the last 10 kDa of the HC-Pro protein are required for interaction with the RNase H-like protein while the N terminal region up to 7 kDa is not necessary.

TABLE 3

HC-Pro segment	Assay	Interaction
Full length ←-----→	Yeast two-hybrid	+++
Full length ←-----→	Farwestern	+++
N 1.3 kDa deletion ←-----→	Yeast two-hybrid	+++
N 7.0 kDa deletion ←-----→	Yeast two-hybrid	+++
C 10 kDa deletion ←-----→	Farwestern	-
C 20 kDa deletion ←-----→	Farwestern	-

+++ in yeast two-hybrid assays indicates that strong blue staining was seen in less than 30 minutes. +++ in farwestern assays indicates that binding was apparent after a short exposure time.

Example 17

Complete Cloning of Partial cDNA Prey Clones

Candidate clones whose corresponding protein sequences interact with SrMV HC-Pro or other bait proteins in vitro

20

under physiological conditions may be completely cloned with 5' RACE procedures or other procedures known to the art.

Example 18

Production of Antiserum

To facilitate in vivo co-immunoprecipitation studies further outlined below, antisera recognizing SrMV HC-Pro or another protein involved in PTGS may be produced. Using a pET or pGEX expression plasmid construct as described above or another 6xhis construct known to the art, one may produce 6Xhis-tagged SrMV HC-Pro or other protein in *E. coli*, purify the fusion protein as described above, and utilize the protein as an antigen with rabbits or other animals to produce polyclonal antisera. Similarly, SrMV P1 and host proteins may also be utilized to produce antisera, depending upon the potential utility of the resultant serum. Other antisera production techniques may also be used to produce polyclonal or, where useful, monoclonal antibodies.

Example 19

Transgenic Plant Studies

Because of its specific mode of action, the SrMV P1/HC-Pro and other proteins identified as involved in PTGS using the above two-hybrid and in vitro methods may target one or more factors that are expressed and functional in silenced tissue. This may be confirmed in planta. Because the cDNA library in the above examples was constructed from a plant harboring a PTGS-silenced SrMV coat protein sequence, and the plant is resistant to SrMV, one cannot readily utilize mechanical transmission of SrMV as a source of SrMV HC-Pro or other viral proteins in such plants, and thus may resort to transgenic methodologies.

Plant transformation studies demonstrate the feasibility of this approach. To date, SrMV HC-Pro transformants and SrMV P1/HC-Pro transformants have been confirmed via Southern and Northern analyses. The plants were produced via particle gun transformation of embryogenic callus, as previously described (Ingelbrecht et al., 1999). Three different plant transformation vectors have been constructed: (1) pIVING1023 which contains the SrMV P1/HC-Pro polyprotein sequence, in which SrMV HC-Pro may be expected to be cleaved out by the SrMV P1 protease in planta, (2) pIVING1002 which contains only the SrMV P1 sequence, and (3) pIVING991-1 which contains only the SrMV HC-Pro sequence. In the present example, two selectable markers for sugarcane transformation have been used, the nptII gene in combination with geneticin, and the bar gene for selection with bialaphos. Bialaphos selection may be used on bar transformed cells, and geneticin selection may be used on bar transgenic cells to select for second transformation events of previously transformed tissue. Other transformation mechanisms and selection systems known to the art may also be used. Such sequentially transformed plants may serve as a source of material for in planta studies.

In a further example, the transgenic plant #16 that was used as a source for construction of the prey cDNA library above, was utilized to generate embryogenic callus. Plant #16 was previously transformed with the bar gene, and in this example was transformed again with the SrMV P1-HC-Pro construct using geneticin selection. Because SrMV P1 has been described as an enhancer of the PTGS-inhibition by SrMV HC-Pro, the capsid message levels may be higher in SrMV

P1-HC-Pro transformed #16 plants as compared to SrMV HC-Pro transformed #16 plants. In addition to having a system of modulated PTGS-suppression, these transgenic plants also allow verification in planta of the interactions between the viral proteins and the identified prey proteins, utilizing SrMV HC-Pro specific antiserum in standard co-immunoprecipitation methodologies.

Sequence homology of the isolated host proteins with known proteins, if present, may be used to postulate their biological function and whether or not they are involved in PTGS in a manner similar to that employed with the genes identified in mutagenesis studies of *Neurospora*, *C. Elegans* and *Arabidopsis*.

One approach for determining functions of the isolated genes may be based on creating suppression phenotypes for these genes in sugarcane via genetic transformation. Efficient triggering of gene suppression can be achieved by simultaneous expression of sense and antisense constructs or constructs designed to produce dsRNA as demonstrated in tobacco, rice and *Arabidopsis* (Waterhouse et al., 1998; Smith et al., 2000; Chuang and Meyerowitz, 2000). See also FIG. 17 for depictions of the basic structure of DNA and resulting mRNA for such constructs. Such constructs may be created for each of the isolated host genes and introduced in plants that display PTGS of a capsid protein transgene, such as plant #16. The transformation methodologies described above can be followed to generate stably transformed plants. Alternatively, a protoplast-based system can be used in combination with electroporation or PEG-mediated transformation. Expression of the capsid protein will be analyzed and reversal of silencing may indicate that the PTGS mechanism in these transgenics is impaired and that the transformed host genes are involved in PTGS.

Example 20

In Planta Studies of HC-Pro and RNase H-like Protein

Embryonic sugarcane calli were bombarded with various constructs according to the above examples in order to provide transient expression to verify the role of HN-Pro and

RNase H-like protein in PTGS. Double stranded mRNA relating to the various proteins was generated using constructs such as those show in FIG. 17 inserted into the vector of FIG. 18.

FIG. 19 shows that GUS under a Ubi promoter alone is expressed in embryonic callus. Introduction of a double stranded GUS mRNA reduces its expression by activating PTGS of GUS mRNA. However expression is restored by both introducing SrMV P1-HC-Pro into the callus and by inducing PTGS of the RNase H-like protein using a double stranded mRNA of that protein. Average results for three separate experiments of this nature are depicted in FIG. 20 and show that statistically significant results are consistently obtained. Overall, these experiments confirm that SrMV P1 and SrMV HC-Pro inhibits PTGS and that the RNase H-like protein is involved in effecting PTGS.

FIG. 21 illustrates the PTGS effect of the RNase H-like protein even more clearly. In these figures calli transiently transformed with GUS exhibit the protein. This protein production is severely disrupted by subsequent or simultaneous transformation with DNA that produces double stranded GUS mRNA, which triggers PTGS. PTGS is not turned off by the transformation of the GUS/double stranded GUS calli with RNase H. However, it is severely hampered by introduction of DNA that produces double stranded mRNA for RNase H-like protein, which induces PTGS of RNase H-like protein itself. These experiments demonstrate that the RNase-H like protein plays a role in PTGS. These results are summarized in Table 4.

TABLE 4

Transformation Constructs	Callus Color
35S-int/GUS	Blue
35S-int/GUS + 35S-int/dsGUS	Not Blue
35S-int/GUS + 35S-int/dsGUS + 35S-int/RNase H-like protein	Not Blue
35S-int/GUS + 35S-int/dsGUS + 35S-int/ds RNase H-like protein	Blue

All references mentioned in any portion of the foregoing specification are incorporated by reference herein.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 13

<210> SEQ ID NO 1

<211> LENGTH: 2103

<212> TYPE: DNA

<213> ORGANISM: Potyvirus Sorghum Mosaic Virus

<400> SEQUENCE: 1

```

gaattcgccc ttatttcagc catggcagga gcatggaaca ctgtgactta caagtggagg      60
ccgaatttgg acaacgcaag agatgttcga aaagtgatgg aacattttgc agcaaagcac    120
caagtttatg atgcaaacg tgcagcagag cataacagta gaatccttcg caggactttt    180
gtacaagaaa ttgcaaaagc acctgaagag aagacttcat acaaacctca ggtgtggggtc   240
gaaaaacaag ataacaatcc aacgatccat ctacactatg ttagattcaa aaacaaggag    300
aaaaagatat taccggagat cactccagga tcggtcgcaa agttaacgcg acggataact    360
gagctcagca aaactacaaa gcttgaagtt gaactcattg gtaagaaaag acgcaaatcg    420
actaaactag caatcaaaag gcgcagaaac agagagtatt tgcattgtga aactagacac    480

```

-continued

```

gaaacaaaca aatttaagcg cgttgacatc aacatagaac gacactgggt tccacttggt 540
aagaagattt caaagtgcta tagtcacata tcaccaagaa tgtacaaaaa catgagcaaa 600
ggcgacagtg ggtaacatt catccagaat ggtgagttat ttataatccg aggaaaaacga 660
gatggcgtcc tacttaatag tatcaccaat gaaactcgaa ttaatgaaat aacttatttt 720
agcgatgctc aggcgaaacga cttctggcga ggttacacag atcatatggt cgaaaatagg 780
ttaatttcta caactcatic agaacacata cccacaataa atttagagaa gtgtggaaag 840
aggatggcat tgttagagat attgtttcac tccacattca aaattacgtg taagcactgt 900
aacaatgacg atcttgaact atcggatgat gagtttgag aaagactata taagaactta 960
atcagaattg aagaaaagca aaaagaatat ttagctgaag atcaaaagct taagcgaatg 1020
atatcctttc tgaaggatag atgcaatcca aaatttgagc atttaccatt attatggcag 1080
gtcgtgaaa caattggaca ttacactgat aatcaagcaa aacagatcct tgaagttaat 1140
gaagcgctca taaaagtga cactctttct gttgaagatg cagtcaaagc tagcgcacgc 1200
ttgctagaga tttcaagatg gtacaagaat aggaaagaat catcgaaaga aggtacactt 1260
agtacattca ggaataaaat ttcacctaaa agtactatta atacagcact gatgtgtgat 1320
aatcagctcg atacaaatgg taacttctca tggggaaaga gagaatatca tgccaagcga 1380
ttctttacaa actattttga agctgttgat ccaaaagaca cgatgaaaa gcatgttact 1440
cggttcaatc caaatggta acgcaactt tcgattggaa aactagttat cccattagac 1500
ttccagaaga ttcgtgaatc atttataggt gttcaagttc aaaaacaagc aattagtaga 1560
gcgtgcttaa gtaaaatcga aaataattac atataccctt gctgttgtgt aactacagaa 1620
tttggtaaac cggtttatc agagatcatt ccaccaacta aaggatcatat tactattgga 1680
aattcgaccg acccaaaaat tgtggatttg cctaattccg acccaaat gatgtacata 1740
gcgaaagatg gttattgta tttgaatata tttttagctg ctctgataaa cgtaaatgaa 1800
gattcagcaa aagattacac aaagttttg cgtgatgaac taattgaaag acttggaag 1860
tggccaaaac tcaagacgt ggcgacagca tgttatgcat tatcagtaat gttccagaa 1920
attaagaacg cggagcttcc acaataacta gtggaccacg aacataaaac catgcatgtg 1980
atagattcgt acggatctct cagtgttggc ttccacatac tcaaagcga cacaatagga 2040
caattaatca aaatgcaata tgaatccatg gaaagtgaag tgagagagta ttagtcggt 2100
tag 2103

```

```

<210> SEQ ID NO 2
<211> LENGTH: 694
<212> TYPE: PRT
<213> ORGANISM: Potyvirus Sorghum Mosaic Virus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (694)..(694)
<223> OTHER INFORMATION: "Xaa" may be any amino acid, not present or may
represent a stop codon in the sequence

```

```

<400> SEQUENCE: 2

```

```

Met Ala Gly Ala Trp Asn Thr Val Thr Tyr Lys Trp Arg Pro Asn Leu
1           5           10          15

```

```

Asp Asn Ala Arg Asp Val Arg Lys Val Met Glu His Phe Ala Ala Lys
20          25          30

```

```

His Gln Val Tyr Asp Ala Lys Arg Ala Ala Glu His Asn Ser Arg Ile
35          40          45

```

```

Leu Arg Arg Thr Phe Val Gln Glu Ile Ala Lys Ala Pro Glu Glu Lys

```

-continued

50					55					60					
Thr	Ser	Tyr	Lys	Pro	Gln	Val	Trp	Val	Glu	Lys	Gln	Asp	Asn	Asn	Pro
65					70					75					80
Thr	Ile	His	Leu	His	Tyr	Val	Arg	Phe	Lys	Asn	Lys	Glu	Lys	Lys	Ile
				85					90					95	
Leu	Pro	Glu	Ile	Thr	Pro	Gly	Ser	Val	Ala	Lys	Leu	Thr	Arg	Arg	Ile
			100					105					110		
Leu	Glu	Leu	Ser	Lys	Thr	Thr	Lys	Leu	Glu	Val	Glu	Leu	Ile	Gly	Lys
		115					120					125			
Lys	Arg	Arg	Lys	Ser	Thr	Lys	Leu	Ala	Ile	Lys	Arg	Arg	Arg	Asn	Arg
	130					135					140				
Glu	Tyr	Leu	His	Cys	Glu	Thr	Arg	His	Glu	Thr	Asn	Lys	Phe	Lys	Arg
145					150					155					160
Val	Asp	Ile	Asn	Ile	Glu	Arg	His	Trp	Phe	Pro	Leu	Val	Lys	Lys	Ile
				165					170					175	
Ser	Lys	Cys	Tyr	Ser	His	Ile	Ser	Pro	Arg	Met	Tyr	Lys	Asn	Met	Ser
			180					185					190		
Lys	Gly	Asp	Ser	Gly	Leu	Thr	Phe	Ile	Gln	Asn	Gly	Glu	Leu	Phe	Ile
		195					200					205			
Ile	Arg	Gly	Lys	Arg	Asp	Gly	Val	Leu	Leu	Asn	Ser	Ile	Thr	Asn	Glu
	210					215					220				
Thr	Arg	Ile	Asn	Glu	Ile	Thr	Tyr	Phe	Ser	Asp	Ala	Gln	Ala	Asn	Asp
225					230					235					240
Phe	Trp	Arg	Gly	Tyr	Thr	Asp	His	Met	Val	Glu	Asn	Arg	Leu	Ile	Ser
				245					250					255	
Thr	Thr	His	Thr	Glu	His	Ile	Pro	Thr	Ile	Asn	Leu	Glu	Lys	Cys	Gly
			260					265					270		
Lys	Arg	Met	Ala	Leu	Leu	Glu	Ile	Leu	Phe	His	Ser	Thr	Phe	Lys	Ile
		275					280					285			
Thr	Cys	Lys	His	Cys	Asn	Asn	Asp	Asp	Leu	Glu	Leu	Ser	Asp	Asp	Glu
	290					295					300				
Phe	Gly	Glu	Arg	Leu	Tyr	Lys	Asn	Leu	Ile	Arg	Ile	Glu	Glu	Lys	Gln
305					310					315					320
Lys	Glu	Tyr	Leu	Ala	Glu	Asp	Gln	Lys	Leu	Lys	Arg	Met	Ile	Ser	Phe
				325				330						335	
Leu	Lys	Asp	Arg	Cys	Asn	Pro	Lys	Phe	Glu	His	Leu	Pro	Leu	Leu	Trp
			340					345					350		
Gln	Val	Ala	Glu	Thr	Ile	Gly	His	Tyr	Thr	Asp	Asn	Gln	Ala	Lys	Gln
		355					360					365			
Ile	Leu	Glu	Val	Asn	Glu	Ala	Leu	Ile	Lys	Val	Asn	Thr	Leu	Ser	Val
	370					375					380				
Glu	Asp	Ala	Val	Lys	Ala	Ser	Ala	Ser	Leu	Leu	Glu	Ile	Ser	Arg	Trp
385					390					395					400
Tyr	Lys	Asn	Arg	Lys	Glu	Ser	Ser	Lys	Glu	Gly	Thr	Leu	Ser	Thr	Phe
				405					410					415	
Arg	Asn	Lys	Ile	Ser	Pro	Lys	Ser	Thr	Ile	Asn	Thr	Ala	Leu	Met	Cys
			420					425					430		
Asp	Asn	Gln	Leu	Asp	Thr	Asn	Gly	Asn	Phe	Leu	Trp	Gly	Lys	Arg	Glu
		435					440					445			
Tyr	His	Ala	Lys	Arg	Phe	Phe	Thr	Asn	Tyr	Phe	Glu	Ala	Val	Asp	Pro
	450					455					460				
Lys	Asp	Thr	Tyr	Glu	Lys	His	Val	Thr	Arg	Phe	Asn	Pro	Asn	Gly	Gln
465					470					475					480

-continued

Arg Lys Leu Ser Ile Gly Lys Leu Val Ile Pro Leu Asp Phe Gln Lys
 485 490 495
 Ile Arg Glu Ser Phe Ile Gly Val Gln Val Gln Lys Gln Ala Ile Ser
 500 505 510
 Arg Ala Cys Leu Ser Lys Ile Glu Asn Asn Tyr Ile Tyr Pro Cys Cys
 515 520 525
 Cys Val Thr Thr Glu Phe Gly Gln Pro Val Tyr Ser Glu Ile Ile Pro
 530 535 540
 Pro Thr Lys Gly His Ile Thr Ile Gly Asn Ser Thr Asp Pro Lys Ile
 545 550 555 560
 Val Asp Leu Pro Asn Ser Asp Pro Pro Met Met Tyr Ile Ala Lys Asp
 565 570 575
 Gly Tyr Cys Tyr Leu Asn Ile Phe Leu Ala Ala Leu Ile Asn Val Asn
 580 585 590
 Glu Asp Ser Ala Lys Asp Tyr Thr Lys Phe Leu Arg Asp Glu Leu Ile
 595 600 605
 Glu Arg Leu Gly Lys Trp Pro Lys Leu Lys Asp Val Ala Thr Ala Cys
 610 615 620
 Tyr Ala Leu Ser Val Met Phe Pro Glu Ile Lys Asn Ala Glu Leu Pro
 625 630 635 640
 Gln Ile Leu Val Asp His Glu His Lys Thr Met His Val Ile Asp Ser
 645 650 655
 Tyr Gly Ser Leu Ser Val Gly Phe His Ile Leu Lys Ala Asn Thr Ile
 660 665 670
 Gly Gln Leu Ile Lys Met Gln Tyr Glu Ser Met Glu Ser Glu Met Arg
 675 680 685
 Glu Tyr Val Val Gly Xaa
 690

<210> SEQ ID NO 3

<211> LENGTH: 937

<212> TYPE: DNA

<213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<400> SEQUENCE: 3

```

ggccattatg gccggggaga acactgtatg aagaacatgg cgccttctta tagtcaagac   60
g'gcagctcg atctcgtact tctgacgct cccgtcgatg ctagcgcgct tcggttctgaa  120
cattcatctc agcttgctag ctetaactgg agatctgtga ttcaaaactc cccacctgat  180
ctcctatgcg gatcggttag accggcaatt aggcgcacgg cagagactgc gaagaacaat  240
ggccgcatct ttcgcacgtg tccggcgtgc aaaatatgga tttggcagga tctgctggac  300
agctatgtga atgctttgat aagctactgt cgtgatgcct ccattgattc ccttcagtca  360
gagcttgaat ctagccgttt attaatttcc gagaagcagg cacagatttc acgtttggag  420
aaacaattgg agacgctgca gccacttata tcaaaatata ctgaacaatc tcgcagcatt  480
gctcaggcct ccattccatc atcaactttt ttcttgggaag cctgcagtct ccgacatcag  540
cggcggaggg tgaatgaaaa tcaaggaggg gctaatttca aaagaagcta ctggagcgat  600
taaagcttcg gttaaaaaata agcaagaatc taacaccagc agcacaatc ccatgagctg  660
gctttttttt gggacaccct ttcatttttc atcaaaaaag ggggggcacc cccagtttcc  720
tccaaaaggc tcccctgtc cgacatcata ggtgatgtga ttacccaaaa acaggttgtc  780
ccgcttctg actcgatgcc aaatttggat tcaatgctgc tcctgttgtt ttaacaatca  840

```

-continued

```
atcattttga ctaaagcat tcccctaaa attgttgta aatttattgt caaacctatt 900
accgcaaagt ccgttgagc gtaatcccc ccttttt 937
```

```
<210> SEQ ID NO 4
<211> LENGTH: 200
<212> TYPE: PRT
<213> ORGANISM: Saccharum hybrid cultivar CP72-1210
```

```
<400> SEQUENCE: 4
```

```
Gly His Tyr Gly Arg Gly Glu His Cys Met Lys Asn Met Ala Ala Ser
1           5           10           15
Tyr Ser Gln Asp Ala Gln Leu Asp Leu Val Leu Pro Asp Ala Pro Val
20           25           30
Asp Ala Ser Ala Ser Arg Ser Glu His Ser Ser Gln Leu Ala Ser Ser
35           40           45
Asn Trp Arg Ser Val Ile Gln Asn Ser Pro Pro Asp Leu Leu Cys Gly
50           55           60
Cys Gly Arg Pro Ala Ile Arg Arg Thr Ala Glu Thr Ala Lys Asn Asn
65           70           75           80
Gly Arg Ile Phe Arg Thr Cys Pro Ala Cys Lys Ile Trp Ile Trp Gln
85           90           95
Asp Leu Leu Asp Ser Tyr Val Asn Ala Leu Ile Ser Tyr Cys Arg Asp
100          105          110
Ala Ser Ile Asp Ser Leu Gln Ser Glu Leu Glu Ser Ser Arg Leu Leu
115          120          125
Ile Ser Glu Lys Gln Ala Gln Ile Ser Arg Leu Glu Lys Gln Leu Glu
130          135          140
Thr Leu Gln Pro Leu Ile Ser Lys Tyr Thr Glu Gln Ser Arg Ser Ile
145          150          155          160
Ala Gln Ala Ser Ile Pro Ser Ser Leu Phe Phe Leu Glu Ala Cys Ser
165          170          175
Leu Arg His Gln Arg Arg Arg Val Asn Glu Asn Gln Gly Gly Ala Asn
180          185          190
Phe Lys Arg Ser Tyr Trp Ser Asp
195          200
```

```
<210> SEQ ID NO 5
<211> LENGTH: 449
<212> TYPE: DNA
<213> ORGANISM: Saccharum hybrid cultivar CP72-1210
```

```
<400> SEQUENCE: 5
```

```
gaattcggcc attatggccg gggcatgtgc agtgaatgcg ccaagtcct gaggtaccaa 60
accactcggg gccccatctg caggcagcct gttgagcgtc tcctcgagat caaagtgagc 120
aacaatctg aagagcagca gcagacgcc caatcgccgc cgctcccagc cccagctctg 180
cagcaggaag aggtgtagcc gtgattaaag tcagttctga gacattatat ggaactagtt 240
tgccgccttc aggcctttcc ctaaagggtt gttctctcat ctgagcaacg gggaatgtaa 300
ccggtacttt acctttagcc tatgtaagct tctggcatcg catggctttg ccgacctctg 360
ctgtacctgc ttatctggag gtcggagacc aagatgccaa gaaagtgtg taccgtatat 420
taaaaaaaaa aaaaaaaaaa aaaaaaaaaa 449
```

```
<210> SEQ ID NO 6
<211> LENGTH: 149
<212> TYPE: PRT
```

-continued

<213> ORGANISM: Saccharum hybrid cultivar CP72-1210
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (66)..(141)
 <223> OTHER INFORMATION: "Xaa" may be any amino acid, not present or may represent a stop codon in the sequence

<400> SEQUENCE: 6

Glu Phe Gly His Tyr Gly Arg Gly Met Cys Ser Glu Cys Ala Lys Val
 1 5 10 15
 Leu Arg Tyr Gln Thr Thr Arg Cys Pro Ile Cys Arg Gln Pro Val Glu
 20 25 30
 Arg Leu Leu Glu Ile Lys Val Ser Asn Lys Ser Glu Glu Gln Gln Gln
 35 40 45
 Thr Pro Gln Ser Pro Pro Leu Pro Ala Pro Ala Leu Gln Gln Glu Glu
 50 55 60
 Val Xaa Pro Xaa Leu Lys Ser Val Leu Arg His Tyr Met Glu Leu Val
 65 70 75 80
 Cys Gly Leu Gln Ala Phe Pro Xaa Arg Phe Val Leu Ser Ser Glu Gln
 85 90 95
 Arg Gly Met Xaa Pro Val Leu Tyr Leu Xaa Pro Met Xaa Ala Ser Gly
 100 105 110
 Ile Ala Trp Leu Cys Arg Pro Leu Leu Tyr Leu Leu Ile Trp Arg Ser
 115 120 125
 Glu Thr Lys Met Pro Arg Lys Val Cys Thr Val Tyr Xaa Lys Lys Lys
 130 135 140
 Lys Lys Lys Lys Lys
 145

<210> SEQ ID NO 7
 <211> LENGTH: 872
 <212> TYPE: DNA
 <213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<400> SEQUENCE: 7

gaattcggcc attatggccg ggaccaggaa ctccaggaat cagatgacaa ctcggggtat 60
 agagctcctg aagtgacat gtcgggtcag tattctcaaa agagtgatgt ttacagcttt 120
 ggtgtcgtca tgcttgagct actgactgga cagaaagcat ttgacagctc tcgggcaagg 180
 tcccagcaat cactagtccg gtgggcttca ccgacagctgc acgacatcga ctcgctagat 240
 cagatggttg atccaacctt agaggggctg taccatgcga aatcactctc tcggttcgca 300
 gacgcaatcg ctctctgtgt ccagcctgaa ccagaattca ggccaccaat gtcggaggtc 360
 gtccagtcac tggctcgtct tgtgcagcga gcaagcatgg ggacagcact aagcagcgag 420
 tgggaattctt gccagttoga tgaatctggt gatcacacgc tctaggggaa aatgatgtgt 480
 atttcctaga gagtctgatg aggaactata gaaggctcac aagtcataga aacttgacgc 540
 ttggcattgt tgtgagtgtg gacggtgtga catgtgccag tgtcagggtga aatgtgactt 600
 ttacctatgc cattttactg agagctctgt gcaacctgaa gttaggggtga aaagaaagt 660
 ccttctttaa aaatatatat ggttcattcg gacgtgtata tgaatatctt ttgaagacaa 720
 tcaactttct gatttcgtct ctgatcgtg tccaaaaatt atcaggggaag atgtagcact 780
 agtctcgcca cagaattagt catctgtata tctcagaaa tccgaacct atccaggaaa 840
 catcaacaga ggacacgtcc acatattcga ac 872

<210> SEQ ID NO 8

-continued

<211> LENGTH: 290
 <212> TYPE: PRT
 <213> ORGANISM: Saccharum hybrid cultivar CP72-1210
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (155)..(248)
 <223> OTHER INFORMATION: "Xaa" may be any amino acid, not present or may represent a stop codon in the sequence

<400> SEQUENCE: 8

Glu Phe Gly His Tyr Gly Arg Asp Gln Glu Leu Gln Glu Ser Asp Asp
 1 5 10 15
 Asn Ser Gly Tyr Arg Ala Pro Glu Val Thr Met Ser Gly Gln Tyr Ser
 20 25 30
 Gln Lys Ser Asp Val Tyr Ser Phe Gly Val Val Met Leu Glu Leu Leu
 35 40 45
 Thr Gly Gln Lys Ala Phe Asp Ser Ser Arg Ala Arg Ser Gln Gln Ser
 50 55 60
 Leu Val Arg Trp Ala Ser Pro Gln Leu His Asp Ile Asp Ser Leu Asp
 65 70 75 80
 Gln Met Val Asp Pro Thr Leu Glu Gly Leu Tyr His Ala Lys Ser Leu
 85 90 95
 Ser Arg Phe Ala Asp Ala Ile Ala Leu Cys Val Gln Pro Glu Pro Glu
 100 105 110
 Phe Arg Pro Pro Met Ser Glu Val Val Gln Ser Leu Val Arg Leu Val
 115 120 125
 Gln Arg Ala Ser Met Gly Thr Ala Leu Ser Ser Glu Trp Asn Ser Cys
 130 135 140
 Gln Phe Asp Glu Ser Gly Asp His Thr Leu Xaa Gly Lys Met Met Cys
 145 150 155 160
 Ile Ser Xaa Arg Val Xaa Xaa Gly Thr Ile Glu Gly Ser Gln Val Ile
 165 170 175
 Glu Thr Cys Ser Leu Ala Leu Leu Xaa Val Val Thr Val Xaa His Val
 180 185 190
 Pro Val Ser Gly Glu Met Xaa Leu Leu Pro Met Pro Phe Tyr Xaa Glu
 195 200 205
 Ser Ala Ala Thr Xaa Ser Arg Gly Glu Lys Lys Val Pro Ser Leu Lys
 210 215 220
 Ile Tyr Met Val His Ser Asp Val Tyr Met Asn Ile Phe Xaa Arg Gln
 225 230 235 240
 Ser Thr Phe Xaa Phe Arg Leu Xaa Ser Leu Ser Lys Asn Tyr Gln Gly
 245 250 255
 Arg Cys Ser Thr Ser Pro Ala Thr Glu Leu Val Ile Cys Ile Ser Ser
 260 265 270
 Glu Ile Arg Thr Ile Ser Arg Lys His Gln Gln Arg Thr Arg Pro His
 275 280 285
 Ile Arg
 290

<210> SEQ ID NO 9
 <211> LENGTH: 1141
 <212> TYPE: DNA
 <213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<400> SEQUENCE: 9

gaattcggcc attatggccg gggaccgcag ttcccccgac cacaccgttc cgccgcgcac 60
 agaggccagc cccgcgccag gagtaagttt gttcttttta acaatatgtc gagggaggag 120

-continued

```

aatgtttaca tggccaagct ggctgagcag gccgaaaggt atgaggagat ggttgagtat 180
atggagaagg tggctaagac tgtagatggt gaagagctca ctgtggagga gcgtaacctc 240
ctgtctgtcg catacaagaa tgtgattggg gctcgccgtg cttcatggcg cattgtctct 300
tccattgaac agaaggagga gtcccgtgtaag aacgaagagc atgtgaacct tatcaaggaa 360
taccgcgggga agattgaggc tgaactgagc aacatctgtg atggcatcct gaaactgctt 420
gactcccacc tagtgccttc ctctactgct gctgaatcaa aggtcttcta cctcaagatg 480
aagggtgact atcacaggta tcttgcgga ttaagactg gtgctgagag gaaggaatct 540
gctgagagca caatggtagc ctacaaggct gctcaggaca ttgctctggc tgagctggca 600
cctacacatc cgataaggct tgggcttgcct cttaacttct cagtgttcta ttatgagatt 660
ctgaactccc cagacaaagc ttgcaacctt gcaaagcagg cgtttgatga agctatctct 720
gagttagaca cccttggggga ggagtcatac aaagatagca ctctgatcat gcagctcctg 780
agggacaact tgaccctttg gacctctgac ctcacggagg atggtgctga tgaggcmeta 840
gaagcctcaa aaggtgatgc tggcgaggga cagtaatctt cggagagggc atgttgttcc 900
agcctggttt tagatgctct atgctgtcga agctgtgccc tgccattatt gtagcagatt 960
tcctctcccc ctcacttcat ttgcctcata ttagtaggct ggtagtggtc gaattagttc 1020
ccattgcttt gtgttgagc tagttggcac taggtccgtg tggactggta ttgttcccct 1080
ggatttgaca agcatgtcct gtggctgctc tagcgtttta ttgagctttg aagcctcgat 1140
t 1141

```

<210> SEQ ID NO 10

<211> LENGTH: 380

<212> TYPE: PRT

<213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (292)..(377)

<223> OTHER INFORMATION: "Xaa" may be any amino acid, not present or may represent a stop codon in the sequence

<400> SEQUENCE: 10

```

Glu Phe Gly His Tyr Gly Arg Gly Pro Gln Phe Pro Arg Pro His Arg
1           5           10           15
Ser Ala Ala His Arg Gly Gln Pro Arg Ala Arg Ser Lys Phe Val Leu
20           25           30
Phe Asn Asn Met Ser Arg Glu Glu Asn Val Tyr Met Ala Lys Leu Ala
35           40           45
Glu Gln Ala Glu Arg Tyr Glu Glu Met Val Glu Tyr Met Glu Lys Val
50           55           60
Ala Lys Thr Val Asp Val Glu Glu Leu Thr Val Glu Glu Arg Asn Leu
65           70           75           80
Leu Ser Val Ala Tyr Lys Asn Val Ile Gly Ala Arg Arg Ala Ser Trp
85           90           95
Arg Ile Val Ser Ser Ile Glu Gln Lys Glu Glu Ser Arg Lys Asn Glu
100          105          110
Glu His Val Asn Leu Ile Lys Glu Tyr Arg Gly Lys Ile Glu Ala Glu
115          120          125
Leu Ser Asn Ile Cys Asp Gly Ile Leu Lys Leu Leu Asp Ser His Leu
130          135          140
Val Pro Ser Ser Thr Ala Ala Glu Ser Lys Val Phe Tyr Leu Lys Met
145          150          155          160

```

-continued

Lys Gly Asp Tyr His Arg Tyr Leu Ala Glu Phe Lys Thr Gly Ala Glu
 165 170 175
 Arg Lys Glu Ser Ala Glu Ser Thr Met Val Ala Tyr Lys Ala Ala Gln
 180 185 190
 Asp Ile Ala Leu Ala Glu Leu Ala Pro Thr His Pro Ile Arg Leu Gly
 195 200 205
 Leu Ala Leu Asn Phe Ser Val Phe Tyr Tyr Glu Ile Leu Asn Ser Pro
 210 215 220
 Asp Lys Ala Cys Asn Leu Ala Lys Gln Ala Phe Asp Glu Ala Ile Ser
 225 230 235 240
 Glu Leu Asp Thr Leu Gly Glu Glu Ser Tyr Lys Asp Ser Thr Leu Ile
 245 250 255
 Met Gln Leu Leu Arg Asp Asn Leu Thr Leu Trp Thr Ser Asp Leu Thr
 260 265 270
 Glu Asp Gly Ala Asp Glu Gly Lys Glu Ala Ser Lys Gly Asp Ala Gly
 275 280 285
 Glu Gly Gln Xaa Ser Ser Glu Arg Ala Cys Cys Ser Ser Leu Val Leu
 290 295 300
 Asp Ala Leu Cys Cys Arg Ser Cys Ala Val Pro Leu Leu Xaa Gln Ile
 305 310 315 320
 Ser Ser Pro Pro His Phe Ile Cys Leu Ile Leu Val Gly Trp Xaa Trp
 325 330 335
 Ser Asn Xaa Phe Pro Leu Leu Cys Val Ala Ala Ser Trp His Xaa Val
 340 345 350
 Arg Val Asp Trp Tyr Cys Ser Pro Gly Phe Asp Lys His Val Leu Trp
 355 360 365
 Ser Leu Xaa Arg Phe Ile Glu Leu Xaa Ser Leu Asp
 370 375 380

<210> SEQ ID NO 11

<211> LENGTH: 829

<212> TYPE: DNA

<213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<400> SEQUENCE: 11

```

gaattcgccc attatggcgc ggctgttcaa gctggaccgt tatatggtat gggacacccat      60
ggatcttcca ccacaattgc ttatggcggg gcatacttgc catattcttc ctcaactgga      120
caatcgagca ataactcatca agagcatgga ttctctgagc ggccagggca gctgagtgt      180
caatatttta tgaggactgg aggttgcaaa tttggaacta tgtgtaaata taaccatcct      240
cgagattgga gcactcctaa gtccaactac atgttcagtc atctctgect tccacttcgt      300
cgggtgctc agccttgtgc gtactatgca caaaatggat attgcagata tggagttgca      360
tgcaaatatg atcaccatg ggttacacta ggctacagtt catctgcttt acccctatct      420
gacatgccaa ttgctcccta ccctatcggc ttctctgttg ccacggtggc tccatcttca      480
tcttccccag aatatatttc aaccaagat ccatcaatca accaagtage atcaccagtg      540
cagcaccgca acatgttggg acaatcttgc caaaaggggt ttcccttcgg atccattatg      600
cgaactcaac ttctacaagt gtcggcagtt caagcctggg gggcgctgat tttctgactg      660
gggatgatc cttaacacaa atttctatac ttgaacagtt tgaagccttc aaggaataaa      720
aactggggcc ttgaaaaacc gggaggggtt cttcccaaat aaaactgtgg tcaaacactca      780
tcttgaattg gtttcctatt caaacggaag aggtttagga gtcacattg      829

```

-continued

<210> SEQ ID NO 12
 <211> LENGTH: 276
 <212> TYPE: PRT
 <213> ORGANISM: Saccharum hybrid cultivar CP72-1210
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (219)..(262)
 <223> OTHER INFORMATION: "Xaa" may be any amino acid, not present or may represent a stop codon in the sequence

<400> SEQUENCE: 12

Glu Phe Gly His Tyr Gly Arg Ala Val Gln Ala Gly Pro Leu Tyr Gly
 1 5 10 15
 Met Gly His His Gly Ser Ser Thr Thr Ile Ala Tyr Gly Gly Ala Tyr
 20 25 30
 Leu Pro Tyr Ser Ser Ser Thr Gly Gln Ser Ser Asn Asn His Gln Glu
 35 40 45
 His Gly Phe Pro Glu Arg Pro Gly Gln Pro Glu Cys Gln Tyr Phe Met
 50 55 60
 Arg Thr Gly Gly Cys Lys Phe Gly Thr Met Cys Lys Tyr Asn His Pro
 65 70 75 80
 Arg Asp Trp Ser Thr Pro Lys Ser Asn Tyr Met Phe Ser His Leu Cys
 85 90 95
 Leu Pro Leu Arg Pro Gly Ala Gln Pro Cys Ala Tyr Tyr Ala Gln Asn
 100 105 110
 Gly Tyr Cys Arg Tyr Gly Val Ala Cys Lys Tyr Asp His Pro Met Gly
 115 120 125
 Thr Leu Gly Tyr Ser Ser Ser Ala Leu Pro Leu Ser Asp Met Pro Ile
 130 135 140
 Ala Pro Tyr Pro Ile Gly Phe Ser Val Ala Thr Leu Ala Pro Ser Ser
 145 150 155 160
 Ser Ser Pro Glu Tyr Ile Ser Thr Lys Asp Pro Ser Ile Asn Gln Val
 165 170 175
 Ala Ser Pro Val Gln His Pro Asn Met Leu Glu Gln Ser Cys Gln Lys
 180 185 190
 Gly Phe Pro Phe Gly Ser Ile Met Arg Thr Gln Leu Leu Gln Val Ser
 195 200 205
 Ala Val Gln Ala Trp Gly Ala Leu Ile Phe Xaa Leu Gly Asp Asp Pro
 210 215 220
 Xaa His Lys Phe Leu Tyr Leu Asn Ser Leu Lys Pro Ser Arg Asn Lys
 225 230 235 240
 Asn Trp Gly Leu Glu Lys Pro Gly Gly Val Leu Pro Lys Xaa Asn Cys
 245 250 255
 Gly Gln His Ser Ser Xaa Ile Gly Phe Leu Phe Lys Arg Lys Arg Phe
 260 265 270
 Arg Ser His Ile
 275

<210> SEQ ID NO 13
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Saccharum hybrid cultivar CP72-1210

<400> SEQUENCE: 13

Val Ile Gln Asn Ser Pro Pro Asp Leu
 1 5

41

What is claimed is:

1. A method of increasing expression of a transgene in a plant the expression of which is at least partially suppressed by PTGS, the method comprising:

transforming the plant with a nucleic acid having a sequence with at least 85% identity with the sequence of SEQ. ID. NO. 1 in its entirety, and encoding a protein operable to at least partially suppress PTGS having a sequence with at least 78% identity with the amino acid sequence of SEQ. ID. NO. 2,

wherein expression of the protein operable to at least partially suppress PTGS increases expression of the transgene in the plant.

2. The method according to claim 1, wherein the protein operable to at least partially suppress PTGS comprises a protein having the amino acid sequence of SEQ. ID. NO. 2.

3. The method according to claim 1, wherein the nucleic acid encoding the protein comprises a nucleic acid having the sequence of SEQ. ID. NO. 1 or the sequence of a degenerate code homolog of SEQ. ID. NO. 1.

4. The method according to claim 1, wherein the transgene is not involved in PTGS.

5. The method according to claim 1, wherein the plant is a monocot.

6. The method according to claim 5, wherein the monocot is sugarcane.

42

7. The method according to claim 5, wherein the monocot is corn.

8. The method according to claim 5, wherein the monocot is sorghum.

9. The method according to claim 5, wherein the monocot is rice.

10. The method according to claim 1, wherein the plant is an embryonic callus.

11. The method according to claim 1, wherein the plant is a mature plant.

12. The method according to claim 1, wherein the protein operable to at least partially suppress PTGS comprises Sorghum Mosaic Virus (SrMV) P1/HC-Pro.

13. A method of increasing expression of a transgene in a plant the expression of which is at least partially suppressed by PTGS, the method comprising:

transforming the plant with a nucleic acid encoding a protein operable to at least partially suppress PTGS,

wherein the protein operable to at least partially suppress PTGS has the amino acid sequence of SEQ. ID. NO. 2 and expression of the protein operable to at least partially suppress PTGS increases expression of the transgene in the plant.

* * * * *