

# Actively Growing Bacteria in the Inland Sea of Japan, Identified by Combined Bromodeoxyuridine Immunocapture and Denaturing Gradient Gel Electrophoresis<sup>∇†</sup>

Koji Hamasaki,<sup>1\*</sup> Akito Taniguchi,<sup>1</sup> Yuya Tada,<sup>1</sup> Richard A. Long,<sup>2‡</sup> and Farooq Azam<sup>3</sup>

Graduate School of Biosphere Science, Hiroshima University, Hiroshima, Japan<sup>1</sup>; Department of Oceanography, College of Geoscience, Texas A&M University, College Station, Texas 77843<sup>2</sup>; and MBRD, Scripps Institution of Oceanography, University of California San Diego, La Jolla, California 92037<sup>3</sup>

Received 6 September 2006/Accepted 21 February 2007

**A fundamental question in microbial oceanography concerns the relationship between prokaryote diversity and biogeochemical function in an ecosystem context. We combined bromodeoxyuridine (BrdU) magnetic bead immunocapture and PCR-denaturing gradient gel electrophoresis (BUMP-DGGE) to examine phylotype-specific growth in natural marine assemblages. We also examined a broad range of marine bacterial isolates to determine their abilities to incorporate BrdU in order to test the validity of the method for application to diverse marine assemblages. We found that 27 of 29 isolates belonging to different taxa could incorporate BrdU. BUMP-DGGE analysis revealed phylogenetic affiliations of DNA-synthesizing, presumably actively growing bacteria across a eutrophic to mesotrophic transect in the Inland Sea of Japan. We found that the BrdU-incorporating (growing) communities were substantially different from the total communities. The majority (34/56) of phylotypes incorporated BrdU and were presumably growing, and these phylotypes comprised 10 alphaproteobacteria, 1 betaproteobacterium, 11 gammaproteobacteria, 11 *Cytophaga-Flavobacterium-Bacteroides* group bacteria, and 1 unclassified bacterium. All BrdU-responsive alphaproteobacteria were members of the *Rhodobacterales*, suggesting that such bacteria were dominant in the growing alphaproteobacterial populations in our samples. The BrdU-responsive gammaproteobacteria belonged to the *Oceanospirillales*, the SAR86 cluster, the *Pseudomonadales*, the *Alteromonadales*, and the *Vibrionales*. Thus, contemporaneous cooccurrence of diverse actively growing bacterial taxa was a consistent pattern in our biogeochemically varied study area.**

Bacteria in seawater play important roles in the ocean's food webs. Their activities and responses to organic substrates significantly influence the flux of organic matter and oceanic biogeochemical cycles (3, 4). Because only a small fraction of bacteria in any seawater sample can be isolated and cultivated, culture-independent methods employing molecular approaches have been used to great advantage during the last two decades to study the natural assemblages of marine bacteria (11, 19, 39). These and other studies have revealed unexpected diversity and dynamics of bacterial community structure in seawater (8, 16, 47). However, the great challenge remains to relate diversity to ecological function and biogeochemical activities of bacteria. In recent studies workers have combined fluorescence in situ hybridization (FISH) with microautoradiography and found differences in the uptake of specific organic substrates by different phylogenetic groups of bacteria (7, 8, 40). Stable-isotope probing has enabled PCR-based DNA analysis of bacteria that incorporate specific substrates (45).

Bromodeoxyuridine (BrdU), a halogenated nucleoside that

can serve as a thymidine (TdR) analog, has been widely used as an alternative to tritiated TdR (<sup>3</sup>H]TdR) incorporation to label proliferating cells in cell biology (2, 59). BrdU incorporated into de novo DNA can be detected using anti-BrdU monoclonal antibody. The applicability of BrdU incorporation as an alternative to <sup>3</sup>H]TdR incorporation for bulk measurement of bacterial secondary production in seawater has been demonstrated (22, 38, 50). Employing antibodies conjugated with magnetic beads, BrdU-labeled DNA can be separated from total DNA, and this immunocapture technique and 16S rRNA gene fingerprinting or sequencing have been used to reveal some of the growing bacterial species in both soil and lake water (1, 5, 58, 60). At the single-cell level, actively growing bacteria can be visualized by using anti-BrdU conjugated with a fluorescent label or an enzymatic reporter (23, 42, 58). In one study of coastal seawater, a significant portion of the 4',6'-diamidino-2-phenylindole (DAPI)-stained bacteria (47%) became BrdU positive in 5 h (23). Pernthaler et al. (42) combined the BrdU immunofluorescence method with 16S rRNA FISH and applied it to North Sea waters. They found that the actively growing community included bacteria belonging to the *Roseobacter*, *Alteromonas*, and SAR86 clades. However, detailed community structures and identities of actively growing bacteria in seawater or even in soils and lake water have not been determined yet.

A fingerprinting approach using reverse-transcribed 16S rRNA fragments has been used to infer potentially active bacterial populations in aquatic environments (14, 34, 48, 53, 57). A comparison of DNA- and RNA-based fingerprints provides

\* Corresponding author. Present address: Department of Marine Ecosystem Dynamics, Ocean Research Institute, The University of Tokyo, 1-15-1 Minamidai, Nakano-ku, Tokyo 164-8639, Japan. Phone: 81-3-5351-6337. Fax: 81-3-5351-6482. E-mail: hamasaki@ori.u-tokyo.ac.jp.

† Supplemental material for this article may be found at <http://aem.asm.org/>.

‡ Present address: Department of Biological Sciences, University of South Carolina, Columbia, SC 29208.

<sup>∇</sup> Published ahead of print on 2 March 2007.

TABLE 1. BrdU incorporation by marine bacterial isolates

Most closely allied taxon	Isolate	Source	Accession no.	BrdU antibody detection	
				Whole cell	DNA dot blot
<i>Alphaproteobacteria</i>					
<i>Erythrobacter</i>	KUR27	Seawater	AB266542	+	+
<i>Erythrobacter</i>	OB10	Seawater	AB266553	-	+
<i>Erythrobacter</i>	ATKR 01	Dinoflagellate	AB266563	+	+
<i>Sulfitobacter</i>	OB4	Seawater	AB266552	+	+
<i>Sulfitobacter</i>	OB14	Seawater	AB266554	+	+
<i>Thalassospira</i>	TAMA2	Dinoflagellate	AB266548	+	+
<i>Gammaproteobacteria</i>					
<i>Alteromonas</i>	KUR4	Seawater	AB266538	+	+
<i>Cobetia</i>	KUR58	Seawater	AB266544	-	+
<i>Colwellia</i>	SL5	Sea ice	AB266558	+	-
<i>Colwellia</i>	SL15	Sea ice	AB266559	+	+
<i>Glaciecola</i>	KUR9	Seawater	AB266539	+	+
<i>Marinobacter</i>	KUR64	Seawater	AB266546	-	+
<i>Pseudoalteromonas</i>	MIDOR1	Green algae	AB266549	+	+
<i>Pseudoalteromonas</i>	CHAKO4	Brown algae	AB266547	+	NA <sup>a</sup>
<i>Pseudoalteromonas</i>	OB0	Seawater	AB266551	+	+
<i>Pseudoalteromonas</i>	KUR2	Seawater	AB266537	+	+
<i>Pseudomonas</i>	KUR18	Seawater	AB266541	-	+
<i>Psychrobacter</i>	KUR35	Seawater	AB266543	-	+
<i>Vibrio</i>	RYO1	Green algae	AB266555	+	+
<i>Betaproteobacteria</i>					
<i>Chromobacterium violaceum</i>	IAM12470 (= ATCC 12472)	Freshwater	AE016825	+	NA
CFB group					
<i>Algibacter</i>	OB15	Seawater	AB266550	+	+
<i>Algoriphagus</i>	SL1	Sea ice	AB266556	+	-
<i>Flavobacterium</i>	SL8	Sea ice	AB266561	-	-
<i>Maribacter</i>	TAMA1	Dinoflagellate	AB266562	+	-
<i>Maribacter</i>	SL4	Sea ice	AB266557	+	+
<i>Pibocella</i>	SL14	Sea ice	AB266560	-	-
Low-G+C-content gram-positive bacteria					
<i>Bacillus</i>	KUR61	Seawater	AB266545	-	+
<i>Actinobacteria</i>					
<i>Microbacterium</i>	KUR17	Seawater	AB266540	-	+
<i>Microbacterium</i>	ATKR 03-1	Dinoflagellate	AB266564	+	+

<sup>a</sup> NA, not analyzed.

insight into total bacterial communities and their potentially active fractions. Recent work using DNA- and RNA-derived clone libraries indicated that the RNA-derived library included distinct phylotypes which were not detected in the DNA library, suggesting that RNA-based analysis could be used to characterize actively metabolizing but less abundant bacteria in a community (33). This concept is based on the assumption that the number of ribosomes per cell is a measure of cell activity, including the growth rate. However, the RNA contents of species vary greatly (17, 28), which makes interpretation problematic for natural mixed-species communities. Also, differences between DNA- and RNA-derived fingerprints may be attributed to random priming and cDNA synthesis in reverse transcription-PCR analyses.

In this study, we used combined BrdU magnetic bead immunocapture and PCR-denaturing gradient gel electrophoresis (BUMP-DGGE) to investigate the community structures and phylogenetic identities of actively growing bacteria across

a eutrophic to mesotrophic transect. This approach is similar to the method developed by Borneman (5) for soil samples. As one of the essential assumptions for the validity of the method is that all taxa in the microbial communities in a sample are able to incorporate BrdU into DNA, we also examined a broad range of marine bacterial isolates to determine whether they could incorporate BrdU.

#### MATERIALS AND METHODS

**Isolation and phylogenetic identification of marine bacteria.** To determine whether bacteria could incorporate BrdU, marine isolates were obtained from various environments, such as coastal seawater, sea ice, marine macroalgae, and dinoflagellate cultures (Table 1). Cultures were grown and bacteria were isolated using 0.5× ZoBell 2216E medium. Isolates were analyzed by performing PCR with partial 16S rRNA gene sequences, followed by DGGE for phylogenetic typing (37). Isolated strains that yielded PCR amplicons which migrated at the same position on a gel were grouped into one phylotype. The phylogenetic affiliation of each representative isolate was examined by partially sequencing the 16S rRNA gene. Bacterial 16S rRNA genes were amplified by colony PCR

using universal forward primer 27F and bacterium-specific reverse primer 1492R. After PCR products were treated with a QIAquick PCR purification kit (QUIAGEN Inc., Valencia, CA), bidirectional sequencing was performed with a Dual CyDye terminator sequencing kit and a Long-Read Tower automated DNA sequencer (Amersham Biosciences Corp., Piscataway, NJ), using primers complementary to the bases at positions 341 to 358 and 517 to 534. Sequences were aligned with known sequences in the DDBJ (DNA Data Bank of Japan) database using BLAST (Basic Local Alignment Search Tool). Since we found no betaproteobacteria, one representative strain of this taxon (*Chromobacterium violaceum* IAM12470 [= ATCC 12472]) was obtained from the IAM culture collection at the Institute of Molecular and Cellular Biosciences, The University of Tokyo.

**Incubation of isolates with BrdU.** Isolates were grown in SWM medium at room temperature for 1 to 2 days. The SWM medium used for BrdU labeling of cultured isolates was prepared by adding 0.0058% (wt/vol) Casamino Acids, 0.69 mM glucose, 0.30 mM  $\text{NH}_4\text{Cl}$ , 0.25 mM  $\text{NaH}_2\text{PO}_4$ , 0.2  $\mu\text{M}$  ferric citrate, and 0.2  $\mu\text{M}$  EDTA to seawater filtered with Whatman GF/F filters (G. Steward and F. Azam, unpublished data) and was modified by adding 0.001% (wt/vol) Bacto peptone (Difco). The seawater was replaced with distilled water for culturing and labeling of *C. violaceum*. An aliquot of each culture was inoculated into fresh SWM medium to obtain an optical density at 600 nm of 0.01 to 0.02. After 30 min of incubation at 23°C, 200 nM (final concentration) BrdU was added to the tubes (9 ml). Control tubes received 200 nM TdR. The cultures were incubated for an additional 12 h, and then the contents were pelleted by centrifugation at  $20,000 \times g$  for 5 min. The supernatants were discarded, and the pelleted cells were fixed with 4% paraformaldehyde, washed with phosphate-buffered saline (PBS), and resuspended in PBS. Each cell suspension was mixed with an equal volume of 100% ethanol and stored at  $-15^\circ\text{C}$ .

**Detection of incorporated BrdU.** BrdU was detected by two immunochemical methods, one employing immunocytochemistry with whole-cell preparations and the other employing a chemiluminescence immunoassay with genomic DNA dot blots. Immunocytochemical fluorescence staining and assessment by fluorescence microscopy were performed by using the procedure described previously (23). Aliquots (5  $\mu\text{l}$ ) of bacterial cell suspensions were spotted onto 6-mm-diameter wells of Teflon printed glass slides (Electron Microscopy Science, Hatfield, PA) and air dried for 1 h. After the cells were permeabilized and the double-stranded DNA was denatured, anti-BrdU antibody was applied to detect incorporated BrdU. The antibody signal was amplified by catalyzed reporter deposition and was observed by using an Olympus BX51 epifluorescence microscope (total magnification,  $\times 1,000$ ; Olympus Corp., Tokyo, Japan). Also, a chemiluminescence immunoassay with genomic DNA dot blots was performed by using the procedure described previously (22). Aliquots (100  $\mu\text{l}$ ) of bacterial cell suspensions were filtered through a nylon membrane (pore size, 0.22  $\mu\text{m}$ ; MagnaGraph; GE Water & Process Technologies, Trevose, PA) with a 96-well-format dot blotter (Minifold I; Schleicher & Schuell, Dassel, Germany). After the cells were lysed, cross-linked DNA on the membrane was detected with anti-BrdU antibody. The chemiluminescence signal was developed with an enzyme-conjugated antibody and then detected by using a cooled charge-coupled device imaging system (VersaDoc5000; Bio-Rad Laboratories, Hercules, CA).

**Field sampling and BrdU labeling.** Coastal seawater was collected at nine stations in the western part of the Inland Sea of Japan from the R/V *Toyoshio-Maru* in July 2002 and 2003 (ST1 at  $34^\circ 19' \text{N}$ ,  $132^\circ 22' \text{E}$ ; ST2 at  $34^\circ 09' \text{N}$ ,  $132^\circ 18' \text{E}$ ; ST3 at  $34^\circ 01' \text{N}$ ,  $132^\circ 30' \text{E}$ ; ST4 at  $33^\circ 40' \text{N}$ ,  $132^\circ 23' \text{E}$ ; ST5 at  $33^\circ 34' \text{N}$ ,  $132^\circ 14' \text{E}$ ; ST7 at  $33^\circ 24' \text{N}$ ,  $132^\circ 14' \text{E}$ ; ST8 at  $33^\circ 06' \text{N}$ ,  $132^\circ 14' \text{E}$ ; ST9 at  $33^\circ 17' \text{N}$ ,  $132^\circ 27' \text{E}$ ; and ST10 at  $32^\circ 15' \text{N}$ ,  $132^\circ 00' \text{E}$ ). Depth profiles for salinity and temperature at each site were obtained with a Sea-Bird SBE9plus conductivity-temperature-depth profiler (Sea-Bird Electronics, Inc., Bellevue, WA). Seawater was collected from a depth of 5 m with a Van Dorn bottle and was filtered with 100- $\mu\text{m}$  nylon mesh to remove large zooplankton. Samples were diluted (1:3) with 0.22- $\mu\text{m}$  filtered seawater collected from the same site to reduce micro- and nanozooplankton grazing on bacteria during incubation for BrdU labeling. BrdU was added to the seawater at a final concentration of 1  $\mu\text{M}$ , and samples were incubated in a 10-liter dark plastic bottle for 1 or 5 h at the in situ temperature  $\pm 2^\circ\text{C}$ . Bacteria were collected on 0.22- $\mu\text{m}$  cartridge filters (Sterivex GS; Millipore Corp., Billerica, MA).

To determine the chlorophyll *a* concentration, duplicate seawater samples were filtered (pressure differential,  $<100$  mm Hg) through Whatman GF/F filters and extracted in the dark with  $N,N$ -dimethyl formamide at  $4^\circ\text{C}$  for 24 h (52), and the chlorophyll *a* concentration was determined fluorometrically (25). Particulate matter collected on precombusted (500°C, 2 h) Whatman GF/F filters was analyzed to determine the particulate organic carbon and particulate organic nitrogen contents using a CHN analyzer (Yanaco Co., Ltd., Tokyo, Japan). Seawater filtrates obtained with Whatman GF/F filters were collected for analysis

of dissolved organic carbon in precombusted glass ampoules and frozen after flame sealing until analysis with a Shimadzu TOC5000 (Shimadzu Corp., Kyoto, Japan) (51). Seawater samples fixed with 2% (final concentration) formalin were stained with DAPI, filtered onto 0.2- $\mu\text{m}$ -pore-size black polycarbonate filters (Millipore), and mounted on glass slides (44). Bacterial abundance was determined by counting the bacteria in 10 fields using epifluorescence microscopy (Olympus BX-51).

**DNA extraction.** DNA was extracted by using the xanthogenate-sodium dodecyl sulfate protocol described by Tillett and Neilan (54), with volume modification. Each Sterivex GS filter unit was thawed and filled with xanthogenate-sodium dodecyl sulfate buffer (1% potassium ethyl xanthogenate [Wako Pure Chemical, Osaka, Japan], 100 mM Tris-HCl [pH 7.4], 20 mM EDTA [pH 8], 1% sodium dodecyl sulfate, 800 mM ammonium acetate). After the filter housing was sealed, the unit was incubated at  $70^\circ\text{C}$  for 120 min. Following incubation, the filter unit was vortexed for 10 s. The lysate was eluted into 2-ml microcentrifuge tubes and then placed on ice for 30 min. Cell debris was pelleted by centrifugation at  $22,000 \times g$  for 15 min at  $4^\circ\text{C}$ . The supernatant was transferred to a clean 1.5-ml microcentrifuge tube and mixed with an equal volume of 100% isopropanol. After incubation at room temperature for 10 min, the precipitated DNA was pelleted by centrifugation at  $22,000 \times g$  for 20 min. The pellet was washed once with 70% ethanol, air dried, and resuspended in 50  $\mu\text{l}$  of water.

**BrdU magnetic bead immunocapture.** Immunocapture was performed as described by Urbach et al. (58), with slight modifications. In the procedures described below, all preparations were incubated at room temperature. Herring sperm DNA (1.25 mg  $\text{ml}^{-1}$  in PBS; Trevigen Inc., Gaithersburg, MD) was boiled for 1 min, quickly frozen in dry-ice ethanol, and thawed. The solution was mixed (9:1) with anti-BrdU monoclonal antibody (diluted 1:10 in PBS; Sigma-Aldrich, St. Louis, MO) and incubated for 30 min. Extracted DNA samples (0.5 to 1  $\mu\text{g}$ ) were boiled for 1 min, quickly frozen in dry-ice ethanol, and thawed. Each denatured sample solution was mixed with 10  $\mu\text{l}$  of the herring sperm DNA-antibody mixture and incubated for 30 min. Paramagnetic beads coated with goat anti-mouse immunoglobulin G (Dynabeads; Dynal Biotech, Oslo, Norway) were washed with PBS containing acetylated bovine serum albumin (BSA) (1 mg  $\text{ml}^{-1}$ ; Nacalai Tesque, Kyoto, Japan), using a magnetic concentrator. The beads were resuspended in PBS-BSA at their initial concentration. The bead suspension (25  $\mu\text{l}$ ) was mixed with each sample (20  $\mu\text{l}$ ). After incubation for 30 min with constant agitation, the beads were washed seven times with 0.5 ml PBS-BSA. BrdU-labeled DNA captured by the beads was eluted by adding 100  $\mu\text{l}$  of 1.7 mM BrdU (in PBS-BSA) and incubating the preparation for 30 min with constant agitation. Glycogen (2  $\mu\text{l}$ ; 20 mg  $\text{ml}^{-1}$ ; Roche Diagnostics, Mannheim, Germany) was added to the bead supernatant, and the DNA was collected by two rounds of ethanol precipitation. The precipitate was resuspended in 10 to 20  $\mu\text{l}$  of water and subjected to PCR amplification and DGGE. To evaluate the specificity of immunocapture, a parallel DNA sample without BrdU labeling was processed in the same manner and subjected to PCR amplification as a negative control.

**DGGE.** Bacterial 16S rRNA genes were amplified from either immunocaptured or nonimmunocaptured DNA samples by a touchdown PCR protocol (12) and then subjected to bacterial community analysis by DGGE. Nonimmunocaptured samples were analyzed before and after incubation. A bacterial primer complementary to bases at positions 341 to 358 containing a 40-bp GC clamp was used as the forward primer. A universal 17-bp primer complementary to bases at positions 517 to 534 was used as the reverse primer for the samples collected in 2002 (37). A different universal primer complementary to bases at positions 907 to 926 was used for the samples collected in 2003 (6).

The PCR products (200 to 500 ng) were loaded onto 8% polyacrylamide gels in  $0.5 \times \text{TAE}$  (20 mM Tris, 10 mM acetate, 0.5 mM  $\text{Na}_2\text{EDTA}$ ; pH 8.2) with a denaturing gradient from 20 to 70% from the top to the bottom. Electrophoresis was performed at 85 V for 16 h at  $60^\circ\text{C}$  in a hot-bath DGGE unit (CBS Scientific, San Diego, CA) with a running buffer consisting of  $0.5 \times \text{TAE}$ . Gels were stained for 30 min with  $1 \times \text{SYBR}$  gold nucleic acid stain in  $0.5 \times \text{TAE}$  (Molecular Probes, Eugene, OR) and were destained for 10 min with  $0.5 \times \text{TAE}$ . The gels were then visualized and documented using a charge-coupled device camera imaging system (Printgraph AE6911CX; ATTO Corp., Tokyo, Japan). DGGE bands were excised, and the DNA was eluted overnight. The eluants were used for PCR reamplification, followed by cloning with a TOPO TA cloning kit (Invitrogen, Carlsbad, CA). The presence of inserts was verified by PCR amplification using primers M13F and M13R. Clones with insertions that were the correct size were analyzed by DGGE to confirm the position of the cloned bands relative to the position of the original sample.

**Cluster analysis of DGGE banding patterns.** To assess the similarity of community structures (DGGE banding patterns), the Jaccard coefficient was calcu-




FIG. 1. Structures of total and active bacterial communities in the Inland Sea of Japan in July 2002. Seawater was incubated with  $1 \mu\text{M}$  BrdU for 5 h to label actively growing bacteria. The PCR products amplified from BrdU-containing and immunocaptured DNA [BrdU-incorporated ( $T_5$ )] were compared with the products amplified from total community DNA [Total ( $T_0$ )] by DGGE analysis. Each excised, cloned, and sequenced band is numbered. The relationships of excised band sequences to other sequences in the DDBJ database are shown in Table 3.

lated (18), and the distance matrix was analyzed by using the between-group average linkage method for clustering with the software SPSS Base 10.0J (SPSS Inc., Chicago, IL). The presence of DGGE bands in a sample was determined by eye, and the results were scored 1 or 0 to generate a binary matrix. This matrix was used to generate a distance matrix and a dendrogram with SPSS.

**Sequencing and phylogenetic analysis.** Bidirectional sequencing was performed with a Dual CyDye terminator sequencing kit using T7 primers and a Long-Read Tower automated DNA sequencer (Amersham Biosciences Corp.). Sequences were aligned with known sequences in the DDBJ database using BLAST. Phylogenetic relationships were inferred by pairwise comparison and the neighbor-joining method using CLUSTAL W (55). Phylogenetic trees were edited using Treeview (41).

**Nucleotide sequence accession numbers.** The nucleotide sequences of the 16S rRNA gene determined in this study have been deposited in the DDBJ nucleotide sequence database under accession numbers AB266537 to AB266564. The nucleotide sequences of DGGE bands have been deposited in the DDBJ nucleotide sequence database under accession numbers AB265970 to AB266026.

## RESULTS

**BrdU incorporation by marine bacterial isolates.** We obtained 98 isolates, which grouped into 28 phylotypes based on DGGE migration of 16S rRNA gene partial fragments. The

abilities of 28 marine isolates plus one betaproteobacterium from a culture collection to incorporate BrdU were examined. The phylogenetic groups and affinities with known genera of our isolates were determined by sequencing 16S rRNA genes (Table 1). These isolates included 6 alphaproteobacteria, 13 gammaproteobacteria, 6 *Cytophaga-Flavobacterium-Bacteroides* (CFB) group bacteria, 2 low-G+C-content gram-positive bacteria, and 1 actinobacterium. Twenty-seven of the 29 isolates incorporated BrdU under the culture conditions that we employed (Table 1).

**Environmental factors.** The near-surface salinity increased along our north-south transect, with lower salinity observed near the Ohta River runoff in Hiroshima Bay in the northern part of the sampling area, while the temperature did not exhibit such a pattern (see Table S1 in the supplemental material). The concentration of chlorophyll *a* varied from 0.35 to 14  $\text{mg m}^{-3}$ , and the concentrations at ST1 were high (11 and 14  $\text{mg m}^{-3}$ ). The particulate organic carbon and particulate organic nitrogen concentrations were also highest at ST1 and




FIG. 2. Structures of total and active bacterial communities in the Inland Sea of Japan in July 2003. Seawater was incubated with 1  $\mu$ M BrdU for 1 h to label actively growing bacteria. The PCR products amplified from BrdU-containing and immunocaptured DNA [BrdU-incorporated ( $T_1$ )] were compared with the products amplified from total community DNA before [Total ( $T_0$ )] and after [Total ( $T_1$ )] incubation by DGGE analysis. Each excised, cloned, and sequenced band is numbered. The relationships of excised band sequences to other sequences in the DDBJ database are shown in Table 3.

lowest at the southern end of transect. The same pattern was observed for the dissolved organic carbon concentration. The total bacterial counts ranged from  $3.3 \times 10^5$  to  $4.6 \times 10^6$  cells  $ml^{-1}$ , and a regular pattern was not observed.

**BUMP-DGGE analysis.** To confirm the specificity of immunocapture, control samples were processed like the BrdU-treated samples and then subjected to PCR. The control samples were obtained from parallel incubation mixtures to which BrdU was not added, as well as mixtures amended with BrdU. No PCR products were obtained with the control unlabeled samples, while PCR products that were the appropriate size (ca. 200 or 500 bp) were obtained with the BrdU-treated samples.

To rapidly compare the growing fraction of the bacterial community with the total bacterial community, we performed a DGGE analysis of the 16S rRNA genes. BrdU labeling and DGGE analysis of 16S rRNA gene fragments produced banding patterns for the BrdU-incorporating bacterial community that were used for comparisons with the total bacterial community for both 2002 and 2003 samples (Fig. 1 and 2). In both transects the bacterial community structure varied, and the variation was greater in 2003 (Fig. 1 and 2). Two distinct communities were observed for the 2002 transect; the communities in Hiroshima Bay (ST1a, ST2, and ST3) and the outer samples (ST4 and ST5) exhibited about 80% similarity, as determined by using Jaccard's index. When the total bacterial community was compared to the BrdU-incorporating phylotypes, the levels of dissimilarity were greater, and the differences between the community at ST1a, ST2, and ST3 and the community at ST4 and ST5 were reinforced. In 2003, a similar pattern was observed; the community at the Hiroshima Bay station (ST1b) was different from the community at the re-




FIG. 3. Relationship between total and active bacterial community structures in the Inland Sea of Japan in 2002 (A) and 2003 (B). The structure of the total community (Total) was compared with the structure of the BrdU-incorporating community (BrdU-incorporated) before ( $T_0$ ) and after ( $T_1$  or  $T_5$ ) incubation by DGGE analysis. The tree was constructed by using the between-group average linkage method with the Jaccard coefficient distance matrix for clustering with the software SPSS Base 10.0J (SPSS Inc.).

TABLE 2. Comparisons of total and BrdU-incorporating operational taxonomic units in the Inland Sea of Japan<sup>a</sup>

Sampling location	No. of DGGE bands							% of BrdU-positive bands	
	Total (T <sub>0</sub> )	Total (T <sub>1</sub> )	BrdU	In common <sup>b</sup>		Unique <sup>c</sup>		Total (T <sub>0</sub> ) and BrdU	Total (T <sub>1</sub> ) and BrdU
				Total (T <sub>0</sub> ) and BrdU	Total (T <sub>1</sub> ) and BrdU	Total (T <sub>0</sub> ) and BrdU	Total (T <sub>1</sub> ) and BrdU		
ST1a	16		17	4		29		59	
ST2	16		16	4		28		57	
ST3	13		17	4		26		65	
ST4	22		15	7		30		50	
ST5	15		16	6		25		64	
ST1b	13	11	13	6	7	20	17	65	76
ST7	10	12	10	4	5	16	17	63	59
ST8	11	10	7	3	4	15	13	47	54
ST9	11	11	13	7	7	17	17	76	76
ST10	12	12	10	3	3	19	19	53	53

<sup>a</sup> Total (T<sub>0</sub>), total community before incubation; Total (T<sub>1</sub>), total community after incubation; BrdU, BrdU-incorporating community.

<sup>b</sup> Number of bands appearing at the same position for the total community and the BrdU-incorporating community samples.

<sup>c</sup> Number of unique bands found at each sampling location.

maining stations. In 2003, the total communities at the beginning and at the end of the incubations were similar (<10% difference) as determined by a pairwise analysis of samples from the same site; however, these communities differed by 25% or more from the BrdU-incorporating communities at the same site (Fig. 2 and 3B). Comparisons of DGGE bands revealed that 15 to 30 phylotypes were present at each location and that 47 to 76% of these phylotypes were present in BrdU-incorporating fractions (Table 2).

**Sequencing and phylogenetic analysis of 16S rRNA gene fragments.** To investigate which groups were represented in the total community fractions and the BrdU-incorporating fractions, we excised 57 DGGE bands or phylotypes (34 bands in 2002 and 23 bands in 2003) (Fig. 1 and 2) from the gels and sequenced them; these phylotypes included 24 phylotypes from the BrdU-incorporating community (Table 3). We found 51 phylotypes related to 15 alphaproteobacteria, 1 betaproteobacterium, 15 gammaproteobacteria, 17 CFB group bacteria, 2 cyanobacteria, and 1 unclassified bacterium (Fig. 4 and 5). Forty of these phylotypes were found in BrdU-incorporating fractions. These phylotypes were related to 11 alphaproteobacteria, 1 betaproteobacterium, 13 gammaproteobacteria, 14 CFB group bacteria, and 1 unclassified bacterium. All of the phylotypes that were related to the *Alphaproteobacteria* except one unclassified phylotype were clustered in two major subgroups, the *Rhodobacterales* and SAR11 clusters, and all of the BrdU-incorporating phylotypes were clustered in the *Rhodobacterales*. The phylotypes that were related to the *Gammaproteobacteria* were clustered in five major subgroups, the *Oceanospirillales*, the SAR86 cluster, the *Pseudomonadales*, the *Alteromonadales*, and the *Vibrionales*, and BrdU-positive phylotypes were present in every cluster. In particular, the phylotype corresponding to *Vibrio splendidus* (Fig. 1, bands 25 and 33) was present at all locations from ST1a to ST5, and the signals of this phylotype at ST2 and ST5 were intense. Also, the two phylotypes corresponding to uncultured CFB group bacteria (Fig. 2, bands 40, 38, and 57) were present at all locations from ST1b to ST10, as determined by BrdU-positive bands. Almost the same phylotype (band 7) as the band 38 and 57 phylotype was found to be present at ST1, ST4, and ST5, as determined

by BrdU-positive bands (Fig. 1). Eleven phylotypes did not appear in BrdU-incorporating fractions. These phylotypes included three SAR11 group bacteria, one bacterium belonging to the *Rhodobacterales*, one bacterium belonging to the *Oceanospirillales*, one unclassified gammaproteobacterium, three CFB group bacteria, and two cyanobacteria (Table 3).

## DISCUSSION

A fundamental question in microbial oceanography concerns the relationship between prokaryote diversity and biogeochemical function in an ecosystem context. While culture-independent methods for determining diversity are well established, it is still methodologically challenging to determine phylotype-specific in situ activities of bacteria that have not been cultured yet. Phylotype-specific growth performance and phenome expression in natural marine assemblages are influenced by physicochemical and nutritional conditions, as well as inter- and intraspecies interactions. Insight at the broad phylogenetic group level has been obtained by combining [<sup>3</sup>H]TdR incorporation and FISH (8, 9, 40). RNA-based DGGE fingerprints (reverse transcription-PCR-DGGE) have revealed phylotypes with active gene expression and whether a gene of interest was expressed (13, 57).

BUMP-DGGE complements these methods by addressing phylotype-specific growth (growing organisms versus nongrowing organisms). It allows workers to determine phylogenetic affiliations of DNA-synthesizing, presumably actively growing bacteria in seawater, as in this study. Thus, it is possible to determine the phylotypes contributing to bacterial production and biogeochemical cycles at a given time and locale. As suggested by Borneman (5), this approach could also help determine phylotype-specific functional gene expression (e.g., genes important in biogeochemical cycles) in a given ecological scenario, although this was not tested here. Furthermore, if pelagic archaeon-specific primers are used, the protocol should reveal archaeal growth and help address currently cutting-edge questions (e.g., whether some or all shallow-water or deep-sea *Archaea* are chemoautotrophic) (24, 26). This approach should

TABLE 3. Affiliations of excised DGGE band sequences with other sequences in the DDBJ database

DGGE band <sup>a</sup>	Clone	Accession no. <sup>b</sup>	DNA fraction <sup>c</sup>	BrdU <sup>d</sup>	Phylogenetic group
From ST1a to ST5 in 2002					
1	HB02-12	AB265970	Total	+	<i>Gammaproteobacteria</i> , SAR86
2 (17)	HB02-13	AB265971	Total	+	CFB
3	HB02-14	AB265972	Total	+	<i>Gammaproteobacteria</i> , SAR86
4	HB02-15	AB265973	Total	-	CFB
5 (32)	HB02-16	AB265974	Total	+	<i>Gammaproteobacteria</i> , <i>Pseudomonadales</i>
6	HB02-17	AB265975	Total	+	CFB
7	HB02-18	AB265976	Total	+	CFB
8	HB02-19	AB265977	Total	+	<i>Gammaproteobacteria</i>
9	HB02-20	AB265978	Total	-	<i>Alphaproteobacteria</i> , SAR11
10	HB02-27	AB265979	Total	+	<i>Alphaproteobacteria</i>
11	SIS02-21	AB265980	Total	+	<i>Gammaproteobacteria</i> , SAR86
12	HB02-22	AB265981	Total	+	<i>Gammaproteobacteria</i>
13	SIS02-30	AB265982	Total	-	Cytnobacteria
14 (27)	SIS02-31	AB265983	Total	+	<i>Gammaproteobacteria</i> , <i>Alteromonadales</i>
15	SIS02-95	AB265984	Total	-	<i>Alphaproteobacteria</i> , SAR11
16	SIS02-26	AB265985	Total	-	<i>Alphaproteobacteria</i> , SAR11
17 (2)	HB02-5b	AB265986	BrdU	+	CFB
18	HB02-6b	AB265987	BrdU	+	CFB
19	HB02-7b	AB265988	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
20	HB02-8b	AB265989	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
21	HB02-9b	AB265990	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
22	HB02-10b	AB265991	BrdU	+	<i>Alphaproteobacteria</i>
23	HB02-11b	AB265992	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
24	HB02-2b	AB265993	BrdU	+	<i>Gammaproteobacteria</i> , <i>Oceanospirillales</i>
25 (33)	HB02-4b	AB265994	BrdU	+	<i>Gammaproteobacteria</i> , <i>Vibrionales</i>
26	HB02-1b	AB265995	BrdU	+	CFB
27 (14)	HB02-3b	AB265996	BrdU	+	<i>Gammaproteobacteria</i> , <i>Alteromonadales</i>
28	SIS02-35b	AB265997	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
29	SIS02-28b	AB265998	BrdU	+	<i>Gammaproteobacteria</i> , <i>Vibrionales</i>
30	SIS02-92b	AB265999	BrdU	+	<i>Gammaproteobacteria</i> , <i>Vibrionales</i>
31	SIS02-29b	AB266000	BrdU	+	<i>Gammaproteobacteria</i> , <i>Pseudomonadales</i>
32	SIS02-33b	AB266001	BrdU	+	<i>Gammaproteobacteria</i> , <i>Pseudomonadales</i>
33 (25)	SIS02-36b	AB266002	BrdU	+	<i>Gammaproteobacteria</i> , <i>Vibrionales</i>
34	SIS02-34b	AB266003	BrdU	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
From ST1b to ST10 in 2003					
35 (37)	HB03-70	AB266004	Total	-	CFB
36	HB03-3	AB266005	Total	+	CFB
37 (35)	SIS03-9	AB266006	Total	-	CFB
38 (57)	SIS03-10	AB266007	Total	+	CFB
39	SIS03-11	AB266008	Total	-	<i>Gammaproteobacteria</i> , <i>Oceanospirillales</i>
40	SIS03-13	AB266009	Total	+	CFB
41	SIS03-19	AB266010	Total	+	CFB
42	SIS03-20	AB266011	Total	+	<i>Alphaproteobacteria</i>
43	SIS03-21	AB266012	Total	+	<i>Alphaproteobacteria</i>
44	SIS03-23	AB266013	Total	+	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
45	SIS03-24	AB266014	Total	+	<i>Gammaproteobacteria</i>
46	SIS03-25	AB266015	Total	+	Unclassified
47	SIS03-26	AB266016	Total	-	<i>Gammaproteobacteria</i>
48	SIS03-27	AB266017	Total	-	Cyanobacteria
49	SIS03-18	AB266018	Total	+	CFB
50	SIS03-34	AB266019	Total	-	<i>Alphaproteobacteria</i> , <i>Rhodobacterales</i>
51	HB03-36	AB266020	Total	-	CFB
52	HB03-62b	AB266021	BrdU	+	CFB
53	HB03-102b	AB266022	BrdU	+	<i>Betaproteobacteria</i>
54	SIS03-64b	AB266023	BrdU	+	CFB
55	SIS03-108b	AB266024	BrdU	+	CFB
56	SIS03-55b	AB266025	BrdU	+	CFB
57	SIS03-65b	AB266026	BrdU	+	CFB

<sup>a</sup> The number in parentheses indicates a band that had the same sequence.

<sup>b</sup> Accession numbers for sequences obtained in this study.

<sup>c</sup> Origin of the excised and sequenced bands. Total, total community DNA fraction; BrdU, BrdU-incorporating community DNA fraction.

<sup>d</sup> Presence (+) or absence (-) of the band in samples representing BrdU-incorporating fractions.




FIG. 4. Distribution of phylotypes found in the Inland Sea of Japan: phylogenetic tree showing the similarities of sequences from bands (bold type) to sequences from members of the *Alpha*- and *Gammaproteobacteria*. The tree was inferred by the neighbor-joining method using CLUSTAL W. The bands found in lanes representing the BrdU-incorporating community are indicated by asterisks. The scale bar indicates the number of base pair substitutions per nucleotide position.

also guide the development of enrichment or pure cultures of bacteria and archaea that have not been cultured yet.

In laboratory-based studies workers have found that not all bacteria incorporate TdR (27), so the bacteria that are not responsive to TdR may also not incorporate BrdU because of the lack of a transporter or thymidine kinase (43). In previous studies researchers found that 34/36 isolates were responsive to

BrdU (23, 42, 58). We tested 29 additional isolates, and only 2 were not responsive to BrdU (Table 1). Cumulatively then, 61/65 isolates belonging to diverse phylogenetic groups (i.e., *Alphaproteobacteria*, *Gammaproteobacteria*, *Betaproteobacteria*, CFB group, low-G+C-content gram-positive bacteria, and *Actinobacteria*) incorporated BrdU. Thus, the BUMP-DGGE approach is potentially broadly applicable to phylogenetically






FIG. 5. Distribution of phylotypes found in the Inland Sea of Japan: phylogenetic tree showing the similarities of sequences from bands (bold type) to sequences from members of the CFB group bacteria, cyanobacteria, and *Betaproteobacteria*. The tree was inferred by the neighbor-joining method using CLUSTAL W. The bands found in lanes representing the BrdU-incorporating community are indicated by asterisks. The scale bar indicates the number of base pair substitutions per nucleotide position.

diverse assemblages in pelagic marine environments; however, three of the four negative bacterial isolates were affiliated with the CFB group, which reportedly can dominate bacterial assemblages in seawater environments and especially associate with organic particles. The possible significance of bacteria that are not responsive to BrdU must not be dismissed, and this phenomenon needs to be examined further. Also, there is a lower limit for retrieving and detecting BrdU-labeled DNA, mainly because of a loss of labeled DNA during immunocapture. Hence, some bacteria may be overlooked due to their very low densities in a community even if they are actively growing and incorporating BrdU.

The bacterial community structure at low-salinity stations was quite different from that at other stations. ST1 was located near an area where there was river runoff, which presumably supported a community with a high concentration of chlorophyll *a* (see Table S1 in the supplemental material). Changes in physicochemical properties of estuarine waters, including salinity, have been found to be associated with community shifts

(7, 10). Micro-FISH previously showed that [<sup>3</sup>H]TdR assimilation due to major bacterial phylogenetic groups changed along a salinity gradient from 0.1 to 29 practical salinity units (7). In our study, the DGGE patterns at low-salinity stations ST1, ST2, and ST3 were generally different from those at ST4 and ST5. One might expect the presence of freshwater phylotypes in BUMP-DGGE profiles at low-salinity stations. While the salinity was not very low (31.5 practical salinity units), freshwater influence was reflected in communities that incorporated BrdU, and one band obtained in 2003 (ST1b) was closely related to *Delftia acidovorans* belonging to the *Betaproteobacteria*, the dominant group in freshwater environments.

Two incubation times (1 and 5 h) were employed for BrdU labeling in this study. Although a shorter incubation time is preferable to avoid bias due to protistan grazing or viral lysis, 5 h was chosen to achieve enough labeling for immunocapture in 2002; however, a shorter incubation (1 h) was found to be enough in 2003. DGGE banding patterns for 1- and 5-h incubations were compared in 2003, and no significant differences

were observed (not shown). When the method is used for coastal seawater samples, 1 h of incubation is recommended for BrdU labeling.

Micro-FISH studies have shown that the correlation between abundance and growth (assimilation of TdR) was not significant for alphaproteobacteria, gammaproteobacteria, and CFB group bacteria in the Delaware estuary (9). The results implied that the impact of mortality was greater than the impact of growth in structuring the bacterial community. We found that BrdU-incorporating (growing) communities were largely different than the "total" communities. The percentages of overlapping bands for the total and BrdU-labeled communities were  $\leq 40\%$  (Table 2). Indeed, many bands (86/134) for growing phylotypes were not detectable in the gels for the "total" community. The difference could not be explained by a shift in the total community since the 1-h incubation time was too short to significantly change the "total" community banding (Fig. 3). It is tempting to propose that BUMP-DGGE was able to detect a number of rapidly growing but low-abundance (hence undetected in "total" community gels) phylotypes. However, the minimum BrdU incorporation intensity for positive immunocapture is not known, and further work is needed to test this hypothesis. One motivation for a future test of this hypothesis is that phylotypes that are present at low levels but have high growth rates can make large contributions to community bacterial production and organic matter fluxes. Rapidly growing bacteria might be subject to intense grazing by protists or viral lysis and thus prevented from becoming abundant enough to be detected in PCR-DGGE.

The *Roseobacter* and SAR11 clusters are two major subgroups of *Alphaproteobacteria* found in marine environments (19). In our study, these two clusters exhibited contrasting characteristics with respect to BrdU incorporation. We found 11 phylotypes related to *Roseobacter* but no phylotypes related to the SAR11 cluster in BrdU-incorporating fractions. We thought that since *Roseobacter*-related phylotypes were reportedly rapid growers, they might dominate actively growing alphaproteobacterial populations in our samples. Gonzalez et al. (20) reported that *Roseobacter* and SAR11 were numerically important in the heterotrophic bacterial community (average,  $>20\%$  of the 16S rRNA genes sampled) and important in cycling of organic sulfur compounds during a North Atlantic algal bloom. The importance of these clusters was also shown by bacterial production and dimethylsulfoniopropionate consumption during an algal bloom in the North Sea (61). Members of the *Roseobacter* cluster are commonly found in association with algal cells and have relatively high growth rates in nutrient-rich conditions (21, 47). As suggested previously (20, 31, 32, 35, 61), *Roseobacter*-related bacteria are well adapted to nutrient-rich conditions, such as coastal waters, algal blooms, and microscale "hot spots" in oceanic waters.

In contrast, SAR11 bacteria might be relatively slow growers in coastal waters, because they did not respond well to BrdU even though at least three representatives were present. This hypothesis is consistent with the results of culture studies of *Pelagibacter ubique*, the cultured representative of the SAR11 group, which grows in a pristine seawater medium at very low rates (46). However, it is inconsistent with the conclusion reached by Malmstrom et al. (30). These workers examined in situ assimilation of leucine by SAR11 bacteria in the Northwest

Atlantic Ocean and suggested that the specific growth rates of SAR11 bacteria either equaled or exceeded the growth rates of the total prokaryotic community. One possible explanation for these findings is that there are metabolically different subpopulations of SAR11 cluster bacteria, because this cluster is a phylogenetically, and probably metabolically, diverse group of bacteria (19). Alternatively, members of the SAR11 cluster may not be able to incorporate BrdU and thus may have escaped detection as growing bacteria in this study. Methodology needs to be developed to test for the possible presence of uncultured bacteria that are growing but are not responsive to BrdU.

A phylotype similar to *V. splendidus* was found in BrdU-incorporating fractions at all sampling locations from ST1 to ST5. Quantification of the *V. splendidus* population by quantitative PCR in a previous study (56) indicated that it was consistently present as a member of the coastal bacterioplankton community. *Vibrio* species, including many pathogenic strains, are widely distributed in marine environments. This group can comprise  $\sim 10\%$  of the readily culturable bacteria, whereas culture-independent molecular analyses suggested that it accounts for  $<1\%$  of total bacterial populations (15). However, the molecularly based estimates to date do not account for the difference between active and inactive populations. Our previous work examining BrdU incorporation at the single-cell level in 1-h incubations suggested that  $\sim 20\%$  of the total bacteria in coastal waters are robustly growing (23). Hence, *Vibrio* spp. that are active growers in coastal waters may contribute disproportionately more to bacterial production than the amount expected from calculations based on their biomass.

In conclusion, this study revealed the consistent cooccurrence of diverse actively growing bacterial taxa in rather small seawater samples in all natural assemblages examined over biogeochemically varied areas in coastal waters. While cooccurrence of diverse taxa is commonly reported, there is only a limited amount of culture-independent data on contemporaneous growth of diverse bacterial taxa. Microscale patchiness of organic matter distribution might be invoked as a mechanism creating microniches in the pelagic realm (29). Experimental perturbation experiments involving BUMP-DGGE might be used to determine the environmental mechanisms for maintenance and regulation of the diversity and the phylogenetic identity of the growing taxa. Further application of the approach to various marine environments can help determine key players in bacterium-mediated biogeochemical processes, as well as bacterial productivity. There may be some universal active growers, considering the wide distribution of particular types of bacteria, such as SAR11, RCA, and *Silicibacter*-like bacteria (35, 36, 49). Once the key players in a particular environment are determined, specific probes can be designed and used to obtain quantitative estimates of their dynamics by FISH and/or quantitative PCR of immunocaptured DNA. Such analyses should lead to linkages of microbial community dynamics and microbial metabolic activity in ocean biogeochemical modeling.

#### ACKNOWLEDGMENTS

We thank the officers and crew of the R/V *Toyoshio-maru* for their assistance and support during the cruise.

This research was supported by grants-in-aid 15688006 and 16651007 from JSPS and by an Environmental Research Grant from the Showa Shell Sekiyu Foundation to K.H. F.A. was supported by NSF grants ANT04-44134 and OCE04-28900. R.A.L. was supported by NSF grant MCB 0453877.

## REFERENCES

1. Artursson, V., and J. K. Jansson. 2003. Use of bromodeoxyuridine immunocapture to identify active bacteria associated with arbuscular mycorrhizal hyphae. *Appl. Environ. Microbiol.* **69**:6208–6215.
2. Asquith, B., C. Debecq, D. Macallan, L. Willems, and C. Bangham. 2002. Lymphocyte kinetics: the interpretation of labeling data. *Trends Immunol.* **23**:596–601.
3. Azam, F. 1998. Microbial control of oceanic carbon flux: the plot thickens. *Science* **280**:694–696.
4. Azam, F., and R. Long. 2001. Sea snow microcosms. *Nature* **414**:495–498.
5. Borneman, J. 1999. Culture-independent identification of microorganisms that respond to specific stimuli. *Appl. Environ. Microbiol.* **65**:3398–3400.
6. Casamayor, E., H. Schafer, L. Baneras, C. Pedros-Alio, and G. Muyzer. 2000. Identification of and spatio-temporal differences between microbial assemblages from two neighboring sulfurous lakes: comparison by microscopy and denaturing gradient gel electrophoresis. *Appl. Environ. Microbiol.* **66**:499–508.
7. Cottrell, M., and D. Kirchman. 2003. Contribution of major bacterial groups to bacterial biomass production (thymidine and leucine incorporation) in the Delaware estuary. *Limnol. Oceanogr.* **48**:168–178.
8. Cottrell, M., and D. Kirchman. 2000. Natural assemblages of marine proteobacteria and members of the *Cytophaga-Flavobacter* cluster consuming low- and high-molecular-weight dissolved organic matter. *Appl. Environ. Microbiol.* **66**:1692–1697.
9. Cottrell, M., and D. Kirchman. 2004. Single-cell analysis of bacterial growth, cell size, and community structure in the Delaware estuary. *Aquat. Microb. Ecol.* **34**:139–149.
10. Crump, B., C. Hopkinson, M. Sogin, and J. Hobbie. 2004. Microbial biogeography along an estuarine salinity gradient: combined influence of bacterial growth and residence time. *Appl. Environ. Microbiol.* **70**:1494–1505.
11. DeLong, E. 2002. Microbial population genomics and ecology. *Curr. Opin. Microbiol.* **5**:520–524.
12. Don, R. H., P. T. Cox, B. J. Wainwright, K. Baker, and J. S. Mattick. 1991. 'Touchdown' PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Res.* **19**:4008.
13. Ebie, Y., N. Noda, H. Miura, M. Matsumura, S. Tsuneda, A. Hirata, and Y. Inamori. 2004. Comparative analysis of genetic diversity and expression of amoA in wastewater treatment processes. *Appl. Microbiol. Biotechnol.* **64**:740–744.
14. Eichler, S., R. Ghristen, C. Holtje, P. Westphal, J. Botel, I. Brettar, A. Mehling, and M. Hoffe. 2006. Comparison and dynamics of bacterial communities of a drinking water supply system as assessed by RNA- and DNA-based 16S rRNA gene fingerprinting. *Appl. Environ. Microbiol.* **72**:1858–1872.
15. Eilers, H., J. Pernthaler, F. Glockner, and R. Amann. 2000. Culturability and in situ abundance of pelagic bacteria from the North Sea. *Appl. Environ. Microbiol.* **66**:3044–3051.
16. Fandino, L., L. Riemann, G. Steward, R. Long, and F. Azam. 2001. Variations in bacterial community structure during a dinoflagellate bloom analyzed by DGGE and 16S rDNA sequencing. *Aquat. Microb. Ecol.* **23**:119–130.
17. Fegatella, F., J. Lim, S. Kjelleberg, and R. Cavicchioli. 1998. Implications of rRNA operon copy number and ribosome content in the marine oligotrophic ultramicrobacterium *Sphingomonas* sp. strain RB2256. *Appl. Environ. Microbiol.* **64**:4433–4438.
18. Ferrarri, V., and J. Hollibaugh. 1999. Distribution of microbial assemblages in the central Arctic ocean basin studied by PCR/DGGE: analysis of a large data set. *Hydrobiologia* **401**:55–68.
19. Giovannoni, S., and M. Rappe. 2000. Evolution, diversity, and molecular ecology of marine prokaryotes, p. 47–84. *In* D. L. Kirchman (ed.), *Microbial ecology of the oceans*. Wiley-Liss, New York, NY.
20. Gonzalez, J., R. Simo, R. Massana, J. Covert, E. Casamayor, C. Pedros-Alio, and M. Moran. 2000. Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom. *Appl. Environ. Microbiol.* **66**:4237–4246.
21. Grossart, H., F. Levold, M. Allgaier, M. Simon, and T. Binkhoff. 2005. Marine diatom species harbour distinct bacterial communities. *Environ. Microbiol.* **7**:860–873.
22. Hamasaki, K. 2006. Comparison of bromodeoxyuridine immunoassay with tritiated thymidine radioassay for measuring bacterial productivity in oceanic waters. *J. Oceanogr.* **62**:793–799.
23. Hamasaki, K., R. Long, and F. Azam. 2004. Individual cell growth rates of marine bacteria, measured by bromodeoxyuridine incorporation. *Aquat. Microb. Ecol.* **35**:217–227.
24. Herndl, G. J., T. Reinthaler, E. Teira, H. van Aken, C. Veth, A. Pernthaler, and J. Pernthaler. 2005. Contribution of *Archaea* to total prokaryotic production in the deep Atlantic Ocean. *Appl. Environ. Microbiol.* **71**:2303–2309.
25. Holm-Hansen, O., C. J. Lorenzen, R. W. Holems, and J. D. H. Strickland. 1965. Fluorometric determination of chlorophyll. *J. Cons. Perm. Int. Explor. Mer* **30**:3–15.
26. Ingalls, A. E., S. R. Shah, R. L. Hansman, L. I. Aluwihare, G. M. Santos, E. R. M. Druffel, and A. Pearson. 2006. Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon. *Proc. Natl. Acad. Sci. USA* **103**:6442–6447.
27. Jeffrey, W., and J. Paul. 1990. Thymidine uptake, thymidine incorporation, and thymidine kinase activity in marine bacterium isolates. *Appl. Environ. Microbiol.* **56**:1367–1372.
28. Lee, S., and P. Kemp. 1994. Single-cell RNA content of natural marine planktonic bacteria measured by hybridization with multiple 16S rRNA-targeted fluorescent probes. *Limnol. Oceanogr.* **39**:869–879.
29. Long, R., and F. Azam. 2001. Microscale patchiness of bacterioplankton assemblage richness in seawater. *Aquat. Microb. Ecol.* **26**:103–113.
30. Malmstrom, R., M. Cottrell, H. Elifantz, and D. Kirchman. 2005. Biomass production and assimilation of dissolved organic matter by SAR11 bacteria in the Northwest Atlantic Ocean. *Appl. Environ. Microbiol.* **71**:2979–2986.
31. Malmstrom, R., R. Kiene, M. Cottrell, and D. Kirchman. 2004. Contribution of SAR11 bacteria to dissolved dimethylsulfoniopropionate and amino acid uptake in the North Atlantic Ocean. *Appl. Environ. Microbiol.* **70**:4129–4135.
32. Malmstrom, R., R. Kiene, and D. Kirchman. 2004. Identification and enumeration of bacteria assimilating dimethylsulfoniopropionate (DMSP) in the North Atlantic and Gulf of Mexico. *Limnol. Oceanogr.* **49**:597–606.
33. Moeseneder, M., J. Arrieta, and G. Herndl. 2005. A comparison of DNA- and RNA-based clone libraries from the same marine bacterioplankton community. *FEMS Microbiol. Ecol.* **51**:341–352.
34. Moeseneder, M., C. Winter, and G. Herndl. 2001. Horizontal and vertical complexity of attached and free-living bacteria of the eastern Mediterranean Sea, determined by 16S rDNA and 16S rRNA fingerprints. *Limnol. Oceanogr.* **46**:95–107.
35. Moran, M. A., A. Buchan, J. M. Gonzalez, J. F. Heidelberg, W. B. Whitman, R. P. Kiene, J. R. Henriksen, G. King, R. Belas, C. Fuqua, L. Brinkac, M. Lewis, S. Johri, B. Weaver, G. Pai, J. A. Eisen, E. Rahe, W. M. Sheldon, W. Ye, T. R. Miller, J. Carlton, D. A. Rasko, I. T. Paulsen, Q. Ren, S. C. Daugherty, R. T. Deboy, R. J. Dodson, A. S. Durkin, R. Madupu, W. C. Nelson, S. A. Sullivan, M. J. Rosovitz, D. H. Haft, J. Selengut, and N. Ward. 2004. Genome sequence of *Silicibacter pomeroyi* reveals adaptations to the marine environment. *Nature* **432**:910–913.
36. Morris, R., M. Rappe, S. Connon, K. Vergin, W. Siebold, C. Carlson, and S. Giovannoni. 2002. SAR11 clade dominates ocean surface bacterioplankton communities. *Nature* **420**:806–810.
37. Muyzer, G., E. Del Waal, and A. Uitterlinden. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* **59**:695–700.
38. Nelson, C. E., and C. Carlson. 2005. A nonradioactive assay of bacterial productivity optimized for oligotrophic pelagic environments. *Limnol. Oceanogr. Methods* **3**:211–220.
39. Olsen, G. J., D. J. Lane, S. J. Giovannoni, N. R. Pace, and D. A. Stahl. 1986. Microbial ecology and evolution: a ribosomal RNA approach. *Annu. Rev. Microbiol.* **40**:337–365.
40. Ouverney, C., and J. Fuhrman. 1999. Combined microautoradiography-16S rRNA probe technique for determination of radioisotope uptake by specific microbial cell types in situ. *Appl. Environ. Microbiol.* **65**:1746–1752.
41. Page, R. 1996. TREEVIEW: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**:357–358.
42. Pernthaler, A., J. Pernthaler, M. Schattner, and R. Amann. 2002. Identification of DNA-synthesizing bacterial cells in coastal North Sea plankton. *Appl. Environ. Microbiol.* **68**:5728–5736.
43. Pollard, P., and D. Moriarty. 1984. Validity of the tritiated thymidine method for estimating bacterial growth rates: measurement of isotope dilution during DNA synthesis. *Appl. Environ. Microbiol.* **48**:1076–1083.
44. Porter, K. G., and Y. S. Feig. 1980. The use of DAPI for identifying and counting aquatic microflora. *Limnol. Oceanogr.* **25**:943–948.
45. Radajewski, S., P. Ineson, N. Parekh, and J. Murrell. 2000. Stable-isotope probing as a tool in microbial ecology. *Nature* **403**:646–649.
46. Rappe, M., S. Connon, K. Vergin, and S. Giovannoni. 2002. Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**:630–633.
47. Riemann, L., G. Steward, and F. Azam. 2000. Dynamics of bacterial community composition and activity during a mesocosm diatom bloom. *Appl. Environ. Microbiol.* **66**:578–587.
48. Schafer, H., L. Bernard, C. Courties, P. Lebaron, P. Servais, R. Pukall, E. Stackebrandt, M. Troussellier, T. Guindulain, J. Vives-Rego, and G. Muyzer. 2001. Microbial community dynamics in Mediterranean nutrient-enriched seawater mesocosms: changes in the genetic diversity of bacterial populations. *FEMS Microbiol. Ecol.* **34**:243–253.
49. Selje, N., M. Simon, and T. Brinkhoff. 2004. A newly discovered Roseobacter cluster in temperate and polar oceans. *Nature* **427**:445–448.

50. Steward, G., and F. Azam. 1999. Bromodeoxyuridine as an alternative to <sup>3</sup>H-thymidine for measuring bacterial productivity in aquatic samples. *Aquat. Microb. Ecol.* **19**:57–66.
51. Sugimura, Y., and Y. Suzuki. 1988. A high-temperature catalytic oxidation method for the determination of non-volatile dissolved organic carbon in seawater by direct injection of a liquid sample. *Mar. Chem.* **24**:105–131.
52. Suzuki, R., and T. Ishimaru. 1990. An improved method for the determination of phytoplankton chlorophyll using N,N-dimethylformamide. *J. Oceanogr.* **46**:190–194.
53. Teske, A., C. Wawer, G. Muyzer, and N. Ramsing. 1996. Distribution of sulfate-reducing bacteria in a stratified fjord (Mariager Fjord, Denmark) as evaluated by most-probable-number counts and denaturing gradient gel electrophoresis of PCR-amplified ribosomal DNA fragments. *Appl. Environ. Microbiol.* **62**:1405–1415.
54. Tillett, D., and B. Neilan. 2000. Xanthogenate nucleic acid isolation from cultured and environmental cyanobacteria. *J. Phycol.* **36**:251–258.
55. Thompson, J., D. Higgins, and T. Gibson. 1994. Clustal W: improving the sensitivity of progressive multiple-sequence alignments through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
56. Thompson, J. R., S. Pacocha, C. Pharino, V. Klepac-Ceraj, D. E. Hunt, J. Benoit, R. Sarma-Rupavtarm, D. L. Distel, and M. F. Polz. 2005. Genotypic diversity within a natural coastal bacterioplankton population. *Science* **307**:1311–1313.
57. Troussellier, M., H. Schafer, N. Batailler, L. Bernard, C. Courties, P. Lebaron, G. Muyzer, P. Servais, and J. Vivas-Rego. 2002. Bacterial activity and genetic richness along an estuarine gradient (Rhône River plume, France). *Aquat. Microb. Ecol.* **28**:13–24.
58. Urbach, E., K. Vergin, and S. Giovanoni. 1999. Immunochemical detection and isolation of DNA from metabolically active bacteria. *Appl. Environ. Microbiol.* **65**:1207–1213.
59. Van Furth, R., and T. Van Zwet. 1988. Immunocytochemical detection of 5-bromo-2-deoxyuridine incorporation in individual cells. *J. Immunol. Methods* **108**:45–51.
60. Yin, B., D. Crowley, G. Sparovek, W. De Melo, and J. Borneman. 2000. Bacterial functional redundancy along a soil reclamation gradient. *Appl. Environ. Microbiol.* **66**:4361–4365.
61. Zubkov, M., B. Fuchs, S. Archer, R. Kiene, R. Amann, and P. Burkill. 2001. Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulphoniopropionate in an algal bloom in the North Sea. *Environ. Microbiol.* **3**:304–311.