## Probing Dark Matter at the LHC using Vector Boson Fusion Processes

Andres G. Delannoy<sup>2</sup>, Bhaskar Dutta<sup>1</sup>, Alfredo Gurrola<sup>2</sup>, Will Johns<sup>2</sup>, Teruki Kamon<sup>1,3</sup>,

Eduardo Luiggi<sup>4</sup>, Andrew Melo<sup>2</sup>, Paul Sheldon<sup>2</sup>, Kuver Sinha<sup>1</sup>, Kechen Wang<sup>1</sup>, and Sean Wu<sup>1</sup>

<sup>1</sup> Mitchell Institute for Fundamental Physics and Astronomy,

Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843-4242, USA

Department of Physics and Astronomy, Vanderbilt University, Nashville, TN, 37235, USA

<sup>3</sup> Department of Physics, Kyungpook National University, Daegu 702-701, South Korea

Department of Physics, University of Colorado, Boulder, CO 80309-0390, USA

Vector boson fusion (VBF) processes at the Large Hadron Collider (LHC) provide a unique opportunity to search for new physics with electroweak couplings. A feasibility study for the search of supersymmetric dark matter in the final state of two VBF jets and large missing transverse energy is presented at 14 TeV. Prospects for determining the dark matter relic density are studied for the cases of Wino and Bino-Higgsino dark matter. The LHC could probe Wino dark matter with mass up to approximately 600 GeV with a luminosity of 1000 fb<sup>-1</sup>.

Nearly 80% of the matter of the Universe is dark matter (DM) [1]. The identity of DM is one of the most profound questions at the interface of particle physics and cosmology. Weakly interacting massive particles (WIMPs) are particularly promising DM candidates that can explain the observed relic density and are under investigation in a variety of direct and indirect searches. Within the context of *R*-parity conserving supersymmetric extensions of the standard model (SM), the WIMP DM candidate is the lightest supersymmetric particle (LSP), typically the lightest neutralino ( $\tilde{\chi}_1^0$ ), which is a mixture of Bino, Wino, and Higgsino states.

The DM relic density is typically determined by its annihilation cross section at the time of thermal freeze-out. For supersymmetric WIMP DM, the annihilation cross section depends on the mass of  $\tilde{\chi}_1^0$  and its couplings to various SM final states, for which a detailed knowledge of the composition of  $\tilde{\chi}_1^0$  in gaugino/Higgsino states is required. Moreover, other states in the electroweak sector, such as sleptons, staus, or charginos can enter the relic density calculation.

It is important to probe the electroweak sector of supersymmetric models directly in order to study their DM connection. The main challenge to a direct probe of the electroweak sector at the Large Hadron Collider (LHC) is the small production cross section of neutralinos, charginos, and sleptons [2].

In this Letter we explore supersymmetric DM produced directly at the LHC using vector boson fusion (VBF) processes [3, 4]. This appears particularly promising since some of the present authors recently showed that VBF production is quite effective in probing the chargino-neutralino system [5]. It has also been suggested that VBF processes might be useful in both Higgs boson and supersymmetry studies [6–10]. VBF production is characterized by the presence of two tagging jets with large dijet invariant mass in the forward region in opposite hemispheres. As shown in [5], the requirement of tagging jets along with missing transverse energy ( $\not\!\!\!E_T$ ) is very efficient in reducing SM background.

We note that the production of squarks  $(\tilde{q})$  or gluinos  $(\tilde{g})$  through gluon fusion, followed by cascade decay ending in the production of DM, is the classic setting for DM searches in final states with appreciable missing energy, multiple jets and leptons. However, determining the content of the neutralino and the masses of the superpartners without any color charges requires specific model dependent correlation between masses of colored and non-colored superpartners. In very specific settings, it is possible to determine the composition of  $\tilde{\chi}_1^0$  [11], as well as the mass of light staus or sleptons [12]. In general, the combinatoric background poses a major problem for such attempts.

Recently, experiments at the 8-TeV LHC (LHC8) have put lower bounds on the masses of the  $\tilde{g}$  and  $\tilde{q}$ . For comparable masses, the exclusion limits are approximately 1.5 TeV at 95% CL with 13 fb<sup>-1</sup> of integrated luminosity [13–15]. There are also active searches for the lightest top squark ( $\tilde{t}$ ), and exclusion limits in the  $m_{\tilde{t}}$ - $m_{\tilde{\chi}_1^0}$  plane have been obtained in certain decay modes [16, 17].

A direct probe of the electroweak sector using VBF processes is complementary to such searches. A variety of possibilities exist for the colored sector (compressed spectra, mildly fine-tuned split scenarios [18], non-minimal supersymmetric extensions, etc.) with varying implications for existing and future searches. Experimental constraints (e.g. triggering) significantly affect the ability to probe supersymmetric DM in some of the above scenarios, for example those with compressed spectra. The important point to note is that a direct probe of the electroweak sector is largely agnostic about the fate of the colored sector and provides a direct window to DM physics.

The strategy pursued in this Letter will be to investigate direct DM production by VBF processes in events with  $2j + \not\!\!\!E_T$  in the final state. Such an approach has several advantages. The  $2j + \not\!\!\!E_T$  final state configuration provides a search strategy that is free from trigger bias. This is reinforced as the  $p_T$  thresholds for triggering objects are raised by ATLAS and CMS experiments.

In order to probe DM directly, the following processes are investigated:

$$pp \to \tilde{\chi}_1^0 \, \tilde{\chi}_1^0 \, jj, \ \tilde{\chi}_1^\pm \, \tilde{\chi}_1^\mp \, jj, \ \tilde{\chi}_1^\pm \, \tilde{\chi}_1^0 \, jj \ .$$
 (1)

The main sources of SM background are: (i)  $pp \rightarrow Zjj \rightarrow \nu\nu \nu jj$  and (ii)  $pp \rightarrow Wjj \rightarrow l\nu jj$ . The former is an irreducible background with the same topology as the signal. The  $\not{E}_{\rm T}$  comes from the neutrinos. The latter arises from events which survive a lepton veto; (iii)  $pp \rightarrow t\bar{t}$ +jets: This background may be reduced by vetoing b-jets, light leptons,  $\tau$  leptons and light-quark/gluon jets.

Distributions of  $p_{\rm T}(j_1)$ ,  $p_{\rm T}(j_2)$ ,  $M_{j_1j_2}$ , and  $E_{\rm T}$  for background as well as VBF pair production of DM are studied at  $\sqrt{s} = 8$  TeV and 14 TeV. In the case of pure Wino or Higgsino DM,  $\tilde{\chi}_1^{\pm}$  is taken to be outside the exclusion limits for ATLAS' disappearing track analysis [21] and thus VBF production of  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\pm}$ ,  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp}$ , and  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0$  also contribute. The  $\tilde{\chi}_1^0$  masses chosen for this study are in the range 100 GeV to 1 TeV. The colored sector is assumed to be much heavier. There is no contribution to the neutralino production from cascade decays of colored particles.

Events are preselected by requiring  $\not\!\!\!E_T > 50 \text{ GeV}$ and the two leading jets  $(j_1, j_2)$  each satisfying  $p_T \geq 30$ GeV with  $|\Delta \eta(j_1, j_2)| > 4.2$  and  $\eta_{j_1} \eta_{j_2} < 0$ . The preselected events are used to optimize the final selections to achieve maximal signal significance  $(S/\sqrt{S}+B)$ . For the final selections, the following cuts are employed: (i)The tagged jets are required to have  $p_T > 50$  GeV and  $M_{j_1j_2} > 1500$  GeV; (ii) Events with loosely identified leptons  $(l = e, \mu, \tau_h)$  and b-quark jets are rejected, reducing the  $t\bar{t}$  and  $Wjj \rightarrow l\nu jj$  backgrounds by approximately  $10^{-2}$  and  $10^{-1}$ , respectively, while achieving 99% efficiency for signal events. The *b*-jet tagging efficiency used in this study is 70% with a misidentification probability of 1.5%, following Ref. [22]. Events with a third jet (with  $p_T > 50 \text{ GeV}$ ) residing between  $\eta_{j_1}$  and  $\eta_{j_2}$  are also rejected; (iii) The  $\not\!\!\!E_{\rm T}$  cut is optimized for each different value of the DM mass. For  $m_{\tilde{\chi}^0_1} = 100 \text{ GeV}$  (1 TeV),  $\not\!\!E_{\rm T} \ge 200 \text{ GeV} (450 \text{ GeV})$  is chosen, reducing the  $Wjj \rightarrow l\nu jj$  background by approximately  $10^{-3} (10^{-4})$ .

The production cross section as a function of  $m_{\tilde{\chi}_1^0}$  after requiring  $|\Delta \eta(j_1, j_2)| > 4.2$  is displayed in Fig. 1. The left and right panels show the cross sections for LHC8 and LHC14, respectively. For the pure Wino and Higgsino cases, inclusive  $\tilde{\chi}_1^0 \tilde{\chi}_1^0$ ,  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\pm}$ ,  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp}$ , and  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0$  production cross sections are displayed. The green (solid) curve corresponds to the case where  $\tilde{\chi}_1^0$  is 99% Wino. The inclusive production cross section is ~ 40 fb for a 100 GeV Wino at LHC14, and falls steadily with increasing mass. The cross section is approximately 5 – 10 times smaller for the pure Higgsino case, represented by the green (dashed) curve. As the Higgsino fraction in  $\tilde{\chi}_1^0$  decreases for a given mass, the cross section is ~ 10<sup>-2</sup> fb for  $m_{\tilde{\chi}_1^0} = 100$  GeV at LHC14.



FIG. 1: Production cross section as a function of  $m_{\tilde{\chi}_1^0}$  after requiring  $|\Delta \eta(j_1, j_2)| > 4.2$ , at LHC8 and LHC14. For the pure Wino and Higgsino cases, inclusive  $\tilde{\chi}_1^0 \tilde{\chi}_1^0$ ,  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\pm}$ ,  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp}$ , and  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0$  production cross sections are displayed.

Figure 2 shows the dijet invariant mass distribution  $M_{j_1j_2}$  for the tagging jet pair  $(j_1, j_2)$  and main sources of background, after the pre-selection cuts and requiring  $p_T > 50$  GeV for the tagging jets at LHC14. The dashed black curves show the distribution for the case of a pure Wino DM, with  $m_{\tilde{\chi}_1^0} = 50$  and 100 GeV. The dijet invariant mass distribution for W+ jets, Z+ jets, and  $t\bar{t}+$ jets background are also displayed. Clearly, requiring  $M_{j_1j_2} > 1500$  GeV is effective in rejecting background events, resulting in a reduction rate between  $10^{-4}$  and  $10^{-2}$  for the backgrounds of interest.

Figure 3 shows the  $\not\!\!\!E_T$  distribution for an integrated luminosity of 500 fb<sup>-1</sup> at LHC14 after all final selections except the  $\not\!\!\!E_T$  requirement. There is a significant enhancement of signal events in the high  $\not\!\!\!E_T$  region.

The significance as a function of  $\tilde{\chi}_1^0$  mass is plotted in Fig. 4 for different luminosities at LHC14. The blue, red, and black curves correspond to luminosities of 1000, 500, and 100 fb<sup>-1</sup>, respectively. At 1000 fb<sup>-1</sup>, a significance of  $5\sigma$  can be obtained up to a Wino mass of approximately 600 GeV.

Determining the composition of  $\tilde{\chi}_1^0$  for a given mass is very important in order to understand early universe cosmology. For example, if  $\tilde{\chi}_1^0$  has a large Higgsino or Wino component, the annihilation cross section is too



FIG. 2: Distribution of the dijet invariant mass  $M_{j_1j_2}$  normalized to unity for the tagging jet pair  $(j_1, j_2)$  and main sources of background after pre-selection cuts and requiring  $p_T > 50$ GeV for the tagging jets at LHC14. The dashed black curves show the distribution for the case where  $\tilde{\chi}_1^0$  is a nearly pure Wino with  $m_{\tilde{\chi}_1^0} = 50$  and 100 GeV. Inclusive  $\tilde{\chi}_1^0 \tilde{\chi}_1^0, \tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\pm},$  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^{\mp}$ , and  $\tilde{\chi}_1^{\pm} \tilde{\chi}_1^0$  production is considered.



large to fit the observed relic density for  $m_{\tilde{\chi}_1^0}$  mass less than ~ 1 TeV for Higgsinos [23] and ~ 2.5 TeV for Winos. On the other hand if  $\tilde{\chi}_1^0$  is mostly Bino, the annihilation cross section is too small. In the first case one has under-abundance whereas in the second case one has over-abundance of DM. Both problems can be solved if the DM is non-thermal [24] (in the case of thermal DM, addressing the over abundance problem requires addition effects like resonance, coannihilation etc. in the cross section, while the under-abundance problem can be addressed by having multi-component DM [25]). If  $\tilde{\chi}_1^0$  is a suitable mixture of Bino and Higgsino, the observed DM relic density can be satisfied.

From Figs. 1 and 3, it is clear that varying of the rate and the shape of the  $\not\!\!E_T$  distribution can be used to solve



FIG. 4: Significance curves for the case where  $\tilde{\chi}_1^0$  is 99% Wino as a function of  $m_{\tilde{\chi}_1^0}$  mass for different luminosities at LHC14. The green lines correspond to  $3\sigma$  and  $5\sigma$  significances.

In conclusion, this Letter has investigated the direct production of supersymmetric DM by VBF processes at the LHC. The cases of pure Wino, pure Higgsino, and mixed Bino-Higgsino DM have been studied in the  $2j + E_{\rm T}$  final state at 14 TeV. The presence of the energetic VBF jets with large dijet invariant mass as well as the large  $\not\!\!\!E_T$  due to DM production have been used to reduce SM background. It has been shown that broad enhancements in the  $E_{\rm T}$  and VBF dijet mass distributions provide a smoking gun signature for VBF production of supersymmetric DM. By optimizing the  $E_{\rm T}$  cut for a given  $m_{\tilde{\chi}_{*}^{0}}$ , one can simultaneously fit the  $\not\!\!\!E_{\mathrm{T}}$  shape and observed rate in data to extract the mass and composition of  $\tilde{\chi}^0_1$ , and hence solve for the DM relic density. At an integrated luminosity of  $1000 \text{ fb}^{-1}$ , a significance of  $5\sigma$  can be obtained up to a Wino mass of approximately 600 GeV. The relic density can be determined to within



FIG. 5: Contour lines in the relic density- $m_{\tilde{\chi}_1^0}$  plane for 99% Wino (blue dashed) and 99% Higgsino (grey dotted) DMs expected with 500 fb<sup>-1</sup> of luminosity at LHC14. The relic density is normalized to its value at  $m_{\tilde{\chi}_1^0} = 100$  GeV.

- Planck Collaboration, arXiv:1303.5076 [astro-ph.CO]; WMAP Collaboration, arXiv:1212.5226 [astro-ph.CO].
- [2] ATLAS Collaboration, ATLAS-CONF-2012-154. ATLAS-CONF-2013-028.
- [3] R. N. Cahn and S. Dawson, Phys. Lett. B 136, 196 (1984)
   [Erratum-ibid. B 138, 464 (1984)].
- [4] J. D. Bjorken, Phys. Rev. D 47, 101 (1993).
- [5] B. Dutta, A. Gurrola, W. Johns, T. Kamon, P. Sheldon and K. Sinha, Phys. Rev. D 87, 035029 (2013) arXiv:1210.0964 [hep-ph].
- [6] D. L. Rainwater, D. Zeppenfeld and K. Hagiwara, Phys. Rev. D 59, 014037 (1998) [hep-ph/9808468].
- [7] D. Choudhury, A. Datta, K. Huitu, P. Konar, S. Moretti and B. Mukhopadhyaya, Phys. Rev. D 68, 075007 (2003) [hep-ph/0304192]. A. Datta and K. Huitu, Phys. Rev. D 67, 115006 (2003) [hep-ph/0211319].
- [8] G. -C. Cho, K. Hagiwara, J. Kanzaki, T. Plehn, D. Rainwater and T. Stelzer, Phys. Rev. D 73, 054002 (2006) [hep-ph/0601063].
- [9] A. Datta, P. Konar and B. Mukhopadhyaya, Phys. Rev. D 65, 055008 (2002) [hep-ph/0109071]; Phys. Rev. Lett. 88, 181802 (2002) [hep-ph/0111012]; P. Konar and B. Mukhopadhyaya, Phys. Rev. D 70, 115011 (2004) [hep-ph/0311347]; R. C. Cotta, J. L. Hewett, M. P. Le and T. G. Rizzo, arXiv:1210.0525 [hep-ph].
- [10] G. F. Giudice, T. Han, K. Wang and L. -T. Wang, Phys. Rev. D 81, 115011 (2010) [arXiv:1004.4902 [hep-ph]].
- [11] B. Dutta, T. Kamon, N. Kolev, K. Sinha, K. Wang and S. Wu, arXiv:1302.3231 [hep-ph].
- [12] R. L. Arnowitt, B. Dutta, A. Gurrola, T. Kamon, A. Krislock and D. Toback, Phys. Rev. Lett. **100**, 231802 (2008); B. Dutta, A. Gurrola, T. Kamon, A. Krislock, A. B. Lahanas, N. E. Mavromatos and D. V. Nanopoulos, Phys. Rev. D **79**, 055002 (2009); B. Dutta, T. Kamon, A. Krislock, N. Kolev and Y. Oh, Phys. Rev. D

20% (40%) for the case of a pure Wino (Higgsino) for 500 fb<sup>-1</sup> at LHC14, for  $m_{\tilde{\chi}_1^0} = 100$  GeV. We note that our study does not include the effect of large multiple interactions at high luminosity operations at the LHC. This is a very important subject, but outside the scope of the present work, because the final performance will depend on the planned upgrade of ATLAS and CMS detectors.

This work is supported in part by DOE Grant No. DE-FG02-95ER40917 and DE-FG02-04ER41290, NSF Award PHY-1206044, and by the World Class University (WCU) project through the National Research Foundation (NRF) of Korea funded by the Ministry of Education, Science, and Technology (Grant No. R32-2008-000-20001-0). K.S. would like to thank Nathaniel Craig for helpful discussions.

82, 115009 (2010); B. Dutta, T. Kamon, A. Krislock,
K. Sinha and K. Wang, Phys. Rev. D 85, 115007 (2012).

- [13] ATLAS Collaboration, arXiv:1208.0949 [hep-ex].
- [14] ATLAS Collaboration, J. High Energy Phys. 07, 167 (2012) [arXiv:1206.1760 [hep-ex]].
- [15] CMS Collaboration, arXiv:1207.1898 [hep-ex].
- [16] ATLAS Collaboration, ATLAS-CONF-2012-166.
- [17] ATLAS Collaboration, ATLAS-CONF-2012-167.
- [18] N. Arkani-Hamed, A. Gupta, D. E. Kaplan, N. Weiner and T. Zorawski, arXiv:1212.6971 [hep-ph]. A. Arvanitaki, N. Craig, S. Dimopoulos and G. Villadoro, arXiv:1210.0555 [hep-ph].
- [19] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer and T. Stelzer, J. High Energy Phys. 06, 128 (2011) [arXiv:1106.0522 [hep-ph]].
- [20] PGS4 is a parameterized detector simulator. We use version 4 (http://www.physics.ucdavis.edu/~conway/ research/software/pgs/pgs4-general.htm) in the LHC detector configuration.
- [21] ATLAS Collaboration, J. High Energy Phys. 01, 131 (2013) [arXiv:1210.2852 [hep-ex]]. ALEPH Collaboration, Phys. Lett. B 533, 223 (2002) [hep-ex/0203020].
  OPAL Collaboration, Eur. Phys. J. C 29, 479 (2003) [hep-ex/0210043]. DELPHI Collaboration, Eur. Phys. J. C 34, 145 (2004) [hep-ex/0403047].
- [22] CMS Collaboration, JINST 8, P04013 (2013) [arXiv:1211.4462 [hep-ex]].
- [23] R. Allahverdi, B. Dutta and K. Sinha, Phys. Rev. D 86, 095016 (2012) [arXiv:1208.0115 [hep-ph]]. H. Baer, V. Barger and D. Mickelson, arXiv:1303.3816 [hep-ph].
- [24] R. Allahverdi, B. Dutta and K. Sinha, arXiv:1212.6948 [hep-ph].
- [25] H. Baer, V. Barger, P. Huang, D. Mickelson, A. Mustafayev and X. Tata, arXiv:1212.2655 [hep-ph].