

Metabolism of Aflatoxin B₁ by Rat Hepatic Microsomes Induced by Polyhalogenated Biphenyl Congeners

MICHAEL R. HALVORSON,¹ TIMOTHY D. PHILLIPS,^{1*} STEVEN H. SAFE,² AND L. W. ROBERTSON²

Departments of Veterinary Public Health¹ and Veterinary Physiology and Pharmacology,² Texas A&M University, College Station, Texas 77843

Received 14 May 1984/Accepted 2 January 1985

The metabolism of aflatoxin B₁ to aflatoxins M₁ and Q₁ by rat liver microsomes from animals pretreated with polychlorinated or polybrominated biphenyl congeners depended on the structure of the halogenated biphenyl inducers. Microsomes from rats treated with phenobarbital (PB) or halogenated biphenyls that exhibit PB-type activity preferentially enhanced the conversion of aflatoxin B₁ to aflatoxin Q₁. In contrast, microsomes from rats treated with 3-methylcholanthrene (MC) or halogenated biphenyls that exhibit MC-type induction activity increased the metabolism of aflatoxin B₁ to aflatoxin M₁. The coadministration of PB and MC produced microsomes that exhibited both types of induction activity (mixed type) in catalyzing the oxidative metabolism of diverse xenobiotic agents. However, PB-plus-MC-induced hepatic microsomes from immature male Wistar rats preferentially increased the metabolism of aflatoxin B₁ to aflatoxin M₁ but did not enhance the conversion of aflatoxin B₁ to aflatoxin Q₁. Comparable results were observed with microsomes from rats pretreated with halogenated biphenyls classified as mixed-type inducers; moreover, in some cases there was a significant decrease in the conversion of aflatoxin B₁ to aflatoxin Q₁ (compared with that of controls treated with corn oil).

Aflatoxin B₁ (AFB₁) is a mold metabolite that has been characterized as a potent hepatotoxin and hepatocarcinogen in several animal species. As with many other toxins, the evidence suggests that AFB₁ requires oxidative metabolic activation to elicit the observed toxic and carcinogenic effects (7, 27). The active metabolite of AFB₁ is presumed to be 2,3-epoxy-2,3-dihydro-AFB₁. This metabolite is formed by the introduction of oxygen at the C-2-C-3 position via the mixed-function oxidase system (3, 9, 27). 2,3-Epoxy-2,3-dihydro-AFB₁ can alkylate critical cellular macromolecules or form a number of other metabolic products, including 2,3-dihydroxy-2,3-dihydro-AFB₁ and metabolic conjugates (3, 17).

Some reports demonstrate that the cytochrome P-450-dependent oxidation of AFB₁ can also occur at several other positions to produce a wide spectrum of metabolites that still retain their 2,3-olefinic functionality (4, 9, 11). Some of these compounds include aflatoxin M₁ (AFM₁) (oxidation at C-4), aflatoxin Q₁ (AFQ₁) (oxidation at C-9), and aflatoxin P₁ (O-dealkylation). Many of these AFB₁ metabolites are highly toxic and carcinogenic. AFM₁ is a major fungal and mammalian metabolite of AFB₁ and has been identified in animal livers, urine, and milk, including milk from cows (10). AFQ₁ is about 18 times less toxic than AFB₁ is to chicken embryos, and it was not mutagenic in a bacterial assay that included rat liver microsomes (10). In vitro studies indicate that AFM₁, 2,3-dihydroxy-2,3-dihydro-AFB₁, and AFQ₁ are the major metabolic products formed after incubation with microsomal monooxygenases (10, 13, 17).

Several reports show that xenobiotic agent-induced microsomal monooxygenases can markedly alter the metabolic profile of AFB₁ (8, 9, 13). The preferential induction of specific cytochrome P-450 isozymes catalyzes the oxidation of AFB₁ at different rates or sites or both, depending on the specific isozyme induced. Both polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs) are highly persistent pollutants which have been widely identified in

human and animal tissues and are potent inducers of cytochrome P-450-dependent monooxygenases (1, 19, 20, 26). Thus, the interactive effects of these environmental pollutants (PCBs and PBBs) and AFB₁ may be an important consideration in populations exposed to both types of toxins.

PCB and PBB congeners have recently been classed into three major categories (1, 2, 19, 22, 23). One category is typified by phenobarbital (PB)-induced, cytochrome P-450-dependent monooxygenase activities. Another category, typified by 3-methylcholanthrene (MC), induces cytochrome P-448-dependent monooxygenases. A third category represents a mixed pattern of induction such as is produced by coadministering PB and MC. In the present study, we report the effects of several PCB- and PBB-induced rat hepatic microsomal enzyme preparations on the metabolism of AFB₁ to the toxic metabolite AFM₁ and the less toxic metabolite AFQ₁.

MATERIALS AND METHODS

Chemicals. AFB₁, AFM₁, and AFQ₁ were obtained from Sigma Chemical Co., St. Louis, Mo. Tritium-labeled AFB₁ was purchased from Moravek Chemicals, Brea, Calif. A working solution of radiolabeled AFB₁ was prepared by mixing 2.0 mg of unlabeled AFB₁ with 100 μl of [³H]AFB₁ (specific activity, 15.0 Ci/mmol) and diluting with 0.9 ml of methanol. The solutions were stored at -80°C until ready for use. Sodium PB (Harvey Laboratories, Inc., Philadelphia, Pa.) was diluted in saline to a concentration of 30 mg/ml. Firemaster BP-6 and Aroclor 1254 were generous gifts from the Michigan Chemical Corp., St. Louis, Mich., and the Monsanto Chemical Co., St. Louis, Mo., respectively.

Individual PCB and PBB congeners were synthesized from the appropriate precursors and purified by column chromatography on alumina-Florisil and by silica gel thin-layer chromatography (1, 2). The purities were determined by gas-liquid chromatography with a chromatograph (model 5710; Hewlett-Packard Co., Avondale, Calif.) equipped with a ⁶³Ni electron-capture detector and a glass column (0.6 cm

* Corresponding author.

TABLE 1. Effects of microsomal enzyme inducers on cytochrome P-450-mediated metabolism of AFB₁^a

Animal pretreatment	AFB ₁ remaining ^b (nmol/mg of protein)	AFQ ₁ (nmol/ mg of protein)	AFM ₁ (nmol/ mg of protein)	4-(Dimethyl- amino)- antipyrine N- demethylase activity ^c	Benzo[<i>a</i>]- pyrene hydroxylase activity ^d
Corn oil (control)	2.66 ± 0.28	0.07 ± 0.02	0.15 ± 0.03	8.52 ± 0.90	0.43 ± 0.03
PB	2.18 ± 0.26 ^e	0.13 ± 0.02 ^e	0.15 ± 0.01	14.18 ± 0.59 ^e	0.47 ± 0.02
MC	0.65 ± 0.15 ^e	0.03 ± 0.00 ^e	1.03 ± 0.15 ^e	6.69 ± 0.38	5.53 ± 0.65 ^e
PB + MC	1.17 ± 0.22 ^e	0.07 ± 0.01	0.81 ± 0.25 ^e	11.58 ± 0.94 ^e	4.71 ± 0.33 ^e
Aroclor 1254	0.97 ± 0.12 ^e	0.04 ± 0.00 ^e	1.10 ± 0.20 ^e	12.68 ± 0.78 ^e	3.76 ± 0.21 ^e
Firemaster BP-6	1.42 ± 0.20 ^e	0.07 ± 0.02	0.85 ± 0.26 ^e	13.01 ± 0.73 ^e	3.94 ± 0.10 ^e

^a Three animals per group. Values are expressed as mean ± standard deviation.

^b 64 nmol of AFB₁ per incubation tube. AFB₁ remaining = AFB₁ extracted after a 30-min incubation.

^c Nanomoles of HCHO formed per milligram of protein per minute.

^d Nanomoles of benzo[*a*]pyrene metabolized per milligram of protein per minute.

^e Significantly different from control group mean (Student's *t* test, *P* < 0.05).

by 1.2 m) packed with 3% OV101 on Ultrabound Carbowax 20 M (80/100 mesh; RFR Corp., Hope, R.I.). Structures of purified PCB and PBB congeners were confirmed by their nuclear magnetic resonance and their mass spectra.

NADP, glucose 6-phosphate, glucose-6-phosphate dehydrogenase, magnesium chloride, and benzo[*a*]pyrene were obtained from Sigma. 4-(Dimethylamino)antipyrine was supplied by Aldrich Chemical Co., Inc., Milwaukee, Wis. [³H]benzo[*a*]pyrene was obtained from New England Nuclear Corp., Boston, Mass., and was purified by Florisil column chromatography.

Animal treatment and isolation of microsomes. One-month-old male Wistar rats (Timco Breeding Laboratories, Houston, Tex.), approximately 100 g in weight, were housed in stainless steel or plastic cages and allowed free access to feed (Purina Rat Chow no. 5002; Ralston Purina Co., St. Louis, Mo.) and water. Animals were maintained on a diurnal cycle of 12 h of light and 12 h of darkness. Animals were injected intraperitoneally with the compounds to be tested at a level of 150 μmol/kg of body weight on days 1 and 3 as previously described (19, 20). The compounds were administered in corn oil solution, and the animals were killed by cervical dislocation on day 6. PB (400 μmol/kg, dissolved in isotonic saline) and MC (100 μmol/kg, dissolved in corn oil) were administered individually as well as coadministered to the animals on days 1 and 2, and the animals were killed on day 3. These animals served as positive controls. Another control group was injected with corn oil (0.5 ml) on days 1 and 3 and then killed by cervical dislocation on day 6. To lower liver glycogen levels, all animals were fasted for 24 h before being killed.

Microsomes were prepared as described by Parkinson and Safe (21). The rat livers were perfused with ice-cold isotonic saline via the hepatic portal vein. The livers were removed and placed in preweighed beakers of ice-cold 0.25 M sucrose-0.1 mM EDTA solution for the determination of liver weights. The livers were then homogenized, and the homogenate was centrifuged at 10,000 × *g* for 20 min. The microsomal fraction was collected as a 100,000 × *g* pellet by further centrifugation (1 h) of the 10,000 × *g* supernatant fraction. The protein content of the microsomal fraction was determined by the method of Lowry et al. (12).

Assays. The metabolism of AFB₁ by isolated microsomes was carried out under conditions similar to those used by Roebuck and co-workers (24), using 12.5 mg of microsomal protein per tube. Negative controls were prepared by heating selected control tubes in boiling water for 10 min. To each tube, 10 μl of a 2-mg/ml solution of AFB₁ in methanol

was added, and the tubes were transferred to a water bath and incubated with gentle shaking for 30 min at 37°C (16). The total volume in each tube was 5 ml. The reaction was terminated by adding 2.0 ml of ice-cold methanol to each tube.

The rate of oxidative N-demethylation of 4-(dimethylamino)antipyrine was measured by quantifying the production of formaldehyde (6). The formaldehyde, trapped as the semicarbazone, was developed in double-strength Nash reagent (14). The rate of benzo[*a*]pyrene hydroxylation was measured by the radiometric assay of DePierre et al. (5) as improved by Nesnow et al. (18), by quantifying the base-soluble metabolites after hexane extraction of the unreacted benzo[*a*]pyrene.

Aflatoxin analysis. Aflatoxins were extracted from *in vitro* incubations by the following procedure. Hexane (2 ml) was added to each test tube to remove residual lipids. The hexane and the aqueous phases were mixed well and allowed to separate, and the upper hexane layer was discarded. The aflatoxins were extracted from the reaction tubes with three successive 5-ml portions of chloroform. The chloroform extracts were combined and evaporated to dryness under nitrogen. The extracts were redissolved in 200 μl of the high-pressure liquid chromatography mobile phase (25).

The chloroform extracts were analyzed by normal-phase high-pressure liquid chromatography by a previously reported method (25). Aflatoxin peaks were detected by UV absorbance at 365 nm. Fluorescence detection (365-nm excitation; 425-nm emission) was used for confirmation.

Statistics. Statistical comparisons between control and treatment groups were made by using the Student *t* test. Differences from controls were considered significant at *P* < 0.05.

RESULTS

Table 1 summarizes the *in vitro* metabolism of AFB₁ by rat hepatic microsomes from animals pretreated with corn oil, PB, MC, PB plus MC (coadministered), and commercial PCB and PBB (Aroclor 1254 and Firemaster BP-6, respectively). Compared with microsomes from the corn oil controls, hepatic microsomes from all the xenobiotic agent-treated animals increased the amount of AFB₁ metabolized (as determined by the AFB₁ remaining). PB-induced microsomes increased the amount of AFQ₁ formed but did not alter the AFM₁ levels; in contrast, MC-induced microsomes increased the amount of AFM₁ formed but decreased AFQ₁ formation. Microsomes from rats treated with the mixed-type inducers (PB plus MC, Aroclor 1254, and Firemaster

TABLE 2. Effects of polyhalogenated biphenyl congeners on cytochrome P-450-mediated metabolism of AFB₁^a

Animal pretreatment	AFB ₁ remaining ^b (nmol/mg of protein)	AFQ ₁ (nmol/ mg of protein)	AFM ₁ (nmol/ mg of protein)	4-(Dimethylamino)anti- pyrine <i>N</i> - demethylase activity ^c	Benzo[<i>a</i>]pyrene hydroxylase activity ^d
Corn oil (control)	2.66 ± 0.28	0.07 ± 0.02	0.15 ± 0.03	8.52 ± 0.90	0.43 ± 0.03
2,2',4,4'-Tetrachlorobiphenyl	2.31 ± 0.59	0.14 ± 0.05 ^e	0.24 ± 0.06 ^e	10.86 ± 0.88 ^e	0.44 ± 0.02
2,2',4,5,5'-Pentabromobiphenyl	2.54 ± 0.42	0.11 ± 0.02 ^e	0.21 ± 0.15	11.09 ± 0.82 ^e	0.47 ± 0.02
3,3',4,4'-Tetrachlorobiphenyl	0.08 ± 0.39 ^e	0.04 ± 0.01 ^e	1.74 ± 0.45 ^e	5.83 ± 0.63 ^e	5.03 ± 0.44 ^e
3,3',4,4'-Tetrabromobiphenyl	0.79 ± 0.29 ^e	0.05 ± 0.02	1.96 ± 0.44 ^e	3.67 ± 0.48 ^e	4.40 ± 0.16 ^e
2,3,4,4',5-Pentachlorobiphenyl	0.69 ± 0.12 ^e	0.03 ± 0.00 ^e	1.30 ± 0.17 ^e	8.30 ± 0.35	5.11 ± 0.18 ^e
2,3,4,4',5-Pentabromobiphenyl	0.76 ± 0.38 ^e	0.04 ± 0.00 ^e	1.77 ± 0.31 ^e	7.76 ± 0.07	4.82 ± 0.16 ^e

^a Three animals per group. Values are expressed as mean ± standard deviation.

^b 64 nmol of AFB₁ per incubation tube. AFB₁ remaining = AFB₁ extracted after a 30-min incubation.

^c Nanomoles of HCHO formed per milligram of protein per minute.

^d Nanomoles of benzo[*a*]pyrene metabolized per milligram of protein per minute.

^e Significantly different from control group mean (Student's *t* test, *P* < 0.05).

BP-6) all increased the amount of AFM₁ formed but did not affect (or slightly decreased) the levels of AFQ₁ produced.

Table 2 and Fig. 1 summarize the metabolism of AFB₁ by hepatic microsomes from rats pretreated with several PCB and PBB congeners. 2,2',4,4'-Tetrachlorobiphenyl and 2,2',4,5,5'-pentabromobiphenyl resemble PB in their mode of induction of cytochrome P-450-dependent monooxygenases. The microsomes from rats pretreated with these two compounds did not significantly increase the amount of AFB₁ metabolized, but they markedly increased the production of AFQ₁. 3,3',4,4'-Tetrachloro- and tetrabromobiphenyl resemble MC in their mode of induction of cytochrome P-450-dependent monooxygenases. Microsomes from 3,3',4,4'-tetrachloro- and tetrabromobiphenyl-pretreated rats enhanced AFB₁ metabolism, slightly reduced the levels of

AFQ₁ produced, and increased the amount of AFB₁ converted to AFM₁. Microsomes from rats pretreated with 2,3,4,4',5-pentachloro- and pentabromobiphenyl gave results similar to those of microsomes from rats pretreated with 3,3',4,4'-tetrachloro- and tetrabromobiphenyl. 2,3,4,4',5-Pentachloro- and pentabromobiphenyl are classified as mixed-type inducers.

Tables 1 and 2 also compare the in vitro metabolism of AFB₁ to AFM₁ and AFQ₁ with that of two other cytochrome P-450-dependent monooxygenases, 4-(dimethylamino)antipyrine *N*-demethylase and benzo[*a*]pyrene hydroxylase. The *N*-demethylase is a PB-inducible enzyme, whereas benzo[*a*]pyrene hydroxylase is primarily induced by MC. Microsomes from rats pretreated with PB and PB-type inducers increased the levels of hepatic microsomal 4-




FIG. 1. Effect of various microsomal enzyme inducers on the in vitro metabolism of AFB₁ to AFM₁ and AFQ₁. The inducers are classified as PB and PB type (2,2',4,4'-PCB and 2,2',4,5,5'-PBB), MC and MC type (3,3',4,4'-PCB and -PBB), and mixed type (PB plus MC, Firemaster BP-6, Aroclor 1254, and 2,3,4,4',5-PCB and -PBB). Results are reported as nanomoles produced per milligram of protein. A total of 64 nmol of AFB₁ was added per incubation tube. Results are the mean ± the standard deviation (bars) of independent analysis of microsomes from different animals, each assayed in triplicate.

(dimethylamino)antipyrene *N*-demethylase. Microsomes from rats pretreated with MC or MC-type inducers showed increased levels of benzo[*a*]pyrene hydroxylase. Mixed-type inducers increased the levels of both enzymes. These results are consistent with those of earlier studies (1, 19, 23, 26).

DISCUSSION

Previous reports have shown that microsomes from animals pretreated with MC exhibit a 10-fold increase in the formation of AFM₁ from AFB₁ without affecting the levels of AFQ₁ formed (13, 15). Not surprisingly, the AFB₁-4-hydroxylase activities of microsomes from 3,3',4,4'-tetrachlorobiphenyl- and 3,3',4,4'-tetrabromobiphenyl-pretreated rats and MC-pretreated rats were similar. All three xenobiotic agents induce a comparable pattern of cytochrome P-450 isozymes (22). Both 2,2',4,4'-tetrachlorobiphenyl and 2,2',4,5,5'-pentabromobiphenyl resemble PB in their mode of induction of rat hepatic cytochrome P-450 isozymes and cytochrome P-450-dependent monooxygenases (22). This study confirms that PB and the two PB-type inducers preferentially enhance the conversion of AFB₁ to AFQ₁ (8, 9, 13).

The effects of coadministered PB and MC as monooxygenase enzyme inducers are additive, and similar activities have been observed after the administration of Aroclor 1254, Firemaster BP-6, 2,3,4,4',5-pentachlorobiphenyl, 2,3,4,4',5-pentabromobiphenyl, and several other halogenated biphenyls (1, 19, 22, 23). This mixed-type induction activity is characterized by the induction of the microsomal monooxygenase enzymes and the cytochrome P-450 isozymes induced by both PB and MC. Thus, the mixed-type inducers used in this study induced both 4-(dimethylamino)antipyrene *N*-demethylase and benzo[*a*]pyrene hydroxylase. However, the additivity expected for the formation of the metabolites AFM₁ and AFQ₁ was not observed. Microsomes from animals treated with all the mixed-type inducers significantly increased (5 to 12-fold) the amount of AFB₁ converted into AFM₁ but they did not increase the formation of AFQ₁. Moreover, microsomes from rats pretreated with Aroclor 1254, 2,3,4,4',5-pentachlorobiphenyl, and 2,3,4,4',5-pentabromobiphenyl exhibited significantly lower levels of AFQ₁ formed after incubation with AFB₁. Although mixed-type and MC-type inducers elicit markedly different microsomal monooxygenase enzyme induction responses for most substrates, their effects on AFB₁ conversion into AFM₁ and AFQ₁ were similar.

Our results show that approximately 40% of the AFB₁ metabolized by microsomes from rats pretreated with MC and the MC-type inducers 3,3',4,4'-tetrachloro- and tetrabromobiphenyl was accounted for by the metabolites AFQ₁ and AFM₁. Approximately 30% of the AFB₁ metabolized by microsomes from rats pretreated with PB plus MC, Aroclor 1254, Firemaster BP-6, and 2,3,4,4',5-pentachloro- and pentabromobiphenyl was accounted for by AFQ₁ and AFM₁. With microsomes from rats pretreated with PB, 2,2',4,4'-tetrachlorobiphenyl, and 2,2',4,5,5'-pentabromobiphenyl, only about 13% of the metabolized AFB₁ was accounted for by AFQ₁ and AFM₁. The remainder of the AFB₁ metabolized is probably involved in binding to microsomal macromolecules and in the formation of minor metabolites (13, 15, 17).

The results of this study show that rat microsomes induced with mixed-type inducers significantly increased the conversion of AFB₁ to AFM₁ and caused little change in AFQ₁ formation. In a recent study (E. C. Shepherd, Ph.D. dissertation, Texas A&M University, College Station, 1982),

AFB₁ was administered to lactating rats pretreated with halogenated biphenyls. The results showed that AFM₁ is the major metabolite excreted in the milk. The suppression of AFQ₁ formation may also be related to the *in vivo* mixed-type activity of the induced microsomes. The rationale for these unique observations is not apparent, and current research in our laboratories is focused on understanding the interactive effects of PCBs, PBBs, and aflatoxins.

ACKNOWLEDGMENTS

This investigation was supported by TAES H6215, Title XII AID Project 02-50305-2, Department of Defense grant DAAG29-83-G-0088, Public Health Service grant R01 ES02798 from the National Institutes of Health, Center for Comparative Medicine grant 18820-4.

LITERATURE CITED

1. Andres, J., I. Lambert, L. Robertson, S. Bandiera, T. Sawyer, S. Lovering, and S. Safe. 1983. The comparative biologic and toxic potencies of polychlorinated biphenyls and polybrominated biphenyls. *Toxicol. Appl. Pharmacol.* **70**:204-215.
2. Bandiera, S., S. Safe, and A. B. Okey. 1982. Binding of polychlorinated biphenyls classified as either phenobarbitone-, 3-methylcholanthrene-, or mixed-type inducers to cytosolic Ah receptor. *Chem.-Biol. Interact.* **39**:259-277.
3. Croy, R. G., J. M. Essigman, V. N. Reinhold, and G. N. Wogan. 1978. Identification of the principal aflatoxin B₁-DNA adduct formed *in vivo* in rat liver. *Proc. Natl. Acad. Sci. U.S.A.* **75**:1745-1749.
4. Dahms, R., and H. L. Gurtoo. 1976. Metabolism of aflatoxin B₁ to aflatoxins Q₁, M₁ and P₁ by mouse and rat. *Res. Commun. Chem. Pathol. Pharmacol.* **15**:11-20.
5. DePierre, J. W., M. S. Moron, K. A. M. Johnnesen, and L. Evaster. 1975. A reliable, sensitive and convenient radioactive assay for benzo[*a*]pyrene monooxygenase. *Anal. Biochem.* **63**:470-484.
6. DeWaide, J. H., and P. T. Henderson. 1968. Hepatic *N*-demethylation of aminopyrine in rat and trout. *Biochem. Pharmacol.* **17**:1901-1907.
7. Garner, R. C., E. C. Miller, and J. A. Miller. 1972. Liver microsomal metabolism of aflatoxin B₁ to a reactive derivative toxic to *Salmonella typhimurium* TA 1530. *Cancer Res.* **32**:2058-2066.
8. Gurtoo, H. L. 1980. Genetic expression of aflatoxin B₁ metabolism: effects of 3-methylcholanthrene and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin on the metabolism of aflatoxins B₁ and B₂ by various inbred strains of mice. *Mol. Pharmacol.* **18**:296-303.
9. Gurtoo, H. L., and R. P. Dahms. 1979. Effects of inducers and inhibitors on the metabolism of aflatoxin B₁ by rat and mouse. *Biochem. Pharmacol.* **28**:3441-3449.
10. Heathcote, J. G., and J. R. Hibbert. 1978. Aflatoxins: chemical and biological aspects, p. 131-150. Elsevier Scientific Publishing Co., Amsterdam.
11. Hsieh, D. P. H., Z. A. Wong, J. J. Wong, C. Michas, and B. H. Ruebner. 1977. Comparative metabolism of aflatoxin, p. 37-50. *In* J. V. Rodricks, C. M. Hesseltine, and M. A. Mehlman (ed.), *Mycotoxins in human and animal health*. Pathotox Publishing, Park Forest South, Ill.
12. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**:265-275.
13. Metcalf, S. A., P. J. Colley, and G. E. Neal. 1981. A comparison of the effects of pretreatment with phenobarbitone and 3-methylcholanthrene on the metabolism of aflatoxin B₁ by rat liver microsomes and isolated hepatocytes *in vitro*. *Chem.-Biol. Interact.* **35**:145-157.
14. Nash, T. 1958. The colorimetric estimation of formaldehyde by means of the Hantzsch reaction. *Biochem. J.* **55**:416-421.
15. Neal, G. E., and P. J. Colley. 1978. Some high-performance liquid chromatographic studies of the metabolism of aflatoxins by rat liver microsomal preparations. *Biochem. J.* **174**:839-851.

16. Neal, G. E., and P. J. Colley. 1979. The formation of 2,3-dihydroxy-2,3-dihydro aflatoxin B₁ by the metabolism of aflatoxin B₁ *in vitro* by rat liver microsomes. *FEBS Lett.* **101**:382-386.
17. Neal, G. E., D. J. Judah, F. Stirpe, and D. S. P. Patterson. 1981. The formation of 2,3-dihydroxy-2,3-dihydro-aflatoxin B₁ by the metabolism of aflatoxin B₁ by liver microsomes isolated from certain avian and mammalian species and the possible role of the metabolite in the acute toxicity of aflatoxin B₁. *Toxicol. Appl. Pharmacol.* **58**:431-437.
18. Nesnow, S., W. E. Fahl, and C. R. Jefcoate. 1977. An improved radiochemical assay for benzo[a]pyrene monooxygenase. *Anal. Biochem.* **80**:258-266.
19. Parkinson, A., R. Cockerine, and S. Safe. 1980. Polychlorinated biphenyl isomers and congeners as inducers of both 3-methylcholanthrene- and pheno-barbitone-type microsomal enzyme activity. *Chem.-Biol. Interact.* **29**:277-289.
20. Parkinson, A., L. Robertson, L. Safe, and S. Safe. 1980. Polychlorinated biphenyls as inducers of hepatic microsomal enzymes: structure-activity rules. *Chem.-Biol. Interact.* **30**:271-285.
21. Parkinson, A., and S. Safe. 1979. The detection of enzyme induction by rat liver microsome prepared by isoelectric precipitation. *J. Pharm. Pharmacol.* **31**:444-447.
22. Parkinson, A., S. H. Safe, L. W. Robertson, P. E. Thomas, O. E. Ryan, L. M. Reik, and W. Levin. 1983. Immunochemical quantitation of cytochrome P-450 isozymes and epoxide hydrolase in liver microsomes from polychlorinated and polybrominated biphenyl-treated rats: a study of structure-activity relationships. *J. Biol. Chem.* **258**:5967-5976.
23. Robertson, L. W., A. Parkinson, and S. Safe. 1980. Induction of both cytochromes P-450 and P-448 by 2,3',4,4',5-pentabromobiphenyl, a component of Firemaster. *Biochem. Biophys. Res. Commun.* **92**:175-182.
24. Roebuck, B. C., W. G. Siegel, and G. N. Wogan. 1978. *In vitro* metabolism of aflatoxin B₂ by animal and human liver. *Cancer Res.* **38**:999-1002.
25. Shepherd, E. C., T. D. Phillips, N. D. Heidelbaugh, and A. W. Hayes. 1982. High pressure liquid chromatographic determination of aflatoxins by using radial compression separation. *J. Assoc. Off. Anal. Chem.* **65**:665-671.
26. Snyder, R., and H. Remmer. 1979. Classes of hepatic microsomal mixed function oxidase inducers. *Pharmacol. Ther.* **7**:203-244.
27. Swenson, D. H., J. A. Miller, and E. C. Miller. 1975. The reactivity and carcinogenicity of aflatoxin B₁-2,3-dichloride, a model for the putative 2,3-oxide metabolite of aflatoxin B₁. *Cancer Res.* **35**:3811-3823.