

Exclusion of an Exotic Top Quark with $-4/3$ Electric Charge Using Soft Lepton Tagging

T. Aaltonen,²⁴ J. Adelman,¹⁴ B. Álvarez González^{w,12} S. Amerio^{ee,44} D. Amidei,³⁵ A. Anastassov,³⁹ A. Annovi,²⁰ J. Antos,¹⁵ G. Apollinari,¹⁸ J. Appel,¹⁸ A. Apresyan,⁴⁹ T. Arisawa,⁵⁸ A. Artikov,¹⁶ J. Asaadi,⁵⁴ W. Ashmanskas,¹⁸ A. Attal,⁴ A. Aurisano,⁵⁴ F. Azfar,⁴³ W. Badgett,¹⁸ A. Barbaro-Galtieri,²⁹ V.E. Barnes,⁴⁹ B.A. Barnett,²⁶ P. Barria^{gg,47} P. Bartos,¹⁵ G. Bauer,³³ P.-H. Beauchemin,³⁴ F. Bedeschi,⁴⁷ D. Beecher,³¹ S. Behari,²⁶ G. Bellettini^{ff,47} J. Bellinger,⁶⁰ D. Benjamin,¹⁷ A. Beretvas,¹⁸ A. Bhatti,⁵¹ M. Binkley*,¹⁸ D. Bisello^{ee,44} I. Bizjak^{kk,31} R.E. Blair,² C. Blocker,⁷ B. Blumenfeld,²⁶ A. Bocci,¹⁷ A. Bodek,⁵⁰ V. Boisvert,⁵⁰ D. Bortoletto,⁴⁹ J. Boudreau,⁴⁸ A. Boveia,¹¹ B. Brau^{a,11} A. Bridgeman,²⁵ L. Brigliadori^{dd,6} C. Bromberg,³⁶ E. Brubaker,¹⁴ J. Budagov,¹⁶ H.S. Budd,⁵⁰ S. Budd,²⁵ K. Burkett,¹⁸ G. Busetto^{ee,44} P. Bussey,²² A. Buzatu,³⁴ K. L. Byrum,² S. Cabrera^{y,17} C. Calancha,³² S. Camarda,⁴ M. Campanelli,³¹ M. Campbell,³⁵ F. Canelli^{14,18} A. Canepa,⁴⁶ B. Carls,²⁵ D. Carlsmith,⁶⁰ R. Carosi,⁴⁷ S. Carrillo^{n,19} S. Carron,¹⁸ B. Casal,¹² M. Casarsa,¹⁸ A. Castro^{dd,6} P. Catastini^{gg,47} D. Cauz,⁵⁵ V. Cavaliere^{gg,47} M. Cavalli-Sforza,⁴ A. Cerri,²⁹ L. Cerrito^{q,31} S.H. Chang,²⁸ Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁷ G. Chlachidze,¹⁸ F. Chlebana,¹⁸ K. Cho,²⁸ D. Chokheli,¹⁶ J.P. Chou,²³ K. Chung^{o,18} W.H. Chung,⁶⁰ Y.S. Chung,⁵⁰ T. Chwalek,²⁷ C.I. Ciobanu,⁴⁵ M.A. Ciocci^{gg,47} A. Clark,²¹ D. Clark,⁷ G. Compostella,⁴⁴ M.E. Convery,¹⁸ J. Conway,⁸ M. Corbo,⁴⁵ M. Cordelli,²⁰ C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{ff,47} C. Cuenca Almenar,⁶¹ J. Cuevas^{w,12} R. Culbertson,¹⁸ J.C. Cully,³⁵ D. Dagenhart,¹⁸ N. d'Ascenzo^{v,45} M. Datta,¹⁸ T. Davies,²² P. de Barbaro,⁵⁰ S. De Cecco,⁵² A. Deisher,²⁹ G. De Lorenzo,⁴ M. Dell'Orso^{ff,47} C. Deluca,⁴ L. Demortier,⁵¹ J. Deng^{f,17} M. Deninno,⁶ M. d'Errico^{ee,44} A. Di Canto^{ff,47} B. Di Ruzza,⁴⁷ J.R. Dittmann,⁵ M. D'Onofrio,⁴ S. Donati^{ff,47} P. Dong,¹⁸ T. Dorigo,⁴⁴ S. Dube,⁵³ K. Ebina,⁵⁸ A. Elagin,⁵⁴ R. Erbacher,⁸ D. Errede,²⁵ S. Errede,²⁵ N. Ershaidat^{cc,45} R. Eusebi,⁵⁴ H.C. Fang,²⁹ S. Farrington,⁴³ W.T. Fedorko,¹⁴ R.G. Feild,⁶¹ M. Feindt,²⁷ J.P. Fernandez,³² C. Ferrazza^{hh,47} R. Field,¹⁹ G. Flanagan^{s,49} R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²³ J.C. Freeman,¹⁸ I. Furic,¹⁹ M. Gallinaro,⁵¹ J. Galyardt,¹³ F. Garbersen,¹¹ J.E. Garcia,²¹ A.F. Garfinkel,⁴⁹ P. Garosi^{gg,47} H. Gerberich,²⁵ D. Gerdes,³⁵ A. Gessler,²⁷ S. Giagu^{ii,52} V. Giakoumopoulou,³ P. Giannetti,⁴⁷ K. Gibson,⁴⁸ J.L. Gimmell,⁵⁰ C.M. Ginsburg,¹⁸ N. Giokaris,³ M. Giordani^{jj,55} P. Giromini,²⁰ M. Giunta,⁴⁷ G. Giurgiu,²⁶ V. Glagolev,¹⁶ D. Glenzinski,¹⁸ M. Gold,³⁸ N. Goldschmidt,¹⁹ A. Golossanov,¹⁸ G. Gomez,¹² G. Gomez-Ceballos,³³ M. Goncharov,³³ O. González,³² I. Gorelov,³⁸ A.T. Goshaw,¹⁷ K. Goulianos,⁵¹ A. Gresele^{ee,44} S. Grinstein,⁴ C. Grosso-Pilcher,¹⁴ R.C. Group,¹⁸ U. Grundler,²⁵ J. Guimaraes da Costa,²³ Z. Gunay-Unalan,³⁶ C. Haber,²⁹ S.R. Hahn,¹⁸ E. Halkiadakis,⁵³ B.-Y. Han,⁵⁰ J.Y. Han,⁵⁰ F. Happacher,²⁰ K. Hara,⁵⁶ D. Hare,⁵³ M. Hare,⁵⁷ R.F. Harr,⁵⁹ M. Hartz,⁴⁸ K. Hatakeyama,⁵ C. Hays,⁴³ M. Heck,²⁷ J. Heinrich,⁴⁶ M. Herndon,⁶⁰ J. Heuser,²⁷ S. Hewamanage,⁵ D. Hidas,⁵³ C.S. Hill^{c,11} D. Hirschbuehl,²⁷ A. Hocker,¹⁸ S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,²⁹ R.E. Hughes,⁴⁰ M. Hurwitz,¹⁴ U. Husemann,⁶¹ M. Hussein,³⁶ J. Huston,³⁶ J. Incandela,¹¹ G. Introzzi,⁴⁷ M. Iori^{ii,52} A. Ivanov^{p,8} E. James,¹⁸ D. Jang,¹³ B. Jayatilaka,¹⁷ E.J. Jeon,²⁸ M.K. Jha,⁶ S. Jindariani,¹⁸ W. Johnson,⁸ M. Jones,⁴⁹ K.K. Joo,²⁸ S.Y. Jun,¹³ J.E. Jung,²⁸ T.R. Junk,¹⁸ T. Kamon,⁵⁴ D. Kar,¹⁹ P.E. Karchin,⁵⁹ Y. Kato^{m,42} R. Kephart,¹⁸ W. Ketchum,¹⁴ J. Keung,⁴⁶ B. Kietzman,¹⁸ V. Khotilovich,⁵⁴ B. Kilminster,¹⁸ D.H. Kim,²⁸ H.S. Kim,²⁸ H.W. Kim,²⁸ J.E. Kim,²⁸ M.J. Kim,²⁰ S.B. Kim,²⁸ S.H. Kim,⁵⁶ Y.K. Kim,¹⁴ N. Kimura,⁵⁸ L. Kirsch,⁷ S. Klimenko,¹⁹ K. Kondo,⁵⁸ D.J. Kong,²⁸ J. Konigsberg,¹⁹ A. Korytov,¹⁹ A.V. Kotwal,¹⁷ M. Kreps,²⁷ J. Kroll,⁴⁶ D. Krop,¹⁴ N. Krumnack,⁵ M. Kruse,¹⁷ V. Krutelyov,¹¹ T. Kuhr,²⁷ N.P. Kulkarni,⁵⁹ M. Kurata,⁵⁶ S. Kwang,¹⁴ A.T. Laasanen,⁴⁹ S. Lami,⁴⁷ S. Lammel,¹⁸ M. Lancaster,³¹ R.L. Lander,⁸ K. Lannon^{u,40} A. Lath,⁵³ G. Latino^{gg,47} I. Lazzizzera^{ee,44} T. LeCompte,² E. Lee,⁵⁴ H.S. Lee,¹⁴ J.S. Lee,²⁸ S.W. Lee^{x,54} S. Leone,⁴⁷ J.D. Lewis,¹⁸ C.-J. Lin,²⁹ J. Linacre,⁴³ M. Lindgren,¹⁸ E. Lipeles,⁴⁶ A. Lister,²¹ D.O. Litvintsev,¹⁸ C. Liu,⁴⁸ T. Liu,¹⁸ N.S. Lockyer,⁴⁶ A. Loginov,⁶¹ L. Lovas,¹⁵ D. Lucchesi^{ee,44} J. Lueck,²⁷ P. Lujan,²⁹ P. Lukens,¹⁸ G. Lungu,⁵¹ J. Lys,²⁹ R. Lysak,¹⁵ D. MacQueen,³⁴ R. Madrak,¹⁸ K. Maeshima,¹⁸ K. Makhoul,³³ P. Maksimovic,²⁶ S. Malde,⁴³ S. Malik,³¹ G. Manca^{e,30} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁹ C. Marino,²⁷ C.P. Marino,²⁵ A. Martin,⁶¹ V. Martin^{k,22} M. Martínez,⁴ R. Martínez-Ballarín,³² P. Mastrandrea,⁵² M. Mathis,²⁶ M.E. Mattson,⁵⁹ P. Mazzanti,⁶ K.S. McFarland,⁵⁰ P. McIntyre,⁵⁴ R. McNulty^{j,30} A. Mehta,³⁰ P. Mehtala,²⁴ A. Menzione,⁴⁷ C. Mesropian,⁵¹ T. Miao,¹⁸ D. Mietlicki,³⁵ N. Miladinovic,⁷ R. Miller,³⁶ C. Mills,²³ M. Milnik,²⁷ A. Mitra,¹ G. Mitselmakher,¹⁹ H. Miyake,⁵⁶ S. Moed,²³ N. Moggi,⁶ M.N. Mondragon^{n,18} C.S. Moon,²⁸ R. Moore,¹⁸ M.J. Morello,⁴⁷ J. Morlock,²⁷ P. Movilla Fernandez,¹⁸ J. Müllenstädt,²⁹ A. Mukherjee,¹⁸ Th. Muller,²⁷ P. Murat,¹⁸ M. Mussini^{dd,6} J. Nachtman^{o,18} Y. Nagai,⁵⁶ J. Naganoma,⁵⁶ K. Nakamura,⁵⁶ I. Nakano,⁴¹ A. Napier,⁵⁷

J. Nett,⁶⁰ C. Neu^{aa},⁴⁶ M.S. Neubauer,²⁵ S. Neubauer,²⁷ J. Nielsen^g,²⁹ L. Nodulman,² M. Norman,¹⁰
 O. Norriella,²⁵ E. Nurse,³¹ L. Oakes,⁴³ S.H. Oh,¹⁷ Y.D. Oh,²⁸ I. Oksuzian,¹⁹ T. Okusawa,⁴² R. Orava,²⁴
 K. Osterberg,²⁴ S. Pagan Griso^{ee},⁴⁴ C. Pagliarone,⁵⁵ E. Palencia,¹⁸ V. Papadimitriou,¹⁸ A. Papaikonomou,²⁷
 A.A. Paramanov,² B. Parks,⁴⁰ S. Pashapour,³⁴ J. Patrick,¹⁸ G. Pauletta^{jj},⁵⁵ M. Paulini,¹³ C. Paus,³³ T. Peiffer,²⁷
 D.E. Pellett,⁸ A. Penzo,⁵⁵ T.J. Phillips,¹⁷ G. Piacentino,⁴⁷ E. Pianori,⁴⁶ L. Pinera,¹⁹ K. Pitts,²⁵ C. Plager,⁹
 L. Pondrom,⁶⁰ K. Potamianos,⁴⁹ O. Poukhov*,¹⁶ F. Prokoshin^z,¹⁶ A. Pronko,¹⁸ F. Ptohosⁱ,¹⁸ E. Pueschel,¹³
 G. Punzi^{ff},⁴⁷ J. Pursley,⁶⁰ J. Rademacker^c,⁴³ A. Rahaman,⁴⁸ V. Ramakrishnan,⁶⁰ N. Ranjan,⁴⁹ I. Redondo,³²
 P. Renton,⁴³ M. Renz,²⁷ M. Rescigno,⁵² S. Richter,²⁷ F. Rimondi^{dd},⁶ L. Ristori,⁴⁷ A. Robson,²² T. Rodrigo,¹²
 T. Rodriguez,⁴⁶ E. Rogers,²⁵ S. Rolli,⁵⁷ R. Roser,¹⁸ M. Rossi,⁵⁵ R. Rossin,¹¹ P. Roy,³⁴ A. Ruiz,¹²
 J. Russ,¹³ V. Rusu,¹⁸ B. Rutherford,¹⁸ H. Saarikko,²⁴ A. Safonov,⁵⁴ W.K. Sakumoto,⁵⁰ L. Santi^{jj},⁵⁵
 L. Sartori,⁴⁷ K. Sato,⁵⁶ V. Saveliev^v,⁴⁵ A. Savoy-Navarro,⁴⁵ P. Schlabach,¹⁸ A. Schmidt,²⁷ E.E. Schmidt,¹⁸
 M.A. Schmidt,¹⁴ M.P. Schmidt*,⁶¹ M. Schmitt,³⁹ T. Schwarz,⁸ L. Scodellaro,¹² A. Scribano^{gg},⁴⁷ F. Scuri,⁴⁷
 A. Sedov,⁴⁹ S. Seidel,³⁸ Y. Seiya,⁴² A. Semenov,¹⁶ L. Sexton-Kennedy,¹⁸ F. Sforza^{ff},⁴⁷ A. Sfyrla,²⁵
 S.Z. Shalhout,⁵⁹ T. Shears,³⁰ P.F. Shepard,⁴⁸ M. Shimojima^t,⁵⁶ S. Shiraishi,¹⁴ M. Shochet,¹⁴ Y. Shon,⁶⁰
 I. Shreyber,³⁷ A. Simonenko,¹⁶ P. Sinervo,³⁴ A. Sisakyan,¹⁶ A.J. Slaughter,¹⁸ J. Slaunwhite,⁴⁰ K. Sliwa,⁵⁷
 J.R. Smith,⁸ F.D. Snider,¹⁸ R. Snihur,³⁴ A. Soha,¹⁸ S. Somalwar,⁵³ V. Sorin,⁴ P. Squillacioti^{gg},⁴⁷ M. Stanitzki,⁶¹
 R. St. Denis,²² B. Stelzer,³⁴ O. Stelzer-Chilton,³⁴ D. Stentz,³⁹ J. Strologas,³⁸ G.L. Strycker,³⁵ J.S. Suh,²⁸
 A. Sukhanov,¹⁹ I. Suslov,¹⁶ A. Taffard^f,²⁵ R. Takashima,⁴¹ Y. Takeuchi,⁵⁶ R. Tanaka,⁴¹ J. Tang,¹⁴ M. Tecchio,³⁵
 P.K. Teng,¹ J. Thom^h,¹⁸ J. Thome,¹³ G.A. Thompson,²⁵ E. Thomson,⁴⁶ P. Tipton,⁶¹ P. Ttito-Guzmán,³²
 S. Tkaczyk,¹⁸ D. Toback,⁵⁴ S. Tokar,¹⁵ K. Tollefson,³⁶ T. Tomura,⁵⁶ D. Tonelli,¹⁸ S. Torre,²⁰ D. Torretta,¹⁸
 P. Totaro^{jj},⁵⁵ M. Trovato^{hh},⁴⁷ S.-Y. Tsai,¹ Y. Tu,⁴⁶ N. Turini^{gg},⁴⁷ F. Ukegawa,⁵⁶ S. Uozumi,²⁸ N. van Remortel^b,²⁴
 A. Varganov,³⁵ E. Vataga^{hh},⁴⁷ F. Vázquezⁿ,¹⁹ G. Velev,¹⁸ C. Vellidis,³ M. Vidal,³² I. Vila,¹² R. Vilar,¹²
 M. Vogel,³⁸ I. Volobouev^x,²⁹ G. Volpi^{ff},⁴⁷ P. Wagner,⁴⁶ R.G. Wagner,² R.L. Wagner,¹⁸ W. Wagner^{bb},²⁷
 J. Wagner-Kuhr,²⁷ T. Wakisaka,⁴² R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³⁴ D. Waters,³¹ M. Weinberger,⁵⁴
 J. Weinelt,²⁷ W.C. Wester III,¹⁸ B. Whitehouse,⁵⁷ D. Whiteson^f,⁴⁶ A.B. Wicklund,² E. Wicklund,¹⁸ S. Wilbur,¹⁴
 G. Williams,³⁴ H.H. Williams,⁴⁶ P. Wilson,¹⁸ B.L. Winer,⁴⁰ P. Wittich^h,¹⁸ S. Wolbers,¹⁸ C. Wolfe,¹⁴ H. Wolfe,⁴⁰
 T. Wright,³⁵ X. Wu,²¹ F. Würthwein,¹⁰ A. Yagil,¹⁰ K. Yamamoto,⁴² J. Yamaoka,¹⁷ U.K. Yang^r,¹⁴
 Y.C. Yang,²⁸ W.M. Yao,²⁹ G.P. Yeh,¹⁸ K. Yi^o,¹⁸ J. Yoh,¹⁸ K. Yorita,⁵⁸ T. Yoshida^l,⁴² G.B. Yu,¹⁷
 I. Yu,²⁸ S.S. Yu,¹⁸ J.C. Yun,¹⁸ A. Zanetti,⁵⁵ Y. Zeng,¹⁷ X. Zhang,²⁵ Y. Zheng^d,⁹ and S. Zucchelli^{dd6}

(CDF Collaboration[†])

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439, USA*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798, USA*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^{dd}University of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254, USA*

⁸*University of California, Davis, Davis, California 95616, USA*

⁹*University of California, Los Angeles, Los Angeles, California 90024, USA*

¹⁰*University of California, San Diego, La Jolla, California 92093, USA*

¹¹*University of California, Santa Barbara, Santa Barbara, California 93106, USA*

¹²*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹³*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

¹⁴*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

¹⁵*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁶*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁷*Duke University, Durham, North Carolina 27708, USA*

¹⁸*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

¹⁹*University of Florida, Gainesville, Florida 32611, USA*

²⁰*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

²¹*University of Geneva, CH-1211 Geneva 4, Switzerland*

²²*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²³*Harvard University, Cambridge, Massachusetts 02138, USA*

²⁴*Division of High Energy Physics, Department of Physics,*

University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland

²⁵*University of Illinois, Urbana, Illinois 61801, USA*

- ²⁶The Johns Hopkins University, Baltimore, Maryland 21218, USA
- ²⁷Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany
- ²⁸Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea
- ²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA
- ³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³¹University College London, London WC1E 6BT, United Kingdom
- ³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- ³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
- ³⁴Institute of Particle Physics: McGill University, Montréal, Québec, Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6; University of Toronto, Toronto, Ontario, Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³⁵University of Michigan, Ann Arbor, Michigan 48109, USA
- ³⁶Michigan State University, East Lansing, Michigan 48824, USA
- ³⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁸University of New Mexico, Albuquerque, New Mexico 87131, USA
- ³⁹Northwestern University, Evanston, Illinois 60208, USA
- ⁴⁰The Ohio State University, Columbus, Ohio 43210, USA
- ⁴¹Okayama University, Okayama 700-8530, Japan
- ⁴²Osaka City University, Osaka 588, Japan
- ⁴³University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴⁴Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^{ee} University of Padova, I-35131 Padova, Italy
- ⁴⁵LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁴⁷Istituto Nazionale di Fisica Nucleare Pisa, ^{ff} University of Pisa,
- ⁴⁹University of Siena and ^{hh} Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
- ⁴⁹Purdue University, West Lafayette, Indiana 47907, USA
- ⁵⁰University of Rochester, Rochester, New York 14627, USA
- ⁵¹The Rockefeller University, New York, New York 10021, USA
- ⁵²Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, ⁱⁱ Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵³Rutgers University, Piscataway, New Jersey 08855, USA
- ⁵⁴Texas A&M University, College Station, Texas 77843, USA
- ⁵⁵Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-34100 Trieste, ^{jj} University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁶University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁷Tufts University, Medford, Massachusetts 02155, USA
- ⁵⁸Waseda University, Tokyo 169, Japan
- ⁵⁹Wayne State University, Detroit, Michigan 48201, USA
- ⁶⁰University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁶¹Yale University, New Haven, Connecticut 06520, USA

We present a measurement of the electric charge of the top quark using $p\bar{p}$ collisions corresponding to an integrated luminosity of 2.7 fb^{-1} at the CDF II detector. We reconstruct $t\bar{t}$ events in the lepton+jets final state and use kinematic information to determine which b -jet is associated with the leptonically- or hadronically-decaying t -quark. Soft lepton taggers are used to determine the b -jet flavor. Along with the charge of the W boson decay lepton, this information permits the reconstruction of the top quark's electric charge. Out of 45 reconstructed events with 2.4 ± 0.8 expected background events, 29 are reconstructed as $t\bar{t}$ with the standard model $+2/3$ charge, whereas 16 are reconstructed as $t\bar{t}$ with an exotic $-4/3$ charge. This is consistent with the standard model and excludes the exotic scenario at 95% confidence level. This is the strongest exclusion of the exotic charge scenario and the first to use soft leptons for this purpose.

PACS numbers: 12.90.+b, 14.65.Ha

*Deceased

†With visitors from ^aUniversity of Massachusetts Amherst,

Since the discovery of the top quark in 1995 [1], the CDF and D0 collaborations have scrutinized its properties. Measurements of the properties of the top quark all present a consistent picture of the top quark as the third-generation standard model (SM) weak-isospin partner of the bottom quark [2]. However a $+2/3$ -electric-charged top quark has yet to be experimentally confirmed, and an exotic $-4/3$ -charged scenario has been proposed [3]. In this theoretical scenario, the observed excess of events historically attributed to the top quark are instead attributed to an exotic particle, called “XM top quark,” which is identical to the SM top quark except that it decays to W^-b rather than to the SM W^+b . In order to preserve anomaly cancellation, the weak-isospin partner of the bottom (i.e. the “true” top quark) is assumed to exist but is too massive to be observed experimentally.

In this Letter, we present a measurement of the electric charge of the top quark. We analyze data corresponding to an integrated luminosity, $\int \mathcal{L} dt = 2.7 \text{ fb}^{-1}$, collected from February 2002 to April 2008.

We measure the top-quark charge by reconstructing $t\bar{t}$ pairs in the $\ell\bar{\nu}bq\bar{q}'b$ final state. The b -quarks associated with the leptonically (hadronically) decaying W are called the “leptonic” (“hadronic”) b -quarks. Reconstructing the top-quark charge involves identifying either the leptonic or hadronic b -quark and determining its flavor, either as b or \bar{b} . We use a soft electron tagger (SLT_e) [4] and a soft muon tagger (SLT_μ) [5] (collectively referred to as the SLT taggers) to identify the b -jets. The charges of the soft leptons are used to infer the flavor of

the b -jets. A secondary vertex tagger (SECVTX) [6] is also used to identify the b -jets and suppress SM backgrounds. A kinematic fitter [7] determines which b -jet is leptonic and which is hadronic. An event is considered SM if the lepton from the W and the SLT lepton from the leptonic (hadronic) b -jet have the opposite (same) charge sign. The event is considered to be XM otherwise. For the purposes of this measurement, we assume that the XM top quark has identical properties to the SM top quark, except for its electric charge.

This binary event reconstruction implies that if both the kinematic fitter and the SLT tagger are incorrect, then the correct top-quark charge is still reconstructed. From a Monte Carlo (MC) simulation of $t\bar{t}$ events, the fraction of b -jets for which the SLT taggers give the correct flavor assignment is approximately 69%. The fraction of events for which the kinematic fitter properly determines whether a b -jet is the leptonic or hadronic b is approximately 76%. This method reconstructs a SM (XM) charge in approximately 60% (40%) of simulated SM $t\bar{t}$ events.

This technique complements the measurement of the top-quark charge in Ref. [8] which uses the curvature and momentum of tracks within a b -jet cone to determine its charge. The SLT method is much less efficient than this technique since the semileptonic branching fraction for b -jets is only $\sim 10\%$ per lepton flavor; however, the b -jet flavor determination is much more reliable because of the higher b -jet flavor reconstruction purity. The overall reduction in sensitivity with the SLT technique is therefore only a factor of 2 – 3 lower.

The CDF II detector [9] is an azimuthally and forward-backward symmetric general-purpose detector [10] with silicon tracking [11] and drift chamber tracking [12] immersed in a 1.4 T solenoidal magnetic field. Projective electromagnetic- and hadronic-sampling calorimeters [13] lie beyond the solenoid and provide jet and missing E_T (\cancel{E}_T) reconstruction. Muon chambers [14] lie beyond the calorimeter and provide coverage up to $|\eta| \leq 1.0$. Gaseous Cherenkov counters [15] measure the average number of inelastic $p\bar{p}$ collisions to determine the luminosity with a 6% relative uncertainty.

Events are identified with central ($|\eta| \lesssim 1$), high- p_T ($-E_T$) muon (electron) triggers. We select events with a $p_T > 20 \text{ GeV}/c$ ($E_T > 20 \text{ GeV}$) muon (electron), which we call the “primary” lepton. At least four jets [16] with corrected $E_T > 20 \text{ GeV}$ [17] and $|\eta| \leq 2.0$ must be present in the event. To increase our acceptance for $t\bar{t}$ events, we allow one of the four jets to pass a looser selection ($E_T > 12 \text{ GeV}$ and $|\eta| \leq 2.4$), but we do not consider the looser fourth jet for tagging, either by the SLT or SECVTX algorithms. We explicitly reject cosmic muons, electrons from photon conversions, leptons from Z boson decay, and events with more than one energetic and isolated lepton. We also require $H_T > 250 \text{ GeV}$ and $\cancel{E}_T > 30 \text{ GeV}$, where H_T is the scalar sum of the

Amherst, Massachusetts 01003, ^bUniversiteit Antwerpen, B-2610 Antwerp, Belgium, ^cUniversity of Bristol, Bristol BS8 1TL, United Kingdom, ^dChinese Academy of Sciences, Beijing 100864, China, ^eIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^fUniversity of California Irvine, Irvine, CA 92697, ^gUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^hCornell University, Ithaca, NY 14853, ⁱUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^jUniversity College Dublin, Dublin 4, Ireland, ^kUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^lUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017, ^mKinki University, Higashi-Osaka City, Japan 577-8502, ⁿUniversidad Iberoamericana, Mexico D.F., Mexico, ^oUniversity of Iowa, Iowa City, IA 52242, ^pKansas State University, Manhattan, KS 66506, ^qQueen Mary, University of London, London, E1 4NS, England, ^rUniversity of Manchester, Manchester M13 9PL, England, ^sMuons, Inc., Batavia, IL 60510, ^tNagasaki Institute of Applied Science, Nagasaki, Japan, ^uUniversity of Notre Dame, Notre Dame, IN 46556, ^vObninsk State University, Obninsk, Russia, ^wUniversity de Oviedo, E-33007 Oviedo, Spain, ^xTexas Tech University, Lubbock, TX 79609, ^yIFIC(CSIC-Universitat de Valencia), 56071 Valencia, Spain, ^zUniversidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile, ^{aa}University of Virginia, Charlottesville, VA 22906, ^{bb}Bergische Universität Wuppertal, 42097 Wuppertal, Germany, ^{cc}Yarmouk University, Irbid 211-63, Jordan, ^{kk}On leave from J. Stefan Institute, Ljubljana, Slovenia,

transverse energy of the primary lepton, \cancel{E}_T , and jets.

We require each event to have ≥ 1 SLT (either e or μ) tag, and ≥ 1 SECVTX tag. We do not require that different jets in the same event are tagged by the different taggers. To suppress cascade decays of b -jets (i.e. $b \rightarrow c \rightarrow \ell\nu X$) that result in flavor mis-identification, we require the SLT track $p_T > 6$ GeV/ c , since leptons from cascade decays tend to be softer than those from direct semileptonic decays. We further require $p_T^{\text{rel}} > 1.5$ GeV/ c where p_T^{rel} is the SLT $_{\mu}$ track p_T relative to the jet axis.

We use a kinematic fitter described in detail in Ref. [7] which minimizes a reduced χ^2 -like function to fit to the $t\bar{t}$ event hypothesis. The experimental resolution of the final state particles is accounted for, and the particles are kinematically constrained to the W mass and top-quark mass (assumed to be 175 GeV/ c^2), within the theoretical decay widths. Jets are assigned uniquely to each of the four final-state quarks, and those jets tagged by either the SLT or SECVTX algorithms are constrained to be either of the two b -jets. All possible permutations are considered and the one which results in the lowest χ^2 value is chosen. If two different jets are both tagged, then we require that the lowest $\chi^2 < 27$; however if only one jet in the event is tagged, by both SECVTX and the SLT, then we require $\chi^2 < 9$. The tighter requirement on the χ^2 enforces a higher top-quark charge reconstruction purity since there is a greater ambiguity when only one jet is identified as a b -jet by the taggers.

The requirement on the χ^2 , SLT track p_T , and SLT $_{\mu}$ p_T^{rel} variables is determined by optimizing on total expected ϵD^2 , where ϵ is the event-reconstruction efficiency, $D = 2P - 1$ is the dilution, where P is the purity, which is defined as the fraction of reconstructed events that are determined to have an SM charge. Table I presents the expected ϵD^2 using the PYTHIA MC generator [18] to model $t\bar{t}$ and assuming $\sigma_{t\bar{t}} = 6.7 \pm 0.8$ pb [19], $M_t = 175$ GeV/ c^2 , and $\int \mathcal{L} dt = 2.7$ fb $^{-1}$. We choose the pretag expectation as the denominator of the efficiency, although for the optimization that choice is arbitrary. The figure of merit, ϵD^2 , is shown for the combined result, as well as separately for events with one or two tagged jets and with the SLT $_e$ or SLT $_{\mu}$ only. We expect 30.0 ± 5.9 events from $t\bar{t}$ in the tag sample, where the uncertainty is dominated by the theoretical cross section uncertainty and the jet energy scale uncertainty.

Due to the requirement of at least two b -tags, the contribution from non- $t\bar{t}$ backgrounds to the data is very small. Backgrounds from W +jets, WW , WZ , ZZ , single top, Z +jets, Drell-Yan+jets, and multijet production are all considered. The dominant background is due to $W+b\bar{b}$ production, and the total expected contribution to the tag sample from all backgrounds is 2.4 ± 0.8 events, where the uncertainty is dominated by the uncertainty on the jet energy scale and the multijet background estimate. The background estimate uses the same technique as in Ref. [4].

	ϵ (%)	P (%)	ϵD^2 (%)	$\langle N_{SM} \rangle$	$\langle N_{XM} \rangle$
Total	3.26	60.8	0.152	18.3	11.8
1 tagged jet	0.92	58.2	0.025	4.9	3.5
≥ 2 tagged jets	2.34	61.8	0.130	13.4	8.3
SLT $_e$ only	1.62	61.9	0.092	9.2	5.7
SLT $_{\mu}$ only	1.69	59.4	0.060	9.3	6.3

TABLE I: Expected efficiency (ϵ), purity (P), ϵD^2 , and number of events reconstructed as SM and XM, assuming $\sigma_{t\bar{t}} = 6.7$ pb for $\int \mathcal{L} dt = 2.7$ fb $^{-1}$ in PYTHIA simulation.

The measurement of the b -jet flavor determination purity is estimated with simulation but is calibrated by comparing a sample of pure $b\bar{b}$ events in both data and MC. The sample is constructed from events with a $p_T > 8$ GeV/ c ($E_T > 8$ GeV) muon (electron) close to a jet. A recoiling jet must also be found in the event in which an SLT tag is present, and both jets must be tagged by SECVTX. We measure a dilution scale factor, $SF_D = \sqrt{D_{\text{data}}/D_{\text{MC}}}$ where $D = (\epsilon_{\text{OS}} - \epsilon_{\text{SS}})/(\epsilon_{\text{OS}} + \epsilon_{\text{SS}})$, and $\epsilon_{\text{OS/SS}}$ is the tagging efficiency when the trigger lepton and the SLT have the opposite sign (OS) or same sign (SS) charge. The square root originates from the assumption that since both b -jets decay semileptonically they are subject to the same dilution factor. This scale factor accounts for differences between the data and MC such as mis-estimated branching ratios or mis-modeling of neutral B mixing.

We measure SF_D to be 0.92 ± 0.11 . The uncertainty is dominated by the statistical uncertainty and covers dependencies on other variables, such as the jet E_T . We use this to correct the simulation estimate for the $t\bar{t}$ charge reconstruction purity, for which our final estimate is $(60 \pm 3)\%$. The uncertainty in the purity is dominated by uncertainties in the simulation of the QCD radiation both in the initial (ISR) and final-state (FSR), the uncertainty in SF_D , and that arising when an alternate MC generator, HERWIG [20], is used. Simulation confirms the naive expectation that the background is reconstructed symmetrically between SM and XM events. Although simulation may mis-model the background reconstruction, the total background contribution is small, so we apply a very conservative systematic uncertainty, corresponding to twice the uncertainty on the signal purity estimate. Therefore, the background has a ‘‘purity’’ of $(50 \pm 6)\%$.

We observe 45 tagged and reconstructed events in data, of which 29 are reconstructed as SM and 16 are reconstructed as XM, a ratio consistent with the SM hypothesis. Three events have two SLT tags, although only one of these events has both SLT tags close to jets identified as b -jets by the kinematic fit (in this case, both SLT tags are consistent with the SM). Table II shows the number

Subsample	N	N_{SM}	N_{XM}
Primary Electron	25	16	9
Primary Muon	20	13	7
1 Tagged Jet	7	4	3
≥ 2 Tagged Jets	38	25	13
SLT _e	25	15	10
SLT _{μ}	21	15	6
All	45	29	16

TABLE II: Tag configurations in various subsamples of the data, including divisions according to the primary lepton flavor, the number of tagged b -jets, and the SLT flavor. Shown are the number of SM and XM tags as well as the total.

of tags by subsample, including the flavor of the primary lepton, the number of tagged b -jets, and the SLT flavor. Note that there is no significantly different SM/XM admixture in any of the subsamples.

The statistical significance of the measurement is given by the p -value for the test statistic

$$A \equiv \frac{1}{D_S} \frac{N_{SM} - N_{XM} - \langle B \rangle D_B}{N_{SM} + N_{XM} - \langle B \rangle} \quad (1)$$

where N_{SM} (N_{XM}) is the number of SM (XM) events, D_S and D_B are the signal and background dilution, respectively, and $\langle B \rangle$ is the total background expectation. This asymmetry, A , has been normalized so that the median expectation of the SM (XM) hypothesis is $+1.0$ (-1.0). In the data, we measure a normalized asymmetry $A_0 = 1.53 \pm 0.75$ (stat), which clearly favors a SM hypothesis.

We use MC pseudo experiments to determine the p -value. This is done by drawing from Poisson and binomial distributions to model the expected number of events and purities, respectively. Uncertainties are treated as Gaussian distributions. We measure $p_{SM} = p(A \leq A_0 | SM) = 0.69$ and $p_{XM} = p(A \geq A_0 | XM) = 0.0094$ for the SM and XM hypotheses, respectively, while we expect $p_{SM} = 0.50$ and $p_{XM} = 0.028$, assuming the SM. Figure 1 shows the distribution of p -values under the SM and XM hypotheses from pseudo-experiments. We choose the type-I error rate, α , *a priori* by using the standard threshold for exclusion of exotica: $\alpha = 0.05$. From this we exclude the exotic $-4/3$ -charged top quark at 95% confidence level. Table III shows the expected and measured XM p -value with the significant systematic errors added cumulatively.

We can also quantify the result of this measurement with a Bayes Factor (BF), which can be interpreted as the posterior odds in favor of the SM when the prior odds are neutral (equal to unity). This quantity is equal to the ratio, $p(A = A_0 | SM) / p(A = A_0 | XM)$. We evaluate a BF for this measurement to be 85.8 which is considered

Source	Expected p -value	Observed p -value
Stat. only	0.020	0.0054
Dilution Scale Factor	0.021	0.0058
ISR/FSR	0.022	0.0062
Cross Sections	0.023	0.0069
Jet Energy Scale	0.026	0.0080
MC Generator	0.028	0.0094

TABLE III: The p -values obtained for the XM hypothesis and how they are affected by the cumulative addition of systematic uncertainties. Other sources of systematic uncertainties are negligible.




FIG. 1: The SM and XM p -values for the normalized asymmetry test statistic, A , from pseudo experiments shown with all uncertainties combined and statistical uncertainties only.

“strong” evidence [21] for a $+2/3$ -charged top quark.

Figure 2 shows the distribution of the event H_T and the SLT tag p_T . Both the sum and difference of the events classified as SM and XM are shown. The total $t\bar{t}$ contribution (SM+XM) from simulation is normalized to the data and divided between SLT contributions from direct semileptonic b decay, cascade semileptonic decay, and other sources. The expected distribution assuming a $-4/3$ charge XM top quark is shown as a dashed line in the SM-XM plots. These figures demonstrate the preference of the asymmetry for the SM expectation as a function of the event kinematics.

In conclusion, we have presented the strongest exclusion of an exotic top quark with $-4/3$ charge to date (at 95% C.L.), while observing strong evidence for the SM $+2/3$ electric charge of the top quark. This measurement improves on both the expected and measured p -values reported in Ref. [8]. For purposes of comparison, we note that what is labeled as “expected C.L.” in Ref. [8] corresponds to one minus the expectation value of our p_{XM} under the SM hypothesis. This is the first time soft leptons tags have been used to accomplish such a measurement.

We thank the Fermilab staff and the technical staffs

of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

- [1] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1995); S. Abachi *et al.* (D0 Collaboration), Phys. Rev. Lett. **74**, 2632 (1995).
 [2] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **74**, 072006 (2006); V. Abazov *et al.* (D0 Collaboration), Phys. Lett. B **626**, 35 (2005); D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 142001 (2004); V. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **76**, 052006 (2007); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **103**, 092002 (2009); V. Abazov *et al.*




FIG. 2: Distribution of the SLT track p_T for SM+XM tags (a), track p_T for SM-XM tags (b), event H_T for SM+XM tags (c), and event H_T for SM-XM tags (d). The expectation for a hypothetical XM top quark is shown in the dotted line.

- (D0 Collaboration), Phys. Rev. Lett. **103**, 092001 (2009).
 [3] D. Chang, W. Chang, and E. Ma, Phys. Rev. D **59**, 091503 (1999).
 [4] J. P. Chou, Ph.D. thesis, Harvard University, FERMILAB-THESIS-2008-93, 2008; T. Aaltonen, *et al.* (CDF Collaboration), Phys. Rev. D **81**, 092002 (2010).
 [5] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **72**, 032002 (2005); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **79**, 502007 (2009).
 [6] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
 [7] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **73**, 032003 (2006).
 [8] A. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **98**, 041801 (2007).
 [9] A. Abulencia *et al.* (CDF Collaboration), J. Phys. G Nucl. Part. Phys. **34**, (2007).
 [10] We use a transverse coordinate system, where $p_T = p \sin(\theta)$ and $E_T = E \sin(\theta)$ are the momentum and energy measured transverse to the beamline, respectively. We define the pseudorapidity variable, $\eta \equiv -\ln \tan(\theta/2)$. We define \cancel{E}_T as the negative vector sum of the transverse energies in all the calorimeter cells, $-\sum E_T^i$.
 [11] C. Hill *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **530**, 1 (2004); A. Sill *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **447**, 1 (2000); A. Affolder *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **453**, 84 (2000).
 [12] T. Affolder *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **526**, 249 (2004).
 [13] L. Balka *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 272 (1988); S. Bertolucci *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 301 (1988).
 [14] G. Ascoli *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **268**, 33 (1998).
 [15] D. Acosta *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **494**, 57 (2002).
 [16] We reconstruct jets using a fixed-cone algorithm with a cone size of $\Delta R \equiv \sqrt{\Delta\eta^2 + \Delta\phi^2} \leq 0.4$, where ϕ is the azimuthal angle.
 [17] A. Bhatti *et al.* (CDF Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **566**, 2 (2006). We correct jet energies for detector variation, multiple interactions, and un-instrumented regions of the detector.
 [18] T. Sjostrand, S. Mrenna, and P. Skands, J. High Energy Phys. 0605 (2006) 026. We use PYTHIA version 6.216.
 [19] M. Cacciari *et al.*, J. High Energy Phys. 0809 (2008) 127; N. Kidonakis and R. Vogt, Phys. Rev. D **78**, 074005 (2008); S. Moch and P. Uwer, Nucl. Phys. Proc. Suppl. **183**, 75-80 (2008).
 [20] G. Corcella *et al.*, J. High Energy Phys. 01 (2001) 010. We use HERWIG version 6.510.
 [21] R. E. Kass and A. E. Raftery, J. Amer. Statist. Assoc. **90**, 773 (1995).