### Critical Points of D-Dimensional Extended Gravities

#### S. Deser\*

Physics Department, Brandeis University, Waltham, MA 02454, USA and Lauritsen Laboratory, California Institute of Technology, Pasadena, CA 91125, USA

### Haishan Liu<sup>†</sup>

Zheijiang Institute of Modern Physics, Department of Physics, Zheijiang University, Hangzhou 310027, China

#### H. Lü<sup>‡</sup>

China Economics and Management Academy, Central University of Finance and Economics, Beijing 100081 and Institute for Advanced Study, Shenzhen University, Nanhai Ave 3688, Shenzhen 518060, China

# C.N. Pope§

George P. & Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA and DAMTP, Centre for Mathematical Sciences, Cambridge University, Wilberforce Road, Cambridge CB3 OWA, UK

Tahsin Çağrı Şişman and Bayram Tekin\*\*

Department of Physics, Middle East Technical University, 06531, Ankara, Turkey

(Dated: May 28, 2018)

We study the parameter space of D-dimensional cosmological Einstein gravity together with quadratic curvature terms. In D>4 there are in general two distinct (anti)-de Sitter vacua. We show that for appropriate choice of the parameters there exists a critical point for one of the vacua, for which there are only massless tensor, but neither massive tensor nor scalar, gravitons. At criticality, the linearized excitations have vanishing energy (as do black hole solutions). A further restriction of the parameters gives a one-parameter cosmological Einstein plus Weyl<sup>2</sup> model with a unique vacuum, whose  $\Lambda$  is determined.

#### 1. Introduction

Adding new types of terms—and their associated parameters—to an action can lead, in the extended parameter space, to solutions with qualitatively different properties than in the component pieces. In the gravitational models we consider here, such phenomena have already been found, even at the linearized level. Historical examples include multiple maximally-symmetric vacua, one flat and the other (A)dS in Einstein-Hilbert gravity with quadratic curvature terms in dimensions  $D \neq 4$  [1]; "partial masslessness" [2], in which degrees of freedom are lost and gauge invariance "emerges" for massive higher spins s > 1 in AdS backgrounds with suitable mass<sup>2</sup>/ $\Lambda$  ratios; and the tuning of masses in the "new massive gravity" (NMG) in D = 3 [3, 4]. Another example where the tuning of parameters leads to new effects is the so-called "chiral gravity" [5], which is a special case of topologically massive gravity (TMG) in AdS [6]. Most recently [7], still wider critical effects were found in D = 4 systems combining the Einstein-Hilbert action, a cosmological term, and terms quadratic in the curvature. The latter work motivates our extension to generic dimensions, where the interplay between the multiple parameters will provide novel effects, some relying on the explicit cosmological term. For example, as we shall see, multiple vacua and "tuned" masslessness will now interact. We will find these critical models by studying their conserved charges, especially the energy, of black holes, as well as their linearized excitation spectrum.

<sup>\*</sup>Electronic address: deser@brandeis.edu

<sup>†</sup>Electronic address: liu@zimp.zju.edu.cn

<sup>&</sup>lt;sup>‡</sup>Electronic address: mrhonglu@gmail.com

<sup>§</sup>Electronic address: pope@physics.tamu.edu

<sup>¶</sup>Electronic address: sisman@metu.edu.tr

<sup>\*\*</sup>Electronic address: btekin@metu.edu.tr

The most general quadratic gravity model is

$$I = \int d^D x \sqrt{-g} \left[ \frac{1}{\kappa} \left( R - 2\Lambda_0 \right) + \alpha R^2 + \beta R^{\mu\nu} R_{\mu\nu} + \gamma \left( R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - 4R^{\mu\nu} R_{\mu\nu} + R^2 \right) \right]. \tag{1}$$

We follow the notation and conventions of [8, 9], where relevant details of field equations, curvature linearizations, and conserved charges may also be found. For generic values of the parameters, there are two distinct (A)dS vacua<sup>1</sup>, with metrics satisfying  $R_{\mu\nu} = \frac{2\Lambda}{D-2} g_{\mu\nu}$ , where  $\Lambda$  is determined by the quadratic equation

$$\frac{\Lambda - \Lambda_0}{2\kappa} + f\Lambda^2 = 0, \qquad f \equiv (D\alpha + \beta) \frac{(D-4)}{(D-2)^2} + \gamma \frac{(D-3)(D-4)}{(D-1)(D-2)}.$$
 (2)

This phenomenon was first observed [1] in Einstein-Lovelock gravity ( $\alpha = \beta = 0$ ,  $\Lambda_0 = 0$ ), where Minkowski and (A)dS vacua coexist. The source-free linearized equation of motion for the metric fluctuations around an (A)dS vacuum,  $h_{\mu\nu} \equiv g_{\mu\nu} - \bar{g}_{\mu\nu}$ , becomes

$$c\,\mathcal{G}_{\mu\nu}^{L} + (2\alpha + \beta)\left(\bar{g}_{\mu\nu}\bar{\Box} - \bar{\nabla}_{\mu}\bar{\nabla}_{\nu} + \frac{2\Lambda}{D-2}\bar{g}_{\mu\nu}\right)R^{L} + \beta\left(\bar{\Box}\mathcal{G}_{\mu\nu}^{L} - \frac{2\Lambda}{D-1}\bar{g}_{\mu\nu}R^{L}\right) = 0,\tag{3}$$

where c is given by

$$c \equiv \frac{1}{\kappa} + \frac{4\Lambda D}{D-2}\alpha + \frac{4\Lambda}{D-1}\beta + \frac{4\Lambda (D-3)(D-4)}{(D-1)(D-2)}\gamma,$$

and the linearization of the Einstein tensor  $\mathcal{G}_{\mu\nu} \equiv R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu}$  (defined using the effective  $\Lambda$  of the system's vacuum state(s), not the bare parameter  $\Lambda_0$  in the action) is given by

$$\mathcal{G}_{\mu\nu}^{L} = R_{\mu\nu}^{L} - \frac{1}{2}\bar{g}_{\mu\nu}R^{L} - \frac{2\Lambda}{D-2}h_{\mu\nu}.$$

Here, the linearized Ricci tensor  $R^L_{\mu\nu}$  and scalar curvature  $R^L=\left(g^{\mu\nu}R_{\mu\nu}\right)^L$  are given by

$$R^{L}_{\mu\nu} = \frac{1}{2} \left( \bar{\nabla}^{\sigma} \bar{\nabla}_{\mu} h_{\nu\sigma} + \bar{\nabla}^{\sigma} \bar{\nabla}_{\nu} h_{\mu\sigma} - \bar{\Box} h_{\mu\nu} - \bar{\nabla}_{\mu} \bar{\nabla}_{\nu} h \right), \qquad R^{L} = -\bar{\Box} h + \bar{\nabla}^{\sigma} \bar{\nabla}^{\mu} h_{\sigma\mu} - \frac{2\Lambda}{D-2} h.$$

Taking the trace of (3) gives

$$\left[ \left( 4\alpha \left( D - 1 \right) + D\beta \right) \bar{\Box} - \left( D - 2 \right) \left( \frac{1}{\kappa} + 4f\Lambda \right) \right] R^{L} = 0. \tag{4}$$

We see that the D'Alembertian operator is removed if  $\alpha$  and  $\beta$  are chosen so that

$$4\alpha \left( D - 1 \right) + D\beta = 0,\tag{5}$$

and so  $R^L$  is constrained to vanish, provided that

$$\frac{1}{\kappa} + 4f\Lambda \neq 0. \tag{6}$$

(See [9], and the discussion below.) As in [7], we shall choose the gauge  $\bar{\nabla}^{\mu}h_{\mu\nu} = \bar{\nabla}_{\nu}h$ , which then leads to  $R^{L} = -\frac{2\Lambda}{D-2}h$ . Imposing the condition (5), which eliminates the scalar mode (see equation (30) of [9]), one has h = 0 from (4), and hence  $h_{\mu\nu}$  satisfies the transverse and traceless conditions

$$\bar{\nabla}^{\mu}h_{\mu\nu} = 0, \qquad h = 0. \tag{7}$$

<sup>&</sup>lt;sup>1</sup> Depending upon the choice of parameters, and assuming  $\Lambda_0 < 0$ , there can be either two AdS vacua or else one AdS and one dS. If  $\Lambda_0$  is positive, there will be either two dS vacua or one dS and one AdS. If  $\Lambda_0 = 0$  there will be one Minkowski vacuum and either an AdS or a dS vacuum.

The linearized Ricci tensor  $R_{\mu\nu}^L$  and the linearized Einstein tensor  $\mathcal{G}_{\mu\nu}^L$  in this transverse-traceless gauge become

$$R^{L}_{\mu\nu} = \frac{2D\Lambda}{\left(D-1\right)\left(D-2\right)}h_{\mu\nu} - \frac{1}{2}\bar{\Box}h_{\mu\nu}, \qquad \mathcal{G}^{L}_{\mu\nu} = \frac{2\Lambda}{\left(D-1\right)\left(D-2\right)}h_{\mu\nu} - \frac{1}{2}\bar{\Box}h_{\mu\nu}.$$

The equations of motion simplify to

$$-\frac{\beta}{2} \left( \bar{\Box} - \frac{4\Lambda}{(D-1)(D-2)} - M^2 \right) \left( \bar{\Box} - \frac{4\Lambda}{(D-1)(D-2)} \right) h_{\mu\nu} = 0, \tag{8}$$

where

$$M^{2} \equiv -\frac{1}{\beta} \left( c + \frac{4\Lambda\beta}{(D-1)(D-2)} \right). \tag{9}$$

The massless and massive modes satisfy

$$\left(\bar{\Box} - \frac{4\Lambda}{(D-1)(D-2)}\right)h_{\mu\nu}^{(m)} = 0, \qquad \left(\bar{\Box} - \frac{4\Lambda}{(D-1)(D-2)} - M^2\right)h_{\mu\nu}^{(M)} = 0, \tag{10}$$

respectively. Stability requires  $M^2 \geq 0$ , and the case where  $M^2 = 0$ , i.e

$$c + \frac{4\Lambda\beta}{(D-1)(D-2)} = 0, (11)$$

defines the critical point.

## 3. Energy

We now show that at the critical point, the mass and angular momenta of all asymptotically Kerr-AdS and Schwarzschild-AdS black holes vanish.<sup>2</sup> We can calculate the conserved charges of any such asymptotically Kerr-AdS space, as discussed in [11]. Letting  $\bar{\xi}_{\mu}$  be a Killing vector of AdS, the conserved charges associated with this Killing vector will be given by [8]

$$Q^{\mu}(\bar{\xi}) = \left(c + \frac{4\Lambda\beta}{(D-1)(D-2)}\right) \int_{\mathcal{M}} d^{D-1}x \sqrt{-\bar{g}}\bar{\xi}_{\nu}\mathcal{G}_{L}^{\mu\nu} + (2\alpha + \beta) \int_{\partial\mathcal{M}} dS_{i}\sqrt{-g} \left\{\bar{\xi}^{\mu}\bar{\nabla}^{i}R_{L} + R_{L}\bar{\nabla}^{\mu}\bar{\xi}^{i} - \bar{\xi}^{i}\bar{\nabla}^{\mu}R_{L}\right\} + \beta \int_{\partial\mathcal{M}} dS_{i}\sqrt{-g} \left\{\bar{\xi}_{\nu}\bar{\nabla}^{i}\mathcal{G}_{L}^{\mu\nu} - \bar{\xi}_{\nu}\bar{\nabla}^{\mu}\mathcal{G}_{L}^{i\nu} - \mathcal{G}_{L}^{\mu\nu}\bar{\nabla}^{i}\bar{\xi}_{\nu} + \mathcal{G}_{L}^{i\nu}\bar{\nabla}^{\mu}\bar{\xi}_{\nu}\right\}.$$
(12)

Here,  $\mathcal{M}$  is a spacelike hypersurface in the asymptotically AdS space, and  $\partial \mathcal{M}$  denotes its boundary. The first integral can also be written as a boundary term, see [8, 12]. In asymptotically AdS spaces, only the first term survives. (We recall that the charges in (12) are measured at infinity; their detailed form as volume integrals is given by the spatial integral of the field equations' nonlinear terms.) At the critical point then, the charges, and in particular the energy  $(Q^0)$  vanishes. For example, the energy of the Schwarzschild-AdS solution, with asymptotics  $h_{00} = h^{rr} = \left(\frac{r_0}{r}\right)^{D-3}$ , is [8]

$$E_{BH} = \left(c + \frac{4\Lambda\beta}{(D-1)(D-2)}\right) \frac{(D-2)}{2} \Omega_{D-2} r_0^{D-3},\tag{13}$$

where  $\Omega_{D-2}$  is the solid angle on the D-2 sphere. Specifically, in four dimensions,  $\kappa=8\pi G_N$  and  $r_0=2G_N m$ , and then  $E_{BH}=\left[1+16\pi G_N\left(4\alpha+\beta\right)\right]m$ . At the critical point (11), we see that in all dimensions,  $E_{BH}=0$ .

Note that in D=3, the critical theory reduces to the NMG at the "Proca" point. Namely,  $8\alpha+3\beta=0$  gives the NMG theory, and the criticality condition  $M^2=0$  gives  $\Lambda_0=\frac{3}{\kappa\beta}$  which is exactly the point where the linearized NMG can be written explicitly as a unitary massive spin-1 theory with mass squared  $-\frac{8}{\kappa\beta}$  (see [4], and Sec. V.A in [3]). (For AdS and with  $\beta>0$ , one should set  $\kappa<0$  since  $\Lambda=\frac{2}{\kappa\beta}$ .)

<sup>&</sup>lt;sup>2</sup> Actually, unlike for cosmological Einstein theory [10], explicit Kerr-AdS type solutions for these models are as yet unknown. Such solutions presumably exist, and they will approach the standard Kerr-AdS metrics at large distance where the effects of the higher-order curvature terms become negligible.

#### 4. Linear excitations

Let us now consider the energy of the excitations, expressed as explicit volume integrals, by constructing the Hamiltonian for the theory, as in [5, 7]. The quadratic Lagrangian for the metric perturbations of (1) around an AdS vacuum is given by simply multiplying the left-hand side of (3) by  $-\frac{1}{2}h^{\mu\nu}$ ; hence it vanishes on-shell. With the parameter choice (5), and in the transverse-traceless gauge (7), the quadratic action therefore takes the form

$$I_{2} = -\frac{1}{2} \int d^{D}x \sqrt{-\bar{g}} \left[ -\frac{\beta}{2} \bar{\Box} h^{\mu\nu} \bar{\Box} h_{\mu\nu} - \frac{1}{2} \left( \frac{4\Lambda\beta}{(D-1)(D-2)} - c \right) \bar{\nabla}^{\rho} h^{\mu\nu} \bar{\nabla}_{\rho} h_{\mu\nu} + \frac{2\Lambda c}{(D-1)(D-2)} h^{\mu\nu} h_{\mu\nu} \right]. \quad (14)$$

The canonical momenta, using the Ostrogradsky formalism for higher-order Lagrangians, are defined and computed as follows:

$$\Pi_{(1)}^{\mu\nu} \equiv \frac{\delta \mathcal{L}_2}{\delta \dot{h}_{\mu\nu}} - \bar{\nabla}_0 \left( \frac{\delta \mathcal{L}_2}{\delta \left[ \frac{\partial}{\partial t} \left( \bar{\nabla}_0 h_{\mu\nu} \right) \right]} \right) = -\frac{\sqrt{-\bar{g}}}{2} \bar{\nabla}^0 \left[ -\left( \frac{4\Lambda\beta}{(D-1)(D-2)} - c \right) h^{\mu\nu} + \beta \bar{\Box} h^{\mu\nu} \right],$$

$$\Pi_{(2)}^{\mu\nu} \equiv \frac{\delta \mathcal{L}_2}{\delta \left[ \frac{\partial}{\partial t} \left( \bar{\nabla}_0 h_{\mu\nu} \right) \right]} = \frac{\sqrt{-\bar{g}}}{2} \beta \bar{g}^{00} \bar{\Box} h^{\mu\nu}.$$

The Hamiltonian can be written as

$$H \equiv \int d^{D-1}x \left[ \Pi_{(1)}^{\mu\nu} \dot{h}_{\mu\nu} + \Pi_{(2)}^{\mu\nu} \frac{\partial}{\partial t} \left( \bar{\nabla}_0 h_{\mu\nu} \right) \right] - \int \sqrt{-g} \mathcal{L}_2 d^{D-1}x, \tag{15}$$

where we use for the implicit AdS background metric the time-independent form

$$d\bar{s}^2 = \frac{(D-1)(D-2)}{2(-\Lambda)} \left[ -\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\Omega_{D-2}^2 \right].$$

Substituting the canonical momenta into (15) gives the Hamiltonian for the quadratic fluctuations. The energies of the various (linearized) excitations could be obtained directly from (15), but a more convenient starting point is to note that the Hamiltonian H is time-independent, since the Lagrangian has no explicit time dependence. We are therefore free to write H as its time average,  $H = \langle H \rangle = T^{-1} \int_0^T dt H$ . The advantage of doing this is that we can then perform integrations by parts for time derivatives. This allows us to manipulate (15) into a more convenient form. The Lagrangian term  $\int \sqrt{-g} \mathcal{L}_2 d^{D-1} x$ , being proportional to the equation of motion (8), does not contribute to the on-shell energies. Substituting the expressions (10) for the massless and the massive modes into the time average of the Hamiltonian (15), we find, after an integration by parts, that their energies are given by

$$E_{m} = -\frac{1}{2T} \left( c + \frac{4\Lambda\beta}{(D-1)(D-2)} \right) \int d^{D}x \sqrt{-\bar{g}} \left( \dot{h}_{\mu\nu}^{(m)} \bar{\nabla}^{0} h_{(m)}^{\mu\nu} \right), \tag{16}$$

$$E_{M} = \frac{1}{2T} \left( c + \frac{4\Lambda\beta}{(D-1)(D-2)} \right) \int d^{D}x \sqrt{-\bar{g}} \left( \dot{h}_{\mu\nu}^{(M)} \bar{\nabla}^{0} h_{(M)}^{\mu\nu} \right), \tag{17}$$

where the time integrations are understood to be over the interval T. The excitation energies  $E_m$  and  $E_M$  have opposite signs, as expected from their fourth-order origin [13], but at the critical point (11), they vanish. The integral itself for the massless modes is known to be negative, since in pure Einstein gravity for which  $\alpha = \beta = \gamma = 0$ , the energies  $E_m$  are known to be positive. The integral for the massive modes is also expected to be negative, away from  $M^2 = 0$ , at least for small mass.

We started with a theory with four parameters, namely  $\Lambda_0$ ,  $\alpha$ ,  $\beta$  and  $\gamma$  (we are not counting  $\kappa$ , since it can be scaled out). The critical point is defined by the two conditions (5) and (11), implying that any two of the four parameters can be eliminated. Note, however, that we must also require that (6) hold at the critical point, since otherwise,  $R_L$ , a gauge invariant object, would be left undetermined in the theory.

As we already remarked, there are in general two distinct AdS vacua in the theory with given values for the four parameters  $\Lambda_0$ ,  $\alpha$ ,  $\beta$  and  $\gamma$ , corresponding to the two roots of the quadratic equation (2). The specialisation to the critical case that we have been discussing turns the massive spin-2 modes massless for just one of these vacua. In the other vacuum, the spectrum still contains massive as well as massless spin-2 modes, with excitation energies of opposite signs. Whether it is the massive spin-2 modes or the massless spin-2 modes that have the negative energy depends upon the detailed choice of parameters. Note that if the parameters are chosen such that the massless spin-2 modes have positive energy, then it will also be the case that black holes in the non-critical vacuum will have positive mass.

For some purposes, in order to avoid solving the quadratic equation (2) for  $\Lambda$ , it is convenient instead to view it as a linear equation for the parameters of the theory expressed in terms of the critical value of  $\Lambda$ . It seems to be most convenient to take  $\alpha$  and  $\gamma$  as the free parameters for D > 4. From (5), one determines  $\beta$  in terms of  $\alpha$ . Then, from (11), one obtains the unique critical vacuum with a cosmological constant

$$\Lambda_{\text{crit}} = -\frac{D(D-1)(D-2)}{4\kappa \left[ (D-1)(D-2)^2 \alpha + D(D-3)(D-4) \gamma \right]}.$$
 (18)

To determine the corresponding critical  $\Lambda_0$ , we can use (2) to get

$$\Lambda_0 = -\frac{D^2 (D-1) (D-2) [(D-1) (D-2) \alpha + (D-3) (D-4) \gamma]}{8\kappa \left[ (D-1) (D-2)^2 \alpha + D (D-3) (D-4) \gamma \right]^2}.$$
 (19)

The other (non-critical) vacuum has the cosmological constant

$$\Lambda_{\text{noncrit}} = \frac{D[(D-1)(D-2)\alpha + (D-3)(D-4)\gamma] \Lambda_{\text{crit}}}{(D-4)[(D-1)(D-2)\alpha + D(D-3)\gamma]}.$$
 (20)

As a result, the two-parameter critical theory is given by the action

$$I = \int d^{D}x \sqrt{-g} \left[ \frac{1}{\kappa} (R - 2\Lambda_{0}) + \alpha R^{2} - \frac{4(D-1)}{D} \alpha R^{\mu\nu} R_{\mu\nu} + \gamma \left( R^{\mu\nu\rho\sigma} R_{\mu\nu\rho\sigma} - 4R^{\mu\nu} R_{\mu\nu} + R^{2} \right) \right], \tag{21}$$

where the bare cosmological constant is given by (19). The theory has a critical vacuum (18), while there is a noncritical vacuum (20).

Finally, one can overcome the two-vacuum problem, that one of them necessarily allows negative-energy excitations, using one last allowed (consistent as we shall see) reduction, to a single parameter. We require f = 0 in (2), whose only solution is thereby  $\Lambda = \Lambda_0$ . Our four parameters  $(\alpha, \beta, \gamma, \Lambda_0)$ , are now constrained by the three conditions (2, 11), plus scalar mode suppression (5), to obey

$$\beta = -\frac{4\alpha (D-1)}{D}, \qquad \Lambda = \Lambda_0 = -\frac{D}{8\kappa \alpha}, \qquad \alpha = -\frac{\gamma D (D-3)}{(D-1)(D-2)}.$$
 (22)

Note that  $\Lambda_0$  is also fixed in terms of the remaining parameter. The general action (1) now reduces to Einstein-Weyl form with a single arbitrary constant  $\gamma$ ,

$$I = \int d^D x \sqrt{-g} \left[ \frac{1}{\kappa} \left( R - 2\Lambda_0 \right) + \gamma C^{\mu\nu\rho\sigma} C_{\mu\nu\rho\sigma} \right], \qquad \Lambda = \Lambda_0 = \frac{(D-1)(D-2)}{8\kappa\gamma(D-3)}. \tag{23}$$

Here,  $C_{\mu\nu\rho\sigma}$  is the Weyl tensor, for which

$$C^{\mu\nu\rho\sigma}C_{\mu\nu\rho\sigma} = R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma} - \frac{4}{D-2}R^{\mu\nu}R_{\mu\nu} + \frac{2}{(D-1)(D-2)}R^2.$$

Physically, it is clear a priori that only the combination (23) has a single vacuum: the Weyl tensor, being insensitive to conformally related metrics, such as different (A)dS or flat ones, does not contribute directly to (2). We also emphasize that while condition (6) is automatically satisfied by taking f = 0, attempting instead to force the two roots of (2) to be equal would violate it.

#### 5. Conclusions

We have constructed a two-parameter theory of gravity in D dimensions, which admits a critical AdS vacuum in which there are only massless spin-2 modes. For D=3, the critical theory reduces to the NMG at the Proca point, whilst for D=4, it reduces to the critical theory constructed recently in [7]. In four dimensions, once the condition (5) for the elimination of the scalar mode is imposed, the theory has a unique AdS vacuum solution. For dimensions  $D\neq 4$ , by contrast, the theory in general has two distinct AdS vacua, which cannot simultaneously be rendered critical for any choice of parameters. For  $D \geq 5$ , a further specialisation of parameters can be made, leading to a one-parameter critical theory with a unique AdS vacuum. The surprising confluence of so many unexpected properties at our models' critical points may repay further investigation.

### I. ACKNOWLEDGMENTS

S.D. is supported by NSF PHY 07-57190 and DOE DE-FG02-164 92ER40701 grants. H. Liu is supported in part by the National Science Foundation of China (10425525,10875103), National Basic Research Program of China (2010CB833000) and Zhejiang University Group Funding (2009QNA3015). The research of C.N.P. is supported in part by DOE grant DE-FG03-95ER40917. The work of T.C.S. and B.T. is supported by the TÜBİTAK Grant No. 110T339, and METU Grant BAP-07-02-2010-00-02. H. Lü is grateful to the Mitchell Institute for Fundamental Physics and Astronomy at Texas A&M University for hospitality during the course of this work.

- [1] D. G. Boulware and S. Deser, Phys. Rev. Lett. 55, 2656 (1985).
- [2] S. Deser and A. Waldron, Phys. Rev. Lett. 87, 031601 (2001); Nucl. Phys. B 607, 577 (2001).
- [3] E. A. Bergshoeff, O. Hohm and P. K. Townsend, Phys. Rev. D 79, 124042 (2009).
- [4] Y. Liu and Y. W. Sun, JHEP **0904**, 106 (2009).
- [5] W. Li, W. Song and A. Strominger, JHEP **0804**, 082 (2008).
- [6] S. Deser, R. Jackiw and S. Templeton, Annals Phys. 140, 372 (1982); Phys. Rev. Lett. 48, 975 (1982).
- [7] H. Lü and C. N. Pope, arXiv:1101.1971 [hep-th].
- [8] S. Deser and B. Tekin, Phys. Rev. Lett. 89, 101101 (2002); Phys. Rev. D 67, 084009 (2003).
- [9] I. Gullu and B. Tekin, Phys. Rev. D 80, 064033 (2009).
- [10] G. W. Gibbons, H. Lü, D. N. Page and C. N. Pope, J. Geom. Phys. 53, 49 (2005); Phys. Rev. Lett. 93, 171102 (2004).
- [11] S. Deser, I. Kanik and B. Tekin, Class. Quant. Grav. 22, 3383 (2005).
- [12] L. F. Abbott and S. Deser, Nucl. Phys. B **195**, 76 (1982).
- [13] K. S. Stelle, Phys. Rev. **D16**, 953 (1977); Gen. Rel. Grav. **9**, 353 (1978).