

Accidents in (0,2) Landau-Ginzburg theories

Marco Bertolini,^{a,b} Ilarion V. Melnikov,^c and M. Ronen Plesser^a

^a*Center for Geometry and Theoretical Physics, Box 90318
Duke University, Durham, NC 27708-0318, USA*

^b*Kavli Institute for Theoretical Physics
University of California, Santa Barbara, CA 93106-4030, USA*

^c*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M
University, College Station, TX 77843, USA*

E-mail: mb266@phy.duke.edu, ilarion@physics.tamu.edu,
plesser@cgtp.duke.edu

ABSTRACT: We study the role of accidental symmetries in two-dimensional (0,2) superconformal field theories obtained by RG flow from (0,2) Landau-Ginzburg theories. These accidental symmetries are ubiquitous, and, unlike in the case of (2,2) theories, their identification is key to correctly identifying the IR fixed point and its properties. We develop a number of tools that help to identify such accidental symmetries in the context of (0,2) Landau-Ginzburg models and provide a conjecture for a toric structure of the SCFT moduli space in a large class of models. We also give a self-contained discussion of aspects of (0,2) conformal perturbation theory.

Contents

1	Introduction	1
2	A glance at (0,2) Landau-Ginzburg theories	4
3	Accidents	10
3.1	Accidents in (2,2) Landau-Ginzburg orbifolds	10
3.2	A contrived (2,2) example	10
3.3	A simple (0,2) example	11
3.4	Puzzles from enhanced symmetries	12
3.5	Enhanced symmetries of (2,2) LG theories	13
3.6	Subtleties for heterotic vacua	15
4	Marginal deformations of a unitary (2,0) SCFT	16
4.1	Basic results	16
4.2	A few consequences	18
4.3	Deformations and left-moving abelian currents	21
5	Toric geometry of the deformation space	22
5.1	The toric conjecture	23
5.2	Enhanced toric symmetries	26
5.3	Examples	28
5.4	Summary and further thoughts	31
6	Outlook	32
A	An F-term obstruction	33
B	A D-term obstruction in a heterotic vacuum	34

1 Introduction

The construction and classification of conformal field theories (CFTs) plays a key role in modern quantum field theory. One approach to this problem is to solve the conformal bootstrap.¹ Another approach that has proven useful is to study the low energy (or IR) limits of

¹This is difficult in practice and can be carried out analytically only in theories with enormously enlarged symmetries like the W_N algebras of the minimal models [1]. Recently progress has been made with numeric techniques, for example in applications to the three-dimensional Ising model [2, 3].

renormalization group (RG) flows from known CFTs. This has challenges of its own, since the IR dynamics often involves emergent degrees of freedom and interactions.

Nevertheless, as already indicated in the seminal work of [4–7], it is often possible to identify certain classes of operators and their OPEs and correlators of an IR CFT with corresponding objects in terms of the UV degrees of freedom. This is especially useful when the UV theory is asymptotically free, since then perturbative computations can provide information about a non-trivial CFT without a notion of a weak coupling. The identification of UV and IR data is simplified when some amount of supersymmetry is preserved along the RG flow: SUSY constraints lead to well-known simplifications such as the relation between dimensions and R-charges of chiral operators and non-renormalization theorems. For instance, in two dimensional theories with (2,2) SUSY these structures are responsible for many well-known phenomena such as mirror symmetry and the Calabi-Yau (CY) / Landau-Ginzburg (LG) correspondence [8], and the identification of UV and IR data is a key tool in exploration and exploitation of these two-dimensional gems.

Such techniques rely on the assumption that accidental symmetries that might emerge in the IR limit do not invalidate the identification of operators in the IR with their UV avatars. This assumption is well-tested in (2,2) theories but is also often applied to theories with only (0,2) supersymmetry. For instance, it is key to various gauged linear sigma model constructions of (0,2) CFTs corresponding to heterotic string vacua [9–12].

In this note we show that the assumption cannot be taken for granted in (0,2) theories, and the resulting “accidents” have drastic consequences for the IR physics and the relation between UV parameters and IR data. The examples we consider are (0,2) Landau-Ginzburg theories, and we identify a class of accidental symmetries of (0,2) LG RG flows by studying the space of F-term UV couplings modulo field redefinitions. We find that these accidental symmetries significantly modify the analysis of the IR theory. For instance, the spectrum of chiral operators and even the IR central charge are in general modified. This invalidates certain UV theories from giving good models for (0,2) SCFTs appropriate for a heterotic string vacuum—we examine an example taken from [10].

A classic (2,2) example

To describe the challenges of (0,2) accidents more precisely, it is useful to review the successes of the (2,2) theories. Consider the quintic (2,2) LG model with chiral superpotential

$$W = \alpha_0 X_0^5 + \alpha_1 X_1^5 + \cdots + \alpha_4 X_4^5 - 5\alpha_5 X_0 X_1 \cdots X_4 .$$

Here the X_i are (2,2) chiral superfields and the α_a are complex parameters. This theory flows to a $c = \bar{c} = 9$ (2,2) SCFT.² The complex parameter

$$\psi = \alpha_5^5 (\alpha_0 \alpha_1 \cdots \alpha_4)^{-1}$$

is invariant under \mathbb{C}^* rescalings of the chiral superfields X_i and labels a one-parameter family of IR CFTs. At generic values of ψ the IR fixed point is a well-behaved CFT, and small changes in ψ correspond to small marginal deformations of the CFT, where “small” refers to the distance in the Zamolodchikov metric. At special values of ψ the CFT can become singular. For instance, $\psi = 1$ is a finite distance singularity —the analogue of a conifold point. This can be detected in the UV description: the theory develops a family of supersymmetric vacua with $X_i = \text{const.}$, and these signal a non-compact CFT: a theory with a continuum spectrum of conformal dimensions. Another point, $\psi = \infty$ is an infinite-distance singularity.

The quotient of the UV parameter space by field redefinitions is a complicated object [13, 14] with singularities and non-separated points. For instance, we can take the limit $\alpha_0 \rightarrow 0$ and $\alpha_5 \rightarrow 0$ while keeping ψ constant so as to obtain a product of four minimal models coupled to a free chiral superfield X_0 , with $c = \bar{c} = 3(3 + \frac{2}{5})$. Fortunately, all such bad points are singular CFTs. The bad points corresponding to various infinite distance singularities and “wrong” central charges are easily identified in terms of the UV data: they all correspond to singular superpotentials with a continuum of supersymmetric vacua and therefore a continuum of states in the IR CFT. Away from such points the quotient is sensible and describes (2,2) marginal deformations of the IR theory.

Similar considerations apply to more general (2,2) gauged linear sigma models, and with the parametrization of the smooth CFTs in terms of the UV parameters in hand, localization and topological field theory techniques can be used to compute certain correlators and chiral spectra in the CFT in terms of the weakly coupled UV Lagrangian.

(0,2) challenges

As we will show in section 3, the situation is more delicate in (0,2) theories, even in the relatively simple class of LG models (we review these in section 2). The essential difference is that we lack the simple diagnostic we had for a “bad” point in (2,2) theories. It is not sufficient to exclude UV parameters that lead to flat directions in the potential, and the identification of UV parameters with marginal deformations of the CFT requires (at least) a study of loci with enhanced symmetry. Unlike in (2,2) examples, accidental symmetries can emerge for non-singular UV potential, thereby complicating the description of IR physics in terms of the UV data. Unlike in (2,2) theories a family of smooth UV potentials with each potential preserving the same R-symmetry along the RG flow need not correspond to a family of CFTs related by truly marginal deformations.

²A \mathbb{Z}_5 orbifold of the theory describes a (2,2) non-linear sigma model with target space the quintic CY hypersurface in \mathbb{P}^4 at a special value of the complexified Kähler parameter. A complex structure parameter of the geometry is then related to the LG parameter ψ .

Fortunately, at least in (0,2) LG models it appears that we have enough control to identify accidental symmetries and special loci in the parameter space by generalizing the (2,2) paradigm of parametrizing the IR fixed points by the space of UV parameters modulo field redefinitions. This uncovers a rich structure of (0,2) RG flows and of the space of marginal deformations of (0,2) fixed points and will undoubtedly play a role in quantitative descriptions of (0,2) moduli spaces.

Once we have identified a (0,2) LG theory with some particular IR fixed point, it is useful to develop the correspondence between deformations of the UV Lagrangian and (0,2) conformal perturbation theory. In section 4 we describe some properties of (0,2) conformal perturbation theory independent of any embedding of the CFT in a critical heterotic string. This section can be read independently from the rest of the paper. We use these observations in section 5 to describe a conjecture for the global structure of the moduli space of (0,2) SCFTs with expected central charge in terms of the UV data for what we term *plain* (0,2) LG models.

Acknowledgments

IVM would like to thank the Enrico Fermi Institute for hospitality while this work was being completed and especially David Kutasov for discussions. This work is supported in part by the NSF Focused Research Grant DMS-1159404 and Texas A&M (IVM), by NSF Grant PHY-1217109 (MB and MRP) and NSF Grant PHY11-25915 (MB). MRP thanks the Aspen Center for Physics (and NSF Grant 1066293) for hospitality during the initial stage of this work. We thank the organizers of the workshop on Heterotic Strings and (0,2) QFT at Texas A&M University in May, 2014 for hosting MB and MRP while some final steps were completed, and for allowing MRP to present a preliminary version. We also thank Paul Aspinwall and Eric Sharpe for some useful conversations, and IVM and MRP thank Ido Adam for discussions and collaboration on aspects of (0,2) conformal perturbation theory.

2 A glance at (0,2) Landau-Ginzburg theories

We begin with a quick review of (0,2) LG theories following [10, 15, 16].³ We work in Euclidean signature and a (0,2) superspace with coordinates $(z, \bar{z}, \theta, \bar{\theta})$. The UV theory consists of n bosonic chiral multiplets $\Phi_i = \phi_i + \dots$, and N fermionic chiral multiplets $\Gamma^A = \gamma^A + \dots$, as well as their conjugate anti-chiral multiplets. These are given a free kinetic term and a (0,2) SUSY potential term as interactions:

$$\mathcal{L}_{\text{int}} = \int d\theta \mathcal{W} + \text{h.c.} , \quad \mathcal{W} = m_0 \sum_A \Gamma^A J_A(\Phi) , \quad (2.1)$$

where m_0 has mass dimension 1 and the J_A are polynomial in the Φ_i . This is the simplest example of a (0,2) SUSY asymptotically free theory: for energies $E \gg m_0$ the theory is

³Our superspace conventions are those of [16].

well-described by the set of free fields. Conversely, when $E \ll m_0$ the interactions become important and lead to non-trivial IR dynamics that depend on n, N , as well as the choice of ideal $\mathcal{J} = \langle J_1, \dots, J_N \rangle \subset \mathbb{C}[\Phi_1, \dots, \Phi_n]$. What can we say about the IR limit of this theory?

A basic constraint comes from the gravitational anomaly. In the UV the central charges are easy to determine: each Φ multiplet contains a complex boson and a right-moving Weyl fermion, while each Γ contains a left-moving Weyl fermion and an auxiliary field. Hence, we have $c_{UV} = 2n + N$, $\bar{c}_{UV} = 3n$. The RG flow induced by \mathcal{W} will decrease the central charges, but since it is Lorentz-invariant, it will preserve the difference $c - \bar{c} = N - n$.

Another basic property of the theory is the set of global symmetries. The free theory has a large global symmetry that commutes with (0,2) SUSY: namely, $U(n) \times U(N)$ rotations of the chiral superfields. In addition we have the R-symmetry that rotates θ and leaves the lowest components of the superfields ϕ_i and γ^A invariant. The interactions break these symmetries. For completely generic J_A the remaining symmetry just $U(1)_R^0$ – an R-symmetry that assigns charge +1 to θ and γ^A and charge 0 to ϕ_i .

Properties of the superpotential

A key feature of (0,2) LG theories is that the holomorphic superpotential obeys the same non-renormalization properties as the, perhaps more familiar N=1 d=4 Wess-Zumino model’s superpotential. The kinetic term, on the other hand, is a full superspace derivative and will receive complicated corrections along the RG flow. However, just as in (2,2) theories, we expect these corrections to be irrelevant provided that the fields Φ and Γ all acquire non-trivial scaling dimensions. In order to relate these scaling dimensions to properties of the UV theory, we will assume that the interactions preserve an additional global $U(1)$ symmetry, which we will call $U(1)_L$, under which Γ^A have charges Q_A , while the Φ_i carry charges q_i . This will be the case if and only if the ideal is quasi-homogeneous, i.e.

$$J_A(t^{q_i} \Phi_i) = t^{-Q_A} J_A(\Phi) \quad (2.2)$$

for all $t \in \mathbb{C}^*$. We will demand that the ideal is zero-dimensional, i.e. $J_A(\Phi) = 0$ for all A if and only if $\Phi = 0$. If it is not then the theory necessarily has a non-compact set of supersymmetric vacua labeled by vevs of the bosonic fields. We will call such superpotentials singular. We are interested in “compact” CFTs and exclude this possibility.⁴

Another important property of the superpotential is that typically some of its parameters are not, in fact, F-terms. To see this, consider a perturbation of the form $\delta\mathcal{W} = \sum_A \Gamma^A \delta J_A$ around a theory with $\mathcal{W}_0 = \sum_A \Gamma^A J_A$. In the undeformed theory, if we assume canonical kinetic terms, the equations of motion read

$$\overline{\mathcal{D}}\Gamma^A = J_A(\Phi) , \quad \overline{\mathcal{D}}\partial_z \bar{\Phi}_i = \sum_A \Gamma^A \frac{\partial J_A}{\partial \Phi_i} , \quad (2.3)$$

⁴A CFT is compact if its spectrum is such that for every fixed real Δ there is a finite number of fields with dimension less than Δ .

where $\overline{\mathcal{D}}$ is the antichiral superspace derivative $\overline{\mathcal{D}} = \partial_{\overline{\theta}} + \theta \partial_{\overline{z}}$. A more general kinetic term leads to more complicated expressions under the $\overline{\mathcal{D}}$ derivative of the left-hand sides of the equations.⁵ Hence, a first order deformation of \mathcal{W} of the form

$$\delta J_A = \sum_B M_A^B(\Phi) J_B(\Phi) + \sum_i \frac{\partial J_A}{\partial \Phi_i} F_i(\Phi) \quad (2.4)$$

is equivalent up to equations of motion to a D-term deformation.

The LG assumption that the D-terms are irrelevant along the flow implies that any two UV theories with superpotentials related by a holomorphic field redefinition lead to the same IR fixed point. Hence, any two UV superpotentials that are related by a holomorphic field redefinition belong to the same universality class.

(2,2) LG theories

The (0,2) theory will have an enhanced left-moving SUSY when $N = n$, so that in the free limit we can combine (Γ^i, Φ_i) into (2,2) chiral multiplets X^i , and when $J_i = \partial W / \partial \Phi_i$ for some potential W . In that case, we can rewrite the theory in a manifestly (2,2) supersymmetric fashion with a chiral superpotential $W(X)$. The quasi-homogeneity conditions set $Q_i = q_i - 1$, and the resulting central charge is given by the famous

$$\overline{c} = 3 \sum_i (1 - 2q_i) . \quad (2.5)$$

IR consequences of UV symmetries

Returning to the more general (0,2) setting, if we assume that $U(1)_R^0$ and $U(1)_L$ are the only symmetries along the whole RG flow, then we can determine the linear combination of charges corresponding to the IR R-symmetry $U(1)_R^{\text{IR}}$ as well as those of a left-moving $U(1)_L^{\text{IR}}$. The charges of the latter are fixed up to normalization by the quasi-homogeneity condition, and the normalization is fixed by

$$-\sum_A Q_A - \sum_i q_i = \sum_A Q_A^2 - \sum_i q_i^2 . \quad (2.6)$$

This ensures that $U(1)_L^{\text{IR}}$ and $U(1)_R^{\text{IR}}$ have no mixed anomalies and become, respectively, left-moving and right-moving Kac-Moody symmetries in the IR theory. The central charge is determined from the two-point function of the $U(1)_R^{\text{IR}}$ current. The result is

$$\overline{c} = 3(r + n - N) , \quad r = -\sum_A Q_A - \sum_i q_i . \quad (2.7)$$

⁵In a NLSM such total derivatives are more subtle than in this LG setting, as they are usually only sensible patch by patch in target space. Indeed, marginal deformations of NLSMs are such F-terms that cannot be globally recast as D-terms [17, 18].

By studying the cohomology of the supercharge \overline{Q} of the theory, we can also describe chiral operators and their charges. More details can be found in [16], but for our purposes it will be sufficient to note the charges and corresponding dimensions of ϕ_i and γ^A . Denoting the $U(1)_L^{\text{IR}}$ and $U(1)_R^{\text{IR}}$ charges by, respectively, \mathbf{q} and $\overline{\mathbf{q}}$, we have

$$\begin{array}{ccc}
& \phi_i & \gamma^A \\
\mathbf{q} & q_i & Q_A \\
\overline{\mathbf{q}} & q_i & 1 + Q_A \\
h & \frac{q_i}{2} & \frac{2+Q_A}{2} \\
\overline{h} & \frac{q_i}{2} & \frac{1+Q_A}{2}
\end{array}$$

Since these are chiral operators, the right-moving weights are determined in the usual fashion $\overline{h} = \overline{\mathbf{q}}/2$, and the left-moving weights are fixed since RG flow preserves the spin of the operators.

This structure determines many properties of the IR theory such as the elliptic genus [15] and the topological heterotic ring [16]. As for (2,2) theories, there is also a simple prescription for using orbifolds of such (0,2) LG theories to build spacetime SUSY heterotic string vacua [10, 19]. For instance, the elliptic genus is given by [15]

$$\begin{aligned}
Z(\tau, z) &= \text{Tr}_{RR}(-)^F y^{J_{L0}^{\text{IR}}} e^{2\pi i \tau H_L} e^{-2\pi i \overline{\tau} H_R} \\
&= i^{N-n} e^{2\pi i \tau (N-n)/12} y^{-r/2} [\chi(y) + O(e^{2\pi i \tau})] \ ,
\end{aligned} \tag{2.8}$$

where $y = e^{2\pi i z}$, and

$$\chi(y) = \frac{\prod_A (1 - y^{-Q_A})}{\prod_i (1 - y^{q_i})} \Big|_{y^{\text{integer}}} \ . \tag{2.9}$$

The remaining τ -dependent terms are determined by modular properties of $Z(\tau, z)$.

Enhanced symmetry and c -extremization

For special values of the superpotential the UV theory will acquire enhanced symmetries that commute with the (0,2) SUSY algebra. In two dimensions these cannot be spontaneously broken, and, as in four dimensions, the abelian component $U(1)^M$ can mix with $U(1)_R^0$ and $U(1)_L$ symmetries. Fortunately, as in the four-dimensional case we can still find candidate $U(1)_L^{\text{IR}}$ and $U(1)_R^{\text{IR}}$ symmetries by applying the analogue of a -maximization [20] known as c -extremization [21]. We can summarize the results of [21] as follows. Let \mathcal{J}_0 denote the $U(1)_R^0$ R-symmetry current, and let \mathcal{J}_α , $\alpha = 1, \dots, M$ be the currents for $U(1)^M$. Assuming that the correct $U(1)_R^{\text{IR}}$ symmetry is a linear combination of \mathcal{J}_0 and the \mathcal{J}_α , [21] construct

the trial current and trial central charge

$$\mathcal{J} = \mathcal{J}_0 + \sum_{\alpha} t^{\alpha} \mathcal{J}_{\alpha} , \quad \frac{1}{3} \overline{\mathcal{C}} = n - N + 2 \sum_{\alpha} t_{\alpha} K^{\alpha} + \sum_{\alpha, \beta} t^{\alpha} t^{\beta} L_{\alpha\beta} , \quad (2.10)$$

where

$$K^{\alpha} = - \sum_A Q_A^{\alpha} - \sum_i q_i^{\alpha} , \quad L^{\alpha\beta} = \sum_i q_i^{\alpha} q_i^{\beta} - \sum_A Q_A^{\alpha} Q_A^{\beta} . \quad (2.11)$$

Here Q_A^{α} and q_i^{α} denote the $U(1)^{\alpha}$ charges of Γ^A and Φ_i , respectively. The $U(1)_{\text{R}}^{\text{IR}}$ is then identified by extremizing $\overline{\mathcal{C}}$ with respect to t_{α} , leading to $U(1)_{\text{L}}^{\text{IR}}$ charges

$$q_i = \sum_{\alpha} q_i^{\alpha} t_{\alpha*} , \quad Q_A = \sum_{\alpha} Q_A^{\alpha} t_{\alpha*} , \quad (2.12)$$

where $t_{\alpha} = t_{\alpha*}$ is the extremum point. The central charge is then also fixed as $\overline{\mathcal{c}} = \overline{\mathcal{C}}(t_*)$.

The symmetric form L has a real spectrum, and the sign of an eigenvalue has the following significance in the IR theory. We decompose the UV currents according to the sign of the eigenvalues as $\mathcal{J}_{\alpha} \rightarrow \{\mathcal{J}^+, \mathcal{J}^0, \mathcal{J}^-\}$. If we assume that there are no accidental symmetries in the IR, then unitarity of the SCFT implies that in the IR the \mathcal{J}^+ currents must correspond to right-moving Kac-Moody (KM) currents and the \mathcal{J}^- must flow to left-moving KM currents. Finally, the \mathcal{J}^0 must decouple from the SCFT degrees of freedom. The last point has two consequences: on one hand, we should treat a theory with $\ker L \neq 0$ with some care; on the other hand, if we can be certain that the IR limit is nevertheless a unitary CFT, we can without loss of generality restrict to symmetries orthogonal to $\ker L$.

In typical examples of (0,2) LG theories L is negative definite; we do not know of a non-singular model where L has a positive eigenvalue. In fact, as far as the extremization procedure goes, symmetries corresponding to the positive eigenspace of L cannot be broken in the SCFT. More precisely, a UV deformation away from an RG trajectory with a “positive” symmetry is irrelevant — in the IR the “positive” symmetry will be restored. To understand this, we consider the change in the extremized central charge upon breaking a symmetry. Assuming $\ker L = 0$, the extremum central charge is

$$\frac{1}{3} \overline{\mathcal{C}}_0 = n - N - K^T L^{-1} K . \quad (2.13)$$

Now suppose we change parameters so that some of the symmetries are broken. We can characterize the unbroken symmetries by a vector v^{α} , so that the unbroken symmetries satisfy $t^T v = 0$. The modified extremization is then easily carried out with the aid of a Lagrange multiplier s :

$$\frac{1}{3} \overline{\mathcal{C}}_v(t, s) = n - N + 2t^T K + t^T L t + 2st^T v . \quad (2.14)$$

Extremizing with respect to t leads to

$$\frac{1}{3}\overline{C}_v(s) = \frac{1}{3}\overline{C}_0 - 2sv^T L^{-1}K - s^2 v^T L^{-1}v . \quad (2.15)$$

This may be extremized for s if and only if $v^T L^{-1}v \neq 0$, in which case we obtain

$$\frac{1}{3}\overline{C}_v = \frac{1}{3}\overline{C}_0 + \frac{(v^T L^{-1}K)^2}{v^T L^{-1}v} . \quad (2.16)$$

The first observation is that the deformation changes the IR central charge if and only if the original symmetry, with charges determined from $t_* = -L^{-1}K$ is broken. Next, we see that if in addition v belongs to the positive eigenspace of L , then the central charge of the deformed theory is strictly greater than that of the undeformed theory — this means the deformation must be irrelevant in the IR, and we expect the deformed theory to flow to the original undeformed fixed point. Once we eliminate these irrelevant deformations from the parameter space, the symmetries corresponding to the positive eigenvalues of L are never broken, and we can restrict to v in the negative eigenspace of L .

We stress that in all examples we considered L is negative definite. In that case (2.16) shows that when a deformation breaks a symmetry the central charge changes if and only if the deformation breaks the R-symmetry, and whenever that happens the central charge decreases.

Constraints on UV data

The structure relating UV and IR physics sketched above assumes that for a given set of charges (q_i, Q_A) there exists a non-singular potential with a zero-dimensional ideal \mathbf{J} and of course that $U(1)_L$ and $U(1)_R^0$ are the only symmetries all along the RG flow. Both of these are non-trivial assumptions.

It is an open problem to classify all sets of charges consistent with (2.7) and some fixed \overline{c} that can be realized by a non-singular \mathbf{J} .⁶ Demanding that $\chi(y)$ is a polynomial rules out many choices of charges, but while being a necessary condition, it is not sufficient to show that there exists a zero-dimensional \mathbf{J} realizing the charge assignment.

The second assumption, which amounts to the statement that there are no accidental symmetries in the IR, also leads to some necessary conditions. For instance, just as in $N=1$ $d=4$ SQCD [24], violation of unitarity bounds on the charges can indicate an inconsistency in the assumption. In particular, we have the unitarity bounds

$$0 < q_i \leq \overline{c}/3 , \quad 0 < (1 + Q_A) \quad \sum_A (1 + Q_A) \leq \overline{c}/3 . \quad (2.17)$$

These arise by demanding that ϕ_i , γ^A , and $\prod_A \gamma^A$ are chiral primary operators of a unitary $N=2$ superconformal algebra. The latter is particularly strong and eliminates many possible

⁶This should be contrasted with (2,2) LG models, for which such a classification exists [22, 23] and yields a finite set of quasi-homogeneous potentials at fixed central charge.

candidate charges.⁷ While these criteria are important and will certainly play a role in any attempt to classify (0,2) LG theories, they are not sufficient to rule out accidents.

3 Accidents

Having reviewed the basic structure of (0,2) LG theories, we will now study it in a few examples that will illustrate some of the subtleties in their analysis.

3.1 Accidents in (2,2) Landau-Ginzburg orbifolds

There are two familiar examples of accidental symmetries in (2,2) flows. Consider LG orbifolds with potentials

$$W_3 = X_1^3 + X_2^3 + X_3^3 - \psi X_1 X_2 X_3, \quad W_4 = X_1^4 + X_2^4 + X_3^4 + X_4^4 - \psi X_1 X_2 X_3 X_4, \quad (3.1)$$

For W_3 (W_4) we take the orbifold by $\mathbb{Z}_3 \subset U(1)_R$ ($\mathbb{Z}_4 \subset U(1)_R$). The endpoint of the flow in each case has accidental symmetries. In the case of W_3 , which is a special point in the moduli space of a (2,2) compactification on T^2 , there is an accidental N=2 Kac-Moody $U(1)$ algebra for both left and right movers, corresponding to the isometries of the torus. In the case of W_4 the IR theory is actually a (4,4) SCFT, and there are additional currents that enhance $U(1)_L \times U(1)_R$ to $SU(2)_L \times SU(2)_R$. Of course this is the case for any Landau-Ginzburg orbifold (or more generally linear sigma model) that corresponds to a locus in the moduli space of T^2 or K3 compactification.

3.2 A contrived (2,2) example

Consider a (2,2) LG theory with

$$W = X^3 + Y^4. \quad (3.2)$$

There is a unique assignment of R-charge $\bar{q}(X) = 1/3$ and $\bar{q}(Y) = 1/4$, and the IR fixed point is the E_6 minimal model. On the other hand, we can make a field redefinition $\tilde{X} = X - Y$ and $\tilde{Y} = Y$. This is certainly non-singular and leads to a superpotential

$$\tilde{W} = \tilde{X}^3 + 3\tilde{X}^2\tilde{Y} + 3\tilde{X}\tilde{Y}^2 + \tilde{Y}^3 + \tilde{Y}^4. \quad (3.3)$$

If we also perform the field redefinition in the kinetic terms, we have of course done nothing; however, if we assume the D-terms are indeed irrelevant, then taking standard kinetic terms and either W or \tilde{W} interactions should lead to the same IR fixed point. Unlike the original theory, the \tilde{W} theory has no manifest R-symmetry along the flow — the symmetry emerges accidentally in the IR.

⁷In the (2,2) case this translates to the known bound $\sum_i q_i \leq n/3$ [25].

The example is very contrived, but it illustrates the basic issue: field redefinitions can obscure the UV fields that should be identified with IR operators of some fixed scaling dimension. As we show in 3.5, if we restrict to (2,2) theories with a quasi-homogeneous potential, this ambiguity turns out to be harmless. As the next example shows, in (0,2) theories this is not the case.

3.3 A simple (0,2) example

Consider a theory with $N = 3$, $n = 2$ and superpotential

$$\mathcal{W}_0 = \begin{pmatrix} \Gamma^1 & \Gamma^2 & \Gamma^3 \end{pmatrix} \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} \\ \alpha_{21} & \alpha_{22} & \alpha_{23} \\ \alpha_{31} & \alpha_{32} & \alpha_{33} \end{pmatrix} \begin{pmatrix} \Phi_1^6 \\ \Phi_2^2 \\ \Phi_1^3 \Phi_2 \end{pmatrix}. \quad (3.4)$$

For generic values of the 9 parameters α the potential preserves a unique $U(1)_L$ symmetry, and normalizing the charges as in (2.6) leads to $r = 2$, $\bar{c} = 3$, and charge assignments

$$\begin{array}{ccc} \Phi_1 & \Phi_2 & \Gamma^{1,2,3} \\ \mathbf{q} & \frac{1}{7} & \frac{3}{7} & -\frac{6}{7} \\ \bar{\mathbf{q}} & \frac{1}{7} & \frac{3}{7} & \frac{1}{7} \end{array} \quad (3.5)$$

To obtain a description of the parameter space of the IR theory we consider the α modulo field redefinitions consistent with (0,2) SUSY and the $U(1)_L$ symmetry:

$$\Gamma^A \rightarrow \sum_B \Gamma^B M_B^A, \quad \Phi_1 \rightarrow x\Phi_1, \quad \Phi_2 \rightarrow y\Phi_2 + z\Phi_1^3. \quad (3.6)$$

These transformations are invertible if and only if $M \in GL(3, \mathbb{C})$ and $x, y \in \mathbb{C}^*$. The induced action on the Φ monomials is then

$$\begin{pmatrix} \Phi_1^6 \\ \Phi_2^2 \\ \Phi_1^3 \Phi_2 \end{pmatrix} \rightarrow S \begin{pmatrix} \Phi_1^6 \\ \Phi_2^2 \\ \Phi_1^3 \Phi_2 \end{pmatrix}, \quad S = \begin{pmatrix} x^6 & 0 & 0 \\ x^3 z & x^3 y & 0 \\ z^2 & 2yz & y^2 \end{pmatrix}, \quad (3.7)$$

and hence the action on the parameters α is $\alpha \mapsto M\alpha S$.

A bit of algebra shows that every non-singular ideal \mathbf{J} described by α is equivalent by a field redefinition to one of three superpotentials:

$$\begin{aligned} \mathcal{W}_1 &= \Gamma^1 \Phi_1^6 + \Gamma^2 \Phi_2^2 + \Gamma^3 \Phi_1^3 \Phi_2, \\ \mathcal{W}_2 &= \Gamma^1 (\Phi_1^6 + \Phi_2^2) + \Gamma^2 \Phi_1^3 \Phi_2, \\ \mathcal{W}_3 &= \Gamma^1 \Phi_1^6 + \Gamma^2 \Phi_2^2. \end{aligned} \quad (3.8)$$

The UV parameter space is stratified to three points, and we consider each in turn.

1. \mathcal{W}_1 has a $U(1)^2$ global symmetry that acts independently on Φ_1 and Φ_2 ; extremization picks out the following charges.

	θ	Φ_1	Φ_2	Γ^1	Γ^2	Γ^3	
\mathbf{q}	0	$\frac{26}{167}$	$\frac{64}{167}$	$-\frac{156}{167}$	$-\frac{128}{167}$	$-\frac{142}{167}$	$\bar{c} = 3 \left(1 + \frac{2}{167}\right)$
$\bar{\mathbf{q}}$	1	$\frac{26}{167}$	$\frac{64}{167}$	$\frac{11}{167}$	$\frac{39}{167}$	$\frac{25}{167}$	

2. \mathcal{W}_2 has a free Γ^3 multiplet. The interacting part of the theory has no extra global symmetries and $U(1)_L^{\text{IR}} \times U(1)_R^{\text{IR}}$ charges

	θ	Φ_1	Φ_2	Γ^1	Γ^2	
\mathbf{q}	0	$\frac{4}{31}$	$\frac{12}{31}$	$-\frac{24}{31}$	$-\frac{24}{31}$	$\bar{c} = 3 \left(1 + \frac{1}{31}\right)$.
$\bar{\mathbf{q}}$	1	$\frac{4}{31}$	$\frac{12}{31}$	$\frac{7}{31}$	$\frac{7}{31}$	

3. \mathcal{W}_3 has a free Γ^3 multiplet, and the interacting part of the theory is a product of (2,2) minimal models with (2,2) superpotential $W = X_1^7 + X_2^3$ and charges

	θ	Φ_1	Φ_2	Γ^1	Γ^2	
\mathbf{q}	0	$\frac{1}{7}$	$\frac{1}{3}$	$-\frac{6}{7}$	$-\frac{2}{3}$	$\bar{c} = 3 \left(1 + \frac{1}{21}\right)$.
$\bar{\mathbf{q}}$	1	$\frac{1}{7}$	$\frac{1}{3}$	$\frac{1}{7}$	$\frac{1}{3}$	

If we assume that there are no accidental symmetries for the \mathcal{W}_1 , \mathcal{W}_2 and \mathcal{W}_3 theories, we obtain a consistent picture of the RG flows starting with the UV theory in (3.4). There are three basins of attraction; each has a central charge $\bar{c} > 3$, a set of charges consistent with unitarity bounds and no marginal deformations. Moreover, we can construct interpolating RG flows $\mathcal{W}_3 \rightarrow \mathcal{W}_2 \rightarrow \mathcal{W}_1$ by adding relevant deformations to the superpotentials. However, \mathcal{W}_1 has no $U(1)_L$ -invariant relevant deformations that make it flow to a putative $\bar{c} = 3$ theory described by \mathcal{W}_0 .

We conclude that (0,2) LG RG flows have accidental symmetries, and identifying these is key in order to correctly pinpoint even basic properties of the IR theory. For instance, we see in the example at hand that no point in the UV parameter space leads to an IR theory with $\bar{c} = 3$ and $r = 2$.

3.4 Puzzles from enhanced symmetries

There are two questions that probably occur to our erudite reader. First, what's the big deal? One has to take account of field redefinitions when discussing the parameter space of a theory, and it seems that all we learned here is that the parameter space is smaller than one may have naively thought. Second, is it not perverse to discover some accidental symmetries

associated to $\mathcal{W}_{1,2,3}$ versus \mathcal{W}_0 but then blithely assume that $\mathcal{W}_{1,2,3}$ do not themselves suffer from accidents?

There is a pragmatic answer to the second question: we assume there are no accidents unless we are able to identify some paradox in the putative description of the IR physics in terms of the UV parameters. In our example we find such a paradox: while a generic \mathcal{W} has a unique global symmetry in the UV, there are special points with enhanced symmetries and a central charge that exceeds the putative $\bar{c} = 3$ of the generic \mathcal{W} ! Once we take into account the accidental symmetries, we discover that the enhanced symmetries are unavoidable, and there is no $\bar{c} = 3$ theory that can be reached within the parameter space of these UV theories. It is also easy to construct paradoxical examples that would violate unitarity bounds unless one takes accidents into account [16].

The answer to the first question is contained in this pragmatic perspective. The “big deal” is that in the examples with which we are most familiar, namely the (2,2) LG theories, one never encounters these enhanced symmetry puzzles: although there are plenty of points with enhanced symmetries, these never mix with $U(1)_{\mathbb{R}}^{\text{IR}}$, and the central charge does not jump for any choice of non-singular (2,2) superpotential. We discuss this in detail in the next section.

3.5 Enhanced symmetries of (2,2) LG theories

Consider a (2,2) LG theory with a quasi-homogeneous (2,2) superpotential $W(X)$ obeying $W(t^{q_i} X_i) = tW(X)$. Unless W satisfies an independent quasi-homogeneity condition, the (2,2) R-symmetries are fixed uniquely, giving charge $\bar{q} = q_i$ to Φ_i and Γ^i , where (Φ_i, Γ^i) are the (0,2) components of the (2,2) multiplet X_i . Without loss of generality we can restrict attention to $0 < q_i < 1/2$.⁸ A special case occurs when we can split the fields $\{X_i\} \rightarrow \{X_a\} \cup \{X_p\}$ so that $W = W^1(X_a) + W^2(X_p)$. This leads to an enhanced symmetry, but the enhancement is very large: on both the left and right we obtain two $N = 2$ superconformal algebras with \bar{c}_1 and \bar{c}_2 that add up to the total \bar{c} . The enhanced right-moving $U(1)$ symmetry is not part of an $N=2$ Kac-Moody algebra: there are two commuting $N=2$ superconformal algebras, and each $U(1)$ is the lowest component of a different $N = 2$ algebra. Thinking of this theory as a (0,2) LG model and carrying out c -extremization leads to the same result for \bar{c} and charges of the chiral fields.

We will now show that in non-singular (2,2) theories this is the only way that enhanced symmetries occur. Hence, there are no (2,2) accidents.

A necessary and sufficient condition to be able to perform the split $\{X_i\} \rightarrow \{X_a\} \cup \{X_p\}$ and $W = W^1(X_a) + W^2(X_p)$ is that the matrix of second derivatives W_{ki} is block diagonal in the two sets of variables.⁹ Since we understand the symmetry enhancement in that case, we

⁸We assume $q_i > 0$. In that case for a non-singular potential any fields with $q_i \geq 1/2$ can be eliminated by their equations of motion.

⁹We use the shorthand $W_i = \partial W / \partial X_i$, $W_{ki} = \partial W / \partial X_k \partial X_i$, etc.

assume that W_{ki} has no non-trivial block.¹⁰ We will now show that no additional symmetry is possible when W is non-singular. The argument uses three facts.

1. A non-singular W can satisfy at most one linearly independent quasi-homogeneous relation. To see this, suppose the contrary. By taking linear combinations of two relations we arrive at $\sum_i \alpha_i X_i W_i = 0$. We can now split the fields X_i according to $\alpha_i > 0$, $\alpha_i < 0$, or $\alpha_i = 0$: $\{X_i\} \rightarrow \{Y_a\} \cup \{Z_s\} \cup \{U_\alpha\}$ and recast the relation as

$$\sum_a \beta_a Y_a W_a = \sum_s \gamma_s Z_s W_s , \quad (3.9)$$

where $\beta_a, \gamma_m > 0$. Without loss of generality we may assume $\beta_1 = 1 \geq \beta_a$ for $a \neq 1$. Every monomial in W that contains Y_1 must contain at least one Z . Hence, W will be singular unless $W \supset Y_1^m Z_s$ for some s , say $s = 1$. Similarly, $dW|_{Y=0}$ will be independent of Z_1 unless $W \supset Z_1^p Y_a$ for some a , which requires $\beta_a = \gamma_1 p = pm > 1$, where the last inequality follows since W has no quadratic terms in the fields. That is in contradiction with $\beta_a \leq 1$, so the theory must be singular.

2. Suppose we have a symmetry of the (2,2) theory that commutes with the (0,2) SUSY algebra. This means that there are charges Q'_i and q'_i such that

$$-Q'_i W_i = \sum_j q'_j X_j W_{ij} \quad \implies \quad -Q'_i W_{ik} = q'_k W_{ki} + \sum_j q_j X_j W_{jik} .$$

Exchanging i and k in the second equation and taking the difference, we obtain

$$(Q'_k - q'_k) W_{ki} = W_{ki} (Q'_i - q'_i) .$$

This means that whenever $W_{ki} \neq 0$ we need $Q'_k - q'_k = Q'_i - q'_i$.¹¹

3. The (2,2) superpotential satisfies

$$(q'_i - Q'_i) W - \sum_j q'_j X_j W_j = U^i ,$$

where U^i is independent of X^i . This follows by integrating the quasi-homogeneity condition obeyed by W_i .

Using these observations, we now complete the argument as follows. Since W_{ki} does not contain a non-trivial block, we see from the second fact that for all k, i $Q'_k - q'_k = Q'_i - q'_i$. Combining this with the third fact, we find that W satisfies a quasi-homogeneity relation $W(t^{q'_j} X_j) = t^{q'_i - Q'_i} W(X)$; the first fact then implies that either $q'_i = cq_i$ and $q'_i - Q'_i = c$, or W is singular.

¹⁰Take the $n \times n$ matrix W_{ki} and set to 1 all non-zero components. The result is a symmetric matrix A_{ki} that is the adjacency matrix for a graph G on n nodes, with each $A_{ki} \neq 0$ specifying a path in the graph from node k to node i . The statement that there is no non-trivial block is simply that G is connected.

¹¹This is trivially satisfied for the usual (2,2) $U(1)_L$, where $Q_i = q_i - 1$.

3.6 Subtleties for heterotic vacua

We have seen that the identification of UV parameters with a deformation space of an IR CFT, while reasonably well understood for (2,2) theories, is more subtle for (0,2) theories. The difference is that while in non-singular (2,2) theories enhanced symmetries are always associated to a decomposition of the UV theory into non-interacting components, this is not the case for (0,2) models. An enhanced symmetry of a (0,2) model does generically mix with the naive $U(1)_R$, so that the enhanced symmetry point has a different central charge from what one might expect naively. As illustrated by the example in section 3.3, the RG fixed points of a (0,2) model need not realize any CFT with the naive central charge.

There are situations where the consequences are more benign: there is a choice of UV parameters that leads to a CFT with the expected IR symmetries, but even then the identification of UV parameters with marginal deformations of the IR theory requires a careful study of the field redefinition orbits on the space of UV parameters. The following familiar example illustrates the issue.

An $SO(10)$ heterotic Landau-Ginzburg orbifold

Consider a (0,2) theory with the following field content and charge assignment

$$\begin{array}{ccccc}
 & \theta & \Phi_{1,2} & \Phi_{3,\dots,6} & \Gamma^{1,\dots,7} \\
 \mathbf{q} & 0 & \frac{2}{5} & \frac{1}{5} & -\frac{4}{5} \\
 \bar{\mathbf{q}} & 1 & \frac{2}{5} & \frac{2}{5} & \frac{1}{5}
 \end{array} \tag{3.10}$$

It is easy to see that this symmetry leads to $r = 4$ and $\bar{c} = 9$. The orbifold of this theory by $e^{2\pi i J_0}$ is a candidate for an internal SCFT of an $SO(10)$ heterotic vacuum. As described in [10] that does seem to be the case: the massless spectrum is organized into sensible $SO(10)$ multiplets, and there is a reasonable large radius interpretation in terms of a rank 4 holomorphic bundle on a complete intersection CY manifold in $\mathbb{C}\mathbb{P}_{111122}^5$. The generic superpotential for this theory is

$$\mathcal{W} = \sum_A \Gamma^A J_A, \tag{3.11}$$

where each J_A has charge $\mathbf{q} = 4/5$. We can choose the UV parameters of the theory to produce the following non-singular potential:

$$\mathcal{W}_1 = \Gamma^1 \Phi_1^2 + \Gamma^2 \Phi_2^2 + \sum_{i=3}^6 \Gamma^i \Phi_i^4 + \Gamma^7 \times 0. \tag{3.12}$$

This is a product of (2,2) minimal models and a free left-moving fermion. The resulting central charge is $\bar{c} = 3(3 + \frac{1}{15})$. Thus, this choice of UV parameters does not correspond to a point in the moduli space of the $\bar{c} = 9$ CFTs. Of course the orbit of field redefinitions of

this point yields a large basin of attraction of UV theories that flow to the same CFT with $\bar{c} = 3(3 + \frac{1}{15})$. In this case we can identify another point that does lead to $\bar{c} = 9$:

$$\mathcal{W}_2 = \Gamma^1 \Phi_1^2 + \Gamma^2 \Phi_2^2 + \sum_{i=3}^6 \Gamma^i \Phi_i^4 + \Gamma^7 \Phi_1 \Phi_2 . \quad (3.13)$$

While this superpotential still has a $U(1)^6$ global symmetry, c -extremization leads to $\bar{c} = 9$ and R-charges as in the table above. Clearly there is a relevant deformation by $\Gamma^7 \Phi_1 \Phi_2$ that leads to an RG flow from the $\bar{c} = 3(3 + \frac{1}{15})$ theory to the $\bar{c} = 9$ CFT.

The general lesson is clear: field redefinitions stratify the space of UV parameters into orbits, and in general these orbits correspond to different IR fixed points that are not related by marginal deformations — in particular they can have different central charges. The orbits may or may not include an IR fixed point for which the manifest symmetry of the generic superpotential becomes the $U(1)_{\mathbb{R}}^{\text{IR}}$: in this example they do, while in that of section 3.3 they do not.

4 Marginal deformations of a unitary (2,0) SCFT

This section contains a number of results on (2,0) SCFTs. Many if not all of these are well-known in the context of heterotic compactifications, but the derivations given here are more general and give a useful alternative perspective.

4.1 Basic results

Consider a unitary compact (2,0) SCFT with the usual superconformal algebra generators $J(z)$, $G^\pm(z)$, and $T(z)$, with modes given respectively by J_n , G_r^\pm and L_n .¹²

We will show that marginal Lorentz-invariant and supersymmetric deformations of this theory by a local operator take the form

$$\Delta S = \int d^2 z \Delta \mathcal{L} , \quad \Delta \mathcal{L} = \{G_{-1/2}^-, \mathcal{U}\} + \text{h.c.} , \quad (4.1)$$

where \mathcal{U} is a chiral primary operator with $U(1)_L$ -charge $\mathbf{q} = 1$ and weights $(h, \bar{h}) = (1/2, 1)$.¹³ In string theory, where one considers (0,2) SCFTs with quantized \mathbf{q} charges, this is a classic result [26]. Here we will apply the point of view developed for $N = 1$ $d = 4$ SCFTs [27] to arrive at the statement without any assumptions of charge integrality.

Constraints from supersymmetry

Without loss of generality we can consider deformations $\delta \mathcal{L} = \mathcal{O}(z, \bar{z})$ by a quasi-Virasoro primary operator \mathcal{O} , since a descendant would just be a total derivative. Lorentz invariance

¹²While for many purposes it is very convenient to treat the supersymmetric side of the theory as anti-holomorphic, in the discussion that follows it leads to a great profusion of bars. Hence, in this section the SUSY side will be taken to be holomorphic.

¹³Some of the arguments given here were developed by IVM and MRP in collaboration with Ido Adam.

requires \mathcal{O} to have spin 0, i.e. $h_{\mathcal{O}} = \bar{h}_{\mathcal{O}}$. In order for $\delta\mathcal{L}$ to be supersymmetric, we need $[G_{-1/2}^{\mp}, \mathcal{O}]$ to be a total derivative, i.e. $G_{-1/2}^{\mp}|\mathcal{O}\rangle = L_{-1}|\mathcal{M}^{\mp}\rangle$. Applying $G_{-1/2}^{\pm}$ to both sides of the equation and using the N=2 algebra, we obtain

$$\begin{aligned} G_{-1/2}^{\pm}G_{-1/2}^{\mp}|\mathcal{O}\rangle &= L_{-1}G_{-1/2}^{\pm}|\mathcal{M}^{\mp}\rangle, \\ \implies L_{-1}\left[|\mathcal{O}\rangle - G_{-1/2}^{+}|\mathcal{M}^{-}\rangle - G_{-1/2}^{-}|\mathcal{M}^{+}\rangle\right] &= 0. \end{aligned} \quad (4.2)$$

Hence, up to a constant multiple of the identity operator, which does not lead to a deformation of the theory, we can write $|\mathcal{O}\rangle$ as

$$|\mathcal{O}\rangle = G_{-1/2}^{-}|\mathcal{M}^{+}\rangle + G_{-1/2}^{+}|\mathcal{M}^{-}\rangle, \quad (4.3)$$

and hence, without loss of generality, any non-trivial deformation corresponds to a state

$$|\mathcal{O}\rangle = G_{-1/2}^{-}|\mathcal{U}\rangle + G_{-1/2}^{+}|\mathcal{V}\rangle + \left[G_{-1/2}^{+}G_{-1/2}^{-} - \left(1 + \frac{q_{\mathcal{K}}}{2h_{\mathcal{K}}}\right)L_{-1}\right]|\mathcal{K}\rangle, \quad (4.4)$$

where $|\mathcal{U}\rangle$, $|\mathcal{V}\rangle$ and $|\mathcal{K}\rangle$ are all quasi-primary with respect to the N=2 superconformal algebra, i.e. annihilated by the lowering modes of the global N=2 algebra, L_1 and $G_{1/2}^{\pm}$. The linear combination of operators in the last term is fixed by $L_1|\mathcal{O}\rangle = 0$. The spins of the fields are

$$\bar{h}_{\mathcal{U}} - h_{\mathcal{U}} = 1/2, \quad \bar{h}_{\mathcal{V}} - h_{\mathcal{V}} = 1/2, \quad \bar{h}_{\mathcal{K}} - h_{\mathcal{K}} = 1. \quad (4.5)$$

The remaining constraints from supersymmetry are

$$G_{-1/2}^{+}G_{-1/2}^{-}|\mathcal{U}\rangle = L_{-1}|X\rangle, \quad G_{-1/2}^{-}G_{-1/2}^{+}|\mathcal{V}\rangle = L_{-1}|Y\rangle \quad (4.6)$$

for some states $|X\rangle$ and $|Y\rangle$. We will now show that the only solution to these equations is that \mathcal{U} (\mathcal{V}) is a chiral primary (anti-chiral primary) state. It suffices to work out the constraint on $|\mathcal{U}\rangle$ — the one on $|\mathcal{V}\rangle$ follows by exchanging G^{+} and G^{-} .

Without loss of generality we decompose

$$|X\rangle = a|\mathcal{U}\rangle + |\chi\rangle, \quad (4.7)$$

where a is real and $|\chi\rangle$ is orthogonal to $|\mathcal{U}\rangle$. The condition now becomes

$$(G_{-1/2}^{+}G_{-1/2}^{-} - aL_{-1})|\mathcal{U}\rangle = L_{-1}|\chi\rangle. \quad (4.8)$$

Applying $\langle\mathcal{U}|L_1$ to both sides and using orthogonality of $|\mathcal{U}\rangle$ and $|\chi\rangle$, we find $a = 1 + \frac{q_{\mathcal{U}}}{2h_{\mathcal{U}}}$. Application of $\langle\chi|L_1$ to both sides shows $L_{-1}|\chi\rangle = 0$, so we are left with

$$G_{-1/2}^{+}G_{-1/2}^{-}|\mathcal{U}\rangle = \left(1 + \frac{q_{\mathcal{U}}}{2h_{\mathcal{U}}}\right)L_{-1}|\mathcal{U}\rangle. \quad (4.9)$$

Finally, applying $G_{-1/2}^-$, we find

$$(1 + \frac{1}{2h_{\mathcal{U}}})(2h_{\mathcal{U}} - \mathbf{q}_{\mathcal{U}}) = 0 . \quad (4.10)$$

The only solution of this equation consistent with unitarity is $2h_{\mathcal{U}} = \mathbf{q}_{\mathcal{U}}$, i.e. $|\mathcal{U}\rangle$ is a chiral primary state of the N=2 superconformal algebra.

Combining the preceding results and applying them to deformations by real operators, we conclude that real Lorentz-invariant supersymmetric deformations take the form

$$\mathcal{O}(z, \bar{z}) = \left[\{G_{-1/2}^-, \mathcal{U}(z, \bar{z})\} + \text{h.c.} \right] + \{G_{-1/2}^+, [G_{-1/2}^-, \mathcal{K}(z, \bar{z})]\} , \quad (4.11)$$

where \mathcal{U} is a fermionic chiral primary operator with $\bar{h}_{\mathcal{U}} = \frac{1}{2} + h_{\mathcal{U}}$, $h_{\mathcal{U}} = \mathbf{q}_{\mathcal{U}}/2$, and \mathcal{K} is a real bosonic quasi-primary operator with $\bar{h}_{\mathcal{K}} = 1 + h_{\mathcal{K}}$. As in four dimensions [27], we recognize the familiar superpotential and Kähler deformations.

Marginal operators

If we impose in addition that the perturbation is marginal, we obtain the constraints $\mathbf{q}_{\mathcal{U}} = 1$ and $h_{\mathcal{K}} = 0$. The latter implies that $L_{-1}|\mathcal{K}\rangle = 0$, i.e. $\mathcal{K}(\bar{z})$ is an anti-holomorphic conserved current that leads to a trivial deformation of the action. We arrive at the result (4.1).

4.2 A few consequences

The preceding analysis, when combined with some basic assumptions about superconformal perturbation theory, leads to important constraints on (2,0) SCFTs. The key feature is that we can use a (2,0) superspace to recast the marginal deformations into the form

$$\Delta S = \int d^2z \int d\theta \alpha^i \mathcal{U}_i + \text{h.c.} , \quad (4.12)$$

where α^i denote the couplings and \mathcal{U}_i are denote the chiral primary marginal fermi superfields. Assuming there exists a manifestly supersymmetric regularization scheme for conformal perturbation theory, the renormalized action at a renormalization scale μ must take the form

$$\begin{aligned} \Delta \mathcal{L}_{\text{ren}} = & \int d^2\theta \left[Z^a(\alpha, \bar{\alpha}; \mu) \bar{\mathcal{J}}_a + \sum_A \mu^{2-d_A} \bar{K}_A \right] \\ & + \left\{ \int d\theta [(\alpha_i + \delta\alpha^i(\alpha; \mu)) \mathcal{U}_i + \zeta^I(\alpha; \mu) \mathcal{U}_I] + \text{h.c.} \right\} . \end{aligned} \quad (4.13)$$

At the conformal point ($\alpha = 0$) the $\bar{\mathcal{J}}_a$ and \bar{K}_A are real operators of dimension $\Delta_a = 2$ and $\Delta_A > 2$, while the \mathcal{U}_i and \mathcal{U}_I are chiral primary operators with $\mathbf{q} = 1$ and $\mathbf{q} > 1$ respectively.¹⁴ The first line is parallels the $N = 1$ $d = 4$ situation; however, the second line is new, following from the fact that the $\mathcal{U}_i \mathcal{U}_j$ OPE will in general have singular \bar{z} dependence.

¹⁴Compactness of the CFT ensures a gap in dimensions between $\bar{\mathcal{J}}_a$ and \bar{K}_A , as well as between \mathcal{U}_i and \mathcal{U}_I .

Of course supersymmetry still requires that the renormalization of the superpotential should be holomorphic in the parameters.

Marginal irrelevance

Marginal deformations preserve the R-symmetry of the original SCFT. Hence, the unitarity bound $h_{\mathcal{U}} \geq \mathbf{q}/2$ implies that a marginal deformation is at worst marginally irrelevant and never marginally relevant.

D-terms and F-terms

Assuming that conformal perturbation theory is renormalizable, the terms involving the \overline{K}_A and \mathcal{U}_I do not arise, and scale invariance of the theory is equivalent to

$$D^a(\alpha, \overline{\alpha}) \equiv \mu \frac{\partial}{\partial \mu} Z^a = 0 \quad \text{and} \quad \overline{F}^i(\alpha) \equiv \mu \frac{\partial}{\partial \mu} \delta \alpha^i = 0. \quad (4.14)$$

A two-dimensional unitary compact scale-invariant theory is automatically conformal [28], so every deformation satisfying these ‘‘D-term’’ and ‘‘F-term’’ constraints is exactly marginal.

The ‘‘D-term’’ obstructions to marginality are exactly the same as in the $d = 4$ case studied in [27] — such a scale dependence requires the breaking of a global right-moving symmetry. This is easy to understand at leading order in conformal perturbation theory. In the presence of abelian currents \overline{J}_a , the OPE of \mathcal{U} with its conjugate takes the form

$$\mathcal{U}_i(z, \overline{z}) \overline{\mathcal{U}}_{\overline{J}}(w, \overline{w}) \sim \frac{g_{i\overline{J}}}{(z-w)(\overline{z}-\overline{w})^2} + \frac{g_{i\overline{J}} q_a^i \overline{J}^a(w, \overline{w})}{(z-w)(\overline{z}-\overline{w})} + \dots, \quad (4.15)$$

where $\overline{J}^a = \gamma^{ab} \overline{J}_b$, and $z\overline{z} \langle \overline{J}_a(\overline{z}) \overline{J}_b(0) \rangle = \gamma_{ab}$ in the undeformed theory. This leads to a logarithmic divergence in conformal perturbation theory proportional to

$$\int d^2w G_{-1/2}^+ G_{-1/2}^- \int d^2z \frac{|\alpha^i|^2 q_a^i \overline{J}^a(w, \overline{w})}{(z-w)(\overline{z}-\overline{w})} \sim \log \mu \times |\alpha^i|^2 q_a^i \overline{J}^a(w, \overline{w}), \quad (4.16)$$

which corresponds to the leading order D-term proportional to

$$D_a = \sum_i |\alpha^i|^2 q_a^i. \quad (4.17)$$

In applications to heterotic compactifications such a symmetry necessarily corresponds to a gauge boson in the space-time theory, and the space-time picture of the breaking is just the Higgs mechanism: the obstruction to marginality of a coupling α that breaks a right-moving symmetry is encoded in a space-time D-term potential.

We believe that in (0,2) LG models without an orbifold there are no F-term obstructions. The reason is simple: the free field UV presentation of the theory comes with the usual non-renormalization theorems for the superpotential, so the only divergences we expect to encounter will correspond to D-term counter-terms.

The two sources of obstruction are in one to one correspondence with the two ways in which a short chiral primary multiplet can combine into a long multiplet of (2,0) SUSY. Suppose we consider an infinitesimal (2,0) SUSY deformation under which a marginal chiral primary state $|\mathcal{U}\rangle$ acquires weights $(h, \bar{h}) = (\frac{1}{2} + \frac{\epsilon}{2}, 1 + \frac{\epsilon}{2})$. In this case $|\mathcal{U}\rangle$ is no longer chiral primary, and by a choice of basis we can consider two separate cases:

$$G_{-1/2}^+|\mathcal{U}\rangle \neq 0, \quad G_{1/2}^-|\mathcal{U}\rangle = 0, \quad \text{or} \quad G_{-1/2}^+|\mathcal{U}\rangle = 0, \quad G_{1/2}^-|\mathcal{U}\rangle \neq 0. \quad (4.18)$$

In other words, $|\mathcal{U}\rangle$ remains primary but is no longer chiral, or it remains chiral but fails to be primary. The first case corresponds to an F-term obstruction, where at $\epsilon = 0$ we have two chiral primary superfields $(\mathcal{U}, \mathcal{F})$ with $\mathbf{q}_{\mathcal{U}} = 1$ and $\mathbf{q}_{\mathcal{F}} = 2$ and $\bar{h} = 1$, while for $\epsilon > 0$ we find a complex long multiplet with lowest component $|\mathcal{U}\rangle$ and $G_{-1/2}^+|\mathcal{U}\rangle = \sqrt{\epsilon}|\mathcal{F}\rangle$.¹⁵ The second case corresponds to a D-term obstruction, where at $\epsilon = 0$ we have chiral primary superfields \mathcal{U} , its anti-chiral conjugate $\bar{\mathcal{U}}$, and a Kac-Moody current $\bar{\mathcal{J}}$; for $\epsilon > 0$ we obtain a long real multiplet with lowest component $\bar{\mathcal{J}}$ and descendants $G_{-1/2}^+|\bar{\mathcal{J}}\rangle = \sqrt{\epsilon}|\mathcal{U}\rangle$ and $G_{-1/2}^-|\bar{\mathcal{J}}\rangle = \sqrt{\epsilon}|\bar{\mathcal{U}}\rangle$.

In particular, we see that there are no F-term obstructions if the undeformed theory has no chiral primary operators with $\mathbf{q} = 2$ and $\bar{h} = 1$. This is the case, for instance, in every (2,0) SCFT with $c < 6$. If there are also no left-moving Kac-Moody symmetries then every (2,0) marginal deformation must remain exactly marginal. In appendix A we mention a simple example illustrating an F-term obstruction at $c = 9$.

Kähler geometry of the moduli space

One can use the same reasoning as in [27] to argue that the space of truly marginal deformations of a (2,0) SCFT must be a Kähler manifold. This is because the D-term constraints and the quotient by global symmetries lead to a toric quotient on the space of marginal couplings, while the F-term constraints are manifestly holomorphic constraints, restricting the truly marginal directions to a Kähler subvariety of the toric variety. In heterotic compactification this can of course be argued either from the space-time heterotic supergravity or by using additional assumptions of a (2,0) SCFT with integral charges [31]. The argument given here is more direct and general.

Application to (2,2) theories

The case of a (2,2) SCFT and its (2,2)-preserving deformations is much simpler. There are two types of superpotential deformations: the chiral and the twisted chiral. The former corresponds to deformations by chiral primary (c,c) ring operators, while the latter by the (a,c) ring operators. Supersymmetry implies that twisted chiral parameters can never show up in the renormalized chiral superpotential and vice-versa. Moreover, the OPE of the (c,c) and (a,c) chiral primaries with themselves is non-singular, so that neither potential is

¹⁵In a $c = 9$ theory with spectral flow there is a canonical \mathcal{F} for every \mathcal{U} in the theory. Indeed, as observed in [26, 29, 30], the \mathcal{F} (2,0) superfields can be used to construct vertex operators for the space-time auxiliary fields residing in chiral multiplets of the associated four-dimensional theory.

corrected—there are no F-term obstructions to marginality. Hence, all marginal (c,c) and (a,c) symmetry-preserving deformations are truly marginal. This is again a familiar story in string applications [26, 32].

Accidents beyond field redefinitions

We can now see that the field redefinitions of (0,2) LG theories do not describe all accidents. It is not the case that every direction transverse to field redefinition orbits corresponds to a marginal deformation of the IR theory. This is due to the possibility that marginal deformations of a (0,2) theory can turn out to be marginally irrelevant. In (0,2) LG theories this is due to D-term obstructions where a U(1) symmetry is broken by turning on operators with a definite sign of the U(1) charge. We give an example of this phenomenon in a well-known heterotic vacuum in appendix B.

4.3 Deformations and left-moving abelian currents

As a final application of the preceding results, we consider the interplay between deformations of a (2,0) SCFT and left-moving currents.

A (2,0) SCFT may possess a KM algebra on the SUSY side of the world-sheet in addition to the $U(1)_L$ current J_L in the $N = 2$ multiplet. Such structures are familiar from heterotic compactifications preserving 8 space-time supercharges in four dimensions — when realized geometrically these correspond to geometries $\pi : X \rightarrow K3$ — principal T^2 fibrations over a base $K3$ [33, 34]. In each such case we can use a Sugawara-like decomposition to decompose the $N = 2$ world-sheet superconformal algebra (SCA) into two commuting sets of generators, one associated to the KM algebra, and the other corresponding to the remaining degrees of freedom.

Suppose we have an abelian current algebra $U(1)$ with current J_1 . There are two ways that the decomposition can work. If J does not belong to a multiplet of the \mathcal{A}_c SCA, then we must have a decomposition

$$\mathcal{A}_c = \mathcal{A}'_{c'} \oplus \mathcal{A}''_{c''} , \quad (4.19)$$

where $c = c' + c''$, and the lowest components of the $N=2$ multiplets of \mathcal{A}' and \mathcal{A}'' are obtained by appropriate linear combinations of J_L and J_1 . We are familiar with such examples from above: this happens whenever the LG theory decomposes into a product of two non-interacting theories.

If J does belong to a multiplet of \mathcal{A} , then it must be accompanied by a second $U(1)$ Kac-Moody current J_2 , as well as weight $h = 1/2$ operators ψ_1 and ψ_2 . Together these arrange themselves into a well-known $c = 3$ unitary representation of $N = 2$:

$$J_L = : \psi \bar{\psi} : , \quad G^+ = \sqrt{2} \psi \bar{\mathcal{J}} , \quad G^- = \sqrt{2} \bar{\psi} \mathcal{J} , \quad , \quad T = : \mathcal{J} \bar{\mathcal{J}} : - \frac{1}{2} (: \bar{\psi} \partial \psi : + : \psi \partial \bar{\psi} :) , \quad (4.20)$$

where ψ and \mathbf{j} have the free-field OPEs

$$\overline{\psi}(z)\psi(w) \sim (z-w)^{-1}, \quad \overline{\mathbf{j}}(z)\mathbf{j}(w) \sim (z-w)^{-2}. \quad (4.21)$$

This is equivalent to the holomorphic sector of a T^2 (1,0) non-linear sigma model, and we will call it $\mathcal{A}_3^{\text{free}}$.

There is a key difference between these two generalizations. In the first case, there are generally deformations that can break the extra left-moving symmetry — in (2,2) LG this happens when we move away from a Gepner point to a more generic theory. In the second case such breaking is impossible. To see this, we just need to apply what we learned about the structure of SUSY deformations in conformal perturbation theory. Since our algebra splits as

$$\mathcal{A}_c = \mathcal{A}'_{c-3} \oplus \mathcal{A}_3^{\text{free}}, \quad (4.22)$$

a marginal deformation has a similar decomposition

$$\mathcal{U} = \mathcal{U}' + S\psi, \quad (4.23)$$

where \mathcal{U}' is a chiral primary operator with $\overline{h} = 1$ and $\mathbf{q}' = 1$, while S is a (0,1) current. The deformation of the action is then

$$G_{-1/2}^- \cdot \mathcal{U} = G_{-1/2}^- \cdot \mathcal{U}' + \sqrt{2}S\mathbf{j}. \quad (4.24)$$

This is neutral with respect to J_L , \mathbf{j} and $\overline{\mathbf{j}}$. More generally, any relevant deformation must be of the form $\mathcal{U} = \mathcal{U}'$ with $\mathbf{q}' < 1$.

5 Toric geometry of the deformation space

In the previous sections we saw that accidental symmetries play an important role in (0,2) Landau-Ginzburg theories, and more generally, in (0,2) SCFTs. In this section we will describe a conjecture that allows us to account for these accidents in a certain class of (0,2) LG theories. In that context our goal is to describe the moduli space \mathcal{M} of IR fixed points corresponding to a class of UV data determined by a choice of charges q_i and Q_A which have the expected central charge

$$\overline{c} = 3(n - N + r), \quad r = - \sum_A Q_A - \sum_i q_i. \quad (5.1)$$

To do so, we need to perform two steps:

1. decompose the UV parameter space into orbits under the action of field redefinitions;
2. determine which orbits contribute to \mathcal{M} .

The result is expected to be a (typically singular) Kähler space. In general these are rather formidable tasks. The group of field redefinitions is rather large and the space of orbits is non-separable. A reasonable geometry can only emerge after implementing the second task. This involves excluding two types of orbits:

- Along a *discriminant locus* Δ in parameter space, the superpotential is singular. The discriminant will clearly be invariant under field redefinitions, and orbits contained in Δ will not contribute to \mathcal{M} .
- For some non-singular values of the parameters, the theory will have accidental symmetries in the IR. As we have seen, in some cases these symmetries will mix nontrivially with the R-symmetry and the central charge of the IR fixed point will be larger than \bar{c} . Thus, these orbits as well need to be excluded from \mathcal{M} .

In general, the second step is difficult even if one restricts attention to symmetries which act diagonally on the UV fields. Detecting the basin of attraction of some component of the IR moduli space with central charge $\bar{c}' > \bar{c}$ requires a determination of the R-symmetry along each such component to find which deformations away from this locus are in fact irrelevant.

5.1 The toric conjecture

There is a simpler version of both of these problems that may be tractable. The group of field redefinitions always contains an abelian subgroup, the complexification of the $U(1)^n \times U(1)^N$ subgroup of the global symmetry of the free kinetic terms, that corresponds to rescaling the chiral fields of the theory. In particular, if we write the most general superpotential in our class as

$$\mathcal{W} = \sum_A \Gamma^A \sum_{m \in \Delta_A} \alpha_{Am} \prod_i \Phi_i^{m_i} , \quad (5.2)$$

where

$$\Delta_A = \{m \in \mathbb{Z}^n \mid \sum_i m_i q_i = -Q_A\} \quad (5.3)$$

describes the lattice points in the Newton polytope for J_A , then the field redefinitions

$$\Phi_i \mapsto t_i \Phi_i , \quad \Gamma^A \mapsto \tau_A \Gamma^A \quad (5.4)$$

lead to a $T_{\mathbb{C}} = (\mathbb{C}^*)^{N-n+1}$ action¹⁶ on the space of UV parameters $Y = \mathbb{C}^{\sum_A |\Delta_A|}$

$$\alpha_{Am} \mapsto \tau_A \prod_i t_i^{m_i} \times \alpha_{Am} . \quad (5.5)$$

We will refer to these as toric field redefinitions.

¹⁶The rank of the \mathbb{C}^* action is reduced by 1 due to the quasi-homogeneity of \mathcal{W} .

We now restrict attention to these toric actions in *both* of the tasks listed above. Namely, we decompose the parameter space into $T_{\mathbb{C}}$ orbits and exclude those orbits that either lie in Δ or exhibit accidental symmetries contained in $T_{\mathbb{C}}$ and lead to $\bar{c}' > \bar{c}$. The result, which we will call \mathcal{M}_T , will in some cases be equivalent to \mathcal{M} , but in general the two will differ. We will comment on this further below.

The action of the compact torus $T \subset T_{\mathbb{C}}$, given by restricting to $|t| = |\tau| = 1$, determines a moment map $\mu = (\lambda; \Lambda) : Y \rightarrow \mathbb{R}^{N+n}$ with

$$\lambda_i = \sum_A \sum_{m \in \Delta_A} m_i |\alpha_{Am}|^2, \quad \Lambda_A = \sum_{m \in \Delta_A} |\alpha_{Am}|^2. \quad (5.6)$$

Quasi-homogeneity of \mathcal{W} implies that the image lies in the hyperplane

$$\sum_i q_i \lambda_i + \sum_A Q_A \Lambda_A = 0. \quad (5.7)$$

The image of μ is the intersection of this hyperplane with a cone, determined by the charges, inside the positive orthant in \mathbb{R}^{N+n} . This intersection is itself a cone $\widehat{\Sigma}$, of dimension $N+n-1$. The level sets of μ determine a selection of orbits: generic orbits will be $N+n-1$ dimensional, but the action will degenerate along points with a non-trivial stabilizer subgroup, leading to orbits of smaller dimension.¹⁷ More precisely, the cone $\widehat{\Sigma}$ can be subdivided into a fan Σ , such that the collection of orbits containing a point for which $\mu(\alpha) = \mu^*$ is determined by the cone of Σ that contains μ^* . This is the *secondary fan* for the T action. We now have enough structure to state our conjecture.

Conjecture

The toric moduli space \mathcal{M}_T is the complement of the discriminant subvariety Δ in a *toric variety*

$$V = \mu^{-1}(\lambda^*, \Lambda^*)/T, \quad (5.8)$$

where

$$\lambda_i^* = 1 - q_i, \quad \Lambda_A^* = 1 + Q_A. \quad (5.9)$$

This is a rather strong statement, and we will not provide a complete proof but rather some evidence for it. Some ideas on a possible derivation are discussed in section 5.4. We will motivate the conjecture by combining our results and observations from above with some facts about toric varieties.

¹⁷There are orbifold singularities when the subgroup is discrete; we will focus on continuous stabilizer subgroups.

We should note a few important points. First, V may turn out to be empty. Second, while we claim that $V \setminus \Delta$ describes \mathcal{M}_T as a variety, we do not make any statement about the relation between the Zamolodchikov metric on the space of marginal couplings and the metric on V obtained by the Kähler quotient. Finally, in this paper we will be concerned with orbits of continuous field redefinitions. In general there will be additional discrete quotients that identify points in \mathcal{M}_T .

Combinatorics of the secondary fan

Codimension-one cones in Σ are associated with orbits containing a point at which a single $\mathbb{C}^* \subset T_{\mathbb{C}}$ is unbroken. More precisely, $G_{(q',Q')} = \mathbb{C}^* \subset T_{\mathbb{C}}$ acting with charges q'_i, Q'_A on the chiral superfields will fix points at which

$$|\alpha_{Am}|^2 (Q'_A + \sum_i m_i q'_i) = 0 \quad \text{for all } A, m \in \Delta_A . \quad (5.10)$$

The μ -image of the $T_{\mathbb{C}}$ orbits of such points will lie in a cone $\sigma_{(q',Q')}$ generated by the charge vectors of the α_{Am} fixed by $G_{(q',Q')}$. Thus, the codimension-one cones of Σ are determined by one-dimensional subgroups for which $\sigma_{(q',Q')}$ has dimension $N + n - 2$ and lies in the hyperplane

$$\sum_A Q'_A \Lambda_A + \sum_i q'_i \lambda_i = 0 . \quad (5.11)$$

In terms of \mathcal{W} the codimension-one cones of Σ correspond to subgroups for which we can write a (possibly singular) family of models fixed precisely by $(\mathbb{C}^*)^2$. Cones of higher codimension in the fan are boundaries of these cones and arise at the intersections of these hyperplanes.¹⁸ Points in the interior of some cone of the fan (of any codimension) lie in the image of a collection of orbits determined by that cone. Our conjecture is thus equivalent to the statement that the μ -image of the $T_{\mathbb{C}}$ orbits of models with central charge \bar{c} intersects the cone σ^* containing μ^* in its interior.

A cone $\sigma \in \Sigma$ can be specified by its relation to the codimension-one cones $\sigma_{(q',Q')}$. For each of these, σ either lies inside $\sigma_{(q',Q')}$, in which case $\mu \in \sigma$ satisfy (5.11), or it lies on one side or the other, meaning (5.11) is satisfied as a strict inequality for all $\mu \in \sigma$. To prove our claim we thus need to show that orbits of points in parameter space corresponding to models with central charge \bar{c} are precisely those containing in their image points satisfying the inequalities satisfied by μ^* . To do this we must consider all codimension-one cones of Σ . We classify these by the nature of the models exhibiting the enhanced symmetry.

¹⁸Note that this does not imply that cones of higher codimension correspond to models with larger unbroken symmetry: values of μ at the intersection of two codimension-one cones can be in the image of two distinct $T_{\mathbb{C}}$ orbits, each of which is fixed by a different subgroup.

5.2 Enhanced toric symmetries

Symmetries realized by a non-singular potential

Consider first the case of one-parameter subgroups of $T_{\mathbb{C}}$ for which the generic point in the locus they fix corresponds to a nonsingular model with a $U(1)^2$ global symmetry. The IR R-symmetry can then be determined by \bar{c} -extremization as

$$\hat{q}_i = tq_i + sq'_i, \quad \hat{Q}_A = tQ_A + sQ'_A, \quad (5.12)$$

where

$$L \begin{pmatrix} t-1 \\ s \end{pmatrix} = \begin{pmatrix} 0 \\ \sum_i q'_i \lambda_i^* + \sum_A Q'_A \Lambda_A^* \end{pmatrix}. \quad (5.13)$$

L is the negative-definite 2×2 matrix defined in (2.11) and (q, Q) are normalized as in (2.6). We now distinguish two situations.

1. $\bar{c}' = \bar{c}$. If

$$\sum_i q'_i \lambda_i^* + \sum_A Q'_A \Lambda_A^* = 0, \quad (5.14)$$

then the IR symmetry is given by $(q; Q)$, and the $T_{\mathbb{C}}$ orbit of the model with enhanced symmetry is a point in V . In this case, we can apply conformal perturbation theory to deformations of this theory. The symmetry-breaking couplings α_{Am} (those vanishing on the locus exhibiting enhanced symmetry) parameterize classically marginal deformations away from the symmetric theory. The analysis of section 4 shows that in fact some of these will be marginally irrelevant, and the moduli space is given to first order in the symmetry-breaking couplings by the vanishing of the D-term for the broken symmetry. We can write this explicitly here as

$$\begin{aligned} D &= \sum_{A, n \in \Delta_A} (Q'_A + \sum_i m_i q'_i) |\alpha_{Am}|^2 \\ &= \sum_i q'_i \lambda_i + \sum_A Q'_A \Lambda_A. \end{aligned} \quad (5.15)$$

This holds at leading order in conformal perturbation theory about the symmetric point, and our conjecture amounts here to the statement that higher order corrections do not qualitatively modify the structure of the symplectic quotient that leads to the variety V : while the metric may be modified, which orbits are kept and which are excluded is not changed by higher order corrections. This implies that points in V are $T_{\mathbb{C}}$ orbits containing points whose image under μ lies in the cone $\sigma_{(q', Q')}$. We see from (5.14) that this condition is satisfied by μ^* .

2. $\bar{c}' > \bar{c}$. If (5.14) is not satisfied, the central charge \bar{c}' determined by extremization will be larger than \bar{c} , and the $T_{\mathbb{C}}$ orbit of the model with enhanced symmetry is not a point of V . Moreover, the symmetry-breaking parameters α_{Am} are not marginal couplings in this theory. Solving (5.13) we find

$$s = -\frac{r}{\det L} \left(\sum_i q'_i \lambda_i^* + \sum_A Q'_A \Lambda_A^* \right). \quad (5.16)$$

Without loss of generality we can choose the sign of (q', Q') so that s is negative. Since by construction all our couplings are invariant under $(q; Q)$, and by assumption L is negative definite, the sign of the charge under the IR R-symmetry is then the opposite of the charge under (q', Q') . Thus, couplings α_{Am} for which $Q'_A + \sum_i m_i q'_i > 0$ will be *relevant* deformations of the model with enhanced symmetry, while couplings with the opposite charge will be irrelevant; couplings preserving the enhanced symmetry are marginal. The $T_{\mathbb{C}}$ orbits of points in parameter space corresponding to irrelevant deformations of the symmetric model will not be points in V : as discussed above they will exhibit an accidental symmetry in the IR and a central charge \bar{c}' . Orbits for which at least one relevant coupling is nonzero are characterized precisely by the fact that they contain points for which the moment map satisfies

$$\sum_i q'_i \lambda_i + \sum_A Q'_A \Lambda_A > 0. \quad (5.17)$$

This specifies one side of the hyperplane associated to the enhanced symmetry, and, as we have shown, this is the side on which the point μ^* lies.

Symmetries without a smooth realization

If every enhanced symmetry were realized by a non-singular \mathcal{W} the discussion above would suffice. In general, however, there are codimension-one cones in Σ associated to one-parameter subgroups of $T_{\mathbb{C}}$ for which it is not possible to construct a non-singular \mathcal{W} exhibiting the symmetry. In these cases the RG trajectories exhibiting the enhanced symmetry along the flow are singular, and we cannot use their properties to determine the local structure of the moduli space.

A simple example of this is given by the symmetry acting as $\Gamma^A \rightarrow \tau \Gamma^A$ for some A with all the other fields invariant. This fixes the locus $J_A = 0$ which will in general be singular (it will always be singular when $n = N$). In this case, the corresponding hyperplane is $\Lambda_A = 0$, and the associated codimension-one cone lies on the boundary of $\widehat{\Sigma}$.

More interesting is the case of a codimension-one cone in the interior of $\widehat{\Sigma}$ to which the methods of the previous section do not apply. Our conjecture here is that whenever the enhanced symmetry does not satisfy (5.14), non-singular models will only exist when at least one symmetry-breaking coupling whose charge under the broken symmetry is in accord with the sense of the inequality is non-zero. In parallel with the second discussion in the previous

subsection, $T_{\mathbb{C}}$ orbits associated to points in V will be those containing points whose image under μ lies on the side of the hyperplane which contains the point μ^* .

There will also be codimension-one cones in Σ associated to one-parameter subgroups for which there is no non-singular model exhibiting the symmetry, but which satisfy (5.14). Here as well we can classify the symmetry-breaking couplings by their charge under the broken symmetry. In this case, we conjecture that non-singular models will have nonzero values for at least one coupling of *each* sign. Restricting to models with non-zero couplings of only one sign (as well as the neutral couplings) will produce a singular model. The space of $T_{\mathbb{C}}$ orbits associated to points in V in this case will not be toric. It can, however, be described as the complement of the symmetric locus (a component of Δ) in a (singular) toric variety. This contains orbits containing points whose image under μ lies *in* the cone $\sigma_{(q',Q')}$. When we exclude the singular symmetric locus here, we find precisely orbits that have nonzero symmetry-breaking couplings with both signs of the broken charge. The point μ^* clearly lies in this hyperplane.

5.3 Examples

A few examples may be helpful at this point. We proceed from a simple example for which our methods produce correctly the actual moduli space to models demonstrating their limitations.

A plain model

Consider first the class of models with $n = N = 2$ and charges

$$\begin{array}{cccc} \Phi_1 & \Phi_2 & \Gamma^1 & \Gamma^2 \\ \mathbf{q} & \frac{1}{4} & \frac{1}{6} & -\frac{3}{4} & -\frac{5}{6} \end{array} \quad (5.18)$$

and $\bar{c}/3 = r = 1 + \frac{1}{6}$. The most general superpotential is

$$\mathcal{W} = \Gamma^1 (\alpha_{11}\phi_1^3 + \alpha_{12}\phi_1\phi_2^3) + \Gamma^2 (\alpha_{21}\phi_2^5 + \alpha_{22}\phi_1^2\phi_2^2) , \quad (5.19)$$

and the discriminant is

$$\Delta = \alpha_{11}\alpha_{21} (\alpha_{11}\alpha_{21} - \alpha_{12}\alpha_{22}) . \quad (5.20)$$

This is an example of what we call a *plain* model: the torus $T_{\mathbb{C}}$ includes all field redefinitions consistent with the symmetry, so our toric considerations will in fact generate the moduli space \mathcal{M} itself.

The torus $T = \text{U}(1)^3$ action on \mathbb{C}^4 is characterized by the charges and moment map

components

$$\begin{array}{ccccc}
D & \alpha_{11} & \alpha_{12} & \alpha_{21} & \alpha_{22} \\
\lambda_1 & 3 & 1 & 0 & 2 \\
\lambda_2 & 0 & 3 & 5 & 2 \\
\Lambda_1 & 1 & 1 & 0 & 0 \\
\Lambda_2 & 0 & 0 & 1 & 1
\end{array} \tag{5.21}$$

where the latter satisfy

$$3\lambda_1 + 2\lambda_2 = 9\Lambda_1 + 10\Lambda_2 . \tag{5.22}$$

There are six codimension-one cones in Σ . Only two of these are realized by non-singular models; the remaining four comprise the boundaries of $\widehat{\Sigma}$ given by $\Lambda_1 > 0$ and $\Lambda_2 > 0$, as well as $\lambda_2 - 2\Lambda_2 > 0$ and $\lambda_1 - \Lambda_1 > 0$.

There are two codimension-one cones in the interior of $\widehat{\Sigma}$. Consider first the symmetry $(q'; Q') = (1, 0; -3, 0)$, which satisfies (5.14). The non-singular models realizing this symmetry have $\alpha_{12} = \alpha_{22} = 0$. In fact the model reduces to a product of two (2,2) minimal models and, as expected, the central charge is \bar{c} . The symmetry determined by $(q', Q') = (-1, 1; -2, 0)$, for which $\sum_i q'_i \lambda_i^* + \sum_A Q'_A \Lambda_A^* < 0$, fixes models with $\alpha_{11} = \alpha_{22} = 0$.¹⁹ Under the broken symmetry, α_{11} is negatively charged and α_{22} positively charged. We see from (5.20) that, in accordance with the conjecture, non-singular models require a non-zero value for the negatively charged coupling.

The moduli space \mathcal{M} is thus determined. We can fix two of the generators of $T_{\mathbb{C}}$ by setting $\alpha_{11} = \alpha_{21} = 1$, and the remaining couplings parameterize the toric variety $V = \mathbb{C}$ with invariant coordinate $z = \alpha_{12}\alpha_{22}$. The moduli space is $\mathcal{M} = V \setminus \widetilde{\Delta}$ where the discriminant reduces in these coordinates to $1 - z$.

A non-plain model

We can also consider the model with $n = N = 2$ and charges given by

$$\begin{array}{cccc}
\Phi_1 & \Phi_2 & \Gamma^1 & \Gamma^2 \\
\mathbf{q} & \frac{46}{471} & \frac{115}{471} & -\frac{460}{471} & -\frac{230}{471}
\end{array} \tag{5.23}$$

with $\bar{c}/3 = r = 1 + \frac{58}{471}$. The most general superpotential invariant under this symmetry is

$$\mathcal{W} = \Gamma^1(\alpha_{11}\Phi_1^{10} + \alpha_{12}\Phi_1^5\Phi_2^2 + \alpha_{13}\Phi_2^4) + \Gamma^2(\alpha_{21}\Phi_1^5 + \alpha_{22}\Phi_2^2) , \tag{5.24}$$

¹⁹Note that this symmetry leads a 2×2 L matrix that is not negative-definite; however, the corresponding superpotential is singular.

and the discriminant is

$$\Delta = \alpha_{11}\alpha_{22}^2 - \alpha_{12}\alpha_{21}\alpha_{22} + \alpha_{13}\alpha_{21}^2 . \quad (5.25)$$

The torus $T = \text{U}(1)^3$ action on \mathbb{C}^5 is characterized by the charges and moment map components

$$\begin{array}{rcccccc} D & \alpha_{11} & \alpha_{12} & \alpha_{13} & \alpha_{21} & \alpha_{22} \\ \lambda_1 & 10 & 5 & 0 & 5 & 0 \\ \lambda_2 & 0 & 2 & 4 & 0 & 2 \\ \Lambda_1 & 1 & 1 & 1 & 0 & 0 \\ \Lambda_2 & 0 & 0 & 0 & 1 & 1 \end{array} \quad (5.26)$$

where the latter satisfy

$$2\lambda_1 + 5\lambda_2 = 20\Lambda_1 + 10\Lambda_2 . \quad (5.27)$$

The cone $\widehat{\Sigma}$ is the intersection of this with the positive orthant. This is bounded, in this case, by the coordinate hyperplanes. There are four codimension-one cones in the interior of $\widehat{\Sigma}$ here, none of which satisfy (5.14).

The symmetries acting with charges $(q', Q') = (1, 0; -5, -5)$ and $(q', Q') = (1, 0, -5, 0)$ are preserved by singular models, and non-singular models, as per the conjecture, lie in orbits containing points for which $\lambda_1 - 5\Lambda_1 - 5\Lambda_2 < 0 < \lambda_1 - 5\Lambda_1$. The symmetry acting with charges $(q', Q') = (1, 0; -10, 0)$ fixes the locus $\alpha_{12} = \alpha_{13} = \alpha_{21} = 0$ where we find a product of (2,2) minimal models: up to a rescaling

$$\mathcal{W}_1 = \Gamma^1\Phi_1^{10} + \Gamma^2\Phi_2^2 , \quad (5.28)$$

with central charge $\bar{c}_1/3 = 1 + \frac{5}{33} > r$. At this point, the operators associated to α_{12} and α_{13} are *irrelevant* but the operator associated to α_{21} is *relevant*. We conclude that models with $\alpha_{11}\alpha_{22} \neq 0$ and $\alpha_{21} = 0$ flow to this IR fixed point and orbits containing such models do not contribute to V . Orbits that do contribute have a point for which $\lambda_1 > 10\Lambda_1$.

The symmetry acting with charges $(q', Q') = (0, 1; -4, 0)$ fixes the locus $\alpha_{11} = \alpha_{12} = \alpha_{22} = 0$ where we find a product of (2,2) minimal models: up to a rescaling

$$\mathcal{W}_2 = \Gamma^1\Phi_2^4 + \Gamma^2\Phi_1^5 , \quad (5.29)$$

with central charge $\bar{c}_2/3 = 1 + \frac{4}{15} > \bar{c}_1/3$. At this point, the operators associated to α_{11} and α_{12} are *irrelevant* but the operator associated to α_{22} is *relevant*. We conclude that models with $\alpha_{13}\alpha_{21} \neq 0$ and $\alpha_{22} = 0$ flow to this IR fixed point and orbits containing such models do not contribute to V . The orbits that do contribute have a point for which $\lambda_2 > 4\Lambda_1$.

Our toric model V of the moduli space is thus determined here by the cone

$$\lambda_1 > 10\Lambda_1, \quad \lambda_2 > 4\Lambda_1, \quad \Lambda_1 > 0. \quad (5.30)$$

Applying (5.9) we find that, as expected, the point

$$\mu^* = \left(\frac{435}{471}, \frac{356}{471}, \frac{11}{471}, \frac{241}{471}\right) \quad (5.31)$$

lies in this cone. Points in the preimage of this have α_{21} and α_{22} both non-zero. We can use two of our rescalings to fix $\alpha_{21} = \alpha_{22} = 1$, and under the remaining symmetry the three coefficients in J_1 transform homogeneously, so we have $V = \mathbb{P}^2$. Of course, this is an overparametrization. This is not a plain model, and we can use the remaining field redefinitions $\Gamma^2 \rightarrow \Gamma^2 + \Gamma^1(a\Phi_1^5 + b\Phi_2^2)$ to show that these theories flow to a unique IR fixed point. Not unrelated to this is the fact that there is no discriminant here: any point in \mathbb{P}^2 corresponds to a non-singular model.

A model with $N > n$

The model discussed in section 3.3 shows more of the limitations of toric methods. Here we have $Y = \mathbb{C}^9$ and $T_{\mathbb{C}} = (\mathbb{C}^*)^4$ acts on the couplings. $\widehat{\Sigma}$ is the intersection of $\lambda_1 + 3\lambda_2 = 6(\Lambda_1 + \Lambda_2 + \Lambda_3)$ with the positive orthant. There are a total of 18 codimension-one cones in the interior of $\widehat{\Sigma}$. Proceeding with our method we find a five-dimensional toric variety V determined by the moment map values $\mu^* = (\frac{6}{7}, \frac{4}{7}, \frac{1}{7}, \frac{1}{7}, \frac{1}{7})$. This is a puzzle, since we found previously that there are no models in this class with $\bar{c} = 3$. The resolution is that the model

$$\mathcal{W}'_3 = \Gamma^1\Phi_1^6 + \Gamma^2\Phi_2^2 + \Gamma^3\Phi_2^2, \quad (5.32)$$

which is in the inverse image under μ of the point μ^* , is fixed by a $U(1)$ rotation in the $\Gamma^{2,3}$ plane that is not contained in $T_{\mathbb{C}}$. This is a symmetry which arises as an accidental symmetry for all points in V , and is manifest for \mathcal{W}'_3 . This mixes with the IR R-symmetry leading to the central charge found above. This phenomenon in which a Fermi field is in fact free in the IR can occur in non-plain models with $N > n$. For models with $N = n$ a model with a free Fermi field will be singular.

5.4 Summary and further thoughts

We have provided evidence for a strong conjecture on the structure of the space of $T_{\mathbb{C}}$ orbits contributing to V . For models in which these are the only field redefinitions consistent with the UV symmetry this produces the moduli space \mathcal{M} of SCFTs with central charge \bar{c} . By analogy with studies of (0,2) GLSM parameter spaces [35, 36], we call these *plain* models. For models with larger groups of field redefinitions, our discussion is partial in two ways: we have overparametrized the moduli space, and we have failed, in general, to exclude the basins of attraction of models in which a symmetry in the complement of $T_{\mathbb{C}}$ mixes with the IR R-symmetry.

Our evidence, while suggestive, falls short of a derivation of the result. The key difficulties in a proof are twofold. First, the consequences of enhanced symmetries that are only realized by singular superpotentials are difficult to grasp, since we do not have conformal perturbation theory as a guide. For these our evidence is based on the analysis of many examples that all turned out to be consistent with the conjecture. The second difficulty lies in extending the leading order conformal perturbation theory result for enhanced symmetry loci with $\bar{c}' = \bar{c}$. It may be possible to improve this by a more detailed study of the combinatorial structures involved.

A more satisfactory derivation can be imagined, which proceeds by constructing a \bar{c} function along the RG flow and showing that this can be written in terms of α through the combinations forming $(\lambda; \Lambda)$, along the lines of [37–39].²⁰ In that work, global symmetries broken by couplings were incorporated into a -maximization in four-dimensional theories by imposing constraints on the space of symmetries over which one maximized a trial a -function. The Lagrange multipliers implementing the constraints could then be used to parameterize the flow. In our case the symmetry-breaking couplings are the superpotential couplings which break the global symmetry $U(1)^{N+n}$ of the (free) UV theory to $U(1)$ and one can introduce Lagrange multipliers to constrain the symmetries over which \bar{c} is extremized. Of course, imposing $N+n-1$ constraints is a formal procedure, because this is tantamount to specifying the outcome. However, if one proceeds formally, one finds an expression for \bar{c} in terms of the Lagrange multipliers and the *values* of these at the extremum — which reproduces (2.7) — are precisely the values of the moment map given by (5.9). The relation between this formal result and the values of the moment map is not clear to us.

6 Outlook

This project began as an attempt to classify IR fixed points of (0,2) LG theories — a generalization of the results obtained for (2,2) LG theories in [22, 23]. This beautiful work shows that for fixed $c = \bar{c}$ there is a finite set of families of superpotentials $W(X_1, \dots, X_n)$, or equivalently charges $\mathbf{q}(X_i)$ that lead to a non-singular (2,2) SCFT of desired central charge. Having the (0,2) generalization would be very useful: we would have a new class of heterotic vacua and more generally (0,2) SCFTs with many properties computable in terms of the simple UV description. These would naturally fit into the class of (0,2) gauged linear sigma models and could be used to produce a large class of hybrid models along the lines of [40].

What we learned is that, in contrast to the (2,2) case, it is not enough to classify non-singular (0,2) potentials realizing a particular set of $U(1)_L \times U(1)_R$ charges. For instance, the model studied section 3.3 would naively realize a $c = 4$, $\bar{c} = 3$ (0,2) SCFT that could correspond to some rather exotic 8-dimensional heterotic vacuum. In fact no such IR fixed point is obtained for any choice of the UV parameters. This is a general lesson for building UV models of (0,2) SCFTs: a check of UV R-symmetry anomalies is not enough, and while

²⁰A conversation with D. Kutasov, in which he suggested this idea, was instrumental in leading us to the results of this section.

the UV theory may well flow somewhere (i.e. to an SCFT with $\bar{c} > 0$), it may wind up far (i.e. at infinite distance) from the expectations of the model builder. We expect this to be a general lesson applicable to the wider class of gauged linear sigma models. In exploring that latter point it should be interesting to study in detail GLSMs with LG phases that exhibit accidents and extrapolate their consequences to large radius geometries.

For a class of models—the plain LG theories—we were able to obtain a compelling conjecture for a global description of the (0,2) moduli space \mathcal{M} realizing the expected central charge. While the resulting combinatorial structure is consistent with a case-by-case analysis of field redefinitions and their orbits in examples, we were not able to prove it in generality. Progress on both testing and proving the conjecture could be made by developing a better understanding of the combinatorial structure of quasi-homogeneous (0,2) superpotentials, as well as developing Lagrange multiplier techniques and trial \bar{c} functions. A classification of plain LG theories seems achievable; this would yield a large playground to explore LG RG flows and could give hints to the more general classification problem. Finally, it should be illuminating to relate our work to studies of RG flows with redundant couplings, e.g. [41].

A An F-term obstruction

In this section we give an example, taken from [42], that illustrates both D-term and F-term obstructions to marginal couplings. The setting is a (2,2) LG orbifold (LGO) compactification of the heterotic string with a superpotential

$$W = X_0^4 + X_1^4 + X_2^4 + X_3^8 + X_4^8 + \psi X_0 X_1 X_2 X_3 X_4 + \epsilon \Delta W . \quad (\text{A.1})$$

Here ψ and ϵ are parameters and ΔW is a generic polynomial with $\mathbf{q} = 1$. Marginal (2,0) deformations of the LGO correspond to massless E_6 -neutral space-time chiral multiplets. We can compute the massless spectrum exactly as a function of the complex parameters in the superpotential using the technique developed in [43]. This leads to the following results.

1. Setting $\psi = \epsilon = 0$ leads to a $U(1)^4$ right-moving Kac-Moody algebra and 298 marginal (2,0) deformations. We now turn on the (2,2)-preserving ψ and ϵ deformations and investigate what happens to the remaining (2,0) deformations. From above we know that at worst the marginal (2,0) deformations can become marginally irrelevant.
2. With $\psi \neq 0$ but $\epsilon = 0$ the $U(1)^4$ symmetry is broken, and the number of marginal (2,0) deformations is $298 - 4 - 6 = 288$. While 4 of the 10 marginally irrelevant deformations are associated to the broken symmetries the 6 others are not.
3. Finally, turning on $\epsilon \neq 0$ does not break any continuous symmetries, but the number of marginal (2,0) deformations decreases to $288 - 6 = 282$.

Note however, that all the singlets lifted by F-terms correspond to twisted sectors of the LGO. This is consistent with there being no F-term obstructions in pure LG theories.

B A D-term obstruction in a heterotic vacuum

Consider now the (2,2) quintic LG theory coupled to a free left-moving fermion with

$$\mathcal{W} = \sum_{i=1}^5 \Gamma^i J_i(\Phi) = \sum_{i=1}^5 \Gamma^i (\Phi_i^4 + \psi \prod_{j \neq i} \Phi_j) + \Gamma^6 \times 0. \quad (\text{B.1})$$

The interacting fields have their usual $U(1)_L$ charges $q_i = 1/5$ and $Q_i = -4/5$, and for $\psi \neq 0$ there are no extra $U(1)$ symmetries in addition to $U(1)_L \times U(1)_R \times U(1)_6$, where $U(1)_6$ is the symmetry associated to the free Γ^6 . This flows to a conformal field theory with $r = 4$ and $\bar{c} = 9$, and we can now consider deformations of the IR theory from the general perspective of deforming by chiral primary operators. In the (2,2) theory we have a good understanding of the map between the IR chiral primary marginal operators and the UV data, so we can identify the marginal deformations of the IR theory with the space of possible \mathcal{W} modulo field redefinitions. If we keep Γ^6 free, we find a 301-dimensional space of deformations.

We can also include deformations of the form $\Gamma^6 J_6$, where J_6 is some generic degree 5 polynomial. Although these break the $U(1)_6$ symmetry, they preserve the central charge and the $U(1)_L \times U(1)_R$ quantum numbers of the fields. In particular, Γ^6 has the quantum numbers of a free field. Including these J_6 deformations yields a 402-dimensional space of marginal deformations away from the (2,2) $r = 4$ $\bar{c} = 9$ fixed point.

Are all of these 402 marginal deformations exactly marginal? While all of the 301 deformations of the J_i are truly marginal, the 101 extra deformations associated to $J_6 \neq 0$ are marginally irrelevant. This is completely clear from the conformal perturbation theory discussion we gave in the text. All of these break the $U(1)_6$ symmetry, and every symmetry-breaking coupling has the same sign of $U(1)_6$ charge. Let us now see how the same result is recovered from a heterotic space-time perspective.

Heterotic insights

The \mathbb{Z}_5 orbifold of the LG theory just described, combined with an appropriate heterotic GSO projection leads to a well-understood heterotic vacuum: the LG point in the moduli space of the quintic compactification with standard embedding. The massless fields of the resulting space-time $N=1$ $d = 4$ supergravity theory consist of the supergravity multiplet, the axio-dilaton chiral multiplet, the $\mathfrak{e}_6 \oplus \mathfrak{e}_8$ vector multiplets, 326 gauge-neutral chiral multiplets, and a \mathfrak{e}_6 charged chiral spectrum $\mathbf{27} \oplus \overline{\mathbf{27}}^{\oplus 101}$. The 301 deformations of the J_i described above correspond to \mathfrak{e}_6 -preserving marginal deformations in the untwisted sector of the orbifold. These remain truly marginal for any value of the Kähler modulus (itself in a twisted sector), and at large radius they are the 101 complex structure deformations of the CY quintic, as well as 200 of the 224 deformations of the tangent bundle. As reviewed in [44], there are many arguments for why these deformations are truly marginal.

The 101 deformations associated with the J_6 couplings also have a simple space-time interpretation: they correspond to $\mathfrak{so}(10)$ -singlet components of the $\overline{\mathbf{27}}^{\oplus 101}$. Turning on

these deformations corresponds to Higgsing $\mathfrak{e}_6 \rightarrow \mathfrak{so}(10)$. This makes it obvious that the deformations are marginally irrelevant. Under the decomposition of $\mathfrak{e}_6 \supset \mathfrak{so}(10) \oplus \mathfrak{u}(1)$, we have

$$\overline{\mathbf{27}} = \overline{\mathbf{16}}_{-1/2} \oplus \mathbf{10}_1 \oplus \mathbf{1}_{-2} . \quad (\text{B.2})$$

The $\mathfrak{so}(10)$ singlets all have charge -2 under the broken $\mathfrak{u}(1)$, and hence have a D-term space-time potential. This is an example of a ‘‘D-term’’ obstruction to a marginal coupling being exactly marginal.

Since the deformation only involves world-sheet fields in untwisted sector of the orbifold, it is clear by the orbifold inheritance principle that this obstruction to marginality lifts to the un-orbifolded quintic LG model and matches the conformal perturbation theory result. In the orbifold theory it is possible to find exactly marginal deformations that Higgs $\mathfrak{e}_6 \rightarrow \mathfrak{so}(10)$ [45], but they involve an interplay between marginal couplings in twisted and untwisted sectors [44].

References

- [1] A. Belavin, A. M. Polyakov, and A. Zamolodchikov, ‘‘Infinite conformal symmetry in two-dimensional quantum field theory,’’ *Nucl.Phys.* **B241** (1984) 333–380.
- [2] S. El-Showk, M. F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin, *et al.*, ‘‘Solving the 3d Ising Model with the Conformal Bootstrap II. c-Minimization and Precise Critical Exponents,’’ [arXiv:1403.4545 \[hep-th\]](#).
- [3] S. El-Showk, M. F. Paulos, D. Poland, S. Rychkov, D. Simmons-Duffin, *et al.*, ‘‘Solving the 3D Ising Model with the Conformal Bootstrap,’’ *Phys.Rev.* **D86** (2012) 025022, [arXiv:1203.6064 \[hep-th\]](#).
- [4] A. Zamolodchikov, ‘‘Conformal symmetry and multicritical points in two-dimensional quantum field theory. (In Russian),’’ *Sov.J.Nucl.Phys.* **44** (1986) 529–533.
- [5] D. A. Kastor, E. J. Martinec, and S. H. Shenker, ‘‘RG Flow in N=1 Discrete Series,’’ *Nucl. Phys.* **B316** (1989) 590–608.
- [6] E. J. Martinec, ‘‘Algebraic Geometry and Effective Lagrangians,’’ *Phys. Lett.* **B217** (1989) 431.
- [7] C. Vafa and N. P. Warner, ‘‘Catastrophes and the Classification of Conformal Theories,’’ *Phys. Lett.* **B218** (1989) 51.
- [8] K. Hori, S. Katz, A. Klemm, R. Pandharipande, R. Thomas, C. Vafa, R. Vakil, and E. Zaslow, *Mirror symmetry*, vol. 1 of *Clay Mathematics Monographs*. American Mathematical Society, Providence, RI, 2003. With a preface by Vafa.
- [9] E. Witten, ‘‘Phases of N = 2 theories in two dimensions,’’ *Nucl. Phys.* **B403** (1993) 159–222, [arXiv:hep-th/9301042](#).
- [10] J. Distler and S. Kachru, ‘‘(0,2) Landau-Ginzburg theory,’’ *Nucl. Phys.* **B413** (1994) 213–243, [hep-th/9309110](#).
- [11] E. Silverstein and E. Witten, ‘‘Criteria for conformal invariance of (0,2) models,’’ *Nucl. Phys.* **B444** (1995) 161–190, [arXiv:hep-th/9503212](#).

- [12] J. Distler, “Notes on (0,2) superconformal field theories,” [hep-th/9502012](#).
- [13] P. S. Aspinwall, B. R. Greene, and D. R. Morrison, “The monomial divisor mirror map,” *Internat. Math. Res. Notices* (1993) no. 12, 319–337, [arXiv:alg-geom/9309007](#).
- [14] D. A. Cox and S. Katz, “Mirror symmetry and algebraic geometry,”. Providence, USA: AMS (2000) 469 p.
- [15] T. Kawai, Y. Yamada, and S.-K. Yang, “Elliptic genera and N=2 superconformal field theory,” *Nucl.Phys.* **B414** (1994) 191–212, [arXiv:hep-th/9306096](#) [[hep-th](#)].
- [16] I. V. Melnikov, “(0,2) Landau-Ginzburg models and residues,” *JHEP* **09** (2009) 118, [arXiv:0902.3908](#) [[hep-th](#)].
- [17] C. Beasley and E. Witten, “New instanton effects in supersymmetric QCD,” *JHEP* **0501** (2005) 056, [arXiv:hep-th/0409149](#) [[hep-th](#)].
- [18] I. V. Melnikov and E. Sharpe, “On marginal deformations of (0,2) non-linear sigma models,” *Phys.Lett.* **B705** (2011) 529–534, [arXiv:1110.1886](#) [[hep-th](#)].
- [19] R. Blumenhagen, R. Schimmrigk, and A. Wisskirchen, “The (0,2) Exactly Solvable Structure of Chiral Rings, Landau-Ginzburg Theories, and Calabi-Yau Manifolds,” *Nucl. Phys.* **B461** (1996) 460–492, [arXiv:hep-th/9510055](#).
- [20] K. A. Intriligator and B. Wecht, “The exact superconformal R-symmetry maximizes a,” *Nucl. Phys.* **B667** (2003) 183–200, [arXiv:hep-th/0304128](#).
- [21] F. Benini and N. Bobev, “Exact two-dimensional superconformal R-symmetry and c-extremization,” *Phys.Rev.Lett.* **110** (2013) 061601, [arXiv:1211.4030](#) [[hep-th](#)].
- [22] M. Kreuzer and H. Skarke, “No mirror symmetry in Landau-Ginzburg spectral,” *Nucl.Phys.* **B388** (1992) 113–130, [arXiv:hep-th/9205004](#) [[hep-th](#)].
- [23] A. Klemm and R. Schimmrigk, “Landau-Ginzburg string vacua,” *Nucl.Phys.* **B411** (1994) 559–583, [arXiv:hep-th/9204060](#) [[hep-th](#)].
- [24] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl. Phys.* **B435** (1995) 129–146, [arXiv:hep-th/9411149](#).
- [25] M. Kreuzer and H. Skarke, “On the classification of quasihomogeneous functions,” *Commun.Math.Phys.* **150** (1992) 137, [arXiv:hep-th/9202039](#) [[hep-th](#)].
- [26] L. J. Dixon, “Some world sheet properties of superstring compactifications, on orbifolds and otherwise,”. Lectures given at the 1987 ICTP Summer Workshop in High Energy Physics and Cosmology, Trieste, Italy, Jun 29 - Aug 7, 1987.
- [27] D. Green, Z. Komargodski, N. Seiberg, Y. Tachikawa, and B. Wecht, “Exactly marginal deformations and global symmetries,” *JHEP* **1006** (2010) 106, [arXiv:1005.3546](#) [[hep-th](#)].
- [28] J. Polchinski, “Scale and conformal Invariance in quantum field theory,” *Nucl.Phys.* **B303** (1988) 226.
- [29] J. J. Atick, L. J. Dixon, and A. Sen, “String Calculation of Fayet-Iliopoulos d Terms in Arbitrary Supersymmetric Compactifications,” *Nucl.Phys.* **B292** (1987) 109–149.
- [30] M. Dine, I. Ichinose, and N. Seiberg, “F Terms and D Terms in String Theory,” *Nucl.Phys.* **B293** (1987) 253.

- [31] V. Periwal and A. Strominger, “Kähler geometry of the space of $N=2$ superconformal field theories,” *Phys.Lett.* **B235** (1990) 261.
- [32] M. Dine, N. Seiberg, X. G. Wen, and E. Witten, “Nonperturbative Effects on the String World Sheet,” *Nucl. Phys.* **B278** (1986) 769.
- [33] K. Becker, M. Becker, J.-X. Fu, L.-S. Tseng, and S.-T. Yau, “Anomaly cancellation and smooth non-Kaehler solutions in heterotic string theory,” *Nucl. Phys.* **B751** (2006) 108–128, [arXiv:hep-th/0604137](#).
- [34] I. V. Melnikov, R. Minasian, and S. Theisen, “Heterotic flux backgrounds and their IIA duals,” *JHEP* **1407** (2014) 023, [arXiv:1206.1417 \[hep-th\]](#).
- [35] M. Kreuzer, J. McOrist, I. V. Melnikov, and M. Plesser, “(0,2) deformations of linear sigma models,” *JHEP* **1107** (2011) 044, [arXiv:1001.2104 \[hep-th\]](#).
- [36] I. V. Melnikov and M. R. Plesser, “A (0,2) mirror map,” *JHEP* **1102** (2011) 001, [arXiv:1003.1303 \[hep-th\]](#).
- [37] D. Kutasov, “New results on the ‘a theorem’ in four-dimensional supersymmetric field theory,” [arXiv:hep-th/0312098 \[hep-th\]](#).
- [38] D. Kutasov and A. Schwimmer, “Lagrange multipliers and couplings in supersymmetric field theory,” *Nucl.Phys.* **B702** (2004) 369–379, [arXiv:hep-th/0409029 \[hep-th\]](#).
- [39] D. Erkal and D. Kutasov, “a-Maximization, Global Symmetries and RG Flows,” [arXiv:1007.2176 \[hep-th\]](#).
- [40] M. Bertolini, I. V. Melnikov, and M. R. Plesser, “Hybrid conformal field theories,” [arXiv:1307.7063 \[hep-th\]](#).
- [41] N. Behr and A. Konechny, “Renormalization and redundancy in 2d quantum field theories,” *JHEP* **1402** (2014) 001, [arXiv:1310.4185 \[hep-th\]](#).
- [42] P. S. Aspinwall, I. V. Melnikov, and M. R. Plesser, “(0,2) Elephants,” *JHEP* **1201** (2012) 060, [arXiv:1008.2156 \[hep-th\]](#).
- [43] S. Kachru and E. Witten, “Computing the complete massless spectrum of a Landau- Ginzburg orbifold,” *Nucl. Phys.* **B407** (1993) 637–666, [arXiv:hep-th/9307038](#).
- [44] J. McOrist and I. V. Melnikov, “Old issues and linear sigma models,” *Adv.Theor.Math.Phys.* **16** (2012) 251–288, [arXiv:1103.1322 \[hep-th\]](#).
- [45] M. Dine and N. Seiberg, “Are (0,2) models string miracles?,” *Nucl.Phys.* **B306** (1988) 137.