

Aspergillus nidulans *verA* Is Required for Production of the Mycotoxin Sterigmatocystin

NANCY P. KELLER,^{1*} N. JEANNINE KANTZ,¹ AND THOMAS H. ADAMS²

*Department of Plant Pathology and Microbiology¹ and Department of Biology,²
Texas A&M University, College Station, Texas 77843*

Received 18 November 1993/Accepted 4 February 1994

Aspergillus nidulans produces the carcinogenic mycotoxin sterigmatocystin (ST), the next-to-last precursor in the aflatoxin (AF) biosynthetic pathway found in the closely related fungi *Aspergillus flavus* and *Aspergillus parasiticus*. We identified and characterized an *A. nidulans* gene, *verA*, that is required for converting the AF precursor versicolorin A to ST. *verA* is closely related to several polyketide biosynthetic genes involved in polyketide production in *Streptomyces* spp. and exhibits extended sequence similarity to *A. parasiticus* *ver-1*, a gene proposed to encode an enzyme involved in converting versicolorin A to ST. By performing a sequence analysis of the region 3' to *verA*, we identified two additional open reading frames, designated ORF1 and ORF2. ORF2 is closely related to a number of cytochrome P-450 monooxygenases, while ORF1 shares identity with the gamma subunit of translation elongation factor 1. Given that several steps in the ST-AF pathway may require monooxygenase activity and that AF biosynthetic genes are clustered in *A. flavus* and *A. parasiticus*, we suggest that *verA* may be part of a cluster of genes required for ST biosynthesis. We disrupted the *verA* coding region by inserting the *A. nidulans* *argB* gene into the center of the coding region and transformed an *A. nidulans* *argB2* mutant to arginine prototrophy. Seven transformants that produced DNA patterns indicative of a *verA* disruption event were grown under ST-inducing conditions, and all of the transformants produced versicolorin A but negligible amounts of ST (200-fold to almost 1,000-fold less than the wild type), confirming the hypothesis that *verA* encodes an enzyme necessary for converting versicolorin A to ST.

Toxic fungal metabolites, which are called mycotoxins, are produced by numerous fungi, including fungi commonly found as contaminants of food or feed products and fungi used for industrial production of food additives and pharmaceuticals. Many mycotoxins are classified as polyketides, which are bioreactive secondary metabolites that are synthesized like fatty acids (20). Although some polyketides (e.g., antibiotics and cancer therapeutic drugs) are beneficial, the mycotoxigenic polyketides present serious health and economic problems. Of particular concern are the carcinogenic polyketides aflatoxin (AF) and sterigmatocystin (ST), which are produced by many species of the genus *Aspergillus*. It has been proposed that both of these compounds are end products of the same lengthy biosynthetic pathway: initial polyketide precursor→norsolorinic acid→averantin→averufanin→averufin→versiconal hemiacetal acetate→versicolorin B→versicolorin A→demethylsterigmatocystin→ST→O-methylsterigmatocystin→AF B₁ (5, 41). AF and ST cause mammalian liver cancers (1a, 16) and are found in foods, including peanuts, corn, tree nuts, cheese, milk, and meat (8, 14, 21, 28, 32).

In recent studies workers have identified several of the enzymes and genes required for AF biosynthesis in *Aspergillus flavus* and *Aspergillus parasiticus* (10, 11, 23, 29, 34, 41, 43). However, little effort has been made to investigate ST biosynthesis in ST-producing *Aspergillus* spp. As many as 20 *Aspergillus* species, including *Aspergillus nidulans* (2, 17), a filamentous fungus widely used to study basic biological problems (39), have been reported to produce ST (12). Although current federal guidelines on AF contamination do not cover ST, the possible effects of ST on human health cannot be ignored. It has been shown that many of ST's bioreactive activities parallel

those of AF, although slightly higher concentrations of ST are usually needed to produce the same effect (3, 33). Because ST can be produced in greater quantities than AF in certain foods (19) and may be found in a larger variety of foods and environments, it is important to understand parameters that control ST biosynthesis (15, 24, 28, 32, 37). In this paper we describe the isolation and characterization of an *A. nidulans* gene required for ST production, *verA*. Disruption of the *verA* coding region results in an inability to convert versicolorin A to ST.

MATERIALS AND METHODS

Fungal strains and growth conditions. *A. nidulans* FGSC 26 (*biA1 veA1*) and rM31 (*pabaA1 biA1 argB2 veA1*) were maintained as silica stocks and were grown on minimal medium for the production of conidia. Strain rM31 was isolated as the meiotic progeny of strains PW1 (*biA1 argB2 methG1 veA1*) and FGSC 237 (*pabaA1 yA2 trpC801 veA1*). To induce ST biosynthesis, the fungal isolates were grown either on oatmeal porridge or in YEC (0.2% yeast extract, 5.0% corn steep liquor).

Nucleic acid manipulations. An *A. nidulans* genomic cosmid library (1, 7) was screened with an internal fragment from the *A. parasiticus* *ver-1* gene, pBSV2, which has been predicted to encode a keto-reductase required for conversion of versicolorin A to ST (34). All hybridization experiments with ³²P-labeled *ver-1* were carried out at 68°C, and the membranes were washed twice in 2× SSPE–0.1% sodium dodecyl sulfate (SDS) (1× SSPE is 0.18 M NaCl, 10 mM NaPO₄, and 1 mM EDTA [pH 7.7]) at room temperature.

The chromosome containing the *A. nidulans* *ver-1* homolog was identified by probing a nylon membrane containing the separated chromosomes from *A. nidulans* FGSC 4 (kindly provided by Xiaoling Xuei and Paul Skatrud, Eli Lilly Corp.,

* Corresponding author. Phone: (409) 845-0963. Fax: (409) 845-6483. Electronic mail address: npk3325@venus.tamu.edu.




FIG. 1. Restriction map of the *verA* genomic region. pNK10 contains the entire *verA* gene within a 6.6-kb *Hind*III fragment. pJK1 was constructed from pNK10 by removing a 1.1-kb fragment that contained polylinker sequences including an *Eco*RI site. A 1.8-kb *Eco*RI fragment containing *argB*⁺ was inserted into the *verA* coding region to give pJK2. pNK8 was used as a DNA probe in genomic blots to demonstrate disruption of *verA* and contains 509 bp of *verA* in a 1.8-kb *Pst*I fragment. *verA* is indicated by the diagonal lines, and the arrows indicate the directions of transcription for *verA*, *argB*, ORF1, and ORF2 (described in the text). Abbreviations: E, *Eco*RI; EV, *Eco*RV; H, *Hind*III; P, *Pst*I; S, *Sal*I; Ss, *Sst*II; X, *Xba*I.

Indianapolis, Ind.) with radioactively labeled pNK10 DNA, using standard methods (31). After prehybridization for 3 h in a solution containing $5 \times$ SSC, 50 mM NaPO_4 (pH 7.2), 0.4% SDS, $5 \times$ Denhardt's solution, 2.5 mM EDTA, 0.5% dextran sulfate, and 0.1 mg of salmon sperm DNA per ml ($1 \times$ SSC is 0.15 M NaCl plus 0.015 M sodium citrate), the denatured probe was added, and hybridization was carried out for 24 h at 60°C. The membrane was washed at 60°C for 15 min in $2 \times$ SSC and then for 15 min in $1 \times$ SSC and then subjected to autoradiography.

The *verA* disruption plasmid, pJK2, was constructed in two steps (Fig. 1). First, pJK1 was constructed by digesting pNK10 with *Xba*I and religating. This removed a 1.1-kb fragment that contained a polylinker sequence including an *Eco*RI site. Next, pJK1 was digested with *Eco*RI and religated in the presence of a 1.8-kb *Eco*RI fragment containing *argB*⁺ (obtained from pSalArgB) to give pJK2. pJK2 was linearized by double digestion with *Xba*I and *Pst*I, and the vector sequence was removed prior to transformation. Transformation, extraction of transformant genomic DNA, restriction enzyme digestion, gel electrophoresis, Southern blotting and hybridization were performed by using previously described protocols (31). All transformants were isolated from single spores prior to analysis.

Analysis of ST production. Beakers containing 3 g of oatmeal porridge were inoculated with 100 μ l of a suspension containing 1×10^6 spores per ml. After 6 days of growth at 30°C, the cultures were killed by adding 15 ml of acetone. The beakers were gently agitated for 30 min, and then 15 ml of chloroform was added to each beaker. After an additional 30

min of agitation, each solution was passed through anhydrous sodium sulfate and collected in a beaker. After evaporation, each extract was resuspended in 1.5 ml of chloroform, and a 10- μ l portion of each extract was spotted onto a thin-layer chromatography (TLC) plate (Analtech). An ST standard (Sigma) and a versicolorin A standard (a semipure standard previously prepared from extracts of *A. parasiticus* SRRS 164) were also spotted onto the TLC plates. The plates were developed with either benzene-glacial acetic acid (95:5, vol/vol) or toluene-ethyl acetate-glacial acetic acid (80:10:10, vol/vol/vol). After the plates were developed, they were sprayed with aluminum chloride, which enhanced the visualization of ST under long-wavelength (365-nm) UV light (38). A nonreplicated high-performance liquid chromatography (HPLC) analysis of the ST in culture extracts was performed by F. Neeley, Eli Lilly Corp., Indianapolis, Ind.

Nucleotide sequence accession number. The GenBank accession number for *A. nidulans verA* and flanking sequences encoding open reading frames (ORFs) ORF1 and ORF2 is L27825.

RESULTS

Isolation of *verA* and determination of the genomic sequence. The *A. nidulans verA* gene was isolated by using a fragment of the *A. parasiticus ver-1* gene as a heterologous probe to screen an *A. nidulans* cosmid genomic library, and one positive clone, pL24B3, was identified. The *ver-1* hybridizing region was localized to a 6.6-kb *Hind*III fragment and was subcloned into pBluescript SK⁻ to give pNK10 (Fig. 1).

1 ATGCCCCAGAAGGGAGCACCGTGTAGGCGCAAGCTGACCGAGTTGGGAGCGCAAAGCGTGGCCGCGGCTGCGATTGCGCTCGCATACCTGCTTACAAG 100
 101 TCGCCGTACGTTTTCCCGGTGATCGGGTGCCCGACGGTCGAGCAGCTGGAGGCGAATATACGCCCTCGGTGTAGAGCTCAGTGATGAGGAAATGTACGAGA 200
 201 TTGAAGACACGATCCCTTTTGATGTCGGCTTCCCATGGCTTCTTATTTCGAATCGCCCCAGCAGAAGTACCGTAGTGATATGACGACCAGGCATATCTG 300
 301 GCAGGTTACCTGCAATGCCCGGATCGAGAGTGTGCCTAAGCCGAGAGTATGTATCTCTCAACCTGAATTTATGATTTGCTAATCGAACTTACCAGCCTA 400
 401 TCGAGCCAAAGCAGGGTACAAGCAGATGGATCGGAAGTAGTTCTCGGTAGCATTAGCCAAGCATCGGGTCCCGAGCGTTCAAGTATTTTATATATGAGCC 500
 501 TTGTTTCCTTCTATGTCATGGTAGCCAGTATCCATAAGGTATAGGAATCAACCATGCTCCTCCGATAATTACCGTCTCGATGAAAAGTCGCTCTGG 600
 verA-> M S S S D N Y R L D G K V A L V
 - - - - - E R - - - - - H E - R P R
 T G A G R G I G A A I A V A L G Q P G A K V V V N Y A N S R E A A
 601 TAATGGGGTGGCCGCGGCATCGGAGCAGCCATCGCCGTAGCCCTCGGTACGCCGGGCGGAAGTCTGCTCAACTACGCTAACTCCCCTGAGGCCGC 700
 R - W L N R S R R M V P M L S Q S - - - - - E - T A - - - A E
 E K V V D E I K S N A Q S A I S I Q A D V G D P D A V T K L M D Q
 701 AGAAAAGTCTGCGACGAAATCAAGTCGAACGCTCAGAGCGCCATTTCCATTCAAGCCGATGTCCGGTACCCTGATGCCGTCACCAAAGTATGATGATCAG 800
 T - R - - - - - L - - - - - E
 A V E H F G Y L D I V S S N A G I V S F G H V K D V T P D
 801 GCCGTTGAGCACTTCGGATACCTGGATATAGTCTCATCTAACCGGGAATTGTCTCGTTCGGGATGTCAAGGACGTTACGCCAGATGTATGCGTCCCAT 900
 E F D R V F R V N T R G Q F F V A R E A Y
 901 CTCCTTACGAAAGTCTGTAGAGCTCTGACCTCAGCAGGAATTCGACCGAGTATTTCCGGTCAACACGCGGGACAGATTTTTTCGTCGCCCGGAGGCCGA 1000
 - - M - - - - - C - - - - - K - - - - -
 R H L R E G G R I I L T S S N T A S V K G V P R H A V Y S G S K G
 1001 TCGCCATCTGCGTGAAGGCGGACGCATCATCTCACAAAGTCCAACACAGCCAGCGTCAAGGCGTCCCAGGCAGCGTGTGTACTCGGGCTTAAGGGG 1100
 - - - - - M - - - - -
 A I D T F V R C L A I D C G D K K I T V N A V A P G A I K T D M F L
 1101 GCGATTGACACCTTTGTGCGGTGCCTAGCTATCGACTGCGCGGACAAGAAGATCACGGTCAACCGGTCGCTCCCAGGCATCAAGACCGATATGTTTC 1200
 A - - - - -
 S V S R E Y I P N G E T F T D E Q V D E
 1201 TATCCGTGTCGCGAGATATATCCCAATGGGGAGACTTTTACTGATGAGCAGGTGGATGAGGTACGTTTTGTCTTTGTGTAGTATCTACGGCGGCTGC 1300
 C A A W L S P L N R V G L P V D V A R V V S F L A S D A A E
 1301 TAATGGACAGTGTGCCGCTGGTGTGCGCCGCTAAATCGGGTTCGGATTACCGGTTACGTTGGCCCGGTTGGTCAAGTCTTCTAGCTTACAGTGGCGCGA 1400
 - V - - - - -
 W I S G K I I G V D G G A F R *
 1401 ATGGATCAGTGGAAAGATTATTGGCGTTGATGGGGGGCCTTTAGATAAGTACATCATATACTTGAACATATAGGGTAGACATGCAATGTTGCTCCC 1500
 1501 CGCTCGCTTACCGATATCTGCCGATCATCGTCAGCAACCATTAGGTCACGAAAAAGAGTATACTAAGAGTAAACATCCGTGCATGGTATGAACTTAGT 1600
 1601 TGGGTACACCGAGTTAGTACACCGTACTTAAGTACACTCAGCGATTCACTTAGCGGCTGAATCGGCATTTTCATACTCTGCCAGCACCGGAGGCCAG 1700
 * K P P Q I P M E Y E A L V P P G
 1701 CAACATCAACAACAATAGGCAAAGCATGCACACGCTCAAACCAGGTAGATAGGTTGCGGTGCTCATCCCTCCAGCGTTTATCAAGGAAAAACCGGAATGC 1800
 A V D V V I P L A H V R E F W T S L N R H E D R W R K D L F F R F A
 1801 GCCCTGCACAATCCCAGCACAACAGATCAGCTAGGCTGAGGGTTTCCCGACCAAGTACTCTCGCCCAAAAGATGGTTGTCAAGAATCTTTAGCCGT 1900
 G Q V I G L V F L D A L S L T E G V L Y E R G C L H N D L I K L R
 1901 GCTAAAGTGTATCTTTGCTTTGATATATGTTGTGTCAGCATTGAAGTTGGCTCGTCCGATGAGCGGGTTGAACCAGCCCTAACGCTGGGAGGATTTCCGG 2000
 A L T D D K S Q Y I N D A N F N A R G I L P N F W G G L A P L I E
 2001 TGATCCCGAAGGCCATCCAGCGAATGATGGAGGCATATCTTGTCCGGTAGTCCCAAGTAAAGTCGATTTGAACTCTTGAGATGTTACTATACCTCTTAG 2100
 T I G F A M W R I I S A Y E Q G T T G L L T T N S D
 2101 TCAGGAATTGAATAGATGGAATTGCAGTAGCAGCATGGTACCATAGAGAGCAATAGCAATAGATTCCGTCAATACGTACGCGTCCGCCCCACAACGTA 2200
 Y L A I A I S E T L V Y A D A G V F T
 2201 GGAATCTGCTAGAGGTTGAGCTGGAGATACTCTTCGGTAGCATCTTGAATGAAGTATGATGCTGATTTTCAGAGGCAAAATGTTGCTTTTGC 2300
 P I K G L P N L Q L Y E E T A D K F S T I T K I K L P L N N A K A
 2301 TCGCAAGAAATCGCCAGCAGCGGGGTTGAACGGCGAGTGTACAGAGTCCGCAACGGCATTGCGAATAATTTCAATTCAGAGCTGATTTCTCGTATTG 2400
 I A L I A L S R P N F P R T Y L T G F P M <-ORF1

2401 TATGCTTGTGGCAACCTGCTAAATACAAATACTGACAGCAAATCAACTATATGTCAAGACCATGCCCTTCAGCTGTCCGCGTAACCTTAACCTCCCCCAG 2500

2501 GACAACGGCCTTCATCTTTCCCGCATCCGTGAAACGGTCTCGTCCGCATAACTTCGGGGCTGCTCATGACGGGGACAACTCCTCGAAGGTGGCTTGG 2600

2601 CTCGCAAATTGGACAACAGCAAATGCGTCAAGGTCAAATCGATCGAGTCGCCGGCCAGCGGGGTGACCGGTTGCTGCAGGTAGTGTGGGTGTGGCTGA 2700

2701 CTGGAAGGCCCTCCCGCCGAGTCGTTGCAGCAGGGGATATGTTCCGCTCTCCAGTGGTTACGAAATTCGCTGGGTGTGAGGTGCCCGCGACGGGCTAC 2800

2801 AAGAATCAAGACAGTGAACATGGTGGAGTAAAGTGTGTGTATGTTTGCACACTTGTCTCCAGAAATTCGCGCAATACGCCCTTATATATGGCTGT 2900

2901 CCCTATCTCGGTCCGCGAACGAACATAACAATTATTAGAGAGACTCTTCTTACATTTTTGTCAATTTGTTGCCAAAGTCACTTCACTCATTGCTGTCTCTCC 3000

3001 AACCATGTACACAACATATCATCACAGCGGTATGCGTGCATTTCGCTCTTCCACTCTGGACAGCTTCTATCAAGCGCGGCAGGAGGTATGGGCCCTCCAG 3100

ORF2-> M P S F S L L T G H

3101 CGGGCAAACCTAGTACGAGCCCTCTGACCCAATGATTGGCTAGAGGACGATTAAGTGGTATACAAGCCCATGCCTTCTTTCAGCCTGCTGACCGGCCAC 3200

F G A L K Q T I D G M P P N A T L H S I M L K L S Q K F R S G M F Y

3201 TTTGGTCCCTCAAACAACCATCGATGGCATGCCGCCAACGCAACCTGCATAGCATTATGCTGAAATGTGCGAAAAGTTCGGCTCAGGGATGTTCT 3300

I N M W P F S G T W L V V A T P S G A A Q I Q S L N L S K P P L V

3301 ACATCAACATGTGGCCATTACAGCGGTACATGGCTCGTGGTTCGCAACACCGTCTGGCGCGGCCAGATCCAGAGTCTGAATCTTTCGAAGCCCGCTGTT 3400

R R P L E T I T G G P S L M S M H G E T W K R W R A L F N P G F N

3401 GCGAAGACCGCTGGAGACTATCACGGGGCCCAAGCTTGATGAGTATGCATGGTAAACATGGAACCGTGGAGGGCACTGTTTAAATCCAGGCTTAAAC 3500

P N Y L I G L A P L I A D E V V V F C E Q L R Q K A R T G T V F Q L

3501 CCCAACTACTTGATTGGGCTGGCGCCGCTGATCGCCGATGAGGTCGTTGTTTTTGGCAGCAGCTACGGCAGAAGGCCAGAACAGGAACAGTTTCCAGC 3600

E P L T L R L T V D T I C S V T L Y V V T P V G R W P F L T P D L

3601 TTGAACCGCTCACTCTGAGGTTGACAGTTGATACGATTTGCTCTGTGACGTTGATGTGGTTACTCCCGTTGGCGATGGCCCTTCTAACCCCTGACTT 3700

E I H S S T T K L R T T P L P Q R C N G R S N G P R L E L P S T P

3701 AGAGATTCACAGCTCCACCACCAAACCTCAGGACCACCCCTTGCCCTCAGCGCTGCAACGGCAGATCGAATGGGCTCGTTTGGAACTACCTTCAACCCCT 3800

L R R Y L T V R P L V M W Y N N R L M N R F I D Q E V D R A Y R E Q

3801 TTAAGCGGTACCTGACCGTCCGCGCTCTGGTGTGTGGTACAATAACCGCCTTATGAACCGCTTCATCGACCAAGAGGTGACCGAGCGTACCGGGAGC 3900

S G R Q S K S V I S L A L R D Y M K E K D G S L E D F K R R V A P

3901 AGTCTGGCCGTCAAGTCCCGTATCCCTCGCCCTCAGAGATTACATGAAAGAGAAGATGGAAGTCTGGAAGACTTCAAACGACGTGTTGGCGCC 4000

Q L R V F L F A G R D T T S S T L L Y A F Y L L S R H P E A L A K

4001 ACAGTTACGGGTCTTCTCTTCGCGAGGTAGAGATAACAACGAGCAGTACTGCTCTATGCATTCTACTGCTTTCGCCGACATCCAGAGGCCCTAGCTAAG 4100

V R L E H D Q V F G P Y H Q Q V H E K I H Q D A K L L N Q L P Y T T

4101 GTGCGCTTAGAGCAGCAGGCTCTCGGCCATATCATCAACAAGTACACGAGAAAATCCACCAAGATGCGAAACTCCTCAACCAACTCCCTACACAAA 4200

A V L K E T L R L F P P S A S M R E A D P A L K S P T T T A K Y I

4201 CAGCTGTCCCTAAAGAGACTCTGAGGCTCTTCCCTCCGCTGCTCCATGCGTGAAGCGGACCGCGTTGAAATCACCGACGACAACGGCCAAGTATAT 4300

P L Q

4301 CCCACTGCAG 4310

FIG. 2. Nucleotide sequence of a 4,310-bp region containing *verA*, ORF1, and ORF2. The putative translation start site of *verA* begins at nucleotide 555, and the termination codon starts at nucleotide 1446. A comparison of the amino acid sequences of the products of *A. nidulans verA* (bottom line) and *A. parasiticus ver-1* (top line) (34) is shown above the nucleotide sequence; dashes indicate that the amino acids in the two peptides were identical. The comparison shows the high degree of conservation between the two gene products. A putative NADPH binding site, GXGXXA, is underlined (starting at residue 20 for *verA* and residue 18 for *ver-1*). The intron positions in *verA* were predicted on the basis of sequence similarity to a *ver-1* cDNA. Two additional ORFs found downstream of *verA*, one similar to the gamma subunit of elongation factor 1 (ORF1) and the other related to several cytochrome P-450 monooxygenases (ORF2), are also shown.

Further analysis revealed that the similarity spanned the central *EcoRI* site, and by performing sequence analysis of the region we identified the *verA* gene and the coding region (Fig. 2). In addition, a sequence analysis of several thousand base pairs downstream from *verA* revealed two additional potential ORFs, designated ORF1 and ORF2 (Fig. 1). ORF1 and *verA* read in opposite orientations, and ORF1 extends from nucleotides 1648 to 2361 and is predicted to encode a 200-codon polypeptide that exhibits ~40% identity over the entire sequence to the N-terminal half of the gamma subunit of elongation factor 1 (25). ORF2 begins at nucleotide 3170 and extends in the same direction as *verA* through the end of the sequenced region; the 3' end of the gene has not been

sequenced yet. The predicted >380-codon ORF exhibited ~30% identity to several cytochrome P-450 monooxygenases (42).

The predicted amino acid sequence of the *A. nidulans verA* gene product and a comparison with the *A. parasiticus ver-1* gene product are shown in Fig. 2 (34). As Fig. 2 shows, the *verA* and *ver-1* gene products of *A. nidulans* and *A. parasiticus* are closely related. A search of various protein and DNA data bases revealed that the *A. nidulans* gene product exhibited homologies to products of several genes that encode NADPH-dependent reductase activities involved in polyketide biosynthetic processes. The results of previous studies had suggested that an NADPH-requiring reductase was required for conver-




FIG. 3. *verA* is located on chromosome IV. Chromosomes of *A. nidulans* FGSC 4 (17a) were separated. The chromosomal band sizes (in megabases) are indicated on the left; chromosome IV is 2.9 Mb long. (A) Ethidium bromide-stained gel. (B) Hybridization of chromosomes to pNK10.

sion of versicolorin A to ST in the ST-AF pathway (5). The putative NADPH binding site, GXGXXA, is underlined in Fig. 2.

pNK10 (Fig. 1) was used to probe a contour-clamped homogeneous electric field gel blot of the eight separated *A. nidulans* chromosomes to determine the genomic position of *verA*. Figure 3 shows that the *verA* probe hybridized most strongly to chromosome IV, the smallest chromosome of wild-type *A. nidulans* (6). We also observed very weak hybridization to other chromosomes, most notably the chromosome doublets at 3.8 and 3.5 Mb (6), and this hybridization was thought to be nonspecific. A nearly complete overlapping cosmid map of *A. nidulans* chromosome IV has been constructed (29a), and further analysis revealed that the *verA*-containing cosmid is located within a contig that includes another cloned gene, *bimD* (13), which has been mapped to the designated left end of chromosome IV (25a).

***verA* is required for ST production.** To determine whether *verA* is required for ST production, we constructed an *A. nidulans* strain containing an insertion in the *verA* coding region. An *argB2* mutant (rM31) was transformed with pJK2 or pSalArgB, and *argB*⁺ transformants were selected. Genomic DNAs from 51 pJK2 transformants were digested with *Hind*III and probed with a 1.8-kb *Pst*I fragment from pNK8 (Fig. 1). Seven transformants contained a *Hind*III fragment that was 3.4 kb long (the size of the *Hind*III fragment in pJK2) and hybridized to the probe, as predicted for a *verA* gene disruption




FIG. 4. Southern analysis of disrupted *verA* transformants. Genomic DNAs were digested with *Eco*RV and probed with pNK8 (Fig. 1), which contains the 3' end of *verA*. Fragment sizes for wild-type *verA* (lanes A and B) were predicted to be 7.5 and 2.5 kb. Fragment sizes for disrupted *verA* were predicted to be 7.5 and 1.7 kb (lanes C through I). Lanes A through I contained isolates rM31, TJK1, TJK2, TJK6, TJK8, TJK10, TJK4, TJK11, and TJK7, respectively. The positions of lambda *Hind*III markers are indicated on the left.

event, while the remainder of the transformants contained the 6.6-kb *Hind*III fragment predicted for wild-type *verA* (6.6 kb was the size of the pNK10 *Hind*III fragment). Genomic DNAs from the seven putative *verA*-disrupted transformants were then digested with *Eco*RV and probed with pNK8. As Fig. 4 shows, all seven of these strains (TJK2, TJK4, TJK6, TJK7, TJK8, TJK10, and TJK11) contained two hybridizing fragments (1.7 and 7.5 kb), as predicted for a *verA* gene disruption. Two strains containing wild-type *verA*, rM31 and TJK1, contained the expected 2.5- and 7.5-kb fragments.

A TLC analysis of extracts from fungal strains grown under ST-inducing conditions was performed with strains FGSC 26 and TJK1 and the seven *verA* disruption strains described above. As Fig. 5 shows, none of the *verA*-disrupted transformants produced visible ST; instead, all of the strains accumulated an orange pigment that migrated to the same spot as the versicolorin A standard on TLC plates developed in benzene-glacial acetic acid (95:5, vol/vol). Only strains TJK1 and FGSC 26 produced visible ST on the TLC plates. An HPLC analysis of the same extracts revealed that there was a 200-fold to nearly 1,000-fold reduction in the ST concentration in the seven *verA* disruption strains. In this experiment the isolates were grown on 3 g of oatmeal for 1 week, and strain TJK1, which contained the wild-type *verA* gene, produced 924.0 μ g of ST, while strains TJK2, TJK4, TJK6, TJK7, TJK8, TJK10, and TJK11, which contained a disrupted *verA* gene, produced 4.5, 4.0, <0.5, 2.0, 5.5, 5.0, and <0.5 μ g of ST, respectively (the limit of detection was 0.5 μ g of ST).

DISCUSSION

Aspergillus spp. constitute a major portion of the mycotoxin-producing fungi found worldwide. Several food- and feed-contaminating species, including *A. flavus*, *A. nidulans*, and *A. parasiticus*, are also important industrial fermentation organ-




FIG. 5. *verA* disruptants do not make ST. The TLC plate was developed in benzene-glacial acetic acid (95:5, vol/vol). *A. nidulans* FGSC 26 (lane 3) and *A. nidulans* transformant TJK1 (lane 4) produced large quantities of ST, whereas transformants TJK2, TJK4, TJK6, TJK7, TJK8, TJK10, and TJK11 (lanes 5 through 11, respectively) did not produce ST but instead accumulated versicolorin A (VA). A pigment that migrated to the same spot as ST in lanes 5 through 11 was not ST as it fluoresced blue or green, whereas ST fluoresced yellow after treatment with AlCl_3 (38). In addition, the blue or green pigment was also produced along with ST in strains FGSC 26 and TJK1. Lanes 1 and 13 contained the ST standard and lanes 2 and 12 contained the VA standard.

isms (9, 30). An examination of the results of epidemiological studies, feeding studies, and biochemical and chemical studies has established that ST, AF, and related *Aspergillus* polyketides cause mammalian hepatocarcinomas (1, 27) and should be classified as carcinogens (16, 33). Consequently, strict national and international standards of allowable AF levels (more commonly analyzed than ST levels) in food supplies have been imposed to protect public health.

One approach to controlling AF and ST production is to understand the mechanisms that regulate production of enzymes in the pathway. Recent molecular studies have led to identification and isolation of genes and enzymes implicated in the AF biosynthetic pathway in *A. flavus* and *A. parasiticus* (10, 11, 23, 29, 34, 41, 43). However, molecular manipulations of genes in these asexual species are not easy. In fact, gene disruption techniques have only recently been described for *A. parasiticus* (40), and such techniques have not yet been described for *A. flavus*. To overcome these difficulties, we studied AF-ST regulation in *A. nidulans*, a genetically tractable relative of *A. parasiticus* and *A. flavus* for which molecular approaches are well developed (26, 39). In this study we found that *A. parasiticus ver-1*, a gene implicated in the conversion of versicolorin A to ST, has a homolog, *verA*, in *A. nidulans* and that *verA* exhibits 85% amino acid identity with *ver-1*. Disruption of *verA* in *A. nidulans* resulted in a loss of ST production and accumulation of versicolorin A, the immediate precursor of ST. This result is particularly important because, although *ver-1* was isolated from *A. parasiticus* on the basis of its ability to complement a mutation resulting in accumulation of versicolorin A, it has not been demonstrated that *ver-1* is required for AF biosynthesis by gene disruption.

A number of genes required for production of secondary metabolites in bacteria (e.g., polyketides in *Streptomyces* spp. [20]) and in fungi (18, 36) are found in clusters. Recent data support the notion that ST and AF genes are also probably clustered in *A. flavus*, *A. nidulans*, and *A. parasiticus* (10, 17a, 35). A sequence analysis of the DNA region immediately downstream of *verA* revealed two additional potential ORFs

(Fig. 2). One of these ORFs is predicted to encode a polypeptide that is closely related to several cytochrome P-450 monooxygenases (42). Monooxygenase activities are proposed to be required for several steps in the ST-AF pathway (5), leading to the suggestion that this putative gene is also important for ST biosynthesis. The second ORF identified in this region exhibits a high degree of similarity to the gamma subunit of elongation factor 1 (25). What, if any, role this ORF plays in ST-AF biosynthesis is not clear. We have recently identified transcripts which accumulate during ST production from these DNA regions (21a) and will determine their function in future experiments.

Our placement of *verA* on linkage group IV has helped in directing further linkage searches of *verA* to other known genes on this chromosome. Linkage data traditionally have been obtained in *A. nidulans* through sexual crosses, but recent efforts directed toward creating an *A. nidulans* genomic contig map have allowed detailed physical placement of genes on individual chromosomes (6, 7). On the physical map *verA* is next to *bimD* on one end of chromosome IV. The location of *verA* can be confirmed by sexual crosses. The linkage information obtained from *verA* and additional ST genes in *A. nidulans* will be helpful in directing ST-AF linkage studies in *A. flavus* and *A. parasiticus*, in which gene mapping studies are limited to parasexual studies and electrophoretic karyotyping (4, 22).

ACKNOWLEDGMENTS

We thank Frank Neeley, Paul Skatrud, Xiaoling Xuei, and Roland Hodges for their cooperation in this project and Deepak Bhatnagar for his review of the manuscript. We also thank Jonathan Arnold, John Linz, Rolf Prade, and Bill Timberlake for providing unpublished information, Edward Pestovic for technical assistance, and Larry Haller-Harris, Julie McCollum, and Susan Hardin for their work in the DNA sequencing laboratory.

This work was supported by Texas Agricultural Experiment Station funds awarded to N.P.K., by Texas Advanced Research Program grant 010366-79 to T.H.A., and by National Peanut Foundation funds awarded to N.P.K. and T.H.A.

REFERENCES

- Adams, T. H., and W. E. Timberlake. Unpublished data.
- Adams, R. H. 1989. Induction of hepatocellular carcinoma in nonhuman primates by chemical carcinogens. *Cancer Detect. Prev.* 14:215-219.
- Aucamp, P. J., and C. W. Holzappel. 1970. Polyhydroxyanthraquinones from *Aspergillus versicolor*, *Aspergillus nidulans* and *Bipolaris* sp.; their significance in relation to biogenetic theories on aflatoxin B₁. *J. S. Afr. Chem. Inst.* 23:40-56.
- Baertschi, S. W., K. D. Raney, T. Shimada, T. M. Harris, and F. P. Guengerich. 1989. Comparison of rates of enzymatic oxidation of aflatoxin B₁, aflatoxin G₁, and sterigmatocystin and activities of the epoxidases in forming guanyl-N7 adducts and inducing different genetic responses. *Chem. Res. Toxicol.* 2:114-120.
- Bennett, J. W., and K. E. Papa. 1988. The aflatoxigenic *Aspergillus* species. *Adv. Plant Pathol.* 6:263-280.
- Bhatnagar, D., K. C. Erhlich, and T. E. Cleveland. 1992. Oxidation-reduction reactions in biosynthesis of secondary metabolites, p. 255-286. In D. Bhatnagar, E. B. Lillehoj, and D. K. Arora (ed.), *Handbook of applied mycology*. Marcel Dekker, Inc., New York.
- Brody, H., and J. Carbon. 1989. Electrophoretic karyotype of *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. USA* 86:6260-6263.
- Brody, H., J. Griffith, A. J. Cuticchia, J. Arnold, and W. E. Timberlake. 1991. Chromosome-specific recombinant DNA libraries from the fungus *Aspergillus nidulans*. *Nucleic Acids Res.* 19:3105-3109.
- Bullerman, L. B., P. A. Hartman, and J. C. Ayres. 1969. Aflatoxin production in meats. II. Aged dry salamis and aged country cured hams. *Appl. Microbiol.* 18:718-722.
- Carrez, D., W. Janssens, P. Degraeve, W. Fiers, C. A. M. J. J. van

- den Hondel, and J. R. Kinghorn. 1990. Heterologous gene expression by filamentous fungi: secretion of human interleukin-6 by *Aspergillus nidulans*. *Gene* **94**:147–154.
10. Chang, P.-K., J. W. Cary, D. Bhatnagar, T. E. Cleveland, J. W. Bennett, J. E. Linz, C. P. Woloshuk, and G. A. Payne. 1993. Cloning of the *Aspergillus parasiticus* *apa-2* gene associated with the regulation of aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **59**:3273–3279.
 11. Chang, P.-K., C. D. Skory, and J. E. Linz. 1992. Cloning of a gene associated with aflatoxin B₁ biosynthesis in *Aspergillus parasiticus*. *Curr. Genet.* **21**:231–233.
 12. Cole, R. J., and R. H. Cox. 1981. Sterigmatocystins, p. 67–93. *In* R. J. Cole and R. H. Cox (ed.), *Handbook of toxic fungal metabolites*. Academic Press, New York.
 13. Denison, S. H., E. Kafer, and G. S. May. 1993. Mutation in the *bimD* gene of *Aspergillus nidulans* confers a conditional mitotic block and sensitivity to DNA damaging agents. *Genetics* **134**:1085–1096.
 14. Diener, U. F., R. J. Cole, T. H. Sanders, G. A. Payne, L. S. Lee, and M. A. Klich. 1987. Epidemiology of aflatoxin formation by *Aspergillus flavus*. *Annu. Rev. Phytopathol.* **25**:249–270.
 15. Francis, O. J., Jr., G. M. Ware, A. S. Carman, G. P. Kirschenheuter, S. S. Kuan, and R. F. Newell. 1987. Thin-layer chromatographic determination of sterigmatocystin in cheese: interlaboratory study. *J. Assoc. Off. Anal. Chem.* **70**:842–844.
 16. Fujii, K., H. Kurata, S. Odashima, and Y. Hatsuda. 1976. Tumor induction by a single subcutaneous injection of sterigmatocystin in newborn mice. *Cancer Res.* **36**:1615–1618.
 17. Hajjar, J. D., J. W. Bennett, D. Bhatnagar, and R. Bahu. 1989. Sterigmatocystin production by laboratory strains of *Aspergillus nidulans*. *Mycol. Res.* **94**:548–551.
 - 17a. Hodges, R., et al. Unpublished data.
 18. Hohn, T. M., S. P. McCormick, and A. E. Desjardins. 1993. Evidence for a gene cluster involving trichothecene pathway biosynthetic genes in *Fusarium sporotrichioides*. *Curr. Genet.* **24**:291–295.
 19. Holzapfel, C. W., I. F. H. Purchase, P. S. Steyn, and L. Gouws. 1966. The toxicity and chemical assay of sterigmatocystin, a carcinogenic mycotoxin, and its isolation from two new fungal sources. *S. Afr. Med. J.* **40**:1100–1101.
 20. Hopwood, D. A., and D. H. Sherman. 1990. Molecular genetics of polyketides and its comparison to fatty acid biosynthesis. *Annu. Rev. Genet.* **24**:37–66.
 21. Jelinek, C. F., A. E. Pohland, and G. E. Wood. 1989. Review of mycotoxin contamination. Worldwide occurrence of mycotoxins in foods and feeds—an update. *J. Assoc. Off. Anal. Chem.* **72**:223–230.
 - 21a. Keller, N. P., and T. H. Adams. Unpublished data.
 22. Keller, N. P., T. E. Cleveland, and D. Bhatnagar. 1992. Variable electrophoretic karyotypes in members of *Aspergillus* section *flavi*. *Curr. Genet.* **21**:371–375.
 23. Keller, N. P., H. C. Dischinger, Jr., D. Bhatnagar, T. E. Cleveland, and A. H. J. Ullah. 1993. Purification of a 40-kilodalton methyltransferase active in the aflatoxin biosynthetic pathway. *Appl. Environ. Microbiol.* **59**:479–484.
 24. Lafont, P., M. G. Siriwardana, and E. DeBoer. 1990. Contamination of dairy products by fungal metabolites. *J. Environ. Pathol. Toxicol. Oncol.* **10**:99–102.
 25. Maessen, G. D. F., R. Amons, J. P. Zeelen, and W. Moeller. 1987. Primary structure of elongation factor 1-gamma from *Artemia*. *FEBS Lett.* **223**:181–186.
 - 25a. May, G. Personal communication.
 26. Miller, B., K. Miller, and W. E. Timberlake. 1985. Direct and indirect gene replacements in *Aspergillus nidulans*. *Mol. Cell. Biol.* **5**:1714–1721.
 27. Mori, H. 1992. Genotoxicity of naturally occurring metabolites: structural analogs of aflatoxin and related chemicals, p. 231–253. *In* D. Bhatnagar, E. B. Lillehoj, and D. K. Arora (ed.), *Handbook of applied mycology*. Marcel Dekker, Inc., New York.
 28. Northolt, M. D., H. P. Van Egmond, P. Soentoro, and E. Deijll. 1980. Fungal growth and the presence of sterigmatocystin in hard cheese. *J. Assoc. Off. Anal. Chem.* **63**:115–119.
 29. Payne, G. A., G. J. Nystrom, D. Bhatnagar, T. E. Cleveland, and C. P. Woloshuk. 1993. Cloning of the *aft-2* gene involved in aflatoxin biosynthesis from *Aspergillus flavus*. *Appl. Environ. Microbiol.* **59**:156–162.
 - 29a. Prade, R., J. Arnold, and W. E. Timberlake. Personal communication.
 30. Salvo, J. J., D. P. Mobley, D. W. Brown, L. A. Caruso, A. P. Yake, J. L. Spivack, and D. K. Dietrich. 1990. Biosynthesis of ρ -hydroxylated aromatics. *Biotechnol. Prog.* **6**:193–197.
 31. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 32. Scott, P. M., W. van Walbeek, B. Kennedy, and D. Anyeti. 1972. Mycotoxins (ochratoxin A, citrinin and sterigmatocystin) and toxigenic fungi in grains and other agricultural products. *J. Agric. Food Chem.* **20**:1103–1109.
 33. Shimada, T., and F. P. Guengerich. 1989. Evidence for cytochrome P-450NF, the nifedipine oxidase, being the principal enzyme involved in the bioactivation of aflatoxins in human liver. *Proc. Natl. Acad. Sci. USA* **86**:462–465.
 34. Skory, C. D., P.-K. Chang, J. Cary, and J. E. Linz. 1992. Isolation and characterization of a gene from *Aspergillus parasiticus* associated with the conversion of versicolorin A to sterigmatocystin in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **58**:3527–3537.
 35. Skory, C. D., P.-K. Chang, and J. E. Linz. 1993. Regulated expression of the *nor-1* and *ver-1* genes associated with aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **59**:1642–1646.
 36. Smith, D. J., M. K. R. Burnham, J. H. Bull, J. E. Hodgson, M. J. Ward, P. Browne, J. Brown, B. Barton, A. J. Earl, and G. Turner. 1990. β -Lactam antibiotic biosynthetic genes have been conserved in clusters in prokaryotes and eukaryotes. *EMBO J.* **9**:741–747.
 37. Sram, R. J., B. Binkova, L. Dobias, P. Rossner, J. Topinka, D. Vesela, D. Vesely, J. Stejskalova, H. Bavorova, and V. Rericha. 1993. Monitoring genotoxic exposure in uranium miners. *Environ. Health Perspect.* **99**:303–305.
 38. Stack, M., and J. V. Rodricks. 1971. Method for analysis and chemical confirmation of sterigmatocystin. *J. Assoc. Off. Anal. Chem.* **54**:86–90.
 39. Timberlake, W. E. 1991. Cloning and analysis of fungal genes, p. 51–83. *In* M. Alic, J. W. Bennett, and L. L. Lasure (ed.), *More gene manipulations in fungi*. Academic Press, San Diego, Calif.
 40. Wu, T.-S., and J. E. Linz. 1993. Recombinational inactivation of the gene encoding nitrate reductase in *Aspergillus parasiticus*. *Appl. Environ. Microbiol.* **59**:2998–3002.
 41. Yabe, K., and T. Hamasaki. 1993. Stereochemistry during aflatoxin biosynthesis: cyclase reaction in the conversion of versiconal to versicolorin B and racemization of versiconal hemiacetal acetate. *Appl. Environ. Microbiol.* **59**:2493–2500.
 42. Yokotani, N., R. Bernhardt, K. Sogawa, E. Kusunose, O. Gotoh, M. Kusunose, and Y. Fujii-Kuriyama. 1989. Two forms of omega-hydroxylase toward prostaglandin A and laurate. cDNA cloning and their expression. *J. Biol. Chem.* **264**:21665–21669.
 43. Yu, J., J. W. Cary, D. Bhatnagar, T. E. Cleveland, N. P. Keller, and F. S. Chu. 1993. Molecular cloning and characterization of an *Aspergillus parasiticus* gene involved in the conversion of sterigmatocystin to O-methylsterigmatocystin in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* **59**:3564–3571.