



# MIT Open Access Articles

## *Search for Resonant Top-Antitop Production in the Lepton Plus Jets Decay Mode Using the Full CDF Data Set*

The MIT Faculty has made this article openly available. ***Please share*** how this access benefits you. Your story matters.

<b>Citation</b>	Aaltonen, T., S. Amerio, D. Amidei, et al. 2013 Search for Resonant Top-Antitop Production in the Lepton Plus Jets Decay Mode Using the Full CDF Data Set. Physical Review Letters 110(12). © 2013 American Physical Society.
<b>As Published</b>	<a href="http://dx.doi.org/10.1103/PhysRevLett.110.121802">http://dx.doi.org/10.1103/PhysRevLett.110.121802</a>
<b>Publisher</b>	American Physical Society
<b>Version</b>	Final published version
<b>Accessed</b>	Wed Sep 12 09:31:38 EDT 2018
<b>Citable Link</b>	<a href="http://hdl.handle.net/1721.1/81294">http://hdl.handle.net/1721.1/81294</a>
<b>Terms of Use</b>	Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.
<b>Detailed Terms</b>	

## Search for Resonant Top-Antitop Production in the Lepton Plus Jets Decay Mode Using the Full CDF Data Set

T. Aaltonen,<sup>21</sup> S. Amerio,<sup>40a</sup> D. Amidei,<sup>32</sup> A. Anastassov,<sup>15,w</sup> A. Annovi,<sup>17</sup> J. Antos,<sup>12</sup> G. Apollinari,<sup>15</sup> J. A. Appel,<sup>15</sup> T. Arisawa,<sup>53</sup> A. Artikov,<sup>13</sup> J. Asaadi,<sup>48</sup> W. Ashmanskas,<sup>15</sup> B. Auerbach,<sup>2</sup> A. Aurisano,<sup>48</sup> F. Azfar,<sup>39</sup> W. Badgett,<sup>15</sup> T. Bae,<sup>25</sup> A. Barbaro-Galtieri,<sup>26</sup> V. E. Barnes,<sup>44</sup> B. A. Barnett,<sup>23</sup> P. Barria,<sup>42c,42a</sup> P. Bartos,<sup>12</sup> M. Baucus,<sup>40b,40a</sup> F. Bedeschi,<sup>42a</sup> S. Behari,<sup>15</sup> G. Bellettini,<sup>42b,42a</sup> J. Bellinger,<sup>55</sup> D. Benjamin,<sup>14</sup> A. Beretvas,<sup>15</sup> A. Bhatti,<sup>46</sup> K. R. Bland,<sup>5</sup> B. Blumenfeld,<sup>23</sup> A. Bocci,<sup>14</sup> A. Bodek,<sup>45</sup> D. Bortoletto,<sup>44</sup> J. Boudreau,<sup>43</sup> A. Boveia,<sup>11</sup> L. Brigliadori,<sup>6b,6a</sup> C. Bromberg,<sup>33</sup> E. Brucken,<sup>21</sup> J. Budagov,<sup>13</sup> H. S. Budd,<sup>45</sup> K. Burkett,<sup>15</sup> G. Busetto,<sup>40b,40a</sup> P. Bussey,<sup>19</sup> P. Butti,<sup>42b,42a</sup> A. Buzatu,<sup>19</sup> A. Calamba,<sup>10</sup> S. Camarda,<sup>4</sup> M. Campanelli,<sup>28</sup> F. Canelli,<sup>11,15,dd</sup> B. Carls,<sup>22</sup> D. Carlsmith,<sup>55</sup> R. Carosi,<sup>42a</sup> S. Carrillo,<sup>16,1</sup> B. Casal,<sup>9j</sup> M. Casarsa,<sup>49a</sup> A. Castro,<sup>6b,6a</sup> P. Catastini,<sup>20</sup> D. Cauz,<sup>49a</sup> V. Cavaliere,<sup>22</sup> M. Cavalli-Sforza,<sup>4</sup> A. Cerri,<sup>26,e</sup> L. Cerrito,<sup>28,r</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>42a</sup> G. Chlachidze,<sup>15</sup> K. Cho,<sup>25</sup> D. Chokheli,<sup>13</sup> M. A. Ciocci,<sup>42c,42a</sup> A. Clark,<sup>18</sup> C. Clarke,<sup>54</sup> M. E. Convery,<sup>15</sup> J. Conway,<sup>7</sup> M. Corbo,<sup>15</sup> M. Cordelli,<sup>17</sup> C. A. Cox,<sup>7</sup> D. J. Cox,<sup>7</sup> M. Cremonesi,<sup>42a</sup> D. Cruz,<sup>48</sup> J. Cuevas,<sup>9,y</sup> R. Culbertson,<sup>15</sup> N. d'Ascenzo,<sup>15,v</sup> M. Datta,<sup>15,ff</sup> P. De Barbaro,<sup>45</sup> L. Demortier,<sup>46</sup> M. Deninno,<sup>6a</sup> F. Devoto,<sup>21</sup> M. d'Errico,<sup>40b,40a</sup> A. Di Canto,<sup>42b,42a</sup> B. Di Ruzza,<sup>15,p</sup> J. R. Dittmann,<sup>5</sup> M. D'Onofrio,<sup>27</sup> S. Donati,<sup>42b,42a</sup> M. Dorigo,<sup>49b,49a</sup> A. Driutti,<sup>49a</sup> K. Ebina,<sup>53</sup> R. Edgar,<sup>32</sup> A. Elagin,<sup>48</sup> R. Erbacher,<sup>7</sup> S. Errede,<sup>22</sup> B. Esham,<sup>22</sup> R. Eusebi,<sup>48</sup> S. Farrington,<sup>39</sup> J. P. Fernández Ramos,<sup>29</sup> R. Field,<sup>16</sup> G. Flanagan,<sup>15,t</sup> R. Forrest,<sup>7</sup> M. Franklin,<sup>20</sup> J. C. Freeman,<sup>15</sup> H. Frisch,<sup>11</sup> Y. Funakoshi,<sup>53</sup> A. F. Garfinkel,<sup>44</sup> P. Garosi,<sup>42c,42a</sup> H. Gerberich,<sup>22</sup> E. Gerchtein,<sup>15</sup> S. Giagu,<sup>47a</sup> V. Giakoumopoulou,<sup>3</sup> K. Gibson,<sup>43</sup> C. M. Ginsburg,<sup>15</sup> N. Giokaris,<sup>3</sup> P. Giromini,<sup>17</sup> G. Giurgiu,<sup>23</sup> V. Glagolev,<sup>13</sup> D. Glenzinski,<sup>15</sup> M. Gold,<sup>35</sup> D. Goldin,<sup>48</sup> A. Golossanov,<sup>15</sup> G. Gomez,<sup>9</sup> G. Gomez-Ceballos,<sup>30</sup> M. Goncharov,<sup>30</sup> O. González López,<sup>29</sup> I. Gorelov,<sup>35</sup> A. T. Goshaw,<sup>14</sup> K. Goulianos,<sup>46</sup> E. Gramellini,<sup>6a</sup> S. Grinstein,<sup>4</sup> C. Grosso-Pilcher,<sup>11</sup> R. C. Group,<sup>52,15</sup> J. Guimaraes da Costa,<sup>20</sup> S. R. Hahn,<sup>15</sup> J. Y. Han,<sup>45</sup> F. Happacher,<sup>17</sup> K. Hara,<sup>50</sup> M. Hare,<sup>51</sup> R. F. Harr,<sup>54</sup> T. Harrington-Taber,<sup>15,m</sup> K. Hatakeyama,<sup>5</sup> C. Hays,<sup>39</sup> J. Heinrich,<sup>41</sup> M. Herndon,<sup>55</sup> A. Hocker,<sup>15</sup> Z. Hong,<sup>48</sup> W. Hopkins,<sup>15,f</sup> S. Hou,<sup>1</sup> R. E. Hughes,<sup>36</sup> U. Husemann,<sup>56</sup> M. Hussein,<sup>33</sup> J. Huston,<sup>33</sup> G. Introzzi,<sup>42e,42a</sup> M. Iori,<sup>47b,47a</sup> A. Ivanov,<sup>7,o</sup> E. James,<sup>15</sup> D. Jang,<sup>10</sup> B. Jayatilaka,<sup>15</sup> E. J. Jeon,<sup>25</sup> S. Jindariani,<sup>15</sup> M. Jones,<sup>44</sup> K. K. Joo,<sup>25</sup> S. Y. Jun,<sup>10</sup> T. R. Junk,<sup>15</sup> M. Kambeitz,<sup>24</sup> T. Kamon,<sup>25,48</sup> P. E. Karchin,<sup>54</sup> A. Kasmi,<sup>5</sup> Y. Kato,<sup>38,n</sup> W. Ketchum,<sup>11,gg</sup> J. Keung,<sup>41</sup> B. Kilminster,<sup>15,dd</sup> D. H. Kim,<sup>25</sup> H. S. Kim,<sup>25</sup> J. E. Kim,<sup>25</sup> M. J. Kim,<sup>17</sup> S. B. Kim,<sup>25</sup> S. H. Kim,<sup>50</sup> Y. K. Kim,<sup>11</sup> Y. J. Kim,<sup>25</sup> N. Kimura,<sup>53</sup> M. Kirby,<sup>15</sup> K. Knoepfel,<sup>15</sup> K. Kondo,<sup>53,a</sup> D. J. Kong,<sup>25</sup> J. Konigsberg,<sup>16</sup> A. V. Kotwal,<sup>14</sup> M. Kreps,<sup>24</sup> J. Kroll,<sup>41</sup> M. Kruse,<sup>14</sup> T. Kuhr,<sup>24</sup> M. Kurata,<sup>50</sup> A. T. Laasanen,<sup>44</sup> S. Lammel,<sup>15</sup> M. Lancaster,<sup>28</sup> K. Lannon,<sup>36,x</sup> G. Latino,<sup>42c,42a</sup> H. S. Lee,<sup>25</sup> J. S. Lee,<sup>25</sup> S. Leo,<sup>42a</sup> S. Leone,<sup>42a</sup> J. D. Lewis,<sup>15</sup> A. Limosani,<sup>14,s</sup> E. Lipeles,<sup>41</sup> H. Liu,<sup>52</sup> Q. Liu,<sup>44</sup> T. Liu,<sup>15</sup> S. Lockwitz,<sup>56</sup> A. Loginov,<sup>56</sup> D. Lucchesi,<sup>40b,40a</sup> J. Lueck,<sup>24</sup> P. Lujan,<sup>26</sup> P. Lukens,<sup>15</sup> G. Lungu,<sup>46</sup> J. Lys,<sup>26</sup> R. Lysak,<sup>12,d</sup> R. Madrak,<sup>15</sup> P. Maestro,<sup>42c,42a</sup> S. Malik,<sup>46</sup> G. Manca,<sup>27,b</sup> A. Manousakis-Katsikakis,<sup>3</sup> F. Margaroli,<sup>47a</sup> P. Marino,<sup>42d,42a</sup> M. Martínez,<sup>4</sup> K. Matera,<sup>22</sup> M. E. Mattson,<sup>54</sup> A. Mazzacane,<sup>15</sup> P. Mazzanti,<sup>6a</sup> R. McNulty,<sup>27,i</sup> A. Mehta,<sup>27</sup> P. Mehtala,<sup>21</sup> C. Mesropian,<sup>46</sup> T. Miao,<sup>15</sup> D. Mietlicki,<sup>32</sup> A. Mitra,<sup>1</sup> H. Miyake,<sup>50</sup> S. Moed,<sup>15</sup> N. Moggi,<sup>6a</sup> C. S. Moon,<sup>15,z</sup> R. Moore,<sup>15,ee</sup> M. J. Morello,<sup>42d,42a</sup> A. Mukherjee,<sup>15</sup> Th. Muller,<sup>24</sup> P. Murat,<sup>15</sup> M. Mussini,<sup>6b,6a</sup> J. Nachtman,<sup>15,m</sup> Y. Nagai,<sup>50</sup> J. Naganoma,<sup>53</sup> I. Nakano,<sup>37</sup> A. Napier,<sup>51</sup> J. Nett,<sup>48</sup> C. Neu,<sup>52</sup> T. Nigmanov,<sup>43</sup> L. Nodulman,<sup>2</sup> S. Y. Noh,<sup>25</sup> O. Norniella,<sup>22</sup> L. Oakes,<sup>39</sup> S. H. Oh,<sup>14</sup> Y. D. Oh,<sup>25</sup> I. Oksuzian,<sup>52</sup> T. Okusawa,<sup>38</sup> R. Orava,<sup>21</sup> L. Ortolan,<sup>4</sup> C. Pagliarone,<sup>49a</sup> E. Palencia,<sup>9,e</sup> P. Palni,<sup>35</sup> V. Papadimitriou,<sup>15</sup> W. Parker,<sup>55</sup> G. Pauletta,<sup>49c,49a</sup> M. Paulini,<sup>10</sup> C. Paus,<sup>30</sup> T. J. Phillips,<sup>14</sup> G. Piacentino,<sup>42a</sup> E. Pianori,<sup>41</sup> J. Pilot,<sup>36</sup> K. Pitts,<sup>22</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>55</sup> S. Poprocki,<sup>15,f</sup> K. Potamianos,<sup>26</sup> F. Prokoshin,<sup>13,aa</sup> A. Pranko,<sup>26</sup> F. Ptohos,<sup>17,g</sup> G. Punzi,<sup>42b,42a</sup> N. Ranjan,<sup>44</sup> I. Redondo Fernández,<sup>29</sup> P. Renton,<sup>39</sup> M. Rescigno,<sup>47a</sup> T. Riddick,<sup>28</sup> F. Rimondi,<sup>6a,a</sup> L. Ristori,<sup>42a,15</sup> A. Robson,<sup>19</sup> T. Rodriguez,<sup>41</sup> S. Rolli,<sup>51,h</sup> M. Ronzani,<sup>42b,42a</sup> R. Roser,<sup>15</sup> J. L. Rosner,<sup>11</sup> F. Ruffini,<sup>42c,42a</sup> A. Ruiz,<sup>9</sup> J. Russ,<sup>10</sup> V. Rusu,<sup>15</sup> A. Safonov,<sup>48</sup> W. K. Sakumoto,<sup>45</sup> Y. Sakurai,<sup>53</sup> L. Santi,<sup>49c,49a</sup> K. Sato,<sup>50</sup> V. Saveliev,<sup>15,v</sup> A. Savoy-Navarro,<sup>15,z</sup> P. Schlabach,<sup>15</sup> E. E. Schmidt,<sup>15</sup> T. Schwarz,<sup>32</sup> L. Scodellaro,<sup>9</sup> F. Scuri,<sup>42a</sup> S. Seidel,<sup>35</sup> Y. Seiya,<sup>38</sup> A. Semenov,<sup>13</sup> F. Sforza,<sup>42b,42a</sup> S. Z. Shalhout,<sup>7</sup> T. Shears,<sup>27</sup> P. F. Shepard,<sup>43</sup> M. Shimojima,<sup>50,u</sup> M. Shochet,<sup>11</sup> I. Shreyber-Tecker,<sup>34</sup> A. Simonenko,<sup>13</sup> P. Sinervo,<sup>31</sup> K. Sliwa,<sup>51</sup> J. R. Smith,<sup>7</sup> F. D. Snider,<sup>15</sup> V. Sorin,<sup>4</sup> H. Song,<sup>43</sup> M. Stancari,<sup>15</sup> R. St. Denis,<sup>19</sup> B. Stelzer,<sup>31</sup> O. Stelzer-Chilton,<sup>31</sup> D. Stentz,<sup>15,w</sup> J. Strologas,<sup>35</sup> Y. Sudo,<sup>50</sup> A. Sukhanov,<sup>15</sup> I. Suslov,<sup>13</sup> K. Takemasa,<sup>50</sup> Y. Takeuchi,<sup>50</sup> J. Tang,<sup>11</sup> M. Tecchio,<sup>32</sup> P. K. Teng,<sup>1</sup> J. Thom,<sup>15,f</sup> E. Thomson,<sup>41</sup> V. Thukral,<sup>48</sup> D. Toback,<sup>48</sup> S. Tokar,<sup>12</sup> K. Tollefson,<sup>33</sup> T. Tomura,<sup>50</sup> D. Tonelli,<sup>15,e</sup> S. Torre,<sup>17</sup> D. Torretta,<sup>15</sup> P. Totaro,<sup>40a</sup> M. Trovato,<sup>42d,42a</sup> F. Ukegawa,<sup>50</sup> S. Uozumi,<sup>25</sup> F. Vázquez,<sup>16,l</sup> G. Velev,<sup>15</sup> C. Vellidis,<sup>15</sup> C. Vernieri,<sup>42d,42a</sup> M. Vidal,<sup>44</sup> R. Vilar,<sup>9</sup> J. Vizán,<sup>9,cc</sup> M. Vogel,<sup>35</sup> G. Volpi,<sup>17</sup>

P. Wagner,<sup>41</sup> R. Wallny,<sup>8</sup> S. M. Wang,<sup>1</sup> A. Warburton,<sup>31</sup> D. Waters,<sup>28</sup> W. C. Wester III,<sup>15</sup> D. Whiteson,<sup>41,c</sup> A. B. Wicklund,<sup>2</sup> S. Wilbur,<sup>11</sup> H. H. Williams,<sup>41</sup> J. S. Wilson,<sup>32</sup> P. Wilson,<sup>15</sup> B. L. Winer,<sup>36</sup> P. Wittich,<sup>15,f</sup> S. Wolbers,<sup>15</sup> H. Wolfe,<sup>36</sup> T. Wright,<sup>32</sup> X. Wu,<sup>18</sup> Z. Wu,<sup>5</sup> K. Yamamoto,<sup>38</sup> D. Yamato,<sup>38</sup> T. Yang,<sup>15</sup> U. K. Yang,<sup>11,q</sup> Y. C. Yang,<sup>25</sup> W.-M. Yao,<sup>26</sup> G. P. Yeh,<sup>15</sup> K. Yi,<sup>15,m</sup> J. Yoh,<sup>15</sup> K. Yorita,<sup>53</sup> T. Yoshida,<sup>38,k</sup> G. B. Yu,<sup>14</sup> I. Yu,<sup>25</sup> A. M. Zanetti,<sup>49a</sup> Y. Zeng,<sup>14</sup> C. Zhou,<sup>14</sup> and S. Zucchelli<sup>6b,6a</sup>

(CDF Collaboration)<sup>bb</sup>

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*

<sup>3</sup>*University of Athens, 157 71 Athens, Greece*

<sup>4</sup>*Institut de Física d'Altes Energies, ICREA, Universitat Autònoma de Barcelona, E-08193 Bellaterra (Barcelona), Spain*

<sup>5</sup>*Baylor University, Waco, Texas 76798, USA*

<sup>6a</sup>*Istituto Nazionale di Fisica Nucleare Bologna, I-40127 Bologna, Italy*

<sup>6b</sup>*University of Bologna, I-40127 Bologna, Italy*

<sup>7</sup>*University of California, Davis, Davis, California 95616, USA*

<sup>8</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*

<sup>9</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

<sup>10</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*

<sup>11</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*

<sup>12</sup>*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

<sup>13</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

<sup>14</sup>*Duke University, Durham, North Carolina 27708, USA*

<sup>15</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*

<sup>16</sup>*University of Florida, Gainesville, Florida 32611, USA*

<sup>17</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

<sup>18</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*

<sup>19</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

<sup>20</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

<sup>21</sup>*Department of Physics, Division of High Energy Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

<sup>22</sup>*University of Illinois, Urbana, Illinois 61801, USA*

<sup>23</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*

<sup>24</sup>*Institut für Experimentelle Kernphysik, Karlsruhe Institute of Technology, D-76131 Karlsruhe, Germany*

<sup>25</sup>*Center for High Energy Physics: Kyungpook National University, Daegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea; Sungkyunkwan University, Suwon 440-746, Korea; Korea Institute of Science and Technology Information, Daejeon 305-806, Korea; Chonnam National University, Gwangju 500-757, Korea; Chonbuk National University, Jeonju 561-756, Korea; Ewha Womans University, Seoul 120-750, Korea*

<sup>26</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

<sup>27</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*

<sup>28</sup>*University College London, London WC1E 6BT, United Kingdom*

<sup>29</sup>*Centro de Investigaciones Energéticas Medioambientales y Tecnológicas, E-28040 Madrid, Spain*

<sup>30</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

<sup>31</sup>*Institute of Particle Physics, McGill University, Montréal, Québec H3A 2T8, Canada; Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada; University of Toronto, Toronto, Ontario M5S 1A7, Canada; and TRIUMF, Vancouver, British Columbia V6T 2A3, Canada*

<sup>32</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*

<sup>33</sup>*Michigan State University, East Lansing, Michigan 48824, USA*

<sup>34</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*

<sup>35</sup>*University of New Mexico, Albuquerque, New Mexico 87131, USA*

<sup>36</sup>*The Ohio State University, Columbus, Ohio 43210, USA*

<sup>37</sup>*Okayama University, Okayama 700-8530, Japan*

<sup>38</sup>*Osaka City University, Osaka 588, Japan*

<sup>39</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*

<sup>40a</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy*

<sup>40b</sup>*University of Padova, I-35131 Padova, Italy*

<sup>41</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*

<sup>42a</sup>*Istituto Nazionale di Fisica Nucleare Pisa, I-56127 Pisa, Italy*

<sup>42b</sup>*University of Pisa, I-56127 Pisa, Italy*

- <sup>42c</sup>University of Siena, I-56127 Pisa, Italy  
<sup>42d</sup>Scuola Normale Superiore, I-56127 Pisa, Italy,  
<sup>42e</sup>INFN Pavia and University of Pavia, I-27100 Pavia, Italy  
<sup>43</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA  
<sup>44</sup>Purdue University, West Lafayette, Indiana 47907, USA  
<sup>45</sup>University of Rochester, Rochester, New York 14627, USA  
<sup>46</sup>The Rockefeller University, New York, New York 10065, USA  
<sup>47a</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, I-00185 Roma, Italy  
<sup>47b</sup>Sapienza Università di Roma, I-00185 Roma, Italy  
<sup>48</sup>Texas A&M University, College Station, Texas 77843, USA  
<sup>49a</sup>Istituto Nazionale di Fisica Nucleare Trieste/Udine, I-33100 Udine, Italy  
<sup>49b</sup>University of Trieste, I-34127 Trieste, Italy  
<sup>49c</sup>University of Udine, I-33100 Udine, Italy  
<sup>50</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan  
<sup>51</sup>Tufts University, Medford, Massachusetts 02155, USA  
<sup>52</sup>University of Virginia, Charlottesville, Virginia 22906, USA  
<sup>53</sup>Waseda University, Tokyo 169, Japan  
<sup>54</sup>Wayne State University, Detroit, Michigan 48201, USA  
<sup>55</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>56</sup>Yale University, New Haven, Connecticut 06520, USA  
(Received 25 November 2012; published 20 March 2013)

This Letter reports a search for a narrow resonant state decaying into two  $W$  bosons and two  $b$  quarks where one  $W$  boson decays leptonically and the other decays into a quark-antiquark pair. The search is particularly sensitive to top-antitop resonant production. We use the full data sample of proton-antiproton collisions at a center-of-mass energy of 1.96 TeV collected by the CDF II detector at the Fermilab Tevatron, corresponding to an integrated luminosity of  $9.45 \text{ fb}^{-1}$ . No evidence for resonant production is found, and upper limits on the production cross section times branching ratio for a narrow resonant state are extracted. Within a specific benchmark model, we exclude a  $Z'$  boson with mass,  $M_{Z'}$ , below  $915 \text{ GeV}/c^2$  decaying into a top-antitop pair at the 95% credibility level assuming a  $Z'$  boson decay width of  $\Gamma_{Z'} = 0.012M_{Z'}$ . This is the most sensitive search for a narrow  $q\bar{q}$ -initiated  $t\bar{t}$  resonance in the mass region below  $750 \text{ GeV}/c^2$ .

DOI: [10.1103/PhysRevLett.110.121802](https://doi.org/10.1103/PhysRevLett.110.121802)

PACS numbers: 13.85.Rm, 14.65.Ha, 14.70.Pw, 14.80.Tt

The large mass of the top quark, compared to that of the other fundamental particles, gives it a special position within the standard model (SM). Since its discovery [1], the top quark has played an important role in theoretical extensions beyond the standard model [2]. Recently, renewed interest has been directed toward searches including top quark final states for beyond the standard model physics due to discrepancies reported in the  $t\bar{t}$  forward-backward asymmetry [3,4]. Moreover, the most recent search for resonant  $t\bar{t}$  production from D0 [5] reports an approximately  $2\sigma$  excess of events at resonant-mass values around  $950 \text{ GeV}/c^2$ . Many beyond the standard model theories [6–10] predict heavy resonances that add a resonant component to the SM  $t\bar{t}$  production mechanism.

Top quarks decay via the weak interaction, nearly always into a  $W$  boson and a  $b$  quark. The  $W$  boson then decays into a lighter fermion-antifermion pair [11]. We search for resonant production of top quark pairs followed by decays into a final state with one charged lepton (electron or muon) and multiple jets, where one of the  $W$  bosons decays leptonically and the other  $W$  boson decays hadronically. This lepton + jets channel features a distinctive final

state due to the presence of a charged lepton and has a branching ratio of 29% [12].

Unlike previous searches at CDF [13–16], we do not apply constraints based on the presence of top quarks in the event. While we focus the discussion on  $t\bar{t}$  resonances, we construct the top-antitop mass  $M_{t\bar{t}}$  used as a final search discriminant by taking the invariant mass of all objects (lepton, jets, and missing  $E_T$ ) in the event including those that may not originate from top quark production. Other than the event selection defined below, which provides a sample primarily composed of  $t\bar{t}$  events, there are no requirements that the event be consistent with  $t\bar{t}$  production. This results in a more general search that is sensitive not only to  $t\bar{t}$ , but also to any heavy narrow resonance decaying into a final state with a  $W$  boson and three or more jets with one or two jets originating from a  $b$  quark.

As a benchmark model, we consider a specific SM extension, top-color-assisted technicolor [17,18]. This model explains the large mass of the top quark through the introduction of new strong dynamics and also predicts a vector particle ( $Z'$  boson), which couples primarily to the third generation of quarks and has no significant couplings to leptons. The existence of a narrow-width  $Z'$  boson

resonance ( $\Gamma_{Z'} = 0.012M_{Z'}$ ) decaying to  $t\bar{t}$  pairs, using the leptophobic top-color model [18,19], has been searched for by both the CDF [13–16] and D0 [5,20,21] experiments at the Tevatron and also by the ATLAS [22–24] and CMS [25,26] experiments at the LHC. For resonance searches at the highest masses, the LHC experiments have superior sensitivity to the Tevatron due to the higher center-of-mass energy. However, in the lower-mass regions ( $m_{Z'} < 750 \text{ GeV}/c^2$ ) the Tevatron experiments have competitive sensitivity in searches for particles produced in  $q\bar{q}$ -initiated states, such as the  $Z'$  boson. While the production rate for the main background from SM  $t\bar{t}$  is approximately 25 times larger at a center-of-mass energy of 7 TeV, no valence antiquarks are available in the LHC  $pp$  collisions, so the signal production rate increases by a smaller factor relative to the  $p\bar{p}$  collisions of the Tevatron (between four and eight in the lower-mass region).

The collision events discussed in this Letter were produced at the Tevatron  $p\bar{p}$  collider at a center-of-mass energy of 1.96 TeV and were recorded by the CDF II detector [27]. The data sample corresponds to the full data set of the Tevatron, which comes from an integrated luminosity of  $9.45 \text{ fb}^{-1}$ . The CDF II detector consists of high-precision tracking systems for vertex and charged-particle track reconstruction, surrounded by electromagnetic and hadronic calorimeters for energy measurement, and muon subsystems outside the calorimeter for muon detection. CDF II uses a cylindrical coordinate system with azimuthal angle  $\phi$ , polar angle  $\theta$  measured with respect to the positive  $z$  direction along the proton beam, and the distance  $r$  measured from the beam line. The pseudorapidity, transverse energy, and transverse momentum are defined as  $\eta = -\ln[\tan(\frac{\theta}{2})]$ ,  $E_T = E\sin\theta$ , and  $p_T = p\sin\theta$ , respectively, where  $E$  and  $p$  are the energy and momentum of an outgoing particle, respectively. The missing transverse energy  $\vec{\cancel{E}}_T$  is defined by  $\vec{\cancel{E}}_T = -\sum_i E_T^i \hat{n}_i$ , where  $\hat{n}_i$  is a unit vector perpendicular to the beam axis that points to the  $i$ th calorimeter tower ( $\vec{\cancel{E}}_T = |\vec{\cancel{E}}_T|$ ).

The event selection and background estimation methods summarized below closely follow those that were employed in the observation of single top quark production [28] and in the search for the Higgs boson in the  $WH \rightarrow \ell\nu b\bar{b}$  final state [29]. The main difference is that the single top and  $WH$  analyses select events consistent with  $W$  + two or three jets and the current search requires three or more jets.

The data were collected by using online event selections (triggers) requiring one of the following energetic-lepton signatures: a high transverse momentum ( $p_T$ ) electron candidate, a high- $p_T$  muon candidate, or large  $\vec{\cancel{E}}_T$ . Significant  $\vec{\cancel{E}}_T$  can be produced when the neutrino from a leptonic  $W$  boson decay escapes detection.

Candidate events are selected by requiring a lepton candidate with  $p_T^\ell > 20 \text{ GeV}/c$ ,  $\vec{\cancel{E}}_T > 20 \text{ GeV}$ , and three or more jets with  $|\eta| < 2.0$  and  $E_T > 20 \text{ GeV}$  after

correcting the jet energies for instrumental effects [30,31]. One or two jets must be identified as being likely to have originated from a  $b$  quark according to the SECVTX [32] algorithm. This algorithm searches in the jet for a secondary vertex which results from the displaced decay of  $B$  hadrons. Events are rejected if more than one lepton candidate is reconstructed or if they are kinematically inconsistent with leptonic  $W$  boson decays [33]. Events with severely misreconstructed jets or leptons are removed based on angular correlations between the jet or lepton candidate and the  $\vec{\cancel{E}}_T$ .

Models for background processes are derived from a mixture of simulation- and data-driven techniques [28]. Important backgrounds in this final state include SM  $t\bar{t}$  production and other processes that include a  $W$  or  $Z$  boson in association with three or more jets. The events can include true  $b$ -quark jets, as in  $W$  boson +  $b\bar{b}j$  events, or jets that have been misidentified as  $b$ -quark jets, such as in  $W$  boson +  $c\bar{c}j$  and  $W$  boson +  $j\bar{j}j$  events, where  $j$  refers to jets not originating from heavy-flavor quarks. Multijet events without  $W$  bosons also contribute to the sample composition. Additional small background contributions are included from  $Z$  boson production with additional jets, diboson production, and single top quark production.

The expected rate for the SM  $t\bar{t}$  background is taken to be  $7.04 \pm 0.70 \text{ pb}$  [34] as calculated at approximate next-to-next-to-leading order using Martin-Stirling-Thorne-Watt 2008 parton distribution functions [35]. In order to predict the acceptance for nonresonant SM  $t\bar{t}$  events and their kinematic distributions, we use a sample of Monte Carlo events generated by using POWHEG [36] and assuming a top quark mass of  $172.5 \text{ GeV}/c^2$  [37] with parton showering provided by PYTHIA [38] followed by simulation of the CDF II detector [39,40]. The detection efficiency predicted by the simulation is corrected based on measurements using data for lepton identification, trigger efficiencies, and  $b$ -jet tagging efficiencies. The normalization for the QCD multijet and  $W$  boson + jet processes is obtained from a fit to the  $\vec{\cancel{E}}_T$  distribution. The background from events with mistakenly  $b$ -tagged light-flavor jets,  $W$  boson +  $j\bar{j}j$ , for example, is estimated by measuring the rate of such mistags in multijet data [32] and applying this rate to the  $W$  boson + jets data samples before  $b$  tagging. The contribution from true heavy-flavor production in the  $W$  boson + jets event sample is determined from measurements of the heavy-flavor event fraction in a  $W$  boson + 1 jet sample that is independent of the sample used in the resonance search. We model the kinematic distributions of  $W$  boson + jets events by using a combination of ALPGEN [41] matrix-element generation and PYTHIA parton showering. The QCD multijet background is modeled by using a sample of collision events in which one of the lepton identification requirements is inverted to obtain an enriched sample of QCD multijet events.

TABLE I. Summary of the background prediction and observed data for three-jet and four-or-more-jet events. The uncertainties include statistical and systematic contributions.

Process	3-jet events	$\geq 4$ -jet events
$t\bar{t}$	$1930 \pm 200$	$2570 \pm 270$
W/Z boson + jets	$2280 \pm 610$	$570 \pm 190$
Multijets	$150 \pm 60$	$130 \pm 100$
Total background	$4360 \pm 870$	$3270 \pm 560$
Observed	4254	3049

The background predictions are summarized in Table I. In this table and the following figures, we have divided the sample into events that include three jets and events that include four or more jets. For the statistical interpretation of the data, we further subdivide the events based on the number of  $b$ -tagged jets (one or two  $b$  tags) and based on the lepton type (lepton types that can be directly identified by the trigger, or leptons in events selected with the  $\cancel{E}_T$ -based trigger), yielding eight independent channels used to search for a resonance in the  $M_{\ell\bar{\ell}}$  distributions. The sensitivity of the search benefits from this subdivision, because the search subchannels have different background compositions, signal-to-background ratios, and invariant mass resolutions.

We use the invariant mass of all reconstructed objects in the event to discriminate between SM background and  $Z'$  boson signal events. For each event we calculate  $M_{\ell\bar{\ell}}$  by using the momenta of the three or more jets, the charged lepton, and the neutrino. The transverse momentum of the

neutrino is estimated from the  $\vec{\cancel{E}}_T$ . However, because the  $z$  component of the momenta of the scattering partons from the  $p\bar{p}$  collision is unknown, the final-state reconstructed energy need not be balanced in the  $z$  direction. The longitudinal component of the neutrino momentum ( $p_z^\nu$ ) is determined by solving  $M_W^2 = (p^l + p^\nu)^2$ , where  $M_W$ ,  $p^l$ , and  $p^\nu$  are the  $W$  boson mass, the lepton momentum, and the neutrino momentum, respectively. The smaller solution of the resulting quadratic equation is chosen for the  $p_z^\nu$ . If there is no real solution, we set  $p_z^\nu = 0$ . This approach is found to select the correct  $p_z^\nu$  in about 70% of simulated  $t\bar{t}$  events.

For the benchmark model, the  $Z'$  boson cross sections times branching fraction are based on leading-order predictions from Ref. [19] with an additional scaling factor of 1.3 applied to account for next-to-leading-order effects [42]. Signal  $Z'$  boson events are modeled with simulated events generated by PYTHIA in order to study the signal acceptance and to predict the  $M_{\ell\bar{\ell}}$  distributions.

A total of 4254 (3049) events survive the selection criteria for the three-jet (four-or-more-jet) category. The SM  $t\bar{t}$  contribution is estimated to be 43% (78%) for three-jet (four-or-more-jet) events. The remaining events are contributed primarily from the  $W$  boson + jet and QCD multijet processes plus a potential signal contribution from  $Z'$  boson events. The  $M_{\ell\bar{\ell}}$  distributions for the background model and events observed in the data are shown in Fig. 1. We fit the  $M_{\ell\bar{\ell}}$  distributions to what is observed in the data, allowing the background predictions to float within their systematic uncertainties. The  $M_{\ell\bar{\ell}}$  distribution for the




FIG. 1 (color online). Reconstructed  $M_{\ell\bar{\ell}}$  for the three-jet events (left) and four-or-more-jet events (right). The distribution of the  $M_{\ell\bar{\ell}}$  spectrum is on a linear scale, and the inset shows a logarithmic scale of the same distribution. The background expectation is normalized to the best fit to the data. The red histogram shows the expectation for a 600  $\text{GeV}/c^2$  mass hypothesis for the leptophobic top-color resonance normalized to the predicted cross section.

$Z'$  boson signal for the  $600 \text{ GeV}/c^2$  mass hypothesis is also included in Fig. 1.

We calculate a Bayesian credibility level (C.L.) limit on resonant  $t\bar{t}$  production for each mass hypothesis based on the binned observed  $M_{t\bar{t}}$  spectrum using the combined likelihood which includes the priors  $\pi(\vec{\theta})$  on the systematic uncertainties  $\vec{\theta}$ :

$$\mathcal{L}(R, \vec{s}, \vec{b}|\vec{n}, \vec{\theta})\pi(\vec{\theta}) = \prod_{i=1}^{N_C} \prod_{j=1}^{N_{\text{bins}}} \mu_{ij}^{n_{ij}} \frac{e^{-\mu_{ij}}}{n_{ij}!} \prod_{k=1}^{n_{\text{sys}}} e^{-\theta_k^2/2}. \quad (1)$$

In this expression, the first product is over the eight channels,  $N_C$ , we consider in this analysis. The second product is over histogram bins containing  $n_{ij}$  events. The predictions for the bin contents are  $\mu_{ij} = R \times s_{ij}(\vec{\theta}) + b_{ij}(\vec{\theta})$  for channel  $i$  and histogram bin  $j$ , where  $s_{ij}$  represents the potential resonant signal,  $b_{ij}$  is the expected background in the bin, and  $R$  is a scaling factor applied to the signal.

Systematic uncertainties are parametrized by the dependence of  $s_{ij}$  and  $b_{ij}$  on  $\vec{\theta}$ . Each of the  $n_{\text{sys}}$  components of  $\vec{\theta}$ ,  $\theta_k$ , corresponds to a single independent source of systematic uncertainty. We account for correlations by allowing each parameter to have an impact on several sources of signal and background in different channels. Gaussian priors are assumed for the  $\theta_k$ , which are truncated so that no prediction is negative. The likelihood function, multiplied by the  $\theta_k$  priors,  $\pi(\theta_k)$ , is then integrated over  $\theta_k$  including correlations [11]:

$$\mathcal{L}'(R) = \int \mathcal{L}(R, \vec{s}, \vec{b}|\vec{n}, \vec{\theta})\pi(\vec{\theta})d\vec{\theta}. \quad (2)$$

We assume a uniform prior in  $R$  to obtain its posterior distribution. The observed 95% C.L. upper limit on  $R$ ,  $R_{95}^{\text{obs}}$ , satisfies  $0.95 = \int_0^{R_{95}^{\text{obs}}} \mathcal{L}'(R)dR$ . The expected distribution of  $R_{95}$  is computed in an ensemble of pseudoexperiments generated without a signal. In each pseudoexperiment, values of the nuisance parameters are drawn from their priors. The median expected value of  $R_{95}$  in this ensemble is quoted as the expected limit. This statistical procedure is repeated for each resonance mass hypothesis from 350 to  $1200 \text{ GeV}/c^2$ .

We consider uncertainties that affect the normalization as well as uncertainties that affect the  $M_{t\bar{t}}$  distributions. The same set of uncertainties on the dominant background (SM  $t\bar{t}$  production) and the resonant signal are considered: They arise from the uncertainty in the jet energy scale [31], the  $b$ -tagging efficiency, the luminosity measurement [43], the lepton identification and trigger efficiency, and the rate of initial- and final-state radiation from the parton shower model. The jet energy scale,  $b$ -tag, and initial- and final-state radiation variations also affect the shape of the  $M_{t\bar{t}}$  distributions. The background normalizations for the QCD multijet background and for events with a  $W$  boson and heavy-flavor jet ( $b$  or  $c$ ) are assigned uncertainties due




FIG. 2 (color online). Expected and observed upper limits on the production cross section times the branching ratio for a leptophobic top-color  $Z'$  boson in  $9.45 \text{ fb}^{-1}$  of CDF data. The dashed line is the median expected upper limit with the assumption of no signal, the black points are the observed limit, and the blue line is the cross section prediction for leptophobic top-color  $Z'$  boson [19] production.

to limitations in their data-driven estimations [28]. Uncertainties on the renormalization and factorization scale used in the ALPGEN sample affect the shape of the  $M_{t\bar{t}}$  distributions from  $W$  boson + jets.

The resulting 95% C.L. upper limits on the cross section times the branching ratio for the leptophobic top-color  $Z'$  [19] signal hypotheses,  $\sigma_{Z'}\text{BR}(Z' \rightarrow t\bar{t})$ , as a function of  $M_{t\bar{t}}$  are shown in Fig. 2 and Table II together with expected limits derived from pseudoexperiments that include the SM

TABLE II. Expected, observed 95% C.L. upper limits and theoretical prediction for the production cross section times the branching ratio for a narrow  $t\bar{t}$  resonance, given as a function of  $Z'$  boson mass.

$M_{Z'}$ [GeV/ $c^2$ ]	Expected [pb]	Observed [pb]	$\sigma_{Z'}\text{BR}(Z' \rightarrow t\bar{t})$ [pb]
350	0.772	0.687	8.91
400	0.575	0.652	12.3
450	0.670	0.585	8.24
500	0.520	0.427	5.53
550	0.354	0.530	3.51
600	0.245	0.472	2.30
650	0.199	0.269	1.43
700	0.159	0.145	0.917
750	0.137	0.112	0.566
800	0.115	0.099	0.355
850	0.106	0.103	0.208
900	0.097	0.116	0.134
950	0.091	0.118	0.080
1000	0.092	0.129	0.049
1100	0.098	0.132	0.017
1200	0.134	0.166	0.006




FIG. 3 (color online). Tevatron and LHC sensitivity comparison to new physics for narrow  $t\bar{t}$  resonant final states, using the benchmark leptophobic top-color  $Z'$  model. The figure shows the expected upper limits on  $\sigma_{Z'}\text{BR}(Z' \rightarrow t\bar{t})$  divided by the theoretical prediction of  $\sigma_{Z'}\text{BR}(Z' \rightarrow t\bar{t})$  as a function of  $M_{Z'}$ . Dashed, dotted, and solid lines correspond to the expected upper limits of the most sensitive analyses to date from CDF, ATLAS, and CMS experiments, respectively.

background hypothesis only. A benchmark leptophobic top-color model [19] is excluded at 95% C.L. for  $Z'$  boson masses smaller than 915  $\text{GeV}/c^2$  assuming the width of the resonance is  $\Gamma_{Z'} = 0.012M_{Z'}$ . In addition, the limits reported here can be applied to any resonance producing the same final state as long as the decay width is significantly smaller than the reconstruction mass resolution ( $\Gamma_{Z'} \ll 0.15M_{Z'}$ ).

In conclusion, we have performed a search for a heavy resonance decaying into  $t\bar{t}$  using the lepton + jets decay channel in data from  $9.45 \text{ fb}^{-1}$  of integrated luminosity. The data are found to be consistent with the background expectation, and upper limits are set on the production cross section times branching ratio at the 95% C.L. For a specific benchmark model (leptophobic top-color), we exclude  $Z'$  bosons with masses up to 915  $\text{GeV}/c^2$ . As shown in Fig. 3, for masses smaller than approximately 750  $\text{GeV}/c^2$ , this search yields the most constraining limits to date on  $q\bar{q}$ -produced narrow  $t\bar{t}$  resonant states in the lepton + jets decay mode.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, United Kingdom; the Russian Foundation

for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; the Academy of Finland; and the Australian Research Council (ARC).

<sup>a</sup>Deceased.

<sup>b</sup>Visitor from Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy.

<sup>c</sup>Visitor from University of California Irvine, Irvine, CA 92697, USA.

<sup>d</sup>Visitor from Institute of Physics, Academy of Sciences of the Czech Republic, 182-21, Czech Republic.

<sup>e</sup>Visitor from CERN, CH-1211 Geneva, Switzerland.

<sup>f</sup>Visitor from Cornell University, Ithaca, NY 14853, USA.

<sup>g</sup>Visitor from University of Cyprus, Nicosia CY-1678, Cyprus.

<sup>h</sup>Visitor from Office of Science, U.S. Department of Energy, Washington, DC 20585, USA.

<sup>i</sup>Visitor from University College Dublin, Dublin 4, Ireland.

<sup>j</sup>Visitor from ETH, 8092 Zürich, Switzerland.

<sup>k</sup>Visitor from University of Fukui, Fukui City, Fukui Prefecture 910-0017, Japan.

<sup>l</sup>Visitor from Universidad Iberoamericana, Lomas de Santa Fe, México, C.P. 01219, Distrito Federal.

<sup>m</sup>Visitor from University of Iowa, Iowa City, IA 52242, USA.

<sup>n</sup>Visitor from Kinki University, Higashi-Osaka City 577-8502, Japan.

<sup>o</sup>Visitor from Kansas State University, Manhattan, KS 66506, USA.

<sup>p</sup>Visitor from Brookhaven National Laboratory, Upton, NY 11973, USA.

<sup>q</sup>Visitor from University of Manchester, Manchester M13 9PL, United Kingdom.

<sup>r</sup>Visitor from Queen Mary, University of London, London E1 4NS, United Kingdom.

<sup>s</sup>Visitor from University of Melbourne, Victoria 3010, Australia.

<sup>t</sup>Visitor from Muons, Inc., Batavia, IL 60510, USA.

<sup>u</sup>Visitor from Nagasaki Institute of Applied Science, Nagasaki 851-0193, Japan.

<sup>v</sup>Visitor from National Research Nuclear University, Moscow 115409, Russia.

<sup>w</sup>Visitor from Northwestern University, Evanston, IL 60208, USA.

<sup>x</sup>Visitor from University of Notre Dame, Notre Dame, IN 46556, USA.

<sup>y</sup>Visitor from Universidad de Oviedo, E-33007 Oviedo, Spain.

<sup>z</sup>Visitor from CNRS-IN2P3, Paris, F-75205 France.

<sup>aa</sup>Visitor from Universidad Tecnica Federico Santa Maria, 110v Valparaiso, Chile.

<sup>bb</sup>Visitor from Yarmouk University, Irbid 211-63, Jordan.

<sup>cc</sup>Visitor from Universite catholique de Louvain, 1348 Louvain-La-Neuve, Belgium.



- <sup>dd</sup>Visitor from University of Zürich, 8006 Zürich, Switzerland.
- <sup>ee</sup>Visitor from Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA.
- <sup>ff</sup>Visitor from Hampton University, Hampton, VA 23668, USA.
- <sup>gg</sup>Visitor from Los Alamos National Laboratory, Los Alamos, NM 87544, USA.
- [1] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **74**, 2626 (1995); S. Abachi *et al.* (D0 Collaboration), *ibid.* **74**, 2632 (1995).
- [2] C. T. Hill and S. J. Parke, *Phys. Rev. D* **49**, 4454 (1994).
- [3] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **83**, 112003 (2011).
- [4] J. F. Kamenik, J. Shu, and J. Zupan, *Eur. Phys. J. C* **72**, 2102 (2012).
- [5] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **85**, 051101 (2012).
- [6] R. Frederix and F. Maltoni, *J. High Energy Phys.* **01** (2009) 047.
- [7] B. A. Dobrescu, K. Kong, and R. Mahbubani, *J. High Energy Phys.* **06** (2009) 001.
- [8] M. I. Gresham, I. W. Kim, and K. M. Zurek, *Phys. Rev. D* **83**, 114027 (2011).
- [9] K. Agashe, A. Belyaev, T. Krupovnickas, G. Perez, and J. Virzi, *Phys. Rev. D* **77**, 015003 (2008).
- [10] Y. Bai, J. L. Hewett, J. Kaplan, and T. G. Rizzo, *J. High Energy Phys.* **03** (2011) 003.
- [11] K. Nakamura *et al.* (Particle Data Group), *J. Phys. G* **37**, 075021 (2010); J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012).
- [12] In addition to the 29% branching ratio, tau leptons are included when their decay produces an isolated electron or muon candidate.
- [13] T. Affolder *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **85**, 2062 (2000).
- [14] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **77**, 051102 (2008).
- [15] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **84**, 072003 (2011).
- [16] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **84**, 072004 (2011).
- [17] C. T. Hill, *Phys. Lett. B* **345**, 483 (1995).
- [18] R. M. Harris, C. T. Hill, and S. J. Parke, [arXiv:hep-ph/9911288](https://arxiv.org/abs/hep-ph/9911288).
- [19] R. M. Harris and S. Jain, *Eur. Phys. J. C* **72**, 2072 (2012); (private communication).
- [20] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **92**, 221801 (2004).
- [21] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **668**, 98 (2008).
- [22] G. Aad *et al.* (ATLAS Collaboration), *J. High Energy Phys.* **09** (2012) 041.
- [23] G. Aad *et al.* (ATLAS Collaboration), *Eur. Phys. J. C* **72**, 2083 (2012).
- [24] G. Aad *et al.* (ATLAS Collaboration), [arXiv:1211.2202](https://arxiv.org/abs/1211.2202).
- [25] S. Chatrchyan *et al.* (CMS Collaboration), *J. High Energy Phys.* **07** (2012) 029; [arXiv:1211.3338](https://arxiv.org/abs/1211.3338).
- [26] S. Chatrchyan *et al.* (CMS Collaboration), [arXiv:1209.4397](https://arxiv.org/abs/1209.4397).
- [27] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 032001 (2005).
- [28] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **82**, 112005 (2010).
- [29] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **86**, 032011 (2012).
- [30] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. D* **45**, 1448 (1992).
- [31] A. Bhatti *et al.* (CDF Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **566**, 2 (2006).
- [32] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. D* **71**, 052003 (2005).
- [33] F. Sforza, V. Lippi, and G. Chiarelli, *J. Phys. Conf. Ser.* **331**, 032045 (2011).
- [34] U. Langenfeld, S. Moch, and P. Uwer, *Phys. Rev. D* **80**, 054009 (2009).
- [35] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, *Eur. Phys. J. C* **63**, 189 (2009).
- [36] S. Frixione, P. Nason, and C. Oleari, *J. High Energy Phys.* **11** (2007) 070.
- [37] T. Aaltonen *et al.* (CDF and D0 Collaborations), *Phys. Rev. D* **86**, 092003 (2012).
- [38] T. Sjöstrand, P. Eden, C. Friberg, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, *Comput. Phys. Commun.* **135**, 238 (2001).
- [39] R. Brun, F. Bruyant, M. Maire, A. C. McPherson, and P. Zancarini, CERN Report No. CERN-DD-EE/84-1, 1987.
- [40] G. Grindhammer, M. Rudowicz, and S. Peters, *Nucl. Instrum. Methods Phys. Res., Sect. A* **290**, 469 (1990).
- [41] M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. D. Polosa, *J. High Energy Phys.* **07** (2003) 001.
- [42] J. Gao, C. S. Li, B. H. Li, H. X. Zhu, and C.-P. Yuan, *Phys. Rev. D* **82**, 014020 (2010); (private communication).
- [43] S. Klimenko, J. Konigsberg, and T. M. Liss, Report No. FERMILAB-FN-0741, 2003.