

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

PION PRODUCTION AS A PROBE OF THE NUCLEAR MATTER EQUATION OF STATE

### Permalink

<https://escholarship.org/uc/item/2gw629dt>

### Author

Harris, J.W.

### Publication Date

1984-02-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

MAY 29 1984


LIBRARY AND  
DOCUMENTS SECTION

Submitted to Physical Review Letters

PION PRODUCTION AS A PROBE OF THE NUCLEAR  
MATTER EQUATION OF STATE

J.W. Harris, R. Stock, R. Bock, R. Brockmann,  
A. Sandoval, H. Ströbele, G. Odyniec, H.G. Pugh,  
L.S. Schroeder, R.E. Renfordt, D. Schall,  
D. Bangert, W. Rauch, and K.L. Wolf

February 1984



LBL-17404  
c-2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Pion Production as a Probe of the Nuclear Matter Equation of State

J. W. Harris, R. Stock, R. Bock, R. Brockmann,  
A. Sandoval, H. Ströbele  
Gesellschaft für Schwerionenforschung  
D-6100, Darmstadt, West Germany

G. Odyniec, H. G. Pugh, L. S. Schroeder  
Lawrence Berkeley Laboratory, University of California  
Berkeley, California 94720

R. E. Renfordt, D. Schall  
Institut für Hochenergiephysik, Universität Heidelberg  
Heidelberg, West Germany

D. Bangert, W. Rauch  
Fachbereich Physik, Universität Marburg  
D-3550, Marburg, West Germany

and

K. L. Wolf  
Cyclotron Laboratory, Texas A&M University  
College Station, Texas 77843

Abstract:

It is shown that thermal and chemical equilibrium are approached during the high density stage in central nucleus-nucleus collisions and that the yield of produced pions is determined at that time. A chemical model with Rankine-Hugoniot compression is used to extract a nuclear matter equation of state from the observed pion yield assuming a partition of the internal energy per nucleon into thermal and compressional energy fractions.

PACS numbers: 25.70Bc, 23.70 Fg

The primary objective in studying relativistic nucleus-nucleus collisions is to determine the response of nuclear matter under conditions of high temperature and compression.<sup>1-3</sup> In order to relate observables to state variables, it is necessary to consider the rapid variation in space and time of the collision process. It has been suggested<sup>4</sup> that the time dependence of a reaction may be determined by studying the properties of particles emitted preferentially at various stages in the collision process. In our earlier work<sup>5</sup> the total pion multiplicity was proposed as an observable that is sensitive to the high density stage of central collisions where densities several times that of normal nuclear matter are expected to be reached. Since this stage occurs at the turning point between initial compression and later expansion when nuclear conditions and variables are relatively constant, it may survive long enough for equilibrium to be established. If so, the total pion multiplicity should be a sensitive measure<sup>6,7</sup> of the nuclear matter equation of state.

In this Letter the conditions for establishing chemical and thermal equilibrium in various stages of the collision are examined. A chemical model is used to study pion production and extract the nuclear compressional energy as a function of density. The results are in close agreement with our previous analysis which depended upon the validity of intranuclear cascade calculations, and lend credence to the extracted equation of state. The present method offers possibilities for extension to higher energies where cascade calculations face increasing difficulties.

In nucleus-nucleus collisions below  $E_{lab} \sim 2 \text{ GeV/u}$  the system can be considered as a mixture of nucleons, deltas, and pions. The approach to chemical equilibrium (in which the abundances of the constituents are in equilibrium) is determined by the inelastic cross sections for  $NN \leftrightarrow N\Delta \leftrightarrow NN\pi$ . The approximate rate constant for approach to chemical equilibrium of the  $\Delta$ 's is given by<sup>8</sup>

$$\lambda = \langle \sigma v \rangle \rho_N^2 / 2\rho_\Delta^{eq} \quad (1)$$

where  $\rho_N$  and  $\rho_\Delta^{eq}$  are the nucleon and  $\Delta$  densities in equilibrium.

The rate constant  $\lambda$  is proportional to the thermal energy average  $\langle \sigma v \rangle$  of the cross section for  $NN \leftrightarrow N\Delta$  times the relative pair velocity  $v$ . To

estimate  $\lambda$  for the high density stage it is assumed that  $T = 100 \text{ MeV}$ ,  $\rho_\Delta^{eq} = 0.2 \rho_N \approx 0.6 \rho_0$  with  $\rho_N = 3 \rho_0$ , and  $\langle \sigma v \rangle \approx 1.5 \times 10^{23} \text{ fm}^3 \text{ s}^{-1}$ .

The resulting equilibration time constant is  $\tau \equiv \lambda^{-1} \approx 6 \times 10^{-24} \text{ s}$ . Intra-nuclear cascade calculations have been used<sup>5,10</sup> to estimate the duration of the high density stage which is found to be  $2\tau$  and  $3.5\tau$  for central collisions of  $\text{Ar} + \text{KCl}$  and  $\text{La} + \text{La}$ , respectively, long enough for chemical equilibrium to be reached. To make a similar analysis for thermal equilibrium, i.e., equipartition of energy among the various kinetic degrees of freedom, the total cross sections must be used. The resulting time constant is even shorter, so that both thermal and chemical equilibrium exist at the end of the high density stage.

To discuss the expansion phase of the interaction it is again necessary to establish a time scale. If the equilibration time constants are short compared with this time scale, chemical and thermal equilibrium can be

considered to exist at each time during the expansion, the composition and temperature of the system adjusting continuously. In cascade calculations the central density falls a factor of two every  $10^{-23}$ s during expansion. The equilibration time constants at a later stage in the expansion will be larger. For example at  $\rho = 0.7 \rho_0$ , assuming an expansion somewhere between isoergic and isentropic,<sup>11</sup> the chemical equilibration time constant from Eq. 1 lies between the values  $3 \times 10^{-23}$ s  $< \tau < 2 \times 10^{-22}$ s, much longer than the characteristic time of the expansion  $10^{-23}$ s. Thus chemical equilibrium cannot be maintained, contrary to the assumption made in the fireball<sup>12</sup> and hydrodynamical model<sup>13</sup> studies of pion production, and the pion yield will freeze out near the high density stage rather than late in the expansion. This is the effect noted in the cascade calculations which formed the basis of our previous analysis of the pion yield.<sup>5</sup> The thermal equilibration time is determined by the large  $\pi N$  cross section and is much shorter, so that thermal equilibrium is maintained until later in the expansion. The pion yield is established by the  $\pi + \Delta$  abundance at chemical freeze-out, and is unaltered by the pion-number-conserving interchange  $\pi N \leftrightarrow \Delta$ . Furthermore, during expansion the rapidly diminishing number of  $\Delta$ 's (due to  $\Delta \rightarrow \pi N$  decay) make it progressively less likely for " $\pi$  absorption" to occur via  $\Delta N \rightarrow NN$ .

The measured pion multiplicity can now be used to determine the thermal energy content of the system at the end of the high density stage. To accomplish this, the idealization of complete equilibrium at the end of the high density stage and its freeze-out with the onset of expansion will be assumed. The total c.m. excitation energy per baryon in a system at temperature  $T$  and density  $\rho$  is given by<sup>8,12,14</sup>

$$E \div m_N = \frac{3}{2} T + \langle n_\pi \rangle \langle E_\pi \rangle / B + \langle n_\Delta \rangle \langle m^* \rangle / B + W(\rho, T = 0) + E_{\text{flow}} \quad (2)$$

where the left-hand side is the kinetic energy per baryon in the c.m. frame. The first three terms on the right-hand side describe the thermal excitation energy contained in baryons, pions, and in resonance mass excess  $m^*$ . The fourth term is the ground state compressional energy and the last term is the flow kinetic energy, both at density  $\rho$ . At the time of maximum density when the final pion multiplicity is established, the compressional energy is a maximum and there is no kinetic flow energy<sup>15</sup> which will appear only as the system expands. Thus, at the time of maximum density Eq. 2 can be written in a simplified form

$$\epsilon = E_T(\rho, n_\pi, n_\Delta) + E_C(\rho, T = 0) \quad (3)$$

where  $\epsilon$  is the incident c.m. kinetic energy,  $E_T$  the thermal excitation, and  $E_C$  the compressional energy each taken per participant. Here medium effects<sup>16</sup> on the thermal baryon spectrum are ignored, the medium being assumed to exert its influence through the compressional energy only.

In our previous approach<sup>5</sup> the observed pion multiplicity was used to calculate  $E_T$  by means of the cascade model. The difference between the bombarding energy  $\epsilon$  and  $E_T$  then gave the compressional energy  $E_C$ . This quantity was plotted as a function of the density  $\rho$ , also extracted from cascade model calculations, to derive the nuclear equation of state. A similar relationship will now be derived without reference to the cascade model.



In the new approach a relativistic chemical model<sup>12</sup> which assumes an equilibrium mixture of  $N$ ,  $\Delta$  and  $\pi$  was used to predict the total pion multiplicity. This model requires the density and total c.m. energy as input. The density is also necessary to derive a nuclear equation of state. It was obtained using the relativistic Rankine-Hugoniot relation<sup>6</sup>

$$\rho/\rho_0 = \gamma_{c.m.} [1 - \rho_0 E_{lab}/2P]^{-1} \quad (4)$$

where  $\rho/\rho_0$  describes the shock compression in a medium at rest in the center of mass, and  $\gamma_{c.m.}$  is the beam Lorentz factor with respect to the c.m. frame for an incident lab kinetic energy per nucleon  $E_{lab}$ . The quantity  $P$  is the pressure of the medium, related to the equation of state  $W(\rho)$  and temperature  $T$  by the relation

$$P = \rho^2 \partial W(\rho) / \partial \rho + \rho T \quad (5)$$

Equations (4) and (5) were solved within the framework of the chemical model, by the following iterative technique:

1. Assume a trial density dependence  $\rho(\epsilon)$ .
2. Using this density, calculate the total  $\pi + \Delta$  multiplicity at energy  $\epsilon$  using the chemical model.
3. By comparing the experimental and calculated values of the  $\pi + \Delta$  multiplicity as a function of  $\epsilon$ , extract the compressional energy  $E_C(\epsilon)$ .
4. With the trial density and the extracted compressional energy, the function  $W(\rho)$  is obtained.

5. Using this  $W(\rho)$  and the temperature  $T$  obtained from the chemical model in step 2, obtain the pressure  $P$  from equation (5).
6. Using this value of the pressure, obtain a new value  $\rho(\epsilon)$  from equation (4). Repeat the procedure using  $\rho(\epsilon)$  in step 1 until no further change in  $\rho$  is found.

Use of the trial value  $\rho/\rho_0 = 2\gamma_{c.m.}^2$  led to rapid convergence of the procedure. The results are shown in Fig. 1, together with the results previously obtained from cascade calculations and the experimental data. The experimental points are slightly different from those presented in Ref. 5, which were for impact parameters up to  $b = 2.4$  fm, defined by a central trigger. The present data were obtained from measurements in the minimum bias trigger mode. At each bombarding energy the ratio of the pion multiplicity to the observed proton participant number was found to be a constant as shown in Fig. 2. The yields at  $Z_1 + Z_2$ , where the proton participant number equals the total nuclear charge, were taken to correspond to  $b = 0$ . They are plotted in Fig. 1 where  $^{17} \langle n_{\pi+\Delta} \rangle = 3 \langle n_{\pi-} \rangle$ . The cascade and chemical results for the pion yields are seen to be in close agreement. The compressional energies indicated by the horizontal arrows are therefore also similar. Fig. 3 shows the compressional energies for both models, plotted against the density. The results are again rather similar. Also shown in Fig. 3 is an equation of state calculated by Boguta and Stöcker,<sup>1</sup> using a relativistic field theory approach for nuclear matter with an incompressibility constant  $K = 240$  MeV and effective mass  $m_{eff} = 0.75 m_N$ .

In conclusion, it is reasonable to use chemical and thermal equilibrium calculations for the high density stage of nucleus-nucleus collisions, when considering nucleons, pions and deltas. A distinction must be made between

chemical freeze-out which for  $\pi + \Delta$  occurs at the end of the high density phase and thermal freeze-out which occurs later. It will be of interest to investigate further the expansion stage to determine how much the temperature changes and whether there really is a well-defined thermal freeze-out. In the present analysis of the total pion yield, the density and compressional energy were obtained using the chemical model and the Rankine-Hugoniot relationship. The results are similar to those previously obtained using the cascade model and lend confidence to the equation of state derived from the experiments.

The authors wish to thank M. Gyulassy, J. Knoll, P. J. Siemens, and J. Cugnon for stimulating discussions. Particular thanks are due to G. D. Westfall for providing the thermal model results.

This work was supported by the Director, Office of Energy Research, Division of Nuclear Physics of the Office of High Energy and Nuclear Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

References

1. J. Boguta and H. Stöcker, Phys. Lett. 120B (1983) 289;  
J. D. Walecka, Proc. 5th High Energy Heavy Ion Study, LBL Report 12652 (1981) p. 91.
2. H. A. Bethe et al., Nucl. Phys. A324 (1979) 487.
3. M. Nauenberg and G. Chapline, Astrophys. Journ 179 (1973) 277.
4. S. Nagamiya, Phys. Rev. Lett. 49 (1982) 1383.
5. R. Stock et al., Phys. Rev. Lett. 49 (1982) 1236.
6. G. F. Chapline et al., Phys. Rev. D8 (1973) 4302.
7. H. Stöcker, W. Greiner, and W. Scheid, Z. Phys. A286 (1978) 121;
8. J. I. Kapusta, Phys. Rev. C16 (1977) 1493;  
A. Z. Mekjian, Nucl. Phys. A312 (1978) 491;  
S. DasGupta and A. Z. Mekjian, Phys. Reports 72 (1981) 131.
9. J. Cugnon, T. Mitzutani, and J. Vandermeulen, Nucl. Phys. A352 (1981) 505;  
J. Cugnon, D. Kinet, and J. Vandermeulen, Nucl. Phys. A379 (1982) 553.
10. J. W. Harris and R. Stock, LBL Report 17054 (1984).
11. G. Bertsch and J. Cugnon, Phys. Rev. C24 (1981) 2514.
12. J. Gosset et al., Phys. Rev. C16 (1977) 629;  
J. Gosset, J. I. Kapusta and G. D. Westfall, Phys. Rev. C18 (1978) 844.
13. H. Stöcker, LBL Report 12302 (1981).
14. M. I. Sobel et al., Nucl. Phys. A251 (1975) 502.
15. J. I. Kapusta and D. Strottman, Phys. Rev. C23 (1981) 1282;  
H. Stocker et al., Phys. Rev. Lett. 47 (1981) 1807.

16. I. N. Mishustin, F. Myhrer, and P. J. Siemens, Phys. Lett. 95B (1980) 361.
17. This is derived assuming pion production via the  $\Delta(1232)$  resonance.
18. Y. Yariv and Z. Fraenkel, Phys. Rev. C20 (1979) 2227 and C24 (1981) 488. Y. Yariv, private communication.

Figure Captions

1. The  $\pi + \Delta$  yield per participant nucleon is plotted as a function of c.m. energy (lower axis) and laboratory energy (upper axis) for Ar + KCl. The solid circles are the experimental results for zero impact parameter. The dashed and dotted curves correspond to the Cugnon cascade and the chemical model predictions, respectively, and the triangles are predictions of the Yariv-Fraenkel<sup>18</sup> cascade. The horizontal arrows represent the values of the compressional energy per nucleon  $E_c$  determined at each experimental point.
2. The mean  $\pi^-$ -multiplicity observed in Ar + KCl reactions at 1.0, 1.2, 1.4, 1.6 and 1.8 GeV/u plotted as a function of the observed number of proton participants  $Q$ . Only the interpolating lines are shown except for 1.0 and 1.8 GeV/u.
3. Values of the compressional energy, minus 10 MeV binding energy, plotted as a function of the density in units of normal nuclear density  $\rho_0$  derived from the data using the cascade and chemical model, respectively. Only statistical errors are given. The curve is a relativistic field theoretical prediction (Ref. 1) with  $K = 240$  MeV and  $m_{\text{eff}} = 0.75 m_N$ .




Fig. 1

XCG 8312-4933





Fig. 2

XBL804-747A





This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720