

Searches for heavy Higgs bosons in two-Higgs-doublet models and for $t \rightarrow ch$ decay using multilepton and diphoton final states in pp collisions at 8 TeV

V. Khachatryan *et al.**

(CMS Collaboration)

(Received 10 October 2014; published 23 December 2014)

Searches are presented for heavy scalar (H) and pseudoscalar (A) Higgs bosons posited in the two doublet model (2HDM) extensions of the standard model (SM). These searches are based on a data sample of pp collisions collected with the CMS experiment at the LHC at a center-of-mass energy of $\sqrt{s} = 8$ TeV and corresponding to an integrated luminosity of 19.5 fb^{-1} . The decays $H \rightarrow hh$ and $A \rightarrow Zh$, where h denotes an SM-like Higgs boson, lead to events with three or more isolated charged leptons or with a photon pair accompanied by one or more isolated leptons. The search results are presented in terms of the H and A production cross sections times branching fractions and are further interpreted in terms of 2HDM parameters. We place 95% C.L. cross section upper limits of approximately 7 pb on $\sigma\mathcal{B}$ for $H \rightarrow hh$ and 2 pb for $A \rightarrow Zh$. Also presented are the results of a search for the rare decay of the top quark that results in a charm quark and an SM Higgs boson, $t \rightarrow ch$, the existence of which would indicate a nonzero flavor-changing Yukawa coupling of the top quark to the Higgs boson. We place a 95% C.L. upper limit of 0.56% on $\mathcal{B}(t \rightarrow ch)$.

DOI: [10.1103/PhysRevD.90.112013](https://doi.org/10.1103/PhysRevD.90.112013)

PACS numbers: 14.80.Ec, 13.85.-t, 14.80.Bn, 14.80.Fd

I. INTRODUCTION

The standard model (SM) has an outstanding record of consistency with experimental observations. It is not a complete theory, however, and since the recent discovery of a Higgs boson [1–3], attaining a better understanding of the mechanism responsible for electroweak symmetry breaking (EWSB) has become a central goal in particle physics. The experimental directions to pursue this goal include improved characterization of the Higgs boson properties, searches for new particles such as the members of an extended Higgs sector or the partners of the known elementary particles predicted by supersymmetric models, and searches for unusual processes such as rare decays of the top quark. Since the Higgs boson plays a critical role in EWSB, searches and studies of decays with the Higgs boson in the final state have become particularly attractive.

In many extensions of the SM, the Higgs sector includes two scalar doublets [4]. The two Higgs doublet model (2HDM) [5] is a specific example of such a SM extension. In this model five physical Higgs sector particles survive EWSB: two neutral CP-even scalars (h, H), one neutral CP-odd pseudoscalar (A), and two charged scalars (H^+, H^-) [6]. For masses at or below the 1 TeV scale these particles can be produced at the LHC. Both the heavy scalar H and the pseudoscalar A can decay into electroweak

bosons, including the recently discovered Higgs boson. The branching fractions of H and A into final states containing one or more Higgs bosons h often dominate when kinematically accessible. For heavy scalars with masses below the top pair production threshold, the $H \rightarrow hh$ and $A \rightarrow Zh$ decays typically dominate over competing Yukawa decays to bottom quarks, while for heavy scalars with masses above the top pair production threshold, these decays are often comparable in rate to decays into top pairs and are potentially more distinctive.

We describe a search for two members of the extended Higgs sector, H and A , via their decays $H \rightarrow hh$ and $A \rightarrow Zh$, where h denotes the recently discovered SM-like Higgs boson [1–3]. The final states used in this search consist of three or more charged leptons or a resonant photon pair accompanied by at least one charged lepton. (In the remainder of this paper, “lepton” refers to a charged lepton, e, μ , or hadronic decay of the τ lepton, τ_h .) The $H \rightarrow hh$ and $A \rightarrow Zh$ decays can yield multileptonic final states when h decays to WW^*, ZZ^* , or $\tau\tau$. Similarly, the resonant decay $h \rightarrow \gamma\gamma$ can provide a final state that contains a photon pair and one or more leptons from the decay of the other daughter particle.

Using the same data set and technique, we also investigate the process $t \rightarrow ch$, namely the flavor-changing rare decay of the top quark to a Higgs boson accompanied by a charm quark in the $t\bar{t} \rightarrow (bW)(ch)$ decay. The $t \rightarrow ch$ process can occur at an observable rate for some parameters of the 2HDM [7]. Depending on how the h boson and t quark decay, both the multilepton and the lepton + diphoton final states can be produced. Both ATLAS [8]

* Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 3.0 License](https://creativecommons.org/licenses/by/3.0/). Further distribution of this work must maintain attribution to the author(s) and the published articles title, journal citation, and DOI.

and CMS [9] have searched for this process using complementary techniques. The CMS upper limit for the branching fraction of 1.3% at 95% confidence level (C.L.) comes from an inclusive multilepton search that uses the data set analyzed here. We describe here a $t \rightarrow ch$ search using lepton + diphoton events and combine the results of the previously reported multilepton search with the present lepton + diphoton search. This combination results in a considerable improvement in the $t \rightarrow ch$ search sensitivity.

In this paper, we first briefly describe the CMS detector, data collection, and the detector simulation scheme in Sec. II. We then describe in Sec. III the selection of events that are relevant for the search signatures followed by the event classification in Sec. IV, which calls for the data sample to be subdivided in a number of mutually exclusive channels based on the number and flavor of leptons, the number of hadronically decaying τ leptons, photons, the tagged flavors of the jets, as well as the amount of missing transverse energy (E_T^{miss}). A description of the SM background estimation in Sec. V precedes the channel-by-channel comparison of the observed number of events with the background estimation in Sec. VI. We next interpret in Sec. VII these observations in terms of the stand-alone production and decay rates for H and A . Since these rates follow from the parameters of the 2HDM, we reexpress these results in terms of the appropriate 2HDM parameters. Finally, we selectively redeploy the H and A analysis procedure to search for the rare $t \rightarrow ch$ decay.

The multilepton component of this analysis closely follows the previously mentioned CMS inclusive multilepton analysis [9]. In particular, the lepton reconstruction, SM background estimation procedures as well as the data set used are identical in the two analyses and are therefore described minimally here.

II. DETECTOR, DATA COLLECTION, AND SIMULATION

The central feature of the CMS detector is a superconducting solenoidal magnet of field strength 3.8 T. Within the field volume are a silicon pixel and strip tracker, a lead tungstate crystal calorimeter, and a brass-and-scintillator hadron calorimeter. The tracking detector covers the pseudorapidity region $|\eta| < 2.5$ and the calorimeters $|\eta| < 3.0$. Muon detectors based on gas-ionization detectors lie outside the solenoid, covering $|\eta| < 2.4$. A steel-and-quartz-fiber forward calorimeter provides additional coverage between $3 < |\eta| < 5.0$. A detailed description of the detector as well as a description of the coordinate system and relevant kinematical variables can be found in Ref. [10].

The data sample used in this search corresponds to an integrated luminosity of 19.5 fb^{-1} recorded in 2012 with the CMS detector at the LHC. Dilepton triggers (dielectron, dimuon, muon electron) and diphoton triggers are used for

data collection. The transverse momentum (p_T) threshold for dilepton triggers is 17 GeV for the leading lepton and 8 GeV for the subleading lepton. Similarly, the p_T thresholds for the diphoton trigger are 36 and 22 GeV.

The dominant SM backgrounds for this analysis such as $t\bar{t}$ quark pairs and diboson production are simulated using the MadGraph (version 5.1.3.30) [11] generator. We use the CTEQ6L1 leading-order parton distribution function (PDF) set [12]. For the diboson + jets simulation, up to two jets are selected at the matrix element level in MadGraph. The detector simulation is performed with GEANT4 [13]. The generation of signal events is performed using both the MadGraph and PYTHIA generators, with the description of detector response based on the CMS fast simulation program [14].

III. PARTICLE RECONSTRUCTION AND PRELIMINARY EVENT SELECTION

The CMS experiment uses a particle-flow (PF) based event reconstruction [15,16], which takes into account information from all subdetectors, including charged-particle tracks from the tracking system and deposited energy from the electromagnetic and hadronic calorimeters. All particles in the event are classified into mutually exclusive types: electrons, muons, τ leptons, photons, charged hadrons, and neutral hadrons.

Electron and muon candidates used in this search are reconstructed from the tracker, calorimeter, and muon system measurements. Details of reconstruction and identification can be found in Refs. [17,18] for electrons and in Refs. [19,20] for muons. The electron and muon candidates are required to have $p_T \geq 10$ GeV and $|\eta| < 2.4$. For events triggered by the dilepton trigger, the leading electron or muon must have $p_T > 20$ GeV in order to ensure maximal efficiency of the dilepton trigger. Hadronic decays of the τ lepton (τ_h) are reconstructed using the hadron-plus-strips method [21] and must have the measured jet p_T of the jet tagged as a τ_h candidate to be greater than 20 GeV and $|\eta| \leq 2.3$.

Photon candidates are reconstructed using the energy deposit clusters in the electromagnetic calorimeter [18,22]. Candidate photons are required to satisfy shower shape requirements. In order to reject electrons misidentified as photons, the photon candidate must not match any of the tracks reconstructed with the pixel detector. Photon candidates are required to have $p_T \geq 20$ GeV and $|\eta| < 2.5$. For events triggered by the diphoton trigger, the leading (subleading) photon must have $p_T > 40(25)$ GeV.

Jets are reconstructed by clustering PF particles using the anti- k_T algorithm [23] with a distance parameter of 0.5 and are required to have $|\eta| \leq 2.5$. Jets are further characterized as being “ b tagged” using the medium working point of the CMS combined secondary-vertex algorithm [24]. They typically result from the decays of the b quark. The total hadronic transverse energy, H_T , is the scalar sum of the p_T

of all jets with $p_T > 30$ GeV. The E_T^{miss} in an event is defined to be the magnitude of the vectorial p_T sum of all the PF candidates.

The primary vertex for a candidate event is defined as the reconstructed collision vertex with the highest p_T^2 sum of the associated tracks. It also must be within 24 cm from the center of the detector, along the beam axis (z direction), and within 2 cm in a direction transverse to the beam line [25]. We require the candidate leptons to originate from within 0.5 cm in z of the primary vertex and that their impact parameters d_{xy} between the track and the primary vertex in the plane transverse to the beam axis be at most 0.02 cm.

For electrons and muons, we define the relative isolation I_{rel} of the candidate leptons to be the ratio of the p_T sum of all other PF candidates that are reconstructed in a cone defined by $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 0.3$ around the candidate to the p_T of the candidate, and require $I_{\text{rel}} < 0.15$. The photon isolation requirement is similar, but varies as a function of the candidate p_T and η [26]. For the isolation of the τ_h candidates, we require that the p_T sum of all other particles in a cone of $\Delta R < 0.5$ be less than 2 GeV. The isolation variable for leptons and photons is corrected for the contributions from pileup interactions [27]. The combined efficiency for trigger, reconstruction, and identification are approximately 75% for electrons and 80% for muons. The identification and isolation efficiency for prompt leptons is measured in data using a “tag-and-probe” method based on an inclusive sample of Z + jets events [28]. The ratio of the efficiency in data and simulation parametrized by the different p_T and η values of the probed lepton is used to correct the selection efficiency in the simulated samples.

A leptonically decaying Z boson can lead to a trilepton event when the final-state radiation undergoes (internal or external) conversion and one of the leptons escapes detection. Therefore, we reject trilepton events with low missing transverse energy ($E_T^{\text{miss}} < 30$ GeV) when their three body invariant mass is consistent with the Z mass (i.e. $m_{\ell^+\ell^-\ell^\pm}$ or $m_{\ell^+\ell^-\ell^\pm}$ is between 75 and 105 GeV), even if $m_{\ell^+\ell^-}$ is not ($\ell = e, \mu$). Finally, SM background from abundant low-mass Drell-Yan production and low-mass resonances like J/ψ and Υ is suppressed by rejecting an event if it contains a dilepton pair with $m_{\ell^+\ell^-}$ below 12 GeV.

IV. EVENT CLASSIFICATION

We perform searches using a multichannel counting experiment approach. A multilepton event consists of at least three isolated and prompt leptons (e, μ, τ_h), of which at least two must be electrons or muons (“light” leptons). A photon pair together with at least one lepton makes a lepton + diphoton event. The relatively low rates for multilepton and lepton + diphoton final states in SM allow this search to target rare signals.

A. The $H \rightarrow hh$, $A \rightarrow Zh$, and $t \rightarrow ch$ signals

In the $H \rightarrow hh$ search, seven combinations of the hh decays (WW^*WW^* , WW^*ZZ^* , $WW^*\tau\tau$, ZZ^*ZZ^* , $ZZ^*\tau\tau$, ZZ^*bb , and $\tau\tau\tau\tau$) can result in a final state containing multileptons and three combinations ($\gamma\gamma WW^*$, $\gamma\gamma ZZ^*$, and $\gamma\gamma\tau\tau$) can result in lepton + diphoton final states with appreciable rates.

In the $A \rightarrow Zh$ search, the multilepton and diphoton signal events can result from the WW , ZZ , $\tau\tau$, and $\gamma\gamma$ decays of the h , when accompanied by the appropriate decays of the W and Z bosons and the τ lepton. Five combinations of the Zh decays ($Z \rightarrow \ell\ell$, $h \rightarrow WW^*$; $Z \rightarrow \ell\ell$, $h \rightarrow ZZ^*$; $Z \rightarrow \ell\ell$, $h \rightarrow \tau\tau$; $Z \rightarrow \nu\nu$, $h \rightarrow ZZ^*$; $Z \rightarrow qq$, $h \rightarrow ZZ^*$) can result in a final state containing multileptons and one combination ($Z \rightarrow \ell\ell$, $h \rightarrow \gamma\gamma$) leads to lepton + diphoton states with substantial rates.

For the $t \rightarrow ch$ search, three combinations in the decay chain $t\bar{t} \rightarrow (bW)(ch) \rightarrow (b\ell\nu)(ch)$ can lead to multilepton final states, namely $h \rightarrow WW^*$, $h \rightarrow ZZ^*$, and $h \rightarrow \tau\tau$. The $bWch$ channel can also result in a lepton + diphoton final state when the Higgs boson decays to a photon pair. Finally, given the parent $t\bar{t}$ state, the amount of hadronic activity in the $t \rightarrow ch$ signal events is expected to be quite large.

B. Multilepton search channels

A three-lepton event must contain exactly three isolated and prompt leptons (e, μ, τ_h), of which two must be electrons or muons. Similarly, a four-lepton event must contain at least four leptons, of which three must be electrons or muons. With the goal of segregating SM backgrounds, these events are classified on the basis of the lepton flavor, their relative charges, as well as charge and flavor combinations and other kinematic quantities such as dilepton invariant mass and E_T^{miss} , as follows.

Events with τ_h are grouped separately because narrow jets are frequently misidentified as τ_h , leading to larger SM backgrounds for channels with τ_h . Similarly, the presence of a b -tagged jet in an event calls for a separate grouping in order to isolate the $t\bar{t}$ background.

The next classification criterion is the maximum number of opposite-sign and same-flavor (OSSF) dilepton pairs that can be constructed in an event using each light lepton only once. For example, both $\mu^+\mu^-\mu^-$ and $\mu^+\mu^-e^-$ are said to be OSSF1, and a $\mu^+e^-\tau_h$ would be OSSF0. Both $e^+e^+\mu^-$ and $\mu^+\mu^+\tau_h$ are OSSF0(SS), where SS additionally indicates the presence of same-signed electron or muon pairs. Similarly, $\mu^+\mu^-e^+e^-$ is OSSF2. An event with an OSSF pair is said to be “on Z ” if the invariant mass of at least one of the OSSF pair is between 75 and 105 GeV, otherwise it is “off Z .” An OSSF1 off- Z event is “below Z ” or “above Z ” depending on whether the mass of the pair is less than 75 or more than 105 GeV, respectively. An on- Z OSSF2 event may be a “one on- Z ” or a “two on- Z ” event.

Finally, the three-lepton events are classified in five E_T^{miss} bins: < 50 , 50–100, 100–150, 150–200, and > 200 GeV

and the four-lepton events are classified in four E_T^{miss} bins: < 30 , $30\text{--}50$, $50\text{--}100$, and > 100 GeV. This results in a total of 70 three-lepton channels and 72 four-lepton channels which are listed explicitly when we later present the tables of event yields and background predictions (Tables I and II).

C. Lepton+diphoton search channels

A diphoton pair together with at least one lepton makes a lepton + diphoton event. The diphoton invariant mass of the $h \rightarrow \gamma\gamma$ candidates must be between 120 and 130 GeV. The search channels are $\gamma\gamma\ell\ell$, $\gamma\gamma\ell\tau_h$, $\gamma\gamma\ell$, and $\gamma\gamma\tau_h$. Depending on the relative dilepton flavor and invariant mass, the $\gamma\gamma\ell\ell$ events can be OSSF0, OSSF1 on Z, or OSSF1 off Z. The SM background decreases with increasing E_T^{miss} , therefore the events are further classified, when appropriate, in three bins: $E_T^{\text{miss}} < 30$, $30\text{--}50$, and > 50 GeV.

The $t \rightarrow ch$ signal populates the $\gamma\gamma\ell$ and $\gamma\gamma\tau_h$ channels but not the dilepton + diphoton channels. Since the $t \rightarrow ch$ signal events always contain a b quark from the conventional bW decay of one of the top quarks, the $\gamma\gamma\ell$ and $\gamma\gamma\tau_h$ search channels are further classified based on the presence of a b -tagged jet. For these channels, we also split the last E_T^{miss} bin into two: $50\text{--}100$ GeV and > 100 GeV.

The overall lepton + diphoton channel count in this search is seven for $\gamma\gamma\ell\ell$, three for $\gamma\gamma\ell\tau_h$, and eight each for $\gamma\gamma\ell$ and $\gamma\gamma\tau_h$. They are listed explicitly when we present the tables of event yields and background predictions later (Tables VI and VII).

V. BACKGROUND ESTIMATION

A. Multilepton background estimation

Significant sources of multilepton SM background are Z + jets, diboson production (VV + jets; $V = W, Z$), $t\bar{t}$ production, and rare processes such as $t\bar{t}V$ + jets. The techniques we use here to estimate these backgrounds are identical to those used in Ref. [9] and are described briefly below.

WZ and ZZ diboson production can yield events with three or four intrinsically prompt and isolated leptons that can be accompanied by significant E_T^{miss} and H_T . To estimate these background contributions, we use a simulation validated after kinematic comparisons with appropriately enriched data samples.

Processes such as Z + jets and W^+W^- + jets can yield events with two prompt leptons. These can be accompanied by jets that may also contain leptons from the semileptonic decays of hadrons, or other objects that are misreconstructed as prompt leptons, leading to a three-lepton SM background. Since the simulation of the rare fluctuations that lead to such a misidentified prompt lepton can be unreliable, we use the data with two reconstructed leptons to estimate this SM background using the number of isolated prompt tracks in the dilepton data set.

The $t\bar{t}$ decay can result in two prompt leptons and is a source of background when the decay of one of the daughter b quarks reconstructs as the third prompt lepton candidate. This background is estimated from a $t\bar{t}$

TABLE I. Observed (Obs.) yields and SM expectations (Exp.) for three-lepton events. See text for the description of event classification by the number and invariant mass of opposite-sign, same-flavor lepton pairs that are on or below Z (see Sec. 4.2), presence of τ_h , tagged b jets, and the E_T^{miss} in the event. The 70 channels are exclusive.

3 leptons	$m_{\ell\ell}$	E_T^{miss} (GeV)	$N_{\tau_h} = 0, N_b = 0$		$N_{\tau_h} = 1, N_b = 0$		$N_{\tau_h} = 0, N_b \geq 1$		$N_{\tau_h} = 1, N_b \geq 1$	
			Obs.	Exp.	Obs.	Exp.	Obs.	Exp.	Obs.	Exp.
OSSF0(SS)	...	(200, ∞)	1	1.3 ± 0.6	2	1.4 ± 0.5	0	0.70 ± 0.36	0	0.7 ± 0.5
OSSF0(SS)	...	(150, 200)	2	2.1 ± 0.9	0	3.0 ± 1.1	1	2.1 ± 1.0	0	1.5 ± 0.6
OSSF0(SS)	...	(100, 150)	9	10.0 ± 4.9	4	9.9 ± 3.0	12	12.0 ± 5.9	4	6.3 ± 2.8
OSSF0(SS)	...	(50, 100)	34	37 ± 15	54	66 ± 14	32	32 ± 15	24	22 ± 10
OSSF0(SS)	...	(0, 50)	47	46 ± 11	196	221 ± 51	28	24 ± 11	21	31.0 ± 9.6
OSSF0	...	(200, ∞)	5	4.8 ± 2.4	6	5.9 ± 3.1
OSSF0	...	(150, 200)	12	18.0 ± 9.1	21	20 ± 10
OSSF0	...	(100, 150)	94	96 ± 47	91	121 ± 61
OSSF0	...	(50, 100)	351	329 ± 173	300	322 ± 163
OSSF0	...	(0, 50)	682	767 ± 207	230	232 ± 118
OSSF1	Below Z	(200, ∞)	2	2.5 ± 0.9	4	2.1 ± 1.0	1	1.9 ± 0.7	2	2.4 ± 1.2
OSSF1	On Z	(200, ∞)	17	19.0 ± 6.3	4	5.6 ± 1.9	1	2.4 ± 0.8	3	2.1 ± 0.9
OSSF1	Below Z	(150, 200)	7	4.4 ± 1.7	11	9.3 ± 4.6	3	4.7 ± 2.1	7	11.0 ± 5.9
OSSF1	On Z	(150, 200)	38	32.0 ± 8.5	10	11.0 ± 3.6	4	5.4 ± 1.7	2	5.7 ± 2.7
OSSF1	Below Z	(100, 150)	21	26.0 ± 9.9	45	56 ± 27	20	23 ± 11	56	66 ± 33
OSSF1	On Z	(100, 150)	134	129 ± 29	43	51 ± 16	20	18 ± 6	24	28 ± 14
OSSF1	Below Z	(50, 100)	157	129 ± 30	383	380 ± 104	58	60 ± 28	166	173 ± 87
OSSF1	On Z	(50, 100)	862	732 ± 141	1360	1230 ± 323	80	62 ± 17	117	101 ± 48
OSSF1	Below Z	(0, 50)	543	559 ± 93	10200	9170 ± 2710	40	52 ± 14	257	256 ± 79
OSSF1	On Z	(0, 50)	4040	4060 ± 691	51400	51400 ± 15300	181	181 ± 28	1000	1010 ± 286

TABLE II. Observed (Obs.) yields and SM expectation (Exp.) for four-lepton events. See text for the description of event classification by the number and invariant mass of opposite-sign, same-flavor lepton pairs that are on or off Z , presence of τ_h , tagged b jets, and the total E_T^{miss} in the event. The 72 channels are exclusive.

≥ 4 leptons	$m_{\ell^+\ell^-}$	E_T^{miss} (GeV)	$N_{\tau_h} = 0, N_b = 0$		$N_{\tau_h} = 1, N_b = 0$		$N_{\tau_h} = 0, N_b \geq 1$		$N_{\tau_h} = 1, N_b \geq 1$	
			Obs.	Exp.	Obs.	Exp.	Obs.	Exp.	Obs.	Exp.
OSSF0	...	(100, ∞)	0	0.07 ± 0.07	0	0.18 ± 0.09	0	0.05 ± 0.05	0	0.16 ± 0.10
OSSF0	...	(50, 100)	0	0.07 ± 0.06	2	0.80 ± 0.35	0	$0.00^{+0.03}_{-0.00}$	0	0.43 ± 0.22
OSSF0	...	(30, 50)	0	$0.001^{+0.020}_{-0.001}$	0	0.47 ± 0.24	0	$0.00^{+0.02}_{-0.00}$	0	0.11 ± 0.09
OSSF0	...	(0, 30)	0	$0.007^{+0.020}_{-0.007}$	1	0.40 ± 0.16	0	$0.001^{+0.020}_{-0.001}$	0	$0.02^{+0.04}_{-0.02}$
OSSF1	Off Z	(100, ∞)	0	0.07 ± 0.04	4	1.00 ± 0.33	0	0.14 ± 0.09	0	0.46 ± 0.20
OSSF1	On Z	(100, ∞)	2	0.6 ± 0.2	2	3.4 ± 0.8	1	0.80 ± 0.41	0	0.60 ± 0.26
OSSF1	Off Z	(50, 100)	0	0.21 ± 0.09	5	2.6 ± 0.6	0	0.21 ± 0.11	1	0.70 ± 0.32
OSSF1	On Z	(50, 100)	2	1.30 ± 0.39	10	12.0 ± 2.5	2	0.60 ± 0.33	1	0.8 ± 0.3
OSSF1	Off Z	(30, 50)	1	0.16 ± 0.07	4	2.4 ± 0.5	0	0.06 ± 0.06	0	0.47 ± 0.21
OSSF1	On Z	(30, 50)	3	1.20 ± 0.35	11	14.0 ± 3.1	0	0.22 ± 0.12	0	0.80 ± 0.31
OSSF1	Off Z	(0, 30)	1	0.38 ± 0.18	11	5.7 ± 1.7	0	0.05 ± 0.04	0	0.50 ± 0.26
OSSF1	On Z	(0, 30)	1	2.0 ± 0.5	32	30.0 ± 9.2	1	0.19 ± 0.11	3	1.30 ± 0.42
OSSF2	Two on Z	(100, ∞)	0	0.02 ± 0.15	0	0.21 ± 0.13
OSSF2	One on Z	(100, ∞)	1	0.43 ± 0.15	0	0.50 ± 0.29
OSSF2	Off Z	(100, ∞)	0	0.06 ± 0.03	0	0.09 ± 0.07
OSSF2	Two on Z	(50, 100)	3	2.8 ± 2.1	0	0.33 ± 0.11
OSSF2	One on Z	(50, 100)	1	2.0 ± 0.7	1	0.50 ± 0.28
OSSF2	Off Z	(50, 100)	2	0.20 ± 0.14	0	0.12 ± 0.10
OSSF2	Two on Z	(30, 50)	19	22 ± 9	2	0.70 ± 0.24
OSSF2	One on Z	(30, 50)	6	6.5 ± 2.4	0	0.32 ± 0.12
OSSF2	Off Z	(30, 50)	3	1.4 ± 0.6	1	0.15 ± 0.08
OSSF2	Two on Z	(0, 30)	118	109 ± 28	3	2.0 ± 0.5
OSSF2	One on Z	(0, 30)	24	29.0 ± 7.6	1	0.60 ± 0.17
OSSF2	Off Z	(0, 30)	5	7.8 ± 2.3	0	0.18 ± 0.06

Monte Carlo sample and using the probability of occurrence of a misidentified third lepton derived from data.

For search channels that contain τ_h , we estimate the probability of a (sparse) jet misidentified as a τ_h candidate by extrapolating the isolation distribution of the τ_h candidates. Since the shape of this distribution is sensitive to the extent of jet activity, the extrapolation is carried out as a function of the hadronic activity in the sample as determined by the summed p_T of all tracks as well as the leading jet p_T in the event.

Finally, minor backgrounds from rare processes such as $t\bar{t}V$ + jets or SM Higgs production including its associated production with W , Z , or $t\bar{t}$ are estimated using simulation.

B. Lepton+diphoton background estimation

We use a 120–130 GeV diphoton invariant mass window to capture the $h \rightarrow \gamma\gamma$ signal. With the requirement of at least one lepton in these lepton + diphoton channels, the SM background tends to be small and is estimated by interpolating the diphoton mass sidebands of the signal window. We assume the background distribution shape to be a falling exponential as a function of the diphoton invariant mass over the 100–200 GeV mass range.

Figure 1 (top panel) shows the exponential fit to the 100–120 and 130–200 GeV sidebands in the mass

distribution for $\gamma\gamma\tau_h$ events with $E_T^{\text{miss}} < 30$ GeV. We choose this sample to determine the exponent because it is a high-statistics sample. This exponent is used for background determination in all diphoton channels, allowing only the normalization to float from channel to channel. Figure 1 (bottom panel) shows an example of such a fit for the $\gamma\gamma\ell$ sample with a 30–50 GeV E_T^{miss} requirement along with an exponential fit where both the exponent and the normalization are allowed to float. We assign a 50% systematic uncertainty for background determination in the 120–130 GeV Higgs boson mass region. The figure also shows the expected signal multiplied by a factor of three for clarity for $m_H = 300$ GeV, assuming that the production cross section σ for $m_H = 300$ GeV is equal to the standard model Higgs gluon fusion value of 3.59 pb at this mass given by the LHC Higgs Cross Section Working Group in Ref. [29], and a branching fraction $\mathcal{B}(H \rightarrow hh) = 1$.

VI. OBSERVATIONS

Tables I and II list the observed number of events for the three-lepton and four-lepton search channels, respectively. The number of expected events from SM processes is also shown together with the combined statistical and systematic uncertainties. Table III lists the sources of systematic effects and the resultant uncertainties in estimating the




FIG. 1 (color online). (Top panel) Diphoton invariant mass distribution for $\gamma\gamma\tau_h$ events with $E_T^{\text{miss}} < 30$ GeV with an exponential fit derived from the 100–120 and 130–200 GeV sidebands regions. (Bottom panel) The same distribution for the $\gamma\gamma\ell$ events with E_T^{miss} in 30–50 GeV range with an exponential fit (blue curve) where the exponent is fixed to the value obtained from the fit shown in the top figure. Also shown for comparison purposes is an actual fit (magenta curve) to the shown data distribution. An example signal distribution (in red), assuming $\sigma\mathcal{B}(pp \rightarrow H \rightarrow hh)$ to be equal to three times 3.59 pb, as described in the text, shows that the signal is well contained in the 120–130 GeV window.

expected events from the SM. All search channels share systematic uncertainties for luminosity, renormalization scale, PDF, and trigger efficiency.

The observations listed in the tables generally agree with the expectations within the uncertainties. Given the large number of channels being investigated simultaneously, certain deviations between observations and expected values are to be anticipated. We discuss one such deviation later in the context of the H search.

Figure 2 shows observations and background decomposition for some of the most sensitive channels for the $H \rightarrow hh$ search. The amount of signal for $m_H = 300$ GeV, as described above in the context of Fig. 1, is also shown. This information is also listed in Table IV. Figure 3 and Table V show the same for the $A \rightarrow Zh$

TABLE III. A compilation of significant sources of systematic uncertainties in the event yield estimation. Note that a given uncertainty may pertain only to specific sources of background. The listed values are representative and the impact of an uncertainty varies from search channel to channel.

Source of uncertainty	Magnitude (%)
Luminosity	2.6
PDF	10
$E_T^{\text{miss}}(> 50 \text{ GeV})$ resolution correction	4
Jet energy scale	0.5
b -tag scale factor ($\bar{t}\bar{t}$)	6
$e(\mu)$ ID/isolation (at $p_T = 30$ GeV)	0.6 (0.2)
Trigger efficiency	5
$\bar{t}\bar{t}$ misidentification	50
$\bar{t}\bar{t}, WZ, ZZ$ cross sections	10, 15, 15
τ_h misidentification	30
Diphoton background	50

search for $m_A = 300$ GeV, assuming the same cross section and $\mathcal{B}(A \rightarrow Zh) = 1$.

The lepton + diphoton results are summarized in Tables VI and VII. The observations agree with the expectations within the uncertainties.




FIG. 2 (color online). The E_T^{miss} distributions for four-lepton events with an off-Z OSSF1 dilepton pair, no b -tagged jet, no τ_h (top panel), and one τ_h (bottom panel). These nonresonant (off-Z) channels are sensitive to the $H \rightarrow hh$ signal which is shown stacked on top of the background distributions, assuming $\sigma\mathcal{B}(pp \rightarrow H \rightarrow hh) = 3.59$ pb, as described in the text.

TABLE IV. Observed (Obs.) yields and SM expectation (Exp.) for selected four-lepton channels in the $H \rightarrow hh$ search. These are also shown in Fig. 2. See text for the description of event classification. The $H \rightarrow hh$ signal (Sig.) is also listed, assuming $\sigma\mathcal{B}(pp \rightarrow H \rightarrow hh) = 3.59$ pb.

Channel	E_T^{miss} (GeV)	Obs.	Exp.	Sig.
4ℓ (OSSF1, off Z) (no τ_h , no b jets)	(0, 30)	1	0.38 ± 0.18	0.30
	(30, 50)	1	0.16 ± 0.07	0.43
	(50, 100)	0	0.21 ± 0.09	0.39
	(100, ∞)	0	0.07 ± 0.04	0.14
4ℓ (OSSF1, off Z) (1- τ_h , no b jets)	(0, 30)	11	5.7 ± 1.7	0.91
	(30, 50)	4	2.4 ± 0.5	0.98
	(50, 100)	5	2.6 ± 0.6	1.31
	(100, ∞)	4	1.00 ± 0.33	0.25

VII. INTERPRETATION OF RESULTS

A. Statistical procedure

No significant disagreement is found between our observations and the corresponding SM expectations. We derive limits on the production cross section times branching fraction for the new physics scenarios under consideration, and use them to constrain parameters of the




FIG. 3 (color online). The E_T^{miss} distributions for four-lepton events without b -tagged jets which contain an on-Z OSSF1 dilepton pair and one τ_h (top panel), and an OSSF2 dilepton pair with one Z candidate and no τ_h (bottom panel). These resonant (containing a Z) channels are sensitive to the $A \rightarrow Zh$ signal which is shown stacked on top of the background distributions, assuming $\sigma\mathcal{B}(pp \rightarrow A \rightarrow Zh) = 3.59$ pb, as described in the text.

TABLE V. Observed (Obs.) yields and SM expectation (Exp.) for selected four-lepton channels in the $A \rightarrow Zh$ search. These are also shown in Fig. 3. See text for the description of event classification. The $A \rightarrow Zh$ signal (Sig.) is also listed, assuming $\sigma\mathcal{B}(pp \rightarrow A \rightarrow Zh) = 3.59$ pb.

Channel	E_T^{miss} (GeV)	Obs.	Exp.	Sig.
4ℓ (OSSF1, on Z) (1 τ_h , no b jets)	(0, 30)	32	30.0 ± 9.2	6.46
	(30, 50)	11	14.0 ± 3.1	6.72
	(50, 100)	10	12.0 ± 2.5	7.05
	(100, ∞)	2	3.4 ± 0.8	1.12
4ℓ (OSSF2, one on Z) (no τ_h , no b jets)	(0, 30)	24	29.0 ± 7.6	3.15
	(30, 50)	6	6.5 ± 2.4	2.91
	(50, 100)	1	2.0 ± 0.7	4.92
	(100, ∞)	1	0.43 ± 0.15	0.82

TABLE VI. Observed yields and SM expectations for dilepton + diphoton events. The diphoton invariant mass is required to be in the 120–130 GeV window. The ten channels are exclusive.

Channel	E_T^{miss} (GeV)	Obs.	Exp.
$\gamma\gamma\ell\ell$ (OSSF1, off Z)	(50, ∞)	0	$0.19^{+0.25}_{-0.19}$
	(30, 50)	1	$0.17^{+0.25}_{-0.17}$
	(0, 30)	1	1.20 ± 0.74
$\gamma\gamma\ell\ell$ (OSSF1, on Z)	(50, ∞)	0	$0.10^{+0.17}_{-0.10}$
	(30, 50)	1	0.33 ± 0.28
	(0, 30)	0	1.01 ± 0.55
$\gamma\gamma\ell\ell$ (OSSF0)	All	0	$0.00^{+0.17}_{-0.00}$
$\gamma\gamma\ell\tau_h$	(50, ∞)	0	$0.16^{+0.66}_{-0.16}$
	(30, 50)	0	$0.50^{+0.57}_{-0.50}$
	(0, 30)	0	0.76 ± 0.60

TABLE VII. Observed yields and SM expectations for single lepton + diphoton events. The diphoton invariant mass is required to be in the 120–130 GeV window. The eight channels are exclusive.

Channel	E_T^{miss} (GeV)	Obs.	$N_b = 0$		$N_b \geq 1$	
			Obs.	Exp.	Obs.	Exp.
$\gamma\gamma\ell$	(100, ∞)	1	2.2 ± 1.0	0	0.5 ± 0.4	
	(50, 100)	7	9.5 ± 4.4	1	2.3 ± 1.2	
	(30, 50)	29	21 ± 10	2	1.1 ± 0.6	
$\gamma\gamma\tau_h$	(0, 30)	72	77 ± 38	2	2.1 ± 1.1	
	(100, ∞)	1	$0.24^{+0.25}_{-0.24}$	0	0.35 ± 0.28	
	(50, 100)	14	9.3 ± 4.7	1	1.5 ± 0.8	
	(30, 50)	71	67 ± 34	2	2.1 ± 1.2	
	(0, 30)	229	235 ± 117	6	6.4 ± 3.3	

models. We set 95% C.L. upper limits on the cross sections using the modified frequentist construction C.L. [30,31]. We compute the single-channel C.L. limits for each channel and then obtain the combined upper limit.

B. $H \rightarrow hh$ and $A \rightarrow Zh$ model-independent interpretations

Figure 4 (top panel) shows 95% C.L. observed and expected $\sigma\mathcal{B}$ upper limits for the gluon fusion production of heavy scalar H , with the decay $H \rightarrow hh$ along with one and two standard deviation bands around the expected limits using only the multilepton channels. Figure 4 (bottom panel) shows the same using both multilepton and diphoton channels. In placing these model-independent limits, we explicitly assume that h is the recently discovered SM-like Higgs boson [1–3] particularly in regards to the branching fraction of its various decay modes as predicted in the SM.

For low masses, there is an almost two standard deviation discrepancy between the expected and observed 95% C.L. limits in Fig. 4. Its origin traces back to three four-lepton channels listed in Table II, which can also be located in Fig. 2 (bottom panel). They consist of events with a τ_h and three light leptons containing an off- Z OSSF dilepton pair, but not a b -tagged jet. The $H \rightarrow hh$ signal

resides almost entirely in the 0–100 GeV range in E_T^{miss} which is spanned by these three channels collectively. The observed (expected) number of events is 11 (5.7 ± 1.7), 4 (2.4 ± 0.5), and 5 (2.6 ± 0.6) for E_T^{miss} in ranges 0–30, 30–50, and 50–100 GeV, respectively. Summing over the three channels, the observed count is 20 with an expectation of 10.7 ± 1.9 , giving the probability of observing 20 events over the 0–100 GeV E_T^{miss} range to be approximately 2.2%. Systematic uncertainties and their correlations are taken into account when evaluating this probability. The observed discrepancy in the limits shown in Fig. 4 is thus consistent with a broad fluctuation in the observed E_T^{miss} distribution. Given the large number of channels under scrutiny in this search, fluctuations at this level are to be expected. No such deviation is observed in the E_T^{miss} distribution for other search channels.

Next we probe the sensitivity to gluon fusion production of the heavy pseudoscalar A with the decay $A \rightarrow Zh$. Figure 5 (top panel) shows 95% C.L. upper limits on $\sigma\mathcal{B}$ for $A \rightarrow Zh$ search along with one and two standard




FIG. 4 (color online). (Top panel) Observed and expected 95% C.L. $\sigma\mathcal{B}$ limits for gluon fusion production of H and the decay $H \rightarrow hh$ with one and two standard deviation bands shown. These limits are based only on multilepton channels. The h branching fractions are assumed to have SM values. (Bottom panel) The same, but also including lepton + diphoton channels.




FIG. 5 (color online). (Top panel) Observed and expected 95% C.L. $\sigma\mathcal{B}$ limits for gluon fusion production of A and the decay $A \rightarrow Zh$ with one and two standard deviation bands shown. These limits are based only on multilepton channels. The h branching fraction are assumed to have SM values. (Bottom panel) The same, but also including lepton + diphoton channels.

deviation bands around the expected contour using only the multilepton channels. Figure 5 (bottom panel) shows the same signal probed with both multilepton and diphoton channels. The observed and expected exclusions are consistent.

C. Interpretations in the context of two-Higgs-doublet models

General models with two Higgs doublets may exhibit new tree-level contributions to flavor-changing neutral currents that are strongly constrained by low-energy experiments. Prohibitive flavor violation is avoided in a 2HDM if all fermions of a given representation receive their masses through renormalizable Yukawa couplings to a single Higgs doublet, as in the case of supersymmetry. There are four such possible distinct assignments of fermion couplings in models with two Higgs doublets, the most commonly considered of which are called type I and type II models. In type I models all fermions receive their masses through Yukawa couplings to a single Higgs doublet, while in type II models the up-type quarks receive their masses

through couplings to one doublet and down-type quarks and leptons couple to the second doublet. In either type, after electroweak symmetry breaking the physical Higgs scalars are linear combinations of these two electroweak Higgs doublets, so that fermion couplings to the physical states depend on the type of 2HDM, the mixing angle α , and the ratio of vacuum expectation values $\tan\beta$. We next present search interpretations in the context of type I and type II 2HDMs [5]. In these models, the production cross sections for H and A as well as the branching fractions for them to decay to two SM-like Higgs bosons depend on parameters α and $\tan\beta$. The mixing angle between H and h is given by α , whereas $\tan\beta$ gives the relative contribution of each Higgs doublet to electroweak symmetry breaking. In obtaining these model-dependent limits, the daughter h is assumed to be the recently discovered SM-like Higgs boson, but the branching fractions to its various decay modes are assumed to be dictated by the parameters α and $\tan\beta$ of the 2HDM, as described below.

We use the SusHi [32] program to obtain the 2HDM cross sections. The branching fraction for SM-like Higgs




FIG. 6 (color online). (Left panels) Observed and expected 95% C.L. upper limits for gluon fusion production of a heavy Higgs boson H of mass 300 GeV as a function of parameters $\tan\beta$ and $\cos(\beta-\alpha)$ of the types I (upper panel) and II (lower panel) 2HDM. The parameters determine the H production cross section as well as the branching fractions $\mathcal{B}(H \rightarrow hh)$ and $\mathcal{B}(h \rightarrow WW^*, ZZ^*, \tau\tau, \gamma\gamma)$, which are relevant to this search. (Right panel) The $\sigma\mathcal{B}(H \rightarrow hh)$ contours for types I (upper panel) and II (lower panel) 2HDM adopted from Ref. [35]. The excluded regions are either below the open limit contours or within the closed ones.




FIG. 7 (color online). (Left panels) Observed and expected 95% C.L. upper limits for gluon fusion production of an A boson of mass 300 GeV as a function of parameters $\tan \beta$ and $\cos(\beta - \alpha)$ of the types I (upper panel) and II (lower panel) 2HDM. The parameters determine the A production cross section as well as the branching fractions $\mathcal{B}(A \rightarrow Zh)$ and $\mathcal{B}(h \rightarrow WW^*, ZZ^*, \tau\tau, \gamma\gamma)$ which are relevant to this search. (Right panels) The $\sigma\mathcal{B}(A \rightarrow Zh)$ contours for types I (upper panel) and II (lower panel) 2HDM adopted from Ref. [35]. The excluded regions are below the open limit contours.

boson are calculated using the 2HDMC [33] program. The 2HDMC results are consistent with those provided by the LHC Higgs Cross Section Working Group [34]. A detailed list of couplings of H and A to SM fermions and massive gauge bosons in types I and II 2HDMs has been tabulated in Ref. [6]. Figure 6 (top left and bottom left panels) shows observed and expected 95% C.L. upper limits for gluon fusion production of a heavy Higgs boson H of mass 300 GeV for type I and type II 2HDMs, respectively, along with the $\sigma\mathcal{B}$ theoretical predictions (right panels) adopted from Ref. [35] for the two models. Figure 7 (top left and bottom left panels) shows similar results for the pseudo-scalar A Higgs boson of mass 300 GeV. The branching fractions for the SM-like Higgs boson daughters of the H and A vary across the $\tan \beta$ versus $\cos(\beta - \alpha)$ plane and are incorporated in the upper limit calculations.

Finally, we further improve constraints on the 2HDM parameters using the simultaneous null findings for the H and A . Figure 8 shows exclusion in $\tan \beta$ versus $\cos(\beta - \alpha)$ plane for the combined gluon fusion signal for type I (top panel) and type II (bottom panel) 2HDMs, assuming H and

A to be mass degenerate with a mass of 300 GeV. Once again, the branching fractions of the SM-like h daughters are allowed to vary across the plane.

D. $t \rightarrow ch$ search results

The $t \rightarrow ch$ signal predominantly populates lepton + diphoton channels with a b tag and $\ell\ell\ell$ (no τ_h) multilepton channels that lack an OSSF dilepton pair or have an off- Z OSSF pair. Beyond the fact that the presence of charm quark increases the likelihood of an event being classified as being b tagged, no special effort is made to identify the charm quark present in the signal. The observations and SM expectations for the ten most sensitive channels are listed in Table VIII along with the signal yield for a nominal value of $\mathcal{B}(t \rightarrow ch) = 1\%$. No significant excess is observed.

The statistical procedure yields an observed limit of $\mathcal{B}(t \rightarrow ch) = 0.56\%$ and an expected limit of $\mathcal{B}(t \rightarrow ch) = (0.65^{+0.29}_{-0.19})\%$ from SM $t\bar{t}$ production followed by either $t \rightarrow ch$ or its charge-conjugate decay. The $t \rightarrow ch$ branching




FIG. 8 (color online). Combined observed and expected 95% upper limits for gluon fusion production of a heavy Higgs boson H and A of mass 300 GeV for type I (top panel) and type II (bottom panel) 2HDM as a function of parameters $\tan\beta$ and $\cos(\beta-\alpha)$. The parameters determine the H and A production cross sections as well as the branching fractions $\mathcal{B}(H \rightarrow hh)$, $\mathcal{B}(A \rightarrow Zh)$, and $\mathcal{B}(h \rightarrow WW^*, ZZ^*, \tau\tau, \gamma\gamma)$, which are relevant to this search.

fraction is related to the left- and right-handed top-flavor-changing Yukawa couplings λ_{tc}^h and λ_{ct}^h , respectively, by $\mathcal{B}(t \rightarrow ch) \simeq 0.29(|\lambda_{tc}^h|^2 + |\lambda_{ct}^h|^2)$ [7], so that the observed limit corresponds to a limit on the couplings of $\sqrt{|\lambda_{tc}^h|^2 + |\lambda_{ct}^h|^2} < 0.14$.

To facilitate interpretations in broader contexts [36], we also provide limits on $\mathcal{B}(t \rightarrow ch)$ from individual Higgs

TABLE IX. Comparison of the observed and expected 95% C.L. limits on $\mathcal{B}(t \rightarrow ch)$ from individual Higgs boson decay modes along with the 68% C.L. uncertainty ranges.

Higgs boson decay mode		Upper limits on $\mathcal{B}(t \rightarrow ch)$		
		Obs.	Exp.	68% C.L. range
$\mathcal{B}(h \rightarrow WW^*)$	= 23.1%	1.58%	1.57%	(1.02–2.22)%
$\mathcal{B}(h \rightarrow \tau\tau)$	= 6.15%	7.01%	4.99%	(3.53–7.74)%
$\mathcal{B}(h \rightarrow ZZ^*)$	= 2.89%	5.31%	4.11%	(2.85–6.45)%
Combined multileptons (WW^* , $\tau\tau$, ZZ^*)		1.28%	1.17%	(0.85–1.73)%
$\mathcal{B}(h \rightarrow \gamma\gamma)$	= 0.23%	0.69%	0.81%	(0.60–1.17)%
Combined multileptons + diphotons		0.56%	0.65%	(0.46–0.94)%

TABLE VIII. The ten most sensitive search channels for $t \rightarrow ch$, along with the number of observed (Obs.), expected SM background (Exp.), and expected signal (Sig.) events [assuming $\mathcal{B}(t \rightarrow ch) = 1\%$]. The three-lepton channels are taken from Ref. [9], have $H_T < 200$ GeV, and do not contain a τ_h . The stated uncertainties contain both systematic and statistical components.

Channel	E_T^{miss} (GeV)	N_b	Obs.	Exp.	Sig.
$\gamma\gamma\ell$	(50, 100)	≥ 1	1	2.3 ± 1.2	2.88 ± 0.39
	(30, 50)	≥ 1	2	1.1 ± 0.6	2.16 ± 0.30
	(0, 30)	≥ 1	2	2.1 ± 1.1	1.76 ± 0.24
	(50, 100)	0	7	9.5 ± 4.4	2.22 ± 0.31
	(100, ∞)	≥ 1	0	0.5 ± 0.4	0.92 ± 0.14
	(100, ∞)	0	1	2.2 ± 1.0	0.94 ± 0.17
$\ell\ell\ell(\text{OSSF1, below } Z)$	(50, 100)	≥ 1	48	48 ± 23	9.5 ± 2.3
	(0, 50)	≥ 1	34	42 ± 11	5.9 ± 1.2
$\ell\ell\ell(\text{OSSF0})$	(50, 100)	≥ 1	29	26 ± 13	5.9 ± 1.3
	(0, 50)	≥ 1	29	23 ± 10	4.3 ± 1.1

boson decay modes. For this purpose, we assume the SM branching fraction for the Higgs boson decay mode [37] under consideration, and ignore other decay modes. Table IX shows the results, illustrating the analysis sensitivity for the $t \rightarrow ch$ decay in each of the Higgs boson decay modes.

VIII. SUMMARY

We have performed a search for the $H \rightarrow hh$ and $A \rightarrow Zh$ decays of heavy scalar (H) and pseudoscalar (A) Higgs bosons, respectively, which occur in the extended Higgs sector described by the 2HDM. This is the first search for these decays carried out at the LHC. We used multilepton and diphoton final states from a data set corresponding to an integrated luminosity of 19.5 fb^{-1} of data recorded in 2012 from pp collisions at a center-of-mass energy of 8 TeV. We find no significant deviation from the SM expectations and place 95% C.L. cross section upper limits of approximately 7 pb on $\sigma\mathcal{B}$ for $H \rightarrow hh$ and 2 pb for $A \rightarrow Zh$. We further interpret these limits in the context of type I and type II 2HDMs, presenting exclusion contours in the $\tan\beta$ versus $\cos(\beta-\alpha)$ plane.

Using diphoton and multilepton search channels that are sensitive to the decay $t \rightarrow ch$, we place an upper limit of 0.56% on $\mathcal{B}(t \rightarrow ch)$, where the expected limit is 0.65%. This is a significant improvement over the earlier limit of 1.3% from the multilepton search alone [9].

ACKNOWLEDGMENTS

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: the Austrian Federal Ministry of Science, Research and Economy and the Austrian Science Fund; the Belgian Fonds de la Recherche Scientifique and Fonds voor Wetenschappelijk Onderzoek; the Brazilian Funding Agencies (CNPq, CAPES, FAPERJ, and FAPESP); the Bulgarian Ministry of Education and Science; CERN; the Chinese Academy of Sciences, Ministry of Science and Technology, and National Natural Science Foundation of China; the Colombian Funding Agency (COLCIENCIAS); the Croatian Ministry of Science, Education and Sport and the Croatian Science Foundation; the Research Promotion Foundation, Cyprus; the Ministry of Education and Research, Estonian Research Council via IUT23-4 and IUT23-6 and European Regional Development Fund, Estonia; the Academy of Finland, Finnish Ministry of Education and Culture and Helsinki Institute of Physics; the Institut National de Physique Nucléaire et de Physique des Particules/CNRS and Commissariat à l'Énergie Atomique et aux Énergies Alternatives/CEA, France; the Bundesministerium für Bildung und Forschung, Deutsche Forschungsgemeinschaft, and Helmholtz-Gemeinschaft Deutscher Forschungszentren, Germany; the General Secretariat for Research and Technology, Greece; the National Scientific Research Foundation and National Innovation Office, Hungary; the Department of Atomic Energy and the Department of Science and Technology, India; the Institute for Studies in Theoretical Physics and Mathematics, Iran; the Science Foundation, Ireland; the Istituto Nazionale di Fisica Nucleare, Italy; the Korean Ministry of Education, Science and Technology and the World Class University program of NRF, Republic of

Korea; the Lithuanian Academy of Sciences; the Ministry of Education and University of Malaya (Malaysia); the Mexican Funding Agencies (CINVESTAV, CONACYT, SEP, and UASLP-FAI); the Ministry of Business, Innovation and Employment, New Zealand; the Pakistan Atomic Energy Commission; the Ministry of Science and Higher Education and the National Science Centre, Poland; the Fundação para a Ciência e a Tecnologia, Portugal; JINR, Dubna; the Ministry of Education and Science of the Russian Federation, the Federal Agency of Atomic Energy of the Russian Federation, Russian Academy of Sciences, and the Russian Foundation for Basic Research; the Ministry of Education, Science and Technological Development of Serbia; the Secretaría de Estado de Investigación, Desarrollo e Innovación and Programa Consolider-Ingenio 2010, Spain; the Swiss Funding Agencies (ETH Board, ETH Zurich, PSI, SNF, UniZH, Canton Zurich, and SER); the Ministry of Science and Technology, Taipei; the Thailand Center of Excellence in Physics, the Institute for the Promotion of Teaching Science and Technology of Thailand, Special Task Force for Activating Research, and the National Science and Technology Development Agency of Thailand; the Scientific and Technical Research Council of Turkey and Turkish Atomic Energy Authority; the National Academy of Sciences of Ukraine and State Fund for Fundamental Researches, Ukraine; the Science and Technology Facilities Council, U.K.; the U.S. Department of Energy, and the U.S. National Science Foundation. Individuals have received support from the Marie-Curie programme and the European Research Council and EPLANET (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Council of Science and Industrial Research, India; the HOMING PLUS program of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund; the Compagnia di San Paolo (Torino); the Consorzio per la Fisica (Trieste); MIUR project 20108T4XTM (Italy); the Thalís and Aristeia programs cofinanced by EU-ESF and the Greek NSRF; and the National Priorities Research Program by Qatar National Research Fund.

- [1] ATLAS Collaboration, *Phys. Lett. B* **716**, 1 (2012).
 [2] CMS Collaboration, *Phys. Lett. B* **716**, 30 (2012).
 [3] CMS Collaboration, *J. High Energy Phys.* **06** (2013) 081.
 [4] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva, *Phys. Rep.* **516**, 1 (2012).
 [5] N. Craig and S. Thomas, *J. High Energy Phys.* **11** (2012) 083.
 [6] N. Craig, J. A. Evans, R. Gray, C. Kilic, M. Park, and S. Thomas, *J. High Energy Phys.* **02** (2013) 033.
 [7] N. Craig, J. A. Evans, R. Gray, M. Park, S. Somalwar, S. Thomas, and M. Walker, *Phys. Rev. D* **86**, 075002 (2012).
 [8] ATLAS Collaboration, *J. High Energy Phys.* **06** (2014) 008.
 [9] CMS Collaboration, *Phys. Rev. D* **90**, 032006 (2014).
 [10] CMS Collaboration, *JINST* **3**, S08004 (2008).
 [11] F. Maltoni and T. Stelzer, *J. High Energy Phys.* **02** (2003) 027.
 [12] S. Kretzer, H. L. Lai, F. I. Olness, and W. K. Tung, *Phys. Rev. D* **69**, 114005 (2004).
 [13] S. Agostinelli *et al.* (GEANT4), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
 [14] CMS Collaboration, *J. Phys. Conf. Ser.* **219**, 032053 (2010).
 [15] CMS Collaboration, CMS Physics Analysis Summary, Report No. CMS-PAS-PFT-09-001, 2009.
 [16] CMS Collaboration, CMS Physics Analysis Summary, Report No. CMS-PAS-PFT-10-001, 2010.
 [17] CMS Collaboration, CMS Physics Analysis Summary, Report No. CMS-PAS-EGM-10-004, 2010.
 [18] CMS Collaboration, *JINST* **8**, P09009 (2013).
 [19] CMS Collaboration, CMS Physics Analysis Summary, Report No. CMS-PAS-MUO-10-002, 2010.
 [20] CMS Collaboration, *JINST* **7**, P10002 (2012).
 [21] CMS Collaboration, *JINST* **7**, P01001 (2012).
 [22] CMS Collaboration, CMS Physics Analysis Summary, Report No. CMS-PAS-EGM-10-005, 2010.
 [23] M. Cacciari, G. P. Salam, and G. Soyez, *J. High Energy Phys.* **04** (2008) 063.
 [24] CMS Collaboration, *JINST* **8**, P04013 (2013).
 [25] CMS Collaboration, *J. High Energy Phys.* **11** (2012) 067.
 [26] CMS Collaboration, *Eur. Phys. J. C* **74**, 3076 (2014).
 [27] M. Cacciari and G. P. Salam, *Phys. Lett. B* **659**, 119 (2008).
 [28] CMS Collaboration, *J. High Energy Phys.* **10** (2011) 132.
 [29] S. Heinemeyer *et al.*, CERN, Report No. CERN-2013-004, 2013.
 [30] T. Junk, *Nucl. Instrum. Methods Phys. Res., Sect. A* **434**, 435 (1999).
 [31] A. L. Read, *J. Phys. G* **28**, 2693 (2002).
 [32] R. V. Harlander, S. Liebler, and H. Mantler, *Comput. Phys. Commun.* **184**, 1605 (2013).
 [33] D. Eriksson, J. Rathsmann, and O. Stal, *Comput. Phys. Commun.* **181**, 189 (2010).
 [34] S. Dittmaier *et al.* (LHC Higgs Cross Section Working Group), “Handbook of LHC Higgs cross sections: 1. Inclusive observables,” 2011 (unpublished).
 [35] N. Craig, J. Galloway, and S. Thomas, [arXiv:1305.2424](https://arxiv.org/abs/1305.2424).
 [36] K.-F. Chen, W.-S. Hou, C. Kao, and M. Kohda, *Phys. Lett. B* **725**, 378 (2013).
 [37] CMS Collaboration, *Phys. Rev. D* **89**, 092007 (2014).

V. Khachatryan,¹ A. M. Sirunyan,¹ A. Tumasyan,¹ W. Adam,² T. Bergauer,² M. Dragicevic,² J. Erö,² C. Fabjan,^{2b} M. Friedl,² R. Frühwirth,^{2b} V. M. Ghete,² C. Hartl,² N. Hörmann,² J. Hrubec,² M. Jeitler,^{2b} W. Kiesenhofer,² V. Knünz,² M. Krammer,^{2b} I. Krätschmer,² D. Liko,² I. Mikulec,² D. Rabady,^{2c} B. Rahbaran,² H. Rohringer,² R. Schöfbeck,² J. Strauss,² A. Taurok,² W. Treberer-Treberspurg,² W. Waltenberger,² C.-E. Wulz,^{2b} V. Mossolov,³ N. Shumeiko,³ J. Suarez Gonzalez,³ S. Alderweireldt,⁴ M. Bansal,⁴ S. Bansal,⁴ T. Cornelis,⁴ E. A. De Wolf,⁴ X. Janssen,⁴ A. Knutsson,⁴ S. Luyckx,⁴ S. Ochesanu,⁴ B. Roland,⁴ R. Rougny,⁴ M. Van De Klundert,⁴ H. Van Haevermaet,⁴ P. Van Mechelen,⁴ N. Van Remortel,⁴ A. Van Spilbeeck,⁴ F. Blekman,⁵ S. Blyweert,⁵ J. D’Hondt,⁵ N. Daci,⁵ N. Heracleous,⁵ J. Keaveney,⁵ S. Lowette,⁵ M. Maes,⁵ A. Olbrechts,⁵ Q. Python,⁵ D. Strom,⁵ S. Tavernier,⁵ W. Van Doninck,⁵ P. Van Mulders,⁵ G. P. Van Onsem,⁵ I. Vilella,⁵ C. Caillol,⁶ B. Clerbaux,⁶ G. De Lentdecker,⁶ D. Dobur,⁶ L. Favart,⁶ A. P. R. Gay,⁶ A. Grebenyuk,⁶ A. Léonard,⁶ A. Mohammadi,⁶ L. Perniè,^{6,c} T. Reis,⁶ T. Seva,⁶ L. Thomas,⁶ C. Vander Velde,⁶ P. Vanlaer,⁶ J. Wang,⁶ V. Adler,⁷ K. Beernaert,⁷ L. Benucci,⁷ A. Cimmino,⁷ S. Costantini,⁷ S. Crucy,⁷ S. Dildick,⁷ A. Fagot,⁷ G. Garcia,⁷ J. McCartin,⁷ A. A. Ocampo Rios,⁷ D. Ryckbosch,⁷ S. Salva Diblen,⁷ M. Sigamani,⁷ N. Strobbe,⁷ F. Thyssen,⁷ M. Tytgat,⁷ E. Yazgan,⁷ N. Zaganidis,⁷ S. Basegmez,⁸ C. Beluffi,^{8,d} G. Bruno,⁸ R. Castello,⁸ A. Caudron,⁸ L. Ceard,⁸ G. G. Da Silva,⁸ C. Delaere,⁸ T. du Pree,⁸ D. Favart,⁸ L. Forthomme,⁸ A. Giammanco,^{8,e} J. Hollar,⁸ P. Jez,⁸ M. Komm,⁸ V. Lemaitre,⁸ C. Nuttens,⁸ D. Pagano,⁸ L. Perrini,⁸ A. Pin,⁸ K. Piotrkowski,⁸ A. Popov,^{8,f} L. Quertenmont,⁸ M. Selvaggi,⁸ M. Vidal Marono,⁸ J. M. Vizan Garcia,⁸ N. Belyi,⁹ T. Caeberts,⁹ E. Daubie,⁹ G. H. Hammad,⁹ W. L. Aldá Júnior,¹⁰ G. A. Alves,¹⁰ L. Brito,¹⁰ M. Correa Martins Junior,¹⁰ T. Dos Reis Martins,¹⁰ C. Mora Herrera,¹⁰ M. E. Pol,¹⁰ W. Carvalho,¹¹ J. Chinellato,^{11,g} A. Custódio,¹¹ E. M. Da Costa,¹¹ D. De Jesus Damiao,¹¹ C. De Oliveira Martins,¹¹ S. Fonseca De Souza,¹¹ H. Malbouisson,¹¹ D. Matos Figueiredo,¹¹ L. Mundim,¹¹ H. Nogima,¹¹ W. L. Prado Da Silva,¹¹ J. Santaolalla,¹¹ A. Santoro,¹¹ A. Sznajder,¹¹ E. J. Tonelli Manganote,^{11,g} A. Vilela Pereira,¹¹ C. A. Bernardes,^{12b} S. Dogra,^{12a} T. R. Fernandez Perez Tomei,^{12a} E. M. Gregores,^{12b} P. G. Mercadante,^{12b} S. F. Novaes,^{12a} Sandra S. Padula,^{12a} A. Aleksandrov,¹³ V. Genchev,^{13,c} P. Iaydjiev,¹³ A. Marinov,¹³ S. Piperov,¹³ M. Rodozov,¹³ S. Stoykova,¹³ G. Sultanov,¹³

V. Tcholakov,¹³ M. Vutova,¹³ A. Dimitrov,¹⁴ I. Glushkov,¹⁴ R. Hadjiiska,¹⁴ V. Kozuharov,¹⁴ L. Litov,¹⁴ B. Pavlov,¹⁴ P. Petkov,¹⁴ J. G. Bian,¹⁵ G. M. Chen,¹⁵ H. S. Chen,¹⁵ M. Chen,¹⁵ R. Du,¹⁵ C. H. Jiang,¹⁵ S. Liang,¹⁵ R. Plestina,^{15,h} J. Tao,¹⁵ X. Wang,¹⁵ Z. Wang,¹⁵ C. Asawatangtrakuldee,¹⁶ Y. Ban,¹⁶ Y. Guo,¹⁶ Q. Li,¹⁶ W. Li,¹⁶ S. Liu,¹⁶ Y. Mao,¹⁶ S. J. Qian,¹⁶ D. Wang,¹⁶ L. Zhang,¹⁶ W. Zou,¹⁶ C. Avila,¹⁷ L. F. Chaparro Sierra,¹⁷ C. Florez,¹⁷ J. P. Gomez,¹⁷ B. Gomez Moreno,¹⁷ J. C. Sanabria,¹⁷ N. Godinovic,¹⁸ D. Lelas,¹⁸ D. Polic,¹⁸ I. Puljak,¹⁸ Z. Antunovic,¹⁹ M. Kovac,¹⁹ V. Brigljevic,²⁰ K. Kadija,²⁰ J. Letic,²⁰ D. Mekterovic,²⁰ L. Sudic,²⁰ A. Attikis,²¹ G. Mavromanolakis,²¹ J. Mousa,²¹ C. Nicolaou,²¹ F. Ptochos,²¹ P. A. Razis,²¹ M. Bodlak,²² M. Finger,²² M. Finger Jr.,^{22,i} Y. Assran,^{23,j} A. Ellithi Kamel,^{23,k} M. A. Mahmoud,^{23,l} A. Radi,^{23,m,n} M. Kadastik,²⁴ M. Murumaa,²⁴ M. Raidal,²⁴ A. Tiko,²⁴ P. Eerola,²⁵ G. Fedi,²⁵ M. Voutilainen,²⁵ J. Härkönen,²⁶ V. Karimäki,²⁶ R. Kinnunen,²⁶ M. J. Kortelainen,²⁶ T. Lampén,²⁶ K. Lassila-Perini,²⁶ S. Lehti,²⁶ T. Lindén,²⁶ P. Luukka,²⁶ T. Mäenpää,²⁶ T. Peltola,²⁶ E. Tuominen,²⁶ J. Tuominiemi,²⁶ E. Tuovinen,²⁶ L. Wendland,²⁶ T. Tuuva,²⁷ M. Besancon,²⁸ F. Couderc,²⁸ M. DeJardin,²⁸ D. Denegri,²⁸ B. Fabbro,²⁸ J. L. Faure,²⁸ C. Favaro,²⁸ F. Ferri,²⁸ S. Ganjour,²⁸ A. Givernaud,²⁸ P. Gras,²⁸ G. Hamel de Monchenault,²⁸ P. Jarry,²⁸ E. Locci,²⁸ J. Malcles,²⁸ J. Rander,²⁸ A. Rosowsky,²⁸ M. Titov,²⁸ S. Baffioni,²⁹ F. Beaudette,²⁹ P. Busson,²⁹ C. Charlot,²⁹ T. Dahms,²⁹ M. Dalchenko,²⁹ L. Dobrzynski,²⁹ N. Filipovic,²⁹ A. Florent,²⁹ R. Granier de Cassagnac,²⁹ L. Mastrolorenzo,²⁹ P. Miné,²⁹ C. Mironov,²⁹ I. N. Naranjo,²⁹ M. Nguyen,²⁹ C. Ochando,²⁹ P. Paganini,²⁹ S. Regnard,²⁹ R. Salerno,²⁹ J. B. Sauvan,²⁹ Y. Sirois,²⁹ C. Veelken,²⁹ Y. Yilmaz,²⁹ A. Zabi,²⁹ J.-L. Agram,^{30,o} J. Andrea,³⁰ A. Aubin,³⁰ D. Bloch,³⁰ J.-M. Brom,³⁰ E. C. Chabert,³⁰ C. Collard,³⁰ E. Conte,^{30,o} J.-C. Fontaine,^{30,o} D. Gelé,³⁰ U. Goerlach,³⁰ C. Goetzmann,³⁰ A.-C. Le Bihan,³⁰ P. Van Hove,³⁰ S. Gadrat,³¹ S. Beauceron,³² N. Beaupere,³² G. Boudoul,^{32,c} E. Bouvier,³² S. Brochet,³² C. A. Carrillo Montoya,³² J. Chasserat,³² R. Chierici,³² D. Contardo,^{32,c} P. Depasse,³² H. El Mamouni,³² J. Fan,³² J. Fay,³² S. Gascon,³² M. Gouzevitch,³² B. Ille,³² T. Kurca,³² M. Lethuillier,³² L. Mirabito,³² S. Perries,³² J. D. Ruiz Alvarez,³² D. Sabes,³² L. Sgandurra,³² V. Sordini,³² M. Vander Donckt,³² P. Verdier,³² S. Viret,³² H. Xiao,³² Z. Tsamalaidze,^{33,i} C. Autermann,³⁴ S. Beranek,³⁴ M. Bontenackels,³⁴ M. Edelhoff,³⁴ L. Feld,³⁴ O. Hindrichs,³⁴ K. Klein,³⁴ A. Ostapchuk,³⁴ A. Perieanu,³⁴ F. Raupach,³⁴ J. Sammet,³⁴ S. Schael,³⁴ H. Weber,³⁴ B. Wittmer,³⁴ V. Zhukov,^{34,f} M. Ata,³⁵ E. Dietz-Laursonn,³⁵ D. Duchardt,³⁵ M. Erdmann,³⁵ R. Fischer,³⁵ A. Güth,³⁵ T. Hebbeker,³⁵ C. Heidemann,³⁵ K. Hoepfner,³⁵ D. Klingebiel,³⁵ S. Knutzen,³⁵ P. Kreuzer,³⁵ M. Merschmeyer,³⁵ A. Meyer,³⁵ P. Millet,³⁵ M. Olschewski,³⁵ K. Padeken,³⁵ P. Papacz,³⁵ H. Reithler,³⁵ S. A. Schmitz,³⁵ L. Sonnenschein,³⁵ D. Teysier,³⁵ S. Thüer,³⁵ M. Weber,³⁵ V. Cherepanov,³⁶ Y. Erdogan,³⁶ G. Flügge,³⁶ H. Geenen,³⁶ M. Geisler,³⁶ W. Haj Ahmad,³⁶ A. Heister,³⁶ F. Hoehle,³⁶ B. Kargoll,³⁶ T. Kress,³⁶ Y. Kuessel,³⁶ J. Lingemann,^{36,c} A. Nowack,³⁶ I. M. Nugent,³⁶ L. Perchalla,³⁶ O. Pooth,³⁶ A. Stahl,³⁶ I. Asin,³⁷ N. Bartosik,³⁷ J. Behr,³⁷ W. Behrenhoff,³⁷ U. Behrens,³⁷ A. J. Bell,³⁷ M. Bergholz,^{37,p} A. Bethani,³⁷ K. Borras,³⁷ A. Burgmeier,³⁷ A. Cakir,³⁷ L. Calligaris,³⁷ A. Campbell,³⁷ S. Choudhury,³⁷ F. Costanza,³⁷ C. Diez Pardos,³⁷ S. Dooling,³⁷ T. Dorland,³⁷ G. Eckerlin,³⁷ D. Eckstein,³⁷ T. Eichhorn,³⁷ G. Flucke,³⁷ J. Garay Garcia,³⁷ A. Geiser,³⁷ P. Gunnellini,³⁷ J. Hauk,³⁷ M. Hempel,³⁷ D. Horton,³⁷ H. Jung,³⁷ A. Kalogeropoulos,³⁷ M. Kasemann,³⁷ P. Katsas,³⁷ J. Kieseler,³⁷ C. Kleinwort,³⁷ D. Krücker,³⁷ W. Lange,³⁷ J. Leonard,³⁷ K. Lipka,³⁷ A. Lobanov,³⁷ W. Lohmann,^{37,p} B. Lutz,³⁷ R. Mankel,³⁷ I. Marfin,³⁷ I.-A. Melzer-Pellmann,³⁷ A. B. Meyer,³⁷ G. Mittag,³⁷ J. Mnich,³⁷ A. Mussgiller,³⁷ S. Naumann-Emme,³⁷ A. Nayak,³⁷ O. Novgorodova,³⁷ F. Nowak,³⁷ E. Ntomari,³⁷ H. Perrey,³⁷ D. Pitzl,³⁷ R. Placakyte,³⁷ A. Raspereza,³⁷ P. M. Ribeiro Cipriano,³⁷ E. Ron,³⁷ M. Ö. Sahin,³⁷ J. Salfeld-Nebgen,³⁷ P. Saxena,³⁷ R. Schmidt,^{37,p} T. Schoerner-Sadenius,³⁷ M. Schröder,³⁷ C. Seitz,³⁷ S. Spannagel,³⁷ A. D. R. Vargas Trevino,³⁷ R. Walsh,³⁷ C. Wissing,³⁷ M. Aldaya Martin,³⁸ V. Blobel,³⁸ M. Centis Vignali,³⁸ A. R. Draeger,³⁸ J. Erfle,³⁸ E. Garutti,³⁸ K. Goebel,³⁸ M. Görner,³⁸ J. Haller,³⁸ M. Hoffmann,³⁸ R. S. Höing,³⁸ H. Kirschenmann,³⁸ R. Klanner,³⁸ R. Kogler,³⁸ J. Lange,³⁸ T. Lapsien,³⁸ T. Lenz,³⁸ I. Marchesini,³⁸ J. Ott,³⁸ T. Peiffer,³⁸ N. Pietsch,³⁸ J. Poehlsen,³⁸ T. Poehlsen,³⁸ D. Rathjens,³⁸ C. Sander,³⁸ H. Schettler,³⁸ P. Schleper,³⁸ E. Schlieckau,³⁸ A. Schmidt,³⁸ M. Seidel,³⁸ V. Sola,³⁸ H. Stadie,³⁸ G. Steinbrück,³⁸ D. Troendle,³⁸ E. Usai,³⁸ L. Vanelderen,³⁸ C. Barth,³⁹ C. Baus,³⁹ J. Berger,³⁹ C. Böser,³⁹ E. Butz,³⁹ T. Chwalek,³⁹ W. De Boer,³⁹ A. Descroix,³⁹ A. Dierlamm,³⁹ M. Feindt,³⁹ F. Frensch,³⁹ M. Giffels,³⁹ F. Hartmann,^{39,c} T. Hauth,^{39,c} U. Husemann,³⁹ I. Katkov,^{39,f} A. Kornmayer,^{39,c} E. Kuznetsova,³⁹ P. Lobelle Pardo,³⁹ M. U. Mozer,³⁹ Th. Müller,³⁹ A. Nürnberg,³⁹ G. Quast,³⁹ K. Rabbertz,³⁹ F. Ratnikov,³⁹ S. Röcker,³⁹ H. J. Simonis,³⁹ F. M. Stober,³⁹ R. Ulrich,³⁹ J. Wagner-Kuhr,³⁹ S. Wayand,³⁹ T. Weiler,³⁹ R. Wolf,³⁹ G. Anagnostou,⁴⁰ G. Daskalakis,⁴⁰ T. Geralis,⁴⁰ V. A. Giakoumopoulou,⁴⁰ A. Kyriakis,⁴⁰ D. Loukas,⁴⁰ A. Markou,⁴⁰ C. Markou,⁴⁰ A. Psallidas,⁴⁰ I. Topsis-Giotis,⁴⁰ A. Agapitos,⁴¹ A. Panagiotou,⁴¹ N. Saoulidou,⁴¹ E. Stiliaris,⁴¹ X. Aslanoglou,⁴² I. Evangelou,⁴² G. Flouris,⁴² C. Foudas,⁴² P. Kokkas,⁴² N. Manthos,⁴² I. Papadopoulos,⁴² E. Paradas,⁴² G. Bencze,⁴³ C. Hajdu,⁴³ P. Hidas,⁴³ D. Horvath,^{43,q} F. Sikler,⁴³

V. Veszpremi,⁴³ G. Vesztergombi,^{43,r} A. J. Zsigmond,⁴³ N. Beni,⁴⁴ S. Czellar,⁴⁴ J. Karacsi,^{44,s} J. Molnar,⁴⁴ J. Palinkas,⁴⁴ Z. Szillasi,⁴⁴ P. Raics,⁴⁵ Z. L. Trocsanyi,⁴⁵ B. Ujvari,⁴⁵ S. K. Swain,⁴⁶ S. B. Beri,⁴⁷ V. Bhatnagar,⁴⁷ N. Dhirra,⁴⁷ R. Gupta,⁴⁷ U. Bhawandeep,⁴⁷ A. K. Kalsi,⁴⁷ M. Kaur,⁴⁷ M. Mittal,⁴⁷ N. Nishu,⁴⁷ J. B. Singh,⁴⁷ Ashok Kumar,⁴⁸ Arun Kumar,⁴⁸ S. Ahuja,⁴⁸ A. Bhardwaj,⁴⁸ B. C. Choudhary,⁴⁸ A. Kumar,⁴⁸ S. Malhotra,⁴⁸ M. Naimuddin,⁴⁸ K. Ranjan,⁴⁸ V. Sharma,⁴⁸ S. Banerjee,⁴⁹ S. Bhattacharya,⁴⁹ K. Chatterjee,⁴⁹ S. Dutta,⁴⁹ B. Gomber,⁴⁹ Sa. Jain,⁴⁹ Sh. Jain,⁴⁹ R. Khurana,⁴⁹ A. Modak,⁴⁹ S. Mukherjee,⁴⁹ D. Roy,⁴⁹ S. Sarkar,⁴⁹ M. Sharan,⁴⁹ A. Abdulsalam,⁵⁰ D. Dutta,⁵⁰ S. Kailas,⁵⁰ V. Kumar,⁵⁰ A. K. Mohanty,^{50,c} L. M. Pant,⁵⁰ P. Shukla,⁵⁰ A. Topkar,⁵⁰ T. Aziz,⁵¹ S. Banerjee,⁵¹ S. Bhowmik,^{51,t} R. M. Chatterjee,⁵¹ R. K. Dewanjee,⁵¹ S. Dugad,⁵¹ S. Ganguly,⁵¹ S. Ghosh,⁵¹ M. Guchait,⁵¹ A. Gurtu,^{51,u} G. Kole,⁵¹ S. Kumar,⁵¹ M. Maity,^{51,t} G. Majumder,⁵¹ K. Mazumdar,⁵¹ G. B. Mohanty,⁵¹ B. Parida,⁵¹ K. Sudhakar,⁵¹ N. Wickramage,^{51,v} H. Bakhshiansohi,⁵² H. Behnamian,⁵² S. M. Etesami,^{52,w} A. Fahim,^{52,x} R. Goldouzian,⁵² A. Jafari,⁵² M. Khakzad,⁵² M. Mohammadi Najafabadi,⁵² M. Naseri,⁵² S. Pakinat Mehdiabadi,⁵² B. Safarzadeh,^{52,y} M. Zeinali,⁵² M. Felcini,⁵³ M. Grunewald,⁵³ M. Abbrescia,^{54a,54b} L. Barbore,^{54a,54b} C. Calabria,^{54a,54b} S. S. Chhibra,^{54a,54b} A. Colaleo,^{54a} D. Creanza,^{54a,54c} N. De Filippis,^{54a,54c} M. De Palma,^{54a,54b} L. Fiore,^{54a} G. Iaselli,^{54a,54c} G. Maggi,^{54a,54c} M. Maggi,^{54a} S. My,^{54a,54c} S. Nuzzo,^{54a,54b} A. Pompili,^{54a,54b} G. Pugliese,^{54a,54c} R. Radogna,^{54a,54b,c} G. Selvaggi,^{54a,54b} L. Silvestris,^{54a,c} G. Singh,^{54a,54b} R. Venditti,^{54a,54b} P. Verwilligen,^{54a} G. Zito,^{54a} G. Abbiendi,^{55a} A. C. Benvenuti,^{55a} D. Bonacorsi,^{55a,55b} S. Braibant-Giacomelli,^{55a,55b} L. Brigliadori,^{55a,55b} R. Campanini,^{55a,55b} P. Capiluppi,^{55a,55b} A. Castro,^{55a,55b} F. R. Cavallo,^{55a} G. Codispoti,^{55a,55b} M. Cuffiani,^{55a,55b} G. M. Dallavalle,^{55a} F. Fabbri,^{55a} A. Fanfani,^{55a,55b} D. Fasanella,^{55a,55b} P. Giacomelli,^{55a} C. Grandi,^{55a} L. Guiducci,^{55a,55b} S. Marcellini,^{55a} G. Masetti,^{55a,c} A. Montanari,^{55a} F. L. Navarria,^{55a,55b} A. Perrotta,^{55a} F. Primavera,^{55a,55b} A. M. Rossi,^{55a,55b} T. Rovelli,^{55a,55b} G. P. Siroli,^{55a,55b} N. Tosi,^{55a,55b} R. Travaglini,^{55a,55b} S. Albergo,^{56a,56b} G. Cappello,^{56a} M. Chiorboli,^{56a,56b} S. Costa,^{56a,56b} F. Giordano,^{56a,c} R. Potenza,^{56a,56b} A. Tricomi,^{56a,56b} C. Tuve,^{56a,56b} G. Barbagli,^{57a} V. Ciulli,^{57a,57b} C. Civinini,^{57a} R. D'Alessandro,^{57a,57b} E. Focardi,^{57a,57b} E. Gallo,^{57a} S. Gonzi,^{57a,57b} V. Gori,^{57a,57b,c} P. Lenzi,^{57a,57b} M. Meschini,^{57a} S. Paoletti,^{57a} G. Sguazzoni,^{57a} A. Tropiano,^{57a,57b} L. Benussi,⁵⁸ S. Bianco,⁵⁸ F. Fabbri,⁵⁸ D. Piccolo,⁵⁸ F. Ferro,^{59a} M. Lo Vetere,^{59a,59b} E. Robutti,^{59a} S. Tosi,^{59a,59b} M. E. Dinardo,^{60a,60b} S. Fiorendi,^{60a,60b,c} S. Gennai,^{60a,c} R. Gerosa,^{60a,60b,c} A. Ghezzi,^{60a,60b} P. Govoni,^{60a,60b} M. T. Lucchini,^{60a,60b,c} S. Malvezzi,^{60a} R. A. Manzoni,^{60a,60b} A. Martelli,^{60a,60b} B. Marzocchi,^{60a,60b} D. Menasce,^{60a} L. Moroni,^{60a} M. Paganoni,^{60a,60b} D. Pedrini,^{60a} S. Ragazzi,^{60a,60b} N. Redaelli,^{60a} T. Tabarelli de Fatis,^{60a,60b} S. Buontempo,^{61a} N. Cavallo,^{61a,61c} S. Di Guida,^{61a,61d,c} F. Fabozzi,^{61a,61c} A. O. M. Iorio,^{61a,61b} L. Lista,^{61a} S. Meola,^{61a,61d,c} M. Merola,^{61a} P. Paolucci,^{61a,c} P. Azzi,^{62a} N. Bacchetta,^{62a} D. Bisello,^{62a,62b} A. Branca,^{62a,62b} R. Carlin,^{62a,62b} P. Checchia,^{62a} M. Dall'Osso,^{62a,62b} T. Dorigo,^{62a} M. Galanti,^{62a,62b} F. Gasparini,^{62a,62b} U. Gasparini,^{62a,62b} P. Giubileo,^{62a,62b} F. Gonella,^{62a} A. Gozzelino,^{62a} K. Kanishchev,^{62a,62c} S. Lacaprara,^{62a} M. Margoni,^{62a,62b} A. T. Meneguzzo,^{62a,62b} J. Pazzini,^{62a,62b} N. Pozzobon,^{62a,62b} P. Ronchese,^{62a,62b} F. Simonetto,^{62a,62b} E. Torassa,^{62a} M. Tosi,^{62a,62b} P. Zotto,^{62a,62b} A. Zucchetta,^{62a,62b} G. Zumerle,^{62a,62b} M. Gabusi,^{63a,63b} S. P. Ratti,^{63a,63b} C. Riccardi,^{63a,63b} P. Salvini,^{63a} P. Vitulo,^{63a,63b} M. Biasini,^{64a,64b} G. M. Bilei,^{64a} D. Ciangottini,^{64a,64b} L. Fanò,^{64a,64b} P. Lariccia,^{64a,64b} G. Mantovani,^{64a,64b} M. Menichelli,^{64a} F. Romeo,^{64a,64b} A. Saha,^{64a} A. Santocchia,^{64a,64b} A. Spiezia,^{64a,64b,c} K. Androsov,^{65a,z} P. Azzurri,^{65a} G. Bagliesi,^{65a} J. Bernardini,^{65a} T. Boccali,^{65a} G. Broccolo,^{65a,65c} R. Castaldi,^{65a} M. A. Ciocci,^{65a,z} R. Dell'Orso,^{65a} S. Donato,^{65a,65c} F. Fiori,^{65a,65c} L. Foà,^{65a,65c} A. Giassi,^{65a} M. T. Grippo,^{65a,z} F. Ligabue,^{65a,65c} T. Lomtadze,^{65a} L. Martini,^{65a,65b} A. Messineo,^{65a,65b} C. S. Moon,^{65a,aa} F. Palla,^{65a,c} A. Rizzi,^{65a,65b} A. Savoy-Navarro,^{65a,bb} A. T. Serban,^{65a} P. Spagnolo,^{65a} P. Squillacioti,^{65a,z} R. Tenchini,^{65a} G. Tonelli,^{65a,65b} A. Venturi,^{65a} P. G. Verdini,^{65a} C. Vernieri,^{65a,65c,c} L. Barone,^{66a,66b} F. Cavallari,^{66a} G. D'imperio,^{66a,66b} D. Del Re,^{66a,66b} M. Diemoz,^{66a} M. Grassi,^{66a,66b} C. Jorda,^{66a} E. Longo,^{66a,66b} F. Margaroli,^{66a,66b} P. Meridiani,^{66a} F. Micheli,^{66a,66b,c} S. Nourbakhsh,^{66a,66b} G. Organtini,^{66a,66b} R. Paramatti,^{66a} S. Rahatlou,^{66a,66b} C. Rovelli,^{66a} F. Santanastasio,^{66a,66b} L. Soffi,^{66a,66b,c} P. Traczyk,^{66a,66b} N. Amapane,^{67a,67b} R. Arcidiacono,^{67a,67c} S. Argiro,^{67a,67b,c} M. Arneodo,^{67a,67c} R. Bellan,^{67a,67b} C. Biino,^{67a} N. Cartiglia,^{67a} S. Casasso,^{67a,67b,c} M. Costa,^{67a,67b} A. Degano,^{67a,67b} N. Demaria,^{67a} L. Finco,^{67a,67b} C. Mariotti,^{67a} S. Maselli,^{67a} E. Migliore,^{67a,67b} V. Monaco,^{67a,67b} M. Musich,^{67a} M. M. Obertino,^{67a,67c} G. Ortona,^{67a,67b} L. Pacher,^{67a,67b} N. Pastrone,^{67a} M. Pelliccioni,^{67a} G. L. Pinna Angioni,^{67a,67b} A. Potenza,^{67a,67b} A. Romero,^{67a,67b} M. Ruspa,^{67a,67c} R. Sacchi,^{67a,67b} A. Solano,^{67a,67b} A. Staiano,^{67a} U. Tamponi,^{67a} S. Belforte,^{68a} V. Candelise,^{68a,68b} M. Casarsa,^{68a} F. Cossutti,^{68a} G. Della Ricca,^{68a,68b} B. Gobbo,^{68a} C. La Licata,^{68a,68b} M. Marone,^{68a,68b} D. Montanino,^{68a,68b} A. Schizzi,^{68a,68b,c} T. Umer,^{68a,68b} A. Zanetti,^{68a} S. Chang,⁶⁹ A. Kropivnitskaya,⁶⁹ S. K. Nam,⁶⁹ D. H. Kim,⁷⁰ G. N. Kim,⁷⁰ M. S. Kim,⁷⁰ D. J. Kong,⁷⁰ S. Lee,⁷⁰ Y. D. Oh,⁷⁰ H. Park,⁷⁰ A. Sakharov,⁷⁰ D. C. Son,⁷⁰ T. J. Kim,⁷¹ J. Y. Kim,⁷² S. Song,⁷² S. Choi,⁷³ D. Gyun,⁷³ B. Hong,⁷³ M. Jo,⁷³ H. Kim,⁷³ Y. Kim,⁷³ B. Lee,⁷³

K. S. Lee,⁷³ S. K. Park,⁷³ Y. Roh,⁷³ M. Choi,⁷⁴ J. H. Kim,⁷⁴ I. C. Park,⁷⁴ S. Park,⁷⁴ G. Ryu,⁷⁴ M. S. Ryu,⁷⁴ Y. Choi,⁷⁵ Y. K. Choi,⁷⁵ J. Goh,⁷⁵ D. Kim,⁷⁵ E. Kwon,⁷⁵ J. Lee,⁷⁵ H. Seo,⁷⁵ I. Yu,⁷⁵ A. Juodagalvis,⁷⁶ J. R. Komaragiri,⁷⁷ M. A. B. Md Ali,⁷⁷ H. Castilla-Valdez,⁷⁸ E. De La Cruz-Burelo,⁷⁸ I. Heredia-de La Cruz,^{78,cc} R. Lopez-Fernandez,⁷⁸ A. Sanchez-Hernandez,⁷⁸ S. Carrillo Moreno,⁷⁹ F. Vazquez Valencia,⁷⁹ I. Pedraza,⁸⁰ H. A. Salazar Ibarguen,⁸⁰ E. Casimiro Linares,⁸¹ A. Morelos Pineda,⁸¹ D. Krofcheck,⁸² P. H. Butler,⁸³ S. Reucroft,⁸³ A. Ahmad,⁸⁴ M. Ahmad,⁸⁴ Q. Hassan,⁸⁴ H. R. Hoorani,⁸⁴ S. Khalid,⁸⁴ W. A. Khan,⁸⁴ T. Khurshid,⁸⁴ M. A. Shah,⁸⁴ M. Shoaib,⁸⁴ H. Bialkowska,⁸⁵ M. Bluj,⁸⁵ B. Boimska,⁸⁵ T. Frueboes,⁸⁵ M. Górski,⁸⁵ M. Kazana,⁸⁵ K. Nawrocki,⁸⁵ K. Romanowska-Rybinska,⁸⁵ M. Szleper,⁸⁵ P. Zalewski,⁸⁵ G. Brona,⁸⁶ K. Bunkowski,⁸⁶ M. Cwiok,⁸⁶ W. Dominik,⁸⁶ K. Doroba,⁸⁶ A. Kalinowski,⁸⁶ M. Konecki,⁸⁶ J. Krolikowski,⁸⁶ M. Misiura,⁸⁶ M. Olszewski,⁸⁶ W. Wolszczak,⁸⁶ P. Bargassa,⁸⁷ C. Beirão Da Cruz E Silva,⁸⁷ P. Faccioli,⁸⁷ P. G. Ferreira Parracho,⁸⁷ M. Gallinaro,⁸⁷ F. Nguyen,⁸⁷ J. Rodrigues Antunes,⁸⁷ J. Seixas,⁸⁷ J. Varela,⁸⁷ P. Vischia,⁸⁷ S. Afanasiev,⁸⁸ M. Gavrilenko,⁸⁸ I. Golutvin,⁸⁸ A. Kamenev,⁸⁸ V. Karjavin,⁸⁸ V. Konoplyanikov,⁸⁸ A. Lanev,⁸⁸ A. Malakhov,⁸⁸ V. Matveev,^{88,dd} P. Moisenz,⁸⁸ V. Palichik,⁸⁸ V. Perelygin,⁸⁸ M. Savina,⁸⁸ S. Shmatov,⁸⁸ S. Shulha,⁸⁸ N. Skatchkov,⁸⁸ V. Smirnov,⁸⁸ A. Zarubin,⁸⁸ V. Golovtsov,⁸⁹ Y. Ivanov,⁸⁹ V. Kim,^{89,ee} P. Levchenko,⁸⁹ V. Murzin,⁸⁹ V. Oreshkin,⁸⁹ I. Smirnov,⁸⁹ V. Sulimov,⁸⁹ L. Uvarov,⁸⁹ S. Vavilov,⁸⁹ A. Vorobyev,⁸⁹ An. Vorobyev,⁸⁹ Yu. Andreev,⁹⁰ A. Dermenev,⁹⁰ S. Gninenko,⁹⁰ N. Golubev,⁹⁰ M. Kirsanov,⁹⁰ N. Krasnikov,⁹⁰ A. Pashenkov,⁹⁰ D. Tlisov,⁹⁰ A. Toropin,⁹⁰ V. Epshteyn,⁹¹ V. Gavrillov,⁹¹ N. Lychkovskaya,⁹¹ V. Popov,⁹¹ G. Safronov,⁹¹ S. Semenov,⁹¹ A. Spiridonov,⁹¹ V. Stolin,⁹¹ E. Vlasov,⁹¹ A. Zhokin,⁹¹ V. Andreev,⁹² M. Azarkin,⁹² I. Dremin,⁹² M. Kirakosyan,⁹² A. Leonidov,⁹² G. Mesyats,⁹² S. V. Rusakov,⁹² A. Vinogradov,⁹² A. Belyaev,⁹³ E. Boos,⁹³ V. Bunichev,⁹³ M. Dubinin,^{93,ff} L. Dudko,⁹³ A. Ershov,⁹³ V. Klyukhin,⁹³ O. Kodolova,⁹³ I. Lokhtin,⁹³ S. Obraztsov,⁹³ S. Petrushanko,⁹³ V. Savrin,⁹³ A. Snigirev,⁹³ I. Azhgirey,⁹⁴ I. Bayshev,⁹⁴ S. Bitiukov,⁹⁴ V. Kachanov,⁹⁴ A. Kalinin,⁹⁴ D. Konstantinov,⁹⁴ V. Krychkin,⁹⁴ V. Petrov,⁹⁴ R. Ryutin,⁹⁴ A. Sobol,⁹⁴ L. Tourtchanovitch,⁹⁴ S. Troshin,⁹⁴ N. Tyurin,⁹⁴ A. Uzunian,⁹⁴ A. Volkov,⁹⁴ P. Adzic,^{95,gg} M. Ekmedzic,⁹⁵ J. Milosevic,⁹⁵ V. Rekovic,⁹⁵ J. Alcaraz Maestre,⁹⁶ C. Battilana,⁹⁶ E. Calvo,⁹⁶ M. Cerrada,⁹⁶ M. Chamizo Llatas,⁹⁶ N. Colino,⁹⁶ B. De La Cruz,⁹⁶ A. Delgado Peris,⁹⁶ D. Domínguez Vázquez,⁹⁶ A. Escalante Del Valle,⁹⁶ C. Fernandez Bedoya,⁹⁶ J. P. Fernández Ramos,⁹⁶ J. Flix,⁹⁶ M. C. Fouz,⁹⁶ P. Garcia-Abia,⁹⁶ O. Gonzalez Lopez,⁹⁶ S. Goy Lopez,⁹⁶ J. M. Hernandez,⁹⁶ M. I. Josa,⁹⁶ G. Merino,⁹⁶ E. Navarro De Martino,⁹⁶ A. Pérez-Calero Yzquierdo,⁹⁶ J. Puerta Pelayo,⁹⁶ A. Quintario Olmeda,⁹⁶ I. Redondo,⁹⁶ L. Romero,⁹⁶ M. S. Soares,⁹⁶ C. Albajar,⁹⁷ J. F. de Trocóniz,⁹⁷ M. Missiroli,⁹⁷ D. Moran,⁹⁷ H. Brun,⁹⁸ J. Cuevas,⁹⁸ J. Fernandez Menendez,⁹⁸ S. Folgueras,⁹⁸ I. Gonzalez Caballero,⁹⁸ L. Lloret Iglesias,⁹⁸ J. A. Brochero Cifuentes,⁹⁹ I. J. Cabrillo,⁹⁹ A. Calderon,⁹⁹ J. Duarte Campderros,⁹⁹ M. Fernandez,⁹⁹ G. Gomez,⁹⁹ A. Graziano,⁹⁹ A. Lopez Virto,⁹⁹ J. Marco,⁹⁹ R. Marco,⁹⁹ C. Martinez Rivero,⁹⁹ F. Matorras,⁹⁹ F. J. Munoz Sanchez,⁹⁹ J. Piedra Gomez,⁹⁹ T. Rodrigo,⁹⁹ A. Y. Rodríguez-Marrero,⁹⁹ A. Ruiz-Jimeno,⁹⁹ L. Scodellaro,⁹⁹ I. Vila,⁹⁹ R. Vilar Cortabitarte,⁹⁹ D. Abbaneo,¹⁰⁰ E. Auffray,¹⁰⁰ G. Auzinger,¹⁰⁰ M. Bachtis,¹⁰⁰ P. Baillon,¹⁰⁰ A. H. Ball,¹⁰⁰ D. Barney,¹⁰⁰ A. Benaglia,¹⁰⁰ J. Bendavid,¹⁰⁰ L. Benhabib,¹⁰⁰ J. F. Benitez,¹⁰⁰ C. Bernet,^{100,h} G. Bianchi,¹⁰⁰ P. Bloch,¹⁰⁰ A. Bocci,¹⁰⁰ A. Bonato,¹⁰⁰ O. Bondu,¹⁰⁰ C. Botta,¹⁰⁰ H. Breuker,¹⁰⁰ T. Camporesi,¹⁰⁰ G. Cerminara,¹⁰⁰ S. Colafranceschi,^{100,hh} M. D'Alfonso,¹⁰⁰ D. d'Enterria,¹⁰⁰ A. Dabrowski,¹⁰⁰ A. David,¹⁰⁰ F. De Guio,¹⁰⁰ A. De Roeck,¹⁰⁰ S. De Visscher,¹⁰⁰ M. Dobson,¹⁰⁰ M. Dordevic,¹⁰⁰ N. Dupont-Sagorin,¹⁰⁰ A. Elliott-Peisert,¹⁰⁰ J. Eugster,¹⁰⁰ G. Franzoni,¹⁰⁰ W. Funk,¹⁰⁰ D. Gigi,¹⁰⁰ K. Gill,¹⁰⁰ D. Giordano,¹⁰⁰ M. Girone,¹⁰⁰ F. Glege,¹⁰⁰ R. Guida,¹⁰⁰ S. Gundacker,¹⁰⁰ M. Guthoff,¹⁰⁰ J. Hammer,¹⁰⁰ M. Hansen,¹⁰⁰ P. Harris,¹⁰⁰ J. Hegeman,¹⁰⁰ V. Innocente,¹⁰⁰ P. Janot,¹⁰⁰ K. Kousouris,¹⁰⁰ K. Krajczar,¹⁰⁰ P. Lecoq,¹⁰⁰ C. Lourenço,¹⁰⁰ N. Magini,¹⁰⁰ L. Malgeri,¹⁰⁰ M. Mannelli,¹⁰⁰ J. Marrouche,¹⁰⁰ L. Masetti,¹⁰⁰ F. Meijers,¹⁰⁰ S. Mersi,¹⁰⁰ E. Meschi,¹⁰⁰ F. Moortgat,¹⁰⁰ S. Morovic,¹⁰⁰ M. Mulders,¹⁰⁰ P. Musella,¹⁰⁰ L. Orsini,¹⁰⁰ L. Pape,¹⁰⁰ E. Perez,¹⁰⁰ L. Perrozzi,¹⁰⁰ A. Petrilli,¹⁰⁰ G. Petrucciani,¹⁰⁰ A. Pfeiffer,¹⁰⁰ M. Pierini,¹⁰⁰ M. Pimiä,¹⁰⁰ D. Piparo,¹⁰⁰ M. Plagge,¹⁰⁰ A. Racz,¹⁰⁰ G. Rolandi,^{100,ii} M. Rovere,¹⁰⁰ H. Sakulin,¹⁰⁰ C. Schäfer,¹⁰⁰ C. Schwick,¹⁰⁰ A. Sharma,¹⁰⁰ P. Siegrist,¹⁰⁰ P. Silva,¹⁰⁰ M. Simon,¹⁰⁰ P. Spicas,^{100,ji} D. Spiga,¹⁰⁰ J. Steggemann,¹⁰⁰ B. Stieger,¹⁰⁰ M. Stoye,¹⁰⁰ D. Treille,¹⁰⁰ A. Tsiros,¹⁰⁰ G. I. Veres,^{100,r} J. R. Vlimant,¹⁰⁰ N. Wardle,¹⁰⁰ H. K. Wöhri,¹⁰⁰ H. Wollny,¹⁰⁰ W. D. Zeuner,¹⁰⁰ W. Bertl,¹⁰¹ K. Deiters,¹⁰¹ W. Erdmann,¹⁰¹ R. Horisberger,¹⁰¹ Q. Ingram,¹⁰¹ H. C. Kaestli,¹⁰¹ D. Kotlinski,¹⁰¹ U. Langenegger,¹⁰¹ D. Renker,¹⁰¹ T. Rohe,¹⁰¹ F. Bachmair,¹⁰² L. Bäni,¹⁰² L. Bianchini,¹⁰² P. Bortignon,¹⁰² M. A. Buchmann,¹⁰² B. Casal,¹⁰² N. Chanon,¹⁰² A. Deisher,¹⁰² G. Dissertori,¹⁰² M. Dittmar,¹⁰² M. Donegà,¹⁰² M. Dünser,¹⁰² P. Eller,¹⁰² C. Grab,¹⁰² D. Hits,¹⁰² W. Lustermann,¹⁰² B. Mangano,¹⁰² A. C. Marini,¹⁰² P. Martinez Ruiz del Arbol,¹⁰² D. Meister,¹⁰² N. Mohr,¹⁰² C. Nägeli,^{102,kk} F. Nessi-Tedaldi,¹⁰² F. Pandolfi,¹⁰² F. Pauss,¹⁰² M. Peruzzi,¹⁰² M. Quittnat,¹⁰² L. Rebane,¹⁰² M. Rossini,¹⁰² A. Starodumov,^{102,ll} M. Takahashi,¹⁰² K. Theofilatos,¹⁰² R. Wallny,¹⁰² H. A. Weber,¹⁰² C. Amsler,^{103,mm} M. F. Canelli,¹⁰³

V. Chiochia,¹⁰³ A. De Cosa,¹⁰³ A. Hinzmann,¹⁰³ T. Hreus,¹⁰³ B. Kilminster,¹⁰³ C. Lange,¹⁰³ B. Millan Mejias,¹⁰³ J. Ngadiuba,¹⁰³ P. Robmann,¹⁰³ F. J. Ronga,¹⁰³ S. Taroni,¹⁰³ M. Verzetti,¹⁰³ Y. Yang,¹⁰³ M. Cardaci,¹⁰⁴ K. H. Chen,¹⁰⁴ C. Ferro,¹⁰⁴ C. M. Kuo,¹⁰⁴ W. Lin,¹⁰⁴ Y. J. Lu,¹⁰⁴ R. Volpe,¹⁰⁴ S. S. Yu,¹⁰⁴ P. Chang,¹⁰⁵ Y. H. Chang,¹⁰⁵ Y. W. Chang,¹⁰⁵ Y. Chao,¹⁰⁵ K. F. Chen,¹⁰⁵ P. H. Chen,¹⁰⁵ C. Dietz,¹⁰⁵ U. Grundler,¹⁰⁵ W.-S. Hou,¹⁰⁵ K. Y. Kao,¹⁰⁵ Y. J. Lei,¹⁰⁵ Y. F. Liu,¹⁰⁵ R.-S. Lu,¹⁰⁵ D. Majumder,¹⁰⁵ E. Petrakou,¹⁰⁵ Y. M. Tzeng,¹⁰⁵ R. Wilken,¹⁰⁵ B. Asavapibhop,¹⁰⁶ N. Srimanobhas,¹⁰⁶ N. Suwonjandee,¹⁰⁶ A. Adiguzel,¹⁰⁷ M. N. Bakirci,^{107,nn} S. Cerci,^{107,oo} C. Dozen,¹⁰⁷ I. Dumanoglu,¹⁰⁷ E. Eskut,¹⁰⁷ S. Girgis,¹⁰⁷ G. Gokbulut,¹⁰⁷ E. Gurpinar,¹⁰⁷ I. Hos,¹⁰⁷ E. E. Kangal,¹⁰⁷ A. Kayis Topaksu,¹⁰⁷ G. Onengut,^{107,pp} K. Ozdemir,¹⁰⁷ S. Ozturk,^{107,nn} A. Polatoz,¹⁰⁷ K. Sogut,^{107,qq} D. Sunar Cerci,^{107,oo} B. Tali,^{107,oo} H. Topakli,^{107,nn} M. Vergili,¹⁰⁷ I. V. Akin,¹⁰⁸ B. Bilin,¹⁰⁸ S. Bilmis,¹⁰⁸ H. Gamsizkan,¹⁰⁸ G. Karapinar,^{108,rr} K. Ocalan,¹⁰⁸ S. Sekmen,¹⁰⁸ U. E. Surat,¹⁰⁸ M. Yalvac,¹⁰⁸ M. Zeyrek,¹⁰⁸ E. Gülmez,¹⁰⁹ B. Isildak,^{109,ss} M. Kaya,^{109,tt} O. Kaya,^{109,uu} H. Bahtiyar,^{110,vv} E. Barlas,¹¹⁰ K. Cankocak,¹¹⁰ F. I. Vardarlı,¹¹⁰ M. Yücel,¹¹⁰ L. Levchuk,¹¹¹ P. Sorokin,¹¹¹ J. J. Brooke,¹¹² E. Clement,¹¹² D. Cussans,¹¹² H. Flacher,¹¹² R. Frazier,¹¹² J. Goldstein,¹¹² M. Grimes,¹¹² G. P. Heath,¹¹² H. F. Heath,¹¹² J. Jacob,¹¹² L. Kreczko,¹¹² C. Lucas,¹¹² Z. Meng,¹¹² D. M. Newbold,^{112,ww} S. Paramesvaran,¹¹² A. Poll,¹¹² S. Senkin,¹¹² V. J. Smith,¹¹² T. Williams,¹¹² K. W. Bell,¹¹³ A. Belyaev,^{113,xx} C. Brew,¹¹³ R. M. Brown,¹¹³ D. J. A. Cockerill,¹¹³ J. A. Coughlan,¹¹³ K. Harder,¹¹³ S. Harper,¹¹³ E. Olaiya,¹¹³ D. Petyt,¹¹³ C. H. Shepherd-Themistocleous,¹¹³ A. Thea,¹¹³ I. R. Tomalin,¹¹³ W. J. Womersley,¹¹³ S. D. Worm,¹¹³ M. Baber,¹¹⁴ R. Bainbridge,¹¹⁴ O. Buchmuller,¹¹⁴ D. Burton,¹¹⁴ D. Colling,¹¹⁴ N. Cripps,¹¹⁴ M. Cutajar,¹¹⁴ P. Dauncey,¹¹⁴ G. Davies,¹¹⁴ M. Della Negra,¹¹⁴ P. Dunne,¹¹⁴ W. Ferguson,¹¹⁴ J. Fulcher,¹¹⁴ D. Futyan,¹¹⁴ A. Gilbert,¹¹⁴ G. Hall,¹¹⁴ G. Iles,¹¹⁴ M. Jarvis,¹¹⁴ G. Karapostoli,¹¹⁴ M. Kenzie,¹¹⁴ R. Lane,¹¹⁴ R. Lucas,^{114,ww} L. Lyons,¹¹⁴ A.-M. Magnan,¹¹⁴ S. Malik,¹¹⁴ B. Mathias,¹¹⁴ J. Nash,¹¹⁴ A. Nikitenko,^{114,ll} J. Pela,¹¹⁴ M. Pesaresi,¹¹⁴ K. Petridis,¹¹⁴ D. M. Raymond,¹¹⁴ S. Rogerson,¹¹⁴ A. Rose,¹¹⁴ C. Seez,¹¹⁴ P. Sharp,^{114,a} A. Tapper,¹¹⁴ M. Vazquez Acosta,¹¹⁴ T. Virdee,¹¹⁴ J. E. Cole,¹¹⁵ P. R. Hobson,¹¹⁵ A. Khan,¹¹⁵ P. Kyberd,¹¹⁵ D. Leggat,¹¹⁵ D. Leslie,¹¹⁵ W. Martin,¹¹⁵ I. D. Reid,¹¹⁵ P. Symonds,¹¹⁵ L. Teodorescu,¹¹⁵ M. Turner,¹¹⁵ J. Dittmann,¹¹⁶ K. Hatakeyama,¹¹⁶ A. Kashi,¹¹⁶ H. Liu,¹¹⁶ T. Scarborough,¹¹⁶ O. Charaf,¹¹⁷ S. I. Cooper,¹¹⁷ C. Henderson,¹¹⁷ P. Rumerio,¹¹⁷ A. Avetisyan,¹¹⁸ T. Bose,¹¹⁸ C. Fantasia,¹¹⁸ P. Lawson,¹¹⁸ C. Richardson,¹¹⁸ J. Rohlf,¹¹⁸ D. Sperka,¹¹⁸ J. St. John,¹¹⁸ L. Sulak,¹¹⁸ J. Alimena,¹¹⁹ E. Berry,¹¹⁹ S. Bhattacharya,¹¹⁹ G. Christopher,¹¹⁹ D. Cutts,¹¹⁹ Z. Demiragli,¹¹⁹ A. Ferapontov,¹¹⁹ A. Garabedian,¹¹⁹ U. Heintz,¹¹⁹ G. Kukartsev,¹¹⁹ E. Laird,¹¹⁹ G. Landsberg,¹¹⁹ M. Luk,¹¹⁹ M. Narain,¹¹⁹ M. Segala,¹¹⁹ T. Sinthuprasith,¹¹⁹ T. Speer,¹¹⁹ J. Swanson,¹¹⁹ R. Breedon,¹²⁰ G. Breto,¹²⁰ M. Calderon De La Barca Sanchez,¹²⁰ S. Chauhan,¹²⁰ M. Chertok,¹²⁰ J. Conway,¹²⁰ R. Conway,¹²⁰ P. T. Cox,¹²⁰ R. Erbacher,¹²⁰ M. Gardner,¹²⁰ W. Ko,¹²⁰ R. Lander,¹²⁰ T. Miceli,¹²⁰ M. Mulhearn,¹²⁰ D. Pellett,¹²⁰ J. Pilot,¹²⁰ F. Ricci-Tam,¹²⁰ M. Searle,¹²⁰ S. Shalhout,¹²⁰ J. Smith,¹²⁰ M. Squires,¹²⁰ D. Stolp,¹²⁰ M. Tripathi,¹²⁰ S. Wilbur,¹²⁰ R. Yohay,¹²⁰ R. Cousins,¹²¹ P. Everaerts,¹²¹ C. Farrell,¹²¹ J. Hauser,¹²¹ M. Ignatenko,¹²¹ G. Rakness,¹²¹ E. Takasugi,¹²¹ V. Valuev,¹²¹ M. Weber,¹²¹ J. Babb,¹²² K. Burt,¹²² R. Clare,¹²² J. Ellison,¹²² J. W. Gary,¹²² G. Hanson,¹²² J. Heilman,¹²² M. Iova Rikova,¹²² P. Jandir,¹²² E. Kennedy,¹²² F. Lacroix,¹²² H. Liu,¹²² O. R. Long,¹²² A. Luthra,¹²² M. Malberti,¹²² H. Nguyen,¹²² M. Olmedo Negrete,¹²² A. Shrinivas,¹²² S. Sumowidagdo,¹²² S. Wimpenny,¹²² W. Andrews,¹²³ J. G. Branson,¹²³ G. B. Cerati,¹²³ S. Cittolin,¹²³ R. T. D'Agnolo,¹²³ D. Evans,¹²³ A. Holzner,¹²³ R. Kelley,¹²³ D. Klein,¹²³ M. Lebourgeois,¹²³ J. Letts,¹²³ I. Macneill,¹²³ D. Olivito,¹²³ S. Padhi,¹²³ C. Palmer,¹²³ M. Pieri,¹²³ M. Sani,¹²³ V. Sharma,¹²³ S. Simon,¹²³ E. Sudano,¹²³ M. Tadel,¹²³ Y. Tu,¹²³ A. Vartak,¹²³ C. Welke,¹²³ F. Würthwein,¹²³ A. Yagil,¹²³ J. Yoo,¹²³ D. Barge,¹²⁴ J. Bradmiller-Feld,¹²⁴ C. Campagnari,¹²⁴ T. Danielson,¹²⁴ A. Dishaw,¹²⁴ K. Flowers,¹²⁴ M. Franco Sevilla,¹²⁴ P. Geffert,¹²⁴ C. George,¹²⁴ F. Golf,¹²⁴ L. Gouskos,¹²⁴ J. Incandela,¹²⁴ C. Justus,¹²⁴ N. Mccoll,¹²⁴ J. Richman,¹²⁴ D. Stuart,¹²⁴ W. To,¹²⁴ C. West,¹²⁴ A. Apresyan,¹²⁵ A. Bornheim,¹²⁵ J. Bunn,¹²⁵ Y. Chen,¹²⁵ E. Di Marco,¹²⁵ J. Duarte,¹²⁵ A. Mott,¹²⁵ H. B. Newman,¹²⁵ C. Pena,¹²⁵ C. Rogan,¹²⁵ M. Spiropulu,¹²⁵ V. Timciuc,¹²⁵ R. Wilkinson,¹²⁵ S. Xie,¹²⁵ R. Y. Zhu,¹²⁵ V. Azzolini,¹²⁶ A. Calamba,¹²⁶ B. Carlson,¹²⁶ T. Ferguson,¹²⁶ Y. Iiyama,¹²⁶ M. Paulini,¹²⁶ J. Russ,¹²⁶ H. Vogel,¹²⁶ I. Vorobiev,¹²⁶ J. P. Cumalat,¹²⁷ W. T. Ford,¹²⁷ A. Gaz,¹²⁷ E. Luiggi Lopez,¹²⁷ U. Nauenberg,¹²⁷ J. G. Smith,¹²⁷ K. Stenson,¹²⁷ K. A. Ulmer,¹²⁷ S. R. Wagner,¹²⁷ J. Alexander,¹²⁸ A. Chatterjee,¹²⁸ J. Chu,¹²⁸ S. Dittmer,¹²⁸ N. Eggert,¹²⁸ N. Mirman,¹²⁸ G. Nicolas Kaufman,¹²⁸ J. R. Patterson,¹²⁸ A. Ryd,¹²⁸ E. Salvati,¹²⁸ L. Skinnari,¹²⁸ W. Sun,¹²⁸ W. D. Teo,¹²⁸ J. Thom,¹²⁸ J. Thompson,¹²⁸ J. Tucker,¹²⁸ Y. Weng,¹²⁸ L. Winstrom,¹²⁸ P. Wittich,¹²⁸ D. Winn,¹²⁹ S. Abdullin,¹³⁰ M. Albrow,¹³⁰ J. Anderson,¹³⁰ G. Apollinari,¹³⁰ L. A. T. Bauerick,¹³⁰ A. Beretvas,¹³⁰ J. Berryhill,¹³⁰ P. C. Bhat,¹³⁰ K. Burkett,¹³⁰ J. N. Butler,¹³⁰ H. W. K. Cheung,¹³⁰ F. Chlebana,¹³⁰ S. Cihangir,¹³⁰ V. D. Elvira,¹³⁰ I. Fisk,¹³⁰ J. Freeman,¹³⁰ Y. Gao,¹³⁰ E. Gottschalk,¹³⁰ L. Gray,¹³⁰ D. Green,¹³⁰ S. Grünendahl,¹³⁰ O. Gutsche,¹³⁰ J. Hanlon,¹³⁰ D. Hare,¹³⁰ R. M. Harris,¹³⁰ J. Hirschauer,¹³⁰ B. Hooberman,¹³⁰

S. Jindariani,¹³⁰ M. Johnson,¹³⁰ U. Joshi,¹³⁰ K. Kaadze,¹³⁰ B. Klima,¹³⁰ B. Kreis,¹³⁰ S. Kwan,¹³⁰ J. Linacre,¹³⁰ D. Lincoln,¹³⁰ R. Lipton,¹³⁰ T. Liu,¹³⁰ J. Lykken,¹³⁰ K. Maeshima,¹³⁰ J. M. Marraffino,¹³⁰ V. I. Martinez Outschoorn,¹³⁰ S. Maruyama,¹³⁰ D. Mason,¹³⁰ P. McBride,¹³⁰ K. Mishra,¹³⁰ S. Mrenna,¹³⁰ Y. Musienko,^{130,dd} S. Nahn,¹³⁰ C. Newman-Holmes,¹³⁰ V. O'Dell,¹³⁰ O. Prokofyev,¹³⁰ E. Sexton-Kennedy,¹³⁰ S. Sharma,¹³⁰ A. Soha,¹³⁰ W. J. Spalding,¹³⁰ L. Spiegel,¹³⁰ L. Taylor,¹³⁰ S. Tkaczyk,¹³⁰ N. V. Tran,¹³⁰ L. Uplegger,¹³⁰ E. W. Vaandering,¹³⁰ R. Vidal,¹³⁰ A. Whitbeck,¹³⁰ J. Whitmore,¹³⁰ F. Yang,¹³⁰ D. Acosta,¹³¹ P. Avery,¹³¹ D. Bourilkov,¹³¹ M. Carver,¹³¹ T. Cheng,¹³¹ D. Curry,¹³¹ S. Das,¹³¹ M. De Gruttola,¹³¹ G. P. Di Giovanni,¹³¹ R. D. Field,¹³¹ M. Fisher,¹³¹ I. K. Furic,¹³¹ J. Hugon,¹³¹ J. Konigsberg,¹³¹ A. Korytov,¹³¹ T. Kypreos,¹³¹ J. F. Low,¹³¹ K. Matchev,¹³¹ P. Milenovic,^{131,yy} G. Mitselmakher,¹³¹ L. Muniz,¹³¹ A. Rinkevicius,¹³¹ L. Shchutska,¹³¹ M. Snowball,¹³¹ J. Yelton,¹³¹ M. Zakaria,¹³¹ S. Hewamanage,¹³² S. Linn,¹³² P. Markowitz,¹³² G. Martinez,¹³² J. L. Rodriguez,¹³² T. Adams,¹³³ A. Askew,¹³³ J. Bochenek,¹³³ B. Diamond,¹³³ J. Haas,¹³³ S. Hagopian,¹³³ V. Hagopian,¹³³ K. F. Johnson,¹³³ H. Prosper,¹³³ V. Veeraraghavan,¹³³ M. Weinberg,¹³³ M. M. Baarmand,¹³⁴ M. Hohlmann,¹³⁴ H. Kalakhety,¹³⁴ F. Yumiceva,¹³⁴ M. R. Adams,¹³⁵ L. Apanasevich,¹³⁵ V. E. Bazterra,¹³⁵ D. Berry,¹³⁵ R. R. Betts,¹³⁵ I. Bucinskaite,¹³⁵ R. Cavanaugh,¹³⁵ O. Evdokimov,¹³⁵ L. Gauthier,¹³⁵ C. E. Gerber,¹³⁵ D. J. Hofman,¹³⁵ S. Khalatyan,¹³⁵ P. Kurt,¹³⁵ D. H. Moon,¹³⁵ C. O'Brien,¹³⁵ C. Silkworth,¹³⁵ P. Turner,¹³⁵ N. Varelas,¹³⁵ E. A. Albayrak,^{136,vv} B. Bilki,^{136,zz} W. Clarida,¹³⁶ K. Dilsiz,¹³⁶ F. Duru,¹³⁶ M. Haytmyradov,¹³⁶ J.-P. Merlo,¹³⁶ H. Mermerkaya,^{136,aaa} A. Mestvirishvili,¹³⁶ A. Moeller,¹³⁶ J. Nachtman,¹³⁶ H. Ogul,¹³⁶ Y. Onel,¹³⁶ F. Ozok,^{136,vv} A. Penzo,¹³⁶ R. Rahmat,¹³⁶ S. Sen,¹³⁶ P. Tan,¹³⁶ E. Tiras,¹³⁶ J. Wetzel,¹³⁶ T. Yetkin,^{136,bbb} K. Yi,¹³⁶ B. A. Barnett,¹³⁷ B. Blumenfeld,¹³⁷ S. Bolognesi,¹³⁷ D. Fehling,¹³⁷ A. V. Gritsan,¹³⁷ P. Maksimovic,¹³⁷ C. Martin,¹³⁷ M. Swartz,¹³⁷ P. Baringer,¹³⁸ A. Bean,¹³⁸ G. Benelli,¹³⁸ C. Bruner,¹³⁸ J. Gray,¹³⁸ R. P. Kenny III,¹³⁸ M. Malek,¹³⁸ M. Murray,¹³⁸ D. Noonan,¹³⁸ S. Sanders,¹³⁸ J. Sekaric,¹³⁸ R. Stringer,¹³⁸ Q. Wang,¹³⁸ J. S. Wood,¹³⁸ A. F. Barfuss,¹³⁹ I. Chakaberia,¹³⁹ A. Ivanov,¹³⁹ S. Khalil,¹³⁹ M. Makouski,¹³⁹ Y. Maravin,¹³⁹ L. K. Saini,¹³⁹ S. Shrestha,¹³⁹ N. Skhirtladze,¹³⁹ I. Svintradze,¹³⁹ J. Gronberg,¹⁴⁰ D. Lange,¹⁴⁰ F. Rebassoo,¹⁴⁰ D. Wright,¹⁴⁰ A. Baden,¹⁴¹ A. Belloni,¹⁴¹ B. Calvert,¹⁴¹ S. C. Eno,¹⁴¹ J. A. Gomez,¹⁴¹ N. J. Hadley,¹⁴¹ R. G. Kellogg,¹⁴¹ T. Kolberg,¹⁴¹ Y. Lu,¹⁴¹ M. Marionneau,¹⁴¹ A. C. Mignerey,¹⁴¹ K. Pedro,¹⁴¹ A. Skuja,¹⁴¹ M. B. Tonjes,¹⁴¹ S. C. Tonwar,¹⁴¹ A. Apyan,¹⁴² R. Barbieri,¹⁴² G. Bauer,¹⁴² W. Busza,¹⁴² I. A. Cali,¹⁴² M. Chan,¹⁴² L. Di Matteo,¹⁴² V. Dutta,¹⁴² G. Gomez Ceballos,¹⁴² M. Goncharov,¹⁴² D. Gulhan,¹⁴² M. Klute,¹⁴² Y. S. Lai,¹⁴² Y.-J. Lee,¹⁴² A. Levin,¹⁴² P. D. Luckey,¹⁴² T. Ma,¹⁴² C. Paus,¹⁴² D. Ralph,¹⁴² C. Roland,¹⁴² G. Roland,¹⁴² G. S. F. Stephans,¹⁴² F. Stöckli,¹⁴² K. Sumorok,¹⁴² D. Velicanu,¹⁴² J. Veverka,¹⁴² B. Wyslouch,¹⁴² M. Yang,¹⁴² M. Zanetti,¹⁴² V. Zhukova,¹⁴² B. Dahmes,¹⁴³ A. Gude,¹⁴³ S. C. Kao,¹⁴³ K. Klapoetke,¹⁴³ Y. Kubota,¹⁴³ J. Mans,¹⁴³ N. Pastika,¹⁴³ R. Rusack,¹⁴³ A. Singovsky,¹⁴³ N. Tambe,¹⁴³ J. Turkewitz,¹⁴³ J. G. Acosta,¹⁴⁴ S. Oliveros,¹⁴⁴ E. Avdeeva,¹⁴⁵ K. Bloom,¹⁴⁵ S. Bose,¹⁴⁵ D. R. Claes,¹⁴⁵ A. Dominguez,¹⁴⁵ R. Gonzalez Suarez,¹⁴⁵ J. Keller,¹⁴⁵ D. Knowlton,¹⁴⁵ I. Kravchenko,¹⁴⁵ J. Lazo-Flores,¹⁴⁵ S. Malik,¹⁴⁵ F. Meier,¹⁴⁵ G. R. Snow,¹⁴⁵ J. Dolen,¹⁴⁶ A. Godshalk,¹⁴⁶ I. Iashvili,¹⁴⁶ A. Kharchilava,¹⁴⁶ A. Kumar,¹⁴⁶ S. Rappoccio,¹⁴⁶ G. Alverson,¹⁴⁷ E. Barberis,¹⁴⁷ D. Baumgartel,¹⁴⁷ M. Chasco,¹⁴⁷ J. Haley,¹⁴⁷ A. Massironi,¹⁴⁷ D. M. Morse,¹⁴⁷ D. Nash,¹⁴⁷ T. Orimoto,¹⁴⁷ D. Trocino,¹⁴⁷ R.-J. Wang,¹⁴⁷ D. Wood,¹⁴⁷ J. Zhang,¹⁴⁷ K. A. Hahn,¹⁴⁸ A. Kubik,¹⁴⁸ N. Mucia,¹⁴⁸ N. Odell,¹⁴⁸ B. Pollack,¹⁴⁸ A. Pozdnyakov,¹⁴⁸ M. Schmitt,¹⁴⁸ S. Stoynev,¹⁴⁸ K. Sung,¹⁴⁸ M. Velasco,¹⁴⁸ S. Won,¹⁴⁸ A. Brinkerhoff,¹⁴⁹ K. M. Chan,¹⁴⁹ A. Drozdetskiy,¹⁴⁹ M. Hildreth,¹⁴⁹ C. Jessop,¹⁴⁹ D. J. Karmgard,¹⁴⁹ N. Kellams,¹⁴⁹ K. Lannon,¹⁴⁹ W. Luo,¹⁴⁹ S. Lynch,¹⁴⁹ N. Marinelli,¹⁴⁹ T. Pearson,¹⁴⁹ M. Planer,¹⁴⁹ R. Ruchti,¹⁴⁹ N. Valls,¹⁴⁹ M. Wayne,¹⁴⁹ M. Wolf,¹⁴⁹ A. Woodard,¹⁴⁹ L. Antonelli,¹⁵⁰ J. Brinson,¹⁵⁰ B. Bylsma,¹⁵⁰ L. S. Durkin,¹⁵⁰ S. Flowers,¹⁵⁰ C. Hill,¹⁵⁰ R. Hughes,¹⁵⁰ K. Kotov,¹⁵⁰ T. Y. Ling,¹⁵⁰ D. Puigh,¹⁵⁰ M. Rodenburg,¹⁵⁰ G. Smith,¹⁵⁰ B. L. Winer,¹⁵⁰ H. Wolfe,¹⁵⁰ H. W. Wulsin,¹⁵⁰ O. Driga,¹⁵¹ P. Elmer,¹⁵¹ P. Hebda,¹⁵¹ A. Hunt,¹⁵¹ S. A. Koay,¹⁵¹ P. Lujan,¹⁵¹ D. Marlow,¹⁵¹ T. Medvedeva,¹⁵¹ M. Mooney,¹⁵¹ J. Olsen,¹⁵¹ P. Piroué,¹⁵¹ X. Quan,¹⁵¹ H. Saka,¹⁵¹ D. Stickland,^{151,c} C. Tully,¹⁵¹ J. S. Werner,¹⁵¹ S. C. Zenz,¹⁵¹ A. Zuranski,¹⁵¹ E. Brownson,¹⁵² H. Mendez,¹⁵² J. E. Ramirez Vargas,¹⁵² E. Alagoz,¹⁵³ V. E. Barnes,¹⁵³ D. Benedetti,¹⁵³ G. Bolla,¹⁵³ D. Bortoletto,¹⁵³ M. De Mattia,¹⁵³ Z. Hu,¹⁵³ M. K. Jha,¹⁵³ M. Jones,¹⁵³ K. Jung,¹⁵³ M. Kress,¹⁵³ N. Leonardo,¹⁵³ D. Lopes Pegna,¹⁵³ V. Maroussov,¹⁵³ P. Merkel,¹⁵³ D. H. Miller,¹⁵³ N. Neumeister,¹⁵³ B. C. Radburn-Smith,¹⁵³ X. Shi,¹⁵³ I. Shipsey,¹⁵³ D. Silvers,¹⁵³ A. Svyatkovskiy,¹⁵³ F. Wang,¹⁵³ W. Xie,¹⁵³ L. Xu,¹⁵³ H. D. Yoo,¹⁵³ J. Zablocki,¹⁵³ Y. Zheng,¹⁵³ N. Parashar,¹⁵⁴ J. Stupak,¹⁵⁴ A. Adair,¹⁵⁵ B. Akgun,¹⁵⁵ K. M. Ecklund,¹⁵⁵ F. J. M. Geurts,¹⁵⁵ W. Li,¹⁵⁵ B. Michlin,¹⁵⁵ B. P. Padley,¹⁵⁵ R. Redjimi,¹⁵⁵ J. Roberts,¹⁵⁵ J. Zabel,¹⁵⁵ B. Betchart,¹⁵⁶ A. Bodek,¹⁵⁶ R. Covarelli,¹⁵⁶ P. de Barbaro,¹⁵⁶ R. Demina,¹⁵⁶ Y. Eshaq,¹⁵⁶ T. Ferbel,¹⁵⁶ A. Garcia-Bellido,¹⁵⁶ P. Goldenzweig,¹⁵⁶ J. Han,¹⁵⁶ A. Harel,¹⁵⁶ A. Khukhunaishvili,¹⁵⁶ G. Petrillo,¹⁵⁶ D. Vishnevskiy,¹⁵⁶ R. Ciesielski,¹⁵⁷ L. Demortier,¹⁵⁷ K. Goulios,¹⁵⁷ G. Lungu,¹⁵⁷ C. Mesropian,¹⁵⁷ S. Arora,¹⁵⁸ A. Barker,¹⁵⁸ J. P. Chou,¹⁵⁸ C. Contreras-Campana,¹⁵⁸ E. Contreras-Campana,¹⁵⁸ N. Craig,¹⁵⁸

D. Duggan,¹⁵⁸ J. Evans,¹⁵⁸ D. Ferencek,¹⁵⁸ Y. Gershtein,¹⁵⁸ R. Gray,¹⁵⁸ E. Halkiadakis,¹⁵⁸ D. Hidas,¹⁵⁸ A. Lath,¹⁵⁸ S. Panwalkar,¹⁵⁸ M. Park,¹⁵⁸ R. Patel,¹⁵⁸ S. Salur,¹⁵⁸ S. Schnetzer,¹⁵⁸ S. Somalwar,¹⁵⁸ R. Stone,¹⁵⁸ S. Thomas,¹⁵⁸ P. Thomassen,¹⁵⁸ M. Walker,¹⁵⁸ K. Rose,¹⁵⁹ S. Spanier,¹⁵⁹ A. York,¹⁵⁹ O. Bouhali,^{160,ccc} R. Eusebi,¹⁶⁰ W. Flanagan,¹⁶⁰ J. Gilmore,¹⁶⁰ T. Kamon,^{160,ddd} V. Khotilovich,¹⁶⁰ V. Krutelyov,¹⁶⁰ R. Montalvo,¹⁶⁰ I. Osipenkov,¹⁶⁰ Y. Pakhotin,¹⁶⁰ A. Perloff,¹⁶⁰ J. Roe,¹⁶⁰ A. Rose,¹⁶⁰ A. Safonov,¹⁶⁰ T. Sakuma,¹⁶⁰ I. Suarez,¹⁶⁰ A. Tatarinov,¹⁶⁰ N. Akchurin,¹⁶¹ C. Cowden,¹⁶¹ J. Damgov,¹⁶¹ C. Dragoiu,¹⁶¹ P. R. Duderø,¹⁶¹ J. Faulkner,¹⁶¹ K. Kovitanggoon,¹⁶¹ S. Kunori,¹⁶¹ S. W. Lee,¹⁶¹ T. Libeiro,¹⁶¹ I. Volobouev,¹⁶¹ E. Appelt,¹⁶² A. G. Delannoy,¹⁶² S. Greene,¹⁶² A. Gurrola,¹⁶² W. Johns,¹⁶² C. Maguire,¹⁶² Y. Mao,¹⁶² A. Melo,¹⁶² M. Sharma,¹⁶² P. Sheldon,¹⁶² B. Snook,¹⁶² S. Tuo,¹⁶² J. Velkovska,¹⁶² M. W. Arenton,¹⁶³ S. Boutle,¹⁶³ B. Cox,¹⁶³ B. Francis,¹⁶³ J. Goodell,¹⁶³ R. Hirosky,¹⁶³ A. Ledovskoy,¹⁶³ H. Li,¹⁶³ C. Lin,¹⁶³ C. Neu,¹⁶³ J. Wood,¹⁶³ C. Clarke,¹⁶⁴ R. Harr,¹⁶⁴ P. E. Karchin,¹⁶⁴ C. Kottachchi Kankanamge Don,¹⁶⁴ P. Lamichhane,¹⁶⁴ J. Sturdy,¹⁶⁴ D. A. Belknap,¹⁶⁵ D. Carlsmith,¹⁶⁵ M. Cepeda,¹⁶⁵ S. Dasu,¹⁶⁵ L. Dodd,¹⁶⁵ S. Duric,¹⁶⁵ E. Friis,¹⁶⁵ R. Hall-Wilton,¹⁶⁵ M. Herndon,¹⁶⁵ A. Hervé,¹⁶⁵ P. Klabbers,¹⁶⁵ A. Lanaro,¹⁶⁵ C. Lazaridis,¹⁶⁵ A. Levine,¹⁶⁵ R. Loveless,¹⁶⁵ A. Mohapatra,¹⁶⁵ I. Ojalvo,¹⁶⁵ T. Perry,¹⁶⁵ G. A. Pierro,¹⁶⁵ G. Polese,¹⁶⁵ I. Ross,¹⁶⁵ T. Sarangi,¹⁶⁵ A. Savin,¹⁶⁵ W. H. Smith,¹⁶⁵ C. Vuosalo,¹⁶⁵ and N. Woods¹⁶⁵

(CMS Collaboration)

¹*Yerevan Physics Institute, Yerevan, Armenia*

²*Institut für Hochenergiephysik der OeAW, Wien, Austria*

³*National Centre for Particle and High Energy Physics, Minsk, Belarus*

⁴*Universiteit Antwerpen, Antwerpen, Belgium*

⁵*Vrije Universiteit Brussel, Brussel, Belgium*

⁶*Université Libre de Bruxelles, Bruxelles, Belgium*

⁷*Ghent University, Ghent, Belgium*

⁸*Université Catholique de Louvain, Louvain-la-Neuve, Belgium*

⁹*Université de Mons, Mons, Belgium*

¹⁰*Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

¹¹*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*

^{12a}*Universidade Estadual Paulista, São Paulo, Brazil*

^{12b}*Universidade Federal do ABC, São Paulo, Brazil*

¹³*Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria*

¹⁴*University of Sofia, Sofia, Bulgaria*

¹⁵*Institute of High Energy Physics, Beijing, China*

¹⁶*State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

¹⁷*Universidad de Los Andes, Bogota, Colombia*

¹⁸*University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia*

¹⁹*University of Split, Faculty of Science, Split, Croatia*

²⁰*Institute Rudjer Boskovic, Zagreb, Croatia*

²¹*University of Cyprus, Nicosia, Cyprus*

²²*Charles University, Prague, Czech Republic*

²³*Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt*

²⁴*National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*

²⁵*Department of Physics, University of Helsinki, Helsinki, Finland*

²⁶*Helsinki Institute of Physics, Helsinki, Finland*

²⁷*Lappeenranta University of Technology, Lappeenranta, Finland*

²⁸*DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France*

²⁹*Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France*

³⁰*Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg,*

Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France

³¹*Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France*

³²*Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3,*

Institut de Physique Nucléaire de Lyon, Villeurbanne, France

³³*Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia*

³⁴*RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany*

- ³⁵*RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
³⁶*RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany*
³⁷*Deutsches Elektronen-Synchrotron, Hamburg, Germany*
³⁸*University of Hamburg, Hamburg, Germany*
³⁹*Institut für Experimentelle Kernphysik, Karlsruhe, Germany*
⁴⁰*Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece*
⁴¹*University of Athens, Athens, Greece*
⁴²*University of Ioánnina, Ioánnina, Greece*
⁴³*Wigner Research Centre for Physics, Budapest, Hungary*
⁴⁴*Institute of Nuclear Research ATOMKI, Debrecen, Hungary*
⁴⁵*University of Debrecen, Debrecen, Hungary*
⁴⁶*National Institute of Science Education and Research, Bhubaneswar, India*
⁴⁷*Panjab University, Chandigarh, India*
⁴⁸*University of Delhi, Delhi, India*
⁴⁹*Saha Institute of Nuclear Physics, Kolkata, India*
⁵⁰*Bhabha Atomic Research Centre, Mumbai, India*
⁵¹*Tata Institute of Fundamental Research, Mumbai, India*
⁵²*Institute for Research in Fundamental Sciences (IPM), Tehran, Iran*
⁵³*University College Dublin, Dublin, Ireland*
^{54a}*INFN Sezione di Bari, Bari, Italy*
^{54b}*Università di Bari, Bari, Italy*
^{54c}*Politecnico di Bari, Bari, Italy*
^{55a}*INFN Sezione di Bologna, Bologna, Italy*
^{55b}*Università di Bologna, Bologna, Italy*
^{56a}*INFN Sezione di Catania, Catania, Italy*
^{56b}*Università di Catania, Catania, Italy*
^{56c}*CSFNSM, Catania, Italy*
^{57a}*INFN Sezione di Firenze, Firenze, Italy*
^{57b}*Università di Firenze, Firenze, Italy*
⁵⁸*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
^{59a}*INFN Sezione di Genova, Genova, Italy*
^{59b}*Università di Genova, Genova, Italy*
^{60a}*INFN Sezione di Milano-Bicocca, Milano, Italy*
^{60b}*Università di Milano-Bicocca, Milano, Italy*
^{61a}*INFN Sezione di Napoli, Napoli, Italy*
^{61b}*Università di Napoli 'Federico II', Napoli, Italy*
^{61c}*Università della Basilicata (Potenza), Napoli, Italy*
^{61d}*Università G. Marconi (Roma), Napoli, Italy*
^{62a}*INFN Sezione di Padova, Padova, Italy*
^{62b}*Università di Padova, Padova, Italy*
^{62c}*Università di Trento (Trento), Padova, Italy*
^{63a}*INFN Sezione di Pavia, Pavia, Italy*
^{63b}*Università di Pavia, Pavia, Italy*
^{64a}*INFN Sezione di Perugia, Perugia, Italy*
^{64b}*Università di Perugia, Perugia, Italy*
^{65a}*INFN Sezione di Pisa, Pisa, Italy*
^{65b}*Università di Pisa, Pisa, Italy*
^{65c}*Scuola Normale Superiore di Pisa, Pisa, Italy*
^{66a}*INFN Sezione di Roma, Roma, Italy*
^{66b}*Università di Roma, Roma, Italy*
^{67a}*INFN Sezione di Torino, Torino, Italy*
^{67b}*Università di Torino, Torino, Italy*
^{67c}*Università del Piemonte Orientale (Novara), Torino, Italy*
^{68a}*INFN Sezione di Trieste, Trieste, Italy*
^{68b}*Università di Trieste, Trieste, Italy*
⁶⁹*Kangwon National University, Chunchon, Korea*
⁷⁰*Kyungpook National University, Daegu, Korea*
⁷¹*Chonbuk National University, Jeonju, Korea*
⁷²*Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea*
⁷³*Korea University, Seoul, Korea*

- ⁷⁴University of Seoul, Seoul, Korea
⁷⁵Sungkyunkwan University, Suwon, Korea
⁷⁶Vilnius University, Vilnius, Lithuania
⁷⁷National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia
⁷⁸Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico
⁷⁹Universidad Iberoamericana, Mexico City, Mexico
⁸⁰Benemerita Universidad Autonoma de Puebla, Puebla, Mexico
⁸¹Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico
⁸²University of Auckland, Auckland, New Zealand
⁸³University of Canterbury, Christchurch, New Zealand
⁸⁴National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan
⁸⁵National Centre for Nuclear Research, Swierk, Poland
⁸⁶Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland
⁸⁷Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal
⁸⁸Joint Institute for Nuclear Research, Dubna, Russia
⁸⁹Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia
⁹⁰Institute for Nuclear Research, Moscow, Russia
⁹¹Institute for Theoretical and Experimental Physics, Moscow, Russia
⁹²P.N. Lebedev Physical Institute, Moscow, Russia
⁹³Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia
⁹⁴State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia
⁹⁵University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia
⁹⁶Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain
⁹⁷Universidad Autónoma de Madrid, Madrid, Spain
⁹⁸Universidad de Oviedo, Oviedo, Spain
⁹⁹Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain
¹⁰⁰CERN, European Organization for Nuclear Research, Geneva, Switzerland
¹⁰¹Paul Scherrer Institut, Villigen, Switzerland
¹⁰²Institute for Particle Physics, ETH Zurich, Zurich, Switzerland
¹⁰³Universität Zürich, Zurich, Switzerland
¹⁰⁴National Central University, Chung-Li, Taiwan
¹⁰⁵National Taiwan University (NTU), Taipei, Taiwan
¹⁰⁶Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand
¹⁰⁷Cukurova University, Adana, Turkey
¹⁰⁸Middle East Technical University, Physics Department, Ankara, Turkey
¹⁰⁹Bogazici University, Istanbul, Turkey
¹¹⁰Istanbul Technical University, Istanbul, Turkey
¹¹¹National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine
¹¹²University of Bristol, Bristol, United Kingdom
¹¹³Rutherford Appleton Laboratory, Didcot, United Kingdom
¹¹⁴Imperial College, London, United Kingdom
¹¹⁵Brunel University, Uxbridge, United Kingdom
¹¹⁶Baylor University, Waco, Texas 76798, USA
¹¹⁷The University of Alabama, Tuscaloosa, Alabama 35487, USA
¹¹⁸Boston University, Boston, Massachusetts 02215, USA
¹¹⁹Brown University, Providence, Rhode Island 02912, USA
¹²⁰University of California, Davis, Davis, California 95616, USA
¹²¹University of California, Los Angeles, California 90095, USA
¹²²University of California, Riverside, Riverside, California 92521, USA
¹²³University of California, San Diego, La Jolla, California 92093, USA
¹²⁴University of California, Santa Barbara, Santa Barbara, California 93106, USA
¹²⁵California Institute of Technology, Pasadena, California 91125, USA
¹²⁶Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA
¹²⁷University of Colorado at Boulder, Boulder, Colorado 80309, USA
¹²⁸Cornell University, Ithaca, New York 14853, USA
¹²⁹Fairfield University, Fairfield, Connecticut 06430, USA
¹³⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
¹³¹University of Florida, Gainesville, Florida 32611, USA
¹³²Florida International University, Miami, Florida 33199, USA
¹³³Florida State University, Tallahassee, Florida 32306, USA

- ¹³⁴Florida Institute of Technology, Melbourne, Florida 32901, USA
¹³⁵University of Illinois at Chicago (UIC), Chicago, Illinois 60607, USA
¹³⁶The University of Iowa, Iowa City, Iowa 52242, USA
¹³⁷Johns Hopkins University, Baltimore, Maryland 21218, USA
¹³⁸The University of Kansas, Lawrence, Kansas 66045, USA
¹³⁹Kansas State University, Manhattan, Kansas 66506, USA
¹⁴⁰Lawrence Livermore National Laboratory, Livermore, California 94551, USA
¹⁴¹University of Maryland, College Park, Maryland 20742, USA
¹⁴²Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
¹⁴³University of Minnesota, Minneapolis, Minnesota 55455, USA
¹⁴⁴University of Mississippi, Oxford, Mississippi 38677, USA
¹⁴⁵University of Nebraska-Lincoln, Lincoln, Nebraska 68588, USA
¹⁴⁶State University of New York at Buffalo, Buffalo, New York 14260, USA
¹⁴⁷Northeastern University, Boston, Massachusetts 02115, USA
¹⁴⁸Northwestern University, Evanston, Illinois 60208, USA
¹⁴⁹University of Notre Dame, Notre Dame, Indiana 46556, USA
¹⁵⁰The Ohio State University, Columbus, Ohio 43210, USA
¹⁵¹Princeton University, Princeton, New Jersey 08542, USA
¹⁵²University of Puerto Rico, Mayaguez, Puerto Rico 00681, USA
¹⁵³Purdue University, West Lafayette, Indiana 47907, USA
¹⁵⁴Purdue University Calumet, Hammond, Indiana 46323, USA
¹⁵⁵Rice University, Houston, Texas 77251, USA
¹⁵⁶University of Rochester, Rochester, New York 14627, USA
¹⁵⁷The Rockefeller University, New York, New York 10021, USA
¹⁵⁸Rutgers, The State University of New Jersey, Piscataway, New Jersey 08854, USA
¹⁵⁹University of Tennessee, Knoxville, Tennessee 37996, USA
¹⁶⁰Texas A&M University, College Station, Texas 77843, USA
¹⁶¹Texas Tech University, Lubbock, Texas 79409, USA
¹⁶²Vanderbilt University, Nashville, Tennessee 37235, USA
¹⁶³University of Virginia, Charlottesville, Virginia 22904, USA
¹⁶⁴Wayne State University, Detroit, Michigan 48202, USA
¹⁶⁵University of Wisconsin, Madison, Wisconsin 53706, USA

^aDeceased.

^bAlso at Vienna University of Technology, Vienna, Austria.

^cAlso at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

^dAlso at Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France.

^eAlso at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia.

^fAlso at Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia.

^gAlso at Universidade Estadual de Campinas, Campinas, Brazil.

^hAlso at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France.

ⁱAlso at Joint Institute for Nuclear Research, Dubna, Russia.

^jAlso at Suez University, Suez, Egypt.

^kAlso at Cairo University, Cairo, Egypt.

^lAlso at Fayoum University, El-Fayoum, Egypt.

^mAlso at British University in Egypt, Cairo, Egypt.

ⁿAlso at Ain Shams University, Cairo, Egypt.

^oAlso at Université de Haute Alsace, Mulhouse, France.

^pAlso at Brandenburg University of Technology, Cottbus, Germany.

^qAlso at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.

^rAlso at Eötvös Loránd University, Budapest, Hungary.

^sAlso at University of Debrecen, Debrecen, Hungary.

^tAlso at University of Visva-Bharati, Santiniketan, India.

^uAlso at King Abdulaziz University, Jeddah, Saudi Arabia.

^vAlso at University of Ruhuna, Matara, Sri Lanka.

^wAlso at Isfahan University of Technology, Isfahan, Iran.

^xAlso at Sharif University of Technology, Tehran, Iran.

^yAlso at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^zAlso at Università degli Studi di Siena, Siena, Italy.

- ^{aa} Also at Centre National de la Recherche Scientifique (CNRS) - IN2P3, Paris, France.
- ^{bb} Also at Purdue University, West Lafayette, USA.
- ^{cc} Also at Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico.
- ^{dd} Also at Institute for Nuclear Research, Moscow, Russia.
- ^{ee} Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia.
- ^{ff} Also at California Institute of Technology, Pasadena, USA.
- ^{gg} Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia.
- ^{hh} Also at Facoltà Ingegneria, Università di Roma, Roma, Italy.
- ⁱⁱ Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy.
- ^{jj} Also at University of Athens, Athens, Greece.
- ^{kk} Also at Paul Scherrer Institut, Villigen, Switzerland.
- ^{ll} Also at Institute for Theoretical and Experimental Physics, Moscow, Russia.
- ^{mm} Also at Albert Einstein Center for Fundamental Physics, Bern, Switzerland.
- ⁿⁿ Also at Gaziosmanpasa University, Tokat, Turkey.
- ^{oo} Also at Adiyaman University, Adiyaman, Turkey.
- ^{pp} Also at Cag University, Mersin, Turkey.
- ^{qq} Also at Mersin University, Mersin, Turkey.
- ^{rr} Also at Izmir Institute of Technology, Izmir, Turkey.
- ^{ss} Also at Ozyegin University, Istanbul, Turkey.
- ^{tt} Also at Marmara University, Istanbul, Turkey.
- ^{uu} Also at Kafkas University, Kars, Turkey.
- ^{vv} Also at Mimar Sinan University, Istanbul, Istanbul, Turkey.
- ^{ww} Also at Rutherford Appleton Laboratory, Didcot, United Kingdom.
- ^{xx} Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- ^{yy} Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia.
- ^{zz} Also at Argonne National Laboratory, Argonne, USA.
- ^{aaa} Also at Erzincan University, Erzincan, Turkey.
- ^{bbb} Also at Yildiz Technical University, Istanbul, Turkey.
- ^{ccc} Also at Texas A&M University at Qatar, Doha, Qatar.
- ^{ddd} Also at Kyungpook National University, Daegu, Korea.