



1 **Contrasting watershed-scale trends in runoff and sediment**  
2 **yield complicate rangeland water resources planning**

3

4 **M. D. Berg**<sup>1,\*</sup>, **F. Marcantonio**<sup>2</sup>, **M. A. Allison**<sup>3,4</sup>, **J. McAlister**<sup>5</sup>, **B. P. Wilcox**<sup>1</sup>, and  
5 **W. E. Fox**<sup>1,5</sup>

6 [1]{Texas A&M University Department of Ecosystem Science and Management, College  
7 Station, Texas}

8 [2]{Texas A&M University Department of Geology and Geophysics, College Station, Texas}

9 [3]{The Water Institute of the Gulf, Baton Rouge, Louisiana}

10 [4]{Tulane University Department of Earth and Environmental Sciences, New Orleans,  
11 Louisiana}

12 [5]{Texas A&M AgriLife Blackland Research & Extension Center, Temple, Texas}

13 [\*]{now at: Save Water Co, Houston, Texas}

14 Correspondence to: M. D. Berg (mbergtamu@gmail.com)

15

16 **Abstract**

17 Rangelands cover a large portion of the earth's land surface and are undergoing dramatic  
18 landscape changes. At the same time, these ecosystems face increasing expectations to meet  
19 growing water supply needs. To address major gaps in our understanding of rangeland  
20 hydrologic function, we investigated historical watershed-scale runoff and sediment yield in a  
21 dynamic landscape in central Texas, USA. We quantified the relationship between  
22 precipitation and runoff and analyzed reservoir sediment cores dated using Cesium-137 and  
23 Lead-210 radioisotopes. Local rainfall and streamflow showed no directional trend over a  
24 period of 85 years, resulting in a rainfall-runoff ratio that has been resilient to watershed  
25 changes. Reservoir sedimentation rates generally were higher before 1963, but have been  
26 much lower and very stable since that time. Our findings suggest that (1) rangeland water  
27 yields may be stable over long periods despite dramatic landscape changes while (2) these  
28 same landscape changes influence sediment yields that impact downstream reservoir storage.



1 Relying on rangelands to meet water needs demands an understanding of how these dynamic  
2 landscapes function and a quantification of the physical processes at work.

3

#### 4 **1 Introduction**

5 Diverse rangeland ecosystems falling along a grassland–forest continuum cover roughly half  
6 of the earth’s land surface (Breshears, 2006). Generally precipitation-limited, they are  
7 typically used for livestock grazing and harvesting of woody products rather than crop  
8 production. But rangelands worldwide face numerous challenges, including (1) conversion to  
9 urban development or cultivation; (2) shifting plant cover, such as encroachment by woody  
10 plants and invasion by non-native species; and (3) demands for increased production without  
11 sacrificing sustainability (Tilman et al., 2002; Van Auken, 2000; Wilcox et al., 2012b).

12 As growing populations look to these dynamic landscapes to provide critical ecosystem  
13 services—including water supply and water storage—their ability to keep pace with these  
14 demands is uncertain (Havstad et al., 2007; Jackson et al., 2001). Some of this uncertainty is  
15 due to the tremendous variability of runoff and erosion through time and space, which can  
16 vary by orders of magnitude even between portions of a single small field (Gaspar et al.,  
17 2013; Ritchie et al., 2005). Landscape changes affect these processes further still; and water  
18 and sediment yields depend on interactions between climate, vegetation, and local geology.  
19 These complex interactions make predictions difficult; and the influence of human activity  
20 adds yet another compounding layer of difficulty (Peel, 2009; Boardman, 2006; Vorosmarty  
21 and Sahagian, 2000). As a result, major gaps remain in our understanding of rangeland  
22 ecosystems. Further interdisciplinary study is imperative to develop a coherent picture of the  
23 linkages between hydrological, ecological, and geological processes (Newman, 2006; Wilcox  
24 and Thurow, 2006).

25 Some rangeland investigations have focused on the potential of these landscapes to provide  
26 augmented water yields or storage in small reservoirs. Economic and modeling studies have  
27 identified vegetation management as a possible means of increasing runoff and streamflow  
28 (Griffin and McCarl, 1989; Afinowicz et al., 2005), and government agencies have  
29 incorporated these goals into their programs (Texas State Soil and Water Conservation Board,  
30 2005; USDA-NRCS, 2006). Other concerns center on sediment yield, which threatens  
31 downstream surface water storage (Bennett et al., 2002; Dunbar et al., 2010). To determine



1 how to respond to these issues and whether related investments are worthwhile, we must gain  
2 a better understanding of how rangeland systems function with respect to water resources.

3 To date, most research has been based on extrapolation of findings from relatively small-scale  
4 studies to larger scales or on modeled results. However, because runoff and sediment  
5 production are scale-dependent processes, such extrapolation is often unreliable (de Vente and  
6 Poesen, 2005; Wilcox et al., 2003). Since they more accurately reveal the true water and  
7 sediment yields of watersheds, studies of these processes conducted at the catchment scale are  
8 much more relevant to water planning efforts. But whereas catchment-scale data on  
9 precipitation and streamflow are somewhat widely available, corresponding sediment data are  
10 lacking. Since they serve as archives of historical watershed conditions, the use of reservoir  
11 sediments provides one means of filling this data gap and of investigating the impact of  
12 human activity (Edwards and Whittington, 2001; Winter et al., 2001). Linking the findings of  
13 such investigations with observed changes at the watershed scale will greatly facilitate the  
14 development of effective strategies for managing rangeland water resources.

15 In this study, we investigated the hydrological and sediment transport dynamics of rangeland  
16 watersheds. Our main objectives were to (1) quantify long-term trends in precipitation and  
17 streamflow using historical data; (2) estimate historical sedimentation rates through  
18 radioisotope analysis of reservoir sediment cores; and (3) explore the potential effects of  
19 drought conditions on sediment production with historical data. Addressing these objectives  
20 not only improves our understanding of rangeland processes but also provides much-needed  
21 information on the potential of these landscapes to provide for growing global water needs.

22

## 23 **2 Methods**

### 24 **2.1 Study area**

25 As part of a broader study of landscape change and ecosystem function, we examined  
26 rangeland processes in the Lampasas Cut Plain of central Texas, USA. This savanna  
27 landscape is characterized by low buttes and mesas separated by broad, flat valleys. Local  
28 prevailing geology is Cretaceous limestone; soils are loamy and clayey, with occasional sandy  
29 loams, and are susceptible to sheet and gully erosion (Allison, 1991; Clower, 1980). The area  
30 is drained by the Lampasas River. Streamflow in the upper reaches of the river is runoff-  
31 dominated, with localized contributions from springflow (Prein et al., 2013), and has been



1 recorded at two primary stations (Figure 1). Annual precipitation averages approximately 800  
2 mm, decreasing to the north and west (Figure 2).

3 For the sediment study, we examined eight flood-control reservoirs and their watersheds  
4 within the Lampasas River basin. Reservoirs L1, L2, L3, L4, L9, and LX are located in  
5 Lampasas County and were constructed between 1958 and 1961. Before impoundment, the  
6 parallel watersheds of L1, L2, and L3, contributed to the downstream watershed of LX.  
7 Reservoirs M1 and M4, in Mills County, were completed in 1974. Basic attributes of the  
8 reservoirs and their watersheds are compiled in Table 1.

9 Current local land use is predominantly rangeland, and livestock numbers have fluctuated  
10 over the last several decades (Figure 3a) while remaining among the highest in the region  
11 (Wilcox et al., 2012a). Cropland was widespread early in the 20<sup>th</sup> century (Figure 3b) but had  
12 declined by nearly 80% by 2012 (Berg, M. D., manuscript in review, 2015). Amid this  
13 shifting land use, the area has been characterized by large fluctuations in the extent of woody  
14 plant cover, due to brush management and regrowth (Figure 3c), and a dramatic increase in  
15 the density of farm ponds (Figure 3d) over the last several decades (Berg et al., 2015a).

## 16 **2.2 Rainfall and runoff trends**

17 To investigate local hydrological trends, we analyzed historical precipitation and streamflow  
18 data for the Lampasas River basin. We created a composite record of annual precipitation  
19 using a Thiessen polygon approach, centering polygons on available NWS stations (Figure 2).  
20 Streamflow data were derived from the two USGS stream gage stations downstream from the  
21 study watersheds. The lower Youngsport station, with a drainage area of 3,212 km<sup>2</sup>, operated  
22 between 1924 and 1980; the Kempner station, with a drainage area of 2,119 km<sup>2</sup> has remained  
23 active from 1963 to the present.

24 We performed an automated baseflow separation of streamflow data from each station  
25 (Arnold and Allen, 1999). This digital filter approach is objective and reproducible and  
26 partitions annual baseflow and stormflow with high efficiency (Arnold et al., 1995)—  
27 enabling these components to be interpreted in light of changing landscape conditions.

28 Using the precipitation and two streamflow datasets (1924—1980; 1963—2010), we applied a  
29 nonparametric Mann-Kendall trend test to detect directional changes (Lettenmaier et al.,  
30 1994). We performed two-tailed statistical tests for significance, with  $\alpha = 0.10$ .



### 1 **2.3 Reservoir sedimentation rates**

2 To shed light on sediment transport processes, we extracted cores from each of the eight  
3 reservoirs and analyzed sediments using Cesium-137 ( $^{137}\text{Cs}$ ) and Lead-210 ( $^{210}\text{Pb}$ ) tracers.  
4  $^{137}\text{Cs}$  is present in the environment as a result of atomic weapons testing and accidental  
5 emissions.  $^{210}\text{Pb}$  occurs naturally. Both can be used to estimate sedimentation rates and  
6 interpret transport history in a variety of environments (Walling et al., 2003; Ritchie and  
7 McHenry, 1990; Appleby and Oldfield, 1978). Coring sites were selected by locating the  
8 thickest sediment deposits through exploratory hydroacoustic surveys (U.S. Army Corps of  
9 Engineers, 2013, 1989; Dunbar et al., 2002). In each reservoir, we extracted sediment cores at  
10 identified sites near the dam structure, from locations corresponding to the pre-impoundment  
11 floodplain (Figure 4). Taking cores from these areas reduces the likelihood of capturing  
12 mixed profiles, which skew analysis (Sanchez-Cabeza and Ruiz-Fernández, 2012). It also  
13 ensures the collection of fine sediments, to which the radioisotopes preferentially adsorb  
14 (Bennett et al., 2002). We extracted cores using a portable vibracoring system suspended from  
15 a floating platform. This method captures unconsolidated, saturated sediments with minimal  
16 disturbance and compaction (Lanesky et al., 1979). The cores were collected with an  
17 aluminum pipe lowered to the point of refusal, penetrating the pre-impoundment surface.  
18 Retrieved cores were sealed and transported upright to cold storage ( $\sim 5^\circ\text{C}$ ).

19 We sectioned each core vertically in 3-cm intervals, drying each section for analysis  
20 according to IAEA (2003) protocols. A subsample of each core section was ground to  
21 homogenize its contents, sealed in a 50 mm x 9 mm Petri dish, and allowed to ingrow for at  
22 least 21 days so that  $^{210}\text{Pb}$  supported levels reached equilibrium. Counts for  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$   
23 were performed according to Hanna et al. (2014) using a Canberra low-energy germanium  
24 gamma spectrometer. Radioisotope activity was indicated by photopeaks at 46 keV (total  
25  $^{210}\text{Pb}$ ) and 661.6 keV ( $^{137}\text{Cs}$ ). Excess  $^{210}\text{Pb}$  was calculated by subtracting the supported  
26 activity of the  $^{226}\text{Ra}$  parent—obtained by averaging the 295, 351.9, and 609.3 keV peaks of  
27 the  $^{214}\text{Pb}$  and  $^{214}\text{Bi}$  daughter products—from total measured  $^{210}\text{Pb}$  activity at the 46 keV peak.  
28 Activity measurements were validated with IAEA-300 standard reference material.

29 To determine historical linear sedimentation rates, we used as a chronological marker the  
30 depth of peak  $^{137}\text{Cs}$  activity (corresponding to the 1963 peak in global atmospheric fallout)  
31 (Ritchie et al., 1973). We calculated average linear sedimentation rates for the post-1963  
32 period by dividing this depth by the time elapsed between 1963 and the coring date for each



1 reservoir; we calculated the pre-1963 rates by dividing the depth of sediment below the  
2 activity peak by the time elapsed between reservoir impoundment and 1963.

3 To complement  $^{137}\text{Cs}$  analysis, we used excess  $^{210}\text{Pb}$  activities to calculate the linear  
4 sedimentation rate for each core (Krishnaswamy et al., 1971; Bierman et al., 1998). We also  
5 searched for changing deposition rates within each core, as plots of the natural log of excess  
6  $^{210}\text{Pb}$  versus depth indicate stable sedimentation rates over time when  $R^2$  approaches 1.0.

7 Finally, we obtained historical annual Palmer Modified Drought Index (PMDI) data for the  
8 region to identify potential climatic drivers of sedimentation during different periods. We  
9 plotted PMDI and annual peak flows (from USGS data) between 1924 and 2010, identifying  
10 episodes conducive to increased sediment production (in particular, a wet year or years  
11 following a period of intense drought).

12

### 13 **3 Results**

#### 14 **3.1 Rainfall and runoff trends**

15 Despite a great deal of interannual variability, there was no directional change in local  
16 precipitation 1924–1980 ( $p = 0.90$ ) or 1963–2010 ( $p = 0.22$ ), which has remained near a  
17 long-term average of 800 mm (Figure 5a). The same is true of total streamflow (1924–1980:  
18  $p = 0.98$ , 1963–2010:  $p = 0.34$ ), which has averaged between 60 and 70 mm (Figure 5b). As  
19 a result, the rainfall–runoff ratio also remained unchanged, at approximately 8% (1924–  
20 1980:  $p = 0.90$ , 1963–2010:  $p = 0.45$ ). Moreover, neither baseflow nor stormflow exhibited a  
21 directional change over either period of record. However, baseflow as a proportion of total  
22 streamflow did increase 1924–1980 ( $p = 0.02$ ) despite minimal change in overall flow—  
23 almost doubling its contribution (Figure 5c).

#### 24 **3.2 Reservoir sedimentation rates**

25 Sediment core profiles varied widely in depth between reservoirs—from less than 3 cm in LX  
26 to 162 cm in L1 (Figure 6). Activity peaks of  $^{137}\text{Cs}$  supported the analysis of pre-1963  
27 sedimentation rates for reservoirs L1, L2, L3, and L9. Overall, linear sedimentation rates were  
28 higher before 1963 (Table 2; Figure 7). Except in the case of L3, sediment deposition has  
29 slowed since 1963—by 54% in L1, 76% in L2, and 84% in L9. In reservoir L3, it increased



1 by 49% after 1963. Reservoir L1 exhibited the highest sedimentation rate both before and  
2 after 1963. However, when normalized by catchment area, sedimentation rates varied much  
3 more widely. That in L9 was by far the highest—surpassing the next highest reservoir by  
4 nearly 1400% for the pre-1963 period and by 423% for the post-1963 period.

5 Cores from L4, LX, M1, and M4 did not display a  $^{137}\text{Cs}$  peak. For these cores, sedimentation  
6 was assumed to be post-1963 and was estimated by dividing sediment depth by time since  
7 impoundment. For cores L4 and M4, which did not capture the entire sediment profile, actual  
8 rates likely are higher than those calculated.

9 Cores from reservoirs LX and M1 showed vertical mixing that prohibited  $^{210}\text{Pb}$  analysis.  
10 However, remaining cores displayed high correlation between  $^{210}\text{Pb}$  activities and depth,  
11 indicating linear sedimentation rates have remained quite stable over time (Table 2).  $^{210}\text{Pb}$ -  
12 based estimates generally resembled those based on  $^{137}\text{Cs}$  activities. In addition, rates  
13 calculated from  $^{210}\text{Pb}$  activities were similar to the post-1963 rates based on  $^{137}\text{Cs}$  activities ( $p$   
14 = 0.84), suggesting good agreement between the two methods for the period since 1963.

15 Chronological data revealed periods of drought of varying intensity and occasional years of  
16 very high streamflow (Figure 8). The historic 1950s drought was longer and more severe than  
17 any other over the last century; it was followed by periods of very high flow in 1957 and  
18 1960. Comparable high flows in 1965 occurred in the middle of a multi-year drought, and the  
19 severe drought beginning in 2006 featured occasional elevated peak flows. In 1992, very high  
20 flows occurred during a prolonged wet period.

21

## 22 **4 Discussion**

### 23 **4.1 Rainfall and runoff trends**

24 Given the varying trends in precipitation and streamflow observed in many regions (Lins and  
25 Slack, 1999; Andreadis and Lettenmaier, 2006), the dynamic hydrological stability in our  
26 study area is surprising. At the same time, such consistency sheds light on the effects of  
27 watershed changes on local water budgets. Studies at small spatial scales frequently indicate  
28 that landscape changes have important water resource impacts, with the specific response  
29 depending on the relative importance of evapotranspiration, recharge, and runoff (Foley et al.,  
30 2005; Kim and Jackson, 2012). Such changes affect local water budgets and influence water



1 yields (Petersen and Stringham, 2008; Huxman et al., 2005; Farley et al., 2005). However,  
2 complicated feedbacks make effects at larger scales highly uncertain and often overwhelmed  
3 by climatic and physical characteristics (Peel, 2009; Wilcox et al., 2006; Kuhn et al., 2007).  
4 Our rainfall–runoff ratio of 8% is essentially identical to early estimates of 7% for the area  
5 (Tanner, 1937). The lack of a directional trend in streamflows suggests that this region, like  
6 many semiarid landscapes dominated by surface runoff, is largely hydrologically insensitive  
7 to shifting watershed characteristics (Wilcox, 2002). Changes in land use and land cover—  
8 and even the impoundment of small reservoirs—have had negligible impacts on streamflow.

9 It is still not understood why baseflow showed a proportional increase 1924—1980. In some  
10 landscapes, improving range conditions have led to increased infiltration (Wilcox and Huang,  
11 2010). However, local livestock numbers have remained high, and karst features are limited—  
12 unlike other regions where baseflow increases have been attributed to rangeland recovery. It  
13 is possible that infiltration from local impoundments has added to baseflows. Despite minimal  
14 effects on total streamflow, even small dams can create localized groundwater recharge (Graf,  
15 1999; Smith et al., 2002), and Lampasas River tributaries are characterized by a high degree of  
16 connectivity between surface water and local aquifers (Mills and Rawson, 1965).

17 Perennial flow in this part of the Lampasas River is maintained by isolated springs fed by an  
18 aquifer extending beyond the basin (Mills and Rawson, 1965). As a result, the effective  
19 catchment of the river is larger than it appears, and springflow contributions complicate the  
20 interpretation of streamflows. At the same time, it is clear that the fundamental relationship  
21 between rainfall and streamflow has not changed over more than 85 years—suggesting that  
22 the Lampasas River is hydrologically resilient in the face of changing land use and land cover.

#### 23 **4.2 Reservoir sedimentation rates**

24 Because sediment deposition affects reservoir storage and flood detention, understanding  
25 sedimentation rates over time is critical to managing rangeland water resources. Though  
26 questions do remain regarding the opposing trend in reservoir L3, changes in rates make it  
27 clear that sedimentation was more rapid before 1963. The period since that time has been  
28 characterized by stable and lower yields. But what explains the higher rates seen during the  
29 earlier period? Additional historical landscape data may offer a key interpretive lens.

30 Livestock can be powerful instruments of landscape change, both directly (trampling soils)  
31 and indirectly (disturbing protective vegetation). When grazing is prolonged or intense,





1 sediment yield can be great (Trimble and Mendel, 1995). The high animal densities in this  
2 area around the time of reservoir impoundment doubtless contributed to erosion (Figure 3a).

3 Crop production also can result in accelerated erosion by damaging soil structure and  
4 depleting organic matter (Quine et al., 1999). Cropland is a major source of sediment in many  
5 landscapes (Foster and Lees, 1999; Blake et al., 2012). In our study area, cropland acreage has  
6 declined dramatically since the 1930s (Figure 3b). Further, nationwide improvements in soil  
7 conservation have reduced sediment yield from many agricultural lands (Knox, 2001).

8 While woody plant encroachment influences soil loss, removing undesirable shrubs and trees  
9 also elevates short-term sediment yields (Porto et al., 2009). Since the time of initial  
10 settlement, woody plant management has resulted in major land cover changes (Figure 3c).  
11 Most early removal was done manually, and the first mechanical control methods were very  
12 destructive, leading to high erosion rates (Hamilton and Hanselka, 2004). In recent decades,  
13 however, brush removal has declined with shifting landowner priorities (Sorice et al., 2014).

14 Changes in precipitation frequency, duration, or intensity also affect sediment transport (Xie  
15 et al., 2002; Allen et al., 2011). Similarly, drought is an important driver of sediment dynamics  
16 in many rangelands. Extended dry periods can cause long-term shifts in plant cover, leading  
17 to sediment pulses when rains return (Allen and Breshears, 1998; Nearing et al., 2007). The  
18 Lampasas River experienced very high flows in 1957, 1960, 1965, and 1992, and some of  
19 these were associated in time with severe droughts (Figure 8). Just before the impoundment of  
20 most of the reservoirs we examined, the region was in the grip of drought conditions  
21 unmatched since European settlement (Bradley and Malstaff, 2004). Our sediment records  
22 cover only the end of this drought but show pre-1963 deposition 220–630% faster than  
23 subsequent rates. However, any direct effects of the 1957 drought-breaking floods would not  
24 be found in the sediments of the reservoirs, which were impounded beginning in 1958.  
25 Interestingly, we also did not find spikes in sedimentation associated with high flows or  
26 droughts later in the study period. The apparent low importance of drought and floods in  
27 sediment delivery in these watersheds is surprising.

28 Together, these factors have acted over multiple temporal and spatial scales to influence  
29 sediment yields in the study area. Yet because there is no clear link between contemporary  
30 land use, land cover, and sedimentation rates, it is possible that another process has reduced  
31 sediment yields.



### 1 **4.3 Sediment storage**

2 To truly understand the local sediment processes at work, it is important to understand what  
3 our findings actually show. Sedimentation rates are poor indicators of in-field soil erosion and  
4 redistribution (Nearing et al., 2000; Ritchie et al., 2009); what they do reflect is more closely  
5 related to net watershed sediment yield. Sediment yield is buffered by internal storage.  
6 Especially at larger scales, watersheds can have a great deal of internal storage, so that very  
7 little eroded soil actually leaves the watershed, even in the presence of extreme erosion  
8 (Bennett et al., 2005; Porto et al., 2011).

9 In this study area, the increasing density of farm ponds (Figure 3d) represents a key potential  
10 sink for watershed sediments. These ponds retain material that otherwise would be  
11 transported downstream, reducing sediment yields. Because of their smaller contributing  
12 watersheds, ponds have high trap efficiencies, magnifying their effects (Brainard and  
13 Fairchild, 2012). Indeed, impoundments may be the single greatest anthropogenic modifier of  
14 sediment transport; globally, most sedimentation now takes place in aquatic settings and will  
15 be retained therein for long periods (Renwick et al., 2005; Verstraeten et al., 2006).

16 In addition to this storage of eroded sediments in local ponds, a vast amount of sediment from  
17 past erosion likely remains on the landscape (Beach, 1994; Meade, 1982). The initial decades  
18 after European settlement in this area saw intensive cultivation and very high livestock  
19 densities (Jordan-Bychkov et al., 1984; Wilcox et al., 2012a). This destructive combination  
20 remained in place for nearly a century in the Lampasas Cut Plain. By the 1930s, many  
21 rangelands were already seriously degraded (Mitchell, 2000; Bentley, 1898; Box, 1967). While  
22 the methods we used do not allow us to determine whether reservoir sediments result from  
23 contemporary erosion or are a legacy of earlier land use, stabilizing sediment yields and  
24 observations of local gully erosion suggest that deposits from prior erosion continue to be a  
25 source of sediment (Bartley et al., 2007; Mukundan et al., 2011; Phillips, 2003).

26 The lack of sediments in LX appears to lend support to the importance of internal deposits.  
27 This reservoir's watershed is comparable in size to those of L2, L3, and M4, yet  
28 sedimentation rates were only 3%–14% of those in the other reservoirs. When L1, L2, and L3  
29 were impounded, the effective catchment area of LX decreased by 86%. Without the  
30 historical streamflows and sediment loads from those tributaries, deposits are no longer  
31 mobilized and transported downstream.



1 Given this complexity, we suggest that radioisotope tracers have great potential to elucidate  
2 the dynamics of rangeland systems, particularly as their use evolves from primarily research  
3 applications to use as a management and decision-support tool (Mukundan et al., 2012).  
4 Further strides can be made in understanding rangeland processes by (1) incorporating  
5 historical climate, land use, and land cover information to interpret sediment data (Venteris et  
6 al., 2004; Boardman, 2006) and (2) including sediment surveys of the farm ponds that are  
7 much smaller yet far more abundant than the reservoirs we examined (Downing et al., 2006).

8

## 9 **5 Conclusion**

10 We examined long-term trends in rainfall, runoff, and sediment yield in rangeland watersheds  
11 with a dynamic land use history. Over more than 85 years, neither precipitation nor  
12 streamflow showed any directional trend, suggesting a lack of hydrological sensitivity to  
13 landscape change. This raises doubts over efforts to increase runoff by directing land cover  
14 changes. Reservoir sedimentation rates generally were higher before 1963, and then stabilized  
15 at a lower level over the 50 years since 1963. We believe that this decline in sediment yield is  
16 related to long-term landscape changes and an increase in internal storage. As a result, future  
17 changes in land use or sediment storage may impact downstream reservoir capacity. These  
18 findings challenge simplistic assumptions about streamflow and sediment yield in dynamic  
19 rangelands. Determining the role of these landscapes in meeting growing water resource  
20 demands requires a creative approach. Integrating multiple techniques with historical  
21 information enables a more complete understanding of rangeland processes and holds the key  
22 to informed water planning.

23

## 24 **Data availability**

25 Streamflow data are available at the USGS National Water Information System. Stream  
26 gages: 08103800 (Kempner) and 08104000 (Youngsport). Drought data are available at the  
27 NOAA National Climate Data Center. Texas Climate Division: CD 3 (North Central) and CD  
28 6 (Edwards Plateau).

29

## 30 **Acknowledgements**



1 Dan Duncan, Andrea Hanna, and Diana di Leonardo performed activity counts of sediment  
2 samples. This work was supported by USDA-NIFA Managed Ecosystems grant 2011-68002-  
3 30015, USDA-NIFA National Needs Program grant 2009-38420-05631, NSF-CNH grant  
4 413900, and a Tom Slick Graduate Research Fellowship from the Texas A&M University  
5 College of Agriculture and Life Sciences.

6



## 1 **References**

- 2 Afinowicz, J. D., Munster, C. L., and Wilcox, B. P.: Modeling effects of brush management  
3 on the rangeland water budget: Edwards Plateau, Texas, *J. Am. Water Resour. Assoc.*, 41,  
4 181-193, 10.1111/j.1752-1688.2005.tb03727.x, 2005.
- 5 Allen, C. D., and Breshears, D. D.: Drought-induced shift of a forest-woodland ecotone:  
6 Rapid landscape response to climate variation, *Proc. Natl. Acad. Sci. U. S. A.*, 95, 14839-  
7 14842, 10.1073/pnas.95.25.14839, 1998.
- 8 Allen, P. M., Harmel, R. D., Dunbar, J. A., and Arnold, J. G.: Upland contribution of  
9 sediment and runoff during extreme drought: A study of the 1947-1956 drought in the  
10 Blackland Prairie, Texas, *Journal of Hydrology*, 407, 1-11, 10.1016/j.jhydrol.2011.04.039,  
11 2011.
- 12 Allison, J. E.: Soil Survey of Lampasas County, Texas, in, US Department of Agriculture Soil  
13 Conservation Service, Washington, D.C., 154, 1991.
- 14 Andreadis, K. M., and Lettenmaier, D. P.: Trends in 20th century drought over the continental  
15 United States, *Geophys. Res. Lett.*, 33, L10403, 10.1029/2006gl025711, 2006.
- 16 Appleby, P. G., and Oldfield, F.: The calculation of lead-210 dates assuming a constant rate  
17 of supply of unsupported 210Pb to the sediment, *Catena*, 5, 1-8, 10.1016/s0341-  
18 8162(78)80002-2, 1978.
- 19 Arnold, J. G., Allen, P. M., Muttiyah, R., and Bernhardt, G.: Automated base flow separation  
20 and recession analysis techniques, *Groundwater*, 33, 1010, 10.1111/j.1745-  
21 6584.1995.tb00046.x, 1995.
- 22 Arnold, J. G., and Allen, P. M.: Automated methods for estimating baseflow and ground  
23 water recharge from streamflow records, *J. Am. Water Resour. Assoc.*, 35, 411-424,  
24 10.1111/j.1752-1688.1999.tb03599.x, 1999.
- 25 Bartley, R., Hawdon, A., Post, D. A., and Roth, C. H.: A sediment budget for a grazed semi-  
26 arid catchment in the Burdekin basin, Australia, *Geomorphology*, 87, 302-321,  
27 10.1016/j.geomorph.2006.10.001, 2007.
- 28 Beach, T.: The fate of eroded soil - sediment sinks and sediment budgets of agrarian  
29 landscapes in southern Minnesota, 1851-1988, *Ann. Assoc. Am. Geogr.*, 84, 5-28,  
30 10.1111/j.1467-8306.1994.tb01726.x, 1994.
- 31 Bennett, S. J., Cooper, C. M., Ritchie, J. C., Dunbar, J. A., Allen, P. M., Caldwell, L. W., and  
32 McGee, T. M.: Assessing sedimentation issues within aging flood control reservoirs in  
33 Oklahoma, *J. Am. Water Resour. Assoc.*, 38, 1307-1322, 2002.



- 1 Bennett, S. J., Rhoton, F. E., and Dunbar, J. A.: Texture, spatial distribution, and rate of  
2 reservoir sedimentation within a highly erosive, cultivated watershed: Grenada Lake,  
3 Mississippi, *Water Resour. Res.*, 41, 10.1029/2004wr003645, 2005.
- 4 Bentley, H. L.: Cattle ranges of the Southwest: a history of the exhaustion of the pasturage  
5 and suggestions for its restoration, in, edited by: United States Department of Agriculture  
6 Government Printing Office, Washington, D.C., 1898.
- 7 Berg, M. D., Popescu, S. C., Wilcox, B. P., Angerer, J. P., Rhodes, E. C., McAlister, J., and  
8 Fox, W. E.: Small farm ponds: overlooked features with important impacts on watershed  
9 sediment transport, *JAWRA Journal of the American Water Resources Association*, n/a-n/a,  
10 10.1111/1752-1688.12369, 2015a.
- 11 Berg, M. D., Sorice, M. G., Wilcox, B. P., Angerer, J. P., Rhodes, E. C., and Fox, W. E.:  
12 Demographic Changes Drive Woody Plant Cover Trends—An Example from the Great  
13 Plains, *Rangel. Ecol. Manag.*, 68, 315-321, 10.1016/j.rama.2015.05.004, 2015b.
- 14 Bierman, P. R., Albrecht, A., Bothner, M. H., Brown, E. T., Bullen, T. D., Gray, L. B., and  
15 Turpin, L.: Erosion, weathering, and sedimentation, in: *Isotope Tracers in Catchment  
16 Hydrology*, edited by: Kendall, C., and McDonnell, J. J., Elsevier, Amsterdam, 647-678,  
17 1998.
- 18 Blake, W. H., Ficken, K. J., Taylor, P., Russell, M. A., and Walling, D. E.: Tracing crop-  
19 specific sediment sources in agricultural catchments, *Geomorphology*, 139, 322-329,  
20 10.1016/j.geomorph.2011.10.036, 2012.
- 21 Boardman, J.: Soil erosion science: Reflections on the limitations of current approaches,  
22 *Catena*, 68, 73-86, 10.1016/j.catena.2006.03.007, 2006.
- 23 Box, T. W.: Range deterioration in West Texas, *The Southwestern Historical Quarterly*, 71,  
24 37-45, 10.2307/30237942, 1967.
- 25 Bradley, R. G., and Malstaff, G.: Dry periods and drought events of the Edwards Plateau,  
26 Texas, Austin, Texas, 201-210, 2004.
- 27 Brainard, A. S., and Fairchild, G. W.: Sediment characteristics and accumulation rates in  
28 constructed ponds, *J. Soil Water Conserv.*, 67, 425-432, 10.2489/jswc.67.5.425, 2012.
- 29 Breshears, D. D.: The grassland-forest continuum: trends in ecosystem properties for woody  
30 plant mosaics?, *Front. Ecol. Environ.*, 4, 96-104, 10.1890/1540-  
31 9295(2006)004[0096:tgctie]2.0.co;2, 2006.
- 32 Clower, D. F.: Soil Survey of Brown and Mills Counties, Texas, in, US Department of  
33 Agriculture Soil Conservation Service, Washington, D.C., 181, 1980.



- 1 de Vente, J., and Poesen, J.: Predicting soil erosion and sediment yield at the basin scale:  
2 Scale issues and semi-quantitative models, *Earth-Science Reviews*, 71, 95-125,  
3 10.1016/j.earscirev.2005.02.002, 2005.
- 4 Downing, J. A., Prairie, Y. T., Cole, J. J., Duarte, C. M., Tranvik, L. J., Striegl, R. G.,  
5 McDowell, W. H., Kortelainen, P., Caraco, N. F., Melack, J. M., and Middelburg, J. J.: The  
6 global abundance and size distribution of lakes, ponds, and impoundments, *Limnol.*  
7 *Oceanogr.*, 51, 2388-2397, 10.2307/3841076, 2006.
- 8 Dunbar, J. A., Higley, P. D., and Bennett, S. J.: Acoustic Imaging of Sediment Impounded  
9 Within USDA-NRCS Flood Control Dams, Wisconsin, US Department of Agriculture,  
10 Agricultural Research Service, National Sedimentation Laboratory, Channel and Watershed  
11 Processes Research Unit, Oxford, Mississippi Research Report No. 30, 2002.
- 12 Dunbar, J. A., Allen, P. M., and Bennett, S. J.: Effect of multiyear drought on upland  
13 sediment yield and subsequent impacts on flood control reservoir storage, *Water Resour. Res.*,  
14 46, W05526, 10.1029/2008wr007519, 2010.
- 15 Edwards, K. J., and Whittington, G.: Lake sediments, erosion and landscape change during  
16 the Holocene in Britain and Ireland, *Catena*, 42, 143-173, 10.1016/s0341-8162(00)00136-3,  
17 2001.
- 18 Farley, K. A., Jobbágy, E. G., and Jackson, R. B.: Effects of afforestation on water yield: a  
19 global synthesis with implications for policy, *Glob. Change Biol.*, 11, 1565-1576,  
20 10.1111/j.1365-2486.2005.01011.x, 2005.
- 21 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S.,  
22 Coe, M. T., Daily, G. C., Gibbs, H. K., Helkowski, J. H., Holloway, T., Howard, E. A.,  
23 Kucharik, C. J., Monfreda, C., Patz, J. A., Prentice, I. C., Ramankutty, N., and Snyder, P. K.:  
24 Global Consequences of Land Use, *Science*, 309, 570-574, 10.1126/science.1111772, 2005.
- 25 Foster, I. D. L., and Lees, J. A.: Changing headwater suspended sediment yields in the LOIS  
26 catchments over the last century: a paleolimnological approach, *Hydrol. Process.*, 13, 1137-  
27 1153, 10.1002/(sici)1099-1085(199905)13:7<1137::aid-hyp794>3.3.co;2-d, 1999.
- 28 Gaspar, L., Navas, A., Walling, D. E., Machin, J., and Arozamena, J. G.: Using Cs-137 and  
29 Pb-210(ex) assess soil redistribution on slopes at different temporal scales, *Catena*, 102, 46-  
30 54, 10.1016/j.catena.2011.01.004, 2013.
- 31 Graf, W. L.: Dam nation: A geographic census of American dams and their large-scale  
32 hydrologic impacts, *Water Resour. Res.*, 35, 1305-1311, 10.1029/1999wr900016, 1999.



- 1 Griffin, R. C., and McCarl, B. A.: Brushland management for increased water yield in Texas,
- 2 J. Am. Water Resour. Assoc., 25, 175-186, 10.1111/j.1752-1688.1989.tb05679.x, 1989.
- 3 Hamilton, W. T., and Hanselka, C. W.: Mechanical practices prior to 1975, in: Brush
- 4 Management, edited by: Hamilton, W. T., McGinty, A., Ueckert, D. N., Hanselka, C. W., and
- 5 Lee, M. R., Texas A&M University Press, College Station, Texas, 17-32, 2004.
- 6 Hanna, A. J. M., Allison, M. A., Bianchi, T. S., Marcantonio, F., and Goff, J. A.: Late
- 7 Holocene sedimentation in a high Arctic coastal setting: Simpson Lagoon and Colville Delta,
- 8 Alaska, Continental Shelf Research, 74, 11-24, 10.1016/j.csr.2013.11.026, 2014.
- 9 Havstad, K. M., Peters, D. P. C., Skaggs, R., Brown, J., Bestelmeyer, B., Fredrickson, E.,
- 10 Herrick, J., and Wright, J.: Ecological services to and from rangelands of the United States,
- 11 Ecol. Econ., 64, 261-268, 10.1016/j.ecolecon.2007.08.005, 2007.
- 12 Huxman, T. E., Wilcox, B. P., Breshears, D. D., Scott, R. L., Snyder, K. A., Small, E. A.,
- 13 Hultine, K., Pockman, W. T., and Jackson, R. B.: Ecohydrological implications of woody
- 14 plant encroachment, Ecology, 86, 308-319, 10.1890/03-0583, 2005.
- 15 IAEA: Collection and preparation of bottom sediment samples for analysis of radionuclides
- 16 and trace elements in, edited by: Section, N. a. H.-R. E. S., IAEA, Vienna, Austria, 2003.
- 17 Jackson, R. B., Carpenter, S. R., Dahm, C. N., McKnight, D. M., Naiman, R. J., Postel, S. L.,
- 18 and Running, S. W.: Water in a changing world, Ecol. Appl., 11, 1027-1045,
- 19 10.2307/3061010, 2001.
- 20 Jordan-Bychkov, T. G., Bean, J. L., and Holmes, W. M.: Texas, a Geography, Geographies of
- 21 the United States, Westview Press, Boulder, Colorado, 288 pp., 1984.
- 22 Kim, J. H., and Jackson, R. B.: A Global Analysis of Groundwater Recharge for Vegetation,
- 23 Climate, and Soils, Vadose Zone Journal, 11, 10.2136/vzj2011.0021RA, 2012.
- 24 Knox, J. C.: Agricultural influence on landscape sensitivity in the Upper Mississippi River
- 25 Valley, Catena, 42, 193-224, 10.1016/s0341-8162(00)00138-7, 2001.
- 26 Krishnaswamy, S., Lal, D., Martin, J. M., and Meybeck, M.: Geochronology of lake
- 27 sediments, Earth and Planetary Science Letters, 11, 407-414, 10.1016/0012-821X(71)90202-
- 28 0, 1971.
- 29 Kuhn, T. J., Tate, K. W., Cao, D., and George, M. R.: Juniper removal may not increase
- 30 overall Klamath River Basin water yields, California Agriculture, 61, 166-171, 2007.
- 31 Lanesky, D. E., Logan, B. W., Brown, R. G., and Hine, A. C.: A new approach to portable
- 32 vibracoring underwater and on land, Journal of Sedimentary Research, 49, 654-657, 1979.





- 1 Lettenmaier, D. P., Wood, E. F., and Wallis, J. R.: Hydro-Climatological Trends in the
- 2 Continental United States, 1948-88, *J. Clim.*, 7, 586-607, 10.1175/1520-
- 3 0442(1994)007<0586:hctitc>2.0.co;2, 1994.
- 4 Lins, H. F., and Slack, J. R.: Streamflow trends in the United States, *Geophys. Res. Lett.*, 26,
- 5 227-230, 10.1029/1998gl900291, 1999.
- 6 Meade, R. H.: Sources, Sinks, and Storage of River Sediment in the Atlantic Drainage of the
- 7 United States, *The Journal of Geology*, 90, 235-252, 10.2307/30066396, 1982.
- 8 Mills, W. B., and Rawson, J.: Base Flow Studies: Lampasas River, Texas, in, edited by: U.S.
- 9 Geological Survey in Cooperation with the Texas Water Commission, Texas Water
- 10 Commission, Austin, Texas, 1965.
- 11 Mitchell, J. E.: Rangeland resource trends in the United States, in: General Technical Report
- 12 RMRS-GTR-68, edited by: Agriculture, U. S. D. o., USDA, Forest Service, Rocky Mountain
- 13 Research Station, Fort Collins, Colorado, 2000.
- 14 Mukundan, R., Radcliffe, D. E., and Ritchie, J. C.: Channel stability and sediment source
- 15 assessment in streams draining a Piedmont watershed in Georgia, USA, *Hydrol. Process.*, 25,
- 16 1243-1253, 10.1002/hyp.7890, 2011.
- 17 Mukundan, R., Walling, D. E., Gellis, A. C., Slattery, M. C., and Radcliffe, D. E.: Sediment
- 18 source fingerprinting: Transforming from a research tool to a management tool, *J. Am. Water*
- 19 *Resour. Assoc.*, 48, 1241-1257, 10.1111/j.1752-1688.2012.00685.x, 2012.
- 20 Nearing, M. A., Romkens, M. J. M., Norton, L. D., Stott, D. E., Rhoton, F. E., Laflen, J. M.,
- 21 Flanagan, D. C., Alonso, C. V., Binger, R. L., Dabney, S. M., Doering, O. C., Huang, C. H.,
- 22 McGregor, K. C., and Simon, A.: Measurements and models of soil loss rates, *Science*, 290,
- 23 1300-1301, 10.1126/science.290.5495.1300b, 2000.
- 24 Nearing, M. A., Nichols, M. H., Stone, J. J., Renard, K. G., and Simanton, J. R.: Sediment
- 25 yields from unit-source semiarid watersheds at Walnut Gulch, *Water Resour. Res.*, 43,
- 26 W06426, 10.1029/2006wr005692, 2007.
- 27 Newman, B., Wilcox, B., Archer, S., Breshears, D., Dahm, C., Duffy, C., McDowell, N.,
- 28 Phillips, F., Scanlon, B., and E. Vivoni: Ecohydrology of water-limited environments: A
- 29 scientific vision, *Water Resour. Res.*, 42, W06302, 10.1029/2005WR004141, 2006.
- 30 Peel, M. C.: Hydrology: catchment vegetation and runoff, *Prog. Phys. Geogr.*, 33, 837-844,
- 31 10.1177/0309133309350122, 2009.



- 1 Petersen, S. L., and Stringham, T. K.: Infiltration, Runoff, and Sediment Yield in Response to
- 2 Western Juniper Encroachment in Southeast Oregon, *Rangel. Ecol. Manag.*, 61, 74-81,
- 3 10.2111/07-070R.1, 2008.
- 4 Phillips, J.: Alluvial storage and the long-term stability of sediment yields, *Basin Research*,
- 5 15, 153-163, 10.1046/j.1365-2117.2003.00204.x, 2003.
- 6 Porto, P., Walling, D. E., and Callegari, G.: Investigating the effects of afforestation on soil
- 7 erosion and sediment mobilisation in two small catchments in Southern Italy, *Catena*, 79, 181-
- 8 188, 10.1016/j.catena.2009.01.007, 2009.
- 9 Porto, P., Walling, D. E., and Callegari, G.: Using Cs-137 measurements to establish
- 10 catchment sediment budgets and explore scale effects, *Hydrol. Process.*, 25, 886-900,
- 11 10.1002/hyp.7874, 2011.
- 12 Prein, L. J., Srinivasan, R., and Casebolt, P.: Lampasas River Watershed Protection Plan, The
- 13 Lampasas River Watershed Partnership, 207, 2013.
- 14 Quine, T. A., Walling, D. E., Chakela, Q. K., Mandiringana, O. T., and Zhang, X.: Rates and
- 15 patterns of tillage and water erosion on terraces and contour strips: evidence from caesium-
- 16 137 measurements, *Catena*, 36, 115-142, 10.1016/s0341-8162(99)00006-5, 1999.
- 17 Renwick, W. H., Smith, S. V., Bartley, J. D., and Buddemeier, R. W.: The role of
- 18 impoundments in the sediment budget of the conterminous United States, *Geomorphology*,
- 19 71, 99-111, doi:10.1016/j.geomorph.2004.01.010, 2005.
- 20 Ritchie, J., and McHenry, J. R.: Application of Radioactive Fallout Cesium-137 for
- 21 Measuring Soil Erosion and Sediment Accumulation Rates and Patterns: A Review, *J.*
- 22 *Environ. Qual.*, 19, 215-233, 10.2134/jeq1990.00472425001900020006x, 1990.
- 23 Ritchie, J. C., McHenry, J. R., and Gill, A. C.: Dating Recent Reservoir Sediments, *Limnol.*
- 24 *Oceanogr.*, 18, 254-263, 10.4319/lo.1973.18.2.0254, 1973.
- 25 Ritchie, J. C., Nearing, M. A., Nichols, M. H., and Ritchie, C. A.: Patterns of soil erosion and
- 26 redeposition on Lucky Hills Watershed, Walnut Gulch experimental watershed, Arizona,
- 27 *Catena*, 61, 122-130, 10.1016/j.catena.2005.03.012, 2005.
- 28 Ritchie, J. C., Nearing, M. A., and Rhoton, F. E.: Sediment budgets and source determinations
- 29 using fallout Cesium-137 in a semiarid rangeland watershed, Arizona, USA, *Journal of*
- 30 *Environmental Radioactivity*, 100, 637-643, 10.1016/j.jenvrad.2009.05.008, 2009.
- 31 Sanchez-Cabeza, J. A., and Ruiz-Fernández, A. C.: <sup>210</sup>Pb sediment radiochronology: An
- 32 integrated formulation and classification of dating models, *Geochimica et Cosmochimica*
- 33 *Acta*, 82, 183-200, 10.1016/j.gca.2010.12.024, 2012.



- 1 Smith, S. V., Renwick, W. H., Bartley, J. D., and Buddemeier, R. W.: Distribution and  
2 significance of small, artificial water bodies across the United States landscape, *The Science*  
3 of *The Total Environment*, 299, 21-36, 10.1016/s0048-9697(02)00222-x, 2002.
- 4 Sorice, M. G., Kreuter, U. P., Wilcox, B. P., and Fox III, W. E.: Changing landowners,  
5 changing ecosystem? Land-ownership motivations as drivers of land management practices,  
6 *Journal of Environmental Management*, 133, 144-152, 10.1016/j.jenvman.2013.11.029, 2014.
- 7 Tanner, B. M.: A study of the feasibility of water power development at proposed flood  
8 control dams on the Leon and Lampasas Rivers, Master of Science, Civil Engineering,  
9 Agricultural and Mechanical College of Texas, College Station, Texas, 57 pp., 1937.
- 10 Texas State Soil and Water Conservation Board: State Brush Control Plan, in, Texas State  
11 Soil and Water Conservation Board, Temple, Texas, 53, 2005.
- 12 Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R., and Polasky, S.: Agricultural  
13 sustainability and intensive production practices, *Nature*, 418, 671-677, 10.1038/nature01014,  
14 2002.
- 15 Trimble, S. W., and Mendel, A. C.: The cow as a geomorphic agent - a critical review,  
16 *Geomorphology*, 13, 233-253, 10.1016/0169-555x(95)00028-4, 1995.
- 17 U.S. Army Corps of Engineers: Sedimentation Investigations of Rivers and Reservoirs, in,  
18 *Engineering and Design*, U.S. Army Corps of Engineers, Washington, D.C., 1989.
- 19 U.S. Army Corps of Engineers: Hydrographic Surveying, in, *Engineering and Design*, U.S.  
20 Army Corps of Engineers, Washington, D.C., 2013.
- 21 USDA-NRCS: Conservation practice standard: brush management practice code 314, in,  
22 edited by: USDA-NRCS, US Department of Agriculture, NRCS, Washington, DC, 3, 2006.
- 23 Van Auken, O. W.: Shrub invasions of North American semiarid grasslands, *Annu. Rev.*  
24 *Ecol. Syst.*, 31, 197-215, 10.1146/annurev.ecolsys.31.1.197, 2000.
- 25 Venteris, E. R., McCarty, G. W., Ritchie, J. C., and Gish, T.: Influence of management  
26 history and landscape variables on soil organic carbon and soil redistribution, *Soil Sci.*, 169,  
27 787-795, 10.1097/01.ss.0000148742.75369.55, 2004.
- 28 Verstraeten, G., Bazzoffi, P., Lajczak, A., Rădoane, M., Rey, F., Poesen, J., and de Vente, J.:  
29 Reservoir and pond sedimentation in Europe, in: *Soil Erosion in Europe*, John Wiley & Sons,  
30 Ltd, 757-774, 2006.
- 31 Vorosmarty, C. J., and Sahagian, D.: Anthropogenic disturbance of the terrestrial water cycle,  
32 *Bioscience*, 50, 753-765, 10.1641/0006-3568(2000)050[0753:adottw]2.0.co;2, 2000.



- 1 Walling, D. E., Collins, A. L., and Sichingabula, H. M.: Using unsupported lead-210
- 2 measurements to investigate soil erosion and sediment delivery in a small Zambian
- 3 catchment, *Geomorphology*, 52, 193-213, 10.1016/s0169-555x(02)00244-1, 2003.
- 4 Wilcox, B. P.: Shrub control and streamflow on rangelands: a process based viewpoint, *J.*
- 5 *Range Manage.*, 55, 318-326, 2002.
- 6 Wilcox, B. P., Breshears, D. D., and Allen, C. D.: Ecohydrology of a resource-conserving
- 7 semiarid woodland: effects of scale and disturbance, *Ecol. Monogr.*, 73, 223-239,
- 8 10.1890/0012-9615(2003)073[0223:EOARSW]2.0.CO;2, 2003.
- 9 Wilcox, B. P., Dowhower, S. L., Teague, W. R., and Thurow, T. L.: Long-Term Water
- 10 Balance in a Semiarid Shrubland, *Rangel. Ecol. Manag.*, 59, 600-606, 10.2111/06-014R3.1,
- 11 2006.
- 12 Wilcox, B. P., and Thurow, T. L.: Emerging issues in rangeland ecohydrology - Preface,
- 13 *Hydrol. Process.*, 20, 3155-3157, 10.1002/hyp.6324, 2006.
- 14 Wilcox, B. P., and Huang, Y.: Woody plant encroachment paradox: Rivers rebound as
- 15 degraded grasslands convert to woodlands, *Geophys. Res. Lett.*, 37, L07402,
- 16 10.1029/2009gl041929, 2010.
- 17 Wilcox, B. P., Sorice, M. G., Angerer, J., and Wright, C. L.: Historical changes in stocking
- 18 densities on Texas rangelands, *Rangel. Ecol. Manag.*, 65, 313-317, 10.2111/rem-d-11-
- 19 00119.1, 2012a.
- 20 Wilcox, B. P., Turnbull, L., Young, M. H., Williams, C. J., Ravi, S., Seyfried, M. S.,
- 21 Bowling, D. R., Scott, R. L., Germino, M. J., Caldwell, T. G., and Wainwright, J.: Invasion of
- 22 shrublands by exotic grasses: ecohydrological consequences in cold versus warm deserts,
- 23 *Ecohydrology*, 5, 160-173, 10.1002/eco.247, 2012b.
- 24 Winter, L. T., Foster, I. D. L., Charlesworth, S. M., and Lees, J. A.: Floodplain lakes as sinks
- 25 for sediment-associated contaminants - a new source of proxy hydrological data?, *Sci. Total*
- 26 *Environ.*, 266, 187-194, 10.1016/s0048-9697(00)00745-2, 2001.
- 27 Xie, Y., Liu, B., and Nearing, M. A.: Practical thresholds for separating erosive and non-
- 28 erosive storms, *Trans. ASAE*, 45, 1843-1847, 2002.
- 29



1 Table 1. Sediment study reservoirs and watershed characteristics.

Reservoir	Primary Inflow	Surface Area (km <sup>2</sup> )	Watershed Area (km <sup>2</sup> )	Year Impounded	Year Cored	Min. Elev. (m)	Max. Elev. (m)
L1	Donalson Creek	0.20	50.9	1959	2010	367	500
L2	Pitt Creek	0.18	23.2	1959	2010	362	458
L3	Espy Branch	0.11	27.5	1958	2010	355	459
L4	Pillar Bluff Creek	0.07	41.2	1960	2012	345	467
L9	Cemetery Creek	0.02	1.2	1960	2012	322	363
LX	Bean Creek	0.20	23.1	1961	2012	338	420
M1	Middle Bennett Creek	0.14	34.6	1974	2012	422	536
M4	Mustang Creek	0.15	28.0	1974	2012	432	534

2



1 Table 2. Linear sedimentation rates derived from radioisotope activities.


Core	<sup>137</sup> Cs				<sup>210</sup> Pb		R <sup>2</sup> ln dpm g <sup>-1</sup>
	Pre-1963		Post-1963		Core mean		
	cm y <sup>-1</sup>	cm y <sup>-1</sup> km <sup>-2</sup>	cm y <sup>-1</sup>	cm y <sup>-1</sup> km <sup>-2</sup>	cm y <sup>-1</sup>	cm y <sup>-1</sup> km <sup>-2</sup>	
L1	6.4	0.13	2.9	0.06	3.1	0.06	0.90
L2	3.4	0.15	0.8	0.03	0.9	0.04	0.97
L3	1.4	0.05	2.1	0.08	1.3	0.04	0.96
L4	<sup>a</sup>	<sup>a</sup>	0.5 <sup>b</sup>	0.01 <sup>b</sup>	1.2	0.03	0.93
L9	2.5	2.02	0.4	0.32	0.4	0.19	0.94
LX	<sup>a</sup>	<sup>a</sup>	0.1	< 0.01	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>
M1	<sup>a</sup>	<sup>a</sup>	1.5	0.04	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>
M4	<sup>a</sup>	<sup>a</sup>	0.4 <sup>b</sup>	0.01 <sup>b</sup>	0.8	0.01	1.00

2 <sup>a</sup>Core did not display a <sup>137</sup>Cs peak, and rates were calculated using the time elapsed since  
 3 impoundment.

4 <sup>b</sup>Core did not capture the pre-impoundment surface and likely underestimates true values.

5 <sup>c</sup>Core showed significant vertical mixing, preventing calculation of sedimentation rate.


6



1

2 Figure 1. Study area in Texas, USA. Each study watershed encloses a flood control reservoir  
3 from which sediment cores were collected. All watersheds contribute flow to the Lampasas  
4 River.


5



1

2 Figure 2. Average annual precipitation gradient and location of National Weather Service  
3 (NWS) stations used to construct historical precipitation record.






1

2 Figure 3. Historical landscape changes in the study area. (a) Livestock numbers in the  
3 Lampasas Cut Plain. Recreated from Wilcox et al. (2012a). (b) Extent of active cropland in  
4 1937-40 and 2012 (Berg, M. D., manuscript in review, 2015). (c) Historical extent of woody  
5 plant cover in the study watersheds (Berg et al., 2015b). (d) Pond density over time in the  
6 study watersheds (Berg et al., 2015a).




1  
2 Figure 4. Reservoir sediment coring apparatus (top) and representative sediment profile  
3 (bottom).  
4




1

2 Figure 5. Precipitation and streamflow trends of the Lampasas River basin. (a) Precipitation  
3 showed no directional trend. (b) Streamflow showed no directional trend at either the  
4 Youngsport (Y) or Kempner (K) station, despite being highly variable. (c) Baseflow as a  
5 proportion of total streamflow displayed an upward trend over the first portion of the study  
6 period.

7




1  
 2 Figure 6. Sediment core profiles of bulk density and radioisotope activities from the eight  
 3 reservoirs. Solid horizontal lines indicate the pre-impoundment surface (no line indicates the  
 4 core did not capture the pre-impoundment surface). Dashed lines in <sup>137</sup>Cs graphs represent the  
 5 depth of peak activity. The <sup>210</sup>Pb profile for L3 is from a second core collected at the same  
 6 location.



1

2 Figure 6 (continued). Sediment core profiles of bulk density and radioisotope activities from  
 3 the eight reservoirs. Solid horizontal lines indicate the pre-impoundment surface (no line  
 4 indicates the core did not capture the pre-impoundment surface). Dashed lines in <sup>137</sup>Cs graphs  
 5 represent the depth of peak activity.


6



1

2 Figure 7. Linear sedimentation rates derived from <sup>137</sup>Cs activities. Summary comparison of  
3 pre-1963 and post-1963 rates.

4



1

2 Figure 8. Chronology of regional drought (annual Palmer Modified Drought Index) and peak

3 flows on the Lampasas River.