


PUBLICATIONS

1998

**FORAGE-LIVESTOCK
FIELD DAY REPORT - 1998**

**TEXAS A&M UNIVERSITY AGRICULTURAL
RESEARCH and EXTENSION CENTER
at OVERTON**

**Texas Agricultural Experiment Station
Texas Agricultural Extension Service**



April 16, 1998

Research Center Technical Report 98-1

All programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark or a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural Experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

SOIL NITRATE LEVELS IN BERMUDAGRASS PASTURES UNDER LONG-TERM STOCKING RATES AND FERTILITY REGIMENS

F. M. Rouquette, Jr., J. L. Kerby, G. H. Nimr, I. J. Pemberton, G. R. Smith, and V. A. Haby

Background. Grazing of 'Coastal' and common bermudagrass research pastures at TAMU-Overton began in 1969. Related Field Day reports on pH, P, and K detail treatment of pastures during the past 28 years. Nitrogen applications from 1985-1989 were split at 6 to 8-week intervals from Dec. to Sept. From 1990-1996, N was split-applied from Dec. through May only during the ryegrass growing season with no N during the exclusive bermudagrass growing season (Table 1). Soil nitrate-N was determined by conventional soil analysis methodology. The objective of this experiment was to quantify soil nitrate-N levels after 28 years of continuous N fertilizer application, which included 12 years (1985-1996) of no-N fertilization to bermudagrass pastures.

Table 1. Fertilization of bermudagrass pastures during 28-year period, 1969 through 1996.

Date	Annual Fertilization Rate (N-P ₂ O ₅ -K ₂ O)	
	Ryegrass + N	Clover + K
1969 through 1984 ¹	200-100-100	200-100-100
1985-1989 ²	408-0-0	0-0-114
1990-1994 ³	238-0-0	0-0-121
1995-1996 ^{3,4}	289-0-0	0-0-108

¹Pastures not divided into fertility treatments from 1969 through 1984; therefore, all pastures were fertilized identically and were overseeded with both ryegrass and clover.

²Initiation of N vs no-N treatments (1985). N applied at 6 to 8-week intervals from Dec. to Sept.

³N fertilization only during ryegrass growing period, Dec. to May, with none on bermudagrass.

⁴Used rye + ryegrass during 1995 and 1996 with N fertilization from Nov. to May.

Research Findings. Soil nitrate-N levels, across all three stocking rates, at the initiation of the fertility regimens (1985) were low in the 0-6" depth, but with higher levels for common bermudagrass than for Coastal (Table 2). At both 0-6" and 6-18" depths, nitrate-N levels were predictably higher in N-fertilized pastures. Soil tests in 1989, following 5 years of 408 lbs N/ac per year showed higher levels of soil nitrate-N. Similar trends were noted for 1985 (Table 2). Lower soil nitrate-N in Coastal bermudagrass pastures was likely due to differences in forage uptake of N and forage production since Coastal is likely to produce nearly twice as much total dry matter as common bermudagrass.

Soil nitrate-N analysis in 1996 after 6 years of applying fertilizer N only during the ryegrass growing period showed reduced levels compared to 1985 and 1989. Data (Table 3) show differences between fertility levels at each soil depth (0-60"). Differences in soil nitrate-N between

bermudagrasses primarily occurred in the 0 to 18" depths. There was essentially no effect of stocking rate on soil nitrate-N except in the 0-6" soil depth. Even in this top soil layer, stocking rate did not have a major role in influencing soil nitrate-N levels when soil was sampled about 3 months following the last N fertilization. Nitrate-N rapidly declined with soil depth and stabilized at 3 to 4 ppm from 18 to 60" soil depth.

Application. Sandy soils and improved bermudagrass pastures in East Texas have sufficient forage production to prevent significant accumulations of soil nitrate-N under fertilization rates used in this study. Without N-fertilization, contributions from clovers via excreta-based nutrient cycling have been adequate to maintain stands of bermudagrass, and offers fertilization management alternatives for producers.

Table 2. Soil nitrate-nitrogen at two soil depths in bermudagrass fertilized with N (RYG + N) or K (CLV + K) at initiation of fertility x stocking rate regimens.

Item	Nitrate-N (ppm) at Soil Depths			
	0-6"		6"-18"	
	RYG + N	CLV + K	RYG + N	CLV + K
1985 ¹				
Coastal	14.8 b	2.8 b	4.4 a	2.3 a
Common	22.7 a	5.5 a	3.6 a	1.7 a
1989 ²				
Coastal	24.8 b	4.9 b	22.0 a	2.3 a
Common	33.2 a	10.9 a	25.1 a	5.0 a

¹Numbers with different letters: Bermudagrasses differ (P<.05); at 0-6", N vs K (P<.0001); at 6-18", N vs K (P<.04).

²Numbers with different letters: Bermudagrasses differ (P<.004); at both depths, R vs K (P<.0001).

Table 3. Soil nitrate-nitrogen at various soil depths in bermudagrass pastures after 28 years of N application which included 12 years of fertility x stocking rate treatments.

1996	Nitrate-N (ppm) at Soil Depths				
	0-6"	6-18"	18-36"	36-48"	48-60"
<u>Bermudagrass</u>					
Coastal	16.4 a ¹	10.7 a	4.7 a	4.0 a	3.8 a
Common	3.5 b	3.0 b	2.7 a	1.5 b	2.8 a
<u>Fertility</u>					
Clover + K	6.0 b	3.5 b	1.0 b	0.9 b	1.2 b
Ryegrass + N	13.9 a	10.1 a	6.3 a	4.6 a	5.4 a
<u>Stocking Rate</u>					
High	9.1 ab	6.9 a	2.3 a	2.0 a	3.1 a
Medium	6.3 b	8.0 a	4.1 a	3.3 a	3.2 a
Low	14.5 a	5.6 a	4.7 a	2.9 a	3.6 a

¹Numbers within a soil depth and major treatment category followed by a different letter, differ (P<.01).