

# **PUBLICATIONS**

## **1993**

# **FIELD DAY REPORT - 1993**

## **Texas A&M University Agricultural Research and Extension Center at Overton**

**Texas Agricultural Experiment Station  
Texas Agricultural Extension Service**

**Overton, Texas**

**May 28, 1993**

**Research Center Technical Report 93-1**

---

All Programs and information of the Texas Agricultural Experiment Station and Texas Agricultural Extension Service are available to everyone without regard to race, color, religion, sex, age, or national origin.

Mention of trademark of a proprietary product does not constitute a guarantee or a warranty of the product by the Texas Agricultural Experiment Station or Texas Agricultural Extension Service and does not imply its approval to the exclusion of other products that also may be suitable.

---

## EFFECT OF AMMONIUM:NITRATE RATIOS ON SWEETCORN YIELD AND SOIL NITROGEN CONCENTRATIONS

V. A. Haby, F. M. Hons, J. V. Davis, and A. T. Leonard

**Background.** Environmental agencies and the population in general are becoming concerned about the potential of fertilizers to contaminate surface and ground waters with nitrates. Numerous shallow aquifers throughout Texas are used by people who live in rural areas. Some of these aquifers are at risk for nitrate ( $\text{NO}_3^-$ ) contamination. Methodologies are needed which improve crop N uptake and decrease the potential  $\text{NO}_3^-$  contamination of shallow aquifers.

One possible means for lowering the risk of  $\text{NO}_3^-$  pollution in ground waters may be to increase the proportion of plant available N in the ammonium ( $\text{NH}_4^+$ ) form. The main objective of this study is to determine the effect of variable  $\text{NH}_4^+:\text{NO}_3^-$  fertilization on sweet corn yield, N uptake, and soil residual inorganic N.

**Research Findings.** This study is being conducted on a Bowie fine sandy loam near Overton. Supplemental P, K, Mg, S, B, Cu, and Zn were added to the soil. The experimental site was subdivided into 4 replications of 7 treatments applied to 20 x 13.33 ft plots. A control plot received no N. One plot received all the N preplant in the form of ammonium nitrate fertilizer. Treatments 3 through 7 were treated with solutions containing  $\text{NH}_4^+:\text{NO}_3^-$  ratios that varied from zero ammonium nitrogen with 100% of the N applied as calcium nitrate, to 100%  $\text{NH}_4^+$  with no  $\text{NO}_3^-$ . Ratios were 0:100, 25:75, 50:50, 75:25, and 100:0  $\text{NH}_4^+:\text{NO}_3^-$ . Solutions at these ratios were applied by pressure injection through leaky pipe 6 times at 25 lb increments of N/ac at each application. Soil samples were collected from the 0- to 15-, 15- to 30-, and by 30-cm depths to 150-cm deep in each plot. Samples were dried and analyzed for residual  $\text{NH}_4^+$  and  $\text{NO}_3^-$ -N. Harvests were made June 22 and 29.

Yield of the whole plant, ear weight, and number of ears per plant were not significantly affected by N treatment or  $\text{NH}_4^+:\text{NO}_3^-$  ratio (data not shown). Nitrate-N content in the 0- to 15- and 15- to 30-cm depths was increased by increasing the percent  $\text{NO}_3^-$  in the mixture (Table 1). Control plots were low in  $\text{NO}_3^-$ -N. The total N applied preplant as  $\text{NH}_4\text{NO}_3$  increased  $\text{NO}_3^-$ -N in the 0- to 15-cm depth to a level that approximated  $\text{NO}_3^-$  in the soil from the 50:50  $\text{NH}_4^+:\text{NO}_3^-$  mixture. A low concentration of  $\text{NO}_3^-$  was found at all depths below 30 cm.

Soil ammonium concentrations increased as the percentage of  $\text{NH}_4^+$ -N in the blended solutions increased (Table 2). More  $\text{NH}_4^+$ -N was found in the lower soil depths than was  $\text{NO}_3^-$ -N. The sum of the  $\text{NH}_4^+$  and  $\text{NO}_3^-$ -N levels by depths and treatments showed the reason no

differences in yield were detected. The levels of  $\text{NH}_4^+$  plus  $\text{NO}_3^-$ -N in the total soil depth were high and similar for all treatments, including the unfertilized check plot soil.

**Application.** One interesting observation from this study is that soil  $\text{NH}_4^+$ -N levels were generally higher than the  $\text{NO}_3^-$ -N levels in the post-harvest soil samples. There was substantially more soil  $\text{NH}_4^+$  than  $\text{NO}_3^-$  in the soil profile from 30- to 150-cm deep. Leaching of  $\text{NO}_3^-$  in these sandy soils in spring and summer may not be as serious as was anticipated. Accumulation of  $\text{NH}_4^+$  in these deeper depths may be due to soil acidity preventing the increase of the nitrifying bacteria.

Table 1. Effect of  $\text{NH}_4^+:\text{NO}_3^-$  ratio on the  $\text{NO}_3^-$ -N content in the 0- to 15- and 15- to 30-cm depths at 3 sampling times and on profile  $\text{NO}_3^-$ -N after the final harvest. TVA study - 1992.

Treatment $\text{NH}_4^+:\text{NO}_3^-$ %	0-15 cm			15-30 cm			30-150 cm
	5/29	6/19	Postharvest	5/29	6/19	Postharvest	
	-----ppm-----						
Control (0 N)	0.5 c†	0.2B	3.5 B	1.0 d	1.4 c	1.1 b	1.4 NS
Preplant $\text{NH}_4\text{NO}_3$	7.7 bc	12.1 b	13.1 a	10.1 bc	8.1 bc	2.6 b	2.3 NS
0:100	23.6 a	33.7 a	12.9 a	22.4 a	34.7 a	11.9 a	4.2 NS
25:75	21.1 a	15.2 b	8.5 ab	23.9 a	15.8 b	7.0 b	4.1 NS
50:50	10.8 b	12.9 b	9.8 ab	13.1 b	10.7 bc	6.5 b	3.2 NS
75:25	5.4 bc	6.8 b	4.5 b	9.4 bc	7.4 bc	4.5 b	1.6 NS
100:0	1.2 c	3.3 b	4.3 b	3.9 cd	3.4 c	5.1 b	2.8 NS
R <sup>2</sup>	0.86	0.79	0.65	0.85	0.85	0.66	
C.V.	43.9	55.4	44.8	35.3	51.8	54.2	

† $\text{NO}_3^-$ -N levels followed by a similar letter within a column are not different statistically at the  $p = 0.05$  level.

Table 2. Effect of  $\text{NH}_4^+:\text{NO}_3^-$  ratio on the  $\text{NH}_4$ -N content in the 0- to 15- and 15- to 30-cm depths at 3 sampling times and on profile  $\text{NH}_4$ -N after the final harvest. TVA study - 1992.

Treatment $\text{NH}_4^+:\text{NO}_3^-$ %	0-15 cm			15-30 cm			30-150 cm
	5/29	6/19	Postharvest	5/29	6/19	Postharvest	
	-----ppm-----						
Control (0 N)	5.0 c†	7.7‡	21.3‡	7.5 b	4.9 b	4.4‡	41.3‡
Preplant $\text{NH}_4\text{NO}_3$	7.1 c	13.1	5.4	6.8 b	4.7 b	3.5	49.8
0:100	7.5 c	7.0	4.3	5.6 b	13.7 ab	5.9	32.7
25:75	14.6 bc	3.7	6.0	9.3 b	6.3 b	7.0	19.9
50:50	14.9 bc	6.8	24.0	11.0 b	6.3 b	6.0	21.5
75:25	22.9 ab	23.2	11.0	13.6 ab	12.7 ab	0.2	62.2
100:0	29.8 a	23.5	15.3	21.3 a	19.8 a	8.0	30.0
R <sup>2</sup>	0.76	0.42	0.48	0.63	0.61	0.2	
C.V.	42.7	96.4	102.4	50.8	57.1	83.0	

† $\text{NH}_4$ -N levels followed by a similar letter within a column are not different statistically at the  $p = 0.05$  level.

‡Indicates no statistically significant differences due to treatment.