

# **PUBLICATIONS**

## **1990**

## SUSTAINED PRODUCTION FROM COMMON BERMUDAGRASS PASTURES USING CLOVER-POTASSIUM OR RYEGRASS-NITROGEN

F. M. Rouquette, Jr., M. J. Florence, V. A. Haby, and G. R. Smith

### SUMMARY

Common bermudagrass pastures which had been fertilized annually with 200-100-100 lbs N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O/ac for 15 years and grazed at one of three stocking rates were divided into equal sized pastures. Within each stocking rate treatment, one pasture was overseeded with 'Yuchi' arrowleaf clover and fertilized with a single application of approximately 100 lbs K<sub>2</sub>O/ac during each fall of the 5-year study. The other paddock in each stocking rate treatment was overseeded with 'Marshall' ryegrass and received split applications of N. A total annual rate of approximately 400 lbs/ac was applied to the ryegrass and bermudagrass. Cows and calves were used to measure animal performance and gains per unit area. Stocking rates were similar for both the clover-K<sub>2</sub>O and ryegrass-N systems and averaged about 2.0, 1.4, and .85 animal-units/acre, respectively, for high, medium, and low stocking rates. Both calf and cow daily gains were suppressed at the high stocking rates; however, the gains from ryegrass-N were substantially greater than those from the clover-K<sub>2</sub>O. Animal performance from the medium and low stocked pastures was similar for the clover-K<sub>2</sub>O and ryegrass-N systems. Stocking rates during the spring period (Feb.-May) were similar to stocking rates during the summer (June-Sept.). This indicated that nutrient recycling via excreta was relatively effective on the non-N fertilized pastures. Annual fertilizer costs per pound of calf gain were optimized at the medium stocking rates for both K<sub>2</sub>O and N. Actual fertilizer costs ranged from less than a nickel per pound of calf gain for K<sub>2</sub>O to more than 15 cents per pound of calf gain for the N-fertilized pastures.

### INTRODUCTION

Fertilizer applied to exclusive hay meadows is largely removed from the system via hay production. Under grazing systems, however, there is an opportunity for plant food nutrients to be recycled via excreta and used to maintain some level of production. Forage consumed by the grazing animal undergoes microbial degradation in the rumen and the undigested portion is deposited as dung or urine. Various factors affect the extent of recycled plant food nutrients but two primary components include quality of the diet and stocking density. The primary objective of this trial was to determine the influence of previous stocking rate and

fertilizer regimens on common bermudagrass pastures overseeded and fertilized with either clover plus  $K_2O$  or ryegrass plus N.

## PROCEDURES

Common bermudagrass pastures used in this 5-year nutrient recycling study had previously received annual fertilizer rates of 200-100-100 lbs N- $P_2O_5$ - $K_2O$ /ac during each year of a 15-year period. In addition, specific pastures had been grazed at a low, medium, or high stocking rate for the same 15-year period. Thus, for each stocking rate paddock, a total of at least 3000-1500-1500 lbs N- $P_2O_5$ - $K_2O$ /ac had been applied in split applications. The 1500 lb of  $P_2O_5$ /ac combined with nutrient recycling from bovine fecal material had elevated soil test phosphorus to 30 ppm, or very high. Phosphorus fertilizer could be withheld for a few years. Limestone was applied to maintain soil acidity at a level favorable for optimum ryegrass and clover production. In the fall of 1984, each of the stocking rate pastures was subdivided into two equal sized areas. Through random selection, one area was designated to receive only  $K_2O$  and the other area was to receive only N. The pasture that received approximately 100 lbs/ac  $K_2O$  in a single, fall application was overseeded each October with 10 lbs 'Yuchi' arrowleaf clover/ac. During the last 3 years of this study, boron was also applied at the rate of 1.5 to 2 lbs/ac. The pasture that received split applications of a total of approximately 400 lbs N/ac was overseeded each October with 30 lbs 'Marshall' ryegrass/ac. The annual and total fertilizer-lime quantities are shown in Table 1.

Forage in each pasture was sampled every two weeks for nutritive value and at monthly intervals for forage availability. Within any one grazing pressure treatment (High, Medium, or Low), forage availability was regulated as closely as possible between clover- $K_2O$  and ryegrass-N pastures. Brahman X Hereford (F-1) cows and their Simmental-sired calves were used to monitor animal performance and total gains per unit land area. Fall calving cow-calf pairs were grazed from February to early June, and winter-calving cow-calf pairs were grazed from early June to late September. A variable stocking rate was used via the Put-and-Take technique to maintain as closely as possible equivalent grazing pressures among the clover and ryegrass overseeded pastures at each stocking rate. Because of differences in body weight of cows and calves, stocking rates were calculated based on total body weight per acre with one animal-unit being equivalent to 1500 lbs. Animals were weighed at 28-day intervals throughout the 180 to 200-day grazing period.

## RESULTS AND DISCUSSION

Individual cow and calf performance, calf gain per acre, and stocking rate for each of the five years are presented in Tables 2-4 for high, medium, and low stocked pastures, respectively. Table 5 presents a summary of the advantage for the ryegrass-N pastures. Daily calf gains from the high stocked (2.15 AU/ac) ryegrass-N pastures were .64 lbs/day greater than the daily calf gains from the clover-N pastures (1.48 vs .84). This difference in daily gains was due primarily to diversity in stand density and grazing pressure or forage availability rather than to nutritive value. Thus, the impact of calf gain, multiplied by differences in overall stocking rate, resulted in calf gains per acre from the ryegrass-N paddocks that were about 2.5 times greater than that from clover-K<sub>2</sub>O (624 vs 279). During the entire 174-days, 5-year average grazing period, cows on the high stocked bermudagrass-clover pastures lost 354 pounds (-2.33 lbs/day) compared to a loss of 230 pounds (-1.16 lbs/day) for the lactating cows grazing the bermudagrass-ryegrass-N pastures.

At the medium stocking rate of approximately 1.4 AU/ac, cow weights remained at about the same level throughout the grazing period. Daily calf gains were essentially identical at 2.27 for clover-bermudagrass and 2.29 for ryegrass-bermudagrass. The 22 days earlier grazing that ryegrass-N pastures produced was responsible for nearly an extra 100 lb of calf gain/ac. It is noteworthy that these two treatments produced approximately identical individual animal gains and similar stocking rates of 1.40 and 1.45 AU/ac. Thus, there is ample evidence to indicate that nutrient recycling from an existing nutrient pool via excreta had nearly maintained the productivity of common bermudagrass pastures that received no additional fertilizer N during a 5-year period. Performance from the low stocked pastures of clover or ryegrass was also similar with respect to daily calf and cow gains as well as stocking rate (.83 vs .88). The additional 24 days of grazing from the ryegrass-N pastures resulted in an extra 100 pounds of calf gain/ac. The relatively small decrease in stocking rate from 1.4 AU/ac on medium stocked pastures to .85 AU/ac on low stocked pastures was responsible for sufficiently more forage available to provide for cow daily gains of nearly 1 lb/day and total grazing period cow gains of approximately 175 lbs.

Five-year average daily calf gains at each of three stocking rates during the spring and summer periods are presented in Table 6. Within a stocking rate treatment, there were similar animal-units per acre during the spring and summer periods. In addition, there were similar stocking rates between the clover-K<sub>2</sub>O and

ryegrass-N treatments. Obviously from the depressed calf gains that resulted from the clover-K<sub>2</sub>O pastures compared to the ryegrass-N pastures, grazing pressures were inadvertently heavier on the clover pastures. Thus, the extent of nutrient cycling and the resultant impact on the fertilizer-forage system appears to be very effective in the non-N fertilized pastures of common bermudagrass. In a companion paper that deals with Coastal bermudagrass, this degree of similarity between N and non-N fertilized pastures was not as apparent. Thus, with the inherently lower forage producing common bermudagrass pastures, the higher rates of N fertilizer had little production impact as compared to the clover-K<sub>2</sub>O treatment.

With the exception of the high stocking rate which has already been mentioned, the total 5-year calf gain per acre was also similar (Table 7). Figure 1 shows a classical example of the relationship between gain per acre and stocking rate for both the clover-K<sub>2</sub>O and ryegrass-N systems. There was a dramatic decline in gain per acre from the bermudagrass-clover-K<sub>2</sub>O pastures with increasing stocking rate. An optimum relationship was apparently achieved between 1.0 and 1.5 animal-units per acre. The ryegrass-N treated bermudagrass pastures, however, allowed optimum calf gains per acre at approximately 2 animal units per acre. The ryegrass-N treated bermudagrass pastures maintained higher levels of animal production at all levels of stocking, but this was even more dramatic at the higher levels of stocking. The relationships of daily gains of both cows and calves with changes in stocking rate also shows the disparity between clover-K<sub>2</sub>O and ryegrass-N treated pastures (Fig. 2). Calf response from the clover-K<sub>2</sub>O pastures appears to be negatively curvilinear as expected; however, calf response was negatively linear from the ryegrass-N pastures. Certainly forage availability or forage density at these similar stocking rates may explain these differences. The F-1 (Brahman x Hereford) cow continued to lactate even though the weight losses were appreciable. This sometimes small quantity of milk, however, was responsible for the positive calf gains obtained at the high stocking rates. Thus, the importance of milk as a "buffer system" for sub-optimal nutritional regimens of quantity or quality is clearly evident.

The annual cost of fertilizer and the fertilizer costs per pound of calf gain are presented in Table 8. From these economic assessments, the K<sub>2</sub>O costs were optimized at about 3 cents per pound of calf gain at the medium stocking rate. The other two stocking rates projected a fertilizer cost of about a nickel per pound of calf gain. The medium stocking rate was also the optimum rate for fertilizer costs associated with applying 408 lbs N/ac per year. Again, in pasture systems in which

utilization efficiency is low, such as the low stocked pastures, there is a dramatic percentage increase in fertilizer cost per pound of gain. In this study, fertilizer N costs increased from 15 cents to 21 cents on medium (1.40) and low (.83) stocked pastures, respectively.

Additional input data would be necessary in order to construct a detailed cash flow analyses, but the data presented here may provide some base line information on fertilizer and stocking rate management strategies. With the exception of the high stocked clover-K<sub>2</sub>O pastures in which some stand thinning occurred as well as some invasion of bahiagrass, common bermudagrass pastures exhibited no forage-animal production losses by the absence of fertilizer nitrogen. Thus, nutrient recycling appears to be a biological and economic asset in sustaining common bermudagrass grazing systems. Additional data on stand loss and soil fertility are currently being assessed and summaries of this information will be forthcoming.

TABLE 1. FERTILIZER RATES APPLIED TO COMMON BERMUDAGRASS PASTURES DURING A FIVE-YEAR PERIOD

| Year         | Date                                      | Clover +<br>K <sub>2</sub> O | Ryegrass +<br>Nitrogen |
|--------------|---|------------------------------|------------------------|
|              |   | -----lbs/ac-----             |                        |
| 1            | 11-29-84<br>2-20-85 to 9-17-85<br>9-26-85 | 0-0-100                      | 390-0-0                |
|              |   | -----2 Tons Lime/Ac-----     |                        |
| 2            | 11-22-85<br>11-26-85 to 9-8-86            | 0-0-100                      | 400-0-0                |
| 3            | 11-20-86<br>1-27-87 to 8-25-87<br>9-15-87 | 0-0-100 + 2B                 | 400-0-0                |
|              |   | -----1 Ton Lime/Ac-----      |                        |
| 4            | 11-18-87<br>12-1-87 to 8-30-88            | 0-0-150 + 1.5B               | 450-0-0                |
| 5            | 11-9-88<br>12-14-88 to 7-5-89             | 0-0-120 + 1.7B               | 400-0-0                |
| 5-YEAR TOTAL |   | 0-0-570 + 5.2B               | 2040-0-0               |
| YEARLY AVG.  |   | 0-0-114 + 1B                 | 408-0-0                |

TABLE 2. ANIMAL PERFORMANCE FROM HIGH STOCKED COMMON BERMUDAGRASS PASTURES OVERSEEDED WITH EITHER ARROWLEAF CLOVER OR RYEGRASS

| Year                  | Annual Forage | Grazing Days | Average Daily Gain |       | Gain/Animal   |      | Gain/Acre Calf<br>-lbs- | Stocking Rate*<br>AU/ac |
|-----------------------|---------------|--------------|--------------------|-------|---------------|------|-------------------------|-------------------------|
|                       |               |              | Calf               | Cow   | Calf          | Cow  |                         |                         |
|                       |               |              | -----lbs/d-----    |       | -----lbs----- |      |                         |                         |
| 1                     | ARL†          | 175          | 1.30               | -1.51 | 228           | -265 | 436                     | 1.91                    |
| 1                     | RYG           | 187          | 1.81               | - .69 | 340           | -128 | 721                     | 2.13                    |
| 2                     | ARL           | 176          | .79                | -2.24 | 139           | -395 | 244                     | 1.77                    |
| 2                     | RYG           | 211          | 1.41               | -1.45 | 298           | -306 | 584                     | 1.96                    |
| 3                     | ARL           | 208          | .83                | -2.27 | 172           | -474 | 314                     | 1.83                    |
| 3                     | RYG           | 219          | 1.27               | -1.59 | 279           | -349 | 617                     | 2.22                    |
| 4                     | ARL           | 142          | .90                | -2.86 | 129           | -407 | 247                     | 1.97                    |
| 4                     | RYG           | 173          | 1.78               | -.96  | 308           | -166 | 592                     | 1.90                    |
| 5                     | ARL           | 170          | .40                | -1.36 | 68            | -231 | 153                     | 2.17                    |
| 5                     | RYG           | 204          | 1.21               | -.98  | 247           | -201 | 608                     | 2.50                    |
| <u>5-YEAR AVERAGE</u> |               |              |                    |       |               |      |                         |                         |
|                       | ARL           | 174          | .84                | -2.33 | 147           | -354 | 279                     | 1.92                    |
|                       | RYG           | 199          | 1.48               | -1.16 | 294           | -230 | 624                     | 2.15                    |

\*1 AU = 1500-lb body weight.

†ARL = arrowleaf; RYG = ryegrass.

TABLE 3. ANIMAL PERFORMANCE FROM MEDIUM STOCKED COMMON BERMUDAGRASS PASTURES OVERSEEDED WITH EITHER ARROWLEAF CLOVER OR RYEGRASS

| Year                  | Annual Forage | Grazing Days | Average Daily Gain |      | Gain/Animal   |      | Gain/Acre Calf -lbs- | Stocking Rate* AU/ac |
|-----------------------|---------------|--------------|--------------------|------|---------------|------|----------------------|----------------------|
|                       |               |              | Calf               | Cow  | Calf          | Cow  |                      |                      |
|                       |               |              | -----lbs/d-----    |      | -----lbs----- |      |                      |                      |
| 1                     | ARL†          | 175          | 2.30               | .48  | 403           | 85   | 566                  | 1.40                 |
| 1                     | RYG           | 187          | 2.51               | .51  | 471           | 95   | 670                  | 1.42                 |
| 2                     | ARL           | 176          | 2.05               | -.67 | 360           | -118 | 492                  | 1.37                 |
| 2                     | RYG           | 211          | 2.05               | .21  | 433           | -45  | 614                  | 1.41                 |
| 3                     | ARL           | 219          | 2.08               | -.60 | 456           | -132 | 654                  | 1.43                 |
| 3                     | RYG           | 211          | 2.30               | -.70 | 487           | -148 | 713                  | 1.46                 |
| 4                     | ARL           | 160          | 2.73               | .11  | 438           | -17  | 600                  | 1.37                 |
| 4                     | RYG           | 189          | 2.60               | -.06 | 492           | -12  | 661                  | 1.35                 |
| 5                     | ARL           | 162          | 2.27               | .40  | 369           | 66   | 530                  | 1.44                 |
| 5                     | RYG           | 204          | 2.06               | .12  | 419           | 26   | 662                  | 1.60                 |
| <u>5-YEAR AVERAGE</u> |               |              |                    |      |               |      |                      |                      |
|                       | ARL           | 178          | 2.27               | -.06 | 405           | -23  | 568                  | 1.40                 |
|                       | RYG           | 200          | 2.29               | .02  | 460           | -17  | 664                  | 1.45                 |

\*1 AU = 1500-lb body weight.

†ARL = arrowleaf; RYG = ryegrass.





TABLE 5. FIVE-YEAR COMPARISON OF CALF PERFORMANCE FROM COMMON BERMUDAGRASS PASTURES OVERSEEDED WITH EITHER ARROWLEAF CLOVER OR RYEGRASS AND STOCKED AT EACH OF 3 LEVELS

| Grazing Pressure                 | Annual Forage | Grazing Days | CALF  |                 |               | Stocking Rate*<br>AU/ac |
|----------------------------------|---------------|--------------|-------|-----------------|---------------|-------------------------|
|                                  |               |              | ADG   | Gain/<br>Animal | Gain/<br>Acre |                         |
| HIGH                             | ARL†          | 174          | .84   | 147             | 279           | 1.92                    |
| HIGH                             | RYG           | 199          | 1.48  | 294             | 624           | 2.15                    |
| RYEGRASS + NITROGEN<br>ADVANTAGE |               | +25          | + .64 | +147            | +345          | + .23                   |
| MEDIUM                           | ARL           | 178          | 2.27  | 405             | 568           | 1.40                    |
| MEDIUM                           | RYG           | 200          | 2.29  | 460             | 664           | 1.45                    |
| RYEGRASS + NITROGEN<br>ADVANTAGE |               | +22          | + .02 | + 55            | + 96          | + .05                   |
| LOW                              | ARL           | 178          | 2.52  | 450             | 371           | .83                     |
| LOW                              | RYG           | 202          | 2.69  | 544             | 482           | .88                     |
| RYEGRASS + NITROGEN<br>ADVANTAGE |               | +24          | + .17 | + 94            | +111          | + .05                   |

\*1 AU = 1500-lb body weight.

†ARL = arrowleaf; RYG = ryegrass.

TABLE 6. FIVE-YEAR AVERAGE OF CALF DAILY GAINS DURING THE SPRING AND SUMMER PERIODS FROM OVERSEEDED COMMON BERMUDAGRASS PASTURES

| Clover + K <sub>2</sub> O |       |         |       | Ryegrass + N |       |         |       |
|---------------------------|-------|---------|-------|--------------|-------|---------|-------|
| Spring                    |       | Summer* |       | Spring       |       | Summer* |       |
| SR                        | ADG   | SR      | ADG   | SR           | ADG   | SR      | ADG   |
| AU/ac                     | lb/da | AU/ac   | lb/da | AU/ac        | lb/da | AU/ac   | lb/da |
| 1.93                      | .96   | 1.97    | .72   | 2.10         | 1.71  | 2.20    | 1.28  |
| 1.42                      | 2.75  | 1.39    | 1.87  | 1.48         | 2.65  | 1.42    | 1.94  |
| .80                       | 2.99  | .85     | 2.15  | .92          | 3.07  | .85     | 2.32  |

\*Pastures consist of exclusive bermudagrass.

TABLE 7. TOTAL CALF GAIN PER ACRE DURING A FIVE-YEAR PERIOD OF GRAZING COMMON BERMUDAGRASS PASTURES

| Stocking Rate | Clover + K <sub>2</sub> O Bermudagrass | Ryegrass + N Bermudagrass |
|---------------|--|---------------------------|
|               | -----lbs/ac-----                       | -----lbs/ac-----          |
| High          | 1394                                   | 3122                      |
| Medium        | 2842                                   | 3320                      |
| Low           | 1856                                   | 2409                      |






Figure 1. Relationship of calf gain per acre with stocking rate on common bermudagrass pastures in combination with either nitrogen plus ryegrass or potassium plus clover.




Figure 2. Relationship of daily gain of cows and calves at different stocking rates on common bermudagrass (BMG) pastures in combination with either potassium (K) plus clover (CLV) or nitrogen (N) plus ryegrass (RYG).