

I 19.5/1:  
36  
C.2  
Oversize  
Section

TEXAS A&M UNIVERSITY LIBRARY


DEPARTMENT OF THE INTERIOR  
UNITED STATES GEOLOGICAL SURVEY  
CHARLES D. WALCOTT, DIRECTOR

# GEOLOGIC ATLAS

## OF THE UNITED STATES

### PUEBLO FOLIO COLORADO

INDEX MAP



SCALE: 40 MILES = 1 INCH

AREA OF THE PUEBLO FOLIO

AREA OF OTHER PUBLISHED FOLIOS

LIBRARY  
TEXAS A&M UNIVERSITY  
NOV 14 1967

DOCUMENTS

#### LIST OF SHEETS

| DESCRIPTION | TOPOGRAPHY  | AREAL GEOLOGY  | ECONOMIC GEOLOGY  | STRUCTURE SECTIONS    |
|-------------|-------------|----------------|-------------------|-----------------------|
|             | DEFORMATION | ARTESIAN WATER | COLUMNAR SECTIONS | SPECIAL ILLUSTRATIONS |
| FOLIO 36    |             |                | LIBRARY EDITION   | PUEBLO                |

WASHINGTON, D. C.

ENGRAVED AND PRINTED BY THE U.S. GEOLOGICAL SURVEY

BAILEY WILLIS, EDITOR OF GEOLOGIC MAPS      S. J. K. REL., CHIEF ENGRAVER

1897



# EXPLANATION.

The Geological Survey is making a geologic map of the United States, which necessitates the preparation of a topographic base map. The two are being issued together in the form of an atlas, the parts of which are called folios. Each folio consists of a topographic base map and geologic maps of a small area of country, together with explanatory and descriptive texts.

## THE TOPOGRAPHIC MAP.

The features represented on the topographic map are of three distinct kinds: (1) inequalities of surface, called *relief*, as plains, plateaus, valleys, hills, and mountains; (2) distribution of water, called *drainage*, as streams, lakes, and swamps; (3) the works of man, called *culture*, as roads, railroads, boundaries, villages, and cities.

**Relief.**—All elevations are measured from mean sea-level. The heights of many points are accurately determined, and those which are most important are given on the map in figures. It is desirable, however, to give the elevation of all parts of the area mapped, to delineate the horizontal outline, or contour, of all slopes, and to indicate their grade, or degree of steepness. This is done by lines connecting points of equal elevation above mean sea-level, the lines being drawn at regular vertical intervals. These lines are called *contours*, and the uniform vertical space between each two contours is called the *contour interval*. Contours and elevations are printed in brown.

The manner in which contours express elevation, form, and grade is shown in the following sketch and corresponding contour map:




Fig. 1.—Ideal sketch and corresponding contour map.

The sketch represents a river valley between two hills. In the foreground is the sea, with a bay which is partly closed by a hooked sand-bar. On each side of the valley is a terrace. From the terrace on the right a hill rises gradually, while from that on the left the ground ascends steeply to a precipice. Contrasted with this precipice is the gentle descent of the left-hand slope. In the map each of these features is indicated, directly beneath its position in the sketch, by contours. The following explanation may make clearer the manner in which contours delineate elevation, form, and grade:

1. A contour indicates approximately a certain height above sea-level. In this illustration the contour interval is 50 feet; therefore the contours are drawn at 50, 100, 150, 200 feet, and so on, above sea-level. Along the contour at 250 feet lie all points of the surface 250 feet above sea; and similarly with any other contour. In the space between any two contours are found all elevations above the lower and below the higher contour. Thus the contour at 150 feet falls just below the edge of the terrace, while that at 200 feet lies above the terrace; therefore all points on the terrace are shown to be more than 150 but less than 200 feet above sea. The summit of the higher hill is stated to be 670 feet above sea; accordingly the contour at 650 feet surrounds it. In this illustration nearly all the contours are numbered. Where this is not possible, certain contours—say every fifth one—are accentuated and numbered; the heights of others may then be ascertained by counting up or down from a numbered contour.

2. Contours define the forms of slopes. Since contours are continuous horizontal lines conforming to the surface of the ground, they wind smoothly about smooth surfaces, recede into all reentrant angles of ravines, and project in passing about prominences. The relations of contour curves and angles to forms of the landscape can be traced in the map and sketch.

3. Contours show the approximate grade of any slope. The vertical space between two contours is the same, whether they lie along a cliff or on a gentle slope; but to rise a given height on a gentle slope one must go farther than on a steep slope, and therefore contours are far apart on gentle slopes and near together on steep ones.

For a flat or gently undulating country a small contour interval is used; for a steep or mountainous country a large interval is necessary. The smallest interval used on the atlas sheets of the Geological Survey is 5 feet. This is used for regions like the Mississippi delta and the Dismal Swamp. In mapping great mountain masses, like those in Colorado, the interval may be 250 feet. For intermediate relief contour intervals of 10, 20, 25, 50, and 100 feet are used.

**Drainage.**—Watercourses are indicated by blue lines. If the stream flows the year round the line is drawn unbroken, but if the channel is dry a part of the year the line is broken or dotted. Where a stream sinks and reappears at the surface, the supposed underground course is shown by a broken blue line. Lakes, marshes, and other bodies of water are also shown in blue, by appropriate conventional signs.

**Culture.**—The works of man, such as roads, railroads, and towns, together with boundaries of townships, counties, and States, and artificial details, are printed in black.

**Scales.**—The area of the United States (excluding Alaska) is about 3,025,000 square miles. On a map with the scale of 1 mile to the inch this would cover 3,025,000 square inches, and to accommodate it the paper dimensions would need to be about 240 by 180 feet. Each square mile of ground surface would be represented by a square inch of map surface, and one linear mile on the ground would be represented by a linear inch on the map. This relation between distance in nature and corresponding distance on the map is called the scale of the map. In this case it is "1 mile to an inch." The scale may be expressed also by a fraction, of which the numerator is a length on the map and the denominator the corresponding length in nature expressed in the same unit. Thus, as there are 63,360 inches in a mile, the scale "1 mile to an inch" is expressed by  $\frac{1}{63,360}$ . Both of these methods are used on the maps of the Geological Survey.

Three scales are used on the atlas sheets of the Geological Survey; the smallest is  $\frac{1}{250,000}$ , the intermediate  $\frac{1}{125,000}$ , and the largest  $\frac{1}{62,500}$ . These correspond approximately to 4 miles, 2 miles, and 1 mile on the ground to an inch on the map. On the scale  $\frac{1}{62,500}$  a square inch of map surface represents and corresponds nearly to 1 square mile; on the scale  $\frac{1}{125,000}$  to about 4 square miles; and on the scale  $\frac{1}{250,000}$  to about 16 square miles. At the bottom of each atlas sheet the scale is expressed in three different ways, one being a graduated line representing miles and parts of miles in English inches, another indicating distance in the metric system, and a third giving the fractional scale.

**Atlas sheets and quadrangles.**—The map is being published in atlas sheets of convenient size, which are bounded by parallels and meridians. The corresponding four-cornered portions of territory are called *quadrangles*. Each sheet on the scale of  $\frac{1}{250,000}$  contains one square degree, i. e., a degree of latitude by a degree of longitude; each sheet on the scale of  $\frac{1}{125,000}$  contains one-quarter of a square degree; each sheet on the scale of  $\frac{1}{62,500}$  contains one-sixteenth of a square degree. The areas of the corresponding quadrangles are about 4000, 1000, and 250 square miles, respectively.

The atlas sheets, being only parts of one map of the United States, are laid out without regard to the boundary lines of the States, counties or townships. To each sheet, and the quadrangle it represents, is given the name of some well-known

town or natural feature within its limits, and at the sides and corners of each sheet the names of adjacent sheets, if published, are printed.

**Uses of the topographic sheet.**—Within the limits of scale the topographic sheet is an accurate and characteristic delineation of the relief, drainage, and culture of the district represented. Viewing the landscape, map in hand, every characteristic feature of sufficient magnitude should be recognizable. It should guide the traveler; serve the investor or owner who desires to ascertain the position and surroundings of property to be bought or sold; save the engineer preliminary surveys in locating roads, railways, and irrigation ditches; provide educational material for schools and homes; and serve many of the purposes of a map for local reference.

## THE GEOLOGIC MAP.

The areal geologic map represents by colors and conventional signs, on the topographic base map, the distribution of rock formations on the surface of the earth, and the structure-section map shows their underground relations, as far as known, and in such detail as the scale permits.

### KINDS OF ROCKS.

Rocks are of many kinds. The original crust of the earth was probably composed of *igneous rocks*, and all other rocks have been derived from them in one way or another.

Atmospheric agencies gradually break up igneous rocks, forming superficial, or *surficial*, deposits of clay, sand, and gravel. Deposits of this class have been formed on land surfaces since the earliest geologic time. Through the transporting agencies of streams the surficial materials of all ages and origins are carried to the sea, where, along with material derived from the land by the action of the waves on the coast, they form *sedimentary rocks*. These are usually hardened into conglomerate, sandstone, shale, and limestone, but they may remain unconsolidated and still be called "rocks" by the geologist, though popularly known as gravel, sand, and clay.

From time to time in geologic history igneous and sedimentary rocks have been deeply buried, consolidated, and raised again above the surface of the water. In these processes, through the agencies of pressure, movement, and chemical action, they are often greatly altered, and in this condition they are called *metamorphic rocks*.

**Igneous rocks.**—These are rocks which have cooled and consolidated from a liquid state. As has been explained, sedimentary rocks were deposited on the original igneous rocks. Through the igneous and sedimentary rocks of all ages molten material has from time to time been forced upward to or near the surface, and there consolidated. When the channels or vents into which this molten material is forced do not reach the surface, it either consolidates in cracks or fissures crossing the bedding planes, thus forming dikes, or else spreads out between the strata in large bodies, called sills or laccoliths. Such rocks are called *intrusive*. Within their rock enclosures they cool slowly, and hence are generally of crystalline texture. When the channels reach the surface the lavas often flow out and build up volcanoes. These lavas cool rapidly in the air, acquiring a glassy or, more often, a partially crystalline condition. They are usually more or less porous. The igneous rocks thus formed upon the surface are called *extrusive*. Explosive action often accompanies volcanic eruptions, causing ejections of dust or ash and larger fragments. These materials when consolidated constitute breccias, agglomerates, and tuffs. The ash when carried into lakes or seas may become stratified, so as to have the structure of sedimentary rocks.

The age of an igneous rock is often difficult or impossible to determine. When it cuts across a sedimentary rock, it is younger than that rock, and when a sedimentary rock is deposited over it, the igneous rock is the older.

Under the influence of dynamic and chemical forces an igneous rock may be metamorphosed. The alteration may involve only a rearrangement of its minute particles or it may be accompanied by a change in chemical and mineralogical composition. Further, the structure of the rock may be

changed by the development of planes of division, so that it splits in one direction more easily than in others. Thus a granite may pass into a gneiss, and from that into a mica-schist.

**Sedimentary rocks.**—These comprise all rocks which have been deposited under water, whether in sea, lake, or stream. They form a very large part of the dry land.

When the materials of which sedimentary rocks are made are carried as solid particles by the water and deposited as gravel, sand, or mud, the deposit is called a mechanical sediment. These may become hardened into conglomerate, sandstone, or shale. When the material is carried in solution by the water and is deposited without the aid of life, it is called a chemical sediment; if deposited with the aid of life, it is called an organic sediment. The more important rocks formed from chemical and organic deposits are limestone, chert, gypsum, salt, iron ore, peat, lignite, and coal. Any one of the above sedimentary deposits may be separately formed, or the different materials may be intermingled in many ways, producing a great variety of rocks.

Sedimentary rocks are usually made up of layers or beds which can be easily separated. These layers are called *strata*. Rocks deposited in successive layers are said to be stratified.

The surface of the earth is not fixed, as it seems to be; it very slowly rises or sinks over wide expanses, and as it rises or subsides the shore-lines of the ocean are changed: areas of deposition may rise above the water and become land areas, and land areas may sink below the water and become areas of deposition. If North America were gradually to sink a thousand feet the sea would flow over the Atlantic coast and the Mississippi and Ohio valleys from the Gulf of Mexico to the Great Lakes; the Appalachian Mountains would become an archipelago, and the ocean's shore would traverse Wisconsin, Iowa, and Kansas, and extend thence to Texas. More extensive changes than this have repeatedly occurred in the past.

The character of the original sediments may be changed by chemical and dynamic action so as to produce metamorphic rocks. In the metamorphism of a sedimentary rock, just as in the metamorphism of an igneous rock, the substances of which it is composed may enter into new combinations, or new substances may be added. When these processes are complete the sedimentary rock becomes crystalline. Such changes transform sandstone to quartzite, limestone to marble, and modify other rocks according to their composition. A system of parallel division planes is often produced, which may cross the original beds or strata at any angle. Rocks divided by such planes are called slates or schists.

Rocks of any period of the earth's history may be more or less altered, but the younger formations have generally escaped marked metamorphism, and the oldest sediments known, though generally the most altered, in some localities remain essentially unchanged.

**Surficial rocks.**—These embrace the soils, clays, sands, gravels, and boulders that cover the surface, whether derived from the breaking up or disintegration of the underlying rocks by atmospheric agencies or from glacial action. Surficial rocks that are due to disintegration are produced chiefly by the action of air, water, frost, animals, and plants. They consist mainly of the least soluble parts of the rocks, which remain after the more soluble parts have been leached out, and hence are known as residual products. Soils and subsoils are the most important. Residual accumulations are often washed or blown into valleys or other depressions, where they lodge and form deposits that grade into the sedimentary class. Surficial rocks that are due to glacial action are formed of the products of disintegration, together with boulders and fragments of rock rubbed from the surface and ground together. These are spread irregularly over the territory occupied by the ice, and form a mixture of clay, pebbles, and boulders which is known as till. It may occur as a sheet or be bunched into hills and ridges, forming moraines, drumlins, and other special forms. Much of this mixed material was washed away from the ice, assorted by water, and redeposited as beds or trains of sand and clay, thus

(Continued on third page of cover.)



# DESCRIPTION OF THE PUEBLO QUADRANGLE.

## INTRODUCTION.

It is assumed by the writer of this text that the facts and deductions recorded in the folio are of interest not only to professional geologists but to those residents of Pueblo and vicinity who care to know the origin of rocks and hills or who desire to make use of the mineral products of the district. Endeavor has therefore been made to avoid technical language, so far as may be, and where avoidance is impracticable, to explain the terms used. The layman is advised to read the "Explanation" printed on the inner pages of the cover, and his attention is invited to the supplementary explanations in the paragraphs immediately following.

A stratum, layer, bed, or other sedimentary formation having great horizontal extent as compared to its thickness may lie level or may be inclined. If inclined, the amount of its inclination or slope is called its *dip*, and the measure of the dip is the angle between the surface of the formation and a horizontal plane. The direction of its steepest slope is called the direction of dip, and the direction at right angles to this the *strike*. Thus if the dip of a stratum is northeast its strike is northwest. The stratum is also said to *dip* northeast and to *strike* northwest.

When a stratum or other body of rock is broken across and one part has slid past the other, the dislocation is called a *fault*, and the rock is said to be *faulted*. The amount of dislocation, or the distance separating corresponding parts of the severed mass is called the *displacement* of the fault. In fig. 1 the distance *ab*, separating parts of the faulted stratum 2,2, is the displacement.




Fig. 1.—Ideal section showing a fault. The mass of strata on the left has moved down as compared to the mass at the right. The numbers show corresponding strata that were once continuous.

## GEOGRAPHY.

The Pueblo quadrangle extends in longitude from 104° 30' to 105°, and in latitude from 38° to 38° 30'. It is 34.5 miles long, north and south, 27 miles wide, and contains 938 square miles. It includes part of Pueblo County and the southeast corner of Fremont County, Colorado.

The great features of the continent to which it is related are the Rocky Mountains and the Great Plains. The surface of the Plains rises gradually from east to west and is more diversified by hills in its western portion, so that the transition from plain to mountain is not abrupt. There is usually a belt of intermediate character, called foothills, and this belt is well developed in the vicinity of the Pueblo quadrangle. At several places the front line of the mountains is broken by embayments into which extend tongues of the plain, and one of these embayments is traversed from west to east by the Arkansas River. The Pueblo district lies near the western edge of the Plains, opposite the Arkansas embayment. It belongs chiefly to the plains province, but includes also two portions of the foothills belt, one along the western part of its north boundary, the other along the southern part of its west boundary.

Broadly viewed, the surface is a plateau, ranging in height from 4700 to 5500 or 6000 feet. It is lowest at the east, near the city of Pueblo, and rises toward the north, south, and west. The streams run below the general level, following narrow valleys or canyons a few scores or a few hundreds of feet in depth, and the plateau is overlooked by the foothills already mentioned as well as by a few outlying hills and mesas. Baculite Mesa, northeast of Pueblo, rises 400 feet from the plain; the so-

called "Sand hill," southwest of the city, is about half as high; and there are several unnamed tables in the southern part of the quadrangle.

The drainage system corresponds in its principal lines with the general slopes of the surface. Its main artery is the Arkansas River, which enters the district from the west a few miles north of the middle, runs southeasterly to the center, and thence easterly. The general course of tributaries on the north is southerly, the chief being Fountain Creek, which joins the Arkansas at Pueblo. South of the Arkansas the general course of the drainage is toward the northeast, and the most important stream is the St. Charles River, which joins the Arkansas a few miles east of the boundary of the quadrangle. The Arkansas carries a large body of water at all seasons. Each of the other streams is at certain seasons and at certain places lost, its water being absorbed by the sands of its bed.

There is a considerable range of climate, corresponding chiefly with differences of altitude. The eastern part shares the heat and aridity of the western belt of the Plains; the foothill tracts at the west are cooler and moister because higher; and the western part has an additional advantage from summer showers generated in neighboring mountains.

Vegetation is of several types, closely related to climate and soil. A forest of yellow pine occupies most uplands above 6500 feet, and there are straggling pine groves on sandy hillsides down to 5300 feet. In the same zone rocky slopes are sometimes covered with aspens; on sandy soils the lower pines are accompanied by thickets of dwarf oak; and moist canyons shelter the hackberry and other hardwood trees of moderate size. Cottonwoods occupy the bottom lands of all permanent and many intermittent streams. Below the yellow pines are junipers and piñon pines, and these extend down to 5000 feet altitude, with stragglers beyond. They grow only on rocky and gravelly soils where the slope is steep, and are chiefly associated with ledges of limestone and sandstone. The remainder of the land, including much the greater part, is prairie, with an open growth of low bushes and grass.

The moisture necessary for the pine forest is adequate also for cultivated crops, and a few small tracts of favorable soil are farmed without irrigation. The associated climate gives but a short season, and only the hardier cereals and vegetables are grown. There is also a certain amount of natural irrigation of bottom lands, but little agriculture is based on it. Recourse is usually had to artificial irrigation, and for nearly all the land this is essential to successful farming. A large canal—the Bessemer ditch—carries water from the Arkansas River to a broad mesa on the south side and embraces several square miles of arable land, but irrigation is otherwise limited to narrow belts following the river and larger creeks. The cultivated area is but a small fraction of the district, and the remainder is used only for grazing.

## GENERAL GEOLOGY.














The rocks comprise a number of formations, each with its individual character, and these stand in certain definite relations one to another. They are also related in an intimate way to the forms of the surface. Clearly to understand these relations it is necessary to take account of the ways in which the rocks were made, of the ways in which their original arrangement has been disturbed, and of the ways in which the existing forms of hill and valley have been produced. In the investigation of the district the rocks and hills have been questioned and made to tell the story of their origin, and in this way the long sequence of physical events constituting the geologic history of the district has been gradually worked out. Knowledge of the character and arrangement of the rocks has thus preceded knowledge of their history, but it is thought that the subject can be more clearly presented by reversing this order. The history of the physical changes will first be outlined, and will then be used in explaining the

character and arrangement of the rocks and the forms of the surface.

## HISTORY OF PHYSICAL CHANGES.

At the earliest date which need be considered here the surface of the district was a plain, more even than the present surface, and the rocks composing its floor were granite, gneiss, mica-schist, and similar formations, known collectively as "crystalline schists." There was, of course, a history of origin for these formations as for all others, but present purposes do not demand its consideration. The individual rock masses of the series were in the form of irregular plates, dipping steeply downward and fitted together so as to form a continuous mass, and the upper edges of the plates formed the continuous surface of the plain. The date at which this plain existed is so remote that its antiquity is not measured or known in years, and if the number of years could be written it would probably be too large to convey a definite impression to the mind.

IDEAL SECTIONS ILLUSTRATING THE GEOLOGIC HISTORY OF THE PUEBLO DISTRICT.  
(The line W marks the position of the water surface.)



The floor of crystalline rocks, at the epoch assumed as initial, was part of the dry land, just as now, but that condition has not been continuous from the assumed beginning to the present time. From time to time the level of the ocean

has been raised so as to cover the land (or the land has subsided so as to sink beneath the ocean), and there have also been times when the district was covered by the waters of an inland sea.

The first change was a submergence, the plain becoming the bed of an ocean, and on this ocean bed were spread various sediments brought by currents from neighboring lands. First came a deposit of white sand, and then one of limy ooze. In the course of time the sand was cemented and hardened into sandstone, and the limy ooze became limestone. We do not know how evenly these deposits were spread, but it is probable that the limestone at least covered the entire district, forming a continuous sheet or layer, so that the floor of crystalline rock was wholly concealed. The duration of the submergence corresponds to three periods of the geologic time scale, the Silurian, Devonian, and Carboniferous. Fig. 2 represents in a diagrammatic way the floor of crystalline rocks, as seen in section; fig. 3, the same floor with a covering of sediments.

The next event was an unequal uplift of the ocean floor. Parts of it were pushed up from below, so that the even expanse came to be broken in places by heights. At least one of the uplifts was raised above the surface of the ocean (fig. 4), making a new island, or perhaps part of a continent, and in consequence of this exposure was subjected to the dashing of waves, the beating of rain, and the washing of streams, so that the new deposit was locally worn away and even the crystalline rocks were eroded. Another effect of the uprisings which occurred at that time was to partition off a part of the ocean, so that the waters of the Pueblo district became part of an interior sea or salt lake. The sand, gravel, and mud washed by the streams from the surrounding lands accumulated in this sea, forming a new series of deposits. By the waves and currents the material was separated into kinds, so that the deposits in different places were different; and it happened that this district no longer received limy ooze, but clay or mud, and with it sand. Afterward, when the deposits were cemented and hardened, they became strata of shale and sandstone, and they are further distinguished from the older strata on which they rest by having a deep-red color. The general condition of the district at this time is shown in fig. 5. The time represented by the red sediments is the earlier part of the Juratrias period.

There followed a general but not uniform subsidence, so that the land was once more covered by a sea, and this continued during the deposition of other layers of mud and some layers of sand. The sea was completely shut off from the ocean, and became at times a dense brine, like the Dead Sea or Great Salt Lake. When its brine was most concentrated gypsum and perhaps other minerals were separated by evaporation, just as salt is separated by the evaporation of ocean water. There were also disturbances of the bottom by uplift from below, so that part of the new deposit was lifted above the water and again washed away. The hardening of these sediments produced shales of many brilliant colors, with minor layers of sandstone and gypsum. The time of their accumulation was the later part of the Juratrias period.

A change then occurred in the character of the water. The inland sea in which strong brine had gathered became connected with the ocean, so as to be a bay or estuary, and through this connection its waters were rendered as fresh as those of the ocean. At times they were probably even fresher, but on this point the evidence is not clear. In connection with these changes the water first receded from a portion or advanced so as to completely cover it and extend far beyond. The advance was not at a uniform rate, nor even continuous, but alternated with retreat, so that the coast-line passed and repassed the district several times. The mud and sand washed by the rivers from neighboring lands were sorted by the waves on this shifting coast, the result being that the sand was accumulated



at the shore-line and the mud was carried by the currents to deeper water. Each part of the quadrangle, being covered alternately by the shoal water of the coast and the deeper water off shore, received alternating deposits of sand and mud which were afterward compacted into sandstone and shale. The manner in which these sands were deposited is of great practical importance to man, for their long agitation by the waves washed out all the finer particles, so that the resulting sandstone is made up of coarse grains which do not fit closely together and it is therefore porous. Being porous it absorbs water freely and also permits it to flow through, making an underground circulation available for artesian wells.

After sand with occasional layers of mud had accumulated to a depth of several hundred feet, the Pueblo district sank much lower, so as to be deeply covered by water. <sup>Other Cretaceous epochs.</sup> Neighboring regions also went down, and the bay was converted into a broad, deep ocean in which mud slowly gathered for a very long period. There was one short interval when sand came instead of mud, and there were two epochs during which limy ooze alternated with clayey mud, but these changes were not accompanied by any local warplings. The floor of the ocean remained smooth and even in the Pueblo district, and the deposits were gathered in flat, uniform layers. Afterward the mud was converted into shale, the limy ooze into limestone, and the sand into sandstone.

These conditions, including the advancing coast-line with its rapid accumulation of sand and the broad, deep ocean with its slow accumulation of mud, belong to the Cretaceous period. The state of the district at the close of that period is shown in fig. 6, where the heavy black line represents the porous sandstone; the shales above are indicated by broken lines, and the principal limestone bed is represented by a notation of blocks.

The whole quadrangle, together with a much larger territory, was now lifted above the ocean and also warped and deformed. A great lifting of the mountains took place at the same time, and this affected those <sup>Eocene and Neocene periods.</sup> parts of the district which belong to the belt of foothills. A number of wave-like ridges were also produced, and in some places the strata were broken across and faulted. Fig. 7 illustrates the general character of these changes, but it is misleading in that it implies that the formations remained intact while they were deformed. In fact, the great mountain indicated at the left of that diagram never existed, for the changes of form were slow, and as soon as the sea-bed became land and steep slopes were produced rain and streams began their work of erosion. Uplift and erosion thus went on together, and the actual height of the land at any time represented only the difference between these two modifying factors. Fig. 8 exhibits the same internal structure as fig. 7, but after the removal of large portions of the various formations by erosion. It corresponds in a general way to the actual condition of the district at the present time. The time during which this deformation and erosion took place belongs chiefly to the Eocene period but partly also to the Neocene. The work of erosion is still in progress. Every storm that beats on the surface of the district washes a quantity of mud and sand into the creeks, and the turbid creeks carry it to the Arkansas River. The Arkansas, though already loaded with detritus by the storms in the mountains, receives the tribute of the creeks and sweeps it onward toward the ocean.

This is the process by which the land was eroded during the great uplifting, and it has been nearly continuous from that time to the present. Still there have been local interruptions, and one of these belongs to the history of the district. There was an epoch in the Neocene period when the eastward slope of the plains was less than now, so that the Arkansas and such of its branches as headed in the mountains were too sluggish to carry onward the sands that were given them by the mountain storms. The slackened current washed along the mud but dropped some of the sand, so as to build up its bed. There was thus produced on the plain a formation of sand without the intervention of any body of standing water, either lake or ocean. This deposit was not deep and did not cover the

entire surface of the district. It was spread only over the parts which were then lowest and it rested on the worn surfaces of various older formations (fig. 9). It was never hardened into firm rock, but retained the condition of loose sand. When changes of height and changes of slope gave greater speed to the rivers and creeks, so as to renew the erosion of the plain, the soft sands were easily attacked, and little is left of them at the present time. The stream beds, and even the general surface of the country, are worn several hundred feet below their plane, and such remnants of the sands as survive are the caps of mesas. Such a mesa is shown at the right in fig. 10.

As a result of this long and complex history the rocks of the district are broad plates or strata, of great horizontal extent and comparatively small thickness. For the most part they are parallel and lie one upon another like the leaves of a book. They are not flat, but are warped in various ways. They differ in horizontal extent chiefly because the upper <sup>Present condition.</sup> and newer have been more widely eroded than the lower and older. The breadth of any particular formation at the surface, or, in other words, the width of its outcrop, depends partly on the extent to which the formation has been removed and partly on the extent to which the next overlying formation has been removed. If a deep well be drilled from any point of the surface the various formations will be penetrated in a definite order, but the only formations reached will be those which have not been eroded from that place, and the depth at which each will be found will also depend on the amount of local erosion.

#### THE ROCKS.

In this section the various formations are described in the order of their origin, beginning with the oldest. They are classed, according to the periods in which they were formed, as Archean, Silurian, Carboniferous, Juratrias, Cretaceous, Neocene, and Pleistocene. The areas they occupy within the quadrangle are marked on the Historical Geology sheet, and their sequence and relative thicknesses are graphically shown on the Columnar Section sheet.

#### ARCHEAN.

In the southwest part of the quadrangle are two small tracts of crystalline schists to which no formation name has been given. The more abundant kinds of rock are mica-schist and mica-gneiss, colored gray and pink, and a pink variety of granite. The schist and gneiss exhibit schistose structure, the plates of mica lying nearly parallel, so that the mass splits most readily in one direction. The plane of this direction varies in course or "trend" from north to northwest, and is nearly vertical. The granite occurs in large bodies of irregular form, and also in veins penetrating other rocks. The character and relations of the veins show that they were injected in liquid <sup>Schists and granite.</sup> condition after the making of the other rocks, and it is thus known that the granite is of igneous origin. The origin of the other rocks was not determined. The larger of the two tracts occupies the southwest corner of the quadrangle; the smaller lies west of Beulah. Their total area is about 7 square miles. Wherever younger formations are seen to rest on the Archean rocks they are unconformable.

The southwestern tract is an upland, rising 800 to 1500 feet above the adjacent lowland, and is deeply trenched by the canyons of the St. Charles River and its tributaries, so as to have a decidedly rugged and mountainous character. Its higher parts, however, are comparatively <sup>Pre-Silurian topography.</sup> smooth, and the distant view from a favorable point shows that they are regularly related to one another. They are remnants of an earlier plain, originally level but now inclined toward the east and northeast. This plain was at one time the upper limit of the Archean mass, and the modern canyons, some of which are nearly 1000 feet deep, have since been carved from that mass by the streams which occupy them. The plain of the uplands is not itself smooth, though in places it is sufficiently level for cultivation. The different rocks have weathered away unequally, and the granite, resisting erosion better than the others, projects above the general plain in rounded knobs. As will be explained in subsequent paragraphs, the plain was originally carved from the

Archean mass in the early part of the Juratrias period, and was then nearly horizontal. It was afterward buried beneath the Morrison formation and other deposits having a total thickness of thousands of feet. While still buried it was warped and tilted along with the overlying strata, and it has since been denuded or resurrected by the washing away of its cover.

#### SILURIAN.

*Harding sandstone.*—The oldest sedimentary rock of the quadrangle is a white, sugar-like sandstone, 30 feet thick. It appears only in two small tracts west of Beulah, where it rests on the Archean schists. The best exposures are in the walls of the canyon of the north branch of the St. Charles River.

#### CARBONIFEROUS.

*Millsap limestone.*—Above the Harding sandstone are 200 feet of gray and purple limestones, with some shale, especially in the lower part, and on these are 30 feet of coarse gray and red sandstone. Like the Harding, they occupy two small areas west of Beulah, separated by the St. Charles Canyon. It is evident that strata originally continuous were divided by the erosion of the canyon, and they are, in fact, still united a little west of the boundary of the quadrangle. In the northern area, where they crown a ridge of Archean schists, they are much warped, so as to dip in various directions, but their general slope is toward the south. In the southern area they lie lower, and there are several mesas of limited extent capped by a massive bed of limestone. These mesas are not flat, but have warped surfaces which seem to copy the warping of the limestone. <sup>Structure of the older rocks.</sup>

The Millsap beds are parallel and conformable to the Harding below, but their relation to higher beds does not appear, as none rest on them. At the southeast they adjoin the Fountain formation, from which they are separated by a fault.

#### JURATRIAS.

*Fountain formation.*—Two formations represent this period, the Fountain and the Morrison. The Fountain consists chiefly of coarse, deep-red sandstones containing a considerable admixture of clay. The greater number of sand grains are of the mineral feldspar, and the sandstone is consequently of the variety called arkose. In the upper half of the series are many beds of red and chocolate-brown shale; in the lower part are conglomerates. The lowest bed seen is a coarse conglomerate, containing pebbles and boulders of gneiss, schist, and granite, similar to those of the adjacent Archean. The thickness, as measured 2 miles north of Beulah, is 2100 feet. The top of the series is not there seen, but the missing beds are probably thin.

The original relation of the formation to the older rocks was not directly observed, the line of contact being concealed except at a few points where a fault intervenes. The Archean boulders in the lower conglomerate indicate that when the conglomerate was formed there was a neighboring land area occupied by Archean rocks, and this idea is supported by the great abundance of feldspar in the sandstone, for feldspar is an abundant mineral in all the Archean rocks. It is believed that the resurrected plain appearing in the southwest corner of the quadrangle was shaped during the Fountain epoch, the pebbles, sand, and clay which resulted from the wearing-down of the schistose land mass being deposited in a contiguous sea as the Fountain formation. <sup>Gravels of a Juratrias coast.</sup>

The area occupied by these rocks is a belt from 1 to 2 miles broad, extending 3 miles southwestward from Beulah, where it ends against the Archean upland, and 7 miles northward to the valley of Red Creek, where it passes into the Canyon quadrangle at the west. The strata dip eastward at angles varying from 10 to 20 degrees, and pass under the Dakota sandstone, which there forms the crest and eastern slope of a high ridge or "hogback." Opposing feeble resistance to erosion than do the schists and limestones at the west or the firm sandstones at the east, the Fountain beds have been <sup>The valley behind the hogbacks.</sup> more extensively removed and their outcrop constitutes a valley. The southern part of this valley is drained by South Creek and North Creek, tributary to the St. Charles River, and

the northern part by Red Creek and its branches. Near Beulah the slopes are gentle and there are smooth, soil-covered terraces suited for agriculture; but farther north is a labyrinth of small ridges and canyons whose bare sides afford little foothold even to grasses and give a characteristic redness to the scenery.

*Morrison formation.*—The Morrison beds, though broadly exposed in the adjacent quadrangles to the north and northwest, are represented in the Pueblo only by three small areas. One adjoins the northern boundary 3 or 4 miles west of Turkey Creek and is continuous with a much larger tract in the Colorado Springs quadrangle. The beds exposed include only the upper part of the formation. They are chiefly clays or shales of brilliant hues, the lower 200 feet being mostly white and the upper 100 chocolate and green; near the base are several beds of gray and white gypsum; above the middle is a yellowish sandstone, and toward the top are several thin beds of limestone. In the second area, near the northwest corner of the quadrangle, the strata are poorly exposed. <sup>Variegated clays and gypsum.</sup> The third area, near the southwest corner, is a narrow belt crossing the western slope of Hogback Mountain at a high level, and then turning southeast and following the base of the Archean upland. The part exposed on Hogback Mountain consists chiefly of red shale, but contains a few layers of hard red sandstone. Its total thickness is about 70 feet. The part bordering the Archean area is paler, white and orange predominating, and none of the harder strata were observed. The formation there dips steeply to the northeast, and its apparent thickness changes rapidly from point to point by reason of faulting. At one point it is seen to rise with diminishing dip toward the back of the Archean upland, and on the continuation of that upland in the Walsenburg quadrangle at the south are outlying remnants of the Morrison shale protected by caps of Dakota sandstone. The Morrison shale was the first formation spread over the old land-surface from which <sup>The Morrison overlies the Fountain.</sup> the material of the Fountain formation had been eroded. At the end of the Fountain epoch the area of submergence was enlarged so as to include much that had before been land, and the deposits of the Morrison sea accumulated not only on the newly gathered Fountain sediments but on the ancient Archean rocks.

As the Morrison beds are weaker than all their neighbors, they have yielded readily to erosive attack and their outcrops are marked by valleys; but their extent is so small that these valleys belong only to the minor features of the topography.

#### CRETACEOUS.

The outcrops of the formations thus far described occupy about one-fiftieth of the quadrangle; the Cretaceous formations constitute nine-tenths of the entire surface. From an economic point of view the relative importance of the Cretaceous strata is equally pronounced, for they include the limestones, building-stones, fire-clays, and artesian waters of the district. The order of the Cretaceous formations, beginning with the lowest, is: Dakota, <sup>Importance of the Cretaceous.</sup> Graneros, Greenhorn, Carlile, Niobrara, Pierre. The Dakota consists chiefly of sandstone; all the others chiefly of shale. In the Greenhorn is some limestone, in the Carlile a little sandstone, and in the Niobrara an important body of limestone. The broadest outcrop of the Dakota formation is in the southwest part of the quadrangle, and the Pierre occurs at the northeast. The intermediate members constitute a great belt extending from northwest to southeast, but their individual boundaries are irregular and sinuous, and there are many insular tracts. Some of these islands are *outliers*, or remnants cut off from the main belts by the erosion of intervening parts, and these appear in the landscape as buttes or mesas; others are *inliers*, or limited tracts from which overlying beds have been eroded, and these occur usually in lowlands. The outlier of a formation is completely surrounded by outcrops of the next lower formation; the inlier by outcrops of the next higher formation. <sup>Outliers and inliers.</sup>

*Dakota sandstone.*—Mention has already been made of the Dakota sandstone as a porous rock whose sand was washed clean by waves in the process of formation. It is now necessary to



qualify this statement by describing the formation more in detail. Wherever its full section is exposed to view it is found to consist chiefly of a series of thick sandstone beds separated by comparatively thin shale beds. The sandstone changes rapidly in thickness from point to point, so that no two measurements at different places show close agreement, and when comparison is made between localities as much as 10 miles apart it is usually impossible to recognize the identity of individual beds. At base the formation is sharply limited by a surface of unconformity. Usually it rests on Morrison shales, but in the vicinity of Beulah, and thence northward to Red Creek, it rests on the Fountain sandstone, the Morrison formation having disappeared by erosion before the Dakota was deposited. At top it passes into the Graneros shale without any sharp line of separation, the shaly members of the Dakota becoming gradually more numerous and the sandy members thinner until the latter cease altogether. The highest sandstone is usually a single layer of dense, brittle rock having a vertical fracture. It is in the upper part that shales occur of the peculiar quality necessary for use as fire-clays.

The grains composing the Dakota sandstone are chiefly of quartz, and the rock is thus contrasted with the neighboring Fountain sandstone, in which feldspar predominates. It is further contrasted by its color, which is ordinarily light-gray, pink, or white, weathering at the surface to various shades of yellow, orange, and brown. Some of the lower members are locally so coarse-grained as to merit the name conglomerate. The larger pebbles are of quartzite. There are also white grains of some soft material, which may be a kaolin resulting from the decomposition of feldspar. The fractured surface often has a speckled appearance, flecks of yellowish brown dotting a gray ground. The different beds show considerable difference in porosity, and doubtless the same bed varies from place to place. The highest sandstones are not so porous as to convey artesian water, but the heavier beds of the middle and lower parts of the formation have usually an open texture. The greatest measured thickness, near Beulah, is 650 feet, and nearly the whole of this is sandstone. In the northwestern and southeastern parts of the quadrangle the thickness is from 300 to 350 feet, and the sandstone is more interrupted by beds of shale. Elsewhere the formation has been found to contain leaves of plants in great abundance and variety, and also shells indicative of brackish water. The only fossils discovered in this district were a few leaves and the trails of undetermined animals.

The formation underlies nearly the whole of the quadrangle. It is exposed at the surface over a large area in the southwest part and in other important areas near the northwest and southeast corners. There are also three small inliers: one on the Arkansas River at Rock Canyon, where the formation is trenched by the river for a few rods; another east of Greenhorn Creek, not far from the junction of its principal branches; the third, east of the eastern branch of Greenhorn Creek, not far from the south boundary of the quadrangle.

In relation to the forces of erosion the Dakota is the most resistant formation of the district, with the possible exception of the Archean granite. Where uplift has given it a steep dip it rises above the weaker rocks by which it is surrounded in a bold ridge, ordinarily called a "hogback," and the Dakota hogbacks are among the most important features of the foothill belt along the base of the Rocky Mountains. In this quadrangle a well-characterized hogback forms the eastern wall of the Beulah and Red Creek valleys, and there is another, less characteristic, near the northwest corner.

**Graneros shale.**—Resting conformably on the Dakota sandstone is the Graneros shale, from 200 to 220 feet thick. Its color includes various shades of bluish gray, being lightest in the lower part and darkest near the middle of the mass. At most localities several thin beds of white clay were seen, but it is not known whether these are continuous for great distances. Some of them include crystals of selenite, and one occurs in immediate contact with a thin limestone layer

wholly composed of oyster shells. Oyster beds were observed at several other localities, but they are probably very local features. The most persistent of the contained hard beds is a calcareous sandstone, 1 or 2 inches thick, found about 50 feet below the top of the shale. It contains fossil shells of several kinds. In the southern part of the quadrangle a line of concretions is usually found about 30 feet above the base of the formation. They are from 6 to 12 inches thick, calcareous, and often fossiliferous. The shale passes so gradually into the Greenhorn formation above that the line of separation had to be arbitrarily drawn.

The outcrop is a belt of moderate width, very crooked and irregular, following the margin of the Dakota sandstone. The associated topographic forms include valleys, lowlands, and long slopes descending from terraces of Graneros limestone to basements of Dakota sandstone. The formation is notably infertile, so that its slopes, unless overwashed by debris from other beds, are nearly destitute of vegetation.

**Greenhorn limestone.**—In the Greenhorn formation limestone beds from 3 to 12 inches in thickness alternate with shale beds 10 to 20 inches thick. The shale factor is thus, in one sense, the more important, but the limestone ledges resist decay so strongly that their fragments usually cover the surface of the outcrop, concealing the shale from view and giving the impression of a thick sheet of limestone. The limestone is pale-blue and of fine texture. Most of the layers are divided by vertical cracks into smooth, flat flakes, from one-fourth inch to 2 inches thick. Some of them have abundant fossil shells, especially a thin form, of oval outline, marked with concentric waves or ridges (*Inoceramus labiatus*). The shale is bluish-gray and darker than the limestone. It contains the same shells in abundance, but so poorly preserved as to escape casual observation. There are also thin layers of white clay. The thickness of the whole series is from 35 to 50 feet. The Illustrations sheet contains a typical view of the limestone and drawings of its characteristic fossil.

The Greenhorn outcrop is a narrow belt, usually but a few hundred feet across, but occasionally expanding to a half mile. Its course is winding and there are many outliers and inliers. Occasionally the limestone caps a hill or forms the crest of a ridge, but usually it constitutes a terrace interrupting the slope from a cliff of Niobrara limestone above to a valley of Graneros shale below. The shattered limestone stores water better than the adjacent shales, thus favoring the growth of trees, and its zone is usually marked by a belt of junipers and piñons.

**Carlile shale.**—The Carlile formation is from 180 to 210 feet thick, and consists chiefly of argillaceous shale. The lower 50 feet are medium-gray; then come 25 feet of dark-gray, including bands that are nearly black, and above the color is medium-gray. At 50 or 60 feet below the top of the formation the shale becomes sandy, and within the sandy part are lenses of friable sandstone. Many localities show from 10 to 20 feet of yellow sandstone at the very top, and in the southwest part of the quadrangle there is a bed of sandstone 40 feet below the top. The sandy shale contains also many concretions, which are more or less globular and range in diameter from 1 foot to 5 or 6 feet. Within the larger are cavities in the form of ramifying cracks, and these have been partly or wholly filled by white and wine-colored crystals of calcite.

The outcrops of the shale are irregular in plan and comprise a large number of inliers. The greatest development is at the south and southeast. There are considerable areas in the valleys of Rock and Pecks creeks; exposures follow the banks of the Arkansas River from Goodnight to Beaver; and there are several lines of outcrop between Beaver Creek and Wild Horse Park. The sandstone at top usually unites with the Niobrara limestone resting on it in the formation of a cliff, and the shale below either constitutes a steep slope under this cliff or a valley between two facing cliffs.

**Niobrara formation.**—The Niobrara formation consists chiefly of shale and has a thickness of 600 or 700 feet, but is parted from shales above and below by limestones. Considering its ele-

ments in detail, the lowest stratum is a peculiar bed, from 1 foot to 2 feet thick, intermediate in character between limestone and sandstone. Its original color is dark-gray, but its weathered surface is yellowish-brown. It contains many teeth of sharks and other fishes, a few fossil shells, and numerous dark pebbles a half inch in diameter. The presence of the pebbles is noteworthy, as all the rocks for hundreds of feet above and below are composed of fine material. The bed is remarkably persistent, having been found through a broad territory.

The next element is a limestone about 50 feet thick, consisting of strata ordinarily 1 or 2 feet thick, separated by shale layers 1 or 2 inches thick. Where its interior is exposed by quarrying the limestone is seen to have a pale-blue or gray color, but the natural surface is nearly white and somewhat chalky in texture. It breaks under the action of the weather into rough flakes whose longer dimensions are horizontal, and this character serves to distinguish it readily from the Greenhorn limestone, which is split into plates by vertical cracks. It contains fossil remains of various kinds, the most common and characteristic being a rotund bivalve shell (*Inoceramus deformis*; see Illustrations sheet) from 5 to 10 inches across. The shell itself is not often seen complete, but molds of its interior are commonly found and attract attention from their resemblance to the hoof of a horse. At top the limestone passes gradually into calcareous shale, which is more than 100 feet thick and often contains beds of limestone. Thin limestone layers occurring near its top are white or cream-colored, and contain fossil shells, especially a small oyster attached to fragments of a larger shell (See Illustrations sheet). Above this is the body of the formation, a medium-gray shale, which often splits under the action of the weather into paper-like layers. The surfaces of these layers are roughened by minute white crystals of selenite, and the same mineral often occurs in thin veins crossing the rocks in various directions. Fish-scales from a half-inch to an inch in diameter are so abundant that a few minutes' search will usually discover them in the unweathered rock. At the top of the formation are 10 to 20 feet of calcareous shales, including one or two layers of impure limestone.

The calcareous shales are characterized by a somewhat regular variation of texture and hardness, the amount of calcareous material becoming alternately greater and less. These alternating phases resist the weather unequally, so that where the rock is exposed in a cliff its face is barred across by obscure ribs. The space from rib to rib is usually 18 to 24 inches.

The outcrop of the Niobrara is broader than that of any other formation of the quadrangle. It occupies nearly one-half of the entire area. Its belt runs from northwest to southeast, having a simple boundary on the east and a very irregular one on the west. There are also at the west many outliers, usually containing only the basal limestone.

The shale mass appears in the landscape only as a plain of gentle slope. The upper limestone caps a series of small ridges running from Pueblo west-northwest up the valley of Dry Creek. The lower calcareous shales are usually masked in the general plain, but are occasionally betrayed in cliffs. The lower limestone forms the back or top of many mesas and inclined tables, and usually ends in a cliff overlooking exposures of the Carlile shale.

**Pierre formation.**—Above the Niobrara is a still greater formation composed almost entirely of shale. A thickness of 2200 feet appears in the district, but the top is not seen. Different parts of this shale mass exhibit diverse characters, but the gradation from one to another is so complete that it was not found practicable to divide the formation into distinct parts. Still, there is advantage in recognizing a series of zones, even though their boundaries and precise thickness can not be indicated. The Barren zone, so-called on account of the rarity of its fossil remains, lies at the bottom of the series and is 400 to 500 feet thick. It is of bluish-gray color, and its lower part resembles the Niobrara shale in the tendency to divide into papery layers, rough from the crystallization of selenite. The Rusty zone, 600 feet thick, is also bluish-gray in

color but is comparatively free from gypsum. It contains many concretions composed of lime carbonate and iron carbonate, and these are of oval form, measuring, usually, from 1 foot to 2 feet across. Their material is originally dark-gray, but under the action of the weather turns a rusty brown, and the soil derived from the formation is usually so strewn with their angular fragments as to appear reddish-brown. The Baculite zone, 100 to 200 feet thick, is pale-gray, and is so called from the abundance of a fossil shell of that name. The shell has the form of a flattened cylinder, tapering slightly toward one end, and is usually a half-inch to an inch in diameter and 4 or 5 inches long. The Tepee zone, 1000 feet thick, includes the upper part of the formation as exhibited in this quadrangle. It is pale-gray, and contains numerous oval concretions ranging from 1 foot to 4 or 5 feet in diameter. In these concretions are many fossil shells, including several large forms which are also conspicuous by reason of the pearly luster of their white or gray walls. There are also large masses of rough gray limestone, called *tepee cores*, and these, though not abundant, constitute a peculiar and striking feature. They are from 10 to 30 feet in horizontal diameter and extend vertically through the shale to unknown distances, probably 50 or 100 feet. In general form they are cylindrical, but their surfaces are in detail quite irregular. They also contain numerous fossil shells, especially a small bivalve (*Lucina*).

The principal Pierre area is triangular, occupying the northeast corner of the quadrangle, and there is a single outlying tract extending southeast from near Bessemer Junction. The total area is not far from 120 square miles, and exceeds that of any other formation excepting the Niobrara. The surface is in general a plain of gentle inclination, but about the Baculite Mesa and in some other localities there are steep slopes. These are furrowed by numerous gullies, so as to have the character of bad lands. Where the tepee cores occur their positions are marked by steep-sided conical buttes 20 to 50 feet high. The limestone of the core constitutes the apex of the butte, and its fragments sheathe the sides. The resemblance of these buttes to the conical lodges or tepees of northern Indians suggested their name. Figures of Pierre fossils and a sketch of a tepee butte may be found in the Illustrations sheet.

#### NEOCENE.

**Nussbaum formation.**—The beds composing this formation are chiefly of sand, and have an extreme thickness of 100 feet. At a few points the sand is overlain by a fine silt. At various levels it contains pebbles, and near the bottom the pebbles are usually so numerous as to constitute gravel. All these beds are *alluvial*, having been spread where they lie by the flowing water of streams. They have never been submerged under a sea or lake, and as a rule the particles do not cohere, but lie loose, just as when first deposited. The lowest part, including a thickness of 2 or 3 feet, is in a number of places bound together by calcareous or ferruginous cement, so as to constitute a rock of some firmness; and as this part was originally gravel the resulting rock is a conglomerate.

The Nussbaum is not conformable with any underlying formation, but rests on eroded surfaces of the Pierre, Niobrara, and other Cretaceous strata. The greater part of it has been eroded, so that the existing beds are merely remnants. Usually they cap buttes and mesas and lie at a considerable height above the surrounding country (fig. 11). The tops of the mesas slope in various easterly directions, conforming in a general way to the modern drainage, and there can be little question that these slopes show the direction of the streams by which the formation was made.

The largest and thickest body is on Baculite Mesa. A second body occupies the mesa north of Blue Hill, and there are others between that and Turkey Creek. A group of small remnants occurs on the upland between the headwaters of Pecks and Rush creeks, and several are found on the St. Charles and Greenhorn a few miles above their junction.

Variability of the Dakota sandstone.

Graneros shale and concretions.

Fish-tooth bed in the Niobrara.

Niobrara limestones and shells.

Tepee cores in the Pierre shale.

Texture, color, and porosity.

Greenhorn limestone and fossils.

Shales with fish-scales in the Niobrara.

Thickness of the Dakota sandstone.

Areas of Pierre shales.

Extent of the Dakota sandstone.

Carlile shale and sandstone.

Limestone mesas.

Gravel and sand of the higher mesas.

Dakota hogbacks.

Zones of the Pierre shales.

Baculite Mesa and Blue Hill.



PLEISTOCENE.

Other alluvial deposits are found at lower levels in all the principal valleys. Usually they constitute terraces overlooking the streams, and these are often in series, one above another (fig. 11). The lowest of all is the deposit in which the stream now flows and which it modifies from year to year in time of flood. Each terrace was, in fact, once the flood-plain or bottom land of the stream, and their arrangement in steps is a record of the gradual deepening of the valley by erosion. Their order of position is also their order of age, but in an inverse way, as the lowest is the latest. They thus belong to several epochs which might perhaps be distinguished, but in mapping only two colors or patterns have been used, the one indicating the modern alluvium and the other all the earlier deposits.



Fig. 11.—Ideal section showing the arrangement of alluvial deposits.

K, eroded surface of Cretaceous shale; Nn, remnant of Nussbaum beds, capping high mesas; Pe, remnants of earlier Pleistocene beds, capping terraces; Pl, later Pleistocene beds, making flood-plain.

The material as originally deposited includes gravel at bottom with sand and silt above, but the finer beds have been washed away from all the narrower terraces, leaving only gravel; and in the modern alluvium, along the streams, the gravel is often wholly concealed by the overlying sand. As all the material has been brought to its present position by the streams, and as different streams head in regions occupied by different rocks, the composition of the deposits varies from valley to valley. This is specially noticeable when the pebbles are compared. For example: the gravel in the bed of Fountain Creek, as well as in the bordering terraces, consists chiefly of gray quartzite and pink granite, while the gravel of the Arkansas and its terraces includes a wide range of rocks, among which andesites are conspicuous.

The Pleistocene deposits, like the Neocene, rest unconformably on all the Cretaceous formations.

FORMATION NAMES, ETC.

In describing the geology of the district the various formations must frequently be mentioned, and it is important that each have a name. For this purpose it matters little what particular name is used; but as soon as attention is extended to other districts questions of consistency arise. All formations of the Pueblo quadrangle occur also elsewhere, and so far as their identity can be established the same names should be used. Much care has therefore been taken in the selection of the names employed in this folio; but as their use is not in all cases free from doubt, and as some of the names are new, it seems proper to place on record certain qualifications and explanations.

The reference of the schists and granites to the Archean period is provisional only. The rocks are broadly exposed in the adjacent district at the west, and when they shall have been there studied it is possible that some other classification will be found better.

The names Harding, Millsap, Fountain, and Morrison are all derived from the Pikes Peak folio, where they were introduced and defined by Dr. Whitman Cross. The Pikes Peak quadrangle lies northwest of the Pueblo, touching it at one corner, and the continuity of the various formations has not yet been established by direct tracing. The sandstone here named Harding contains no fossils and is connected with the Harding sandstone of the Pikes Peak quadrangle only through similarity of physical character and relation.

In a bed near the middle of the overlying limestone were found a few fossil shells of *Spirifera rockymontana*, a Carboniferous species found also in the Millsap limestone of the Pikes Peak quadrangle, but nothing was found to mark the presence of the Fremont limestone (Silurian), which in the Pikes Peak district separates the Harding and Millsap formations. The application of the name Millsap to the whole limestone series of the

Pueblo district is a somewhat arbitrary procedure and is subject to correction when more facts are available.

There is yet another doubt as to the name Millsap. The fact has recently been brought to the attention of Dr. Cross and myself that Mr. W. F. Cummins applied the same name to a formation in Texas in 1891, but ignored it and apparently abandoned it in 1893. Should the name be retained for the Texas formation the rule of priority will require the substitution of another name for the formation occurring in the Pikes Peak-Pueblo region.

In each district the Fountain formation lacks fossils, and the determination of its period depends wholly on physical characters. These are inconclusive, and the time relations of the formation are in doubt. By Cross it has been provisionally placed in the Carboniferous; by the writer, in the Juratrias.

When the geology of this region was first studied by Dr. F. V. Hayden, in 1868, the beds lying between the Dakota and Niobrara formations were called the "Fort Benton formation." They comprise a series of shales about 450 feet thick, interrupted near the middle by a number of limestone layers. The limestone strata have so important an influence on the topography and afford so much aid in the study of artesian problems that it seemed best, in connection with the present work, to give them a special name, and this led to the separate designation of the shale beds above and below them. The name Graneros, applied to the lower shale, was suggested by Mr. R. C. Hills, and is derived from a village and creek a short distance south of the district. The name Greenhorn, applied to the limestone, is derived from Greenhorn Creek and Greenhorn station. The name Carlile, applied to the upper shale, is derived from Carlile Spring and Carlile station on the Arkansas River. At the localities to which the names refer the several formations are well exposed for study.

In later publications by the survey under Dr. Hayden's direction, the Niobrara and Fort Benton were united under the name "Colorado Group," and this usage has been largely followed. In the atlas of Colorado published by the Hayden Survey the color representing the Colorado group is erroneously made to include a large tract now known to be occupied by the Pierre shale.

The name Nussbaum is here used for the first time, being derived from Nussbaum Spring, which flows from the Nussbaum sands near the south end of Baculite Mesa.

The diversities of usage here mentioned, as well as other discrepancies to be found in the literature of the region, are arranged in tabular form at the bottom of the sheet of columnar sections.

STRUCTURE OF THE ROCKS.

The strata of sediment formed on the bottoms of lakes and oceans are nearly level. Steep dips of strata result from uplift or other disturbance by underground forces. The process by which level strata are transformed into dipping strata is called *deformation*. The resulting forms include *arches* and *domes*, which are convex upward, and *troughs* and *basins*, which are concave upward. Associated with these are faults.

As already stated, the strata of the district were deformed at three epochs: first, after the making of the Millsap limestone; second, after the Morrison shale was formed; third, after the Pierre shale had accumulated. The older formations were affected by two or three of these disturbances; the Cretaceous formations—Dakota to Pierre—only by the last. Thus the oldest are most deformed, but as the area of their exposure within the quadrangle is small little is known of their structure. The Cretaceous formations are so widely exposed to view that their structure is comparatively well determined.

THE DEFORMATION SHEET.

The dips, arches, and troughs given to the Cretaceous strata by deformation are exhibited graphically in the Structure Section sheet and the Deformation sheet. The uppermost section on the Structure Section sheet shows the structure of the formations along the line A A. If a single

rock bed be traced from side to side it will be seen to exhibit several flexures and to be dislocated at two points by faults. Take, for example, the Dakota sandstone (Kd), which lies at the surface near Red Creek and also to the left of White Butte. Between these points it passes beneath the surface, being flexed down into a trough, and various other formations lie above it. To the right, also, it passes down beneath other formations, and although somewhat flexed in detail, descends so rapidly that at the margin of the quadrangle it is nearer to sea-level than to the surface of the land. It is, in fact, about 3000 feet underground. The other sections exhibit similar facts along their respective lines. In the Deformation sheet a different mode of representation is employed. Let us imagine that all the rocks lying above the Dakota sandstone are dug away, so as to lay bare the surface of that formation. Let us suppose, further, the Dakota sandstone to be restored in those small portions of the quadrangle from which erosion has removed it. There will result from this denudation and reconstruction an uneven surface exhibiting the shapes given to the Dakota sandstone by the processes of deformation. The work thus imagined is altogether too stupendous to be actually accomplished, but our knowledge of the thickness and extent of the overlying formations and of other geologic facts makes it possible to determine with considerable accuracy many details of the form which would result. To represent this form a model has been constructed on the same scale as that of the map. A photograph was made of the model, and the engraving constituting the Deformation sheet was prepared from that photograph. The arches and domes of the deformed surface there appear as ridges and mounds, the troughs as valleys, and the faults as cliffs.

When one looks at the photograph of such a model it is important that the light illuminating the photograph come from the same side as the light which illuminated the model when the picture was made. If these relations are reversed the hills may appear as hollows and the hollows as hills. The general effect is also more easily obtained at a distance of a few yards than if the sheet is held in the hand. In the making of this photograph the illumination was from the right, as shown by the shadow of the object placed on the model, and the picture should therefore be so held as to be illuminated from the right.

STRUCTURE OF THE CRETACEOUS ROCKS.

In the central part of the quadrangle and extending thence toward the northwest is a neutral tract where the deformation is only of moderate amount. At the north and northwest, at the southwest, and at the southeast are uplifts which encroach on the quadrangle but lie principally beyond its boundaries. Westward, in the vicinity of the Arkansas River, the strata descend toward a large basin in which lie the oil-bearing rocks of the Florence region. Eastward and northeastward they descend more rapidly toward a still greater basin.

The northern and northwestern uplifts.—The Front Range of the Rocky Mountains, lying to the north-northwest of the district, is itself a great uplift, and much of its margin is fringed by a series of smaller uplifts which appear as arches in the strata of the Plains region. Two of these arches enter the Pueblo quadrangle, pitching down toward the south and losing themselves in the neutral tract. The more westerly is approximately bounded by Beaver Creek and Pierce Gulch; the more easterly is traversed by Turkey Creek and is impressed on the landscape as a high ridge of Dakota sandstone. From the latter project two minor arches, one running southward to Pumpkin Hollow, and the other including Wild Horse Park and a cedar-covered ridge to the east and south of it.

The Beaver arch has gentle slopes, the dips to the southwest and southeast ranging from 1 foot in 40 to 1 in 15. Near its west margin it is crossed by a fault. The Turkey arch is 1000 feet high. It has gentle dips to the east and steeper ones to the southwest, the latter averaging 1 foot in 5.

The Pumpkin arch resembles the Turkey in form but is only one-third as high. The Wild Horse arch is almost independent of the Turkey, joining it at a single point. It has the form of an inverted canoe, 2½ miles wide, 6 or 7 miles long, and 500 feet high. Section AA of the Structure Section sheet crosses three of these arches.

The southwestern uplifts.—The southwest corner of the quadrangle is occupied by the northern part of a high arch, the St. Charles, which runs east as a spur from the great uplift of Greenhorn Mountain.

The top of the arch is comparatively flat, dipping eastward at 1 foot in 10, but its northeastern side is steep and in places vertical. The strata are not only turned sharply down along its edge, but are faulted in a complex way, so that the outcrop does not show their full thickness. At the south end of Hogback Mountain this zone of steep slope branches, one part continuing west and the other turning sharply to the north and curving around the east end of the Beulah arch, which adjoins the St. Charles. The form of the Beulah arch is revealed only in part, as the Cretaceous formations do not occur west of the Dakota hogback. It is a spur of the Wet Mountain uplift and merges with a broader spur, the Red arch, which lies just north and is outlined by the Dakota hogback. The height of the steep northeast face of the St. Charles arch is 1800 feet. The east face of the Beulah arch rises 1400 feet in the first mile, and probably continues with gentler dips. The general dip of the Red arch is about 1 foot in 10, and its height within the district is 1800 feet.

The curved bases of the St. Charles and Beulah arches are followed by a narrow trough, the Three-R trough, which separates them from a low dome somewhat triangular in form. The general depth of the trough is about 400 feet.

From the St. Charles, Beulah, and Red arches there is a general descent east-northeast to the neutral tract, the prevailing dip being 1 foot in 30. Its continuity is broken by a series of parallel faults which cross it at right angles. These are clearly shown on the Deformation sheet and do not require individual mention. In some the eastern side was uplifted; in others it was dropped. The greatest dislocation is but little more than 200 feet, and in some faults it is barely 100 feet. They are minor features as compared to the great fault about the St. Charles arch, but have important influence on the configuration of the surface, as will be explained later. Several of them are represented in sections CC and DD of the Structure Section sheet.

The general northeastward descent from the Red arch is also traversed by a flexure running nearly at right angles to the fault system. This flexure resembles a fault in that it forms the boundary between two great bodies of strata which are at different levels. The strata of the southeastern body are flexed downward so as to remain continuous with those of the northwestern body. This is the Rush flexure, and the amount of its displacement ranges from 200 to 600 feet.

The southeastern uplift.—The Walsenburg and Apishapa quadrangles, lying south and southeast of the Pueblo, include a broad dome of strata; and a spur from this dome, the Carlos arch, enters the Pueblo quadrangle near its southeast corner. Its general height is 400 feet, and it is unsymmetric, being steepest on the west side. In common with the general rock slope on which it rests, it rises toward the south. Six miles south of San Carlos it is sharply interrupted or indented by a lozenge-shaped basin, a definitely limited block of strata, about 1½ miles across, having gone down instead of up.

Arch and basins of the neutral tract.—In a general way the rocks rise gradually from the center of the district toward the north and southwest, so that the middle part might be called a broad, shallow trough. Athwart this trough rises a low arch, the Rock Canyon arch. It is about 10 miles long, north and south, and from 3 to 4 miles broad. Its crest line is uniformly about 350 feet above its base, but, sharing the general curvature of the central trough, is higher near the ends than in the middle. The character of the arch is well shown at Rock Canyon, where the Arkansas River cuts across it, but both ends are obscured by surficial deposits, and it is an open question whether the arch



stands altogether by itself or is somewhat connected with the Wild Horse arch at the north and the Carlos at the south. It appears in sections BB and DD of the Structure Section sheet, and its southern part is expressed in the topography of the district as the "Sand hill" west of Pueblo.

West of the Rock Canyon arch is a shallow and ill-defined hollow, 6 or 8 miles long, north and south, and two-thirds as broad.

**The northeastern depression.**—From the eastern limits of the Wild Horse, Rock Canyon, and Carlos arches there is a general descent of the rocks toward the east and northeast, the average dip being 1 foot in 25. The slope is interrupted by a long, flat arch, 100 to 150 feet high, which passes through the eastern part of the city of Pueblo and runs thence to the north and southwest; and the companion of this arch is a shallow trough just west of it.

**Minor faults and flexures.**—Wherever the Cretaceous strata are quite free from soil and other surficial material, they show many waves and small faults. These can not be traced out and mapped, because they are largely covered from view, but it is believed that they abound everywhere in the district, modifying the greater flexures as ripples modify the broad swells of the ocean.

#### ORIGIN OF THE TOPOGRAPHY.

The hills and valleys, mesas and canyons, and all the details of topographic form that diversify the district have been wrought by the eroding action of flowing streams and beating rain. Had there been no erosion the deformed surface of the highest Cretaceous formation would exhibit a system of smooth arches, domes, troughs, and basins, with here and there a fault-cliff; but that formation was long ago all washed away, and with it disappeared the simple structural shapes. Along with that formation portions of all the lower beds were carried away, and the loss was greatest where the structural arches would otherwise be highest. The general lowering of the surface has amounted to thousands of feet, and during its progress the details of topographic form have been carved out. They have, in fact, been remodeled many times, the pattern being gradually modified as the conditions of the work were varied, and the existing forms are only one phase of a changing scene.

The chief conditions affecting the sculpture of the land are the positions of streams, the slopes of stream beds, and the rock structure, or the extent and arrangement of resistant and yielding rocks. Where the water is gathered in streams it carves deeper than on the interstream areas, and the divides are comparatively high because they escape the strongest action of the water. The down-cutting of the streams is limited by the fact that they are also the carriers of all the waste from the land; if their channels become too flat their current slackens and the waste is not carried off. In this and similar ways the slopes of all the stream beds are automatically adjusted in a harmonious system, so that the wasting of the whole district is nearly uniform, and its entire surface is gradually reduced. The chief variations from uniformity are occasioned by differences in rocks, some of which resist erosion better than others. Nine-tenths of all that has been eroded from the district since the deformation of the Cretaceous system of strata was shale, one of the most yielding of rocks, but in this mass of shale were embedded sandstones and limestones, which are comparatively resistant. As the shale was slowly pared away, the harder beds were from time to time laid bare, appearing first from under the tops of high arches. Whenever so exposed the hard rock retarded erosion and was soon left as a hill projecting above the plain of shale. After a time the hard bed was eaten through at the top of the arch, and a valley was eroded from the core of shale, the worn edge of the hard bed becoming the crest of a circling ridge. Then the valley grew broader and the ridge was moved farther back; but though changing its position it persisted as a topographic feature, marking the outcrop of the resistant rock. The accompanying diagrams (figs. 12-14) illustrate three stages in the erosion of an arch.

Pueblo—5.




Fig. 12.—Ideal section of an arch of strata, including a thick limestone and a thinner limestone, both embedded in shale.

At a certain stage of erosion the thicker limestone projects above the plain at the crest of the arch.



Fig. 13.—The same at a later stage of erosion. The crest of the arch is occupied by a valley in shale. The cut edges of the thicker limestone crown parallel ridges enclosing the valley.




Fig. 14.—The same at a still later stage of erosion. The valley is broadened by the wearing back of the limestone ridges, and a second pair of ridges mark the outcrops of the thinner limestone.

#### THE ARCHEAN UPLAND.

The upland in the southwestern corner of the quadrangle stands higher than any other. As already stated, it is part of a mosaic pavement of various hard rocks which had been first ground off to a level, then buried by Juratrias shales and other strata, and finally uplifted in the St. Charles arch and denuded. Its sloping plain is uneven because the granites of its tessellated pattern are more resistant than the schists, and it is divided into parts because it lies in the track of the St. Charles River and its tributaries. Here, as elsewhere, the streams have carved out the canyons in which they run; and they have carved deeply because the steep slopes of their beds give great power to their waters in time of flood.

#### THE DAKOTA HOGBACKS.

The most resistant formation of the local sedimentary series is the Dakota sandstone. Below it are more than 2000 feet of shale and weak sandstone; above is a still greater depth of shale, containing only one strong bed of notable thickness. Wherever the Dakota is laid bare on the flank of an arch or other uplift it is carved into bold relief by the deep erosion of the enclosing shales, and the ridge to which it gives rise is called a hogback. The outer slope of the ridge (the slope away from the arch) is composed chiefly of the upper layers of sandstone, which dip with the inclination of the surface. The inner slope, which is steeper, exposes the edges of various formations, the Dakota sandstone appearing at the top and the Morrison and Fountain beds below.

The hogback shows its typical character in the southeastern part of the quadrangle, where it curves about the Beulah and Red arches. Its height above the adjacent lowlands is more than 1000 feet, the westward slope being very steep and the eastward more moderate. It is crossed by two streams, Red Creek and the north fork of St. Charles River, which divide it to the base. It ends where the Beulah arch joins the St. Charles, because the sandstone, after its outcrop turns eastward along the latter arch, is so much broken by faults as to have lost much of its resistant character.

In the crest of the hogback between Red Creek and Beulah are three notches, marking the places of streams that once crossed the ridge but have been diverted. They were probably small creeks rising west of the Pueblo quadrangle and flowing eastward, and being small they could not deepen their channels so rapidly as Red Creek and the St. Charles. They fell a prey to the branches of the larger streams, which enlarged their valleys in the yielding beds of the Fountain formation and finally drew off the headwaters of the smaller streams. These old channels, stranded high on the sandstone ridge, are identical in origin with the so-called wind-gaps of the Appalachian region. The northernmost, near Red Creek, and the southernmost, opposite Wells Canyon, are each more than 300 feet deep and nearly half a mile broad; the middle one is smaller. The course of the northern stream is indicated at a few points by remnants of Neocene gravel perched on the mesa between Rush and Pecks creeks, and it thus appears that the stream did not coincide in position with any of the modern waterways. It is noteworthy that these gravels include fragments of granite as well as Dakota sandstone, thus

showing that the creek which brought them rose west of the hogback and in the Archean area beyond the valley now carved from the Fountain beds. The diversion of these streams because they were too weak to keep pace with their neighbors in carving channels through the resistant sandstone is part of a general process of rearrangement by which the minor elements of drainage are turned away from resistant rocks. The arrangement of small waterways is continually adjusted to the arrangement of resistant rocks, which changes as the face of the land is worn down.

At the north the hogback is flexed by two arches, the Beaver and the Turkey. In the Turkey arch it makes an abrupt turn, doubling on itself, and the parts on opposite sides coalesce at top instead of being separated by a valley. The hogback form is thus locally lost, and in its stead is a great hill shaped like the toe of a slipper and cased on three sides by dipping sandstone. The fourth side breaks off in a cliff overlooking a valley in the Morrison shale, but this lies north of the quadrangle boundary. The sandstone hill is furrowed by many gorges and trenched near its eastern base by the canyon of Turkey Creek.

#### LIMESTONE MESAS AND TERRACES.

Above the Dakota sandstone are three limestone beds whose resistance to erosion has diversified the topography. These are the Greenhorn limestone, the heavy limestone at the base of the Niobrara formation, and the thin limestone at the top of the same formation. The first two are closely associated, being separated by only 200 feet of shale, but the third is independent. In going outward from any outcrop of Dakota sandstone, one usually crosses a valley marking the position of the Graneros shale, ascends a low cliff capped by the Greenhorn limestone, crosses a narrow terrace of the same limestone, ascends a second slope marking the outcrop of Carlile shale, climbs a higher cliff whose face is composed of the sandstone beds at top of the Carlile and the limestone beds at base of the Niobrara, and then stands upon a plain constituted by various beds of the Niobrara formation. This topographic stairway is illustrated by the diagram, fig. 15, and is one of the most characteristic features of the district.




Fig. 15.—Typical profile of the cliff capped by the Niobrara limestone and the terrace capped by the Greenhorn limestone.

Where the rock dips gently backward from the head of the stairway the plain at top usually begins with a long slope composed of the upper limestone, and the edge of the limestone is then the crest of a ridge analogous to the Dakota hogback. This general character obtains in the southern part of the district, but the upland is crossed by so many lines of drainage that its limestone cap is cut into a multitude of insular buttes or mesas, and the ridge character is greatly masked. In the region of parallel faults erosion has been further influenced by the dislocations, with the result that the limestone mesas lie at different heights and are even more numerous. On the Historical Geology sheet it will be observed that the pattern representing the Niobrara limestone and indicated by the letters Kn is there broken up into many small patches which coincide with the tops of hills, as indicated by the contours.

On the east side of Rush Creek, where the rocks are flexed downward to the northwest, the Niobrara limestone, by resisting erosion, has preserved a hill-slope with the general profile of the flexure; but there are many places where the limestone has been eaten through by the streams, and cliff-bounded coves have there been opened in the Carlile shale below. As one ascends the creek valley one finds these coves successively larger, and at last merged together.

In the "Sand hill" south of Rock Canyon are many similar valleys eroded from the Carlile shale and surrounded by cliffs of limestone. The

canyon of the Arkansas from Beaver to the Goodnight ranch is everywhere bounded by cliffs of the Niobrara limestone, but is in general not deep enough to reach the Greenhorn ledge. In crossing the Rock Canyon arch the river cuts down to the Dakota sandstone, and the Niobrara cliff recedes on each side so as to open a sort of amphitheater. The steep slopes of this opening are greatly diversified by spurs and canyons, but among these can be traced the Greenhorn terrace, forming an interrupted arch on either side.

On the gentle slopes of the Beaver arch the Niobrara limestone occupies a belt from 1 to 2 miles wide, and is dissected by a plexus of streams. Pierce Gulch divides it on the east and the canyon of Beaver Creek on the west, and a dozen waterways leading from the axis outward to these main channels cut the limestone table into an archipelago of mesa buttes.

The Pumpkin arch is outlined by low ridges of the Niobrara and Greenhorn limestones, somewhat as represented in fig. 14, and the westward ridge, holding the same character, follows the base of Turkey arch also.

The Wild Horse arch is represented in the landscape by a hill, sometimes called the Cedar Ridge. The stage of erosion is here approximately that shown by fig. 12, the arching hard rock being the lower Niobrara limestone; but in part of the arch the stage of fig. 13 has been reached, for four of the small streams draining the ridge have cut through the limestone and opened coves in the Carlile shale. Three of the coves coalesce, making Wild Horse Park, a beautiful valley 2 miles long, whose circle of limestone cliffs is broken only by the narrow gateways through which the eroding streams escape.

South of the Arkansas Canyon the Rock Canyon arch exhibits the same transitional stage of erosion. The resistant Niobrara bed has delayed the wearing down of the land so that the miscalled "Sand hill" rises above the plain, a dome sheathed by limestone, but the draining streams have here and there worn through the limestone, as already mentioned.

The upper limestone of the Niobrara, though thin and too weak to make a serviceable building stone, is yet so much stronger than the shales enclosing it that its outcrop usually caps a ridge or hill 30 to 50 feet high. A line of these hills faces Fountain Creek opposite Pueblo, and another line, starting at the Arkansas River in the western suburb of the city, follows the valley of Dry Creek northward to the vicinity of Blue Hill. These lines have many irregularities occasioned by small faults and flexures.

#### TEEP E BUTTES.

The cylindrical masses of limestone standing vertically in the Pierre shale not only project from the general plain but preserve conical hills of shale. At the top of each hill the limestone core is exposed to the weather and is slowly broken up by frost. The fragments, falling to the slope below, lodge on it and form a stony or gravelly mantle by which its erosion is retarded. Though only 25 to 60 feet high, the hills are conspicuous by reason of their symmetry. They occur only in the northeast part of the quadrangle, abounding on the southern part of Baulite Mesa and on the plain east of Pinyon. The form of the tepee butte and the appearance of a cluster of them are shown by a view in the Illustrations sheet.

#### GRAVEL MESAS AND TERRACES.

Loose gravel and coarse sand, by absorbing storm water as it falls, retard its flow and interfere with its erosive action. Gravel also resists the force of rills by the weight of its pebbles. Thus the gravel and sand beds of the Nussbaum and later formations, though to be counted as weak beds in relation to creeks and rivers, are resistant when occupying interstream areas. In such situations they are more resistant than the various shales of the Cretaceous, and they often protect bodies of shale from erosion. The Baulite Mesa is a large mass of Pierre shale protected from erosion by a capping of Nussbaum sand and gravel. Of the same character are "The Mesa," in Pueblo, and the flat hills southeast of Bessemer Junction, except that the protective gravels are of later date.

Topography of the Rock Canyon and other arches.

Adjustment of drainage to structure.

Sandstone of the Turkey arch.

Uplands of the resistant granites of the St. Charles arch.

Geologic stairway of limestone and shale.

Hogbacks of the resistant Dakota sandstone.

Characteristics of the hogbacks.

Mesas of Niobrara limestone.

Wind-gaps.

Coves in the Carlile shale.

Structure of the Tepee Buttes.

Mesas protected by sand and gravel.



Since the Nussbaum epoch the general level of the land has been reduced by erosion several hundred feet, so that all surviving deposits of that date are the caps of hills. The alluvial deposits of the earlier Pleistocene were made during the same wasting, and their remnants were stranded at various heights above the modern streams, where they protect hills and terraces of shale.

#### ECONOMIC GEOLOGY.

The mineral resources of the district which have already received some development are clay, limestone for flux and quicklime, building stone, artesian water, iron ore, and gypsum. Most of these will have greater development in the future, and there are possibilities of fire-clay and hydraulic cement. Each material is definitely associated with one or more of the geologic formations, so that the areas in which it occurs can be pointed out with the aid of the map of formations, the Historical Geology sheet; but in certain cases more specific information can be given, and to this end two special sheets have been added: The Economic Geology sheet shows the distribution of building stone, limestone, fire-clay, and gypsum. The Artesian Water sheet gives the principal facts about underground water.

#### SANDSTONE.

Sandstones occur in the Carlile, Dakota, Fountain, and Harding formations. Those of the Dakota and Carlile are available for building stone. The Fountain sandstones may include useful beds, but so far as examined contain too much clay to be either strong or durable. The quality of the Harding sandstone was not determined. It has but a small area within the quadrangle and was not seen to be well situated for quarrying.

*Sandstone of the Carlile formation.*—This rock is of fine grain and rather close texture. The cementing material is probably calcite, but there is always enough clay present to make the rock soft and weak. The color is light-yellow and permanent. Some strata are found 1 to 2 feet thick, but the chief product of the quarries is thinner. The total thickness of workable layers of this sandstone rarely exceeds 5 feet. It is found at and near the top of the Carlile formation, and being covered by the heavy limestone of the Niobrara comes to the surface in the cliff that usually marks the edge of that limestone. It is only in places where most of the limestone has been removed by natural erosion that the quarryman can afford to strip off the remainder in order to obtain the sandstone. Such a locality is found in the "Sand hill" south of Rock Canyon, and there are others of less extent in many parts of the district. On the Economic Geology sheet no attempt has been made to show either these places of special availability or variations of quality, but the whole line of outcrop is marked. It is a sinuous line winding through the southern and western parts of the sheet.

*Dakota sandstone.*—Most of the sandstones of the Dakota formation are in thick strata, ranging from 2 to 20 feet. In grain and texture they vary from fine to coarse and from open to close. Their colors—white, yellow, orange, and speckled-brown—are probably permanent. They contain little clay, and the cementing material is usually calcite. Their strength has not been practically tested, but their resistant quality, in virtue of which they project above the plain in hogbacks, betokens architectural durability. Their quantity is inexhaustible; with a net thickness of 200 to 500 feet, they cover 85 square miles within the quadrangle.

With its broad exposure, its great thickness, and its variety of color and texture, there can be little question that the Dakota sandstone has a much greater value for building purposes than the Carlile. The Carlile was first used because most accessible, but the ultimate source of supply is the Dakota, and only the Dakota can furnish a product for shipment.

#### LIMESTONE.

Workable beds of limestone are found in three formations, the Niobrara, the Greenhorn, and the

Millsap. The upper limestone of the Niobrara formation is hardly worthy of mention in this connection, although its convenient occurrence in the suburbs of Pueblo has led to its occasional employment where neither strength, hardness, nor durability is essential.

*Lower limestone of the Niobrara formation.*—This rock is fine-grained, of a pale-gray color, and on the whole is remarkably uniform. It is a typical lime carbonate, with only a trace of magnesia and less than 10 per cent of argillaceous impurity. Its exact composition is shown in the table of analyses. For most purposes its only deleterious constituent is marcasite, or iron sulphide, which forms small nodules in certain layers. Its mode of occurrence is exceptionally convenient for quarrying. Beds from 1 foot to 3 feet in thickness are naturally separated by thin partings of shale, and the whole mass, 35 to 50 feet deep, usually lies in the face of a hill, so that gravity can be used in the handling. It is extensively employed as a flux in the reduction of silver and iron ores, and it is equally qualified for the making of quicklime. Although readily obtainable in blocks of convenient dimensions, it is not serviceable for building purposes because its brittleness leads to cracking and spalling from changes of temperature.

The area in which it occurs at the surface, or so near the surface as to be readily accessible, is very large, and constitutes a complicated series of belts in the northwestern, central, and southern parts of the quadrangle. On the accompanying Economic Geology sheet a special color is assigned to the representation of these belts.

*Limestone of the Greenhorn formation.*—This limestone is also widely distributed, its outcrop forming a terrace on the slope under the Niobrara cliff. Its strata are thinner than those of the Niobrara limestone, and are separated by shale beds of such thickness as to render quarrying expensive. Being less accessible, it has remained practically untested, and its qualities can not be described further than to say that its brittleness and tendency to vertical cleavage disqualify it for building purposes.

*Limestone of the Millsap formation.*—This limestone also is untested, except that a moderate amount has been burned for lime. It differs from both the others in its variability, color and texture showing considerable range. So far as may be judged from its behavior under the influence of the weather, it contains beds which would be available for construction. It occurs only in two small areas near Beulah.

#### MARBLE.

Just west of the boundary of the Pueblo quadrangle is a quarry of marble. It was not visited during the survey of the district because it had not then been opened; but from information since obtained the marble is supposed to occur in the Millsap formation, and it is therefore possible that similar discoveries may be made within the quadrangle in the vicinity of Beulah. The single specimen seen is white and of rather coarse grain.

#### GYPSUM.

Near the northwestern corner of the quadrangle are two small areas of the Morrison formation. These are parts of a much larger area occurring in the Colorado Springs quadrangle, adjacent to the north, so that they belong rather to that quadrangle than to the Pueblo. The formation contains an important series of gypsum beds, and a few of those beds were observed in the more easterly of the Pueblo areas. They are overlooked from the south by a high cliff of Dakota sandstone, and are accessible only from the Colorado Springs side. The gypsum is massive and is mottled with gray and white.

A granular or earthy gypsum, used in Pueblo for the manufacture of plaster, is said to have been obtained from an alluvial flat near Greenhorn station. I am unable to describe its mode of occurrence, as the locality was overlooked during the progress of the survey. Gypsum also occurs in the form of selenite crystals in many of the Cretaceous shales, especially the Niobrara and the lower part of the Pierre, where the shales have been much broken

in the process of deformation. This gypsum is concentrated in veins, but no veins were seen of such thickness as to constitute deposits of practical importance.

#### SHALE AND CLAY.

These argillaceous materials occupy fully one-half of the surface of the quadrangle, and they afford much variety of composition and quality. Shales occur in the Millsap, Fountain, Dakota, and Greenhorn formations, and they constitute the chief part of the Morrison, Graneros, Carlile, Niobrara, and Pierre. Clays occur in the various alluvial formations, sparingly in the Nussbaum and earlier Pleistocene, abundantly in the fresh-formed deposits along the streams that traverse the shale districts.

The clays have been used in Pueblo for the manufacture of bricks and tiles, and trials have been made of the availability of the various clays and shales for the manufacture of pottery.

The beds of shale are so extensive and they afford such variety of texture that it may fairly be expected that they will eventually constitute the basis of important industries. For this reason analyses were made of specimens representing various types, and these are here published (See table of analyses, p. 7) as a matter of record, although it is at present impossible to point out their full economic bearing.

*Hydraulic cement; Portland cement.*—The hydraulic cements, or cements having the property of setting under water, are partly derived from the burning of certain argillaceous limestones and partly from the burning of mixtures of limestone with clay or slag. The successful prosecution of the industry requires much technical knowledge, and in each locality depends on experimentation with the particular materials available; but some preliminary indications may be based on the composition of the materials as determined by chemical analyses, and it appears quite possible that the necessary combinations may be found in this district.

The concretions in the upper part of the Pierre formation are essentially of argillaceous limestone, and closely resemble in composition certain concretions from which hydraulic cement has been made in England without the admixture of other material. They occur in considerable abundance about the flanks of Baculite Mesa and thence northward to the limit of the quadrangle. In the same region occur the tepee cores, which also consist of argillaceous limestone and might be available for the manufacture of cement if mixed with a small amount of the enclosing shale.

In the Niobrara formation, just above the limestone bed quarried for flux, is a series of beds of varying texture containing unequal mixtures of limy and argillaceous material, and it is quite possible that trial may discover in this series individual beds which have the necessary composition. Analysis No. 4 gives the composition of one of the more argillaceous of these beds, and a comparison with analysis No. 1, representing the underlying limestone, shows that a mixture in the ratio of about 1 to 4 would have approximately the composition of some Portland cement before burning.

#### FIRE-CLAY.

Clays and shales which resist great heat without fusing are said to be refractory and are called fire-clays. They differ in composition from ordinary clays, having large amounts of silica and alumina and relatively little iron, lime, magnesia, potash, and soda. In the Pueblo and adjacent quadrangles they have been found only in the Dakota sandstone. The upper part of that formation contains a variable number of shale beds interstratified with the sandstone beds, and some of these are in certain places refractory. The writer collected shale samples from the outcrop of the formation in the Pueblo, Apishapa, Canyon, and Colorado Springs quadrangles, and six of these were shown by proper tests to have the refractory property in greater or less degree. Prospecting will doubtless discover a number of deposits equal to the best of these samples, and may discover still higher grades.

The most satisfactory and valuable test of fire-clay is the fire test. Samples properly prepared are heated in a furnace by the side of standard compounds and the results are compared. On the Seger scale of refractoriness, eleven grades, numbered from 26 to 36, cover the range of fire-clays, the best being No. 36. This test was applied to the six samples mentioned above by Prof. H. O. Hofman, of the Massachusetts Institute of Technology, and their several grades were found to be 29, 30½, 31, 33½, 34½, 34½. Two of the samples were obtained from the Pueblo district, and these are further reported in the table of analyses, Nos. 9 and 11. It is judged from the analysis that No. 10 of that table is also a fire-clay, but the sample in hand was too small for the application of the fire test. To give the above figures a more intelligible meaning, I quote also Professor Hofman's rating of two well-known Colorado fire-clays: a sample from Carbondale gave 33½ on the Seger scale, and one from Golden 31½.

The areas occupied by the upper part of the Dakota formation, or the areas to which the fire-clays are restricted, are shown on the Economic Geology sheet. The largest is in the southwest quarter; others at the northwest and southeast connect with still larger tracts outside the quadrangle; a small tract surrounds Rock Canyon; and two small tracts lie in the valley of Greenhorn Creek and its eastern branch.

The refractory shales are all below the highest sandstone layer. They differ so widely in color that no general description can be given. The one represented by analysis No. 10 is nearly black and full of fossil twigs converted to coal. No. 9 is dark blue-gray; No. 11, light-gray. One of the samples ranking highest by fire test is nearly white, and gritty from the presence of much sand.

#### IRON ORE.

In the Rusty zone of the Pierre shale are many concretions carrying iron. They consist of lime carbonate and iron carbonate with some clay. Where exposed to the air the iron carbonate is slowly converted to limonite, the color changing from gray to reddish-brown. The zone is best exposed on the slope between Baculite Mesa and the Arkansas River, and its belt of outcrop runs thence north-northwest past Overton and Steele Hollow. The concretions were at one time used in combination with other ores in the manufacture of steel.

#### PETROLEUM, ROCK GAS, COAL, PRECIOUS METALS.

None of these substances are to be counted among the mineral resources of the district, but they should nevertheless be mentioned, as there have been false impressions in regard to them.

The only rocks of the district that are even slightly bituminous are certain parts of the Graneros and Carlile shales. If petroleum or gas were naturally distilled from these it would tend to accumulate in the Greenhorn and Niobrara limestones. As the limestones have been repeatedly penetrated by the drill without the discovery of more than a trace of gas, there is no reason to expect that a valuable accumulation will be discovered. The petroleum obtained in the Florence oil field, a few miles west of the district, comes from sandy layers in the upper part of the Pierre shale, but the Pueblo district includes only the lower part of that formation.

In the calcareous shales below the middle of the Niobrara formation fossil logs consisting of coal are occasionally found, but there are no coal beds. Some layers of the Dakota sandstone contain many shreds of vegetal matter in the condition of coal, and so do some of the interbedded shales. It is within the range of possibility that the formation contains local seams of coal, but the occurrence of workable coal beds is not probable. The formation yielding coal in the adjacent Canyon quadrangle does not extend to the Pueblo.

Metalliferous veins often occur on faults, and they are sometimes associated with metamorphic schists. There is therefore warrant for prospecting among the schists and along the faults in the southwest part of the district, and frequent openings show that intelligent search has already been made in that area. The writer is not acquainted with the history of that

Behavior of samples under fire test.

Plastic clays.

Economic importance of shales.

Materials for Portland cement.

Concretions for cement.

Limestones and shales.

Dakota shales available as fire-clay.

Metalliferous veins.

Flux and lime rock of the Niobrara.

The Economic and Artesian sheets.

Extent of outcrop.

Clay-iron concretions.

Quarry rock of the "Sand hill."

Dakota sandstone as building material.

Gypsum in the Morrison formation.

Alluvial gypsum.



search, but the general abandonment of claims seems to show that it was unsuccessful.

#### ARTESIAN WATER.

Rock reservoirs are usually not open chambers, but beds of sand or porous sandstone, and the water occupies the small spaces between the grains. The only important water-bearing beds in the district are the sandstones of the Dakota formation.

Water confined in the rocks so as to press on the cover of its reservoir is said to be artesian. When tapped by boring, it rises in the bore-hole, making an artesian well. The highest level to which it will rise is called its head. If the head is above the surface of the ground the well is said to flow; otherwise it is called a pumping well.

*The question of depth.*—The Dakota formation is altogether wanting from certain small areas; in other, larger areas it lies at the surface, and thence it dips down below the surface. Through the greater part of the quadrangle it is not visible, but is buried under other rocks, chiefly shales. Owing to the deformation of the rocks and the resulting unequal erosion, its depth below the surface differs from place to place, ranging from nothing to more than 3000 feet.

If a person starts to sink a well where the upper stratum of the Dakota formation lies at the surface, he has to go down only 100 to 200 feet to reach one of the artesian beds. If he starts his well where the surface rock is the top of the Greenhorn limestone he must penetrate the Greenhorn limestone, 50 feet, and the Graneros shale, 200 feet, in addition to the 100 feet or more of Dakota beds. If he starts on the Niobrara limestone he must go still deeper, penetrating in addition the Niobrara limestone and the Carlile shale, and the depth of his well will be 600 or 700 feet. The thicknesses of the various formations penetrated in making artesian wells are graphically shown on the Columnar Section sheet by the "Artesian Section," in which the scale of feet has its zero at the horizon of the highest water-bearing bed. If one can determine what member of the rock sys-

Pueblo—7.

tem forms the surface of the ground at any spot, one can read in this section the depth to water. The relations of surface rock to water depth are also illustrated by sections of the Structure Section sheet, and they are still further illustrated by the accompanying ideal section (fig. 16), where




Fig. 16. — Ideal profile and section illustrating artesian conditions.

Kn, limestone of Niobrara formation; Kc, Carlile shale; Kgn, Greenhorn limestone; Kgr, Graneros shale; Kd, Dakota sandstone; H, hogback; GI, line of head; A, barren ground; B, D, F, pumping wells; C, E, flowing wells.

wells at B and C start on the Dakota sandstone, at E on the Greenhorn limestone, at D on the Niobrara limestone, and at F in Niobrara shales. A boring at A, starting in rocks which underlie the Dakota sandstone, would not find the water.

It happens that the formations above the Dakota are remarkably uniform in thickness from place to place, so that a thorough knowledge of the rock formations makes it possible to predict with tolerable accuracy the depth at which the first Dakota stratum will be encountered. The Dakota formation itself is much more variable, not only in thickness but in the order and number of its water-bearing beds. The thickness varies from 650 feet at the southwest to 300 feet at the northeast. The water-bearing beds occur at different distances from the top in different localities, the depth of the first or highest being in places less than 100 feet, and elsewhere more than 150 feet. The number of such beds probably ranges from one to three or four, but this has not been tested by the drill, as the borer usually stops at the first good stream.

Assuming the first artesian water to lie 150 feet below the top of the formation, the writer has estimated its depth below the surface of the ground for the whole artesian area, and by the aid of these estimates has drawn the contours of the Artesian Water sheet. Each contour is a line drawn through points where the estimated depth of water below

the surface of the ground is the same; for example, the water is estimated to lie 800 feet below the surface at all points of the district corresponding to the 800-foot artesian contour. To make practical use of the map the enquirer should locate on it the point as to which he wishes information, and note the numbers of the artesian contours between which it falls. The estimated depth in feet is between the numbers attached to the two contours.

While the estimates are everywhere subject to some error, especially from the variability of the Dakota formation, it is believed that they will in general come within 100 to 200 feet of the fact, and thus prove practically serviceable. Their greatest uncertainty is on Boggs Flat, where for several miles there are no good rock exposures, and on a similar flat 10 or 12 miles farther north; and to indicate this uncertainty the contours are there drawn as broken instead of full lines.

The known variability of the Dakota formation suggests that in some localities there may be no rock bed so porous as to be freely traversed by water; but as no boring known to penetrate the sandstone in the Pueblo and adjoining districts has heretofore failed to find a supply of water, great weight need not be given to the possibility of barren tracts.

The more general facts of distribution shown by the contours are that the depth of artesian water is moderate in the southern, central, and western parts of the territory, and great in the northeastern corner of the quadrangle. Within the artesian area the crests of arches are in general more favorable for wells than the troughs, because erosion has there left less rock to be penetrated. Over much of the Rock Canyon arch the water horizon is within 600 feet of the surface, and the crests of the St. Charles, Wild Horse, Pumpkin, and Beaver arches afford equally favorable sites.

*The question of head.*—The question whether the artesian water when found will rise to the top of the well and flow out is often as important as the question of depth, and unfortunately can not be answered in an equally satisfactory way. If the water-bearing rock were merely a reservoir

in which the water lay motionless the head would be the same at all points; that is, the water would in all wells reach the same level, and flowing territory would be separated from pumping territory by a horizontal line contouring the hillsides. But the water is really flowing at a slow rate through the rock, and its head varies from point to point. The mode and rate of variation depend on the source of supply, the direction of flow, the resistance to flowage through small pores, and various other factors. On these various points there is little information, and all prediction as to head is correspondingly uncertain. Moreover, the head is usually not the same for different water-bearing beds, and it is reduced in each locality by every draft on the supply through a flowing or pumping well. The line on the map separating the supposed flowing from the supposed pumping territory was drawn with much less confidence than the contours of depth.

These two territories are distinguished by colors. A third color shows the territory occupied by the Dakota sandstone but not believed to carry water under notable pressure; this is part of the gathering ground where rain water is absorbed by the rock. A fourth color shows territory in which the sandstone does not occur. For practical purposes these may be classed together as barren ground.

In fig. 16 the broken line, GI, represents the plane of head, a plane sloping away from the Dakota hogback, H. Where this line passes above the profile of the land, as in crossing the valleys at C and E, flowing wells may be obtained; where it passes below the surface, as in the regions B, D, F, only pumping wells are possible. In selecting a site for boring, the water-seeker should bear in mind that the local level of head, or its height above the ocean, is independent of the shape of the ground. The chance of a flowing well is always better in a valley than on a neighboring hill or mesa.

GROVE KARL GILBERT,  
Geologist.

June, 1897.

#### COMPOSITION OF ROCKS AS SHOWN BY CHEMICAL ANALYSIS.

|  | 1. Limestone, lower part of Niobrara formation. | 2. Shale; 30 feet above base of Graneros formation. | 3. Shale; 70 feet below top of Carlile formation. | 4. Shale; 120 feet above base of Niobrara formation. | 5. Shale; Risty zone of Pierre formation. | 6. Shale; Top zone of Pierre formation. | 7. Earthy limestone; Top zone of Pierre formation. | 8. Earthy limestone; middle zone of Pierre formation. | 9. Fire-clay; Dakota formation; Davis ranch. | 10. Fire-clay; Dakota formation; near head of Rock Creek Canyon. | 11. Fire-clay; Dakota formation; near head of Pierre ditch. |
|--|---|---|---|--|---|---|--|---|--|--|---|
| Silica (Si O <sub>2</sub> )                        | 6.4   | 63.60   | 60.60   | 45.89  | 51.69                                     | 60.80                                   | 7.46   | 12.47   | 63.52  | 76.56  | 86.79   |
| Titanium dioxide (Ti O <sub>2</sub> )              |   | .66   | .35   | .52  | .66                                       | .47                                     | 1.78   | 3.30  | .68  | .60  | 8.29  |
| Alumina (Al <sub>2</sub> O <sub>3</sub> )          | 1.3   | 16.74   | 16.42   | 13.24  | 16.50                                     | 15.63                                   |  | 24.72   | 8.30   |  |   |
| Iron sesquioxide (Fe <sub>2</sub> O <sub>3</sub> ) | 2.1   | 4.63  | 4.95  | 3.88   | 7.90                                      | 4.62                                    | .94  | 1.37  | .43  | .38  | 75  |
| Lime (Ca O)  | 50.4  | .68   | 1.61  | 12.09  | 4.41                                      | 1.63                                    | 46.98  | 42.26   | .30  | .12  | .34   |
| Magnesia (Mg O)                                    | trace   | 1.19  | 1.43  | 2.12   | 2.10                                      | 2.73                                    | 2.36   | 2.61  | .13  | .24  | .13   |
| Potash (K <sub>2</sub> O)                          |   | 2.92  | 2.98  | 2.31   | 2.68                                      | 2.55                                    | .37  | trace   | trace  |  | .25   |
| Soda (Na <sub>2</sub> O)                           |   | .29   | .92   | .47  | 2.07                                      | 1.45                                    |  | trace   | trace  |  |   |
| Phosphoric oxide (P <sub>2</sub> O <sub>5</sub> )  |   | .16   | .31   | .17  | .22                                       | .10                                     | undet.   | undet.  | trace  | .06  | .05   |
| Carbon dioxide (C O <sub>2</sub> )                 | 39.5  |   |   | 10.38  | 3.19                                      |   | 39.25  | 35.57   |  |  |   |
| Water lost at 100° C.                              |   | 2.88  | 3.91  | 1.38   | 3.02                                      | 3.19                                    | .16  | .52   | 1.58   | 1.26   |   |
| Water lost above 100° C.                           |   | 5.99  | 5.72  | 4.16   | 6.00                                      | 4.16                                    | .70  | 1.31  | 8.41   | 4.40   | 3.78  |
| Organic material                                   |   | .46   | .84   | 3.47   | .53                                       | 2.87                                    |  | .40   | 8.31   |  |   |
| Total  | 99.7  | 100.20  | 100.04  | 100.08   | 100.97                                    | 100.20                                  | 100.00   | 100.00  | 100.17                                       | 100.23   | 100.38  |
| Fire test  |   |   |   |  |   |   |  |   | 31   |  | 29  |


NOTE.—Analysis No. 1 was made by the chemist of the Pueblo Smelting Company from a carload sample obtained from Harp's quarry. The other analyses were made in the chemical laboratory of the United States Geological Survey, Nos. 2, 3, 4, 5, 6, 9, 10, and 11 by Mr. George Steiger, Nos. 7 and 8 by Dr. W. F. Hillebrand.






LEGEND


RELIEF  
 (printed in brown)




DRAINAGE  
 (printed in blue)



CULTURE  
 (printed in black)



Henry Gannett, Chief Topographer.  
 E.M. Douglas, Topographer in charge.  
 Triangulation by A.H. Thompson.  
 Topography by R.O. Gordon and W.J. Lloyd.  
 Surveyed in 1894.



Scale 1:25000  
 Contour Interval 50 feet.  
 Datum is mean Sea level.  
 Edition of Jan. 1897.





SURFICIAL ROCKS

(Areas of Surficial rocks are shown by patterns of dots and circles.)

- LATER  
Pi  
Alluvium  
(bottom lands)
- EARLIER  
Pe  
Alluvium  
(gravel, sand, and silt  
supplying terraces and mesas)

PLEISTOCENE

SEDIMENTARY ROCKS

(Areas of Sedimentary rocks are shown by patterns of parallel lines.)

- NEOCENE  
Nn  
Nussbaum  
Formation  
(sand, gravel, and silt)
- Kp  
Pierre  
shale  
(gray shale with  
concretions)
- Kn  
Niobrara  
formation  
(shale and limestone)
- Kcr  
Carlile  
shale  
(shale and sandstone)
- Kgn  
Greenhorn  
limestone  
(limestone and shale  
in alternate strata)
- Kgs  
Graneros  
shale  
(gray shale)
- Kd  
Dakota  
sandstone  
(gray sandstone  
and shale)

CRETACEOUS

(Vegetation of the Bentons formation)

- Jm  
Morrison  
formation  
(variegated clay and  
sandstone, gypsum)

JURATRIAS

- Jf  
Fountain  
formation  
(red shale, sandstone,  
and conglomerate)

JURATRIAS ?

- Cm  
Millsap  
limestone  
(limestone and shale)

CARBONIFEROUS

- Sh  
Harding  
sandstone  
(white sandstone)

SILURIAN

(Areas of ancient, crystalline rocks and/or metamorphic rocks of unknown origin are shown by patterns of short dashes.)

- Ar  
Schist and  
Granite  
(igneous and metamorphic)


ARCHEAN ?

Faults

Sections



Henry Gannett, Chief Topographer.  
E.M. Douglas, Topographer in charge.  
Triangulation by A.H. Thompson.  
Topography by R.O. Gordon and W.J. Lloyd.  
Surveyed in 1894.



Contour Interval 50 feet.  
Datum is mean Sea level.  
Edition of Jan. 1897.

Geology by G.K. Gilbert.  
Assisted by Robt. T. Hill.  
Surveyed in 1893.



LEGEND  
(continued)

✕ Quarries

Known productive formations

**Kn**

Niobrara formation  
(limestone for lime and flux)

**Kcr**

Carlisle shale  
(sandstone used for foundations)

**Kd**

Dakota sandstone  
(largely available for building; upper part contains fire clay)

**Jm**

Morrison formation  
(local deposits of gypsum)

**Gm**

Millsap limestone  
(local developments of marble)

LEGEND

SURFICIAL ROCKS

(Areas of Surficial rocks are shown by patterns of dots and circles.)

**Pl**

Alluvium  
(bottom-lands)

**Pe**

Alluvium  
(gravel, sand, and silt capping terraces and mesas)

SEDIMENTARY ROCKS

(Areas of Sedimentary rocks are shown by patterns of parallel lines.)

**Nn**

Nussbaum formation  
(sand, gravel and silt)

**Kp**

Pierre shale  
(gray shale with concretions)

**Kn**

Niobrara formation  
(shale and limestone)

**Kcr**

Carlisle shale  
(shale and sandstone)

**Kgn**

Greenhorn limestone  
(limestone and shale in alternate strata)

**Kgs**

Graneros shale  
(gray shale)

**Kd**

Dakota sandstone  
(gray sandstone and shale)

**Jm**

Morrison formation  
(variegated clay and sandstone; gypsum)

**Jf**

Fountain formation  
(red shale, sandstone, and conglomerate)

**Gm**

Millsap limestone  
(limestone and shale)

**Sh**

Harding sandstone  
(white sandstone)


(Areas of ancient crystalline rocks and metamorphic rocks of unknown origin are shown by patterns of short dashes.)

**As**


Schist and granite  
(igneous and metamorphic)

Faults

Sections



Henry Gannett, Chief Topographer.  
E. M. Douglas, Topographer in charge.  
Triangulation by A. H. Thompson.  
Topography by R. O. Gordon and W. J. Lloyd.  
Surveyed in 1894.



Scale 1:25000  
Contour Interval 50 feet.  
Datum is mean sea level.  
Edition of Jan. 1897.

Geology by G. K. Gilbert.  
Assisted by Robt. T. Hill.  
Surveyed in 1893.

Legend is continued on the left margin.



(Pikes Peak) 38°30' 105°00'

(Boerforno Park) 38°00' 105°00'

SURFICIAL ROCKS






- LATER  
Alluvium (bottom lands)  
Pe  
EARLIER  
Alluvium (gravel, sand, and silt capping terraces and mesas)

SEDIMENTARY ROCKS

- NEOCENE  
Nn Nussbaum formation (sand, gravel, sand, silt)  
Kp Pierre shale (gray shale with concretions)  
Kn Niobrara formation (shale and limestone)  
Kcr Carlile shale (shale and sandstone)  
Kgn Greenhorn limestone (limestone and shale in alternate strata)  
Kgs Graneros shale (gray shale)  
Kd Dakota sandstone (gray sandstone and shale)
- CRETACEOUS  
Subdivisions of the Benton formation
- JURATRIAS  
Jm Morrison formation (variegated clay and sandstone, gypsum)  
Jf Fountain formation (red shale, sandstone, and conglomerate)  
Cm Millsap limestone (limestone and shale)  
Sh Harding sandstone (white sandstone)
- CARBONIFEROUS  
SILURIAN  
Rg Schist and granite (gneiss and metamorphic)


Faults

- Known productive formations  
Kn Niobrara formation (limestone for lime and flour)  
Kcr Carlile shale (sandstone used for foundations)  
Kd Dakota sandstone (largely available for building; upper part contains fire clay)  
Jm Morrison formation (local deposits of gypsum)  
Cm Millsap limestone (local developments of marble)




Henry Gannett, Chief Topographer.  
E.M. Douglas, Topographer in charge.  
Triangulation by A.H. Thompson.  
Topography by R.O. Gordon and W.J. Lloyd.  
Surveyed in 1894.

Geology by G.K. Gilbert.  
Assisted by Robt. T. Hill.  
Surveyed in 1893.








Modeled by Edwin E. Howell.  
Engraved by A. Hoen & Co.

This sheet is a view from above of the flexed and faulted surface of the Dakota sandstone as it would appear if restored wherever it has been removed by erosion and laid bare wherever it is covered by other formations. It is based on the photograph of a plaster model in which the vertical and horizontal scales are the same,  $\frac{1}{125000}$ . To give the proper effect the sheet should be so placed that the light, from window or lamp, falls on it from the right.

Geology by G. K. Gilbert.  
Surveyed in 1893.





LEGEND

- Area of artesian water which will probably yield flowing wells
- Area of artesian water which will probably yield pumping wells
- Area of artesian water formation which will probably not yield water freely
- Area not containing artesian water formation
- Contours showing approximate depth below the surface of the ground of the highest water-bearing bed of the Dakota sand stone. Contour interval is 200 feet. Figures show depth in feet
- Artesian wells

Henry Gannett, Chief Topographer.  
E.M. Douglas, Topographer in charge.  
Triangulation by A.H. Thompson.  
Topography by R.O. Gordon and W.J. Lloyd.  
Surveyed in 1894.


Scale 1:25000  
Miles  
Kilometers

Contour Interval 50 feet.  
Datum is mean Sea level.  
Edition of Jan. 1897.


Geology by G.K. Gilbert.  
Surveyed in 1893.



# COLUMNAR SECTIONS



NOTE:—The right hand edge of the section column is drawn as the profile of an ideal cliff; the stronger beds, chiefly sandstones and limestones, project beyond the general face.



**NAMES OF FORMATIONS—COMPARATIVE TABLE.**

| PERIOD.          | NAMES AND SYMBOLS USED IN THIS FOLIO. | NAMES USED IN REPORT ON UNDERGROUND WATER, 17TH ANNUAL REPORT, U. S. GEOL. SURVEY, 1896. | CROSS: PIKES PEAK FOLIO, U. S. GEOLOGICAL SURVEY, 1894. | ELDRIDGE: REPORT ON THE FLORENCE OIL FIELD, AM. INSTITUTE OF MINEING ENGINEERS, 1891. | HAYDEN: ATLAS OF COLORADO, 1881.                |
|------------------|---------------------------------------|--|---|---|---|
| PLEISTOCENE      | Later alluvium.                       | Pl   | Alluvium.   | Quaternary.   | Alluvium.                                       |
|                  | Earlier alluvium.                     | Pe   | Terrace sands and gravels.                              |   |   |
| NEOCENE          | Nussbaum.                             | Nn   | Upland sands and gravels.                               |   |   |
| CRETACEOUS       | Pierre.                               | Kp   | Pierre.   | Montana (comprising Fox Hills and Pierre).  | Fox Hills (including also Fort Pierre).         |
|                  | Niobrara.                             | Kn   | Apishapa. Timpas.                                       |   |   |
|                  | Carlile.                              | Kcr  | Carlile.  | Colorado.   | Colorado (comprising Niobrara and Benton).      |
|                  | Greenhorn.                            | Kgn  | Greenhorn.  |   | Colorado (comprising Niobrara and Fort Benton). |
|                  | Graneros.                             | Kgs  | Graneros.   |   |   |
|                  | Dakota.                               | Kd   | Dakota.   | Dakota.   | Dakota.   |
| JURATRIAS        | Morrison.                             | Jm   | Morrison.   | Jura.   | Variegated Beds, etc.                           |
|                  | Fountain.                             | Jf   | Juratrias.  | Fountain (classed as Carboniferous).  | Trias.  |
| ARCH. SIL. CARB. | Millsap.                              | Cm   | Millsap.  | Carboniferous.  | Carboniferous.                                  |
|                  | Harding.                              | Sh   | Harding.  | Silurian.   |   |
| ARCH. ?          | Archean ?                             | Rs   |   | Archean.  | Archean.  |






Fig. 17.




Fig. 18.

FIGS. 17, 18.—*INOCERAMUS DEFORMIS*.

Single valves of a bivalve shell found fossil in the limestone at the base of the Niobrara formation. Fig. 17 shows the side of a specimen of ordinary size; fig. 18 the edge of a small individual. The mud which once filled the shells was hardened to stone and the shells were afterward broken away, leaving molds of the interior.



FIG. 19.—OYSTER SHELLS ATTACHED TO A LARGER SHELL (*INOCERAMUS*).

From a photograph; natural size.

Such groups are found in several formations, but they are peculiarly abundant in the calcareous shales near the bottom of the Niobrara.




Fig. 20.




Fig. 21.

FIGS. 20, 21.—*INOCERAMUS LABIATUS*.

A fossil bivalve shell occurring in abundance in certain layers of the Greenhorn limestone. Fig. 20 shows an individual of moderate size; fig. 21 a small individual in which the concentric ridges are unusually strong.



FIG. 22.—*HETERO CERAS NEBRASCENSE*.

From a photograph; natural size.

A fossil shell occurring in the Tepee zone of the Pierre shale and best preserved in concretions. This specimen, which includes two-thirds of the whole individual, is more nearly complete than the specimens usually found. Fragments 3 or 4 inches in length are comparatively common.



Fig. 23.




Fig. 24.

FIGS. 23, 24.—*PLACENTICERAS PLACENTA*.

Two views, natural size, of a small individual; specimens often have a diameter several times greater. The fossil shell is found only in the Tepee zone of the Pierre shale. Its nearest relative among living shells is the Nautilus.



FIG. 25.—THE GREENHORN LIMESTONE.

From a photograph.

A characteristic outcrop, showing the alternation of limestone ledges with softer layers of shale.



FIG. 26.—*LUCINA OCCIDENTALIS*.

From a photograph; natural size.

This fossil shell is a bivalve, occurring in the Tepee zone of the Pierre shale. It is abundant in the limestone cores of the Tepee buttes (fig. 27), and is the most characteristic fossil of those cores.




FIG. 27.—A TEPEE BUTTE.

Part of the limestone core is seen in the crest. These buttes are characteristic of the Tepee zone of the Pierre shale.



ary  
deposited  
forms char-  
and gravel,  
kames. The  
called glacial  
to the adjacent  
is usual also to  
sits of the sea and  
made at the same

#### OF ROCKS.

er distinguished according to  
ages, for rocks were not formed all  
time, but from age to age in the earth's  
Classification by age is independent of  
igneous, sedimentary, and surficial rocks  
of the same age.

the predominant material of a rock mass  
initially the same, and it is bounded by rocks  
erent materials, it is convenient to call the  
throughout its extent a *formation*, and such  
ation is the unit of geologic mapping.

eral formations considered together are  
ated a *system*. The time taken for the  
sition of a formation is called an *epoch*, and  
time taken for that of a system, or some  
fraction of a system, a *period*. The rocks  
napped by formations, and the formations are  
ified into systems. The rocks composing a  
em and the time taken for its deposition are  
ven the same name, as, for instance, Cambrian  
system, Cambrian period.

As sedimentary deposits or strata accumulate  
the younger rest on those that are older, and the  
relative ages of the deposits may be discovered  
by observing their relative positions. This rela-  
tionship holds except in regions of intense distur-  
bance; sometimes in such regions the disturb-  
ance of the beds has been so great that their  
position is reversed, and it is often difficult to  
determine the relative ages of the beds from their  
positions; then *fossils*, or the remains of plants  
and animals, are a guide to show which of two  
or more formations is the oldest.

Strata often contain the remains of plants and  
animals which lived in the sea or were washed  
from the land into lakes or seas or were buried in  
surficial deposits on the land. Rocks that con-  
tain the remains of life are called *fossiliferous*.  
By studying these remains, or fossils, it has been  
found that the species of each period of the earth's  
history have to a great extent differed from those  
of other periods. Only the simpler kinds of  
marine life existed when the oldest fossiliferous  
rocks were deposited. From time to time more  
complex kinds developed, and as the simpler ones  
lived on in modified forms life became more  
varied. But during each period there lived pecu-  
liar forms, which did not exist in earlier times  
and have not existed since; these are charac-  
teristic types, and they define the age of any bed of  
rock in which they are found. Other types  
passed on from period to period, and thus linked  
the systems together and formed a chain of life  
from the time of the oldest fossiliferous rocks to  
the present.

When two formations are remote one from the  
other and it is impossible to observe their relative  
positions, the characteristic fossil types found in  
them may determine which was deposited first.

Fossil remains found in the rocks of different  
areas, provinces, and continents, afford the most  
important means for combining local histories  
into a general earth history.

*Colors and patterns.*—To show the relative ages  
of strata, the history of the sedimentary rocks is  
divided into periods. The names of the periods  
in proper order (from new to old), with the color  
or colors and symbol assigned to each, are here  
given. The names of certain subdivisions of the  
periods, frequently used in geologic writings, are  
bracketed against the appropriate period name.

To distinguish the sedimentary formations of  
any one period from those of another the patterns  
for the formations of each period are printed in  
the appropriate period-color, with the exception  
of the first (Pleistocene) and the last (Archean).  
The formations of any one period, with the  
exception of Pleistocene and Archean, are distin-

guished from one another by different patterns,  
made of parallel straight lines. Two tints of the

| PERIOD.                                | SYMBOL. | COLOR.         |
|--|---------|----------------|
| Pleistocene                            | P       | Any colors.    |
| Neocene { Pliocene }<br>{ Miocene }    | N       | Bufs.          |
| Eocene (including Oligocene)           | E       | Olive-browns.  |
| Cretaceous                             | K       | Olive-greens.  |
| Juratrias { Jurassic }<br>{ Triassic } | J       | Blue-greens.   |
| Carboniferous (including Permian)      | C       | Blues.         |
| Devonian                               | D       | Blue-purples.  |
| Silurian (including Ordovician)        | S       | Red-purples.   |
| Cambrian                               | C       | Pinks.         |
| Algonkian                              | A       | Orange-browns. |
| Archean                                | R       | Any colors.    |

period-color are used: a pale tint (the underprint)  
is printed evenly over the whole surface represent-  
ing the period; a dark tint (the overprint) brings  
out the different patterns representing formations.  
Each formation is furthermore given a letter-  
symbol of the period. In the case of a sedimentary  
formation of uncertain age the pattern is  
printed on white ground in the color of the period  
to which the formation is supposed to belong,  
the letter-symbol of the period being omitted.

The number and extent of surficial formations  
of the Pleistocene render them so important that,  
to distinguish them from those of other periods  
and from the igneous rocks, patterns of dots and  
circles are used. These may be printed in any  
colors.

The origin of the Archean rocks is not fully  
settled. Many of them are certainly igneous.  
Whether sedimentary rocks are also included is  
not determined. The Archean rocks, and all meta-  
morphic rocks of unknown origin, of whatever age,  
are represented on the maps by patterns consisting  
of short dashes irregularly placed. These are  
printed in any color, and may be darker or lighter  
than the background. If the rock is a schist the  
dashes or hachures may be arranged in wavy par-  
allel lines. If the rock is known to be of sedi-  
mentary origin the hachure patterns may be com-  
bined with the parallel-line patterns of sedi-  
mentary formations. If the metamorphic rock is  
recognized as having been originally igneous, the  
hachures may be combined with the igneous  
pattern.

Known igneous formations are represented by  
patterns of triangles or rhombs printed in any  
brilliant color. If the formation is of known age  
the letter-symbol of the formation is preceded by  
the capital letter-symbol of the proper period.  
If the age of the formation is unknown the letter-  
symbol consists of small letters which suggest the  
name of the rocks.

#### THE VARIOUS GEOLOGIC SHEETS.

*Areal sheet.*—This sheet shows the areas occu-  
pied by the various formations. On the margin is  
a *legend*, which is the key to the map. To ascer-  
tain the meaning of any particular colored pat-  
tern and its letter-symbol on the map the reader  
should look for that color, pattern, and symbol in  
the legend, where he will find the name and  
description of the formation. If it is desired to  
find any given formation, its name should be  
sought in the legend and its color and pattern  
noted, when the areas on the map corresponding  
in color and pattern may be traced out.

The legend is also a partial statement of the  
geologic history. The formations are arranged  
according to origin into surficial, sedimentary, and  
igneous, and within each class are placed in the  
order of age, so far as known, the youngest at the  
top.

*Economic sheet.*—This sheet represents the dis-  
tribution of useful minerals, the occurrence of  
artesian water, or other facts of economic interest,  
showing their relations to the features of topog-  
raphy and to the geologic formations. All the  
formations which appear on the areal sheet are  
shown on this sheet by fainter color-patterns.  
The areal geology, thus printed, affords a subdued  
background upon which the areas of productive  
formations may be emphasized by strong colors.  
A symbol for mines is introduced at each occur-  
rence, accompanied by the name of the principal  
mineral mined or of the stone quarried.

*Structure-section sheet.*—This sheet exhibits the  
relations of the formations beneath the surface.

In cliffs, canyons, shafts, and other natural and

artificial cuttings, the relations of different beds  
to one another may be seen. Any cutting which  
exhibits those relations is called a *section*, and the  
same name is applied to a diagram representing  
the relations. The arrangement of rocks in the  
earth is the earth's *structure*, and a section exhibit-  
ing this arrangement is called a *structure section*.

The geologist is not limited, however, to the  
natural and artificial cuttings for his information  
concerning the earth's structure. Knowing the  
manner of the formation of rocks, and having  
traced out the relations among beds on the sur-  
face, he can infer their relative positions after  
they pass beneath the surface, draw sections  
which represent the structure of the earth to a  
considerable depth, and construct a diagram  
exhibiting what would be seen in the side of a  
cutting many miles long and several thousand feet  
deep. This is illustrated in the following figure:




Fig. 2.—Sketch showing a vertical section in the front of the picture, with a landscape above.

The figure represents a landscape which is cut  
off sharply in the foreground by a vertical plane  
that cuts a section so as to show the underground  
relations of the rocks.

The kinds of rock are indicated in the section  
by appropriate symbols of lines, dots, and dashes.  
These symbols admit of much variation, but the  
following are generally used in sections to repre-  
sent the commoner kinds of rock:




Fig. 3.—Symbols used to represent different kinds of rock.

The plateau in fig. 2 presents toward the lower  
land an escarpment, or front, which is made up  
of sandstones, forming the cliffs, and shales, con-  
stituting the slopes, as shown at the extreme left  
of the section.

The broad belt of lower land is traversed by  
several ridges, which are seen in the section to  
correspond to beds of sandstone that rise to the  
surface. The upturned edges of these beds form  
the ridges, and the intermediate valleys follow the  
outcrops of limestone and calcareous shales.

Where the edges of the strata appear at the  
surface their thickness can be measured and the  
angles at which they dip below the surface can be  
observed. Thus their positions underground can  
be inferred.

When strata which are thus inclined are traced  
underground in mining, or by inference, it is fre-  
quently observed that they form troughs or arches,  
such as the section shows. But these sandstones,  
shales, and limestones were deposited beneath the  
sea in nearly flat sheets. That they are now bent  
and folded is regarded as proof that forces exist  
which have from time to time caused the earth's  
surface to wrinkle along certain zones.

On the right of the sketch the section is com-  
posed of schists which are traversed by masses of  
igneous rock. The schists are much contorted  
and their arrangement underground can not be  
inferred. Hence that portion of the section  
delineates what is probably true but is not  
known by observation or well-founded inference.

In fig. 2 there are three sets of formations, dis-

tinguished by their underground relations. The  
first of these, seen at the left of the section, is the  
set of sandstones and shales, which lie in a hori-  
zontal position. These sedimentary strata are  
now high above the sea, forming a plateau, and  
their change of elevation shows that a portion  
of the earth's mass has swelled upward from a  
lower to a higher level. The strata of this set are  
parallel, a relation which is called *conformable*.

The second set of formations consists of strata  
which form arches and troughs. These strata  
were once continuous, but the crests of the arches  
have been removed by degradation. The beds,  
like those of the first set, are conformable.

The horizontal strata of the plateau rest upon  
the upturned, eroded edges of the beds of the  
second set at the left of the section. The over-  
lying deposits are, from their position, evidently  
younger than the underlying formations, and the  
bending and degradation of the older strata must  
have occurred between the deposition of the older  
beds and the accumulation of the younger. When  
younger strata thus rest upon an eroded surface  
of older strata the relation between the two is an  
*unconformable* one, and their surface of contact is  
an *unconformity*.

The third set of formations consists of crystal-  
line schists and igneous rocks. At some period  
of their history the schists were plicated by pres-  
sure and traversed by eruptions of molten rock.  
But this pressure and intrusion of igneous rocks  
have not affected the overlying strata of the second  
set. Thus it is evident that an interval of consid-  
erable duration elapsed between the formation of  
the schists and the beginning of deposition of the  
strata of the second set. During this interval the  
schists suffered metamorphism; they were the  
scene of eruptive activity; and they were deeply  
eroded. The contact between the second and  
third sets, marking a time interval between two  
periods of rock formation, is another uncon-  
formity.

The section and landscape in fig. 2 are ideal,  
but they illustrate relations which actually occur.  
The sections in the structure-section sheet are  
related to the maps as the section in the figure is  
related to the landscape. The profiles of the sur-  
face in the section correspond to the actual slopes  
of the ground along the section line, and the  
depth of any mineral-producing or water-bearing  
stratum which appears in the section may be  
measured from the surface by using the scale of  
the map.

*Columnar-section sheet.*—This sheet contains a  
concise description of the rock formations which  
occur in the quadrangle. The diagrams and  
verbal statements form a summary of the facts  
relating to the character of the rocks, to the thick-  
nesses of the formations, and to the order of  
accumulation of successive deposits.

The rocks are described under the correspond-  
ing heading, and their characters are indicated in  
the columnar diagrams by appropriate symbols.  
The thicknesses of formations are given under  
the heading "Thickness in feet," in figures which  
state the least and greatest measurements. The  
average thickness of each formation is shown in  
the column, which is drawn to a scale—usually  
1000 feet to 1 inch. The order of accumulation  
of the sediments is shown in the columnar arrange-  
ment: the oldest formation is placed at the bot-  
tom of the column, the youngest at the top,  
and igneous rocks or other formations, when  
present, are indicated in their proper relations.

The formations are combined into systems  
which correspond with the periods of geologic  
history. Thus the ages of the rocks are shown,  
and also the total thickness of each system.

The intervals of time which correspond to  
events of uplift and degradation and constitute  
interruptions of deposition of sediments may be  
indicated graphically or by the word "unconform-  
ity," printed in the columnar section.

Each formation shown in the columnar section  
is accompanied by its name, a description of its  
character, and its letter-symbol as used in the  
maps and their legends.

CHARLES D. WALCOTT,

Revised Me