

dup



**United States
Department of
Agriculture**

Agricultural
Research
Service

Index-Catalogue of Medical and Veterinary Zoology

Supplement 23, Part 6,
Section C. Treatment

Parasite-Subject Catalogue
Subject Headings and Treatment

**United States
Department of
Agriculture**

Agricultural
Research
Service

May 1982

Index-Catalogue of Medical and Veterinary Zoology

Supplement 23, Part 6,
Section C. Treatment

Parasite-Subject Catalogue Subject Headings and Treatment

By

Shirley J. Edwards, In Charge

Martha W. Hood, Zoologist

Judith H. Shaw, Zoologist

Jane D. Rayburn, Technical Information Specialist

Margie D. Kirby, Technical Information Specialist

Deborah T. Hanfman, Technical Information Specialist

Judith A. Zidar, Technical Information Specialist

Preface

The Index-Catalogue of Medical and Veterinary Zoology is an index to the world's literature on animal parasites of animals, including man. The Catalogue is distributed to qualified individuals and libraries throughout the world without charge. It has been maintained in cumulative files since 1892. Only the Author Catalogue has been published in its entirety. A revision of the Author Catalogue of the Index-Catalogue of Medical and Veterinary Zoology, consisting of Parts 1 to 18, was published during the period 1932-52. Beginning in 1953, a series of supplements designed to publish the backlog was initiated. This was completed with Supplement 6, published in 1956. From 1956 to 1964, supplements covering authors A to Z were issued on an annual basis.

Beginning with Supplement 15, the Parasite-Subject Catalogues, containing indices to the author references, have been issued. The Author Catalogues of Supplements 15-21 continued the format of previous supplements. Users should note that for each reference in the Author Catalogues of these supplements the author(s) plus the date and letter (e.g., Smith, J.; and Doe, L., 1978 b) are the key to all items in the Parasite-Subject Catalogues derived from that reference. In other words, when using the Parasite-Subject Catalogues of Supplements 15-21, it is necessary to consult the Author Catalogue of the corresponding supplement for complete bibliographic information.

Commencing with Supplement 22, basic bibliographic information is included with each entry in Parts 2-7. It should be emphasized, however, that it will still be useful to consult the Author Catalogue for a variety of other information that may be found there: Title of the reference, translated title, language of text and summaries, issue date, library from which the original may be obtained, published corrections, related references by the same author, and other miscellaneous information.

Each supplement consists of the following parts:

- Part 1, Authors: A-Z
- Part 2, Parasite-Subject Catalogue: Parasites: Protozoa
- Part 3, Parasite-Subject Catalogue: Parasites: Trematoda and Cestoda
- Part 4, Parasite-Subject Catalogue: Parasites: Nematoda and Acanthocephala
- Part 5, Parasite-Subject Catalogue: Parasites: Arthropoda and Miscellaneous Phyla
- Part 6, Parasite-Subject Catalogue: Subject Headings and Treatment
- Part 7, Parasite-Subject Catalogue: Hosts

Users should bear in mind that this is an Index-Catalogue, not a treatise, and should not expect to find reasons for any given entry. Nor does citing of synonymy mean that it is necessarily correct. The same statement holds for hosts, locations, localities, authorship of taxa, designation of new taxa, etc. These items are cited as given by the author(s) of the publication being indexed.

The information included in any given supplement represents only the publications that have been indexed in that supplement; and therefore, exclusion of, or limited entries for, any given author or parasite has no significance. No pretension is made for completeness, and assistance in correcting errors or obtaining additional information is appreciated. Reprints of papers on parasitology are requested.


Explanatory Note

Author Catalogue

The Author Catalogue (Part 1 of each supplement) contains full bibliographic information for each publication indexed during the compilation of that supplement. A symbol for the library from which the original publication may be obtained is given at the end of each entry, e.g., Wa, Wm, Wc, etc. A key to these library symbols may be found in Supplements 10 and 20. A list of serial abbreviations new to our files is published at the beginning of each Author Catalogue.

Parasite Catalogues

The Parasite Catalogues (Parts 2-5 of each supplement) are divided by parasite phyla (Protozoa, Trematoda, etc.). They are arranged alphabetically by genera, parasitic diseases, and higher taxa and then alphabetically by species within genera. Entries under each heading are in turn arranged alphabetically by authors and then chronologically for each author. Each entry consists of the name of the parasite or parasitic disease, the author(s) of the publication, date, abbreviated title of the publication, volume, number, inclusive pages, and a subheading. Illustrations of parasites are indicated by the word *illus.* following the name of the parasite.



A variety of information is found indented beneath the author line(s) of each entry: Classification, hosts, synonymy, keys, treatment, etc. Subheadings are guides to the subject matter of the publication.

- (1) **Classification:** In entries based on systematic articles, the subheading may give the higher taxa in which the taxon has been placed or it may list the lower taxa included in a higher taxon.
- (2) **Hosts:** The only hosts recorded are those that pertain directly to the author's own work. Scientific host names are used unless the author gives only common names, in which case the host names are given exactly as in the original publication.

Locations of parasites in or on hosts are given in parentheses (). Where a host-parasite relationship is well known, a host may be given under a parasite name and not in the Host Catalogue; this applies particularly to parasites of medical and veterinary importance and of worldwide distribution. A + before the host name on the parasite entry means that no host entry was made for this particular reference.

- (3) **Synonymy:** Usually only those synonyms which the author indicates as new, or which are new to the files of the Index-Catalogue of Medical and Veterinary Zoology, are given.
- (4) **Keys:** The subheading "key" indicates that the name is included in a taxonomic key.
- (5) **Treatment:** When there are several antiparasitic agents mentioned in a publication, a general term is used in the subheading, e.g., anthelmintics, insecticides, protozoocides. However, in the Treatment Catalogue, all agents tested by the investigator(s) are listed.
- (6) **Geographic Distribution:** When there are multiple hosts and geographic localities, the appropriate locality is recorded after each host name; when the hosts of a parasite are all from one locality, they are recorded as "all from" this locality.
- (7) **Other Subject Matter:** Phrases indicate other subject matter discussed (e.g., immunity, metabolism, morphology, etc.).

Subject Headings Catalogue

The Subject Headings Catalogue (the first section of Part 6 of each supplement) is an alphabetic arrangement of entry terms from a controlled list of subject headings. Each entry consists of the subject heading, bibliographic information, and a subheading reflecting the information contained in the paper. Subject headings with numerous entries are separated into alphabetized subdivisions, e.g.,

Immunity
Immunity, Agglutination
Immunity, Allergy

Treatment Catalogue

In the Treatment Catalogue (a section of Part 6 of each supplement), all entries referring to one antiparasitic agent are grouped under one heading (regardless of the name used by the investigator) and are then listed alphabetically by author. Other names for the same agent are cross-referenced to the name used for filing. When generic and chemical names are available, preference is given to those names as headings

However, when host common names are in Cyrillic alphabet languages, host Latin names are assigned and listed instead of the common name; these are in square brackets [].

rather than to trade names or code numbers and letters. Code number designations for compounds are entered in the Number Index in numerical order and cross-referenced to the name under which they are listed in the alphabetical section. Salts of a compound are usually grouped together, e.g., piperazine adipate, piperazine citrate, etc., are all listed under Piperazine. Sometimes verifying synonymy of drug names is impossible; consequently, groupings and cross-references are not always authenticated although as many as possible have been checked with reliable sources. In some instances, the cross-references are based entirely on information in papers indexed and verification was not possible. Foreign language terminology has been anglicized where feasible. Chemosterilants, Molluscicides, and Repellents are entered under these three collective headings and not under the individual chemical. The format is the same as the parasite entries: Entry term (in this case, drug name), bibliographic information, and subheading.

Host Catalogue

The Host Catalogue (Part 7 of each supplement) is arranged alphabetically by genera, common names, and higher taxa and then alphabetically by species within genera. Nominate subspecies are interfiled with the species. Entries under each heading are in turn arranged alphabetically by author(s) and then chronologically for each author. The format is the same as in the other Catalogues, i.e., entry term (in this case, host name), bibliographic information, and subheading. Indented beneath the author line(s) of each host entry are all the parasites of a particular phylum that were reported from this host in the paper in question. Body locations of these parasites will be found in parentheses () either in the subheading or with the host name. Experimental infection is reported as such. When there are multiple parasites and geographic localities, the appropriate locality is recorded after each parasite name; when the parasites from this host are all from one locality, they are recorded as "all from" this locality. When authors use only common names of hosts, scientific names are cautiously supplied from authoritative sources after

careful consideration. Cross-references from the common name used by the author to the scientific name supplied by the Index-Catalogue are filed among the host entries. Such supplied names are given in square brackets []. If a scientific name cannot be supplied, English common names are used. Scientific names or English common names are always supplied for common names in Cyrillic alphabet languages, and no cross-references are made. Surveys of parasites of humans and domestic animals are often indexed under geographic headings and entered in Part 6, Subject Headings, in addition to appearing in the Host Catalogue. In this case, all parasite phyla are grouped under the same host entry, and individual parasite entries are not included in the Parasite Catalogue.

Visitors are welcome to come to the Animal Parasitology Institute to use the cumulative files. Arrangements should be made in advance for lengthy visits.

All correspondence should be addressed to:

Index-Catalogue of Medical and Veterinary Zoology
Animal Parasitology Institute
USDA, ARS, BARC-East, Building 1180
Beltsville, Maryland 20705 U.S.A.

It is hoped that these Catalogues will serve as a useful tool to workers in the field of parasitology. Users are requested to preserve the Catalogues, since they are not designed for general distribution and the edition is limited.

The compilers thank the staffs of the National Agricultural Library, the National Library of Medicine, and all other libraries who have aided us invaluablely by making publications available to us.

Trade names are used in this publication solely for the purpose of providing specific information. Mention of a trade name does not constitute a guarantee or warranty of the product by the U.S. Department of Agriculture or an endorsement by the Department over other products not mentioned.

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...
...the ...

NUMBER INDEX

1-Adamantamine. See Amantadine.

PJT 1. See Cypermethrin.

Uroporphyrin I. See Uroporphyrin I.

Aerol-2 aerosol. See Aerol-2 aerosol.

Banminth II. See Morantel.

Aza 3-emetine. See Aza 3-emetine.

Cibacron blue F3GA. See Cibacron blue F3GA.

Esb₃. See Sulfachloropyrazine.

BB-04. See Piperazine or Thiabendazole.

IA-4. See 8-Chloro-2-[(diethylamino)ethyl]-2H-[1]-benzothiopyrano[4,3,2-cd]indazole-5-methanol monomethane-sulfonate.

IA-4 N-oxide. See 8-Chloro-2-[(diethylamino)ethyl]-2H-[1]-benzothiopyrano[4,3,2-cd]indazole-5-methanol monomethane-sulfonate.

Atgard V. See Dichlorvos.

5-Azacytidine. See 5-Azacytidine..

5-Chloroethylthiamine. See Beclotiamine.

Flagyl V. See Metronidazole.

5-FUDR. See Floxuridine.

6-Aminonicotinamide. See 6-Aminonicotinamide.

6-Azauracil. See 6-Azauracil.

Hipolen-6. See Crufomate.

Ruelene 6-R. See Crufomate.

DL-7-Azatryptophan. See DL-7-Azatryptophan.

Benzazon VII. See Benzazon VII.

NUMBER INDEX -- Continued.

7-Chlorolincomycin. See Clindamycin.

7-Deazaadenosine. See Tubercidin.

TN-7. See Nitrimidazine.

8-Aza-adenine. See 8-Aza-adenine.

8-Azaguanine. See Guanazolo.

Acetoxy-9-ellipticine. See Acetoxy-9-ellipticine.

Bischloroethyldeuteroporphyrin IX. See Bischloroethyldeuteroporphyrin IX.

Protoporphyrin IX. See Protoporphyrin IX.

Linton-X. See Toxaphene.

RC12. See 5-Bromo-4[bis-(2-diethylamino ethyl)-amino] veratrole.

K-15. See 2-Phenyl-4H-1,3,2-benzodioxaphosphorin-2-sulfide.

Nemicide L 15. See Tetramisole.

Lindatox-20. See Benzene hexachloride.

Lypor 20. See Temephos.

Poron 20. See Phosmet.

BW 21Z. See Permethrin.

U-21. See Clindamycin.

U-24. See N-Demethyl-4'-pentyl clindamycin hydrochloride.

Ardenone-25. See Ardenone-25.

Coyden 25. See Meticlorpindol.

Gasil 35. See Gasil 35.

Carbetox 37. See Carbetox 37.

Dursban 44 Insecticide Formulation. See Chlorpyrifos.

NUMBER INDEX -- Continued.

- COBAN 45. See Monensin.
- Abi-Zet₅₀. See Pyrimethamine or Sulfamethazine.
- Asunto1 50. See Coumaphos.
- Baytex-50. See Fenthion.
- Liquamycin injectable L-50. See Oxytetracycline.
- Liquamycin (T-50). See Oxytetracycline.
- Malathion 50 EC. See Malathion.
- Paramar M-50. See Paramar M-50.
- Perdix PE 50. See Butonate.
- Sumithion 50 EC. See Fenitrothion.
- Malathion 57. See Malathion.
- VUAgT-71. See VUAgT-71.
- C-076. See Avermectin B_{1a}.
- Marvex Super-100. See Dichlorvos.
- Nuvan 100 EC. See Dichlorvos.
- Terramycin 100. See Oxytetracycline.
- Schistocide T-109. See 2,8-Dimethyl-5-β-diethyl-amino-ethylamino-thiochromone.
- GX-118. See Phosmet.
- Starbar GX-118. See Phosmet.
- NRDC 143. See Permethrin.
- NRDC 149. See Cypermethrin.
- F 151. See (Diamino-4,6-triazinyl-1,3,5-amino-2)-4-phenyl-arsino di (D-thio-3-amino-2-methyl-3-butyric acid).
- F 159. See F 159.

NUMBER INDEX -- Continued.

- ICRF 159. See Razoxane.
- NRDC 161. See Decamethrin.
- NF-180. See Furazolidone.
- ICRF 192. See ICRF 192.
- Gasil 200. See Gasil 200.
- Liquamycin/LA-200. See Oxytetracycline.
- Nivaquine 200. See Chloroquine.
- Terramycin/LA (T-200). See Oxytetracycline.
- Hoechst S-201. See Bis-(β-carbohydrazide-ethyl) sulfone.
- Bayer 205. See Suramin.
- SPA-S-222. See Mepartricin or Sodium lauryl sulfate.
- MK-302. See Arprinocid.
- TF-302. See Crufomate.
- SAN 322 I. See Propetamphos.
- EQ-335. See Benzene hexachloride.
- G-418. See G-418.
- Pyractone M429. See Bucarpolate or Pyrethrins.
- MK436. See 3-(1-Methyl-5-nitroimidazol-2-yl)-3α,4,5,6,7α-hexahydro-1,2-benzisoxazole.
- A 445. See Thiophanate.
- WR 448. See Dapsone.
- Antibiotic X-464. See Nigericin.
- TM-481. See Lonomycin.
- Diethylaminoethyl-dextran 500. See Diethylaminoethyl-dextran 500.
- Fasigyn 500. See Tinidazole.

NUMBER INDEX -- Continued.

Zoecon 515. See Methoprene.

SKF 525-A. See Proadifen hydrochloride.

MW 591. See Lasalocid.

NF-602. See Furaltadone.

Hoechst S-616. See N'-(3-Chloro-2,4,6-trimethylphenyl)-N,N-diethylenediamine hydrochloride.

Searle SN 654. See Mepartricin.

Ciba-Geigy B 663. See Clofazimine.

Preparation-665. See Mekarzole.

HOE 668. See p-(4-Amidino-phenoxy)-benzaldehyde-p-amidino-phenylhydrazone dihydrochloride.

CI-679. See 2,4-Diamino-6-[(3,4-dichlorobenzyl)-nitrosoamino]-quinazoline.

Hoechst S-688. See Maleinyl-4-(3-chloro-p-tolyl)piperazide.

MW 764. See Narasin.

Pyractone M817. See Piperonyl butoxide or Pyrethrins.

8-Aza 26-diaminopurine. See 8-Aza 26-diaminopurine.

HOE-881. See Fenbendazole.

Hoe 881V. See Fenbendazole.

NF-902. See Furaltadone.

MK 905. See Cambendazole.

MK-910. See 1-Methyl-2-(p-fluorophenyl)-5-nitroimidazole.

MK915. See Flunidazole.

L. 13/59. See Trichlorfon.

WR 1,544. See Chloroquine.

NUMBER INDEX -- Continued.

Blue dextran (2000). See Blue dextran (2000).

Amoebicide 2004. See Chlorphenoxamide or Tinidazole.

Bayer 2349. See Trichlorfon.

Bayer 2353. See Niclosamide.

C-C 2481. See Cyclobendazole.

Wander compound 2495. See 4,4'-(4-Methyl-2-imidazolin-2-yl) terephthalanilide dimethane sulphate.

Bay 2502. See Nifurtimox.

Bayer 2502. See Nifurtimox.

S-2539 Forte. See Phenothrin.

U.K. 2679. See Pyrantel.

Wander compound 2783. See [4-(4-Methyl-2-imidazolin-2-yl)-4'-[(p-4-methyl-2-imidazolin-2-yl)phenyl]carbamoyl]-carbanilide-dimethanesulphonate.

Hoechst 2910. See Benoxafos.

WR 2,975. See Primaquine.

WR 2,976. See Quinine.

WR 2978. See Pyrimethamine.

Hoe 2982. See Heptenophos.

3024 I.C.I. See C,C-Diallyl-bis-(4-amino-2-methyl-6-quinolyl) malonamide.

Dow M3615. See Chlorpyrifos.

ME 3625. See Niclofolan.

UK 3883. See 2-(N-Isopropylaminomethyl)-6-methyl-7-nitro-1,2,3,4-tetrahydroquinoline.

DE-3936. See Lonomycin.

Dow M3983. See Chlorpyrifos.

NUMBER INDEX -- Continued.

Compound 4072. See Chlorfenvinphos.

GC 4072. See Chlorfenvinphos.

Pfizer UK 4271. See Oxamniquine.

UK 4271. See Oxamniquine.

CGP-4540. See Nitroscanate.

Preparation 5006. See Propoxur.

U. K. 5066. See U. K. 5066.

Bay b 5097. See Clotrimazole.

U. K. 5378. See U. K. 5378.

WR 5,473. See Cycloguanil.

U. K. 5574. See U. K. 5574.

U. K. 5704. See U. K. 5704.

5757. See Fenbautel.

Bay h 5757. See Febantel.

Bay Vh 5757. See Febantel.

U. K. 5876. See U. K. 5876.

U. K. 5925. See U. K. 5925.

WR 5,949. See Trimethoprim.

WR 5 990. See WR 5 990.

WR 6 007. See WR 6 007.

WR 6,020. See Isopentaquine.

WR 6,021. See Pentaquine.

WR 6 025. See WR 6 025.

WR 6026. See 8-(6-Diethylamino-hexylamino)-6-methoxy-4-methylquinoline.

NUMBER INDEX -- Continued.

WR 6 028. See WR 6 028.

WR 6,527. See Tetracycline.

WR-6,798. See Diformyl dapsone.

WR 7,557. See Sulfadiazine.

Bayer 7602 Ac. See C,C-Diallyl-bis-(4-amino-2-methyl-6-quinolyl) malonamide.

WR-7903. See WR-7903.

SN-8137. See Oxychloroquine.

R-8299. See Tetramisole.

Embay 8440. See Praziquantel.

Embay 8440-Bayer. See Praziquantel.

Shell SD-8447. See Tetrachlorvinphos.

Shell SD-8448. See 2-Chloro-1-(2,4,5-trichlorophenyl) vinyl diethylphosphate.

8,823 R.P. See Metronidazole.

RS-8858. See Oxfendazole.

Bay 9007. See Fenthion.

Bayer 9015. See Niclofolan.

Bayer 9037. See Quintiofos.

Bayer 9053. See Phoxim.

L 9053. See L 9053.

GO 9333. See Nitroscanate.

SN-9584. See SN-9584.

9.712RP. See Benzoyl metronidazole.

SN-10274. See SN-10274.

NUMBER INDEX -- Continued.

Merck SN 10,275. See (6,8-Dichloro-2-phenyl-4-quinolyl)- α -piperidyl carbinol hydrochloride.

RP 11074 [i.e. RP 11974]. See Phosalone.

12-278. See Dichlorquinazine.

SQ 12,525. See Lonomycin.

CP 12,574. See Tinidazole.

RD-12,869. See 6-Chloro-5- β -diethylaminoethyl-amino-8-methylquinoline.

Am 13,146. See Teclozan.

Win 13.146. See Teclozan.

SN-13,465. See 6-Bromo- α -diheptylaminoethyl-9-phenanthrenemethanol.

SD-14114. See Hexakis (β , β -dimethylphenethyl) distannoxane.

CP-14,445. See Oxantel.

14.539 RP. See Secnidazole.

SN-15,068. See 6,8-Dichloro-2-(3',4'-dichlorophenyl)- α -(di-n-butylaminomethyl)-4-quinoline methanol.

153C51. See 1,7-Bis(p-aminophenoxy)heptane.

Wellcome 153C51. See 1,7,Bis(p-aminophenoxy) heptane.

Bayer 15 922. See Trichlorfon.

A-16,612. See Teroxalene.

Compound 16,842. See Bitoscanate.

Hoechst 16.842. See Bitoscanate.

Ciba 17'581. See 1- β -Diethylamino-ethylamino-4,6,8-trimethyl-5-azathioxanthone.

R-17,635. See Mebendazole.

SQ 18,506. See trans-5-Amino-3-[2-(5-nitro-2-furyl)vinyl]- Δ^2 -1,2,4-oxadiazole.

SQ 18,506 analogues. See trans-5-Amino-3-[2-(5-nitro-2-furyl)vinyl]- Δ^2 -1,2,4-oxadiazole.

NUMBER INDEX -- Continued.

Sch 18545. See N,N'-Bis (3,4 ditrifluoromethyl-phenyl) methylmalonamide.

Cebegine RU 19110. See Halofuginone.

Sch 20350. See p-Chlorophenyl isothiocyanate.

R-20458. See (E)-6,7-Epoxy-1-(p-ethylphenoxy)-3,7-dimethyl-2-octene.

Sch 21480. See Tioxidazole.

SQ 21,704. See SQ 21,704.

Sch 23154. See 1-(2-Nitro-4-propyl-oxyphenyl)-3-carbomethoxy-S-methyl isothio-urea.

CGA-23'654. See Nitroscanate.

GS-23'654. See Nitroscanate.

WR 27,653. See 5-Bromo-4[bis-(2-diethylamino ethyl)-amino] veratrole.

SKF 29044. See Parbendazole.

Bay 29493. See Fenthion.

WR 29 597. See WR 29 597.

F30066. See Nitrofurylacrylamide.

WR-30,090. See 6,8-Dichloro-2-(3',4'-dichlorophenyl)- α -(di-n-butylaminomethyl)-4-quinoline methanol.

31,559 R.P. See Lonomycin.

WR-33,063. See 6-Bromo- α -diheptylaminoethyl-9-phenanthrenemethanol.

Hoechst 33258. See 2-[2-(4-Hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazyl) benzimidazole.

FMC 33297. See Permethrin.

Aldrich C 3360-3. See Aldrich C 3360-3.

349C59. See Moxipraquine.

MW 356.48. See Hycanthone.

Cyanamid-38023. See Famphur.

NUMBER INDEX -- Continued.

- WR-38,839. See 4,6-Diamino-1,2-dihydro-2,2-dimethyl-1-(3,4-dichlorobenzoyloxy)-1,3,5-triazine hydrochloride.
- Bayer 39007. See Propoxur.
- WR-40,070. See 2,4-Diamino-5-piperonyl-pyrimidine.
- FMC 41655. See Permethrin.
- Upjohn U-42,564. See N-[(2,4-Dimethylphenyl)imino]methyl-N-methyl-benzenesulfenamide.
- SD 43775. See Fenvalerate.
- CGA 50439. See Tifato1.
- CL-64,855. See 2-Amino-5-(1-methyl-5-nitro-2-imidazolyl)-1,3,4-thiadiazole.
- 67-255. See 3-Ethyl-8-methyl-1,3,8-triazabicyclo[4,4,0] decan-2-one.
- Ro 7-0207. See Ornidazole.
- Ro 7-1051. See Benznidazole.
- CL 71.366. See Piperamide.
- WR-74,106. See Terephthalic acid.
- WR-77,135. See WR-77,135.
- WR-81,844. See 1-(3,4-Dichlorophenyl)-3-[4-(N-ethyl-3-piperidyl-amino)-6-methyl-2-pyrimidinyl]guanidine.
- WR-91,808. See WR-91,808.
- WR 93,133. See 2-(4-Chlorophenyl)-2-(4-piperidyl)-tetrahydrofuran hydrochloride.
- WR-98,057. See WR-98,057.
- 98/202. See 6-Amidino-2-(4'-amidinophenyl)-thionaphthene dilactate.
- WR-99,210. See WR-99,210.
- WR 99,662. See 2-[3-(Dimethylamino) propyl-amino]-4-(trichloromethyl)-6-(α,α,α -trichlorom-tolyl)-2-triazine.
- WR-99,682. See WR-99,682.
- 102/198. See 4'6-Diamidino-2-phenylindole.

NUMBER INDEX -- Continued.

- Ro 11-3128. See (+)-5-(o-Chlorophenyl)-1,3-dihydro-3-methyl-7-nitro-2H-1,4-benzodiazepine-2-one.
- ICI 120645. See 9-(2,6-Dichlorobenzyl) adenine.
- ICI 120688. See 9-(2,6-Dichlorobenzyl) adenine-1-N-oxide.
- WR 122,455. See α -(2-Piperidyl)-3,6-bis (trifluoromethyl)-9-phenanthrene methanol.
- ICI 123487. See Arprinocid.
- ICI 125752. See 9-(2-Chloro-6-fluorobenzyl) adenine-1-N-oxide.
- WR 127 854. See WR 127 854.
- WR-131,834. See WR-131,834.
- WR-135,403. See WR-135,403.
- WR 136 479. See WR 136 479.
- WR 138 720. See Floxuridine.
- WR 139 007. See Imidazole-4-carboxamide, 5-(3,3-dimethyl-1-triazeno).
- WR 139 502. See Streptozotocin.
- WR 142,490. See Mefloquine.
- WR-143,803. See WR-143,803.
- WR-146,459. See WR-146,459.
- WR-148,763. See WR-148,763.
- WR-148,946. See WR-148,946.
- WR-149,809. See WR-149,809.
- WR-150,726. See WR-150,726.
- WR-151,312. See WR-151,312.
- WR 154,904. See α -(2-Piperidyl)-2,6-di-(p-trifluoromethylphenyl)-4-pyridinemethanol.
- WR-154,928. See 2,4-Diamino-6-(2-naphthyl)-thioquinazoline.
- WR-158,121. See WR-158,121.

NUMBER INDEX -- Continued.

- WR-158,122. See 2,4-Diamino-6-(2-naphthylsulfonyl)-quinazoline.
- WR 158 221. See Coralyne sulfoacetate.
- WR-159,412. See 2,4-Diamino-6-(5-trifluoromethylphenyl)-thioquinazoline.
- WR-162,878. See WR-162,878.
- WR-165,355. See WR-165,355.
- WR-165,533. See WR-165,533.
- WR-171,669. See 1-(1,3-Dichloro-6-trifluoromethyl-9-phenanthryl) 3-di-(n-butyl) aminopropanol.
- WR-172,435. See 3-(Dibutylamino)-1[2,6-bis(trifluoromethylphenyl)-4-pyridyl]propanol.
- WR-173,060. See WR-173,060.
- WR-175,039. See WR-175,039.
- WR-175,412. See WR-175,412.
- WR-177,504. See α -(Di-N-butylaminoethyl)-2,8-(bis-trifluoromethyl)-4-quinolinemethanol, hydrochloride.
- WR 177 529. See cis-Diamminedichloroplatinum.
- WR-177,602. See WR-177,602.
- WR-178,919. See WR-178,919.
- WR-178,979. See WR-178,979.
- WR-180,117. See WR-180,117.
- WR-180,153. See WR-180,153.
- WR-180,409. See α -(2-Piperidyl)-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridine-methanol.
- WR 180,409.H₃PO₄. See α -(2-Piperidyl)-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridine-methanol.
- WR-181,018. See WR-181,018.
- WR 181 023. See 4-Methylprimaquine.
- WR-182,231. See WR-182,231.

NUMBER INDEX -- Continued.

- WR 182 234. See 2-Methylprimaquine dihydrochloride monohydrate.
- WR-183,544. See WR-183,544.
- WR-183,545. See WR-183,545.
- WR-183,546. See WR-183,546.
- WR-183,606. See WR-183,606.
- WR-184,806. See α -(tert-Butylaminoethyl)-2,8-bis-(trifluoromethyl)-4-quinolinemethanol.
- WR 184,806.H₃PO₄. See DL-2,8-Bis(trifluoromethyl)-4-[1-hydroxy-3-(N-t-butylamino)propyl]-quinoline phosphate.
- WR-185,020. See WR-185,020.
- PM-185184. See Secnidazole.
- WR-190,420. See WR-190,420.
- WR 192 515. See WR 192 515.
- WR-194,965. See 2-(t-Butylaminomethyl)-4-t-butyl-6-(4-chlorophenyl)-phenol.
- WR-199,426. See WR-199,426.
- WR 201 678. See WR 201 678.
- WR-204,165. See 3,6-Bis-(t-butyl)-8-(4-chlorophenyl)-2H,4H-1,3-benzooxazine.
- WR 206 027. See WR 206 027.
- WR 208 442. See WR 208 442.
- WR 210 810. See WR 210 810.
- WR 211 077. See WR 211 077.
- WR 211 663. See WR 211 663.
- WR 211 666. See WR 211 666.
- WR 211 672. See WR 211 672.
- WR 211 784. See WR 211 784.
- WR 211 789. See 8-(6-Ethylaminoethylamino)-6-methoxy-4 methyl-quinoline.
- WR 211 814. See 3-Methylprimaquine diphosphate.

NUMBER INDEX -- Continued.

WR 211 819. See WR 211 819.
WR 212 293. See WR 212 293.
WR 213640. See WR 213640.
WR 214 198. See WR 214 198.
WR 215 300. See WR 215 300.
WR 215 732. See WR 215 732.
WR 216 837. See WR 216 837.
WR 217 271. See WR 217 271.
WR 218 574. See WR 218 574.
WR 218 669. See WR 218 669.
WR 218 805. See WR 218 805.
WR 218 806. See WR 218 806.
WR 220 078. See Inosine diglycolaldehyde.
WR 222 849. See WR 222 849.
WR 223 658. See WR 223 658.
WR 223 747. See WR 223 747.
WR 223 756. See WR 223 756.
WR 224 387. See WR 224 387.
WR 225 503. See WR 225 503.

NUMBER INDEX -- Continued.

WR 225 742. See WR 225 742.
WR-226,253. See α -(2-Piperidyl)-2-trifluoromethyl-6,8-dichloro-4-quinolinemethanol.
WR 226 257. See WR 226 257.
WR 226 292. See WR 226 292.
WR 226 619. See WR 226 619.
WR 227 175. See WR 227 175.
WR 227 495. See WR 227 495.
WR 228 327. See WR 228 327.
WR 230 220. See WR 230 220.
WR 231 163. See WR 231 163.
Win 24,933-2. See Hycanthone.
WR 295 595. See WR 295 595.
BPRL-5344-16. See BPRL-5344-16.
U. K. 5585-11. See U. K. 5585-11.
U. K. 5778/11. See U. K. 5778/11.
Dichloro analog L-628,914. See 6-Amino-9-(2,6-dichlorobenzyl) purine.
Ro 1-9334/20. See Dehydroemetine.
C9333-Go/CGP 4540. See Nitroscanate.

Aabomycin A

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Aabomycin S

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Abate. See Temephos.

Abi-Zetso. See Pyrimethamine or Sulfamethazine.

Absinthium -- Wormwood.

Wormwood, single styled, racemes and leaves
Slepnev, N. K., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 107-109
Ascaris suum, in vitro testing of anthelmintic activity of water extracts of some plants

Acaprin. See 1,3-Di-6-quinolyurea.

Acaricides

Sinclair, A. N., 1978, Vet. Rec., v. 103 (11), 247 [Letter]
sheep scab control with use of showers

Acaricides

Whitehead, G. B., 1973, South African Med. J., v. 47 (8), 342-344
ticks, development of resistance to acaricides, review: Eastern Cape Province

Acedist. See Bromophenophos.

Acemidophene

Vishniauskas, A.; and Rudaitis, A., 1978, Veterinariia, Moskva (4), 68-69
Fasciola hepatica, cattle (exper.), acemidophene, dertil and sulfene compared, acemidophene insufficiently effective

Acephate -- O,S-Dimethyl acetyl-phosphoramidothioate.

Acephate

Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661
flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico

2-Acetamido-5-nitrothiazole. See Aminitroazole.

Acetaminophen -- p-Acetylamino-phenol.

p-Acetylamino-phenol

Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective

Acetarsone

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Acetic acid -- Glacial acetic acid.

Glacial acetic acid

Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, Current Sc., Bangalore, v. 47 (8), 283-284 [Letter]
Trichodina [sp.] on exterior of *Cyprinus carpio*, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

Acetophenetidin -- p-Acetophenetidine; Phenacetine.

Phenacetine (p-Acetophenetidine)

Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective

p-Acetophenetidine. See Acetophenetidin.

2-Acetoxy-3-bromo-5-chloro-N-(4-bromophenyl)-thiobenzamide. See Brotianide.

Acetoxy-9-ellipticine

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

p-Acetylamino-phenol. See Acetaminophen.

Acetyl-p-aminophenolacetate

Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective

Achromycin. See Tetracycline.

Acranil

Haak, W.; et al., 1972, Ang. Parasitol., v. 13 (4), 200-207
cestodes of humans, recommended drug therapy

- Acridinium chloride**
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- 4'-(9-Acridinylamino) methanesulfonyl-m-anisidide
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Acriflavine** -- Acriflavine hydrochloride; Trypaflavine.
- Acriflavin hydrochloride**
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- Acriflavine**
Hajduk, S. L., 1979, *J. Cell Sc.*, v. 35, 185-202
Crithidia fasciculata, Trypanosoma equiperdum, observations on dyskinetoplasty, possible mechanisms of acriflavine action
- Acriflavine**
Nagington, J., 1975, *Tr. Ophth. Soc. United Kingdom*, v. 95 (2), 207-209
Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity
- Trypaflavine**
Nikol'skii, S. N.; Nikiforenko, V. I.; and Pozov, S. A., 1977, *Veterinariia, Moskva* (4), 71-75
Piroplasma jakimovi, cattle, morphological and biological comparison with P. bigeminum, epizootiology (Ixodes ricinus as main vector; frequent association with leptospiriosis), treatment: Siberia
- Acriflavine**
Ono, T.; and Inoki, S., 1976, *Kiseichugaku Zasshi (Japan. J. Parasitol.)*, v. 25 (2), 47-58
Trypanosoma gambiense-infected mice, interaction between trypanosomes and peritoneal cells after treatment with human serum, anti-trypanosome mouse-serum, and acriflavine, electron microscopy
- Acriflavin**
Pandey, N. N.; and Mishra, S. S., 1978, *Indian Vet. J.*, v. 55 (2), 144-148
Babesia bigemina, indigenous cow calves, berenil and acriflavin effective, sulfadimethoxine ineffective: India
- Acriflavin**
Razzaque, A.; and Mishra, S. S., 1977, *Bull. Animal Health and Prod. Africa*, v. 25 (4), 409-414
Trypanosoma evansi, buffalo calves (exper.), berenil, acriflavin, and antrycide prosalt, berenil most effective
- Acriflavine** -Continued.
- Acriflavine**
Shirshov, A. T.; et al., 1975, *Izvest. Akad. Nauk SSSR, s. Biol.* (5), 759-765
Crithidia oncopelti, acriflavine, effect on structure of kinetoplast, kinetoplast DNA, protein synthesis in kinetoplast and cytoplasmic ribosomes; suggests that information required for synthesis of kinetoplast ribosomes is contained in kinetoplasts
- Trypaflavin**
Vulchovski, Ia., 1977, *Vet. Med. Nauki*, v. 14 (3), 79-87
Babesia, cattle, efficacy of trypanblau, trypaflavin, acaprin, berenil, and pyrodia (berenil and pyrodia superior to other drugs)
- Acriflavine**
Warton, A.; and Modlinska, M., 1975, *Acta Parasitol. Polon.*, v. 23 (1-11), 127-133
Trypanosoma spp., mice, rats, effect of acriflavine on dynamics of trypanosome population size and formation of dyskinetoplastic forms in host blood
- Acriflavine**
Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
-
- Acriflavine hydrochloride.** See **Acriflavine.**
- 4'-(3-Acrylyl)benzo-15-crown-5
Brown, G. R.; and Foubister, A. J., 1979, *J. Med. Chem.*, v. 22 (8), 997-999
benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against Eimeria tenella
- Actamer.** See **Bithionol.**
- ACTH** -- Adrenocorticotropin.
- Adrenocorticotropin**
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
-
- Actidione.** See **Cycloheximide.**
- Actinomycin D.** See **Dactinomycin.**

Actinomycin D. See Dactinomycin.

Adenine arabinoside. See Vidarabine.

Adenosine

Dewey, V. C.; Kidder, G. W.; and Nolan, L. L., 1978, *Biochem. Pharmacol.*, v. 27 (10), 1479-1485

Crithidia fasciculata, mechanism of inhibition of growth by adenosine and adenosine analogs

Adenosine N-oxide

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Adrenocorticotropin. See ACTH.

Aerol-2 aerosol

Kan, P. T.; and Kholodov, I. Ia., 1978, *Veterinariia, Moskva* (6), 61-63

H[yalomma] anaticum, aerol-2 aerosol for control in animal quarters, various dosages, dispersals and exposure times tested

Agaric acid -- Agaricin.

Agaricin

Nathan, H. C.; et al., 1979, *J. Protozool.*, v. 26 (4), 657-660

Trypanosoma brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Agaricin. See Agaric acid.

Aklomide -- Aklomix; 2-Chloro-4-nitrobenzamide; Novostat-W (with Sulfantran).

Aklomide

Karlsson, T.; and Reid, W. M., 1978, *Avian Dis.*, v. 22 (3), 487-495

Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Aklomide (Aklomix)

McQuiston, T. E.; and McDougald, L. R., 1979, *Ztschr. Parasitenk.*, v. 59 (2), 107-113

Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Aklomide + Sulfantran (=Novostat-W)

Manuel, M. F.; and Neri, R. A., 1975, *Philippine J. Vet. Med.*, v. 14 (1), 106-116

Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Aklomide -- Continued.

Aklomide

Winchester, R. V., 1978, *N. Zealand J. Sc.*, Wellington, v. 21 (4), 553-555
coccidiostats aklomide and zoalene, residues in poultry meat, determination method

Aklomix. See Aklomide.

Albendazole -- Albendazole carbamate; Albendazole sulphone; Albendazole sulphoxide; Methyl [5-(propylthio)-1H-benzimidazol-2-yl] carbamate; Valbazen.

Albendazole

Benz, G. W.; and Ernst, J. V., 1978, *Am. J. Vet. Research*, v. 39 (7), 1107-1108

Dictyocaulus viviparus, calves (exper.), albendazole administered orally, significant reduction of adults, elimination of first-stage larvae passed in feces

Albendazole (Valbazen)

Borgsteede, F. H. M., 1979, *Vet. Quart.*, v. 1 (4), 181-188

gastrointestinal nematodes, calves, albendazole administered at end of grazing season: Netherlands

Albendazole

Campbell, N. J.; and Hall, C. A., 1979, *Research Vet. Sc.*, v. 26 (1), 90-93

Fasciola hepatica, benzimidazole-resistant strains of *Haemonchus contortus* and *Trichostrongylus colubriformis*, sheep (exper.), efficacy of albendazole

Albendazole

Giordia, H.; McCampbell, H. C.; and Stuedemann, J. A., 1978, *Am. J. Vet. Research*, v. 39 (3), 517-518

Moniezia benedeni, *M. expansa*, calves, albendazole, anthelmintic efficacy at 4 dose levels, no signs of toxicosis

Albendazole

Coles, G. C.; and Briscoe, M. G., 1978, *Vet. Rec.*, v. 103 (16), 360-361 [Letter]

Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for *Haemonchus contortus* eggs

Albendazole carbamate

Craig, T. M.; and Ronald, N. C., 1978, *Southwest Vet.*, v. 31 (2), 121-124

Taenia saginata, calves (exper.), albendazole in single oral dose had apparent deleterious action against cysticerci, preliminary study

Albendazole

Cummins, L. J.; and Callinan, A. P. L., 1979, *Austral. Vet. J.*, v. 55 (7), 348-349

[Letter]
nematodes, cattle, albendazole and fenbendazole, effect of oesophageal groove reflex on anthelmintic efficiency

Albendazole -- Continued.

Albendazole

Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (11), 675-679
Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Albendazole sulphone

Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (11), 675-679
Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Albendazole sulphoxide

Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (11), 675-679
Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Albendazole (Valbazen)

Downey, N. E., 1978, *Vet. Rec.*, v. 103 (19), 427-428
 gastrointestinal nematodes, calves, efficacy of albendazole, compared with levamisole: Ireland

Albendazole

Dubey, J. P.; et al., 1978, *Am. J. Vet. Research*, v. 39 (6), 1027-1031
Paragonimus kellicotti, specific-pathogen-free cats, albendazole, excellent results (reduced ova production, changed worm morphology, reduced pulmonic lesions)

Albendazole

Foreyt, W. J.; and Drawe, D. L., 1978, *Am. J. Vet. Research*, v. 39 (12), 1901-1903
 gastrointestinal nematodes and *Fascioloides magna*, white-tailed deer, albendazole highly effective against nematodes, 38% effective against *F. magna*: Victoria, Texas

Albendazole

Georgi, J. R.; et al., 1979, *Am. J. Vet. Research*, v. 40 (6), 829-831
Filaroides hirthei, beagle dogs, coprophagia is principal mechanism of transmission from dams to pups and among pups, control by albendazole treatment of brood bitches coupled with isolation of weanlings and older pups from sources of contaminated feces

Albendazole

Georgi, J. R.; Slauson, D. O.; and Theodorides, V. J., 1978, *Am. J. Vet. Research*, v. 39 (5), 803-806
Filaroides hirthei, dogs (exper.), albendazole, killed all but small proportion of worms and sterilized the ones that survived

Albendazole (Valbazen)

Gunawan, M.; et al., 1979, *Research Vet. Sc.*, v. 27 (1), 111-115
Haemonchus contortus, *Trichostrongylus colubriformis*, efficacies of fenbendazole and albendazole against developing and adult stages of benzimidazole-resistant strains, sheep (exper.)

Albendazole -- Continued.

Albendazole

Hall, C. A.; et al., 1978, *Research Vet. Sc.*, v. 25 (3), 364-367
Haemonchus contortus, *Trichostrongylus colubriformis*, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Albendazole

Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, *Research Vet. Sc.*, v. 25 (3), 360-363
Haemonchus contortus, *Trichostrongylus colubriformis*, levels of benzimidazole resistance recorded from an egg hatch test procedure

Albendazole

Ireland, C. M.; et al., 1979, *Biochem. Pharmacol.*, v. 28 (17), 2680-2682
 relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiroides dubius* in mice

Albendazole (Valbazen)

Le Jambre, L. F., 1979, *Austral. Vet. J.*, v. 55 (2), 65-67
Ostertagia circumcincta, *O. trifurcata*, sheep (exper.), effectiveness of levamisole, thiabendazole, albendazole, and oxfendazole against levamisole-resistant strains

Albendazole

Lloyd, S.; Soulsby, E. J. L.; and Theodorides, V. J., 1978, *Experientia*, v. 34 (6), 723-724
Taenia saginata, calves (exper.), albendazole effective against metacestodes

Albendazole

McCracken, R. O., 1978, *J. Parasitol.*, v. 64 (2), 214-219
Trichinella spiralis, mice, mebendazole and albendazole more effective against immature than mature worms, despite this decline in drug sensitivity it is postulated that these benzimidazoles have potential therapeutic value

Albendazole (Valbazen)

Marriner, S.; and Bogan, J. A., 1979, *Vet. Rec.*, v. 105 (11), 261
 benzimidazole anthelmintics, sheep, oral vs. intraruminal vs. intra-abomasal administration

Albendazole

Rendano, V. T.; et al., 1979, *Equine Vet. J.*, v. 11 (4), 223-231
Strongylus vulgaris, foals (exper.), albendazole, sequential arteriography for evaluation of larvicidal activity

Albendazole

Ronald, N. C.; Craig, T. M.; and Bell, R. R., 1979, *Am. J. Vet. Research*, v. 40 (9), 1299-1300
Fasciola hepatica, *Fascioloides magna*, cattle, albendazole, controlled evaluation: Texas

Albendazole -- Continued.

Albendazole

Ross, D. B.; Eichler, D. A.; and Cameron, D., 1978, Vet. Rec., v. 102 (25), 556-557
Dictyocaulus filaria, *Ostertagia circumcincta*, *Trichostrongylus colubriformis*, lambs (exper.), albendazole, very effective at a relatively low dose level

Albendazole

van Schaalkwyk, P. C.; et al., 1979, J. South African Vet. Ass., v. 50 (1), 31-35
 helminths, sheep (nat. and exper.), albendazole

Albendazole (Valbazen)

Todd, K. S., jr., 1978, Vet. Med. and Small Animal Clin., v. 73 (4), 453-454
Mesocostoides corti, dogs (exper.), albendazole completely effective in removing adults

Albendazole (Valbazen)

Todd, K. S., jr.; Howland, T. P.; and Macy, D. W., 1978, Canine Pract., Santa Barbara, v. 5 (6), 11-12, 14
Paragonimus kellicotti, dog, albendazole may be effective, case report

Albendazole (Valbazen)

Todd, K. S., jr.; and Mansfield, M. E., 1978, Mod. Vet. Pract., v. 59 (5), 377
 gastrointestinal nematodes, cattle, albendazole, highly effective, no adverse reactions: Illinois

Albendazole

Wescott, R. B.; et al., 1979, Am. J. Vet. Research, v. 40 (3), 369-371
 cattle parasites, efficacy of albendazole in controlled and field trials: Washington

Albendazole (Valbazen)

Williams, J. C.; et al., 1979, Vet. Rec., v. 105 (5), 98-100
Ostertagia ostertagi, fourth stage larvae, cattle, albendazole

Albendazole carbamate. See Albendazole.

Albendazole sulphone. See Albendazole.

Albendazole sulphoxide. See Albendazole.

Alcohol, Anhydrous -- Ethanol; Ethyl alcohol.

Ethanol

Campbell, W. C., 1977, Proc. Helminth. Soc. Washington, v. 44 (2), 120-125
Trichinella spiralis, susceptibility to deleterious effect of ethanol in vitro and in infected rats and pigs

Ethanol

Kunstyr, I.; and Ammerpohl, E., 1978, Lab. Animals, v. 12 (2), 95-97
Spironucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Alcohol, Anhydrous -- Continued.

Ethyl alcohol

Slepnev, N. K.; and Zen'kov, A. V., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 135-137
E[chinococcus] granulosus, *protoscolices*, destructive action of high and low temperatures; lysol and creolin most destructive of chemicals tested

Ethanol

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Alcopar. See Bephenium.

Aldrich C 3360-3

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Alginic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Alkyl-(5-acyl-1-H-benzimidazol-2-yl) carbamates
 Raeymaekers, A. H. M.; et al., 1978, Arzneimittel-Forsch., v. 28 (4), 586-594
Syphacia muris, *Strongyloides ratti*, synthesis and anthelmintic activity of mebendazole, flubendazole and other alkyl-(5-acyl-1-H-benzimidazol-2-yl) carbamates in rats

8- ω -Alkylamino-6-methoxy quinolines, basically-substituted

Kinnamon, K. E.; et al., 1979, Mil. Med., v. 144 (10), 660-664
 leishmaniasis, military significance, laboratory trials in *Mesocricetus auratus* using aminoquinolines significantly more active than currently used antileishmanial agents

8- ω -Alkylamino-6-methoxy quinolines, heterocyclic substituted

Kinnamon, K. E.; et al., 1979, Mil. Med., v. 144 (10), 660-664
 leishmaniasis, military significance, laboratory trials in *Mesocricetus auratus* using aminoquinolines significantly more active than currently used antileishmanial agents

S-(Alkyl-1)-isothiuronium hydrohalides

Payares, G.; and Ercoli, N., 1978, *Exper. Parasitol.*, v. 45 (1), 1-7

Schistosoma mansoni, drug-immobilized cercariae have reduced virulence but are not dead, cercariae become avirulent only when flame cell is affected, no protection against reinfection in mice injected with immobilized cercariae of reduced virulence

Allethrin

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Allopurinol -- Allopurinol ribonucleoside; 4-Hydroxypyrazolo(3,4-d)pyrimidine; 4-Hydroxy-1-β-D-ribofuranosylpyrazolo(3,4-d)pyrimidine.

Allopurinol

Irvin, A. D.; and Young, E. R., 1978, *Research Vet. Sc.*, v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Allopurinol

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Allopurinol (4-Hydroxypyrazolo(3,4-d)pyrimidine)

Marr, J. J.; Berens, R. L.; and Nelson, D. J., 1978, *Science* (4360), v. 201, 1018-1020

Trypanosoma cruzi, antiprotozoal effect of allopurinol can be accounted for by its in vivo transformation into a toxic adenine analog by the parasite

Allopurinol

Nelson, D. J.; et al., 1979, *J. Biol. Chem.*, v. 254 (10), 3959-3964

Leishmania braziliensis, L. donovani, pyrazolo(3,4-d)pyrimidines, metabolism, possible explanation for antileishmanial activity

Allopurinol ribonucleoside

Nelson, D. J.; et al., 1979, *J. Biol. Chem.*, v. 254 (22), 11544-11549

Leishmania spp., allopurinol ribonucleoside as an antileishmanial agent: biological effects, metabolism, and enzymatic phosphorylation

Allopurinol

Pfaller, M. A.; and Marr, J. J., 1974, *Antimicrob. Agents and Chemotherapy*, v. 5 (5), 469-472

Leishmania braziliensis, allopurinol inhibits growth in vitro at concentrations which are attainable in human tissues and body fluids

Allopurinol -- Continued.

Allopurinol

Spector, T.; Jones, T. E.; and Elion, G. B., 1979, *J. Biol. Chem.*, v. 254 (17), 8422-8426
Leishmania donovani promastigotes, adenylosuccinate synthetase and adenylosuccinate lyase, purification, properties, substrate and inhibitor specificities, selective amination of allopurinol ribonucleotide may be related to its antileishmanial activity

Allopurinol

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Allopurinol ribonucleoside. See Allopurinol.3-Allyl-3,4-dihydro-2,2-dimethyl-2H-naphtho [1,2-b] pyran-5,6-dione. See Allyl-β-lapachone.

D-C-Allyl-glycine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Allyl-β-lapachone -- 3-Allyl-3,4-dihydro-2,2-dimethyl-2H-naphtho [1,2-b] pyran-5,6-dione; 3-Allyl-β-lapachone.

3-Allyl-β-lapachone

Docampo, R.; et al., 1978, *Ztschr. Parasitenk.*, v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

Allyl-β-lapachone

Lopes, J. N.; et al., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

3-Allyl-β-lapachone. See Allyl-β-lapachone.Allyl thiourea. See Thiosinamine.

- Allylcarb
Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides
- Allylcarb
Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies
- Alon. See 3-4-Isopropylphenyl-1,1-dimethylurea.
- Altik. See Dioxathion or Toxaphene.
- Alugan -- 5-Bromomethyl 1-1,2,3,4,4-7,7-hexachlorobicyclo-(2,2,1)hepten-(2); Bromocyclen.
- Alugan
Bajric, A.; Levi, I.; and Hlubna, D., 1978, Vet. Glasnik, v. 32 (10), 843-846
Cnemidocoptes pilae in Melopsittacus undulatus, clinical picture, alugan: Buiatric Clinic, Sarajevo
- Bromocyclen
Bonin, W., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (2), 34-37
Amblyomma hebraeum, Psoroptes cuniculi, Melophagus ovinus, Dermanyssus gallinae, heptenophos, rapid mode of action, broad range of efficacy, short residual effect and effective as a vapour poison, compared with other standard drug preparations
- Alugan
Kamyszek, F., 1977, Med. Wet., v. 33 (6), 354-356
Sarcoptes scabiei, pigs, alugan, neguvon
- Alugan
Pfeiffer, H., 1979, Ztschr. Parasitenk., v. 59 (1), 95-106
Cheyletiella blakei, cats, clinical symptoms also appeared in owner, alugan successfully removed mites from cats and owner's skin affection then disappeared; morphology, differential diagnosis from C. parasitivorax: Osterreich
- Alugan
Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran
- Alugan (Bromocyclen)
Ribbeck, R.; Schroeder, E.; and Schumann, H., 1979, Monatsh. Vet.-Med., v. 34 (10), 383-384
Lucilia sericata, dogs, cats, myiasis, surgical removal and drug treatment
- Amantadine -- 1-Adamantamine.
- Amantadine
Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances
- Amantadine
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity
- Amantidine
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Ambilhar. See Niridazole.
- Amicarbalide -- 3:3'-Diamidinocarbanilide; 3,3'-Diaminocarbanilide diisethionate; Diapron; Pirodia; Pyrodia.
- Amicarbalide
Adam, K. M. G.; et al., 1978, Brit. Vet. J., v. 134 (5), 428-433
babesiasis, cattle, 2 outbreaks compared (clean cattle introduced to tick-infested endemic area; tick-infested cattle introduced to clean area), percentage seropositive for babesial antibody, degree of manifest disease, treatment with amicarbalide, results indicate that vaccination against Babesia divergens is feasible and desirable: Scotland
- Amicarbalide
Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo
- Amicarbalide
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Amicarbalide -- Continued.

Amicarbalide (Diapron)
Nathan, H. C.; et al., 1979, J. Protozool.,
v. 26 (4), 657-660

Trypanosoma brucei brucei, mice, effect of
amicarbalide, imidocarb, and several other
agents

Pyrodisia

Vulchovski, Ia., 1977, Vet. Med. Nauki, v. 14
(3), 79-87

Babesia, cattle, efficacy of trypan blue,
trypaflavin, acaprin, berenil, and pyrodisia
(berenil and pyrodisia superior to other drugs)

Amides

Botero R., D., 1978, Ann. Rev. Pharmacol. and
Toxicol., v. 18, 1-15

antiparasitic drugs in current use for human
intestinal protozoa and helminths, brief re-
view of pharmacology, secondary effects, tox-
icity and contraindications

6-Amidino-2-(4'-amidinophenyl)-thionaphthene di-
lactate -- 98/202; 2-Phenyl-thionaphthene diamid-
dine.

2-Phenyl-thionaphthene diamidine (98/202)

Ercoli, N., 1978, Proc. Soc. Exper. Biol. and
Med., v. 157 (3), 397-401

Trypanosoma venezuelense, rats, comparison of
"true" and "false" prophylaxis using pentami-
dine, suramin and 98/202

p-(4-Amidino-phenoxy)-benzaldehyde-p-amidino-
phenylhydrazine dihydrochloride (HOE 668)

Raether, W.; Seidenath, H.; and Loewe, H., 1978,
Ann. Trop. Med. and Parasitol., v. 72 (6), 543-
547

Leishmania donovani, golden hamsters, HOE 668
compared with known antileishmanial drugs,
toxicity precludes further development but
very good anti-leishmanial action qualifies
it as standard compound in screening tests

Amidophos

Rastegaev, Iu. M., 1978, Vestnik Sel'skokhoz.
Nauki Kazakhstana (3), 73-76

myiasis, helminths, horses, chlorophos,
amidophos, economic effectiveness of treat-
ment: northern Kazakhstan

Amidophos

Talanov, G. A.; and Nikolaev, P. I., 1978,
Veterinariia, Moskva (9), 59-60

Parascaris, [Strongylata], Gasterophilus,
horses, use of chlorophos and amidophos:
Iakutsk ASSR

Aminitrozole -- 2-Acetamido-5-nitrothiazole;

Nitazol.

Aminitrozole

Benazet, F.; et al., 1970, Scand. J. Infect.
Dis., v. 2 (2), 139-143

intestinal and hepatic parasites, nitrohetero-
cyclic antiparasitics, laboratory studies
of chemotherapeutic activity and toxicity in
exper. animals

Aminitrozole -- Continued.

Aminitrozole

Campbell, W. C.; Bartels, E.; and Cuckler,
A. C., 1978, J. Parasitol., v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid
assay suitable for routine screening of com-
pounds for antischistosome activity, reduc-
tion in severity of hepatic lesions used as
chief criterion of efficacy

Nitazol

Parre, J.; and Simovart, H. E., 1977, Eesti
Pollumaj. Akad. Teadusl. Toode Kogum. (104),
114-118

histomoniasis, turkeys, age and seasonal
dynamics in relation to epizootiology; dis-
ease outbreaks in young birds under stress
conditions; nitazol satisfactory, trichopol
good prophylactic and therapeutic effect

Aminoacrichine

Rusak, L. V.; and Kovchur, V. N., 1972, Para-
zitologiya, Leningrad, v. 6 (1), 85-87

Hymenolepis nana, glycogen content in para-
site tissues decreased after treatment of
infected mice with aminoacrichine, phenasal,
or trichlorophen, implications for mechanism
of drug action

D- α -Aminobutyric acid

Williamson, J.; and Scott-Finnigan, T. J.,
1978, Antimicrob. Agents and Chemotherapy,
v. 13 (5), 735-744

trypanocidal activity of antitumor anti-
biotics and other metabolic inhibitors,
microtest for rapid preliminary assay in
vitro, parasite motility and infectivity
for mice are indexes respectively of respi-
ration and glycolysis and of cell division,
implications of results for combination
chemotherapy and deposit prophylaxis (with
polyanions)

γ -Aminobutyric acid. See GABA.

3-Amino-9-p-carbethoxyaminophenyl-10-methyl-
phenanthridinium ethano-sulphonate. See Car-
bidium sulfate.

6-Amino-9-(2-chloro-6-fluorobenzyl) purine. See
Arprinocid.

1-Aminocyclopentane carboxylic acid -- Cyclo-
leucine.

Cycloleucine

Williamson, J.; and Scott-Finnigan, T. J.,
1978, Antimicrob. Agents and Chemotherapy,
v. 13 (5), 735-744

trypanocidal activity of antitumor anti-
biotics and other metabolic inhibitors,
microtest for rapid preliminary assay in
vitro, parasite motility and infectivity
for mice are indexes respectively of respi-
ration and glycolysis and of cell division,
implications of results for combination
chemotherapy and deposit prophylaxis (with
polyanions)

6-Amino-9-(2,6-dichlorobenzyl) purine -- Dichloro analog L-628,914.

Dichloro analog L-628,914

Tamas, T.; et al., 1978, Poultry Science, v. 57 (2), 381-385

Eimeria acervulina, *E. maxima*, *E. necatrix*, *E. tenella*, arprinocid and dichloro analog L-628,914 in feed, decrease in oocysts' capacity to sporulate

Aminoethanephosphorothioates, Disulfide, Isothiourea and other thiol derivatives

Nelson, D. L.; and Pellegrino, J., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (6), 443-449

Schistosoma mansoni, mice, chemical structure-activity relationship study of 85 active derivatives of aminoethanephosphorothioate, disulfide and isothioureia

Aminoethanethiols, Mono-, di-, and tri-substituted

Nelson, D. L.; and Pellegrino, J., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (4), 264-267

Schistosoma mansoni, mice, experimental chemotherapy with substituted aminoethanethiols

Aminoethanethiosulfuric acids, Mono-, di- and trisubstituted

Nelson, D. L.; and Pellegrino, J., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (5), 365-370

Schistosoma mansoni, mice, aminoethanethiosulfuric acids derivatives, chemotherapeutic trials

S-2-Aminoethyl-L-cysteine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

8-Amino-6-methoxyquinolines, 4-Substituted

Carroll, F. I.; et al., 1979, J. Med. Chem., v. 22 (6), 694-699

Plasmodium spp. in mice and rhesus monkeys, synthesis and antimalarial activity of some 4-substituted 8-amino-6-methoxyquinolines, 4-ethylprimaquine was approximately as active and was less toxic than primaquine

8-[(4'-Amino-1'-methylbutyl)amino]-4-ethyl-6-methoxyquinoline. See 4-Ethylprimaquine.

4-[(4-Amino-1-methylbutyl)amino]-2-methoxyacridine

Scovill, J. P.; et al., 1979, J. Med. Chem., v. 22 (10), 1164-1167

Plasmodium cynomolgi in rhesus monkeys, derivatives of 4-amino-2-methoxyacridine

4-[(4-Amino-1-methylbutyl)amino]-2-methoxy-9-methylacridine

Scovill, J. P.; et al., 1979, J. Med. Chem., v. 22 (10), 1164-1167

Plasmodium cynomolgi in rhesus monkeys, derivatives of 4-amino-2-methoxyacridine

8-(4-Amino-1-methylbutylamino)-6-methoxyquinoline. See Primaquine.

2-Amino-5-(1-methyl-5-nitro-2-imidazolyl)-1,3,4-thiadiazole -- CL-64,855.

CL-64,855

Galhotra, A. P.; et al., 1979, Haryana Agric. Univ. J. Research, v. 9 (1), 69-74

Trypanosoma evansi, dogs (exper.), clinico-pathological changes, therapeutic trials with berenil and CL-64,855

6-Aminonicotinamide

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

trans-5-Amino-3-[2-(5-nitro-2-furyl)vinyl]- Δ^2 -1,2,4-oxadiazole -- SQ 18,506; SQ 18,506 analogues.

SQ 18506

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

SQ 18,506

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 339-347

Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

SQ 18,506 analogues

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 339-347

Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

trans-5-Amino-3-[2-(5-nitro-2-furyl)vinyl]- Δ^2 -1,2,4-oxadiazole (SQ18506)

Sims, P.; and Gutteridge, W. E., 1978, Biochem. Pharmacol., v. 27 (24), 2815-2820

Trypanosoma cruzi, inhibitory action of SQ18506 against nucleic acid synthesis

trans-5-Amino-3-[2-(5-nitro-2-furyl)vinyl]- Δ^2 -1,2,4-oxadiazole (SQ18506)

Sims, P.; and Gutteridge, W. E., 1979, Internat. J. Parasitol., v. 9 (1), 61-67

Trypanosoma cruzi, mode of action of SQ18506 considered to be inhibition of nucleic acid synthesis

2-Amino-5-nitropyridine

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonocides using Coulter Counter

- 2-Amino-5-nitropyrimidine
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- Aminonitrothiazole
Antunes, L. J.; et al., 1974, *Rev. Soc. Brasil.
Med. Trop.*, v. 8 (1), 9-13
human intestinal schistosomiasis mansoni be-
fore and after treatment with aminonitro-
thiazole, immunoglobulin levels, immediate
and delayed cutaneous hypersensitivity
- 2-Amino-5-nitrothiazole
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- 1-(4-Amino-2-n-propyl-5-pyrimidinyl-methyl)-2-
picolinium chloride hydrochloride. See
Amprolium.
- Aminopterin
Gutteridge, W. E.; Cover, B.; and Gaborak, M.,
1978, *Ann. Trop. Med. and Parasitol.*, v. 72
(4), 329-338
Trypanosoma cruzi, rapid, simple primary
screen to test compounds for activity as
potential trypanocides using infected A/JAX
inbred mice
- Aminopterin
Irvin, A. D.; and Young, E. R., 1978, *Research
Vet. Sc.*, v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxan-
thine uptake in vitro could be used as
primary screen for babesicidal drugs but
drugs showing in vitro activity are not
necessarily active in vivo
- 4-Aminopyrazolo(3,4-d)pyrimidine
Nelson, D. J.; et al., 1979, *J. Biol. Chem.*, v.
254 (10), 3959-3964
Leishmania braziliensis, L. donovani, pyrazolo
(3,4-d)pyrimidines, metabolism, possible ex-
planation for antileishmanial activity
- 4-Aminopyrazolo-(2,3-d)-pyrimidine beta-D-ribo-
furanoside. See Tubercidin.
- 3- and 5-Aminoquinolines
Khan, M. Sami; and LaMontagne, M. P., 1979, *J.
Med. Chem.*, v. 22 (8), 1005-1008
3- and 5-aminoquinolines, potential antimalar-
ials, synthesis, testing of some against
Plasmodium berghei in mice, Leishmania dono-
vani in hamsters, or P. cynomolgi in rhesus
monkeys
- 4-Aminoquinolines
Schmidt, L. H.; et al., 1977, *Antimicrob.
Agents and Chemotherapy*, v. 11 (5), 826-843
Plasmodium falciparum in Aotus trivirgatus,
activities of various 4-aminoquinolines
against chloroquine-resistant and -suscepti-
ble strains, observations confirm cross-
resistance among 4-aminoquinolines but in-
dicate that some derivatives may be thera-
peutically effective against infections
refractory to maximally tolerated doses of
chloroquine
- 7-Amino-3(β -D-ribofuranosyl)pyrazolo-[4,3-d]
pyrimidine. See Formycin.
- Aminosidine. See Paromomycin.
- Aminosidine sulfate. See Paromomycin.
- 4-Amino-6-trichloroethenyl-1,3-benzenedisulfon-
amide
Schulman, M. D.; et al., 1979, *J. Parasitol.*,
v. 65 (4), 555-561
Fasciola hepatica, rats, 4-amino-6-trichloro-
ethenyl-1,3-benzenedisulfonamide, pharmaco-
kinetic basis for efficacy
- Amitraz -- 1,5-Di-(2,4-dimethylphenyl)-3-methyl-
1,3,5-triazapenta-1,4-diene; N'-(2,4-Dimethyl-
phenyl)-N-[[2,4-dimethylphenyl]imino]-
methyl]-N-methylmethanimidamide; Taktic.
- Amitraz
Dawkins, C. C.; and Gladney, W. J., 1978, *J.
Econ. Entom.*, v. 71 (4), 657-660
3-host ticks, guinea pigs, amitraz, Upjohn
U-42,564, chlordimeform, detachment response
and mortality
- Amitraz (Taktic)
Gothe, R.; and Guizar, R. C., 1979, *Zentralbl.
Vet.-Med., Reihe B*, v. 26 (4), 290-303
Argas walkerae, chickens (exper.), in vivo
model for evaluating detaching potential of
various acaricides
- Amitraz (Taktic)
Roberts, M. C.; and Seawright, A. A., 1979,
Austral. Vet. J., v. 55 (11), 553-554
amitraz, horses, toxicity experiments, large
intestinal impaction: Australia
- Amitraz
Schuntner, C. A.; and Thompson, P. G., 1978,
Austral. J. Biol. Sc., v. 31 (2), 141-148
Boophilus microplus, larvae, ¹⁴C labelled
amitraz, metabolism, only amitraz and N-2,4-
dimethylphenyl-N'-methylformamide (metabo-
lite) toxic to larvae, piperonyl butoxide
applied simultaneously with amitraz had
slight effect on metabolism, three-fold syn-
ergistic effect; SKF 525-A similarly applied
had negligible effect on both metabolism and
toxicity to ticks
- Amni majus (water decoction or petroleum ether
extract)
Abdulla, W. A.; et al., 1977, *Egypt. J. Bil-
harz.*, v. 4 (1), 19-26
S[chistosoma] mansoni, anti-schistosomal
activity of the plant Amni majus and bergapten
(furocoumarin present in A. majus), compared
with tarter emetic, screening in mice
- Ammonia
Chevranova, Iu. A.; Bukhtoiarov, A. I.; and
Iastrebov, E. E., 1978, *Gig. i Sanitariia*
(4), 100-102
Ascaris suum, A. lumbricoides, use of ammo-
nia to destroy ascarid eggs in sewage sludge

Ammonium nitrate

Antonenko, A. E., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 121-126
nematode eggs, coccidian oocysts, non-litter pig manure, substances tested as disinfectants

Ammonium sulfate

Antonenko, A. E., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 121-126
nematode eggs, coccidian oocysts, non-litter pig manure, substances tested as disinfectants

Ammonium sulphate

Venkateswara Rao, P.; et al., 1977, Riv. Parassitol., Roma, v. 38 (1), 13-21
Prosthogonimus sp. cercariae, cercaricidal effect of certain common fertilizers, ammonium sulphate may be cercaricide of choice

Amodiaquine -- Amodiaquine hydrochloride; Basoquine; Camoprimum (with Primaquine); Camoquine; 4-[(7-Chloro-4-quinolyl)amino]- α -(diethylamino)-o-cresol.

Camoprimum + Maloprimum

Botelho, A., 1974, South African Med. J., v. 48 (31), 1351-1352
malariae, humans, clinical trials using sulphones and sulphonamides with a pyrimidine derivative

Amodiaquine (Basoquine)

Cedillos, R. A.; Warren, M.; and Jeffery, G. M., 1978, Am. J. Trop. Med. and Hyg., v. 27 (3), 466-472
Plasmodium vivax, humans, 2 regimens of primaquine (5-day; single dose) in combination with amodiaquine compared with amodiaquine alone, field evaluation: Zapotitan Valley, El Salvador

Camoquine

Dutta, G. P.; and Narain, L., 1978, Indian J. Exper. Biol., v. 16 (7), 838-840
Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Amodiaquin

Fitch, C. D.; Chevli, R.; and Gonzalez, Y., 1974, Antimicrob. Agents and Chemotherapy, v. 6 (6), 757-762
Plasmodium falciparum, Aotus trivirgatus erythrocytes infected with chloroquine-susceptible vs. chloroquine-resistant strain, effect of substrate (glucose) on chloroquine and amodiaquin accumulation

Amodiaquine

McMahon, J. E., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 465-472
Wuchereria bancrofti, humans, preliminary screening of levamisole and amodiaquine for antifilarial activity; some trials combined with diethylcarbamazine

Camoquin

Mishra, A. K.; and Sharma, N. N., 1979, Trop. Animal Health and Prod., v. 11 (4), 222-226
Anaplasma marginale, calves (exper.), comparative efficacy of several drugs

Amodiaquine -- Continued.

Amodiaquine

Montgomerie, A. M.; Proctor, G. R.; and Green, B., 1979, Biochem. Soc. Tr., v. 7 (6), 1251-1253
binding to DNA of indolobenzazepine analogue of antimalarial drug amodiaquine

Amodiaquin

Schmidt, L. H.; et al., 1977, Antimicrob. Agents and Chemotherapy, v. 11 (5), 826-843
Plasmodium falciparum in Aotus trivirgatus, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

Amodiaquine hydrochloride. See Amodiaquine.

Amoebicide 2004. See Chlorphenoxamide or Tinidazole.

Amoebicides

Arnold, K., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (2), 177-185
human parasitic diseases, trends in development of chemotherapy, review

Amoebicides

Fernex, M., 1977, Schweiz. Med. Wchnschr., v. 107 (18), 643-644
human malariae and amoebiasis, brief review of current therapeutic and prophylactic measures

Amoebicides

Noemi, I., 1975, Rev. Chilena Pediat., v. 46 (1), 63-67
therapeutic recommendations for treating intestinal parasites in infants and small children, drug and dosage review

Amoebicides

Ocaranza M., J., 1972, Rev. Gastroenterol. Mexico (219), v. 37, 180-188
Entamoeba histolytica, humans, classification of amoebicides according to their chemotherapeutic actions, compilation of results obtained by various researchers

Amoebicides

Perpignano, G.; Argiolas, N.; and Pintus, A., 1974, Rassegna Med. Sarda, v. 77, n.s., v. 26 (2), 143-167
Entamoeba histolytica, human hepatic abscess, current clinical aspects reviewed, diagnosis by X-ray and scintigraphy, recommendations for therapy

Amoebicides

Rev. Med. Chile, 1978, v. 106 (4), 319-331
Entamoeba histolytica, current status of human amoebiasis (physiopathology, clinical forms, diagnosis, treatment, epidemiology), anniversary presentation at the Academy of Medicine of Chile

Amopyroquin

- Schmidt, L. H.; et al., 1977, Antimicrob. Agents and Chemotherapy, v. 11 (5), 826-843
Plasmodium falciparum in *Aotus trivirgatus*, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

Amphotalide

- Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124
 preparation and testing of 6 long-acting schistosomicidal resinsates

Amphotericin B -- Fungizone.**Amphotericin B (Fungizone)**

- Bjorvatn, B.; and Neva, F. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 480-485
Leishmania tropica, white mouse model, experimental therapy using sodium stibogluconate, amphotericin B, metronidazole and WR 6026

Amphotericin B

- Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Amphotericin B

- Castro, R. M.; et al., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (5), 338-339
 leishmaniasis, human, recent infection, amphotericin B treatment, Montenegro test positive only after 42 days, independent of treatment

Amphotericin B (Fungizon)

- Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113
 human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Amphotericin B

- Hermanne, J.; Jadin, J. B.; and Martin, J. J., 1972, Semaine Hop. Paris, v. 48 (21), 1539-1550 (Ann. Pediat., Paris, v. 19 (5), 425-436)
Naegleria gruberi, human, fatal meningoencephalitis after swimming in public pools, amoeba discovered in spinal fluid, amphotericin B studied as possible therapy: Belgium

Amphotericin B

- Kasprzak, W.; Mazur, T.; and Rucka, A., 1974, Ann. Soc. Belge Med. Trop., v. 54 (4-5), 351-357
 4 strains of free-living amoebae isolated from lakes in Poland, pathogenicity for mice, response to several drugs, identified as *Acanthamoeba* spp. on basis of morphology and protein disc electrophoretic patterns

Amphotericin B (Fungizone)

- Lee, K. K.; et al., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (2), 217-220
Naegleria fowleri, in vitro susceptibility to selected antimicrobial agents singly and in combination

Amphotericin B -- Continued.**Amphotericin B + Tetracycline**

- Moggian, G.; Tamburini, F.; and Visona, E., 1975, Quad. Clin. Ostet. e Ginec., v. 30 (4), 143-154
T[richomonas] vaginalis, human vulvovaginitis, clinical trials comparing mepartricin with amphotericin B combined with tetracycline in topical creams, good results

Amphotericin B

- Roizenblatt, J., 1979, Am. J. Opth., Chicago, v. 87 (2), 175-179
 American (mucocutaneous) leishmaniasis, human (eye), interstitial keratitis, case report, differential diagnosis, amphotericin B: Brazil

Amphotericin B + Tetracycline

- Ruggeri, E., 1976, Quad. Clin. Ostet. e Ginec., v. 31 (4), 211-223
Trichomonas vaginalis, vaginal trichomoniasis, oral mepartricin showed significant therapeutic superiority when compared in clinical trials with patients who received nimorazole therapy orally or with patients treated with a vaginal cream containing amphotericin B and tetracycline

Amphotericin B

- Thong, Y. H.; Rowan-Kelly, B.; and Ferrante, A., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (3), 336-337
Naegleria fowleri-infected mice, amphotericin B potentiated by tetracycline shows promise in treating meningoencephalitis

Amphotericin B methyl ester

- Lee, K. K.; et al., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (2), 217-220
Naegleria fowleri, in vitro susceptibility to selected antimicrobial agents singly and in combination

Amprol. See Amprolium.**Amprolium -- 1-(4-Amino-2-n-propyl-5-pyrimidinyl-**

methyl)-2-picolinium chloride hydrochloride; Amprol; Amprolmix plus (with Ethopabate); Amprol plus (with Ethopabate); Amprolsol; Amprolvet; Amprosol; Coccidiovit; Corid; Pancoxin (with Ethopabate and Sulfaquinoxaline); Pancoxin plus (with Ethopabate, Pyrimethamine, and Sulfaquinoxaline); Supacox (with Ethopabate, Pyrimethamine, and Sulfaquinoxaline); Thiaccoclide.

Amprolium

- Augustine, P. C.; Vetterling, J. M.; and Doran, D. J., 1977, Proc. Helminth. Soc. Washington, v. 44 (2), 147-149
Eimeria tenella, comparison of amprolium- and buquinolate-resistant strains to a drug-sensitive strain with respect to (1) oocyst production in chicks and (2) infectivity, rate of development, and oocyst production in primary chick kidney cell cultures

Amprolium -- Continued.

Coccidiovit

Avakian, A. A.; et al., 1978, Veterinaria, Moskva (11), 76-77
coccidiosis, chickens (broilers), prophylactic control by various preparations, pharmacococci recommended, treatment economics: Krymsk oblast

Amprolium

Bajwa, R. S.; and Gill, B. S., 1977, Acta Vet. Brno, v. 46 (1-2), 149-158
Eimeria tenella, chickens (exper.), decoquinate vs. amprolium used prophylactically vs. therapeutically, anticoccidial activity against different levels of infection, effect on development of immunity

Amprol Plus

Bedrnik, P., 1977, Veterinarstvi, v. 27 (10), 458-459
coccidiosis, chickens, current prevalence, amprol plus treatment

Amprol plus

Bedrnik, P.; et al., 1979, Veterinarstvi, v. 29 (8), 353-355
coccidiosis, broiler chicken fattening, lasalocid compared with amprol plus and monensin: Czechoslovakia

Amprolium (Amprolvet)

Brunthaler, F., 1977, Prakt. Tierarzt, v. 58 (11), 849-851
coccidiosis, dogs, treatment with various drugs, best results with amprolium

Amprolium plus Ethopabate (= Amprol plus)

Buys, S. B.; and Rasmussen, R. W., 1978, J. South African Vet. Ass., v. 49 (2), 127-128
significantly higher mortalities found in nicarbazine fed chickens than in amprolium fed chickens when exposed to same heat stress conditions

Amprolium

Chapman, H. D., 1978, Parasitology, v. 76 (2), 177-183
Eimeria tenella, Houghton strain, experimental development of resistance to amprolium, clopidol, and methyl benzoquate

Coccidiovit

Gobzem, V. R.; and Nazarov, V. G., 1978, Veterinaria, Moskva (3), 67-69
Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Amprolium + Ethopabate (=Amprolmix plus)

Hamet-Foure, N.; Macar, C.; and Robin, B., 1979, Avian Path., v. 8 (1), 107-113
Eimeria meleagrimitis, *E. adenoides*, turkeys, activity of clopidol with methylbenzoquate and amprolium with ethopabate: France

Amprolium

Horton, G. M. J.; and Stockdale, P. H. G., 1979, Am. J. Vet. Research, v. 40 (7), 966-970
Eimeria spp., lambs, amprolium and monensin, oocyst discharge, feed utilization, and ruminal metabolism

Amprolium -- Continued.

Amprolium

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidostats in feeds, qualitative identification test

Amprolium + Ethopabate

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Amprolium

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Amprolium + Sulphaquinoxaline + Ethopabate (= Pancocin)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Amprolium + Sulphaquinoxaline + Ethopabate + Pyrimethamine (= Supacox)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Amprolium (Amprol)

McDougald, L. R., 1979, Poultry Science, v. 58 (1), 76-80
coccidiosis, histomoniasis, turkeys, tests for efficacy and compatibility indicate that amprolium and carbarsone can be used in combination

Amprolium

McDougald, L. R.; and Johnson, J. K., 1979, Poultry Science, v. 58 (1), 72-75
Eimeria spp., turkeys, efficacy of arprinocid compared with amprolium, floor pen studies

Amprolium + Ethopabate

McDougald, L. R.; Karlsson, T.; and Reid, W. M., 1979, Avian Dis., v. 23 (4), 999-1005
coccidiosis, chickens (exper.), natural outbreak of infectious bursal disease (IBD) during comparison of anticoccidials for their effect on development of immunity, interaction between diseases, immunity to coccidiosis not blocked by IBD

Amprolium

McDougald, L. R.; and McQuiston, T. E., 1978, Avian Dis., v. 22 (4), 765-770
Eimeria spp., turkeys, coccidiosis management, innate (age) resistance and acquired immunity vs. anticoccidial medication

Amprolium

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Amprolium -- Continued.

- Amprolium (Amprol)
McQuistion, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anti-coccidial drugs
- Amprol plus
Manuel, M. F., 1972, Philippine J. Vet. Med., v. 11 (2), 92-100
Eimeria tenella, broilers (exper.), evaluation of amprol plus and coyden-25 with or without 3-nitro-50 (roxarsone)
- Amprol plus
Manuel, M. F.; and Buduan, R. J., 1972, Philippine J. Vet. Med., v. 11 (2), 73-91
battery-raised broilers, effects of amprol plus or cycostat with or without payzone on weight gain and feed efficiency, no coccidiosis found
- Amprolium (Amprol)
Manuel, M. F.; and Neri, R. A., 1975, Philippine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats
- Amprolium
Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups
- Pancoxin plus
Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups
- Amprosol
Mogollon, J. D.; and Mofina, T. de P., 1977, Rev. Inst. Colomb. Agropec, v. 14 (4), 473-488
Eimeria tenella field strain, sensitivity against 3 anticoccidial drugs
- Amprolium + Ethopabate
Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products
- Amprolium (Corid)
Sangster, L. T.; et al., 1978, Vet. Med. and Small Animal Clin., v. 73 (10), 1317-1319
coccidiosis, baby pigs, cause of scours, clinical and pathologic features, treatment with amprolium: southern Georgia
- Amprolium
Sansi, K. A. O., 1976, Bull. Animal Health and Prod. Africa, v. 24 (1), 93-96
haemorrhagic syndrome in chickens, possibly associated with use of amprolium as coccidiostat, case report: Vom, Nigeria

Amprolium -- Continued.

- Amprolium (Amprosol)
Santiago, M. A. M.; da Costa, U. C.; and da Silva, O. L., 1977, Rev. Centro Cien. Rurais, v. 7 (3), 297-301
Eimeria spp., sheep, natural infection, amprolium, furoxone, not sufficient control for sheep grazing on pasture
- Pancoxin
Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazin, and pancoxin: England; France; Germany
- Amprol plus
Sevcik, B.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21) (4), 375-383
coccidiosis, broilers, field trials of anti-coccidials: CSSR
- Amprolium
Sherkov, S., 1977, Vet. Med. Nauki, v. 14 (2), 37-44
Eimeria tenella, chickens (exper.), amprolium alone and with additional amounts of thiamine in feed mixture, evaluation of prophylactic use, thiamine contributes to lowered activity of amprolium
- Pancoxin
Sherkov, Sh., 1977, Vet. Sbirka, v. 75 (3), 35-38
Eimeria tenella, pancoxin, chickens raised under conditions of high and low temperatures; influence of thiamine on development of coccidiosis
- Pancoxin plus
Sherkov, S. N.; Leitch, B.; and Kokash, L., [1977], Egypt. J. Vet. Sc., v. 13 (1), 1976, 37-43
Sarcocystis tenella, kittens (exper.), development in intestines, life cycle; attempted parasite suppression using statyl and pancoxin plus
- Amprolium (Amprosol)
Singh, J.; and Gill, B. S., 1976, Riv. Parasitol., Roma, v. 37 (1), 63-70
Eimeria necatrix, different levels of infection, chicks, activity of amprolium used prophylactically and therapeutically, effect on development of immunity
- Amprolium (Amprosol)
Singh, J.; and Hussain, O., 1978, Indian Vet. J., v. 55 (1), 56-60
Eimeria tenella, chicks (exper.), amprolium provided better protection than codrinal, both drugs interfered to some extent with development of immunity
- Amprolium
Stockdale, P. H. G.; and Yates, W. D. G., 1978, Vet. Parasitol., v. 4 (3), 209-214
Eimeria zuernii, calves (exper.), successful chemotherapy with amprolium or monensin, resistance to reinfection after chemotherapy

Amprolium -- Continued.

Pancoxin plus

Stoianov, P.; et al., 1978, Vet. Med. Nauki, v. 15 (8), 105-114
E[imeria] tenella, broiler chickens, varied temperature and moisture regimes, blood biochemistry, host resistance, efficacy of pancoxin plus

Amprolium

Villadelgado, M. A.; and Abuso, O. T., [1977], Philippine J. Vet. Med., v. 15 (1-2), 1976, 123-128
furazolidone as supplement to commercial broiler feed (already containing penicillin and amprolium), results indicate supplementation unwarranted

Amprolium + Ethopabate (= Amprol Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Amprolium + Ethopabate + Sulfaquinoxaline + Pyrimethamine (= Pancoxin Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Amprolium (Thiacoccide)

Volont, L. A.; Rudakov, V. V.; and Nikol'skaia, E. B., 1979, Khimiko Farm. Zhurnal, v. 13 (6), 10-12
amprolium, inhibition of activity of some cholinesterases, possible role in molecular mechanism of side effects in livestock

Amprolium

Zaprianov, M. Ia., 1978, Vet. Med. Nauki, v. 15 (4), 103-107
Eimeria kofoidi in Alectoris graeca cypriatis (nat. and exper.), amprolium, DOT-soluble, and Esb; tested

Amprolmix plus. See Amprolium + Ethopabate.

Amprol plus. See Amprolium or Ethopabate.

Amprolsol. See Amprolium.

Amprolvet. See Amprolium.

Amprosol. See Amprolium.

Amylosubtylene

Mustaikis, B.; and Medzevicius, A., 1978, Lietuvos TSR Moks. Akad. Darbai, s. C (82), (2), 75-83
Trichocephalus suis, suckling pigs, influence of amylosubtylene on host carbohydrate metabolism

Ancaris. See Piperazine or Thenium.

Ancylo1. See Disophenol.

Antabuse. See Disulfiram.

Anthelcide-EQ. See Oxibendazole.

Anthelmintics

Alvarez Chacon, R., 1975, Semana Med. Mexico (1071), an. 22, v. 84 (11), 328-332
parasites of children, drugs in current use, treatment recommendations and contraindications

Anthelmintics

Arnold, K., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (2), 177-185
human parasitic diseases, trends in development of chemotherapy, review

Anthelmintics

Behm, C. A.; and Bryant, C., 1979, Vet. Parasitol., v. 5 (1), 39-49
anthelmintic action--a metabolic approach, a review

Anthelmintics

Boersema, J. H., 1977, Tijdschr. Diergeneesk., v. 102 (24), 1450-1455
trichostrongylids and lungworms in ruminants, review of important data on spectrum, mechanism of action, resistance, and toxicity of anthelmintics used in the Netherlands

Anthelmintics

Corwin, R. M., 1979, Equine Pract., v. 1 (1), 51-55
equine parasites, guide to anthelmintics for control and eradication

Anthelmintics

Dasgupta, N. K., 1976, Modern pharmacology and therapeutics, 2. ed., 860 pp., illus.

Anthelmintics

Do Duong Thai; and Nguyen Tuyet Mai, 1973, Rev. Med., Hanoi, 165-168
Ascaris lumbricoides, laboratory studies on drug resistance to anthelmintics (ascaridin used in experiments), finding that sex of worm, maturity of worm, and drug concentrations all influenced reactions to drugs

Anthelmintics

Dorchies, P., 1979, Bull. G.T.V. (3), 43-50
strongylosis, anthelmintics, mechanisms of action, factors causing variation in activity, limitations, review

Anthelmintics

Enigk, K., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (3), 77-81
nematodes, pigs, treatment and prophylaxis (tetramisole; pyrantel tartrate; panacur; mebendazole)

Anthelmintics

Epel'dimov, L. S., 1977, Sborn. Nauch. Rabot SibNIVI (28), 76-83
ascariasis, swine, dictyocaulosis, sheep, research and industrial trials of new anthelmintics, review

- Anthelmintics**
 Euzeby, J., 1978, Cahiers Med. Vet., v. 47, spec. no., 165-184
 fascioliasis and strongylosis of ruminants, cestodiasis of all species of animals, anthelmintic therapy, review
- Anthelmintics**
 Euzeby, J., 1978, Bull. Soc. Sc. Vet. et Med. Comp. Lyon, v. 80 (3), 123-137
 cestodiasis, review of advances in chemotherapy and the possibilities of immunotherapy
- Anthelmintics**
 Euzeby, J., 1979, Rev. Med. Vet., Toulouse, v. 130 (2), 241-246, 249-256
 Dirofilaria immitis, dogs, treatment and prophylaxis, extensive review
- Anthelmintics**
 Gandhi, T. P.; and Shah, G. F., 1978, Indian Drugs, v. 15 (8), 137-142
 recent trends in the treatment of intestinal worms, review
- Anthelmintics**
 Goldsmith, R. S., 1974, Rev. Med. Pharmacol., 4 ed., 621-649
 anthelmintics in current use, extensive review of modes of action, epidemiologic factors, clinical administration, contraindications and cautions
- Anthelmintics**
 Kimura, S.; and Shimizu, A., 1977, Nippon Zyuishi-Kai Zassi (J. Japan Vet. Med. Ass.), v. 30 (12), 642-645
 Fasciola hepatica, cattle, recommended timing of anthelmintic treatment: Kobe area
- Anthelmintics**
 Lengyel, A.; and Janko, M., 1979, Orvosi Hetilap, v. 120 (49), 2969-2972
 human intestinal parasites, current therapeutic regimens, review: Hungary
- Anthelmintics**
 Miller, R. F., 1979, Proc. 1979 Maryland Nutrition Conf. Feed Mfr. (University of Maryland, Mar. 15-16), 87-96
 parasitism, ruminants, influence on utilization of nutrients, timing of treatment, milk production, reproduction, review
- Anthelmintics**
 Niemand, H. G., 1976, Prakt. Tierarzt, v. 57 (5), 321-323
 intestinal helminths and coccidia, dogs, therapy, review
- Anthelmintics**
 Noemi, I., 1975, Rev. Chilena Pediat., v. 46 (1), 63-67
 therapeutic recommendations for treating intestinal parasites in infants and small children, drug and dosage review
- Anthelmintics**
 Popov, M. A., 1978, Veterinariia, Moskva (8), 65-68
 [Strongylata], sheep, goats, specialized farms, economic benefits and seasonal timing of anthelmintic treatment
- Anthelmintics**
 Schroeder, J.; and Swan, G. E., 1979, Rhodesian Vet. J., v. 10 (1), 11-16
 anthelmintic treatment of cattle, epizootiological and economic considerations, review
- Anthelmintics**
 Seo, B. S., 1977, Taehan Uihak Hyophoe Chi (J. Korean Med. Ass.), v. 20 (6), 503-508
 human intestinal parasitic disease, current therapy, review
- Anthelmintics**
 Willgeroth, F.; and Rummel, W., 1979, Fortschr. Med., v. 97 (47), 2180-2186
 drugs in pregnancy and lactation, includes anthelmintics, adverse effects, review
- Anthiolimine** -- Anthiomaline; Antimony lithium thiomalate; Antiomaline; Lithium antimony thiomalate.
- Antimony lithium thiomalate (Anthiomaline)**
 Ghazal, A.; Ismail, M.; and Sharabi, F. M., 1978, Egypt. J. Pharm. Sc., v. 17 (1), 1976, 43-52
 Schistosoma mansoni-infected mice, tissue histamine content before and after treatment with several antischistosomal drugs
- Anthiomaline**
 Minor Franco, A.; and Sanchez Granados, P., 1974, Rev. Med. Hosp. Gen., Mexico, v. 37 (6), 383-387
 human cutaneous leishmaniasis, case report, unsuccessfully treated with metronidazole but cured with anthiomaline; high percentage of localization on ears and ear lobes in areas of Mexico
- Anthiomaline**
 Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
 Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared
- Anthiomaline**
 Singh, D. V.; Joshi, H. C.; and Shivnani, G. A., 1972, Philippine J. Vet. Med., v. 11 (2), 101-109
 Setaria cervi, Onchocerca armillata, buffaloes, clinical, hematological, and blood biochemical studies, efficacy of anthiomaline
-
- Anthiomaline.** See Anthiolimine.
- (2,6-Anthraquinonylene)bis[N',N'''-diethylacetamide]
 Burden, E. J.; et al., 1979, Experientia, v. 35 (1), 33-35
 Entamoeba histolytica, rats, hamsters, potent activity of bisamidines of 2,6-diaminoanthraquinone
- Antibiotic X-464.** See Nigericin.

Anticoccidials -- Coccidiostats.

Anticoccidials

Niemand, H. G., 1976, *Prakt. Tierarzt*, v. 57 (5), 321-323
intestinal helminths and coccidia, dogs, therapy, review

Coccidiostats

Plisek, K., 1974, *Veterinaria, Praha*, v. 16 (5-6), 589-640
coccidiostats, survey of currently used substances, toxicity, modes of application, review

Coccidiostats

Sevcik, B.; et al., 1974, *Veterinaria, Praha*, v. 16 (5-6), 421-588
Eimeria tenella, chickens, 613 substances screened as coccidiostats, extensive detailed statistical results

Anticoccidials

Zaionts, V. I.; et al., 1976, *Khimiko-Farm. Zhurnal*, v. 10 (11), 20-29
molecular mechanisms of action of some anticoccidial preparations, review

Antimalarials

Arnold, K., 1978, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 9 (2), 177-185
human parasitic diseases, trends in development of chemotherapy, review

Antimalarials

Fernex, M., 1977, *Schweiz. Med. Wchnschr.*, v. 107 (18), 643-644
human malarial and amoebiasis, brief review of current therapeutic and prophylactic measures

Antimalarials

Gear, J. H. S.; et al., 1978, *South African Med. J.*, v. 53 (23), 921 [Letter]
malaria, human, treatment recommendations

Antimalarials

Guyer, B.; and Candy, D., 1979, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 73 (2), 230-232
malaria, febrile children, problems associated with administration of injectable quinine compounds over oral chloroquine preparations: tropical Africa

Antimalarials

Hansford, C. F., 1974, *South African Med. J.*, v. 48 (30), 1314-1316
antimalarials, general review of drugs currently available with special reference to those of benefit in South Africa

Antimalarials

Jopling, W. H., 1979, *Lancet*, London (8130), v. 1, 1340 [Letter]
malaria, travellers, chemoprophylaxis, recommendations

Antimalarials

Loos, E., 1974, *Bull. Soc. Sc. Med. Gr.-Duche Luxembourg*, v. 111 (1), 19-25
synthetic antimalarials, ocular complications that can result from prolonged use

Antimalarials

Powell, R. D., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (1, pt. 1), 1-5
Plasmodium spp., servicemen returning from Vietnam, relationships between chemoprophylaxis history and occurrence and timing of post-departure episodes of malaria: Iowa City VA Hospital

Antimalarials

Rev. Prat., Paris, v. 27 (37), 1977, 2341-2409
Plasmodium spp., issue devoted to current status of human malarial with discussion on: life cycle, geographic distribution, immunopathology, diagnosis, clinical management, treatment and prophylaxis

Antiminth. See Pyrantel.

Antimonial compounds

Woolhouse, N. M., 1979, *Biochem. Pharmacol.*, v. 28 (16), 2413-2418
antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Antimony dimercapto-potassium succinate

Furtado, T., 1974, *Rev. AMMG*, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Antimony dimercapto-succinate. See Stibocaptate.Antimony lithium thiomalate. See Anthiolimine.Antimony potassium tartrate -- Potassium antimonyl tartrate; Potassium antimony tartrate; Tartar emetic.

Tartar emetic

Abdel Samad, M. M.; et al., 1977, *Tropenmed. u. Parasitol.*, v. 28 (4), 554-559
Schistosoma mansoni, mice, liver monoamine oxidase activity during course of infection and after chemotherapy, may be useful index for progression or regression of liver fibrosis

Tartar emetic

Abdulla, W. A.; et al., 1977, *Egypt. J. Bilharz.*, v. 4 (1), 19-26
S[chistosoma] mansoni, anti-schistosomal activity of the plant *Ammi majus* and bergapten (furocoumarin present in *A. majus*), compared with tartar emetic, screening in mice

Tartar-emetic (Potassium-antimony tartrate)

Abdulla, W. A.; et al., 1977, *Egypt. J. Bilharz.*, v. 4 (2), 129-140
bilharzid much less toxic than tartar emetic when given in same dose level to guinea pigs, dogs, and monkeys

Potassium antimony tartrate

Coles, G. C.; and Chappell, L. H., 1979, *Exper. Parasitol.*, v. 47 (1), 49-53
Schistosoma mansoni, effects of potassium antimony tartrate on immature vs. adult worms in vitro and in vivo, if antimony acts by inhibition of phosphofructokinase it is not clear why young worms are more resistant to chemotherapy than adults

Antimony potassium tartrate -- Continued.

- Potassium antimony tartrate
Doenhoff, M. J.; and Bain, J., 1978, Clin. and Exper. Immunol., v. 33 (2), 232-238
Schistosoma mansoni-infected mice deprived of their T-cells, relative lack of efficacy of potassium antimony tartrate, demonstration of drug-antiserum synergy
- Antimony potassium tartrate (Tartar emetic)
Ghazal, A.; Ismail, M.; and Sharabi, F. M., 1978, Egypt. J. Pharm. Sc., v. 17 (1), 1976, 43-52
Schistosoma mansoni-infected mice, tissue histamine content before and after treatment with several antischistosomal drugs
- Antimony potassium tartrate
Khayyal, M. T.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 149-156
S[chistosoma] mansoni, mice, antimony potassium tartrate therapy given with penicillamine as adjuvant gives same therapeutic results with fewer side effects; ameliorates lipid changes in host but not in parasites
- Antimony potassium tartrate
Lee, H. S.; Lee, H. B.; and Moon, M. H., 1979, Korean J. Animal Sc., v. 21 (4), 281-288
Besnoitia besnoiti, goats (exper.), antimony potassium tartrate, pacprim
- Tartar emetic
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity
- Potassium antimony tartrate (Tartar emetic)
Mahran, S. G.; et al., 1976, Egypt. J. Bilharz., v. 3 (2), 239-245
Schistosoma mansoni, in vivo and in vitro trials comparing efficacy of bilharcid with that of tartar emetic
- Potassium antimony tartrate
New, R. R. C.; et al., 1978, Nature, London (5648), v. 272, 55-56
Leishmania donovani, mice, reduced parasite count in liver after treatment with drug-loaded liposomes (potassium antimony tartrate, sodium antimony gluconate), enhanced activity as compared to drug injected alone
- Tartar emetic
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared
- Tartar emetic (Potassium antimonyl tartrate)
Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1978, Egypt. J. Bilharz., v. 4 (1), 1977, 9-18
schistosomicides, comparative study on effects of bilharcid, tartar emetic and piperazine hexahydrate on activity of oxidoreductase enzymes of rat liver preparations

Antimony potassium tartrate -- Continued.

- Tartar-emetic (Potassium antimony tartrate)
Shehata, H.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 117-128
S[chistosoma] mansoni, mice with anemia and leucocytosis, bilharcid safer therapy than tartar emetic
- Tartar emetic
Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method
-
- Antimony sodium gluconate -- Pentostam; Sodium antimony gluconate; Sodium antimonyl gluconate; Sodium stibogluconate; Triostam; Triostib.
- Sodium stibogluconate (Pentostam)
Alving, C. R.; et al., 1978, Proc. National Acad. Sc., v. 75 (6), 2959-2963
Leishmania donovani, hamsters, superior efficacies of liposome-encapsulated meglumine antimoniate and sodium stibogluconate, efficacy of treatment influenced by lipid composition and charge of liposomes, morphologic evidence that liposomes travel to intracellular site of parasite, encapsulation and reduction of dose should minimize toxic reactions to antimonials
- Sodium stibogluconate (Pentostam)
Bjorvatn, B.; and Neva, F. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 480-485
Leishmania tropica, white mouse model, experimental therapy using sodium stibogluconate, amphotericin B, metronidazole and WR 6026
- Sodium antimony gluconate (Triostib)
Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy
- Sodium stibogluconate (Pentostam)
Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy
- Sodium stibogluconate (Pentostam)
Garett, A., 1978, Southwest. Vet., v. 31 (2), 125-128
Leishmania donovani in Doberman Pinscher dog (bone marrow macrophage), case report, inconclusive treatment with sodium stibogluconate: Texas, imported from Greece
- Sodium stibogluconate (Pentostam)
Ghosn, S., 1979, Current Med. Research and Opin., v. 6 (4), 280-283
cutaneous leishmaniasis, 31 patients with positive smears, evaluation of sodium stibogluconate intramuscularly or infiltrated around lesions: Syria

Antimony sodium gluconate -- Continued.

- Sodium antimony gluconate (Pentostam)
Kaplan, R. J.; Wilkin, J. K.; and Hartman, D. L., 1978, *South. Med. J.*, v. 71 (4), 469-470
cutaneous leishmaniasis, man, sodium antimony gluconate, transient electrocardiogram abnormalities which cleared after termination of therapy: Tennessee (had lived in Central and South America for previous 18 months)
- Sodium stibogluconate (Pentostam)
Maru, M., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (1), 15-18
visceral leishmaniasis, hospitalized patients, clinical and laboratory aspects, pentostam or neostibosam therapy: northwestern Ethiopia
- Sodium stibogluconate
Naik, S. R.; et al., 1979, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 73 (1), 61-65
Leishmania donovani, human, 24 sporadic cases of kala-azar, epidemiology, clinical features, complications and associations, hematological and biochemical findings, diagnosis, treatment: north-western India
- Sodium stibogluconate (Pentostam)
Neal, R. A., 1972, *Rev. Inst. Med. Trop. S. Paulo*, v. 14 (6), 341-351
Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate
- Sodium antimony gluconate (Pentostam)
New, R. R. C.; et al., 1978, *Nature*, London (5648), v. 272, 55-56
Leishmania donovani, mice, reduced parasite count in liver after treatment with drug-loaded liposomes (potassium antimony tartrate, sodium antimony gluconate), enhanced activity as compared to drug injected alone
- Triostam (Sodium antimony gluconate)
Pereira, L. H.; et al., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (3), 123-126
Schistosoma mansoni, mice, simple and rapid method for mass screening of prophylactic agents using peritoneal schistosomula
- Sodium stibogluconate
Raether, W.; Seidenath, H.; and Loewe, H., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 543-547
Leishmania donovani, golden hamsters, HOE 668 compared with known antileishmanial drugs, toxicity precludes further development but very good anti-leishmanial action qualifies it as standard compound in screening tests
- Sodium antimony gluconate
Sakamoto, T., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice
- Sodium antimony gluconate
Sapire, D. W.; and Silverman, N. H., 1970, *South African Med. J.*, v. 44 (33), 948-950
urinary schistosomiasis, 10-year-old boy, sodium antimony gluconate treatment caused acute poisoning with myocardial involvement, suggestions for management

Antimony sodium gluconate -- Continued.

- Sodium antimony gluconate (Triostam)
Silverman, N. H.; Katz, J. S.; and Levin, S. E., 1973, *South African Med. J.*, v. 47 (6), 219-221
Capillaria hepatica in 17-month-old female, clinical presentation of visceral larva migrans, liver biopsy confirmed diagnosis, diethylcarbamazine and sodium antimony gluconate improved but did not cure infection: district of Bethal, South Africa
- Sodium antimony gluconate
Swarup-Mitra, S.; Chowdhury, A. K. R.; and Sarkar, M., 1979, *Indian J. Med. Research*, v. 69, 571-576
Leishman[ia] donovan[i], human, erythrocytic enzymes inhibited, haematological changes, normalization following sodium antimony gluconate therapy
-
- Antimony tartrate
Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, *J. Parasitol.*, v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy
- Antiomaline. See Anthiolimine.
- Antiparasitics
Dorchies, P.; Franc, M.; and Ducos de Lahitte, J., 1978, *Prat. Vet. Equine*, v. 10 (4), 161-164
review of antiparasitics used in treatment of equine digestive parasitoses
- Antiparasitics
Fisher, M. H.; and Wang, C. C., 1978, *Ann. Rep. Med. Chem.*, v. 13, 130-138
antiparasitic agents, review
- Antiparasitics
Laemmler, G., 1977, *Pharmakol. u. Toxikol.* (Frimmer), 2. ed., 75-106
includes extensive review of antiparasitics used in veterinary medicine, toxicity
- Antiparasitics
Van den Bossche, H., 1978, *Nature*, London (5664), v. 273, 626-630
chemotherapy of parasitic infections, review
- Antrycide. See Quinapyramine.
- Antrycide dimethylsulphate. See Quinapyramine.
- Antrycide prosalt. See Quinapyramine.
- Antrypol. See Suramin.

Arabinosyl adenine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Arabinosyl-6-mercaptapurine

Senft, A. W.; and Crabtree, G. W., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1847-1856

Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Aralen. See Chloroquine.

Ardenone-25

Avakian, A. A.; et al., 1978, *Veterinariia*, Moskva (11), 76-77

coccidiosis, chickens (broilers), prophylactic control by various preparations, pharmacococci recommended, treatment economics; Krymsk oblast

Ardenone-25

Krylov, V. F., 1978, *Veterinariia*, Moskva (10), 68-69

Eimeria tenella strain resistant to pharmacocid after 35 laboratory passages in chickens, cross-resistance only to rigecocin

Areco-Canine Tablets. See Arecoline.Arecoline -- Areco-Canine Tablets; Arecoline hydrobromide.

Arecoline hydrobromide

Gemmell, M. A., 1978, *Bull. World Health Organ.*, v. 56 (3), 433-443

tapeworm control in dogs for prevention of hydatidosis and cysticercosis in sheep, monthly drug treatment program, age-specific prevalence of *Taenia hydatigena* in lambs used as principal indicator, 13-year assessment: Styx Valley and Maniototo County, South Island, New Zealand

Arecoline hydrobromide (Areco-Canine Tablets)

Todd, K. S., jr.; Howland, T. P.; and Woerpel, R. W., 1978, *Am. J. Vet. Research*, v. 39 (2), 315-316

Mesocostoides corti, dogs (exper.), bunamidine hydrochloride and uredofos (good results), arecoline hydrobromide and niclosamide (variable results)

Arecoline hydrobromide. See Arecoline.Aristocort. See Triamcinolone.Arpalit. See Trichlorfon.Arpocox. See Arprinocid.Arprinocid -- 6-Amino-9-(2-chloro-6-fluorobenzyl) purine; Arpocox; 9-(2-Chloro-6-fluorobenzyl) adenine; 9-(2-Chloro-6-fluorophenylmethyl)-9H-purin-6-amine; ICI 123487; MK-302.

Arprinocid

Karlsson, T.; and Reid, W. M., 1978, *Avian Dis.*, v. 22 (3), 487-495

Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Arprinocid (MK-302; Arpocox)

Kilgore, R. L.; et al., 1978, *Poultry Science*, v. 57 (4), 907-911

Eimeria spp., broiler chickens (exper.), arprinocid, effects on weight, feed efficiency, and histological lesions

Arprinocid (MK-302)

Kutzer, E.; et al., 1979, *Wien. Tierarztl. Monatsschr.*, v. 66 (6-7), 197-202

Eimeria spp., broiler chickens, arprinocid, drug efficacy compared with monensin

Arprinocid (MK-302; ICI 123487)

Latter, V. S.; and Wilson, R. G., 1979, *Parasitology*, v. 79 (1), 169-175

Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Arprinocid

Long, P. L.; Millard, B. J.; and Smith, K. M., 1979, *Avian Path.*, v. 8 (4), 453-467

Eimeria spp., chickens, effect of 4 anticoccidial drugs on development of immunity, field and laboratory conditions

Arprinocid (Arpocox)

McDougald, L. R.; and Johnson, J. K., 1979, *Poultry Science*, v. 58 (1), 72-75

Eimeria spp., turkeys, efficacy of arprinocid compared with amprolium, floor pen studies

Arprinocid

McDougald, L. R.; Karlsson, T.; and Reid, W. M., 1979, *Avian Dis.*, v. 23 (4), 999-1005

coccidiosis, chickens (exper.), natural outbreak of infectious bursal disease (IBD) during comparison of anticoccidials for their effect on development of immunity, interaction between diseases, immunity to coccidiosis not blocked by IBD

Arprinocid (Arpocox)

McQuistion, T. E.; and McDougald, L. R., 1979, *Ztschr. Parasitenk.*, v. 59 (2), 107-113

Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Arprinocid -- Continued.

Arprinocid (ARPOCOX)

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Arprinocid (MK-302; Arpocox)

Ruff, M. D.; et al., 1978, Avian Dis., v. 22 (1), 32-41
Eimeria spp., broilers (exper.), arprinocid, drug efficacy in both battery and floor-pen trials

Arprinocid (MK-302)

Ruff, M. D.; Anderson, W. I.; and Reid, W. M., 1978, J. Parasitol., v. 64 (2), 306-311
Eimeria spp. in broilers, arprinocid decreased number of oocysts produced, fewer of the oocysts sporulated, and those oocysts which did sporulate were less infective than those from unmedicated birds

Arprinocid (Arpocox; MK-302)

Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazine, and pancosin: England; France; Germany

Arprinocid (Arpocox)

Tamas, T.; et al., 1978, Poultry Science, v. 57 (2), 381-385
Eimeria acervulina, *E. maxima*, *E. necatrix*, *E. tenella*, arprinocid and dichloro analog L-628,914 in feed, decrease in oocysts' capacity to sporulate

Arprinocid

Wang, C. C.; et al., 1979, Biochem. Pharmacol., v. 28 (15), 2249-2260
 arprinocid inhibits hypoxanthine-guanine transport, may be mode of anticoccidial action

Arprinocid (MK-302)

Wang, C. C.; Simashkevich, P. M.; and Stotish, R. L., 1979, Biochem. Pharmacol., v. 28 (15), 2241-2248
Eimeria tenella, mode of anticoccidial action of arprinocid

Arprocarb. See Propoxur.

Arsanilic acid

Henry, S. C., 1979, J. Am. Vet. Med. Ass., v. 174 (6), 601-603
Eperythrozoon suis, swine, clinical signs of infection confirmed by indirect hemagglutination and measuring packed cell volume; oxytetracycline and arsanilic acid combined with lice control, arsenic toxicosis

Arsenamide -- Caparsolate; Caparsolate sodium; Filaramide; Sodium thiacetarsamide; Thiacetarsamide sodium.

Sodium thiacetarsamide (Caparsolate sodium)
 Hawe, R. S., 1979, J. Am. Animal Hosp. Ass., v. 15 (5), 577-582
Dirofilaria immitis, cat (cranial vena cava), clinical diagnosis in absence of circulating microfilariae, sodium thiacetarsamide

Thiacetarsamide sodium (Caparsolate sodium)
 Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Thiacetarsamide sodium (Caparsolate)
 Stokhof, A. A.; and Wolvekamp, W. T. C., 1978, Tijdschr. Diergeneesk., v. 103 (2), 1121-1129
Dirofilaria immitis, dogs, 4 case reports, chemotherapy: Netherlands (imported from United States, South America, or South Africa)

Thiacetarsamide sodium
 Watson, A. D. J.; Farrow, B. R. H.; and Hoskins, L. P., 1978, Austral. Vet. Practitioner, v. 8 (3), 129-132
Haemobartonella felis, cats (nat. and exper.), thiacetarsamide sodium therapy; prednisolone, tetracycline, and chloramphenicol given sequentially

Thiacetarsamide sodium
 Wong, M. M.; and Suter, P. F., 1979, Am. J. Vet. Research, v. 40 (3), 414-420
Dirofilaria immitis, dogs (exper.) without microfilaremia, indirect fluorescent antibody titers, degree of eosinophilia, and radiologic findings before and after treatment, reinfection, necropsy findings, significance of tests, application to diagnosis

Thiacetarsamide sodium (Caparsolate; Filaramide)
 Zech, W. H., 1978, Canine Pract., Santa Barbara, v. 5 (5), 4 [Letter]
Dirofilaria immitis, dogs, one 'pre-injection' with caparsolate prior to initiation of full treatment regimen eliminates host toxic reaction to drug

Arsenic

Hammant, C. A., 1977, Rhodesian Vet. J., v. 8 (4), 67-70
 ticks, resistance survey of field strains to commonly used ixodicides, changeover from arsenic to dioxathion dipping of cattle, dramatic improvement in tick control, necessitates complete overhaul of dipping facilities and retraining of personnel: Tribal Trust Lands of Rhodesia

Arsenic

Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73
Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Arsenic trioxide
Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Arsenicals
Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

1-Aryl-2-(alkylamino)ethanol antimalarials
Kim, K. H.; et al., 1979, J. Med. Chem., v. 22 (4), 366-391
P[lasmodium] berghei, 646 1-aryl-2-(alkylamino)ethanol antimalarials, quantitative structure-activity relationship, mathematical analysis

6-[[Aryl and aralkyl]amino]methyl]-2,4-pteridine-diamines and -pteridinediamine 8-oxides
Worth, D. F.; et al., 1978, J. Med. Chem., v. 21 (4), 331-337
Plasmodium gallinaceum, P. berghei, 6-[[aryl and aralkyl]amino]methyl]-2,4-pteridinediamines and -pteridinediamine 8-oxides, synthesized for antimalarial evaluation, laboratory trials with experimental animals

1-Arylethyl quaternary ammonium bromides
Dimmock, J. R.; Smith, P. J.; and Tsui, S. K., 1979, J. Pharm. Sc., v. 68 (7), 866-871
antimicrobial and antineoplastic activity of benzylfluorenyl and 1-arylethyl quaternary ammonium salts, synthesis and evaluation

5-Arylfuryl-2-quinoxalines
Oleinik, A. F.; et al., 1978, Khimiko-Farm. Zhurnal, v. 12 (7), 71-77
5-aryl-2-quinoxalines, synthesis, anti-leishmanial activity in white mice

6-[[Aryloxy- and arylthio-)methyl]-2,4-pteridine-diamines and -pteridinediamine 8-oxides
Werbil, L. M.; et al., 1978, J. Med. Chem., v. 21 (4), 337-339
Plasmodium berghei, 6-[[aryloxy- and arylthio-)methyl]-2,4-pteridinediamines and -pteridinediamine 8-oxides, synthesis for antimalarial evaluation, none showed significant activity in trials with mice

Ascabiol. See Benzyl benzoate.

Ascarel. See Cascara sagrada or Santonin.

Ascaridil. See Tetramisole.

Ascaridin. See Chenopodium oil.

L-Asparaginase
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Aspidium -- Filixan; Male fern extract.

Male fern extract
Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91
cestodes, pathomorphology resulting from action of various anthelmintics

Filixan
Klenov, A. P., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 152-155
[Bothriocephalus], white amur, effectiveness of various anthelmintics

Aspidium
Mello, E. B. F.; et al., 1978, Zentralbl. Bakteriolog., 1. Abt. Orig., Reihe A, v. 241 (3), 384-387
Taenia solium, T. saginata, humans, oral treatment with aspidium preceded by hypertonic magnesium sulphate solution

Aspirin
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Astiban. See Stibocaptate.

Asuntol. See Coumaphos.

Asuntol 50. See Coumaphos.

Atabrine. See Quinacrine.

Atebrin. See Quinacrine.

Atgard. See Dichlorvos.

Atgard V. See Dichlorvos.

Atrican -- 2- α -Thenoyl-amino-5-nitrothiazole.

2- α -Thenoyl-amino-5-nitrothiazole
Benazet, F.; et al., 1970, Scand. J. Infect. Dis., v. 2 (2), 139-143
intestinal and hepatic parasites, nitroheterocyclic antiparasitics, laboratory studies of chemotherapeutic activity and toxicity in exper. animals

Atropine

Hillman, G. R.; Gibler, A. M.; and Anderson, J. W., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 992-997

Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

Aureomycin. See Chlortetracycline.

Avatec. See Lasalocid.

Avermectin fraction B_{1a} (C-076)

Benz, G. W.; and Ernst, J. V., 1979, *Am. J. Vet. Research*, v. 40 (8), 1187-1188
gastrointestinal nematodes, calves (exper.), evaluation of avermectin fraction B_{1a}, controlled experiment

Avermectin B_a

Blair, L. S.; and Campbell, W. C., [1979], *J. Parasitol.*, v. 64 (6), 1978, 1032-1034
Dirofilaria immitis, pre-cardiac stages in *Mustela putorius furo*, trials of avermectin B_a, mebendazole, and melarsoprol, possible value of *Dirofilaria-Mustela* model for chemotherapeutic studies

Avermectin B_{1a}

Blair, L. S.; and Campbell, W. C., 1979, *Am. J. Vet. Research*, v. 40 (7), 1031-1032
Dirofilaria immitis, dogs, avermectin B_{1a} rapidly removed microfilariae from blood, administered with adulticide (melarsoprol) removal was permanent

Avermectin B_{1a}

Egerton, J. R.; et al., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (3), 372-378
nematodes of cattle, sheep, dogs, and chickens, efficacy of avermectin B_{1a}

Avermectin B_{1a}

Fritz, L. C.; Wang, C. C.; and Gorio, A., 1979, *Proc. National Acad. Sc.*, v. 76 (4), 2062-2066
avermectin B_{1a}, effect on neuromuscular preparations of lobster, *Ascaris lumbricoides*, frog, and crayfish

Avermectin B_{1a}

Williams, M.; and Yarbrough, G. G., 1979, *European J. Pharmacol.*, v. 56 (3), 273-276
avermectin B_{1a}, enhancement of in vitro binding of ³H-diazepam to rat and mouse brain membranes, can also enhance some pharmacological actions of diazepam

Avermectin B_{2a}

Campbell, W. C.; Blair, L. S.; and Lotti, V. J., 1979, *J. Helminth.*, v. 53 (3), 254-256
Trichinella spiralis, mice, efficacy of avermectin B_{2a} not suppressed by agents known to block cholinergic neurotransmission

Avermectins

Blair, L. S.; and Campbell, W. C., 1978, *J. Helminth.*, v. 52 (4), 305-307
Ancylostoma caninum, dogs (exper.), avermectin B_{2a} more effective than B_{1a}, A_{1a} and A_{2a} are less effective than either; small trial indicates B_{1a} is also active against natural infections of *A. braziliense* in dogs

Avermectins

Burg, R. W.; et al., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (3), 361-367
avermectins, characteristics of the producing culture *Streptomyces avermitilis*, production of avermectin by fermentation, *Nematospiroides dubius*-mouse assay

Avermectins

Campbell, W. C.; and Blair, L. S., 1978, *J. Helminth.*, v. 52 (4), 308-310
Dirofilaria immitis, dogs (exper.), avermectins, activity against precardiac larvae, profound suppressive effect on circulating microfilariae, inactive against adult stage

Avermectins

Miller, T. W.; et al., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (3), 368-371
avermectins, isolation of complex from broth, separation of all its components, chromatographic properties of each

Avermectins (A₁, A₂, B₁, B₂)

Ostlind, D. A.; Cifelli, S.; and Lang, R., 1979, *Vet. Rec.*, v. 105 (8), 168
Cuterebra spp., mice, avermectins

Avlochlor. See Chloroquine.

Avlochlor. See Chloroquine.

Avlothane. See Hexachloroethane.

Axenomycin D

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Axenomycins

Della Bruna, C.; Ricciardi, M. L.; and Sanfilippo, A., 1973, *Antimicrob. Agents and Chemotherapy*, v. 3 (6), 708-710
axenomycins, effectiveness against several cestode spp. in nat. and exper. infections of various animals

8-Aza-adenine

Irvin, A. D.; and Young, E. R., 1978, *Research Vet. Sc.*, v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Azacrine hydrochloride

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

5-Azacytidine

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

8-Aza 26-diaminopurine

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Aza-3 emetine

Gilbert, J.; et al., 1978, Farmaco, Pavia, ed. Scient., v. 33 (4), 237-252

Entamoeba histolytica-infected mice, aza-3 emetine inactive in comparative laboratory trials with known amoebicides

8-Azaguanine. See Guanazolo.

Azaserine

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Azaserine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Azathioprine -- Imuran.

Azathioprine

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Azathioprine -- Continued.

Imuran

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

DL-7-Azatriptophan

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

6-Azauracil

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Azidine. See Berenil.

Azuntol

Ueno, H.; and Chibana, T., 1978, Japan Agric. Research Quart., v. 12 (3), 152-156

Stephanofilaria okinawaensis, cattle, distribution, clinical signs, chemotherapy, intermediate host determined

Azure eosin

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65

Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

- Babesan. See 1,3-Di-6-quinolylurea.
- Bacdip. See Quintiofos.
- Bacillus amyloliquefasciens
Guida, V. O.; et al., 1974, Rev. Brasil. Med., v. 31 (7), 465-470
Schistosoma mansoni, humans with intestinal, hepato-intestinal and hepatosplenic compensated forms of infection, Bacillus amyloliquefasciens resulted in clinical and parasitologic cure, apparent enzymatic action on parasites
- Bac[illus] thuringiensis exotoxin (Turingin; Miazol)
Tonkonozhenko, A. P.; et al., 1977, Veterinariia, Moskva (6), 41-42
[Wohlfahrtia] myiasis, sheep, treatment with Bac[illus] thuringiensis exotoxin as dust
- Bacitracin MD + Halofuginone
Edgar, S. A.; and Flanagan, C., 1979, Poultry Science, v. 58 (6), 1476-1482
Eimeria spp. (recent field isolates resistant to various drugs), halofuginone with roxarsone and/or bacitracin MD
- Bacitracin MD + Halofuginone + Roxarsone
Edgar, S. A.; and Flanagan, C., 1979, Poultry Science, v. 58 (6), 1476-1482
Eimeria spp. (recent field isolates resistant to various drugs), halofuginone with roxarsone and/or bacitracin MD
- Baker's antifol
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Banminth. See Pyrantel.
- Banminth II. See Morantel.
- Banminth-C
Epel'dimov, L. S.; and Plotnikova, L. M., 1977, Sborn. Nauch. Rabot SibNIVI (28), 140-142
[Trichostrongylus], rabbits infected with sheep species as models for anthelmintic study, tests of nilverm, banminth-C, cupric carbonate
- Banminth D. See Diethylcarbamazine or Morantel.
- Banminth paste. See Pyrantel.
- Banocide. See Diethylcarbamazine.
- Basoquine. See Amodiaquine.
- Basudin. See Diazinon.
- Batestan. See Benoxafos.
- Baygon. See Propoxur.
- Baygon MEB. See Plifenate.
- Baymix Crumbles. See Coumaphos.
- Bayrena. See Sulfameter.
- Baytex. See Fenthion.
- Baytex-50. See Fenthion.
- Beclotiamine (5-Chloroethylthiamine; Clotiamine; Cocciden)
Matsuzawa, T., 1978, Parasitology, v. 77 (2), 235-241
Eimeria tenella, chickens, beclotiamine, mode of action studies; attempts to potentiate or antagonize its activity revealed that pyri-thiamine and 2,4-dinitrophenol also showed slight anticoccidial activity and that a combination of 2,4-DNP and beclotiamine was effective but weight gain was not as good as with beclotiamine alone
- Belmet. See Sulfadiazine or Sulfamethazine or Sulfamerazine.
- Bemarsal. See Diphetarson.
- Benacil
Iakubovskii, M. V., 1979, Veterinariia, Moskva (2), 41-42
ascariasis, trichocephaliasis, oesophagostomiasis, swine, comparative effectiveness and economic value of various drugs: Minsk oblast
- Benlate. See Benomyl.
- Benomy1 -- Benlate; Methyl-1-(butyl-carbamoyl)-2-benzimidazole carbamate.
- Benomyl (Benlate)
Brooks, W. M.; Cranford, J. D.; and Pearce, L. W., 1978, J. Invert. Path., v. 31 (2), 239-245
Nosema heliothidis, benomyl, at concentrations tested, not effective in eliminating infection in Heliothis zea in laboratory colonies, but definitely deleterious to microsporidium, potential should be further evaluated
- Methyl-1-(butyl-carbamoyl)-2-benzimidazole carbamate
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Benoxafos -- Batestan; Benoxophos; O,O-Diethyl-s-(5,7-dichlorobenzoxazol-2-ylmethyl)-dithio-phosphate; Hoechst 2910.

Benoxafos

Bonin, W., 1977, Berl. u. Munchen. Tierarztl. Wchnschr., v. 90 (2), 34-37
Amblyomma hebraeum, Psoroptes cuniculi, Melophagus ovinus, Dermanyssus gallinae, heptenophos, rapid mode of action, broad range of efficacy, short residual effect and effective as a vapour poison, compared with other standard drug preparations

Batestan (Hoechst 2910)

Chemtai, A. A. K., 1977, Bull. Animal Health and Prod. Africa, v. 25 (3), 299-306
acaricide-treated zebu cattle, blood cholinesterase, radiometric assay

Benoxophos (Batestan)

Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73
Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Benoxophos. See **Benoxafos.**

Bensalkon. See **Benzalkonium chloride.**

Benzalkonium chloride -- **Bensalkon.**

Bensalkon

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

5-Benzamido-2 (4-thiazolyl) benzimidazole
Aboul-Atta, N.; and Denham, D. A., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 671-672

Trichinella spiralis-infected mice, 5-benzamido-2 (4-thiazolyl) benzimidazole, effect on different life-cycle stages

5-Benzamido-2-(4-thiazolyl)-benzimidazole
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Benzazon VII -- 5-Nitro-2-furaldehyde thiosemicarbazone.

Benzazon VII

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Microbiol. (6), 35-50
Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Benzene hexachloride -- Benzene hexachloride, Gamma; BHC; -BHC; EQ-335; Gamatox; Gamexane; Gamma benzene hexachloride; Gamma BHC; Gamma-isomer hexachlorocyclohexane; Gammexane powder; α -HCH; γ -HCH; Hexachlorane; Hexachlorane, gamma isomer; 1,2,3,4,5,6-Hexachlorocyclohexane; Hexicide; Kwellada; Lindane; Lindatox-20; Neo-scabexaan; Neoscabexan.

Gamatox (BHC)

Atef, M.; and El-Say, A., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 73-84
Hyalomma dromedarii, Rhipicephalus s. sanguineus, laboratory tests (immersion technique) with delnav, supona, toxaphene, BHC

Gamma benzene hexachloride (Kwellada)

Belle, E. A.; et al., 1979, Canad. J. Pub. Health, v. 70 (2), 133-135
Sarcoptes scabiei, outbreak among hospital patients and staff, gamma benzene hexachloride; Demodex folliculorum also found: Hamilton, Ontario, probably imported from USA

Lindane

Blommers, L., 1979, J. Med. Entom., v. 16 (1), 82-83
Pediculus capitis, nymph rearing technique, insecticide tests against nymphs

Lindane

Blommers, L.; and van Lennep, M., 1978, Entom. Exper. et Applic., v. 23 (3), 243-251
Pediculus humanus capitis, school children, field trials with lindane, laboratory tests with lindane, dieldrin, and malathion, presence of resistance to lindane confirmed: Netherlands

Lindane (Neo-scabexaan)

Boersema, J. H., 1978, Tijdschr. Diergeneesk., v. 103 (7), 377-380
Chorioptes bovis, horse with foot-mange, resistant to coumaphos, lindane successful

Lindane (Benzene hexachloride)

Chakrabarti, A.; and Misra, S. K., 1979, Indian Vet. J., v. 56 (6), 497-500
Demodex canis, dogs, incidence in relation to season, host age, sex, and breed, clinical manifestations, in vivo and in vitro activity of several acaricides: India

Benzene hexachloride -- Continued.

Lindane

Cole, M. M.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 118-119
Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

 α -HCH

Delak, M.; and Radokovic, M., 1978, *Vet. Arhiv, Zagreb*, v. 48 (1), 17-22
Fasciola hepatica from bile ducts of cattle, and bovine liver samples, amount of contamination with DDT, γ -HCH, and α -HCH

 γ -HCH

Delak, M.; and Radokovic, M., 1978, *Vet. Arhiv, Zagreb*, v. 48 (1), 17-22
Fasciola hepatica from bile ducts of cattle, and bovine liver samples, amount of contamination with DDT, γ -HCH, and α -HCH

Lindane (Neoscabexan)

Dorrestein, G. M.; and Van Bronswijk, J. E. M. H., 1979, *Vet. Parasitol.*, v. 5 (4), 389-398
Trixacarus caviae as cause of mange in *Cavia porcellus* (nat. and exper.), clinical symptoms, pathology, treatment; papular urticaria in humans associated with mangy guinea-pigs: The Netherlands

Lindane

Drummond, R. O.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 130-133
Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Gammexane powder

El-Bahay, G. H.; et al., 1978, *Vet. Med. J., Giza*, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, *Argas persicus*, evaluation of 10 insecticides

Lindane

Gina, A.; et al., 1978, *Shendetesia Populiore* (144) (2), 15-18
Ornithonyssus bacoti, rodents, epidemiology, control with lindane: Tiranés

Lindane (EQ-335)

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, *J. Med. Entom.*, v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Lindane

Goszczynska, K.; and Styczynska, B., 1972, *Roczniki Panstw. Zakl. Hig.*, v. 23 (2), 245-251
Pediculus humanus humanus, selection of laboratory strain reared through several generations aimed at induction of resistance to DDT and lindane; resistance developed to DDT but not to lindane

Gamma BHC

Khan, M. H.; and Srivastava, S. C., 1977, *Indian J. Animal Health*, v. 16 (2), 137-140
Boophilus microplus engorged females, in vitro tests with dursban, gamma BHC, sumithion, supona, dimecron, egg production and viability; supona most effective

Benzene hexachloride -- Continued. γ HCH

Kirkwood, A. C.; Quick, M. P.; and Page, K. W., 1978, *Vet. Rec.*, v. 102 (3), 50-54
ectoparasites, sheep, showers and plunge dipping, efficacy using diazinon against *Lucilia sericata*, γ HCH against *Psoroptes communis ovis*

Gamma-benzene hexachloride (Hexicide)

Knudsen, E. A.; and Pedersen, T. G., 1979, *Ugeskr. Laeger*, v. 141 (43), 2965-2966
scabies, humans, therapeutic trials of hexicide vs. dioxanthogen: Denmark

Lindane

Lourens, J. H. M., 1979, *J. Econom. Entom.*, v. 72 (5), 790-793
Amblyomma variegatum, *A. lepidum*, baseline data on susceptibility to organochlorine acaricides, genetic basis for resistance in *A. variegatum*

Lindane

Lourens, J. H. M.; and van de Klashorst, G., 1979, *Ztschr. Ang. Entom.*, v. 87 (3), 230-238
Rhipicephalus appendiculatus, organochlorine susceptible and tolerant populations: East Africa

 γ -BHC

Lourens, J. H. M.; and Tatchell, R. J., 1979, *Bull. Entom. Research*, v. 69 (2), 235-242
Rhipicephalus evertsi evertsi, identification and inheritance of resistance factors to organochlorine acaricides, experimental hybrids between susceptible and resistant strains

Gamma BHC

Meermann, A., 1978, *Vet.-Med. Nachr.* (2), 212-215
Psoroptes ovis, sheep, phoxim, gamma BHC, laboratory and field trials: Nordfriesland

Benzene hexachloride

Needham, J. R., 1978, *J. Inst. Animal Techn.*, v. 29 (1), 1-15
Myobia musculi, *Myocoptes musculus*, conventional mouse colony, acaricides

Lindane

Ottenschot, T. R. F.; and Gil, D., 1978, *Tijdschr. Diergeneesk.*, v. 103 (2), 1104-1108
cheyletiellosis in long-haired cats, chronic pruritis, treatment with diazinon flea collars and lindane baths: Netherlands; Western Germany

Gamma-isomer hexachlorocyclohexane (Lindane)

Privora, M.; Rupes, V.; and Cerny, V., 1970, *Folia Parasitol.*, v. 17 (1), 81-84
Dermacentor marginatus, laboratory trials testing six insecticides

Gamatox

Rak, H., 1976, *J. Vet. Fac. Univ. Tehran*, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Benzene hexachloride -- Continued.

Lindane

- Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Lindane

- Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Lindane

- Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Hexachlorane

- Reshetniak, V. Z.; Bartenev, V. S.; and Firsov, N. F., 1977, Veterinariia, Moskva (10), 79
 anaplasmosis, sheep outbreak, pathology, oxytetracycline effective treatment; *Hyalomma scupense* as probable vector, hexachlorane in creolin emulsion for control; no further anaplasmosis or *H. scupense* found: Rostovsk oblast

Lindane (Gamexane)

- Schenone, H.; et al., 1971, Rev. Chilena Pediat., v. 42 (8), 561-566
 human scabies, incidence survey showed children under 15 to be most heavily infested, apparent spread through overcrowding of sleeping facilities of families, lindane therapy recommended as drug of choice: Santiago, Chile

Hexachlorane, gamma isomer

- Suiunchaliev, R. S., 1978, Veterinariia, Moskva (6), 63-65
 psoroptic mange, sheep, method for determining creolin concentration in dip containing emulsion of creolin, gamma isomer of hexachlorane, and water

Benzene hexachloride (B.H.C.)

- Vihan, V. S.; Sinha, N. K.; and Sahni, K. L., 1979, Indian Vet. Med. J., v. 3 (3), 197-198
Sarcoptes scabiei, sheep, tetmosol and benzene hexachloride compared

Hexachlorane, gamma isomer

- Vodianov, A. A., 1979, Veterinariia, Moskva (5), 46-48
 psoroptosis, sheep, hexachlorane, gamma isomer, mineral oil emulsion compared with creolin base preparations

BHC

- Young, E.; Zumpt, F.; and Whyte, I. J., 1972, J. South African Vet. Ass., v. 43 (2), 226
Sarcoptes scabiei in *Panthera leo*, skin lesions, successful treatment with BHC and/or malathion-containing preparations: Kruger National Park

Benzene hexachloride -- Continued.

Lindatox-20

- Zarzara, C., 1979, Rev. Crest. Animalelor, v. 29 (5), 42-45
Cimex lectularius, lindatox-20, neguvon, and carbetox 37, laboratory tests; carbetox 37 for control in poultry sheds, application methods

Benzene hexachloride, Gamma. See Benzene hexachloride.

Benzimidazoles

- Friedman, P. A.; and Platzer, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
 benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Benzimidazoles

- Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
 relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiriodes dubius* in mice

Benzimidazoles

- Kelly, J. D.; et al., 1978, Research Vet. Sc., v. 25 (3), 376-385
Haemonchus contortus, effect of changes in genetic constitution associated with development of benzimidazole resistance on physiological characteristics of parasitic and free-living stages (infectivity, pathogenicity, exsheathment response, etc.)

Benzimidazoles

- Simpkin, K. G.; and Coles, G. C., 1978, Research Vet. Sc., v. 25 (2), 249-250
Haemonchus and *Trichostrongylus colubriformis* in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parabendazole, oxbendazole)

Benznidazole -- Benzonidazole; N-Benzyl-2-nitroimidazoleacetamide; N-Benzyl-2-nitro-1-imidazoleacetate; Radanil; Ro 7-1051.

N-Benzyl-2-nitro-imidazoleacetamide (Ro 7-1051)

- Andrade, S. G.; and Figueira, R. M., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (5), 335-341
Trypanosoma cruzi-infected mice, therapeutic action of Ro 7-1051 against 2 parasite strains

Benznidazole

- Barclay, C. A., 1977, Rev. Neurol. Argentina, v. 3 (3), 477-482
[Trypanosoma] cruzi, humans with acute and chronic infections, therapy trials with benznidazole

Benznidazole -- Continued.

N-Benzyl-2-nitro-1-imidazoleacetamide (Ro 7-1051)

Brener, Z.; Costa, C. A. G.; and Chiari, C., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (6), 450-455

Trypanosoma cruzi, mice treated with nitrofurazone, nifurtimox, or Ro 7-1051, differences in susceptibility of 4 parasite strains to active drugs attributed to biological characteristics of strains rather than to mode of drug action

Benzonidazole (Ro 7-1051)

Ferreira, H. de O., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (5), 357-364

Trypanosoma cruzi, humans, benzonidazole, therapeutic trials

Ro 7-1051

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Ro 7-1051

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 339-347

Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

N-Benzyl-2-nitro-1-imidazolacetamide (Ro 7-1051)

Levi, G. C.; Amato Neto, V.; and Sant'anna, I. F. de A. B., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (1), 49-54

Trypanosoma cruzi, humans, Ro 7-1051 therapy, side effects

Benznidazole (Ro 7-1051; Radanil)

Polak, A.; and Richle, R., 1978, Ann. Trop. Med. and Parasitol., v. 72 (1), 45-54

Trypanosoma cruzi in vivo and in vitro, benznidazole, effect on growth and viability, aerobic and anaerobic respiration, and synthesis of protein, RNA, and DNA

N-Benzyl-2-nitro-1-imidazoleacetate (R07-1051)

Prata, A., 1978, AMB, Rev. Ass. Med. Brasil., v. 24 (4), 140-142

Chagas disease, humans, nifurtimox, R07-1051, comparative study

Benznidazole (Radanil)

Raaflaub, J.; and Ziegler, W. H., 1979, Arzneimittel-Forsch., v. 29 (1), 1611-1614

benznidazole, human, single-dose pharmacokinetics

(5-Benzoyl-1H-benzimidazole-2-yl)-carbamic acid methyl ester. See Mebendazole.

Benzoyl metronidazole -- Flagyl suspension; 9.712RP.

Benzoylmetronidazole (Flagyl suspension)

Roos, R. F., 1978, South African Med. J., v. 54 (21), 869-870

Trichomonas vaginalis, pregnant women with vaginal infections, clinical trials, single dose therapy with benzoylmetronidazole: Durban, South Africa

Benzoyl-metronidazole (9.712RP)

Sankale, M.; Coly, D.; and Niang, I., 1974, Therapie, v. 29 (3), 411-415

amoebiasis, children, oral suspension of benzoyl-metronidazole particularly useful therapy

Benzyl benzoate -- Ascabiol.

Benzyl benzoate lotion

Azulay, R. D.; et al., 1975, Rev. Brasil. Med., v. 32 (12), 800-802

scabies, human, clinical trials, efficacy of thiabendazole vs. benzyl benzoate as topical lotions

Benzyl benzoate

Belda, W., 1975, Rev. Brasil. Clin. e Terap., v. 4 (8), 309-312

human scabies, increasing incidence, successful management with monosulfiram: Brazil

Ascabiol (Benzyl benzoate emulsion)

Chakrabarti, A.; and Misra, S. K., 1979, Indian Vet. J., v. 56 (6), 497-500

Demodex canis, dogs, incidence in relation to season, host age, sex, and breed, clinical manifestations, in vivo and in vitro activity of several acaricides: India

Benzyl benzoate

Rajan, U., 1975, Singapore Med. J., v. 16 (4), 297-300

head lice, school children, pyrethrum the drug of choice in comparison clinical trials with benzyl benzoate

Benzylfluorenyl quaternary ammonium bromides

Dimmock, J. R.; Smith, P. J.; and Tsui, S. K., 1979, J. Pharm. Sc., v. 68 (7), 866-871

antimicrobial and antineoplastic activity of benzylfluorenyl and 1-arylethyl quaternary ammonium salts, synthesis and evaluation

3-Benzylideme amino-4-phenylthiazoline-2-thione

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

N-Benzyl-2-nitro-imidazoleacetamide. See Benznidazole.

Benzonidazole. See Benznidazole.

Benzophosphate. See Phosalone.

Benzo[b]thien-4-yl methylcarbamate. See Mobam.

N-Benzyl-2-nitro-1-imidazoleacetate. See Benzimidazole.

Benzyltriphenylphosphonium salts

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *J. Med. Chem.*, v. 22 (4), 452-455

Trypanosoma rhodesiense, mice, activity of benzyltriphenylphosphonium salts, toxicity at higher dose levels

Bephenium -- Alcopar; Bephenium hydroxynaphthoate; Naphthamon.

Bephenium hydroxynaphthoate

Botero R., D., 1978, *Ann. Rev. Pharmacol. and Toxicol.*, v. 18, 1-15

antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Bephenium hydroxynaphthoate (Alcopar)

Gatti, F.; et al., 1972, *Ann. Soc. Belge Med. Trop.*, v. 52 (1), 19-31

intestinal nematodes, African schoolchildren, trimestrial administration of levamisole compared with other anthelmintics: Kimwenza, Republique Democratique du Congo

Bephenium hydroxynaphthoate

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, *Rev. Brasil. Med.*, v. 31 (11), 791-794

human soil transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Naphthamon

Kadyrov, N. T., 1978, *Veterinariia, Moskva* (7), 57-58

Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Bephenium hydroxynaphthoate

Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (1), 55-58

Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (*Mesocricetus auratus*)

Bephenium hydroxynaphthoate (Alcopar)

Sarup, S.; Banerjee, D. P.; and Gautam, O. P., 1978, *Indian Vet. J.*, v. 55 (6), 497-498

Ancylostoma caninum, ova and infective larvae, in vitro evaluation of fenbendazole, helatac, alcopar, banminth II, only banminth II effective; banminth II-treated larvae administered orally to mice, none recovered from lungs or liver

Naphthamon

Shedivtsova, A., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Bephenium -- Continued.

Bephenium hydroxynaphthoate (Alcopar)

Zaman, V.; and Loh, Y. P., 1974, *Singapore Med. J.*, v. 15 (2), 147-148

Necator americanus, humans, clinical trials testing efficacy of pyrantel embonate, levotetramisole, and bephenium hydroxynaphthoate: Singapore

Bephenium hydroxynaphthoate. See Bephenium.

Bercotox. See Dioxathion.

Berenil -- Azidine; Berenil diacetate; Di-

acetate of 4,4-diazoaminodibenzamidine; 4,4'-Diamidino-diazoaminobenzene diacetate; 4-4'-Diamidino-diazo-amino benzol diacetate; Diminazene; Diminazene acetate; Diminazene salt; Diminazene suraminates; Diminazine acetate; Ganaseg.

Berenil (Diminazine acetate)

Aiyedun, B. A., 1977, *Bull. Animal Health and Prod. Africa*, v. 25 (3), 245-250

Trypanosoma gambiense, *T. brucei*, rats, induction of high level of immunity by administration of Freund's complete adjuvant a week before infection followed by cure with berenil

Berenil (Diminazene)

Bauer, F.; Raether, W.; and Seeger, K., 1978, *Cahiers Bleus Vet.* (27), 265-271

protozoal disease in exper. hosts, enhanced effect of berenil + reverin vs. berenil alone

Diminazene acetate

Bourn, D.; and Scott, M., 1978, *Trop. Animal Health and Prod.*, v. 10 (4), 191-203

trypanosomiasis, successful use of Zebu work oxen in agricultural development of tsetse infested land, environmental conditions, epizootiology of trypanosomiasis in oxen and in *Glossina morsitans*, strategic drug use (alternation of diminazene acetate and isometamidium to control trypanosomes; radoxanide to control helminths): Wollega province, western Ethiopia

Azidine

Dubovyi, S. Z.; et al., 1977, *Veterinariia, Moskva* (3), 71-72

babesiosis, piroplasmosis, cattle, dimidine as effective chemoprophylaxis under pasture conditions with presence of vector, *Boophilus calcaratus*, comparison with azidine

Berenil

Dwivedi, S. K.; Mallick, K. P.; and Malhotra, M. N., 1979, *Indian Vet. J.*, v. 56 (4), 333-335

Babesia bovis, Indian water buffaloes, 2 clinical case reports, treatment with berenil effective: Rithora, India

Berenil (Diminazine acetate)

Galhotra, A. P.; et al., 1979, *Haryana Agric. Univ. J. Research*, v. 9 (1), 69-74

Trypanosoma evansi, dogs (exper.), clinico-pathological changes, therapeutic trials with berenil and CL-64,855

Berenil -- Continued.

Diminazene

Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested

Diminazene suraminat

Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested

Berenil + Chlorpromazine

Gretillat, S., 1978, Bull. Acad. Vet. France, n.s., v. 51 (3), 351-358
Haemobartonella sp., horses, symptoms, diagnosis, pathology, treatment with chlorpromazine + berenil: Niger

Berenil diacetate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Diminazene

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Diminazene acetate (Berenil granules)

James, D. M., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 471-476
T[rypanosoma] congolense, T. brucei, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs

Berenil

Jennings, F. W.; et al., 1979, Internat. J. Parasitol., v. 9 (4), 381-384
Trypanosoma brucei, brain as source of relapsing infection in mice after berenil chemotherapy

Berenil

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Diminazene acetate

Malhotra, D. V.; Gautam, O. P.; and Banerjee, D. P., 1979, Indian J. Animal Sc., v. 49 (1), 75-77
Babesia equi, donkeys (exper), diminazene acetate as effective as diminazene acetate + rolitetracycline, quinuronium sulphate ineffective

Berenil -- Continued.

Diminazene acetate + Rolitetracycline

Malhotra, D. V.; Gautam, O. P.; and Banerjee, D. P., 1979, Indian J. Animal Sc., v. 49 (1), 75-77

Babesia equi, donkeys (exper), diminazene acetate as effective as diminazene acetate + rolitetracycline, quinuronium sulphate ineffective

Berenil

Meshkov, S.; Iotov, I.; and Sirachev, D., 1977, Vet. Sbirka, v. 75 (5), 32-33
piroplasmosis, berenil, cattle, control of Boophilus calcaratus with chlorophos

Berenil

Nathan, H. C.; et al., 1979, J. Protozool., v. 26 (4), 657-660
Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Azidine

Nikol'skii, S. N.; Nikiforenko, V. I.; and Pozov, S. A., 1977, Veterinariia, Moskva (4), 71-75

Piroplasma jakimovi, cattle, morphological and biological comparison with P. bigeminum, epizootiology (Ixodes ricinus as main vector; frequent association with leptospirosis), treatment: Siberia

Azidine

Osmanov, A. R., 1976, Dokl. Vsesoiuz. Akad. Sel'skokhoz. Nauk (12), 24-26
azidine, pharmacodynamics, milk cows, calves, chickens, white mice, negative effects on host require supplementary vitamins, minerals, and methionine

Berenil

Pandey, N. N.; and Mishra, S. S., 1978, Indian Vet. J., v. 55 (2), 144-148
Babesia bigemina, indigenous cow calves, berenil and acriflavin effective, sulfadimethoxine ineffective: India

Berenil

Razzaque, A.; and Mishra, S. S., 1977, Bull. Animal Health and Prod. Africa, v. 25 (4), 409-414
Trypanosoma evansi, buffalo calves (exper.), berenil, acriflavin, and antrycide prosalt, berenil most effective

Berenil (Diminazene acetate)

Roelants, G. E.; et al., 1979, Clin. and Exper. Immunol., v. 37 (3), 457-469
Trypanosoma congolense-infected mice, numbers of parasites in peripheral blood, changes in spleen cell populations, immune depression, suppressor cell activity, changes after berenil treatment

Diminazene acetate

Ruebush, T. K. II; et al., 1979, Am. J. Trop. Med. and Hyg., v. 28 (2), 184-189
Babesia microti, 65-year-old man, treated with diminazene acetate after failure to respond to chloroquine therapy, development of acute idiopathic polyneuritis: Nantucket Island

Berenil -- Continued.

- Berenil**
Ruppel, J. F.; and Burke, J., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 481-495
[Trypanosoma] gambiense, humans, clinical trials with levofuraltadone and levofuraltadone hydrochloride, comparisons with berenil, follow-up reports: Kimpangu, Republique du Zaire
- Diminazene aceturate (Berenil)**
Rurangirwa, F. R.; et al., 1979, Infect. and Immun., v. 26 (3), 822-826
Trypanosoma congolense- or T. vivax-infected Bos indicus, suppression of antibody response to Leptospira biflexa and Brucella abortus and recovery from immunosuppression after berenil treatment
- Berenil**
Saha, A. C.; and Das, S. N., 1978, Indian J. Animal Health, v. 17 (1), 89-90
Babesia [sp.], one-month-old cross-bred Jersey calf (blood), case report, recovery after treatment with berenil
- Diminazene aceturate (Berenil)**
Scott, J. M.; et al., 1978, Research Vet. Sc., v. 25 (1), 115-117
Trypanosoma congolense, zebu cattle, attempted protection using multi-stabilate vaccine given either as live or dead organisms followed by trypanocidal therapy, disappointing results
- Berenil**
Sharma, R. D.; and Gautam, O. P., 1977, Haryana Vet., v. 16 (1), 19-26
Babesia caballi, B. equi, horses, case reports, haematological changes, berenil, some adverse reactions
- Berenil**
Sinha, M.; Goswami, D. N.; and Das Gupta, N. N., 1978, Indian J. Biochem. & Biophys., v. 15 (3), 162-165
daunomycin, berenil, dielectric studies on interaction with DNA
- Ganaseg**
Thompson, K. C.; et al., 1978, Trop. Animal Health and Prod., v. 10 (2), 75-81
Anaplasma marginale, Babesia argentina, B. bigemina, cattle under tropical conditions, immunization with virulent organisms followed by drug therapy (ganaseg; gloxazone; emicina) vs. chemoprophylaxis (imidocarb); tick and gastrointestinal parasite control without haemoparasitic control had advantage over no control system at all
- Berenil**
Vulchovski, Ia., 1977, Vet. Med. Nauki, v. 14 (3), 79-87
Babesia, cattle, efficacy of trypanblue, trypaflavin, acaprin, berenil, and pyrodis (berenil and pyrodis superior to other drugs)
- Berenil**
Wellde, B. T.; et al., 1978, Exper. Parasitol., v. 45 (1), 26-33
Trypanosoma congolense, cattle (exper.), thrombocytopenia, effects of parasite concentration, curative berenil therapy, and immune status on thrombocyte levels; coagulation abnormalities

Berenil -- Continued.

- Diminazene**
Whitelaw, D. D.; et al., 1979, Research Vet. Sc., v. 26 (1), 102-107
Trypanosoma spp., cattle, mice, suppressed antibody response to louping-ill vaccine, value of diminazene therapy in alleviating this effect
- Berenil**
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Diminazene salt (Berenil)**
Worms, M. J.; and Hawking, F., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 548-549
Litomosoides carinii-infected Sigmodon hispidus, effects of suramin, homidium bromide, quinapyramine, diminazene, and isometamidium after observation period of more than 5 weeks
-
- Berenil diacetate.** See Berenil.
- Bergapten**
Abdulla, W. A.; et al., 1977, Egypt. J. Bilharz., v. 4 (1), 19-26
S[chistosoma] mansoni, anti-schistosomal activity of the plant Ammi majus and bergapten (furocoumarin present in A. majus), compared with tarter emetic, screening in mice
- Beryllium sulfate**
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Betamethasone**
Andrade, S. G.; and Macedo, V., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (6), 421-430
Trypanosoma cruzi, mice, human, Bayer 2502, combined use of Bayer 2502 and corticoid (betamethasone) more effective than drug used alone
- Betamethazone**
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity
- BHC.** See Benzene hexachloride.

- γ -BHC. See Benzene hexachloride.
- BHS. See Bithionol.
- Bifuran. See Furazolidone or Nitrofurazone.
- Bigumal. See Chlorguanide.
- Biklotimol
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results
- Bilarcil. See Trichlorfon.
- Bilevon-Injektion. See Hexachlorophene.
- Bilevon-M. See Niclofolan.
- Bilevon-R. See Niclofolan.
- Bilharcid. See Piperazine diantimonyl tartrate.
- Bilharzid. See Piperazine diantimonyl tartrate.
- Bilirubin
Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, Biochem. Pharmacol., v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism
- Biomitsin. See Chlortetracycline.
- Biomycin. See Chlortetracycline.
- Biovetin. See Chlortetracycline.
- Birlane. See Chlorfenvinphos.
- 2,2-Bis(p-acetyl aminophenoxy) diethyl ether.
See Diamphenethide.
- 1,6-Bis(6-amino-2-methyl-4-quinolylamino) hexane dihydrochloride
Kinnamon, K. E.; and Rane, D. S., 1978, Internat. J. Parasitol., v. 8 (6), 515-523
Trypanosoma rhodesiense, mice, greater than 1 year protection from lethal infections by prophylactic drugs and active immunity
- 1,7-Bis(p-aminophenoxy)heptane -- 153C51; Wellcome 153C51.
1:7-Bis-(p-aminophenoxy) heptane (Wellcome 153 C 51)
Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, Rev. Brasil. Pesqui. Med. e Biol., v. 2 (1), 45-49
Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs
- Wellcome 153C51
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared
- 1,7-Bis(p-aminophenoxy)heptane (153C51)
Watts, S. D. M., 1978, Comp. Biochem. and Physiol., v. 60C (2), 109-114
Schistosoma mansoni, 153C51-treated parasites, unlikely that interference with glucose uptake observed in vitro in male parasites could be primary cause of schistosome death or even of physiological distress in vivo
- 1,7-Bis (p-aminophenoxy)heptane (153C51)
Watts, S. D. M.; and Atkins, A. M., 1979, Biochem. Pharmacol., v. 28 (17), 2579-2584
effects of schistosomicide 1,7-bis(p-aminophenoxy)heptane (153C51) on lysosomes and membrane stability
- 1,7-Bis(p-aminophenoxy)heptane (153C51)
Watts, S. D. M.; Orpin, A.; and MacCormick, C., 1979, Parasitology, v. 78 (3), 287-294
Schistosoma mansoni, tegument pathology following chemotherapy with 153C51, lysosomal involvement (accumulation of inclusions with characteristics of residual lysosomes, changes in localization of acid phosphatase), immunological factors probably not involved
-
- 3,6-Bis-(t-butyl)-8-(4-chlorophenyl)-2H,4H-1,3-benzooxazine (WR-204,165)
Schmidt, L. H.; and Crosby, R., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (5), 672-679
Plasmodium falciparum, P. vivax, Aotus trivirgatus griseimembra, antimalarial activities of WR-194,965 and WR-204,165
- Bis-(β -carbohydrazide-ethyl) sulfone -- Hoechst S-201.
Hoechst S-201
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared
-
- 1,3-Bis (p-chlorobenzylideneamino guanidine hydrochloride. See Robenidine.

- Bischloroethyldeuteroporphyrin IX
Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis
- 1,3-Bis(2-chloroethyl)-1-nitrosourea
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Bis[4-(4-chlorophenyl)methylene]-carbonimidic dihydrazide. See Robenidide.
- N,N-Bis (dichloroacetyl)-N,N-bis(2-ethoxy-ethyl)-1,4-bis (aminomethyl) benzene. See Teclozan.
- Bis (dimethyl-dithio-carbamoyl) ethylene diamine
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- N,N'-Bis (3,4 ditrifluoromethylphenyl) methylmalonamide (Sch 18545)
Panitz, E., 1979, Parasitology, v. 78 (1), 33-40
Eimeria spp., chicks, anticoccidial efficacy and cross-resistance patterns of N,N'-bis (3,4 ditrifluoromethylphenyl) methylmalonamide compound have no practical application because of weight gain depression
- 1,2-Bis(3-ethoxycarbonyl-2-thioureido) benzene
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Bis(2-hydroxy-3,5-dichlorophenyl) sulfoxide.
See Bithionol.
- Bis(2-hydroxy-3,5-dichlorophenyl) sulfoxide.
See Bithionol.
- 3,3'-Bis-(2-imidazolin-2-yl) carbanilide dihydrochloride. See Imidocarb.
- 3,3'-Bis(2-imidazolin-2-yl)carbanilide dipropionate. See Imidocarb.
- N-[2-[2,3-Bis (methoxycarbonyl) guanidino]-5-(phenylthio)-phenyl] 2-methoxy-acet-amid. See Febantel.
- 1,2-Bis(3-methoxycarbonyl-2-thioureido) benzene
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Bis-phenyl-(2-chlorophenyl)-1-imidazolyl-methane.
See Clotrimazole.
- 1,4-Bis(trichloromethyl) benzene -- 1,4-Bis-trichloromethylbenzol; Chloxy1; Hetol; Hexachloroxy1; Hexachlorparaxy1.
- Hetol
Araki, K., 1975, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 24 (4), 203-219
Paragonimus ohirai, rats (exper.), effect of bithionol and hetol against various developmental stages, parasitological, pathological, and immunological evaluation
- Hetol
Daniiarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective
- Chloxy1
Kim, N. Kh., 1976, Terap. Arkh., v. 48 (6), 130-132
opisthorchiasis, patients treated with chloxy1, changes in aldolase, transaminases, and blood coagulants during therapy
- Hexachloroxy1
Reinhardt, P., 1978, Monatsh. Vet.-Med., v. 33 (23), 898-901
Dicrocoelium dendriticum, sheep, efficacy of various anthelmintics compared
- Hexachlorparaxy1
Zharikov, I. S.; Orlovskii, V. I.; and Kamin-skii, I. I., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 74-77
paramphistomiasis, bovine, hexachlorparaxy1 and fenasal, ineffective in chronic disease
-
- 1,4-Bis-trichloromethylbenzol. See 1,4-Bis(trichloromethyl) benzene.
- DL-2,8-Bis(trifluoromethyl)-4-[1-hydroxy-3-(N-t-butylamino)propyl]-quinoline phosphate (WR 184,806.H₂PO₄)
Stampfli, H.; et al., 1979, J. Liquid Chromatography, v. 2 (1), 53-65
multi-component solvent system for analysis of candidate antimalarial (WR 180,409.H₂PO₄) and its internal standard (WR 184,806.H₂PO₄) by normal phase high pressure liquid chromatography
- 3,6-Bis-(trifluoromethyl)-α-2-piperidyl-9-phenanthrenemethanol hydrochloride. See α-(2-Piperidyl)-3,6-bis (trifluoromethyl)9-phenanthrene methanol.
- 2,8-Bis-(trifluoromethyl)-α-(2-piperidyl)-4-quinolinemethanol methylsulfonate monohydrate. See Mefloquine.
- Bithionol -- Actamer; BHS; Bis(2-hydroxy-3,5-dichlorophenyl) sulfoxide; Bis(2-hydroxy-3,5-dichlorophenyl) sulfoxide; Bithionol sulfoxide; Bitin-S; Bithirazine (with Piperazine); 2,2'-Dioxy-3,3',5,5'-tetrachlorodiphenyl sulfoxide; Sulfene; 2,2'-Sulfinylbis (4,6-dichlorophenyl); Sulphene; 2,2'-Thiobis (4,6-dichlorophenyl).

Bithionol -- Continued.

Bithionol

- Araki, K., 1975, *Kiseichugaku Zasshi* (Japan. J. Parasitol.), v. 24 (4), 203-219
Paragonimus ohirai, rats (exper.), effect of bithionol and hetol against various developmental stages, parasitological, pathological, and immunological evaluation

Bithionol

- Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10), 28-36
 cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Bithionol + Mepacrine

- Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10), 28-36
 cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Sulphene

- Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10), 28-36
 cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Bithionol

- Bogdanov, V. R.; et al., 1977, *Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (29), 90-91
 cestodes, pathomorphology resulting from action of various anthelmintics

Sulphene

- Bogdanov, V. R.; et al., 1977, *Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.* (29), 90-91
 cestodes, pathomorphology resulting from action of various anthelmintics

BHS

- Chroustova, E.; Willomitzer, J.; and Strouhal, Z., 1977, *Veterinarstvi*, v. 27 (10), 460-462
Fasciola hepatica, wild ruminants, prevalence, BHS treatment, Frescon for snail control

Bithionol (Actamer)

- Chrusciel, T. L.; et al., 1970, *Acta Parasitol. Polon.*, v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances

Bithionol

- Daniiarov, I. A.; et al., 1978, *Veterinariia, Moskva* (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Bitin-S

- Demetriou, A.; Phillips, B. M.; and Hendrickse, R. G., 1978, *Arch. Dis. Childhood*, v. 53 (11), 912-913
Paragonimus diagnosed in child presenting with soft tissue swellings on basis of clinical, radiological, serological, and epidemiological evidence despite failure to demonstrate ova in sputum, treatment with bitin-S followed by complete cure: Britain (had previously lived in eastern Nigeria)

Bithionol -- Continued.

Bithionol

- Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (11), 675-679
Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Bithionol sulphoxide

- Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (11), 675-679
Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Bithirazine

- Golovneva, L. F.; et al., 1970, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 8, 105-106
 ascariasis, chickens, bithirazine tested under production conditions, more study needed

Bithionol

- Grados B., O.; and Berrocal S., L. A., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (6), 425-427
Fasciola hepatica, humans, bithionol: Peru

Bithionol

- Hamajima, F.; et al., 1979, *Internat. J. Parasitol.*, v. 9 (3), 241-249
Clonorchis sinensis, *Metagonimus takahashii*, *Paragonimus miyazakii*, in vitro effects of bithionol and menichlopholan on motility, metabolism, and fine structure

Bithionol

- Klenov, A. P., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 152-155
 [Bothriocephalus], white amur, effectiveness of various anthelmintics

BHS

- Kolar, Z., 1978, *Veterinarstvi*, v. 28 (6), 276-277
Fascioloides magna, red deer, treatment with BHS

Bithionol

- Ono, S., 1977, *Shonika Shonryo (J. Pediat. Pract.)*, v. 40 (5), 47-50
Dipylidium caninum, human, case report, bithionol

Bithionol

- Oshima, T., 1976, *Shinshu Igaku Zasshi (Shinshu Med. J.)*, v. 24 (3), 191-204
diphyllobothriasis, humans, epidemiology, biology, bithionol therapy; recent outbreak in Nagano Prefecture

Bithionol

- Sakamoto, T., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

Bithionol sulfoxide-Diethylcarbamazine

- Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Bithionol -- Continued.

- Bithionol sulfoxide + 8-Hydroxy-quinoline
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Sulfinylbis(4,6-dichlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis(4,6-dichlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- Sulfene
Vishniauskas, A.; and Rudaitis, A., 1978,
Veterinariia, Moskva (4), 68-69
Fasciola hepatica, cattle (exper.), acemidophene, dertil and sulfene compared, acemidophene insufficiently effective
- BHS
Willomitzer, J.; Chroustova, E.; and Kolar, Z.,
1976, Veterinarstvi, v. 26 (9), 417-419
Fascioloides magna, deer, control with BHS for deer and Frescon as molluscicide for snail control

Bithionol sulfoxide. See Bithionol.

Bithirazine. See Bithionol or Piperazine.

Bitin-S. See Bithionol.

Bitoscanate -- Compound 16,842; Hoechst 16.842;
Jonit; p-Phenylene-bis(isothiocyanate);
Phenylene-diisothiocyanate (1,4); 1,4-Phenylene-diisothiocyanate; Phenylene-isothiocyanate 1,4.

Bitoscanate (Compound 16,842; Jonit)
Biagi F., F.; Zavala, J.; and Malagon, F., 1969,
Rev. Inst. Med. Trop. S. Paulo, v. 11 (6), 444-448
Trichuris trichiura, uncinariasis, humans, efficacy of bitoscanate, side effects

Bitoscanate
Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Phenylenediisothiocyanate (1,4)
Camillo-Coura, L., 1974, Rev. Soc. Brasil. Med. Trop., v. 8 (6), 340-356
mass therapy trials using piperazine for ascariasis and phenylenediisothiocyanate for ancylostomiasis; emphasis also on need for social and economic improvements and proper sanitation to achieve some measure of control: States of Rio de Janeiro and Guanabara, Brazil

Bitoscanate -- Continued.

- Jonit (Phenylene diisothiocyanate 1,4)
Chamorro, H.; and Okonsky, L., 1973, Rev. Brasil. Med., v. 30 (6), 399-402
ancylostomiasis, human, evaluation of jonit as therapy for hookworm carriers: residents of Apipe Islands on upper Parana River
- 1,4-Phenylene-diisothiocyanate (Jonit)
Freire, J. dos S.; Bezerra, V. C.; and Prado, A. A., 1973, Rev. Patol. Trop., v. 2 (2), 155-158
human ancylostomiasis, comparative therapeutic trials using jonit and tetrachloroethylene; tetrachloroethylene gave higher cure rates: Brazil
- Jonit (Hoechst 16.842)
Hartung, M.; Doria, M.; and Paredes, E., 1973, Rev. Brasil. Clin. e Terap., v. 2 (10), 505-508
human ancylostomiasis, clinical trials with jonit, evaluation of tolerance and efficacy, some side effects, recommended for therapy
- Phenylene diisothiocyanate (Jonit)
Hasbun, E. R.; and Zepeda, J. R., 1977, Rev. Inst. Invest. Med., San Salvador, v. 6 (2), 190-192
Ancylostomidae, humans, clinical trials testing efficacy of pyrantel pamoate and phenylene diisothiocyanate, medications well tolerated with few side effects: El Salvador
- Jonit (Phenylene-diisothiocyanate (1,4))
Hsieh, H. C.; et al., 1970, Taiwan i Hsueh Hui Tsa Chih (J. Formosan Med. Ass.), v. 69 (8), 405-409
A[ncylostoma] duodenale, N[ecator] americanus, human, single and mixed infections, therapeutic efficacy of jonit: Taiwan
- Jonit (Phenylene-di-iso-thiocyanate-1,4)
Louzada, G. Z.; de Carli, G. A.; and Louzada, J. L. Z., 1972, Rev. Brasil. Clin. e Terap., v. 1 (5), 337-340
Ancylostoma duodenale, humans, clinical trials with jonit
- Phenylene-diisothiocyanate-(1,4) (Jonit)
Nohmi, N.; and Lobato, M. B. P., 1972, Rev. Ass. Med. Minas Gerais, v. 23 (3), 113-118
Ascaris lumbricoides, Trichocephalus trichiura, humans, clinical trials testing the efficacy of jonit
- Bitoscanate
Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (1), 55-58
Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (Mesocricetus auratus)
- Jonit
Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies
- Bitoscanate (Jonit)
Tonelli, E.; et al., 1975, Rev. AMMG, v. 26 (3-4), 118-120
ancylostomiasis, children younger than 5 years of age, therapy trials with jonit, drug well tolerated but showed low efficacy: Belo Horizonte, Brazil

Bitoscanate -- Continued.

Phenylene-isothiocyanate 1,4
 Vasconcelos, W. M. de P.; and Maia, M. A.,
 1972, Rev. Patol. Trop., v. 1 (3), 405-408
 ancylostomiasis, phenylene-isothiocyanate 1,4
 therapy, poor results when treating patients
 with associated pemphigus who are also re-
 ceiving steroids, probably both the dermatosis
 and the steroids are influencing factors

Bitricide. See Praziquantel.

Blasticidin-S-benzylaminobenzen
 Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
 Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
 130
 Echinococcus granulosus, scolical effect
 of 65 antibiotic, antineoplastic, cytostatic,
 and other agents in vitro

Blasticidin-S-methylol
 Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
 Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
 130
 Echinococcus granulosus, scolical effect
 of 65 antibiotic, antineoplastic, cytostatic,
 and other agents in vitro

Blasticidin-S-styreneoxide
 Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
 Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
 130
 Echinococcus granulosus, scolical effect
 of 65 antibiotic, antineoplastic, cytostatic,
 and other agents in vitro

Blattanex. See Propoxur.

Bleomycin hydrochloride
 Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
 Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
 130
 Echinococcus granulosus, scolical effect
 of 65 antibiotic, antineoplastic, cytostatic,
 and other agents in vitro

Blotic. See Propetamphos.

Blue dextran (2000)
 Williamson, J.; and Scott-Finnigan, T. J.,
 1978, Antimicrob. Agents and Chemotherapy,
 v. 13 (5), 735-744
 trypanocidal activity of antitumor anti-
 biotics and other metabolic inhibitors,
 microtest for rapid preliminary assay in
 vitro, parasite motility and infectivity
 for mice are indexes respectively of respi-
 ration and glycolysis and of cell division,
 implications of results for combination
 chemotherapy and deposit prophylaxis (with
 polyanions)

Blue vitriol. See Cupric sulfate.

Bolfo. See Propoxur.

Bonlam. See Cambendazole.

Bovicam. See Cambendazole.

Bovi-douvex. See Niclosulide.

Bovinox
 Camoens, J. K., 1977, Malaysian Vet. J., v. 6
 (3), 111-124
 Boophilus microplus, dairy cattle, various
 control measures discussed but spraying
 acaricides on pastures shows particular
 promise: Air Hitam, Johor, Malaysia

BPRL-5344-16
 Gladney, W. J.; Price, M. A.; and Graham, O.
 H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
 Amblyomma maculatum, cattle, efficacy of
 various insecticides applied as sprays, ear
 smears and dusts, or in slow-release devices,
 field tests

Brilliant green
 Daniyarov, I. A.; et al., 1978, Veterinariia,
 Moskva (2), 64-65
 Echinococcus spp., sheep, 28 anthelmintics
 and dyes tested, none effective

Brobenzoxaldine -- Intestopan (with Broxyquino-
 line); Metrointestopan (with Broxyquinoline
 and Metronidazole).

Intestopan + Metronidazole (= Metrointestopan)
 Ramirez Andrade, R., 1977, Semana Med. Mexico
 (1170), an. 24, v. 93 (2), 41-44
 amoebiasis, human chronic colitis or recto-
 colitis, clinical trials testing efficacy of
 metronidazole combined with intestopan, good
 results, drug well tolerated

5-Bromomethyl 1-1,2,3,4,4-7,7-hexachlorbicyclo-
 (2,2,1)hepten-(2). See Alugan.

5-Bromo-4[bis-(2-diethylamino ethyl)-amino] vera-
 trole (RC12; WR 27,653)
 Rane, D. S.; and Kinnamon, K. E., 1979, Am. J.
 Trop. Med. and Hyg., v. 28 (6), 937-947
 sporozoite-induced Plasmodium berghei in
 mice, development of high volume tissue
 schizonticidal drug screen based upon mor-
 tality of infected mice

7-Bromo-6-chlorofebrifugine-hydrobromide. See
 Halofuginone.

d1-7-Bromo-6-chloro-3-[3-(3-hydroxy-2-piperidyl)-
 acetonyl]-4(3H)-quinazolinone hydrobromide.
See Halofuginone.

Bromochlorophen -- 2,2'-Methylenebis(4-chloro-6-
 bromophenol).

2,2'-Methylenebis(4-chloro-6-bromophenol)
 Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
 J. Vet. Research, v. 23 (3), 81-94
 Echinococcus granulosus in vitro, scolical
 effect of salicylanilide and bisphenol deri-
 vatives

Bromocyclen. See Alugan.

Bromodeoxyuridine

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

4-Bromo-2,5-dichlorophenyl diethyl phosphorothionate. See Bromophos ethyl.

6-Bromo- α -diheptylaminoethyl-9-phenanthrenemethanol -- SN-13,465; WR-33,063.

6-Bromo- α -diheptylaminoethyl-9-phenanthrene methanol

Carroll, F. I.; Berrang, B.; and Linn, C. P., 1978, J. Med. Chem., v. 21 (4), 326-330

Plasmodium berghei, P. cynomolgi, experimental animals, resolution of antimalarial agents via complex formation with α -(2,4,5,7-tetra-nitro-9-fluorenylideneaminoxy)propionic acid, significant differences in toxicity

WR-33,063

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-33,063 (SN-13,465)

Schmidt, L. H.; et al., 1978, Antimicrob.

Agents and Chemotherapy, v. 14 (3), 292-314

Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

3-Bromo-3,4-dihydro-2,2-dimethyl-2H-naphtho [1,2-b]pyran-5,6-dione. See 3-Bromo- β -lapachone.

4'-Bromo-2,6-dihydroxybenzanilide. See Resorantel.

3-Bromo-2,2'-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione. See 3-Bromo- β -xyloidone

3-Bromo-2,2'-dimethyl-2H-naphtho[2,3-b]pyran-5,10-dione. See 3-Bromo- α -xyloidone.

3-Bromo- β -lapachone -- 3-Bromo-3,4-dihydro-2,2-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione.

3-Bromo- β -lapachone

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

3-Bromo- β -lapachone -- Continued.

3-Bromo- β -lapachone

Docampo, R.; et al., 1978, Ztschr. Parasitenk., v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

Bromo- β -lapachone

Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

Bromophenophos -- Acedist; Bromphenophos; 2,2'-Dihydroxy-3,3',5,5'-tetrabromo-1,1'-biphenylmono (dihydrogen phosphate); 4,4',6,6'-Tetrabromo-2,2'-biphenyldiolmono(dihydrogenphosphate).

Acedist

Qadir, A. N. M. A., 1979, Indian Vet. J., v. 56 (5), 429-431

Fasciola gigantica, goats, acedist, comparison with bilevon: Bangladesh

4,4',6,6'-Tetrabromo-2,2'-biphenyldiolmono(dihydrogenphosphate)

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Acedist (Bromphenophos)

Stoianova-Radoeva, S., 1978, Vet. Med. Nauki, v. 15 (4), 79-85

acedist, dovenix, treated sheep, residues in milk, effect on blood biochemical indices

Bromophos -- o,o-Dimethyl-o-2,5-dichloro-4-bromophenyl-thio-phosphate.

Bromophos

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, B. microplus, laboratory tests of insecticides

o,o-Dimethyl-o-2,5-dichloro-4-bromophenyl-thio-phosphate (Bromophos)

Privora, M.; Rupes, V.; and Cerny, V., 1970, Folia Parasitol., v. 17 (1), 81-84

Dermacentor marginatus, laboratory trials testing six insecticides

Bromophos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144

Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Bromophos -- Continued.Bromophos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Bromophos-ethyl -- 4-Bromo-2,5-dichlorophenyl diethyl phosphorothionate; Nexagan.Bromophos-ethyl

Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439
 economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Bromophos-ethyl

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Nexagan (Bromophos ethyl)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105

Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Bromophos-ethyl (Nexagan)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142

Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Bromophos-ethyl

Schuntner, C. A.; and Thompson, P. G., 1978, Austral. J. Biol. Sc., v. 31 (4), 317-325
Boophilus microplus, mechanisms of resistance of 2 strains to bromophos-ethyl

6[2[(5-Bromo-2-pyridyl) amino] vinyl)-1-ethyl-2-picolinium iodide

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

3-Bromo- α -xyloidone -- 3-Bromo-2,2'-dimethyl-2H-naphtho[2,3-b]pyran-5,10-dione.3-Bromo- α -xyloidone

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

3-Bromo- β -xyloidone -- 3-Bromo-2,2'-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione.3-Bromo- β -xyloidone

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

Bromphenophos. See Bromphenophos.Bromsalans -- 3,5-Dibromosalicylanilide; Dibromsalan; Mintes-B; TBS; 3,5,4'-Tribromosalicylanilide; Tribromsalan; Tribromosalicylanilide; Trinoin.Trinoin

Merenyi, L., 1978, Magy. Allat. Lapja, v. 100, v. 33 (2), 132-134
 fascioliasis, control, treatment: Nograd county

3,5-Dibromosalicylanilide

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

3,5-Dibromosalicylanilide + 3,5,4'-Tribromosalicylanilide (= Mintes-B)

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

3,5,4'-Tribromosalicylanilide

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

3,5,4'-Tribromosalicylanilide + 3,5-Dibromosalicylanilide (= Mintes-B)

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

3,5-Dibromosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

3,5,4'-Tribromosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Dibromsalan

Vodrazka, J.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21) (4), 363-374
 tribromsalan, dynamic behavior in sheep blood plasma; antifasciolicides, effects on some sheep plasma enzymes

Bromsalans -- Continued.

- Tribromsalan (Tribromsalicylanide; TBS)
Vodrazka, J.; et al., 1979, Biol. a Chem.
Zivoc. Vyroby, Vet., v. 15 (v. 21) (4), 363-374
tribromsalan, dynamic behavior in sheep blood plasma; antifasciolicides, effects on some sheep plasma enzymes

Brot. See Tetramisole.

Brotianide -- 2-Acetoxy-3-bromo-5-chloro-N-(4-bromophenyl)-thiobenzamide; Dirian.

- Brotianid (Dirian)
Corba, J.; et al., 1978, Veterinarstvi, v. 28 (2), 87-88
Fasciola hepatica, heifers, brotianid, rafoxanide
- Brotianide (Dirian)
Corba, J.; Pacenovsky, J.; and Krupicer, I., 1976, Vet. Med. Rev. (2), 181-189
Fasciola hepatica, sheep; Paramphistomum spp., cattle, brotianide

Broxyquinoline -- Intestopan (with Brobenzoxaldine); Metrointestopan (with Brobenzoxaldine and Metronidazole); Starogyn.

- Broxyquinoline (Starogyn)
Lovgren, T.; and Salmela, I., 1978, Acta Path. et Microbiol. Scand., v. 86B (3), 155-158
Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents
- Intestopan + Metronidazole (= Metrointestopan)
Ramirez Andrade, R., 1977, Semana Med. Mexico (1170), an. 24, v. 93 (2), 41-44
amoebiasis, human chronic colitis or rectocolitis, clinical trials testing efficacy of metronidazole combined with intestopan, good results, drug well tolerated

Buban. See Bunamidine.

Bubulin. See Trichlorfon.

Bucarpolate -- Pyractone M429 (with Pyrethrins).

- Bucarpolate + Pyrethrins
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides
- Pyractone M429
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides

Bunamidine -- Bunamidine hydrochloride; Bunamidine hydroxynaphthoate; N,N-Di-n-butyl-4-hexyloxy-1-naphthamidine; N,N-Dibutyl-4-hexyloxy-1-naphthamidine hydrochloride; Scolaban; Buban.

- Bunamidine
Alaimo, R. J.; et al., 1978, J. Med. Chem., v. 21 (3), 298-300
furodazole, anthelmintic trials with experimental animals, bunamidine and niclosamide used as reference drugs
- Bunamidine hydroxynaphthoate
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results
- Bunamidine hydrochloride
Bankov, D. E., 1977, Vet.-Med. Nachr. (2), 145-148
Echinococcus granulosus, dogs (exper.), bunamidine hydrochloride, cantrodiphen, praziquantel, praziquantel most effective and showed no toxicity
- Bunamidine hydrochloride (Scolaban)
Boray, J. C.; et al., 1979, Austral. Vet. J., v. 55 (2), 45-53
nematodes and cestodes of dogs and cats, efficiency and safety of nitroscanate, comparison with mebendazole, bunamidine hydrochloride, and praziquantel
- Bunamidine HCl
Chatfield, R. C.; and Yeary, R. A., 1979, Vet. Parasitol., v. 5 (2-3), 177-193
Hymenolepis diminuta, bunamidine HCl, applicability of in vitro cultivation in determination of LC₅₀, effect on enzymes involved in energy metabolism and on ultrastructure
- Bunamidine hydrochloride
Gemmell, M. A., 1978, Bull. World Health Organ., v. 56 (3), 433-443
tapeworm control in dogs for prevention of hydatidosis and cysticercosis in sheep, monthly drug treatment program, age-specific prevalence of Taenia hydatigena in lambs used as principal indicator, 13-year assessment: Styx Valley and Maniototo County, South Island, New Zealand
- Bunamidine hydrochloride + Niclosamide
Gemmell, M. A., 1978, Bull. World Health Organ., v. 56 (3), 433-443
tapeworm control in dogs for prevention of hydatidosis and cysticercosis in sheep, monthly drug treatment program, age-specific prevalence of Taenia hydatigena in lambs used as principal indicator, 13-year assessment: Styx Valley and Maniototo County, South Island, New Zealand
- Bunamidine hydroxynaphthoate (BHN, Bunamidin)
Kassai, T.; et al., 1977, Magy Allat. Lapja, v. 99, v. 32 (6), 373-377
Echinococcus granulosus, dogs, bunamidine hydroxynaphthoate in biscuit form effective, lopatol containing cantrodifene unsatisfactory

Bunamidine -- Continued.

Bunamidine hydrochloride

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Bunamidine hydrochloride

Thakur, A. S.; Prezioso, U.; and Marchevisky, N., 1979, Exper. Parasitol., v. 47 (2), 131-133
Echinococcus granulosus, praziquantel did not possess absolute ovicidal activity against eggs either within or outside proglottids, bunamidine hydrochloride did not kill eggs within proglottids

Bunamidine hydrochloride (Scolaban)

Todd, K. S., jr.; Howland, T. P.; and Woerpel, R. W., 1978, Am. J. Vet. Research, v. 39 (2), 315-316
Mesocostoides corti, dogs (exper.), bunamidine hydrochloride and uredofos (good results), arecoline hydrobromide and niclosamide (variable results)

Bunamidine hydrochloride. See Bunamidine.

Bunamidine hydroxynaphthoate. See Bunamidine.

Buquinolate

Augustine, P. C.; Vetterling, J. M.; and Doran, D. J., 1977, Proc. Helminth. Soc. Washington, v. 44 (2), 147-149
Eimeria tenella, comparison of amprolium- and buquinolate-resistant strains to a drug-sensitive strain with respect to (1) oocyst production in chicks and (2) infectivity, rate of development, and oocyst production in primary chick kidney cell cultures

Buquinolate

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Busulfan

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Butamisol hydrochloride -- (-)-2-Methyl-N-[3-(2,3,5,6-tetrahydroimidazo-[2,1-b]thiazol-6-yl)phenyl] propanamide monohydrochloride.

Butamisol hydrochloride

Grieve, R. B.; Moore, B. G.; and Bradley, R. E., 1979, Am. J. Vet. Research, v. 40 (1), 139-141
gastrointestinal parasites, horses and ponies, critical test evaluation of butamisol, compared with efficacy of piperazine-thiabendazole

Butea frondosa -- Palasonin.

Palasonin

Lal, J.; Chandra, S.; and Sabir, M., 1978, Indian J. Pharm. Sc., v. 40 (3), 97-98
modified method for isolation of palasonin, the anthelmintic principle of Butea frondosa seeds, activity tested in vitro against Ascaris lumbricoides

Butea frondosa seeds

Sharma, N. D.; and Sisodia, C. S., 1976, Indian Vet. J., v. 53 (12), 920-922
Ascaridia galli, Butea frondosa seeds compared with piperazine adipate, in vivo (chicks) and in vitro trials

Butonate (Perdix PE 50)

Hiepe, T.; et al., 1978, Monatsh. Vet.-Med., v. 33 (23), 901-904
Caloglyphus berlesei, sheep, enzootic occurrence, pruritis, wool loss, and dermal lesions, controlled by immediate shearing, perdix PE 50, and moving entire herd to clean sheep pen

Butyl aminobenzoate -- PB Dressing (with Piperonyl butoxide and Propylene glycol).

PB Dressing

Nie, I. A.; and Pick, C. R., 1978, J. Inst. Animal Techn., v. 29 (1), 63-68
Otodectes cynotis, ferrets (ears), PB dressing: laboratory colony, Mill Hill, London

t-Butylaminoethanol

McManus, E. C.; et al., 1979, Exper. Parasitol., v. 47 (1), 13-23
Eimeria tenella, chickens (exper.), t-butylaminoethanol alone or in synergistic combination with sulfaquinoxaline and pyrimethamine, anticoccidial efficacy, specific reversal of toxicity for parasite and host by choline and dimethylaminoethanol

t-Butylaminoethanol

McManus, E. C.; and Rogers, E. F., 1979, Exper. Parasitol., v. 48 (2), 235-238
Eimeria tenella, chickens, synergistic interaction of sulfaquinoxaline and t-butylaminoethanol

α-(tert-Butylaminoethyl)-2,8-bis-(trifluoromethyl)-4-quinolinemethanol -- WR-184,806.

WR-184,806

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta

α-(tert-Butylaminoethyl)-2,8-bis-(trifluoromethyl)-4-quinolinemethanol (WR-184,806)

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (5), 680-689
Plasmodium falciparum, P. vivax, Aotus trivirgatus griseimembra, antimalarial activities of WR-184,806 and WR-226,253

2-(*t*-Butylaminomethyl)-4-*t*-butyl-6-(4-chlorophenyl)-phenol (WR-194,965)
Schmidt, L. H.; and Crosby, R., 1978, *Antimicrob. Agents and Chemotherapy*, v. 14 (5), 672-679

Plasmodium falciparum, *P. vivax*, *Aotus trivirgatus griseimembra*, antimalarial activities of WR-194,965 and WR-204,165

4-*tert*-Butyl-2-chlorophenyl methyl-methylphosphoroamidate. See Crufomate.

N-Butylmercurithiosalicylic acid *N*-butylester
Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

γ -Butyrolactone

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Calcium chloride

Antonenko, A. E., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 121-126
nematode eggs, coccidian oocysts, non-litter pig manure, substances tested as disinfectants

Calcium leukovorin. See Folinic acid.

Calcium oxide -- Quick lime.

Quick lime

Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, Current Sc., Bangalore, v. 47 (8), 283-284 [Letter].
Trichodina [sp.] on exterior of *Cyprinus carpio*, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

Calcium phosphate -- Superphosphate.

Superphosphate

Antonenko, A. E., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 121-126
nematode eggs, coccidian oocysts, non-litter pig manure, substances tested as disinfectants

Calf thymus DNA

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Calusterone

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Camben Cattle Paste. See Cambendazole.

Cambendazole -- Bonlam; Bovicam; Camben Cattle Paste; Camvet; Equiben; 5-Isopropoxy-carbonylamino-2-(4-thiazolyl)-benzimidazole; Isopropyl 2-(4-thiazolyl)-5-benzimidazolecarbamate; MK 905; Noviben Suspension; 2-(4-Thiazolyl)-5-carbamoylisopropoxybenzimidazole; 2-(4-Thiazolyl)-5-isopropoxycarbonylamino-benzimidazole; 2-(4-Thiazolyl)-5-isopropylcarbonylamino-benzimidazole.

Cambendazole

Amato Neto, V.; et al., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (3), 161-163
Strongyloides stercoralis, adults and children, cambendazole, clinical trials: Brazil

Cambendazole -- Continued.

Cambendazole

Baranski, M. C.; et al., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (4), 213-218
Strongyloides stercoralis, human, cambendazole, double-blind clinical study of therapeutic action, possible toxicity and adverse reactions: Brazil

Cambendazole (Camben Cattle Paste)

Bowen, F. L., 1979, Austral. J. Exper. Agric. and Animal Husb. (98), v. 19, 269-275
nematodes, weaner cattle, treatment/management procedure involving alternate grazing with sheep and monthly treatment with cambendazole, liveweight changes, financial returns: Northern Tablelands of New South Wales

Cambendazole

Cairns, G. C.; and Holmden, J. H., 1977, N. Zealand Vet. J., v. 25 (1-2), 35-37
cambendazole, efficacy against horse parasites

Cambendazole (Noviben Suspension)

Chevis, R. A. F.; and Kelly, J. D., 1978, N. Zealand Vet. J., v. 26 (9), 218-220
Fasciola hepatica, rats and sheep (both exper.), mebendazole, parbendazole, cambendazole, thiabendazole, anthelmintic activity, molecular structure-activity analyses

Cambendazole

Coles, G. C.; and Briscoe, M. G., 1978, Vet. Rec., v. 103 (16), 360-361 [Letter]
Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for *Haemonchus contortus* eggs

Cambendazol (Bonlam)

Corba, J.; et al., 1978, Veterinarstvi, v. 28 (6), 274-275
Dicrocoelium dendriticum, sheep, trials of cambendazole, fenbendazole, diamfenetid

5-Isopropoxy-carbonylamino-2-(4-thiazolyl)-benzimidazole (Cambendazole)

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Cambendazole (Camvet)

Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1979, Am. J. Vet. Research, v. 40 (4), 590-594
large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Cambendazole

Friedman, P. A.; and Platzer, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Cambendazole -- Continued.

Cambendazole (Camvet)

Greve, J. H.; Hanson, R. L.; and McGill, L. D., 1979, J. Am. Vet. Med. Ass., v. 174 (8), 828-829

larval cestode causing parasitic ascites in dog, intraperitoneal therapy with cambendazole successful: southern California

Cambendazole (Equiben)

Grzywinski, L.; et al., Med. Wet., v. 33 (2), 89-90

Strongylidae, Parascaris equorum, Oxyuris equi, horses, cambendazole

Cambendazole

Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367

Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Cambendazole

Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, Research Vet. Sc., v. 25 (3), 360-363

Haemonchus contortus, Trichostrongylus colubriformis, levels of benzimidazole resistance recorded from an egg hatch test procedure

Cambendazole (Bovicam)

Hogg, R. A., 1978, Vet. Rec., v. 103 (21), 477 [Letter]

death of two Hereford-cross bullocks following cambendazole dosing

Cambendazole

Huggins, D., 1977, Rev. Brasil. Med., v. 34 (6), 373-375

Strongyloides stercoralis, children, cambendazole given as single dose

Cambendazole

Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682

relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of Nematospiroides dubius in mice

Cambendazole

Koehler, P.; Bryant, C.; and Behm, C. A., 1978, Internat. J. Parasitol., v. 8 (5), 399-404

Fasciola hepatica, ATP synthesis in succinate decarboxylase system from mitochondria, inhibition in vitro by mebendazole and a soluble derivative of cambendazole

Cambendazole (MK 905)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73

Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Cambendazole -- Continued.

Cambendazole

Lyons, E. T.; Drudge, J. H.; and Tolliver, S. C., 1978, Vet. Med. and Small Animal Clin., v. 73 (7), 921-923

Dictyocaulus viviparus and gastro-intestinal nematodes, calves, cambendazole paste, controlled test, drug efficacy, some activity against Moniezia spp., no toxicosis: Kentucky

Cambendazole

Mango, A. M.; and Rubin, R. O., 1976, Bull. Animal Health and Prod. Africa, v. 24 (1), 69-71

Taenia saginata, calves (exper.), cambendazole, seems to be active against Cysticercus bovis in tissues

Cambendazole

Martirani, I.; and Rodrigues, L. D., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (1), 71-75

intestinal helminths, humans, clinical trials with cambendazole

Cambendazole

Milhaud, G.; Pinault, L.; and Bouchez, N., 1978, Rec. Med. Vet., v. 154 (1), 43-47

cambendazole, slaughterhouse ruminants, cattle, residues in meat and offal

Cambendazole

Prichard, R. K.; Hennessy, D. R.; and Steel, J. W., 1978, Vet. Parasitol., v. 4 (4), 309-315

Haemonchus contortus, Trichostrongylus colubriformis, sheep, Ostertagia ostertagi, cattle, 4 benzimidazoles, mode of action and pharmacokinetic behavior, implications for prolonged administration as a new concept for increasing spectrum and effectiveness of anthelmintics

Cambendazole

Rodrigues, L. D.; et al., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 57-67

Strongyloides stercoralis and other human intestinal parasites, clinical trials with cambendazole

Cambendazole

Salazar, D.; Herrera R., D.; and Quiroz R., H., 1974, Tecn. Pecuaria Mexico (26), 28-30

Ancylostoma caninum, dogs, cambendazole not satisfactory at dosage levels used

Cambendazole

Sanchez Moreno, M.; and Barrett, J., 1979, Parasitology, v. 78 (1), 1-5

Hymenolepis diminuta, adults, monoamine oxidase, occurrence and properties, inhibition by several anthelmintics

Cambendazole

Schmidt, R. L.; et al., 1979, J. Wildlife Management, v. 43 (2), 461-467

Protostrongylus, domesticated, captive, or free-ranging Ovis c. canadensis, evaluation of 5 drugs

Cambendazole -- Continued.

Cambendazole

Simpkin, K. G.; and Coles, G. C., 1978, Research Vet. Sc., v. 25 (2), 249-250
Haemonchus and Trichostrongylus colubriformis in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parbendazole, oxbendazole)

Cambendazole

Slocombe, J. O. D.; and Cote, J. F., 1978, Canad. Vet. J., v. 19 (7), 184-186
gastrointestinal nematodes, horses, field trial with cambendazole paste, effective

Camolar. See Cycloguanil.

Camoprim. See Amodiaquine or Primaquine.

Camoquine. See Amodiaquine.

Camphechlor. See Toxaphene.

Camptothecin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Camvet. See Cambendazole.

Canaural. See Neomycin or Nystatin or Prednisolone.

L-Canavanine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Canesten. See Clotrimazole.

Cannabidiol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

Cannabinol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

Cannabinol, abnormal

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

9-nor-Cannabinol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

Cantrodifene. See Nitroscanate.

Cantrodiphen. See Nitroscanate.

Caparsolate. See Arsenamide.

Caparsolate sodium. See Arsenamide.

Capsicum annum

Frischkorn, C. G. B.; Frischkorn, H. E.; and Carrazzoni, E., 1978, Naturwissenschaften, v. 65 (9), 480-483
Schistosoma mansoni, cercaricidal activity of various plants from northeast Brazil

Captan -- N-Trichloromethyl-thio-4-cyclohexene-1,2-dicarboximide.

N-Trichloromethyl-thio-4-cyclohexene-1,2-dicarboximide

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Carbadox (Mecadox)

Stewart, T. B.; et al., 1979, Am. J. Vet. Research, v. 40 (10), 1472-1475
Strongyloides dentatus and other parasites, pigs on contaminated lots, pyrantel tartrate and carbadox in feed

4-Carbamidophenyl arsenic acid. See Carbarsone.

2-Carbamomethoxy-5-benzoylbenzimidazole. See Mebendazole.

2-Carbamomethoxy-5-butylbenzimidazole. See Parbendazole.

2-Carbamomethoxy-5-phenylthiobenzimidazole. See Fenbendazole.

2-Carbamomethoxy-5-propoxybenzimidazole. See
Oxibendazole.

Carbanolate

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Carbarsone -- 4-Carbamidophenyl arsenic acid;
Carbosep.

Carbarsone (Carbosep)

McDougald, L. R., 1979, Poultry Science, v. 58 (1), 76-80
coccidiosis, histomoniasis, turkeys, tests for efficacy and compatibility indicate that amprolium and carbarsone can be used in combination

4-Carbamidophenyl arsenic acid (Carbarsone)

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriol., 1 Abt. Orig., Reihe A, v. 241 (3), 358-367
Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Carbaryl -- N-Methyl-1-naphthylcarbamate; 1-Naphthalenol methylcarbamate; 1-Naphthyl methyl carbamate; Opigal; Sevin.

Carbaryl (Sevin)

Abdel Rahman, M. S.; El Gendi, A. Y. I.; and Moursi, H. S. A., 1977, J. Egypt. Vet. Med. Ass., v. 37 (2), 55-69
nematodes and cestodes, dogs, effectiveness of trichlorfon, oxinotriphos, and carbaryl compared: Giza and Cairo Governorates, Egypt

Carbaryl (Sevin)

Barnes, A. M.; et al., 1974, J. Med. Entom., v. 11 (1), 83-87
flea vectors of plague, control on wild Peromyscus maniculatus by use of 2% carbaryl dust in bait stations: Buckhorn Mountain, Larimer County, Colorado

Carbaryl

Bonin, W., 1977, Berl. u. Munchen. Tierarztl. Wchenschr., v. 90 (2), 34-37
Amblyomma hebraeum, Psoroptes cuniculi, Melophagus ovinus, Dermanyssus gallinae, heptenophos, rapid mode of action, broad range of efficacy, short residual effect and effective as a vapour poison, compared with other standard drug preparations

Carbaryl

Cole, M. M.; et al., 1973, J. Econom. Entom., v. 66 (1), 118-119
Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

Carbaryl

Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439
economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Carbaryl -- Continued.

Carbaryl

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Carbaryl

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Carbaryl

Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, J. Econom. Entom., v. 71 (2), 315-318
Ornithonyssus sylviarum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban permethrin and SD-43775 also effective; mites displayed tolerance to malathion

Carbaryl

Loomis, F. C.; Bramhall, E. L.; and Dunning, L. L., 1979, J. Econom. Entom., v. 72 (6), 856-859
Ornithonyssus sylviarum, White Leghorn hens, carbaryl and fenvalerate compared

Carbaryl (Sevin)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Sevin

Puchkova, E. A., 1977, Veterinariia, Moskva (7), 19-22
D[ermanyssus] gallinae, C[imex] lectularius, lice, chickens on industrial scale farms, control, sevin, dicresil, chloronphos, carbophos; other complex sanitation measures

Opigal

Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Carbaryl

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Carbaryl

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Carbaryl -- Continued.

Carbaryl

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Sevin

Sanchez Moreno, M.; Monteoliva, M.; and Her-moso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Carbathion

Krasnonos, L. N., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 186-189
ascaris microfocus, eggs from soil treated with carbathion, invasive capacity reduced

Carbetox 37

Zarzara, C., 1979, Rev. Crest. Animalelor, v. 29 (5), 42-45
Cimex lectularius, lindatox-20, neguvon, and carbetox 37, laboratory tests; carbetox 37 for control in poultry sheds, application methods

Carbidium ethane sulphonate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Carbidium sulfate -- 3-Amino-9-p-carbethoxyamino-phenyl-10-methyl-phenanthridinium ethano-sulphonate.

Carbidium sulphate

Brener, Z., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (4), 302-306
Trypanosoma cruzi, description of method allowing study of drug action on trypomastigotes in mice

Carbocyclic-adenosine. See Cycloadenosine.

Carbocyclic-inosine

Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Carbocyclic-6-methyl-mercaptapurine ribonucleoside

Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Carbolic acid. See Phenol.

3'-Carbomethoxybenzo-15-crown-5

Brown, G. R.; and Foubister, A. J., 1979, J. Med. Chem., v. 22 (8), 997-999
 benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against *Eimeria tenella*

4'-Carbomethoxybenzo-15-crown-5

Brown, G. R.; and Foubister, A. J., 1979, J. Med. Chem., v. 22 (8), 997-999
 benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against *Eimeria tenella*

Carbon disulfide + Piperazine (= Mascyl)

Olsson, T., 1977, Svensk Vet.-Tidn., v. 29 (20), 795-800
Ascaris suum, slaughter swine, treatment with neguvon or mascyl

Carbon disulphide

Saxena, S. N., 1974, U. P. Vet. J., v. 2 (4), 132-137
*Ascarops strongylin*a, piglets (exper.), critical trials of efficacy of carbon disulphide, thiabendazole, and sodium fluoride against mature worms

Carbon tetrachloride -- CCl_4 ; Fasciolin [of Furmaga, S.; Gundlach, J. L.; and Sobieszewski, K., 1974]; Tetrafinol; Red Spot.

Carbon tetrachloride

Angus, K. W.; and Greig, A., 1979, J. Comp. Path., v. 89 (4), 605-607
 anthelmintic dose of carbon tetrachloride, lamb, acute poisoning, renal and hepatic calcification

Carbon tetrachloride (Red Spot)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
Haemonchus contortus, *Trichostrongylus colubriformis*, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Carbon tetrachloride

Cawthorne, M. A.; et al., 1971, Research Vet. Sc., v. 12 (6), 516-520
 sheep, carbon tetrachloride toxicity markedly increased by prior administration of DDT, ethoxyquin given 24-48 hr before CCl_4 prevented toxicity, ethoxyquin did not diminish fasciolicidal action of CCl_4

Carbon tetrachloride

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Carbon tetrachloride -- Continued.Fasciolin (CCl₄)

Furmaga, S.; Gundlach, J. L.; and Sobieszewski, K., 1974, *Acta Parasitol. Polon.*, v. 22 (12-21), 229-247

Fasciola hepatica-infected and uninfected sheep treated and not treated with 3 fasciolicides (fasciolin, distolon, and zanil), blood serum levels of Ca, P, Na, K, and Mg

Carbon tetrachloride

Parshad, V. R.; and Guraya, S. S., 1978, *Vet. Parasitol.*, v. 4 (2), 111-120

4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces

Carbon tetrachloride

Petrov, Iu. F., 1978, *Veterinariia, Moskva* (5), 64-66

[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Carbon tetrachloride

Sethuraman, V.; and Verma, B. B., 1978, *Indian J. Animal Sc.*, v. 48 (8), 622-625

carbon tetrachloride toxicity, healthy buffalo calves, clinical, haematological, and biochemical changes

Tetrafinol

Tesarcik, J., 1971, *Prace VURH Vodnany* (9), 99-132

Neoechinorhynchus rutili, *Eimeria* spp., carp, tetrafinol, nitrofurazone, prophylaxis and therapy under aquarium conditions

Tetrafinol

Tesarcik, J., 1972, *Parazitologiya, Leningrad*, v. 6 (2), 190-191

Neoechinorhynchus rutili, localization in intestine of carp, changes during season, anthelmintic introduced per rectum is not effective, better results with tetrafinol fed to fish at 1 mg/kg body weight

Carbon tetrachloride

Vodrazka, J.; et al., 1979, *Biol. a Chem. Zivoc. Vyroby, Vet.*, v. 15 (v. 21) (4), 363-374

tribromsalan, dynamic behavior in sheep blood plasma; antifasciolicides, effects on some sheep plasma enzymes

Carbon tetrachloride

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, *Austral. Vet. J.*, v. 54 (10), 501-502 [Letter]

Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thia-bendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxynil are also useful chemical alternatives

Carbonyl cyanide m-chlorophenyl hydrazone (CCCP)

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Carbophenothion -- S-[[(4-Chlorophenyl)thio]

methyl] 0,0-diethyl phosphorodithioate; Garrathion.

Carbophenothion

Drummond, R. O.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Carbophenothion (Garrathion)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, *PANS*, v. 25 (2), 135-142

Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Carbophos. See Malathion.Carbosep. See Carbarsone.

3-Carboxybenzo-15-crown-5

Brown, G. R.; and Foubister, A. J., 1979, *J. Med. Chem.*, v. 22 (8), 997-999

benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against *Eimeria tenella*

4'-(2-Carboxyethyl)benzo-15-crown-5

Brown, G. R.; and Foubister, A. J., 1979, *J. Med. Chem.*, v. 22 (8), 997-999

benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against *Eimeria tenella*

2-Carboxy-5-fluorovaleric acid derivatives

Sevcik, B.; et al., 1974, *Veterinaria, Praha*, v. 16 (5-6), 421-588

Eimeria tenella, chickens, 613 substances screened as coccidiostats, extensive detailed statistical results

N¹-Carboxymethyleneamino-2-(thiazolyl)5-isopropoxycarbonyl-aminobenzimidazole-triethylamino salt

Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618

Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

2-(4'-Carboxystyryl)-5-nitro-1-vinylimidazole

Morton, D. M.; Fuller, D. M.; and Green, J. N., 1973, *Xenobiotica*, v. 3 (4), 257-266

2-styryl-5-nitroimidazoles, metabolism and excretion in laboratory animals, activity against *Trypanosoma rhodesiense*

Caricide. See Diethylcarbamazine.

Carmine

Daniyarov, I. A.; et al., 1978, Veterinaria, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Carnidazole -- Spartrix.

Spartrix

Hauser, K. W., 1977, Prakt. Tierarzt, v. 58, special no., 56
trichomoniasis, parrots, diagnosis, treatment with ronidazole and spartrix, review

Carrageenan

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cascara sagrada + Sodium santoninate (Ascarel) Oldham, R. R.; et al., 1971, South. Med. J., v. 64 (4), 480-482

possible santonin poisoning (hemolytic crisis) in young child treated for worms with ascarel

Caviphos -- [0-Methyl-0-(2,2-dichlorovinyl) phosphate]₂, Ca[0,0-dimethyl-(2,2-dichlorovinyl) phosphate]₂.

Caviphos

Colglazier, M. L., 1979, Am. J. Vet. Research, v. 40 (3), 384-386

gastrointestinal helminths and stomach bots, ponies, critical trials with oxfendazole and caviphos; observations on spontaneous elimination of small strongylids prior to treatment: Maryland

CaVP. See Dankil.

CCl₄. See Carbon tetrachloride.

Cedrus deodara wood oil

Chandra, S.; et al., 1978, Indian Vet. J., v. 55 (12), 963-965

Cedrus deodara wood oil, rabbits, increase in vascular permeability at site of local application, possible use in mange-infected sheep

Cellocidin

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Cetovex. See trans-1,4-Di-(2-isothiocyanatoethyl) cyclohexane.

Cetrimide. See Cetrimonium bromide.

Cetrimonium bromide -- Cetrimide; Cetyl-trimethyl-ammonium bromide.

Cetrimide

Eslami, A.; Ahrari, H.; and Saadatzaheh, H., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (3), 307-308

Echinococcus granulosus, scolicalidal activity of cetrimide compared to that of sodium chloride; findings suggest that cetrimide can be used successfully during human hydatid surgery

Cetyl-trimethyl-ammonium bromide. See Cetrimonium bromide.

Chaulmoogra oil

Chakrabarti, A.; and Misra, S. K., 1979, Indian Vet. J., v. 56 (6), 497-500

Demodex canis, dogs, incidence in relation to season, host age, sex, and breed, clinical manifestations, in vivo and in vitro activity of several acaricides: India

Chemcoccid

Epllov, N. N.; and Nazarov, V. G., 1977, Veterinaria, Moskva (6), 65-66

Eimeria spp., calves, chemcoccid effective; compared with biomycin and norsulfazole: Belgorodsk oblast

Chemcoccid

Krylov, V. F., 1978, Veterinaria, Moskva (10), 68-69

Eimeria tenella strain resistant to pharmacocid after 35 laboratory passages in chickens, cross-resistance only to rigecocin

Chemosterilants

Adhami, N.; and Khan, N. H., 1976, Indian J. Entom., v. 37 (1), 1975, 60-63

Cimex hemipterus, hempa as chemosterilant, reduction of oviposition and egg viability, more effective against males than females as measured by egg hatchability

Chemosterilants

Osburn, R. L.; and Oliver, J. H., jr., 1978, J. Parasitol., v. 64 (4), 719-726

Dermacenter variabilis, effects of metepa on cytology and fertility of males treated as unfed adults

Chemosterilants

Osburn, R. L.; and Oliver, J. H., jr., 1979, Ann. Entom. Soc. Am., v. 72 (6), 740-745

Dermacenter variabilis, chemosterilization with metepa, effect on cytology and fertility of fed and partially fed males

Chemosterilants

Rawlins, S. C.; Jurd, L.; and Snow, J. W., 1979, J. Econom. Entom., v. 72 (5), 674-677

Cochliomyia hominivorax, antifertility effects of benzylphenols and benzyl-1,3-benzodioxoles

Chenopodium oil -- Ascaridin.

Chenopodium oil

Andrien, J. M.; et al., 1971, Ann. Soc. Belges Med. Trop. Parasitol., v. 51 (3), 299-308
fatal human encephalitis after administration of chenopodium oil as a vermifuge, case reports

Ascaridin

Do Duong Thai; and Nguyen Tuyet Mai, 1973, Rev. Med., Hanoi, 165-168
Ascaris lumbricoides, laboratory studies on drug resistance to anthelmintics (ascaridin used in experiments), finding that sex of worm, maturity of worm, and drug concentrations all influenced reactions to drugs

Chlophenotane spirits. See DDT.

Chloquinat + Chloroquin phosphate + Diiodo-hydroxy quinaline (=Resotren [composite])

Samaddar, J.; Biswas, D. N.; and Ghose, A. N., 1978, Indian Vet. J., v. 55 (7), 572-576
leucocytozoonosis, W[white] L[eg] H[orn] birds, quinine bisulphate and resotren failed to ensure absolute recovery, though general condition of treated birds improved

Chlorambucil

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Chloramine T

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Chloramphenicol -- Chloromycetin; Quemicetina; Sintomycin.

Quemicetina + Nitrimidazine + Nystatin (as a combined vaginal tablet)

Aruta, J.; Galani, M.; and Fertilio, O., 1973, Rev. Chilena Obst. y Ginec., v. 38 (3), 118-120
human vaginal trichomoniasis, comparison therapeutic trials using oral naxogin and vaginal tablets of naxogin combined with nystatin and quemicetina in cases with diagnostic problems after diagnosis confirmed by cytology or colposcopy: Chile

Chloramphenicol + Metronidazole + Chloroquine

Bindschadler, D. D., 1974, Rocky Mountain Med. J., v. 71 (7), 387-389
E[ntamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Chloramphenicol -- Continued.

Sintomycin

Gobzem, V. R.; and Nazarov, V. G., 1978, Veterinariia, Moskva (3), 67-69
Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Chloramphenicol (Chloromycetin)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, Chemotherapy, v. 25 (4), 222-226
Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxy-tetracycline and tetracycline in controlling acute infection

Chloramphenicol

Seilhamer, J. J.; and Byers, T. J., 1978, J. Protozool., v. 25 (4), 486-489
Acanthamoeba castellanii, mutants resistant to erythromycin, chloramphenicol, and oligomycin

Chloramphenicol

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Chloramphenicol

Watson, A. D. J.; Farrow, B. R. H.; and Hoskins, L. P., 1978, Austral. Vet. Practitioner, v. 8 (3), 129-132
Haemobartonella felis, cats (nat. and exper.), thiacetarsamide sodium therapy; prednisolone, tetracycline, and chloramphenicol given sequentially

Chlorchinaldol. See Chlorquinaidol.Chlordimeform. See Chlorphenamidine.Chlorfenvinfos. See Chlorfenvinfos.Chlorfenvinphos -- Birlane; Chlorfenvinfos; 2-Chloro-1-(2,4-dichlorophenyl) vinyl diethyl phosphate; Compound 4072; Diethyl 1-(2,4-dichlorophenyl)-2-chlorovinyl-phosphate; GC 4072; Pfizona; Supona; Supone; Suposan.

Supona

Atef, M.; and El-Say, A., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 73-84
Hyalomma dromedarii, Rhipicephalus s. sanguineus, laboratory tests (immersion technique) with delnav, supona, toxaphene, BHC

Supona (Chlorfenvinphos)

Chemtai, A. A. K., 1977, Bull. Animal Health and Prod. Africa, v. 25 (3), 299-306
acaricide-treated zebu cattle, blood cholinesterase, radiometric assay

Chlorfenvinphos -- Continued.

Pfizona

Chineme, C. N.; Bida, S. A.; and Nuru, S., 1979, Bull. Animal Health and Prod. Africa, v. 27 (1), 41-45

Sarcoptes scabiei var. ovis, sheep, severe infestation, clinical and histopathology, Pfizona dipping: Kaduna State, Nigeria

Chlorfenvinphos

Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439
economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Compound 4072

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Chlorfenvinphos

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Suposan (Supona)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105

Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Chl[or]fenvinphos

Hammant, C. A., 1977, Rhodesian Vet. J., v. 8 (4), 67-70
ticks, resistance survey of field strains to commonly used ixodicides, changeover from arsenic to dioxathion dipping of cattle, dramatic improvement in tick control, necessitates complete overhaul of dipping facilities and retraining of personnel: Tribal Trust Lands of Rhodesia

Chlorfenvinphos

Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73
Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Supona (Chlorfenvinfos, GC 4072)

Khan, M. H.; and Srivastava, S. C., 1977, Indian J. Animal Health, v. 16 (2), 137-140
Boophilus microplus engorged females, in vitro tests with dursban, gamma BHC, sumithion, supona, dimecron, egg production and viability; supona most effective

Chlorfenvinfos (Birlane; Supona)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Chlorfenvinphos -- Continued.

Chlorfenvinphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Chlorfenvinphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Chlorphenvinphos (Supona)

Rechav, Y.; Whitehead, G. B.; and Terry, S. B., 1978, J. South African Vet. Ass., v. 49 (2), 99-101
ticks, mortality curves of larvae dipped in dioxathion, chlorphenvinphos, and oxionthiophos, time of application, larvae of ticks exhibit diel periodicity in sensitivity to acaricides

Supone

Reich, C. I.; et al., 1978, Exper. Parasitol., v. 44 (1), 50-55
Boophilus microplus, 2 Argentinian strains, one resistant and one sensitive to organophosphate acaricides, differences in cholinesterase system

Chlorguanide -- Bigumal; Paludrine; Proguanil; Proguanil hydrochloride.

Proguanil

Buyst, H., 1975, Ann. Soc. Belge Med. Trop., v. 55 (2), 95-104
Trypanosoma rhodesiense, human, specific treatment with suramin and mel B, adjuvant antimalarial treatment with chloroquine and proguanil; modifications of sleeping sickness therapy advocated on physio-pathological and epidemiological grounds: Luangwa Valley, Zambia

Proguanil

Buyst, H., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 201-212
sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia

Chlorguanide (Paludrine)

Gon, F.; and Reid, F. P., 1975, South African Med. J., v. 49 (4), 120-122
disseminated intravascular coagulation with haemolytic anaemia and thrombocytopenia, fatal illness in man probably caused by malarial prophylaxis (pyrimethamine and chlorguanide): Johannesburg, South Africa (from Zambia)

Bigumal

Levchenko, F. F., 1978, Veterinariia, Moskva (7), 62-64
Theileria annulata, cattle, chinocide and bigumal treatment effective, treatment with hemostimulin and vitamins and microelements given in feed to counteract anemia and atonicity of digestive tract resulting from therapy: Gissarsk valley

Chlorguanide -- Continued.

Proguanil hydrochloride

Marshall, R. J.; and Ojewole, J. A. O., 1978, *Toxicol. and Applied Pharm.*, v. 46 (3), 759-768

quinoline and nonquinoline antimalarial drugs, effects on isolated guinea pig cardiac muscle

Chlorguanide (Proguanil)

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, *Israel J. Med. Sc.*, v. 14 (5), 601-605

Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiprotozoite immunity by natural bites of infected mosquitoes, symposium presentation

Proguanil

Ray, A. P.; Parkinson, A. D.; and Black, R. H., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (1), 19-22

Plasmodium berghei in chloroquine resistant white mice, results of treatment with combinations of proguanil and dapsone

Proguanil

Varnai, F.; and Ecker, A., 1977, *Therap. Hungar.*, v. 25 (4), 131-133

malaria, humans travelling to endemic areas, drug prophylaxis, comparative study, least unwanted side effects and lowest morbidity rate recorded with pyrimethamine: Hungary

Chlorhexidine -- Chlorhexidine gluconate; Hibitane.

Chlorhexidine (Hibitan®)

Ovchinnikov, N. M.; and Skuratovich, A. A., 1978, *Vestnik Dermat. i Venerol.* (6), 49-52
trichomoniasis and other human venereal diseases, experimental trials with chlorhexidine, humans, rabbits

Chlorhexidine gluconate

Stepien-Rukasz, H., 1978, *Terap. i Lek.*, v. 6, v. 28 (10), 350-352

Taenia saginata, *Ascaris lumbricoides*, *Enterobius vermicularis*, eggs, effects of chlorhexidine

Chlorhexidin

Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230

Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Chlorhexidine gluconate. See Chlorhexidine.

Chlorine

Iwanczuk, I.; and Kelus, J., 1971, *Roczniki Panstw. Zakl. Hig.*, v. 22 (2), 179-187

tests for ovicidal activity of iodine and chlorine in swimming pools using *Aspiculuris tetraptera* as model for human *Enterobius vermicularis* and *Ascaris suis* as model for *Ascaris lumbricoides*; neither chemical was an effective ovicide

Chlorine

Uvaliev, I. U.; and Baigaziev, K. K., 1979, *Vestnik Sel'skokhoz. Nauki Kazakhstana* (1), 75-78

besnoitiosis, bovine, disinfection of animals or hides by sodium hydroxide solution spray; disinfection of premises by sodium hydroxide or chlorine solution sprays

Chlormethiuron -- 3-(4-Chloro-2-methyl-phenyl)-1,1-dimethyl-thiuron; Dipofen.

Chlormethiuron (Dipofen)

Gothe, R.; and Guizar, R. C., 1979, *Zentralbl. Vet.-Med.*, Reihe B, v. 26 (4), 290-303

Argas walkerae, chickens (exper.), in vivo model for evaluating detaching potential of various acaricides

Chlorobromofebrifugine (CBF)

Ishii, T.; and Onaga, H., 1978, *Nippon Zyuisei-Kai Zassi (J. Japan Vet. Med. Ass.)*, v. 31 (9), 534-538

Eimeria spp., chicks (exper.) and cell cultures, activity of chlorobromofebrifugine

DL-trans-6-Chloro-7-bromo-3-[3-(3-hydroxy-2-piperidyl) acetyl]-4-(3H)-quinazolinone hydrobromide. See Halofuginone.

2-Chloro-1-(2,4-dichlorophenyl) vinyl diethyl phosphate. See Chlorfenvinphos.

6-Chloro-5-8-diethylaminoethylamino-8-methyl-quinoline -- RD-12,869.

RD-12,869

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (1), 43-46

Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

8-Chloro-2-[(diethylamino)ethyl]-2H-[1]-benzothioopyrano[4,3,2-cd]indazole-5-methanol monomethane-sulfonate -- IA-4; IA-4 N-oxide.

IA-4

Batzinger, R. P.; and Bueding, E., 1977, *J. Pharmacol. and Exper. Therap.*, v. 200 (1), 1-9
mutagenic activities in vitro and in vivo of 5 antischistosomal compounds, comparative anti-*Schistosoma mansoni* activities of hycanthone, IA-4, and IA-4 N-oxide, observations provide evidence that mutagenic activities can be dissociated from desired chemotherapeutic effects by suitable structural modifications

IA-4 N-oxide

Batzinger, R. P.; and Bueding, E., 1977, *J. Pharmacol. and Exper. Therap.*, v. 200 (1), 1-9
mutagenic activities in vitro and in vivo of 5 antischistosomal compounds, comparative anti-*Schistosoma mansoni* activities of hycanthone, IA-4, and IA-4 N-oxide, observations provide evidence that mutagenic activities can be dissociated from desired chemotherapeutic effects by suitable structural modifications

- 0[2-Chloro-2-(diethyl carbamoyl)-1-methyl-vinyl]-0,0-dimethyl phosphate. See Phosphamidon.
- 4'-Chloro-3,5-diiodosalicylanilide. See Clioxanide.
- 4'-Chloro-3,5-diiodosalicylanilide acetate ester. See Clioxanide.
- 1-(2-Chloroethyl)-3-cyclohexyl-1-nitrosourea
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 5-Chloroethylthiamine. See Beclotiamine.
- 9-(2-Chloro-6-fluorobenzyl) adenine. See Arprinocid.
- 9-(2-Chloro-6-fluorobenzyl) adenine-1-N-oxide (ICI 125752)
Latter, V. S.; and Wilson, R. G., 1979, *Parasitology*, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture
- 9-(2-Chloro-6-fluorophenylmethyl)-9H-purin-6-amine. See Arprinocid.
- 4'-Chloro-2'-hydroxy-3,5-diiodosalicylanilide
Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosis in vitro, scolicalidal effect of salicylanilide and bisphenol derivatives
- 2'-Chloro-1-hydroxy-2-naphthanilide-4'-isothiocyanate
Singh, H.; et al., 1978, *Ztschr. Naturforsch.*, Sect. C, *Biosc.*, v. 33 (5-6), 447-448
Hymenolepis nana, rats, mice, 2'-chloro-1-hydroxy-2-naphthanilide-4'-isothiocyanate, synthesis and cestodicidal activity, highly effective and safe, comparative efficacy with yomesan; further tests showed marked activity against H. diminuta in rats and Taenia sp. in dogs
- 7-Chloro-10-hydroxy-3-(4-trifluoromethyl-phenyl)-3,4-dihydroacridine-1,9(2H,10H)-dione. See Floxacrine.
- 7-Chlorolincomycin. See Clindamycin.
- alpha-Chloromethyl-2-methyl-5-nitro-imidazole-1-ethanol. See Ornidazole.
- 0-(3-Chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) 0,0-diethyl phosphorothioate. See Coumaphos.
- 3-(4-Chloro-2-methyl-phenyl)-1,1-dimethyl-thiuron. See Chlormethiuron.
- Chloromycetin. See Chloramphenicol.
- 2-Chloro-4-nitrobenzamide. See Aklomide.
- N-(2'-Chloro-4'-nitro-phenyl)-5' chlorosalicylanilide piperazine salt. See Niclosamide.
- Chlorophenoxamide. See Chlorophenoxamide.
- 1-(p-Chlorophenoxy)-6,7-epoxy-3,7-dimethyl-2-nonene
Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, *J. Econom. Entom.*, v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity
- 1-(p-Chlorophenoxy)-6,7-epoxy-3-ethyl-7-methyl-2-nonene
Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, *J. Econom. Entom.*, v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity
- N-[[4-Chlorophenyl]amino]carbonyl]-2,6-difluorobenzamide. See Diflubenzuron.
- 3-(4-Chlorophenyl)-alpha-[(dibutylamino)methyl]-5,7-dichloro-1-naphthalenemethanol hydrochloride
Shamblee, D. A.; and Gillespie, J. S., jr., 1979, *J. Med. Chem.*, v. 22 (1), 86-89
Plasmodium berghei, mice, activity of trichloronaphthalene amino alcohols
- 1-[3-(4-Chlorophenyl)-5,7-dichloro-1-naphthyl]-3-(di-n-butylamino)propanol hydrochloride
Shamblee, D. A.; and Gillespie, J. S., jr., 1979, *J. Med. Chem.*, v. 22 (1), 86-89
Plasmodium berghei, mice, activity of trichloronaphthalene amino alcohols
- 1-[3-(4-Chlorophenyl)-5,7-dichloro-1-naphthyl]-2-(2-piperidyl)ethanol hydrochloride
Shamblee, D. A.; and Gillespie, J. S., jr., 1979, *J. Med. Chem.*, v. 22 (1), 86-89
Plasmodium berghei, mice, activity of trichloronaphthalene amino alcohols
- 3-(4-Chlorophenyl)-5,7-dichloro-alpha-(2-piperidyl)-1-naphthalenemethanol acetate
Shamblee, D. A.; and Gillespie, J. S., jr., 1979, *J. Med. Chem.*, v. 22 (1), 86-89
Plasmodium berghei, mice, activity of trichloronaphthalene amino alcohols
- (+)-5-(o-Chlorophenyl)-1,3-dihydro-3-methyl-7-nitro-2H-1,4-benzodiazepine-2-one -- Ro 11-3128.
- Ro 11-3128
Baard, A. P.; et al., 1979, *South African Med. J.*, v. 55 (16), 617-618 [Letter]
schistosomiasis, hospitalized patients, open dose-finding study and preliminary clinical trials using Ro 11-3128

(+)-5-(o-Chlorophenyl)-1,3-dihydro-3-methyl-7-nitro-2H-1,4-benzodiazepine-2-one -- Continued.

(+)-5-(o-Chlorophenyl)-1,3-dihydro-3-methyl-7-nitro-2H-1,4-benzodiazepine-2-one (Ro 11-3128)
Pax, R.; Bennett, J. L.; and Fetterer, R., 1978, Arch. Pharmacol., v. 304 (3), 309-315
Schistosoma mansoni, S. japonicum, praziquantel and Ro 11-3128 produce rapid rise in tension of musculature, uptake studies of inorganic cations suggest that interference with inorganic ion transport mechanisms causes contraction of schistosome musculature

RO11-3128

Pax, R.; Fetterer, R.; and Bennett, J. L., 1979, Comp. Biochem. and Physiol., v. 64C (1), 123-127

Schistosoma mansoni, effects of fluoxetine and imipramine on adult males in vitro, interactions with 5-hydroxytryptamine-induced contractile activity, interaction with anti-schistosomal compounds praziquantel and RO11-3128

N-(4'-Chlorophenyl)-3,5-diiodoacetylsalicylamide.
See Clixoxide.

p-Chlorophenyl isothiocyanate (Sch 20350)
Loebenberg, D.; et al., 1979, J. Parasitol., v. 65 (2), 233
dogs naturally infected with various helminths, anthelmintic activity of Sch 20350

2-(4-Chlorophenyl)-2-(4-piperidyl)-tetrahydrofuran hydrochloride (WR 93,133)
Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

S-[[[4-Chlorophenyl]thio]methyl] 0,0-diethyl phosphorodithioate. See Carbophenothion.

Chlorophos. See Trichlorfon.

6-Chloropurine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

N'-(6-Chloro-2 pyrazinyl)-sulfanilamide. See Sulfachloropyrazine.

Chloroquine -- Aralen; Avlochlor; Avloclor;
Chloroquine base; Chloroquine diphosphate;
Chloroquine phosphate; Chloroquine sulfate;
Contramibial (with Diiodohydroxyquin and Tetracycline); Delagil; Malaquin; Nivaquine; Nivaquine 200; Resochin; Resotren [composite] (with Chloquinatone and Diiodohydroxy quinoline); 4-(1'-Methyl-4'-diethylaminobutyl-amino)-7-chloroquinoline; WR 1,544.

Chloroquine -- Continued.

Chloroquine phosphate (Resochin)
Abu-Aisha, H.; Abu-Sabaa, H. M. A.; and Nur, T., 1979, J. Trop. Med. and Hyg., v. 82 (2), 36-37
cardiac arrest, man given undiluted chloroquine intravenously for possible cerebral malaria, cautions on mode of drug administration

Chloroquine

Ajao, O. G., 1978, J. Trop. Med. and Hyg., v. 81 (8), 153-155
malaria, overt attacks in humans as cause of post-operative fever, depressed acquired immunity resulting from stress of surgery, recommends routine administration of chloroquine prior to surgical procedures: Nigeria

Chloroquine diphosphate

Al Tawil, N., 1977, J. Trop. Med. and Hyg., v. 80 (12), 230-237
Plasmodium falciparum, strain resistant to chloroquine therapy discovered in Vientiane, Lao People's Democratic Republic

Chloroquine

Al Tawil, N., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (3), 409-413
Plasmodium falciparum, semi-immune humans, clearance of asexual parasitaemia with single dose sulfadoxine-pyrimethamine, comparison with standard dose of chloroquine over 3 days: Laos

Nivaquine

Andrianjafy, M.; Thuillier, M.; and Gandrille, M. C., 1976, Rev. Franc. Transfus. et Immunohematol., v. 19 (4), 603-604
malaria in young woman given blood transfusions, donor of blood found to have high malarial antibody titers, successful treatment with nivaquine, case report: France

Delagil

Avessalomov, I. S.; and Baenov, N., 1977, Veterinariia, Moskva (8), 76
theileriasis, cattle, delagil combined with vitamins and penicillin, effective and non-toxic

Chloroquine (Aralen)

Ayala, S. C., 1978, Vet. Med. and Small Animal Clin., v. 73 (2), 217-218
Plasmodium brasilianum in pet Cebus capucinus (blood), case report, chloroquine, good results, public health implications: Cali Colombia, acquired from Catatumbo River region along Colombia-Venezuela border, South America

Chloroquine + Chloramphenicol + Metronidazole
Bindschadler, D. D., 1974, Rocky Mountain Med. J., v. 71 (7), 387-389
[Entamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Chloroquine -- Continued.

Chloroquine + Emetine

Bindschadler, D. D., 1974, Rocky Mountain Med. J., v. 71 (7), 387-389

E[ntamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Chloroquine phosphate

Boonpucknavig, V.; Boonpucknavig, S.; and Bhamarapravati, N., 1979, Arch. Path. and Lab. Med., v. 103 (11), 567-572

Plasmodium berghei berghei-infected mice treated with chloroquine phosphate, focal glomerulonephritis in hyperimmune state, clinical, immunopathologic, and histopathologic findings

Chloroquine

Brohult, J.; et al., 1979, Ann. Trop. Med. and Parasitol., v. 73 (4), 327-331

malaria, humans, analysis of chemoprophylactic habits and reasons for breakdowns in therapy, small mining town: Yekepa, Liberia

Chloroquine diphosphate (Avlochlor)

Brohult, J.; et al., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 401-405

Plasmodium falciparum, chloroquine use by non-immunes when in endemic areas, analysis of drug concentrations in human serum during short and long term malaria prophylaxis, recommendations for standard vs. double dosage, clinical implications

Chloroquine diphosphate

Brohult, J.; et al., 1979, Lancet, London (8141), v. 2, 522 [Letter]

malaria prophylaxis, chloroquine diphosphate, recommended dosage for non-immune individuals

Chloroquine diphosphate

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Chloroquine

Buyst, H., 1975, Ann. Soc. Belge Med. Trop., v. 55 (2), 95-104

Trypanosoma rhodesiense, human, specific treatment with suramin and mel B, adjuvant antimalarial treatment with chloroquine and proguanil; modifications of sleeping sickness therapy advocated on physio-pathological and epidemiological grounds: Luangwa Valley, Zambia

Chloroquine

Buyst, H., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 201-212

sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia

Chloroquine -- Continued.

Chloroquine

Cerecedo Cortina, V.; and Gaxiola Gaxiola, R., 1972, Rev. Fac. Med., Univ. Nac. Auton. Mexico, an. 15, v. 15 (1), 25-28

human hepatic amoebic abscess, side effects of currently used amoebicides (emetine, chloroquine, metronidazole, aminosidine)

Chloroquine

Cheah, J. S.; et al., 1970, Singapore Med. J., v. 11 (4), 287-289

Clonorchis sinensis, man, case report, pathology, chloroquine, possibly infected by eating raw fresh-water carp imported from China: Singapore

Chloroquine

Chongsuphajaisiddhi, T.; et al., 1979, Southeast Asian J. Trop. Med. and Pub. Health, v. 10 (1), 132-137

falciparum malaria, children, chloroquine resistance, efficacy of quinine and fansidar, clinical study: Thailand

Chloroquine

Choudhry, V. P.; et al., 1978, Trop. and Geogr. Med., v. 30 (3), 331-335

malaria, chloroquine-induced haemolysis and acute renal failure in children with glucose-6-phosphate dehydrogenase deficiency

Chloroquine

Dutta, G. P.; and Narain, L., 1978, Indian J. Exper. Biol., v. 16 (7), 838-840

Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Chloroquine

Eke, R. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 1074-1075

Plasmodium falciparum, possible chloroquine-resistant strain, recrudescence of infection in 42-year-old hospital worker after chloroquine therapy, radical cure with sulfadiazine and pyrimethamine: Nigeria

Chloroquine

Elslager, E. F.; et al., 1979, J. Med. Chem., v. 22 (10), 1247-1257

Plasmodium spp., antimalarial activity of 2,4-diamino-6-(2-naphthylsulfonyl)quinazoline and related 2,4-diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines

Chloroquine

Ferraroni, J. J.; and Hayes, J., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 909-911

Plasmodium falciparum outbreak among indigenous indian tribe, 3 cases resistant to chloroquine responded favorably to fansidar therapy: Uauaris, Territory of Roraima, Brazil

Chloroquine

Ferraroni, J. J.; Waki, S.; and Suzuki, M., 1977, Acta Amazonica, v. 7 (1), 147-148

Plasmodium falciparum, humans, chloroquine resistance, confirmed in vitro: Manaus, Amazonas

Chloroquine -- Continued.

Chloroquine diphosphate

Field, R. C.; et al., 1978, Brit. J. Pharmacol., v. 62 (2), 159-164
effects of chloroquine, primaquine and ethidium on precursor incorporation into DNA, RNA and protein in mammalian tissues

Chloroquine

Fitch, C. D.; Chan, R. L.; and Chevli, R., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 258-262

Plasmodium berghei, undiminished mefloquine accumulation by erythrocytes infected with chloroquine-resistant strain provides explanation for superiority of mefloquine in treating chloroquine-resistant malaria, but competition observed between chloroquine and mefloquine raises possibility that same process of accumulation serves both drugs

Chloroquine

Fitch, C. D.; Chevli, R.; and Gonzalez, Y., 1974, Antimicrob. Agents and Chemotherapy, v. 6 (6), 757-762

Plasmodium falciparum, *Aotus trivirgatus* erythrocytes infected with chloroquine-susceptible vs. chloroquine-resistant strain, effect of substrate (glucose) on chloroquine and amodiaquin accumulation

Chloroquine

Fitch, C. D.; Ng, R. C. K.; and Chevli, R., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (2), 185-193

Plasmodium berghei, evidence that erythrocyte surface components determine affinity with which chloroquine is accumulated and thereby determine whether or not the malaria parasite will be susceptible to the drug

Chloroquine

Fletcher, K. A.; and Sarikabhuti, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (5), 489-490

Plasmodium berghei, mice, anti-plasmodial activity of chloroquine does not appear to be associated with inhibition of erythrocytic glucose-6-phosphate dehydrogenase

Chloroquine

Fogh, S.; Jepsen, S.; and Effersøe, P., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (2), 228-229

Plasmodium falciparum, chloroquine resistant strain in non-immune male, treatment with fansidar terminated infection: Danish tourist to Kenya

Chloroquine (Resochin; Aralen; Avloclor; Nivaquin)

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98

Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Chloroquine diphosphate

Godoy, G. A.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (1), 38-48

Plasmodium falciparum, humans, grade I chloroquine resistant strains: Bolivar State, Venezuela

Chloroquine -- Continued.

Chloroquine

Golenser, J.; et al., 1978, Israel J. Med. Sc., v. 14 (5), 606-610

Plasmodium berghei, immunization of chloroquinized rats against sporozoites by bites of infected mosquitoes: influence of number of exposures to infected mosquitoes on antibody titers and protection; influence of exposure to different numbers of infective mosquitoes on antibody production and protection; specificity of antiplasmodial antibodies; influence of passive transfer of sera from rats immune to sporozoites or to erythrocytic forms on development of sporozoites, symposium presentation

Chloroquine

Gupta, O. K.; and Banerjee, R. N., 1976, Ann. Soc. Belge Med. Trop., v. 56 (3), 151-157

Plasmodium vivax, humans, possible association of disseminated intravascular coagulation and hyperuricaemia, improvement after chloroquine therapy

Chloroquine

Horrobin, D. F.; and Karmali, R. A., 1979, Nature, London (5740), v. 282, 758

effect of chloroquine on Epstein-Barr virus expression, response to comment of Lenoir, G.; and Geser, A., 1979, Nature, London (5740), v. 282, 758

Contramibial

Ingelet, B., 1974, Rev. Infirm. Afrique Noire (26), 17-19

human amoebic colitis, clinical trials testing contramibial given orally over a 4-day period, good therapeutic results

Chloroquine

Jones, R. L.; Davidson, M. W.; and Wilson, W. D., 1979, Biochim. et Biophys. Acta, v. 561 (1), 77-84

chloroquine does not bind to DNA by classical intercalation mechanism typical of quinacrine and ethidium

Chloroquine

Karmali, R. A.; et al., 1978, Nature, London (5679), v. 275, 444-445

chloroquine enhances Epstein-Barr virus expression and may thus play important part in development of African Burkitt's lymphoma

Chloroquine phosphate

Kean, B. H., 1979, J. Am. Med. Ass., v. 241 (4), 395

Plasmodium falciparum, chloroquine-resistance, 2 case reports of people having travelled in Africa

Chloroquine

Khan, A. A.; and Maguire, M. J., 1978, Brit. Med. J. (6128), v. 1, 1669-1670

P[lasmodium] falciparum, chloroquine-resistant strain reported in young child, therapy with doxycycline + chloroquine resulted in cure: Zambia

Chloroquine -- Continued.

Chloroquine + Doxycycline

Khan, A. A.; and Maguire, M. J., 1978, Brit. Med. J. (6128), v. 1, 1669-1670
P[lasmodium] falciparum, chloroquine-resistant strain reported in young child, therapy with doxycycline + chloroquine resulted in cure: Zambia

Chloroquine

Koehler, P.; and Bachmann, R., 1978, Molec. Pharm., v. 14 (1), 155-163
Ascaris suum muscle tissue, comparison of effects of levamisole, thiabendazole, chloroquine, and praziquantel on electron transport in Ascaris muscle submitochondrial particles

Chloroquine phosphate (Malaquin)

Kumar, B. J.; and Gowda, D. K. V., 1978, Indian Vet. J., v. 55 (9), 729-730
theileriasis, dairy cow treated with chloroquine and oxytetracycline, chloroquine toxicity causing corneal opacity and possibly abortion

Chloroquine

Langreth, S. G.; Nguyen-Dinh, P.; and Trager, W., 1978, Exper. Parasitol., v. 46 (2), 235-238
Plasmodium falciparum, fine structure of merozoite invasion of human erythrocytes in vitro, successful invasion after 3 hr in presence of concentration of chloroquine harmful to feeding stages

Chloroquine

Lenoir, G.; and Geser, A., 1979, Nature, London (5740), v. 282, 758
effect of chloroquine on Epstein-Barr virus expression and on development of Burkitt's lymphoma, comment on Karmali, R. A.; et al., 1978, Nature, London (5679), v. 275, 444-445

Chloroquine

Lopez Antunano, F. J.; and Wernsdorfer, W. H., 1979, Bull. World Health Organ., v. 57 (4), 663-665
chloroquine-resistant Plasmodium falciparum, in vitro response to mefloquine, microtechnique system

Chloroquine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Chloroquine sulfate

Magzoub, M., 1972, Sudan Med. J., v. 10 (3), 145-148
Schistosoma mansoni adult worms removed from mice treated with chloroquine showed reduced exogenous glucose uptake, increased lactic acid production and reduced motility

Chloroquine diphosphate

Marshall, R. J.; and Ojewole, J. A. O., 1978, Toxicol. and Applied Pharm., v. 46 (3), 759-768
quinoline and nonquinoline antimalarial drugs, effects on isolated guinea pig cardiac muscle

Chloroquine -- Continued.

Chloroquine

Maurois, P.; et al., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 491-493
Plasmodium vivax in 33-year-old non-immune patient being treated with chloroquine, changes in serum lipoproteins

Chloroquine

Merchant, J. R.; and Engineer, A. B., 1978, Current Sc., Bangalore, v. 47 (21), 803-804 [Letter]
simplified procedure for synthesis of 7-chloro-2,3-dihydro-4(1H)-quinolone, important intermediate of chloroquine

Chloroquine

Metge, P.; et al., 1979, Bull. Soc. Ophth. France, v. 79 (4-5), 347-351
malaria, humans, chloroquine induced retinopathy, 6 cases

Chloroquine

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, Israel J. Med. Sc., v. 14 (5), 601-605
Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiparasite immunity by natural bites of infected mosquitoes, symposium presentation

Chloroquine + Primaquine

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, Israel J. Med. Sc., v. 14 (5), 601-605
Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiparasite immunity by natural bites of infected mosquitoes, symposium presentation

Chloroquin diphosphate

Mishra, A. K.; and Sharma, N. N., 1979, Trop. Animal Health and Prod., v. 11 (4), 222-226
Anaplasma marginale, calves (exper.), comparative efficacy of several drugs

Chloroquine

Nagaratnam, N.; Chetiyawardana, A. D.; and Rajiyah, S., 1978, Postgrad. Med. J., London (628), v. 54, 108-112
aplastic anaemia and acute myeloblastic leukemia following chloroquine therapy for malaria and discoid lupus erythematosus, case reports

Chloroquine

Neves, J.; and Moura, H. B., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (2), 126-130
Plasmodium falciparum, P. vivax, prevalence survey in hospital patients, discussion of changes in prevalence with introduction of chloroquine resistant strains of P. falciparum, treatment trials with various malarial drugs: Brazil

Chloroquine

Nguyen-Dinh, P.; and Trager, W., 1978, Science (4348), v. 200, 1397-1398
Plasmodium falciparum, African strain, production of chloroquine resistance in vitro

Chloroquine -- Continued.

Chloroquine

Ogawa, S.; et al., 1979, *Lancet*, London (8131), v. 1, 1408 [Letter]
progression of retinopathy long after cessation of chloroquine therapy

Chloroquine

Olaeta Elizalde, R., 1973, *Rev. Cir. Hosp. Juarez*, Mexico (187-188), v. 44, 1972-1973, 59-64
human hepatic amoebic abscess, therapeutic recommendations (emetine, dehydroemetine, chloroquine, metronidazole)

Chloroquine

Olivares Lopez, F.; et al., 1977, *SPM Salud Pub. Mexico*, v. 19 (5), 701-707
Plasmodium malariae, *P. vivax*, survey of blood transfusion-induced infections, diagnostic problems, chloroquine therapy, problems in blood donor control: Mexico

Chloroquine diphosphate

Omer, A. H. S., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (5), 853-857
Plasmodium falciparum, human, clinical trial of response to chloroquine, little evidence of resistance: Gezira and Bor areas, Sudan

Chloroquine diphosphate

Omer, A. H. S., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (5), 858-863
Plasmodium spp., human, prevalence by parasite species and by host age group, dramatic response to mass chemoprophylaxis with chloroquine: Gezira and Bor regions, Sudan

Chloroquine

Pena Yanez, A.; et al., 1972, *Rev. Clin. Espan.*, v. 124 (4), 405-412
pulmonary amoebiasis in man without involvement of liver, diagnosis after expectoration of chocolate colored material, successful therapy of emetine and chloroquine: Spain (had resided previously in Algeria)

Chloroquine

Pillay, N.; and Bhoola, R. L., 1975, *South African Med. J.*, v. 49 (35), 1443-1444
Plasmodium falciparum, woman, case report, probable resistance to chloroquine, successfully treated with quinine: South Africa (had recently returned from Mozambique)

Chloroquine

Ponnampalam, J. T., 1978, *J. Trop. Med. and Hyg.*, v. 81 (10), 198-203
human malarias, comparative study of prophylaxis using chloroquine and a combination of sulfadoxine and pyrimethamine: residents of rubber estate in central Malaysia

Chloroquine

Powers, K. G.; and Jacobs, R. L., 1972, *Antimicrob. Agents and Chemotherapy*, v. 1 (1), 49-53
Plasmodium falciparum, chloroquine-resistant strain in *Aotus trivirgatus*, 2 chlorinated lincomycin analogues cured blood-induced infections

Chloroquine -- Continued.

Chloroquine + Pyrimethamine

du Preez, O.; and Cockcroft, C. R., 1975, *South African Med. J.*, v. 49 (34), 1407-1408
malaria prophylaxis trials, army personnel camping in endemic area, 3 drug combinations, no infections reported in trial groups while local population acting as control reported 250 falciparum cases: Caprivi Strip, South Africa

Chloroquine (Nivaquine 200)

Price Evans, D. A.; Fletcher, K. A.; and Baty, J. D., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (1), 11-17
urinary excretion of chloroquine in different ethnic groups, study of healthy volunteers

Chloroquine (WR 1,544)

Rane, D. S.; and Kinnamon, K. E., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (6), 937-947
sporozoite-induced *Plasmodium berghei* in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Chloroquine

Ray, A. P.; Parkinson, A. D.; and Black, R. H., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (1), 19-22
Plasmodium berghei in chloroquine resistant white mice, results of treatment with combinations of proguanil and dapsone

Chloroquine sulphate

Richards, W. H. G.; and Maples, B. K., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (2), 99-108
Plasmodium falciparum in continuous culture, effects of pyrimethamine and chloroquine on parasite growth and viability

Chloroquine

Rosario, V. E.; et al., 1978, *Lancet*, London (8057), v. 1, 185-187
Plasmodium chabaudi, infection of mice with mixtures of drug-resistant (pyrimethamine or chloroquine) and drug sensitive strains, resulting infections were maintained in absence of drugs with some persistence of resistant forms over sensitive forms

Chloroquine base

Roy, R. G., 1978, *Indian J. Med. Research*, v. 68, 741-743
[Plasmodium] vivax, *P. falciparum*, humans, presumptive treatment with 600 mg. chloroquine base, good response, no RII or RIII type chloroquine resistance seen: Dharmapuri and North Arcot districts, Tamil Nadu State

Chloroquine phosphate

Ruebush, T. K. II; et al., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (2), 184-189
Babesia microti, 65-year-old man, treated with diminazene aceturate after failure to respond to chloroquine therapy, development of acute idiopathic polyneuritis: Nantucket Island

Chloroquine -- Continued.

Chloroquine

Rumans, L. W.; Dennis, D. T.; and Atmosoedjono, S., 1979, *Lancet*, London (8142), v. 2, 580-581 [Letter]
Plasmodium falciparum, fansidar-resistant malaria in case also resistant to chloroquine: Indonesia

Chloroquin phosphate + Diiodohydroxy quinaline + Chloquinat (=Resotren [composite])

Samaddar, J.; Biswas, D. N.; and Ghose, A. N., 1978, *Indian Vet. J.*, v. 55 (7), 572-576
 leucocytozoonosis, W[hite] L[eg] H[orn] birds, quinine bisulphate and resotren failed to ensure absolute recovery, though general condition of treated birds improved

Chloroquine

Schmidt, L. H., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (4), 703-717
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus griseimembra*, responses of established infections to chloroquine, quinine, and pyrimethamine

Chloroquine

Schmidt, L. H., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (5), 793-807
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus griseimembra*, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

Chloroquine

Schmidt, L. H.; et al., 1977, *Antimicrob. Agents and Chemotherapy*, v. 11 (5), 826-843
Plasmodium falciparum in *Aotus trivirgatus*, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

Chloroquine

Sfikakis, P.; et al., 1971, *Therapeutique*, v. 47 (4), 383-385
 chloroquine administered to normal subjects, no cardiotoxicity observed

Chloroquine (base)

Shanmugham, C. A. K.; et al., 1978, *Indian J. Med. Research*, v. 67, 926-928
Plasmodium falciparum, clinical trials with chloroquine base: Tamil Nadu

Chloroquine (Delagil)

Shedivtsova, A., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Chloroquine

Shirley, J. A.; Eykyn, S. J.; and Pearson, T. C., 1979, *Brit. Med. J.* (6194), v. 2, 834-835
Plasmodium falciparum, occurrence of chloroquine resistant infection in Bangladeshi girl with acute lymphoblastic leukaemia

Chloroquine -- Continued.

Chloroquine phosphate

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, *Zentralbl. Bakteriol.*, 1 Abt. Orig., Reihe A, v. 241 (3), 358-367
Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Chloroquine phosphate

Stoskopf, M. K.; and Beier, J., 1979, *J. Am. Vet. Med. Ass.*, v. 175 (9), 944-947
Plasmodium relictum, *P. elongatum* in *Spheniscus demersus*, diagnostic methods evaluated, chloroquine phosphate and primaquine phosphate therapy: Baltimore Zoo

Chloroquine (Aralen; Nivaquine)

Trojan, H. J., 1975, *Rev. Internat. Trachome et Path. Ocul. Trop. et Subtrop.*, v. 52 (3-4), 129-137
 ocular pathology associated with long-term use of chloroquine

Chloroquine

Ugarte, G.; Apt, W.; and Faiguenbaum, J., 1977, *Rev. Med. Chile*, v. 105 (3), 176-178
Plasmodium vivax infection in man thought to have hepatic amoebiasis because of complaints of jaundice, fever, and hepatomegaly, after blood smears revealed evidence of malaria man was cured with chloroquine and daraprim: Chile, had made recent visit to Brazil

Chloroquine

Van Poucke, G., 1979, *East African Med. J.*, v. 56 (4), 158-162
P[lasmodium] falciparum, humans, intramuscular treatment with chloroquine vs. quinimax: East Africa

Chloroquine (Delagil)

Varnai, F.; and Ecker, A., 1977, *Therap. Hungar.*, v. 25 (4), 131-133
 malaria, humans travelling to endemic areas, drug prophylaxis, comparative study, least unwanted side effects and lowest morbidity rate recorded with pyrimethamine: Hungary

Chloroquine phosphate

Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Chloroquine

Ward, W. F.; Lipschutz, F. I.; and Hahn, E. W., 1975, *Thromb. et Diath. Haemorrh.*, v. 34 (2), 545-547 [Letter]
 accelerated hemostasis in chloroquine-treated rats

Chloroquine

Warhurst, D. C.; and Thomas, S. C., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (3), 203-211
Plasmodium berghei, effect of some metabolic inhibitors upon chloroquine-induced pigment clumping

Chloroquine base. See Chloroquine.

Chloroquine diphosphate. See Chloroquine.

Chloroquine phosphate. See Chloroquine.

Chloroquine sulfate. See Chloroquine.

4-[(7-Chloro-4-quinolyloxy)amino]- α -(diethylamino)-
o-cresol. See Amodiaquine.

Chlorothymol

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128

Echinococcus multilocularis, various anthelmintics, scolical effects in vitro and/or therapeutic effects in mice

(4-Chloro-o-toloxyl) acetic acid -- Chwastox.

Chwastox

Moczon, T., 1976, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (5), 289-292

Fasciola hepatica miracidia, inhibitory effect of pesticides on enzyme activity

1-(3-Chloro-p-tolyl) piperazine hydrochloride -- Mirasan.

Mirasan

Foster, R., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 1-9

Schistosoma mansoni, *S. haematobium*, *S. japonicum*, oxamniquine, efficacy and toxicity in various exper. animals, clinical trials in humans

Mirasan

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46

Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

2-Chloro-1-(2,4,5-trichlorophenyl) vinyl diethylphosphate -- Shell SD-8448.

Shell SD-8448

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate. See Tetrachlorvinphos.

Chlorotrimazole. See Clotrimazole.

N'-(3-Chloro-2,4,6-trimethylphenyl)-N,N-diethylenediamine hydrochloride -- Hoechst S-616.

Hoechst S-616

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46

Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

p-Chloro-m-xylene

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Chlorphenamidin -- Chlordimeform.

Chlordimeform

Dawkins, C. C.; and Gladney, W. J., 1978, J. Econom. Entom., v. 71 (4), 657-660

3-host ticks, guinea pigs, amitraz, Upjohn U-42,564, chlordimeform, detachment response and mortality

Chlordimeform

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144

Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Chlordimeform

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960

Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Chlorphenoxamide -- Amoebicide 2004 (with Tinidazole); Chlorphenoxamide; Ethophamide; N-(beta-Ethoxy-ethyl)-N-[p-phenoxy-(4'-nitro)-benzyl]-dichloroacetamide; Ethylchlordifene; Etofamide; Kitnos.

Chlorophenoxamide

Campos, R., 1973, Rev. Brasil. Clin. e Terap., v. 2 (11), 587-588

Entamoeba histolytica, comparison of efficacy of nifuratel and other amoebicides using material cultured from intestinal ulcers of patient with intestinal symptomatic amoebiasis

Amoebicide 2004

Ferreira Tolsa, O., 1978, Semana Med. Mexico (1203), an. 25, v. 95 (4), 85-88

Entamoeba histolytica, school children, mass therapy with amoebicide 2004, well tolerated, good results

Etofamide (Kitnos)

Huggins, D.; and Maciel, M. F., 1975, Rev. Soc. Brasil. Med. Trop., v. 9 (2), 69-72

Entamoeba histolytica, children with chronic intestinal infection, clinical trials testing the efficacy of etofamide

Chlorphenoxamide -- Continued.

Etofamide

- Levi, G. C.; Amato Neto, V.; and Konichi, S. R., 1973, Rev. Soc. Brasil. Med. Trop., v. 7 (6), 335-339
Entamoeba histolytica, humans with asymptomatic, mild or chronic infections, clinical trials testing etofamide in varying dosages and time schedules: state of Sao Paulo

Etofamide (Kitnos)

- Otero, N. B.; et al., 1973, Rev. Brasil. Clin. e Terap., v. 2 (11), 581-586
 human chronic intestinal amoebiasis, clinical trials evaluating etofamide as therapy, 90% cure obtained; no liver damage or other toxic reactions

Chlorproguanil (Lapudrine)

- Brohult, J.; et al., 1979, Ann. Trop. Med. and Parasitol., v. 73 (4), 327-331
 malaria, humans, analysis of chemoprophylactic habits and reasons for breakdowns in therapy, small mining town: Yekepa, Liberia

Chlorpromazine (Largactil) + Berenil

- Gretillat, S., 1978, Bull. Acad. Vet. France, n.s., v. 51 (3), 351-358
Haemobartonella sp., horses, symptoms, diagnosis, pathology, treatment with chlorpromazine + berenil: Niger

Chlorpyrifos -- Chlorpyrifos; O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate; Dow M3615; Dow M3983; Dursban; Dursban 44 Insecticide Formulation; Lorsban; Ridlice.

Chlorpyrifos

- Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765
Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control

Chlorpyrifos + Ronnel

- Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765
Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control

Dursban

- Camoens, J. K., 1977, Malaysian Vet. J., v. 6 (3), 111-124
Boophilus microplus, dairy cattle, various control measures discussed but spraying acaricides on pastures shows particular promise: Air Hitam, Johor, Malaysia

Chlorpyrifos

- Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439
 economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Chlorpyrifos -- Continued.

Chlorpyrifos

- Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Chlorpyrifos

- Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Dursban

- Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105
Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Chlorpyrifos

- Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73
Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Chlorpyrifos

- Ivey, M. C., 1979, J. Econom. Entom., v. 72 (6), 909-911
 chlorpyrifos and its metabolite 3,5,6-trichloro-2-pyridinol, residues in body tissues of cattle wearing impregnated plastic ear tags

Chlorpyrifos

- Ivey, M. C.; et al., 1972, J. Econom. Entom., v. 65 (6), 1647-1649
 chlorpyrifos and oxygen analogue, residues in body tissues of dipped cattle

Chlorpyrifos (Dursban 44 Insecticide Formulation)

- Ivey, M. C.; and Palmer, J. S., 1979, J. Econom. Entom., v. 72 (6), 837-838
 chlorpyrifos and its metabolite 3,5,6-trichloro-2-pyridinol, residues in swine after pour-on application for control of *Haematopinus suis* and *Sarcoptes scabiei*

Chlorpyrifos

- Ivey, M. C.; Palmer, J. S.; and Hooten, E. C., 1978, J. Econom. Entom., v. 71 (4), 697-700
 residues of chlorpyrifos and its metabolite 3,5,6-trichloro-2-pyridinol (pyridinol) were found in body tissues of Hereford yearlings after wearing chlorpyrifos-impregnated ear bands

Chlorpyrifos (Ridlice)

- Kettle, P. R.; and Lukies, J. M., 1979, N. Zealand Vet. J., v. 27 (4), 78-79
Linognathus vituli, cattle, pour-on formulations of phosmet, methidathion, chlorpyrifos, and temephos: Kaitoke, near Upper Hutt, New Zealand

Chlorpyrifos -- Continued.

Dursban (Chlorpyrifos)

Khan, M. H.; and Srivastava, S. C., 1977, Indian J. Animal Health, v. 16 (2), 137-140
Boophilus microplus engorged females, in vitro tests with dursban, gamma BHC, sumithion, supona, dimecron, egg production and viability; supona most effective

Chlorpyrifos (Dursban; Lorsban)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Chlorpyrifos (Dursban; Dow M3615; Dow M3983)

Mount, G. A.; et al., 1978, J. Econom. Entom., v. 71 (1), 27-28
Eutrombicula alfreddugesi, area control, chlorpyrifos, granules and concentrated sprays compared: Gainesville, Florida

Chlorpyrifos (Dursban)

Oba, M. S. P.; de Campos, M. S.; and de Almeida, M. A., 1977, Biologico, S. Paulo, v. 43 (9-10), 218-220
Menacanthus stramineus, Megninia cubitalis, M. gynlimura, chickens and chicken-houses, chlorpyrifos entirely satisfactory

Chlorpyrifos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Chlorpyrifos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Dursban

Yeoman, G. H.; and Bell, T. A., 1978, Vet. Rec., v. 103 (15), 337
Lucilia sericata, sheep, aluminium alkoxide gellants mixed with insecticide and applied to breech area, results suggest that this new control method against cutaneous myiasis gives higher protection than current means of control with no abnormal problems of toxicity, tissue residues, or wool processing

Chlorpyrifos methyl

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Chlorpyrifos. See Chlorpyrifos.Chlorquinaldol -- Chlorchinaldol; Sterosan.

Chlorchinaldol (Sterosan)

Lovgren, T.; and Salmela, I., 1978, Acta Path. et Microbiol. Scand., v. 86B (3), 155-158
Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents

Chlortetracycline -- Aureomycin; Biomitsin; Biomycin; Biovetin; Chlortetracycline hydrochloride.

Chlortetracycline

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Biomycin (Chlortetracyclin)

Evplov, N. N.; and Nazarov, V. G., 1977, Veterinariia, Moskva (6), 65-66
Eimeria spp., calves, chemococcide effective; compared with biomycin and norsulfazol: Belgorodsk oblast

Chlortetracycline

Gill, B. S.; et al., 1978, Internat. J. Parasitol., v. 8 (6), 467-469
Theileria annulata, calves, immunization by treating tick (Hyalomma anatolicum anatolicum) stabilate-induced infections with 1 or 2 doses of long-acting oxytetracycline vs. 8 doses of chlortetracycline

Biovetin

Gobzem, V. R., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 118-120
E[imeria] bovis, E. cylindrica, E. ellipsoidalis, calves, safe prophylaxis with biovetin

Biomycin

Gobzem, V. R.; and Nazarov, V. G., 1978, Veterinariia, Moskva (3), 67-69
Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Aureomycin

Mishra, A. K.; and Sharma, N. N., 1979, Trop. Animal Health and Prod., v. 11 (4), 222-226
Anaplasma marginale, calves (exper.), comparative efficacy of several drugs

Chlortetracycline

Ohshima, S.; Hoshino, M.; and Tanaka, H., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (3), 127-131
Toxoplasma, 11 strains, susceptibility to 6 drugs, mice

Chlortetracycline

Rees, R. G. P.; and Shelley, A. J., 1977, Acta Amazonica, v. 7 (1), 47-49
Balantidium coli, Yanomama-Indian, serious dysentery, case report, Ascaris lumbricoides and Trichuris trichiura also present, chlortetracycline, improved condition: Tootobi, norte do Estado do Amazonas, Brasil

Aureomycin

Samizadeh-Yazd, A.; et al., 1979, Am. J. Vet. Research, v. 40 (8), 1107-1109
Eimeria spp., lambs, efficacy of monensin and aureomycin separately and combined

Aureomycin

Sharma, M. M.; et al., 1979, Indian Vet. Med. J., v. 3 (3), 187-191
theileriasis, Jersey cattle, incidence and treatment with berenil, babesan, nevaquine, and aureomycin: Exotic Nucleus Cattle Farm, Bassi, Jaipur

Chlortetracycline -- Continued.

Chlortetracyclin (Aureomycin)
Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriol., 1 Abt. Orig., Reihe A, v. 241 (3), 358-367
Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Aureomycin (Chlortetracycline)

Sinha, R. P.; and Dubey, R. K., 1978, Indian Vet. J., v. 55 (5), 372-376

Anaplasma marginale outbreak in non-preimmune Jersey cattle imported from United States and Denmark to farm in Bihar, clinical symptoms and pathological findings, epizootiological factors responsible for outbreak (high ambient temperature, stress of vaccination for rinderpest virus, presence of tick vectors), control achieved through chemotherapy of sick and healthy animals, removal of vectors, and housing in cool sheds; outbreaks in exotic herds could be avoided if cattle were imported in early winter: India

Chlortetracycline (Aureomycin)

Sweet, V. H.; and Stauber, E. H., 1978, J. Am. Vet. Med. Ass., v. 172 (11), 1310-1312

Anaplasma marginale, cattle, serologic survey, chlortetracycline, oxytetracycline, good results, concluded that test and treatment program would need to be continued for several years to achieve anaplasmosis-free status: northern Idaho; southeastern Washington

Chlortetracycline

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Chlortetracycline hydrochloride. See Chlortetracycline.

Chloxy1. See 1,4-Bis(trichloromethyl) benzene.

Chondroitin sulfate

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Chromomycin A₃ -- Toyomycin.

Toyomycin

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Chromomycin A₃

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Chwastox. See (4-Chloro-o-toloxyl) acetic acid.

Cibacron blue F3GA

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cibacron brilliant blue BRP

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Ciclobendazole. See Cyclobendazole.

Cidial. See Phenthoate.

Ciplin. See Sulfamethoxazole or Trimethoprim.

Citarin. See Tetramisole.

Citarin-L. See Tetramisole.

Citarin-L spot on. See Tetramisole.

Citrazine. See Piperazine.

Citric acid

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230

Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Citromycin chloride

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Clendrol

Pott, J. M.; and Riley, C. J., 1979, Vet. Rec., v. 104 (25), 579

Otodectes cynotis, dogs, cats (ear canals of both), canaural, comparison with proprietary topical ear preparation (clendrol), controlled trial

Cleocin phosphate. See Clindamycin.

Clindamycin -- 7-Chlorolincomycin; Clindamycin hydrochloride; Clindamycin phosphate; Dalacin-C; U-21; Cleocin phosphate.

Clindamycin

Burke, G. J.; and Mills, A. E., 1979, South African Med. J., v. 55 (5), 156 [Letter] toxoplasmosis, Black Rhodesian youth, unusual presentation with cardiac arrhythmia and low cholinesterase levels, clindamycin

Clindamycin HCl

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Clindamycin

Koontz, L. C.; et al., 1979, Exper. Parasitol., v. 48(2), 206-212

Plasmodium berghei, mice infected with clindamycin-resistant parasites, uptake of clindamycin and its metabolites by erythrocytes, impaired uptake is not mechanism of resistance

Clindamycin

Ohshima, S.; Hoshino, M.; and Tanaka, H., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (3), 127-131

Toxoplasma, 11 strains, susceptibility to 6 drugs, mice

Clindamycin hydrochloride (U-21)

Powers, K. G.; and Jacobs, R. L., 1972, Antimicrob. Agents and Chemotherapy, v. 1 (1), 49-53

Plasmodium falciparum, chloroquine-resistant strain in *Aotus trivirgatus*, 2 chlorinated lincomycin analogues cured blood-induced infections

Clindamycin -- Continued.

Clindamycin phosphate

Tabbara, K. F.; et al., 1979, Arch. Opth., Chicago, v. 97 (3), 542-544

Toxoplasma gondii, rabbits (eye), clindamycin phosphate

Clindamycin hydrochloride (Dalacin-C)

Thiermann, E.; et al., 1977, Rev. Med. Chile, v. 105 (7), 433-435

Toxoplasma gondii, mice, experimental trials comparing efficacy of clindamycin with that of pyrimethamine combined with sulfamethoxy-pyridazine, combination drug cured 100% of mice while mice treated with clindamycin survived during treatment but 50% died from severe infections after therapy had been discontinued

Clindamycin + Sulfamethoxypyridazine

Thiermann, E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens compared during acute and late infections; pyrimethamine + sulfamethoxypyridazine was most effective

Clindamycin hydrochloride. See Clindamycin.

Clindamycin phosphate. See Clindamycin.

Clioquinol. See Iodochlorhydroxyquin.

Clioaxanide -- 4'-Chloro-3,5-diiodosalicylanilide; 4'-Chloro-3,5-diiodosalicylanilide acetate ester; N-(4'-Chlorophenyl)-3,5-diiodoacetyl-salicylamide; Tremerad.

Clioaxanide

Douch, P. G. C., 1979, Xenobiotica, v. 9 (4), 263-268

Moniezia expansa, *Ascaris suum*, metabolism of clioaxanide and resorantel and related compounds

4'-Chloro-3,5-diiodosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

4'-Chloro-3,5-diiodosalicylanilide acetate ester

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Clioaxanide (Tremerad)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]

Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiabendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxylin are also useful chemical alternatives

Clofazimine -- Ciba-Geigy B 663; Lampren.

Lampren

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Clofibrate

McQuiston, T. E., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (1), 12-14

Plasmodium berghei, mice acclimated to 22°C or 5°C before infection, some treated with clofibrate and some briefly exposed to -35°C after infection, parasitemia and plasma free fatty acid levels

Clont. See Metronidazole.

Clopidol. See Meticlorpindol.

Clopidol. See Meticlorpindol.

Clotiamine. See Beclotiamine.

Clotrimazole -- Bay b 5097; Bis-phenyl-(2-chlorophenyl)-1-imidazolyl-methane; Canesten; Chlorotrimazole; Meclo (with Metronidazole).

Chlorotrimazole

Berghella, A., 1973, *Quad. Clin. Ostet. e Ginec.*, v. 28 (5-12), 155-168

Trichomonas vaginalis, human vaginal trichomoniasis, topically applied mepartricin compared favorably with chlorotrimazole used as drug standard in clinical trials

Clotrimazole

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Clotrimazole (Canesten; Bay B 5097)

Facchini, V., 1974, *Riv. Ital. Ginec.*, v. 55 (6), 485-491

Trichomonas, human vulvo-vaginitis, canesten vaginal tablets or cream, trials of therapeutic efficacy

Clotrimazole (Canesten)

Falcone, I.; Wanick, M. C.; and do Nascimento, R., 1975, *Rev. Brasil. Clin. e Terap.*, v. 4 (6), 229-232

human vaginal trichomoniasis, use of clotrimazole to treat pregnant women, drug efficacy of over 93%

Bay b 5097

Gittler Lajchter, S.; and Bernal Carrasco, E., 1974, *Semana Med. Mexico* (1027), an. 21, v. 81 (3), 69-76

Trichomonas vaginalis, pregnant and non-pregnant women and young girls, trichomonal vaginal infections or mixed *Candida* or bacterial infections, clinical trials with Bay b 5097, effective

Clotrimazole -- Continued.

Clotrimazole

Imparato, E.; et al., 1976, *Quad. Clin. Ostet. e Ginec.*, v. 31 (4), 225-239

Trichomonas vaginalis, human vaginal trichomoniasis, evaluation of mepartricin as oral therapy using nimorazole and clotrimazole as reference drugs, best results obtained with mepartricin

Canesten

Lecca, U.; and Canfora, S., 1974, *Riv. Ostet. e Ginec. Prat. e Med. Perinat.*, v. 54 (1), 1973/1974, 77-82

Trichomonas, human cervico-vaginal infections, canesten therapy

Clotrimazole (Canesten)

Lovgren, T.; and Salmela, I., 1978, *Acta Path. et Microbiol. Scand.*, v. 86B (3), 155-158

Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents

Clotrimazole

Nagington, J., 1975, *Tr. Ophth. Soc. United Kingdom*, v. 95 (2), 207-209

Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity

Canesten

Pecori, M.; Vigorito, A.; and Celentano, C., 1974, *Rassegna Internaz. Clin. e Terap.*, Napoli, v. 54 (6), 355-359

Trichomonas vaginalis, human, vaginitis, successful therapy with canesten vaginal tablets

Clotrimazole (Canesten)

Ragni, N.; and Foglia, G., 1974, *Riv. Ostet. e Ginec. Prat. e Med. Perinat.*, v. 54 (7), 1973/1974, 342-350

Trichomonas, human cervico-vaginal infections and mixed fungal infections, therapy with clotrimazole

Canesten (Clotrimazole)

Resch, B.; Altmayer, P.; and Bartfai, G., 1978, *Therap. Hungar.*, v. 26 (4), 185-187

Trichomonas vaginalis, women, single or mixed *Candida* infections, canesten

Canesten (Bay b-5097)

Riccobono, G., 1972, *Riv. Ostet. e Ginec. Prat. e Med. Perinat.*, v. 53 (10), 489-498

Trichomonas vaginalis, human vaginal and cervical infections, canesten topical therapy

Clotrimazole (Canesten)

Saumann, A.; et al., 1975, *Rev. Chilena Obst. y Ginec.*, v. 40 (6), 378-382

human *Trichomonas vaginalis* vaginitis, clinical trials using local applications or suppositories of clotrimazole: Chile

Canesten (Clotrimazole)

Varga, A., 1978, *Therap. Hungar.*, v. 26 (1), 40-42

Trichomonas vaginalis, women, vulvovaginitis, canesten tablets and ointment

Clotrimazole -- Continued.

- Clotrimazole + Metronidazole (= Meclo vaginal cream or suppositories)
Zacutti, A., 1975, *Quad. Clin. Ostet. e Ginec.*, v. 30 (2), 65-96
Trichomonas vaginalis, clinical trials comparing metronidazole and clotrimazole alone or in new combination drug (meclo) as vaginal creams, vaginal suppositories and as oral tablets, pregnant and non-pregnant women and their sexual partners, good clinical results both in prevention and cure of infections

Cobalt protoporphyrin

- Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism

Coban. See Monensin.COBAN 45. See Monensin.Cocciden. See Beclotiamine.Coccidin. See Dinitolmide.Cocciostats. See Anticoccidials.Cocciovit. See Amprolium.Codrinal. See Tetracycline or β -Toluenesulfonyl- β -methoxy-ethyl urethane sodium.Coformycin -- 3- β -D-Ribofuranosyl-6,7,8-trihydroimidazo[3,4-d][1,3]diazepin-8-(R)-ol.

Coformycin

- Senft, A. W.; and Crabtree, G. W., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Colchicine

- Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Colchicine

- Injeyan, H.; Huebner, E.; and Meerovitch, E., 1979, *J. Protozool.*, v. 26 (2), 253-259
Entamoeba sp. (Laredo isolate), morphologically distinct colchicine-resistant variant, properties compared to those of parent strain

Colchicine

- Sinden, R. E.; and Smalley, M. E., 1979, *Parasitology*, v. 79 (2), 277-296
Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

Colchicine

- Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Colistin sulfate

- Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Combantrin. See Pyrantel.ComBot. See Trichlorfon.Comboto. See Trichlorfon.Compound 4072. See Chlorfenvinphos.Compound 16,842. See Bitoscanate.

Compound 'E'. See (Diamino-4,6-triazinyl-1,3,5-amino-2)-4-phenyl-arsino di (D-thio-3-amino-2-methyl-3-butyric acid) or 2-[2-(4-Hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazyl) benzimidazole.

Concurat. See Tetramisole.Contramibial. See Chloroquine or Diiodohydroxyquin or Tetracycline.Coopers Summer Drench. See Haloxon.Cooper-Tox. See Toxaphene.Coopertox [of Atef, M.; and El-Say, A., 1976]. See Toxaphene.Coopertox [of Grillo Torrado, M. M.; and Perez Arrieta, A., 1977]. See Ethion.

Copper hematoporphyrin D

Meshnick, S. R.; et al., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Copper-8-quinolate

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Copper sulfate. See Cupric sulfate.

Co-Ral. See Coumaphos.

Coralox -- Coroxon; Estrella; Umbethion.

Estrella (Umbethion)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, *Rev. Med. Vet.*, Buenos Aires, v. 58 (2), 101-102, 105
Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Coroxon

Waladde, S. M., 1976, *Bull. Animal Health and Prod. Africa*, v. 24 (1), 73-79
Boophilus microplus and *Haemaphysalis longicornis* larvae, comparative toxicological and biochemical study of effects of coumaphos and coroxon in vitro

Coralyne sulfoacetate (WR 158 221)

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, active in screening of antitumor compounds for efficacy against infection

Cordycepin -- 3'-Deoxyadenosine.

Cordycepin

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Cordycepin

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cordycepin -- Continued.

Cordycepin + Erythro-9-(2-hydroxy-3-nonyl)adenine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cordycepin + Guanosine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cordycepin + Guanosine + EHNA

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cordycepin-N-oxide

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Coriban. See Diamphenethide.

Corid. See Amprolium.

Coroxon. See Coralox.

Cortisone

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Co-trimoxazole. See Sulfamethoxazole or Trimethoprim.

Coumaphos -- Asuntol; Asuntol 50; Baymix Crumbles; O-(3-Chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) O,O-diethyl phosphorothioate; Co-Ral; O,O-Diethyl O-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate.

Coumaphos (Asuntol)
Boersema, J. H., 1978, Tijdschr. Diergeneesk., v. 103 (7), 377-380
Chorioptes bovis, horse with foot-mange, resistant to coumaphos, lindane successful

Asuntol
Camoens, J. K., 1977, Malaysian Vet. J., v. 6 (3), 111-124
Boophilus microplus, dairy cattle, various control measures discussed but spraying acaricides on pastures shows particular promise: Air Hitam, Johor, Malaysia

Asuntol
da Costa, U. C.; and Saraiva, D., 1977, Rev. Centro Cien. Rurais, v. 7 (4), 417-419
Myocoptes musculus, white mice [in English title; "camundongos brancos (Wistar)" in Portuguese text], treatment with asuntol and neguvon

Coumaphos
Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Asuntol
El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, Argas persicus, evaluation of 10 insecticides

Coumaphos
Eschle, J. L.; et al., 1973, J. Econom. Entom., v. 66 (1), 290-291
Haematobia irritans, dairy cattle, effectiveness of coumaphos on cable-type back-rubbers, no residues detected in milk

Coumaphos
Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Coumaphos
Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Asuntol (Coumaphos)
Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105
Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Coumaphos -- Continued.

Coumaphos
Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, J. Econom. Entom., v. 71 (2), 315-318
Ornithonyssus sylviarum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban permethrin and SD-43775 also effective; mites displayed tolerance to malathion

Coumaphos
Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73
Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Coumaphos
Harvey, T. L.; and Brethour, J. R., 1979, J. Econom. Entom., v. 72 (4), 516-518
Haematobia irritans, coumaphos-treated and untreated steers, weight gains, economic importance: Hays, Kansas

Asuntol
Liebisch, A.; et al., 1978, Vet.-Med. Nachr. (1), 49-62
Psoroptes ovis, sheep, asuntol emulsion: Nordfriesland

Asuntol
Liebisch, A.; et al., 1978, Vet.-Med. Nachr. (1), 63-76
mange, cattle, asuntol

Asuntol (Coumaphos)
Liebisch, A.; et al., 1978, Vet. Med. Rev. (1), 49-62
Psoroptes ovis, sheep, increasing incidence, controlled field trial with asuntol dip, good results; need for good dipping technique, most frequent sources of failure: Federal Republic of Germany

Asuntol (Coumaphos)
Liebisch, A.; et al., 1978, Vet. Med. Rev. (1), 63-76
Chorioptes bovis, Sarcoptes bovis, Psoroptes ovis, cattle, spray treatment with asuntol effectively controlled mange: Germany

Coumaphos (Asuntol; Co-Ral)
Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Coumaphos (Co-Ral)
Meleney, W. P.; and Roberts, I. H., 1979, J. Med. Entom., v. 16 (1), 52-58
Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Coumaphos (CoRal)
Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661
flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico

Coumaphos -- Continued.

Asuntol + Neguvon

Oba, M. S. P.; Ogassawara, S.; and Costa, A. J. S., 1977, Arq. Inst. Biol., Sao Paulo, v. 44 (1-2), 95-97

Chorioptes bovis var. *bovis*, bovinos, clinical aspects, cure with neguvon + asuntol: Municipio de Sao Carlos, Sao Paulo

Asuntol 50

Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Coumaphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Coumaphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Coumaphos

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Baymix Crumbles

Todd, A. C.; et al., 1978, Vet. Med. and Small Animal Clin., v. 73 (5), 614, 616-619
deworming of dairy cows with Baymix Crumbles increased milk production and fattening, treated cows entered next lactation sooner than controls: Pennsylvania and North Carolina

Asuntol

Vural, A., 1977, Vet.-Med. Nachr. (1), 75-76
Psoroptes communis var. *ovis*, sheep, asuntol dip

Asuntol

Vural, A., 1977, Vet. Med. Rev. (1), 75-76
Psoroptes communis var. *ovis*, sheep, treatment with asuntol highly effective

Coumaphos

Waladde, S. M., 1976, Bull. Animal Health and Prod. Africa, v. 24 (1), 73-79
Boophilus microplus and *Haemaphysalis longicornis* larvae, comparative toxicological and biochemical study of effects of coumaphos and coroxon in vitro

Coumaphos

Wright, F. C.; and Riner, J. C., 1979, South-west. Entom., v. 4 (1), 40-45
Psoroptes ovis, *P. cuniculi*, 10 acaricides evaluated using 'tea-bag' technique

Coxistac. See Salinomycin.

Coyden. See Meticlorpindol.

Coyden 25. See Meticlorpindol.

Creolin

Slepnev, N. K.; and Zen'kov, A. V., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 135-137

E[chinococcus] granulosus, *protoscolices*, destructive action of high and low temperatures; lysol and creolin most destructive of chemicals tested

Creolin

Suiunchaliev, R. S., 1978, Veterinariia, Moskva (6), 63-65

psoroptic mange, sheep, method for determining creolin concentration in dip containing emulsion of creolin, gamma isomer of hexachlorane, and water

Cresol

Sherkov, Sh.; et al., 1978, Vet. Sbirka, v. 76 (6), 39-41

coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat

Crotamiton -- Eurax.

Crotamiton

Burns, B. R.; Lampe, R. M.; and Hansen, G. H., 1979, Am. J. Dis. Child., v. 133 (10), 1031-1034

neonatal *Sarcoptes scabiei*, 5 case reports, distinctive clinical pattern, crotamiton treatment: William Beaumont Army Medical Center, El Paso, Texas

Eurax

Tierney, F.; and Baillie, J., 1979, Vet. Med. and Small Animal Clin., v. 74 (1), 69-70
Cnemidocoptes pilae, Mexican red-headed parrot (beak, face), case report, combined aerosol (malathion solution) and topical treatment (eurax and Goodwinol cream) highly effective and less stressful

Crotoxyphos -- α -Methylbenzyl (E)-3-hydroxycrotonate dimethyl phosphate.

Crotoxyphos

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Crotoxyphos

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886

laboratory-reared *Haematobia irritans*, susceptibility to topically applied insecticides

Crotoxyphos

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Crotoxyphos -- Continued.

Crotoxyphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
 Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Crotoxyphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
 Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Crotoxyphos

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
 Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Crotoxyphos

Wright, F. C.; and Riner, J. C., 1979, Southwest. Entom., v. 4 (1), 40-45
 Psoroptes ovis, P. cuniculi, 10 acaricides evaluated using 'tea-bag' technique

Crufomate -- 4-tert-Butyl-2-chlorophenyl methyl-methylphosphoroamidate; Hipolen-6; Ruelene; Ruelene 6-R; TF-302.

Crufomate (TF-302)

Giordia, H.; and McCampbell, H. C., 1978, Georgia Vet., v. 30 (2), 15-17
 nematodes, beef cattle, efficacy of new pour-on formulation of crufomate

4-tert-Butyl-2-chlorophenyl methyl methyl phosphoroamidate

Costa, H. M. de A.; et al., 1975, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 27 (3), 295-301
 nematodes, cattle, 4-tert-butyl-2-chlorophenyl methyl methyl phosphoroamidate applied by pour-on method

4-tert-Butyl-2-chlorophenyl methyl-methylphosphoroamidate

Costa, J. O.; et al., 1975, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 27 (3), 289-293
 nematodes, cattle, 4-tert-butyl-2-chlorophenyl methyl-methylphosphoroamidate applied by pour-on method

Crufomate

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
 Boophilus annulatus, B. microplus, laboratory tests of insecticides

Crufomate

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
 laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Crufomate

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
 Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Crufomate -- Continued.

Crufomate (Ruelene)

Meleney, W. P.; and Roberts, I. H., 1979, J. Med. Entom., v. 16 (1), 52-58
 Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Ruelene

Oproiu, V.; et al., 1977, Rev. Crest. Animal-elor, v. 27 (7), 47-49
 Hypoderma bovis, cattle, neguvon, ruelene, trichlorphon, curative treatment in spring, prophylactic treatment in autumn more efficient

Ruelene 6-R (Hipolen-6)

Sayin, F.; and Meric, I., 1976, Vet. Fak. Der-gisi, Ankara Univ., v. 23 (3-4), 301-307
 Hypoderma, indigenous cattle, pour-on application of ruelene 6-R, tiguvon, good results: Central Anatolia, Turkey

Crufomate

Smith D. L., 1976, Manitoba Entom., v. 10, 5-8
 Hypoderma spp., calves, weight gains, no significant difference between treated and untreated calves in response to control of cattle grubs with insecticides (trichlorfon and crufomate): Manitoba

Crufomate

Wright, F. C.; and Riner, J. C., 1979, Southwest. Entom., v. 4 (1), 40-45
 Psoroptes ovis, P. cuniculi, 10 acaricides evaluated using 'tea-bag' technique

Cucurbita citrullus seeds

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, Scientia Pharm., v. 47 (2), 114-118
 Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

(E)-1-(p-Cumenyloxy)-6,7-epoxy-3,7-dimethyl-2-octene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
 Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

Cupric carbonate

Epel'dimov, L. S.; and Plotnikova, L. M., 1977, Sborn. Nauch. Rabot SibNIVI (28), 140-142
 [Trichostrongylus], rabbits infected with sheep species as models for anthelmintic study, tests of nilverm, banminth-C, cupric carbonate

Cupric sulfate -- Blue vitriol; Copper sulfate.

Cupric sulfate + Phenothiazine salt

Artem'ev, G. M., 1978, Vestnik Sel'skokhoz. Nauki Kazakhstana (10), 91-93
 helminthiases, sheep, economic losses, phenothiazine salt and cupric sulfate mix: Pavlodarsk oblast

Cupric sulfate -- Continued.

Copper sulphate

Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, *Current Sc.*, Bangalore, v. 47 (8), 283-284 [Letter]

Trichodina [sp.] on exterior of *Cyprinus carpio*, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

Copper sulfate

Bogdanov, V. R.; et al., 1977, *Sborn. Nauch. Rabot SibNIVI*, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91

cestodes, pathomorphology resulting from action of various anthelmintics

Copper sulfate

Bogdanov, V. R.; Koshkina, N. G.; and Logachev, E. D., 1977, *Sborn. Nauch. Rabot SibNIVI*, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 92-93

Dipylidium caninum, various anthelmintics, in vitro action on surface tissues and inactivation of enzymes

Cupric sulfate-Phenothiazine mixture

Nekipelova, R. A.; Kurnikov, V. A.; and Iksanov, S. F., 1978, *Veterinariia*, Moskva (10), 67

Nematodirus and other nematodes, sheep, effects of addition of trace elements to phenothiazine-cupric sulfate mixture: Tselinogradsk oblast

Copper sulfate + Ferrous sulfate

Sverba, V. A.; and Shemchuk, V. R., 1978, *Veterinariia*, Moskva (10), 69-71

Sinergasilus major, white amur, copper sulfate and ferrous sulfate mixture, chlorophos, carbophos, formula for estimating concentrations in relation to temperature and other factors in aquaria or ponds

Copper sulphate

Venkateswara Rao, P.; et al., 1977, *Riv. Parassitol.*, Roma, v. 38 (1), 13-21

Prosthogonimus sp. cercariae, cercaricidal effect of certain common fertilizers, ammonium sulphate may be cercaricide of choice

Cyanamid-38023. See Famphur.

N-(2-Cyanoethyl) chloroacetamide

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

4-Cyano-2 iodo-6 nitrophenol. See Nitroxynil.

(±)-α-Cyano-3-phenoxybenzyl (±)-cis, trans-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropane-carboxylate. See Cypermethrin.

Cyano-(3-phenoxyphenyl)-methyl-4-chloro-α-(1-methylethyl)-benzeneacetate. See Fenvalerate.

Cyazone

Shchelkanov, K. G.; and Epel'dimov, L. S., 1977, *Sborn. Nauch. Rabot SibNIVI*, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 76-80

Dictyocaulus, sheep, divezid, nilverm, and cyazone, changes in lung tissue of sheep and some tissues of parasite resulting from drug action

Cycloadenosine -- 9-(β-DL-2α,3α-Dihydroxy-4β-[hydroxymethyl]-cyclopentyl)-adenine; Carbocyclic-adenosine.

Cycloadenosine

Dewey, V. C.; Kidder, G. W.; and Nolan, L. L., 1978, *Biochem. Pharmacol.*, v. 27 (10), 1479-1485

Crithidia fasciculata, mechanism of inhibition of growth by adenosine and adenosine analogs

Carbocyclic-adenosine

Senft, A. W.; and Crabtree, G. W., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1847-1856

Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Cyclobendazole -- C-C 2481; Ciclobendazole;

5-(Cyclopropyl-carbonyl)-2-(methoxycarbonylamino)-benzimidazole; Methyl-5-(cyclopropyl-carbonyl)-benzimidazol-2-yl-carbamate.

Ciclobendazole (C-C 2481)

Degremont, A.; and Stahel, E., 1978, *Schweiz. Med. Wchnschr.*, v. 108 (37), 1430-1433

mild helminth infections in humans, cyclobendazole tested, drug well tolerated

Ciclobendazole

Guggenmoos, R.; et al., 1978, *Tropenmed. u. Parasitol.*, v. 29 (4), 423-426

nematodes, humans, cyclobendazole, vermicial effect compared with metronidazole in double-blind study, side effects only in small percentage of cases: Bamenda, Cameroon

Cycloguanil -- Camolar; Cycloguanil pamoate; WR 5,473.

Cycloguanil pamoate (Camolar)

Akhound-Zadeh, H., 1976, *Rev. Internat. Serv. Sante Armees*, v. 49 (5), 421-426

chronic cutaneous leishmaniasis, soldier with severe ulcers that did not heal despite 8 years of therapy with various anti-leishmanial drugs, chronicity thought to be result of immuno-deficiency, ulcers finally cured after additional therapy with monomycine: Iran (had travelled to Khouzistan)

Cycloguanil pamoate (Camolar)

Furtado, T., 1974, *Rev. AMMG*, v. 25 (3), 108-113

human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Cycloguanil -- Continued.

Cycloguanil pamoate (Camolar)

Gusmao, H. H.; and de Souza, J. A. G., 1973, Rev. Saude Pub., S. Paulo, v. 7 (4), 335-341
human cutaneous leishmaniasis, single dose treatment with cycloguanil pamoate gave good results: Amapa Territory, Brazil

Cycloguanil

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Cycloguanil pamoate

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351
Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Cycloguanil (WR 5,473)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Cycloguanil pamoate. See Cycloguanil.

Cycloheximide -- Actidione.

Cycloheximide

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Actidione

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Cycloheximide

Sinden, R. E.; and Smalley, M. E., 1979, Parasitology, v. 79 (2), 277-296
Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

2-Cyclohexylcarbonyl-1,3,4,6,7,11b-hexahydro-2H-pyrazino[2,1a]isoquinolin-4-one. See Praziquantel.

Cycloleucine. See 1-Aminocyclopentane carboxylic acid.

Cyclophosphamide

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Cyclophosphamide

Machemer, L.; and Lorke, D., 1978, Arch. Toxicol., v. 39 (3), 187-197
praziquantel, mutagenicity studies on mice and Cricetulus griseus, no indication of mutagenic action, compared with cyclophosphamide and placebo

5-(Cyclopropyl-carbonyl)-2-(methoxycarbonylamino)-benzimidazole. See Cyclobendazole.

[5-[(Cyclopropylmethyl)sulfinyl]-1H-benzimidazol-2-yl] carbamic acid, methyl ester
Cruthers, L. R.; et al., 1978, Experientia, v. 34 (12), 1574
variety of nematodes, cestode, and trematode species in domestic animals, orally active benzimidazole anthelmintics discovered to be active by injection also

Cyclosamide. See Niclosamide.

D-Cycloserine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cycostat. See Robenidine.

Cygon. See Dimethoate.

Cyhexatin (Plictran)

Meleney, W. P.; and Roberts, I. H., 1979, J. Med. Entom., v. 16 (1), 52-58
Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Cymbopogen proximus herb

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, Scientia Pharm., v. 47 (2), 114-118
Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

Cypermethrin -- (\pm)- α -Cyano-3-phenoxybenzyl (\pm)-cis, trans-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate; NRDC 149; PJT 1.

Cypermethrin (NRDC 149)

Hall, C. A., 1978, Austral. Vet. J., v. 54 (10), 471-472

Damalina ovis, sheep (exper.), cypermethrin proved effective in eradicating lice and at 5 and 10 ppm prevented reinfestation for 7 and 19 weeks respectively; addendum briefly gives results for permethrin in similar tests

Cypermethrin (NRDC 149)

Nolan, J.; Roulston, W. J.; and Schnitzerling, H. J., 1979, Austral. Vet. J., v. 55 (10), 463-466

Boophilus microplus, range of resistant strains on naturally and experimentally infected cattle, field and stall spraying trials, efficacy of synthetic pyrethroids for tick control, potentiation of pyrethroids by organo-phosphorus compounds

Cytarabine -- Cytosine arabinoside.

Cytosine arabinoside

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Cytembena

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Cythioate. See 0,0-Dimethyl-0-p-sulfamoylphenyl phosphorothioate.

Cythion. See Malathion.

Cytochalasin B

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Cytosine arabinoside. See Cytarabine.

Dactinomycin -- Actinomycin D; Actinomycin D.

Actinomycin D

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Actinomycin D

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Actinomycin D

Sinden, R. E.; and Smalley, M. E., 1979, *Parasitology*, v. 79 (2), 277-296

Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

Actinomycin D

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Dalacin-C. See Clindamycin.

Dankil (CaVP, Nestyne)

Ronald, N. C.; Bell, R. R.; and Simpson, J. E., 1978, *Southwest. Vet.*, v. 31 (3), 201-203

gastrointestinal nematodes, swine, critical evaluation of dankil, highly effective: east-central Texas

DAPI. See 4'6-Diamidino-2-phenylindole.Dapsone -- DDS; Diaminodiphenylsulfone; Maloprim (with Pyrimethamine); WR 448.

Dapsone + Pyrimethamine (= Maloprim)

Botelho, A., 1974, *South African Med. J.*, v. 48 (31), 1351-1352

malarias, humans, clinical trials using sulphones and sulphonamides with a pyrimidine derivative

Maloprim + Camoprim

Botelho, A., 1974, *South African Med. J.*, v. 48 (31), 1351-1352

malarias, humans, clinical trials using sulphones and sulphonamides with a pyrimidine derivative

Dapsone -- Continued.

Dapsone + Pyrimethamine

Buyst, H., 1977, *Ann. Soc. Belge Med. Trop.*, v. 57 (4-5), 201-212

sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia

Dapsone + Pyrimethamine (= Maloprim)

Hughes, A.; and Gatus, B. J., 1979, *J. Trop. Med. and Hyg.*, v. 82 (6), 120-121

severe megaloblastic anaemia, woman treated with daily dosage of maloprim

Diaminodiphenylsulfone (DDS)

Pellegrino, J.; and Katz, N., 1975, *Rev. Inst. Med. Trop. S. Paulo*, v. 17 (3), 199-205

Schistosoma mansoni, laboratory animals and humans, diaminodiphenylsulfone interfered with parasite egg laying

Dapsone + Pyrimethamine

du Preez, O.; and Cockcroft, C. R., 1975, *South African Med. J.*, v. 49 (34), 1407-1408

malaria prophylaxis trials, army personnel camping in endemic area, 3 drug combinations, no infections reported in trial groups while local population acting as control reported 250 falciparum cases: Caprivi Strip, South Africa

Dapsone (WR 448)

Rane, D. S.; and Kinnamon, K. E., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (6), 937-947

sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Dapsone

Ray, A. P.; Parkinson, A. D.; and Black, R. H., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (1), 19-22

Plasmodium berghei in chloroquine resistant white mice, results of treatment with combinations of proguanil and dapsone

Daraprim. See Pyrimethamine.Darvisul. See Diaveridine or Sulfaquinoxaline.

Daunomycin

Sinha, M.; Goswami, D. N.; and Das Gupta, N. N., 1978, *Indian J. Biochem. & Biophys.*, v. 15 (3), 162-165

daunomycin, berenil, dielectric studies on interaction with DNA

Daunorubicin

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

DBTD. See Dibutyltin dilaurate.

DDS. See Dapsone.

DDT -- Chlophenotane spirits; p,p-Dichlorodiphenyl-trichloromethyl-methane; Ivoran.

DDT
Cole, M. M.; et al., 1973, J. Econom. Entom., v. 66 (1), 118-119
Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

DDT
Delak, M.; and Radokovic, M., 1978, Vet. Arhiv, Zagreb, v. 48 (1), 17-22
Fasciola hepatica from bile ducts of cattle, and bovine liver samples, amount of contamination with DDT, γ -HCH, and α -HCH

DDT
Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

DDT
Goszczyńska, K.; and Styczynska, B., 1972, Roczniki Panstw. Zakl. Hig., v. 23 (2), 245-251
Pediculus humanus humanus, selection of laboratory strain reared through several generations aimed at induction of resistance to DDT and lindane; resistance developed to DDT but not to lindane

Chlophenotane spirits (DDT; Ivoran)
Jensen, O.; Bjerregaard, P.; and Nielsen, A. O., 1979, Ugeskr. Laeger, v. 141 (4), 225-226
head lice, humans, quassia extract vs. chlophenotane: Denmark

DDT
Kiefer, M.; and Cyprich, D., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Zool. (23), 5-12
ticks, susceptibility to acaricides: Slovakia

DDT
Kir'iakova, A. N.; et al., 1974, Parazitologiya, Leningrad, v. 8 (2), 157-163
Xenopsylla skrjabini and Coptopsylla lamellifer on Rhombomys opimus and in its colonies, evaluation of effectiveness of deep-dusting with DDT as possible anti-plague measure: Aral-Kum and Dulan, Priaral'sk Karakum

p,p-Dichlorodiphenyl-trichloromethyl-methane (DDT)
Privora, M.; Rupes, V.; and Cerny, V., 1970, Folia Parasitol., v. 17 (1), 81-84
Dermacentor marginatus, laboratory trials testing six insecticides

ppDDT
Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

DDT -- Continued.

ppDDT
Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

DDT
Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

DDT
Thaug, U.; et al., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (3), 390-397
X[enopsylla] cheopis, X. astia, collection from small mammals during plague outbreak, some resistance to DDT: Hlegu, Burma

p,p'-DDT
Uspenskii, I. V., 1974, Parazitologiya, Leningrad, v. 8 (4), 312-321
Ixodes persulcatus, susceptibility to acaricides

DDT
Uspenskii, I. V.; and Repkina, L. V., 1974, Parazitologiya, Leningrad, v. 8 (1), 3-11
Ixodes persulcatus, physiological age and age structure of natural populations over a season, changes in mean physiological age under influence of DDT solution or of the solvent (methyl-ethyl ketone), comparisons of changes in physiological age and susceptibility to DDT

DDVF. See Dichlorvos.

DDVP. See Dichlorvos.

7-Deazaadenosine. See Tubercidin.

3-Deazauridine
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

DEC. See Diethylcarbamazine.

Decamethrin (NRDC 161)
Nolan, J.; Roulston, W. J.; and Schnitzerling, H. J., 1979, Austral. Vet. J., v. 55 (10), 463-466
Boophilus microplus, range of resistant strains on naturally and experimentally infected cattle, field and stall spraying trials, efficacy of synthetic pyrethroids for tick control, potentiation of pyrethroids by organo-phosphorus compounds

Decaris. See Tetramisole.

Deccox. See Decoquinatate.

Decobald-cobamide

Christow, C., 1978, *Biochem. and Exper. Biol.*, v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂ antagonists upon growth

Decobald-cobinamide

Christow, C. P., 1978, *Riv. Biol.*, v. 71 (1-4), n. s. v. 31, 113-116
Trichomonas foetus, effect of decobald-cobinamide and L-I-Methyl-2-aminoethenole upon growth

Decoquinatate -- Deccox; 6-n-Decyloxy-7-ethoxy-4-hydroxyquinoline-3-carboxylate; 6-Ethyl-(decyloxy)-7-ethoxy-4-hydroxy-3-quinoline-carboxylate.

Decoquinatate

Bajwa, R. S.; and Gill, B. S., 1977, *Acta Vet. Brno*, v. 46 (1-2), 149-158
Eimeria tenella, chickens (exper.), decoquinatate vs. amprolium used prophylactically vs. therapeutically, anticoccidial activity against different levels of infection, effect on development of immunity

Decoquinatate (Deccox)

Fox, J. E., 1978, *Mod. Vet. Pract.*, v. 59 (8), 599-603
 bovine coccidiosis, review, emphasis on prevention and control; field tests, decoquinatate against *Eimeria bovis* and *E. zuernii* prevented clinical signs of disease with no observable signs of toxicity

Decoquinatate

Karlsson, T.; and Reid, W. M., 1978, *Avian Dis.*, v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Decoquinatate

Latter, V. S.; and Wilson, R. G., 1979, *Parasitology*, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Decoquinatate

Lee, E. H., 1979, *Canad. Vet. J.*, v. 20 (4), 102-104
Eimeria tenella, drug-resistant field strains, White Leghorn chickens, single and low-level oocyst infections, treatment with robenidine or decoquinatate

Decoquinatate

McDougald, L. R.; and Galloway, R. B., 1977, *Ztschr. Parasitenk.*, v. 54 (1), 95-100
Eimeria tenella in vitro, development inhibited by serum from chickens fed anticoccidial drugs, technique to assay drug activity and to characterize and quantitate therapeutic effect

Decoquinatate -- Continued.

Decoquinatate

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Decoquinatate

Singh, J.; and Gill, B. S., 1976, *Riv. Parasitol.*, Roma, v. 37 (1), 57-62
Eimeria necatrix, different levels of infection, chicks, activity of decoquinatate used prophylactically and therapeutically, effect on development of immunity

6-n-Decyloxy-7-ethoxy-4-hydroxyquinoline-3-carboxylate. See Decoquinatate.

Dehydroemetine -- Ro 1-9334/20.

Dehydroemetine

Botero R., D., 1978, *Ann. Rev. Pharmacol. and Toxicol.*, v. 18, 1-15
 antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Dehydroemetine (Ro 1-9334/20)

Correa, M. O.; and Fleury, G. C., 1971, *Rev. Soc. Brasil. Med. Trop.*, v. 5 (5), 267-270
Fasciola hepatica, human, case report, cure with dehydroemetine: Estado de Sao Paulo

Dehydroemetine

Dutta, G. P.; and Narain, L., 1978, *Indian J. Exper. Biol.*, v. 16 (7), 838-840
Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Dehydroemetine

Fuchs, P., 1978, *Deutsche Med. Wchnschr.*, v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Dehydroemetine

Katz, N.; and Pellegrino, J., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (5), 245-252
Schistosoma mansoni, Cebus monkeys, correlation of number of eggs per gram of rectal tissue with number of female worms, challenge infection effect, or drug action

Dehydroemetine

Lumbreras, H.; and Uyema, N., 1972, *Rev. Peruana Med. Trop.*, v. 1 (2), 95-98
Balantidium coli, in vitro, comparative action of dehydroemetine and emetine hydrochloride

Dehydroemetine -- Continued.

Dehydroemetine + Di-iodohydroxyquinoline + Oxytetracycline

Masters, D. K.; and Hopkins, A. D., 1979, *J. Trop. Med. and Hyg.*, v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire

Dehydroemetine

Olaeta Elizalde, R., 1973, *Rev. Cir. Hosp. Jua-rez, Mexico* (187-188), v. 44, 1972-1973, 59-64
 human hepatic amoebic abscess, therapeutic recommendations (emetine, dehydroemetine, chloroquine, metronidazole)

Dekelmin. See Methyridine.

Delagil. See Chloroquine.

Delnav. See Dioxathion.

Delnav DFF. See Dioxathion.

Demeclocycline -- Demeclocycline hydrochloride; Ledermycin.

Demeclocycline hydrochloride (Ledermycin)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, *Chemotherapy*, v. 25 (4), 222-226
Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizontocidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Demeclocycline hydrochloride. See Demeclocycline

N-Demethyl-4'-pentyl clindamycin hydrochloride (U-24)

Powers, K. G.; and Jacobs, R. L., 1972, *Antimicrob. Agents and Chemotherapy*, v. 1 (1), 49-53
Plasmodium falciparum, chloroquine-resistant strain in *Aotus trivirgatus*, 2 chlorinated lincomycin analogues cured blood-induced infections

Demodecil

Rebelo, M. E.; Diogo, M. R.; and Serra, J. S., 1977, *Reposit. Trab. Lab. Nac. Invest. Vet., Lisboa*, v. 9, 117-120
Demodex cuniculi, rabbits, treatment with demodecil

3'-Deoxyadenosine. See Cordycepin.

6-Deoxy-6-demethyl-6-methylene-5-hydroxy-tetracycline. See Methacycline.

2-Deoxy-D-glucose (2-DG)

Bunn, M. M.; et al., 1977, *Ztschr. Parasitenk.*, v. 52 (3), 245-256
Herpetomonas samuelpessoai in vitro, 2-deoxy-D-glucose (2-DG) inhibits growth and respiration, modifies ultrastructure of cells; some carbohydrates decrease effect of 2-DG

2-Deoxyglucose

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

5'-Deoxy-5'-S-isobutyl adenosine

Trager, W.; Robert-Gero, M.; and Lederer, E., 1978, *FEBS Letters*, v. 85 (2), 264-266
Plasmodium falciparum, antimalarial activity of S-isobutyl adenosine analogues in culture

4-Deoxypyridine

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Depo-Medrol. See Methylprednisolone acetate.

Dertil. See Niclofolan.

Dertil B. See Niclofolan.

Dertil O. See Niclofolan.

6-Desoxy-5-hydroxytetracycline. See Doxycycline.

Deuteroporphyrin IX

Meshnick, S. R.; et al., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Deuteroporphyrin IX bisglycol

Meshnick, S. R.; et al., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

- Deuteroporphyrin IX disulfonic acid
Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis
- Dexamethasone -- Tresaderm (with Neomycin and Thiabendazole).
- Tresaderm
Faulk, R. H.; and Schwirck, S., 1978, Vet. Med. and Small Animal Clin., v. 73 (3), 307-308
Otodectes cynotis, dogs, cats (ears of both), tresaderm, clinical trial, highly effective
-
- Dextran -- Dextran sulfate.
- Dextran sulphate 500
James, D. M., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 471-476
[Trypanosoma] congolense, T. brucei, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs
- Dextran 500
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Dextran sulfate 500
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Dextran sulfate 2000
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Dextran sulfate. See Dextran.
- Di-[2(4-acetamido phenoxy)ethyl]ether. See Diamphenethide.
- Di-aceturate of 4,4-diazoaminodibenzamidine. See Berenil.
- O-N,N'-Diacetyl-aminobenzole
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- N,N'-Diacetyl-p-aminophenol acetate
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- Diacetylanilide
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- Diacetyl-ethylene-diamine
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- N,N'-Diacetyl urea
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- C,C-Diallyl-bis-(4-amino-2-methyl-6-quinoly1) malonamide -- Bayer 7602 Ac: Diallylmalonyl-(4-amino-2-methyl-quinoly1-6-amide) acetate; 3024 I.C.I.
- Diallylmalonyl-(4-amino-2-methyl-quinoly1-6-amide) acetate (3024 I.C.I.)
Brener, Z., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (4), 302-306
Trypanosoma cruzi, description of method allowing study of drug action on trypomastigotes in mice
- Bayer 7602 Ac
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
-
- Diallylmalonyl-(4-amino-2-methyl-quinoly1-6-amide) acetate. See C,C-Diallyl-bis-(4-amino-2-methyl-6-quinoly1) malonamide.
- Diamfenetide. See Diamphenethide.
- 3:3'-Diamidinocarbanilide. See Amicarbalide.

4,4'-Diamidino-diazoaminobenzene diacetate.
See Berenil.

4-4'-Diamidino-diazo-amino benzol diacetate.
See Berenil.

Diamidino-diphenoxypentane. See Pentamidine.

1,5-Di(4-amidinodiphenoxypentane) di(2-hydroxyethanesulfonate). See Pentamidine.

4'6-Diamidino-2-phenylindole -- 102/198; DAPI.

DAPI (102/198)

Williamson, J.; and McLaren, D. J., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 660-661

Trypanosoma rhodesiense, ultrastructural alterations induced by treatment with DAPI (new diamidine trypanocide)

2,6-Diaminoanthraquinone, bisamidines
Burden, E. J.; et al., 1979, Experientia, v. 35 (1), 33-35

Entamoeba histolytica, rats, hamsters, potent activity of bisamidines of 2,6-diaminoanthraquinone

2,6-Diaminoanthraquinone bisamidines
Fabio, P. F.; et al., 1978, J. Med. Chem., v. 21 (3), 273-276

Entamoeba histolytica, diaminoanthraquinone bisamidines, laboratory trials comparing activity against cecal form in rats and hepatic form in golden hamsters with activity of known amoebicides

3,3'-Diaminocarbanilide diisethionate. See Amicarbalide.

Diamino-dephenoxyalkanes

Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, J. Parasitol., v. 64 (1), 69-77

Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy

2,4-Diamino-6-[(3,4-dichlorobenzyl)-nitroso-amino]-quinazoline (CI-679)

Schmidt, L. H.; and Rossan, R. N., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 781-792

Plasmodium cynomolgi, monkeys infected with Ro or Ro/PM strains, appraisals of various activities of CI-679

4,6-Diamino-1,2-dihydro-2,2-dimethyl-1-(3,4-dichlorobenzoyloxy)-1,3,5-triazine hydrochloride -- WR-38,839.

WR-38,839

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

Diaminodiphenylsulfone. See Dapsone.

2,4-Diamino-6-(2-naphthylsulfonyl)-quinazoline -- WR-158,122.

2,4-Diamino-6-(2-naphthylsulfonyl)-quinazoline (WR-158,122)

Elslager, E. F.; et al., 1979, J. Med. Chem., v. 22 (10), 1247-1257

Plasmodium spp., antimalarial activity of 2,4-diamino-6-(2-naphthylsulfonyl)quinazoline and related 2,4-diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines, laboratory studies; activity of WR-158,122 enhanced by coadministration of sulfadiazine

2,4-Diamino-6-(2-naphthyl)-sulfonylquinazoline (WR-158,122)

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-158,122

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

2,4-Diamino-6-(2-naphthyl)-sulfonyl-quinazoline (WR-158,122)

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 808-818

Plasmodium falciparum, P. vivax, various drug-resistant and drug-susceptible strains in Aotus trivirgatus griseimembra, capacity of sulfadiazine to enhance activities of WR-158,122 and WR-159,412

2,4-Diamino-6-(2-naphthylsulphonyl)quinazoline (WR-158122) + Sulphadiazine (WR-7557)

Wise, D. L.; Gresser, J. D.; and McCormick, G. J., 1979, J. Pharm. and Pharmacol., v. 31 (4), 201-204

dual antimalarial system, sustained release of ³H-labelled WR-7557 and ¹⁴C-labelled WR-158122 in biodegradable carrier, rhesus monkeys, mice

2,4-Diamino-6-(8-naphthylsulfonyl)-5,6,7,8-tetrahydroquinazoline

Carroll, F. I.; Berrang, B.; and Linn, C. P., 1978, J. Med. Chem., v. 21 (4), 326-330

Plasmodium berghei, P. cynomolgi, experimental animals, resolution of antimalarial agents via complex formation with α-(2,4,5,7-tetra-nitro-9-fluorenylideneaminoxy)propionic acid, significant differences in toxicity

2,4-Diamino-6-(2-naphthyl)-thioquinazoline --
WR-154,928.

2,4-Diamino-6-(2-naphthyl)-thioquinazoline (WR-154,928)

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-154,928

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

2,4-Diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines

Elslager, E. F.; et al., 1979, J. Med. Chem., v. 22 (10), 1247-1257

Plasmodium spp., antimalarial activity of 2,4-diamino-6-(2-naphthylsulfonyl)quinazolines and related 2,4-diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines

2,4-Diamino-6-[(phenyl- and naphthyl)thio]quinazolines

Elslager, E. F.; et al., 1978, J. Med. Chem., v. 21 (10), 1059-1070

Plasmodium spp. in laboratory animals, thioquinazoline analogues synthesized and tested under laboratory conditions showed substantial suppressive antimalarial and prophylactic activity when compared with reference compounds, analogues also retained potent antimalarial effects against strains resistant to common antimalarials; the most active compound, 2,4-diamino-6-[(α,α,α -trifluoro-m-tolyl)thio]quinazoline has been designated for preclinical toxicologic studies

2,4-Diamino-5-piperonyl-pyrimidine -- WR-40,070.

WR-40,070

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

2,4-Diaminopyrroloquinazoline derivatives

McCormack, J. J.; et al., 1979, Biochem. Pharmacol., v. 28 (21), 3227-3229

Inhibition of dihydrofolate reductases by derivatives of 2,4-diaminopyrroloquinazoline, Crithidia oncopelti used as one source of reductases

2,4-Diamino-6-substituted quinazolines

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

(Diamino-4,6-triazinyl-1,3,5-amino-2)-4-phenylarsino di (D-thio-3-amino-2-methyl-3-butyric acid) -- Compound 'E' (with 2-[2-(4-Hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazyl) benzimidazole); F 151.

[[Diamino-4,6 triazinyl-1,3,5 amino-2)-4 phenylarsino di (D-thio-3 amino-2 methyl-3 butyric acid)] (F 151)

Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455

Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity

Compound 'E'

Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455

Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity

2-[4-[(4,6-Diamino-1,3,5-triazin-2-yl)-amino]phenyl]-1,3,2-dithiarsolane-4-methanol. See Melarsoprol.

2,4-Diamino-6-(5-trifluoromethylphenyl)-thioquinazoline -- WR-159,412.

WR-159,412

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

2,4-Diamino-6-(5-trifluoromethylphenyl)-thioquinazoline (WR-159,412)

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 808-818

Plasmodium falciparum, P. vivax, various drug-resistant and drug-susceptible strains in Aotus trivirgatus griseimembra, capacity of sulfadiazine to enhance activities of WR-158,122 and WR-159,412

2,4-Diamino-6-[(α,α,α -trifluoro-m-tolyl)thio]quinazoline

Elslager, E. F.; et al., 1978, J. Med. Chem., v. 21 (10), 1059-1070

Plasmodium spp. in laboratory animals, thioquinazoline analogues synthesized and tested under laboratory conditions showed substantial suppressive antimalarial and prophylactic activity when compared with reference compounds, analogues also retained potent antimalarial effects against strains resistant to common antimalarials; the most active compound, 2,4-diamino-6-[(α,α,α -trifluoro-m-tolyl)thio]quinazoline has been designated for preclinical toxicologic studies

2,4-Diamino-5-(3,4,5-trimethoxybenzyl)-pyrimidine. See Trimethoprim.

cis-Diamminedichloroplatinum (WR 177 529)
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, Antimicrob. Agents and Chemotherapy, v.
15 (2), 157-160

Trypanosoma rhodesiense, mice, active in
screening of antitumor compounds for efficacy
against infection

Diamphenethide -- Coriban; Di-[2(4-acetamido
phenoxy)ethyl]ether; Diamfenetide; Diamphene-
tide; N,N'-[Oxybis(2,1-ethanedioxy-4,1-
phenylene)] bisacetamide; 2,2-Bis(p-acetyl
aminophenoxy) diethyl ether.

Diamphenethide (Coriban)
Calamel, M.; Villemin, P.; and Leimbacher, F.,
1979, Rec. Med. Vet., v. 155 (1), 37-46
Dicrocoelium dendriticum, sheep, diamphene-
thide, efficacy in relation to duration,
dosage, host age, and retreatment

Diamfenetid (Coriban)
Corba, J.; et al., 1978, Veterinarstvi, v. 28
(6), 274-275
Dicrocoelium dendriticum, sheep, trials of
cambendazole, fenbendazole, diamfenetid

Diamphenethide (Coriban)
Corba, J.; Hovorka, J.; and Popovic, S., 1973,
Vet. Med., Praha, v. 46, v. 18 (6), 365-370
Fasciola hepatica, sheep, efficacy of diam-
phenethide

Coriban
Didenko, P. P.; et al., 1979, Veterinaria,
Moskva (5), 49-50
Sanguinicola inermis, carp, coriban in feed

Diamphenethide (Coriban)
Evans, J. W.; and Green, P. E., 1978, Austral.
Vet. J., v. 54 (9), 454-455 [Letter]
Platynosomum concinnum, cats, anthelmintics,
drug trials

Diamphenetide (Coriban)
Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot
SibNIVI (28), 143-146
Fasciola hepatica, rabbits, new anthel-
mintics tested, phenacetine highly effective

Diamphenethide
Oleinik, A. P., 1977, Veterinaria, Moskva
(5), 65-67
fascioliasis, sheep, diamphenethide, de-
termining dosage and toxicity levels

Diamphenethide (Coriban)
Over, H. J.; et al., 1978, Tijdschr. Dier-
geneesk., v. 103 (2), 129-139
Fasciola hepatica in sheep grazing on known
contaminated pastures, diamphenethide, good
results: province of Zeeland, the Nether-
lands

Diamfenetide
Rew, R. S.; Colglazier, M. L.; and Enzie, F. D.,
1978, J. Parasitol., v. 64 (2), 290-294
Fasciola hepatica, lambs (exper.), diamfene-
tide, clinical and anthelmintic effects, value
of serum gamma-glutamyl transpeptidase in
detecting hepatobiliary damage

Diamphenethide -- Continued.

Diamphenethide
Rowlands, D. ap T.; and Clampitt, R. B., 1979,
Vet. Parasitol., v. 5 (2-3), 155-175
Fasciola hepatica, sheep, cattle (both
exper.), bodyweight, blood and plasma analy-
ses, emphasis on use of plasma enzyme levels
to detect and monitor liver damage and to
assess efficacy of diamphenethide against
immature flukes

Diamphenetide. See Diamphenethide.

Diapron. See Amicarbalide.

Diaveridine -- Darvisul (with Sulfaquinoxaline);
Saquadil (with Sulfaquinoxaline); Sulphacom-
bine (with Sulphadimidine).

Diaveridine + Sulphadimidine (=Sulphacombine)
Danek, J.; et al., 1978, Biol. a Chem. Zivoc.
Vyroby, Vet., v. 14 (v. 20) (2), 151-169
Eimeria spp., rabbits (nat. and exper.),
sulphacombine, controlled test, subacute
toxicity, no negative effect on followed
indicators

Diaveridine
Latter, V. S.; and Wilson, R. G., 1979, Para-
sitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assess-
ment of anticoccidial activity in cell
culture

Diaveridine
McHardy, N., 1978, Ann. Trop. Med. and Parasi-
tol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected
bovine lymphoblastoid cell cultures used in
in vitro screens to test wide range of
compounds for chemotherapeutic activity

Diaveridine + Sulfaquinoxaline (=Darvisul liquid)
Manuel, M. F.; and Neri, R. A., 1975, Philip-
pine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels,
efficacy of 7 water-soluble coccidiostats

Diaveridine + Sulphadimidine (=Sulphacombine)
Strakova, J.; Sevcik, B.; and Dvorak, M.,
1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14
(v. 20) (2), 171-180
coccidiosis, chicks, sulphacombine, acute and
subacute toxicity studies

Diaveridine + Sulphaquinoxaline (= Saquadil)
Williams, R. B., 1978, Vet. Parasitol., v. 4
(2), 193-197
Isospora mayuri and Eimeria colchici in Pavo
cristatus controlled by sulphaquinoxaline and
diaveridine in drinking water

Diazinon -- Basudin; Diazinon DFF; 0,0-Diethyl-0-(2-isopropyl-6-methyl-4-pyrimidinyl); 0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate; 0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate; 0,0-Diethyl-0-2-isopropyl-4-methyl-6-pyrimidyl phosphorothionate; 0,0-Diethyl-0-(2-isopropyl-4-methyl-pyrimidyl-6) thio-phosphate; 0,0-Diethyl 0-[6-methyl-2-(1-methylethyl)-4-pyrimidinyl] phosphorothioate; Dimpylat; Neocidol.

Diazinon

Drummond, R. O.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 130-133
Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Diazinon DFF

Flower, P. J., 1978, *Trop. Animal Health and Prod.*, v. 10 (4), 207-213
Psoroptes ovis, sheep, course of disease outbreak, problems in attempting to limit its spread, partial control achieved through dipping in diazinon DFF backed by legislation: Lesotho

Diazinon

Foreyt, W. J.; Long, G. G.; and Gates, N. L., 1978, *Vet. Med. and Small Animal Clin.*, v. 73 (4), 503-505
Trichodectes canis on *Canis latrans*, case reports, severe alopecia, diazinon dusted on bedding highly effective

Neocidol (Diazinon)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, *Rev. Med. Vet.*, Buenos Aires, v. 58 (2), 101-102, 105
Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Diazinon

Kirkwood, A. C.; Quick, M. P.; and Page, K. W., 1978, *Vet. Rec.*, v. 102 (3), 50-54
ectoparasites, sheep, showers and plunge dipping, efficacy using diazinon against *Lucilia sericata*, γ HCH against *Psoroptes communis ovis*

Diazinon

Lloyd, J. E.; Olson, E. J.; and Pfadt, R. E., 1978, *J. Econom. Entom.*, v. 71 (3), 548-550
Melophagus ovinus, sheep, diazinon, ronnel, low volume spraying gave rapid control with no adverse effects

Diazinon (Basudin)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, *PANS*, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Diazinon

Naidu, N. V.; and Raghavachar, B., 1978, *Indian J. Animal Health*, v. 17 (2), 117-121
diazinon exposed vs. normal Indian buffaloes, changes in blood cholinesterase activity and clinical response

Diazinon -- Continued.

Diazinon (Dimpylat)

Niemand, H. G.; Niemand, S.; and Wendel, E., 1979, *Kleintier-Praxis*, v. 24 (4), 173-175
discussion of chief effective principles of different flea collars (dichlorvos, diazinon, propoxur), toxicity

Diazinon

Ottenschot, T. R. F.; and Gil, D., 1978, *Tijdschr. Diergeneesk.*, v. 103 (2), 1104-1108
cheyletiellosis in long-haired cats, chronic pruritis, treatment with diazinon flea collars and lindane baths: Netherlands; Western Germany

Diazinon

Palmer, J. S., 1978, *Am. J. Vet. Research*, v. 39 (7), 1231-1232
diazinon, acute toxicity studies of micro-encapsulated vs. wettable powder formulation applied dermally to calves and steers

Diazinon

Rawlins, S. C.; and Mansingh, A., 1978, *J. Econom. Entom.*, v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Diazinon

Rawlins, S. C.; and Mansingh, A., 1978, *J. Econom. Entom.*, v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Diazinon

Rawlins, S. C.; and Mansingh, A., 1979, *J. Econom. Entom.*, v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Diazinon (Neocidol)

Sivertsen, T.; and Sjøli, N. E., 1977, *Norsk Vet.-Tidsskr.*, v. 89 (12), 797-803
Lucilia caesar, sheep, diazinon: Hordaland

Diazinon

Tenquist, J. D.; and Roberts, M., 1978, *N. Zealand Vet. J.*, v. 26 (4), 88-91
Lucilia sericata, *Damalinea ovis*, sheep, 2 new 'Mini-shower' models of dipping with fenthion-ethyl and diazinon compared with plunge and shower dipping, residues in wool samples

Diazinon

Yeoman, G. H.; and Bell, T. A., 1978, *Vet. Rec.*, v. 103 (15), 337
Lucilia sericata, sheep, aluminium alkoxide gellants mixed with insecticide and applied to breech area, results suggest that this new control method against cutaneous myiasis gives higher protection than current means of control with no abnormal problems of toxicity, tissue residues, or wool processing

Diazinon DFF. See Diazinon.

- Dibenzo-18-crown-6
Brown, G. R.; and Foubister, A. J., 1979, J. Med. Chem., v. 22 (8), 997-999
benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against *Eimeria tenella*
- 3,3'-Dibrom-5,5'-dichlor-2,2'-dioxydiphenylsulfide
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,3'-Dibrom-5,5'-dichlor-2,2'-dioxydiphenylsulfone
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,3'-Dibrom-5,5'-dichlor-2,2'-dioxydiphenylsulfide
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- Dibromodulcitol
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 3,5-Dibromo-4'-ethoxysalicylanilide
Sakamoto, T.; and Gemmill, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives
- Dibromomannitol
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 3,5-Dibromosalicylanilide. See Bromsalans.
- 3,5-Dibromo-3'-trifluoromethylsalicylanilide.
See Fluorosalan.
- Dibromsalan. See Bromsalans.
- Dibromsulphen
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results
- 3-(Dibutylamino)-1[2,6-bis(trifluoromethylphenyl)-4-pyridyl]propanol -- WR-172,435.
WR-172,435
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus griseimembra*, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds
-
- α -(Di-N-butylaminoethyl)-2,8-(bis-trifluoromethyl)-4-quinolinemethanol, hydrochloride -- WR-177,504.
WR-177,504
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus griseimembra*, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*
-
- α -(Di-n-butylaminoethyl)-1,3-dichloro-6-trifluoromethyl-9-phenanthrenemethanol
Carroll, F. I.; Berrang, B.; and Linn, C. P., 1978, J. Med. Chem., v. 21 (4), 326-330
Plasmodium berghei, *P. cynomolgi*, experimental animals, resolution of antimalarial agents via complex formation with α -(2,4,5,7-tetra-nitro-9-fluorenylideneaminoxy)propionic acid, significant differences in toxicity
- α -Dibutylaminomethyl-2,6-bis(p-trifluoromethylphenyl)-4-pyridinemethanol
Bouwsmas, O. J.; Stewart, J. T.; and Capomacchia, A. C., 1978, J. Pharm. Sc., v. 67 (9), 1224-1228
 α -dibutylaminomethyl-2,6-bis(p-trifluoromethylphenyl)-4-pyridinemethanol, potential antimalarial agent, characterization of pharmacologically important species derived by electronic absorption and fluorescence spectroscopy
- α -(Di-n-butylaminomethyl)-2,6-bis(4-trifluoromethylphenyl)-4-pyridinemethanol
Carroll, F. I.; Berrang, B.; and Linn, C. P., 1978, J. Med. Chem., v. 21 (4), 326-330
Plasmodium berghei, *P. cynomolgi*, experimental animals, resolution of antimalarial agents via complex formation with α -(2,4,5,7-tetra-nitro-9-fluorenylideneaminoxy)propionic acid, significant differences in toxicity

- α -(Di-n-butylaminomethyl)-6,8-dichloro-2-(3',4'-dichlorophenyl)-4-quinolinemethanol
Carroll, F. I.; Berrang, B.; and Linn, C. P., 1978, J. Med. Chem., v. 21 (4), 326-330
Plasmodium berghei, P. cynomolgi, experimental animals, resolution of antimalarial agents via complex formation with α -(2,4,5,7-tetra-nitro-9-fluorenylideneaminoxy)propionic acid, significant differences in toxicity
- α -(Dibutylaminomethyl)-6,8-dichloro-2-(3',4'-dichlorophenyl)-4-quinolinemethanol
Stella, V.; et al., 1978, J. Pharm. Sc., v. 67 (10), 1375-1377
 α -(dibutylaminomethyl)-6,8-dichloro-2-(3',4'-dichlorophenyl)-4-quinolinemethanol (an antimalarial), enhancement of bioavailability by formulation with oleic acid in soft gelatin capsule
- N,N-Di-n-butyl-4-hexyloxy-1-naphthamide.
See Bunamidine.
- N,N-Dibutyl-4-hexyloxy-1-naphthamide hydrochloride. See Bunamidine.
- Dibutyltin dilaurate -- DBTD; Polystat (with Dinsed or Roxarsone or Sulfantran).
- Polystat
McDougald, L. R.; and McQuiston, T. E., 1978, Avian Dis., v. 22 (4), 765-770
Eimeria spp., turkeys, coccidiosis management, innate (age) resistance and acquired immunity vs. anticoccidial medication
- Dibutyltin dilaurate (DBTD)
Shlosberg, A.; and Egyed, M. N., 1979, Vet. and Human Toxicol., v. 21 (1), 1-3
coccidiostat dibutyltin dilaurate inadvertently introduced into feed, mass poisoning in cattle, palm doves, and mink: Israel
-
- Dicestal. See Dichlorophen.
- 5,5'-Dichlor-3,3'-dinitro-biphenyl-2,2'-diol.
See Niclofolan.
- 5,5'-Dichlor-2,2'-dioxydiphenylsulfide
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- '-Dichlor-2,2'-dioxydiphenylsulfone
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 5,5'-Dichlor-2,2'-dioxydiphenylsulfoxide
Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- Dichloro analog L-628,914. See 6-Amino-9-(2,6-dichlorobenzyl) purine.
- 9-(2,6-Dichlorobenzyl) adenine (ICI 120645)
Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture
- 9-(2,6-Dichlorobenzyl) adenine-1-N-oxide (ICI 120688)
Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture
- 2,4-Dichloro-6-(0-chloroanilino)-1,3,5-triazine
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scoliotic effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- 1,1-Dichloro-2-(o-chlorophenyl)-2-(p-chlorophenyl)ethane. See Mitotane.
- 6,8-Dichloro-2-(3',4'-dichlorophenyl)- α -(di-n-butylaminomethyl)-4-quinoline methanol -- SN-15,068; WR-30,090.
- WR-30,090
Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs
- WR-30,090 (SN-15,068)
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta

2,6-Dichloro-3,5-dicyano-4-phenyl pyridine
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

5,5'-Dichloro-2,2',dihydroxy-3,3'-dinitro-phenyl.
See Niclofolan.

3,5-Dichloro-2,6-dimethyl-4-pyridinol. See Meticlorpindol.

3,3'-Dichloro-5,5'-dinitro-0,0'-biphenol. See Niclofolan.

p,p-Dichlorodiphenyl-trichloromethyl-methane.
See DDT.

0,0-Di(2-chloroethyl)-0-(3-chloro-4-methylcoumarin-7-yl)phosphate. See Haloxon.

N-(Dichlorofluoromethylthio)-N-(dimethyl-sulfamoyl) aniline

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

2',5-Dichloro-4'-nitrosalicylanilide. See Niclosamide.

3,5-Dichloro-4'-nitrosalicylanilide
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolicalidal effect of salicylanilide and bisphenol derivatives

Dichlorophen -- Dicastal; Dichlosal (with Phenasaal); 2,2'-Dihydroxy-5,5'-dichlorophenyl methane; 2,2'-Methylenebis(4-chlorophenol).

Dichlorophen
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36

cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results

Dichlosal (= Phenasaal + Dichlorophen)
Grinenko, N. V.; et al., 1976, Med. Parazitol.

i Parazit. Bolezni, v. 45 (1), 101-103
H[ymenolepis] nana, patients of different age groups, dichlosal or trichlosal in divided doses, 5 or 7 day courses

Dichlosal
Grinenko, N. V.; et al., 1976, Med. Parazitol.

i Parazit. Bolezni, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasaal, trichlorophene, dichlosal, and trichlosal tested in graded doses

Dichlorophen -- Continued.

Dichlorophen (Dicastal)
Pandey, B. B.; and Rai, P., 1976, U. P. Vet. J., v. 4 (2), 74-77

Taenia hydatigena and Multiceps multiceps in puppies (exper.), Embelia ribes alcoholic extract not as effective as dichlorophen

Dichlorophen
Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128

Echinococcus multilocularis, various anthelmintics, scolicalidal effects in vitro and/or therapeutic effects in mice

2,2'-Methylenebis(4-chlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolicalidal effect of salicylanilide and bisphenol derivatives

3-(3,5-Dichlorophenyl)-5,5'-dimethyloxazoline dione-2,4

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

1-(3,4-Dichlorophenyl)-3-[4-(N-ethyl-3-piperidyl-amino)-6-methyl-2-pyrimidinyl] guanidine -- WR-81,844.

WR-81,844

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

(6,8-Dichloro-2-phenyl-4-quinoly)- α -piperidyl carbinoil hydrochloride (Merck SN 10,275)

Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, Rev. Brasil. Pesqui. Med. e Biol., v. 2 (1), 45-49

Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

1-(1,3-Dichloro-6-trifluoromethyl-9-phenanthryl)-3-di-(n-butyl) aminopropanol -- WR-171,669.

WR-171,669

Schmidt, L. H.; et al., 1978, Antimicrob.

Agents and Chemotherapy, v. 14 (3), 292-314

Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

- 2,2-Dichlorovinyl dimethyl phosphate. See
Dichlorvos.
- Dichlorquinazine (12-278)
Schmidt, L. H.; et al., 1977, Antimicrob.
Agents and Chemotherapy, v. 11 (5), 826-843
Plasmodium falciparum in Aotus trivirgatus,
activities of various 4-aminoquinolines
against chloroquine-resistant and -suscepti-
ble strains, observations confirm cross-
resistance among 4-aminoquinolines but in-
dicate that some derivatives may be thera-
peutically effective against infections
refractory to maximally tolerated doses of
chloroquine
- Dichlorvos -- Atgard; Atgard V; DDVF; DDVP; 2,2-
Dichlorovinyl dimethyl phosphate; O,O-Di-
methyl 2,2-dichlorovinyl phosphate; Equigard;
Marvex Super-100; No-Pest strips; Nuvan;
Nuvan 100 EC; PVC-DDVP; Strike Insect Strips;
Task; Vapona; Vaporet dog collars; Vapona
Strips.
- Dichlorvos (DDVP)
Allen, S. D.; VanKampen, K. R.; and Brooks,
D. R., 1978, Feline Pract., v. 8 (3), 9-10,
12-16
feline dichlorvos flea collars evaluated
for toxicity under conditions of high
temperature and low humidity, no hematologic
or neurologic changes
- Dichlorvos (Strike Insect Strips)
Beardles, M. L.; et al., 1978, J. Econom. Entom.,
v. 71 (2), 287-289
Haematobia irritans, cattle, effective con-
trol with dichlorvos-impregnated rear leg
bands, pasture test: Camp Stanley, Texas
- Dichlorvos
Beardles, M. L.; et al., 1979, Southwest.
Entom., v. 4 (1), 70-72
Haematobia irritans, cattle, comparative
efficacy of dichlorvos-impregnated ear tags,
leg bands, and tail tags
- Nuvan (Dichlorvos)
Chakrabarti, A.; and Misra, S. K., 1979, In-
dian Vet. J., v. 56 (6), 497-500
Demodex canis, dogs, incidence in relation
to season, host age, sex, and breed, clinical
manifestations, in vivo and in vitro
activity of several acaricides: India
- Nuvan 100 EC (Dichlorvos; DDVP)
Chellappa, D. J.; Subramanian, R.; and Gopala-
krishnan, C. A., 1977, Indian Poultry Gaz.,
v. 61 (4), 135-137
Menopon gallinae, Lipeurus caponis, poultry,
malathion 50 EC, sumithion 50 EC, nuvan 100
EC, drug trial, sumithion 50 EC most effec-
tive, knapsack sprayer superior to conven-
tional dipping procedure, preliminary report
- Dichlorvos (Vapona)
Cole, M. M.; et al., 1976, J. Med. Entom.,
v. 12 (6), 625-630
rodent fleas, dichlorvos impregnated in
granules coated with rodent bait evaluated
as vapor toxicant for flea control, field
tests, effective flea control on Dipodomys
spectabilis probably due to its habit of
storing food in its burrow: southeast New
Mexico
- Dichlorvos -- Continued.
- Dichlorvos
Coppedge, J. R.; et al., 1978, J. Econom.
Entom., v. 71 (3), 483-486
Cochliomyia hominivorax, development and
field evaluation of bait system (SWASS)
containing dichlorvos and bait for suppres-
sion of adult screwworms
- Dichlorvos
Coppedge, J. R.; et al., 1978, J. Econom.
Entom., v. 71 (4), 579-584
Cochliomyia hominivorax, evaluation of screw-
worm adult suppression system (SWASS), a
bait-toxicant system using swormlure-2 and
dichlorvos, elimination by SWASS coupled
with later release of sterile screwworms:
Curacao
- Dichlorvos
Darrow, D. I., 1973, J. Econom. Entom.,
v. 66 (1), 133-135
Bovicola spp. on goats, control with di-
chlorvos-impregnated resin neck collars
- Dichlorvos
Ducos de Lahitte, J.; Dorchies, P.; and Franc,
M., 1978, Prat. Vet. Equine, v. 10 (4), 157-159
gastro-intestinal strongylosis, foals,
weight gains, effect of treatment
- Dichlorvos
Frazar, E. D.; and Schmidt, C. D., 1979, J.
Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, sus-
ceptibility to topically applied insecticides
- Dichlorvos (No-Pest-Strip; Vapona Strip)
Gibson, L., 1978, Avicult. Mag., v. 84 (1),
56-57
lung mites in Chloebia gouldiae (trachea),
symptoms, successfully treated with No-Pest-
Strip (19 % dichlorvos) hung in cage: im-
ported from England
- Dichlorvos
Gladney, W. J.; Price, M. A.; and Graham, O.
H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of
various insecticides applied as sprays, ear
smears and dusts, or in slow-release devices,
field tests
- Equigard
Hawthorne, H. W., 1978, Zool. Garten, n. F.,
v. 48 (4), 269-270
equigard, effective use as broad spectrum
anthelmintic in exotic and domestic rumi-
nants, methods of administering drug,
precautions: Catskill Game Farm, New York
- Dichlorvos (Equigard)
Keller, H.; and Mueller, R., 1979, Berl. u.
Munchen. Tierarztl. Wchnschr., v. 92 (4), 63-
65
dichlorvos, trichlorfon, horses, decrease in
plasma cholinesterase activity, concluded
that relaxation of horses with succinyl-
choline should not be carried out within 10
days after exposure to organophosphorus type
anthelmintics

Dichlorvos -- Continued.

Nuvan

- Khan, M. H., 1979, Indian Vet. J., v. 56 (9), 739-743
Lipeurus caponis, *Menacanthus stramineus*, White Leghorn fowl, organophosphorus insecticides, costs evaluated

Marvex Super-100

- Khan, M. H., 1979, Indian Vet. J., v. 56 (9), 739-743
Lipeurus caponis, *Menacanthus stramineus*, White Leghorn fowl, organophosphorus insecticides, costs evaluated

DDVP

- Khan, D.; and Haseeb, M. A., 1976, Pakistan J. Zool., v. 8 (2), 173-176
Ganeo micracetabulus and *Cercaria reflexicauda cercariae*, effects of 5 insecticides at various concentrations, toxicity varies but results indicate cercariae are susceptible to insecticides

Dichlorvos (DDVP)

- Kibble, R., 1979, Austral. Vet. J., v. 55 (1), 40-41 [Letter]
 fleas, ticks, dogs, DDVP and propoxur-impregnated collars, recommendations: Australia

Dichlorvos

- Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Dichlorvos

- Lensink, B. M.; Rijpstra, A. C.; and Erken, A. H. M., 1979, Zool. Garten, n. F., v. 49 (2), 121-126
Ollulanus tricuspis in *Panthera tigris tigris* (vomitus, stomach), clinical symptoms in mother and offspring, treatment with various anthelmintics, complete recovery achieved with levamisole: Artis-Zoo, Amsterdam

Dichlorvos (Atgard V)

- Marti, O. G.; Stewart, T. B.; and Hale, O. M., [1979], J. Parasitol., v. 64 (6), 1028-1031
 gastrointestinal nematodes, pigs raised under similar management conditions, comparative efficacy of fenbendazole, dichlorvos, and levamisole HCl

DDVP

- Migunov, I. M.; Timofeev, P. V.; and Grudin, B. L., 1977, Veterinariia, Moskva (6), 68-69
Melophagus ovinus, sheep, DDVP effective, applied as aerosol to group of sheep in sealed enclosure

Dichlorvos (DDVP; Vapona)

- Niemand, H. G.; Niemand, S.; and Wendel, E., 1979, Kleintier-Praxis, v. 24 (4), 173-175
 discussion of chief effective principles of different flea collars (dichlorvos, diazinon, propoxur), toxicity

Dichlorvos -- Continued.

- O,O-Dimethyl 2,2-dichlorovinyl phosphate (DDVP)
 Reiner, E.; et al., 1978, Comp. Biochem. and Physiol., v. 60C (2), 155-157
Metastrongylus apri, cholinesterase, kinetic properties with respect to substrate hydrolysis and inhibition by organophosphorus compounds

Dichlorvos (Task)

- Remfry, J., 1978, Lab. Animals, v. 12 (4), 213-218
 helminth infections in imported *Macaca mulatta*, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Dichlorvos (Atgard)

- Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235
Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parbendazole and dichlorvos

Dichlorvos

- Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Dichlorvos

- Schmidt, R. L.; et al., 1979, J. Wildlife Management, v. 43 (2), 461-467
Protostrongylus, domesticated, captive, or free-ranging *Ovis c. canadensis*, evaluation of 5 drugs

Dichlorvos (DDVP; PVC-DDVP)

- Stanton, H. C.; Albert, J. R.; and Mersmann, H. J., 1979, Am. J. Vet. Research, v. 40 (3), 315-320
 dichlorvos, sows and growing pigs, unformulated vs. slow-release polyvinyl chloride resin formulation, pharmacology and safety

Dichlorvos (Vapona)

- Tharaldsen, J.; and Grøndalen, J., 1978, J. Small Animal Practice, v. 19 (4), 245-250
Pneumonyssus caninum, dog (nostrils), case report, dichlorvos: Norway

Dichlorvos

- Yeoman, G. H.; and Bell, T. A., 1978, Vet. Rec., v. 103 (15), 337
Lucilia sericata, sheep, aluminium alkoxide gellants mixed with insecticide and applied to breech area, results suggest that this new control method against cutaneous myiasis gives higher protection than current means of control with no abnormal problems of toxicity, tissue residues, or wool processing

Dichlorvos (Atgard)

- Young, R., jr.; Hass, D. K.; and Brown, L. J., 1979, J. Animal Sc., v. 48 (1), 45-51
Ascaris suum, *Oesophagostomum dentatum*, dichlorvos administered to non-parasitized and parasitized sows during late gestation, improved reproductive performance found to be independent of anthelmintic activity

Dichlosal. See Dichlorophen or Niclosamide.

Dicofol -- Kelthane.

Dicofol (Kelthane)
El-Bahay, G. H.; et al., 1978, Vet. Med. J.,
Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, Argas persicus,
evaluation of 10 insecticides

Dicofol
Rawlins, S. C.; and Mansingh, A., 1978, J.
Econ. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains,
patterns of resistance to acaricides

Dicresil

Puchkova, E. A., 1977, Veterinariia, Moskva
(7), 19-22
D[ermanyssus] gallinae, C[imex] lectularius,
lice, chickens on industrial scale farms,
control, sevin, dicresil, chlorophos,
carbophos; other complex sanitation measures

Dicrotophos

Rawlins, S. C.; and Mansingh, A., 1978, J.
Econ. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, suscepti-
bility to acaricides: Jamaica; St. Kitts;
Trinidad; Guyana

Dicrotophos

Rawlins, S. C.; and Mansingh, A., 1978, J.
Econ. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains,
patterns of resistance to acaricides

Dicumarol

Brotherton, J., 1978, Arzneimittel-Forsch.,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

2,3-Dicyano-1,4-dithioanthraquinone

Sakamoto, T.; and Gemmill, M. A., 1979, Mem.
Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
130
Echinococcus granulosus, scolicidal effect
of 65 antibiotic, antineoplastic, cytosta-
tic, and other agents in vitro

1,5-Di-(2,4-dimethylphenyl)-3-methyl-1,3,5-tri-
azapenta-1,4-diene. See Amitraz.

Dieldrin -- 1,2,3,4,10,10-Hexachloro-6,7-epoxy-
1,4,4a,5,6,7,8,8a-octahydro-endo-1,4-exo-
5,8-dimethano-naphthalene.

Dieldrin

Blommers, L., 1979, J. Med. Entom., v. 16 (1),
82-83
Pediculus capitis, nymph rearing technique,
insecticide tests against nymphs

Dieldrin

Blommers, L.; and van Lennep, M., 1978, Entom.
Exper. et Applic., v. 23 (3), 243-251
Pediculus humanus capitis, school children,
field trials with lindane, laboratory tests
with lindane, dieldrin, and malathion,
presence of resistance to lindane confirmed:
Netherlands

Dieldrin -- Continued.

Dieldrin

Crampton, P. L.; and Gichanga, M. M., 1979,
Bull. Entom. Research, v. 69 (3), 427-439
economically important Ixodidae from major
cattle-raising areas, survey of resistance
to organochlorine and organophosphorus acari-
cides: Kenya

Dieldrin

Khan, D.; and Haseeb, M. A., 1976, Pakistan J.
Zool., v. 8 (2), 173-176
Ganeo micracetabulus and Cercaria reflexi-
cauda cercariae, effects of 5 insecticides
at various concentrations, toxicity varies
but results indicate cercariae are suscepti-
ble to insecticides

Dieldrin

Kiefer, M.; and Cyprich, D., 1978, Acta Fac.
Rerum Nat. Univ. Comenianae, Zool. (23), 5-12
ticks, susceptibility to acaricides: Slo-
vakia

Dieldrin

Lourens, J. H. M., 1979, J. Econom. Entom.,
v. 72 (5), 790-793
Amblyomma variegatum, A. lepidum, baseline
data on susceptibility to organochlorine
acaricides, genetic basis for resistance in
A. variegatum

Dieldrin

Lourens, J. H. M.; and van de Klashorst, G.,
1979, Ztschr. Ang. Entom., v. 87 (3), 230-238
Rhipicephalus appendiculatus, organochlorine
susceptible and tolerant populations: East
Africa

Dieldrin

Lourens, J. H. M.; and Tatchell, R. J., 1979,
Bull. Entom. Research, v. 69 (2), 235-242
Rhipicephalus evertsi evertsi, identification
and inheritance of resistance factors to
organochlorine acaricides, experimental
hybrids between susceptible and resistant
strains

Dieldrin

Uspenskii, I. V., 1974, Parazitologia, Lenin-
grad, v. 8 (4), 312-321
Ixodes persulcatus, susceptibility to acari-
cides

1-[[2-(Diethylamino) ethyl] amino]-4-(hydroxy-
methyl)-thioxanthen-9-one. See Hycanthone.

5-[[Diethylamino)ethyl]amino]-8-methoxyquinoline
dihydrochloride hemihydrate

Khan, M. Sami; and LaMontagne, M. P., 1979, J.
Med. Chem., v. 22 (8), 1005-1008
3- and 5-aminoquinolines, potential antimalar-
ials, synthesis, testing of some against
Plasmodium berghei in mice, Leishmania dono-
vani in hamsters, or P. cynomolgi in rhesus
monkeys

1-β-Diethylamino-ethylamino-4,6,8-trimethyl-5-azathioxanthone -- Ciba 17'581.

Ciba 17'581

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Diethylaminoethyl-dextran 500

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

2-Diethylaminoethyl-2,2-diphenylvalerate hydrochloride. See Proadifen hydrochloride.

4-[[6-(Diethylamino)hexyl]amino]-2-methoxy-9-methylacridine

Scovill, J. P.; et al., 1979, *J. Med. Chem.*, v. 22 (10), 1164-1167

Leishmania donovani in golden hamster, derivatives of 4-amino-2-methoxyacridine

8-(6-Diethylamino-hexylamino)-6-methoxy-4-methylquinoline -- WR 6026.

WR 6026

Bjorvatn, B.; and Neva, F. A., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (3), 480-485
Leishmania tropica, white mouse model, experimental therapy using sodium stibogluconate, amphotericin B, metronidazole and WR 6026

8-(6-Diethylamino-hexylamino)-6-methoxy-4-methylquinoline (WR 6026)

Kinnamon, K. E.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (4), 751-757

Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

6-(4-Diethylamino-1-methylbutylamino)-5,8-dimethoxyquinoline, 2- and 4-amino derivatives

Nickel, P.; et al., 1978, *Arzneimittel-Forsch.*, v. 28 (4), 578-581

Plasmodium vinckei, synthesis of 2- and 4-amino derivatives of 6-(4-diethylamino-1-methylbutylamino)-5,8-dimethoxyquinoline, laboratory trials in mice for possible anti-malarial activity

5-[[4-(Diethylamino)-1-methylbutyl]amino]-8-methoxyquinoline trihydrochloride dihydrate

Khan, M. Sami; and LaMontagne, M. P., 1979, *J. Med. Chem.*, v. 22 (8), 1005-1008

3- and 5-aminoquinolines, potential antimalarials, synthesis, testing of some against Plasmodium berghei in mice, Leishmania donovani in hamsters, or P. cynomolgi in rhesus monkeys

0-[2-(Diethylamino)-6-methyl-4-pyrimidinyl]-0,0-dimethyl phosphorothioate. See Pirimiphos-methyl.

Diethylcarbamazine -- Banminth D (with Morantel);

Banocide; Caricide; DEC; Diethylcarbamazine citrate; 1-Diethylcarbamoyl-4-methylpiperazine; 1-Diethylcarbamyl-4-methyl-piperazine; Ditrazine; Diroicide; Filarzan; Franocid; Hetrazan; Notezine.

Diethylcarbamazine (Banocide; Hetrazan; DEC)

Allen, G. D.; Goodchild, T. M.; and Weatherly, B. C., 1979, *J. Chromatography (Biomed. Appl.)*, v. 164 (4), 521-526

diethylcarbamazine, determination of concentrations in human plasma and urine

Diethylcarbamazine citrate (Hetrazan)

Baqui, A.; and Ansari, J. A., 1976, *Kiseichugaku Zasshi (Japan. J. Parasitol.)*, v. 25 (5), 409-414

Setaria cervi, white rats, hetrazan, thia-bendazole, tetramisole

Diethylcarbamazine (DEC)

Bartholomew, C. F.; Nathan, M. B.; and Tikasingh, E. S., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (4), 423-424

Mansonella ozzardi, human, diethylcarbamazine (DEC), no microfilaricidal effect; effective against Wuchereria bancrofti microfilariae in mixed infections; infection with M. ozzardi is unlikely to interfere with Mazzotti test for Onchocerca volvulus or with DEC provocative test in daytime surveys of periodic W. bancrofti: Trinidad

Diethylcarbamazine

Cassar, E. J., 1979, *Austral. and N. Zealand J. Med.*, v. 9 (5), 600 [Letter]

A[ngiostrongylus] cantonensis, man, case report, probable cause of eosinophilic meningitis, diethylcarbamazine: Brisbane

Diethylcarbamazine (DEC)

Chandrasekaran, B.; and Harinath, B. C., 1979, *Indian J. Med. Research*, v. 70, 305-308

isolation, characterization, and quantitation of diethylcarbamazine-N-oxide from human urine

Ditrazine

Daniyarov, I. A.; et al., 1978, *Veterinariia, Moskva* (2), 64-65

Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Diethylcarbamazine

Delving, M., 1978, *Vet. Rec.*, v. 102 (25), 562 [Letter]

Setaria labiatopapillosa, infertility in a female Bosman's potto, successful treatment with diethylcarbamazine: imported to zoo near Harrogate, England

Diethylcarbamazine -- Continued.

Diethylcarbamazine

Denham, D. A.; et al., 1978, J. Parasitol., v. 64 (3), 463-468

Brugia pahangi, anthelmintic effect of diethylcarbamazine in vitro, in *Aedes aegypti*, in *Meriones unguiculatus*, and in *Felis catus*, implications for use of primary and secondary screens for filaricidal activity

Diethylcarbamazine

Desowitz, R. S.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (6), 1148-1151

Dirofilaria immitis-infected dogs with severe adverse reactions after diethylcarbamazine treatment, rapid and marked decrease in precipitating and reaginic antibodies, possible model for reactions in human filariasis

Diethylcarbamazine citrate

Ewert, A.; and Emerson, G. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 496-499

Brugia malayi, fourth stage and adult parasites in cats, effect of diethylcarbamazine citrate, effective model to compare efficacy of drugs against adult lymphatic-dwelling filariae

Diethylcarbamazine (Notezine)

Gayral, P.; et al., 1978, Ann. Parasitol., v. 53 (6), 669-674

Dipetalonema dessetae in *Proechimys oris*, microfilaricidal action of diethylcarbamazine

Diethylcarbamazine

Hedjazi, M.; and Mirzayans, A., 1978, Rev. Med. Vet., Toulouse, v. 129 (12), 1685-1691

Parafilaria multipapillosa, horses, clinical aspects, diethylcarbamazine: Teheran, Iran

Diethylcarbamazine citrate

Holt, P. E.; Cooper, J. E.; and Needham, J. R., 1979, J. Small Animal Practice, v. 20 (5), 269-286

Angusticaecum spp., *Tachygonetria* sp., and *Atractis dactyluris* in *Testudo*, treatment: Britain

Diethylcarbamazine (DEC)

Hyodo, T.; et al., 1978, Nippon Zyuisi-Kai Zassi (J. Japan Vet. Med. Ass.), v. 31 (8), 454-458

diethylcarbamazine, dogs, effects of long-term administration

Banminth D

Kadyrov, N. T., 1978, Veterinariia, Moskva (7), 57-58

Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Diethylcarbamazine

Kale, O. O., 1979, J. Helminth., v. 53 (2), 169-174

Onchocera volvulus, human, effect of graded doses of diethylcarbamazine on concentration of microfilariae in hydrocoele fluid, urine, and skin

Diethylcarbamazine -- Continued.

Diethylcarbamazine

Kazacos, K. R.; and Smith, L. E., jr., 1979, Am. J. Trop. Med. and Hyg., v. 28 (2), 213-215

Loa loa, 27-year-old Purdue University student from Sudan who had lived in United States for 4 years, case report, successful treatment with diethylcarbamazine, possible public health significance: Indiana

Diethylcarbamazine (Franocid)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73

Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Diethylcarbamazine

Laigret, J.; Fagneaux, G.; and Tuira, E., 1978, Bull. World Health Organ., v. 56 (6), 985-990

Wuchereria bancrofti var. *pacifica*, humans with lymphatic filariasis, diethylcarbamazine long-term therapy given at widely spaced intervals: Polynesie francaise

Diethylcarbamazine (Banocide; DEC)

Langham, M. E.; Traub, Z. D.; and Richardson, R., 1978, Tropenmed. u. Parasitol., v. 29 (2), 156-162

Onchocerca volvulus, humans, transepidermal administration of diethylcarbamazine, more effective and safer than oral route: Liberia

Diethylcarbamazine

de Leao, R. N. Q.; et al., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (5), 300-306

Lagochilascaris minor, 14-year-old boy, clinical aspects, pathology, successful therapy with diethylcarbamazine

Diethylcarbamazine

McMahon, J. E., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 465-472

Wuchereria bancrofti, humans, preliminary screening of levamisole and amodiaquine for antifilarial activity; some trials combined with diethylcarbamazine

Diethylcarbamazine

Matsuda, H.; Takaoka, M.; and Tanaka, H., 1976, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 25 (2), 94-99

Litomosoides carinii and *Dipetalonema witei* microfilariae, diethylcarbamazine, *Meriones unguiculatus*

Diethylcarbamazine

Meyers, W. M.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (6), 1137-1147

Dipetalonema streptocerca, human, diethylcarbamazine therapy, degeneration of adult worms, may thus produce radical cures and may serve as chemoprophylactic: Zaire

Diethylcarbamazine

Narasimham, M. V. V. L.; et al., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (4), 571-575

Wuchereria bancrofti, human, comparison of levamisole, levamisole + mebendazole, and diethylcarbamazine: Calicut, Kerala, India

Diethylcarbamazine -- Continued.

- Diethylcarbamazine
Nesarajah, M. S., 1975, *Thorax*, v. 30 (5), 574-577
tropical eosinophilia, humans, pulmonary function before and after treatment with diethylcarbamazine
- Diethylcarbamazine (Filarzan, DEC)
Partono, F.; Purnomo; and Soewarta, A., 1979, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 73 (5), 536-542
Brugia timori, mass therapy with diethylcarbamazine followed 1 year later by short term selective re-treatment, simple control method: Flores, Indonesia
- Diethylcarbamazine
Piessens, W. F.; and Beldekas, M., 1979, *Nature*, London (5741), v. 282, 845-847
Brugia malayi, diethylcarbamazine enhances antibody-mediated cellular adherence to microfilariae
- Diethylcarbamazine
Prescott, C. W.; O'Grady, A. S. B.; and English, P. B., 1978, *Austral. Vet. J.*, v. 54 (8), 404-405 [Letter]
Dirofilaria immitis, dose rate of diethylcarbamazine for heartworm prophylaxis
- Diethylcarbamazine + Levamisole
Prod'hon, J.; et al., 1979, *Med. Trop.*, v. 39 (6), 631-635
Onchocerca volvulus, humans, evaluation of 2 therapeutic schemes using diethylcarbamazine combined with levamisole, recommendations for optimum therapy: Haute Volta
- Diethylcarbamazine
Prod'hon, J.; Moreau, J. P.; and Mongin, C., 1979, *Med. Trop.*, v. 39 (6), 619-628
Onchocerca volvulus, humans, mass therapy trials, diethylcarbamazine and levamisole, administered separately or combined: Haute Volta
- Diethylcarbamazine
Rougemont, A.; et al., 1978, *Compt. Rend. Soc. Biol.*, Paris, v. 172 (2), 397-402
Onchocerca volvulus microfilariae under the influence of diethylcarbamazine, ultrastructural observations
- Diethylcarbamazine-Bithionol sulfoxide
Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives
- Diethylcarbamazine
Sethumadhavan, K. V. P.; et al., 1978, *Indian J. Med. Research*, v. 67, 759-762
[*Wuchereria bancrofti*, human carriers, attempted clearance of microfilaremia using diethylcarbamazine: Calicut town, Kerala
- Ditrazine
Shedivtsova, A., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Diethylcarbamazine -- Continued.

- Diethylcarbamazine (DEC)
Shibuya, T.; et al., 1978, *Japan. J. Exper. Med.*, v. 48 (5), 411-418
Litomosoides carinii in *Sigmodon hispidus*, screening filaricides for human filariasis, evaluation of intrathoracic injection method
- Diethylcarbamazine (Hetrazan)
Silverman, N. H.; Katz, J. S.; and Levin, S. E., 1973, *South African Med. J.*, v. 47 (6), 219-221
Capillaria hepatica in 17-month-old female, clinical presentation of visceral larva migrans, liver biopsy confirmed diagnosis, diethylcarbamazine and sodium antimonyl gluconate improved but did not cure infection: district of Bethal, South Africa
- Diethylcarbamazine
Singhal, K. C.; Madan, B. R.; and Saxena, P. N., 1978, *Indian J. Physiol. and Pharmacol.*, v. 22 (1), 93-97
Setaria cervi, in vitro, action of diethylcarbamazine on parasite spontaneous movement and glucose consumption
- Diethylcarbamazine
Sowa, J.; and Sowa, S. C. I., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (1), 79-85
Onchocerca volvulus, children, long-term treatment with low doses of diethylcarbamazine: Ghana
- Diethylcarbamazine (DEC)
Tanaka, H.; et al., 1977, *Japan. J. Exper. Med.*, v. 47 (4), 315-317
Litomosoides carinii-infected cotton rats, improved method for intrapleural injection of anti-filarial drugs to evaluate macrofilaricidal action
- Diethylcarbamazine
Thompson, P. E.; Zeigler, J. B.; and McCall, J. W., 1973, *Antimicrob. Agents and Chemotherapy*, v. 3 (6), 693-697
Litomosoides carinii in Mongolian jirds and cotton rats, 3-ethyl-8-methyl-1,3,8-triazabicyclo[4,4,0]decan-2-one and diethylcarbamazine compared, strong but transient effect against microfilariae but no appreciable effect in killing adult worms
- Diethylcarbamazine (Caricide)
Verma, B. B.; and Sinha, B. P., 1978, *Indian Vet. J.*, v. 55 (5), 411-412
Dracunculus medinensis, dog (left side just behind axilla), case report, effectively treated with diethylcarbamazine
- Diethylcarbamazine
Weller, P. F.; and Ottesen, E. A., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (1), 31-32
Wuchereria bancrofti, in persons with sub-periodic infections diethylcarbamazine fails to provoke marked increase in circulating microfilariae immediately after therapy is initiated; in contrast, in persons with nocturnally periodic infections there is significant increase

Diethylcarbamazine -- Continued.

Dirocide

Yaros, K.; and Hale, J. E., 1978, Vet. Med. and Small Animal Clin., v. 73 (6), 684 [Letter] polydipsia and polyuria in diabetic dog given diroicide syrup (heartworm prophylaxis) containing glucose, condition restabilized when given diroicide in tablet form

Diethylcarbamazine

Zahner, H.; et al., 1978, Tropenmed. u. Parasitol., v. 29 (1), 15-26
Litomosoides carinii in Mastomys natalensis (exper.), effect of diethylcarbamazine against microfilariae in several organs; dynamics of cell adhesion, immobilization, and elimination of microfilariae

Diethylcarbamazine citrate. See Diethylcarbamazine.

1-Diethylcarbamoyl-4-methylpiperazine. See Diethylcarbamazine.

1-Diethylcarbamy-4-methyl-piperazine. See Diethylcarbamazine.

0,0-Diethyl-(6-chlorbenzoxazoliny-3-methyl)dithiophosphate. See Phosalone.

0,0-Diethyl 0-(3-chloro-4-methyl-2-oxo-2H-1-benzopyran-7-yl) phosphorothioate. See Coumaphos.

0,0-Diethyl-s-(5,7-dichlorobenzoxazol-2-ylmethyl)-dithio-phosphate. See Benoxafos.

Diethyl 1-(2,4-dichlorophenyl)-2-chlorovinyl-phosphate. See Chlorfenvinphos.

Diethyldithiocarbamate

Scheibel, L. W.; Adler, A.; and Trager, W., 1979, Proc. National Acad. Sc., v. 76 (10), 5303-5307
Plasmodium falciparum, antimalarial effects of tetraethylthiuram disulfide and its reduction product diethyldithiocarbamate

0,0-Diethyl-0-(2-isopropyl-6-methyl-4-pyrimidinyl). See Diazinon.

0,0-Diethyl 0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate. See Diazinon.

0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate. See Diazinon.

0,0-Diethyl-0-2-isopropyl-4-methyl-6-pyrimidyl phosphorothionate. See Diazinon.

0,0-Diethyl-0-(2-isopropyl-4-methyl-pyrimidyl-6) thiophosphate. See Diazinon.

0,0-Diethyl-0-4(methylmercapto-3-methylphenyl) thiophosphate. See Fenthion ethyl.

0,0-Diethyl 0-[6-methyl-2-(1-methylethyl)-4-pyrimidinyl] phosphorothioate. See Diazinon.

0,0-Diethyl-0-naphthaloximide phosphate. See Phthalophos.

Diethyl 4,4'-0-phenylene bis (3-thioallophanate). See Thiophanate.

3,5-Diethyl-phenyl-N-methyl-carbamate. See Phenetcarb.

Diethylstilbestrol -- Gynben vaginal cream (with Diiodohydroxyquin and Sulfadiazine).

Gynben vaginal cream

Aguiar, L. M.; Bastos, A. da C.; and Salvattore, C. A., 1977, Rev. Brasil. Med., v. 34 (3), 151-152
vaginal trichomoniasis, humans, gynben vaginal cream, excellent antipruritic effect and good tolerance

Diethylstilbestrol

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Diethyl [thio [0-[3-(p-tolylsulphonyl) ureido] phenyl] carbamoyl] phosphoramidate. See Diuredosan.

0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate. See Chlorpyrifos.

Difetarstone. See Diphetarstone.

Difezil. See Diphezil.

Diflubenzuron -- N-[(4-Chlorophenyl)amino]carbonyl-2,6-difluorobenzamide.

Diflubenzuron

Hopkins, D. E.; and Chamberlain, W. F., 1978, Southwest. Entom., v. 3 (4), 292-294
Melophagus ovinus, inhibited maturation on sheep dipped in diflubenzuron (preventing formation of pupae) or methoprene (preventing adult eclosion), possible candidate control agents for sheep ked and other larviparous pests

Diflubenzuron

Hopkins, D. E.; and Chamberlain, W. F., 1978, J. Econom. Entom., v. 71 (1), 25-26
Bovicola limbatus, 3rd instars, relationship between inhibition of ecdysis and time and quantity of ingestion of diflubenzuron (inhibitor of cuticle deposition), inhibition of ecdysis increased progressively as age of nymphs increased, timing of treatment important for control

Diformyl dapsone -- WR-6,798.

WR-6,798

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus griseimembra*, methods employed in search for new blood schizonticidal drugs

2,3-Dihydro-2,2'-dimethyl-2H-naphtho [1,2-b]-furan-4,5-dione. See Nor- β -lapachone.

3,4-Dihydro-2,2'-dimethyl-2H-naphtho [1,2-b]pyran-5-one-6-spiro-2'-oxyrane. See Methylene- β -lapachone.

S(2,3-Dihydro-5-methoxy-2-oxo-1,3,4-thiadiazol-3-ylmethyl) dimethyl phosphorothiothionate. See Methidathion.

4,5-Dihydro-3-methyl-1,2,4-oxadiazole
 Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

2,3-Dihydro-1H-pyrazolo-(2,3-a)imidazole
 Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

2,2'-Dihydroxy-5,5'-dichlorophenyl methane. See Dichlorophen.

9-(8-DL-2 α ,3 α -Dihydroxy-4 β -[hydroxymethyl]-cyclopentyl)-adenine. See Cycloadenosine.

4,6-Dihydropyrazolo(3,4-d)pyrimidine. See Oxypurinol.

2,2'-Dihydroxy-3,3',5,5'-tetrabromo-1,1'biphenylmono (dihydrogen phosphate). See Bromophenophos.

3,5-Diiodo-3'-chloro-4'-(p-chlorophenoxy)-sali-cylanilide. See Rafoxanide.

Diiodohydroxyquin -- Contramibial (with Chloroquine and Tetracycline); Diiodohydroxyquinoline; Di-iodoquine; Gynben vaginal cream (with Diethylstilbestrol and Sulfadiazine); Metodine (with Metronidazole); Resotren [composite] (with Chloquinat and Chloroquin phosphate).

Gynben vaginal cream

Aguiar, L. M.; Bastos, A. da C.; and Salvatore, C. A., 1977, Rev. Brasil. Med., v. 34 (3), 151-152

vaginal trichomoniasis, humans, gynben vaginal cream, excellent antipruritic effect and good tolerance

Diiodohydroxyquin -- Continued.

Di-iodoquine

Bhandari, B.; and Sankhla, K., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (3), 345-346 [Letter]

Entamoeba histolytica-associated rectal prolapse in children, di-iodoquine and metronidazole

Diiodohydroxyquinoline + Metronidazole (= Metodine)

Botero Ramos, D., 1976, Semana Med. Mexico (1127), an. 23, v. 89 (7), 209-212

Entamoeba histolytica, human, clinical trials testing efficacy of metronidazole combined with diiodohydroxyquinoline

Contramibial

Ingelet, B., 1974, Rev. Infirm. Afrique Noire (26), 17-19

human amoebic colitis, clinical trials testing contramibial given orally over a 4-day period, good therapeutic results

Di-iodohydroxyquinoline + Oxytetracycline

Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101

Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire

Di-iodohydroxyquinoline + Oxytetracycline + Dehydroemetine

Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101

Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire

Diiodohydroxy quinoline + Chloroquin phosphate + Chloquinat (=Resotren [composite])

Samaddar, J.; Biswas, D. N.; and Ghose, A. N., 1978, Indian Vet. J., v. 55 (7), 572-576

leucocytozoonosis, W[white] L[eg] H[orn] birds, quinine bisulphate and resotren failed to ensure absolute recovery, though general condition of treated birds improved

Diiodohydroxyquin

Spencer, M. J.; Garcia, L. S.; and Chapin, M. R., 1979, Am. J. Dis. Child., v. 133 (4), 390-393

Dientamoeba fragilis, children with acute and chronic gastrointestinal symptoms, clinical findings, symptomatic recovery after treatment with diiodohydroxyquin or metronidazole indicates pathogenic role

Diiodohydroxyquin

Thacker, S. B.; et al., 1979, Am. J. Pub. Health, v. 69 (12), 1279-1281

Entamoeba histolytica, *Giardia lamblia*, control attempts in a residential facility for mentally retarded persons: Washington, D. C.

Diiodohydroxyquinoline. See Diiodohydroxyquin.

2,6-Diiodo-4-nitrophenol. See Disophenol.

Di-iodoquine. See Diiodohydroxyquin.

trans-1,4-Di-(2-isothiocyanatoethyl) cyclohexane -- Cetovex.

Cetovex

Georgieva, D., 1978, Vet. Sbirka, v. 76 (4), 32-33
anthelmintic treatment, lambs, weight gains

Diloxanide furoate (Furamide)

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Diloxanide furoate (Furamide)

Salaki, J. S.; Shirey, J. L.; and Strickland, G.T., 1979, Am. J. Trop. Med. and Hyg., v. 28 (2), 190-193
Entamoeba polecki, 24-year-old Peace Corps volunteer (feces), symptomatic intestinal infection cured with diloxanide furoate and metronidazole: United States (previously in Upper Volta)

Dimecron. See Phosphamidon.

1,4-Dimethane sulfonyl butane

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Dimethoate -- Cygon; O,O-Dimethyl S-[2-(methylamino)-2-oxoethyl] phosphorodithioate; O,O-Dimethyl S-(N-methylcarbamoylmethyl) phosphorodithioate; Rogor; Roxion.

Dimethoate (Cygon)

Dohany, A. L.; Cromroy, H. L.; and Cole, M. M., 1977, J. Med. Entom., v. 14 (1), 79-81
Eutrombicula alfreddugesi, guinea pigs and cotton rats (Sigmodon hispidus), dimethoate, determination of correct dosage for control

Dimethoate (Rogor)

El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, Argas persicus, evaluation of 10 insecticides

Dimethoate (Rogor; Cygon; Roxion)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142
Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Dimethoate

Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661
flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southern New Mexico

Dimethoate -- Continued.

Dimethoate

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Dimethoate

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Dimethoate

Trosper, J. H., 1979, Southeast Asian J. Trop. Med. and Pub. Health, v. 10 (1), 62-66
Leptotrombidium deliense, rodents, dimethoate for control of chiggers in endemic scrub typhus area, unsatisfactory results: Pescadores Islands of Taiwan

O,O-Dimethyl-S-2-(acetylamino) ethyldithiophosphate

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

O,S-Dimethyl acetyl-phosphoramidothioate. See Acephate.

p-Dimethylaminobenzonitrile + Sulfaquinoxaline (= Nitryl)

Mladenovic, Z.; Movsesijan, M.; and Borojevic, D., 1978, Vet. Glasnik, v. 32 (10), 829-834
Eimeria spp., chickens (exper.), mixed infections, cycostat, nitryl, and stenorol

4-(2-Dimethylaminoethoxy)-2-(5-nitro-1-methyl-2-imidazolylmethylene)-1-andonone sulphate

Brotherton, J., 1978, Arzneimittelforsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

6-(2-Dimethylaminoethoxy)-2-(5-nitro-1-methyl-2-imidazolylmethylene)-1-tetralone sulphate

Brotherton, J., 1978, Arzneimittelforsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

2-[3-(Dimethylamino) propyl-amino]-4-(trichloromethyl)-6-(α,α,α -trichloro-m-tolyl)-2-triazine -- WR-99,662.

WR-99,662

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

1-(3-Dimethylaminopropyl)-4-(p-methoxy-phenyl) piperazine dihydrochloride. See Piperamide.

4-Dimethylamino-m-tolyl methylcarbamate -- Metacil.

Metacil + Iramin
Orekhova, M. M., 1974, Vet. Nauka--Proizvod., Trudy, Minsk, v. 12, 139-140
E[imeria] tenella, chicks, coccidiostatic activity of iramin in combination with metacil

4-Dimethylamino-3,5-xyllyl methylcarbamate -- Zectran.

Zectran
Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Dimethyl benzyl alkyl ammonium chloride
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

N,N-Dimethyldecylamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

o,o-Dimethyl-o-2,5-dichloro-4-bromophenyl-thiophosphate. See Bromophos.

2,6-Dimethyl-3,5-dichloro-4-hydroxy-pyridine. See Meticlorpindol.

2,6-Dimethyl-3,5-dichloro-pyridinol-4. See Meticlorpindol.

O,O-Dimethyl 2,2-dichlorovinyl phosphate. See Dichlorvos.

2,8-Dimethyl-5-β-diethylamino-ethylamino-thiochromone -- Schistocide T-109.

Schistocide T-109
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

O,O-Dimethyl 0-[p-(dimethylsulfamoyl) phenyl] phosphorothioate. See Famphur.

O-Dimethyl-0-(4-dimethyl-sulfamoyl) phenylthiophosphate. See Famphur.

O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate. See Malathion.

N,N-Dimethyldodecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

N,N-Dimethylheptadecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

N,N-Dimethylhexadecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

1,5-Dimethyl-1-6-hydro-pyrido-4,3b-carbazole. See Olivacine.

2,6-Dimethyl-4 hydroxypyridine derivatives
Sevcik, B.; et al., 1974, Veterinaria, Praha, v. 16 (5-6), 421-588
Eimeria tenella, chickens, 613 substances screened as coccidiostats, extensive detailed statistical results

O,O-Dimethyl-(1-hydroxy-2,2,2-trichlorethyl)-phosphate. See Trichlorfon.

Dimethyl-[[2-(2-methoxyacetamido)-4-(phenylthio)anilino]methyl-idin]-dicarbamate. See Febantel.

Dimethyl [[2-[(methoxyacetyl)amino]-4-(phenylthio)phenyl]carbonimidoyl]bis]carbamate. See Febantel.

O,O-Dimethyl S-[2-(methylamino)-2-oxoethyl] phosphorodithioate. See Dimethoate.

O,O-Dimethyl S-(N-methylcarbamoylmethyl) phosphorodithioate. See Dimethoate.

O,O-Dimethyl-0-(3-methyl-4-nitrophenyl) phosphorothioate. See Fenitrothion.

O,O-Dimethyl-0(3-methyl-4-nitrophenyl) thiophosphate. See Fenitrothion.

O,O-Dimethyl 0-[4-(methylthio)-m-tolyl] phosphorothioate. See Fenthion.

2,3-Dimethyl-1-(4-nitrobenzyl) imidazolium iodide
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

1,2-Dimethyl-5-nitroimidazole. See Dimetrida-

- 1,3-Dimethyl-5-nitroimidazolium iodide
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- N,N-Dimethyloctadecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared
- (E)-3,7-Dimethyl-2,6-octadienyl p-(methylthio) phenyl ether
Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity
- 1-[(3,7-Dimethyl-2,6-octadienyl)oxy]-4-methoxybenzene
Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity
- N,N-Dimethylpentadecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared
- N,N-Dimethyl-N-2-phenoxyethyl-N-2'-thenylammonium)-p-chlorobenzensulfonate. See Thienium.
- N'-[(2,4-Dimethylphenyl)-N-[(2,4-dimethylphenyl)imino]methyl]-N-methylmethanimidamide. See Amitraz.
- N-[[[(2,4-Dimethylphenyl)imino]methyl]-N-methylbenzenesulfenamide (Upjohn U-42,564)
Dawkins, C. C.; and Gladney, W. J., 1978, J. Econom. Entom., v. 71 (4), 657-660
3-host ticks, guinea pigs, amitraz, Upjohn U-42,564, chlordimeform, detachment response and mortality
- 2-(2',4'-Dimethyl-phenylimino)-3-methyl-4-thiazolin. See Tifatol.
- 3,6-Dimethyl-6-phenyltetrahydro-2H-1,3-oxazine quaternary salts
Grier, N., 1979, J. Pharm. Sc., v. 68 (4), 407-411
Eimeria tenella, trichostrongyle, synthesis, in vitro and in vivo evaluation of quaternary salts of 4-phenyl-1,2,3,6-tetrahydropyridine and 3,6-dimethyl-6-phenyltetrahydro-2H-1,3-oxazine, none effective
- 0,0-Dimethyl phosphorodithioate S-ester with N-(mercaptomethyl) phthalimide. See Phosmet.
- 5-(0,0-Dimethylphosphoryl)-6-chlorobicyclo (3,2,0)-hepta-1,5-dien. See Heptenophos.
- 0,0-Dimethyl-S(phthalimidomethyl)dithiophosphate. See Phosmet.
- 0,0-Dimethyl phthalimidomethyl phosphorothiothionate. See Phosmet.
- 4-[p-(4,6-Dimethyl-2-pyrimidylsulfamyl)phenylazo]-1-naphthylamine
Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124
preparation and testing of 6 long-acting schistosomicidal resinsates
- N,N-(Dimethylquinolylium-methyl sulphate-6)-urea. See 1,3-Di-6-quinolyurea.
- J,0-Dimethyl-0-p-sulfamoylphenyl phosphorothioate
-- Cythioate.
- Cythioate
Doval, C. P.; and Gupta, I., 1978, Indian Vet. J., v. 55 (11), 890-892
mange and lice in sheep, mange and ticks in dogs, efficacy of cythioate, safe, convenient and acceptable
-
- Dimethylsulphoxide
Chrucieli, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances
- N,N-Dimethyltetradecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared
- 0,0-Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphate. See Trichlorfon.
- Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate. See Trichlorfon.
- 0,0-Dimethyl-2,2,2-trichloro-1-hydroxyethyl-phosphorous. See Trichlorfon.
- 0,0-Dimethyl-0,1,2,4,5-trichlorophenyl (2-chlorovinyl) phosphate. See Vinylphosphate.
- 0,0-Dimethyl-0-(2,4,5-trichlorophenyl) phosphorothioate. See Ronnel.
- N,N-Dimethyltridecanamine
Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

N,N-Dimethylundecanamine

Fisher, W. F.; et al., 1979, *Southwest. Entom.*, v. 4 (3), 249-253
 Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

Dimetridazole -- 1,2-Dimethyl-5-nitroimidazole;
 Dimetridazole methanesulphonate; Emtryl.

Dimetridazole

Chute, M. B.; Chute, A. M.; and Wilkins, G. C., 1978, *Parasitology*, v. 77 (1), 41-48
 Histomonas meleagridis, therapeutic treatment of chickens with dimetridazole may reduce but not eliminate transmission of this protozoan by eggs of Heterakis gallinarum from medicated birds

Dimetridazole (Emtryl)

Flatt, R. E.; Halvorsen, J. A.; and Kemp, R. L., 1978, *Lab. Animal Sc.*, v. 28 (1), 62-65
 Hexamita muris in laboratory mice (small intestine, between epithelial cells, lamina propria), case history, dimetridazole controlled clinical disease but did not eliminate the infection

Dimetridazole (Emtryl)

Kunstyr, I.; and Ammerpohl, E., 1978, *Lab. Animals*, v. 12 (2), 95-97
 Spironucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Dimetridazole

Lindmark, D. G.; and Mueller, M., 1976, *Antimicrob. Agents and Chemotherapy*, v. 10 (3), 476-482
 metronidazole and 11 other nitroimidazoles, antitrichomonad activity against Tritrichomonas foetus and Trichomonas vaginalis, mutagenic action in Salmonella test, reducibility of nitro group by T. foetus homogenates, results underscore role of reduction of nitro group in antitrichomonad and mutagenic activity of nitroimidazoles

Dimetridazole (Emtryl)

Panigrahy, B.; et al., 1978, *Avian Dis.*, v. 22 (4), 815-818
 Giardia sp., parakeets (intestine), case reports, successful treatment with dimetridazole: commercial aviaries in Texas

Dimetridazole

Plisek, K.; Billova, V.; and Malhocka, A., 1977, *Veterinaria, Praha*, v. 19 (6), 49-71
 dimetridazole, mice, chicks, acute and chronic toxicity, administration in therapeutic concentrations is safe in above animal species

Dimetridazole

Rettig, T., 1978, *J. Zoo Animal Med.*, v. 9 (3), 98-100
 Trichomonas gallinae in Haliaeetus leucocephalus (center of palate), case report, successful treatment with dimetridazole: vicinity of Tusten Station, Sullivan County, New York

Dimetridazole -- Continued.

Dimetridazole methanesulphonate (Emtryl)

Stoessel, F. R.; and Haberkorn, S. E. M., 1977, *Gac. Vet.*, Buenos Aires (323), v. 39, 457-461
 Trichomonas foetus, bulls, dimetridazole methanesulphonate injected intramuscularly or subcutaneously, all animals negative to T. foetus in post-treatment analysis

Dimetridazole methanesulphonate (Emtryl)

Stoessel, F. R.; and Haberkorn, S. E. M., 1977, *Gac. Vet.*, Buenos Aires (324), v. 39, 506-510
 Trichomonas foetus, bulls, dimetridazole methanesulphonate by intraruminal injection, most efficient treatment with daily dose of 100 mg per kg during 5 consecutive days

Dimetridazole

Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230
 Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Dimetridazole methanesulphonate. See Dimetridazole.

Dimidine

Dubovyi, S. Z.; et al., 1977, *Veterinariia, Moskva* (3), 71-72
 babesiasis, piroplasmosis, cattle, dimidine as effective chemoprophylaxis under pasture conditions with presence of vector, Boophilus calcaratus, comparison with azidine

Dimidine

Rakhimov, T. Kh.; et al., 1977, *Veterinariia, Moskva* (10), 75-77
 piroplasmosis, franciellosis, cattle, dimidine and imidocarb tested in various doses, recommended for control

Dimidium Br

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
 Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Diminazene. See Berenil.Diminazene aceturate. See Berenil.Diminazene salt. See Berenil.Diminazene suraminat. See Berenil.Diminazine aceturate. See Berenil.Dimpylat. See Diazinon.

Dinitolmide -- Coccidin; 3,5-Dinitro-o-toluamide;
DOT soluble; Zoalen; Zoalene; Zoamix.

Coccidin

Gobzem, V. R.; and Nazarov, V. G., 1978, Veterinaria, Moskva (3), 67-69
Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Zoalene (I)

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidostats in feeds, qualitative identification test

Zoalene (II)

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidostats in feeds, qualitative identification test

Zoalene

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Zoalene

Krylov, V. F., 1978, Veterinaria, Moskva (10), 68-69
Eimeria tenella strain resistant to pharmacocid after 35 laboratory passages in chickens, cross-resistance only to rigecocin

Zoalene

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Zoalene (Zoamix)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Dinitolmide

Long, P. L.; Millard, B. J.; and Smith, K. M., 1979, Avian Path., v. 8 (4), 453-467
Eimeria spp., chickens, effect of 4 anticoccidial drugs on development of immunity, field and laboratory conditions

Zoalene (Zoamix)

McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Zoalen

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Dinitolmide -- Continued.

Zoalene

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Zoalene

Parre, J.; and Olkonen, E., 1977, Eesti Pol-lumaj. Akad. Teadusl. Toode Kogum. (104), 100-108
Eimeria tenella, E. acervulina, E. brunetti, polyvalvaccine tested, chicks maintained in battery cages or deep litter; simultaneous zoalene treatment efficacious in deep litter maintenance

Zoalene

Sevcik, B.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21), (4), 375-383
coccidiosis, broilers, field trials of anticoccidials: CSSR

Zoalene + Ethopabate

Sevcik, B.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21), (4), 375-383
coccidiosis, broilers, field trials of anticoccidials: CSSR

Zoalene

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Zoalene

Winchester, R. V., 1978, N. Zealand J. Sc., Wellington, v. 21 (4), 553-555
coccidiostats akloamide and zoalene, residues in poultry meat, determination method

DOT soluble

Zaprianov, M. Ia., 1978, Vet. Med. Nauki, v. 15 (4), 103-107
Eimeria kofoidi in Alectoris graeca cypriatis (nat. and exper.), amprolium, DOT-soluble, and Esb, tested

3,5-Dinitrobenzamide. See Nitromide.

3,3'-Dinitro-5,5'-dichlor-2,2'-dioxydiphenylsulfide

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitarn. Bolezni, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

3,3'-Dinitro-5,5'-dichlor-2,2'-dioxydiphenylsulfone

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183

Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

3,3'-Dinitro-5,5'-dichlor-2,2'-dioxydiphenylsulfide

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183

Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

2,4-Dinitrophenol

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

2,4-Dinitrophenol

Matsuzawa, T., 1978, Parasitology, v. 77 (2), 235-241

Eimeria tenella, chickens, beclotiamine, mode of action studies; attempts to potentiate or antagonize its activity revealed that pyri-thiamine and 2,4-dinitrophenol also showed slight anticoccidial activity and that a combination of 2,4-DNP and beclotiamine was effective but weight gain was not as good as with beclotiamine alone

2,4-Dinitrophenol

Mueller, M.; et al., 1979, Comp. Biochem. and Physiol., v. 64B (1), 97-100

Tritrichomonas foetus, *Trichomonas vaginalis*, *Entamoeba invadens*, effects of 2,4-dinitrophenol (including effect on accumulation of metronidazole)

3,5-Dinitro-o-toluamide. See Dinitolmide.

Dinocap -- 2-(1-Methyl heptyl)-4,6-dinitrophenyl crotonate.

2-(1-Methyl heptyl)-4,6-dinitrophenyl crotonate Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Dinsed -- Polystat (with Dibutyltin dilaurate and Roxarsone and Sulfantran).

Polystat

McDougald, L. R.; and McQuiston, T. E., 1978, Avian Dis., v. 22 (4), 765-770

Eimeria spp., turkeys, coccidiosis management, innate (age) resistance and acquired immunity vs. anticoccidial medication

Diethyl. See Pyrimithate.

2,3-p-Dioxanedithiol S,S-bis (0,0-diethyl phosphorodithioate). See Dioxathion.

S,S-1,4-Dioxane-2,3-diyl bis-(0,0-diethyl phosphorodithioate). See Dioxathion.

Dioxathion -- Altik (with Toxaphene); Bercotox; Delnav; Delnav DFF; S,S-1,4-Dioxane-2,3-diyl bis-(0,0-diethyl phosphorodithioate; 2,3-p-Dioxanedithiol S,S-bis (0,0-diethyl phosphorodithioate); Navadel; Quimofos.

Delnav (Bercotox)

Atef, M.; and El-Say, A., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 73-84

Hyalomma dromedarii, *Rhipicephalus s. sanguineus*, laboratory tests (immersion technique) with delnav, supona, toxaphene, BHC

Dioxathion

Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439

economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Dioxathion

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Dioxathion

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886

laboratory-reared *Haematobia irritans*, susceptibility to topically applied insecticides

Dioxathion

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586

Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Quimofos (Delnav)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105

Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Dioxathion -- Continued.

Delnav (Quimofos)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (4), 309-310, 313-314, 317-318, 321-322

Boophilus microplus, strain B, inheritance of phosphorus resistance to delnav is genetically conditioned

Dioxathion (Delnav DFF)

Hammant, C. A., 1977, Rhodesian Vet. J., v. 8 (4), 67-70

ticks, resistance survey of field strains to commonly used ixodocides, changeover from arsenic to dioxathion dipping of cattle, dramatic improvement in tick control, necessitates complete overhaul of dipping facilities and retraining of personnel: Tribal Trust Lands of Rhodesia

Dioxathion (Delnav DFF)

Hammant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73

Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Dioxathion + Toxaphene (= Altik)

Keating, M. I., 1978, Bull. Animal Health and Prod. Africa, v. 26 (4), 285-292

excretion of toxaphene and dioxathion in milk of dairy cows

Dioxathion (Delnav)

Lourens, J. H. M.; and Lyaruu, D. M., 1979, PANS, v. 25 (2), 135-142

Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Bercotox

Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96

ectoparasites, veterinary practice, bercotox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Dioxathion

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144

Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Dioxathion

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960

Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Dioxathion

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427

Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Dioxathion -- Continued.

Dioxathion (Delnav)

Rechav, Y.; Whitehead, G. B.; and Terry, S. B., 1978, J. South African Vet. Ass., v. 49 (2), 99-101

ticks, mortality curves of larvae dipped in dioxathion, chlorphenvinphos, and oxionthiophos, time of application, larvae of ticks exhibit diel periodicity in sensitivity to acaricides

Dioxathion

Wright, F. C.; and Riner, J. C., 1979, Southwest. Entom., v. 4 (1), 40-45

Psoroptes ovis, *P. cuniculi*, 10 acaricides evaluated using 'tea-bag' technique

2,2'-Dioxydiphenylsulfide

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183

Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

2,2'-Dioxydiphenylsulfone

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183

Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

2,2'-Dioxydiphenylsulfoxide

Kozhokaru, A. F.; and Topaly, V. P., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 178-183

Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation

2,2'-Dioxy-3,3',5,5'-tetra chlorodiphenyl sulfide. See Bithionol.

Diphetarzone -- Bemarsal; Difetarzone.

Difetarzone

Rubidge, C. J.; O'Dowd, P. B.; and Powell, S. J., 1973, South African Med. J., v. 47 (23), 991-992

Trichuris trichiura, children, difetarzone, clinical trials: King Edward VIII Hospital, Durban

Diphetarsone -- Continued.

Bemarsal

Trzaska, B.; and Leinweber-Noiszewska, H., 1974, Ann. Acad. Med. Stetinensis, suppl. 10, 19-25

Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis

Diphezy1 -- Difezil.

Difezil

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Dipofen. See Chlormethiuron.

Dipterex. See Trichlorfon.

2,2'-Dipyridyl

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

1,3-Di-6-quinolylurea -- Acaprin; Babesan; N,N-(Dimethylquinolylum-methyl sulphate-6)-urea; Quinorium sulphate; Quinuronium sulphate.

Quinorium sulphate [?Quinuronium sulphate]

Hashemi-Fesharki, R.; and Amjadi, A. R., 1977, Arch. Inst. Razi (29), 83-86

Babesia bovis, outbreak in imported Holstein and Red-Danish cattle, imidocarb dihydrochloride more effective than quinorium sulphate: Rasht, northern Iran

Quinuronium SO₄

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Quinuronium sulphate

Malhotra, D. V.; Gautam, O. P.; and Banerjee, D. P., 1979, Indian J. Animal Sc., v. 49 (1), 75-77

Babesia equi, donkeys (exper), diminazene aceturate as effective as diminazene aceturate + rolitetracycline, quinuronium sulphate ineffective

Babesan

Pal, M.; and Verma, J. D., 1975, U. P. Vet. J., v. 3 (4), 222-225

Babesia bigemina infection in Haryana cow following recovery from parturient paresis (milk fever), case report, complete recovery after treatment with babesan and nutrient supplements: India

1,3-Di-6-quinolylurea -- Continued.

Babesan

Sharma, M. M.; et al., 1979, Indian Vet. Med. J., v. 3 (3), 187-191

theileriosis, Jersey cattle, incidence and treatment with berenil, babesan, nevaquine, and aureomycin: Exotic Nucleus Cattle Farm, Bassi, Jaipur

Acaprin

Vulchovski, Ia., 1977, Vet. Med. Nauki, v. 14 (3), 79-87

Babesia, cattle, efficacy of trypan blue, trypaflavin, acaprin, berenil, and pyrodis (berenil and pyrodis superior to other drugs)

Dirian. See Brotianide.

Dirocide. See Diethylcarbamazine.

Disalan

Timofeev, B. A.; et al., 1978, Veterinariia, Moskva (4), 67-68

Fasciola hepatica, cattle, disalan effective, no toxic effect; comparison of anthelmintic effect with dertil

Disodium ethane-1-hydroxy-1,1-diphosphonate

Newsome, A. L.; and Harley, J. P., 1978, Experientia, v. 34 (12), 1567-1568

Trichinella spiralis, rats, acceleration of cyst calcification by administration of vitamin D₃, inhibition of cyst calcification by administration of disodium ethane-1-hydroxy-1,1-diphosphonate, demonstrates that cyst calcification is not an irreversible process and is subject to drug therapy

Disodiumthiobis(4-chloro-6-nitrophenolate)

Sakamoto, T.; and Gemmill, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Disodiumthiobis(3,4,6-trichlorophenolate)

Sakamoto, T.; and Gemmill, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Disophenol -- Ancylo1; 2,6-Diiodo-4-nitrophenol; DNP.

Disophenol (2,6,-Diiodo-4-nitrophenol; DNP)

Aronson, C. E.; and Serlick, E. R., 1977, Biochem. Pharmacol., v. 26 (23), 2297-2305

disophenol, effects on isolated perfused rat heart

Disophenol

Douch, P. G. C.; and Buchanan, L. L., 1979, Xenobiotica, v. 9 (8), 467-473

Ascaris suum, Moniezia expansa, disophenol, nitroxynil, nitrodan, metabolism by intact helminths, by helminth enzyme preparations, and by mouse- and sheep-liver enzymes

Disophenol -- Continued.

Disophenol (Ancylo1)

Retnasabapathy, A.; and Baskaran, G., 1976, v. 53 (10), 806-811
 ancylostomiasis, dogs, morantel tartrate, pyrantel pamoate and disophenol effective

Disophenol (Ancylo1)

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 226-229
 Ancylostoma caninum, dogs, comparative efficacy of disophenol, fenbendazole, pyrantel pamoate, and thiabendazole, clinical trials

Ancylo1

Singh, H.; Singh, R. P.; and Bali, M. K., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 55-58
 Ancylostoma sp., thiabendazole, fenbendazole, ancylo1, and pyrantel pamoate, in vitro effects on development of eggs and infective larvae

Disophenol

Tripathy, S. B.; and Chakrabarti, A., 1978, Indian J. Animal Health, v. 17 (2), 163-166
 Fasciola gigantica, cattle, disophenol: India

Distamycin A

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Distodin. See Hexachlorophene.

Distolon. See Niclofolan.

Disulfiram -- Antabuse; Tetraethylthiuram disulfide.

Disulfiram

Bennett, J. L.; and Gianutsos, G., 1978, Biochem. Pharmacol., v. 27 (5), 817-820
 Schistosoma mansoni in mice, disulfiram reduces norepinephrine levels in both male and female parasites and induces abnormal egg production, varying effects on other schistosome physiological and biochemical parameters, disulfiram also reduces pathological consequences of infection and affects parasite development

Disulfiram

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
 Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Disulfiram -- Continued.

Tetraethylthiuram disulfide (Antabuse; Disulfiram)

Scheibel, L. W.; Adler, A.; and Trager, W., 1979, Proc. National Acad. Sc., v. 76 (10), 5303-5307
 Plasmodium falciparum, antimalarial effects of tetraethylthiuram disulfide and its reduction product diethyldithiocarbamate

Disulfiram

Seed, J. L.; Pratt, M. C.; and Bennett, J. L., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 508-514
 Schistosoma mansoni, mice, chronic administration of disulfiram in diet blocks formation of egg shell in female schistosomes, reduces host mortality, and decreases granuloma formation; these results however are rapidly reversible

Disulfiram (Tetraethylthiuram disulfide)

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Dithiazanine -- Dithiazanine iodide; Dizan.

Dithiazanine iodide

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, Rev. Brasil. Med., v. 31 (11), 791-794
 human soil transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Dithiazanine iodide

Gudimenko, I. I., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 100-104
 trichuriasis, oesophagostomiasis, ascariasis, swine, testing thiabendazole, dithiazanine iodide, dipterex, hygromycin-B, and bubulin

Dithiazanine iodide

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
 Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

Dithiazanine iodide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
 Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Dithiazanine -- Continued.

Dithiazanine

Shedivtsova, A., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Dithiazanine iodide (Dizan)

Stokhof, A. A.; and Wolvekamp, W. T. C., 1978, Tijdschr. Diergeneesk., v. 103 (2), 1121-1129

Dirofilaria immitis, dogs, 4 case reports, chemotherapy: Netherlands (imported from United States, South America, or South Africa)

Dithiazanine iodide

Wong, M. M.; and Suter, P. F., 1979, Am. J. Vet. Research, v. 40 (3), 414-420

Dirofilaria immitis, dogs (exper.) without microfilaremia, indirect fluorescent antibody titers, degree of eosinophilia, and radiologic findings before and after treatment, reinfection, necropsy findings, significance of tests, application to diagnosis

Dithiazanine iodide. See Dithiazanine.

2,2'-Dithiobis(4-chloro-6-nitrophenol)

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

2,2'-Dithiobis(4-chlorophenol)

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

2,2'-Dithiobis(4,6-dichlorophenol)

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Dithiosemicarbazone -- alpha Ethoxyethylglyoxal dithiosemicarbazone; Gloxazone.

Gloxazone

Thompson, K. C.; et al., 1978, Trop. Animal Health and Prod., v. 10 (2), 75-81

Anaplasma marginale, *Babesia argentina*, *B. bigemina*, cattle under tropical conditions, immunization with virulent organisms followed by drug therapy (ganaseg; gloxazone; emicina) vs. chemoprophylaxis (imidocarb); tick and gastrointestinal parasite control without haemoparasitic control had advantage over no control system at all

Ditrazine. See Diethylcarbamazine.

Diuredosan -- Diethyl [thio [O-[3-(p-tolylsulphonyl) ureido] phenyl] carbamoyl] phosphoramidate; Sansalid; Uredofos; Uredofos, Disodium salt.

Diuredosan

Gemmell, M. A.; Johnstone, P. D.; and Oudemans, G., 1978, Research Vet. Sc., v. 25 (1), 111-112

Echinococcus granulosus, *Taenia hydatigena*, dogs, diuredosan, significant activity against *T. hydatigena*, no significant dose response curve against *E. granulosus*

Uredofos

Tetzlaff, R. D.; and Weir, W. D., 1978, Lab. Animal Sc., v. 28 (3), 287-289

concurrent *Hymenolepis nana* and *Syphacia obvelata* infections in mice, uredofos and disodium salt of uredofos, determination of effective dose levels

Uredofos, Disodium salt

Tetzlaff, R. D.; and Weir, W. D., 1978, Lab. Animal Sc., v. 28 (3), 287-289

concurrent *Hymenolepis nana* and *Syphacia obvelata* infections in mice, uredofos and disodium salt of uredofos, determination of effective dose levels

Uredofos (Sansalid)

Todd, K. S., jr.; Howland, T. P.; and Woerpel, R. W., 1978, Am. J. Vet. Research, v. 39 (2), 315-316

Mesocestoides corti, dogs (exper.), bunamidine hydrochloride and uredofos (good results), arecoline hydrobromide and niclosamide (variable results)

Divermin

Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91

cestodes, pathomorphology resulting from action of various anthelmintics

Divezid

Shchelkanov, K. G.; and Epel'dimov, L. S., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 76-80

Dictyocaulus, sheep, divezid, nilverm, and cyazone, changes in lung tissue of sheep and some tissues of parasite resulting from drug action

Divezid

Shevchenko, R. V., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 73-75

divezid, sheep cardiovascular system, modification of heart action followed by restoration of normal function of heart muscle

Dixanthogen

Knudsen, E. A.; and Pedersen, T. G., 1979, Ugeskr. Laeger, v. 141 (43), 2965-2966

scabies, humans, therapeutic trials of hexicide vs. dixanthogen: Denmark

Dixol. See Trichlorfon.

Dizan. See Dithiazanine.

Dizinc bis (dimethyldithiocarbamate) ethylene bis (dithiocarbamate)

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

DNP. See Disophenol.

DOT soluble. See Dinitolmide.

Dovenix. See Nitroxynil.

Doxorubicin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Doxycycline -- 6-Desoxy-5-hydroxytetracycline; Doxycycline hydrochloride; Terradoxyn.

Doxycycline

Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397

Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances

Doxycycline hydrochloride (Terradoxyn)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, Chemotherapy, v. 25 (4), 222-226

Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Doxycycline -- Continued.

Doxycycline + Chloroquine

Khan, A. A.; and Maguire, M. J., 1978, Brit. Med. J. (6128), v. 1, 1669-1670

P[lasmodium] falciparum, chloroquine-resistant strain reported in young child, therapy with doxycycline + chloroquine resulted in cure: Zambia

Doxycycline

Kuttler, K. L.; and Simpson, J. E., 1978, Am. J. Vet. Research, v. 39 (2), 347-349

Anaplasma marginale, splenectomized calves (exper.), 2 formulations of oxytetracycline (T-200 and T-50), doxycycline, drug efficacies, influence of treatment on parasitemia and packed cell volume

Doxycycline hydrochloride. See Doxycycline.

Drofenit. See Tetramisole.

Droncit. See Praziquantel.

DTHP. See Trichlorfon.

Duamycin. See Nigericin.

Duodegran. See Ronidazole.

Dursban. See Chlorpyrifos.

Dursban 44 Insecticide Formulation. See Chlorpyrifos.

Echinon. See Nitroscanate.

Econazole (nitrate)
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Ectiban. See Permethrin.

Ectoral. See Ronnel.

Ectrin. See Fenvalerate.

Efloran. See Metronidazole.

Eglumine
Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

EHNA. See Erythro-9-(2-hydroxy-3-nonyl)adenine.

Elancoban. See Monensin.

Elancoban Premix. See Monensin.

Elimix. See Pyrimithate.

Ellipticine
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Embay 8440. See Praziquantel.

Embay 8440-Bayer. See Praziquantel.

Embazin. See Sulfaquinoxaline.

Embelia ribes extract -- Sonex (with Nicotine and Pomegranate).

Sonex
Matta, S. C.; and Ahluwalia, S. S., 1979, *Indian Vet. J.*, v. 56 (7), 616-617
helminths, poultry, helminta-P, sonex

Embelia ribes alcoholic extract
Pandey, B. B.; and Rai, P., 1976, *U. P. Vet. J.*, v. 4 (2), 74-77
Taenia hydatigena and Multiceps multiceps in puppies (exper.), Embelia ribes alcoholic extract not as effective as dichlorophen

Emericid. See Lonomecin.

Emetine -- Emetine aminoxide; Emetine hydrochloride.

Emetine
Akhound-Zadeh, H., 1976, *Rev. Internat. Serv. Sante Armees*, v. 49 (5), 421-426
chronic cutaneous leishmaniasis, soldier with severe ulcers that did not heal despite 8 years of therapy with various anti-leishmanial drugs, chronicity thought to be result of immuno-deficiency, ulcers finally cured after additional therapy with monomycine: Iran (had travelled to Khouzistan)

Emetine aminoxide
Anorve Carmona, I.; and Rodriguez Sanchez, A., 1974, *Rev. Med. Hosp. Gen.*, Mexico, v. 37 (5), 331-335
human amoebiasis, clinical trials testing efficacy of emetine aminoxide, drug useful for both amoebiasis and trichocephaliasis

Emetine + Chloroquine
Bindschadler, D. D., 1974, *Rocky Mountain Med. J.*, v. 71 (7), 387-389
E[ntamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Emetine
Botero R., D., 1978, *Ann. Rev. Pharmacol. and Toxicol.*, v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Emetine hydrochloride
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Emetine hydrochloride
Campos, R., 1973, *Rev. Brasil. Clin. e Terap.*, v. 2 (11), 587-588
Entamoeba histolytica, comparison of efficacy of nifuratel and other amoebicides using material cultured from intestinal ulcers of patient with intestinal symptomatic amoebiasis

Emetine
Cerecedo Cortina, V.; and Gaxiola Gaxiola, R., 1972, *Rev. Fac. Med., Univ. Nac. Auton. Mexico*, an. 15, v. 15 (1), 25-28
human hepatic amoebic abscess, side effects of currently used amoebicides (emetine, chloroquine, metronidazole, aminosidine)

Emetine hydrochloride
Dutta, G. P.; and Narain, L., 1978, *Indian J. Exper. Biol.*, v. 16 (7), 838-840
Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Emetine -- Continued.

Emetine

Entner, N., 1979, J. Protozool., v. 26 (2), 324-328

Entamoeba histolytica, emetine binding to ribosomes, inhibition of protein synthesis and amebicidal action, capacity to bind emetine is index of drug resistance

Emetine hydrochloride

Koutsaimanis, K. G.; Timms, P. W.; and Ree, G. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (4), 768-769

amoebiasis, Nigerian ship's steward with multiple hepatic abscesses, failure to respond to recommended doses of metronidazole, successful treatment with emetine hydrochloride

Emetine hydrochloride

Lumbreras, H.; and Uyema, N., 1972, Rev. Peruana Med. Trop., v. 1 (2), 95-98

Balantidium coli, in vitro, comparative action of dehydroemetine and emetine hydrochloride

Emetine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Emetine

Olaeta Elizalde, R., 1973, Rev. Cir. Hosp. Juarez, Mexico (187-188), v. 44, 1972-1973, 59-64

human hepatic amoebic abscess, therapeutic recommendations (emetine, dehydroemetine, chloroquine, metronidazole)

Emetine

Pena Yanez, A.; et al., 1972, Rev. Clin. Espan., v. 124 (4), 405-412

pulmonary amoebiasis in man without involvement of liver, diagnosis after expectoration of chocolate colored material, successful therapy of emetine and chloroquine: Spain (had resided previously in Algeria)

Emetine HCl

Sinden, R. E.; and Smalley, M. E., 1979, Parasitology, v. 79 (2), 277-296

Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

Emetine hydrochloride

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriologie, 1 Abt. Orig., Reihe A, v. 241 (3), 358-367

Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Emetine

Skromne-Kadlubik, G.; Medina, J.; and Rauda, L., 1976, SPM Salud Pub. Mexico, v. 18 (2), 395-398

Entamoeba histolytica, human, comparative survey, conventional medications vs. radiation therapy

Emetine -- Continued.

Emetine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Emetine aminoxide. See Emetine.

Emetine hydrochloride. See Emetine.

Emicina. See Oxytetracycline.

Emimycin

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175

Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Emimycin

Ryley, J. F.; and Hardman, L., 1978, Parasitology, v. 76 (1), 11-20

Eimeria spp., chicks (exper.), effects of dietary vitamin K on severity of disease with particular attention to effects of vitamin K on response to anticoccidial drugs, concluded that use of vitamin K deficient diet for experimental work is quite justified

Emtryl. See Dimetridazole.

Endrin

Khan, D.; and Haseeb, M. A., 1976, Pakistan J. Zool., v. 8 (2), 173-176

Ganeo micracetabulus and *Cercaria reflexicauda* cercariae, effects of 5 insecticides at various concentrations, toxicity varies but results indicate cercariae are susceptible to insecticides

Endrin

Pandey, B. B., 1978, Indian Vet. J., v. 55 (3), 253

endrin, toxicity in bullocks treated for tick infestation: Purmoti, Azamgarh (U.P.)

Enteroseptol. See Iodochlorhydroxyquin.

Enterovioform. See Iodochlorhydroxyquin.

Entex. See Fenthion.

Entobex. See Phanquone.

Eosin BA

Daniyarov, I. A.; et al., 1978, Veterinaria, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

(E)-10,11-Epoxy-N,N-diethyl-3,7,11-trimethyl-2-dodecenamide

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

(E)-6,7-Epoxy-3,7-dimethyl-1-[p-(methylthio)phenyl]-2-octene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

4-[(6,7-Epoxy-3,7-dimethyl-2-nonenyl)oxy]-1,2-(methylenedioxy)benzene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

(+)4-[6,7-Epoxy-3,7-dimethyloctyl)oxy]-1,2-(methylenedioxy)benzene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

6,7-Epoxy-3,7-dimethyl-1-(2-propynyloxy)-2-octene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

6,7-Epoxy-7-ethyl-3-methyl-1-(p-nitrophenoxy)-2-nonene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

4-[(6,7-Epoxy-3-ethyl-7-methyl-2-nonenyl)oxy]-1,2-(methylenedioxy)benzene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

(E)-6,7-Epoxy-1-(p-ethylphenoxy)-3,7-dimethyl-2-octene

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

(E)-6,7-Epoxy-1-(p-ethylphenoxy)-3,7-dimethyl-2-octene (R-20458)

Wright, J. E.; et al., 1974, J. Med. Entom., v. 11 (4), 385-389

Cochliomyia hominivorax, methoprene and R-20458 inhibited emergence in vitro when applied topically or incorporated in larval diet, yearling heifers sprayed with each analogue and infested with larvae displayed little change in serum chemistry and adults emerged from the wound-reared larvae

Equiben. See Cambendazole.

Equigard. See Dichlorvos.

Equivurm Plus. See Mebendazole.

Equizole. See Thiabendazole.

Equizole A. See Piperazine or Thiabendazole.

DL-Erythro-2,8-bis(trifluoro-methyl)- α -(2-piperidyl)-4-quinolinemethanol hydrochloride. See Mefloquine.

Erythrocin. See Erythromycin.

Erythro-9-(2-hydroxy-3-nonyl)adenine -- EHNA.

Erythro-9-(2-hydroxy-3-nonyl)adenine + Cordycepin
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

EHNA + Cordycepin + Guanosine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Erythromycin -- Erythrocin; Erythromycin stearate.

Erythromycin stearate (Erythrocin)
Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, *Chemotherapy*, v. 25 (4), 222-226
Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Erythromycin

Seilhamer, J. J.; and Byers, T. J., 1978, *J. Protozool.*, v. 25 (4), 486-489
Acanthamoeba castellanii, mutants resistant to erythromycin, chloramphenicol, and oligomycin

Erythromycin

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Erythromycin stearate. See Erythromycin.

DL-Erythro- α -(2-piperidyl)-2,8-bis-(trifluoromethyl)-4-quinolinemethanol. See Mefloquine.

Esb₃. See Sulfachloropyrazine.

Eserine. See Physostigmine.

Estrella. See Coralox.

Ethanol. See Alcohol, Anhydrous.

Ethidium. See Homidium.

Ethidium bromide. See Homidium.

Ethion -- Coopertox [of Grillo Torrado, M. M.; and Perez Arrieta, A., 1977].

Ethion

Drummond, R. O.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Ethion -- Continued.

Coopertox (Ethion)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, *Rev. Med. Vet.*, Buenos Aires, v. 58 (2), 101-102, 105
Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Ethion

Nolan, J.; Roulston, W. J.; and Schnitzerling, H. J., 1979, *Austral. Vet. J.*, v. 55 (10), 463-466
Boophilus microplus, range of resistant strains on naturally and experimentally infected cattle, field and stall spraying trials, efficacy of synthetic pyrethroids for tick control, potentiation of pyrethroids by organo-phosphorus compounds

Ethion

Waters, K. S., 1978, *Queensland Agric. J.*, v. 104 (3), 215-225
control of buffalo flies on cattle using back rubber charged with ethion, good results, directions for making and maintaining

D-Ethionine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Ethopabate -- Amprol plus (with Amprolium); Amprolmix plus (with Amprolium); Pancoxin (with Amprolium and Sulfaquinoxaline); Pancoxin plus (with Amprolium and Pyrimethamine and Sulfaquinoxaline); Supacox (with Amprolium and Pyrimethamine and Sulfaquinoxaline).

Amprol plus

Bedrnik, P., 1977, *Veterinarstvi*, v. 27 (10), 458-459
coccidiosis, chickens, current prevalence, amprol plus treatment

Amprol plus

Bedrnik, P.; et al., 1979, *Veterinarstvi*, v. 29 (8), 353-355
coccidiosis, broiler chicken fattening, lasalocid compared with amprol plus and monensin: Czechoslovakia

Ethopabate plus Amprolium (= Amprol plus)

Buys, S. B.; and Rasmussen, R. W., 1978, *J. South African Vet. Ass.*, v. 49 (2), 127-128
significantly higher mortalities found in nicarbazine fed chickens than in amprolium fed chickens when exposed to same heat stress conditions

Ethopabate -- Continued.

Ethopabate + Amprolium (=Amprolmix plus)
Hamet-Foure, N.; Macar, C.; and Robin, B., 1979, Avian Path., v. 8 (1), 107-113
Eimeria meleagridis, E. adenoeides, turkeys, activity of clopidol with methylbenzoate and amprolium with ethopabate: France

Ethopabate

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidostats in feeds, qualitative identification test

Ethopabate + Amprolium

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Ethopabate + Amprolium + Sulphaquinoxaline (= Pancoxin)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Ethopabate + Amprolium + Sulphaquinoxaline + Pyrimethamine (= Supacox)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Ethopabate + Amprolium

McDougald, L. R.; Karlsson, T.; and Reid, W. M., 1979, Avian Dis., v. 23 (4), 999-1005
coccidiosis, chickens (exper.), natural outbreak of infectious bursal disease (IBD) during comparison of anticoccidials for their effect on development of immunity, interaction between diseases, immunity to coccidiosis not blocked by IBD

Amprol plus

Manuel, M. F., 1972, Philippine J. Vet. Med., v. 11 (2), 92-100
Eimeria tenella, broilers (exper.), evaluation of amprol plus and coyden-25 with or without 3-nitro-50 (roxarsone)

Amprol plus

Manuel, M. F.; and Buduan, R. J., 1972, Philippine J. Vet. Med., v. 11 (2), 73-91
battery-raised broilers, effects of amprol plus or cycostat with or without payzone on weight gain and feed efficiency, no coccidiosis found

Pancoxin plus

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Ethopabate -- Continued.

Ethopabate + Amprolium

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Pancoxin

Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazin, and pancoxin: England; France; Germany

Amprol plus

Sevcik, B.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21) (4), 375-383
coccidiosis, broilers, field trials of anticoccidials: CSSR

Ethopabate + Zoalene

Sevcik, B.; et al., 1979, Biol. a Chem. Zivoc. Vyroby, Vet., v. 15 (v. 21), (4), 375-383
coccidiosis, broilers, field trials of anticoccidials: CSSR

Pancoxin

Sherkov, Sh., 1977, Vet. Sbirka, v. 75 (3), 35-38
E[imeria] tenella, pancoxin, chickens raised under conditions of high and low temperatures; influence of thiamine on development of coccidiosis

Pancoxin plus

Sherkov, S. N.; Leitch, B.; and Kokash, L., [1977], Egypt. J. Vet. Sc., v. 13 (1), 1976, 37-43
Sarcocystis tenella, kittens (exper.), development in intestines, life cycle; attempted parasite suppression using statyl and pancoxin plus

Pancoxin plus

Stoianov, P.; et al., 1978, Vet. Med. Nauki, v. 15 (8), 105-114
E[imeria] tenella, broiler chickens, varied temperature and moisture regimes, blood biochemistry, host resistance, efficacy of pancoxin plus

Ethopabate + Amprolium (= Amprol Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Ethopabate + Sulphaquinoxaline + Pyrimethamine + Amprolium (= Pancoxin Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

- F 159
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
- Facizine. See Tinidazole.
- Falmonox. See Teclozan.
- Famfur. See Famphur.
- Famophos. See Famphur.
- Famphur** -- Cyanamid-38023; 0,0-Diemthyl 0-[p-(dimethylsulfamoyl) phenyl] phosphorothioate; 0-Diemthyl-0-(4-diemthyl-sulfamoyl) phenylthiophosphate; Famfur; Famophos; Warbex.
- Famphur
Drummond, R. O.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides
- Warbex
Evstaf'ev, M. N., 1978, *Veterinariia*, Moskva (11), 70-72
Hypoderma bovis, cattle, insecticides tested, aerosol method of application more useful for large, specialized farms than for individual treatment: Tiimensk oblast
- Famphur bolus
Hair, J. A.; et al., *J. Econom. Entom.*, v. 72 (1), 135-138
Boophilus spp., Dermacentor albipictus, cattle (exper.), sustained release famphur bolus
- Famphur (Warbex)
Loomis, E. C.; and Schock, R. C., 1978, *J. Med. Entom.*, v. 14 (6), 649-651
Hypoderma lineatum, H. bovis, cattle, famphur, pour-on application compared with intramuscular injection, field tests: California
- Warbex (Famophos; Famfur; Cyanamid-38023)
Nepoklonov, A. A.; and Zabolotnyi, K. F., 1978, *Veterinariia*, Moskva (3), 99-100
warbex, method for analyzing meat and milk for residues
- Famphur bolus
Teel, P. D.; Hair, J. A.; and Stratton, L. G., 1979, *J. Econom. Entom.*, v. 72 (2), 230-233
Amblyomma maculatum, A. americanum, cannulated Hereford heifers (exper.), evaluation of sustained-release famphur bolus administered orally
- Fasciolin [of Furmaga, S.; Gundlach, J. L.; and Sobieszewski, K., 1974]. See Carbon tetrachloride.
- Fasigyn. See Tinidazole.
- Fasigyn 500. See Tinidazole.
- Fasigyne. See Tinidazole.
- Febantel** -- Bay h 5757; Bay Vh 5757; N-[2-[2,3-Bis (methoxycarbonyl) guanidino]-5-(phenylthio)-phenyl] 2-methoxy-acet-amid; Dimethyl-[[2-(2-methoxyacetamido)-4-(phenylthio) anilino]methyl-idin]-dicarbamate; Dimethyl [[2-[(methoxyacetyl)amino]-4-(phenylthio) phenyl]carbonimidoyl]bis]carbamate; Rintal; Rintal paste.
- Febantel (Rintal)
Behrens, H., 1978, *Vet.-Med. Nachr.* (2), 169-173
helminths, sheep, rintal, field trial
- Febantel (Rintal)
Burger, H. J., 1978, *Vet. Rec.*, v. 103 (26-27), 572-574
5 gastrointestinal nematodes, sheep (exper.), febantel, efficacy against fourth-stage larvae, pre-adult fifth, or adult stages
- Febantel (Rintal)
Connan, R. M., 1978, *Vet.-Med. Nachr.* (2), 145-149
nematodes, pigs (exper.), febantel
- Febantel (Rintal paste)
Drudge, J. H.; et al., 1979, *Am. J. Vet. Research*, v. 40 (6), 758-761
Parascaris equorum and other horse parasites, oxibendazole, critical tests and clinical trials; febantel paste
- Febantel (Rintal paste)
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1978, *Am. J. Vet. Research*, v. 39 (9), 1419-1421
nematodes, bots, horses, febantel, activity of paste formulation alone or with trichlorfon paste, critical tests
- Febantel (Rintal)
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1979, *J. Equine Med. and Surg.*, v. 3 (3), 135-140
parasites, horses, critical tests with febantel alone or in combination with trichlorfon
- Rintal (Bay Vh 5757; Febantel)
Enigk, K.; and Dey-Hazra, A., 1978, *Deutsche Tierarztl. Wchnschr.*, v. 85 (9), 350-354
gastrointestinal nematodes, horses, rintal, drug efficacy, egg reduction tests, critical test
- Febantel (Rintal)
Enigk, K.; and Dey-Hazra, A., 1978, *Vet.-Med. Nachr.* (2), 134-144
nematodes, pigs (nat. and exper.), rintal, controlled test
- Fanasil. See Sulfadoxine.
- Fansidar. See Pyrimethamine or Sulfadoxine.

Febantel -- Continued.

- Febantel (Rintal)
Enigk, K.; and Dey-Hazra, A., 1978, Vet.-Med. Nachr. (2), 195-203
nematodes, wild mammals and birds, rintal: zoos and wild animal parks, vicinity of Calcutta
- Febantel (Rintal)
Grelck, H.; Hoerchner, F.; and Woehrl, H., 1978, Vet.-Med. Nachr. (2), 154-159
nematodes, calves (exper.), febantel, efficacy against larval and adult stages, controlled trial
- Febantel (Rintal; Bay Vh 5757)
Hasslinger, M. A.; and Mueller, R., 1978, Vet.-Med. Nachr. (2), 186-191
strongyles, horses, rintal, granular and paste formulations, field trials
- Febantel (Rintal)
Hopkins, T. J.; and Rafferty, M., 1978, Vet.-Med. Nachr. (2), 160-168
nematodes, sheep, febantel, drug trials, cross-resistance of thiabendazole-resistant strains of *Haemonchus contortus* and *Trichostrongylus colubriformis*: Australia
- Febantel (Rintal)
Morrow, G. L., 1978, Vet. Med. and Small Animal Clin., v. 73 (11), 1388, 1393
strongyles, ascarids, bots, horses, febantel and trichlorfon paste formulations, drug trials
- Rintal (Bay Vh 5757)
Reuss, U., 1978, Vet.-Med. Nachr. (2), 174-179
gastrointestinal parasites, sheep, rintal, tolerance, effectiveness, application methods
- Febantel (Rintal; Bay Vh 5757)
Shmidl, J. A.; et al., 1978, Vet. Med. and Small Animal Clin., v. 73 (6), 775-776, 781
febantel paste and suspension, horses, toxicology evaluation
- Febantel (Rintal)
Shmidl, J. A.; et al., 1979, Vet. Med. and Small Animal Clin., v. 74 (12), 1796-1799
safety evaluation of concurrent administration of febantel and trichlorfon in paste and liquid forms to horses, no significant toxic effects
- Febantel (Rintal)
Stiefelhagen, W.; and Uhlemann, F. F., 1978, Vet.-Med. Nachr. (2), 192-194
nematodes, horses, rintal, effectiveness, tolerance
- Febantel (Rintal)
Terblanche, H. J. J., 1978, Vet.-Med. Nachr. (2), 180-185
febantel, rams, effect on fertility, drug trials
- Febantel (Rintal; Bay h 5757)
Thomas, H., 1978, Research Vet. Sc., v. 25 (3), 290-293
gastrointestinal nematodes, sheep (exper.), febantel effective against adult and larval stages using various testing methods

Febantel -- Continued.

- Rintal
Uhlemann, F. F., 1978, Vet.-Med. Nachr. (2), 150-153
nematodes, pigs, rintal, effective against all developmental stages, field trial: Bonn
-
- Fenasal. See Niclosamide.
- Fenbautel (5757)
Georgieva, D., 1978, Vet. Sbirka, v. 76 (4), 32-33
anthelmintic treatment, lambs, weight gains
- Fenbendazole -- 2-Carbamomethoxy-5-phenylthio-benzimidazole; Fenbendazole suspension; HOE-881; Hoe 881V; Methyl-5-(phenylthio)-2-benzimidazole carbamate; Methyl[5-(phenylthio)-1H-benzimidazole-2-yl] carbamate; Panacur; 5 (Phenylthio)-2-benzimidazole-carbamic acid methyl ester; 5-Phenylthio-benzimidazole-2-methylcarbaminate; [5-(Phenylthio)-1H-benzimidazole-2-yl] carbamic acid methyl ester; Rumevite Wormablok containing Panacur.
- Fenbendazole (Panacur)
Anderson, N.; and Lord, V., 1979, Austral. Vet. J., v. 55 (4), 158-162
Ostertagia ostertagi, *Trichostrongylus axei*, cattle, fenbendazole, oxfendazole, and 1e-vamisole compared
- Fenbendazole (Panacur)
Angus, K. W.; Coop, R. L.; and Sykes, A. R., 1979, Research Vet. Sc., v. 26 (1), 120-122
Ostertagia circumcincta, *Trichostrongylus colubriformis*, sheep (exper.), time required for restoration of normal mucosal architecture of small intestine following fenbendazole treatment
- Fenbendazole suspension
Bali, M. K.; and Singh, R. P., 1977, Haryana Agric. Univ. J. Research, v. 7 (3), 155-157
nematodiasis, goats, efficacy of fenbendazole
- Fenbendazole
Bali, M. K.; and Singh, R. P., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 230-232
Haemonchus contortus, in vitro larvicidal effects of fenbendazole, banminth II, and nemafox
- Fenbendazole
Bali, M. K.; Singh, R. P.; and Kaushik, R. K., 1977, Indian J. Animal Research, v. 11 (2), 81-83
Haemonchus contortus, *Oesophagostomum* spp., *Strongylus papillosus*, *Trichuris* sp., sheep, fenbendazole, drug efficacy, good results against all nematodes except *Trichuris* sp.
- Fenbendazole suspension (Panacur)
Barger, I. A., 1979, Austral. Vet. J., v. 55 (2), 68-70
nematodes, grazing dairy cattle, single anthelmintic treatment (fenbendazole) had no significant influence on milk production: Australia

Fenbendazole -- Continued.

Fenbendazole (Panacur)

Batte, E. G., 1978, Vet. Med. and Small Animal Clin., v. 73 (9), 1183-1186

Ascaris suum, *Trichuris suis*, *Stephanurus dentatus*, pigs, fenbendazole, drug trials, good results, no adverse effects

Panacur

Bauer, B.; et al., 1978, Berl. u. Munchen.

Tierarztl. Wchnschr., v. 91 (21), 413-418

endoparasite control with panacur in Criollo x Zebu x Charolais calves under extensive management conditions in the tropics

Fenbendazole (Panacur; Hoe 881V)

Benz, G. W.; and Ernst, J. V., 1978, Am. J.

Vet. Research, v. 39 (7), 1103-1105

gastrointestinal nematodes, calves (exper.), fenbendazole highly effective given orally as a suspension or in medicated feed

Fenbendazole (Panacur)

Boeckeler, W.; and Segebade, R., 1977,

Tierarztl. Umschau, v. 32 (9), 473-478

nematodes, fallow-deer, panacur, good results, recommended that medicated food be given once at beginning of frost-period and at end of winter

Fenbendazole (Panacur)

Burke, T. M.; and Roberson, E. L., 1978, Am.

J. Vet. Research, v. 39 (11), 1799-1801

helminths, dogs, fenbendazole, critical trials, no evidence of drug toxicosis

Fenbendazole (Panacur)

Burke, T. M.; and Roberson, E. L., 1979, Am.

J. Vet. Research, v. 40 (4), 552-554

Toxocara canis, *Ancylostoma caninum*, neonatal pups (exper.), fenbendazole, reduction in worm burdens and marked improvement in clinical condition

Fenbendazole

Callinan, A. P. L.; and Cummins, L. J., 1979,

Austral. Vet. J., v. 55 (8), 370-373

nematodes, cattle (nat. and exper.), efficacies of various anthelmintics against adult and larval stages: western Victoria

Fenbendazole

Coles, G. C.; and Briscoe, M. G., 1978, Vet.

Rec., v. 103 (16), 360-361 [Letter]

Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for *Haemonchus contortus* eggs

Fenbendazole (Panacur)

Corba, J.; et al., 1977, Vet. Med., Praha,

v. 50, v. 22 (4), 201-206

gastro-intestinal nematodes, cattle, fenbendazole 100% effective without undesirable symptoms

Fenbendazole (Panacur)

Corba, J.; et al., 1978, Veterinarstvi, v. 28

(6), 274-275

Dicrocoelium dendriticum, sheep, trials of cambendazole, fenbendazole, diamfenetid

Fenbendazole -- Continued.

Fenbendazole (Panacur)

Corba, J.; et al., 1979, Brit. Vet. J., v. 135

(4), 318-323

helminths of sheep and cattle, efficacy of fenbendazole

Fenbendazole (Panacur)

Craig, T. M.; and Bell, R. R., 1978, Am. J.

Vet. Research, v. 39 (6), 1037-1038

gastrointestinal nematodes, calves, natural infection, fenbendazole compared with levamisole, controlled experiment, varied results

Fenbendazole

Cummins, L. J.; and Callinan, A. P. L.,

1979, Austral. Vet. J., v. 55 (7), 348-349

[Letter]

nematodes, cattle, albendazole and fenbendazole, effect of oesophageal groove reflex on anthelmintic efficiency

Fenbendazole

Douch, P. G. C.; and Buchanan, L. L., 1979,

Xenobiotica, v. 9 (11), 675-679

Moniezia expansa, *Ascaris suum*, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Fenbendazole (Panacur)

Drudge, J. H.; Lyons, E. T.; and Tolliver,

S. C., 1978, J. Equine Med. and Surg., v. 2

(1), 22-26

nematodes, cestodes, horses, controlled tests and clinical trials with suspension and granule formulations of fenbendazole highly effective; dose rate of 10 mg/kg inadequate for effective control of *Strongyloides westeri*; no activity on *Gasterophilus* spp., *Habronema* spp., *Draschia*, *Anoplocephala* spp.

Fenbendazole (Panacur)

Drudge, J. H.; Lyons, E. T.; and Tolliver,

S. C., 1978, Vet. Med. and Small Animal Clin.,

v. 73 (5), 623-625

Dictyocaulus viviparus and gastrointestinal nematodes in calves, controlled-test evaluation of fenbendazole against natural infections, mixed results

Fenbendazole (Panacur)

Drudge, J. H.; Lyons, E. T.; and Tolliver, S.

C., 1979, Am. J. Vet. Research, v. 40 (4), 590-

594

large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Fenbendazole (Panacur)

Dubey, J. P.; Miller, T. B.; and Sharma, S. P.,

1979, J. Am. Vet. Med. Ass., v. 174 (8), 835-

837

Paragonimus kellicotti, dogs (exper.), fenbendazole effective

Fenbendazole (Panacur)

Duwel, D., 1978, Kleintier-Praxis, v. 23 (5),

237-242

helminths, dogs, fenbendazole in granule and powder form, anthelmintic efficacy, no undesirable side effects

Fenbendazole -- Continued.

Fenbendazole

Duewel, D.; and Schleich, H., 1978, Zentralbl. Vet.-Med., Reihe B, v. 25 (10), 800-805
Hymenolepis diminuta, laboratory rats, fenbendazole, mode of action

Fenbendazole (Panacur)

Duewel, D.; and Strasser, H., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (6), 239-241
Ancylostoma caninum, Toxocara canis, pregnant bitches, effect of long-term fenbendazole treatment during and after pregnancy, results show whelps were helminth-free after birth

Fenbendazole (Panacur)

Duewel, D.; and Tiefenbach, B., 1978, Tierarztl. Umschau, v. 33 (5), 252, 254-255
Moniezia expansa, M. benedeni, sheep, mixed infections with nematodes, fenbendazole, various preparation forms proved to be very effective

Fenbendazole

Duncan, J. L.; Armour, J.; and Bairden, K., 1978, Vet. Rec., v. 103 (10), 211-212
gastrointestinal nematodes, calves, high efficiency of fenbendazole against inhibited fourth stage larvae and adults, not affected by seasonal timing of administration (i. e. either in autumn or in winter)

Fenbendazole (Panacur)

Eckert, J.; Barandun, G.; and Pohlenz, J., 1978, Schweiz. Med. Wchnschr., v. 108 (29), 1104-1112
Echinococcus spp., larval stages, laboratory animals, fenbendazole, mebendazole

Fenbendazole (Panacur)

Enigk, K., 1977, Tierarztl. Umschau, v. 32 (8), 414-420
nematodes, pigs, fenbendazole, efficacy at very low doses, drug trials, results demonstrate that effective dose be spread over six consecutive days

Fenbendazole (Panacur)

Enigk, K.; Dey-Hazra, A.; and Batke, J., 1975, Acta Parasitol. Polon., v. 23 (26-40), 367-372
helminths of geese, treatment (Amidostomum anseris with mebendazol, fenbendazol, levamisol, and pyrantel tartrat; Trichostrongylus tenuis with mebendazol and fenbendazol; Drepanidotaenia lanceolata and Hymenolepis setigera with mebendazol)

Fenbendazole

Friedman, P. A.; and Platzer, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Panacur

Furmaga, S.; Gundlach, J. L.; and Filar, J., 1977, Med. Wet., v. 33 (3), 137-141
gastrointestinal helminths, sheep, panacur

Fenbendazole -- Continued.

Fenbendazole

Gautam, O. P.; Bansal, S. R.; and Dey-Hazra, A., 1976, Indian Vet. J., v. 53 (12), 965-966
Neoascaris vitulorum, buffalo-calves, field trials with fenbendazole, 100% effective at 7.5 mg/kg body weight

Fenbendazole (Panacur)

Gunawan, M.; et al., 1979, Research Vet. Sc., v. 27 (1), 111-115
Haemonchus contortus, Trichostrongylus colubriformis, efficacies of fenbendazole and albendazole against developing and adult stages of benzimidazole-resistant strains, sheep (exper.)

Fenbendazole

Guralp, N.; and Tinar, R., 1978, Vet. Fak. Dergisi, Ankara Univ., v. 25 (3), 440-447
nematodes of dogs or cats, fenbendazole

Fenbendazole

Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367
Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Fenbendazole

Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, Research Vet. Sc., v. 25 (3), 360-363
Haemonchus contortus, Trichostrongylus colubriformis, levels of benzimidazole resistance recorded from an egg hatch test procedure

Fenbendazole (Panacur)

Handlos, M.; and Handlos, B., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (7), 290, 292-293
strongyles, horses, fenbendazole, effective against adults, partially effective against immature stages: Sejenane, North Tunisia

Fenbendazole

Hinz, E., 1978, Zentralbl. Bakteriologie, 1. Abt. Orig., Reihe A, v. 240 (4), 542-548
Echinococcus multilocularis, mice, fenbendazole, significant reduction in worm burden and number of protoscolices, degree of efficiency subject to way of application, duration of medication and parasite strain

Fenbendazole

Hinz, E., 1978, Zentralbl. Bakteriologie, 1. Abt. Orig., Reihe A, v. 241 (3), 388-394
Echinococcus multilocularis, HH vs. S strain, mice treated with fenbendazole as emulsion or in feed, serum protein values, compared with untreated and with uninfected mice

Fenbendazole

Hinz, E., 1978, Zentralbl. Bakteriologie, 1. Abt. Orig., Reihe A, v. 242 (2), 268-272
Echinococcus multilocularis, HH vs. S strain, mice treated with fenbendazole as emulsion or in feed, indirect fluorescent antibody titers, compared with untreated mice

Fenbendazole -- Continued.

- Fenbendazole (Panacur)
Inderbitzin, F.; and Eckert, J., 1978, Berl. u. Munchen. Tierarztl. Wchnschr., v. 91 (20), 395-399
Ostertagia ostertagi, Dictyocaulus viviparus, calves (exper.), efficacy of fenbendazole against inhibited and adult stages
- Fenbendazole
Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of Nematospiroides dubius in mice
- Fenbendazole
Kalita, C. C.; Gautam, O. P.; and Banerjee, D. P., 1978, Indian Vet. J., v. 55 (8), 660-662
Haemonchus spp., sheep, fenbendazole, good results, ineffective against Trichuris spp. and Moniezia spp.: India
- Fenbendazole
Kaushik, R. K.; Banerjee, D. P.; and Bali, M. K., 1977, Haryana Vet., v. 16 (2), 61-64
Toxocara canis, dogs (nat. and exper.), fenbendazole very effective, in vitro trials on eggs revealed no ovicidal or larvicidal effect
- Fenbendazole
Kerboeuf, D.; Hubert, J.; and Le Stang, J. P., 1979, Rec. Med. Vet., v. 155 (2), 143-148
Ostertagia ostertagi, cattle, serum pepsinogen levels in relation to worm burden and anthelmintic treatments
- Fenbendazole (Panacur)
Kirsch, R., 1977, Deutsche Tierarztl. Wchnschr., v. 84 (2), 52-54
strongylids and ascarids in horses (nat. and exper.), fenbendazole, excellent results against adult stages, partial effect on immature stages
- Fenbendazole (Panacur)
Kirsch, R., 1978, Research Vet. Sc., v. 25 (3), 263-265
Ostertagia ostertagi, Haemonchus contortus, Trichostrongylus colubriformis, fenbendazole, in vivo and in vitro studies on ovicidal activity
- Fenbendazole (Panacur)
Kirsch, R.; Petri, K.; and Degenhardt, H., 1978, Kleintier-Praxis, v. 23 (6), 291-298
nematodes, pigeons, fenbendazole
- Fenbendazole (Panacur)
Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae
- Fenbendazole
Le Stang, J.-P.; Hubert, J.; and Kerboeuf, D., 1978, Rev. Med. Vet., Toulouse, v. 129 (10), 1355-1362, 1365-1369
gastrointestinal nematodes, cattle, efficacy of fenbendazole and pyrantel tartrate

Fenbendazole -- Continued.

- Panacur
Love, J., 1979, Vet. Rec., v. 104 (1), 22 [Letter]
warble fly, ostertagiasis, young stock, simultaneous prophylactic treatment with rycovet warblecide and panacur
- Fenbendazole
Luethgen, W., 1979, Tierarztl. Umschau, v. 34 (2), 104, 107-112
Ascaridia columbae and Capillaria columbae in Columba livia dom., fenbendazole, laboratory and field trials of effectiveness, reversible disorders in feather development were only adverse side effects
- Fenbendazole (Panacur)
Lweno, M. F.; and Semuguruka, W. D., 1978, Bull. Animal Health and Prod. Africa, v. 26 (1), 11-15
Haemonchus contortus, Oesophagostomum columbianum, Trichostrongylus colubriformis, sheep, goats, fenbendazole, highly effective, no side effects: Tanzania
- Fenbendazole (Panacur)
McBeath, D. G.; et al., 1978, Equine Vet. J., v. 10 (1), 5-8
strongyle parasites, horses, fenbendazole effective against both adult and larval stages permitting anthelmintic treatment at less frequent intervals
- Fenbendazole
McBeath, D. G.; Dean, S. P.; and Preston, N. K., 1979, Vet. Rec., v. 105 (22), 507-509
Ostertagia ostertagi, winter calving dairy cows, fenbendazole administered during the dry period resulted in increases in subsequent lactation yields: farms in north-west England
- Fenbendazole (Rumevite Wormablok containing Panacur)
McBeath, D. G.; Preston, N. K.; and Thompson, F., 1979, Brit. Vet. J., v. 135 (3), 271-278
nematodes, ewes, fenbendazole administered in feed-block formulation prior to lambing reduced peri-parturient faecal egg count rise and subsequent pasture contamination and lamb infection, experimental and field studies
- Panacur
McEwan, A. D.; and Oakley, G. A., 1978, Vet. Rec., v. 102 (14), 314-315
panacur, thiabendazole, and nilzan with dye marker added, oral dosing of cattle showed evidence of rumen by-pass, reduced drug efficacy probably resulting from closure of oesophageal groove
- Fenbendazole (Panacur)
McEwan, A. D.; Oakley, G. A.; and Robinson, M., 1979, Vet. Rec., v. 105 (1), 15-16
Dictyocaulus viviparus, calves (exper.), lung lesions more severe with fenbendazole than with levamisole and certain types more severe in treated calves vs. controls, may be direct result of drug action

Fenbendazole -- Continued.

Fenbendazole (Panacur)

Marriner, S.; and Bogan, J. A., 1979, Vet. Rec., v. 105 (11), 261
benzimidazole anthelmintics, sheep, oral vs. intraruminal vs. intra-abomasal administration

Fenbendazole (Panacur)

Marti, O. G.; Stewart, T. B.; and Hale, O. M., [1979], J. Parasitol., v. 64 (6), 1978, 1028-1031
gastrointestinal nematodes, pigs raised under similar management conditions, comparative efficacy of fenbendazole, dichlorvos, and levamisole HCl

Fenbendazole (Panacur)

Melbourne, C. P., 1978, J. Zoo Animal Med., v. 9 (4), 146-148
Trichuris sp. and 5 strongyloid genera, giraffes, camels, fenbendazole and thiabendazole: Longleat Safari Park, Great Britain

Fenbendazole

Nicholas, W. L.; and Stewart, A. C., 1979, Ann. Trop. Med. and Parasitol., v. 73 (1), 57-62
Toxocara canis, laboratory mouse, fenbendazole and oxfendazole killed larvae in brains and musculature, migratory larvae more susceptible, possible use in preventing pre-natal infection in dogs

Fenbendazole (Panacur)

Pfeiffer, H., 1978, Wien. Tierarztl. Monatschr., v. 65 (11), 343-346
Dictyocaulus viviparus, calves (exper.), fenbendazole, efficacy of repeated administration of small doses against inhibited larvae

Fenbendazole

Prichard, R. K.; Hennessy, D. R.; and Steel, J. W., 1978, Vet. Parasitol., v. 4 (4), 309-315
Haemonchus contortus, Trichostrongylus colubriformis, sheep, Ostertagia ostertagi, cattle, 4 benzimidazoles, mode of action and pharmacokinetic behavior, implications for prolonged administration as a new concept for increasing spectrum and effectiveness of anthelmintics

Fenbendazole

Prichard, R. K.; Kelly, J. D.; and Thompson, H. G., 1978, Vet. Parasitol., v. 4 (3), 243-255
Haemonchus contortus, Trichostrongylus colubriformis, sheep, thiabendazole, fenbendazole, concentrations of anthelmintics or their radiolabelled metabolites in parasite tissues after administration to host, differences between amount of each anthelmintic incorporated by susceptible and resistant parasite strains and between the two parasites, effect of route of administration on anthelmintic concentration in parasite tissue and host plasma

Fenbendazole

Rodriguez Caabeiro, F.; Martinez Fernandez, A. R.; and Sanmartin Duran, M. L., 1978, Rev. Iver. Parasitol., v. 38 (3-4), 551-567
Trichinella spiralis, efficacy of fenbendazole, mice (exper.)

Fenbendazole -- Continued.

Fenbendazole (Panacur)

Samizadeh-Yazd, A.; and Todd, A. C., 1978, Am. J. Vet. Research, v. 39 (10), 1668-1671
Nematodirus helvetianus, calves (exper.), fenbendazole, good results against adult worms, reduced egg production and morphogenesis of ova to infective larval stages, no toxicity

Fenbendazole (HOE-881)

Sanchez-Carrillo, C.; and Beltran-Hernandez, F., 1977, SPM Salud Pub. Mexico, v. 19 (5), 691-700
intestinal parasites, humans, clinical trials testing fenbendazole, effective against Necator americanus, Trichuris trichiura, and Ascaris lumbricoides, not effective against Strongyloides stercoralis: Estado de Chiapas, Mexico

Fenbendazole (Panacur)

Sangster, N. C.; et al., 1979, Research Vet. Sc., v. 26 (1), 85-89
Trichostrongylus colubriformis, Haemonchus contortus, benzimidazole-resistant strains, sheep (exper.), efficacy of fenbendazole given as single or divided dose

Fenbendazole (Panacur)

Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1978, Rev. Centro Cien. Rurais, v. 8 (1), 35-38
Haemonchus contortus, thiabendazole-resistant strain, sheep, fenbendazole

Fenbendazole

Sarup, S.; Banerjee, D. P.; and Gautam, O. P., 1978, Indian Vet. J., v. 55 (6), 497-498
Ancylostoma caninum, ova and infective larvae, in vitro evaluation of fenbendazole, helatac, alcopar, banminth II, only banminth II effective; banminth II-treated larvae administered orally to mice, none recovered from lungs or liver

Fenbendazole

Schmidt, R. L.; et al., 1979, J. Wildlife Management, v. 43 (2), 461-467
Protostrongylus, domesticated, captive, or free-ranging Ovis c. canadensis, evaluation of 5 drugs

Fenbendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Vet., v. 16 (1), 5-7
Ancylostoma caninum, dogs, pyrantel pamoate, fenbendazole, thiabendazole, clinical trials, comparative efficacy

Fenbendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 226-229
Ancylostoma caninum, dogs, comparative efficacy of disophenol, fenbendazole, pyrantel pamoate, and thiabendazole, clinical trials

Fenbendazole -- Continued.

Fenbendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 55-58
 Ancylostoma sp., thiabendazole, fenbendazole, ancylosol, and pyrantel pamoate, in vitro effects on development of eggs and infective larvae

Fenbendazole (Panacur)

Stehle, S., 1977, Kleintier-Praxis, v. 22 (6), 261-266
 helminths in birds of prey, fenbendazole, good results

Fenbendazole (Panacur)

Strasser, H.; and Tiefenbach, B., 1977, Deutsche Tierarztl. Wchnschr., v. 84 (12), 479-480
 Syphacia muris, rat breeding colony, long-term treatment with fenbendazole, reinvasion of colony from outside after treatment was stopped

Fenbendazole (Panacur)

Thomas, R. J., 1978, Vet. Rec., v. 102 (18), 394-397
 gastrointestinal nematodes and lungworms of sheep, fenbendazole in-feed medication highly effective against inhibited larvae and most adults, implications for control of pasture contamination

Fenbendazole (Panacur)

Vindevogel, H.; Duchatel, J. P.; and Fievez, L., 1978, Ann. Med. Vet., v. 122 (2), 109-115
 capillariasis in pigeons (nat. and exper.), fenbendazole, good results

Fenbendazole (Panacur)

Williams, J. C.; et al., 1979, Am. J. Vet. Research, v. 40 (8), 1087-1090
 Ostertagia ostertagi, cattle, fenbendazole, efficacy against adults, developing stages, and inhibited early 4th-stage larvae; includes some results for other worm genera: Louisiana

Fenbendazole suspension. See Fenbendazole.

Fenchlorphos. See Ronnel.

Fenitrothion -- 0,0-Dimethyl-0-(3-methyl-4-nitrophenyl) phosphorothioate; 0,0-Dimethyl-0(3-methyl-4-nitrophenyl) thiophosphate; Sumithion; Sumithion 50 EC.

Sumithion 50 EC (Fenitrothion)

Chellappa, D. J.; Subramanian, R.; and Gopalakrishnan, C. A., 1977, Indian Poultry Gaz., v. 61 (4), 135-137
 Menopon gallinae, Lipeurus caponis, poultry, malathion 50 EC, sumithion 50 EC, nuvan 100 EC, drug trial, sumithion 50 EC most effective, knapsack sprayer superior to conventional dipping procedure, preliminary report

Fenitrothion -- Continued.

Sumithion

Khan, M. H.; and Srivastava, S. C., 1977, Indian J. Animal Health, v. 16 (2), 137-140
 Boophilus microplus engorged females, in vitro tests with dursban, gamma BHC, sumithion, supona, dimecron, egg production and viability; supona most effective

Sumithion

Kumar, A.; et al., 1977, Indian J. Poultry Sc., v. 12 (3), 48-50
 Eomenacanthus stramineus, chicks (exper.), sumithion, malathion, dipterex, comparative efficacy, haematology and blood glucose levels

0,0-Dimethyl-0(3-methyl-4-nitrophenyl) thiophosphate (Fenitrothion)

Privora, M.; Rupes, V.; and Cerny, V., 1970, Folia Parasitol., v. 17 (1), 81-84
 Dermacentor marginatus, laboratory trials testing six insecticides

Fenitrothion

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144
 Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Fenitrothion

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
 Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Fenitrothion

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
 Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Fenothrin. See Phenothrin.

Fenthion -- Baytex; Entex; Bay 9007; Bay 29493; Baytex-50; 0,0-Dimethyl 0-[4-(methylthio)-m-tolyl] phosphorothioate; Lebaycid; Mercaptophos; Queletex; Tiguvon; Tiguvon Spot On.

Fenthion (Tiguvon)

Boulard, C. F., 1979, Vet. Parasitol., v. 5 (4), 379-387
 Hypoderma-infected or uninfected calves, treatment with fenthion or trichlorfon, blood histamine levels, circulating antibody titers to Hypoderma lineatum antigen in infected calves; blood histamine levels in guinea pigs after injection of ground-up Hypoderma lineatum larvae or application of fenthion

Fenthion

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
 Boophilus annulatus, B. microplus, laboratory tests of insecticides

Fenthion -- Continued.

- Baytex**
Evstaf'ev, M. N., 1978, Veterinariia, Moskva (11), 70-72
Hypoderma bovis, cattle, insecticides tested, aerosol method of application more useful for large, specialized farms than for individual treatment: Tiumensk oblast
- Lebaycid**
Guttowa, A.; and Boniecka, B., 1977, Acta Parasitol. Polon., v. 24 (28-34), 315-322
Fasciola hepatica, Triaenophorus nodulosus, embryos, in vitro effects of pesticides Vapam and Lebaycid, implications for effects of environmental pollution on structure of ecosystems
- Fenthion (Tiguvon Spot On)**
Joubert, J. P. J.; and Minne, J. A., 1979, J. South African Vet. Ass., v. 50 (1), 47-48
fenthion, pregnant beef cows, no clinical symptoms of toxicity or abortions observed
- Fenthion**
Kiefer, M.; and Cyprich, D., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Zool. (23), 5-12
ticks, susceptibility to acaricides: Slovakia
- Fenthion (Tiguvon)**
Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae
- Baytex-50 (Fenthion)**
Letunov, V. N., 1977, Veterinariia, Moskva (10), 81
[Hypoderma], reindeer, Baytex-50, effective treatment
- Fenthion (Tiguvon)**
Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661
flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico
- Lebaycid**
Moczon, T., 1976, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (5), 289-292
Fasciola hepatica miracidia, inhibitory effect of pesticides on enzyme activity
- Tiguvon (Bay 29493, Bay 9007, Fenthion, Mercaptophos, Baytex, Entex, Queletex)**
Sayin, F.; and Meric, I., 1976, Vet. Fak. Dergisi, Ankara Univ., v. 23 (3-4), 301-307
Hypoderma, indigenous cattle, pour-on application of ruelene 6-R, tiguvon, good results: Central Anatolia, Turkey

Fenthion -- Continued.

- Fenthion**
Teel, P. D.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 297-300
Amblyomma maculatum feeding on fenthion-infused sheep, evidence of cholinergic neurotransmitter involvement in overall nervous control of ixodid tick salivary fluid secretion, principal cause of death in systemic organophosphate-poisoned ixodid ticks may be disruption of normal salivary gland function
- Fenthion**
Uspenskii, I. V., 1974, Parazitologiya, Leningrad, v. 8 (4), 312-321
Ixodes persulcatus, susceptibility to acaricides
-
- Fenthion ethyl** -- 0,0-Diethyl-0-4(methylmercapto-3-methylphenyl) thiophosphate.
- Fenthion-ethyl**
Tenquist, J. D.; and Roberts, M., 1978, N. Zealand Vet. J., v. 26 (4), 88-91
Lucilia sericata, Damalina ovis, sheep, 2 new 'Mini-shower' models of dipping with fenthion-ethyl and diazinon compared with plunge and shower dipping, residues in wool samples
-
- Fenvalerate** -- Cyano-(3-phenoxyphenyl)-methyl-4-chloro- α -(1-methylethyl)-benzeneacetate; Ectrin; SD 43775.
- Fenvalerate (SD 43775)**
Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765
Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control
- Fenvalerate (SD 43775)**
Ahrens, E. H.; and Cocke, J., 1979, J. Econom. Entom., v. 72 (2), 215
Haematobia irritans, cattle, fenvalerate-impregnated ear tags: Hidalgo Co., Texas
- SD-43775**
Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, J. Econom. Entom., v. 71 (2), 315-318
Ornithonyssus sylviarum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban permethrin and SD-43775 also effective; mites displayed tolerance to malathion
- Fenvalerate (Ectrin)**
Loomis, F. C.; Bramhall, F. L.; and Dunning, L. L., 1979, J. Econom. Entom., v. 72 (6), 856-859
Ornithonyssus sylviarum, White Leghorn hens, carbaryl and fenvalerate compared

Ferrous sulfate + Copper sulfate
Sverba, V. A.; and Shemchuk, V. R., 1978,
Veterinariia, Moskva (10), 69-71
Sinergasilus major, white amur, copper sul-
fate and ferrous sulfate mixture, chlorophos,
carbophos, formula for estimating concentra-
tions in relation to temperature and other
factors in aquaria or ponds

Filaramide. See Arsenamide.

Filaricides
Denham, D. A., 1979, *J. Helminth.*, v. 53 (2),
175-187
methods for testing compounds for filaricidal
activity, review

Filarzan. See Diethylcarbamazine.

Filixan. See Aspidium.

Flagyl. See Metronidazole.

Flagyl V. See Metronidazole.

Flagyl suspension. See Benzoyl metronidazole.

Floxacrine -- 7-Chloro-10-hydroxy-3-(4-trifluoro-
methyl-phenyl)-3,4-dihydroacridine-1,9(2H,
10H)-dione.

Floxacrine
Schmidt, L. H., 1979, *Antimicrob. Agents and
Chemotherapy*, v. 16 (4), 475-485
Plasmodium spp. in monkeys, floxacrine, lack-
ing in radical curative activity, significant
prophylactic activity but with requirement
for daily dosage, untoward host reaction

Floxuridine -- 5-Fluoro-2'-deoxyuridine; 5-FUDR;
WR 138 720.

5-FUDR (5-Fluoro-2'-deoxyuridine; WR 138 720)
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, *Antimicrob. Agents and Chemotherapy*, v.
15 (2), 157-160
Trypanosoma rhodesiense, mice, active in
screening of antitumor compounds for efficacy
against infection

Flubendazole -- Methyl [5-(4-fluorobenzoyl) 1H-
benzimidazole-2-yl] carbamate.

Flubendazole
Canese, A.; et al., 1978, *Rev. Paraguaya
Microbiol.*, v. 13 (1), 31-33
common human intestinal parasites, flubenda-
zole, clinical trials, good results

Flubendazole -- Continued.

Flubendazole
Cavier, R.; and Notteghem, M. J., 1979, *Ann.
Pharm. Franc.*, v. 37 (1-2), 33-36
Hymenolepis nana var. fraterna, mice, com-
parative study of mebendazole and flubenda-
zole

Flubendazole
Notteghem, M. J.; Leger, N.; and Cavier, R.,
1979, *Ann. Pharm. Franc.*, v. 37 (3-4), 153-
156
Echinostoma caproni, mice, flubendazole

Flubendazole
Raeymaekers, A. H. M.; et al., 1978, *Arzneimit-
tel-Forsch.*, v. 28 (4), 586-594
Syphacia muris, Strongyloides ratti, synthesis
and anthelmintic activity of mebendazole,
flubendazole and other alkyl-(5-acyl-1-H-
benzimidazol-2-yl) carbamates in rats

Flubendazole
Schenone, H.; et al., 1977, *Bol. Chileno Para-
sitol.*, v. 32 (3-4), 85-86
Ascaris lumbricoides, Trichuris trichiura,
Enterobius vermicularis, children, treatment
trials with flubendazole, drug well tolerated

Flubendazole
Thienpont, D.; et al., 1978, *Arzneimittel-
Forsch.*, v. 28 (4), 605-612
intestinal helminths, nat. and exper. infec-
tions in laboratory and domestic animals,
critical and controlled tests with flubenda-
zole to establish biological and pharmacologi-
cal properties

Flubendazole
Vanparijs, O.; Hermans, L.; and Thienpont, D.,
1979, *Vet. Parasitol.*, v. 5 (2-3), 237-242
Trichinella spiralis, rats, flubendazole,
anthelmintic activity against intestinal,
migrating, and encysted phase

Flukanide. See Rafoxanide.

Flukicides
Le Bars, H.; and Banting, A. de L., 1979, *Med.
& Chir. Digest.*, v. 8 (5), 435-441
Fasciola hepatica, exper. infection in rab-
bits, sheep, and cattle, variations in blood
parameters that reflect alterations in liver
function compared with normal values in
order to establish standards for studying
toxicity of flukicides

Flunidazole -- (Hydroxy-2'-ethyl)-1 (p-fluoro-
phenyl) 2-nitro-5-imidazole; 1-(2-Hydroxy-
ethyl)-2-(p-fluorophenyl)-5-nitroimidazole;
MK915.

Flunidazole
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Flunidazole -- Continued.

Flunidazole

- Cavier, R.; and Cenac, J., 1972, *Therapie*, v. 27 (4), 733-742
Trichomonas vaginalis, *Entamoeba*, in vitro and in vivo (rats, hamsters), efficacy of flunidazole compared with metronidazole
- 1-(2-Hydroxyethyl)-2-(p-fluorophenyl)-5-nitroimidazole (Flunidazole, MK915)
 Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
-
- 2-Fluoroadenosine
 Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- 2-Fluoro-2'-deoxyadenosine
 Senft, A. W.; and Crabtree, G. W., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs
- 5-Fluoro-2'-deoxyuridine. See Floxuridine.
- Fluorodopan
 Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 4-Fluoro-3-nitroaniline
 Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- DL-m-Fluoro-phenylalanine
 Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Fluorosalan -- 3,5-Dibromo-3'-trifluoromethylsalicylanilide.

- 3,5-Dibromo-3'-trifluoromethylsalicylanilide
 Sakamoto, T., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice
- 3,5-Dibromo-3'-trifluoromethylsalicylanilide
 Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
-
- 4-Fluoro-N-[2-(4-thiazolyl) benzimidazole-5-yl] benzamide
 Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- 5-Fluorouracil
 Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- 5-Fluorouracil
 Irvin, A. D.; and Young, E. R., 1978, *Research Vet. Sc.*, v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo
- 5-Fluorouracil
 Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 5-Fluorouracil
 Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Fluoxymesterone
 Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Folinic acid -- Calcium leukovorin.

Calcium leukovorin

McLeod, R.; et al., 1979, Am. J. Med., v. 67 (4), 711-714

Toxoplasma gondii, immunosuppressed man, brain abscesses, sulfadiazine, pyrimethamine, and calcium leukovorin, case report

Formaldehyde -- Formalin.

Formalin

Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, Current Sc., Bangalore, v. 47 (8), 283-284 [Letter]

Trichodina [sp.] on exterior of Cyprinus carpio, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

Formalin

El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, Argas persicus, evaluation of 10 insecticides

Formalin

Gilbert, J. P.; Gratzek, J. B.; and Brown, J., 1979, J. Fish Dis., v. 2 (3), 191-196
formalin and malachite green-oxalate (alone or in combination) used as model system for testing synergistic action of parasiticides in vitro

Formalin

Imada, R.; and Muroga, K., 1979, Bull. Japan. Soc. Scient. Fish. (Nippon Suisan Gakkaishi), v. 45 (1), 25-29

Pseudodactylogyrus microrchis on Anguilla anguilla (gills), trichlorfon, formalin, and sodium chloride baths compared, 2 trichlorfon baths effective and harmless

Formalin

Kunstyr, I.; and Ammerpohl, E., 1978, Lab. Animals, v. 12 (2), 95-97

Spironucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Formaldehyde

Sherkov, Sh.; et al., 1978, Vet. Sbirka, v. 76 (6), 39-41

coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat

Formaldehyde

Singh, Y., 1977, Indian Bee J., v. 36 (1-4), 1974, 16

Nosema apis, honey bees, formaldehyde (some improvement), thymol (good results): apiary at Jeolikote, India

Formaldehyde -- Continued.

Formalin

Slepnev, N. K.; and Zen'kov, A. V., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 135-137

E[chinococcus] granulosus, protoscolices, destructive action of high and low temperatures; lysol and creolin most destructive of chemicals tested

Formalin

Vlasenko, M. I.; and Meshcheriakova, A. A., 1977, Veterinariia, Moskva (4), 75-78

Chilodonella cyprini, Ichthyophthirius multifiliis, [Costia], fish, formalin solution for control, toxicity tested and safe levels established

Formaldehyde

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230

Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Formalin. See Formaldehyde.

Formycin -- 7-Amino-3(β-D-ribofuranosyl)pyrazolo-[4,3-d]pyrimidine.

Formycin

Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856

Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

4'-Formylbenzo-15-crown-5

Brown, G. R.; and Foubister, A. J., 1979, J. Med. Chem., v. 22 (8), 997-999

benzo-15-crown-5 polyethers, synthesis, in vivo and in vitro tests against Eimeria tenella

2-(1-Formyl-1-phenylhydroazino)-2-thiazoline

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Foschlor. See Trichlorfon.

Fouadin. See Stibophen.

Fowler's solution. See Potassium arsenite solution.

Framycetin. See Neomycin.

Franocid. See Diethylcarbamazine.

Ftalozol. See Phthalylsulfathiazole.

Ftorafur

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Fuadin. See Stibophen.

Fuchsin acid

Daniiarov, I. A.; et al., 1978, *Veterinariia, Moskva* (2), 64-65

Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

5-FUDR. See Floxuridine.

Fumagillin -- Fumidil B.

Fumagillin

Kogan, G. F., 1975, *Vet. Nauka--Proizvod., Trudy, Minsk*, v. 13, 196-198

nosematosis, bees, fumagillin had good therapeutic effect, metronidazole, sulfadimethoxin and enteroseptol showed no substantial effect

Fumagillin

Lehnert, T.; and Shimanuki, H., 1979, *Apidologie*, v. 10 (1), 17-22

Nosema apis, package bees, population change and spore levels in older vs. newly emerged bees, much of disease eliminated with death of older bees, effect of feeding fumagillin at time packages are installed

Fumidil B

Muresan, E.; et al., 1978, *Apicultura Romaniaa*, v. 53 (4), 13-15, 16

Nosema apis, Fumidil B and NOSAN against *Paramecium caudatum* as substitute test agent (having sensitivity similar to *N. apis*)

Fumidil B. See Fumagillin.

Fungi imperfecti

Krizkova, L.; Balanova, J.; and Balan, J., 1979, *Biologia, Bratislava, s. C, Biol.* (1), v. 34 (3), 241-245

antiprotozoal and antinematodal activity of Fungi imperfecti from soil samples collected in Mongolia

Fungizone. See Amphotericin B.

Furacillin. See Nitrofurazone.

Furacin. See Nitrofurazone.

Furadantin. See Nitrofurantoin.

Furaltadone -- Levofuraltadone; Levo-furaltadone hydrochloride; levo-5-Morpholinomethyl-3-(5-nitrofurylidene-amino)-2-oxasolidinone; NF-602; NF-902.

Levofuraltadone

Blandon, R.; Guevara, J. F.; and Johnson, C. M., 1976, *Rev. Med. Panama*, v. 1 (3), 153-162

Chagas disease in children, diagnosis, pathology, therapeutic trials with various drugs, metronidazole was well tolerated and therefore most promising therapy: Panama

Furaltadone

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Levo-furaltadone (NF-602)

Rassi, A.; and Ferreira, H. de O., 1971, *Rev. Soc. Brasil. Med. Trop.*, v. 5 (5), 235-262

Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best response

Levo-furaltadone hydrochloride (NF-902)

Rassi, A.; and Ferreira, H. de O., 1971, *Rev. Soc. Brasil. Med. Trop.*, v. 5 (5), 235-262

Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best response

Levofuraltadone (NF 602)

Ruppel, J. F.; and Burke, J., 1977, *Ann. Soc. Belge Med. Trop.*, v. 57 (4-5), 481-495

[*Trypanosoma*] gambiense, humans, clinical trials with levofuraltadone and levofuraltadone hydrochloride, comparisons with berenil, follow-up reports: Kimpangu, Republique du Zaire

Levofuraltadone hydrochloride (NF 902)

Ruppel, J. F.; and Burke, J., 1977, *Ann. Soc. Belge Med. Trop.*, v. 57 (4-5), 481-495

[*Trypanosoma*] gambiense, humans, clinical trials with levofuraltadone and levofuraltadone hydrochloride, comparisons with berenil, follow-up reports: Kimpangu, Republique du Zaire

Furamide. See Diloxanide furoate.

Furanace. See Nifurpirinol.

Furantoin. See Nitrofurantoin.

Furapromidium. See Nitrofurylacrylamide.

Furazolidone -- Bifuran (with Nitrofurazone); Furoxona; Furoxone; NF-180; 3-(5-Nitrofurfurylidene-amino)-2-oxazolidinone; 3-(5-Nitro-2-furfurylidenamino)-2-oxazolidone; N-(5-Nitro-2-furfurylidene)-3-amino-2-oxazolidone.

Bifuran

Aggarwal, C. K.; et al., 1978, Indian Vet. J., v. 55 (12), 952-957
bifuran, embazin, effect of coccidiostats, antibiotics, and litters (built up vs. fresh) on broiler chick performance

Furazolidone

Benazet, F.; et al., 1970, Scand. J. Infect. Dis., v. 2 (2), 139-143
intestinal and hepatic parasites, nitroheterocyclic antiparasitics, laboratory studies of chemotherapeutic activity and toxicity in exper. animals

Furazolidone

Borland, E. D., 1979, Vet. Rec., v. 105 (10), 169
nervous syndrome in pigs, suspected furazolidone toxicity

Furazolidone

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Furazolidone

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Furoxone (Furazolidone)

Cherian, Z.; Jose, M. P.; and Jayakumar, K. M., 1977, Kerala J. Vet. Sc., v. 8 (1), 71-72
ancylostomiasis in mongrel dogs, clinical treatment trials: decaris highly effective without toxicity, tetracap reduced severity of infection, furoxone had no effect

Furazolidone (Furoxon)

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Microbiol. (6), 35-50
Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Furazolidone -- Continued.

Furazolidone (I)

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidiostats in feeds, qualitative identification test

Furazolidone (II)

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidiostats in feeds, qualitative identification test

Furazolidone

Kluska, J., 1978, Terap. i Leky, v. 6, v. 28 (3), 107-112
Enterobius, lambliasis, institutionalized children, control by improved sanitation and hygiene in conjunction with anthelmintics

Furazolidone (Furoxone)

Laemmler, G.; Saenger, I.; and Wegerhof, P. H., 1978, Tropenmed. u. Parasitol., v. 29 (2), 178-182
Litomosoides carinii in Mastomys natalensis, filaricidal activity of furazolidone

Furazolidone

Levi, G. C.; de Avila, C. A.; and Amato Neto, V., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (6), 422-424
giardiasis, humans, efficacy of various drugs, comparative study, side effects

Furazolidone (NF-180)

Manuel, M. F.; Morales, E.; and Trovela, E., 1977, Philippine J. Vet. Med., v. 16 (1-2), 20-30
Leucocytozoon caulleryi, chickens under natural conditions, clopidol, halofuginone and furazolidone given in feed, clopidol 100% effective, no detrimental effects by any drug on host growth or red and white blood cell count

Furazolidone (NF-180)

Manuel, M. F.; and Trovela, E., 1977, Philippine J. Vet. Med., v. 16 (1-2), 31-39
Leucocytozoon caulleryi, chickens under field conditions, efficacy of halofuginone and furazolidone alone and in combination, given with feed, furazolidone at high dosage showed some adverse host growth effects, neither drug showed adverse effect on blood picture

Furazolidone (Furoxon)

Moursy, A. A. M., 1977, J. Egypt. Vet. Med. Ass., v. 37 (2), 99-110
furazolidone, ducks, toxicity, physiopathological changes

Furazolidone (Furoxone)

Rassi, A.; and Ferreira, H. de O., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (5), 235-262
Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best response

Furazolidone -- Continued.

Furazolidone

St. Omer, V. V., 1978, Vet. Med. and Small Animal Clin., v. 73 (9), 1125-1128, 1132
furazolidone, toxicity in animals, review

Furoxone (NF-180)

Santiago, M. A. M.; da Costa, U. C.; and da Silva, O. L., 1977, Rev. Centro Cien. Rurais, v. 7 (3), 297-301

Eimeria spp., sheep, natural infection, amprolium, furoxone, not sufficient control for sheep grazing on pasture

Furazolidone

Villadegado, M. A.; and Abuso, O. T., [1977], Philippine J. Vet. Med., v. 15 (1-2), 1976, 123-128

furazolidone as supplement to commercial broiler feed (already containing penicillin and amprolium), results indicate supplementation unwarranted

Furazolidone (Furoxone; Furoxona)

Wolfe, M. S.; and Moede, A. L., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 762-765

Giardia lamblia, serum sickness in 2 persons who had received furazolidone therapy, possible incrimination of tartrazine (component of Latin American-produced furazolidone (Furoxona) which is no longer included in United States-produced furazolidone (Furoxone))

Furodazole -- 2-(2-Furyl)-7-methyl-1H-imidazo[4,5-f]quinolin-9-ol.

Furodazole

Alaimo, R. J.; et al., 1978, J. Med. Chem., v. 21 (3), 298-300

furodazole, anthelmintic trials with experimental animals, bunamidine and niclosamide used as reference drugs

Furoxona. See Furazolidone.

Furoxone. See Furazolidone.

2-(2-Furyl)-7-methyl-1H-imidazo[4,5-f]quinolin-9-ol. See Furodazole.

Fusidic acid, sodium salt

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

- G-418
Loebenberg, D.; Counelis, M.; and Waitz, J. A., 1975, *Antimicrob. Agents and Chemotherapy*, v. 7 (6), 811-815
antibiotic G-418, promising activity against a variety of protozoa and helminths in vivo and in vitro
- GABA [γ -Aminobutyric acid]
Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, *Rev. Iber. Parasitol.*, v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility
- Gabroral. See Paromomycine.
- Galactosyl ceramide
Alving, C. R.; et al., 1979, *Science* (4411), v. 205, 1142-1144
Plasmodium berghei, mice, therapeutic effects of glycolipids in liposomes against sporozoite-induced malaria
- Galinid. See Tetramisole.
- Gallium nitrate
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Gamatox. See Benzene hexachloride.
- Gamaverm. See Piperazine or Thiabendazole.
- Gamexane. See Benzene hexachloride.
- Gamma benzene hexachloride. See Benzene hexachloride.
- Gamma BHC. See Benzene hexachloride.
- Gamma-isomer hexachlorocyclohexane. See Benzene hexachloride.
- Gammexane powder. See Benzene hexachloride.
- Ganaseg. See Berenil.
- Gantrisin. See Sulfisoxazole.
- Gardona. See Tetrachlorvinphos.
- Garrathion. See Carbophenothion.
- Gasil 35
Kirkwood, A. C., 1974, *Internat. Pest Control*, v. 16 (6), 12-15
Dermanyssus gallinae, laboratory and field experiments using gasil 35 and gasil 200
- Gasil 200
Kirkwood, A. C., 1974, *Internat. Pest Control*, v. 16 (6), 12-15
Dermanyssus gallinae, laboratory and field experiments using gasil 35 and gasil 200
- Gentamicin (Genticyn)
Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, *Chemotherapy*, v. 25 (4), 222-226
Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizontocidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection
- Gentian violet
Daniiarov, I. A.; et al., 1978, *Veterinariia, Moskva* (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested; none effective
- Gentian violet
Iakubovskii, M. V.; and Zen'kov, A. V., 1977, *Vet. Nauka--Proizvod.*, Trudy, Minsk, v. 15, 72-78
nematodes, swine raising complexes, combined control measures, sanitation, anthelmintics, suiverm most effective
- Gentian violet
Sakamoto, T., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolical effects in vitro and/or therapeutic effects in mice
- Genticyn. See Gentamicin.
- Germanin. See Suramin.
- Gevisol
Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures
- Glacial acetic acid. See Acetic acid.
- Gloxazone. See Dithiosemicarbazone.
- Glucantime. See N-Methylglucamine antimonate.
- Glucosyl ceramide
Alving, C. R.; et al., 1979, *Science* (4411), v. 205, 1142-1144
Plasmodium berghei, mice, therapeutic effects of glycolipids in liposomes against sporozoite-induced malaria

D-Glutamic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

L-Glutamic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Glutaraldehyde

Kunstyr, I.; and Ammerpohl, E., 1978, *Lab. Animals*, v. 12 (2), 95-97

Spirocnucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Glycamide. See Glycarbylamide.

Glycarbylamide -- Glycamide; Imidazole-4,5-dicarboxamide.

Glycamide

Krylov, M. V.; et al., 1975, *Parazitologiya*, Leningrad, v. 9 (1), 82-91

Eimeria tenella, mechanisms of resistance to glycamide

Glycerol

Brohn, F. H.; and Clarkson, A. B., jr., 1978, *Acta Trop.*, v. 35 (1), 23-33

Trypanosoma brucei brucei, effect of glycerol on anaerobic glycolysis in vitro, concomitant administration of salicylhydroxamic acid and glycerol to infected rats results in rapid clearance of parasitemia

Glycerol + Salicylhydroxamic acid

Evans, D. A.; and Holland, M. F., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (2), 203-204

Trypanosoma vivax, mice, effective treatment with salicylhydroxamic acid + glycerol

Glycerol

Nathan, H. C.; et al., 1979, *J. Protozool.*, v. 26 (4), 657-660

Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Glycerol/SHAM

Nathan, H. C.; et al., 1979, *J. Protozool.*, v. 26 (4), 657-660

Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Glycerol + Salicylhydroxamic acid

Van Der Meer, C.; Versluijs-Broers, J. A. M.; and Opperdoes, F. R., 1979, *Exper. Parasitol.*, v. 48 (1), 126-134

Trypanosoma brucei brucei, rats, treatment with salicylhydroxamic acid + glycerol and suramin + glycerol

Glycerol + Suramin

Van Der Meer, C.; Versluijs-Broers, J. A. M.; and Opperdoes, F. R., 1979, *Exper. Parasitol.*, v. 48 (1), 126-134

Trypanosoma brucei brucei, rats, treatment with salicylhydroxamic acid + glycerol and suramin + glycerol

Goodwinol cream

Tierney, F.; and Baillie, J., 1979, *Vet. Med. and Small Animal Clin.*, v. 74 (1), 69-70

Cnemidoptes pilae, Mexican red-headed parrot (beak, face), case report, combined aerosol (malathion solution) and topical treatment (eurax and Goodwinol cream) highly effective and less stressful

Gramicidin

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Guanazolo -- 8-Azaguanine.

8-Azaguanine

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

8-Azaguanine

Irvin, A. D.; and Young, E. R., 1978, *Research Vet. Sc.*, v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

8-Azaguanine

Sinden, R. E.; and Smalley, M. E., 1979, *Parasitology*, v. 79 (2), 277-296

Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

8-Azaguanine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Guanosine + Cordycepin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Guanosine + EHNA + Cordycepin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Gynben vaginal cream. See Diethylstilbestrol or Diiodohydroxyquin or Sulfadiazine.

Halazone

Fripp, P. J.; Armstrong, F. A.; and Jaskulla, E., 1972, South African Med. J., v. 46 (47), 1819-1822

Schistosoma haematobium, S. mansoni, cercaria-
acidal activity, hypochlorite solutions vs.
halazone solution, formulations to make small
amounts of water potable

Halofuginone -- 7-Bromo-6-chlorofebrifugine-
hydrobromide; dl-7-Bromo-6-chloro-3-[3-(3-
hydroxy-2-piperidyl)-acetyl]-4(3H)-quina-
zolinone hydrobromide; Cebegine RU 19110;
DL-trans-6-Chloro-7-bromo-3-[3-(3-hydroxy-2-
piperidyl) acetyl]-4-(3H)-quinazolinone
hydrobromide; Stenorol.

Halofuginone (Stenorol)

Bedrnik, P.; et al., 1979, Arch. Geflugelk.,
v. 43 (1), 7-10

Eimeria spp., chickens (exper.), comparative
efficiency of anticoccidials in combination
with growth promotant nitrovin

Halofuginone (Stenorol)

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1469-1475

Eimeria spp. (recent field isolates resistant
to various drugs), chickens, halofuginone,
action cidal rather than static

Halofuginone (Stenorol)

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant
to various drugs), halofuginone with roxar-
sone and/or bacitracin MD

Halofuginone + Bacitracin MD

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant
to various drugs), halofuginone with roxar-
sone and/or bacitracin MD

Halofuginone + Roxarsone

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant
to various drugs), halofuginone with roxar-
sone and/or bacitracin MD

Halofuginone + Roxarsone + Bacitracin MD

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant
to various drugs), halofuginone with roxar-
sone and/or bacitracin MD

Halofuginone (Stenorol)

Edgar, S. A.; and Flanagan, C., 1979, Poultry
Science, v. 58 (6), 1483-1489

Eimeria spp., turkeys, halofuginone effective

Halofuginone

Latter, V. S.; and Wilson, R. G., 1979, Para-
sitology, v. 79 (1), 169-175

Eimeria tenella, factors influencing assess-
ment of anticoccidial activity in cell
culture

Halofuginone -- Continued.

Halofuginone (Stenorol)

Manuel, M. F.; Morales, E.; and Trovela, E.,
1977, Philippine J. Vet. Med., v. 16 (1-2),
20-30

Leucocytozoon caulleryi, chickens under
natural conditions, clopidol, halofuginone
and furazolidone given in feed, clopidol
100% effective, no detrimental effects by
any drug on host growth or red and white
blood cell count

Halofuginone (Stenorol)

Manuel, M. F.; and Trovela, E., 1977, Philip-
pine J. Vet. Med., v. 16 (1-2), 31-39

Leucocytozoon caulleryi, chickens under
field conditions, efficacy of halofuginone
and furazolidone alone and in combination,
given with feed, furazolidone at high dosage
showed some adverse host growth effects,
neither drug showed adverse effect on blood
picture

Stenorol (Cebegine RU 19110)

Mladenovic, Z.; Movsesijan, M.; and Borojevic,
D., 1978, Vet. Glasnik, v. 32 (10), 829-834

Eimeria spp., chickens (exper.), mixed in-
fections, cycostat, nitryl, and stenorol

Stenorol

Mörch, J., 1978, Nord. Vet. Med., v. 30 (6),
253-259

Eimeria spp., chickens (exper.), efficacy of
coccidiostats in feed, better production
efficiency of medicated groups

Stenorol

Morrison, W. D.; Ferguson, A. E.; and Leeson,
S., 1979, Poultry Science, v. 58 (5), 1160-
1166

Eimeria spp., chicks (exper.), salinomycin
and stenorol compared with other anticoc-
cidials, efficacy and effect on chick per-
formance

Stenorol

Pujic, P.; Krecov, M.; and Simic, V., 1977,
Vet. Glasnik, v. 31 (4), 277-280

[Eimeria] spp., chickens, stenorol prevents
coccidiosis under laboratory and field con-
ditions, during fattening period

Halofuginone

Schindler, P.; et al., 1979, Poultry Science,
v. 58 (1), 23-27

Eimeria spp., broiler chicken pen trials,
arprinocid in feed highly effective pro-
phylaxis, comparison with halofuginone, mo-
nensin, nicarbazin, and pancloxin: England;
France; Germany

Halofuginone

Toriumi, T.; et al., 1978, Scient. Rep. Fac.
Agric. Okayama Univ. (52), 49-53

Leucocytozoon caulleryi, chickens (exper.),
sulfamonomethoxine and halofuginone in feed
prevented infection

Haloxon -- Coopers Summer Drench; 0,0-Di(2-chloroethyl)-0-(3-chloro-4-methylcoumarin-7-yl)phosphate; Eustidil; Loxon.

Haloxon (Eustidil)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Haloxon (Loxon)

Ogunsusi, R. A., 1979, Research Vet. Sc., v. 27 (1), 131-132
Haemonchus contortus, sheep, oxfendazole, haloxon, efficacy against arrested larvae, controlled trial, dry season: northern Nigeria

Loxon

Petrov, Iu. F., 1978, Veterinariia, Moskva (5), 64-66
[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Haloxon

Reiner, E.; et al., 1978, Comp. Biochem. and Physiol., v. 60C (2), 155-157
Metastrongylus apri, cholinesterase, kinetic properties with respect to substrate hydrolysis and inhibition by organophosphorus compounds

Haloxon (Coopers Summer Drench)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thia-bendazole, results confirm the usefulness of levamisole, naphthalophos, and radoxanide for this purpose, haloxon and nitroxynil are also useful chemical alternatives

Halquinol -- Quixalin.

Quixalin

Sharma, S. C.; and Asthana, V. S., 1978, Indian Vet. Med. J., v. 2 (2), 83-84
paramphistomes, buffaloes, cows, and bullocks, clinical aspects, distodin and quixalin: Andala, Aligarh District, Uttar Pradesh

α -HCH. See Benzene hexachloride.

γ -HCH. See Benzene hexachloride.

Helatac. See Parabendazole.

Helmatac. See Parabendazole.

Helmex. See Pyrantel.

Helmintia-P. See Phenothiazine or Piperazine or Senna or Tin or Vernonia anthelmintica.

Helmoral B

Ojukwu, E. M.; and Ikeme, M. M., 1978, Bull. Animal Health and Prod. Africa, v. 26 (3), 242-247 [pages assembled incorrectly]
helminths, Zebu cattle, helmoral B, critical tests and field trials: Nigeria

Hematoporphyrin

Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Hematoporphyrin

Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, Biochem. Pharmacol., v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism

Hematoporphyrin D

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Hematoporphyrin IX diacetate

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Hematoporphyrin IX dimethyl ether

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Heme

Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

- Heme**
Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism
- Hemosporidin**
Nikol'skii, S. N.; Niki forenko, V. I.; and Pozov, S. A., 1977, *Veterinariia*, Moskva (4), 71-75
Piroplasma jakimovi, cattle, morphological and biological comparison with P. bigeminum, epizootiology (Ixodes ricinus as main vector); frequent association with leptospirosis; treatment: Siberia
- Hempa (Hexamethyl phosphoramidate)**
Adhami, N.; and Khan, N. H., 1976, *Indian J. Entom.*, v. 37 (1), 1975, 60-63
Cimex hemipterus, hempa as chemosterilant, reduction of oviposition and egg viability, more effective against males than females as measured by egg hatchability
- Heparin**
Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Heptachlor -- 1,4,5,6,7,8,8-Heptachloro-3a,4,7,7a-tetrahydro-4,7-methano-1H-indene.**
- Heptachlor**
Lourens, J. H. M.; and van de Klashorst, G., 1979, *Ztschr. Ang. Entom.*, v. 87 (3), 230-238
Rhipicephalus appendiculatus, organochlorine susceptible and tolerant populations: East Africa
-
- 1,4,5,6,7,8,8-Heptachloro-3a,4,7,7a-tetrahydro-4,7-methano-1H-indene. See Heptachlor.
- Heptenophos -- 5-(0,0-Dimethylphosphoryl)-6-chlorobicyclo (3,2,0)-hepta-1,5-dien;**
Hoe 2982; Ragadan.
- Heptenophos (Ragadan)**
Bonin, W., 1977, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 90 (2), 34-37
Amblyomma hebraeum, Psoroptes cuniculi, Melophagus ovinus, Dermanyssus gallinae, heptenophos, rapid mode of action, broad range of efficacy, short residual effect and effective as a vapour poison, compared with other standard drug preparations
- Hetol.** See 1,4-Bis(trichloromethyl) benzene.
- Hetolin -- β,β,β -Tris-(4-chlorophenyl)-propionic acid-N'-methylpiperazide.**
- β,β,β -Tris-(4-chlorophenyl)-propionic acid-N'-methylpiperazide
Reinhardt, P., 1978, *Monatsh. Vet.-Med.*, v. 33 (23), 898-901
Dicrocoelium dendriticum, sheep, efficacy of various anthelmintics compared
-
- Hetrazan.** See Diethylcarbamazine.
- Hexachlorane.** See Benzene hexachloride.
- Hexachlorane, gamma isomer.** See Benzene hexachloride.
- 3,5,6,3',5',6'-Hexachlor-2,2'-dioxydiphenylsulfide**
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,5,6,3',5',6'-Hexachlor-2,2'-dioxydiphenylsulfone**
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,5,6,3',5',6'-Hexachlor-2,2'-dioxydiphenylsulf-oxide**
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 1,2,3,4,5,6-Hexachlorocyclohexane.** See Benzene hexachloride.
- 1,2,3,4,10,10-Hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-endo-1,4-exo-5,8-dimethanonaphthalene.** See Dieldrin.

Hexachloroethane -- Avlothane.

Avlothane

- Reddy, R. G., 1979, *Livestock Advis.*, v. 4 (3), 37-40
Balantidium coli and mixed infection with amphistomes, cattle and buffaloes, incidence and treatment

Hexachlorophene -- Bilevon-Injektion; Distodin; 2,2'-Methylenebis(3,4,6-trichlorophenol); Previken.

Bilevon-Injektion

- Andrews, P.; Dorn, H.; and Wirtz, S., 1977, *Vet.-Med. Nachr.* (2), 129-134
Fasciola hepatica, F. gigantica, cattle (nat. and exper.), bilevon-Injektion, bilevon-R tablet formulation, rafoxanide, and nitroxyuil compared

Hexachlorophene

- Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10), 28-36
cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results

Hexachlorophene

- Bath, M. L., 1978, *J. Small Animal Practice*, v. 19 (4), 241-244
hexachlorophene, toxicity, dogs, case reports

Hexachlorophene

- Gadzhev, Ia. G.; et al., 1977, *Veterinariia*, Moskva (5), 63-64
Fasciola gigantica, cattle, acute infection outbreak, related to increased Lymnaea auricularia population in newly irrigated area: Azerbaïdzhan SSR

Previken

- Merenyi, L., 1978, *Magy. Allat. Lapja*, v. 100, v. 33 (2), 132-134
fascioliasis, control, treatment: Nograd county

Hexachlorophene (Distodin)

- Parshad, V. R.; and Guraya, S. S., 1978, *Vet. Parasitol.*, v. 4 (2), 111-120
4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces

2,2'-Methylenebis(3,4,6-trichlorophenol)

- Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Distodin

- Sharma, S. C.; and Asthana, V. S., 1978, *Indian Vet. Med. J.*, v. 2 (2), 83-84
paramphistomes, buffaloes, cows, and bullocks, clinical aspects, distodin and quixalin: Andala, Aligarh District, Uttar Pradesh

Hexachloroxytol. See 1,4-Bis(trichloromethyl) benzene.

Hexachlorparaxylo. See 1,4-Bis (trichloromethyl) benzene.

Hexakis (β,β-dimethylphenethyl) distannoxane -- Hexakis (2-methyl-2-phenylpropyl)distannoxane; SD-14114; Vendex.

Vendex (SD-14114)

- Meleney, W. P.; and Roberts, I. H., 1979, *J. Med. Entom.*, v. 16 (1), 52-58
Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Hexakis (2-methyl-2-phenylpropyl)distannoxane. See Hexakis (β,β-dimethylphenethyl) distannoxane.

Hexamethylmelamine

- Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Hexamethyl phosphoramidate. See Hempa.

Hexanema

- Khan, M. H., 1979, *Indian Vet. J.*, v. 56 (9), 739-743
Lipeurus caponis, Menacanthus stramineus, White Leghorn fowl, organophosphorus insecticides, costs evaluated

Hexicide. See Benzene hexachloride.

Hexylresorcinol

- Goldsmith, R. S., 1978, *South. Med. J.*, v. 71 (12), 1513-1515
Metagonimus yokogawai, American woman traveling in the Orient, case report, chronic diarrhea treated unsuccessfully with hexylresorcinol, cure with tetrachloroethylene: California

Hexyl-resorcinol

- Perez, C.; et al., 1977, *Rev. Med. Chile*, v. 105 (8), 520-522
Trichuris trichiura, massive infections in children, clinical manifestations, diagnostic serology, successfully cured by rectal administration of hexyl-resorcinol: Chile

Hexyl-resorcinol

- Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, *Rev. Iber. Parasitol.*, v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Hibitane. See Chlorhexidine.

Hi-Lo Dip

- Pilarczyk, J. P., 1979, *Canine Pract.*, Santa Barbara, v. 6 (6), 51-52
chemical burn and toxicity in dog treated with flea dip that had been improperly stored

Hipolen-6. See Crufomate.

Homidium -- Ethidium; Ethidium bromide; Homidium bromide.

Ethidium bromide-DNA complex

Avila, J. L.; Bretana, A.; and Avila, A., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (3), 456-460

Trypanosoma cruzi, mice, chemotherapy with ethidium bromide-DNA complex, effectiveness seems limited to early Chagas' disease

Ethidium bromide

Bachrach, U.; et al., 1979, *Exper. Parasitol.*, v. 48 (3), 464-470

Leishmania spp., effect of ethidium, pentamidine, and methylglyoxal-bis (guanyldrazone) on growth and on polyamine, RNA, and DNA synthesis

Ethidium bromide

Benard, J.; Riou, G.; and Saucier, J. M., 1979, *Nucleic Acids Research*, v. 6 (5), 1941-1952

Trypanosoma cruzi at different stages of culture and grown in presence of ethidium, kinetoplast DNA, characterization by sedimentation analysis

Ethidium bromide

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Ethidium bromide

Field, R. C.; et al., 1978, *Brit. J. Pharmacol.*, v. 62 (2), 159-164

effects of chloroquine, primaquine and ethidium on precursor incorporation into DNA, RNA and protein in mammalian tissues

Ethidium Br

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Ethidium

Gutteridge, W. E.; Dave, D.; and Richards, W. H. G., 1979, *Biochim. et Biophys. Acta*, v. 582 (3), 390-401

Kinetoplastida spp., Plasmodium spp., conversion of dihydroorotate to orotate, mechanism of reaction different in these 2 groups of protozoa, possible target of chemotherapeutic attack

Ethidium bromide

Jadin, J. M.; et al., 1977, *Ann. Soc. Belge Med. Trop.*, v. 57 (4-5), 525-531

Trypanosoma cruzi intra- and extracellular forms, *T. brucei* extracellular forms, mice, ethidium bromide vs. ethidium bromide-DNA complexes as therapy

Ethidium

Jones, R. L.; Davidson, M. W.; and Wilson, W. D., 1979, *Biochim. et Biophys. Acta*, v. 561 (1), 77-84

chloroquine does not bind to DNA by classical intercalation mechanism typical of quinacrine and ethidium

Homidium -- Continued.

Ethidium

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Homidium bromide (Ethidium)

Worms, M. J.; and Hawking, F., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (5), 548-549

Litomosoides carinii-infected *Sigmodon hispidus*, effects of suramin, homidium bromide, quinapyramine, diminazene, and isometamidium after observation period of more than 5 weeks

Homidium bromide. See Homidium.

Homomycin. See Hygromycin.

Horseradish, green leaves

Klenov, A. P., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 152-155

[*Bothriocephalus*], white amur, effectiveness of various anthelmintics

Humatin. See Paromomycin.

Hyaluronic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Hycanthone -- 1-[[2-(Diethylamino) ethyl] amino]-4-(hydroxy-methyl)-thioxanthen-9-one; Etrenol; Hycanthon methanesulfonate; Hycanthon methanesulphonate; Hycanthon sulfamate; MW 356.48; Win 24,933-2.

Hycanthon

Abdel Samad, M. M.; et al., 1977, *Tropenmed. u. Parasitol.*, v. 28 (4), 554-559

Schistosoma mansoni, mice, liver monoamine oxidase activity during course of infection and after chemotherapy, may be useful index for progression or regression of liver fibrosis

Hycanthon (Etrenol)

Andrade, Z. A.; et al., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (3), 160-170

schistosomiasis, human, hycanthon, acute hepatic toxicity

Hycanthon -- Continued.

- Etrenol**
Aslamazov, E. G., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 222-224
schistosomiasis, etrenol treatment, literature review
- Hycanthon**
Batzinger, R. P.; and Bueding, E., 1977, *J. Pharmacol. and Exper. Therap.*, v. 200 (1), 1-9
mutagenic activities in vitro and in vivo of 5 antischistosomal compounds, comparative anti-Schistosoma mansoni activities of hycanthon, IA-4, and IA-4 N-oxide, observations provide evidence that mutagenic activities can be dissociated from desired chemotherapeutic effects by suitable structural modifications
- Hycanthon**
Bina, J. C.; and Prata, A., 1974, *Rev. Soc. Brasil. Med. Trop.*, v. 8 (4), 217-222
schistosomiasis mansoni, attempted control using mass therapy with hycanthon, small village population in endemic area: Canabrava, Bahia, Brasil
- Hycanthon**
Buaiz, V.; et al., 1976, *AMB, Rev. Ass. Med. Brasil.*, v. 22 (5), 171-174
S[chistosoma] mansoni, 3,100 patients treated with hycanthon, tolerance, cure rate, toxicity, indications for use, and proposed dosage schedule: Brazil
- Hycanthon**
Buchanan, N., 1977, *South African Med. J.*, v. 52 (21), 834 [Letter]
hycanthon therapy resulting in fatal massive hepatic necrosis, child, case report
- Hycanthon**
Buchanan, N.; et al., 1978, *South African Med. J.*, v. 53 (7), 257-258
Schistosoma haematobium, 11-year-old girl, case report, fatal hepatic necrosis associated with hycanthon therapy: South Africa
- Hycanthon**
Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, *J. Parasitol.*, v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy
- Hycanthon**
de Castro, H. N.; Nohmi, N.; and Marques, M. do R., 1973, *Rev. Ass. Med. Minas Gerais*, v. 24 (4), 223-232
Schistosoma mansoni, young woman, toxic hepatitis after hycanthon therapy for hepatic-intestinal schistosomiasis, combined cumulative effects of hycanthon and oral contraceptive thought to be causative factors: Belo Horizonte, Brazil
- Hycanthon (Etrenol)**
Cohen, C., 1978, *Gastroenterology*, v. 75 (1), 103-106
schistosomiasis, case reports of hepatic toxic hepatitis with massive hepatic necrosis in patients treated with intramuscular hycanthon
- Hycanthon** -- Continued.
Hycanthon (Etrenol)
Coutinho, A. D.; and Barreto, V. S., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (1), 57-70
S[chistosoma] mansoni, human, hycanthon, frequent side effects, possesses therapeutic value if carefully monitored
- Hycanthon**
da Cunha, A. S.; et al., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (2), 131-136
S[chistosoma] mansoni, human, hycanthon, therapeutic evaluation
- Hycanthon**
da Cunha, A. S.; et al., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (3), 213-222
S[chistosoma] mansoni, humans, hycanthon toxicity
- Hycanthon**
Dennis, E. W., 1978, *South African Med. J.*, v. 54 (4), 137-138 [Letter]
reply to report of Buchanan, N. et al. (*South African Med. J.*, v. 53, p. 257) regarding fatal hepatic necrosis in association with hycanthon therapy
- Hycanthon**
Dias, L. C. de S.; et al., 1978, *Rev. Saude Pub.*, S. Paulo, v. 12 (1), 110
Schistosoma mansoni, isolation of strain resistant to hycanthon and to oxamniquine
- Hycanthon**
Ferraz, M. P. T.; et al., 1973, *Rev. Paul. Med.*, v. 81 (5), 275-278
Schistosoma mansoni, symptomatic psychosis in 2 persons who had received hycanthon treatment for schistosomiasis, case reports: Brazil
- Hycanthon**
Foster, R., 1973, *Rev. Inst. Med. Trop. S. Paulo*, v. 15, suppl. 1 (6), 1-9
Schistosoma mansoni, S. haematobium, S. japonicum, oxamniquine, efficacy and toxicity in various exper. animals, clinical trials in humans
- Hycanthon**
Godoy, P.; et al., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (2), 114-120
[Schistosoma] mansoni, human, hycanthon, hepatic failure and death, case report, pathologic findings
- Hycanthon**
Goncalves, C. S.; et al., 1977, *AMB, Rev. Ass. Med. Brasil.*, v. 23 (9), 305-308
schistosomiasis, humans, hepatitis and other hepatic pathology after hycanthon therapy
- Hycanthon methanesulfonate**
Guerra, M. de O.; et al., 1974, *Rev. Ginec. e Obst.*, Sao Paulo, v. 131 (9-10), 241-245
administration of schistosomal drug hycanthon to lactating rats, offspring had decreased liver weight and evidence of liver damage, possible excretion of drug or its metabolites into mother's milk or alteration of milk metabolism by drug

Hycanthone -- Continued.

Hycanthone

Guimaraes, R. X.; et al., 1979, *AMB, Rev. Ass. Med. Brasil.*, v. 25 (2), 48-50
Schistosoma mansoni, human hepato-intestinal form, resistance to hycanthone and oxamniquine: Brazil

Hycanthone

Hillman, G. R.; Gibler, A. M.; and Anderson, J. W., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 992-997
Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

Hycanthone

Hillman, G. R.; Senft, A. W.; and Gibler, W. B., 1978, *J. Parasitol.*, v. 64 (4), 754-756
Schistosoma mansoni, hycanthone, mode of action, possible explanations of some discrepancies in results in published reports

Hycanthone

Huggins, D.; Correia, J. U.; and de Sousa, E. M., 1973, *Rev. Soc. Brasil. Med. Trop.*, v. 7 (5), 305-311
Schistosoma mansoni, humans, evaluation of hycanthone as therapy

Hycanthone

Katz, N., 1971, *Rev. Soc. Brasil. Med. Trop.*, v. 5 (2), 55-60
S[chistosoma] mansoni, hycanthone therapy less effective in treating newly infected persons than those with chronic infections, fecal egg counts were however markedly decreased in persons with new infections

Hycanthone methanesulfonate

Katz, N., 1972, *Rev. Ass. Med. Minas Gerais*, v. 23 (1), 27-34
human schistosomiasis mansoni, outline for therapeutic trials with comparison of effects of hycanthone and experimental drug UK-4271

Hycanthone

Katz, N.; et al., 1973, *Rev. Soc. Brasil. Med. Trop.*, v. 7 (6), 381-387
Schistosoma mansoni, isolation of drug resistant strain (WW strain), reactions in mice to therapy with hycanthone, niridazole and oxamniquine compared with reactions of LE drug sensitive strain

Hycanthone

Katz, N.; and Pellegrino, J., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (5), 245-252
Schistosoma mansoni, Cebus monkeys, correlation of number of eggs per gram of rectal tissue with number of female worms, challenge infection effect, or drug action

Hycanthone sulfamate

Katz, N.; and Pellegrino, J., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (5), 245-252
Schistosoma mansoni, Cebus monkeys, correlation of number of eggs per gram of rectal tissue with number of female worms, challenge infection effect, or drug action

Hycanthone -- Continued.

Hycanthone

Magzoub, M.; and Adam, S. E. I., 1978, *J. Pharmacol. and Exper. Therap.*, v. 206 (2), 382-387
Schistosoma mansoni-infected Arvicanthus niloticus, evaluation of oral therapy with single and multiple doses of hycanthone and oxamniquine alone or in combination

Hycanthone

Marinho, R. P.; et al., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (1), 54-59
Schistosoma mansoni, man treated with hycanthone, development of acute yellow atrophy of liver, fatal illness, case report: Minas Gerais, Brazil

Hycanthone

Marinho, R. P.; et al., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (6), 354-361
S[chistosoma] mansoni, human, yellow atrophy of liver resulting from hycanthone therapy, fatal illness, case report: Caete, Minas Gerais, Brazil

Hycanthone

Marinho, R. P.; and Neves, J., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (2), 70-76
S[chistosoma] mansoni and septicemic salmonellosis, mixed human infection, both infections cured by hycanthone

Hycanthone methane sulphonate

Maritz, J. C., 1970, *South African Med. J.*, v. 44 (5), 126-128
bilharziasis, school children, hycanthone clinical trials: Dennilton, Transvaal

Hycanthone

Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, *Rev. Brasil. Pesqui. Med. e Biol.*, v. 2 (1), 45-49
Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

Hycanthone

Neame, K. D.; et al., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 587-588
Schistosoma mansoni, hycanthone inhibits nucleic acid synthesis in vitro but it seems unlikely that this is mechanism by which drug kills worms in vivo

Hycanthone

Neves, J.; Marinho, R. P.; and Raso, P., 1972, *Rev. Brasil. Pesqui. Med. e Biol.*, v. 5 (3-4), 91-100
S[chistosoma] mansoni, young woman presenting with hepatic and gastrointestinal symptoms, differential diagnostic problems, successfully treated with hycanthone, clinical case report: Belo Horizonte, Brasil

Hycanthone

Noleto, P. A.; et al., 1974, *Rev. Soc. Brasil. Med. Trop.*, v. 8 (6), 335-340
Schistosoma mansoni, clinical trials using hycanthone to treat 1,000 persons from local endemic areas of Brazil

Hycanthone -- Continued.

Hycanthone

de Oliveira, C. A.; et al., 1969, Rev. Inst. Med. Trop. S. Paulo, v. 11 (2), 130-139
Schistosoma mansoni, humans, intramuscular administration of hycanthone, clinical trials

Hycanthone

de Oliveira, C. A.; et al., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (3), 202-212
Schistosoma mansoni, human, acute cases, hycanthone

Hycanthone methanesulfonate (MW 356.48)

Ong, T. M., 1978, Mutation Research, v. 55 (1), 43-70
hycanthone and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Hycanthone

Pedro, R. de J.; et al., 1979, Rev. Brasil. Pesqui. Med. e Biol., v. 12 (2-3), 207-211
Schistosoma mansoni, observations on oxamni-quine therapy: treatment of children, drug resistance of human strain as well as its resistance to hycanthone, hepatic histopathology during therapy, neurotoxic effects, treatment of mixed salmonellosis infection

Hycanthone

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Hycanthone (Etrenol)

Rubidge, C. J.; et al., 1970, South African Med. J., v. 44 (43), 1246
bilharziasis, human urinary tract infections, 3 clinical trials testing hycanthone: Durban, South Africa

Hycanthone methanesulphonate

Saad, A. A.; et al., 1977, Acta Vitaminol. et Enzymol., v. 31 (6), 183-186
Schistosoma mansoni-infected and uninfected mice, β -glucuronidase activity in whole urinary bladder tissue homogenates before and after treatment with hycanthone methanesulphonate

Hycanthone (Etrenol)

Saad, A. A.; et al., 1978, Biochem. Pharmacol., v. 27 (4), 473-474
Schistosoma mansoni-infected and normal mice treated with hycanthone, progressive and prolonged increase in β -glucuronidase activity in liver and spleen homogenates, concluded that hycanthone is hepatotoxic drug and is possibly carcinogenic in mice

Hycanthone

Salgado, J. A.; et al., 1972, Rev. Soc. Brasil. Med. Trop., v. 6 (3), 129-133
S[chistosoma] mansoni, electrocardiographic study of 434 persons before, during, and after therapy with hycanthone, only slight changes recorded, drug therefore recommended for persons with cardiovascular diseases

Hycanthone -- Continued.

Hycanthone

Salih, S. Y.; Abu Sabaa, H.; and Mustafa, D., 1978, J. Trop. Med. and Hyg., v. 81 (7), 126-130
Schistosoma mansoni, S. haematobium, humans, clinical trials testing efficacy of single dose therapy with hycanthone, some gastrointestinal side effects, cure rates of about 90% with significant reduction in egg excretion in those who were not cured: Sudan

Hycanthone

Salih, S. Y.; Voller, A.; and Woodruff, A. W., 1978, Tropenmed. u. Parasitol., v. 29 (3), 269-274 [Erratum v. 30 (1), 1979, 130]
Schistosoma mansoni patients showing different clinical forms of infection and S. haematobium patients before treatment and 6 weeks after treatment with hycanthone, comparison of immunoglobulin levels

Hycanthone

da Silva, L. C.; et al., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (2), 121-130
Schistosoma mansoni, humans, immunodiffusion, hemagglutination, immunofluorescence and eosinophil counts before and after therapy with hycanthone or niridazole

Hycanthone

da Silva, L. C.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (6), 344-349
human schistosomiasis mansoni, immunofluorescence, passive hemagglutination, and immunodiffusion tests used to detect early antibody increases after hycanthone therapy

Hycanthone

da Silva, L. C.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 206-210
S[chistosoma] mansoni, humans parasitologically cured, repeat therapy with niridazole or hycanthone, evaluated by passive hemagglutination, indirect immunofluorescence and immunodiffusion tests

Hycanthone (Etrenol)

Takaoka, L.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (5), 378-386
schistosomiasis mansoni, humans with chronic infection, electrocardiographic alterations after hycanthone therapy

Hycanthone

Vilela, M. P.; et al., 1972, Ann. Soc. Belge Med. Trop., v. 52 (6), 551 [Letter]
Schistosoma mansoni, 13-year-old child, hepatic and psychotic manifestations after hycanthone therapy: Brazil

Hycanthone

Warren, K. S.; et al., 1978, Lancet, London (8060), v. 1, 352-354
Schistosoma mansoni, humans, dose-response to hycanthone established for 3 dosage levels using less drug than that recommended on drug packaging, possible use of these decreased levels for mass therapy especially in endemic areas as a means of avoiding severe toxic reactions

Hycanthone -- Continued.

Hycanthone (Etrenol)

Woolhouse, N. M., 1979, *Biochem. Pharmacol.*, v. 28 (16), 2413-2418
antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Hycanthone methanesulfonate. See Hycanthone.

Hycanthone methanesulphonate. See Hycanthone.

Hycanthone sulfamate. See Hycanthone.

Hydrazones

Molodykh, Zh. V.; et al., 1977, *Khimiko-Farm. Zhurnal*, v. 11 (7), 37-40
Nippostrongylus braziliensis, *Hymenolepis nana*, mice, anthelmintic activity of hydrazones, phthalazones, and phthalazinyhydrazones, relationship to chemical structure

Hydrocortisone

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Hydrogen peroxide

Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Hydrol

El-Bahay, G. H.; et al., 1978, *Vet. Med. J.*, Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, *Argas persicus*, evaluation of 10 insecticides

Hydroprene -- Ethyl 3,7,11-trimethyl-2,3-dodecadienoate.

Ethyl 3,7,11-trimethyl-2,4-dodecadienoate

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, *J. Econom. Entom.*, v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

Hydroxybenzylbenzimidazole

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

9-nor-9-Hydroxycannabinol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

1-Hydroxy-2-(2-chloro-4-isothiocyanato)naphthanalide

Dubey, S. K.; et al., 1978, *J. Med. Chem.*, v. 21 (11), 1178-1181
Hymenolepis spp., *Taenia* sp., laboratory animals, synthesis and screening of substituted 1-hydroxy-2-naphthanalides as potential cestocidal agents, niclosamide used as reference compound

2-Hydroxy-3-(8-cyclohexyloctyl)-1,4-naphthoquinone. See Menoctone.1-(2-Hydroxyethyl)-2-(p-fluorophenyl)-5-nitroimidazole. See Flunidazole.(Hydroxy-2'-ethyl)-1 (p-fluorophenyl) 2-nitro-5-imidazole. See Flunidazole.Hydroxyethylmethyl nitroimidazole. See Metronidazole.1-(2-Hydroxyethyl)-2-methyl-5-nitroimidazole. See Metronidazole.

2-Hydroxy-2',3,4',5,5',6-hexachlorobenzenesulfonanilide

Hayes, T. J.; and Mitrovic, M., 1979, *Experientia*, v. 35 (3), 325-326
Fasciola hepatica, sheep (exper.), potent fasciolicidal activity of 2-hydroxy-2',3,4',5,5',6-hexachlorobenzenesulfonanilide

(±) 9-nor-9 β -Hydroxy hexahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

Hydroxylamine hydrochloride

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

9-Hydroxy- β -lapachone

Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

2-Hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthalenedione. See Lapachol.2-Hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthoquinone. See Lapachol.

- 6-Hydroxymethyl-2-isopropylamino-methyl-7-nitro-1,2,3,4-tetrahydroquinolone. See Oxamniquine.
- 3-Hydroxy-5-methyl isoxazole
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- 6-Hydroxymethyl-2-(2[5-nitro-2-furyl]vinyl)pyridine. See Nifurpirinol.
- N-Hydroxy naphthalimide diethyl phosphate. See Phthalophos.
- 1-Hydroxy-2-naphthanalides, substituted
Dubey, S. K.; et al., 1978, J. Med. Chem., v. 21 (11), 1178-1181
Hymenolepis spp., Taenia sp., laboratory animals, synthesis and screening of substituted 1-hydroxy-2-naphthanalides as potential cestocidal agents, niclosamide used as reference compound
- 2-Hydroxy-5-nitropyridine
Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- DL- β -Hydroxy-norvaline
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- 2-[2-(4-Hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazyl) benzimidazole -- Hoechst 33258; 4-[5-(4-Methyl-1-piperazyl)(2,5'-bi-1-4-benzimidazole-2-yl)] phenol trihydrochloride; Compound 'E' (with (Diamino-4,6-triazinyl-1,3,5-amino-2)-4-phenyl-arsino di (D-thio-3-amino-2-methyl-3-butyric acid)).
- Compound 'E'
Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455
Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity
- 4-[5-(4-Methyl-1-piperazyl)(2,5'-bi-1-4-benzimidazole-2-yl)] phenol trihydrochloride (Hoechst 33258)
Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455
Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity
- (Hydroxy-2-propyl)-1-methyl-2-nitro-5-imidazole. See Secnidazole.
- 4-Hydroxypyrazolo(3,4-d)pyrimidine. See Allopurinol.
- 8-Hydroxyquinoline, copper salt
Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- 8-Hydroxy-quinoline-Bithionol sulfoxide
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives
- 8-Hydroxyquinoline
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- 4-Hydroxy-3 quinoline carbon acid, ethyl esters
Sevcik, B.; et al., 1974, Veterinaria, Praha, v. 16 (5-6), 421-588
Eimeria tenella, chickens, 613 substances screened as coccidiostats, extensive detailed statistical results
- Hydroxyquinolines
Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications
- 4-Hydroxy-1- β -D-ribofuranosylpyrazolo(3,4-d)pyrimidine. See Allopurinol.
- Hydroxystilbamidine isethionate
Nagington, J., 1975, Tr. Ophth. Soc. United Kingdom, v. 95 (2), 207-209
Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity
- 8 β -Hydroxy Δ^9 -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids
- (\pm) 9-nor-9-Hydroxy Δ^6a-10a -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

(±) 11-Hydroxy Δ^{6a-10a} -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, Antimicrob. Agents and Chemotherapy,
v. 16 (5), 674-679

Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids

11-Hydroxy Δ^8 -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, Antimicrob. Agents and Chemotherapy,
v. 16 (5), 674-679

Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids

11-Hydroxy Δ^9 -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, Antimicrob. Agents and Chemotherapy,
v. 16 (5), 674-679

Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids

9-nor-9 β -Hydroxy Δ^{10} -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, Antimicrob. Agents and Chemotherapy,
v. 16 (5), 674-679

Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids

N-Hydroxyurea
Brotherton, J., 1978, Arzneimittel-Forsch.,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Hydroxyurea
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, Antimicrob. Agents and Chemotherapy, v.
15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

Hygromix. See Hygromycin.

Hygromycin -- Homomycin; Hygromix; Hygromycin B;
Hygrovetin.

Hygromycin-B
Gudimenko, I. I., 1970, Nauch. Trudy, Nauchno-
Issled. Vet. Inst., v. 8, 100-104
trichuriasis, oesophagostomiasis, ascariasis,
swine, testing thiabendazole, dithiazanine
iodide, dipterex, hygromycin-B, and
bubulin

Hygromycin -- Continued.

Hygromycin B (Hygromix)
Munoz Davila, A., 1975, Rev. Fac. Cien. Agron.
y Vet., Santo Domingo, v. 3 (3), 47-48
Ascaris suum, Oesophagostomum spp., Trichuris
suis, pigs, action of hygromycin B

Hygrovetin
Parre, J.; Kaarma, A.; and Parts, V., 1977,
Eesti Pollumaj. Akad. Teadusl. Toode Kogum.
(104), 128-136
intestinal helminths, swine farm, control
methods, dehelminthization with hygrovetin;
economic losses, economic benefits of treat-
ment

Homomycin
Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
130
Echinococcus granulosus, scolical effect
of 65 antibiotic, antineoplastic, cytosta-
tic, and other agents in vitro

Hygromycin B. See Hygromycin.

Hygrovetin. See Hygromycin.

Hypocid. See Trichlorfon.

Hypodermacide. See Trichlorfon.

Hypodermin. See Trichlorfon.

Hypodermin-chlorophos. See Trichlorfon.

Hypoxanthine
Irvin, A. D.; and Young, E. R., 1978, Research
Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxan-
thine uptake in vitro could be used as
primary screen for babesicidal drugs but
drugs showing in vitro activity are not
necessarily active in vivo

ICRF 192

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Idoxuridine

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Imidan. See Phosmet.

Imidazole-4-carboxamide, 5-(3, 3-dimethyl-1-triazeno) (WR 139 007)

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, active in screening of antitumor compounds for efficacy against infection

Imidazole-4,5-dicarboxamide. See Glycarbylamide.

Imidazoline hydroiodide, 2-substituted

Ahmad, S.; Kishor, K.; and Shanker, K., 1979, *Indian Drugs*, v. 16 (5), 107-109

Hymenolepis nana, anthelmintic activity of imidazolines in vivo and in vitro, synthesis and acetylcholine esterase inhibitory activity of compounds

Imidazolines

Ahmad, S.; Kishor, K.; and Shanker, K., 1979, *Indian Drugs*, v. 16 (5), 107-109

Hymenolepis nana, anthelmintic activity of imidazolines in vivo and in vitro, synthesis and acetylcholine esterase inhibitory activity of compounds

Imidazo[4,5-f]quinolin-9-ols

Alaimo, R. J.; et al., 1978, *J. Med. Chem.*, v. 21 (3), 298-300

Hymenolepis nana, imidazo[4,5-f]quinolin-9-ols, evaluation of anthelmintic activity using bunamidine as reference drug

Imidocarb -- 3,3'-Bis-(2-imidazolyl) carbanilide dihydrochloride; 3,3'-Bis(2-imidazolyl) carbanilide dipropionate; Imidocarb dihydrochloride; Imidocarb dipropionate; Imizol.

Imidocarb dihydrochloride

El-Abdin, Y. Z.; Hamza, S. M.; and El Refaii, A. H., 1977, *Egypt. J. Vet. Sc.*, v. 13 (2), 1976, 77-84

Theileria annulata, cattle, imidocarb dihydrochloride, serum enzyme activities and chemical constituents before and after treatment

Imidocarb -- Continued.

Imidocarb dipropionate (Imizol)

El Refaii, A. H.; and Michael, S. A., 1979, *Vet. Rec.*, v. 105 (3), 60

urine retention in Egyptian bulls possibly caused by or linked with subclinical *Theileria annulata* infection, imidocarb dipropionate treatment

Imidocarb dihydrochloride

Hashemi-Fesharki, R.; and Amjadi, A. R., 1977, *Arch. Inst. Razi* (29), 83-86

Babesia bovis, outbreak in imported Holstein and Red-Danish cattle, imidocarb dihydrochloride more effective than quinorium sulphate: Rasht, northern Iran

Imidocarb

Irvin, A. D.; and Young, E. R., 1978, *Research Vet. Sc.*, v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Imidocarb

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Imidocarb

Mishra, A. K.; and Sharma, N. N., 1979, *Trop. Animal Health and Prod.*, v. 11 (4), 222-226

Anaplasma marginale, calves (exper.), comparative efficacy of several drugs

Imidocarb (Imizol)

Nathan, H. C.; et al., 1979, *J. Protozool.*, v. 26 (4), 657-660

Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Imidocarb

Rakhimov, T. Kh.; et al., 1977, *Veterinariia, Moskva* (10), 75-77

piroplasmosis, franciellosis, cattle, dimidine and imidocarb tested in various doses, recommended for control

Imidocarb

Thompson, K. C.; et al., 1978, *Trop. Animal Health and Prod.*, v. 10 (2), 75-81

Anaplasma marginale, *Babesia argentina*, *B. bigemina*, cattle under tropical conditions, immunization with virulent organisms followed by drug therapy (ganaseg; gloxazone; emicina) vs. chemoprophylaxis (imidocarb); tick and gastrointestinal parasite control without haemoparasitic control had advantage over no control system at all

Imidocarb -- Continued.

Imidocarb

- Thompson, K. C.; et al., 1978, Trop. Animal Health and Prod., v. 10 (3), 141-144
Anaplasma marginale, *Babesia bigemina*, *B. argentina* (bovis), immunization and chemoprophylaxis of *Bos taurus* calves and subsequent challenge with *Boophilus microplus*, economic gain estimated: Colombia

Imidocarb dipropionate

- Wilson, A. J.; et al., 1979, Austral. Vet. J., v. 55 (2), 71-73
Anaplasma marginale, cattle and splenectomized calves (exper.), long-acting formulation of tetracycline compared with standard tetracycline and imidocarb dipropionate

Imidocarb dihydrochloride. See Imidocarb.

Imidocarb dipropionate. See Imidocarb.

Imizol. See Imidocarb.

Imuran. See Azathioprine.

 β -Indolilacrylic acid

- Christow, C. P., 1974, Rev. Biol. Trop., v. 21 (2), 1973, 187-195
Trichomonas vaginalis, in vitro, reproduction inhibited by anti-amino acids (β -indolilacrylic acid, styrylacetic acid)

Indomethacin

- McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Inimur. See Nifuratel.

Inosine diglycolaldehyde (WR 220 078)

- Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, active in screening of antitumor compounds for efficacy against infection

Insect growth regulators

- Barrett, C. C.; et al., 1978, Southwest. Entom., v. 3 (3), 232-236
Hypoderma lineatum, *H. bovis*, cattle, effect of methoprene on cattle grub eclosion, different times and methods of treatment

Insect growth regulators

- Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, J. Econom. Entom., v. 66 (1), 127-130
Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

Insect growth regulators

- Hopkins, D. E.; and Chamberlain, W. F., 1978, J. Econom. Entom., v. 71 (1), 25-26
Bovicola limbatus, 3rd instars, relationship between inhibition of ecdysis and time and quantity of ingestion of diflubenzuron (inhibitor of cuticle deposition), inhibition of ecdysis increased progressively as age of nymphs increased, timing of treatment important for control

Insect growth regulators

- Hopkins, D. E.; and Chamberlain, W. F., 1978, Southwest. Entom., v. 3 (4), 292-294
Melophagus ovinus, inhibited maturation on sheep dipped in diflubenzuron (preventing formation of pupae) or methoprene (preventing adult eclosion), possible candidate control agents for sheep ked and other larviparous pests

Insect growth regulators

- McDaniel, R. S. II; and Oliver, J. H., jr., 1978, J. Parasitol., v. 64 (3), 571-573
Dermacentor variabilis nymphs, effects of insect growth regulators on certain aspects of developmental and reproductive biology

Insect growth regulators

- Miller, J. A.; et al., 1977, J. Econom. Entom., v. 70 (4), 417-423
Haematobia irritans, evaluation of effectiveness of methoprene in drinking water of cattle to suppress hornfly populations preparatory to release of sterile male flies: island of Molokai, Hawaii

Insect growth regulators

- Norman, J. O.; and Younger, R. L., 1979, J. Med. Entom., v. 16 (1), 43-47
Hypoderma larvae from cattle treated with juvenile hormone analogues, microbial flora, relationships with inflammatory reactions in dorsal subcutis of cattle

Insect growth regulators

- Prasert, V.; et al., 1975, J. Econom. Entom., v. 68 (5), 639-640
Oestrus ovis, sheep, treatment with methoprene nasal spray

Insect growth regulators

- Wright, J. E.; et al., 1974, J. Med. Entom., v. 11 (4), 385-389
Cochliomyia hominivorax, methoprene and R-20458 inhibited emergence in vitro when applied topically or incorporated in larval diet, yearling heifers sprayed with each analogue and infested with larvae displayed little change in serum chemistry and adults emerged from the wound-reared larvae

Insecticides

- Baldry, D. A. T.; et al., 1978, PANS, v. 24 (4), 423-434
[*Trypanosoma brucei gambiense*], control of *Glossina tachinoides* by application of insecticides from helicopters, operational aspects and application techniques: Komoe valley, West Africa

Insecticides

Baldry, D. A. T.; Molyneux, D. H.; and van Wettere, P., 1978, PANS, v. 24 (4), 447-454
Trypanosoma brucei gambiense, control of *Glossina tachinoides*, spray application of decamethrin from helicopters: Komoe valley, Upper Volta, West Africa

Insecticides

Busvine, J. R., 1977, Linn. Soc. Symposium Ser. (5), 145-150
 human African and South American trypanosomiasis, currently used vector control measures

Insecticides

Lee, C. W.; et al., 1978, PANS, v. 24 (4), 404-422
 [Trypanosoma brucei gambiense], control of *Glossina tachinoides* by application of insecticides from helicopters, calibration of equipment and insecticide dispersal: Upper Volta, West Africa

Insecticides

Leveque, C.; Odei, M.; and Pugh Thomas, M., 1977, Linn. Soc. Symposium Ser. (5), 133-143
 onchocerciasis, description of World Health Organization's monitoring program to evaluate effect of abate on non-target organisms: Volta River Basin

Insecticides

Lodha, K. R., 1974, Insecticides and their usage in the control of livestock pests, 104 pp.

Insecticides

Molyneux, D. H.; et al., 1978, PANS, v. 24 (4), 391-403
Trypanosoma brucei gambiense, control of *Glossina tachinoides* with insecticides, first of series of papers (objectives, experimental area, and insecticides): River Komoe valley, West Africa

Insecticides

Service, M. W., 1977, Linn. Soc. Symposium Ser. (5), 151-164
 human malaria, review of current control measures, problems in vector control

Insecticides

Sherlock, I. A.; and Muniz, T. M., 1975, Rev. Brasil. Malariol., v. 26-27, 1974-1975, 93-103
 control of *Panstrongylus megistus* vectors of human Chagas disease, using BHC: S. Felipe, State of Bahia, Brazil

Insecticides

van Wettere, P.; et al., 1978, PANS, v. 24 (4), 435-446
 [Trypanosoma brucei gambiense], control of *Glossina tachinoides*, evaluation of insecticides applied as aerosols from helicopters: Komoe valley, Upper Volta, West Africa

Intestopan. See Brobenzoxaldine or Broxyquinoline.

Iodine -- Lugol's iodine; Lugol solution; Pervinox; Povidone-Iodine.

Lugol solution

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Iodine

Iwanczuk, I.; and Kelus, J., 1971, Roczniki Panstw. Zakl. Hig., v. 22 (2), 179-187
 tests for ovicidal activity of iodine and chlorine in swimming pools using *Aspicularis* tetraptera as model for human *Enterobius vermicularis* and *Ascaris suis* as model for *Ascaris lumbricoides*; neither chemical was an effective ovicide

Povidone-Iodine (Pervinox)

Muhafra, J., 1977, Semana Med. (4994), an. 84, v. 151 (1), 31-33
Pediculus humanus var capitis, children, clinical trials with povidone-iodine, good results: Argentina

Lugol's iodine

Sharma, S. D.; and Bhatia, Y. S., 1978, Indian Vet. Med. J., v. 2 (4), 217-218
 coenurosis, goat (near base of right horn), case report, injection of cyst with Lugol's iodine coupled with paracentesis and antibiotic therapy, complete recovery

Iodochlorhydroxyquin -- Clioquinol; Enteroseptol; Enterovioform; Iodochlorhydroxyquin; Mexaform (with Phanquone and Oxyphenonium bromide).

Enterovioform

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Mexaform

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Enteroseptol

Kogan, G. F., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 196-198
 nosematosis, bees, fumagillin had good therapeutic effect, metronidazole, sulfadimetoxin and enteroseptol showed no substantial effect

Clioquinol + Oxytetracycline

Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire

Iodochlorhydroxyquin -- Continued.

Mexaform

Rostkowska, J., 1970, Acta Parasitol. Polon., v. 18 (27-41), 377-392

Balantidium coli, effect of various bacteria on propagation in vitro, on erythrophagocytic capability of balantidia, and on susceptibility of balantidia to atebirin, entobex, mexaform, and protargol; Trichomonas hominis, Chilomastix mesnili, and Dientamoeba fragilis found to be without effect; effect of balantidia on bacteria

Iodochlorhydroxyquin (Enterovioform)

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriolog., 1 Abt. Orig., Reihe A, v. 241 (3), 358-367

Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Mexaform

Trzaska, B.; and Leinweber-Noiszewska, H., 1974, Ann. Acad. Med. Stetinensis, suppl. 10, 19-25

Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis

Iodochlorhydroxyquin. See Iodochlorhydroxyquin.

3-Iodo-3,4-dihydro-2,2-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione. See 3-Iodo- β -lapachone.

Iodofenphos

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960

Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

3-Iodo-4-hydroxy-5-nitrobenzotrile. See Nitroxynil.

3-Iodo- β -lapachone -- 3-Iodo-3,4-dihydro-2,2-dimethyl-2H-naphtho[1,2-b]pyran-5,6-dione.

3-Iodo- β -lapachone

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

3-Iodo- β -lapachone

Docampo, R.; et al., 1978, Ztschr. Parasitenk., v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

3-Iodo- β -lapachone -- Continued.Iodo- β -lapachone

Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

Iodophene. See Iodophthalein.

3-p-Iodophenylcarbonyl-4-hydroxycoumarin

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Iodophthalein -- Iodophene.

Iodophene

Petrov, Iu. F., 1978, Veterinariia, Moskva (5), 64-66

[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Ipronidazole -- Ipropran; 1-Methyl-2-isopropyl-5-nitroimidazole.

Ipronidazole (Ipropran)

Schildknecht, E. G.; and Squibb, R. L., 1979, Parasitology, v. 78 (1), 19-31

Histomonas meleagridis in turkeys, effects of vitamins A, E, and K (alone and in combination with ipronidazole) on performance and on plasma enzymes, plasma enzyme levels correlated well with progressive pathological changes

Ipropran. See Ipronidazole.

Iramin + Metacyl

Orekhova, M. M., 1974, Vet. Nauka--Proizvod., Trudy, Minsk, v. 12, 139-140

E[imeria] tenella, chicks, coccidiostatic activity of iramin in combination with metacyl

Iron hematoporphyrin D

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050

Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Iron naphthenate

Niazov, A. N.; et al., 1977, Izvest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk (6), 68-70

ixodid ticks, naphthenates tested as acaricides

Isobenzan

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Isobenzan

Gladney, W. J.; et al., 1974, J. Med. Entom., v. 11 (3), 303-306
Amblyomma maculatum, cattle (exper.), male tick pheromone applied to small area on cattle attracted female ticks from other sites, mixture of pheromone and isobenzan attracted and killed female ticks

S-Isobutyl adenosine analogues

Trager, W.; Robert-Gero, M.; and Lederer, E., 1978, FEBS Letters, v. 85 (2), 264-266
Plasmodium falciparum, antimalarial activity of S-isobutyl adenosine analogues in culture

Isohematoporphyrin IX

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Isometamidium -- Isometamidium chloride; Isometamidium chloride hydrochlorate; Isometamidium salt; Samorin; Trypamidium.Isometamidium chloride hydrochlorate (Trypamidium; Samorin)

Balis, J., 1977, Rev. Elevage et Med. Vet. Pays Trop., n.s., v. 30 (4), 373-375
 isometamidium chloride hydrochlorate, dromedary and domestic mammals, toxicity

Isometamidium chloride hydrochlorate (Trypamidium; Samorin)

Balis, J.; and Richard, D., 1977, Rev. Elevage et Med. Vet. Pays Trop., n.s., v. 30 (4), 369-372
Trypanosoma evansi, dromedary, isometamidium chloride hydrochlorate, intravenous and intramuscular injections, toxicity

Isometamidium

Bourn, D.; and Scott, M., 1978, Trop. Animal Health and Prod., v. 10 (4), 191-203
 trypanosomiasis, successful use of Zebu work oxen in agricultural development of tsetse infested land, environmental conditions, epizootiology of trypanosomiasis in oxen and in *Glossina morsitans*, strategic drug use (alternation of diminazene aceturate and isometamidium to control trypanosomes; radoxanide to control helminths): Wollega province, western Ethiopia

Isometamidium chloride (Samorin)

James, D. M., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 471-476
 [*Trypanosoma*] *congolense*, *T. brucei*, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs

Isometamidium -- Continued.Isometamidium salt

Worms, M. J.; and Hawking, F., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 548-549
Litomosoides carinii-infected *Sigmodon hispidus*, effects of suramin, homidium bromide, quinapyramine, diminazene, and isometamidium after observation period of more than 5 weeks

Isometamidium chloride. See Isometamidium.

Isometamidium chloride hydrochlorate. See Isometamidium.

Isometamidium salt. See Isometamidium.

Isopentaquine -- 8-(4-Isopropylamino-1-methylbutylamino)-6-methoxyquinoline oxalate; WR 6,020.Isopentaquine

Blandon, R.; Guevara, J. F.; and Johnson, C. M., 1976, Rev. Med. Panama, v. 1 (3), 153-162
 Chagas disease in children, diagnosis, pathology, therapeutic trials with various drugs, metronidazole was well tolerated and therefore most promising therapy: Panama

Isopentaquine (WR 6,020)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
 sporozoite-induced *Plasmodium berghei* in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

5-Isopropoxy carbonylamino-2-(4-thiazolyl) benzimidazole. See Cambendazole.

1-Isopropoxyphenyl-N-methylcarbamate. See Propoxur.

8-(4-Isopropylamino-1-methylbutylamino)-6-methoxyquinoline oxalate. See Isopentaquine.

2-(N-Isopropylaminomethyl)-6-methyl-7-nitro-1,2,3,4-tetrahydroquinoline -- U.K. 3883.

U.K. 3883

Pellegrino, J.; and Katz, N., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (1), 59-66
Schistosoma mansoni, *S. japonicum*, laboratory animals, U. K. 3883 tested, highly effective against *S. mansoni*, devoid of activity against *S. japonicum* in mice

8-(5-Isopropylaminopentylamino)-6-methoxyquinoline phosphate. See Pentaquine.

Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate. See Methoprene.

N-Isopropyl-3-(5-nitro-2-furyl)-acrylamide. See Nitrofurylacrylamide.

3-4-Isopropylphenyl-1,1-dimethylurea -- Alon.

Alon

Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

2-(4'-Isopropylstyryl)-5-nitro-1-vinylimidazole
Morton, D. M.; Fuller, D. M.; and Green, J. N., 1973, Xenobiotica, v. 3 (4), 257-266
2-styryl-5-nitroimidazoles, metabolism and excretion in laboratory animals, activity against *Trypanosoma rhodesiense*

Isopropyl 2-(4-thiazolyl)-5-benzimidazolecarbamate. See Cambendazole.

Isothiocyanate derivatives

Sevcik, B.; et al., 1974, Veterinaria, Praha, v. 16 (5-6), 421-588
Eimeria tenella, chickens, 613 substances screened as coccidiostats, extensive detailed statistical results

4-Isothiocyanate-4'-nitrodiphenyl ether. See Nitroscanate.

4-Isothiocyanato-4'-nitrodiphenylamine. See Nitroscanate.

Isothiocyanato-2-pyridinyl benzimidazoles
Haugwitz, R. D.; et al., 1979, J. Med. Chem., v. 22 (9), 1113-1118
Nematospiroides dubius, *Hymenolepis nana*, mice, synthesis and activities of 2-pyridinyl-5-isothiocyanatobenzimidazoles

4-Isothiocyano-4'-nitro-diphenylamine. See Nitroscanate.

4-Isothiocyano 4-nitro diphenyl ether. See Nitroscanate.

Ivoran. See DDT.

Ixodicides

Wilson, R. G., 1978, J. South African Vet. Ass., v. 49 (1), 49-51
Boophilus, resistance to ixodicides due to altered site of drug action, altered rate of tick metabolism, and/or altered rate of drug transportation, review

- Jodopax
Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures
- Jonit. See Bitoscanate.
- Kamala
Tani, S., 1976, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 25 (4), 262-273
Echinostoma hortense, humans (feces), treatment with kamala: Akita Prefecture, Japan
- Kamala (*Mallotus philippinensis*)
Yokogawa, M.; et al., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (1), 35-40
Metagonimus yokogawai, hamsters, anthelmintic efficacy of various fractions of kamala, crude rottlerin most effective
- Kasugamycin
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Kelfizine. See Sulfalene.
- Kelthane. See Dicofol.
- Kenalog. See Triamcinolone acetonide.
- Kerosene
El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213
Hyalomma dromedarii, Argas persicus, evaluation of 10 insecticides
- Ketrax. See Tetramisole.
- Kitnos. See Chlorphenoxamide.
- Klion. See Metronidazole.
- Klion-D
Domany, Z., 1976, Therap. Hungar., v. 24 (4), 148-151
Trichomonas vaginalis associated with candidiasis, women with vaginal infections, clinical trials with klion-D vaginal preparations: Hungary
- Korlan. See Ronnel.
- Korsolin
Kunstyr, I.; and Ammerpohl, E., 1978, Lab. Animals, v. 12 (2), 95-97
Spironucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite
- Krimnos
Mallick, K. P.; Mishra, S. S.; and Sahai, B. N., 1977, Indian J. Animal Research, v. 11 (2), 77-80
Haemonchus spp., goats (exper.), thiabendazole, pyrantel tartrate, krimnos, comparative efficacy, blood picture before and after treatment, krimnos not effective
- Kriofos
Getta, G. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 72-75
hypodermatitis, cattle, insecticides, repellents
- Kwellada. See Benzene hexachloride.

L 9053

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Lactosyl ceramide

Alving, C. R.; et al., 1979, *Science* (4411), v. 205, 1142-1144

Plasmodium berghei, mice, therapeutic effects of glycolipids in liposomes against sporozoite-induced malaria

Laevamisole. See Tetramisole.Lampit. See Nifurtimox.Lampren. See Clofazimine.Lapachol -- 2-Hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthalenedione; 2-Hydroxy-3-(3-methyl-2-butenyl)-1,4-naphthoquinone.

Lapachol

Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

Lapachol

Docampo, R.; et al., 1978, *Ztschr. Parasitenk.*, v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

Lapachol

Lopes, J. N.; et al., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on *Crithidia fasciculata*

 α -Lapachone

Boveris, A.; et al., 1977, *Rev. Asoc. Argent. Microbiol.*, v. 9 (2), 54-61

Trypanosoma cruzi, epimastigote forms in saline medium, action of β -lapachone and its metabolite form α -lapachone on parasite growth and production of hydrogen peroxide; β -lapachone inhibited growth but possible effect was mediated by the hydrogen peroxide and related free radicals

 α -Lapachone

Boveris, A.; et al., 1978, *Biochem. J.*, v. 175 (2), 431-439

Trypanosoma cruzi, differential effect of β -lapachone and α -lapachone on superoxide anion and hydrogen peroxide production and on growth of epimastigotes

 α -Lapachone

Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

 α -Lapachone

Docampo, R.; et al., 1978, *Ztschr. Parasitenk.*, v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

 α -Lapachone

Lopes, J. N.; et al., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on *Crithidia fasciculata*

 β -Lapachone

Boveris, A.; et al., 1977, *Rev. Asoc. Argent. Microbiol.*, v. 9 (2), 54-61

Trypanosoma cruzi, epimastigote forms in saline medium, action of β -lapachone and its metabolite form α -lapachone on parasite growth and production of hydrogen peroxide; β -lapachone inhibited growth but possible effect was mediated by the hydrogen peroxide and related free radicals

 β -Lapachone

Boveris, A.; et al., 1978, *Biochem. J.*, v. 175 (2), 431-439

Trypanosoma cruzi, differential effect of β -lapachone and α -lapachone on superoxide anion and hydrogen peroxide production and on growth of epimastigotes

 β -Lapachone

Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

 β -Lapachone

Cruz, F. S.; Docampo, R.; and de Souza, W., 1978, *Acta Trop.*, v. 34 (1), 35-40

Trypanosoma cruzi, β -lapachone markedly increases generation of hydrogen peroxide in intact epimastigotes

 β -Lapachone

Docampo, R.; et al., 1978, *Ztschr. Parasitenk.*, v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

 β -Lapachone

Docampo, R.; et al., 1978, *Arch. Biochem. and Biophys.*, v. 186 (2), 292-297

Trypanosoma cruzi, β -lapachone-treated epimastigotes, lipid peroxidation and generation of free radicals, superoxide anion, and hydrogen peroxide

β -Lapachone

Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531
 Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

 β -Lapachone-6-phenylhydrazone

Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531
 Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

Lapudrine. See Chlorproguanil.

Largactil. See Chlorpromazine.

Lasalocid -- Avatec; Lasalocid sodium; MW 591.Lasalocid (Avatec)

Bedrnik, P.; et al., 1979, Veterinarstvi, v. 29 (8), 353-355
 coccidiosis, broiler chicken fattening, lasalocid compared with amprol plus and monensin: Czechoslovakia

Lasalocid (Avatec)

Bedrnik, P.; et al., 1979, Arch. Geflugelk., v. 43 (1), 7-10
 Eimeria spp., chickens (exper.), comparative efficiency of anticoccidials in combination with growth promotant nitrovin

Lasalocid

Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (2), 304-307
 Eimeria spp., broilers (exper.), salinomycin, monensin, lasalocid, drug toleration and anticoccidial efficacy compared in 5 field trials, commercial facilities

Lasalocid sodium (AVATEC)

Cruthers, L. R.; et al., 1978, Poultry Science, v. 57 (5), 1227-1233
 Eimeria spp., broiler chicks (exper.), lonomycin in feed, high degree of anticoccidial activity, comparison with lasalocid and monensin

Lasalocid

Fitzgerald, P. R.; and Mansfield, M. E., 1979, J. Parasitol., v. 65 (5), 824-825
 Eimeria spp., calves (exper.), lasalocid

Lasalocid sodium (Avatec)

Foreyt, W. J.; Gates, N. L.; and Wescott, R. B., 1979, Am. J. Vet. Research, v. 40 (1), 97-100
 Eimeria spp., confinement-reared lambs (exper.) from weaning to market weight, monensin and lasalocid highly effective in eliminating oocysts, weight gains and feed conversion rates measured

Lasalocid -- Continued.Lasalocid (Avatec)

Kaemmerer, K.; and Fink, J., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (6), 216-220
 monensin-natrium, lasalocid, salinomycin, influence of high dosages on heart of chickens

Lasalocid

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
 Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Lasalocid

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
 Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Lasalocid (Avatec)

McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
 Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Lasalocid

Meingassner, J. G.; et al., 1979, Poultry Science, v. 58 (2), 308-313
 Eimeria tenella, broilers (exper.), monensin, lasalocid, anticoccidial activity enhanced when combined with tiamulin, tiamulin alone was inactive; metabolic fate of monensin and tiamulin separately or in combination monitored using isolated, perfused rat liver

Lasalocid

Migaki, T. T.; Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (5), 1192-1196
 Eimeria spp., chicks (exper.), salinomycin, monensin, lasalocid, efficacy in battery trials

Lasalocid sodium (Avatec)

Mitrovic, M.; Schildknecht, E. G.; and Marusich, W. L., 1979, Poultry Science, v. 58 (5), 1154-1159
 Eimeria spp., turkey poults (exper.), single and mixed infections, lasalocid sodium, controlled battery trials

Lasalocid

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
 Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Lasalocid

Ruff, M. D., 1978, Proc. 1978 Maryland Nutrition Conf. Feed Mfr. (University of Maryland, Mar. 16-17), 32-37
 Eimeria spp., chickens, anticoccidials, safe withdrawal times, effect on nutrient malabsorption

Lasalocid -- Continued.

Lasalocid (MW 591)

Smith, C. K. II; and Strout, R. G., 1979, *Exper. Parasitol.*, v. 48 (3), 325-330
Eimeria tenella, accumulation and retention of lasalocid and narasin by extracellular sporozoites

Lasalocid sodium. See Lasalocid.

Lauryl dimethyl benzyl ammonium saccharinate
 Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Lauryl isoquinolium saccharinate

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Lead arsenate

Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10) 28-36

cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Lebaycid. See Fenthion.

Lederkyn. See Sulfamethoxypridazine.

Ledermycin. See Demeclocycline.

Lerbek. See Methyl benzoate or Meticlorpindol.

Levamisole. See Tetramisole.

Levamisole chlorhydrate. See Tetramisole.

Levamisole hydrochloride. See Tetramisole.

Levamisole phosphate. See Tetramisole.

Levasole. See Tetramisole.

Levasole Injection. See Tetramisole.

Levasole tablets. See Tetramisole.

Levofuraltadone. See Furaltadone.

Levo-furaltadone hydrochloride. See Furaltadone.

Levomycetin

Gobzem, V. R.; and Nazarov, V. G., 1978, *Veterinariia*, Moskva (3), 67-69

Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Levo-tetramisole. See Tetramisole.

Lime-sulfur solution (Orthorix Spray)

Fletcher, K. C., 1978, *J. Am. Vet. Med. Ass.*, v. 173 (9), 1231-1232

Notoedres cati in *Uncia uncia* (skin), intense pruritus with resulting alopecia, lime-sulfur solution and sulfur ointment healed all lesions without complications: zoo

Lime-sulphur solution

Munro, R.; and Munro, H. M. C., 1979, *Austral. Vet. J.*, v. 55 (2), 90 [Letter]

Lynxacarus [sp.], cats, lesions, successful treatment with lime-sulphur solution: Fiji

Lincomycin + Spectinomycin (= Lincospectin)

Bischofova, N.; and Zajicek, D., 1976, *Veterinarstvi*, v. 26 (11), 506-507

Balantidium coli, pigs, large-scale fattening farms, Lincospectin treatment

Lincomycin

Maxey, B. W.; and Page, R. K., 1977, *Poultry Science*, v. 56 (6), 1909-1913

Eimeria brunetti, *E. maxima*, chickens, lincomycin feed medication for control of necrotic enteritis

Lincospectin. See Lincomycin or Spectinomycin.

Lindane. See Benzene hexachloride.

Lindatox-20. See Benzene hexachloride.

Linton-X. See Toxaphene.

Liquamycin. See Oxytetracycline.

Liquamycin injectable L-50. See Oxytetracycline.

Liquamycin/LA-200. See Oxytetracycline.

Liquamycin (T-50). See Oxytetracycline.

Liquor arsenicalis. See Potassium arsenite solution.

Lithium antimony thiomalate. See Anthiolimine.

Lomidine. See Pentamidine.

Lonomycin -- DE-3936; Emericid; 31,559 R.P.;
SQ 12,525; TM-481.

Lonomycin (TM-481; SQ 12,525)
Cruthers, L. R.; et al., 1978, Poultry Science, v. 57 (5), 1227-1233
Eimeria spp., broiler chicks (exper.), lonomycin in feed, high degree of anticoccidial activity, comparison with lasalocid and monensin

Lopatol. See Nitroscanate.

Lorsban. See Chlorpyrifos.

Loxon. See Haloxon.

Lucanthone -- Miracil-D; Nilodin; Tixantone.

Tixantone (Lucanthone)
Abdulla, W. A.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 109-115
S[chistosoma] mansoni, prophylactic activity, antischistosomal drugs, albino mice, most effective within 2 weeks post exposure

Lucanthone
Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, J. Parasitol., v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy

Lucanthone
Foster, R., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 1-9
Schistosoma mansoni, S. haematobium, S. japonicum, oxamniquine, efficacy and toxicity in various exper. animals, clinical trials in humans

Lucanthone (Miracil D)
Ghazal, A.; Ismail, M.; and Sharabi, F. M., 1978, Egypt. J. Pharm. Sc., v. 17 (1), 1976, 43-52
Schistosoma mansoni-infected mice, tissue histamine content before and after treatment with several antischistosomal drugs

Nilodin
Laba, L., 1974, Ann. Acad. Med. Stetinensis, suppl. 10, 33-38
schistosomiasis, case reports of infection in African students living in Poland, unsuccessful therapy with nilodin in one case with cure after use of ambilhar

Lucanthone
Ong, T. M., 1978, Mutation Research, v. 55 (1), 43-70
hycanthone and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Lucanthone -- Continued.

Lucanthone
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Miracil D
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Lucensomycin
Carli, S.; et al., 1979, Riv. Zootecn. e Vet. (1), 7-9
[asciola] hepatica, in vitro, lucensomycin, fluke motility, effect of drug reduced by steroid compounds, not affected by alpha-mercaptopropionylglycine

Lugol's iodine. See Iodine.

Lugol solution. See Iodine.

Lypor 20. See Temephos.

D-Lysine
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

L-Lysine
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

- Lysol
Chauhan, P. P. S.; et al., 1978, Indian Vet. J., v. 55 (12), 1011-1013
Neoscaris vitulorum eggs, action of boiling water, direct sunlight, and lysol on viability, tested by infectivity to albino rats
- Lysol
Slepnev, N. K.; and Zen'kov, A. V., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 135-137
E[chinococcus] granulosus, protoscolices, destructive action of high and low temperatures; lysol and creolin most destructive of chemicals tested
- Lysozyme + Nifuratel + Nystatin (= Macmiror plus)
Martins, A. D., 1974, Rev. Brasil. Med., v. 31 (7), 503-508
human vaginal trichomoniasis alone or in the presence of fungal infections, successful therapy combining oral nifuratel and vaginal treatment with nifuratel combined with nystatin and lysozyme

Macmiror. See Nifuratel.

Macmiror plus. See Lysozyme or Nifuratel or Ny-statin.

Madribon. See Sulfadimethoxine.

Magmilor. See Nifuratel.

Magnesium sulphate

Mello, E. B. F.; et al., 1978, Zentralbl. Bakteriologie, 1. Abt. Orig., Reihe A, v. 241 (3), 384-387

—*Taenia solium*, *T. saginata*, humans, oral treatment with aspidium preceded by hypertonic magnesium sulphate solution

Magnesium sulphate

Venkateswara Rao, P.; et al., 1977, Riv. Parassitol., Roma, v. 38 (1), 13-21

Prosthogonimus sp. cercariae, cercaricidal effect of certain common fertilizers, ammonium sulphate may be cercaricide of choice

Malachite green-oxalate

Gilbert, J. P.; Gratzek, J. B.; and Brown, J., 1979, J. Fish Dis., v. 2 (3), 191-196

formalin and malachite green-oxalate (alone or in combination) used as model system for testing synergistic action of parasiticides in vitro

Malachite green

Iashchuk, V. D., 1977, Veterinarika, Moskva (9), 71-73

Costia necatrix, rainbow trout, severe outbreak, economic benefits of treatment with malachite green: Vinnitsk fish farm

Malalaquin. See Chloroquine.

Malathion -- Carbophos; Cythion; O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate; Malathion 50 EC; Malathion 57; Malathion aerosol solution; Malathion powder; Prioderm.

Malathion (Prioderm)

Blommers, L., 1978, Acta Leidensia, v. 46, 7-8

Pediculus capitis, humans and on human hair in vitro, malathion shampoo

Malathion

Blommers, L., 1979, J. Med. Entom., v. 16 (1), 82-83

Pediculus capitis, nymph rearing technique, insecticide tests against nymphs

Malathion (Prioderm)

Blommers, L.; and van Lennep, M., 1978, Acta Leidensia, v. 46, 9-15

Pediculus capitis, susceptibility of 1st instar nymphs to malathion, propoxur, and plifemate, residual action of 3 compounds applied as lotions to human hair

Malathion -- Continued.

Malathion

Blommers, L.; and van Lennep, M., 1978, Entom. Exper. et Applic., v. 23 (3), 243-251

Pediculus humanus capitis, school children, field trials with lindane, laboratory tests with lindane, dieldrin, and malathion, presence of resistance to lindane confirmed: Netherlands

Malathion

Carsons, J. K., 1977, Malaysian Vet. J., v. 6 (3), 111-124

Boophilus microplus, dairy cattle, various control measures discussed but spraying acaricides on pastures shows particular promise: Air Hitam, Johor, Malaysia

Malathion 50 EC

Chellappa, D. J.; Subramanian, R.; and Gopala-krishnan, C. A., 1977, Indian Poultry Gazz., v. 61 (4), 135-137

Menopon gallinae, *Lipeurus caponis*, poultry, malathion 50 EC, sumithion 50 EC, nuvan 100 EC, drug trial, sumithion 50 EC most effective, knapsack sprayer superior to conventional dipping procedure, preliminary report

Malathion

Cole, M. M.; et al., 1973, J. Econom. Entom., v. 66 (1), 118-119

Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

Malathion

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Malathion

El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213

Hyalomma dromedarii, *Argas persicus*, evaluation of 10 insecticides

Malathion

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886

laboratory-reared *Haematobia irritans*, susceptibility to topically applied insecticides

Malathion

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586

Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Malathion

Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, J. Econom. Entom., v. 71 (2), 315-318

Ornithonyssus sylvaticum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban permethrin and SD-43775 also effective; mites displayed tolerance to malathion

Malathion -- Continued.

Malathion

Khan, D.; and Haseeb, M. A., 1976, Pakistan J. Zool., v. 8 (2), 173-176

Ganeo micracetabulus and Cercaria reflexicauda cercariae, effects of 5 insecticides at various concentrations, toxicity varies but results indicate cercariae are susceptible to insecticides

Malathion

Khan, M. H., 1979, Indian Vet. J., v. 56 (9), 739-743

Lipeurus caponis, Menacanthus stramineus, White Leghorn fowl, organophosphorus insecticides, costs evaluated

Malathion

Kiefer, M.; and Cyprich, D., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Zool. (23), 5-12
ticks, susceptibility to acaricides: Slovakia

Malathion

Kumar, A.; et al., 1977, Indian J. Poultry Sc., v. 12 (3), 48-50

Eomenacanthus stramineus, chicks (exper.), sumithion, malathion, dipterex, comparative efficacy, haematology and blood glucose levels

Malathion 57

McKeever, P. J.; and Allen, S. K., 1979, J. Am. Vet. Med. Ass., v. 174 (7), 718-720

Cheyletiella spp., cats, pruritic dermatitis, pyrethrins, malathion 57

Malathion (Cythion)

Meleney, W. P.; and Roberts, I. H., 1979, J. Med. Entom., v. 16 (1), 52-58

Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Malathion powder

Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15

Myobia musculi, Myocoptes musculinus, conventional mouse colony, acaricides

Malathion

Preston, S.; and Fry, L., 1977, Roy. Soc. Health J., v. 97 (6), 291

head lice in children, clinical trials testing the effectiveness of malathion cream shampoo versus malathion lotion, both forms equally effective

Carbophos

Puchkova, E. A., 1977, Veterinariia, Moskva (7), 19-22

D[ermanyssus] gallinae, C[imex] lectularius, lice, chickens on industrial scale farms, control, sevin, dicresil, chlorophos, carbophos; other complex sanitation measures

Carbophos

Sverba, V. A.; and Shemchuk, V. R., 1978, Veterinariia, Moskva (10), 69-71

Sinergasilus major, white amur, copper sulfate and ferrous sulfate mixture, chlorophos, carbophos, formula for estimating concentrations in relation to temperature and other factors in aquaria or ponds

Malathion -- Continued.

Malathion aerosol solution

Tierney, F.; and Baillie, J., 1979, Vet. Med. and Small Animal Clin., v. 74 (1), 69-70
Cnemidocoptes pilae, Mexican red-headed parrot (beak, face), case report, combined aerosol (malathion solution) and topical treatment (eurax and Goodwinol cream) highly effective and less stressful

Malathion

Uspenskii, I. V., 1974, Parazitologiya, Leningrad, v. 8 (4), 312-321

Ixodes persulcatus, susceptibility to acaricides

Malathion

Wright, F. C.; and Riner, J. C., 1979, South-west. Entom., v. 4 (1), 40-45

Psoroptes ovis, P. cuniculi, 10 acaricides evaluated using 'tea-bag' technique

Malathion

Young, E.; Zumpt, F.; and Whyte, I. J., 1972, J. South African Vet. Ass., v. 43 (2), 226

Sarcoptes scabiei in Panthera leo, skin lesions, successful treatment with BHC and/or malathion-containing preparations: Kruger National Park

Malathion 57. See Malathion.

Malathion aerosol solution. See Malathion.

Malathion 50 EC. See Malathion.

Malathion powder. See Malathion.

Male fern extract. See Aspidium.

1-Maleinyl-4-(3'-chloro-4'-methyl-phenyl) piperazine. See Maleinyl-4-(3-chloro-p-tolyl) piperazide.

Maleinyl-4-(3-chloro-p-tolyl) piperazide --

Hoechst S-688; 1-Maleinyl-4-(3'-chloro-4'-methyl-phenyl) piperazine.

1-Maleinyl-4-(3'-chloro-4'-methyl-phenyl) piperazine (Hoechst S-688)

Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, Rev. Brasil. Pesqui. Med. e Biol., v. 2 (1), 45-49

Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

Hoechst S-688

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46

Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Mallotus philippinensis. See Kamala.

Maloprim. See Dapsone or Pyrimethamine.

Malva aegyptica herb

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118
Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

Manganese ethylene bisdithiocarbamate

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Mansil. See Oxamniquine.

Mansonil. See Niclosamide.

Mapharsemin. See Oxophenarsine.

Maretin. See Phthalophos.

Marvex Super-100. See Dichlorvos.

Mascyl. See Carbon disulfide or Piperazine.

Mebenav. See Mebendazole.

Mebendazole -- (5-Benzoyl-1H-benzimidazole-2-yl)-carbamic acid; 2-Carbamomethoxy-5-benzoyl-benzimidazole; Equivurm Plus; Mebenav; Mebenvet; Mebenvet Granulatum; Mebutar; Menzole; Methyl-5(6)-benzoyl-2-benzimidazole; Methyl-5-benzoyl-2-benzimidazole carbamate; Methyl (5-benzoyl-1-H-benzimidazol-2-yl) carbamate; Methyl-N-[5(6)-benzoyl-2-benzimidazolyl] carbamate; Multispec; Nemasole; Pantelmin; R-17,635; Telmin; Telmin RLT Sheep Drench; Vermox.

Mebendazole

Akusawa, M.; and Deguchi, N., 1975, *Kiseichugaku Zasshi (Japan. J. Parasitol.)*, v. 24 (6), 357-361
Trichuris vulpis, dogs, mebendazole effective with no adverse effects

Mebendazole

Albert, H.; and Hoerchner, F., 1979, *Berl. u. Munchen. Tierarztl. Wchnschr.*, v. 92 (10), 189-193
Taenia saginata, calves (exper.), serum antibodies, enzyme-linked immunosorbent assay, titres following reinfection and drug therapy

Mebendazole -- Continued.

Mebendazole (Mebutar)

Alonso, R. E., 1976, *Semana Med.* (4956), an. 83, v. 149 (7), 234-237
intestinal parasites, school children, clinical studies with tinidazole and mebendazole: Institute Aguirre 'Patronato de la Infancia'

Mebendazole (Vermox)

Alton, K. B.; Patrick, J. E.; and McGuire, J. L., 1979, *J. Pharm. Sc.*, v. 68 (7), 880-882
mebendazole, high-performance liquid chromatographic assay, tested on human plasma with known drug amounts added; possible use with Echinococcus multilocularis patients receiving chronic high dosage

Mebendazole

Amato Neto, V., 1972, *Rev. Brasil. Clin. e Terap.*, v. 1 (4), 237-238
mebendazole, new anthelmintic with wide scope of activity, review

Mebendazole

Amato Neto, V., 1976, *Rev. Brasil. Clin. e Terap.*, v. 5 (7), 269-272
ascariasis, trichocephaliosis, adults and children, clinical trials using single dose of mebendazole

Mebendazole

Amato Neto, V.; et al., 1973, *Rev. Inst. Med. Trop. S. Paulo*, v. 15 (1), 34-37
hookworm, enterobiasis, ascariasis, trichuriasis, children, mebendazole

Mebendazole (Vermox)

Ammann, R.; Akovbiantz, A.; and Eckert, J., 1979, *Schweiz. Med. Wchnschr.*, v. 109 (5), 148-151
Echinococcus granulosus, humans (lungs, liver), mebendazole

Mebendazole (Mebenvet)

Ashton, D. G., 1979, *Vet. Rec.*, v. 105 (5), 109
mebendazole toxicity in parakeets

Mebendazole

Barsanti, J. A.; et al., 1979, *Cornell Vet.*, v. 69 (1), 45-53
Mesocoestoides spp. causing peritonitis in a dog (peritoneal cavity, vaginal tunic of testicle), mebendazole

Mebendazole

Bennet, E. M.; Behm, C.; and Bryant, C., 1978, *Internat. J. Parasitol.*, v. 8 (6), 463-466
Mesocoestoides corti, mice (infected, injected with dead larvae previous to infection, or irradiated), effects of mebendazole and levamisole alone or together on tetrathyridia, concluded that anthelmintic efficacy of mebendazole depends on its anthelmintic activity supplemented by host's immune response and that levamisole stimulates the latter

Mebendazole -- Continued.

Mebendazole

Bernberg, H. C.; Clarke, V. de V.; and Gelfand, M., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (2), 233-234

Dipetalonema perstans-like unsheathed microfilarial infection, humans, combined treatment with levamisole and mebendazole: Rhodesia

Mebendazole

Bina, J. C.; et al., 1977, Rev. Inst. Med.

Trop. S. Paulo, v. 19 (1), 47-51

intestinal helminths, closed rural population, evaluation of mass treatment with mebendazole

Mebendazole

Blair, L. S.; and Campbell, W. C., [1979], J.

Parasitol., v. 64 (6), 1978, 1032-1034

Dirofilaria immitis, pre-cardiac stages in Mustela putorius furo, trials of avermectin B.a, mebendazole, and melarsoprol, possible value of Dirofilaria-Mustela model for chemotherapeutic studies

Mebendazole

Boray, J. C.; et al., 1979, Austral. Vet. J.,

v. 55 (2), 45-53

nematodes and cestodes of dogs and cats, efficiency and safety of nitroscanate, comparison with mebendazole, bunamidine hydrochloride, and praziquantel

Mebendazole (Nemasole; R-17635)

Borda, C. E.; et al., 1978, Bol. Chileno Parasitol., v. 33 (3-4), 57-61

Ancylostoma duodenale, humans, mebendazole, useful in individual and mass control: Argentina

Mebendazole

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15

antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Mebendazole

Boyd, W. P., jr.; Campbell, F. W.; and Trudeau, W. L., 1978, Am. J. Trop. Med. and Hyg., v. 27

(1, pt. 1), 39-41

Strongyloides stercoralis, severe hyperinfection in patient with renal failure, case report, successful therapy with mebendazole

Mebendazole (Telmin)

Bueno, L.; Ruckebusch, Y.; and Dorchie, Ph., 1979, Vet. Parasitol., v. 5 (2-3), 253-260

strongyle-infected ponies (nat. and exper.), disturbances of digestive motility, effect of mebendazole treatment

Mebendazole

Cavier, R.; and Notteghem, M. J., 1979, Ann.

Pharm. Franc., v. 37 (1-2), 33-36

Hymenolepis nana var. fraterna, mice, comparative study of mebendazole and flubendazole

Mebendazole -- Continued.

Mebendazole

Chaia, G.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (4), 239-247

Trichuris trichiura and Necator americanus in humans, Ancylostoma caninum and T. vulpis in dogs, mebendazole: Brazil

Mebendazole (Telmin RLT Sheep Drench)

Chevis, R. A. F.; and Kelly, J. D., 1978, N.

Zealand Vet. J., v. 26 (9), 218-220

Fasciola hepatica, rats and sheep (both exper.), mebendazole, parbendazole, cambendazole, thiabendazole, anthelmintic activity, molecular structure-activity analyses

Mebendazole

Chiari, L.; Araujo, S. M.; and Guerrero, J.,

[1978], Riv. Parasitol., Roma, v. 38 (2-3),

1977, 329-332

helminths in animal tissue, technique for collecting or counting using compression between plastic, useful in anthelmintic evaluation, mebendazole tested

Mebendazole

Chongsuphajaisiddhi, T.; et al., 1978, Ann.

Trop. Med. and Parasitol., v. 72 (1), 59-63

soil-transmitted nematodes of children, mebendazole confirmed as effective and safe therapy, clinical trials in Thailand

Mebendazole (Equivurm Plus; Telmin)

Clayton, H. M.; and Neave, R. M. S., 1979,

Vet. Rec., v. 104 (25), 571-572

Dictyocaulus arnfieldi, donkeys (feces), mebendazole, controlled trial

Mebendazole

Coles, G. C.; and Briscoe, M. G., 1978, Vet.

Rec., v. 103 (16), 360-361 [Letter]

Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for Haemonchus contortus eggs

Mebendazole

Daniyarov, I. A.; et al., 1978, Veterinariia,

Moskva (2), 64-65

Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Mebendazole

Denham, D. A.; Suswillo, R. R.; and Rogers, R.,

1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72

(5), 546-547

Brugia pahangi, mebendazole, anthelmintic activity in vitro and in infected Aedes aegypti, Meriones unguiculatus, and cats

Mebendazole (R 17,635)

Devay, O., 1974, Rev. Patol. Trop., v. 3 (1).

43-49

intestinal helminths, children, clinical trials testing efficacy of mebendazole: Periperi, Brazil

Mebendazole (Telmin)

Drudge, J. H.; Lyons, E. T.; and Tolliver, S.

C., 1979, Am. J. Vet. Research, v. 40 (4), 590-594

large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Mebendazole -- Continued.

Mebenvet

- Dvorakova, L.; and Rachac, L., 1978, Veterinar-stvi, v. 28 (9), 398-399
Echinuria uncinata, domestic ducks, epizootic occurrence, prevalence, necrosis of proventriculus, nilverm and mebendazole, preventive measures: South Bohemia region

Mebendazole (Vermox)

- Eckert, J.; Barandun, G.; and Pohlenz, J., 1978, Schweiz. Med. Wchnschr., v. 108 (29), 1104-1112
Echinococcus spp., larval stages, laboratory animals, fenbendazole, mebendazole

Mebendazole (Vermox)

- Eckert, J.; and Wissler, K., 1978, Therap. Umschau, v. 35 (9), 766-776
echinococcosis, life cycle, current immunodiagnostic methods reviewed; exper. studies with vermax in rodents, compared with previous studies in man

Mebendazol

- Enigk, K.; Dey-Hazra, A.; and Batke, J., 1975, Acta Parasitol. Polon., v. 23 (26-40), 367-372
helminths of geese, treatment (Amidostomum anseris with mebendazol, fenbendazol, levamisol, and pyrantel tartrat; Trichostrongylus tenuis with mebendazol and fenbendazol; Drepanidotaenia lanceolata and Hymenolepis setigera with mebendazol)

Mebendazole (Telmin)

- Evans, W. S.; Gray, B.; and Novak, M., 1979, J. Parasitol., v. 65 (1), 31-34
Hymenolepis spp., effect of pure mebendazole and Telmin on developing larvae in Tribolium confusum

Mebendazole

- Fain, A.; et al., 1977, Ann. Soc. Belge Med. Trop., v. 57 (3), 137-142
Raillietina celebensis, human infant, case report, mixed infection with Trichuris trichiura, complete recovery after treatment with mebendazole: Papeete, Tahiti, emigrated to Belgium

Mebendazole

- Friedman, P. A.; and Platzer, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Mebendazole

- Gallie, G. J.; and Sewell, M. M. H., 1977, Trop. Animal Health and Prod., v. 9 (1), 24
Taenia saginata, calves (exper.), treatment with intraperitoneal injection of mebendazole, not deleterious to either young or fully developed cysticerci

Mebendazole (Vermox)

- Gamble, W. G.; et al., 1979, J. Am. Med. Ass., v. 241 (9), 904-907
Echinococcus multilocularis, human (liver), mebendazole, first case acquired in contiguous United States: Lyon County, Minnesota

Mebendazole -- Continued.

Mebendazole

- Gemmell, M. A.; Johnstone, P. D.; and Oudemans, G., 1978, Research Vet. Sc., v. 25 (1), 107-108
Echinococcus granulosus, Taenia hydatigena, dogs, controlled trial with mebendazole incorporated in prepared food vs. as tablets given with this food, neither treatment fully effective

Mebendazole

- Girardi, C.; et al., 1977, Ann. Fac. Med. Vet. Torino, v. 24, 82-113
nematodes, cestodes, mebendazole, controlled test, naturally infected dogs or cats; toxicity, absorption and elimination of mebendazole, non-infected dogs, cats and guinea pigs

Mebendazole (Vermox; R17,635)

- Goldsmid, J. M., 1974, South African Med. J., v. 48 (54), 2265-2266
intestinal helminths, man, baboons, mebendazole, clinical trials, highly effective broad-spectrum drug: Central Africa

Mebendazole (Vermox)

- Goldsmid, J. M.; and Rogers, S., 1976, South African Med. J., v. 50 (29), 1129-1132
Wuchereria bancrofti, Dipetalonema perstans, humans, various recovery techniques available for laboratory diagnosis, laboratory regimen for routine investigations suggested; preliminary observations on treatment of D. perstans with mebendazole: Rhodesia

Mebendazole (Telmin)

- Guerrero, J.; and Sharp, M. L., 1979, Equine Pract., v. 1 (2), 53-55
nematodes, horses, mebendazole suspension, critical evaluation

Mebendazole

- Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367
Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Mebendazole

- Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, Research Vet. Sc., v. 25 (3), 360-363
Haemonchus contortus, Trichostrongylus colubriformis, levels of benzimidazole resistance recorded from an egg hatch test procedure

Mebendazole

- Heath, D. D., 1978, Vet. Parasitol., v. 4 (1), 11-19
Taenia hydatigena, neonatal lambs, subcutaneous injection of viable eggs induced 100% protection against development of viable larvae from oral challenge but no protection against simultaneous infection with eggs of Taenia ovis and Echinococcus granulosus, maternally derived immunity was not enhanced by hyperimmunization of ewe but did not interfere with development of protection in immunized lambs, immunizing lesion regressed rapidly after treatment of lambs with mebendazole

Mebendazole -- Continued.

Mebendazole

Heath, D. D.; and Lawrence, S. B., 1978, N. Zealand Vet. J., v. 26 (1-2), 11-15
Echinococcus granulosus, *Taenia hydatigena*, *T. ovis*, sheep, effect of extended oral dosing regime with mebendazole compared with one parenteral injection of either mebendazole or praziquantel

Mebendazole

Heath, D. D.; and Lawrence, S. B., 1979, Internat. J. Parasitol., v. 9 (1), 73-76
Taenia crassiceps, rats, mebendazole, single large oral treatment markedly more effective in killing cysts than same amount of drug divided into daily smaller doses, levamisole promoted vigorous host cellular response but did not enhance action of mebendazole

Mebendazole (Telmin)

Hoerchner, F.; and Albert, H., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (6), 107-111
Taenia saginata, calves inoculated with eggs, response to reinfection and/or drug therapy (mebendazole, praziquantel)

Mebendazole (Telmin)

Holt, P. E.; Cooper, J. E.; and Needham, J. R., 1979, J. Small Animal Practice, v. 20 (5), 269-286
Angusticaecum spp., *Tachygonetria* sp., and *Atractis dactyluris* in Testudo, treatment: Britain

Mebendazole (R17.635)

Huggins, D., 1975, Rev. Soc. Brasil. Med. Trop., v. 9 (3), 115-119
 intestinal parasitism, children, clinical trials with mebendazole suspension, results varied from 100% elimination of *Ascaris lumbricoides* to no effect on *Strongyloides stercoralis*

Mebendazole

Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
 relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiroides dubius* in mice

Mebenvet

Kadyrov, N. T., 1978, Veterinariia, Moskva (7), 57-58
Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Mebendazole (Vermox)

Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167
Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria

Mebendazole -- Continued.

Mebendazole (Vermox)

Kan, S. K. P., 1979, Ann. Trop. Med. and Parasitol., v. 73 (4), 397-398
 atypical nematode ova, probably *Ascaris*, 19-year-old girl (feces) after treatment with pyrantel pamoate; mebendazole therapy resulted in ova-free feces: Malaysia

Mebendazole

Katz, N.; and Zicker, F., 1973, Rev. Soc. Brasil. Med. Trop., v. 7 (4), 225-229
Taenia spp., humans, clinical trials with mebendazole

Mebendazole (Mebenvet)

Kavai, A.; et al., 1977, Magy. Allat. Lapja, v. 99, v. 32 (8), 533-535
 nematodes, foals, efficacy of mebendazole, no noticeable difference between mebendazole pulvis and mebenvet 10% granulate

Mebendazole

Kern, P.; Dietrich, M.; and Volkmer, K. J., 1979, Tropenmed. u. Parasitol., v. 30 (1), 65-72
Echinococcus alveolaris, *E. granulosus*, humans, clinical trials with mebendazole, allergic reactions in 2 of 7 treated probably due to spillage of antigenic material, no other serious side effects

Mebendazole

Keystone, J. S.; and Murdoch, J. K., 1979, Ann. Int. Med., v. 91 (4), 582-586
 mebendazole, mode of action, pharmacokinetics, and clinical efficacy of approved and nonapproved uses, review over past 5 years

Mebendazole

Koehler, P.; Bryant, C.; and Behm, C. A., 1978, Internat. J. Parasitol., v. 8 (5), 399-404
Fasciola hepatica, ATP synthesis in succinate decarboxylase system from mitochondria, inhibition in vitro by mebendazole and a soluble derivative of cambendazole

Mebendazole

Krotov, A. I.; Cherniaeva, A. I.; and Budanova, I. S., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 165-168
Alveococcus multilocularis, white mice, effect of thiabendazole, sarcolysin acridine, levamisole, and mebendazole on larval cyst development

Mebendazole

Kumar, V.; et al., 1978, Acta Zool. et Path. Antverpiensia (70), 221-225
 helminthiasis in wild mammals and birds of zoological gardens, need for control through preventive and chemotherapeutic measures; summary of observations on use of mebendazole in mammals and birds and of methyridine in pheasants: Antwerp Zoo

Mebendazole (R 17635)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Mebendazole -- Continued.

- Mebendazole (R 17,635; Vermox)
Lecomte-Ramioul, S., 1975, Rev. Med. Liege, v. 30 (21), 733-736
Enterobius vermicularis, therapeutic trials with mebendazole, establishment of therapeutic scheme suitable for mass therapy in a children's home
- Mebendazole (Vermox)
Lengyel, A.; Albi, I.; and Rovo, J., 1978, Therap. Hungar., v. 26 (2), 64-66
Enterobius vermicularis, boys in primary school, single low dose mebendazole: Children's Town, Fot
- Mebendazole
Lengyel, A.; Levai, J.; and Rovo, J. T., 1975, Therap. Hungar., v. 23 (2), 60-63
Trichuris trichiura, children from both local and tropical areas, mebendazole: Hungary
- Mebendazole
Lensink, B. M.; Rijpstra, A. C.; and Erken, A. H. M., 1979, Zool. Garten, n. F., v. 49 (2), 121-126
Ollulanus tricuspis in Panthera tigris tigris (vomitus, stomach), clinical symptoms in mother and offspring, treatment with various anthelmintics, complete recovery achieved with levamisole: Artis-Zoo, Amsterdam
- Mebendazole
McCracken, R. O., 1978, J. Parasitol., v. 64 (2), 214-219
Trichinella spiralis, mice, mebendazole and albendazole more effective against immature than mature worms, despite this decline in drug sensitivity it is postulated that these benzimidazoles have potential therapeutic value
- Mebendazole (Telmin)
Mason, P. C.; Hodgkinson, N. L.; and McAllum, H. J. F., 1978, N. Zealand Vet. J., v. 26 (5), 131-132
Acuaria skrjabini, mebendazole, caged Uraeginthus bengalus: aviary, Waldronville, Otago, New Zealand
- Mebendazole (Equivurm Plus)
Melbourne, C. P., 1978, J. Zoo Animal Med., v. 9 (4), 146-148
helminths, zebras, mebendazole: Longleat Safari Park, Great Britain
- Mebendazole
Meltzer, R. S.; et al., 1979, Am. J. Med. Sc., v. 277 (1), 91-98
Strongyloides stercoralis, woman, development of hyperinfection syndrome while on high-dose corticosteroids and following splenectomy, central nervous system involvement, antemortem diagnosis, thiabendazole, levamisole, and mebendazole therapy: Memorial Sloan-Kettering Cancer Center, New York (had traveled in Italy and Sicily)

Mebendazole -- Continued.

- Mebendazole (Vermox)
Merdivenci, A.; Sengul, M.; and Baydemir, M., 1976, Turk Hijiy. ve Deneysel Biyol. Dergisi, v. 36 (2), 224-237
Ascaris lumbricoides, Enterobius vermicularis, children, thiabendazole and mebendazole compared, both successful
- Mebendazole
Meuldermans, W. E. G.; et al., 1976, European J. Drug Metab. and Pharmacokinet., v. 1 (1), 35-40
mebendazole, in vitro metabolism by pig, rat, and dog liver fractions
- Mebendazole (Vermox)
Migasena, S.; Suntharasamai, P.; and Harinasuta, T., 1978, Ann. Trop. Med. and Parasitol., v. 72 (2), 199-200
Necator americanus, humans, treatment trials comparing efficacy of mebendazole, tetrachlorethylene and pyrantel pamoate: Bangkok
- Mebendazole
Muangmanee, L.; Aswapokee, N.; and Vanasin, B., 1977, Siriraj Hosp. Gaz., v. 29 (4), 439-445
Capillaria philippinensis, man, intestinal infection with associated malabsorption syndrome, case report, comparison with histories of 3 earlier cases, cured with mebendazole: Surin province, Thailand
- Mebendazole
Murray-Lyon, I. M.; and Reynolds, K. W., 1979, Brit. Med. J. (6198), v. 2, 1111-1112
hydatid cysts, humans, fever following mebendazole treatment possibly a reaction to drug-induced tissue necrosis in cysts
- Mebendazole + Trichlorfon
Muylle, E.; Oyaert, W.; and Rogiers, M., 1979, Vlaams Diergeneesk. Tijdschr., v. 48 (4), 279-282
Gasterophilus intestinalis larvae, horses, mebendazole + trichlorfon paste, endoscopic assessment of efficacy
- Mebendazole
Narasimham, M. V. V. L.; et al., 1978, South-east Asian J. Trop. Med. and Pub. Health, v. 9 (4), 571-575
Wuchereria bancrofti, human, comparison of levamisole, levamisole + mebendazole, and diethylcarbamazine: Calicut, Kerala, India
- Mebendazole (Multispec)
Neave, R. M. S., 1979, Vet. Rec., v. 104 (21), 490 [Letter]
mebendazole highly toxic for psittaciformes and columbiformes
- Mebendazole (Telmin)
Novak, M.; and Evans, W. S., 1978, Canad. J. Zool., v. 56 (4), 604-607
Hymenolepis diminuta in Tribolium confusum, mebendazole, effect on different developmental stages of cysticercoids

Mebendazole -- Continued.

- Mebendazole (Mebenvet)
Oguz, T., 1976, Vet. Fak. Dergisi, Ankara Univ., v. 23 (3-4), 385-395
Cysticercus tenuicollis, lambs (exper.), efficacy of embay 8440 and mebendazole
- Mebendazole (Menzole; Mebutar)
Osimani, J. J.; et al., 1976, Semana Med. (4964), an. 83, v. 149 (15), 503-508
helminthiasis, humans, clinical trials with mebendazole: Montevideo, Uruguay
- Mebendazole (Vermox; Pantelmin)
Otero Dominguez, R.; Galvez Oviedo, M.; and Estevez Cazallas, B., 1977, Rev. Cubana Med. Trop., v. 29 (1), 13-16
Trichuris trichiura, Necator americanus, Enterobius vermicularis, humans, clinical trials using mebendazole: Cuba
- Mebendazole
Patel, A. A.; et al., 1978, Indian J Pharm., v. 40 (1), 20-21
mebendazole, non-aqueous method for estimating quantities, routine control measure for manufacturing laboratories
- Mebendazole
Paul, F. M.; and Zaman, V., 1975, Singapore Med. J., v. 16 (1), 11-18
Trichuris trichiura, young children with associated malnutrition and nutritional anemia, clinical trials with mebendazole, effective, well tolerated, no side effects: Outram Road General Hospital, Singapore
- Mebenvet
Pav, J., 1978, Veterinarstvi, v. 28 (2), 84-86
nematodes, prevalence in hares, seasonal dynamics, treatment with mebenvet
- Mebendazole
Pawlowski, Z.; Kozakiewicz, B.; and Wroblewski, H., 1978, Vet. Sc. Commun., v. 2 (2), 137-139
Taenia saginata cysticerci, bullocks (exper.), mebendazole and praziquantel administered in fodder, drug efficacy
- Mebendazole
Raeymaekers, A. H. M.; et al., 1978, Arzneimittelforsch., v. 28 (4), 586-594
Syphacia muris, Strongyloides ratti, synthesis and anthelmintic activity of mebendazole, flubendazole and other alkyl-(5-acyl-1-H-benzimidazol-2-yl) carbamates in rats
- Mebendazole
Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (1), 55-58
Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (Mesocricetus auratus)
- Mebendazole
Reinhardt, P., 1978, Monatsh. Vet.-Med., v. 33 (23), 898-901
Dicrocoelium dendriticum, sheep, efficacy of various anthelmintics compared

Mebendazole -- Continued.

- Mebendazole (Telmin)
Remfry, J., 1978, Lab. Animals, v. 12 (4), 213-218
helminth infections in imported Macaca mulatta, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University
- Mebendazole (Mebutar)
Repetto, O. M.; and Slaski, F., 1975, Semana Med. (4928), an. 82, v. 147 (23), 673-676
intestinal parasites, children, evaluation of mebendazole as possible broad spectrum anthelmintic
- Mebendazole (Mebenvet Granulatum)
Ribbeck, R.; and Winter, J., 1978, Monatsh. Vet.-Med., v. 33 (18), 706-707
gastrointestinal nematodes, goats, mebendazole
- Mebendazole
Rodriguez Caabeiro, F.; et al., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 249-258
Trichinella spiralis, human, efficacy of mebendazole at various times during infestation
- Mebendazole
Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility
- Mebendazole (Mebenav)
Santiago, M. A. M.; da Costa, U. C., 1978, Rev. Centro Cien. Rurais, v. 8 (1), 7-13
helminths, domestic fowl, mebendazole in feed, anthelmintic activity, no effect on host egg production
- Mebendazole (R17635)
Sargent, R. G.; et al., 1975, South. Med. J., v. 68 (1), 38-40
trichuriasis, humans, single-blind clinical study of mebendazole: South Carolina
- Mebendazole (Vermox)
Scragg, J. N.; and Proctor, E. M., 1978, Am. J. Trop. Med. and Hyg., v. 27 (2, pt. 1), 255-257
Trichuris trichiura, children, mebendazole treatment of symptomatic infections, anti-diarrheal agent (loperamide hydrochloride) enhanced efficacy of therapy
- Mebendazole
Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies
- Mebendazole
Shibuya, T.; et al., 1979, Japan. J. Exper. Med., v. 49 (5), 351-353
Litomosoides carinii in cotton rat (exper.), macro- and microfilaricidal activity, intrathoracic, intraperitoneal, and oral administration

Mebendazole -- Continued.

- Mebendazole
Shiratsuchi, M. S.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (3), 206-209
Trichocephalus trichiurus, humans, mebendazole
- Mebendazole
Shivakumar, A. M.; and Sabir, M., 1979, Indian Vet. J., v. 56 (2), 105-111
mebendazole, pharmacological actions on various organ-systems of animals
- Mebendazole
Simpkin, K. G.; and Coles, G. C., 1978, Research Vet. Sc., v. 25 (2), 249-250
Haemonchus and Trichostrongylus colubriformis in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parbendazole, oxbendazole)
- Mebendazole (Telmin)
Slocombe, J. O. D., 1979, Canad. Vet. J., v. 20 (5), 136-140
Anoplocephala perfoliata, horses, prevalence, efficacy of pyrantel pamoate, mebendazole, and niclosamide in field and critical trials: southern Ontario
- Mebendazole (Multispec)
Smith, G. A., 1979, Vet. Rec., v. 104 (24), 559 [Letter]
mebendazole, toxicity for Columbiformes and Psittaciformes questioned
- Mebendazole (R 17,635)
de Souza, D. W. C.; Souza, M. S. de L.; and Neves, J., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (1), 30-33
human mixed infections, intestinal parasites, mebendazole
- Mebendazole (R 17635)
de Souza, D. W. C.; Souza, M. S. de L.; and Neves, J., 1973, Rev. Soc. Brasil. Med. Trop., v. 7 (4), 237-241
human polyparasitism, clinical trials with mebendazole, cure rates over 90% with all parasites except Taenia: vicinity of Belo Horizonte, Brazil
- Mebendazole (Vermox)
Starke, J., 1979, Deutsche Med. Wchnschr., v. 104 (32), 1132-1135
Echinococcus alveolaris, humans (liver), mebendazole, case reports
- Mebendazole
Tidball, J. S.; Aguas, J. P.; and Aldis, J. W., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (1), 33-40
Capillaria philippinensis, humans (stools), 32 cases, history of eating raw fish, thiabendazole, mebendazole, life cycle discussed: San Antonio and San Narciso, Zambales Province, Western Luzon, Philippines

Mebendazole -- Continued.

- Mebendazole
Verheyen, A.; et al., 1978, J. Parasitol., v. 64 (3), 411-425
Taenia taeniaeformis, mebendazole medication of infected mice induced drastic time-related changes on surface topography of mature cysticerci, difference in susceptibility towards the drug between scolex, pseudoproglottids, and bladder in relation to morphology of their microtrichous covering
- Mebendazole (Vermox)
Wagner, E. D.; and Rexinger, D. D., 1978, Am. J. Trop. Med. and Hyg., v. 27 (1, pt. 1), 203-205
Ascaris lumbricoides, Trichuris trichiura, mebendazole and levamisole tested for possible ovicidal effects in vivo, human: Sierra Leone; Nassau, Bahamas
- Mebendazole
Wikerhauser, T.; Brglez, J.; and Kuticic, V., 1978, Acta Parasitol. Iugoslavica, v. 9 (2), 57-63
Echinococcus granulosus, mice, mebendazole, praziquantel
- Mebendazole (Vermox)
Wilson, J. F.; Davidson, M.; and Rausch, R. L., 1978, Am. Rev. Resp. Dis., v. 118 (4), 747-757
Echinococcus multilocularis, humans, case reports, mebendazole, clinical trial
- Mebendazole (Vermox)
Winter, P. A. D.; and Fripp, P. J., 1978, South African Med. J., v. 54 (14), 556 [Letter]
cutaneous larva migrans, human, topical application of mebendazole in anaesthetic ointment: South Africa
- Mebendazole
Zajicek, D.; Marova, M.; and Zahradnikova, W., 1978, Vet. Med., Praha, v. 51, v. 23 (1), 29-33
Haemonchus contortus, Trichostrongylus colubriformis, sheep (exper.), efficacy of mebendazole, hematological indices before and after treatment
-
- Mebenvet. See Mebendazole.
- Mebenvet Granulatum. See Mebendazole.
- Mebutar. See Mebendazole.
- Mecadox. See Carbadox.
- Mecamylamine
Hillman, G. R.; Gibler, A. M.; and Anderson, J. W., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 992-997
Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

Meclo vaginal cream or suppositories. See Clotrimazole or Metronidazole.

Mefloquine -- 2,8-Bis-(trifluoromethyl)- α -(2-piperidyl)-4-quinolinemethanol methylsulfonate monohydrate; DL-Erythro-2,8-bis(trifluoro-methyl)- α -(2-piperidyl)-4-quinolinemethanol hydrochloride; DL-Erythro- α -(2-piperidyl)-2,8-bis-(trifluoromethyl)-4-quinolinemethanol; Mefloquine hydrochloride; WR 142,490.

Mefloquine hydrochloride
Desjardin, R. E.; et al., 1979, Clin. Pharm. and Therap., v. 26 (3), 372-379
Plasmodium falciparum, healthy and infected humans, mefloquine hydrochloride, kinetics

Mefloquine
Fitch, C. D.; Chan, R. L.; and Chevli, R., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 258-262
Plasmodium berghei, undiminished mefloquine accumulation by erythrocytes infected with chloroquine-resistant strain provides explanation for superiority of mefloquine in treating chloroquine-resistant malaria, but competition observed between chloroquine and mefloquine raises possibility that same process of accumulation serves both drugs

Mefloquine
Lopez Antunano, F. J.; and Wernsdorfer, W. H., 1979, Bull. World Health Organ., v. 57 (4), 663-665
chloroquine-resistant Plasmodium falciparum, in vitro response to mefloquine, microtechnique system

Mefloquine
Mendenhall, D. W.; Higuchi, T.; and Sternson, L. A., 1979, J. Pharm. Sc., v. 68 (6), 746-750
mefloquine, hydrophobic amine antimalarial, low levels in whole blood samples analyzed with plastic ion-selective electrode

Mefloquine
Nakagawa, T.; et al., 1979, J. Pharm. Sc., v. 68 (6), 718-721
antimalarials, whole blood concentrations, gas liquid chromatography determinations, in vivo time course plots

2,8-Bis-(trifluoromethyl)- α -(2-piperidyl)-4-quinolinemethanol methylsulfonate monohydrate (WR 142,490)
Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Mefloquine hydrochloride
Rozman, R. S.; Molek, N. A.; and Koby, R., 1978, Drug Metabolism and Disposition, v. 6 (6), 654-658
mefloquine hydrochloride, absorption, distribution, and excretion in the mouse

Mefloquine -- Continued.

Mefloquine (WR-142,490)
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta

Mefloquine
Thong, Y. H.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (4), 388-390
mefloquine (new antimalarial compound), effect on mitogen-induced human and mouse lymphocyte proliferative responses, effect on antibody responses and delayed-type hypersensitivity responses to sheep red blood cells in treated mice

Mefloquine hydrochloride. See Mefloquine.

Meglumine antimoniate. See N-Methylglucamine antimonate.

Mekarazole (Preparation-665)
Shedivtsova, A., 1976, Med Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Melarsen
Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Melarsenoxide
Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Melarsonyl potassium -- Mel W; Trimelarsen.

Mel W
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Melarsonyl potassium -- Continued.

Mel W (Trimelarsan)

Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
 Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Melarsonyl potassium (Mel W; Trimelarsan)

Tanaka, H.; et al., 1977, Japan. J. Exper. Med., v. 47 (4), 315-317
 Litomosoides carinii-infected cotton rats, improved method for intrapleural injection of anti-filarial drugs to evaluate macrofilaricidal action

Melarsoprol -- 2-[4-[(4,6-Diamino-1,3,5-triazin-2-yl)-amino]phenyl]-1,3,2-dithiarsolane-4-methanol; Mel B.

Melarsoprol

Blair, L. S.; and Campbell, W. C., [1979], J. Parasitol., v. 64 (6), 1978, 1032-1034
 Dirofilaria immitis, pre-cardiac stages in Mustela putorius furo, trials of avermectin B_{1a}, mebendazole, and melarsoprol, possible value of Dirofilaria-Mustela model for chemotherapeutic studies

Melarsoprol

Blair, L. S.; and Campbell, W. C., 1979, Am. J. Vet. Research, v. 40 (7), 1031-1032
 Dirofilaria immitis, dogs, avermectin B_{1a} rapidly removed microfilariae from blood, administered with adulticide (melarsoprol) removal was permanent

Mel B (Melarsoprol)

Buyst, H., 1975, Ann. Soc. Belge Med. Trop., v. 55 (2), 95-104
 Trypanosoma rhodesiense, human, specific treatment with suramin and mel B, adjuvant antimalarial treatment with chloroquine and proguanil; modifications of sleeping sickness therapy advocated on physio-pathological and epidemiological grounds: Luangwa Valley, Zambia

Mel-B

Buyst, H., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 201-212
 sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia

Mel-B

Traub, N.; et al., 1978, East African Med. J., v. 55 (10), 477-481
 Trypanosoma brucei rhodesiense, human congenital, fatal infection in mother, infant successfully treated with suramin and mel-B, immunoglobulin levels at diagnosis, during treatment, and post-treatment, case reports: Zambia

Mel B. See Melarsoprol.

Melphalan

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
 Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Mel W. See Melarsonyl potassium.

Menadione

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329
 Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

Menadione

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
 Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Menadione

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
 Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Menadione

Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579
 Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Menazon

Getta, G. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 72-75
 hypodermatosis, cattle, insecticides, repellents

Menichlopholan. See Niclofolan.

Menoctone -- 2-Hydroxy-3-(8-cyclohexyloctyl)-1,4-naphthoquinone.

Menoctone

Gutteridge, W. E.; Dave, D.; and Richards, W. H. G., 1979, Biochim. et Biophys. Acta, v. 582 (3), 390-401
 Kinetoplastida spp., Plasmodium spp., conversion of dihydroorotate to orotate, mechanism of reaction different in these 2 groups of protozoa, possible target of chemotherapeutic attack

Menoctone

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
 Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Menzole. See Mebendazole.

Mepacrine. See Quinacrine.

Mepacrine hydrochloride. See Quinacrine.

Mepartricin -- Methyl-partricin; Searle SN 654;
SPA-S-222 (with Sodium lauryl sulfate);
Tricandil.

Methyl partricin (Tricandil)
Andriani, A.; and Loiudice, L., 1975, Quad.
Clin. Ostet. e Ginec., v. 30 (1), 21-29
Trichomonas vaginalis, clinical trials test-
ing the efficacy of methyl partricin vaginal
tablets in treating human vaginitis, marked
trichomonacidal effect with certain dose-
effect relationship

Mepartricin (Tricandil)
Berghella, A., 1973, Quad. Clin. Ostet. e Gi-
nec., v. 28 (5-12), 155-168
Trichomonas vaginalis, human vaginal tricho-
moniasis, topically applied mepartricin com-
pared favorably with chlorotrimazole used as
drug standard in clinical trials

Mepartricin + Sodium lauryl sulfate (SPA-S-222)
Imparato, E.; et al., 1976, Quad. Clin. Ostet.
e Ginec., v. 31 (4), 225-239
Trichomonas vaginalis, human vaginal tricho-
moniasis, evaluation of mepartricin as oral
therapy using nimorazole and clotrimazole as
reference drugs, best results obtained with
mepartricin

Mepartricin (Tricandil)
Moggian, G.; Tamburini, E.; and Visona, E.,
1975, Quad. Clin. Ostet. e Ginec., v. 30 (4),
143-154
T[richomonas] vaginalis, human vulvovaginitis,
clinical trials comparing mepartricin with
amphotericin B combined with tetracycline in
topical creams, good results

Methyl-partricin
Ograjensek, Z., 1974, Quad. Clin. Ostet. e
Ginec., v. 29 (5-6), 127-133
Trichomonas vaginalis, human vaginitis, suc-
cessful clinical trials with methyl-partricin
vaginal suppositories using natamycin as re-
ference drug

Methyl-partricin (Tricandil)
Perini, G.; and Bulzomi, R., 1974, Quad. Clin.
Ostet. e Ginec., v. 29 (1-2), 39-46
Trichomonas vaginalis, human vaginal infec-
tions, successful clinical trials with methyl-
partricin vaginal tablets which provided more
prompt resolution of infection than did vagi-
nal suppositories

Mepartricin + Sodium lauryl sulfate (SPA-S-222)
Ruggeri, E., 1976, Quad. Clin. Ostet. e Ginec.,
v. 31 (4), 211-223
Trichomonas vaginalis, vaginal trichomoniasis,
oral mepartricin showed significant
therapeutic superiority when compared in
clinical trials with patients who received
nimorazole therapy orally or with patients
treated with a vaginal cream containing
amphotericin B and tetracycline

Mepartricin -- Continued.

Methylpartricin (Searle SN 654)
Zanella, D.; et al., 1976, Therapiewoche, v. 26
(44), 7244-7250
trichomoniasis, human vaginal, single or
mixed candidiasis infections, methylpartricin
therapy compared with metronidazole

Mercaptophos. See Fenthion.

4-Mercaptopteridin
Christow, C., 1978, Biochem. and Exper. Biol.,
v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂
antagonists upon growth

2-Mercaptopurine
Irvin, A. D.; and Young, E. R., 1978, Research
Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxan-
thine uptake in vitro could be used as
primary screen for babesicidal drugs but
drugs showing in vitro activity are not
necessarily active in vivo

6-Mercaptopurine
Brotherton, J., 1978, Arzneimittel-Forsch.,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

6-Mercaptopurin
Christow, C., 1978, Biochem. and Exper. Biol.,
v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂
antagonists upon growth

6-Mercaptopurine
Irvin, A. D.; and Young, E. R., 1978, Research
Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxan-
thine uptake in vitro could be used as
primary screen for babesicidal drugs but
drugs showing in vitro activity are not
necessarily active in vivo

6-Mercaptopurine
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, Antimicrob. Agents and Chemotherapy, v.
15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

6-Mercaptopurine
McHardy, N., 1978, Ann. Trop. Med. and Parasi-
tol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected
bovine lymphoblastoid cell cultures used in
in vitro screens to test wide range of
compounds for chemotherapeutic activity

6-Mercaptopurine
Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
Fac. Agric. Kagoshima Univ. (24), v. 15, 125-
130
Echinococcus granulosus, scolicedal effect
of 65 antibiotic, antineoplastic, cytosta-
tic, and other agents in vitro

Mercaptotriazoles

Soliman, R.; and Hammouda, N. A., 1979, J. Pharm. Sc., v. 68 (11), 1377-1381
Schistosoma mansoni, Toxocara canis, mice (exper.), synthesis and activity of new mercaptotriazoles

Mercuric acetate

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Mercuric oxide

Benchimol, R.; and Benchimol, E., 1973, Rev. Brasil. Oftal., v. 32 (2), 203-204
Phthirus pubis, human infection of right eyelashes, apparently transmitted to lashes after scratching pubic area, case report, cure with mercuric oxide

Merthiolate. See Thimerosal.**Mesulfen**

Belda, W., 1975, Rev. Brasil. Clin. e Terap., v. 4 (8), 309-312
human scabies, increasing incidence, successful management with monosulfiram: Brazil

Metacil. See 4-Dimethylamino-m-tolyl methyl-carbamate.**Metamidium**

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Metamidium, Double

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Metamidium hydrochloride, Red isomer

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Metasystox. See Methyl demeton.**Methacycline** -- 6-Deoxy-6-demethyl-6-methylene-5-hydroxy-tetracycline; Rdomomycin.**Methacycline**

Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances

Methacycline -- Continued.**Rdomomycin**

Kasprzak, W.; Mazur, T.; and Rucka, A., 1974, Ann. Soc. Belge Med. Trop., v. 54 (4-5), 351-357
4 strains of free-living amoebae isolated from lakes in Poland, pathogenicity for mice, response to several drugs, identified as Acanthamoeba spp. on basis of morphology and protein disc electrophoretic patterns

Metham sodium -- Vapam.**Vapam**

Guttowa, A.; and Boniecka, B., 1977, Acta Parasitol. Polon., v. 24 (28-34), 315-322
Fasciola hepatica, Triaenophorus nodulosus, embryos, in vitro effects of pesticides Vapam and Lebaycid, implications for effects of environmental pollution on structure of ecosystems

Vapam

Moczon, T., 1976, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (5), 289-292
Fasciola hepatica miracidia, inhibitory effect of pesticides on enzyme activity

Methidathion -- S(2,3-Dihydro-5-methoxy-2-oxo-1,3,4-thiadiazol-3-ylmethyl) dimethyl phosphorothiothionate; Somonil.**Methidathion (Somonil)**

Kettle, P. R.; and Lukies, J. M., 1979, N. Zealand Vet. J., v. 27 (4), 78-79
Linognathus vituli, cattle, pour-on formulations of phosmet, methidathion, chlorpyrifos, and temephos: Kaitoke, near Upper Hutt, New Zealand

Methisazone (Methylisatin thiosemicarbazone)

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Methoprene -- Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate; Zoecon 515.**Methoprene**

Barrett, C. C.; et al., 1978, Southwest. Entom., v. 3 (3), 232-236
Hypoderma lineatum, H. bovis, cattle, effect of methoprene on cattle grub eclosion, different times and methods of treatment

Methoprene -- Continued.

Isopropyl 11-methoxy-3,7,11-trimethyl-2,4-decadienoate

Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, *J. Econom. Entom.*, v. 66 (1), 127-130

Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

Methoprene

Hopkins, D. E.; and Chamberlain, W. F., 1978, *Southwest. Entom.*, v. 3 (4), 292-294

Melophagus ovinus, inhibited maturation on sheep dipped in diflubenzuron (preventing formation of pupae) or methoprene (preventing adult eclosion), possible candidate control agents for sheep ked and other larviparous pests

Methoprene

Miller, J. A.; et al., 1977, *J. Econom. Entom.*, v. 70 (4), 417-423

Haematobia irritans, evaluation of effectiveness of methoprene in drinking water of cattle to suppress hornfly populations preparatory to release of sterile male flies: island of Molokai, Hawaii

Methoprene

Miller, J. A.; et al., 1979, *Southwest. Entom.*, v. 4 (3), 195-200

Haematobia irritans, cattle, methoprene sustained-release bolus for control in manure, 3 formulations compared: Camp Stanley, TX

Methoprene

Miller, R. W.; Pickens, L. G.; and Hunt, L. M., 1978, *J. Econom. Entom.*, v. 71 (2), 274-278

Haematobia irritans, cattle, methoprene as feed additive reduced horn fly counts: Beltsville, Md.

Methoprene

Prasert, V.; et al., 1975, *J. Econom. Entom.*, v. 68 (5), 639-640

Oestrus ovis, sheep, treatment with methoprene nasal spray

Methoprene (Zoecon 515)

Wright, J. E.; et al., 1974, *J. Med. Entom.*, v. 11 (4), 385-389

Cochliomyia hominivorax, methoprene and R-20458 inhibited emergence in vitro when applied topically or incorporated in larval diet, yearling heifers sprayed with each analogue and infested with larvae displayed little change in serum chemistry and adults emerged from the wound-reared larvae

Methotrexate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Methotrexate

Jaffe, J. J.; et al., 1978, *J. Parasitol.*, v. 64 (2), 193-197

Brugia pahangi-infected *Aedes aegypti* treated with sulfisoxazole and methotrexate singly or in combination, average number of infective larvae recovered was half of that recovered from controls and many larvae recovered were small and sluggish, most likely mode of action is inhibition of synthesis de novo of dihydrofolate in either parasite or more likely in mosquito host (leading to folate-related nutritional deficiencies inimical to normal filarial larval development)

Methotrexate

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Methotrexate

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511

Theileria parva and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

6-Methoxycarbonylamino-2-(4-thiazolyl)-1-benzimidazoloxoacetic acid di(2-hydroxyethyl)amine salt

Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618

Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Methoxychlor -- 1,1,1-Trichloro-2,2-bis(p-methoxyphenyl) ethane.Methoxychlor

Frazar, E. D.; and Schmidt, C. D., 1979, *J. Econom. Entom.*, v. 72 (6), 884-886

laboratory-reared *Haematobia irritans*, susceptibility to topically applied insecticides

Methoxychlor

Wright, F. C.; and Riner, J. C., 1979, *Southwest Entom.*, v. 4 (1), 40-45

Psoroptes ovis, *P. cuniculi*, 10 acaricides evaluated using 'tea-bag' technique

2-(β-Methoxyethyl)pyridine. See Methyridine.4-[p-6-(4-Methoxyimidylsulfamyl)phenylazo]-1-N-(N',N'-diethylaminoethyl)naphthylamine

Korolkovas, A.; Yang, G. N.; and Manfrinato, E., 1977, *Rev. Farm. e Bioquim. Univ. Sao Paulo*, v. 15 (1-2), 19-25

schistosomal activity and synthesis of 4-[p-6-(4-methoxyimidylsulfamyl)phenylazo]-1-N-(N',N'-diethylaminoethyl)naphthylamine

- 5-Methoxy-2-sulfanilamidepyrimidine. See
Sulfamer.
- 2-Methoxy-3-sulfanilamidopyrazine. See Sulfa-
lene.
- 4-Methoxy-6-sulfanilamido-pyrimidine monohydrate.
See Sulfamonomethoxine.
- 1-Methoxy Δ^8 -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, *Antimicrob. Agents and Chemotherapy*,
v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids
- 11-Methoxy Δ^8 -tetrahydrocannabinol
Pringle, H. L.; Bradley, S. G.; and Harris, L.
S., 1979, *Antimicrob. Agents and Chemotherapy*,
v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -
tetrahydrocannabinol and other cannabinoids
- L-I-Methyl-2-aminoethenole
Christow, C. P., 1978, *Riv. Biol.*, v. 71
(1-4), n. s. v. 31, 113-116
Trichomonas foetus, effect of decobald-co-
binamide and L-I-Methyl-2-aminoethenole
upon growth
- L-2-Methyl-2-aminoethenole
Christow, C., 1978, *Biochem. and Exper. Biol.*,
v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂
antagonists upon growth
- Methylbenzimidazol-2-yl carbamate
Ireland, C. M.; et al., 1979, *Biochem. Pharma-
col.*, v. 28 (17), 2680-2682
relative effectiveness of several benzimida-
zole carbamates and related compounds on
assembly of sheep brain microtubules in vitro
and on infections of Nematospiroides dubius
in mice
- 3'-Methylbenzo-15-crown-5
Brown, G. R.; and Foubister, A. J., 1979, *J.
Med. Chem.*, v. 22 (8), 997-999
benzo-15-crown-5 polyethers, synthesis, in
vivo and in vitro tests against Eimeria
tenella
- Methyl benzoquate -- Lerbek (with Metichlorpindol);
Methyl-7-benzoyloxy-6-butyl-1,4-dihydroxy-4-
oxoquinoline-3-carboxylate; Statyl.
- Methyl benzoquate
Chapman, H. D., 1978, *Parasitology*, v. 76 (2),
177-183
Eimeria tenella, Houghton strain, experimen-
tal development of resistance to amprolium,
clopidol, and methyl benzoquate
- Methylbenzoquate + Metichlorpindol (= Lerbek)
Greuel, E.; and Kuehnhold, W., 1977, *Prakt.
Tierarzt*, v. 58 (5), 338-341
Eimeria spp., chickens (exper.), synergistic
effect of metichlorpindol and methylbenzo-
quate, rotation program with other coccidio-
stats discussed
- Methyl benzoquate -- Continued.
- Methylbenzoquate + Clopidol (=Lerbek)
Hamet-Foure, N.; Macar, C.; and Robin, B., 1979,
Avian Path., v. 8 (1), 107-113
Eimeria meleagritidis, E. adenoides, turkeys,
activity of clopidol with methylbenzoquate
and amprolium with ethopabate: France
- Methyl benzoquate
Joyner, L. P.; and Norton, C. C., 1978, *Parasi-
tology*, v. 76 (3), 369-377
Eimeria maxima, activity of methyl benzoquate
and clopidol, synergy shown to be supra-
additive, collateral sensitivity could not be
demonstrated in resistant lines, effect of
Lerbek against standard and drug-resistant
lines, resistance transfer experiments with
clopidol- and methyl benzoquate-resistant
lines, preparation of bi-resistant lines,
attempts to develop Lerbek-resistant strain
- Methyl benzoquate + Clopidol (= Lerbek)
Joyner, L. P.; and Norton, C. C., 1978, *Parasi-
tology*, v. 76 (3), 369-377
Eimeria maxima, activity of methyl benzoquate
and clopidol, synergy shown to be supra-
additive, collateral sensitivity could not be
demonstrated in resistant lines, effect of
Lerbek against standard and drug-resistant
lines, resistance transfer experiments with
clopidol- and methyl benzoquate-resistant
lines, preparation of bi-resistant lines,
attempts to develop Lerbek-resistant strain
- Statyl
Krylov, V. F., 1978, *Veterinariia, Moskva*
(10), 68-69
Eimeria tenella strain resistant to pharm-
coccid after 35 laboratory passages in
chickens, cross-resistance only to rigeococcin
- Methyl benzoquate
Latter, V. S.; and Wilson, R. G., 1979, *Para-
sitology*, v. 79 (1), 169-175
Eimeria tenella, factors influencing assess-
ment of anticoccidial activity in cell
culture
- Statyl
McHardy, N., 1978, *Ann. Trop. Med. and Parasi-
tol.*, v. 72 (6), 501-511
Theileria parva- and T. annulata-infected
bovine lymphoblastoid cell cultures used in
in vitro screens to test wide range of
compounds for chemotherapeutic activity
- Lerbek
Mørch, J., 1978, *Nord. Vet. Med.*, v. 30 (6),
253-259
Eimeria spp., chickens (exper.), efficacy of
coccidiostats in feed, better production
efficiency of medicated groups
- Methyl benzoquate + Clopidol (= Lerbek)
Norton, C. C.; and Joyner, L. P., 1978, *Parasi-
tology*, v. 77 (3), 243-248
Eimeria maxima, development of resistance to
Lerbek, appearance of and subsequent selec-
tion for abnormal bisporocystic oocysts

Methyl benzoate -- Continued.

Methyl benzoate

Rollinson, D.; Joyner, L. P.; and Norton, C. C., 1979, *Parasitology*, v. 78 (3), 361-367
Eimeria maxima (Weybridge) and *E. maxima* (indentata) were distinguished by electrophoretic mobility of phosphoglucosyltransferase, this enzyme was used as marker to detect genetic transfer of methyl benzoate resistance between resistant and sensitive lines of these parasites

Methyl benzoate

Ryley, J. F.; and Hardman, L., 1978, *J. Parasitology*, v. 64 (5), 878-881
Eimeria acervulina, *E. mivati*, speciation studies (cross-immunity and drug resistance studies), some immunological relationship was demonstrated but the failure of the 2 organisms to interbreed in the drug resistance studies lends support to status of *E. mivati* as distinct species

Methylbenzoate

Ryley, J. F.; and Hardman, L., 1978, *Parasitology*, v. 76 (1), 11-20
Eimeria spp., chicks (exper.), effects of dietary vitamin K on severity of disease with particular attention to effects of vitamin K on response to anticoccidial drugs, concluded that use of vitamin K deficient diet for experimental work is quite justified

Statyl

Sherkov, S. N.; Leitch, B.; and Kokash, L., [1977], *Egypt. J. Vet. Sc.*, v. 13 (1), 1976, 37-43

Sarcocystis tenella, kittens (exper.), development in intestines, life cycle; attempted parasite suppression using statyl and pancocin plus

Methyl-5(6)-benzoyl-2-benzimidazole. See Mebendazole.

Methyl 5-benzoyl-2-benzimidazole carbamate. See Mebendazole.

Methyl (5-benzoyl-1-H-benzimidazol-2-yl) carbamate. See Mebendazole.

Methyl-N-[5(6)-benzoyl-2-benzimidazolyl] carbamate. See Mebendazole.

α -Methylbenzyl (E)-3-hydroxycrotonate dimethyl phosphate. See Crotoxyphos.

Methyl-7-benzoyloxy-6-butyl-1,4-dihydroxy-4-oxoquinoline-3-carboxylate. See Methyl benzoate.

Methyl 5-butyl-2-benzimidazole carbamate. See Parbendazole.

Methyl-5-6-butyl 1-2-benzimidazole carbamate. See Parbendazole.

Methyl-1-(butylcarbamoyl)-2-benzimidazole carbamate. See Benomyl.

1-Methyl-2-carbamoyloxymethyl-5-nitroimidazole. See Ronidazole.

1-Methyl-2-(p-carboxamidophenyl)-5-nitroimidazole
 Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Methyl-1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Methyl-5-(cyclopropylcarbonyl)-benzimidazol-2-yl-carbamate. See Cyclobendazole.

Methyl demeton -- Metasystox.

Metasystox

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, *Rev. Iber. Parasitol.*, v. 38 (1-2), 415-426

Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

[O-Methyl-O-(2,2-dichlorovinyl) phosphate]₂ Ca[O,O-dimethyl-(2,2-dichlorovinyl) phosphate]₂. See Caviphos.

4-(1'-Methyl-4'-diethylaminobutylamino)-7-chloroquinoline. See Chloroquine.

Methyl p-[(1,5-dimethylhexyl)oxy]benzoate
 Chamberlain, W. F.; Hopkins, D. E.; and Gingrich, A. R., 1973, *J. Econom. Entom.*, v. 66 (1), 127-130

Bovicola bovis, evaluation of 21 compounds for juvenile hormone activity

2,2'-Methylenebis(4-chloro-6-aminophenol)
 Sakamoto, T.; and Gemmill, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4-chloro-6-bromophenol). See Bromochlorophen.

2,2'-Methylenebis(4-chloro-6-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4-chlorophenol). See Dichlorophen.

Methylenebis(3,4-dichlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4,6-dichlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4-methylphenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4-nitro-6-bromophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(4-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Methylenebis(3,4,6-trichlorophenol). See Hexachlorophene.

Methylene blue
Brotherton, J., 1978, Arzneimittelforsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Methylene blue
Daniiarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Methylene blue (Tetramethylthionine chloride)
Lacuta, A. Q.; and Newman, A. R., [1977], Philippine J. Vet. Med., v. 15 (1-2), 1976, 169-180
Demodex canis, German Shepherd puppy, treatment with methylene blue positive iontophoresis, good results, case report; iontophoresis, review

Methylene- β -lapachone -- 3,4-Dihydro-2,2'-dimethyl-2H-naphtho [1,2-b] pyran-5-one-6-spiro-2'-oxyrane.

Methylene- β -lapachone -- Continued.

Methylene- β -lapachone
Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

Methylene- β -lapachone
Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531
Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

(E)-1-Methylethyl-3-[[[(ethylamino)methoxyphosphinothioyl]oxyl]-2-butenate. See Propetamphos.

Methyl [5-(4-fluorobenzoyl) 1H-benzimidazole-2-yl] carbamate. See Flubendazole.

1-Methyl-2-(p-fluorophenyl)-5-nitroimidazole (MK 910)
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

N-Methylglucamine antimonate -- Glucantime; Meglumine antimoniate.

Meglumine antimoniate (Glucantime)
Alving, C. R.; et al., 1978, Life Sc., v. 22 (12), 1021-1025
Leishmania donovani in Mesocricetus auratus (exper.), meglumine antimoniate, alone and incorporated into liposomes, results suggest that liposome-encapsulated meglumine antimoniate may be markedly more effective than drug alone in chronic infections

Meglumine antimoniate (Glucantime)
Alving, C. R.; et al., 1978, Proc. National Acad. Sc., v. 75 (6), 2959-2963
Leishmania donovani, hamsters, superior efficacies of liposome-encapsulated meglumine antimoniate and sodium stibogluconate, efficacy of treatment influenced by lipid composition and charge of liposomes, morphologic evidence that liposomes travel to intracellular site of parasite, encapsulation and reduction of dose should minimize toxic reactions to antimonials

N-Methylglucamine antimonate
Chiari, C. de A.; Mayrink, W.; and Magalhaes, P. A., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (5), 298-303
American cutaneous leishmaniasis, humans, N-methylglucamine antimonate therapy evaluated by indirect fluorescent antibody test

N-Methylglucamine antimonate -- Continued.

Methylglucamine antimonate (Glucantime)
Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

N-Methyl-glucamine antimonate
Raether, W.; Seidenath, H.; and Loewe, H., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 543-547
Leishmania donovani, golden hamsters, HOE 668 compared with known antileishmanial drugs, toxicity precludes further development but very good anti-leishmanial action qualifies it as standard compound in screening tests

Methylglyoxal-bis (guanylhydrazone)
Bachrach, U.; et al., 1979, Exper. Parasitol., v. 48 (3), 464-470
Leishmania spp., effect of ethidium, pentamidine, and methylglyoxal-bis (guanylhydrazone) on growth and on polyamine, RNA, and DNA synthesis

Methylglyoxal bis(guanylhydrazone)
Chang, K. P.; et al., 1978, J. Protozool., v. 25 (1), 145-149
methylglyoxal bis(guanylhydrazone) (MGBG), little in vitro effect on Blastocritidia culicis, Crithidia oncopelti, and Leishmania spp. but complete inhibition of growth of Trypanosoma brucei, reduced parasitemia of T. brucei and T. congolense in rats but infections relapsed, tracer studies with T. brucei showed that MGBG interfered with nucleoside incorporation

Methylglyoxal bis(guanyl hydrazone)
Nathan, H. C.; et al., 1979, J. Protozool., v. 26 (4), 657-660
Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

2-(1-Methyl heptyl)-4,6-dinitrophenyl crotonate.
See Dinocap.

[4-(4-Methyl-2-imidazolin-2-yl)-4'-[(p-4-methyl-2-imidazolin-2-yl)phenyl]carbamoyl]-carbanilide-dimethanesulphonate (Wander compound 2783)
Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested

4,4'-(4-Methyl-2-imidazolin-2-yl) terephthalanilide dimethane sulphate (Wander compound 2495)
Gill, B. S., 1972, Ann. Soc. Belge Med Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats chemoprophylactic trials, 8 compounds tested

Methylisatin thiosemicarbazone. See Methisazone.

1-Methyl-2-isopropyl-5-nitroimidazole. See Ipronidazole.

4-[p-(5-Methyl-3-isoxazolylsulfamyl)-phenylazo]-1-naphthylamine
Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124
preparation and testing of 6 long-acting schistosomicidal resins

Methylmercadone. See Nifuratel.

2-Methyl mercapto imidazoline hydroiodide
Ahmad, S.; Kishor, K.; and Shanker, K., 1979, Indian Drugs, v. 16 (5), 107-109
Hymenolepis nana, anthelmintic activity of imidazolines in vivo and in vitro, synthesis and acetylcholine esterase inhibitory activity of compounds

6-Methylmercaptapurine ribonucleoside
Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

4-Methyl-6-methoxy-3-[(4-amino-1-methylbutyl)amino]-quinoline diphosphate
Khan, M. S.; and LaMontagne, M. P., 1979, J. Med. Chem., v. 22 (8), 1005-1008
3- and 5-aminoquinolines, potential antimalarials, synthesis, testing of some against Plasmodium berghei in mice, Leishmania donovani in hamsters, or P. cynomolgi in rhesus monkeys

N-Methyl-1-naphthylcarbamate. See Carbaryl.

3-Methyl-4-(5'-nitrofurfurylidene-amino)-tetrahydrate-4H-1,4-thiazine-1,1-dioxide. See Nifurtimox.

1-Methyl-5-nitroimidazoles, substituted at 2-position
Winkelmann, E.; Raether, W.; and Gebert, U., 1978, Arzneimittelforsch., v. 28 (10), 1682-1684
activity of 16 novel 5-nitroimidazoles against protozoa in mice and golden hamsters, compared with metronidazole and tinidazole, structure-activity relationships

3-(1-Methyl-5-nitroimidazol-2-yl)-3 α ,4,5,6,7 α -hexahydro-1,2-benzisoxazole (MK436)
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

- N^1 -(1-Methyl-5-nitroimidazol-2-ylmethylcarbonyl)- N^2 -dimethylformadinium chloride hydrochloride
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- 1-(1-Methyl-5-nitro-2-imidazolylmethyl)-2-picolinium chloride
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- 3-Methyl-5-(4-nitrophenylazo)rhodanine. See Nitrodan.
- Methyl-2(6)-nitropyridinecarboxamides
Morisawa, Y.; et al., 1978, J. Med. Chem., v. 21 (2), 194-199
Eimeria tenella, synthesis and anticoccidial activity of methyl-2(6)-nitro- and -3(5)-nitropyridinecarboxamides
- Methyl-3(5)-nitropyridinecarboxamides
Morisawa, Y.; et al., 1978, J. Med. Chem., v. 21 (2), 194-199
Eimeria tenella, synthesis and anticoccidial activity of methyl-2(6)-nitro- and -3(5)-nitropyridinecarboxamides
- 3-Methyl-1,2,4-oxadiazole
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- 1-Methyl-2-(2-oxooxazolidin-3-yliminomethyl)-5-nitroimidazole
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618
Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens
- Methyl-5-p-oxy-2-benzimidazole-carbamate. See Oxibendazole.
- Methyl-partricin. See Mepartricin.
- Methyl 5(6)-phenyl-sulfinyl-2-benzimidazole-carbamate. See Oxfendazole.
- Methyl(5-(phenylsulfinyl)-1H-benzimidazol-2-yl) carbamate. See Oxfendazole.
- Methyl 6-(phenylsulfinyl)imidazo[1,2-a] pyridine-2-carbamate
Bochis, R. J.; et al., 1978, J. Med. Chem., v. 21 (2), 235-237
anthelmintic activity of methyl 6-(phenylsulfinyl)imidazo[1,2-a]pyridine-2-carbamate tested against multiple helminths and experimental hosts, results of tests show it to have highly potent broad-spectrum activity
- Methyl-5-(phenylthio)-2-benzimidazole carbamate. See Fenbendazole.
- Methyl[5-(phenylthio)-1H-benzimidazole-2-yl] carbamate. See Fenbendazole.
- 4-[5-(4-Methyl-1-piperazinyl)(2,5'-bi-1-4-benzimidazole-2-yl) phenol trihydrochloride. See 2-[2-(4-Hydroxyphenyl)-6-benzimidazolyl]-6-(1-methyl-4-piperazyl) benzimidazole.
- Methylprednisolone acetate (Depo-Medrol)
Nozik, R. A., 1977, Tr. Am. Acad. Ophth. and Otolaryngol., v. 83 (5), 811-818
toxoplasmic retinochoroiditis, humans, periorbital corticosteroid injections administered alone or in conjunction with systemic antimicrobial agents
- 4-Methylprimaquine -- WR 181 023.
- WR 181 023
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
-
- 2-Methylprimaquine dihydrochloride monohydrate -- WR 182 234.
- WR 182 234
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
-
- 3-Methylprimaquine diphosphate -- WR 211 814.
- WR 211814
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

3-Methylprimaquine diphosphate -- Continued.

WR 211 814

Kinnamon, K. E.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (4), 751-757
Leishmania donovani-*Mesocricetus auratus* model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

Methyl 5-n-propoxy-2-benzimidazole carbamate.
 See Oxibendazole.

Methyl-6-n-propoxybenzothiazole-2-carbamate.
 See Tioxidazole.

[5-[(2-Methylpropyl)sulfinyl]-1H-benzimidazol-2-yl] carbamic acid, methyl ester
 Cruthers, L. R.; et al., 1978, *Experientia*, v. 34 (12), 1574
 variety of nematodes, cestode, and trematode species in domestic animals, orally active benzimidazole anthelmintics discovered to be active by injection also

Methyl [5-(propylthio)-1H-benzimidazol-2-yl] carbamate. See Albendazole.

3-Methyl pyrantel tartrate. See Morantel.

DL- α -Methylserine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

2-(4'-Methylstyryl)-5-nitro-1-vinylimidazole
 Morton, D. M.; Fuller, D. M.; and Green, J. N., 1973, *Xenobiotica*, v. 3 (4), 257-266
 2-styryl-5-nitroimidazoles, metabolism and excretion in laboratory animals, activity against *Trypanosoma rhodesiense*

5-Methyl-3-sulfanilamido-isoxazole. See Sulfamethoxazole.

5-Methyl-tetrahydrohomofolic acid
 Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

(-)-2-Methyl-N-[3-(2,3,5,6-tetrahydroimidazo[2,1-b]thiazol-6-yl)phenyl] propanamide monohydrochloride. See Butamisole hydrochloride.

(1-Methyl-1,1,4,5,6-tetrahydro-2,2-thienyl)vinylpyrimidinium. See Pyrantel.

trans-2-(2-(3-Methyl-2-thienyl)vinyl)1-methyl-1,4,5,6-tetrahydropyrimidine tartrate. See Morantel.

trans-1-Methyl-2-[2-(thienyl)-vinyl]-1,4,5,6-tetrahydropyrimidine. See Pyrantel.

L-0-Methylthreonine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Methyridine -- Dekelmin; 2-(β -Methoxyethyl)pyridine; Mintic; Promintic.

Promintic

Daniyarov, I. A.; et al., 1978, *Veterinariia*, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Promintic

Kashnikov, A. A., 1973, *Vet. Nauka--Proizvod.*, Trudy, Minsk, v. 11, 140-142
 macracanthorhynchosis, swine, action of various anthelmintics

Mintic

Kuchin, A. S., 1970, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 8, 88-92
 S[trongyloides] papillosus, sheep, pathology, mintic, promintic, and phenothiazine, all effective

Promintic

Kuchin, A. S., 1970, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 8, 88-92
 S[trongyloides] papillosus, sheep, pathology, mintic, promintic, and phenothiazine, all effective

Methyridine

Kumar, V.; et al., 1978, *Acta Zool. et Path. Antverpiensia* (70), 221-225
 helminthiasis in wild mammals and birds of zoological gardens, need for control through preventive and chemotherapeutic measures; summary of observations on use of mebendazole in mammals and birds and of methyridine in pheasants: Antwerp Zoo

Methyridine (Dekelmin)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, *Ztschr. Parasitenk.*, v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Methyridine -- Continued.Methyridine

Rao, Y. V. B. G., 1976, Indian Vet. J., v. 53 (10), 776-777
Capillaria obsignata, chickens (exper.), critical tests with methyridine, pyrantel tartrate, and levamisole

Methyridine (Promintic)

Srivastava, K. K.; Varma, T. K.; and Rao, B. V., 1976, Indian Vet. J., v. 53 (10), 772-775
 strongylosis, cattle, promintic 100% effective in field trials after 7-14 days:
 Bareilly district, State of Uttar Pradesh

Metichlorpindol -- Clopidol; Clopindol; Coyden;

Coyden 25; 3,5-Dichloro-2,6-dimethyl-4-pyridinol; 2,6-Dimethyl-3,5-dichloro-4-hydroxy-pyridine; 2,6-Dimethyl-3,5-dichloro-pyridinol-4; Lerbek (with Methyl benzoquate); Rigeccocin.

Clopidol

Chapman, H. D., 1978, Parasitology, v. 76 (2), 177-183
Eimeria tenella, Houghton strain, experimental development of resistance to amprolium, clopidol, and methyl benzoquate

Rigeccocin (Clopidol)

Czegledi-Janko, G.; Balla, J.; and Toth, L., 1977, Acta Vet., Budapest, v. 26 (4), 1976, 445-453
 rigeccocin, very low residues in organs of broiler chickens after long-term treatment

Metichlorpindol + Methylbenzoquate (= Lerbek)

Greuel, E.; and Kuehnhold, W., 1977, Prakt. Tierarzt, v. 58 (5), 338-341
Eimeria spp., chickens (exper.), synergistic effect of metichlorpindol and methylbenzoquate, rotation program with other coccidiostats discussed

Clopidol + Methylbenzoquate (=Lerbek)

Hamet-Fouere, N.; Macar, C.; and Robin, B., 1979, Avian Path., v. 8 (1), 107-113
Eimeria meleagritidis, *E. adenoeides*, turkeys, activity of clopidol with methylbenzoquate and amprolium with ethopabate: France

Clopidol

Joyner, L. P.; and Norton, C. C., 1978, Parasitology, v. 76 (3), 369-377
Eimeria maxima, activity of methyl benzoquate and clopidol, synergy shown to be supra-additive, collateral sensitivity could not be demonstrated in resistant lines, effect of Lerbek against standard and drug-resistant lines, resistance transfer experiments with clopidol- and methyl benzoquate-resistant lines, preparation of bi-resistant lines, attempts to develop Lerbek-resistant strain

Meticlorpindol -- Continued.Clopidol + Methyl benzoquate (= Lerbek)

Joyner, L. P.; and Norton, C. C., 1978, Parasitology, v. 76 (3), 369-377
Eimeria maxima, activity of methyl benzoquate and clopidol, synergy shown to be supra-additive, collateral sensitivity could not be demonstrated in resistant lines, effect of Lerbek against standard and drug-resistant lines, resistance transfer experiments with clopidol- and methyl benzoquate-resistant lines, preparation of bi-resistant lines, attempts to develop Lerbek-resistant strain

Clopidol

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Rigeccocin

Krylov, V. F., 1978, Veterinariia, Moskva (10), 68-69
Eimeria tenella strain resistant to pharmacoid after 35 laboratory passages in chickens, cross-resistance only to rigeccocin

Clopidol

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Clopidol (Coyden)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
 coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Clopidol

McDougald, L. R.; and Galloway, R. B., 1977, Ztschr. Parasitenk., v. 54 (1), 95-100
Eimeria tenella in vitro, development inhibited by serum from chickens fed anticoccidial drugs, technique to assay drug activity and to characterize and quantitate therapeutic effect

Clopidol

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Clopidol (Coyden)

McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Coyden-25

Manuel, M. F., 1972, Philippine J. Vet. Med., v. 11 (2), 92-100
Eimeria tenella, broilers (exper.), evaluation of amprol plus and coyden-25 with or without 3-nitro-50 (roxarsone)

Meticlorpindol -- Continued.

Clopidol (Coyden-25)

Manuel, M. F.; Morales, E.; and Trovela, E., 1977, Philippine J. Vet. Med., v. 16 (1-2), 20-30

Leucocytozoon caulleryi, chickens under natural conditions, clopidol, halofuginone and furazolidone given in feed, clopidol 100% effective, no detrimental effects by any drug on host growth or red and white blood cell count

Lerbek

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259

Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Clopidol (Coyden)

Mogollon, J. D.; and Molina, T. de P., 1977, Rev. Inst. Colomb. Agropec, v. 14 (4), 473-488

Eimeria tenella field strain, sensitivity against 3 anticoccidial drugs

Clopidol + Methyl benzoate (= Lerbek)

Norton, C. C.; and Joyner, L. P., 1978, Parasitology, v. 77 (3), 243-248

Eimeria maxima, development of resistance to Lerbek, appearance of and subsequent selection for abnormal bisporocystic oocysts

Clopidol

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250

Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Clopidol (Coyden; Meticlorpindol; Clopidol)

Ryan, J. J.; et al., 1979, J. Environ. Quality, v. 8 (3), 439-442

clopidol transferred to lambs by feeding chicken waste, did not appear to accumulate in lamb tissues or faeces

Clopidol

Sevcik, B.; et al., 1974, Veterinaria, Praha, v. 16 (5-6), 421-588

coccidiosis; nicarbazine, clopidol, efficacy and toxicity in various animals, review

Clopidol (Coyden)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289

coccidiosis, broilers, anticoccidials, floor pen trials

Metrifonatum. See Trichlorfon.

Metriphosphate. See Trichlorfon.

Metrogyl. See Metronidazole.

Metrointestopan. See Brobenzoxaldine or Broxyquinoline or Metronidazole.

Metronidazole -- Clont; Efloran; Flagyl; Flagyl V; Hydroxyethylmethyl nitroimidazole; 1-(2-Hydroxyethyl)-2-methyl-5-nitroimidazole; Klion; Meclo vaginal cream or suppositories (with Clotrimazole); Metodine (with Diiodohydroxyquinoline); Metrogyl; Metrointestopan (with Intestopan); Mezil; Orvagil; 8,823 R.P.; Trichopol; Vagilen.

Metronidazole

Akashi, E.; et al., 1976, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 30 (7), 581-587
Trichomonas vaginitis, human, tinidazole, metronidazole, comparison clinical studies

Metronidazole

Anderson, R.; et al., 1979, South African Med. J., v. 55 (15), 593-596
metronidazole, no detectable effects on functions of human blood neutrophils and lymphocytes

Metronidazole

Aswapokee, N.; et al., 1974, Siriraj Hosp. Gaz., v. 26 (3), 371-376
Entamoeba histolytica, human hepatic abscess, single course of metronidazole plus closed aspiration of abscess, clinical trials, treatment well tolerated without side effects: Siriraj Hospital

Metronidazole

Atias, A.; et al., 1977, Rev. Med. Chile, v. 105 (7), 456-459
Entamoeba histolytica, invasive intestinal infection in nursing infants, statistics of 30 cases, metronidazole therapy: Chile

Metronidazole

Aure T., M., 1974, Rev. Obst. y Ginec. Venez., v. 34 (4), 437-440
human vaginal trichomoniasis, clinical trials comparing efficacy of tinidazole with metronidazole, concluded that drugs are of equal value in treatment: Venezuela

Metronidazole (Flagyl; 8823 R.P.)

Baines, E. J., 1978, J. Antimicrob. Chemother., v. 4, Suppl. C, 97-111
metronidazole, its past, present, and future, symposium presentation

Metronidazole

Bassett, M. L.; Danta, G.; and Cook, T. A., 1978, Brit. Med. J. (6129), v. 2, 19
Giardia lamblia, humans with associated peripheral neuropathy of unknown etiology, 2 case reports, resolution of giardiasis and neurological symptoms after metronidazole therapy

Metodine. See Diiodohydroxyquin or Metronidazole.

Metrifonate. See Trichlorfon.

Metronidazole -- Continued.

Metronidazole (Mezil)

Belehu, A.; Naafs, B.; and Touw-Langendijk, E., 1978, Brit. J. Dermat., v. 99 (4), 421-422
Leishmania aethiops, metronidazole of no value in treatment of five cases of Ethiopian mucocutaneous leishmaniasis

Metronidazole

Benazet, F.; et al., 1970, Scand. J. Infect. Dis., v. 2 (2), 139-143
intestinal and hepatic parasites, nitroheterocyclic antiparasitics, laboratory studies of chemotherapeutic activity and toxicity in exper. animals

Metronidazole

Bhandari, B.; and Sankhla, K., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (3), 345-346
[Letter]
Entamoeba histolytica-associated rectal prolapse in children, di-iodoquine and metronidazole

Metronidazole

Bindschadler, D. D., 1974, Rocky Mountain Med. J., v. 71 (7), 387-389
E[ntamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Metronidazole + Chloroquine + Chloramphenicol

Bindschadler, D. D., 1974, Rocky Mountain Med. J., v. 71 (7), 387-389
E[ntamoeba] histolytica, man, development of systemic amoebiasis with multiple hepatic abscesses 9 months after successful treatment for amebic dysentery with metronidazole, systemic infection successfully treated with combination of emetine and chloroquine followed by an intensive course of metronidazole, chloroquine and chloramphenicol

Metronidazole (Flagyl)

Bjorvatn, B.; and Neva, F. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 480-485
Leishmania tropica, white mouse model, experimental therapy using sodium stibogluconate, amphotericin B, metronidazole and WR 6026

Metronidazole (Flagyl)

Blandon, R.; Guevara, J. F.; and Johnson, C. M., 1976, Rev. Med. Panama, v. 1 (3), 153-162
Chagas disease in children, diagnosis, pathology, therapeutic trials with various drugs, metronidazole was well tolerated and therefore most promising therapy: Panama

Metronidazole

Boggs, C. H.; and Chakravorty, R. C., 1979, Virginia Med., v. 106 (9), 674-675
Entamoeba histolytica, 56-year-old male, necrotizing amebic colitis and perforated colon, case report, successful treatment with surgery, metronidazole, tetracycline: Virginia

Metronidazole -- Continued.

Metronidazole + Diiodohydroxyquinoline (= Metrodine)

Botero Ramos, D., 1976, Semana Med. Mexico (1127), an. 23, v. 89 (7), 209-212
Entamoeba histolytica, human, clinical trials testing efficacy of metronidazole combined with diiodohydroxyquinoline

Metronidazole

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Klion

Breier, M., 1972, Therap. Hungar., v. 20 (3), 109-111
Trichomonas vaginalis, humans, urological infections, klion: Hungary

Klion

Breier, M.; and Gyarmathy, F., 1971, Therap. Hungar., v. 19 (2), 64-66
Trichomonas vaginalis, males, epididymitis and other urogenital infections, clinical aspects, klion therapy: Hungary

Metronidazole

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Metronidazole (Flagyl)

Bruckner, D. A.; Bueding, E.; and Voge, M., 1979, J. Parasitol., v. 65 (3), 473-474
Trichomonas vaginalis, mice, lack of obligatory association between mutagenic and antitrichomonal effects of metronidazole

Metronidazole

Buttar, H. S.; Siddiqui, W. H.; and Moffatt, J. H., 1979, J. Pharm. and Pharmacol., v. 31 (8), 542-544
metronidazole, absorption, distribution, and elimination following intravaginal and oral administration to female rats

Metronidazole

Campos, R., 1973, Rev. Brasil. Clin. e Terap., v. 2 (11), 587-588
Entamoeba histolytica, comparison of efficacy of nifuratel and other amoebicides using material cultured from intestinal ulcers of patient with intestinal symptomatic amoebiasis

Metronidazole

de Carneri, I., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (5), 321-325
Balantidium coli in culture, nitrimidazine and metronidazole tested

Metronidazole

Cavier, R.; et al., 1979, Ann. Pharm. Franc., v. 37 (7-8), 309-312
Trichomonas vaginalis, Entamoeba histolytica, rats, 2-nitro benzofuran derivatives compared with metronidazole

Metronidazole -- Continued.

Metronidazole

Cavier, R.; and Cenac, J., 1972, *Therapeutique*, v. 48 (5-6), 391-394

Trichomonas vaginalis, nimorazole vs. metronidazole, in vitro and in vivo trials with rats and mice

Metronidazole

Cavier, R.; and Cenac, J., 1972, *Therapie*, v. 27 (4), 733-742

Trichomonas vaginalis, *Entamoeba*, in vitro and in vivo (rats, hamsters), efficacy of flunidazole compared with metronidazole

Metronidazole

Cerecedo Cortina, V.; and Gaxiola Gaxiola, R., 1972, *Rev. Fac. Med., Univ. Nac. Auton. Mexico*, an. 15, v. 15 (1), 25-28

human hepatic amoebic abscess, side effects of currently used amoebicides (emetine, chloroquine, metronidazole, aminosidine)

Metronidazole

Chiari, L.; Guerrero, J.; and dos Santos, C. N., 1978, *Ztschr. Parasitenk.*, v. 56 (2), 107-114

Entamoeba histolytica, experimental muscular infection in hamsters, pathology, metronidazole trial; useful biological model, particularly for chemotherapy studies

Metronidazole and 8 metronidazole analogues

Chien, Y. W.; and Mizuba, S. S., 1978, *J. Med. Chem.*, v. 21 (4), 374-380

Trichomonas vaginalis, metronidazole and 8 analogues, analysis of antimicrobial and electroactivity relationships

Metronidazole (Flagyl)

Das, S. R.; and Ghoshal, S., 1979, *Current Sc.*, Bangalore, v. 48 (2), 69-70 [Letter]

Entamoeba histolytica, axenically grown, amoebicidal activity of metronidazole reduced in vitro by intestinal bacteria

Metronidazole (Flagyl)

DeFord, J. W., 1973, *South. Med. J.*, v. 66 (10), 1149-1153

amebiasis, new methods of diagnosis and treatment with emphasis on the indirect hemagglutination test and metronidazole

Metronidazole

Delgado y Garnica, R.; and Martinez-Murray, R., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (4), 307-309

Entamoeba histolytica, patients, metronidazole: Mexico City

Metronidazole

Diez T., A.; et al., 1972, *Rev. Obst. y Ginec. Venez.*, v. 32 (3), 433-438

human vaginal trichomoniasis, clinical trials comparing efficacy of tinidazole with that of metronidazole: Venezuela

Metronidazole

Dutta, G. P.; and Narain, L., 1978, *Indian J. Exper. Biol.*, v. 16 (7), 838-840

Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Metronidazole -- Continued.

Metronidazole

Dykers, J. R., jr., 1978, *Am. J. Obst. and Gynec.*, v. 132 (5), 579-580

trichomonal vaginitis, treatment of patient and consort, metronidazole, single dose at various dosage levels, some side effects

Metronidazole

Dykers, J. R., jr., 1978, *Am. J. Obst. and Gynec.*, v. 132 (6), 703-704 [Letter]

trichomonal vaginitis, single dose treatment with metronidazole

Metronidazole

Edwards, D. I., 1979, *J. Antimicrob. Chemother.*, v. 5 (5), 499-502

Trichomonas vaginalis and other anaerobes, metronidazole, mechanism of antimicrobial action

Metronidazole

Edwards, D. I.; et al., 1979, *J. Antimicrob. Chemother.*, v. 5 (3), 315-316 [Letter]

metronidazole and misonidazole absorbed and inactivated by bacteria and fungi, mechanism of inactivation and relationship to known mechanism of drug action

Metronidazole

Fabio, P. F.; et al., 1978, *J. Med. Chem.*, v. 21 (3), 273-276

Entamoeba histolytica, diaminoanthraquinone bisamidines, laboratory trials comparing activity against cecal form in rats and hepatic form in golden hamsters with activity of known amoebicides

Metronidazole

Forsgren, A.; and Forssman, L., 1979, *Brit. J. Vener. Dis.*, v. 55 (5), 351-353

Trichomonas vaginalis, 35-year-old woman, symptomatic metronidazole-resistant vaginitis for 10 years, some resistance also to tinidazole and ornidazole: Sweden

Metronidazole (Clont; Flagyl; Orvagil)

Fuchs, P., 1978, *Deutsche Med. Wchnschr.*, v. 103 (3), 97-98

Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Hydroxyethylmethyl nitroimidazole (Flagyl)

Furtado, T., 1974, *Rev. AMMG*, v. 25 (3), 108-113

human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Metronidazole (Flagyl)

Gambardella, A.; et al., 1973, *Riforma Med.*, v. 87 (25), 992-1002

Entamoeba histolytica, human, development of cutaneous infection after surgical treatment of amoebic hepatitis with involvement of the peritoneum, complete recovery after metronidazole therapy: Pomigliano d'Arco, Italy

Metronidazole -- Continued.

Metronidazole

Garg, A. K.; and Jain, A., 1979, Indian Pediat., v. 16 (10), 913-916
E[ntamoeba] histolytica, giardiasis, children, metronidazole: India

Metronidazole

Giannone, R., 1973, Rev. Brasil. Med., v. 30 (9), 604-605
Trichomonas vaginalis, increasing resistance to metronidazole in parasite strains isolated in the Lombardy area of Italy

Metronidazole

Gillin, F. D.; and Diamond, L. S., 1978, J. Protozool., v. 25 (4), 539-543
Entamoeba histolytica, other Entamoeba spp., technique for clonal growth in agar, possible use in drug testing

Metronidazole

Goldstein, F.; Thornton, J. J.; and Szydowski, T., 1978, Am. J. Digest. Dis., n.s., v. 23 (6), 559-560
Giardia lamblia, humans, hepatobiliary form, trophozoites in bile, nonvisualization of gall bladder as important diagnostic feature, relief of biliary symptoms after metronidazole therapy, clinical case report

Metronidazole (Flagyl)

Grant, D.; and Woo, P. T. K., 1978, Canad. J. Zool., v. 56 (6), 1360-1366
Giardia spp. in small mammals, comparative studies, results suggest host specificity of some spp., infectivity of stored cysts varies with temperature, lack of prophylactic effect in rats treated with metronidazole or quinacrine hydrochloride

Metronidazole

Guggenmoos, R.; et al., 1978, Tropenmed. u. Parasitol., v. 29 (4), 423-426
nematodes, humans, ciclo bendazole, vermucidal effect compared with metronidazole in double-blind study, side effects only in small percentage of cases: Bamenda, Cameroon

Metronidazole

Gutierrez Samperio, C.; et al., 1976, Semana Med. Mexico (1106), an. 23, v. 87 (10), 293-297
amoebiasis, human, severe invasive intestinal infection, metronidazole administered parenterally with good results

Metronidazole

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Metronidazole (Flagyl)

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 339-347
Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

Metronidazole -- Continued.

Metronidazole

Hackett, L. P.; and Dusci, L. J., 1979, J. Chromatography, v. 175 (2), 347-349
metronidazole, tinidazole, detection and quantitation in human plasma using high-performance liquid chromatography

Metronidazole (Flagyl)

Hartley-Asp, B., 1979, Lancet, London (8110), v. 1, 275 [Letter]
vaginal trichomoniasis, patients, metronidazole, no chromosome-breaking activity was found, safe drug for short-term treatment

Metronidazole (Flagyl)

Hartley-Asp, B., 1979, Mutation Research, v. 67 (2), 193-196
metronidazole exhibits no cytogenetic effect in micronucleus test in mice or on human lymphocytes in vitro

Metronidazole (Flagyl)

Hartley-Asp, B., 1979, Toxicol. Letters, v. 4 (1), 15-19
T[richomonas] vaginalis, human, metronidazole treatment, absence of lymphocyte chromosomal damage

Metronidazole (Flagyl)

Hartong, W. A.; Gourley, W. K.; and Arvanitakis, C., 1979, Gastroenterology, v. 77 (1), 61-69
Giardia lamblia, patients, clinical spectrum and functional-structural abnormalities of small intestinal mucosa, treatment with metronidazole or quinacrine: Kansas Univ. Medical Center

Metronidazole

Hatchuel, W., 1975, South African Med. J., v. 49 (45), 1879-1881
Entamoeba histolytica, patients with amoebic liver abscesses, tinidazole vs. metronidazole, equally efficacious

Metronidazole (Flagyl V)

Holt, P. E.; Brown, A.; and Brown, B., 1978, Vet. Rec., v. 102 (18), 404-405
Strongyloides [sp.] in Lampropeltis getulus holbrooki, clinical symptoms, unsuccessful treatment with thiabendazole, mixed infection with flagellates and Ochetsomatidae sp., treatment for flagellates with metronidazole also unsuccessful, case report

Metronidazole (Flagyl)

Houghton, G. W.; et al., 1979, J. Antimicrob. Chemother., v. 5 (5), 621-623 [Letter]
metronidazole in healthy men, pharmacokinetics of oral vs. intravenous administration

Metronidazole

Islam, N.; and Hasan, M., 1978, J. Trop. Med. and Hyg., v. 81 (1), 20-22
Entamoeba histolytica, therapeutic efficacy of metronidazole and tinidazole compared in persons with hepatic infections, side effects

Metronidazole -- Continued.

- Metronidazole (Flagyl)
Jokipii, L.; and Jokipii, A. M. M., 1979, Med. Microbiol. and Immunol., v. 167 (1), 61-70
metronidazole, agar-well diffusion bioassay using bacteria, increased sensitivity
- Metronidazole (Flagyl)
Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167
Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria
- Metronidazole (Flagyl)
Kavousi, S., 1979, Am. J. Trop. Med. and Hyg., v. 28 (1), 19-23
giardiasis, infants and children, short and long-term followup after treatment with quinacrine vs. metronidazole
- Metronidazole (Flagyl)
Keystone, J. S.; Krajden, S.; and Warren, M. R., 1978, Canad. Med. Ass. J., v. 119 (3), 241-248
Giardia lamblia, epidemiology of outbreak in day-care nurseries, transmission apparently person-to-person, more Canadian children were symptomatic and infected than were immigrant children attending the nurseries, infections cleared with metronidazole or atabrine, control measures suggested including treatment of all infected children regardless of whether they were symptomatic: Toronto, Canada
- Metronidazole
Kluska, J., 1978, Terap. i Leky, v. 6, v. 28 (3), 107-112
Enterobius, lamblia, institutionalized children, control by improved sanitation and hygiene in conjunction with anthelmintics
- Metronidazole
Koch, R. L.; et al., 1979, Biochem. Pharmacol., v. 28 (24), 3611-3615
appearance of acetamide derived from metronidazole in conventional rats appears to be mediated by intestinal microflora, acetamide is a weak carcinogen
- Metronidazole
Koch, R. L.; and Goldman, P., 1979, J. Pharmacol. and Exper. Therap., v. 208 (3), 406-410
metronidazole forms N-(2-hydroxyethyl)-oxamic acid, anaerobic metabolism
- Metronidazole
Koesters, J.; et al., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (13), 266-268
Chilomastix gallinarum in chickens (exper.) and in vitro, metronidazole
- Metronidazole
Koesters, J.; Cubillos, A.; and Zuelch, S., 1978, Bol. Chileno Parasitol., v. 33 (3-4), 69-72
Chilomastix sp., cultivation in Dobell Laidlaw medium; in vitro and in vivo (chickens) testing of metronidazole

Metronidazole -- Continued.

- Metronidazole
Kogan, G. F., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 196-198
nosematosis, bees, fumagillin had good therapeutic effect, metronidazole, sulfadimethoxin and enteroseptol showed no substantial effect
- Metronidazole
Koutsaimanis, K. G.; Timms, P. W.; and Ree, G. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (4), 768-769
amoebiasis, Nigerian ship's steward with multiple hepatic abscesses, failure to respond to recommended doses of metronidazole, successful treatment with emetine hydrochloride
- Metronidazole
Lambert, B.; Lindblad, A.; and Ringborg, U., 1979, Mutation Research, v. 67 (3), 281-287
metronidazole and two of its urinary metabolites, no direct genotoxic effect on human lymphocytes in vitro
- Metronidazole
Lanbeck, K.; and Lindstrom, B., 1979, J. Chromatography (Biomed. Appl.), v. 162 (1), 117-121
assay of metronidazole and tinidazole in plasma and metronidazole in feces by high-performance liquid chromatography
- Metronidazole (Flagyl)
LaRusso, N. F.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (1), 19-24
absence of strand breaks in DNA treated with metronidazole
- Metronidazole (Flagyl)
LaRusso, N. F.; Lindmark, D. G.; and Mueller, M., 1978, Biochem. Pharmacol., v. 27 (18), 2247-2254
metronidazole, biliary and renal excretion, hepatic metabolism, and hepatic subcellular distribution in bile fistula rats, glucuronide conjugate of metronidazole was devoid of activity against Tritrichomonas foetus
- Metronidazole
Leite, E. V.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (1), 28-35
Giardia lamblia, humans, clinical trials, ornidazole and metronidazole compared: Brazil
- Metronidazole
Levi, G. C.; de Avila, C. A.; and Amato Neto, V., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (6), 422-424
giardiasis, humans, efficacy of various drugs, comparative study, side-effects
- Metronidazole
Lindmark, D. G.; and Mueller, M., 1976, Antimicrob. Agents and Chemotherapy, v. 10 (3), 476-482
metronidazole and 11 other nitroimidazoles, antitrichomonad activity against Tritrichomonas foetus and Trichomonas vaginalis, mutagenic action in Salmonella test, reducibility of nitro group by T. foetus hydrogenates, results underscore role of reduction of nitro group in antitrichomonad and mutagenic activity of nitroimidazoles

Metronidazole -- Continued.

- Metronidazole (Flagyl)
Lovgren, T.; and Salmela, I., 1978, Acta Path. et Microbiol. Scand., v. 86B (3), 155-158
Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents
- Metronidazole
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity
- Metronidazole
Manica, D.; and Peixoto, S., 1972, Rev. Brasil. Clin. e Terap., v. 1 (5), 317-320
human trichomoniasis, clinical trials comparing efficacy of tinidazole and metronidazole
- Metronidazole + Oxytetracycline
Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire
- Metronidazole
Matsuda, S., 1978, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 32 (2), 141-147
tinidazole, metronidazole, absorption, excretion, comparative study
- Metronidazole
Mehta, A. P.; and Guirges, S. Y., 1979, J. Trop. Med. and Hyg., v. 82 (7), 134-136
acute amoebic dysentery tentively diagnosed as caused by Acanthamoeba, 16-year-old-male, case report, successfully treated with metronidazole: India
- Metronidazole
Meingassner, J. G.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (1), 1-3
Tritrichomonas foetus, metronidazole-resistant and susceptible strains, in vitro susceptibility testing, results suggest that the two strains differ in regulation of internal redox systems and underscore the role that testing methods may play in the in vitro detection of nitroimidazole-resistant protozoan parasites
- Metronidazole
Meingassner, J. G.; and Thurner, J., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 254-257
Trichomonas vaginalis, isolation of strain resistant to metronidazole and other 5-nitroimidazoles
- Metronidazole
Minor Franco, A.; and Sanchez Granados, P., 1974, Rev. Med. Hosp. Gen., Mexico, v. 37 (6), 383-387
human cutaneous leishmaniasis, case report, unsuccessfully treated with metronidazole but cured with anthiomaline; high percentage of localization on ears and ear lobes in areas of Mexico

Metronidazole -- Continued.

- Metronidazole (Flagyl)
Moore, J. R., 1979, J. Am. Coll. Health Ass., v. 28 (2), 128
trichomoniasis, women, vaginitis, metronidazole given as 1 gram, single dose: Columbia University, New York
- Metronidazole
Mueller, M.; et al., 1979, Comp. Biochem. and Physiol., v. 64B (1), 97-100
Tritrichomonas foetus, Trichomonas vaginalis, Entamoeba invadens, effects of 2,4-dinitrophenol (including effect on accumulation of metronidazole)
- Metronidazole (Klion)
Muennich, D.; and Molnar, K., 1972, Therap. Hungar., v. 20 (1-2), 65-70
Entamoeba histolytica, adults, metronidazole: Hungary
- Trichopol
Musaev, F. A., 1972, Parazitologiya, Leningrad, v. 6 (2), 185-188
Trichomonas elongata, occurrence in oral cavity of healthy persons vs. those with oral cavity diseases, host age and sex, suitable media for cultivating trichomonads, activity in vitro of several medicinal substances, role of trichomonads in periodontitis confirmed by treatment of patients with trichopol
- Metronidazole (Flagyl)
Nesvadba, J., 1979, Kleintier-Praxis, v. 24 (4), 177-179
Giardia, cat, ornidazole, metronidazole, case report
- Metronidazole
Nseka, K.; and Mueller, M., 1978, Compt. Rend. Soc. Biol., Paris, v. 172 (6), 1094-1098
Tritrichomonas foetus, Entamoeba invadens, effect of glycolysis inhibitors on uptake of metronidazole
- Metronidazole (Flagyl)
Nygaard, B.; et al., 1977, Ugeskr. Laeger, v. 139 (9), 524-526
Trichomonas vaginalis, women with vaginitis, 7-day therapy with metronidazole vs. single-dose ornidazole
- Metronidazole
Olaeta Elizalde, R., 1973, Rev. Cir. Hosp. Juaréz, Mexico (187-188), v. 44, 1972-1973, 59-64
human hepatic amoebic abscess, therapeutic recommendations (emetine, dehydroemetine, chloroquine, metronidazole)
- Metronidazole
Padonou, K. O., 1975, Ang. Parasitol., v. 16 (2), 94-97
Dracunculus medinensis, humans, metronidazole superior to procaine penicillin: Nigeria
- Trichopol
Parre, J.; and Simovart, H. E., 1977, Festi Pollumaj. Akad. Teadusl. Toode Kogum. (104), 114-118
histomoniasis, turkeys, age and seasonal dynamics in relation to epizootiology; disease outbreaks in young birds under stress conditions; nitazol satisfactory, trichopol good prophylactic and therapeutic effect

Metronidazole -- Continued.

Metronidazole

Prakash, P.; and Saxena, S., 1976, Rajasthan Med. J., v. 15 (4), 240-244
giardiasis in children, evaluation of metronidazole using varying dosage schedules, recommendations for therapy

Metronidazole + Intestopan (= Metrointestopan)

Ramirez Andrade, R., 1977, Semana Med. Mexico (1170), an. 24, v. 93 (2), 41-44
amoebiasis, human chronic colitis or rectocolitis, clinical trials testing efficacy of metronidazole combined with intestopan, good results, drug well tolerated

Metronidazole

Richle, R.; et al., 1978, Arzneimittel-Forsch., v. 28 (4), 612-625
trichomoniasis, amoebiasis, lamblia, extensive in vitro and in vivo trials (humans, domestic animals, laboratory animals) with ornidazole to establish chemotherapeutic properties, efficacy slightly superior to metronidazole in comparative trials

Metronidazole

Salaki, J. S.; Shirey, J. L.; and Strickland, G.T., 1979, Am. J. Trop. Med. and Hyg., v. 28 (2), 190-193
Entamoeba polecki, 24-year-old Peace Corps volunteer (feces), symptomatic intestinal infection cured with diloxanide furate and metronidazole: United States (previously in Upper Volta)

Metronidazole

Sandia, O. G.; et al., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 52-56
E[ntamoeba] histolytica, humans with chronic intestinal amoebiasis, clinical trials with ornidazole vs. metronidazole

Metronidazole (Flagyl)

Sandront-Degee, M.; Werbrouck-Navette, J.; and Lambotte, R., 1975, Rev. Med. Liege, v. 30 (17), 560-562
Trichomonas vaginalis, humans, therapeutic trials comparing action of tiberal with metronidazole, fasigyn and tinidazole

Metronidazole (Metrogyl)

Sharma, V. P.; Rathore, H. S.; and Sharma, M. M., 1979, Am. J. Trop. Med. and Hyg., v. 28 (4), 658-660
dracunculiasis, patients, metronidazole in various dose schedules gives symptomatic relief but no preventive or vermifugal action

Metronidazole (Flagyl; Trichopol)

Shutskii, I. V.; et al., 1978, Vestnik Dermat. i Venerol. (9), 71-74
trichomoniasis, women, urogenital infections, clinical trials using various regimens of trichopol

Metronidazole (Flagyl)

Siddiqui, W. H.; and Buttar, H. S., 1979, Arch. Internat. Pharm. et Therap., v. 239 (1), 4-15
metronidazole as a single product or commercial cream, rats, intravaginal absorption, metabolism and disposition

Metronidazole -- Continued.

Metronidazole (Flagyl)

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriologie, 1 Abt. Orig., Reihe A, v. 241 (3), 358-367
Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Metronidazole

Skromne-Kadlubik, G.; Medina, J.; and Rauda, L., 1976, SPM Salud Pub. Mexico, v. 18 (2), 395-398
Entamoeba histolytica, human, comparative survey, conventional medications vs. radiation therapy

Metronidazole

Soyka, E., 1975, Therap. Umschau, v. 32 (9), 599-602
Trichomonas vaginalis, women, vaginal infections, single dose therapy with tinidazole compared with metronidazole: Switzerland

Metronidazole

Spencer, M. J.; Garcia, L. S.; and Chapin, M. R., 1979, Am. J. Dis. Child., v. 133 (4), 390-393
Dientamoeba fragilis, children with acute and chronic gastrointestinal symptoms, clinical findings, symptomatic recovery after treatment with diiodohydroxyquin or metronidazole indicates pathogenic role

Metronidazole

Sugano, H.; and Ando, K., 1978, Nippon Zyuishi-Kai Zasshi (J. Japan Vet. Med. Ass.), v. 31 (11), 635-638
Giardia, dogs (nat. and exper.), diarrhea, metronidazole

Metronidazole

Thacker, S. B.; et al., 1979, Am. J. Pub. Health, v. 69 (12), 1279-1281
Entamoeba histolytica, Giardia lamblia, control attempts in a residential facility for mentally retarded persons: Washington, D. C.

Metronidazole (Flagyl)

Thin, R. N.; et al., 1979, Brit. J. Vener. Dis., v. 55 (5), 354-356
Trichomonas vaginalis, women, vaginitis, double-blind comparison of single dose and a 5-day course of metronidazole: England

Metronidazole

Trzaska, B.; and Leinweber-Noiszewska, H., 1974, Ann. Acad. Med. Stetinensis, suppl. 10, 19-25
Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis

Metronidazole -- Continued.

Metronidazole

Valasco Chiriboga, A.; and Marquez Marino, G., 1975, Rev. Colomb. Obst. y Ginec., v. 26 (5), 291-298

human *Trichomonas vaginalis* vaginitis, comparative clinical trials using tinidazole and metronidazole; tinidazole recommended as drug of choice: Colombia

Metronidazole

Walzer, P. D.; Rutherford, I.; and East, R., 1978, Am. Rev. Resp. Dis., v. 118 (2), 415-418
Trichomonas sp. in man resulting in empyema secondary to presumed aspiration pneumonia, clinical case report, successful metronidazole therapy: University of Kentucky Medical Center Hospital, Lexington, Kentucky

Metronidazole

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Metronidazole

Winkelmann, E.; Raether, W.; and Gebert, U., 1978, Arzneimittel-Forsch., v. 28 (10), 1682-1684

activity of 16 novel 5-nitroimidazoles against protozoa in mice and golden hamsters, compared with metronidazole and tinidazole, structure-activity relationships

Metronidazole

Wyrzykowski, Z.; Wyrzykowska, K.; and Janowska, I., 1977, Med. Wet., v. 33 (5), 289-292

blood picture of turkeys fed fodder containing premix enriched with metronidazole

Metronidazole

Yabur T., J. A., 1972, Rev. Obst. y Ginec. Venez., v. 32 (3), 425-432

urogenital trichomoniasis, women, statistics of case histories, clinical aspects, diagnosis, value of metronidazole therapy: Venezuela

Metronidazole + Clotrimazole (= Meclo vaginal cream or suppositories)

Zacutti, A., 1975, Quad. Clin. Ostet. e Ginec., v. 30 (2), 65-96

Trichomonas vaginalis, clinical trials comparing metronidazole and clotrimazole alone or in new combination drug (meclo) as vaginal creams, vaginal suppositories and as oral tablets, pregnant and non-pregnant women and their sexual partners, good clinical results both in prevention and cure of infections

Metronidazole (Efloran)

Zanella, D.; et al., 1976, Therapiewoche, v. 26 (44), 7244-7250

trichomoniasis, human vaginal, single or mixed candidiasis infections, methylpartricin therapy compared with metronidazole

Mexaform. See Iodochlorhydroxyquin or Phanquone or Oxyphenonium bromide.

Mezil. See Metronidazole.

Miazol. See *Bacillus thuringiensis*.

Miconazole (nitrate)

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Miconazole

Lee, K. K.; et al., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (2), 217-220
Naegleria fowleri, in vitro susceptibility to selected antimicrobial agents singly and in combination

Milian ointment

Najar, H. C. F., 1978, Rev. Brasil. Med., v. 35 (3), 205-207

human scabies resulting in severe pruritis, clinical trials of milian ointment alone and combined with tetracosactide; combined therapy more successful with total eradication in 5 days

Milton solution; Milton tablets

Fripp, P. J.; Armstrong, F. A.; and Jaskulla, E., 1972, South African Med. J., v. 46 (47), 1819-1822

Schistosoma haematobium, *S. mansoni*, cercaricidal activity, hypochlorite solutions vs. halazone solution, formulations to make small amounts of water potable

Mimorazole. See Nitrimidazine.

Minimycin

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Minocin. See Minocycline.

Minocycline -- Minocin; Minocycline hydrochloride.

Minocycline hydrochloride (Minocin)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, Chemotherapy, v. 25 (4), 222-226

Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Minocycline (Minocin)

Lee, K. K.; et al., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (2), 217-220

Naegleria fowleri, in vitro susceptibility to selected antimicrobial agents singly and in combination

- Minocycline hydrochloride. See Minocycline.
- Mintes-B. See Bromsalans.
- Mintesol. See Thiabendazole.
- Mintezol. See Thiabendazole.
- Mintezole. See Thiabendazole.
- Mintic. See Methyridine.
- Minzolum. See Thiabendazole.
- Miracil-D. See Lucanthone.
- Mirasan. See 1-(3-Chloro-p-tolyl) piperazine hydrochloride.
- Mirincamycin
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
- Misonidazole
Edwards, D. I.; et al., 1979, *J. Antimicrob. Chemother.*, v. 5 (3), 315-316 [Letter]
metronidazole and misonidazole absorbed and inactivated by bacteria and fungi, mechanism of inactivation and relationship to known mechanism of drug action
- Mitomycin-C
Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Mitomycin C
Sinden, R. E.; and Smalley, M. E., 1979, *Parasitology*, v. 79 (2), 277-296
Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle
- Mitomycin C
Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Mitotane (1,1-Dichloro-2-(o-chlorophenyl)-2-(p-chlorophenyl)ethane)
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Mobam -- Benzo[b]thien-4-yl methylcarbamate.
- Mobam
Cole, M. M.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 118-119
Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides
-
- Molluscicides
Amin, A. H.; and Fenwick, A., 1978, *Trop. Doctor*, v. 8 (1), 8-12
schistosomiasis vector snail control, recommendations for use on small scale or by poor developing countries
- Molluscicides
Amin, M. A., 1972, *Sudan Med. J.*, v. 10 (2), 75-82
Biomphalaria and Bulinus vector snails, evaluation of drip-feed application of copper sulphate as molluscicide and of use of mechanical barriers for mollusk control: Gezira, Sudan
- Molluscicides
Ayad, N., 1976, *Egypt. J. Bilharz.*, v. 3 (2), 129-155
vector control programs (molluscicides, biological and physical control) being carried out in Egypt in an attempt to achieve permanent results in control of human schistosomiasis
- Molluscicides
Banna, H. B.; and Plummer, J. M., 1978, *Comp. Biochem. and Physiol.*, v. 61C (1), 33-36
Bulinus truncatus (intermediate host of Schistosoma haematobium), effect of N-tritylmorpholine on molluscan heart
- Molluscicides
Christie, J. D.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (3), 616-622
slow-release copper molluscicide tested against Biomphalaria glabrata, laboratory and field trials: St. Lucia
- Molluscicides
Chroustova, E.; Willomitzer, J.; and Strouhal, Z., 1977, *Veterinarstvi*, v. 27 (10), 460-462
Fasciola hepatica, wild ruminants, prevalence, BHS treatment, Frescon for snail control
- Molluscicides
Chu, K. Y., 1978, *Bull. World Health Organ.*, v. 56 (2), 313-322
Schistosoma haematobium, snail control trials by modification of habitat and application of niclosamide: Volta Lake village, Ghana

Molluscicides

- Contreras B., J. A., 1976, Vet. Med. Rev. (2), 190-195
fascioliasis, cattle, high incidence of abortions reduced substantially and permanently following bilevon R treatment and snail control programme using copper sulphate: Venezuela

Molluscicides

- Cotteleer, C.; and Fameree, L., 1976, Vet. Med. Rev. (2), 196-202
fascioliasis, cattle, bilevon-R treatment combined with prophylactic decontamination of pastures with frescon evaluated during 3-year control scheme

Molluscicides

- El Kheir, Y. M.; and El Tohami, M. S., 1979, J. Trop. Med. and Hyg., v. 82 (11-12), 237-241
molluscicidal activity of certain Sudanese plants evaluated, preliminary screening of 78 plants used in folk-medicine

Molluscicides

- El Kheir, Y. M.; and El Tohami, M. S., 1979, J. Trop. Med. and Hyg., v. 82 (11-12), 242-247
molluscicidal activity of certain Sudanese plants evaluated, screening of *Gnidia kraussiana* Meisn leaf, stem and root

Molluscicides

- Gardner, D. R.; and Moreton, R. B., 1978, Pesticide Biochem. and Physiol., v. 8 (3), 225-240
disinhibition caused by the molluscicide N-trityl morpholine in the isolated nervous system of *Lymnaea stagnalis*

Molluscicides

- Goel, H. C.; and Prasad, R., 1978, Indian J. Exper. Biol., v. 16 (5), 620-622
Indoplanorbis exustus (vector of animal schistosomiasis), susceptibility of eggs to certain molluscicides, suggested that snail control might be accomplished more economically and with less damage to ecosystem if low concentrations of molluscicides were directed primarily against eggs rather than high concentrations against adult snails

Molluscicides

- Hanumante, M. M.; and Kulkarni, S. S., 1979, Bull. Environment. Contam. and Toxicol., v. 23 (6), 725-727
acute toxicity of mercuric chloride and pentachlorophenol to *Channa gachua*

Molluscicides

- Jobin, W. R., 1979, Am. J. Trop. Med. and Hyg., v. 28 (1), 142-154
schistosomiasis control, 6 pilot projects, review with emphasis on cost of mollusciciding programs

Molluscicides

- Jordan, P.; et al., 1978, Bull. World Health Organ., v. 56 (1), 139-146
Schistosoma mansoni, evaluation of experimental mollusciciding program to control transmission: St. Lucia

Molluscicides

- Medina, F. R.; and Woodbury, R., 1979, J. Agric. Univ. Puerto Rico, v. 63 (3), 366-376
Lymnaea spp., molluscicidal assay of 200 Puerto Rican terrestrial plants, possible use in biological control of snail vectors of fascioliasis hepatica

Molluscicides

- Over, H. J.; et al., 1977, Tijdschr. Diergeneesk., v. 102 (5), 304-317
Fasciola hepatica, control of intermediate host, *Lymnaea truncatula*, by Frescon (N-tritylmorpholine), field conditions trial, effectiveness determined by snail population measurements and infections of tracer lambs, particularly effective when used in spring

Molluscicides

- Pellegrino, J.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (1), 10-15
molluscicidal activity of 2,5-bis(1-aziridinyl)-p-benzoquinone on *Biomphalaria* snails, very effective in field trials

Molluscicides

- Purchio, A.; and Campos, R., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (4), 236-238
schistosomiasis snail vectors, molluscicidal activity of aflatoxin B-1 in *Aspergillus parasiticus* extract

Molluscicides

- Shiff, C. J., 1970, South African Med. J., v. 44 (7), 167-168
bilharzia control, for rapid reduction of prevalence of parasites, molluscicides are best combined with improved hygiene and sanitation, and suppressive chemotherapy: Africa

Molluscicides

- Shiff, C. J.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (4), 375-380
Schistosoma haematobium, seasonal patterns in transmission, epidemiology in school children, control by winter application of molluscicides: Rhodesia

Molluscicides

- Shoeb, H. A.; and El-Emam, M. A., 1976, Egypt. J. Bilharz., v. 3 (2), 157-167
screening of active chemical constituents of *Ambrosia maritima* for possible molluscicidal activity against snail vectors of human schistosomiasis

Molluscicides

- de Sousa, M. P.; and Rouguayrol, M. Z., 1974, Rev. Brasil. Pesqui. Med. e Biol., v. 7 (4), 389-393
numerous plants native to northeastern Brazil tested for molluscicidal activity against snail vectors of human schistosomiasis

Molluscicides

- Sullivan, J. T.; and Cheng, T. C., 1976, J. Invert. Path., v. 28 (2), 255-257
Biomphalaria glabrata, comparative toxicity of internal (injected) copper and equivalent external concentration

Molluscicides

- Sullivan, J. T.; and Palmieri, J. R., 1979, J. Parasitol., v. 65 (1), 50-54
Echinostoma audyi, effect of duration and intensity of infection on survival of *Lymnaea rubiginosa* exposed to copper sulfate

Molluscicides

- Uhazy, L. S.; Tanaka, R. D.; and MacInnis, A. J., 1978, Science (4359), v. 201, 924-926
Schistosoma mansoni, identification of chemicals that attract or trap its snail vector, *Biomphalaria glabrata*, results indicate that it should be possible to formulate slow- or no-release molluscicides coupled with controlled-release attractants, may attract and kill larval schistosomes as well as snail vectors

Molluscicides

- Willomitzer, J., 1977, Acta Vet. Brno, v. 46 (1-2), 167-174
 molluscicidal efficacy of lastanox, powdered colophony, macerates of spruce and fir needles, and 2'-chloro-4-fluoro-4'-nitrosalicylanilide, toxic effects on aquarium fish

Molluscicides

- Willomitzer, J.; Chroustova, E.; and Kolar, Z., 1976, Veterinarstvi, v. 26 (9), 417-419
Fascioloides magna, deer, control with BHS for deer and Frescon as molluscicide for snail control

Monensin -- Coban; COBAN 45; Elancoban; Elancoban Premix; Monensin sodium; Romensin; Rumensin.

Monensin (Elancoban)

- Bedrnik, P.; et al., 1979, Arch. Geflugelk., v. 43 (1), 7-10
Eimeria spp., chickens (exper.), comparative efficiency of anticoccidials in combination with growth promotant nitrovin

Monensin (Elancoban)

- Bedrnik, P.; et al., 1979, Veterinarstvi, v. 29 (8), 353-355
 coccidiosis, broiler chicken fattening, lasalocid compared with amprol plus and monensin: Czechoslovakia

Monensin

- Chapman, H. D., 1978, Avian Path., v. 7 (2), 269-277
Eimeria maxima, *E. brunetti*, *E. tenella*, chickens, effect of monensin on development of immunity acquired by repeated low-level infections

Monensin

- Chapman, H. D., 1979, Avian Path., v. 8 (2), 181-186
Eimeria maxima in chickens (exper.), sensitivity to monensin of field isolates from monensin-exposed vs. unexposed sites, low level resistance may have developed as a result of previous drug exposure

Monensin

- Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (2), 304-307
Eimeria spp., broilers (exper.), salinomycin, monensin, lasalocid, drug toleration and anticoccidial efficacy compared in 5 field trials, commercial facilities

Monensin -- Continued.

Monensin sodium (Romensin)

- Collins, E. A.; and McCrea, C. T., 1978, Vet. Rec., v. 103 (17), 386 [Letter]
 monensin sodium, cattle, fatal incident following accidental overdosage

Monensin sodium (COBAN 45)

- Cruthers, L. R.; et al., 1978, Poultry Science, v. 57 (5), 1227-1233
Eimeria spp., broiler chicks (exper.), lonomycin in feed, high degree of anticoccidial activity, comparison with lasalocid and monensin

Monensin

- Dykstra, D. D.; and Reid, W. M., 1978, Poultry Science, v. 57 (2), 398-402
Eimeria tenella, gnotobiotic chickens, monensin suppressed bacterial numbers in absence of *Eimeria*, prevented increase in bacterial numbers in presence of *Eimeria*

Monensin (Rumensin)

- Fitzgerald, P. R.; and Mansfield, M. E., 1978, Am. J. Vet. Research, v. 39 (1), 7-10
Eimeria spp., lambs (nat. and exper.), effect of monensin on oocyst discharge, feed consumption, fecal output, and weight gain (by sex of host): Illinois

Monensin sodium (Rumensin)

- Foreyt, W. J.; Gates, N. L.; and Wescott, R. B., 1979, Am. J. Vet. Research, v. 40 (1), 97-100
Eimeria spp., confinement-reared lambs (exper.) from weaning to market weight, monensin and lasalocid highly effective in eliminating oocysts, weight gains and feed conversion rates measured

Monensin

- Greuel, E.; and Kuehnhold, W., 1977, Prakt. Tierarzt, v. 58 (5), 338-341
Eimeria spp., chickens (exper.), synergistic effect of metichlorpindol and methylbenzoate, rotation program with other coccidiostats discussed

Monensin (Rumensin)

- Horton, G. M. J.; and Stockdale, P. H. G., 1979, Am. J. Vet. Research, v. 40 (7), 966-970
Eimeria spp., lambs, amprolium and monensin, oocyst discharge, feed utilization, and ruminal metabolism

Monensin

- Jeffers, T. K., 1978, Avian Dis., v. 22 (1), 157-161
Eimeria tenella isolants obtained from monensin-treated flocks, monensin sensitivity tests revealed no monensin-resistant isolants

Monensin-sodium (Elancoban; Rumensin)

- Kaemmerer, K.; and Fink, J., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (6), 216-220
 monensin-sodium, lasalocid, salinomycin, influence of high dosages on heart of chickens

Monensin -- Continued.

Monensin

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Monensin sodium (Elancoban Premix)

Kemp, J., 1978, Vet. Rec., v. 102 (21), 467
 [Letter]
 monensin sodium toxicity from turkey starter ration in young guinea fowl keats

Monensin

Kutzer, E.; et al., 1979, Wien. Tierarztl. Monatsschr., v. 66 (6-7), 197-202
Eimeria spp., broiler chickens, arprinocid, drug efficacy compared with monensin

Monensin

Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Monensin (Elancoban)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
 coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Monensin

Long, P. L.; Millard, B. J.; and Smith, K. M., 1979, Avian Path., v. 8 (4), 453-467
Eimeria spp., chickens, effect of 4 anticoccidial drugs on development of immunity, field and laboratory conditions

Monensin (Rumensin)

McDougald, L. R., 1978, Am. J. Vet. Research, v. 39 (11), 1748-1749
Eimeria bovis, Holstein-Friesian calves (exper.), monensin added to complete feed ration as preventive medication, control excellent

Monensin (Rumensin)

McDougald, L. R.; and Dunn, W. J., 1978, Am. J. Vet. Research, v. 39 (9), 1459-1462
Eimeria spp., lambs (exper.), efficacy of monensin in feed

Monensin

McDougald, L. R.; and Galloway, R. B., 1977, Ztschr. Parasitenk., v. 54 (1), 95-100
Eimeria tenella in vitro, development inhibited by serum from chickens fed anticoccidial drugs, technique to assay drug activity and to characterize and quantitate therapeutic effect

Monensin

McDougald, L. R.; Karlsson, T.; and Reid, W. M., 1979, Avian Dis., v. 23 (4), 999-1005
 coccidiosis, chickens (exper.), natural outbreak of infectious bursal disease (IBD) during comparison of anticoccidials for their effect on development of immunity, interaction between diseases, immunity to coccidiosis not blocked by IBD

Monensin -- Continued.

Monensin

McDougald, L. R.; and McQuistion, T. E., 1978, Avian Dis., v. 22 (4), 765-770
Eimeria spp., turkeys, coccidiosis management, innate (age) resistance and acquired immunity vs. anticoccidial medication

Monensin (Coban; Elancoban)

McQuistion, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Monensin sodium

Malone, P., 1978, Vet. Rec., v. 103 (21), 477-478
 cattle, nitrite poisoning attributed to feeding monensin sodium, diet of barley straw and grazing turnips, treatment by intravenous injection of methylene blue

Monensin

Meingassner, J. G.; et al., 1979, Poultry Science, v. 58 (2), 308-313
Eimeria tenella, broilers (exper.), monensin, lasalocid, anticoccidial activity enhanced when combined with tiamulin, tiamulin alone was inactive; metabolic fate of monensin and tiamulin separately or in combination monitored using isolated, perfused rat liver

Monensin

Migaki, T. T.; and Babcock, W. E., 1979, Poultry Science, v. 58 (2), 481-482
 salinomycin, anticoccidial, safety in broiler chickens compared with monensin

Monensin

Migaki, T. T.; Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (5), 1192-1196
Eimeria spp., chicks (exper.), salinomycin, monensin, lasalocid, efficacy in battery trials

Monensin

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Monensin (Coban)

Mogollon, J. D.; and Molina, T. de P., 1977, Rev. Inst. Colomb. Agropec, v. 14 (4), 473-488
Eimeria tenella field strain, sensitivity against 3 anticoccidial drugs

Coban

Morrison, W. D.; Ferguson, A. E.; and Leeson, S., 1979, Poultry Science, v. 58 (5), 1160-1166
Eimeria spp., chicks (exper.), salinomycin and stenorol compared with other anticoccidials, efficacy and effect on chick performance

Monensin -- Continued.

Monensin

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Monensin

Onaga, H.; et al., 1978, Nippon Zyuisi-Kai Zasshi (J. Japan Vet. Med. Ass.), v. 31 (10), 592-596
Eimeria spp., chicks and cell cultures, monensin

Monensin

Ordidge, R. M.; Schubert, F. K.; and Stoker, J. W., 1979, Vet. Rec., v. 104 (16), 375 [Letter]
 death of horses accidentally fed cattle feed supplement containing monensin

Monensin

Reid, W. M.; Anderson, W. I.; and McDougald, L. R., 1978, Avian Path., v. 7 (4), 569-576
Eimeria spp., turkey poults, anticoccidial protection and development of immunity while using monensin

Monensin

Ruff, M. D., 1978, Proc. 1978 Maryland Nutrition Conf. Feed Mfr. (University of Maryland, Mar. 16-17), 32-37
Eimeria spp., chickens, anticoccidials, safe withdrawal times, effect on nutrient malabsorption

Monensin

Ruff, M. D.; et al., 1979, Poultry Science, v. 58 (2), 298-303
Eimeria spp., battery raised broilers (exper.), narasin compared with monensin

Monensin

Ryley, J. F.; and Hardman, L., 1978, Parasitology, v. 76 (1), 11-20
Eimeria spp., chicks (exper.), effects of dietary vitamin K on severity of disease with particular attention to effects of vitamin K on response to anticoccidial drugs, concluded that use of vitamin K deficient diet for experimental work is quite justified

Monensin

Samizadeh-Yazd, A.; et al., 1979, Am. J. Vet. Research, v. 40 (8), 1107-1109
Eimeria spp., lambs, efficacy of monensin and aureomycin separately and combined

Monensin

Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazin, and pancoxin: England; France; Germany

Monesin -- Continued.

Elancoban

Sherkov, Sh.; et al., 1978, Vet. Sbirka, v. 76 (6), 39-41
 coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat

Monensin

Stockdale, P. H. G.; and Yates, W. D. G., 1978, Vet. Parasitol., v. 4 (3), 209-214
Eimeria zuernii, calves (exper.), successful chemotherapy with amprolium or monensin, resistance to reinfection after chemotherapy

Monensin (Elancoban)

Stuart, J. C., 1978, Vet. Rec., v. 102 (14), 303-304
 monensin toxicity in turkeys aged 25 weeks and older

Monensin (Elancoban)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
 coccidiosis, broilers, anticoccidials, floor pen trials

Monensin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Monensin sodium. See Monensin.

Monocrotophos -- Nuvacron.

Nuvacron

El-Bahay, G. H.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 205-213
 Hyalomma dromedarii, Argas persicus, evaluation of 10 insecticides

Monomycine. See Paromomycin.

Monopar. See Stilbazium iodide.

Monosulfiram. See Sulfirame.

Morantel -- Banminth II; Banminth D (with Diethyl-carbamazine); Exhelm-E; 3-Methyl pyrantel tartrate; trans-2(2-(3-Methyl-2-thienyl)vinyl)-1-methyl-1,4,5,6-tetrahydropyrimidine tartrate; Morantel citrate; Morantel tartrate; Ovithelm.

Banminth II

Bali, M. K.; and Singh, R. P., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 230-232
Haemonchus contortus, in vitro larvicidal effects of fenbendazole, banminth II, and nemafox

Morantel (Banminth 2)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
Haemonchus contortus, Trichostrongylus colubriformis, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Morantel tartrate (Banminth II)

Chandrasekharan, K.; et al., 1973, Kerala J. Vet. Sc., v. 4 (2), 193-195
gastrointestinal nematodes, zoo animals, morantel tartrate: Trichur Zoo, Kerala

Morantel tartrate (Banminth II)

Chandrasekharan, K.; Sundaram, R. K.; and Peter, C. T., 1973, Kerala J. Vet. Sc., v. 4 (1), 59-62
gastrointestinal helminths, calves and kids, morantel tartrate

Ovithelm

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Morantel tartrate

Jones, R. M.; Pott, J. M.; and Cornwell, R. L., 1978, Brit. Vet. J., v. 134 (2), 166-170
Trichostrongylus colubriformis, lambs; Oster-tagia ostertagi, Dictyocaulus viviparus, calves, low level feed administration of morantel tartrate, good results

Banminth D

Kadyrov, N. T., 1978, Veterinariia, Moskva (7), 57-58
Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Morantel tartrate (Banminth II)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Morantel tartrate (Banminth II)

Lai, M.; et al., 1978, Clin. Vet., Milano, v. 101 (3), 123-130
gastro-intestinal strongyles, sheep, efficacy of morantel-tartrate

Morantel tartrate

Le Jambre, L. F.; and Martin, P. J., 1979, Vet. Sc. Commun., v. 3 (2), 153-158
levamisole resistant Oster-tagia circumcincta and O. trifurcata, sheep, cross resistant to morantel tartrate but not to naphthalophos

Morantel -- Continued.

Morantel tartrate

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Austral. Vet. J., v. 54 (12), 570-574
Trichostrongylus colubriformis, linear dose responses of selected and unselected strains to thiabendazole, levamisole, and morantel tartrate

Morantel tartrate

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Internat. J. Parasitol., v. 8 (6), 443-447
Ostertagia circumcincta, development of simultaneous resistance to thiabendazole, morantel tartrate, and levamisole, multiple selection associated with increase in O. trifurcata in population and increase in larval inhibition

Morantel tartrate

Martin, P. J.; and Le Jambre, L. F., 1979, Vet. Sc. Commun., v. 3 (2), 159-164
Ostertagia circumcincta, O. trifurcata, larval paralysis as in vitro assay of levamisole and morantel tartrate resistance

Banminth II

Nayar, K. N. M.; et al., 1978, Kerala J. Vet. Sc., v. 9 (1), 35-37
Gnathostoma spinigerum, dog, case report, treatment with banminth II not effective, successful surgical removal: India

Morantel tartrate

Oripov, A. O., 1978, Veterinariia, Moskva (4), 74-76
Strongylata, sheep, various anthelmintics in granular form tested for mass dehelminthization, no harmful effects, nilverm most effective

Morantel tartrate

Pott, J. M.; Jones, R. M.; and Cornwell, R. L., 1979, Internat. J. Parasitol., v. 9 (2), 153-157
parasitic gastroenteritis and bronchitis in grazing calves, effect of low level feed incorporation of morantel in early season

Morantel tartrate (Banminth-II)

Retnasabapathy, A.; and Baskaran, G., 1976, v. 53 (10), 806-811
ancylostomiasis, dogs, morantel tartrate, pyrantel pamoate and disophenol effective

Morantel citrate

Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235
Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parabendazole and dichlorvos

Morantel tartrate

Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235
Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parabendazole and dichlorvos

Morantel -- Continued.

Morantel tartrate (Exhelm-E)

Sangster, N. C.; et al., 1979, Research Vet. Sc., v. 27 (1), 106-110

Trichostrongylus colubriformis, Ostertagia circumcincta, Merino and crossbred sheep, field observations and preliminary critical trials showed varying degrees of drug resistance to levamisole hydrochloride, morantel tartrate, and thiabendazole; differences in infectivity and drug efficacy between breeds: Australia

Morantel tartrate (Banminth II)

Sarup, S.; Banerjee, D. P.; and Gautam, O. P., 1978, Indian Vet. J., v. 55 (6), 497-498

Ancylostoma caninum, ova and infective larvae, in vitro evaluation of fenbendazole, helatac, alcopar, banminth II, only banminth II effective; banminth II-treated larvae administered orally to mice, none recovered from lungs or liver

Morantel tartrate (Exhelm E)

Stockdale, P. H. G.; and Harries, W. N., 1979, Canad. Vet. J., v. 20 (9), 223-226

gastrointestinal nematodes, cattle, anthelmintic treatment upon entry into feedlot, no improvement in rate of weight gain or feed conversion efficiency: Alberta

Banminth-II (Morantel tartrate)

Vihan, V. S.; and Sahni, K. L., 1979, Indian J. Animal Research, v. 13 (1), 15-18

gastrointestinal nematodes, lambs, efficacy of helatac, nilverm, thiabendole, and banminth-II: Makhdoom, Mathura (U. P.)

Morantel citrate. See Morantel.

Morantel tartrate. See Morantel.

6-(2-Morpholinoethoxy)-2-(5-nitro-1-methyl-2-imidazolyl-methylene)-tetralone sulphate

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

levo-5-Morpholinomethyl-3-(5-nitrofurylidene-amino)-2-oxasolidinone. See Furaltadone.

1-(3-Morpholinopropyl)-2-(5-nitro-1-methyl-2-imidazolyl)-benzimidazole

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Moxipraquine -- 349C59

349C59

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Moxnidazole -- 3-(5-Nitro-1-methyl-2-imidazolyl-methylene-amino)-5-morpholino-methyl-2-oxasolidinone hydrochloride.

Moxnidazole

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Multispec. See Mebendazole.

Mycodex Pet Shampoo (containing pyrethrins)

McKeever, P. J.; and Allen, S. K., 1979, J. Am. Vet. Med. Ass., v. 174 (7), 718-720

Cheyletiella spp., cats, pruritic dermatitis, pyrethrins, malathion 57

- Nafoxidine hydrochloride
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Naganol. See Suramin.
- Naled
Frazar, E. D.; and Schmidt, C. D., 1979, *J. Econom. Entom.*, v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides
- Naled
Rawlins, S. C.; and Mansingh, A., 1978, *J. Econom. Entom.*, v. 71 (1), 142-144
Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana
- Naled
Rawlins, S. C.; and Mansingh, A., 1978, *J. Econom. Entom.*, v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides
- Naled
Rawlins, S. C.; and Mansingh, A., 1979, *J. Econom. Entom.*, v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies
- NaP. See Sodium antimony dimethylcysteine tartrate.
- 1-Naphthalenol methylcarbamate. See Carbaryl.
- Naphthalophos. See Phthalophos.
- Naphthamon. See Bephenium.
- Naphthenate complex
Niazov, A. N.; et al., 1977, *Izvest. Akad. Nauk Turkmen. SSR, s. Biol. Nauk* (6), 68-70
ixodid ticks, naphthenates tested as acaricides
- 1,4-Naphthoquinone
Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones
- 1,4-Naphthoquinone
Docampo, R.; et al., 1978, *Ztschr. Parasitenk.*, v. 57 (3), 189-198
Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages
- Naphthoquinones
Boveris, A.; et al., 1978, *Comp. Biochem. and Physiol.*, v. 61C (2), 327-329
Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones
- Naphthoquinones
Korolkovas, A.; and Pellegrino, J., 1978, *Rev. Inst. Med. Trop. S. Paulo*, v. 20 (4), 219-223
Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active
- Naphthoquinones
Meshnick, S. R.; et al., 1978, *J. Exper. Med.*, v. 148 (2), 569-579
Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents
- 1,4-Naphthylenediamines
Korolkovas, A.; and Pellegrino, J., 1978, *Rev. Inst. Med. Trop. S. Paulo*, v. 20 (4), 219-223
Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active
- 1,4-Naphthylenediamine salts
Korolkovas, A.; Gloria, T. S.; and Haraguchi, T., 1973, *Rev. Farm. e Bioquim. Univ. Sao Paulo*, v. 11 (2), 247-254
Schistosoma mansoni, study of 3 new naphthylenediamine salts (m-chlorobenzoate; 3,5-dihydroxybenzoate; 2-hydroxycinchoninate) with in vitro schistosomicidal activity
- 1-Naphthyl methyl carbamate. See Carbaryl.
- 1,8-Naphthyridine derivatives
Suzuki, N.; Kato, M.; and Dohmori, R., 1979, *J. Pharm. Soc. Japan*, v. 99 (2), 155-164
Trichomonas vaginalis, synthesis and activity of 1,8-naphthyridine derivatives
- Narasin
Karlsson, T.; and Reid, W. M., 1978, *Avian Dis.*, v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis
- Narasin
Ruff, M. D.; et al., 1979, *Poultry Science*, v. 58 (2), 298-303
Eimeria spp., battery raised broilers (exper.), narasin compared with monensin
- Narasin (MW 764)
Smith, C. K. II; and Strout, R. G., 1979, *Exper. Parasitol.*, v. 48 (3), 325-330
Eimeria tenella, accumulation and retention of lasalocid and narasin by extracellular sporozoites

Natamycin. See Pimaricin.

Navadel. See Dioxathion.

Naxogin. See Nitrimidazine.

Naxogyn. See Nitrimidazine.

Neguvon. See Trichlorfon.

Nemafax. See Thiophanate.

Nemasole. See Mebendazole.

Nematin

Lipova, E.; and Zajicek, D., 1979, Veterinarstvi, v. 29 (3), 125-126
Strongyloides papillosus, calves, intensity of infection, clinical aspects, nilverm and nematin treatment

Nemicide. See Tetramisole.

Nemicide L 15. See Tetramisole.

Neoarsphenamine -- Neosalvarsan; Novarsenol.

Novarsenol

Lochkarev, V. A.; and Evpak, I. D., 1977, Veterinariia, Moskva (6), 64-65
E[imeria] zurni, E. smithi, calves, novarsenol effective

Neosalvarsan

Youssef, A. H., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 147-157
Dipetalonema evansi, camels, filarial orchitis and possible significance as prevalent reproductive disease; surgical treatment and use of neosalvarsan, foudadin, and neguvon, histopathology of gonads: Egypt

Neocidol. See Diazinon.

Neomycin -- Canaural (with Nystatin and Prednisolone); Framycetin; Neosporin (with Polymyxin B); Tresaderm (with Dexamethasone and Thiabendazole).

Tresaderm

Faulk, R. H.; and Schwirck, S., 1978, Vet. Med. and Small Animal Clin., v. 73 (3), 307-308
Otodectes cynotis, dogs, cats (ears of both), tresaderm, clinical trial, highly effective

Neomycin -- Continued.

Neosporin

Nagington, J., 1975, Tr. Ophth. Soc. United Kingdom, v. 95 (2), 207-209
Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity

Canaural

Pott, J. M.; and Riley, C. J., 1979, Vet. Rec., v. 104 (25), 579
Otodectes cynotis, dogs, cats (ear canals of both), canaural, comparison with proprietary topical ear preparation (clendrol), controlled trial

Neo neo arsemin

Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Neosalvarsan. See Neoarsphenamine.

Neo-scabexaan. See Benzene hexachloride.

Neoscabexan. See Benzene hexachloride.

Neosporin. See Neomycin or Polymyxin B.

Neostibosam. See Ethylstibamine.

Nerium oleander herb

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, Scientia Pharm., v. 47 (2), 114-118
Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

Nestyne. See Dankil.

Nevaquine

Sharma, M. M.; et al., 1979, Indian Vet. Med. J., v. 3 (3), 187-191
theileriasis, Jersey cattle, incidence and treatment with berenil, babesan, nevaquine, and aureomycin: Exotic Nucleus Cattle Farm, Bassi, Jaipur

Nexagan. See Bromophos-ethyl.

Nicarb. See Nicarbazin.

Nicarbazine -- Nicarb; Nicrazin.Nicarbazine (Nicrazin)

Buys, S. B.; and Rasmussen, R. W., 1978, J. South African Vet. Ass., v. 49 (2), 127-128 significantly higher mortalities found in nicarbazine fed chickens than in amprolium fed chickens when exposed to same heat stress conditions

Nicarbazine

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236
coccidostats in feeds, qualitative identification test

Nicarbazine

Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Nicarbazine (Nicarb)

McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Nicarbazine

Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, Rev. Brasil. Pesqui. Med. e Biol., v. 2 (1), 45-49
Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

Nicarbazine

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250
Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Nicarbazine

Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazine, and pancosin: England; France; Germany

Nicarbazine

Sevcik, B.; et al., 1974, Veterinaria, Praha, v. 16 (5-6), 421-588
coccidiosis; nicarbazine, clopidol, efficacy and toxicity in various animals, review

Niclofolan -- Bayer 9015; Bilevon-M; Bilevon-R; Dertil; Dertil B; Dertil O; 5,5'-Dichlor-3,3'-dinitro-biphenyl-2,2'-diol; 5,5'-Dichloro-2,2',dihydroxy-3,3'-dinitro-phenyl; 3,3'-Dichloro-5,5'-dinitro-0,0'-biphenol; Distolon; ME 3625; Menichlopholan.Niclofolan -- Continued.Bilevon R

Contreras B., J. A., 1976, Vet. Med. Rev. (2), 190-195
fascioliasis, cattle, high incidence of abortions reduced substantially and permanently following bilevon R treatment and snail control programme using copper sulphate: Venezuela

Bilevon [R] (Niclofolan; Menichlopholan)

Corba, J., 1978, Vet. Med. Rev. (1), 77-81
Fasciola hepatica, cattle, sheep, bilevon injection highly effective and well tolerated; not effective against Dicrocoelium spp.

Bilevon [R] (Niclofolan; Menichlopholan)

Corba, J., 1978, Vet.-Med. Nachr. (1), 77-81
Fasciola hepatica, cattle, sheep, bilevon injection

Bilevon-R (Niclofolan)

Cotteleer, C.; and Fameree, L., 1976, Vet. Med. Rev. (2), 196-202
fascioliasis, cattle, bilevon-R treatment combined with prophylactic decontamination of pastures with frescon evaluated during 3-year control scheme

Dertil O

Daniiarov, I. A.; et al., 1978, Veterinaria, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Bilevon R

Daniiarov, I. A.; et al., 1978, Veterinaria, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Distolon (Bilevon-M; Bayer 9015; ME 3625)

Furmaga, S.; Gundlach, J. L.; and Sobieszewski, K., 1974, Acta Parasitol. Polon., v. 22 (12-21), 229-247
Fasciola hepatica-infected and uninfected sheep treated and not treated with 3 fasciolicides (fasciolin, distolon, and zanil), blood serum levels of Ca, P, Na, K, and Mg

Menichlopholan

Hamajima, F.; et al., 1979, Internat. J. Parasitol., v. 9 (3), 241-249
Clonorchis sinensis, Metagonimus takahashii, Paragonimus miyazakii, in vitro effects of bithionol and menichlopholan on motility, metabolism, and fine structure

Dertil B

Merényi, L., 1978, Magy. Allat. Lapja, v. 100, v. 33 (2), 132-134
fascioliasis, control, treatment: Nograd county

Dertil O

Merényi, L., 1978, Magy. Allat. Lapja, v. 100, v. 33 (2), 132-134
fascioliasis, control, treatment: Nograd county

Niclofolan -- Continued.

Bilevon [R]

Parshad, V. R.; and Guraya, S. S., 1978, Vet. Parasitol., v. 4 (2), 111-120

4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces

Bilevon [R]

Qadir, A. N. M. A., 1979, Indian Vet. J., v. 56 (5), 429-431

Fasciola gigantica, goats, acedist, comparison with bilevon: Bangladesh

5,5'-Dichloro-3,3'-dinitro-2,2'-biphenyldiol

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Dertil

Timofeev, B. A.; et al., 1978, Veterinariia, Moskva (4), 67-68

Fasciola hepatica, cattle, disalan effective, no toxic effect; comparison of anthelmintic effect with dertil

Dertil

Vishniauskas, A. I., 1978, Magy. Allat. Lapja, v. 100, v. 33 (8-9), 551-552

Fasciola, sheep (exper.), dertil injectable

Dertil

Vishniauskas, A.; and Rudaitis, A., 1978, Veterinariia, Moskva (4), 68-69

Fasciola hepatica, cattle (exper.), acemidophene, dertil and sulfene compared, acemidophene insufficiently effective

Niclofolan (Bilevon [R])

Voegel, H., 1977, Vet.-Med. Nachr. (2), 125-128
niclofolan, development of injectable solution

Bilevon [R] (Niclofolan; Dertil)

Ziegler, K., 1979, Vet. Med., Praha, v. 52, v. 24 (7), 391-395

Fasciola hepatica, cattle, dovenix and bilevon highly effective; drug toxicity tests in rats

Niclosamide -- Bayer 2353; N-(2'-Chloro-4'-nitrophenyl)-5' chlorosalicylanilide piperazine salt; Cyclosamide; 2',5-Dichloro-4'-nitrosalicylanilide; Dichlosal (with Dichlorophene); Fenasal; Mansonil; Phenasal; Radeverm; Sagimid; Trichlosal (with Trichlorophen); Yomesan.

Niclosamide

Alaimo, R. J.; et al., 1978, J. Med. Chem., v. 21 (3), 298-300

furodazole, anthelmintic trials with experimental animals, bunamidine and niclosamide used as reference drugs

Niclosamide -- Continued.

Niclosamide

Atias, A.; and Cattán, P. E., 1976, Rev. Med. Chile, v. 104 (4), 216-217

Diphylobothrium pacificum, ova in feces of boy who had eaten raw fish, tapeworm recovered after therapy with niclosamide, child had frequented beach where similar tapeworm had been recovered earlier from sea lion, first reported case in Chile

Cyclosamide

Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36

cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results

Phenasal, α -form

Bekhli, A. F.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 228

Hymenolepis nana, mice, comparative activity of two polymorphic forms of phenasal

Phenasal, β -form

Bekhli, A. F.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 228

Hymenolepis nana, mice, comparative activity of two polymorphic forms of phenasal

Mansonil

Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91

cestodes, pathomorphology resulting from action of various anthelmintics

Phenasal

Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91

cestodes, pathomorphology resulting from action of various anthelmintics

Sagimid

Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91

cestodes, pathomorphology resulting from action of various anthelmintics

Mansonil

Bogdanov, V. R.; Koshkina, N. G.; and Logachev, E. D., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 92-93

Dipylidium caninum, various anthelmintics, in vitro action on surface tissues and inactivation of enzymes

Phenasal

Bogdanov, V. R.; Koshkina, N. G.; and Logachev, E. D., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 92-93

Dipylidium caninum, various anthelmintics, in vitro action on surface tissues and inactivation of enzymes

Niclosamide

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15

antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Niclosamide -- Continued.

- Fenasal
Daniiarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective
- Sagimid
Daniiarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective
- Phenasal
Dovzhenko, V. A.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 169-173
Diphyllobothrium latum, Hymenolepis nana, Taeniarrhynchus saginatus, phenasal effects on various parts of strobila
- Niclosamide + Bunamidine hydrochloride
Gemmell, M. A., 1978, Bull. World Health Organ., v. 56 (3), 433-443
tapeworm control in dogs for prevention of hydatidosis and cysticercosis in sheep, monthly drug treatment program, age-specific prevalence of Taenia hydatigena in lambs used as principal indicator, 13-year assessment: Styx Valley and Maniototo County, South Island, New Zealand
- Phenasal + Dichlorophene (= Dichlosal)
Grinenko, N. V.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (1), 101-103
H[ymenolepis] nana, patients of different age groups, dichlosal or trichlosal in divided doses, 5 or 7 day courses
- Phenasal + Trichlorophen (= Trichlosal)
Grinenko, N. V.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (1), 101-103
H[ymenolepis] nana, patients of different age groups, dichlosal or trichlosal in divided doses, 5 or 7 day courses
- Dichlosal
Grinenko, N. V.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasal, trichlorophene, dichlosal, and trichlosal tested in graded doses
- Phenasal
Grinenko, N. V.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasal, trichlorophene, dichlosal, and trichlosal tested in graded doses
- Trichlosal
Grinenko, N. V.; et al., 1976, Med. Parazitol. i Parazitar. Bolezni, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasal, trichlorophen, dichlosal, and trichlosal tested in graded doses
- Niclosamide (Radeverm)
Haak, W.; et al., 1972, Ang. Parasitol., v. 13 (4), 200-207
cestodes of humans, recommended drug therapy

Niclosamide -- Continued.

- Niclosamide (Yomesan)
Hira, P. R., 1975, Ann. Soc. Belge Med. Trop., v. 55 (4), 321-326
Inermicapsifer madagascariensis, children, case reports, niclosamide, wild rodents as reservoir hosts: Zambia
- Phenasal
Iashchuk, V. D.; and Vasil'kov, G. V., 1977, Veterinariia, Moskva (7), 62-65
[Bothriocephalus], carp, phenasal-feed mix as control measure in fish farms, economic effectiveness analyzed mathematically
- Niclosamide
Jaroovesama, N.; and Harinasuta, T., 1972, Siriraj Hosp. Gaz., v. 24 (7), 1095-1099
taeniasis, human, comparative treatment trials using quinacrine and niclosamide
- Niclosamide (Yomesan)
Jones, W. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (2), 300-302
Hymenolepis diminuta, Dipylidium caninum, man, niclosamide, effective, relatively non-toxic drug for initial therapy
- Fenasal (Bayer 2353, Yomesan)
Kashnikov, A. A., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 140-142
macracanthorhynchosis, swine, action of various anthelmintics
- Fenasal
Klenov, A. P., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 152-155
[Bothriocephalus], white amur, effectiveness of various anthelmintics
- Yomesan
Komma, M. D.; and Santos, V. L. V., 1972, Rev. Patol. Trop., v. 1 (1), 69-72
Taenia solium, T. saginata, humans, comparative therapeutic trials using yomesan and pumpkin seed based taeniafuge; correlation of species of Taenia with type of therapy and recovery of scolex with therapy
- Phenasal + Nilverm
Oripov, A. O.; Bekirov, R. E.; and Dzhumaev, Z., 1978, Veterinariia, Moskva (12), 60
helminths, dogs, phenasal and nilverm given in feed (sausage form)
- Mansonil
Parshad, V. R.; and Guraya, S. S., 1978, Vet. Parasitol., v. 4 (2), 111-120
4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces
- Niclosamide (Yomesan)
Remfry, J., 1978, Lab. Animals, v. 12 (4), 213-218
helminth infections in imported Macaca mulatta, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Niclosamide -- Continued.

Niclosamide

Rizzo, G.; and Ricciardi, G., [1978], Riv. Parassitol., Roma, v. 38 (2-3), 1977, 303-306
Diphyllobothrium latum, woman, case report, niclosamide, ingested broiled eels possible source: Italy

Phenasal

Rusak, L. V.; and Kovchur, V. N., 1972, Parazitologiya, Leningrad, v. 6 (1), 85-87
Hymenolepis nana, glycogen content in parasite tissues decreased after treatment of infected mice with aminoacrichine, phenasal, or trichlorophen, implications for mechanism of drug action

2',5-Dichloro-4'-nitrosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicalidal effect of salicylanilide and bisphenol derivatives

Niclosamide

Sanford, S. E., 1978, Canad. Vet. J., v. 19 (8), 234-236
Polymorphus boschadis cause of death in mute swans, case reports, results of treatment with niclosamide were equivocal

Yomesan

Singh, H.; et al., 1978, Ztschr. Naturforsch., Sect. C, Biosc., v. 33 (5-6), 447-448
Hymenolepis nana, rats, mice, 2'-chloro-1-hydroxy-2-naphthanilide-4'-isothiocyanate, synthesis and cestodicidal activity, highly effective and safe, comparative efficacy with yomesan; further tests showed marked activity against H. diminuta in rats and Taenia sp. in dogs

Niclosamide (Yomesan)

Slocombe, J. O. D., 1979, Canad. Vet. J., v. 20 (5), 136-140
Anoplocephala perfoliata, horses, prevalence, efficacy of pyrantel pamoate, mebendazole, and niclosamide in field and critical trials: southern Ontario

Niclosamide (Yomesan)

Todd, K. S., jr.; Howland, T. P.; and Woerpel, R. W., 1978, Am. J. Vet. Research, v. 39 (2), 315-316
Mesocostoides corti, dogs (exper.), bunamidine hydrochloride and uredofos (good results), arecoline hydrobromide and niclosamide (variable results)

Fenasal

Vibe, P., 1978, Veterinariia, Moskva (5), 21-22
cestodes, sheep, fenasal highly effective for mass dehelminthiazation: southern Kazakhstan

Fenasal

Zharikov, I. S.; Orlovskii, V. I.; and Kaminiskii, I. I., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 74-77
paramphistomiasis, bovine, hexachlorparaxylol and fenasal, ineffective in chronic disease

Niclosamide -- Continued.

Yomesan

Zychowicz, C.; Kostkiewicz, M.; and Kowalczyk, S., 1975, Przegl. Epidemiol., v. 29 (2), 223-227
Hymenolepis nana, children and adult personnel of a state home for children successfully treated with yomesan after discovery that more than 57% of the children had tapeworm infections: Mragowo

Niclosulide (Bovi-douvex)

Federis, M. T.; and Tongson, M. S., 1977, Philippine J. Animal Indust., v. 32 (1-4), 94-114
Fasciola gigantica, cattle, carabaos, niclosulide, anthelmintic efficacy based on egg counts of fecal samples, less effective in young animals, no adverse side effects: Philippines

Nicotine -- Nicotine sulfate; Sonex (with Embelia ribes and Punica granatum).

Sonex

Matta, S. C.; and Ahluwalia, S. S., 1979, Indian Vet. J., v. 56 (7), 616-617
helminths, poultry, helminta-P, sonex

Nicotine sulfate. See Nicotine.

Nicrazin. See Nicarbazin.

Nifuratel -- Inimur; Macmiror; Macmiror plus (with Lysozyme and Nystatin); Magmilor; Methylmercadone; Omnes; Polmiror.

Nifuratel (Methylmercadone)

Benazet, F.; et al., 1970, Scand. J. Infect. Dis., v. 2 (2), 139-143
intestinal and hepatic parasites, nitroheterocyclic antiparasitics, laboratory studies of chemotherapeutic activity and toxicity in exper. animals

Nifuratel (Macmiror)

Campos, R., 1973, Rev. Brasil. Clin. e Terap., v. 2 (11), 587-588
Entamoeba histolytica, comparison of efficacy of nifuratel and other amoebicides using material cultured from intestinal ulcers of patient with intestinal symptomatic amoebiasis

Nifuratel (Macmiror)

de Figueiredo, J. C.; and de Figueiredo, J. A., 1973, Rev. Brasil. Clin. e Terap., v. 2 (4), 139-144
human intestinal amoebiasis, clinical trials with nifuratel, 70% cure with good tolerance and no side effects

Nifuratel -- Continued.

Nifuratel (Methylmercadone; Macmiror)

Garcia Quintero, D., 1972, Rev. Obst. y Ginec. Venez., v. 32 (3), 439-441

human vulvovaginitis resulting from trichomonal infections or mixed infections with candidiasis, clinical trials with nifuratel, drug well tolerated without severe side-effects: Venezuela

Nifuratel (Macmiror)

Martins, A. D., 1974, Rev. Brasil. Med., v. 31 (7), 503-508

human vaginal trichomoniasis alone or in the presence of fungal infections, successful therapy combining oral nifuratel and vaginal treatment with nifuratel combined with nystatin and lysozyme

Nifuratel + Nystatin + Lysozyme (= Macmiror plus)

Martins, A. D., 1974, Rev. Brasil. Med., v. 31 (7), 503-508

human vaginal trichomoniasis alone or in the presence of fungal infections, successful therapy combining oral nifuratel and vaginal treatment with nifuratel combined with nystatin and lysozyme

Nifuratel (Macmiror; Magmilor; Polmiror; Omnes; Inimur)

Rognoni, V.; and Saçone, I., 1976, Riv. Ostet. e Ginec. Prat. e Med. Perinat., v. 56 (6), 544-551

Trichomonas vaginalis, human vaginal infections, nifuratel, results of various therapeutic schemes compared, results from other studies compared, possible toxicity studied

Nifuratel

da Silva, M. A. B., 1973, Rev. Brasil. Med., v. 30 (11), 730-731

Giardia lamblia, children, nifuratel, successful clinical trials

Nifurpirinol -- Furanace; 6-Hydroxymethyl-2-(2[5-nitro-2-furyl]vinyl)pyridine.

Furanace

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Nifurtimox -- Bay 2502; Bayer 2502; Lampit; 5-Methyl-4-(5'-nitrofurfurylidene-amino)-tetrahydrate-4H-1,4-thiazine-1,1-dioxide; Nitrofurfuridine; Nitrofurfurylidene.

Lampit (Bay 2502; Bayer 2502)

Andrade, S. G.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (6), 380-389

Trypanosoma cruzi, mice, Peruvian or Colombian strain, Bay 2502, results varied with strain, suggestions for future research

Nifurtimox -- Continued.

Bay 2502

Andrade, S. G.; and Andrade, Z. A., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (4), 268-275

Trypanosoma cruzi, pathological changes in untreated vs. Bay 2502-treated mice with chronic infections

Bay 2502

Andrade, S. G.; Andrade, Z. A.; and Figueira, R. M., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (2), 124-129

Trypanosoma cruzi, Colombian strain, drug resistance, Bay 2502-treated mice

Bayer 2502

Andrade, S. G.; and Macedo, V., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (6), 421-430

Trypanosoma cruzi, mice, human, Bayer 2502, combined use of Bayer 2502 and corticoid (betamethasone) more effective than drug used alone

Lampit (Bayer 2502)

Blandon, R.; Guevara, J. F.; and Johnson, C. M., 1976, Rev. Med. Panama, v. 1 (3), 153-162

Chagas disease in children, diagnosis, pathology, therapeutic trials with various drugs, metronidazole was well tolerated and therefore most promising therapy: Panama

Nifurtimox

Brener, Z.; Costa, C. A. G.; and Chiari, C., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (6), 450-455

Trypanosoma cruzi, mice treated with nitrofurazone, nifurtimox, or Ro 7-1051, differences in susceptibility of 4 parasite strains to active drugs attributed to biological characteristics of strains rather than to mode of drug action

Bayer 2502

Cancado, J. R.; et al., 1973, Rev. Soc. Brasil. Med. Trop., v. 7 (3), 155-166

Trypanosoma cruzi, method of standardization of processes and selection of patients for drug treatment clinical trials; statistics of trial testing Bayer 2502

Nifurtimox (Lampit; Bayer 2502)

Cancado, J. R.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (2), 111-125

Trypanosoma cruzi, human chronic infections, clinical trials of nifurtimox, toxicity

Nifurtimox (Lampit)

Cancado, J. R.; et al., 1976, Rev. Goiana Med., v. 22 (3-4), 203-233

Trypanosoma cruzi, humans, extensive clinical study of nifurtimox as therapy, nifurtimox considered to be a suppressive rather than curative drug

Nifurtimox (Lampit; Bay 2502)

Cerisola, J. A.; et al., 1977, Bol. Chileno Parasitol., v. 32 (3-4), 51-62

Trypanosoma cruzi, humans, evaluation of efficacy of nifurtimox therapy using follow-up xenodiagnosis (monthly over 11-month period): Argentina, Chile, Brazil

Nifurtimox -- Continued.

Bay 2502

Cerisola, J. A.; Alvarez, M.; and De Rissio, A. M., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (6), 403-411

Trypanosoma cruzi, human, acute disease, 2-year longitudinal study, comparison of results of complement fixation, hemagglutination and fluorescent antibody tests, with and without Bay 2502 treatment

Lampit

Chapuis, Y., 1973, Ann. Soc. Belge Med. Trop., v. 53 (6), 581-593

Trypanosoma cruzi, young children, extremely severe infections with acute cardiovascular involvement and shock, lampit effective but seldom available: Cochabamba, Bolivia

Nifurtimox

Docampo, R.; and Stoppani, A. O. M., 1979, Arch. Biochem. and Biophys., v. 197 (1), 317-321

Trypanosoma cruzi epimastigotes, generation of superoxide anion and hydrogen peroxide induced by nifurtimox

Nifurtimox (Lampit)

Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455

Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity

Lampit (Bayer 2502; Nifurtimox)

Fernandez, J. J.; and Cedillos, R. A., 1976, Rev. Inst. Invest. Med., San Salvador, v. 5 (2), 176-187

Trypanosoma cruzi, analysis of therapy of 35 patients with lampit, discussion of various side effects and minor toxic reactions, review of other therapeutic measures and comparison with results using lampit

Lampit

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comeniana, Microbiol. (6), 35-50

Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Nifurtimox (Lampit)

Fuglsang, H.; and Anderson, J., 1978, Tropenmed. u. Parasitol., v. 29 (3), 355-358

Onchocerca volvulus, patients, preliminary trials with oral nifurtimox, possible effect on adult parasites: savanna of Cameroon

Lampit

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Nifurtimox -- Continued.

Lampit

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 339-347

Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

Nifurtimox (Lampit)

Janssens, P. G.; and de Muyneck, A., 1977, Ann. Soc. Belg. Med. Trop., v. 57 (4-5), 475-480

African trypanosomiasis, humans, clinical trials with nifurtimox

Nifurtimox (Lampit)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73

Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Lampit (Bayer 2502)

Levi, G. C.; and Amato Neto, V., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (5), 369-372

Trypanosoma cruzi, humans with chronic infections, lampit, frequent side effects

Nifurtimox (Lampit)

Marsden, P. D.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (4), 391-394

South American mucocutaneous leishmaniasis, humans, nifurtimox, clinical trials, not recommended for routine use: Brazil

Lampit

Mocelin, A. J.; et al., 1977, Transplantation, v. 23 (2), 163 [Letter]

Trypanosoma cruzi in 29-year-old woman recipient of kidney transplant, negative blood smears after lampit, kidney function preserved: Brazil

Nifurtimox

Pagano, M. A.; O'Neill, E.; and Aristimuno, G., 1977, Rev. Neurol. Argentina, v. 3 (3), 467-470

human Chagas disease, nifurtimox therapy, development in some persons of an axonal neuropathy involving mainly the lower limbs

Nifurtimox

Prata, A., 1978, AMB, Rev. Ass. Med. Brasil., v. 24 (4), 140-142

Chagas disease, humans, nifurtimox, R07-1051, comparative study

Bayer 2502

Rassi, A.; and Ferreira, H. de O., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (5), 235-262

Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best re-

Nifurtimox

Ruggieri, H.; and Piserchia, C. N., 1977, Rev. Neurol. Argentina, v. 3 (3), 474-476

Chagas disease, humans with psychiatric disturbances, symptoms intensified by nifurtimox therapy, disappearance or diminution of symptoms when drug was discontinued

Nifurtimox -- Continued.

Nifurtimox

Schmunis, G. A.; et al., 1978, *J. Infect. Dis.*, v. 138 (3), 401-404

Trypanosoma cruzi, persistence of tissue-reacting (EVI) antibodies in nifurtimox-treated patients followed for several months to 2 years, implications for significance of EVI antibodies in pathogenesis of Chagas disease

Nitrofurfurylidene (Bayer 2502)

da Silva, N. N.; et al., 1974, *Rev. Soc. Brasil. Med. Trop.*, v. 8 (6), 325-334

Trypanosoma cruzi, clinical trials with nitrofurfurylidene in patients with chronic Chagas disease

Bayer 2502

Tippit, T. S., 1978, *Southwest. Vet.*, v. 31 (2), 97-104

Trypanosoma cruzi, dog, Bayer 2502, good results

Nitrofurfuridine (Lampit)

Urrutia, L. E.; Ramirez, R.; and Ruiz, J., 1976, *Rev. Inst. Invest. Med.*, San Salvador, v. 5 (2), 154-159

Trypanosoma cruzi, survey of 40 children with either acute Chagas disease or chagasic myocarditis, 5-year follow-up of relationship between therapy with lampit, electrocardiographic changes, and changes in body weight: San Salvador, El Salvador

Nifurtimox

Viviani, C. A.; et al., 1977, *Rev. Neurol. Argentina*, v. 3 (3), 471-473

Chagas disease, humans with neurologic disturbances, symptoms aggravated by nifurtimox therapy

Lampit (Nifurtimox; Bayer 2502)

Wegner, D. H. G., 1975, *Rev. Inform. Med. Terap.*, v. 50 (5-6), 67-73

Trypanosoma cruzi, humans, clinical review, recommendations for using lampit as therapy

Nigella sativa seeds

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118

Ascaridia galli, *Ascaris vitulorum*, in vitro anthelmintic activity of some Egyptian plants; only *Nerium oleander* caused death of worms

Nigella sativa

Agarwal, R.; Kharya, M. D.; and Shrivastava, R., 1979, *Indian J. Exper. Biol.*, v. 17 (11), 1264-1265

Taenia solium, *Bunostomum trigonocephalum*, *Oesophagostomum columbianum*, anthelmintic activity of essential oil of *Nigella sativa* in vitro

Nigericin (Duamycin; Polytherin A; Antibiotic X-464)

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Nigericin

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Nilodin. See Lucanthone.

Nilverm. See Tetramisole.

Nilzan. See Oxyclozanide or Tetramisole.

Nimorazole. See Nitrimidazine.

Niridazole -- Ambilhar; Niridizole; Nitrothiazole; 5-Nitro-2-thiazolyl-2-imidazolidine; 1-(5-Nitro-2-thiazolyl)-2-imidazolidinone.

Niridazole

Abdel Samad, M. M.; et al., 1977, *Tropenmed. u. Parasitol.*, v. 28 (4), 554-559

Schistosoma mansoni, mice, liver monoamine oxidase activity during course of infection and after chemotherapy, may be useful index for progression or regression of liver fibrosis

Niridizole (Ambilhar)

Al-Mallah, A. K.; Abdel Aziz, F. T.; and Hassanein, R. R., 1977, *Zentralbl. Vet.-Med.*, Reihe A, v. 24 (1), 76-80

Schistosoma haematobium, human males, normal rats and Dutch rabbits, ambilhar, activity of serum choline esterase and transaminases as measure of side effects of drug on liver

Niridazole (Ambilhar)

Bassily, S.; et al., 1979, *Ann. Trop. Med. and Parasitol.*, v. 73 (3), 295-296

Schistosoma mansoni-infected male farmers, niridazole administered at reduced daily dose over a 12-day period, clinical trials

Niridazole

Benazet, F.; et al., 1970, *Scand. J. Infect. Dis.*, v. 2 (2), 139-143

intestinal and hepatic parasites, nitroheterocyclic antiparasitics, laboratory studies of chemotherapeutic activity and toxicity in exper. animals

Niridazole

Blumer, J. L.; et al., 1979, *Molec. Pharm.*, v. 16 (3), 1019-1030

niridazole, aerobic metabolism by rat liver microsomes

Niridazole

Bulay, O.; et al., 1979, *Cancer Research*, v. 39 (12), 4996-5002

niridazole, rats, induction of kidney tumors

Niridazole -- Continued.

Niridazole

Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, *J. Parasitol.*, v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy

Niridazole

Erasmus, D. A.; and Davies, T. W., 1979, *Exper. Parasitol.*, v. 47 (1), 91-106
Schistosoma mansoni, S. haematobium, calcareous corpuscles in vitelline cells, morphological observations, X-ray microanalysis, effect of drug treatment

Niridazole

Ferreira, J. M.; Bassoi, O. N.; and Shiroma, M., 1976, *Rev. Hosp. Clin., S. Paulo*, v. 31 (3), 144-150
Sch[istosoma] mansoni, human infections, associated prolonged infection with salmonellosis, suggests that schistosomes may operate as reservoirs for the multiplication of the Salmonellae, mixed infections cured by use of the antischistosomal drug (niridazole) alone

Niridazole

Foster, R., 1973, *Rev. Inst. Med. Trop. S. Paulo*, v. 15 suppl. 1 (6), 1-9
Schistosoma mansoni, S. haematobium, S. japonicum, oxamniquine, efficacy and toxicity in various exper. animals, clinical trials in humans

Niridazole (Ambilhar)

Furtado, T., 1974, *Rev. AMMG*, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Niridazole

Gateff, C.; et al., 1971, *Ann. Soc. Belges Med. Trop. Parasitol.*, v. 51 (3), 309-324
S[chistosoma] haematobium, influence of infection on economic capacity of young adult population of workers on sugar cane estate, effect of niridazole treatment: Cameroons

Niridazole

Gutteridge, W. E.; Gaborak, M.; and Cover, B., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 339-347
Trypanosoma cruzi, in vivo and in vitro activity of SQ 18,506 compared with that of similar nitroheterocyclic compounds

Ambilhar (Niridazole)

Hafeiz, A. A.; Salah, M. K.; and Abdel Aziz, F. T., 1978, *Vet. Med. J., Giza*, v. 24 (24), 1976, 65-69
ambilhar, rabbits, excretion of iron glucuronic acid, and etherial sulphates in urine

Niridazole

Katz, N.; et al., 1973, *Rev. Soc. Brasil. Med. Trop.*, v. 7 (6), 381-387
Schistosoma mansoni, isolation of drug resistant strain (WW strain), reactions in mice to therapy with hycanthon, niridazole and oxamniquine compared with reactions of LE drug sensitive strain

Niridazole -- Continued.

Niridazole

Khayyal, M. T.; et al., 1978, *Egypt. J. Bilharz.*, v. 4 (1), 1977, 89-96
S[chistosoma] mansoni, effect of niridazole on lipid pattern of worms and serum and liver of infected and non-infected mice

Ambilhar

Labo, L., 1974, *Ann. Acad. Med. Stetinensis*, suppl. 10, 33-38
schistosomiasis, case reports of infection in African students living in Poland, unsuccessful therapy with nilodin in one case with cure after use of ambilhar

Ambilhar

Magzoub, M., 1971, *Sudan Med. J.*, v. 9 (3), 178-182
Schistosoma mansoni, untreated worms and worms treated with ambilhar or astiban, electron microscopy of cuticle, subcuticular region, and gut; possibility that egg formation is interrupted by either treatment

Niridazole

Molyneux, M. E.; and Galatius-Jensen, F., 1978, *South African Med. J.*, v. 54 (21), 871-872
schistosomal myelopathy, man, presentation with quadriplegia, large mass demonstrated in spinal cord, total clinical recovery after niridazole: Malawi

Niridazole

Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, *Rev. Brasil. Pesqui. Med. e Biol.*, v. 2 (1), 45-49
Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

Niridazole

Ong, T. M., 1978, *Mutation Research*, v. 55 (1), 43-70
hycanthon and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Niridazole

Ottesen, E. A.; et al., 1978, *Clin. and Exper. Immunol.*, v. 33 (1), 38-47
Schistosoma mansoni, patients with acute, subacute, and chronic disease before and after niridazole treatment, lymphocyte responsiveness to schistosome antigens, possible implications of diminished cellular immune reactivity in chronic disease state

Niridazole (Ambilhar)

Pugh, R. N. H., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (5), 495-497
Schistosoma haematobium, human, concurrent single-dose therapy with metrifonate and niridazole: Malumfashi District, Nigeria

Niridazole

da Silva, L. C.; et al., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (2), 121-130
Schistosoma mansoni, humans, immunodiffusion, hemagglutination, immunofluorescence and eosinophil counts before and after therapy with hycanthon or niridazole

Niridazole -- Continued.

Niridazole

da Silva, L. C.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 206-210
S[*chistosoma*] *mansoni*, humans parasitologically cured, repeat therapy with niridazole or hycanthone, evaluated by passive hemagglutination, indirect immunofluorescence and immunodiffusion tests

Niridazole

Simitzis-Le Flohic, A. M.; et al., 1977, Semaine Hop. Paris, v. 53 (22-23), 1369-1370
schistosomiasis with minor parasitism of distomiasis, ascariasis, trichocephaliasis, woman with presenting symptoms of adrenal insufficiency, cortisone therapy resulted in aggravated symptoms and asthenia, parasitism diagnosed, piperazine therapy resulted in toxic neurologic reactions, illness resolved after niridazole therapy: France (had resided in Central African Republic)

Niridazole

Tiboldi, T., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 1026-1030
Schistosoma mansoni-infected mice, histopathological changes in ovaries can be reversed by adequate antischistosomal therapy

Ambilhar

Velasquez-Antich, A., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (5), 347-353
Schizotrypanum cruzi, mice, ambilhar

Niridazole

Witham, R. R.; and Mosser, R. S., 1979, Gastroenterology, v. 77 (6), 1316-1318
schistosomiasis *mansoni* in Arabian man as cause of duodenitis, successful treatment with niridazole, case report: United States (emigrated from Yemen)

Niridazole (Ambilhar)

Woolhouse, N. M., 1979, Biochem. Pharmacol., v. 28 (16), 2413-2418
antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Niridazole. See Niridazole.

Nitazol. See Aminitrozole.

Nitidine chloride

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Nitridazole

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Nitrimidazine -- N-beta-Ethyl-morpholino-1-nitro-5 imidazole; Nimorazole; Naxogin; Naxogyn; Nimorazole; TN-7.

Nitrimidazine -- Continued.

Nimorazole (Naxogyn)

Apt, W.; et al., 1976, Rev. Med. Chile, v. 104 (11), 791-793
Entamoeba histolytica, human intestinal amoebiasis, successful treatment with nimorazole

Naxogin (Nitrimidazine)

Aruta, J.; Galani, M.; and Fertilio, O., 1973, Rev. Chilena Obst. y Ginec., v. 38 (3), 118-120
human vaginal trichomoniasis, comparison therapeutic trials using oral naxogin and vaginal tablets of naxogin combined with nystatin and quemicetina in cases with diagnostic problems after diagnosis confirmed by cytology or colposcopy: Chile

Nitrimidazine + Nystatin + Quemicetina (as a combined vaginal tablet)

Aruta, J.; Galani, M.; and Fertilio, O., 1973, Rev. Chilena Obst. y Ginec., v. 38 (3), 118-120
human vaginal trichomoniasis, comparison therapeutic trials using oral naxogin and vaginal tablets of naxogin combined with nystatin and quemicetina in cases with diagnostic problems after diagnosis confirmed by cytology or colposcopy: Chile

Nitrimidazine (TN-7)

Autera, C., 1973, Quad. Clin. Ostet. e Ginec., v. 28 (1), 1-16
Trichomonas vaginalis, human trichomoniasis, clinical aspects, satisfactory trials with nitrimidazine treating both women with vaginitis and their sexual partners

Nitrimidazine

Barbosa J., J.; et al., 1975, Rev. Obst. y Ginec. Venez., v. 35 (3), 259-272
human vaginal trichomoniasis, successful clinical trials testing efficacy of nitrimidazine, few side effects: Venezuela

Nimorazole

Bataillard, J., 1978, Rev. Franc. Gynec. et Obst., v. 73 (1), 67-70
human vaginal trichomoniasis, successful therapy with nimorazole except in instances of continued exposure to reinfection

Nitrimidazine (Nimorazole)

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Nitrimidazine

de Carneri, I., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (5), 321-325
Balantidium coli in culture, nitrimidazine and metronidazole tested

Nimorazole (Naxogyn)

Cavler, R.; and Cenac, J., 1972, Therapeutique, v. 48 (5-6), 391-394
Trichomonas vaginalis, nimorazole vs. metronidazole, in vitro and in vivo trials with rats and mice

Naxogyn

Dajoux, R., 1974, Rev. Franc. Gynec. et Obst., v. 69 (2), 133-136
Trichomonas vaginalis, human vaginal infection associated with punctiforme colpitis, therapy with naxogyn

Nitrimidazine -- Continued.

Nitrimidazine

Fabio, P. F.; et al., 1978, J. Med. Chem., v. 21 (3), 273-276

Entamoeba histolytica, diaminoanthraquinone bisamidines, laboratory trials comparing activity against cecal form in rats and hepatic form in golden hamsters with activity of known amoebicides

Nimorazole (Nitrimidazine; Naxogin)

Huggins, D., 1975, Rev. Brasil. Clin. e Terap., v. 4 (10), 421-424

Giardia lamblia, humans, successful clinical trials with nimorazole

Nitrimidazine (Naxogin)

Huggins, D.; and Maciel, M. F., 1972, Rev. Soc. Brasil. Med. Trop., v. 6 (3), 163-166

Giardia lamblia, children, therapy with nitrimidazine, well tolerated, good results: Brazil

Nimorazole

Imparato, E.; et al., 1976, Quad. Clin. Ostet. e Ginec., v. 31 (4), 225-239

Trichomonas vaginalis, human vaginal trichomoniasis, evaluation of mepartricin as oral therapy using nimorazole and clotrimazole as reference drugs, best results obtained with mepartricin

Nimorazole

Levi, G. C.; de Avila, C. A.; and Amato Neto, V., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (6), 422-424

giardiasis, humans, efficacy of various drugs, comparative study, side-effects

Nimorazole (Nitrimidazine; Naxogin)

Lima, R. S. F., 1975, Rev. Brasil. Clin. e Terap., v. 4 (5), 201-204

human Giardia lamblia, treatment of 32 patients with nimorazole resulted in 100% cure with marked clinical improvement and no side effects

Nimorazole

Lindmark, D. G.; and Mueller, M., 1976, Antimicrob. Agents and Chemotherapy, v. 10 (3), 476-482

metronidazole and 11 other nitroimidazoles, antitrichomonad activity against Trichomonas foetus and Trichomonas vaginalis, mutagenic action in Salmonella test, reducibility of nitro group by T. foetus homogenates, results underscore role of reduction of nitro group in antitrichomonad and mutagenic activity of nitroimidazoles

Nimorazole

Meingassner, J. G.; and Thurner, J., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 254-257

Trichomonas vaginalis, isolation of strain resistant to metronidazole and other 5-nitroimidazoles

Mimorazole [i.e.? Nimorazole] (Naxogin)

Merchan Lopez, D.; and Olmos, R., 1976, Rev. Obst. y Ginec. Venez., v. 36 (1), 181-185

human genital trichomoniasis, clinical trials with mimorazole, good results obtained: Venezuela

Nitrimidazine -- Continued.

Nimorazole

Pesando, P. C.; and Guaschino, S., 1979, Minerva Ginec., v. 31 (3), 155-162

Trichomonas vaginalis, women, vaginal infections, clinical evaluation of nimorazole

Nitrimidazine (Naxogin)

Merdivenci, A.; Baydemir, M.; and Sengul, M., 1976, Turk Hijiy. ve Deneysel Biyol. Dergisi, v. 36 (2), 238-247

Giardia intestinalis, children, tinidazole and nitrimidazine compared, both successful

Nimorazole

Ruggeri, E., 1976, Quad. Clin. Ostet. e Ginec., v. 31 (4), 211-223

Trichomonas vaginalis, vaginal trichomoniasis, oral mepartricin showed significant therapeutic superiority when compared in clinical trials with patients who received nimorazole therapy orally or with patients treated with a vaginal cream containing amphotericin B and tetracycline

2-Nitro 4-acetylaminophenol

Kolesnikov, V. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 143-146

Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective

6-Nitrobenzimidazole

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

p-Nitrobenzyl-6-thioguanosine

Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856

Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Nitrodan -- 3-Methyl-5-(4-nitrophenylazo)rhodanine.

Nitrodan

Douch, P. G. C.; and Buchanan, L. L., 1979, Xenobiotica, v. 9 (8), 467-473

Ascaris suum, Moniezia expansa, disophenol, nitroxylnil, nitrodan, metabolism by intact helminths, by helminth enzyme preparations, and by mouse- and sheep-liver enzymes

Nitro-2 dimethoxy-5,6 benzofuran

Cavier, R.; et al., 1979, Ann. Pharm. Franc., v. 37 (7-8), 309-312

Trichomonas vaginalis, Entamoeba histolytica, rats, 2-nitro benzofuran derivatives compared with metronidazole

Nitro-2 dimethoxy-6,7 benzofuran

Cavier, R.; et al., 1979, Ann. Pharm. Franc., v. 37 (7-8), 309-312

Trichomonas vaginalis, Entamoeba histolytica, rats, 2-nitro benzofuran derivatives compared with metronidazole

5-Nitro-diphenol

Mirck, M. H.; and Bergsma, R. F., 1979, Tijdschr. Diergeneesk., v. 104 (15-16), 633-634 anthelmintic containing '5-nitro-diphenol' as active principle found actually to contain tetramisole-hydrochloride, unsuitable systemic anthelmintic for horses, brief review

Nitrodiphenylaminoisothiocyanate. See Nitros-canate.

5-Nitro-2-furaldehyde-semicarbazone. See Nitro-furazone.

5-Nitro-2-furaldehyde thiosemicarbazone. See Benzazon VII.

Nitrofurantoin -- Furadantin; Furantoin; 1-(5-Nitro-2-furfurylideno)-hydantoin.

Nitrofurantoin

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Furadantin (Furantoin)

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Microbiol. (6), 35-50 Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Nitrofurantoin (Furadantin)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73 Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Nitrofurazone -- Bifuran (with Furazolidone); Furacillin; Furacin; 5-Nitro-2-furaldehyde-semicarbazone; Nitrofurazone Spofa Premix.

Bifuran

Aggarwal, C. K.; et al., 1978, Indian Vet J., v. 55 (12), 952-957 bifuran, embazin, effect of coccidiostats, antibiotics, and litters (built up vs. fresh) on broiler chick performance

Nitrofurazone

Brener, Z., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (4), 302-306 Trypanosoma cruzi, description of method allowing study of drug action on trypanostigotes in mice

Nitrofurazone

Brener, Z.; Costa, C. A. G.; and Chiari, C., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (6), 450-455 Trypanosoma cruzi, mice treated with nitrofurazone, nifurtimox, or Ro 7-1051, differences in susceptibility of 4 parasite strains to active drugs attributed to biological characteristics of strains rather than to mode of drug action

Nitrofurazone -- Continued.

Nitrofurazone

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672 trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Nitrofurazone (Furacin)

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Microbiol. (6), 35-50 Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Nitrofurazone

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338 Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Nitrofurazone

Hrdlicka, J.; and Stolejda, J., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (3), 233-236 coccidiostats in feeds, qualitative identification test

Furacillin

Musaev, F. A., 1972, Parazitologiya, Leningrad, v. 6 (2), 185-188 Trichomonas elongata, occurrence in oral cavity of healthy persons vs. those with oral cavity diseases, host age and sex, suitable media for cultivating trichomonads, activity in vitro of several medicinal substances, role of trichomonads in periodontitis confirmed by treatment of patients with trichopol

Nitrofurazone

Rassi, A.; and Ferreira, H. de O., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (5), 235-262 Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best response

Nitrofurazone Spofa Premix

Tesarcik, J., 1971, Prace VURH Vodnany (9), 99-132 Neoechinorhynchus rutili, Eimeria spp., carp, tetrafinol, nitrofurazone, prophylaxis and therapy under aquarium conditions

Nitrofurazone Spofa Premix. See Nitrofurazone.

Nitrofurfuridine. See Nifurtimox.

1-(5-Nitro-2-furfurylideno)-hydantoin. See Nitrofurantoin.

3-(5-Nitro-2-furfurylideneamino)-2-oxazolidone.
See Furazolidone.

Nitrofurfurylidene. See Nifurtimox.

3-(5-Nitrofurfurylidene-amino)-2-oxazolidinone.
See Furazolidone.

N-(5-Nitro-2-furfurylidene)-3-amino-2-oxazolidone. See Furazolidone.

Nitrofurylacrylamide -- F30066; Furapromidium;
N-Isopropyl-3-(5-nitro-2-furyl)-acrylamide.

Furapromidium (F30066)
Ong, T. M., 1978, Mutation Research, v. 55
(1), 43-70
hycanthone and other antischistosomal drugs,
general properties, teratogenicity, carcinogenicity,
mutagenicity, and other genetically related activities, review

Nitrogen mustard
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, Antimicrob. Agents and Chemotherapy, v. 15
(2), 157-160
Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

3-Nitro-4 hydroxyphenylarsonic acid. See
Roxarsone.

Nitroimidazoles
Lindmark, D. G.; and Mueller, M., 1976, Antimicrob.
Agents and Chemotherapy, v. 10 (3),
476-482
metronidazole and 11 other nitroimidazoles,
antitrichomonad activity against Tritrichomonas
foetus and Trichomonas vaginalis,
mutagenic action in Salmonella test, reducibility
of nitro group by T. foetus homogenates, results
underscore role of reduction of nitro group in
antitrichomonad and mutagenic activity of nitroimidazoles

Nitroimidazoles
Meingassner, J. G.; et al., 1978, Antimicrob.
Agents and Chemotherapy, v. 13 (1), 1-3
Tritrichomonas foetus, metronidazole-resistant
and susceptible strains, in vitro susceptibility
testing, results suggest that the two strains
differ in regulation of internal redox systems
and underscore the role that testing methods
may play in the in vitro detection of nitroimidazole-
resistant protozoan parasites

4-Nitro-4'isothiocyanate-diphenylamine. See
Nitroscanate.

4-Nitro-4-isothiocyanato-diphenyl ether. See
Nitroscanate.

3-(5-Nitro-1-methyl-2-imidazoylethylmethylene-amino)-
5-morpholino-methyl-2-oxazolidinone hydrochloride.
See Moxnidazole.

Nitromide -- 3,5-Dinitrobenzamide.

3,5-Dinitrobenzamide
Orekhova, M. M., 1973, Vet. Nauka--Proizvod.,
Trudy, Minsk, v. 11, 156-159
[Eimeria tenella], chickens, 3,5-dinitrobenzamide
having negative effect on schizonts and gamonts,
but no effect on sporozoites; no effect on nucleic
acids of endogenous stages shown

Nitromin
Sakamoto, T.; and Gemmell, M. A., 1979, Mem.
Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicalidal effect of
65 antibiotic, antineoplastic, cytostatic, and
other agents in vitro

4-(4'-Nitrophenoxy) phenyl isothiocyanate. See
Nitroscanate.

1-(2-Nitro-4-propyl-oxyphenyl)-3-carbomethoxy-5-
methyl isothio-urea (Sch 23154)
Loebenberg, D.; et al., 1979, J. Parasitol.,
v. 65 (5), 823-824
Syphacia obvelata, mice, Sch 23154 compared
with pyrantel pamoate and pyrinium pamoate

Nitroscanate -- Cantrodifene; Cantrodiphen;
CGA-23'654; C9333-Go/CGP 4540; CGP-4540;
Echinon; GO 9333; GS-23'654; 4-Isouthiocyanate-
4'-nitrodiphenyl ether; 4-Isouthiocyanato-4'-
nitrodiphenylamine; 4-Isouthiocyanato-4'-nitro-
diphenylamine; 4-Isouthiocyanato-4-nitro diphenyl
ether; Lopatol; Nitrodiphenylaminoisothiocyanate;
4-Nitro-4'isouthiocyanate-diphenylamine; 4-Nitro-
4-isouthiocyanato-diphenyl ether; 4-(4'-Nitrophenoxy)
phenyl isothiocyanate.

Lopatol
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10),
28-36
cestodes of sheep, drug trials; Stilesia globipunctata,
tested several diagnostic methods with unfavorable
results

Cantrodiphen
Bankov, D. E., 1977, Vet.-Med. Nachr. (2),
145-148
Echinococcus granulosus, dogs (exper.),
bunamidine hydrochloride, cantrodiphen,
praziquantel, praziquantel most effective and
showed no toxicity

Nitroscanate (Lopatol)
Boray, J. C.; et al., 1979, Austral. Vet. J.,
v. 55 (2), 45-53
nematodes and cestodes of dogs and cats, efficiency
and safety of nitroscanate, comparison with
mebendazole, bunamidine hydrochloride, and
praziquantel

4-Isouthiocyanato-4'-nitro-diphenylamine (CGP 4540)
Cha, Y. N.; and Bueding, E., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (6), 1188-1191
Schistosoma mansoni, mice, recovery of hepatic
drug-metabolizing capacity following curative
dose of 4-isouthiocyanato-4'-nitro-diphenylamine

Nitroscanate -- Continued.

Nitroscanate (Lopatol; Cantrodifene)

Evans, J. W.; and Green, P. E., 1978, Austral. Vet. J., v. 54 (9), 454-455 [Letter]
Platynosomum concinnum, cats, anthelmintics, drug trials

Nitroscanate

Gemmell, M. A., 1978, Bull. World Health Organ., v. 56 (3), 433-443
tapeworm control in dogs for prevention of hydatidosis and cysticercosis in sheep, monthly drug treatment program, age-specific prevalence of *Taenia hydatigena* in lambs used as principal indicator, 13-year assessment: Styx Valley and Maniototo County, South Island, New Zealand

4-Nitro-4'-isothiocyanate-diphenylamine (GO 9333)

Gupta, P. S.; et al., 1979, J. Trop. Med. and Hyg., v. 82 (6), 117-119
Ancylostoma duodenale, *Necator americanus*, human, clinical trials with 4-nitro-4'-isothiocyanate-diphenylamine, ideal dosage schedule established

Lopatol (Cantrodifene)

Kassai, T.; et al., 1977, Magy Allat. Lapja, v. 99, v. 32 (6), 373-377
Echinococcus granulosus, dogs, bunamidine hydroxynaphthoate in biscuit form effective, lopatol containing cantrodifene unsatisfactory

Nitroscanate (CGP-4540)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

4-Isouthiocyanato-4'-nitrodiphenylamine (C9333-Go/CGP 4540)

Middleton, K. R.; Schaefer, F. W. III; and Saz, H. J., 1979, *Experientia*, v. 35 (2), 243-244
4-isouthiocyanato-4'-nitrodiphenylamine, anthelmintic activity against *Nematospiroides dubius* and *Hymenolepis nana* in mice and *Hymenolepis diminuta* in rats, no activity against *Spirometra mansonioides* in cats

Nitrodiphenylaminoisothiocyanate (C9333 GO/CGP 4540)

Ong, T. M., 1978, *Mutation Research*, v. 55 (1), 43-70
hycanzone and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Echinon (Nitroscanate)

Rajkovic-Janje, R.; and Ropic, D., 1978, *Vet. Glasnik*, v. 32 (3), 225-229
Echinococcus granulosus, dogs (exper.), echinon

5-Nitro-2-thiazolyl-2-imidazolidine. See Niridazole.

1-(5-Nitro-2-thiazolyl)-2-imidazolidinone. See Niridazole.

2-Nitro 4-trichlor-acetyl-aminophenol

Kolesnikov, V. I., 1977, *Sborn. Nauch. Rabot SibNIVI* (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective

Nitro-2 trimethoxy-4,5,6 benzofuran

Cavier, R.; et al., 1979, *Ann. Pharm. Franc.*, v. 37 (7-8), 309-312
Trichomonas vaginalis, *Entamoeba histolytica*, rats, 2-nitro benzofuran derivatives compared with metronidazole

Nitro-2 trimethoxy-4,6,7 benzofuran

Cavier, R.; et al., 1979, *Ann. Pharm. Franc.*, v. 37 (7-8), 309-312
Trichomonas vaginalis, *Entamoeba histolytica*, rats, 2-nitro benzofuran derivatives compared with metronidazole

Nitroxynil -- 4-Cyano-2 iodo-6 nitrophenol;

Dovenix; 3-Iodo-4-hydroxy-5-nitrobenzonnitrile; Trodax.

Nitroxynil (Dovenix)

Delavenay, R. P., 1978, *Rev. Elevage et Med. Vet. Pays Trop.*, n. s., v. 31 (2), 171-177
parasites of *Camelus dromedarius*, nitroxynil, well tolerated in toxicity assay, very effective against *Cephalopina titillator* and *Haemonchus contortus*: Ethiopia

Nitroxynil

Douch, P. G. C.; and Buchanan, L. L., 1979, *Xenobiotica*, v. 9 (8), 467-473
Ascaris suum, *Moniezia expansa*, disophenol, nitroxynil, nitrodan, metabolism by intact helminths, by helminth enzyme preparations, and by mouse- and sheep-liver enzymes

Nitroxynil (Trodax)

Gupta, R. P.; and Malik, P. D., 1979, *Indian Vet. J.*, v. 56 (10), 834-838
helminths, sheep, trodax, activity under field and controlled conditions

Dovenix

Petrov, D.; Milushev, I.; and Monov, M., 1978, *Vet. Med. Nauki*, v. 15 (4), 33-39
Oestrus ovis, sheep, neguvon, ranide, and dovenix compared; use of ranide economically justified only for mixed infection with *Fasciola hepatica*

4-Cyano-2-iodo-6-nitrophenol

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Dovenix

Stoianova-Radoeva, S., 1978, *Vet. Med. Nauki*, v. 15 (4), 79-85
acedist, dovenix, treated sheep, residues in milk, effect on blood biochemical indices

Nitrothiazole. See Niridazole.

Nitroxylin -- Continued.

Nitroxylin (Trodat)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]

Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiazobenzazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxylin are also useful chemical alternatives

Nitroxylin (Trodat)

Wellington, A. C., 1978, J. South African Vet. Ass., v. 49 (2), 125-126

Fasciola gigantica, Haemonchus placei, Bunostomum phlebotomum and Oesophagostomum radiatum, cattle, nitroxylin injected subcutaneously achieved class A efficacy when evaluated by non parametric method, not effective against Cooperia spp

Nitroxylin (Trodat)

Wellington, A. C., 1978, J. South African Vet. Ass., v. 49 (2), 131-132

Parafilaria bovicola, cattle, efficacy of nitroxylin administered at two dosage levels, reduced lesion areas and carcass lesions, financial implications

Dovenix (Nitroxylin)

Ziegler, K., 1979, Vet. Med., Praha, v. 52, v. 24 (7), 391-395

Fasciola hepatica, cattle, dovenix and bilevon highly effective; drug toxicity tests in rats

Nitryl. See p-Dimethylaminobenzonitrile or Sulfaquinoxaline.

Nivaquine. See Chloroquine.

Nivaquine 200. See Chloroquine.

Noevar. See Trichlorfon.

Nogalamycin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

No-Pest strips. See Dichlorvos.

Nor-β-lapachone -- 2,3-Dihydro-2,2'-dimethyl-2H-naphtho [1,2-b]-furan-4,5-dione.

Nor-β-lapachone

Boveris, A.; et al., 1978, Comp. Biochem. and Physiol., v. 61C (2), 327-329

Trypanosoma cruzi, correlation between superoxide anion production and trypanocidal action of naphthoquinones

Nor-β-lapachone -- Continued.

Nor-β-lapachone

Docampo, R.; et al., 1978, Ztschr. Parasitenk., v. 57 (3), 189-198

Trypanosoma cruzi, naphthoquinones, effect on ultrastructure and superoxide anion and hydrogen peroxide production of different stages

Nor-β-lapachone

Lopes, J. N.; et al., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 523-531

Trypanosoma cruzi, 1,4-naphthoquinone and 1,2-naphthoquinone derivatives, in vitro and in vivo (mice) evaluation of effects on growth, viability, and infectivity; in vitro studies also on Crithidia fasciculata

Norsulfazole. See Sulfathiazole.

D-Norvaline

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

NOSAN

Muresan, E.; et al., 1978, Apicultura Romania, v. 53 (4), 13-15, 16

Nosema apis, Fumidil B and NOSAN against Paramecium caudatum as substitute test agent (having sensitivity similar to N. apis)

Notezine. See Diethylcarbamazine.

Novarsenol. See Neoarsphenamine.

Novastat-W. See Aklomide or Sulfantran.

Noviben Suspension. See Cambendazole.

Noxal. See Sulfaquinoxaline.

Nucleocidin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Nuvacron. See Monocrotophos.

Nuvan. See Dichlorvos.

Nuvan 100 EC. See Dichlorvos.

Nystatin -- Canaural (with Framycetin and Prednisolone); Macmiror plus (with Lysozyme and Nifuratel).

Nystatin

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Nystatin + Quemicetina + Nitrimidazine (as a combined vaginal tablet)

Aruta, J.; Galani, M.; and Fertilio, O., 1973, Rev. Chilena Obst. y Ginec., v. 38 (3), 118-120
human vaginal trichomoniasis, comparison therapeutic trials using oral naxogin and vaginal tablets of naxogin combined with nystatin and quemicetina in cases with diagnostic problems after diagnosis confirmed by cytology or colposcopy: Chile

Nystatin -- Continued.

Nystatin + Lysozyme + Nifuratel (= Macmiror plus) Martins, A. D., 1974, Rev. Brasil. Med., v. 31 (7), 503-508

human vaginal trichomoniasis alone or in the presence of fungal infections, successful therapy combining oral nifuratel and vaginal treatment with nifuratel combined with nystatin and lysozyme

Canaural

Pott, J. M.; and Riley, C. J., 1979, Vet. Rec., v. 104 (25), 579

Otodectes cynotis, dogs, cats (ear canals of both), canaural, comparison with proprietary topical ear preparation (clendrol), controlled trial

Octylthiobenzoate. See Tioctilate.

Oligomycin

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Oligomycin

Seilhamer, J. J.; and Byers, T. J., 1978, *J. Protozool.*, v. 25 (4), 486-489
Acanthamoeba castellanii, mutants resistant to erythromycin, chloramphenicol, and oligomycin

Olivacine -- 1,5-Dimethyl-1-6-hydro-pyrido-4,3b-carbazole.

Olivacine

Leon, L.; et al., 1978, *Exper. Parasitol.*, v. 45 (2), 151-159
Trypanosoma cruzi, effect of olivacine in vitro on growth, on macromolecular synthesis, on ultrastructure, and on respiration of epimastigotes, in vivo activity does not parallel in vitro effects

Olivomycin

Sukhareva-Nemakova, N. N.; Titova, T. S.; and Avakian, A. A., 1975, *Izvest. Akad. Nauk SSSR, s. Biol.* (2), 275-280
Crithidia oncopelti, comparative study of ultrastructure, cultures differing in sensitivity to olivomycin; lipid drops in cytoplasm of resistant protozoa; nature of action of olivomycin on sensitive parasites

Omnes. See Nifuratel.

Omnizole. See Thiabendazole.

Omnizole Wormer Paste. See Thiabendazole.

Oncodazole

Ireland, C. M.; et al., 1979, *Biochem. Pharmacol.*, v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiroides dubius* in mice

Opigal. See Carbaryl.

Orasol

Klenov, A. P., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 152-155
[*Bothriocephalus*], white amur, effectiveness of various anthelmintics

Organophosphate acaricides

Reich, C. I.; et al., 1978, *Exper. Parasitol.*, v. 44 (1), 50-55
Boophilus microplus, 2 Argentinian strains, one resistant and one sensitive to organophosphate acaricides, differences in cholinesterase system

Organophosphorus warblecide

Tordoff, D. B., 1978, *Vet. Rec.*, v. 102 (4), 92 [Letter]
heifers, transient infertility possibly caused by dressing with organophosphorus warblecide

Organosulphur compounds

Abdou, N. A.; et al., 1978, *Egypt. J. Pharm. Sc.*, v. 17 (2), 1976, 153-159
synthesis of some organosulphur compounds structurally related to certain antibilharzial drugs, to be screened for possible activity

Ornidazole -- alpha-Chloromethyl-2-methyl-5-nitroimidazole-1-ethanol; Ro 7-0207; Tiberol.

Ornidazole (Tiberol)

Caprilli, F.; Mercantini, R.; and Barba, C., 1977, *Riv. Ostet. e Ginec. Prat. e Med. Perinat.*, v. 57 (6), 547-551
Trichomonas vaginalis, human vaginal infections, oral and vaginal therapy with ornidazole

Ornidazole (Tiberol)

Chung, S. O.; et al., 1978, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 9 (1), 74-78
Trichomonas vaginalis, human, ornidazole, various dosage regimens, drug safe and effective: Korea

Ornidazole

Forsgren, A.; and Forssman, L., 1979, *Brit. J. Vener. Dis.*, v. 55 (5), 351-353
Trichomonas vaginalis, 35-year-old woman, symptomatic metronidazole-resistant vaginitis for 10 years, some resistance also to tinidazole and ornidazole: Sweden

Ornidazole (Tiberol)

Fuchs, P., 1978, *Deutsche Med. Wchnschr.*, v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Tiberol (Ro 7-0207)

Gross, J., 1976, *Rev. Med. Suisse Rom.*, v. 96 (5), 399-404
Trichomonas, human vaginitis with resulting leukorrhea, treatment with tiberol

Tiberol (Ro-07-0207)

Iyngkaran, N.; Lee, E. L.; and Robinson, M. J., 1978, *Scand. J. Infect. Dis.*, v. 10 (3), 243-246
Giardia lamblia, children, single dose treatment with tiberol, effective in all children treated; diagnosis by mucosal imprint, examination of stools, duodenal juice, or jejunal biopsy material compared, mucosal imprint method most reliable: Malaysia

Ornidazole (Tiberol)

Kornø, B.; Nygaard, B.; and Jensen, R. H., 1978, *Ugeskr. Laeger*, v. 140 (25), 1485-1487
Trichomonas vaginalis, women with vaginal infections, controlled trials with ornidazole given orally in various dosages, some side effects

Ornidazole -- Continued.

- Ornidazole (Tiberal)
Leite, E. V.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (1), 28-35
Giardia lamblia, humans, clinical trials, ornidazole and metronidazole compared: Brazil
- Ornidazol
Lindner, J. G. E. M.; et al., 1979, Chemotherapy, v. 25 (4), 243-248
cervicitis, women, effect of ornidazol on vaginal bacterial flora and Trichomonas vaginalis
- Tiberal (Ro 7-0207; Ornidazole)
Maneschi, M.; et al., 1979, Minerva Ginec., v. 31 (3), 163-168
Trichomonas, human vulvovaginitis, treatment trials with Ro 7-0207
- Ornidazole (Tiberal)
Mettler, L., 1978, Therapiewoche, v. 28 (34), 6095-6098
trichomoniasis, women with vaginitis and their sexual partners, single dose therapy with ornidazole, cure rates, side effects
- Ornidazole (Tiberal)
Nesvadba, J., 1979, Kleintier-Praxis, v. 24 (4), 177-179
Giardia, cat, ornidazole, metronidazole, case report
- Ornidazole (Tiberal)
Nygaard, B.; et al., 1977, Ugeskr. Laeger, v. 139 (9), 524-526
Trichomonas vaginalis, women with vaginitis, 7-day therapy with metronidazole vs. single-dose ornidazole
- Tiberal (Ro 7-0207; Ornidazole)
Restaino, A.; Ferreri, R.; and Ambrosio, M. G., 1977, Riv. Ostet. e Ginec. Prat. e Med. Perinat., v. 57 (6), 454-458
Trichomonas, human vaginitis, tiberal, comparison of 3 therapeutic schemes
- Ro-7-0207
Restaino, A.; Loizzi, P.; and Loverro, G., 1977, Riv. Ostet. e Ginec. Prat. e Med. Perinat., v. 57 (2), 170-174
Trichomonas vaginalis, human vaginitis, oral and vaginal therapy with Ro-7-0207
- Ornidazole (Ro 7-0207; Tiberal)
Richle, R.; et al., 1978, Arzneimittel-Forsch., v. 28 (4), 612-625
trichomoniasis, amoebiasis, lamblia, extensive in vitro and in vivo trials (humans, domestic animals, laboratory animals) with ornidazole to establish chemotherapeutic properties, efficacy slightly superior to metronidazole in comparative trials

Ornidazole -- Continued.

- Ornidazole (Tiberal)
Sandia, O. G.; et al., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 52-56
E[ntamoeba] histolytica, humans with chronic intestinal amoebiasis, clinical trials with ornidazole vs. Metronidazole
- Tiberal (Ro 7-0270)
Sandront-Degee, M.; Werbrouck-Navette, J.; and Lambotte, R., 1975, Rev. Med. Liege, v. 30 (17), 560-562
Trichomonas vaginalis, humans, therapeutic trials comparing action of tiberal with metronidazole, fasigyn and tinidazole
- Ornidazole (Tiberal)
Serup, J.; and Jensen, R. H., 1978, Ugeskr. Laeger, v. 140 (25), 1483-1484
Trichomonas vaginalis, women with vaginal infection and their sexual partners, controlled therapeutic trials, ornidazole vs. tinidazole as single oral dose
- Tiberal (Ro 7-0207)
Sonnino, S., 1975, Rassegna Internaz. Clin. e Terap., Napoli, v. 55 (21), 1331-1334
Trichomonas vaginalis, human vaginal infections, therapy with tiberal both orally and vaginally, sexual partners also treated
-
- Orthorix Spray. See Lime-sulfur solution.
- Orvagil. See Metronidazole.
- Ouabain
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Ovithelm. See Morantel.
- Oxamic acid, sodium salt
Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- Oxamniquine -- 6-Hydroxymethyl-2-isopropylamino-methyl-7-nitro-1,2,3,4-tetrahydroquinolone; Mansil; Oxamniquine embonate; Oxamniquine hydroxynaphthoate; Oxamniquine phthalate; Oxamniquine salicylate; Pfizer UK 4271; UK 4271; Vansil.

Oxamniquine -- Continued.

Oxamniquine

Abaza, H. H.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 602-604

Schistosoma mansoni, S. haematobium, human, associated complication of polyposis of colon, therapy with oxamniquine was effective and safe: Egypt

Oxamniquine (Mansil)

de Almeida, F. M. R.; et al., 1979, AMB, Rev. Ass. Med. Brasil., v. 25 (3), 101-104; Errata, v. 25 (7), at bottom of Indice

Schistosoma mansoni, humans, therapeutic evaluation of oxamniquine, no serious adverse effects: Sao Paulo

Oxamniquine (Pfizer UK 4271)

Axton, J. H. M.; and Garnett, P. A., 1976, South African Med. J., v. 50 (27), 1051-1053
Schistosoma spp., children, clinical trials with oral oxamniquine, effective only in treating S. mansoni

Oxamniquine

Bassily, S.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (6), 1284-1286
Schistosoma mansoni, patients with advanced complicated disease (some with concomitant S. haematobium infection), treatment with oxamniquine, encouraging results

Oxamniquine (UK 4271)

Bina, J. C.; and Prata, A., 1975, Rev. Soc. Brasil. Med. Trop., v. 9 (4), 175-178
[Schistosoma] mansoni, children, clinical trials testing efficacy of oxamniquine syrup in varying dosages; efficacy of dosage rose with age of child treated

Oxamniquine

Brazil, R. P.; and Gilbert, B., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (2), 87-88
Leishmania b. braziliensis, hamsters (exper.), oxamniquine, basis for study of human Leishmania species

Oxamniquine

Chavasse, C. J.; Brown, M. C.; and Bell, D. R., 1978, Ann. Trop. Med. and Parasitol., v. 72 (3), 293-295
Schistosoma mansoni, short term effects of oxamniquine on activity of paired worms in vitro

Oxamniquine

Clarke, V. de V.; et al., 1976, South African Med. J., v. 50 (46), 1867-1871
Schistosoma haematobium, S. mansoni, human, dose-finding trials for oral oxamniquine, effective only in treating S. mansoni: Rhodesia

Oxamniquine (Pfizer UK-4271)

Clarke, V. de V.; Blair, D. M.; and Weber, M., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 73-77
Schistosoma mansoni or S. haematobium or mixed infections, children, oxamniquine, clinical trials: Rhodesia

Oxamniquine -- Continued.

Oxamniquine (U.K. 4271)

Coura, J. R.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 41-46
[Schistosoma] mansoni, human, oxamniquine, efficient, low toxicity

Oxamniquine (UK 4271)

Coutinho, A.; Domingues, A. L. C.; and Bonfim, J. R. A., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 15-34
[Schistosoma] mansoni, human, oxamniquine, possible toxic side effects, clinical and parasitological action

Oxamniquine

Dias, L. C. de S.; et al., 1978, Rev. Saude Pub., S. Paulo, v. 12 (1), 110
Schistosoma mansoni, isolation of strain resistant to hycanthone and to oxamniquine

Oxamniquine

Domingues, A. L. C.; and Coutinho, A., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (3), 164-180
[Schistosoma] mansoni, humans, oral oxamniquine, extensive clinical trials: Brazil

Oxamniquine (UK-4271)

Eyakuse [sic], V. M., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 67-72
[Schistosoma] mansoni, human, oxamniquine, single dose of 30mg/kg (larger dose than previously recommended) intramuscularly, clinical trial

Oxamniquine (Vansil)

Eyakuze, V. M.; Rutasitara, W. K.; and Ndalaha, J. B., 1979, East African Med. J., v. 56 (1), 22-25
Schistosoma mansoni, schoolchildren and adults, oral oxamniquine, large field trial, drug feasible for routine treatment of people living in endemic areas: Nansio, Ukerewe, Tanzania

Oxamniquine

Farid, Z.; et al., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 501-502
Schistosoma mansoni, humans, advanced intestinal infections, oxamniquine therapy

Oxamniquine (UK-4271)

Foster, R., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 1-9
Schistosoma mansoni, S. haematobium, S. japonicum, oxamniquine, efficacy and toxicity in various exper. animals, clinical trials in humans

Oxamniquine

Girgis, N. I.; Mansour, N. S.; and Khalil, S. B., 1979, East African Med. J., v. 56 (9), 428-430
Schistosoma mansoni, hamsters, prophylactic-protective activity of oral oxamniquine

Oxamniquine

Guimaraes, R. X.; et al., 1979, AMB, Rev. Ass. Med. Brasil., v. 25 (2), 48-50
Schistosoma mansoni, human hepato-intestinal form, resistance to hycanthone and oxamniquine: Brazil

Oxamniquine -- Continued.

Oxamniquine

Higashi, G. I.; and Farid, Z., 1979, Brit. Med. J. (6194), v. 2, 830
Schistosoma mansoni, fever in oxamniquine-treated patients, cause uncertain

Oxamniquine (UK 4271)

Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167

Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria

Oxamniquine (U.K. 4271)

Kale, O. O.; and Lucas, A. O., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (1), 55-63
Schistosoma haematobium, humans, clinical trials, oral and parenteral oxamniquine, assessment of tolerance and toxicity: Western Nigeria

UK-4271

Katz, N., 1972, Rev. Ass. Med. Minas Gerais, v. 23 (1), 27-34
human schistosomiasis mansoni, outline for therapeutic trials with comparison of effects of hycanthon and experimental drug UK-4271

Oxamniquine

Katz, N.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (1), 25-29
Schistosoma mansoni, human, oxamniquine, clinical trials

Oxamniquine (U.K. 4271)

Katz, N.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 35-40
Schistosoma mansoni, human early phase or chronic infection, oxamniquine, clinical and laboratory aspects, high therapeutic activity and low toxicity

Oxamniquine

Katz, N.; et al., 1973, Rev. Soc. Brasil. Med. Trop., v. 7 (6), 381-387
Schistosoma mansoni, isolation of drug resistant strain (WW strain), reactions in mice to therapy with hycanthon, niridazole and oxamniquine compared with reactions of LE drug sensitive strain

Oxamniquine (U.K. 4271)

Katz, Naftale; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (5), 371-377
S[chistosoma] mansoni, humans, clinical trials with oral oxamniquine, side effects: Brazil

Oxamniquine (Mansil)

Keystone, J. S., 1978, Am. J. Trop. Med. and Hyg., v. 27 (2, pt. 1), 360-362
Schistosoma mansoni, humans, grand mal seizure and transient electroencephalograph changes associated with oxamniquine therapy, case reports

Oxamniquine embonate

Korolkovas, A.; Itaya, M.; and Coelho, Z. M. P., 1977, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 15 (1-2), 27-34
schistosomiasis, synthesis of 4 oxamniquine derivatives, oxamniquine embonate shows curative and prophylactic activity

Oxamniquine -- Continued.

Oxamniquine hydroxynaphthoate

Korolkovas, A.; Itaya, M.; and Coelho, Z. M. P., 1977, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 15 (1-2), 27-34
schistosomiasis, synthesis of 4 oxamniquine derivatives, oxamniquine embonate shows curative and prophylactic activity

Oxamniquine phthalate

Korolkovas, A.; Itaya, M.; and Coelho, Z. M. P., 1977, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 15 (1-2), 27-34
schistosomiasis, synthesis of 4 oxamniquine derivatives, oxamniquine embonate shows curative and prophylactic activity

Oxamniquine salicylate

Korolkovas, A.; Itaya, M.; and Coelho, Z. M. P., 1977, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 15 (1-2), 27-34
schistosomiasis, synthesis of 4 oxamniquine derivatives, oxamniquine embonate shows curative and prophylactic activity

Oxamniquine

Magzoub, M.; and Adam, S. E. I., 1978, J. Pharmacol. and Exper. Therap., v. 206 (2), 382-387
Schistosoma mansoni-infected Arvicanthus niloticus, evaluation of oral therapy with single and multiple doses of hycanthon and oxamniquine alone or in combination

Oxamniquine (Mansil)

Melo, A. H.; Pereira, L. H.; and Correa, M. C. R., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (2), 158-159
Schistosoma mansoni, mice, high doses of oxamniquine produced inhibition of cercaria-schistosomulum transformation, suggests that drug is active during process of host-larvae adaptation

Oxamniquine

Oliveira, M. A.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (5), 298-300
Schistosoma mansoni, mice treated with oxamniquine vs. untreated mice, effects of drug on parasite migration and development in host

Oxamniquine (Vansil)

Omer, A. H. S., 1978, Brit. Med. J. (6131), v. 2, 163-165
Schistosoma mansoni, humans, clinical trials testing efficacy and acceptability of oral oxamniquine to establish optimal regimen for use in the Sudan

Oxamniquine (UK 4271)

Ong, T. M., 1978, Mutation Research, v. 55 (1), 43-70
hycanthon and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Oxamniquine (U.K. 4271)

Pedro, R. de J.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 63-66
Schistosoma mansoni, human, oxamniquine

Oxamniquine -- Continued.

Oxamniquine

Pedro, R. de J.; et al., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (2), 130-137
Schistosoma mansoni, humans, oxamniquine, clinical trials, undesirable effects monitored

Oxamniquine

Pedro, R. de J.; et al., 1979, Rev. Brasil. Pesqui. Med. e Biol., v. 12 (2-3), 207-211
Schistosoma mansoni, observations on oxamniquine therapy: treatment of children, drug resistance of human strain as well as its resistance to hycanthone, hepatic histopathology during therapy, neurotoxic effects, treatment of mixed salmonellosis infection

Oxamniquine

Pellegrino, J.; Gilbert, B.; and Valadares, T. E., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (6), 456-458
Schistosoma mansoni, mice, topically applied oxamniquine, antischistosomal and chemoprophylactic activity

Oxamniquine

Pellegrino, J.; Katz, N.; and Dias, E. P., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 10-14
Schistosoma mansoni, oxamniquine, mice, hamsters and Cebus monkeys

Oxamniquine (U.K. 4271)

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (2), 97-101
Schistosoma mansoni, mice, preliminary laboratory trials of oxamniquine, antischistosomal activity more pronounced on early developing forms (up to 7 days) than on maturing worms

Oxamniquine

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Oxamniquine

Pereira, L. H.; et al., 1974, Rev. Inst. Med. Trop. S. Paulo, v. 16 (3), 123-126
Schistosoma mansoni, mice, simple and rapid method for mass screening of prophylactic agents using peritoneal schistosomula

Oxamniquine (Vansil)

Pitchford, R. J.; and Lewis, M., 1978, South African Med. J., v. 53 (17), 677-680
schistosomosis infections, human, clinical trials using oxamniquine; effective only with Schistosoma mansoni: lowveld of Eastern Transvaal

Oxamniquine (UK 4271)

Prata, A.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 47-57
Schistosoma mansoni, human acute or chronic infections, oxamniquine as single intramuscular dose, extensive clinical trial, main disadvantage is severe pain at site of injection

Oxamniquine -- Continued.

Oxamniquine

Rees, P. H.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 78-82
Schistosoma mansoni, school children, oxamniquine, clinical trials: Kenya

Oxamniquine

Shafei, A. Z., 1979, J. Trop. Med. and Hyg., v. 82 (1), 18-20
Schistosoma mansoni, humans, oral oxamniquine

Oxamniquine (UK 4271)

da Silva, L. C.; et al., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 suppl. 1 (6), 58-62
S[chistosoma] mansoni, human, non endemic area, oxamniquine, severe pain at injection site

Oxamniquine (UK 4271)

da Silva, L. C.; et al., 1974, Rev. Inst. Med. Trop. S. Paulo, v. 16 (2), 103-109
S[chistosoma] mansoni, human, oxamniquine, clinical trials, oral administration

Oxamniquine (UK 4271)

da Silva, L. C.; et al., 1975, Rev. Inst. Med. Trop. S. Paulo, v. 17 (5), 307-311
S[chistosoma] mansoni, human, oral oxamniquine, large-scale clinical trials

Oxamniquine (Mansil)

Woolhouse, N. M., 1979, Biochem. Pharmacol., v. 28 (16), 2413-2418
antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Oxamniquine embonate. See Oxamniquine.

Oxamniquine hydroxynaphthoate. See Oxamniquine.

Oxamniquine phthalate. See Oxamniquine.

Oxamniquine salicylate. See Oxamniquine.

Oxantel -- CP-14,445; Oxantel pamoate; Oxantel tartrate; OxypyranTEL; OxypyranTEL pamoate; trans-1,4,5,6-Tetrahydro-2-(3-hydroxystyryl)-1-methyl pyrimidine; trans-1,4,5,6-Tetrahydro-2-(trans-3-hydroxystyryl)-1-methyl pyrimidine; TricoCEL.

OxypyranTEL

Amato Neto, V.; et al., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (4), 261-263
Trichocephalus trichiurus, humans, oxypyranTEL, efficacy trials

Oxantel

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Oxantel -- Continued.

Oxantel pamoate

Cho, S. Y., 1976, Taehan Yangnihak Chapchi (Korean J. Pharmacol.), v. 12 (1), 69-73
nematodes, human, evaluation of efficacy of combined therapy of pyrantel pamoate and oxantel pamoate

Oxypyranter (Tricocel)

Cimerman, B.; et al., 1978, Rev. Brasil. Med., v. 35 (3), 201-204
Trichuris trichiura, humans, therapeutic trials with oxypyranter, minor side effects

Oxantel pamoate (CP-14,445)

Lim, J. K., 1974, Taehan Yangnihak Chapchi (Korean J. Pharmacol.), v. 10 (2), 97-101
Trichocephalus trichiurus, soldiers and inpatients at Armed Forces Hospital, evaluation of anthelmintic effect and tolerance of oxantel pamoate: Korea

Oxantel (CP-14-445)

Paul, F. M.; and Zaman, V., 1976, Singapore Med. J., v. 17 (4), 219-224
Trichuris trichiura, children with gastrointestinal symptoms, protein caloric malnutrition, and nutritional anemia, clinical trials with oxantel, well tolerated: Singapore General Hospital

Oxantel tartrate

Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235
Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parbendazole and dichlorvos

Oxypyranter pamoate (Tricocel)

Rodrigues, L. D.; Martirani, I.; and Cabeca, M., 1975, Rev. Brasil. Clin. e Terap., v. 4 (11), 459-462
Trichuris trichiura, humans, double-blind clinical trial with oxypyranter pamoate, good results

Oxantel pamoate. See Oxantel.

Oxantel tartrate. See Oxantel.

Oxfendazole -- Methyl(5-(phenylsulfinyl)-1H-benzimidazol-2-yl) carbamate; Methyl 5(6)-phenyl-sulfinyl-2-benzimidazole-carbamate; RS-8858; Synanthic; Systemex.

Oxfendazole

Anderson, N.; and Lord, V., 1979, Austral. Vet. J., v. 55 (4), 158-162
Ostertagia ostertagi, Trichostrongylus axei, cattle, fenbendazole, oxfendazole, and levamisole compared

Oxfendazole -- Continued.

Oxfendazole (Synanthic;Systemex)

Armour, J.; Duncan, J. L.; and Reid, J. F. S., 1978, Vet. Rec., v. 102 (12), 263-264
Ostertagia ostertagi and Cooperia oncophora, calves, oxfendazole, highly effective against arrested larvae and adults as well as adult stages of Trichostrongylus axei and Nematodirus helvetianus

Oxfendazole

Baker, N. F.; Fisk, R. A.; and Miller, J. E., 1978, Am. J. Vet. Research, v. 39 (8), 1258-1261
gastrointestinal helminths, calves, oxfendazole as drench, paste or bolus, very effective

Oxfendazole

Borgsteede, F. H. M., 1977, Tijdschr. Diergeneesk., v. 102 (14), 801-804
gastro-intestinal helminths, calves, field trials with oxfendazole

Oxfendazole

Chalmers, K., 1978, N. Zealand Vet. J., v. 26 (6), 162-164
nematodes, cattle, efficacy of oxfendazole given to coincide with autumn-winter and winter-spring larval-inhibition peak: New Zealand

Oxfendazole

Chalmers, K., 1979, N. Zealand Vet. J., v. 27 (1-2), 8, 13
Dictyocaulus viviparus, Friesian calves, and D. filaria, sheep, oxfendazole: New Zealand

Oxfendazole (Synanthic)

Chalmers, K., 1979, N. Zealand J. Exper. Agric., v. 7 (2), 111-114
gastrointestinal nematodes, lambs, calves (both exper.), efficacy of oxfendazole

Oxfendazole

Colglazier, M. L., 1979, Am. J. Vet. Research, v. 40 (3), 384-386
gastrointestinal helminths and stomach bots, ponies, critical trials with oxfendazole and caviaphos; observations on spontaneous elimination of small strongylids prior to treatment: Maryland

Oxfendazole (RS-8858)

Corwin, R. M.; Kennedy, J. A.; and Pratt, S. E., 1979, Am. J. Vet. Research, v. 40 (2), 297-298
nematodes, pigs, oxfendazole, dose titration

Oxfendazole

Douch, P. G. C.; and Buchanan, L. L., 1979, Xenobiotica, v. 9 (11), 675-679
Moniezia expansa, Ascaris suum, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Oxfendazole (Synanthic)

Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1979, Am. J. Vet. Research, v. 40 (4), 590-594
large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Oxfendazole -- Continued.

- Oxfendazole (Synanthic; Systamex)
Duncan, J. L.; and Reid, J. F. S., 1978, Vet. Rec., v. 103 (15), 332-334
gastrointestinal nematodes, ponies, oxfendazole, anthelmintic efficacy against adult and immature stages
- Oxfendazole
Gemmell, M. A.; Johnstone, P. D.; and Oudemans, G., 1979, Research Vet. Sc., v. 26 (3), 389-390
Echinococcus granulosus, Taenia hydatigena, dogs, oxfendazole
- Oxfendazole
Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367
Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate
- Oxfendazole
Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of Nematospiroides dubius in mice
- Oxfendazole
Kistner, T. P.; et al., 1979, Vet. Parasitol., v. 5 (2-3), 195-204
gastrointestinal and lungworm helminths, sheep, oxfendazole, dose titration study
- Oxfendazole
Kistner, T. P.; and Wyse, D., 1978, Austral. Vet. J., v. 54 (10), 469-470
Haemonchus contortus benzimidazole-resistant strain, sheep (exper.), oxfendazole 100% effective against 3rd, 4th, early 5th, and adult stages
- Oxfendazole (Systamex)
Le Jambre, L. F., 1979, Austral. Vet. J., v. 55 (2), 65-67
Ostertagia circumcincta, O. trifurcata, sheep (exper.), effectiveness of levamisole, thiabendazole, albendazole, and oxfendazole against levamisole-resistant strains
- Oxfendazole (Synanthic; Systamex)
Marriner, S.; and Bogan, J. A., 1979, Vet. Rec., v. 105 (11), 261
benzimidazole anthelmintics, sheep, oral vs. intraruminal vs. intra-abomasal administration
- Oxfendazole (Systamex)
Michael, S. A.; et al., 1979, Vet. Rec., v. 104 (15), 338-340
nematodes, cestodes, indigenous Egyptian sheep, oxfendazole, field trial: Nile Delta region

Oxfendazole -- Continued.

- Oxfendazole (Systamex)
Michael, S. A.; Higgins, A. J.; and El Refaii, A. H., 1979, Trop. Animal Health and Prod., v. 11 (2), 63-68
gastrointestinal nematodes, Egyptian goats (exper.), oxfendazole
- Oxfendazole
Nerenberg, C.; Runkel, R. A.; and Matin, S. B., 1978, J. Pharm. Sc., v. 67 (11), 1553-1557
oxfendazole in bovine, equine, or canine plasma or serum, radioimmunoassay for determination of certain pharmacokinetic parameters
- Oxfendazole
Nicholas, W. L.; and Stewart, A. C., 1979, Ann. Trop. Med. and Parasitol., v. 73 (1), 57-62
Toxocara canis, laboratory mouse, fenbendazole and oxfendazole killed larvae in brains and musculature, migratory larvae more susceptible, possible use in preventing pre-natal infection in dogs
- Oxfendazole
Ogunsusi, R. A., 1979, Research Vet. Sc., v. 27 (1), 131-132
Haemonchus contortus, sheep, oxfendazole, haloxon, efficacy against arrested larvae, controlled trial, dry season: northern Nigeria
- Oxfendazole (Systamex; Synanthic)
Piercy, D. W. T.; Reynolds, J.; and Brown, P. R. M., 1979, Brit. Vet. J., v. 135 (5), 405-410
oxfendazole, reproductive safety in sheep and cattle
- Oxfendazole
Prichard, R. K.; Hennessy, D. R.; and Steel, J. W., 1978, Vet. Parasitol., v. 4 (4), 309-315
Haemonchus contortus, Trichostrongylus colubriformis, sheep, Ostertagia ostertagi, cattle, 4 benzimidazoles, mode of action and pharmacokinetic behavior, implications for prolonged administration as a new concept for increasing spectrum and effectiveness of anthelmintics
- Oxfendazole
Todd, K. S., jr.; and Mansfield, M. E., 1979, Am. J. Vet. Research, v. 40 (3), 423-424
nematodes, calves, oxfendazole in 4 formulations
- Oxfendazole (Systamex)
Webb, R. F.; and McCully, C. H., 1979, Austral. Vet. J., v. 55 (7), 347-348 [Letter]
Haemonchus contortus, sheep, resistance to oxfendazole: New South Wales
- Oxfendazole
Wescott, R. B.; Shelton, T. A.; and Gates, N. L., 1979, West. Veterinarian, v. 17 (2), 22-23
gastrointestinal nematodes and cestodes, sheep, 3 field trials with oxfendazole, effective; no apparent effect on coccidial oocysts

Oxibendazole -- Anthelcide-EQ; 2-Carbamomethoxy-5-propoxybenzimidazole; Methyl-5-p-oxy-2-benzimidazole-carbamate; Methyl 5-n-propoxy-2-benzimidazole carbamate; Oxibendazole feed premix.

Oxibendazole
Coles, G. C.; and Briscoe, M. G., 1978, Vet. Rec., v. 103 (16), 360-361 [Letter]
Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for Haemonchus contortus eggs

Oxibendazole
Drudge, J. H.; et al., 1979, Am. J. Vet. Research, v. 40 (6), 758-761
Parascaris equorum and other horse parasites, oxibendazole, critical tests and clinical trials; febantel paste

Oxibendazole (Anthelcide-EQ)
Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1979, Am. J. Vet. Research, v. 40 (4), 590-594
large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Oxibendazole
Friedman, P. A.; and Platzner, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Oxibendazole
Girardi, C.; Gallo, M. G.; and Lanfranchi, P., 1977, Ann. Fac. Med. Vet. Torino, v. 24, 201-213
gastrointestinal nematodes, cestodes, sheep, oxibendazole, controlled test, very effective except for Trichuris ovis and cestodes

Oxibendazole
Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367
Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Oxibendazole
Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, Research Vet. Sc., v. 25 (3), 360-363
Haemonchus contortus, Trichostrongylus colubriformis, levels of benzimidazole resistance recorded from an egg hatch test procedure

Oxibendazole
Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of Nematospiroides dubius in mice

Oxibendazole -- Continued.

Oxibendazole
Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Oxibendazole (Anthelcide EQ)
Lock, T. F.; et al., 1979, Vet. Med. and Small Animal Clin., v. 74 (9), 1247, 1250
nematodes, horses, oxibendazole, clinical trials

Oxibendazol
Ramajo Martin, V.; Simon Vicente, F.; and Reyes, R., 1977, Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C., [v. 3], 1976, 53-58
Oesophagostomum dentatum, pigs (exper.), oxibendazol and parbendazol, 100% effective

Oxibendazole
Rodriguez Caabeiro, F.; Martinez Fernandez, A. R.; and Sanmartin Duran, M. L., 1978, Rev. Iber. Parasitol., v. 38 (3-4), 623-638
Trichinella spiralis, mice (exper.), oxibendazole

Oxibendazole
Simpkin, K. G.; and Coles, G. C., 1978, Research Vet. Sc., v. 25 (2), 249-250
Haemonchus and Trichostrongylus colubriformis in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parbendazole, oxibendazole)

Oxibendazole feed premix
Williams, J. C.; Sheehan, D.; and Fuselier, R., 1978, Proc. Helminth. Soc. Washington, v. 45 (1), 129-131
gastrointestinal parasites, cattle, oxibendazole, controlled test

Oxibendazole feed premix. See Oxibendazole.

Oxide
Daniiarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Oxinothiofios. See Quintiofos.

Oxinothiophos. See Quintiofos.

Oxipurinol. See Oxypurinol.

Oxophenarsine -- Mapharsemin.

Mapharsemin
Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Oxophenarsine -- Continued.

Oxophenarsine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

N,N'-[Oxybis(2,1-ethanedioxy-4,1-phenylene)] bisacetamide. See Diamphenethide.

Oxychloroquine -- SN-8137.

SN-8137

Schmidt, L. H.; et al., 1977, *Antimicrob. Agents and Chemotherapy*, v. 11 (5), 826-843
Plasmodium falciparum in *Aotus trivirgatus*, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

Oxyclozanide -- Nilzan (with Tetramisole); 3,3',5,5',6-Pentachloro-2,2'-dihydroxybenzanilide; 3,3',5,5',6-Pentachloro-2'-hydroxysalicylanilide; Zanil.

Oxyclozanide (Zanil)

Foreyt, W. J.; and Todd, A. C., 1974, *Am. J. Vet. Research*, v. 35 (3), 375-377
Fascioloides magna, cattle, efficacies of rafoxanide and oxyclozanide: Texas Gulf Coast Region

Zanil (Oxyclozanid)

Furmaga, S.; Gundlach, J. L.; and Sobieszewski, K., 1974, *Acta Parasitol. Polon.*, v. 22 (12-21), 229-247
Fasciola hepatica-infected and uninfected sheep treated and not treated with 3 fasciolicides (fasciolin, distolon, and zanil), blood serum levels of Ca, P, Na, K, and Mg

Zanil (Oxyclozanid)

Krzyzanowski, J., 1977, *Polskie Arch. Wet.*, v. 20 (1), 17-32
 zanil, bulls, no negative influence on quality or fertilizing capacity of semen, no teratogenic action in offspring of bulls

Nilzan

McEwan, A. D.; and Oakley, G. A., 1978, *Vet. Rec.*, v. 102 (14), 314-315
 panacur, thiabendazole, and nilzan with dye marker added, oral dosing of cattle showed evidence of rumen by-pass, reduced drug efficacy probably resulting from closure of oesophageal groove

Oxyclozanide -- Continued.

Zanil (Oxyclozanide)

Nozdryn-Plotnicki, Z.; and Owczarewicz, A., 1977, *Polskie Arch. Wet.*, v. 20 (3), 185-191
 zanil, rabbits, morphological and histochemical changes in kidneys

Oxyclozanide (Zanil)

Parshad, V. R.; and Guraya, S. S., 1978, *Vet. Parasitol.*, v. 4 (2), 111-120
 4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces

3,3',5,5',6-Pentachloro-2,2'-dihydroxybenzanilide
 Sakamoto, T.; and Gemmill, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Oxyclozanide (Zanil)

Venkat Reddy, P.; Murthy, D. K.; and Subramanyam, S., 1977, *Indian J. Exper. Biol.*, v. 15 (11), 1073-1074
 oxyclozanide, treatment of roots of *Allium cepa* induced mitotic aberrations

Oxyclozanide

Vodrazka, J.; et al., 1979, *Biol. a Chem. Zivoc. Vyroby, Vet.*, v. 15 (v. 21) (4), 363-374
 tribromsalan, dynamic behavior in sheep blood plasma; antifasciolicides, effects on some sheep plasma enzymes

Oxyphenonium bromide -- Mexaform (with Iodochlorhydroxyquin and Phanquone).

Mexaform

Fuchs, P., 1978, *Deutsche Med. Wchnschr.*, v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Mexaform

Rostkowska, J., 1970, *Acta Parasitol. Polon.*, v. 18 (27-41), 377-392
Balantidium coli, effect of various bacteria on propagation in vitro, on erythrophagocytic capability of balantidia, and on susceptibility of balantidia to atebirin, entobex, mexaform, and protargol; *Trichomonas hominis*, *Chilomastix mesnili*, and *Dientamoeba fragilis* found to be without effect; effect of balantidia on bacteria

Mexaform

Trzaska, B.; and Leinweber-Noiszewska, H., 1974, *Ann. Acad. Med. Stetinensis*, suppl. 10, 19-25
Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis

Oxypurinol -- 4,6-Dihydroxypyrazolo(3,4-d)pyrimidine; Oxipurinol.

Oxypurinol

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in *in vitro* screens to test wide range of compounds for chemotherapeutic activity

Oxipurinol

Nelson, D. J.; et al., 1979, J. Biol. Chem., v. 254 (10), 3959-3964

Leishmania braziliensis, *L. donovani*, pyrazolo(3,4-d)pyrimidines, metabolism, possible explanation for antileishmanial activity

Oxypyrantel. See Oxantel.

Oxypyrantel pamoate. See Oxantel.

Oxytetracycline -- Emicina; Liquamycin; Liquamycin injectable L-50; Liquamycin/LA-200; Liquamycin (T-50); Oxytetracycline chloride; Oxytetracycline HCl; Oxytetracycline hydrochloride; Terragluclin; Terramycin; Terramycin 100; Terramycin injectable long acting; Terramycin injectable solution; Terramycin/LA; Terramycin/LA (T-200); Oxytetracycline, long-acting.

Oxytetracycline (Terramycin)

Brown, C. G. D.; et al., 1977, Tropenmed. u. Parasitol., v. 28 (4), 513-520

Theileria parva, *Bos taurus* with patent East Coast fever induced by inoculation of cryopreserved stabilates of tick-derived infective particles, preliminary evaluation of n-pyrrolidinomethyl tetracycline and oxytetracycline

Oxytetracycline

Gill, B. S.; et al., 1978, Internat. J. Parasitol., v. 8 (6), 467-469

Theileria annulata, calves, immunization by treating tick (*Hyalomma anatolicum anatolicum*) stabilate-induced infections with 1 or 2 doses of long-acting oxytetracycline vs. 8 doses of chlortetracycline

Oxytetracycline

Gottschall, J. L.; et al., 1979, Lab. Invest., v. 41 (1), 5-12

morphologic changes of rat type II pneumocytes induced by oxytetracycline, may be important in interpreting studies of experimental infections with *Pneumocystis carinii*

Oxytetracycline

Henry, S. C., 1979, J. Am. Vet. Med. Ass., v. 174 (6), 601-603

Eperythrozoon suis, swine, clinical signs of infection confirmed by indirect hemagglutination and measuring packed cell volume; oxytetracycline and arsenic acid combined with lice control, arsenic toxicosis

Oxytetracycline -- Continued.

Oxytetracycline hydrochloride (Terramycin)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, Chemotherapy, v. 25 (4), 222-226

Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Oxytetracycline (Terramycin)

Kumar, B. J.; and Gowda, D. K. V., 1978, Indian Vet. J., v. 55 (9), 729-730

theileriasis, dairy cow treated with chloroquine and oxytetracycline, chloroquine toxicity causing corneal opacity and possibly abortion

Oxytetracycline

Kuttler, K. L.; and Simpson, J. E., 1978, Am. J. Vet. Research, v. 39 (2), 347-349

Anaplasma marginale, splenectomized calves (exper.), 2 formulations of oxytetracycline (T-200 and T-50), doxycycline, drug efficacies, influence of treatment on parasitemia and packed cell volume

Oxytetracycline (Liquamycin (T-50))

Kuttler, K. L.; Young, M. F.; and Simpson, J. E., 1978, Vet. Med. and Small Animal Clin., v. 73 (2), 187, 190-192

Anaplasma marginale, cows (exper.), relative efficacy of a long-acting oxytetracycline (terracycline) compared with a presently available oxytetracycline (liquamycin), both formulations highly effective and resulted in rapid recovery

Oxytetracycline (Terramycin/LA (T-200))

Kuttler, K. L.; Young, M. F.; and Simpson, J. E., 1978, Vet. Med. and Small Animal Clin., v. 73 (2), 187, 190-192

Anaplasma marginale, cows (exper.), relative efficacy of a long-acting oxytetracycline (terracycline) compared with a presently available oxytetracycline (liquamycin), both formulations highly effective and resulted in rapid recovery

Oxytetracycline

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in *in vitro* screens to test wide range of compounds for chemotherapeutic activity

Oxytetracycline HCl

Magonigle, R. A.; Simpson, J. E.; and Frank, F. W., 1978, Am. J. Vet. Research, v. 39 (9), 1407-1410

Anaplasma marginale, cattle (exper.), efficacy of new long-acting oxytetracycline formulation (Liquamycin/LA-200), 1 intramuscular injection, comparison with Liquamycin injectable L-50 administered on 2 consecutive days

Oxytetracycline -- Continued.

- Oxytetracycline hydrochloride (Terramycin)
Mallick, K. P.; Dwivedi, S. K.; and Malhotra, M. N., 1979, Indian Vet. J., v. 56 (8), 693-694
Anaplasma marginale, indigenous goats, 5 clinical cases, oxytetracycline hydrochloride: India
- Oxytetracycline + Clioquinol
Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire
- Oxytetracycline + Dehydroemetine + Di-iodohydroxyquinoline
Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire
- Oxytetracycline + Di-iodohydroxyquinoline
Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire
- Oxytetracycline + Metronidazole
Masters, D. K.; and Hopkins, A. D., 1979, J. Trop. Med. and Hyg., v. 82 (5), 99-101
Entamoeba histolytica, human, comparative trial of 4 amoebicide regimes, recommendations for use in tropical rural hospital: Zaire
- Oxytetracycline (Terramycin)
Mchinja, S. J.; Young, A. S.; and Omwoyo, P., 1975, Bull. Animal Health and Prod. Africa, v. 25 (2), 158-161
Theileria parva, inoculation of oxytetracycline into rabbits on which infected Rhipicephalus appendiculatus are feeding did not inhibit development of parasites nor affect their subsequent infectivity for cattle, possible use of this system in screening prophylactic drugs against East Coast fever
- Terramycin
Mishra, A. K.; and Sharma, N. N., 1979, Trop. Animal Health and Prod., v. 11 (4), 222-226
Anaplasma marginale, calves (exper.), comparative efficacy of several drugs
- Oxytetracycline (Terragluclin)
Mordasov, P. M.; Golovnev, V. I.; and Zakharik, N. V., 1977, Vet. Nauka--Proizvod., Trudy, Minsk, v. 15, 93-95
anaplasmosis, bovine, treatment with oxytetracycline
- Oxytetracycline
Radley, D. E.; et al., 1979, Vet. Parasitol., v. 5 (2-3), 117-128
Theileria lawrencei, immunization of Bos taurus by infection (with single and multiple Theileria spp. isolates) and chemoprophylaxis (long-acting oxytetracycline)

Oxytetracycline -- Continued.

- Oxytetracycline hydrochloride
Renshaw, H. W.; et al., 1976, Proc. 80. Ann. Meet. U. S. Animal Health Ass., 79-88
Anaplasma marginale, cattle, some long-lasting immunity persists after elimination of carrier status with oxytetracycline hydrochloride
- Oxytetracycline
Reshetniak, V. Z.; Bartenev, V. S.; and Firsov, N. F., 1977, Veterinariia, Moskva (10), 79
anaplasmosis, sheep outbreak, pathology, oxytetracycline effective treatment; Hyalomma scupense as probable vector, hexachlorane in creolin emulsion for control; no further anaplasmosis or H. scupense found: Rostovsk oblast
- Oxytetracycline (Liquamycin; Terramycin)
Roby, T. O.; Simpson, J. E.; and Amerault, T. E., 1978, Am. J. Vet. Research, v. 39 (7), 1115-1116
Anaplasma marginale, calves (exper.), elimination of carrier state using long-acting formulation of oxytetracycline
- Oxytetracycline (Terramycin injectable solution)
Singh, B.; Gautam, O. P.; and Banerjee, D. P., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 35-39
Anaplasma, cattle and sheep (both exper.), blood changes before and after treatment with terramycin injectable solution
- Oxytetracycline (Terramycin injectable long acting; Terramycin 100)
Stewart, C. G.; et al., 1979, J. South African Vet. Ass., v. 50 (2), 83-85
Anaplasma marginale, splenectomized calves (nat. and exper.), oxytetracycline, long- vs. short-acting formulation
- Oxytetracycline (Terramycin)
Sweet, V. H.; and Stauber, E. H., 1978, J. Am. Vet. Med. Ass., v. 172 (11), 1310-1312
Anaplasma marginale, cattle, serologic survey, chlortetracycline, oxytetracycline, good results, concluded that test and treatment program would need to be continued for several years to achieve anaplasmosis-free status: northern Idaho; southeastern Washington
- Oxytetracycline hydrochloride (Liquamycin)
Swift, B. L.; Settlemire, J., jr.; and Thomas, G. M., 1978, Theriogenology, v. 10 (6), 481-485
Anaplasma marginale, pregnant heifers (exper.), oxytetracycline hydrochloride, did not abort and transplacental transmission did not occur
- Emicina
Thompson, K. C.; et al., 1978, Trop. Animal Health and Prod., v. 10 (2), 75-81
Anaplasma marginale, Babesia argentina, B. bigemina, cattle under tropical conditions, immunization with virulent organisms followed by drug therapy (ganaseg; gloxazone; emicina) vs. chemoprophylaxis (imidocarb); tick and gastrointestinal parasite control without haemoparasitic control had advantage over no control system at all

Oxytetracycline -- Continued.

- Oxytetracycline (Terramycin/LA; Emicina)
 Todorovic, R. A.; Gonzalez, E. F.; and Garcia, O., 1979, Tropenmed. u. Parasitol., v. 30 (2), 236-238
Anaplasma marginale, cattle (exper.), new long-acting injectable oxytetracycline (terramycin/LA) compared with commercial preparation (emicina)
- Oxytetracycline
 Uilenberg, G.; van Vorstenbosch, C. J. A. H. V.; and Perie, N. M., 1979, Vet. Quart., v. 1 (1), 14-22
Anaplasma mesaeterum sp. n., sheep (nat. and exper.) and goats (exper.), pathogenicity, oxytetracycline treatment, cross-immunity tests with *A. ovis*: Ameland, the Netherlands
- Oxytetracycline
 Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures
- Oxytetracycline
 Wang, L. T.; and Yang, S. P., 1971, Taiwan i Hsueh Hui Tsa Chih (J. Formosan Med. Ass.), v. 70 (3), 131-134
Entamoeba histolytica, human, oxytetracycline-resistant amoebic dysentery: Taiwan

Oxytetracycline -- Continued.

- Oxytetracycline hydrochloride
 West, H. J., 1979, J. Small Animal Practice, v. 20 (9), 543-549
Haemobartonella canis, dog, case report, oxytetracycline hydrochloride
- Oxytetracycline, long-acting (Terramycin L/A T-200)
 Wilson, A. J.; et al., 1979, Austral. Vet. J., v. 55 (2), 71-73
Anaplasma marginale, cattle and splenectomised calves (exper.), long-acting formulation of tetracycline compared with standard tetracycline and imidocarb dipropionate
-
- Oxytetracycline chlorhydrate. See Oxytetracycline.
- Oxytetracycline HCl. See Oxytetracycline.
- Oxytetracycline hydrochloride. See Oxytetracycline.
- Oxytetracycline, long-acting. See Oxytetracycline.

Pacprim. See Sulfadimethoxine or Sulfisomidine or Sulfisoxazole or Trimethoprim.

Palasonin. See Butea frondosa.

Paludrine. See Chlorguanide.

Pamaquine

Hayashi, T.; et al., 1978, Bull. Fac. Agric. Tottori Univ., v. 30, 82-88

Theileria sergenti, grazing cattle, administration of pamaquine probably causes blood coagulation disorders

Pamaquine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Panacur. See Fenbendazole.

Pancoxin. See Amprolium or Ethopabate or Sulfaquinoxaline.

Pancoxin plus. See Amprolium or Ethopabate or Pyrimethamine or Sulfaquinoxaline.

Pantelmin. See Mebendazole.

Paracetamol

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Paramar M-50

Khan, M. H., 1979, Indian Vet. J., v. 56 (9), 739-743

Lipeurus caponis, Menacanthus stramineus, White Leghorn fowl, organophosphorus insecticides, costs evaluated

Paramite. See Phosmet.

Pararosaniline -- Pararosaniline salts and derivatives; TAC; Tris (p-aminophenyl) carbonium; Tris (p-aminophenyl) carbonium pamoate.

Tris (p-aminophenyl) carbonium

Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124

preparation and testing of 6 long-acting schistosomicidal resins

Pararosaniline salts and derivatives

Korolkovas, A.; and Pellegrino, J., 1977, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 15 (1-2), 5-9

Schistosoma mansoni, mice, pararosaniline salts and derivatives synthesized and tested

Pararosaniline -- Continued.

Tris (p-aminophenyl) carbonium pamoate (TAC) Monteiro, W.; Pellegrino, J.; and da Silva, M. L. H., 1969, Rev. Brasil. Pesqui. Med. e Biol., v. 2 (1), 45-49

Schistosoma mansoni-infected mice, physiological and morphological changes in parasite egg formation after mice were treated with one of 7 known antischistosomal drugs

Pararosaniline salts and derivatives. See Pararosaniline.

Parbendazole -- 2-Carbamomethoxy-5-butylbenzimidazole; Helatac; Helmatac; Methyl 5-butyl-2-benzimidazole carbamate; Methyl-5-6-butyl 1-2-benzimidazole carbamate; Parbendazole premix; SKF 29044; Wormguard.

Parbendazole

Callinan, A. P. L.; and Cummins, L. J., 1979, Austral. Vet. J., v. 55 (8), 370-373
nematodes, cattle (nat. and exper.), efficacies of various anthelmintics against adult and larval stages: western Victoria

Parbendazole (Wormguard)

Chevis, R. A. F.; and Kelly, J. D., 1978, N. Zealand Vet. J., v. 26 (9), 218-220

Fasciola hepatica, rats and sheep (both exper.), mebendazole, parbendazole, cambendazole, thiabendazole, anthelmintic activity, molecular structure-activity analyses

Parbendazole

Coles, G. C.; and Briscoe, M. G., 1978, Vet. Rec., v. 103 (16), 360-361 [Letter]

Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for Haemonchus contortus eggs

Parbendazole

Friedman, P. A.; and Platzer, E. G., 1978, Biochim. et Biophys. Acta, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Helmatac (Parbendazole)

Gaur, S. N. S.; and Dutt, S. C., 1979, Indian Vet. J., v. 56 (1), 62-64

Mecistocirrus digitatus, cattle, comparative efficacy of thibendole, nilverm, and helmatac

Parbendazole

Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367

Haemonchus contortus, Trichostrongylus colubriformis, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Parbendazole

Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, Research Vet. Sc., v. 25 (3), 360-363

Haemonchus contortus, Trichostrongylus colubriformis, levels of benzimidazole resistance recorded from an egg hatch test procedure

Parbendazole -- Continued.

Parbendazole

Ireland, C. M.; et al., 1979, *Biochem. Pharmacol.*, v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiroides dubius* in mice

Parbendazole (SKF 29044)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, *Ztschr. Parasitenk.*, v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Parbendazol

Ramajo Martin, V.; Simon Vicente, F.; and Reyes, R., 1977, *Anuario Cent. Edafol. y Biol. Aplic. C.S.I.C.*, [v. 3], 1976, 53-58
Oesophagostomum dentatum, pigs (exper.), oxibendazol and parbendazol, 100% effective

Parbendazole

Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (1), 55-58
Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (*Mesocricetus auratus*)

Parbendazole (Wormguard)

Robinson, M., 1979, *Vet. Parasitol.*, v. 5 (2-3), 223-235
Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parbendazole and dichlorvos

Parbendazole

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Parbendazole

Sanchez Moreno, M.; and Barrett, J., 1979, *Parasitology*, v. 78 (1), 1-5
Hymenolepis diminuta, adults, monoamine oxidase, occurrence and properties, inhibition by several anthelmintics

Parbendazole

Santos, L. E.; et al., 1977, *Rev. Fac. Med. Vet. e Zootec. Univ. S. Paulo*, v. 14 (1), 149-169
treated and untreated Guzera heifers compared for weight gains, hemoglobin levels, and helminth infestation: region of Sertaozinho, Sao Paulo, Brazil

Parbendazole premix (Helatac)

Sarup, S.; Banerjee, D. P.; and Gautam, O. P., 1978, *Indian Vet. J.*, v. 55 (6), 497-498
Ancylostoma caninum, ova and infective larvae, in vitro evaluation of fenbendazole, helatac, alcopar, banminth II, only banminth II effective; banminth II-treated larvae administered orally to mice, none recovered from lungs or liver

Parbendazole -- Continued.

Parbendazole

Simpkin, K. G.; and Coles, G. C., 1978, *Research Vet. Sc.*, v. 25 (2), 249-250
Haemonchus and *Trichostrongylus colubriformis* in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parbendazole, oxibendazole)

Parbendazole

Spaldonova, R.; and Hovorka, J., 1972, *Ang. Parasitol.*, v. 13 (4), 207-213
Trichinella spiralis, white mice (exper.), parbendazole, doses of 100 mg/kg effective in every stage of infection, reduced efficacy when period of administration was less than 5 days

Parbendazole

Ueno, H.; and Chibana, T., 1978, *Japan Agric. Research Quart.*, v. 12 (3), 152-156
Stephanofilaria okinawaensis, cattle, distribution, clinical signs, chemotherapy, intermediate host determined

Parbendazole (Helatac)

Varshney, T. R.; and Singh, Y. P., 1979, *Indian Vet. J.*, v. 56 (3), 207-210
gastrointestinal helminths, sheep, naphthalophos more effective than parbendazole in controlled study under farm conditions

Helatac (Parbendazole)

Vihan, V. S.; and Sahni, K. L., 1979, *Indian J. Animal Research*, v. 13 (1), 15-18
gastrointestinal nematodes, lambs, efficacy of helatac, nilverm, thiabendole, and banminth-II: Makhdoom, Mathura (U. P.)

Parbendazole (Worm-Guard)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, *Austral. Vet. J.*, v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiabendazole, results confirm the usefulness of levamisole, naphthalophos, and raxofanide for this purpose, haloxon and nitroxylin are also useful chemical alternatives

Parbendazole premix. See Parbendazole.

Paromomycin -- Aminosidine; Aminosidine sulfate; Gabroral; Humatin; Monomycine.

Monomycine

Akhound-Zadeh, H., 1976, *Rev. Internat. Serv. Sante Armees*, v. 49 (5), 421-426
chronic cutaneous leishmaniasis, soldier with severe ulcers that did not heal despite 8 years of therapy with various anti-leishmanial drugs, chronicity thought to be result of immuno-deficiency, ulcers finally cured after additional therapy with monomycine: Iran (had travelled to Khuzistan)

Paromomycin -- Continued.

Paromomycin

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Aminosidine

Cerecedo Cortina, V.; and Gaxiola Gaxiola, R., 1972, Rev. Fac. Med., Univ. Nac. Auton. Mexico, an. 15, v. 15 (1), 25-28
human hepatic amoebic abscess, side effects of currently used amoebicides (emetine, chloroquine, metronidazole, aminosidine)

Aminosidine sulfate (Gaboral)

Fernandez Hernandez, H.; Porro Abdo, Y.; and Ciolli Martinez, M., 1977, Rev. Cubana Med. Trop., v. 29 (3), 153-161
Amoeba histolytica, children with acute diarrhea, clinical trials testing efficacy of aminosidine sulfate: Cuba

Paromomycin (Humatin)

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Paromomycin (Humatin)

Nagington, J., 1975, Tr. Ophth. Soc. United Kingdom, v. 95 (2), 207-209
Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity

Paromomycin

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351
Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Paromomycin

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Paromomycin (Humatin)

Singh, U. S.; Raizada, M. K.; and Rao, V. K. M., 1978, Zentralbl. Bakteriolog., 1 Abt. Orig., Reihe A, v. 241 (3), 358-367
Hartmannella culbertsoni, axenically grown, purification and properties of L-histidine ammonia-lyase, marked inhibitory effect of certain amoebicidal drugs and divalent cations

Peganum harmala alkaloids

Vecherkin, S. S.; et al., 1977, Veterinariia, Moskva (10), 77-78
theileriasis, cattle, Peganum harmala alkaloids effective, no harmful or cumulative effects

Penicillamine

Khayyal, M. T.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 149-156
S[chistosoma] mansoni, mice, antimony potassium tartrate therapy given with penicillamine as adjuvant gives same therapeutic results with less side effects; ameliorates lipid changes in host but not in parasites

Penicillin -- Penicillin G (K salt); Procaine penicillin.

Penicillin

Evans, W. S., 1978, Canad. J. Zool., v. 56 (5), 1210-1211
Hymenolepis microstoma, effect of streptomycin and penicillin on growth and differentiation in vitro

Penicillin G (K salt)

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Procaine penicillin

Padonou, K. O., 1975, Ang. Parasitol., v. 16 (2), 94-97
Dracunculus medinensis, humans, metronidazole superior to procaine penicillin: Nigeria

Penicillin G (K salt). See Penicillin.

Pentachlorobenzylalcohol

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

3,3',5',6'-Pentachloro-2,2'-dihydroxybenzanilide.
See Oxyclozanide.3,3',5',6'-Pentachloro-2'-hydroxysalicylanilide.
See Oxyclozanide.Pentamidine -- Diamidino-diphenoxypentane;

1,5-Di(4-amidinodiphenoxypentane) di(2-hydroxyethanesulfonate); Lomidine; Pentamidine isethionate.

Pentamidine isethionate

Bachrach, U.; et al., 1979, Exper. Parasitol., v. 48 (3), 464-470
Leishmania spp., effect of ethidium, pentamidine, and methylglyoxal-bis (guanylhydrazine) on growth and on polyamine, RNA, and DNA synthesis

PB Dressing. See Butyl aminobenzoate or Piperonyl butoxide or Propylene glycol.

Pentamidine -- Continued.

Pentamidine (Lomidine)

Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455

Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity

Pentamidine

Ercoli, N., 1978, Proc. Soc. Exper. Biol. and Med., v. 157 (3), 397-401

Trypanosoma venezuelense, rats, comparison of "true" and "false" prophylaxis using pentamidine, suramin and 98/202

Pentamidine isethionate

Farre-Sostres, I.; et al., 1973, Rev. Espan. Pediat. (174), v. 29, 827-840

Pneumocystis carinii, pneumonia in infant presenting as eosinophilia, diagnosis through lung biopsy, successfully treated with pentamidine: Spain

Lomidine

Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113

human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Pentamidine isethionate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Pentamidine isethionate

Hughes, W. T.; et al., 1978, J. Pediat., St. Louis, v. 92 (2), 285-291

Pneumocystis carinii, children, comparison of pentamidine isethionate and trimethoprim combined with sulfamethoxazole (TMP-SMZ) in treating *Pneumocystis pneumonia*, results show that TMP-SMZ is as effective as pentamidine, has minimal side effects, offers oral administration and is more readily available

Lomidine (Diamidino-diphenoxypentane)

Kayembe, D.; and Wery, M., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 1-8

Trypanosoma gambiense, 16 strains isolated in Zaire, lomidine, prophylactic and therapeutic trials in white rats

Pentamidine isethionate

Kluge, R. M.; Spaulding, D. M.; and Spain, A. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 975-978

Pneumocystis carinii in steroid-conditioned rats, combination of pentamidine with trimethoprim-sulfamethoxazole, data suggest that combination therapy is no more effective than trimethoprim-sulfamethoxazole alone and may be, in fact, harmful, trimethoprim by itself has no place in treatment of pneumocystosis

Pentamidine -- Continued.

Pentamidine (Lomidine)

Limbos, P.; et al., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 495-500

[Trypanosoma] rhodesiense, 16-year-old girl, renal insufficiency after treatment with pentamidine: European visitor to Kagera park, Rwanda

Pentamidine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Pentamidine isethionate

Naik, S. R.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 61-65

Leishmania donovani, human, 24 sporadic cases of kala-azar, epidemiology, clinical features, complications and associations, hematological and biochemical findings, diagnosis, treatment: north-western India

Pentamidine isethionate

Nathan, H. C.; et al., 1979, J. Protozool., v. 26 (4), 657-660

Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents

Pentamidine (Lomidine)

Pipano, E.; Jeruhm, I.; and Frank, M., 1979, Trop. Animal Health and Prod., v. 11 (1), 13-16

Babesia bigemina, calves, pentamidine, sterilizing dose was at least 5 times as great as that needed for clinical recovery, promising agent for chemoimmunization

Pentamidine isethionate

Raether, W.; Seidenath, H.; and Loewe, H., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 543-547

Leishmania donovani, golden hamsters, HOE 668 compared with known antileishmanial drugs, toxicity precludes further development but very good anti-leishmanial action qualifies it as standard compound in screening tests

Pentamidine isethionate

Western, K. A.; Norman, L.; and Kaufmann, A. F., 1975, J. Infect. Dis., v. 131 (3), 273-276

Pneumocystis carinii in rats, pentamidine isethionate did not exert significant chemoprophylactic effect against proliferation of parasites although treated rats survived longer than controls; possible application for use as chemoprophylactic agent in humans, study results not encouraging

Pentamidine isethionate. See Pentamidine.

Pentaquine -- Pentaquine diphosphate; WR 6,021; 8-(5-Isopropylaminopentylamino)-6-methoxyquinoline.

Pentaquine diphosphate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Pentaquine (WR 6,021)

Rane, D. S.; and Kinnamon, K. E., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (6), 937-947

sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Pentaquine diphosphate. See Pentaquine.

Pentostam. See Antimony sodium gluconate.

Peracetic acid

Radvan, R.; and Merka, V., 1972, *Sborn. Ved. Praci Lek. Fak. Karlovy Univ. Hradci Kralove, Suppl.*, v. 15 (5), 485-490

Ascaris suum, effect of performic, peracetic, and perpropionic acids on eggs at stages of development of larvae and invasive larvae, poor results

Perdix PE 50. See Butonate.

Performic acid

Radvan, R.; and Merka, V., 1972, *Sborn. Ved. Praci Lek. Fak. Karlovy Univ. Hradci Kralove, Suppl.*, v. 15 (5), 485-490

Ascaris suum, effect of performic, peracetic, and perpropionic acids on eggs at stages of development of larvae and invasive larvae, poor results

Perhydrol

Musaev, F. A., 1972, *Parazitologiya, Leningrad*, v. 6 (2), 185-188

Trichomonas elongata, occurrence in oral cavity of healthy persons vs. those with oral cavity diseases, host age and sex, suitable media for cultivating trichomonads, activity in vitro of several medicinal substances, role of trichomonads in periodontitis confirmed by treatment of patients with trichopol

Permethrin -- BW 212; Ectiban; FMC 33297;

FMC 4T655; NRDC 143; Permethrin EC spray; 3-Phenoxybenzyl (+)-cis, trans-3-(2,2-dichlorovinyl)-2,2-dimethyl cyclopropanecarboxylate; m-Phenoxybenzyl cis, trans-(+)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate; 3-Phenoxybenzyl-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate; (3-Phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate; (3-Phenoxyphenyl)methyl cis, trans-(+)-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate.

Permethrin -- Continued.

Permethrin (NRDC 143)

Hall, C. A., 1978, *Austral. Vet. J.*, v. 54 (10), 471-472

Damalinea ovis, sheep (exper.), cypermethrin proved effective in eradicating lice and at 5 and 10 ppm prevented reinfestation for 7 and 19 weeks respectively; addendum briefly gives results for permethrin in similar tests

Ectiban (Permethrin)

Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, *J. Econom. Entom.*, v. 71 (2), 315-318

Ornithonyssus sylviarum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban (permethrin) and SD-43775 also effective; mites displayed tolerance to malathion

Permethrin EC spray (BW 212)

Harvey, T. L.; and Brethour, J. R., 1979, *J. Econom. Entom.*, v. 72 (4), 532-534

Haematobia irritans, permethrin EC spray applied to one individual bull, cow, or steer per herd eliminated all horn flies from herds within 1 day

Permethrin

Hunt, L. M.; Gilbert, B. N.; and Lemeilleur, C. A., 1979, *Poultry Science*, v. 58 (5), 1197-1201

¹⁴C-labeled permethrin, distribution and depletion of radioactivity in hens treated dermally and in their eggs

Permethrin (NRDC 143)

Nolan, J.; Roulston, W. J.; and Schnitzerling, H. J., 1979, *Austral. Vet. J.*, v. 55 (10), 463-466

Boophilus microplus, range of resistant strains on naturally and experimentally infected cattle, field and stall spraying trials, efficacy of synthetic pyrethroids for tick control, potentiation of pyrethroids by organo-phosphorus compounds

m-Phenoxybenzyl cis, trans-(+)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (FMC 33297; FMC 41655)

Schmidt, C. D.; et al., 1976, *J. Econom. Entom.*, v. 69 (4), 484-486

Haematobia irritans, Stomoxys calcitrans, cattle, m-phenoxybenzyl cis, trans-(+)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate, spot tests, large-cage tests, field tests

Permethrin

Schreck, C. E.; Posey, K.; and Smith, D., 1978, *J. Econom. Entom.*, v. 71 (3), 397-400 permethrin as potential clothing treatment against bloodsucking arthropods, minimum effective dosage necessary for quick kill on contact

Permethrin EC spray. See Permethrin.

- Perpropionic acid
Radvan, R.; and Merka, V., 1972, Sborn. Ved. Praci Lek. Fak. Karlovy Univ. Hradci Kralove, Suppl., v. 15 (5), 485-490
Ascaris suum, effect of performic, peracetic, and perpropionic acids on eggs at stages of development of larvae and invasive larvae, poor results
- Pervinox. See Iodine.
- Pfizona. See Chlorfenvinphos.
- Phanquone -- Entobex; Mexaform (with Iodochlorhydroxyquin and Oxyphenonium bromide).
- Mexaform
Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis
- Entobex
Rostkowska, J., 1970, Acta Parasitol. Polon., v. 18 (27-41), 377-392
Balantidium coli, effect of various bacteria on propagation in vitro, on erythrophagocytic capability of balantidia, and on susceptibility of balantidia to atebirin, entobex, mexaform, and protargol; Trichomonas hominis, Chilomastix mesnili, and Dientamoeba fragilis found to be without effect; effect of balantidia on bacteria
- Mexaform
Rostkowska, J., 1970, Acta Parasitol. Polon., v. 18 (27-41), 377-392
Balantidium coli, effect of various bacteria on propagation in vitro, on erythrophagocytic capability of balantidia, and on susceptibility of balantidia to atebirin, entobex, mexaform, and protargol; Trichomonas hominis, Chilomastix mesnili, and Dientamoeba fragilis found to be without effect; effect of balantidia on bacteria
- Mexaform
Trzaska, B.; and Leinweber-Noiszewska, H., 1974, Ann. Acad. Med. Stetinensis, suppl. 10, 19-25
Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis
-
- Pharmacocicide
Avakian, A. A.; et al., 1978, Veterinaria, Moskva (11), 76-77
coccidiosis, chickens (broilers), prophylactic control by various preparations, pharmacocicide recommended, treatment economics; Krymsk oblast
- Pharmacocid
Krylov, V. F., 1978, Veterinaria, Moskva (10), 68-69
Eimeria tenella strain resistant to pharmacocid after 35 laboratory passages in chickens, cross-resistance only to rigecoccin
- Phenacetine. See Acetophenetidin.
- Phenamidine
McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity
- 9-Phenanthrenemethanols
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghoi and these same compounds
- Phenasal. See Niclosamide.
- 1,9-Phenazine-bis(dialkylaminocarboxamides)
Sawhney, S. N.; and Boykin, D. W., 1979, J. Pharm. Sc., v. 68 (4), 524-526
Plasmodium berghei, 1,9-Phenazine-bis(dialkylaminocarboxamides) synthesized and screened as potential antimalarials, no significant activity
- Phenazine-5-oxide
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Phenetcarb -- 3,5-Diethyl-phenyl-N-methyl-carbamate.
- Phenetcarb
Frolov, B. A.; et al., 1978, Veterinaria, Moskva (11), 75-76
Dermanyssus gallinae, Argas persicus, chickens, acaricides tested
-
- Phenol -- Carbolic acid.
- Phenol
Sherkov, Sh.; et al., 1978, Vet. Sbirka. v. 76 (6), 39-41
coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat
- Carbolic acid
Slepnev, N. K.; and Zen'kov, A. V., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 135-137
E[chinococcus] granulosus, protoscolices, destructive action of high and low temperatures; lysol and creolin most destructive of chemicals tested
-

Phenothiazine -- Helmita-P (with Piperazine, Senna, Tin and Vernonia anthelmintica); Phenothiazine Marki A; Phenothiazine salt; Phenothiazine sulphoxide; Phenovis; Phenzeen; Thiodiphenylamin.

Phenovis (Thiodiphenylamin)

Abdel Rahman, M. S.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 39-48
gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt

Phenothiazine salt + Cupric sulfate

Artem'ev, G. M., 1978, Vestnik Sel'skokhoz. Nauki Kazakhstana (10), 91-93
helminthiases, sheep, economic losses, phenothiazine salt and cupric sulfate mix: Pavlodarsk oblast

Phenothiazine (Phenzeen)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
Haemonchus contortus, Trichostrongylus colubriformis, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Phenothiazine

Dalton, S. E., 1978, Vet. Rec., v. 103 (7), 131-134
nematodes of sheep (nat. and exper.), thiophanate in low daily dosage reduced faecal egg output, egg hatchability, worm burdens, and pasture contamination, increased lamb weight gain, effective against Haemonchus contortus and Nematodirus spathiger, partially effective against Trichostrongylus colubriformis; comparison with phenothiazine treatment

Phenothiazine

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Phenothiazine

Douch, P. G. C.; and Buchanan, L. L., 1979, Xenobiotica, v. 9 (11), 675-679
Moniezia expansa, Ascaris suum, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Phenothiazine sulphoxide

Douch, P. G. C.; and Buchanan, L. L., 1979, Xenobiotica, v. 9 (11), 675-679
Moniezia expansa, Ascaris suum, sulphoxidases and sulphoxide reductases, oxidation and reduction of anthelmintics

Phenothiazine Marki A

Kadyrov, N. T., 1978, Veterinariia, Moskva (7), 57-58
Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Phenothiazine

Kauzal, G. P., 1979, Austral. Vet. J., v. 55 (8), 400 [Letter]
daily dose phenothiazine system vs. conventional drenching, pregnant ewes

Phenothiazine -- Continued.

Phenothiazine

Klenov, A. P., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 152-155
[Bothriocephalus], white amur, effectiveness of various anthelmintics

Phenothiazine

Kuchin, A. S., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 88-92
S[trongyloides] papillosus, sheep, pathology, mentic, promentic, and phenothiazine, all effective

Helmita-P

Matta, S. C.; and Ahluwalia, S. S., 1979, Indian Vet. J., v. 56 (7), 616-617
helminths, poultry, helmita-P, sonex

Phenothiazine-Cupric sulfate mixture

Nekipelova, R. A.; Kurnikov, V. A.; and Iksanov, S. F., 1978, Veterinariia, Moskva (10), 67
Nematodirus and other nematodes, sheep, effects of addition of trace elements to phenothiazine-cupric sulfate mixture: Tselingradsk oblast

Phenothiazine

Oripov, A. O., 1978, Veterinariia, Moskva (4), 74-76
Strongylata, sheep, various anthelmintics in granular form tested for mass dehelminthization, no harmful effects, nilverm most effective

Phenothiazine

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Phenothiazine

Velichkin, P. A.; et al., 1977, Veterinariia, Moskva (7), 17-19
[Heterakis], [Ascaridia], control on poultry farms by phenothiazine, piperazine adipate, nilverm, and other control measures

Phenothiazine (Phenovis)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiabendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxynil are also useful chemical alternatives

Phenothiazine Marki A. See Phenothiazine.

Phenothiazine salt. See Phenothiazine.

Phenothiazine sulphoxide. See Phenothiazine.

Phenothrin -- d-Phenothrin; (3-Phenoxyphenyl)methyl cis, trans-(+)-2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate; S-2539 Forte.

d-Phenothrin (S-2539 Forte)
Schechter, M. S.; et al., 1974, *J. Med. Entom.*, v. 11 (2), 231-233
d-phenothrin, aircraft disinsection trials, not effective against *Xenopsylla cheopis* and *Rhipicephalus sanguineus*

d-Phenothrin. See Phenothrin.

Phenovis. See Phenothiazine.

3-Phenoxybenzyl-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate. See Permethrin.

m-Phenoxybenzyl cis, trans-(*)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate. See Permethrin.

3-Phenoxybenzyl (*)-cis, trans-3-(2,2-dichlorovinyl)-2,2-dimethyl cyclopropanecarboxylate. See Permethrin.

(3-Phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate. See Permethrin.

(3-Phenoxyphenyl)methyl cis, trans-(*)-3-(2,2-dichloroethenyl)2,2-dimethylcyclopropanecarboxylate. See Permethrin.

(3-Phenoxyphenyl) methyl cis, trans-(+)-2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate. See Phenothrin.

Phenthoate -- Cidial; Ethyl mercaptophenylacetate S-ester with O,O-dimethyl phosphorodithioate.

Cidial
Cole, M. M.; et al., 1973, *J. Econom. Entom.*, v. 66 (1), 118-119
Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

Phenthoate
Rawlins, S. C.; and Mansingh, A., 1978, *J. Econom. Entom.*, v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

N⁶-Phenyladenosine
Senft, A. W.; and Crabtree, G. W., 1977, *Biochem. Pharmacol.*, v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

2-Phenyl-4H-1,3,2-benzodioxaphosphorin-2-sulfide (K-15)
Tanaka, H.; et al., 1977, *Kiseichugaku Zasshi* (Japan. *J. Parasitol.*), v. 26 (1), 1-5
Litomosoides carinii, cotton rats, antifilarial activity of 2-phenyl-4H-1,3,2-benzodioxaphosphorin-2-sulfide (K-15), potent inhibitory effect on cholinesterase in mice

Phenyl bis-(hydroxy-2 naphthyl-1) methane derivatives
Lacroix, R.; et al., 1979, *Ann. Pharm. Franc.*, v. 37 (3-4), 131-137
Trichomonas vaginalis, phenyl bis-(hydroxy-2-naphthyl-1) methane derivatives, activity in vitro

p-Phenylene-bis(isothiocyanate). See Bitoscanate.

Phenylene-diisothiocyanate (1,4). See Bitoscanate.

1,4-Phenylene-diisothiocyanate. See Bitoscanate.

Phenylene-isothiocyanate 1,4. See Bitoscanate.

2-Phenylhydrazino-2-thiazoline
Denham, D. A.; et al., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (6), 615-618
Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Phenylmercuric acetate
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

4-Phenyl-1,2,3,6-tetrahydropyridine quaternary salts
Grier, N., 1979, *J. Pharm. Sc.*, v. 68 (4), 407-411
Eimeria tenella, trichostrongyle, synthesis, in vitro and in vivo evaluation of quaternary salts of 4-phenyl-1,2,3,6-tetrahydropyridine and 3,6-dimethyl-6-phenyltetrahydro-2H-1,3-oxazine, none effective

5 (Phenylthio)-2-benzimidazole-carbamic acid methyl ester. See Fenbendazole.

5-Phenylthio-benzimidazole-2-methylcarbaminate. See Fenbendazole.

[5-(Phenylthio)-1H-benzimidazole-2-yl] carbamic acid methyl ester. See Fenbendazole.

2-Phenyl-thionaphthene diamidine. See 6-Amidino-2-(4'-amidinophenyl)-thionaphthene dilactate.

Phenzeen. See Phenothiazine.

- Philadelphus spp. extracts
Fazakas, B.; and Racz, G., 1976, Rev. Med.,
Tirgu-Mures, v. 22 (1), 25-28
Trichomonas vaginalis in culture medium,
trichomonacidal action of extracts from flow-
ers of various species of the genus Philadel-
phus, P. coronarius most active as it de-
stroyed all parasites in 3-5 minutes
- Phosalone -- Benzophosphate; O,O-Diethyl-(6-
chlorobenzoxazoliny-3-methyl) dithiophos-
phate; RP 11074 [i.e. RP 11974]; Zolone.
- Benzophosphate (Zolone; Phosalone; RP 11074)
Frolov, B. A.; et al., 1978, Veterinariia,
Moskva (11), 75-76
Dermanyssus gallinae, Argas persicus,
chickens, acaricides tested
-
- Phosmet -- 0,0-Dimethyl phosphorodithioate S-
ester with N-(mercaptomethyl) phthalimide;
0,0-Dimethyl-S(phthalimidomethyl)dithiophos-
phate; 0,0-Dimethyl phthalimidomethyl phos-
phorothiolothionate; GX-118; Imidan; Paramite;
Poron 20; Prolate; Starbar GX-118.
- Imidan
Drummond, R. O.; et al., 1973, J. Econom. En-
tom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory
tests of insecticides
- Phosmet (Starbar GX-118)
Fisher, W. F.; et al., 1979, Southwest. En-
tom., v. 4 (3), 249-253
Psoroptes cuniculi, rabbits, phosmet, toxa-
phene, and 10 experimental alkyl amines com-
pared
- Phosmet
Frazar, E. D.; and Schmidt, C. D., 1979, J.
Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, sus-
ceptibility to topically applied insecticides
- Phosmet (Imidan)
Gladney, W. J.; Price, M. A.; and Graham, O.
H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of
various insecticides applied as sprays, ear
smears and dusts, or in slow-release devices,
field tests
- Phosmet (Poron 20)
Kettle, P. R.; and Lukies, J. M., 1979, N.
Zealand Vet. J., v. 27 (4), 78-79
Linognathus vituli, cattle, pour-on formula-
tions of phosmet, methidathion, chlorpyrifos,
and temephos: Kaitoke, near Upper Hutt, New
Zealand
- Paramite (Phosmet)
Knapp, F. W., 1978, Vet. Med. and Small Animal
Clin., v. 73 (9), 1161-1163
Sarcoptes scabiei var. canis, dogs, paramite,
good results
- Phosmet (Prolate, Starbar GX-118)
Meleney, W. P.; and Roberts, I. H., 1979, J.
Med. Entom., v. 16 (1), 52-58
Psoroptes ovis, cattle, acaricides, dipping,
spraying, or spray-dipping trials
- 0,0-Dimethyl-S(phthalimidomethyl)dithiophosphate
(Imidan)
Privora, M.; Rupes, V.; and Cerny, V., 1970,
Folia Parasitol., v. 17 (1), 81-84
Dermacentor marginatus, laboratory trials
testing six insecticides
- Phosmet (Prolate; Imidan)
Roberts, I. H.; Wilson, G. I.; and Meleney, W.
P., 1978, J. Am. Vet. Med. Ass., v. 173 (7),
840-842
Psoroptes ovis, cattle (exper.), phosmet,
efficacy of single and double treatments at
various concentrations, some signs of toxico-
sis
- Phosmet
Wright, F. C.; and Riner, J. C., 1979, South-
west. Entom., v. 4 (1), 40-45
Psoroptes ovis, P. cuniculi, 10 acaricides
evaluated using 'tea-bag' technique
-
- Phosphamidon -- 0[2-Chloro-2-(diethyl carbamoyl)-
1-methyl-vinyl]-0,0-dimethyl phosphate;
Dimecron.
- Phosphamidon
Drummond, R. O.; et al., 1973, J. Econom. En-
tom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory
tests of insecticides
- Phosphamidon
Khan, D.; and Haseeb, M. A., 1976, Pakistan J.
Zool., v. 8 (2), 173-176
Ganeo micracetabulus and Cercaria reflexi-
cauda cercariae, effects of 5 insecticides
at various concentrations, toxicity varies
but results indicate cercariae are suscepti-
ble to insecticides
- Dimecron
Khan, M. H., 1979, Indian Vet. J., v. 56 (9),
739-743
Lipeurus caponis, Menacanthus stramineus,
White Leghorn fowl, organophosphorus insecti-
cides, costs evaluated
- Dimecron (Phosphamidon)
Khan, M. H.; and Srivastava, S. C., 1977,
Indian J. Animal Health, v. 16 (2), 137-140
Boophilus microplus engorged females, in
vitro tests with dursban, gamma BHC, sumith-
ion, supona, dimecron, egg production and
viability; supona most effective
- Phosphamidon
Rawlins, S. C.; and Mansingh, A., 1978, J.
Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains,
patterns of resistance to acaricides
-
- Phoxim (Bayer 9053)
Meermann, A., 1978, Vet.-Med. Nachr. (2),
212-215
Psoroptes ovis, sheep, phoxim, gamma BHC,
laboratory and field trials: Nordfriesland

Phthalazinylhydrazones

Molodykh, Zh. V.; et al., 1977, Khimiko-Farm. Zhurnal, v. 11 (7), 37-40

Nippostrongylus braziliensis, *Hymenolepis nana*, mice, anthelmintic activity of hydrazones, phthalazones, and phthalazinylhydrazones, relationship to chemical structure

Phthalazones

Molodykh, Zh. V.; et al., 1977, Khimiko-Farm. Zhurnal, v. 11 (7), 37-40

Nippostrongylus braziliensis, *Hymenolepis nana*, mice, anthelmintic activity of hydrazones, phthalazones, and phthalazinylhydrazones, relationship to chemical structure

Phthalophos -- 0,0-Diethyl-0-naphthaloximide phosphate; N-Hydroxy naphthalimide diethyl phosphate; Maretin; Naphthalophos; Rametin; Rametin H.Rametin (Naphthalophos)

Abdel Rahman, M. S.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 39-48

gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt

Naphthalophos (Rametin H)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25

Haemonchus contortus, *Trichostrongylus colubriformis*, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Phthalophos

Kanaev, A. I.; et al., 1977, Veterinariia, Moskva (10), 103-104

diagnosis of phthalophos poisoning of fish

Naphthalophos (Rametin H)

Le Jambre, L. F.; and Barger, I. A., 1979, Austral. Vet. J., v. 55 (7), 346-347 [Letter]

benzimidazole resistant *Haemonchus contortus*, lambs, rafoxanide and naphthalophos, efficiency against adults and inhibited larvae: New South Wales

Naphthalophos

Le Jambre, L. F.; and Martin, P. J., 1979, Vet. Sc. Commun., v. 3 (2), 153-158

levamisole resistant *Ostertagia circumcincta* and *O. trifurcata*, sheep, cross resistant to morantel tartrate but not to naphthalophos

Maretin

Oripov, A. O., 1978, Veterinariia, Moskva (4), 74-76

Strongylata, sheep, various anthelmintics in granular form tested for mass dehelminthization, no harmful effects, nilverm most effective

Naphthalophos (Rametin)

Varshney, T. R.; and Singh, Y. P., 1979, Indian Vet. J., v. 56 (3), 207-210

gastrointestinal helminths, sheep, naphthalophos more effective than parabendazole in controlled study under farm conditions

Phthalophos -- Continued.Naphthalophos (Rametin H)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]

Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiabendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxylin are also useful chemical alternatives

Phthalylsulfathiazole -- Ftalozol.Ftalozol

Gobzem, V. R.; and Nazarov, V. G., 1978, Veterinariia, Moskva (3), 67-69

Eimeria spp., calves, diagnostic difficulties, clinical symptoms, chemoprophylactic substances tested at various dosages and in various combinations

Physostigmine -- Eserine.Eserine

Hillman, G. R.; Gibling, A. M.; and Anderson, J. W., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 992-997

Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

Pimafucin. See Pimaricin.Pimaricin -- Natamycin; Pimafucin.Natamycin (Pimafucin)

Lovgren, T.; and Salmela, I., 1978, Acta Path. et Microbiol. Scand., v. 86B (3), 155-158

Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents

Natamycin

Ograjensek, Z., 1974, Quad. Clin. Ostet. e Ginec., v. 29 (5-6), 127-133

Trichomonas vaginalis, human vaginitis, successful clinical trials with methyl-partricin vaginal suppositories using natamycin as reference drug

Pine needles

Slepnev, N. K., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 107-109

Ascaris suum, in vitro testing of anthelmintic activity of water extracts of some plants

Piperamide -- CL 71.366; 1-(3-Dimethylaminopropyl)-4-(p-methoxy-phenyl) piperazine dihydrochloride.

CL 71.366
Andrade, S. G.; Figueira, R. M.; and Carvalho, M. L., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (2), 135-145
Trypanosoma cruzi, mice, activity of CL 71.366

Piperamide
Brener, Z., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (4), 302-306
Trypanosoma cruzi, description of method allowing study of drug action on trypomastigotes in mice

Piperazine -- Ancaris (with Thienium closylate); BB-04 (with Thiabendazole); Bithirazine (with Bithionol); Citrazine; Equizole A (with Thiabendazole); Gamaverm (with Thiabendazole); Helmita-P (with Phenothiazine, Senna, Tin, and Vernonia anthelmintica); Mascyl (with Carbon disulfide); Piperazine adipate; Piperazine adipinate; Piperazine aminohexanate; Piperazine chlorhydrate; Piperazine citrate; Piperazine hexahydrate; Piperazine hydrate; Piperazine monophosphate; Piperazine phosphate; Piperazine S; Piperazine salt; Piperazine sulfate; Suiverm (with Thiabendazole); Ta-Verm; Vermex.

Piperazine phosphate
Abdel Hamid, Y. M., 1971, Vet. Med. J., Giza, v. 19 (19), 391-392
Neoascaris vitulorum adults, 3-week old buffalo calf (faeces, rectum), maturity of worms indicated intrauterine transmission, treatment with piperazine phosphate, case report: Assiut University

Piperazine monophosphate
Abdulla, W. A.; et al., 1977, Egypt. J. Ri-lharz., v. 4 (2), 109-115
S[chistosoma] mansoni, prophylactic activity, antischistosomal drugs, albino mice, most effective within 2 weeks post exposure

Piperazine
Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Piperazine
Camillo-Coura, L., 1974, Rev. Soc. Brasil. Med. Trop., v. 8 (6), 340-356
mass therapy trials using piperazine for ascariasis and phenylenediisothiocyanate for ancylostomiasis; emphasis also on need for social and economic improvements and proper sanitation to achieve some measure of control: States of Rio de Janeiro and Guanabara, Brazil

Piperazine -- Continued.

Piperazine hexahydrate + Thiabendazole (= BB-04)
Camillo-Coura, L.; et al., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (2), 103-113
human Ascaris lumbricoides, Trichuris trichiura, Ancylostomidae, treatment trials with piperazine combined with thiabendazole in silicone base (BB-04), mixed results

Piperazine
Cardenas Santiuste, C.; Rizo, F.; and Brooks, R. W., 1972, Rev. Cubana Cirug., v. 11 (1), 19-24
Ascaris lumbricoides, infection in child resulting in intestinal occlusion and perforation of Meckel's diverticulum, post-surgical therapy with piperazine unsuccessful as Ascaris continued to appear in feces up to one year later: Cuba

Piperazine
Chaia, G.; and da Cunha, A. S., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (2), 152-160
intestinal helminths, children, efficacy of anthelmintic therapy: Vila Sao Vicente, Belo Horizonte

Piperazine citrate
Cho, S. Y.; Bae, J. H.; and Park, S. C., 1973, Taehan Uihak Hyophoe Chi (J. Korean Med. Ass.), v. 16 (8), 649-656
Ascaris lumbricoides, incidence in ROK army, mass therapy with piperazine: Korea

Piperazine phosphate
Corwin, R. M.; and Miller, T. A., 1978, Am. J. Vet. Research, v. 39 (2), 263-265
Toxocara canis, young dogs and weaned pups, thienium closylate, piperazine phosphate, efficacy when administered alone or in combination, critical controlled trials

Piperazine
Daniliarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Piperazine citrate
Dudley, B. W.; and Lease, E. J., 1973, South. Med. J., v. 66 (10), 1100, 1126
ascariasis, schoolchildren, disease control using repetitive treatment (piperazine or pyrantel pamoate) on a mass therapy basis: South Carolina

Piperazine
Faraco, B. A.; and Faraco, B. F. C., 1973, Rev. Brasil. Med., v. 30 (2), 79-81
Ascaris lumbricoides, pregnant women, piperazine well tolerated, no evidence of transplacental transmission of infection to fetus

Piperazine + Thiabendazole
Fernandes, J. L.; and Garcia, E., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (3), 155-158
human intestinal parasites, clinical trials with combined piperazine and thiabendazole: Sao Paulo

Piperazine -- Continued.

- Piperazine hexahydrate + Thiabendazole (= Gama-
verm)
Ferriolli Filho, F.; et al., 1971, Rev. Soc.
Brasil. Med. Trop., v. 5 (4), 209-212
Ascaris lumbricoides, Strongyloides stercora-
lis, Ancylostomidae, mixed human infections,
clinical trials testing efficacy of piperaza-
zine and thiabendazole in combination therapy
- Piperazine adipate
Furtunescu, G.; et al., 1973, Rev. Med.-Chir.
Soc. Med. si Nat. Iasi, v. 77 (1), 179-183
in vitro and in vivo trials (dogs) testing
efficacy of new piperazine aminohexanate,
comparison with piperazine adipate and
piperazine hydrate, results good with Ascaris
suum and Toxocara canis but unresponsive with
Trichuris vulpis and Ancylostoma caninum
- Piperazine aminohexanate (Piperazine S)
Furtunescu, G.; et al., 1973, Rev. Med.-Chir.
Soc. Med. si Nat. Iasi, v. 77 (1), 179-183
in vitro and in vivo trials (dogs) testing
efficacy of new piperazine aminohexanate,
comparison with piperazine adipate and
piperazine hydrate, results good with Ascaris
suum and Toxocara canis but unresponsive with
Trichuris vulpis and Ancylostoma caninum
- Piperazine hydrate
Furtunescu, G.; et al., 1973, Rev. Med.-Chir.
Soc. Med. si Nat. Iasi, v. 77 (1), 179-183
in vitro and in vivo trials (dogs) testing
efficacy of new piperazine aminohexanate,
comparison with piperazine adipate and
piperazine hydrate, results good with Ascaris
suum and Toxocara canis but unresponsive with
Trichuris vulpis and Ancylostoma caninum
- Bithirazine
Golovneva, L. F.; et al., 1970, Nauch. Trudy,
Nauchno-Issled. Vet. Inst., v. 8, 105-106
ascariasis, chickens, bithirazine tested
under production conditions, more study
needed
- Piperazine citrate
Gonzalez Gutierrez, J.; Pavon Lebrero, R.; and
Castro Cauto, A., 1978, Rev. Inform. Med.
Terap., v. 53 (1-3), 54-56
Ascaris lumbricoides, children, clinical
management, piperazine therapy
- Piperazine chlorhydrate
Goulart, E. G.; de Arruda, M. E.; and Jourdan,
M. C., 1974, Rev. Brasil. Med., v. 31 (11),
791-794
human soil transmitted nematodes, laboratory
trials testing ovicidal and larvacidal ef-
fects of selected anthelmintic drugs; prophyl-
actic treatment of organic fertilizer or
contaminated soil by these drugs seemed to be
ineffective
- Piperazine hexahydrate
Graf, W.; Haldimann, B.; and Flury, W., 1978,
Schweiz. Med. Wchnschr., v. 108 (5), 177-181
oxyuriasis, man with terminal renal failure
receiving long-term hemodialysis, severe
toxic reaction from piperazine therapy, case
report; piperazine contraindicated in pa-
tients with renal failure

Piperazine -- Continued.

- Piperazine + Thiabendazole (= Equizole A)
Grieve, R. B.; Moore, B. G.; and Bradley,
R. E., 1979, Am. J. Vet. Research, v. 40 (1),
139-141
gastrointestinal parasites, horses and
ponies, critical test evaluation of butami-
sole, compared with efficacy of piperazine-
thiabendazole
- Piperazine citrate
Hatchuel, W.; Isaacson, M.; and de Villiers,
D. J., 1973, South African Med. J., v. 47 (3),
91-93
Ascaris lumbricoides, children, pyrantel
pamoate vs. piperazine citrate, clinical
trials: Alexandra Township near Johannes-
burg
- Piperazine adipinate
Iakubovskii, M. V., 1979, Veterinariia, Moskva
(2), 41-42
ascariasis, trichocephaliosis, oesophagosto-
miasis, swine, comparative effectiveness and
economic value of various drugs: Minsk
oblast
- Piperazine salt
Iakubovskii, M. V.; and Zen'kov, A. V., 1977,
Vet. Nauka--Proizvod., Trudy, Minsk, v. 15,
72-78
nematodes, swine raising complexes, combined
control measures, sanitation, anthelmintics,
suiverm most effective
- Suiverm
Iakubovskii, M. V.; and Zen'kov, A. V., 1977,
Vet. Nauka--Proizvod., Trudy, Minsk, v. 15,
72-78
nematodes, swine raising complexes, combined
control measures, sanitation, anthelmintics,
suiverm most effective
- Piperazine adipinate
Kadyrov, N. T., 1978, Veterinariia, Moskva
(7), 57-58
Delafondia vulgaris, horses, anthelmintics
tested; preventive dehelminthization every
15 days recommended for horses on pasture
- Piperazine hexahydrate
Levi, G. C.; et al., 1972, Rev. Inst. Med.
Trop. S. Paulo, v. 14 (6), 392-396
ascariasis, human, clinical trials comparing
piperazine, tetramisole (or levamisole) and
pyrantel pamoate: Brazil
- Helmint-P
Matta, S. C.; and Ahluwalia, S. S., 1979,
Indian Vet. J., v. 56 (7), 616-617
helminths, poultry, helmint-P, sonex
- Piperazine citrate
Miller, M. J.; et al., 1978, South. Med. J.,
v. 71 (2), 137-140
ascariasis, children throughout the world,
levamisole vs. known anthelmintics, levami-
sole recommended for mass chemotherapy: Iran,
Brazil, Mississippi, Louisiana

Piperazine -- Continued.

Piperazine

Nistri, A.; and Arenson, M. S., 1978, *Experientia*, v. 34 (8), 1046-1047
piperazine, effect on central and peripheral cholinergic synapses of the frog

Piperazine + Carbon disulfide (= Mascyl)

Olsson, T., 1977, *Svensk Vet.-Tidn.*, v. 29 (20), 795-800
Ascaris suum, slaughter swine, treatment with nequvon or mascyl

Piperazine citrate (Citrazine)

Owen, D.; and Turton, J. A., 1979, *Lab. Animals*, v. 13 (2), 115-118
Syphacia obvelata, laboratory mice and rats, thiabendazole and piperazine citrate in feed and water

Piperazine

Parodi Hueck, L.E.; et al., 1975, *Rev. Fac. Med. Univ. Zulia*, v. 7 (1-2), 1974-1975, 193-197
ascariasis, children, case reviews, recommendations for piperazine and conservative medical treatment except in cases of surgical emergencies as volvulus or perforation

Piperazine hexahydrate (Vermex)

Parshad, V. R.; and Guraya, S. S., 1978, *Vet. Parasitol.*, v. 4 (2), 111-120
4 helminth spp., comparison of phosphatases, effects of pH, various chemicals, and some anthelmintics on enzyme activity, anthelmintics may affect absorptive process in worms by virtue of their effect on phosphatase system at absorptive surfaces

Piperazine adipinate

Petrov, Iu. F., 1978, *Veterinariia*, Moskva (5), 64-66
[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Piperazine hexahydrate

Petrov, Iu. F., 1978, *Veterinariia*, Moskva (5), 64-66
[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Piperazine

Phillips, L.; Sturman, G.; and West, G. B., 1975, *Agents and Actions*, v. 5 (5), 467
Ascaris suum, variations in histamine content, effect of piperazine

Piperazine phosphate + Thienium closylate (= Ancaris)

Remfry, J., 1978, *Lab. Animals*, v. 12 (4), 213-218
helminth infections in imported Macaca mulatta, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Piperazine

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, *Rev. Iber. Parasitol.*, v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Piperazine -- Continued.

Piperazine citrate (Ta-Verm)

Sargent, R. G.; et al., 1972, *South. Med. J.*, v. 65 (3), 294-298
intestinal helminths, children, prevalence survey, clinical trials to determine effectiveness of 2 different repetitive treatment programs using piperazine citrate syrup to treat Ascaris: Horry County, South Carolina

Piperazine

Sasi, P. K.; and Raj, R. K., 1978, *Experientia*, v. 34 (9), 1156-1157
Ascaris lumbricoides has necessary mechanism for biosynthesis and degradation of phospholipids and triacylglycerols, piperazine decreases level of triacylglycerols of this parasite by stimulating activity of lipase and partially inhibiting activity of phosphatidate phosphatase

Piperazine hexahydrate

Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1978, *Egypt. J. Bilharz.*, v. 4 (1), 1977, 9-18
schistosomicides, comparative study on effects of bilharzid, tartar emetic and piperazine hexahydrate on activity of oxidoreductase enzymes of rat liver preparations

Piperazine adipate

Sharma, N. D.; and Sisodia, C. S., 1976, *Indian Vet. J.*, v. 53 (12), 920-922
Ascaridia galli, Butea frondosa seeds compared with piperazine adipate, in vivo (chicks) and in vitro trials

Piperazine adipinate

Shedivtsova, A., 1976, *Med. Parazitol. i Parazit. Bolezni*, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Piperazine

Simitzis-Le Flohic, A. M.; et al., 1977, *Semaine Hop. Paris*, v. 53 (22-23), 1369-1370
schistosomiasis with minor parasitism of distomiasis, ascariasis, trichocephaliasis, woman with presenting symptoms of adrenal insufficiency, cortisone therapy resulted in aggravated symptoms and asthenia, parasitism diagnosed, piperazine therapy resulted in toxic neurologic reactions, illness resolved after niridazole therapy: France (had resided in Central African Republic)

Piperazine adipate

Slepnev, N. K., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 108-112
ascariasis, trichuriasis, oesophagostomiasis, swine, comparative effectiveness of various anthelmintics

Piperazine phosphate

Slepnev, N. K., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 108-112
ascariasis, trichuriasis, oesophagostomiasis, swine, comparative effectiveness of various anthelmintics

Piperazine -- Continued.

Piperazine sulfate

Slepnev, N. K., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 108-112
ascariasis, trichuriasis, oesophagostomiasis, swine, comparative effectiveness of various anthelmintics

Piperazine hexahydrate

Swartzwelder, J. C.; et al., 1972, Rev. Biol. Trop., v. 20 (2), 295-307
tetrachloroethylene administered prophylactically to population of coffee plantation to reduce incidence of hookworm, piperazine hexahydrate administered simultaneously reduced incidence of *Ascaris lumbricoides* during initial phase of treatment only: Costa Rica

Piperazine + Thiabendazole (= Gamaverm)

Velho Soli, A. S.; et al., 1972, Rev. Soc. Bras. Med. Trop., v. 6 (4), 207-210
human helminths, clinical trials using combined piperazine and thiabendazole: Brazil

Piperazine adipate

Velichkin, P. A.; et al., 1977, Veterinariia, Moskva (7), 17-19
[Heterakis], [Ascaridia], control on poultry farms by phenothiazine, piperazine adipate, nilverm, and other control measures

Piperazine adipate. See Piperazine.

Piperazine adipinate. See Piperazine.

Piperazine aminohexanate. See Piperazine.

Piperazine chlorhydrate. See Piperazine.

Piperazine citrate. See Piperazine.

Piperazine diantimonyl tartrate -- Bilharzid;
Bilharzid.

Bilharzid

Abdulla, W. A.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 109-115
S[*chistosoma*] mansoni, prophylactic activity, antischistosomal drugs, albino mice, most effective within 2 weeks post exposure

Bilharzid (Piperazine diantimonyl tartrate)

Abdulla, W. A.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 129-140
bilharzid much less toxic than tartar emetic when given in same dose level to guinea pigs, dogs, and monkeys

Piperazine-di-antimonyl tartrate (Bilharzid)

El-Hawary, M. F. S.; et al., 1976, Egypt. J. Physiol. Sc., v. 2 (1-2), 1975, 69-78
bilharziasis, serum and urinary amino acid patterns in Egyptian children at different stages of the disease and before and after bilharzid therapy

Piperazine diantimonyl tartrate -- Continued.

Piperazine diantimonyl tartrate (Bilharzid)

Mahran, S. G.; et al., 1976, Egypt. J. Bilharz., v. 3 (2), 239-245
Schistosoma mansoni, in vivo and in vitro trials comparing efficacy of bilharzid with that of tartar emetic

Bilharzid (Piperazine diantimonyl tartrate)

Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1976, Egypt. J. Bilharz., v. 3 (2), 247-254
inhibition of citrate oxidation in rat liver by the antischistosomal drug bilharzid, inhibition decreased by time, addition of citrate protected the enzyme against inhibitory action of the drug if administered before the bilharzid

Bilharzid (Piperazine diantimonyl tartrate)

Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1978, Egypt. J. Bilharz., v. 4 (1), 1977, 9-18
schistosomicides, comparative study on effects of bilharzid, tartar emetic and piperazine hexahydrate on activity of oxidoreductase enzymes of rat liver preparations

Bilharzid (Piperazine diantimonyl tartrate)

Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1978, Egypt. J. Bilharz., v. 4 (1), 1977, 27-34
antischistosomal drug bilharzid, inhibition of malate and pyruvate oxidation in rat liver

Bilharzid (Piperazine diantimonyl tartrate)

Sharaf, A. A.; El-Sherbini, A.; and Abdulla, W. A., 1978, Egypt. J. Bilharz., v. 4 (1), 1977, 35-45
antischistosomal drug bilharzid, inhibitory action on succinate oxidation in normal rat liver

Bilharzid (Piperazine diantimonyl tartrate)

Shehata, H.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 117-128
S[*chistosoma*] mansoni, mice with anemia and leucocytosis, bilharzid safer therapy than tartar emetic

Piperazine hexahydrate. See Piperazine.

Piperazine hydrate. See Piperazine.

Piperazine monophosphate. See Piperazine.

Piperazine phosphate. See Piperazine.

Piperazine S. See Piperazine.

Piperazine salt. See Piperazine.

Piperazine sulfate. See Piperazine.

α -(2-Piperidyl)-3,6-bis (trifluoromethyl)-9-phenanthrene methanol -- 3,6-Bis-(trifluoromethyl)- α -2-piperidyl-9-phenanthrenemethanol hydrochloride; WR 122,455

3,6-Bis-(trifluoromethyl)- α -2-piperidyl-9-phenanthrenemethanol hydrochloride (WR 122,455)
Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

WR-122,455

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-122,455

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

α -(2-Piperidyl)-2,6-di-(p-trifluoromethylphenyl)-4-pyridinemethanol -- WR 154,904.

WR 154,904

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

α -(2-Piperidyl)-2-trifluoromethyl-6,8-dichloro-4-quinolinemethanol -- WR-226,253.

WR-226,253

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta

α -(2-Piperidyl)-2-trifluoromethyl-6,8-dichloro-4-quinolinemethanol (WR-226,253)

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (5), 680-689
Plasmodium falciparum, P. vivax, Aotus trivirgatus griseimembra, antimalarial activities of WR-184,806 and WR-226,253

α -(2-Piperidyl)-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridinemethanol -- DL-Threo- α -(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate; WR-180,409; WR 180,409.H₃PO₄; DL-Threo- α -2'-piperidyl-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridinemethanol.

DL-Threo- α -2'-piperidyl-2-(4-trifluoro-methylphenyl)-6-trifluoromethyl-4-pyridinemethanol
Nakagawa, T.; et al., 1979, J. Pharm. Sc., v. 68 (6), 718-721
antimalarials, whole blood concentrations, gas liquid chromatography determinations, in vivo time course plots

WR-180,409

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

DL-Threo- α -(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate (WR 180,409.H₃PO₄)
Stampfli, H.; et al., 1979, J. Liquid Chromatography, v. 2 (1), 53-65
multi-component solvent system for analysis of candidate antimalarial (WR 180,409.H₃PO₄) and its internal standard (WR 184,806.H₃PO₄) by normal phase high pressure liquid chromatography

Piper marginatum

Frischkorn, C. G. B.; Frischkorn, H. E.; and Carrazoni, E., 1978, Naturwissenschaften, v. 65 (9), 480-483
Schistosoma mansoni, cercaricidal activity of various plants from northeast Brazil

Piperonyl butoxide -- PB Dressing (with Butyl aminobenzoate and Propylene glycol); Pyractone M817 (with Pyrethrins).

Piperonyl butoxide + Pyrethrum

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econ. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Pyractone M817

Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides

PB Dressing

Nie, I. A.; and Pick, C. R., 1978, J. Inst. Animal Techn., v. 29 (1), 63-68
Otodectes cynotis, ferrets (ears), PB dressing: laboratory colony, Mill Hill, London

Piperonyl butoxide -- Continued.

Piperonyl butoxide

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosis, scolicidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Piperonyl butoxide

Schuntner, C. A.; and Thompson, P. G., 1978, Austral. J. Biol. Sc., v. 31 (2), 141-148

Boophilus microplus, larvae, ¹⁴C labelled amitraz, metabolism, only amitraz and N-2,4-dimethylphenyl-N¹-methylformamidine (metabolite) toxic to larvae, piperonyl butoxide applied simultaneously with amitraz had slight effect on metabolism, three-fold synergistic effect; SKF 525-A similarly applied had negligible effect on both metabolism and toxicity to ticks

Pirimiphos ethyl

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960

Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Pirimiphos-methyl -- 0-[2-(Diethylamino)-6-

methyl-4-pyrimidinyl]-0,0-dimethyl phosphorothioate.

Pirimiphos-methyl

Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661

flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico

Pirimiphos methyl

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960

Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Pirodia. See Amicarbalide.

Pletil. See Tinidazole.

Plictran. See Cyhexatin.

Plifenate (Baygon MEB)

Blommers, L.; and van Lennep, M., 1978, Acta Leidensia, v. 46, 9-15

Pediculus capitis, susceptibility of 1st instar nymphs to malathion, propoxur, and plifenate, residual action of 3 compounds applied as lotions to human hair

Polmiror. See Nifuratel.

Polyethylene sulfonate

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Polyflavine

Nagington, J., 1975, Tr. Ophth. Soc. United Kingdom, v. 95 (2), 207-209

Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity

Polygalacturonic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Poly-D-glutamic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Poly-L-glutamic acid

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Poly-D-lysine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Poly-L-lysine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Polymyxin B -- Neosporin (with Neomycin).**Polymyxin B**

Kasprzak, W.; Mazur, T.; and Rucka, A., 1974, *Ann. Soc. Belge Med. Trop.*, v. 54 (4-5), 351-357

4 strains of free-living amoebae isolated from lakes in Poland, pathogenicity for mice, response to several drugs, identified as *Acanthamoeba* spp. on basis of morphology and protein disc electrophoretic patterns

Neosporin

Nagington, J., 1975, *Tr. Ophth. Soc. United Kingdom*, v. 95 (2), 207-209

Acanthamoeba spp. isolated from human eye infections, clinical report, in vitro trials of compounds for possible amoebicidal and cysticidal activity

Polyoxin

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130

Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Polystat. See Dibutyltin dilaurate or Dinsed or Roxarsone or Sulfantran.**Polytherin A.** See Nigericin.**Polyuridylic acid**

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Polyvinylsulfate

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Pomegranate -- *Punica granatum*; Sonex (with *Embelia ribes* and Nicotine).***Punica granatum* (rind of fruit)**

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118
Ascaridia galli, *Ascaris vitulorum*, in vitro anthelmintic activity of some Egyptian plants; only *Nerium oleander* caused death of worms

Sonex

Matta, S. C.; and Ahluwalia, S. S., 1979, *Indian Vet. J.*, v. 56 (7), 616-617
helminths, poultry, helmintha-P, sonex

Poquil. See Pyrvinium.**Poron 20.** See Phosmet.**Portulaco oleracea seeds**

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118
Ascaridia galli, *Ascaris vitulorum*, in vitro anthelmintic activity of some Egyptian plants; only *Nerium oleander* caused death of worms

Potassium antimonyl tartrate. See Antimony potassium tartrate.**Potassium antimony tartrate.** See Antimony potassium tartrate.**Potassium arsenite solution** -- Fowler's solution; *Liquor arsenicalis*.**Liquor arsenicalis (Fowler's solution)**

Reddy, R. G., 1979, *Livestock Advis.*, v. 4 (3), 37-40

Balantidium coli and mixed infection with amphistomes, cattle and buffaloes, incidence and treatment

Potassium cyanide

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Potassium dichromate

Kunstyr, I.; and Ammerpohl, E., 1978, *Lab. Animals*, v. 12 (2), 95-97

Spiroplasma muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Potassium permanganate

Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, *Current Sc.*, Bangalore, v. 47 (8), 283-284 [Letter]

Trichodina [sp.] on exterior of *Cyprinus carpio*, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

- potassium permanganate
Musaev, F. A., 1972, Parazitologiya, Lenin-grad, v. 6 (2), 185-188
Trichomonas elongata, occurrence in oral cavity of healthy persons vs. those with oral cavity diseases, host age and sex, suitable media for cultivating trichomonads, activity in vitro of several medicinal substances, role of trichomonads in periodontitis confirmed by treatment of patients with trichopol
- ovan. See Pyrvinium.
- ovidone-Iodine. See Iodine.
- praziquantel -- Bitricide; 2-Cyclohexylcarbonyl-1,3,4,6,7,11b-hexahydro-2H-pyrazino[2,1a]isoquinolin-4-one; Droncit; Embay 8440; Embay-8440-Bayer.
- praziquantel
Albert, H.; and Hoerchner, F., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (10), 189-193
Taenia saginata, calves (exper.), serum antibodies, enzyme-linked immunosorbent assay, titres following reinfection and drug therapy
- Praziquantel (Droncit)
Andersen, F. L.; Conder, G. A.; and Marsland, W. P., 1978, Am. J. Vet. Research, v. 39 (11), 1861-1862
Echinococcus granulosus, dogs (exper.), praziquantel, efficacy of injectable and tablet formulations against mature tapeworms, no signs of toxicosis
- Droncit (Praziquantel)
Andrews, P., 1976, Vet. Med. Rev. (2), 154-165
droncit, animals, plasma concentrations and distribution in host body determined using biological assay with helminth
- Praziquantel
Andrews, P., 1978, Ztschr. Parasitenk., v. 56 (1), 99-106
Schistosoma mansoni, praziquantel, effects on miracidia and their hatching; effects on cercarial swimming, penetration and subsequent development in final host
- Praziquantel
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results
- Praziquantel (Droncit)
Bankov, D. E., 1977, Vet.-Med. Nachr. (2), 145-148
Echinococcus granulosus, dogs (exper.), bunamidine hydrochloride, cantrodiphen, praziquantel, praziquantel most effective and showed no toxicity
- Praziquantel -- Continued.
- Praziquantel (Embay 8440; Droncit)
Bartsch, H.; et al., 1978, Mutation Research, v. 58 (2-3), 133-142
praziquantel, absence of mutagenic activity in bacteria, yeasts, insects, and mammalian cells, short-term assays, anti-schistosomal effectiveness of this drug is not related to mutagenic activity
- Praziquantel (Droncit)
Boray, J. C.; et al., 1979, Austral. Vet. J., v. 55 (2), 45-53
nematodes and cestodes of dogs and cats, efficiency and safety of nitroscanate, comparison with mebendazole, bunamidine hydrochloride, and praziquantel
- Praziquantel
Buehring, K. U.; et al., 1978, European J. Drug Metab. and Pharmacokinet., v. 3 (3), 179-190
praziquantel-¹⁴C, metabolism in man and Rhesus monkey, in vitro studies with rat liver homogenates
- Praziquantel (Embay 8440)
Chavasse, C. J.; Brown, M. C.; and Bell, D. R., 1979, Ztschr. Parasitenk., v. 58 (2), 169-174
Schistosoma mansoni, activity response in vitro to praziquantel
- Praziquantel (EMBAY 8840; Droncit; Bitricide)
Chubb, J. M.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (2), 284-293
praziquantel, effects on electromechanical properties of isolated rat atria
- Praziquantel
Coles, G. C., 1979, J. Helminth., v. 53 (1), 31-33
Schistosoma mansoni, praziquantel, effect on different life cycle stages, possible mechanism of action
- Droncit (Praziquantel)
Dey-Hazra, A., 1976, Vet. Med. Rev. (2), 134-141
tapeworms, dogs and cats, droncit: India
- Praziquantel (Droncit)
Diekmann, H. W., 1979, European J. Drug Metab. and Pharmacokinet., v. 4 (3), 139-141
quantitative determination of praziquantel in body fluids by gas chromatography
- Praziquantel
Diekmann, H. W.; and Buehring, K. U., 1976, European J. Drug Metab. and Pharmacokinet., v. 1 (2), 107-112
praziquantel, metabolism in rat, beagle dog, and rhesus monkey
- Praziquantel (Droncit)
Evans, J. W.; and Green, P. E., 1978, Austral. Vet. J., v. 54 (9), 454-455 [Letter]
Platynosomum concinnum, cats, anthelmintics, drug trials

Praziquantel -- Continued.

- Praziquantel (Droncit)
Gallie, G. J.; and Sewell, M. M. H., 1978, Trop. Animal Health and Prod., v. 10 (1), 36-38
Taenia saginata, calves (exper.), praziquantel, ineffective against 4-week-old cysticerci, advisable to let at least 3 months elapse after initial infection for successful treatment
- Praziquantel (EMBAY 8440)
Goennert, R.; and Andrews, P., 1977, Ztschr. Parasitenk., v. 52 (2), 129-150
Schistosoma mansoni in mice, Mastomys, and hamster, praziquantel, various routes of administration compared, all effective; fractional doses double efficacy of single dose; more effective against invading and mature stages than against juveniles
- Droncit
Guralp, N.; et al., 1976, Vet. Fak. Dergisi, Ankara Univ., v. 23 (1-2), 171-174
tapeworms, dogs and cats, droncit 100% effective
- Droncit (Praziquantel)
Guralp, N.; et al., 1976, Vet. Med. Rev. (2), 129-133
tapeworms, dogs, cats, droncit
- Praziquantel (Droncit; Embay 8440)
Guralp, N.; Oguz, T.; and Zeybek, H., 1977, Vet. Fak. Dergisi, Ankara Univ., v. 24 (1), 85-89
Dicrocoelium dendriticum, sheep, chemotherapeutic trials with embay 8440
- Praziquantel (Embay 8440)
Heath, D. D.; and Lawrence, S. B., 1978, N. Zealand Vet. J., v. 26 (1-2), 11-15
Echinococcus granulosus, Taenia hydatigena, T. ovis, sheep, effect of extended oral dosing regime with mebendazole compared with one parenteral injection of either mebendazole or praziquantel
- Praziquantel (Droncit)
Hoerchner, F.; and Albert, H., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (6), 107-111
Taenia saginata, calves inoculated with eggs, response to reinfection and/or drug therapy (mebendazole, praziquantel)
- Praziquantel
James, C.; Webbe, G.; and Nelson, G. S., 1977, Ztschr. Parasitenk., v. 52 (2), 179-194
Schistosoma haematobium and S. mansoni in baboons, S. japonicum in vervet monkeys, praziquantel in various dosages, good results
- Droncit (Praziquantel)
Kobulej, T.; and Varga, I., 1978, Magy. Allat. Lapja, v. 33 (5), 306-308
Echinococcus granulosus, dogs (exper.), droncit, highly effective, no side effects

Praziquantel -- Continued.

- Praziquantel
Koebler, P.; and Bachmann, R., 1978, Molec. Pharm., v. 14 (1), 155-163
Ascaris suum muscle tissue, comparison of effects of levamisole, thiabendazole, chloroquine, and praziquantel on electron transport in Ascaris muscle submitochondrial particles
- Praziquantel (Droncit)
Koudela, B.; and Schanzel, H., 1978, Acta Vet. Brno, v. 47 (1-2), 87-90
Cysticercus pisiformis, rabbits, praziquantel, negative result
- Praziquantel
Machemer, L.; and Lorke, D., 1978, Arch. Toxicol., v. 39 (3), 187-197
praziquantel, mutagenicity studies on mice and Cricetulus griseus, no indication of mutagenic action, compared with cyclophosphamide and placebo
- Droncit (Praziquantel)
Muermann, P.; von Eberstein, M.; and Froberg, H., 1976, Vet. Med. Rev. (2), 142-153
droncit, animals and man, toxicity and local tolerance after single and repeated doses, sensitising and teratogenic properties
- Embay 8440 (Droncit, Praziquantel)
Oguz, T., 1976, Vet. Fak. Dergisi, Ankara Univ., v. 23 (3-4), 385-395
Cysticercus tenuicollis, lambs (exper.), efficacy of embay 8440 and mebendazole
- Praziquantel
Patzschke, K.; et al., 1979, European J. Drug Metab. and Pharmacokinet., v. 4 (3), 149-156
praziquantel, serum concentrations and renal excretion in humans after oral administration, results of 3 determination methods
- Praziquantel
Pawlowski, Z.; Kozakiewicz, B.; and Wroblewski, H., 1978, Vet. Sc. Commun., v. 2 (2), 137-139
Taenia saginata cysticerci, bullocks (exper.), mebendazole and praziquantel administered in fodder, drug efficacy
- Praziquantel
Pax, R.; Bennett, J. L.; and Fetterer, R., 1978, Arch. Pharmacol., v. 304 (3), 309-315
Schistosoma mansoni, S. japonicum, praziquantel and Ro 11-3128 produce rapid rise in tension of musculature, uptake studies of inorganic cations suggest that interference with inorganic ion transport mechanisms causes contraction of schistosome musculature
- Praziquantel
Pax, R.; Fetterer, R.; and Bennett, J. L., 1979, Comp. Biochem. and Physiol., v. 64C (1), 123-127
Schistosoma mansoni, effects of fluoxetine and imipramine on adult males in vitro, interactions with 5-hydroxytryptamine-induced contractile activity, interaction with anti-schistosomal compounds praziquantel and R011-3128

Praziquantel -- Continued.

- Praziquantel (EMBAY 8440)
Pellegrino, J.; et al., 1977, Ztschr. Parasitenk., v. 52 (2), 151-168
Schistosoma mansoni in mice, hamsters, and Cebus monkeys, praziquantel effective, results compared in relation to routes of administration, dosages, degree of hepatic shift
- Praziquantel (Droncit)
Preiss, H.; and Luebke, R., 1978, Kleintier-Praxis, v. 23 (7), 345-346, 349-350
Echinococcus granulosus, dog, praziquantel, clinical aspects
- Praziquantel (Embay 8440)
Puetter, J., 1979, European J. Drug Metab. and Pharmacokinet., v. 4 (3), 143-148
fluorometric method for determining praziquantel in blood-plasma and urine
- Praziquantel (Droncit)
Sakamoto, T., 1977, Vet. Med. Rev. (1), 64-74
adult tapeworms in cats or dogs, praziquantel highly effective, dosage range
- Droncit (Praziquantel)
Sakamoto, T., 1977, Vet.-Med. Nachr. (1), 64-74
adult tapeworms, dogs, cats, droncit, drug efficacy at various doses
- Praziquantel (Droncit)
Sakamoto, T., 1977, Vet.-Med. Nachr. (2), 153-162
Cysticercus fasciolaris, Mesocestoides corti, Echinococcus multilocularis, laboratory mice and rats, praziquantel, evaluation of activity against larval stages
- Praziquantel
Sakamoto, T.; et al., 1979, Bull. Fac. Agric. Kagoshima Univ. (29), 81-87
cestodes, cats, dogs, goats, praziquantel
- Praziquantel (Embay 8440)
Steiner, K.; et al., 1976, European J. Drug Metab. and Pharmacokinet., v. 1 (2), 85-95
praziquantel, pharmacokinetics in animals
- Praziquantel (Embay 8440)
Steiner, K.; and Garbe, A., 1976, European J. Drug Metab. and Pharmacokinet., v. 1 (2), 97-106
praziquantel, distribution in rats
- Droncit (Praziquantel)
Thakur, A. S.; Prezioso, U.; and Marchevsky, N., 1978, Am. J. Vet. Research, v. 39 (5), 859-860
Echinococcus granulosus, dogs (exper.), droncit, 100% effective against both immature and gravid worms, no adverse reactions

Praziquantel -- Continued.

- Praziquantel (Droncit)
Thakur, A. S.; Prezioso, U.; and Marchevsky, N., 1979, Exper. Parasitol., v. 47 (2), 131-133
Echinococcus granulosus, praziquantel did not possess absolute ovicidal activity against eggs either within or outside proglottids, bunamidine hydrochloride did not kill eggs within proglottids
- Praziquantel (EMBAY 8440; Droncit)
Thomas, H.; and Goennert, R., 1977, Ztschr. Parasitenk., v. 52 (2), 117-127
Hymenolepis nana and H. microstoma in mice, H. diminuta in rats, good results with praziquantel, in vivo mode of action (immobilization followed by paralysis)
- Praziquantel
Thomas, H.; and Goennert, R., 1978, Research Vet. Sc., v. 24 (1), 20-25
cestodes of cats, dogs, and sheep, praziquantel highly effective in one oral or subcutaneous dose
- Praziquantel (EMBAY 8440)
Thomas, H.; and Goennert, R., 1978, Ztschr. Parasitenk., v. 55 (2), 165-179
cestodes, praziquantel activity tested
- Praziquantel (Droncit; Embay 8440)
Tinar, R.; and Burgu, A., 1978, Vet. Fak. Dergisi, Ankara Univ., v. 25 (3), 366-371
Hymenolepis nana, mice, praziquantel orally and subcutaneously
- Droncit (Praziquantel)
Vasilev, I.; Denev, I.; and Kostov, R., 1977, Vet.-Med. Nachr. (2), 149-152
cestodes of poultry, droncit, highly effective
- Praziquantel (Embay 8440)
Walther, M.; and Sanitz, W., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (7), 131-135
Taenia saginata, calves (exper.), enzyme-linked immunosorbent assay using T. saginata and T. crassiceps antigens, comparison with indirect hemagglutination during course of infection and following praziquantel treatment
- Praziquantel (EMBAY 8440)
Webbe, G.; and James, C., 1977, Ztschr. Parasitenk., v. 52 (2), 169-177
Schistosoma spp. in hamsters, praziquantel effective against 5 species
- Praziquantel (Droncit)
Wikerhauser, T.; Brglez, J.; and Kuticic, V., 1978, Acta Parasitol. Iugoslavica, v. 9 (2), 57-63
Echinococcus granulosus, mice, mebendazole, praziquantel

Prednisolone -- Canaural (with Neomycin and Ny-
statin).

Prednisolone

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, *Antimicrob. Agents and Chemotherapy*, v.
15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

Canaural

Pott, J. M.; and Riley, C. J., 1979, *Vet. Rec.*,
v. 104 (25), 579

Otodectes cynotis, dogs, cats (ear canal of
both), canaural, comparison with proprietary
topical ear preparation (clendrol), con-
trolled trial

Prednisolone

Watson, A. D. J.; Farrow, B. R. H.; and
Hoskins, L. P., 1978, *Austral. Vet. Practitioner*,
v. 8 (3), 129-132

Haemobartonella felis, cats (nat. and exper.),
thiacetarsamide sodium therapy; prednisolone,
tetracycline, and chloramphenicol given se-
quentially

Prednisone

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, *Antimicrob. Agents and Chemotherapy*, v.
15 (2), 157-160

Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

Preparation-665. See Mekarzole.

Preparation 5006. See Propoxur.

Previken. See Hexachlorophene.

Primaquine -- 8-(4-Amino-1-methylbutylamino)-6-
methoxyquinoline; Camoprims (with Amodiaquine);
Primaquine diphosphate; Primaquine phosphate;
WR 2,975.

Primaquine

Blandon, R.; Guevara, J. F.; and Johnson, C.
M., 1976, *Rev. Med. Panama*, v. 1 (3), 153-162

Chagas disease in children, diagnosis, pa-
thology, therapeutic trials with various
drugs, metronidazole was well tolerated and
therefore most promising therapy: Panama

Camoprims + Maloprims

Botelho, A., 1974, *South African Med. J.*, v. 48
(31), 1351-1352

malarias, humans, clinical trials using sul-
phones and sulphonamides with a pyrimidine
derivative

Primaquine

Brener, Z., 1971, *Rev. Inst. Med. Trop. S.
Paulo*, v. 13 (4), 302-306

Trypanosoma cruzi, description of method
allowing study of drug action on trypomasti-
gotes in mice

Primaquine -- Continued.

Primaquine diphosphate

Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Primaquine

Carroll, F. I.; et al., 1979, *J. Med. Chem.*,
v. 22 (6), 694-699

Plasmodium spp. in mice and rhesus monkeys,
synthesis and antimalarial activity of some
4-substituted 8-amino-6-methoxyquinolines,
4-ethylprimaquine was approximately as active
and was less toxic than primaquine

Primaquine

Carroll, F. I.; Berrang, B.; and Linn, C. P.,
1978, *J. Med. Chem.*, v. 21 (4), 326-330

Plasmodium berghei, *P. cynomolgi*, experiment-
al animals, resolution of antimalarial agents
via complex formation with α -(2,4,5,7-tetra-
nitro-9-fluorenylideneaminoxy)propionic
acid, significant differences in toxicity

Primaquine

Cedillos, R. A.; Warren, M.; and Jeffery, G.
M., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27
(3), 466-472

Plasmodium vivax, humans, 2 regimens of
primaquine (5-day; single dose) in combina-
tion with amodiaquine compared with amodia-
quine alone, field evaluation: Zapotitan
Valley, El Salvador

Primaquine diphosphate

Field, R. C.; et al., 1978, *Brit. J. Pharmacol.*,
v. 62 (2), 159-164

effects of chloroquine, primaquine and ethid-
ium on precursor incorporation into DNA,
RNA, and protein in mammalian tissues

Primaquine diphosphate

Gutteridge, W. E.; Cover, B.; and Gaborak, M.,
1978, *Ann. Trop. Med. and Parasitol.*, v. 72
(4), 329-338

Trypanosoma cruzi, rapid, simple primary
screen to test compounds for activity as
potential trypanocides using infected A/JAX
inbred mice

WR 2 975

Kinnamon, K. E.; et al., 1978, *Am. J. Trop.
Med. and Hyg.*, v. 27 (4), 751-757

Leishmania donovani-*Mesocricetus auratus*
model, antileishmanial activity of lepidines
(6-methoxy-4-methyl-8-aminoquinoline deriv-
atives)

Primaquine

McHardy, N., 1978, *Ann. Trop. Med. and Parasi-
tol.*, v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected
bovine lymphoblastoid cell cultures used in
in vitro screens to test wide range of
compounds for chemotherapeutic activity

Primaquine phosphate

Marshall, R. J.; and Ojewole, J. A. O., 1978,
Toxicol. and Applied Pharm., v. 46 (3), 759-
768

quinoline and nonquinoline antimalarial
drugs, effects on isolated guinea pig car-
diac muscle

Primaquine -- Continued.

Primaquine

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, Israel J. Med. Sc., v. 14 (5), 601-605

Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiparasite immunity by natural bites of infected mosquitoes, symposium presentation

Primaquine + Chloroquine

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, Israel J. Med. Sc., v. 14 (5), 601-605

Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiparasite immunity by natural bites of infected mosquitoes, symposium presentation

Primaquine diphosphate

Pirson, P.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (3), 347 [Letter]

Plasmodium berghei, mice (exper.), liposome-entrapped primaquine used as therapy against pre-erythrocytic stage infection, less toxic than free primaquine

Primaquine (WR 2,975)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947

sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Primaquine

Rassi, A.; and Ferreira, H. de O., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (5), 235-262

Trypanosoma cruzi, extensive clinical trials testing efficacy of various nitrofuranes (singly, mixed nitrofuranes, or in association with primaquine), therapeutic response as based mainly on xenodiagnosis and the Guerreiro-Machado test showed nitrofurazone and especially Bayer 2502 to give best response

Primaquine phosphate

Stoskopf, M. K.; and Beier, J., 1979, J. Am. Vet. Med. Ass., v. 175 (9), 944-947

Plasmodium relictum, P. elongatum in Spheniscus demersus, diagnostic methods evaluated, chloroquine phosphate and primaquine phosphate therapy: Baltimore Zoo

Primaquine

Tanabe, K.; et al., 1978, J. Med. Chem., v. 21 (1), 133-136

Plasmodium cynomolgi-rhesus monkey antimalarial screening of 5-phenylthio and 5-anilino derivatives of primaquine in an attempt to diminish toxicity of primaquine

Primaquine

Thong, Y. H.; Ferrante, A.; and Rowan-Kelly, B., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 537-539

primaquine possesses potent immunosuppressive activity at concentrations within therapeutic range for vivax malaria

Primaquine -- Continued.

Primaquine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Primaquine analogues, 2-substituted

Shetty, R. V.; and Blanton, C. D., jr., 1978, J. Med. Chem., v. 21 (9), 995-998

2-substituted primaquine analogues synthesized and evaluated in laboratory animals against Plasmodium berghei, P. cynomolgi, Trypanosoma rhodesiense, Leishmania donovani and Schistosoma mansoni; significant activity was observed against P. berghei and L. donovani

Primaquine derivatives

Tanabe, K.; et al., 1978, J. Med. Chem., v. 21 (1), 133-136

Plasmodium cynomolgi-rhesus monkey antimalarial screening of 5-phenylthio and 5-anilino derivatives of primaquine in an attempt to diminish toxicity of primaquine

Primaquine diphosphate. See Primaquine.

Primaquine phosphate. See Primaquine.

Prioderm. See Malathion.

Proadifen hydrochloride -- 2-Diethylaminoethyl-2,2-diphenylvalerate hydrochloride; SKF 525-A.

Proadifen hydrochloride (SKF 525-A)

Schuntner, C. A.; and Thompson, P. G., 1978, Austral. J. Biol. Sc., v. 31 (2), 141-148

Boophilus microplus, larvae, ¹⁴C labelled amitraz, metabolism, only amitraz and N-2,4-dimethylphenyl-N'-methylformamide (metabolite) toxic to larvae, piperonyl butoxide applied simultaneously with amitraz had slight effect on metabolism, three-fold synergistic effect; SKF 525-A similarly applied had negligible effect on both metabolism and toxicity to ticks

Procaine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Procaine penicillin. See Penicillin.

Proflavin hemisulphate
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Progesterone
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S.,
1979, *Antimicrob. Agents and Chemotherapy*, v.
15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in
screening of antitumor compounds for efficacy
against infection

Proguanil. See Chlorguanide.

Proguanil hydrochloride. See Chlorguanide.

Prolate. See Phosmet.

Promecarb
Rawlins, S. C.; and Mansingh, A., 1978, *J.*
Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, suscepti-
bility to acaricides: Jamaica; St. Kitts;
Trinidad; Guyana

Promecarb
Rawlins, S. C.; and Mansingh, A., 1978, *J.*
Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains,
patterns of resistance to acaricides

Promintic. See Methyridine.

Propetamphos -- Blotic; (E)-1-Methylethyl-3-
[[[ethylamino)methoxyphosphinothioyl]oxyl]-
2-butenate; SAN 322 I.

SAN 322 I (Blotic)
Gothe, R.; and Mieth, H., 1978, *Berl. u.*
Munchen. Tierarztl. Wchnschr., v. 91 (10),
193-196
Boophilus microplus strains, Rhipicephalus
appendiculatus, R. evertsi evertsi, SAN
322 I, drug efficacy in vitro

Propolis
Suchy, H.; et al., 1974, *Przegl. Lek.*, v. 31
(6), 646-648
Trichomonas vaginalis, treatment of human
vaginal infections with ethanol solutions of
propolis with good results and quick healing

Propoxur -- Arprocarb; Bayer 39007; Baygon;
Blattanex; Bolfo; 1-Isopropoxyphenyl-N-
methylcarbamate; Preparation 5006; Uden.

Propoxur
Blommers, L., 1979, *J. Med. Entom.*, v. 16 (1),
82-83
Pediculus capitis, nymph rearing technique,
insecticide tests against nymphs

Propoxur -- Continued.

Propoxur
Blommers, L.; and van Lennep, M., 1978, *Acta*
Leidensia, v. 46, 9-15
Pediculus capitis, susceptibility of 1st
instar nymphs to malathion, propoxur, and
plifenate, residual action of 3 compounds
applied as lotions to human hair

Propoxur
Drummond, R. O.; et al., 1973, *J. Econom. En-*
tom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory
tests of insecticides

Propoxur
Frazar, E. D.; and Schmidt, C. D., 1979, *J.*
Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, sus-
ceptibility to topically applied insecticides

Baygon (Arprocarb; Blattanex; Preparation 5006;
Propoxur; Uden; Bayer 39007)
Frolov, B. A.; et al., 1978, *Veterinariia*,
Moskva (11), 75-76
Dermanyssus gallinae, Argas persicus,
chickens, acaricides tested

Propoxur
Gladney, W. J.; Price, M. A.; and Graham, O.
H., 1977, *J. Med. Entom.*, v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of
various insecticides applied as sprays, ear
smears and dusts, or in slow-release devices,
field tests

Propoxur
Kibble, R., 1979, *Austral. Vet. J.*, v. 55 (1),
40-41 [Letter]
fleas, ticks, dogs, DDVP and propoxur-impreg-
nated collars, recommendations: Australia

Propoxur (Bolfo)
Niemand, H. G.; Niemand, S.; and Wendel, E.,
1979, *Kleintier-Praxis*, v. 24 (4), 173-175
discussion of chief effective principles
of different flea collars (dichlorvos,
diazinon, propoxur), toxicity

Bolfo
Rak, H., 1976, *J. Vet. Fac. Univ. Tehran*,
v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-
tox, asuntol 50, alon, bolfo, alugan, opigal,
gamatox, tetmosol, neguvon: Iran

Propoxur
Rawlins, S. C.; and Mansingh, A., 1978, *J.*
Econom. Entom., v. 71 (1), 142-144
Boophilus microplus, five strains, suscepti-
bility to acaricides: Jamaica; St. Kitts;
Trinidad; Guyana

Propoxur
Rawlins, S. C.; and Mansingh, A., 1978, *J.*
Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains,
patterns of resistance to acaricides

Propoxur
Rawlins, S. C.; and Mansingh, A., 1979, *J.*
Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal
residues on grass surfaces, greenhouse and
pasture studies

Propylene glycol -- PB Dressing (with Butyl amino-benzoate and Piperonyl butoxide).

PB Dressing

Nie, I. A.; and Pick, C. R., 1978, J. Inst. Animal Techn., v. 29 (1), 63-68
Otodectes coynotis, ferrets (ears), PB dressing: laboratory colony, Mill Hill, London

Prostaglandin E.1

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Prostaglandin E.2

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Protargol

Rostkowska, J., 1970, Acta Parasitol. Polon., v. 18 (27-41), 377-392
Balantidium coli, effect of various bacteria on propagation in vitro, on erythrophagocytic capability of balantidia, and on susceptibility of balantidia to atebirin, entobex, mexaform, and protargol; *Trichomonas hominis*, *Chilomastix mesnili*, and *Dientamoeba fragilis* found to be without effect; effect of balantidia on bacteria

Prothidium -- Prothidium dimethyl Br; Pyrithidium bromide.

Prothidium dimethyl Br

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Pyrithidium bromide (Prothidium)

James, D. M., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 471-476
[Trypanosoma] congolense, *T. brucei*, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs

Prothidium dimethyl Br. See Prothidium.

Protium heptaphyllum

Frischkorn, C. G. B.; Frischkorn, H. E.; and Carrazzoni, E., 1978, Naturwissenschaften, v. 65 (9), 480-483
Schistosoma mansoni, cercaricidal activity of various plants from northeast Brazil

Protoporphyrin IX

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Protoporphyrin IX

Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, Biochem. Pharmacol., v. 26 (20), 1923-1928
Trypanosoma brucei, *T. congolense*, heme lysis of bloodstream forms, *T. brucei*, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism

Protozoacides

Alvarez Chacon, R., 1975, Semana Med. Mexico (1071), an. 22, v. 84 (11), 328-332
 parasites of children, drugs in current use, treatment recommendations and contraindications

Protozoacides

Amato Neto, V., 1973, Rev. Paul. Med., v. 81 (2), 93-104
 human protozoan infections, drugs in current use, dosage recommendations, review

Protozoacides

Catchpool, J. F., 1974, Rev. Med. Pharmacol., 4. ed., 594-620
 antiprotozoal drugs in current use, extensive review of modes of action, epidemiologic factors, clinical administration, contraindications and cautions

Protozoacides

Dasgupta, N. K., 1976, Modern pharmacology and therapeutics, 2. ed., 860 pp., illus.

Protozoacides

Lengyel, A.; and Janko, M., 1979, Orvosi Hetilap, v. 120 (49), 2969-2972
 human intestinal parasites, current therapeutic regimens, review: Hungary

Protozoacides

Noemi, I., 1975, Rev. Chilena Pediat., v. 46 (1), 63-67
 therapeutic recommendations for treating intestinal parasites in infants and small children, drug and dosage review

Protozoacides

Seo, B. S., 1977, Taehan Uihak Hyophoe Chi (J. Korean Med. Ass.), v. 20 (6), 503-508
 human intestinal parasitic disease, current therapy, review

Protozoacides

Timofeev, B. A.; and Karpenko, I. G., 1977, [Chemotherapy of protozoan diseases of livestock], 99 pp.
 protozoan diseases of livestock, manual of drugs and dosages

Pteridines

Peroutka, M.; and Cihar, R., 1978, Apidologie, v. 9 (4), 291-304
Nosema apis in drones (exper.), effect of pollen, pharyngeal gland secretions, and pteridines (stimulation, inhibition, or no effect) on parasite reproduction

- Pulicaria dysenterica** herb
Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118
Ascaridia galli, *Ascaris vitulorum*, in vitro anthelmintic activity of some Egyptian plants; only *Nerium oleander* caused death of worms
- Pulicaria undulata** herb
Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, *Scientia Pharm.*, v. 47 (2), 114-118
Ascaridia galli, *Ascaris vitulorum*, in vitro anthelmintic activity of some Egyptian plants; only *Nerium oleander* caused death of worms
- Pumpkin seed based taeniafuge**
Komma, M. D.; and Santos, V. L. V., 1972, *Rev. Patol. Trop.*, v. 1 (1), 69-72
Taenia solium, *T. saginata*, humans, comparative therapeutic trials using yomesan and pumpkin seed based taeniafuge; correlation of species of *Taenia* with type of therapy and recovery of scolex with therapy
- Punica granatum.** See Pomegranate.
- Purinribosid**
Christow, C., 1978, *Biochem. and Exper. Biol.*, v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂ antagonists upon growth
- Puromycin** -- Puromycin dihydrochloride; Puromycin 2HCL.
- Puromycin**
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter
- Puromycin 2HCl**
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
- Puromycin dihydrochloride**
Sinden, R. E.; and Smalley, M. E., 1979, *Parasitology*, v. 79 (2), 277-296
Plasmodium falciparum, modified microculture technique used as bioassay for various antimetabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle
- Puromycin**
Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Puromycin aminonucleoside**
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
- Puromycin aminonucleoside**
Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- Puromycin dihydrochloride.** See Puromycin.
- Puromycin 2HCl.** See Puromycin.
- PVC-DDVP.** See Dichlorvos.
- Pyractone M429.** See Bucarpolate or Pyrethrins.
- Pyractone M817.** See Piperonyl butoxide or Pyrethrins.
- Pyrantel** -- Antiminth; Banminth; Banminth paste; Combantrin; Helmex; (1-Methyl-1,1,4,5,6-tetrahydro-2,2-thienyl)vinyl-pyrimidinium; trans-1-Methyl-2-[2-(thienyl)-vinyl]-1,4,5,6-tetrahydropyrimidine; Pyrantel citrate; Pyrantel embonate; Pyrantel pamoate; Pyrantel tartrate; Pyrantel tartrate; Strongid-P granules or paste; Strongid-T; U.K. 2679.
- Banminth (Pyrantel tartrate)**
Abdel Rahman, M. S.; et al., 1978, *Vet. Med.* J., Giza, v. 24 (24), 1976, 39-48
gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt
- Pyrantel pamoate (U.K. 2679)**
Amato Neto, V.; Levi, G. C.; and Campos, L. L., 1970, *Rev. Inst. Med. Trop. S. Paulo*, v. 12 (3), 207-210
Ascaris lumbricoides, human carriers, treatment trials using pyrantel pamoate showed cure rate of 95%: Brazil
- Pyrantel tartarate**
Anosa, V. O., 1978, *Vet. Parasitol.*, v. 4 (2), 103-109
seasonal variations in blood picture of white Fulani calves naturally exposed to helminthiasis, effect of pyrantel tartarate treatment, results indicate that low hematological values in African cattle are not inherited characteristic but are rather due to seasonal effects of parasitism and nutrition inherent in indigenous husbandry practices

Pyrantel -- Continued.

Pyrantel pamoate

Araki, T.; Nakazato, H.; and Ikoma, K., 1976, *Kiseichugaku Zasshi* (Japan. J. Parasitol.), v. 25 (3), 153-160
helminthiasis, human, serum IgE levels before and after treatment with pyrantel pamoate, radioimmunosorbent technique and single radial immunodiffusion method

Pyrantel pamoate

Baranski, M. C.; et al., 1971, *Rev. Inst. Med. Trop. S. Paulo*, v. 13 (6), 422-427
Enterobius vermicularis, humans, pyrantel pamoate compared with pyrvinium pamoate: Brazil

Pyrantel pamoate

Baranski, M. C.; et al., 1976, *AMB, Rev. Ass. Med. Brasil.*, v. 22 (7), 257-260
Necator americanus, 60 patients, anthelmintic activity of pyrantel pamoate: Brazil

Pyrantel embonate

Behnke, J. M.; and Parish, H. A., 1979, *Exper. Parasitol.*, v. 47 (1), 116-127
Nematospiroides dubius, arrested development of larvae in immune mice, resumption of development after cortisone treatment, arrested larvae were insusceptible to activity of pyrantel embonate

Pyrantel pamoate + Trichlorfon

Bentley, O. E.; et al., 1978, *Vet. Med. and Small Animal Clin.*, v. 73 (1), 70-73
horses, pyrantel pamoate + trichlorfon, field trials, results show that drugs are safe for horses when preceded by a feeding of grain or a complete horse ration, mild side effects: Kentucky; Alabama; Texas; Kansas

Pyrantel pamoate

Bhaibulaya, M.; et al., 1977, *Southeast Asian J. Trop. Med. and Pub. Health*, v. 8 (3), 329-334
human soil-transmitted helminths, clinical trials with pyrantel pamoate given prophylactically to villagers in agricultural areas in an attempt to control spread of infections, mixed results in infection rate findings post treatment: village near Bangkok, Thailand

Pyrantel pamoate

Botero R., D., 1978, *Ann. Rev. Pharmacol. and Toxicol.*, v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Pyrantel pamoate (Combantrin)

Carlos, E. T.; and Dizon, O. R., 1972, *Philippine J. Vet. Med.*, v. 11 (2), 156-159
ascarids, *Ancylostoma*, puppies, pyrantel pamoate

Pyrantel pamoate (Combantrin)

Chiu, J. K.; et al., 1975, *Taiwan i Hsueh Hui Tsa Chih* (J. Formosan Med. Ass.), v. 74 (11), 728-733
Enterobius vermicularis, pre-school and first-grade children, prevalence survey, clinical trials with pyrantel pamoate: Taipei City, Taiwan

Pyrantel -- Continued.

Pyrantel pamoate

Cho, S. Y., 1976, Taehan Yangnihak Chapchi (Korean J. Pharmacol.), v. 12 (1), 69-73
nematodes, human, evaluation of efficacy of combined therapy of pyrantel pamoate and oxantel pamoate

Pyrantel embonate (Strongid-P granules or paste)

Clayton, H. M.; Duncan, J. L.; and Gilbert, G. A., 1979, *Vet. Rec.*, v. 105 (17), 389-391
ascarids, strongyles, horses and ponies, pyrantel embonate, field trials for long-term control

Pyrantel pamoate (Combantrin)

da Costa, F. C., 1976, *Rev. Portug. Clin. e Terap.*, v. 2 (3), 131-140
Ascaris lumbricoides, *Trichocephalus dispar*, humans, statistics of prevalence survey, therapeutic trials with pyrantel pamoate: Portugal

Pyrantel pamoate

Dudley, B. W.; and Lease, E. J., 1973, *South. Med. J.*, v. 66 (10), 1100, 1126
ascariasis, schoolchildren, disease control using repetitive treatment (piperazine or pyrantel pamoate) on a mass therapy basis: South Carolina

Pyrantel tartrate (Banminth)

Enigk, K.; Dey-Hazra, A.; and Batke, J., 1975, *Acta Parasitol. Polon.*, v. 23 (26-40), 367-372
helminths of geese, treatment (Amidostomum anseris with mebendazol, fenbendazol, levamisole, and pyrantel tartrate; *Trichostrongylus tenuis* with mebendazol and fenbendazol; *Drepanidotaenia lanceolata* and *Hymenolepis setigera* with mebendazol)

Pyrantel pamoate (Helmex)

Felder, J., 1975, *Therap. Gegenw.*, v. 114 (9), 1438-1444
oxyuriasis, children, therapy with helmex

Pyrantel pamoate (Combantrin)

Goldsmid, J. M.; and Saunders, C. R., 1973, *South African Med. J.*, v. 47 (6), 205-206
hookworms, humans, pyrantel pamoate, clinical trials: Rhodesian Lowveld

Pyrantel pamoate

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, *Rev. Brasil. Med.*, v. 31 (11), 791-794
human soil transmitted nematodes, laboratory trials testing ovicidal and larvicidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Pyrantel pamoate (Combantrin)

Hasbun, E. R.; and Zepeda, J. R., 1977, *Rev. Inst. Invest. Med.*, San Salvador, v. 6 (2), 190-192
Ancylostomidae, humans, clinical trials testing efficacy of pyrantel pamoate and phenylene diisothiocyanate, medications well tolerated with few side effects: El Salvador

Pyrantel -- Continued.

- Pyrantel pamoate (Combantrin)
Hatchuel, W.; Isaaacson, M.; and de Villiers, D. J., 1973, South African Med. J., v. 47 (3), 91-93
Ascaris lumbricoides, children, pyrantel pamoate vs. piperazine citrate, clinical trials: Alexandra Township near Johannesburg
- Pyrantel pamoate (Combantrin)
Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167
Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria
- Pyrantel pamoate (Combantrin)
Kan, S. K. P., 1979, Ann. Trop. Med. and Parasitol., v. 73 (4), 397-398
atypical nematode ova, probably Ascaris, 19-year-old girl (feces) after treatment with pyrantel pamoate; mebendazole therapy resulted in ova-free feces: Malaysia
- Pyrantel pamoate
Katz, N.; et al., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (3), 212-221
nematodes, human, clinical trials of pyrantel pamoate
- Pyrantel pamoate
Katz, N.; Antunes, C. M. F.; and Zicker, F., 1973, Rev. Inst. Med. Trop. S. Paulo, v. 15 (5), 331-339
ancylostomiasis, trichuriasis, human, clinical trials with pyrantel pamoate; Nematospiroides dubius-infected white mice, pyrantel pamoate is ineffective against larval Ancylostomidae
- Pyrantel pamoate (Banminth paste)
Kaushik, R. K.; Banerjee, D. P.; and Singh, R. P., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 50-54
Toxocara canis, dogs (nat. and exper.), pyrantel pamoate effective with no side effects; no ovicidal or larvicidal effect in vitro
- Pyrantel tartrate
Kerboeuf, D.; Hubert, J.; and Le Stang, J. P., 1979, Rec. Med. Vet., v. 155 (2), 143-148
Ostertagia ostertagi, cattle, serum pepsinogen levels in relation to worm burden and anthelmintic treatments
- Pyrantel pamoate
Klein, J. B.; Bradley, R. E., sr.; and Conway, D. P., 1978, Vet. Med. and Small Animal Clin., v. 73 (8), 1011-1013
Toxocara canis and Ancylostoma caninum, dogs, pyrantel pamoate effective against adult stages, not effective against Dipylidium caninum, Trichuris vulpis, or Dirofilaria immitis
- Pyrantel tartrate (Banminth)
Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in Mastomys natalensis, efficacy of various anthelmintics against third stage larvae

Pyrantel -- Continued.

- Pyrantel tartrate
Le Stang, J.-P.; Hubert, J.; and Kerboeuf, D., 1978, Rev. Med. Vet., Toulouse, v. 129 (10), 1355-1362, 1365-1369
gastrointestinal nematodes, cattle, efficacy of fenbendazole and pyrantel tartrate
- Pyrantel pamoate (U.K. 2679)
Levi, G. C.; et al., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (5), 343-346
Enterobius vermicularis, Ascaris lumbricoides, humans, pyrantel pamoate: Brazil
- Pyrantel pamoate
Levi, G. C.; et al., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 392-396
ascariasis, human, clinical trials comparing piperazine, tetramisole (or levamisole) and pyrantel pamoate: Brazil
- Pyrantel pamoate
Lindquist, W. D., 1978, Am. J. Vet. Research, v. 39 (11), 1868-1869
Baylisascaris procyonis, mice (exper.), pyrantel tartrate and pyrantel pamoate administered in feed prevented cerebral migration, concluded that parasite might be useful model to test anthelmintics against migratory ascarids
- Pyrantel tartrate
Lindquist, W. D., 1978, Am. J. Vet. Research, v. 39 (11), 1868-1869
Baylisascaris procyonis, mice (exper.), pyrantel tartrate and pyrantel pamoate administered in feed prevented cerebral migration, concluded that parasite might be useful model to test anthelmintics against migratory ascarids
- Pyrantel pamoate (Antiminth)
Loebenberg, D.; et al., 1979, J. Parasitol., v. 65 (5), 823-824
Syphacia obvelata, mice, Sch 23154 compared with pyrantel pamoate and pyrvinium pamoate
- Pyrantel pamoate (Combantrin)
Louzada, G. Z.; et al., 1972, Rev. Brasil. Clin. e Terap., v. 1 (4), 247-252
human intestinal parasites, clinical trials with pyrantel pamoate, considered drug of choice for ascariasis, enterobiasis and ancylostomiasis: Brazil
- Pyrantel pamoate (Combantrin)
Lynch, P.; and Parry, E., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 658-659
hookworm, whipworm, roundworm, prevalence in Gurkhas recruited into British Army, efficacy of pyrantel pamoate: native to Nepal
- Pyrantel tartrate (Banminth)
Mallick, K. P.; Mishra, S. S.; and Sahai, B. N., 1977, Indian J. Animal Research, v. 11 (2), 77-80
Haemonchus spp., goats (exper.), thiabendazole, pyrantel tartrate, krimnos, comparative efficacy, blood picture before and after treatment, krimnos not effective

Pyrantel -- Continued.

Pyrantel pamoate (Combantrin)

Migasena, S.; Suntharasamai, P.; and Harinasuta, T., 1978, Ann. Trop. Med. and Parasitol., v. 72 (2), 199-200

Necator americanus, humans, treatment trials comparing efficacy of mebendazole, tetrachlorethylene and pyrantel pamoate: Bangkok

Pyrantel pamoate

Miller, M. J.; et al., 1978, South. Med. J., v. 71 (2), 137-140

ascariasis, children throughout the world, levamisole vs. known anthelmintics, levamisole recommended for mass chemotherapy: Iran, Brazil, Mississippi, Louisiana

Pyrantel pamoate (Combantrin)

Noemi, I.; et al., 1975, Rev. Chilena Pediat., v. 46 (2), 107-110

Enterobius vermicularis, *Ascaris lumbricoides*, children, therapeutic trials with a single oral dose of pyrantel pamoate, few side effects: Santiago, Chile

Pyrantel tartrate

Rao, Y. V. B. G., 1976, Indian Vet. J., v. 53 (10), 776-777

Capillaria obsignata, chickens (exper.), critical tests with methyridine, pyrantel tartrate, and levamisole

Pyrantel pamoate (Combantrin)

Retnasabapathy, A.; and Baskaran, G., 1976, v. 53 (10), 806-811

ancylostomiasis, dogs, morantel tartrate, pyrantel pamoate and disophenol effective

Pyrantel citrate

Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235

Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parabendazole and dichlorvos

Pyrantel tartrate

Robinson, M., 1979, Vet. Parasitol., v. 5 (2-3), 223-235

Trichuris suis, pigs (exper.), oxantel tartrate alone or in combination with tartrate or citrate salts of pyrantel or morantel, comparison with parabendazole and dichlorvos

Pyrantel

Sanchez Moreno, M.; and Barrett, J., 1979, Parasitology, v. 78 (1), 1-5

Hymenolepis diminuta, adults, monoamine oxidase, occurrence and properties, inhibition by several anthelmintics

Pyrantel pamoate

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426

Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Pyrantel pamoate (Combantrin)

Santadvoot, C.; et al., 1975, Siriraj Hosp. Gaz., v. 27 (7), 957-962

enterobiasis, children, clinical trials with pyrantel pamoate

Pyrantel -- Continued.

Pyrantel pamoate

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Pyrantel tartrate (Banminth)

Shelton, M. E.; and Draper, D. D., 1978, Vet. Med. and Small Animal Clin., v. 73 (11), 1423, 1425-1426, 1428

nematodes, dogs, pyrantel tartrate, efficacy of continuous low-level feeding, therapeutic and prophylactic activity

Pyrantel pamoate

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Vet., v. 16 (1), 5-7

Ancylostoma caninum, dogs, pyrantel pamoate, fenbendazole, thiabendazole, clinical trials, comparative efficacy

Pyrantel pamoate

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 226-229

Ancylostoma caninum, dogs, comparative efficacy of disophenol, fenbendazole, pyrantel pamoate, and thiabendazole, clinical trials

Pyrantel pamoate

Singh, H.; Singh, R. P.; and Bali, M. K., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 55-58

Ancylostoma sp., thiabendazole, fenbendazole, ancylool, and pyrantel pamoate, in vitro effects on development of eggs and infective larvae

Pyrantel pamoate (Strongid-T)

Slocombe, J. O. D., 1979, Canad. Vet. J., v. 20 (5), 136-140

Anoplocephala perfoliata, horses, prevalence, efficacy of pyrantel pamoate, mebendazole, and niclosamide in field and critical trials: southern Ontario

Pyrantel pamoate

de Souza, D. W. C.; Neves, J.; and de Lemos, M. S., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (1), 67-72

A[scaris] lumbricoides, humans, pyrantel pamoate and laevo isomer of tetramisole, clinical trials, satisfactory results: Brazil

Pyrantel tartrate (Banminth)

Stewart, T. B.; et al., 1979, Am. J. Vet. Research, v. 40 (10), 1472-1475

Strongyloides dentatus and other parasites, pigs on contaminated lots, pyrantel tartrate and carbadox in feed

Pyrantel pamoate (Combantrin)

Yamaguchi, T.; et al., 1975, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 24 (3), 93-98

Trichostrongylus orientalis, human, pyrantel pamoate

Pyrantel -- Continued.

- Pyrantel pamoate (Combantrin)
Yu, J. C.; and Chiu, J. K., 1976, Taiwan i Hsueh Hui Tsa Chih (J. Formosan Med. Ass.), v. 75 (11), 639-646
Enterobius vermicularis, Ascaris lumbricoides, school children, clinical trials with pyrantel pamoate
- Pyrantel embonate (Combantrin)
Zaman, V.; and Loh, Y. P., 1974, Singapore Med. J., v. 15 (2), 147-148
Necator americanus, humans, clinical trials testing efficacy of pyrantel embonate, levotetramisole, and bephenium hydroxynaphthoate: Singapore

Pyrantel citrate. See Pyrantel.

Pyrantel embonate. See Pyrantel.

Pyrantel pamoate. See Pyrantel.

Pyrantel tartarate. See Pyrantel.

Pyrantel tartrate. See Pyrantel.

Pyrethrins -- Pyractone M429 (with Bucarpolate); Pyractone M817 (with Piperonyl butoxide).

Pyractone M429
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides

Pyractone M817.
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides

Pyrethrins + Bucarpolate
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculus, conventional mouse colony, acaricides

Pyrethrum
Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Pyrethrum + Piperonyl butoxide
Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Pyrethrum
Rajan, U., 1975, Singapore Med. J., v. 16 (4), 297-300
head lice, school children, pyrethrum the drug of choice in comparison clinical trials with benzyl benzoate

4-Pyridinemethanols
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

2-Pyridinyl-5-isothiocyanatobenzimidazoles
Haugwitz, R. D.; et al., 1979, J. Med. Chem., v. 22 (9), 1113-1118
Nematospiroides dubius, Hymenolepis nana, mice, synthesis and activities of 2-pyridinyl-5-isothiocyanatobenzimidazoles

2-Pyridyl-S-adenosine
Trager, W.; Robert-Gero, M.; and Lederer, E., 1978, FEBS Letters, v. 85 (2), 264-266
Plasmodium falciparum, antimalarial activity of S-isobutyl adenosine analogues in culture

Pyrikelfizine. See Pyrimethamine or Sulfalene.

Pyrimethamine -- Abi-Zet₅₀ (with Sulfadimidine); Daraprim; 6-Ethyl-2,4-diamino-5-(p-chlorophenyl)-pyrimidine; Fansidar (with Sulfadoxine); Maloprim (with Dapsone); Pancoxin plus (with Amprolium, Ethopabate and Sulfadoxine); Pyrikelfizine (with Sulfamethoxyprazine); Pyrimethamine isethionate; Supacox (with Amprolium, Ethopabate and Sulfadoxine); Tindurin; WR 2978.

Pyrimethamine
Agarwal, K.; Puri, S. K.; and Dutta, G. P., 1979, Indian J. Med. Research, v. 69, 577-582
Plasmodium berghei, rats, selection of pyrimethamine resistant strain by interrupted subcurative therapy (gradually increasing doses during serial passage); cross sensitivity to other antimalarials

Pyrimethamine + Sulfadoxine (= Fansidar)
Al Tawil, N., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (3), 409-413
Plasmodium falciparum, semi-immune humans, clearance of asexual parasitaemia with single dose sulfadoxine-pyrimethamine, comparison with standard dose of chloroquine over 3 days: Laos

Maloprim + Camoprim
Botelho, A., 1974, South African Med. J., v. 48 (31), 1351-1352
malarias, humans, clinical trials using sulfones and sulphonamides with a pyrimidine derivative

Pyrimethamine + Dapsone (= Maloprim)
Botelho, A., 1974, South African Med. J., v. 48 (31), 1351-1352
malarias, humans, clinical trials using sulfones and sulphonamides with a pyrimidine derivative

Pyrimethamine -- Continued.

Pyrimethamine + Fansil

Botelho, A., 1974, South African Med. J., v. 48 (31), 1351-1352
malarias, humans, clinical trials using sulphones and sulphonamides with a pyrimidine derivative

Pyrimethamine + Dapsone

Buyst, H., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 201-212
sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia

Pyrimethamine + Sulfadoxine (= Fansidar)

Chongsuphajaisiddhi, T.; et al., 1979, South-east Asian J. Trop. Med. and Pub. Health, v. 10 (1), 132-137
falciparum malaria, children, chloroquine resistance, efficacy of quinine and fansidar, clinical study: Thailand

Pyrimethamine

Doberstyn, E. B.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 15-17
Plasmodium vivax, humans, pyrimethamine alone or combined with sulfadoxine is not effective therapy for acute episode: Thailand

Pyrimethamine + Sulfadoxine (= Fansidar)

Doberstyn, E. B.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 15-17
Plasmodium vivax, humans, pyrimethamine alone or combined with sulfadoxine is not effective therapy for acute episode: Thailand

Pyrimethamine

Eke, R. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 1074-1075
Plasmodium falciparum, possible chloroquine-resistant strain, recrudescence of infection in 42-year-old hospital worker after chloroquine therapy, radical cure with sulfadiazine and pyrimethamine: Nigeria

Pyrimethamine

Eislager, E. F.; et al., 1979, J. Med. Chem., v. 22 (10), 1247-1257
Plasmodium spp., antimalarial activity of 2,4-diamino-6-(2-naphthylsulfonyl)quinazoline and related 2,4-diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines

Pyrimethamine + Sulfadoxine (= Fansidar)

Ferraroni, J. J.; and Hayes, J., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 909-911
Plasmodium falciparum outbreak among indigenous indian tribe, 3 cases resistant to chloroquine responded favorably to fansidar therapy: Uauaris, Territory of Roraima, Brazil

Pyrimethamine + Sulfadoxine (= Fansidar)

Fogh, S.; Jepsen, S.; and Effersøe, P., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (2), 228-229
Plasmodium falciparum, chloroquine resistant strain in non-immune male, treatment with fansidar terminated infection: Danish tourist to Kenya

Pyrimethamine -- Continued.

Pyrimethamine (Daraprim)

Furtado, T., 1974, Rev. AMMG, v. 25 (3), 108-113
human cutaneous and mucocutaneous leishmaniasis, recommendations for therapy

Pyrimethamine (Daraprim)

Gon, F.; and Reid, F. P., 1975, South African Med. J., v. 49 (4), 120-122
disseminated intravascular coagulation with haemolytic anaemia and thrombocytopenia, fatal illness in man probably caused by malarial prophylaxis (pyrimethamine and chlor-guanide): Johannesburg, South Africa (from Zambia)

Pyrimethamine

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Pyrimethamine

Herman, Z.; Sokoła, A.; and Szaflarski, J., 1970, Acta Parasitol. Polon., v. 18 (42-50), 483-486
Toxoplasma gondii, mice, comparison of pyrimethamine and sulfamethoxy-pyrazine used separately or in combination and of a long-acting preparation of sulfamethoxy-pyrazine

Pyrimethamine + Sulfamethoxy-pyrazine (= Pyrikel-fizine)

Herman, Z.; Sokoła, A.; and Szaflarski, J., 1970, Acta Parasitol. Polon., v. 18 (42-50), 483-486
Toxoplasma gondii, mice, comparison of pyrimethamine and sulfamethoxy-pyrazine used separately or in combination and of a long-acting preparation of sulfamethoxy-pyrazine

Pyrimethamine + Dapsone (= Maloprim)

Hughes, A.; and Gatus, B. J., 1979, J. Trop. Med. and Hyg., v. 82 (6), 120-121
severe megaloblastic anaemia, woman treated with daily dosage of maloprim

Pyrimethamine

Jadin, J.; Timperman, G.; and De Ruysser, F., 1975, Ann. Soc. Belge Med. Trop., v. 55 (6), 603-608
Plasmodium b. berghei, pyrimethamine-resistant strain preserved at very low temperature for 11 years, maintained virulence but lost drug resistance, gametocytogenesis increased, cyclical transmission was successful, parasites crossed blood-brain barrier indicating this strain could serve as laboratory model for P. falciparum cerebral malaria

Pyrimethamine + Amprolium + Sulphaquinoxaline + Ethopabate (= Supacox)

Long, P. L.; and Millard, B. J., 1978, Avian Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on oocyst output of various treatment regimens

Pyrimethamine -- Continued.

Pyrimethamine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Pyrimethamine

McLeod, R.; et al., 1979, Am. J. Med., v. 67 (4), 711-714

Toxoplasma gondii, immunosuppressed man, brain abscesses, sulfadiazine, pyrimethamine, and calcium leukovorin, case report

Pyrimethamine

McManus, E. C.; et al., 1979, Exper. Parasitol., v. 47 (1), 13-23

Eimeria tenella, chickens (exper.), t-butylaminoethanol alone or in synergistic combination with sulfaquinoxaline and pyrimethamine, anticoccidial efficacy, specific reversal of toxicity for parasite and host by choline and dimethylaminoethanol

Pyrimethamine + Sulfadimidine (=Abi-Zet₅₀)

Manuel, M. F.; and de Leon, M. L., 1978, Philippine J. Vet. Med., v. 17 (1-2), 129-141
Eimeria tenella, White Leghorn chicks (exper.), comparative efficacy of Esb₃ and Abi-Zet₅₀

Pyrimethamine + Sulfamonomethoxine

Manuel, M. F.; Morales, E. G.; and Trovela, E., [1977], Philippine J. Vet. Med., v. 15 (1-2), 1976, 87-95

Leucocytozoon caulleryi, White Leghorn cockerels, superior prophylactic value of sulfamonomethoxine + pyrimethamine administered in feed under field conditions, no detrimental effects on growth rate or blood picture

Pyrimethamine

Marshall, R. J.; and Ojewole, J. A. O., 1978, Toxicol. and Applied Pharm., v. 46 (3), 759-768

quinoline and nonquinoline antimalarial drugs, effect on isolated guinea pig cardiac muscle

Pyrimethamine

Meuwissen, J. H. E. T.; Golenser, J.; and Verhave, J. P., 1978, Israel J. Med. Sc., v. 14 (5), 601-605

Plasmodium berghei, rats under prophylactic treatment with various drug regimens, development of effective antiparasite immunity by natural bites of infected mosquitoes, symposium presentation

Pancoxin plus

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259

Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Pyrimethamine

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351

Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Pyrimethamine -- Continued.

Pyrimethamine + Sulfadiazine

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351

Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Pyrimethamine

Neves, J.; and Moura, H. B., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (2), 126-130

Plasmodium falciparum, P. vivax, prevalence survey in hospital patients, discussion of changes in prevalence with introduction of chloroquine resistant strains of P. falciparum, treatment trials with various malarial drugs: Brazil

Pyrimethamine

Ohshima, S.; Hoshino, M.; and Tanaka, H., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (3), 127-131

Toxoplasma, 11 strains, susceptibility to 6 drugs, mice

Pyrimethamine + Sulfadiazine

de Paillerets, F.; et al., 1975, Semaine Hop. Paris, v. 51 (43), 2589-2592 (Ann. Pediat., Paris, v. 22 (11), 801-804)

toxoplasmosis, infant, congenital infection, pyrimethamine-sulfadiazine therapy supplemented with folic acid

Pyrimethamine + Sulfadoxine (= Fansidar)

Ponnampalam, J. T., 1978, J. Trop. Med. and Hyg., v. 81 (10), 198-203

human malarias, comparative study of prophylaxis using chloroquine and a combination of sulfadoxine and pyrimethamine: residents of rubber estate in central Malaysia

Pyrimethamine

du Preez, O.; and Cockcroft, C. R., 1975, South African Med. J., v. 49 (34), 1407-1408

malaria prophylaxis trials, army personnel camping in endemic area, 3 drug combinations, no infections reported in trial groups while local population acting as control reported 250 falciparum cases: Caprivi Strip, South Africa

Pyrimethamine + Chloroquine

du Preez, O.; and Cockcroft, C. R., 1975, South African Med. J., v. 49 (34), 1407-1408

malaria prophylaxis trials, army personnel camping in endemic area, 3 drug combinations, no infections reported in trial groups while local population acting as control reported 250 falciparum cases: Caprivi Strip, South Africa

Pyrimethamine + Dapsone

du Preez, O.; and Cockcroft, C. R., 1975, South African Med. J., v. 49 (34), 1407-1408

malaria prophylaxis trials, army personnel camping in endemic area, 3 drug combinations, no infections reported in trial groups while local population acting as control reported 250 falciparum cases: Caprivi Strip, South Africa

Pyrimethamine -- Continued.

Daraprim (Pyrimethamine)

Quintero Monasterios, R.; and Castro R., G., 1972, Rev. Obst. y Ginec. Venez., v. 32 (3), 483-488
human urogenital toxoplasmosis, statistics of cases studied because of secondary sterility, clinical management, therapy with daraprim and supronal

Pyrimethamine (WR 2,978)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Pyrimethamine isethionate

Richards, W. H. G.; and Maples, B. K., 1979, Ann. Trop. Med. and Parasitol., v. 73 (2), 99-108
Plasmodium falciparum in continuous culture, effects of pyrimethamine and chloroquine on parasite growth and viability

Pyrimethamine

Rosario, V. E.; et al., 1978, Lancet, London (8057), v. 1, 185-187
Plasmodium chabaudi, infection of mice with mixtures of drug-resistant (pyrimethamine or chloroquine) and drug sensitive strains, resulting infections were maintained in absence of drugs with some persistence of resistant forms over sensitive forms

Pyrimethamine + Sulphadoxine (= Fansidar)

Rumans, L. W.; Dennis, D. T.; and Atmosoedjono, S., 1979, Lancet, London (8142), v. 2, 580-581 [Letter]
Plasmodium falciparum, fansidar-resistant malaria in case also resistant to chloroquine: Indonesia

Pyrimethamine

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 703-717
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, responses of established infections to chloroquine, quinine, and pyrimethamine

Pyrimethamine

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

Pyrimethamine

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 808-818
Plasmodium falciparum, P. vivax, various drug-resistant and drug-susceptible strains in Aotus trivirgatus griseimembra, capacity of sulfadiazine to enhance activities of WR-158,122 and WR-159,412

Pyrimethamine -- Continued.

Pancoxin plus

Sherkov, S. N.; Leitch, B.; and Kokash, L., [1977], Egypt. J. Vet. Sc., v. 13 (1), 1976, 37-43
Sarcocystis tenella, kittens (exper.), development in intestines, life cycle; attempted parasite suppression using statyl and pancoxin plus

Pancoxin plus

Stoianov, P.; et al., 1978, Vet. Med. Nauki, v. 15 (8), 105-114
E[imeria] tenella, broiler chickens, varied temperature and moisture regimes, blood biochemistry, host resistance, efficacy of pancoxin plus

Pyrimethamine + Sulfamethoxy-pyridazine

Thiermann, E.; et al., 1977, Rev. Med. Chile, v. 105 (7), 433-435
Toxoplasma gondii, mice, experimental trials comparing efficacy of clindamycine with that of pyrimethamine combined with sulfamethoxy-pyridazine, combination drug cured 100% of mice while mice treated with clindamycine survived during treatment but 50% died from severe infections after therapy had been discontinued

Pyrimethamine + Sulfamethoxy-pyridazine

Thiermann, E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 747-750
Toxoplasma gondii, mice, 5 treatment regimens compared during acute and late infections; pyrimethamine + sulfamethoxy-pyridazine was most effective

Daraprim

Ugarte, G.; Apt, W.; and Faiguenbaum, J., 1977, Rev. Med. Chile, v. 105 (3), 176-178
Plasmodium vivax infection in man thought to have hepatic amoebiasis because of complaints of jaundice, fever, and hepatomegaly, after blood smears revealed evidence of malaria man was cured with chloroquine and daraprim: Chile, had made recent visit to Brazil

Pyrimethamine (Tindurin)

Varnai, F.; and Ecker, A., 1977, Therap. Hungar., v. 25 (4), 131-133
malaria, humans travelling to endemic areas, drug prophylaxis, comparative study, least unwanted side effects and lowest morbidity rate recorded with pyrimethamine: Hungary

Pyrimethamine

Vincke, I. H., 1970, Ann. Soc. Belges Med. Trop. Parasitol., v. 50 (3), 339-358
Plasmodium berghei berghei, mice, action of pyrimethamine and sulphomethoxine on pre-erythrocytic and sporogonous cycles

Pyrimethamine + Amprolium + Ethopabate + Sulfamethoxaline (= Pancoxin Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Pyrimethamine -- Continued.

Pyrimethamine

Werner, H.; et al., 1977, Tropenmed. u. Parasitol., v. 28 (4), 528-532

Toxoplasma gondii, latent infected mice, substantial reduction in brain cysts obtained by administration of hyperimmune serum, pyrimethamine, and SDDS in various combinations; effectiveness of therapy varied with parasite strain

Pyrimethamine

Williams, R. L.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (2, pt. 1), 226-231

Plasmodium falciparum, humans with chloroquine-resistant infection (exper.), acetylase phenotype does not influence therapeutic response to sulfalene or sulfalene combined with pyrimethamine

Pyrimethamine+Sulfamonomethoxine

Yoshida, Y.; et al., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (6), 367-375

Pneumocystis carinii pneumonia, rats, comparative efficacy of pyrimethamine+sulfamonomethoxine vs. trimethoprim+sulfamethoxazole

Pyrimethamine isethionate. See Pyrimethamine.

Pyrimido[5,4-c]quinolines and derivatives

Nasr, M.; Nabih, I.; and Burckhalter, J. H., 1978, J. Med. Chem., v. 21 (3), 295-298

Plasmodium gallinaceum, Plasmodium berghei, pyrimido[5,4-c]quinolines and derivatives, laboratory trials, inactive as antimalarials

4-[p-(2-Pyrimidylsulfamyl)phenylazo]-1-naphthylamine

Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124

preparation and testing of 6 long-acting schistosomicidal resins

Pyrimithate -- Diethyl; Elimix.

Elimix (Diethyl)

Grillo Torrado, J. M.; and Perez Arrieta, A., 1977, Rev. Med. Vet., Buenos Aires, v. 58 (2), 101-102, 105

Boophilus microplus, new strain 22, strain G, phosphorus-resistance to various acaricides compared; acetylcholinesterase activity of strain 22 was markedly less than that of strain G

Pyriethamine

Matsuzawa, T., 1978, Parasitology, v. 77 (2), 235-241

Eimeria tenella, chickens, beclotiamine, mode of action studies; attempts to potentiate or antagonize its activity revealed that pyriethamine and 2,4-dinitrophenol also showed slight anticoccidial activity and that a combination of 2,4-DNP and beclotiamine was effective but weight gain was not as good as with beclotiamine alone

Pyriethidium bromide. See Prothidium.

Pyrodis. See Amicarbalide.

n-Pyrrolidinomethyl tetracycline. See Roli-tetracycline.

Pyrvinium -- Poquil; Povan; Pyrvinium pamoate; Vankin.

Pyrvinium pamoate

Baranski, M. C.; et al., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (6), 422-427

Enterobius vermicularis, humans, pyrantel pamoate compared with pyrvinium pamoate: Brazil

Pyrvinium pamoate

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15

antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Pyrvinium pamoate

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, Rev. Brasil. Med., v. 31 (11), 791-794

human soil transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Pyrvinium pamoate (Poquil)

Hayashi, S.; et al., 1976, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 25 (2), 99-108

Enterobius vermicularis, children; Syphacia obvelata, mice, tablets F and S of pyrvinium pamoate compared

Pyrvinium pamoate (Povan)

Loebenberg, D.; et al., 1979, J. Parasitol., v. 65 (5), 823-824

Syphacia obvelata, mice, Sch 23154 compared with pyrantel pamoate and pyrvinium pamoate

Pyrvinium pamoate

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426

Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Vankin

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Pyrvinium pamoate. See Pyrvinium.

Quassia extract

Jensen, O.; Bjerregaard, P.; and Nielsen, A. O., 1979, Ugeskr. Laeger, v. 141 (4), 225-226

head lice, humans, quassia extract vs. chlorphenotane: Denmark

Queletex. See Fenthion.

Quemicetina. See Chloramphenicol.

Quercetin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Quick lime. See Calcium oxide.

Quimofos. See Dioxathion.

Quinacrine -- Atabrine; Atebrin; Mepacrine;

Mepacrine hydrochloride; Quinacrine hydrochloride.

Mepacrine

Ansdell, V. E.; and Common, J. D. A., 1979, J. Trop. Med. and Hyg., v. 82 (9-10), 206-207

Giardia lamblia, 21-year-old Kenyan Asian, corneal damage after therapy with mepacrine: London

Mepacrine + Bithionol

Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36

cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results

Quinacrine

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15

antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Quinacrine hydrochloride

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Mepacrine

Dutta, G. P.; and Narain, L., 1978, Indian J. Exper. Biol., v. 16 (7), 838-840

Entamoeba histolytica, influence of pH on amoebicidal activity of 6 systemically active amoebicides against axenically grown parasites, results indicate that acidic pus in amoebic liver abscesses may account for some therapeutic failures

Quinacrine -- Continued.**Quinacrine hydrochloride (Atabrine)**

Grant, D.; and Woo, P. T. K., 1978, Canad. J. Zool., v. 56 (6), 1360-1366

Giardia spp. in small mammals, comparative studies, results suggest host specificity of some spp., infectivity of stored cysts varies with temperature, lack of prophylactic effect in rats treated with metronidazole or quinacrine hydrochloride

Quinacrine (Atabrine)

Gutierrez, Y.; Buchino, J. J.; and Schubert, W. K., 1978, J. Pediat., St. Louis, v. 93 (2), 245-247

Mesocestoides sp. infection in 12-year-old girl (stools), case report, quinacrine therapy: United States

Quinacrine hydrochloride (Atabrine)

Hartong, W. A.; Gourley, W. K.; and Arvanitakis, C., 1979, Gastroenterology, v. 77 (1), 61-69

Giardia lamblia, patients, clinical spectrum and functional-structural abnormalities of small intestinal mucosa, treatment with metronidazole or quinacrine: Kansas Univ. Medical Center

Quinacrine

Jaroonvesama, N.; and Harinasuta, T., 1972, Siriraj Hosp. Gaz., v. 24 (7), 1095-1099

taeniasis, human, comparative treatment trials using quinacrine and niclosamide

Quinacrine

Jones, R. L.; Davidson, M. W.; and Wilson, W. D., 1979, Biochim. et Biophys. Acta, v. 561 (1), 77-84

chloroquine does not bind to DNA by classical intercalation mechanism typical of quinacrine and ethidium

Quinacrine (Atabrine)

Kavousi, S., 1979, Am. J. Trop. Med. and Hyg., v. 28 (1), 19-23

giardiasis, infants and children, short and long-term followup after treatment with quinacrine vs. metronidazole

Mepacrine hydrochloride (Quinacrine; Atabrine)

Keystone, J. S.; Krajden, S.; and Warren, M. R., 1978, Canad. Med. Ass. J., v. 119 (3), 241-248

Giardia lamblia, epidemiology of outbreak in day-care nurseries, transmission apparently person-to-person, more Canadian children were symptomatic and infected than were immigrant children attending the nurseries, infections cleared with metronidazole or atabrine, control measures suggested including treatment of all infected children regardless of whether they were symptomatic: Toronto, Canada

Mepacrine

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511

Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Quinacrine -- Continued.

Quinacrine hydrochloride

Thacker, S. B.; et al., 1979, Am. J. Pub. Health, v. 69 (12), 1279-1281

Entamoeba histolytica, Giardia lamblia, control attempts in a residential facility for mentally retarded persons: Washington, D. C.

Quinacrine hydrochloride. See Quinacrine.

Quinapyramine -- Antrycide; Antrycide prosalt; Quinapyramine chloride; Quinapyramine dimethylsulphate; Quinapyramine prophylactic; Quinapyramine prosalt; Quinapyramine salt; Quinapyramine sulphate; Antrycide dimethylsulphate.

Quinapyramine prophylactic (Quinapyramine sulphate + Quinapyramine chloride; Quinapyramine prosalt)

Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44

Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested

Antrycide dimethylsulphate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Quinapyramine dimethylsulphate (Antrycide)

James, D. M., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 471-476

T[rypanosoma] congolense, T. brucei, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs

Quinapyramine chloride + Quinapyramine sulphate (= Antrycide prosalt)

Razzaque, A.; and Mishra, S. S., 1977, Bull. Animal Health and Prod. Africa, v. 25 (4), 409-414

Trypanosoma evansi, buffalo calves (exper.), berenil, acriflavin, and antrycide prosalt, berenil most effective

Quinapyramine sulphate + Quinapyramine chloride (= Antrycide prosalt)

Razzaque, A.; and Mishra, S. S., 1977, Bull. Animal Health and Prod. Africa, v. 25 (4), 409-414

Trypanosoma evansi, buffalo calves (exper.), berenil, acriflavin, and antrycide prosalt, berenil most effective

Antrycide prosalt

Thakur, D. K.; and Sinha, K. P., 1978, Indian Vet. J., v. 55 (3), 237-238

Trypanosoma evansi, bovines, 3 lines of treatment tested, best results with antrycide prosalt along with tranquilizer and antihistamine

Quinapyramine -- Continued.

Antrycide

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Quinapyramine salt (Antrycide)

Worms, M. J.; and Hawking, F., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (5), 548-549

Litomosoides carinii-infected Sigmodon hispidus, effects of suramin, homidium bromide, quinapyramine, diminazene, and isometamidium after observation period of more than 5 weeks

Quinapyramine chloride. See Quinapyramine.

Quinapyramine dimethylsulphate. See Quinapyramine.

Quinapyramine prophylactic. See Quinapyramine.

Quinapyramine prosalt. See Quinapyramine.

Quinapyramine salt. See Quinapyramine.

Quinapyramine sulphate. See Quinapyramine.

Quinidine

Katz, N.; and Pellegrino, J., 1974, Rev. Inst. Med. Trop. S. Paulo, v. 16 (5), 245-252

Schistosoma mansoni, Cebus monkeys, correlation of number of eggs per gram of rectal tissue with number of female worms, challenge infection effect, or drug action

Quinidine monohydrate

Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonocides using Coulter Counter

Quinidine sulfate

Marshall, R. J.; and Ojewole, J. A. O., 1978, Toxicol. and Applied Pharm., v. 46 (3), 759-768

quinoline and nonquinoline antimalarial drugs, effects on isolated guinea pig cardiac muscle

Quinimax. See Quinine.

Quinine -- Quinimax; Quinine dihydrochloride;
Quinine sulfate; WR 2,976.

Quinine sulphate
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter

Quinine sulfate
Charoenlarp, P.; Vanijanonta, S.; and Chat-
panyaporn, P., 1979, *Southeast Asian J. Trop.
Med. and Pub. Health*, v. 10 (1), 127-131
falciparum malaria, humans, prednisolone
administered with quinine sulfate did not
increase red cell survival

Quinine
Chongsuphajaisiddhi, T.; et al., 1979, *South-
east Asian J. Trop. Med. and Pub. Health*, v. 10
(1), 132-137
falciparum malaria, children, chloroquine re-
sistance, efficacy of quinine and fansidar,
clinical study: Thailand

Quinine
Elslager, E. F.; et al., 1979, *J. Med. Chem.*,
v. 22 (10), 1247-1257
Plasmodium spp., antimalarial activity of
2,4-diamino-6-(2-naphthylsulfonyl)quinazoline
and related 2,4-diamino-6-[(phenyl and naph-
thyl)sulfinyl and sulfonyl]quinazolines

Quinine
Geddes, A. M., 1976, *J. Antimicrob. Chemother.*,
v. 2 (1), 106 [Letter]
chloroquine-resistant malaria, intravenous
infusion of quinine for treatment of severe
life-threatening infections, given by mouth
for less seriously ill patients [correction
of omission in Geddes, A. M., 1975, *J. Anti-
microb. Chemother.*, v. 1 (4), 349-350]

Quinine
Glew, R. H.; Collins, W. E.; and Miller, L. H.,
1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (1, pt.
1), 9-13
Plasmodium falciparum, selection of increased
quinine resistance in Aotus monkeys

Quinine sulfate
Marshall, R. J.; and Ojewole, J. A. O., 1978,
Toxicol. and Applied Pharm., v. 46 (3), 759-
768
quinoline and nonquinoline antimalarial
drugs, effects on isolated guinea pig car-
diac muscle

Quinine
Moretti, G.; et al., 1974, *Semaine Hop. Paris*,
v. 50 (15), 989-997
Plasmodium falciparum, woman, fatal infec-
tion accompanied by jaundice and renal
failure, unresponsive to quinine therapy,
review of clinical aspects: Cayenne (Guyane)

Quinine
Neves, J.; and Moura, H. B., 1972, *Rev. Inst.
Med. Trop. S. Paulo*, v. 14 (2), 126-130
Plasmodium falciparum, P. vivax, prevalence
survey in hospital patients, discussion of
changes in prevalence with introduction of
chloroquine resistant strains of P. falci-
parum, treatment trials with various malarial
drugs: Brazil

Quinine -- Continued.

Quinine dihydrochloride
Patrick, I. T., 1979, *Brit. Med. J.* (6163),
v. 1, 619-620 [Letter]
human cerebral malaria, successful treatment
regimen using intravenous quinine dihydro-
chloride

Quinine (and quinine isomers)
Pellegrino, J.; and Katz, N., 1974, *Rev. Inst.
Med. Trop. S. Paulo*, v. 16 (5), 301-304
Schistosoma mansoni, mice, hamsters, blind
screening trials, quinine and its isomers
(quinine, quinine valerate, quinine gluco-
nate, quinine ethyl carbonate, quinine phos-
phate, quinidine, quinidine sulfate, D-cin-
chonine, cinchonine hydrochloride and cincho-
nidine sulfate)

Quinine
Pillay, N.; and Bhoola, R. L., 1975, *South
African Med. J.*, v. 49 (35), 1443-1444
Plasmodium falciparum, woman, case report,
probable resistance to chloroquine, success-
fully treated with quinine: South Africa
(had recently returned from Mozambique)

Quinine (WR 2,976)
Rane, D. S.; and Kinnamon, K. E., 1979, *Am. J.
Trop. Med. and Hyg.*, v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in
mice, development of high volume tissue
schizonticidal drug screen based upon mor-
tality of infected mice

Quinine
Roncoroni, A. J.; and Martino, O. A., 1979,
Am. J. Trop. Med. and Hyg., v. 28 (3), 440-444
Plasmodium falciparum, 36-year-old male
after travel to Senegal, case report, severe
infection cured by exchange blood trans-
fusion in conjunction with classical drug
therapy: Argentina

Quinine
Schmidt, L. H., 1978, *Am. J. Trop. Med. and
Hyg.*, v. 27 (4), 703-717
Plasmodium falciparum and P. vivax in Aotus
trivirgatus griseimembra, responses of
established infections to chloroquine,
quinine, and pyrimethamine

Quinine
Schmidt, L. H., 1979, *Am. J. Trop. Med. and
Hyg.*, v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus
trivirgatus griseimembra, strains resistant
to chloroquine, quinine, or pyrimethamine,
antimalarial properties of selected 2,4-
diamino-6-substituted quinazolines

Quinine sulfate
Sucharit, P.; et al., 1979, *Southeast Asian J.
Trop. Med. and Pub. Health*, v. 10 (1), 138-141
Plasmodium falciparum still sensitive to
quinine, in vitro and in vivo (humans):
Thailand

Quinine
Thong, Y. H.; and Ferrante, A., 1978, *Am. J.
Trop. Med. and Hyg.*, v. 27 (2, pt. 1), 354-356
quinine, inhibition of mitogen-induced human
lymphocyte proliferative responses, this
immunosuppressive property may be undesirable
side effect in treatment of malaria

Quinine -- Continued.

Quinine (Quinimax)

Van Poucke, G., 1979, East African Med. J., v. 56 (4), 158-162

Plasmodium falciparum, humans, intramuscular treatment with chloroquine vs. quinimax: East Africa

Quinine dihydrochloride. See Quinine.

Quinine sulfate. See Quinine.

Quinocide

Levchenko, F. F., 1978, Veterinariia, Moskva (7), 62-64

Theileria annulata, cattle, quinocide and bigumal treatment effective, treatment with hemostimulin and vitamins and microelements given in feed to counteract anemia and atonicity of digestive tract resulting from therapy: Gissarsk valley

4-Quinolinemethanols

Schmidt, L. H.; et al., 1978, Antimicrob.

Agents and Chemotherapy, v. 13 (6), 1011-1030

Plasmodium falciparum in *Aotus trivirgatus griseimembra*, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

5-(8-Quinolinoxymethyl)-4-substituted-3-mercaptop-1,2,4(4H)-triazoles

Soliman, R.; and Hammouda, N. A., 1979, J. Pharm. Sc., v. 68 (11), 1377-1381

Schistosoma mansoni, *Toxocara canis*, mice (exper.), synthesis and activity of new mercaptotriazoles

Quinorium sulphate. See 1,3-Di-6-quinolyurea.

N'-(2-Quinoxaliny) sulfanilamide. See Sulfaquinoxaline.

Quintiofos -- Bacdip; Bayer 9037; 0-Ethyl 0-(8-hydroxy-quinoline)-phenyl phosphorothioate; 0-Ethyl-0-(8-hydroxyquinolyl)-phenylthionophosphate; 0-Ethyl-0-(8-quinolyl)-benzenethionophosphonate; Oxinothiofos; Oxinothiofos.

Oxinothiophos (Bacdip; Bayer 9037)

Abdel Rahman, M. S.; El Gendi, A. Y. I.; and Moursi, H. S. A., 1977, J. Egypt. Vet. Med. Ass., v. 37 (2), 55-69

nematodes and cestodes, dogs, effectiveness of trichlorfon, oxinothiophos, and carbaryl compared: Giza and Cairo Governorates, Egypt

Quintiofos -- Continued.

Oxinothiophos

Bonin, W., 1977, Berl. u. Munchen. Tierarztl.

Wchnschr., v. 90 (2), 34-37

Amblyomma hebraeum, *Psoroptes cuniculi*, *Melophagus ovinus*, *Dermanyssus gallinae*, *heptenophos*, rapid mode of action, broad range of efficacy, short residual effect and effective as a vapour poison, compared with other standard drug preparations

Bacdip (Oxinothiofos)

Chemtai, A. A. K., 1977, Bull. Animal Health

and Prod. Africa, v. 25 (3), 299-306

acaricide-treated zebu cattle, blood cholinesterase, radiometric assay

Bacdip

El-Bahay, G. H.; et al., 1978, Vet. Med. J.,

Giza, v. 24 (24), 1976, 205-213

Hyalomma dromedarii, *Argas persicus*, evaluation of 10 insecticides

Oxinothiophos (Bacdip)

Hammant, C. A.; and Matthewson, M. D., 1977,

Rhodesian Vet. J., v. 8 (4), 71-73

Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Quintiofos (Bacdip; Bayer 9037)

Lourens, J. H. M.; and Lyaru, D. M., 1979,

PANS, v. 25 (2), 135-142

Rhipicephalus appendiculatus, susceptibility of organochlorine susceptible and resistant East African strains to ten cholinesterase inhibiting acaricides

Oxionthiophos (Bacdip)

Rechav, Y.; Whitehead, G. B.; and Terry,

S. B., 1978, J. South African Vet. Ass., v. 49 (2), 99-101

ticks, mortality curves of larvae dipped in dioxathion, chlorphenvinphos, and oxionthiophos, time of application, larvae of ticks exhibit diel periodicity in sensitivity to acaricides

Quinuronium sulphate. See 1,3-Di-6-quinolyurea.

Quixalin. See Halquinol.

Quixoline

Reddy, R. G., 1979, Livestock Advis., v. 4 (3), 37-40

Balantidium coli and mixed infection with amphistomes, cattle and buffaloes, incidence and treatment

Rabon. See Tetrachlorvinphos.

Radanil. See Benzimidazole.

Radeverm. See Niclosamide.

Rafoxanide -- 3,5-Diiodo-3'-chloro-4'-(p-chloro-phenoxy)-salicylanilide; Flukanide; Ranide; Ursovermit.

Rafoxanide (Flukanide)

Altaif, K. I., 1979, Trop. Animal Health and Prod., v. 11 (4), 241-245
helminths, Awassi sheep, tetramisole, rafoxanide, body weight gains, wool growth: Iraq

Rafoxanide

Bourn, D.; and Scott, M., 1978, Trop. Animal Health and Prod., v. 10 (4), 191-203
trypanosomiasis, successful use of Zebu work oxen in agricultural development of tsetse infested land, environmental conditions, epizootiology of trypanosomiasis in oxen and in *Glossina morsitans*, strategic drug use (alternation of diminazene aceturate and isometamidium to control trypanosomes; rafoxanide to control helminths): Wollega province, western Ethiopia

Rafoxanide (Ranide)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
Haemonchus contortus, *Trichostrongylus colubriformis*, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Rafoxanide

Collado-Torres, M. L.; and de Leon, D., 1979, J. Agric. Univ. Puerto Rico, v. 63 (1), 78-79
Platynosomum fastosum, cats, rafoxanide ineffective: Puerto Rico

Rafoxanide (Flukanid, Ranid)

Corba, J.; et al., 1978, Veterinarstvi, v. 28 (2), 87-88
Fasciola hepatica, heifers, brotianid, rafoxanide

Rafoxanide (Ursovermit)

Dedek, W.; et al., 1978, Arch. Exper. Vet.-Med., v. 32 (6), 951-955
rafoxanide, metabolism, residues and excretion in blood, milk, meat, and urine of lactating cows

Rafoxanide

Foreyt, W. J.; and Todd, A. C., 1974, Am. J. Vet. Research, v. 35 (3), 375-377
Fascioloides magna, cattle, efficacies of rafoxanide and oxclozanide: Texas Gulf Coast Region

Ranide

Georgieva, D., 1978, Vet. Sbirka, v. 76 (4), 32-33
anthelmintic treatment, lambs, weight gains

Rafoxanide

Hillyer, G. V.; and Santiago de Weil, N., 1979, J. Parasitol., v. 65 (5), 680-684
Fasciola hepatica, rats, rabbits, enzyme linked immunosorbent assay can be used for serodiagnosis and for prediction of chemotherapeutic success

Rafoxanide -- Continued.

Rafoxanide (Ranide)

Johnstone, I. L.; et al., 1979, Austral. J. Exper. Agric. and Animal Husb. (98), v. 19, 303-311
parasites, sheep, effect of four control schemes in two environments on wool production and liveweight gains

Rafoxanide (Ranide)

Johnstone, I. L.; Coote, B. G.; and Smart, K. E., 1979, Austral. J. Exper. Agric. and Animal Husb. (99), v. 19, 414-418
pre- and/or post-lambing anthelmintic treatment, effect on lamb birth weight and liveweight gain, fecal egg counts, pasture contamination: New South Wales

Rafoxanide

Lamberg, K. J., 1978, Norsk Vet.-Tidsskr., v. 90 (4), 223-227
Fasciola hepatica, sheep, outbreak described, possible snail hosts, rafoxanide treatment: Rogaland

Rafoxanide (Ranide)

Le Jambre, L. F.; and Barger, I. A., 1979, Austral. Vet. J., v. 55 (7), 346-347 [Letter]
benzimidazole resistant *Haemonchus contortus*, lambs, rafoxanide and naphthalophos, efficiency against adults and inhibited larvae: New South Wales

Ranide

Petrov, D.; Milushev, I.; and Monov, M., 1978, Vet. Med. Nauki, v. 15 (4), 33-39
Oestrus ovis, sheep, neguvon, ranide, and downix compared; use of ranide economically justified only for mixed infection with *Fasciola hepatica*

Rafoxanide

Prichard, R. K., 1978, Parasitology, v. 76 (3), 277-288
Fasciola hepatica, metabolic profile of adult flukes obtained from rafoxanide-treated sheep, concluded that mode of action of rafoxanide in vivo is by uncoupling oxidative phosphorylation

Rafoxanide (Flukanide; Ranide)

Razafindrakoto, C.; Ranaivoson, A.; and Megard, J. P., 1978, Rev. Elevage et Med. Vet. Pays Trop., n. s., v. 31 (2), 165-169
Fasciola gigantica, Malagasy zebu cattle, rafoxanide injectable, recommended for curative and prophylactic treatment

Rafoxanide

Reinhardt, P., 1978, Monatsh. Vet.-Med., v. 33 (23), 898-901
Dicrocoelium dendriticum, sheep, efficacy of various anthelmintics compared

Rafoxanide (Ranide)

Schillhorn Van Veen, T. W., 1978, Vet. Rec., v. 102 (16), 364-365
Haemonchus contortus, lambs (abomasum, faeces), casualties of young lambs following prolonged rainy season, further casualties at end of following dry season associated with inhibited *H. contortus* larvae suggest chronic haemonchosis syndrome (lambs had been previously treated with thiabendazole and rafoxanide); high pasture infection: Shika, near Zaria

Rafoxanide -- Continued.Rafoxanide (Ranide)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiazobenzazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxylnil are also useful chemical alternatives

Ragadan. See Heptenophos.

Rametin. See Phthalophos.

Rametin H. See Phthalophos.

Ranide. See Rafoxanide.

Razoxane -- ICRF 159.

ICRF 159

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Red clover blossoms

Slepnev, N. K., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 107-109
Ascaris suum, in vitro testing of anthelmintic activity of water extracts of some plants

Red Spot. See Carbon tetrachloride.

Repellents

Bar-Zeev, M.; and Gothilf, S., 1974, J. Med. Entom., v. 11 (4), 389-392
Ornithodoros tholozani, rats (exper.), laboratory and field evaluation of 9 repellents, pyrethrum far superior, o-vanillin and ephedrine more effective than other remaining repellents but toxic to host

Repellents

Getta, G. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 72-75
 hypodermatosis, cattle, insecticides, repellents

Repellents

Kulkarni, S. M., 1977, J. Med. Entom., v. 14 (1), 64-70
Leptotrombidium deliense, *L. akamushi*, laboratory evaluation of repellents dibutylphthalate, dimethylphthalate, N,N-diethylm-toluamide (deet), deet + dimethylphthalate, pyrethrum impregnated on filter paper and nylon/cotton fabric

Repellents

Sixl, W.; and Stuenzner, D., 1975, Ang. Parasitol., v. 16 (2), 106-108
 autan spray, effective repellent against ticks, field and laboratory trials

Repellents

Smirnova, S. N.; and Dremova, V. P., 1971, Parazitologia, Leningrad, v. 5 (4), 357-360
Rhipicephalus turanicus, *Hyalomma* spp., sensitivity to various repellents

Resochin. See Chloroquine.

Resorantel -- 4'-Bromo-2,6-dihydroxybenzanilide; Terenol.

Resorantel

Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
 cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Resorantel

Douch, P. G. C., 1979, Xenobiotica, v. 9 (4), 263-268
Moniezia expansa, *Ascaris suum*, metabolism of clixoxanide and resorantel and related compounds

Resotren [composite]. See Chloquinatone or Chloroquine or Diiodohydroxyquin.

Resulfon. See Sulfaguanidine.

Reverin. See Rolitetracline.

Ribavirin -- Virazole.

Ribavirin

McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
Theileria parva and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Virazole

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

3- β -D-Ribofuranosyl-6,7,8-trihydroimidazo[3,4-d][1,3]diazepin-8-(R)-ol. See Coformycin.

Ricifon. See Trichlorfon.

Ridlice. See Chlorpyrifos.

Rifaldin. See Rifampin.

Rifampicin. See Rifampin.

Rifampin -- Rifaldin; Rifampicin.

Rifampicin (Rifaldin)
 Dourado, H. V.; et al., 1975, Rev. Brasil. Clin. e Terap., v. 4 (1), 1-6
 human tegumentary leishmaniasis, therapeutic effectiveness of rifampicin in 55 patients presenting with cutaneous lesions: State of Amazonas, Brazil

Rifampicin
 Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
 Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Rifampicin
 Sinden, R. E.; and Smalley, M. E., 1979, Parasitology, v. 79 (2), 277-296
 Plasmodium falciparum, modified microculture technique used as bioassay for various anti-metabolites by examining their ability to inhibit gametocytogenesis; characterization of sexual cell-cycle

Rigecoccin. See Metichlorpindol.

Rintal. See Febantel.

Rintal paste. See Febantel.

Ripercol. See Tetramisole.

Ripercol-L. See Tetramisole.

Robenidine -- 1,3-Bis (p-chlorobenzylideneamino guanidine hydrochloride; Bis[(4-chlorophenyl)methylene]-carbonimidic dihydrazide; Cycostat; Robenz.

Robenidine
 Greuel, E.; and Kuehnhold, W., 1977, Prakt. Tierarzt, v. 58 (5), 338-341
 Eimeria spp., chickens (exper.), synergistic effect of metichlorpindol and methylbenzoate, rotation program with other coccidiostats discussed

Robenidine -- Continued.

Robenidine
 Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
 Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis

Robenidine
 Latter, V. S.; and Wilson, R. G., 1979, Parasitology, v. 79 (1), 169-175
 Eimeria tenella, factors influencing assessment of anticoccidial activity in cell culture

Robenidine
 Lee, E. H., 1979, Canad. Vet. J., v. 20 (4), 102-104
 Eimeria tenella, drug-resistant field strains, White Leghorn chickens, single and low-level oocyst infections, treatment with robenidine or decoquinate

Robenidine
 Long, P. L.; and Millard, B. J., 1978, Parasitology, v. 76 (1), 1-9
 Eimeria grenieri in Numida meleagris (intestine, caeca) (nat. and exper.), life cycle, reproduction rate, pathogenicity (severe depression of body weight gain), immunity to reinfection, treatment with sulphaquinoxaline in drinking water and robenidine in food: Britain

Robenidine
 Long, P. L.; Millard, B. J.; and Smith, K. M., 1979, Avian Path., v. 8 (4), 453-467
 Eimeria spp., chickens, effect of 4 anti-coccidial drugs on development of immunity, field and laboratory conditions

Robenidine
 McDougald, L. R.; and Galloway, R. B., 1977, Ztschr. Parasitenk., v. 54 (1), 95-100
 Eimeria tenella in vitro, development inhibited by serum from chickens fed anti-coccidial drugs, technique to assay drug activity and to characterize and quantitate therapeutic effect

Robenidine
 McHardy, N., 1978, Ann. Trop. Med. and Parasitol., v. 72 (6), 501-511
 Theileria parva- and T. annulata-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Robenidine
 McLoughlin, D. K.; and Chute, M. B., 1978, J. Parasitol., v. 64 (5), 874-877
 Eimeria tenella, chickens, robenidine protected against cecal coccidiosis initiated by parasite strain with no previous drug exposure, no cross resistance found with 13 strains resistant to other anticoccidials, when the sensitive strain was serially propagated in chickens medicated with robenidine it became resistant, no cross resistance found when this experimental strain was tested against 12 other anticoccidials

Robenidine -- Continued.

Robenidine (Robenz, Cycostat)

McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113

Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs

Cycostat

Manuel, M. F.; and Buduan, R. J., 1972, Philippine J. Vet. Med., v. 11 (2), 73-91
battery-raised broilers, effects of amprol plus or cycostat with or without payzone on weight gain and feed efficiency, no coccidiosis found

Cycostat (Robenidine)

Mladenovic, Z.; Movsesijan, M.; and Borojevic, D., 1978, Vet. Glasnik, v. 32 (10), 829-834
Eimeria spp., chickens (exper.), mixed infections, cycostat, nitryl, and stenorol

Cycostat

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259

Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Robenidine

Olson, G.; et al., 1978, Poultry Science, v. 57 (5), 1245-1250

Eimeria spp. field isolates, chickens (exper.), arprinocid in comparison trials with marketed drugs, effective against all isolates tested including those refractory to many of the other products

Robenidine (Cycostat)

Peeters, J. E.; Halen, P.; and Meulemans, G., 1979, Brit. Vet. J., v. 135 (4), 349-354

Eimeria spp., rabbits (nat. and exper.), efficacy of robenidine

Robenidine

Ryley, J. F.; and Hardman, L., 1978, J. Parasitol., v. 64 (5), 878-881

Eimeria acervulina, *E. mivati*, speciation studies (cross-immunity and drug resistance studies), some immunological relationship was demonstrated but the failure of the 2 organisms to interbreed in the drug resistance studies lends support to status of *E. mivati* as distinct species

Robenidine

Ryley, J. F.; and Hardman, L., 1978, Parasitology, v. 76 (1), 11-20

Eimeria spp., chicks (exper.), effects of dietary vitamin K on severity of disease with particular attention to effects of vitamin K on response to anticoccidial drugs, concluded that use of vitamin K deficient diet for experimental work is quite justified

Robenidine (Cycostat)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289

coccidiosis, broilers, anticoccidials, floor pen trials

Robenz. See Robenidine.

Rogor. See Dimethoate.

Rolitetracycline -- n-Pyrrolidinomethyl tetracycline; Reverin.

Reverin (Rolitetracycline)

Bauer, F.; Raether, W.; and Seeger, K., 1978, Cahiers Bleus Vet. (27), 265-271
protozoal disease in exper. hosts, enhanced effect of berenil + reverin vs. berenil alone

n-Pyrrolidinomethyl tetracycline (Reverin)

Brown, C. G. D.; et al., 1977, Tropenmed. u. Parasitol., v. 28 (4), 513-520

Theileria parva, *Bos taurus* with patent East Coast fever induced by inoculation of cryopreserved stabilates of tick-derived infective particles, preliminary evaluation of n-pyrrolidinomethyl tetracycline and oxytetracycline

Rolitetracycline (Reverin)

Jagdish, S.; et al., 1979, Vet. Rec., v. 104 (7), 140-142

Theileria annulata, immunising infection in calves by injecting ground up infected *Hyalomma anatolicum anatolicum* supernate, severity of reactions in rolitetracycline-treated vs. non-treated calves compared, adequate protection, durable immunity to subsequent severe homologous challenge

Rolitetracycline + Diminazene aceturate

Malhotra, D. V.; Gautam, O. P.; and Banerjee, D. P., 1979, Indian J. Animal Sc., v. 49 (1), 75-77

Babesia equi, donkeys (exper), diminazene aceturate as effective as diminazene aceturate + rolitetracycline, quinuronium sulphate ineffective

Romensin. See Monensin.

Randomycin. See Methacycline.

Ronidazole -- Duodegran; 1-Methyl-2-carbamoyloxy-methyl-5-nitroimidazole.

1-Methyl-2-carbamoyloxymethyl-5-nitroimidazole (Ronidazole)

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Ronidazole (Duodegran)

Hauser, K. W., 1977, Prakt. Tierarzt, v. 58, special no., 56

trichomoniasis, parrots, diagnosis, treatment with ronidazole and spartrix, review

Ronnel -- 0,0-Dimethyl-0-(2,4,5-trichlorophenyl) phosphorothioate; Ectoral; Fenchlorphos; Korlan; Trolene.

Ronnel

Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765

Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control

Ronnel + Chlorpyrifos

Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765

Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control

Ronnel

Crow, S. E., 1978, Mod. Vet. Pract., v. 59 (3), 184-186

Demodex canis in nasal lesion of dog, treated with ronnel

Fenchlorphos (Ectoral)

Dorrestein, G. M.; and Van Bronswijk, J. E. M. H., 1979, Vet. Parasitol., v. 5 (4), 389-398

Trixacarus caviae as cause of mange in Cavia porcellus (nat. and exper.), clinical symptoms, pathology, treatment; papular urticaria in humans associated with mangy guinea-pigs: The Netherlands

Ronnel

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, B. microplus, laboratory tests of insecticides

Ronnel

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886

laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Ronnel

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586

Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Ronnel (Korlan)

Lloyd, J. E.; Olson, E. J.; and Pfadt, R. E., 1978, J. Econom. Entom., v. 71 (3), 548-550

Melophagus ovinus, sheep, diazinon, ronnel, low volume spraying gave rapid control with no adverse effects

Ronnel (Trolene)

Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661

flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico

Ronnel -- Continued.

Ronnel

Rumsey, T. S., 1979, J. Animal Sc., v. 49 (4), 1059-1065

ronnel, positive effect on weight gains of feedlot steers not related to anthelmintic or insecticidal action or to significant changes in ruminal measurements, rumen microbial morphology, or plasma amino acid patterns

Ronnel

Wright, F. C.; and Riner, J. C., 1979, South-west. Entom., v. 4 (1), 40-45

Psoroptes ovis, P. cuniculi, 10 acaricides evaluated using 'tea-bag' technique

Rotenone

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, B. microplus, laboratory tests of insecticides

Rovamycin. See Spiramycin.

Roxarsone -- 3-Nitro-4 hydroxyphenylarsonic acid; Polystat (with Dibutyltin dilaurate, Dinsed, and Sulfantran).

Roxarsone + Bacitracin MD + Halofuginone

Edgar, S. A.; and Flanagan, C., 1979, Poultry Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant to various drugs), halofuginone with roxarsone and/or bacitracin MD

Roxarsone + Halofuginone

Edgar, S. A.; and Flanagan, C., 1979, Poultry Science, v. 58 (6), 1476-1482

Eimeria spp. (recent field isolates resistant to various drugs), halofuginone with roxarsone and/or bacitracin MD

Polystat

McDougald, L. R.; and McQuiston, T. E., 1978, Avian Dis., v. 22 (4), 765-770

Eimeria spp., turkeys, coccidiosis management, innate (age) resistance and acquired immunity vs. anticoccidial medication

3 Nitro-4 hydroxyphenylarsonic acid

Morrison, W. D.; Ferguson, A. E.; and Leeson, S., 1979, Poultry Science, v. 58 (5), 1160-1166

Eimeria spp., chicks (exper.), salinomycin and stenorol compared with other anticoccidials, efficacy and effect on chick performance

3 Nitro-4 hydroxyphenylarsonic acid + Salinomycin

Morrison, W. D.; Ferguson, A. E.; and Leeson, S., 1979, Poultry Science, v. 58 (5), 1160-1166

Eimeria spp., chicks (exper.), salinomycin and stenorol compared with other anticoccidials, efficacy and effect on chick performance

Roxion. See Dimethoate.

Ruelene. See Crufomate.

Ruelene 6-R. See Crufomate.

Rumensin. See Monensin.

Rumevite Wormablok containing Panacur. See Fenbendazole.

Rycovet warblecide pour-on
Love, J., 1979, Vet. Rec., v. 104 (1), 22
[Letter]
warble fly, ostertagiasis, young stock,
simultaneous prophylactic treatment with
rycovet warblecide and panacur

- Safranine**
 Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
 Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective
- Sagimid.** See Niclosamide.
- Salicylhydroxamic acid** -- SHAM
- Salicylhydroxamic acid**
 Brohn, F. H.; and Clarkson, A. B., jr., 1978, Acta Trop., v. 35 (1), 23-33
 Trypanosoma brucei brucei, effect of glycerol on anaerobic glycolysis in vitro, concomitant administration of salicylhydroxamic acid and glycerol to infected rats results in rapid clearance of parasitemia
- Salicylhydroxamic acid (SHAM) + Glycerol**
 Evans, D. A.; and Holland, M. F., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (2), 203-204
 Trypanosoma vivax, mice, effective treatment with salicylhydroxamic acid + glycerol
- Salicylhydroxamic acid (SHAM)**
 Nathan, H. C.; et al., 1979, J. Protozool., v. 26 (4), 657-660
 Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents
- SHAM/Glycerol**
 Nathan, H. C.; et al., 1979, J. Protozool., v. 26 (4), 657-660
 Trypanosoma brucei brucei, mice, effect of amicarbalide, imidocarb, and several other agents
- Salicylhydroxamic acid + Glycerol**
 Van Der Meer, C.; Versluijs-Broers, J. A. M.; and Opperdoes, F. R., 1979, Exper. Parasitol., v. 48 (1), 126-134
 Trypanosoma brucei brucei, rats, treatment with salicylhydroxamic acid + glycerol and suramin + glycerol
-
- Salinomycin**
 Benz, G. W.; and Ernst, J. V., 1979, Am. J. Vet. Research, v. 40 (8), 1180-1186
 Eimeria bovis, calves (exper.), efficacy of salinomycin
- Salinomycin (Coxistac)**
 Chappel, L. R., 1979, J. Parasitol., v. 65 (1), 137-143
 Eimeria spp., chicks, site of action of salinomycin
- Salinomycin (Coxistac)**
 Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (2), 304-307
 Eimeria spp., broilers (exper.), salinomycin, monensin, lasalocid, drug toleration and anticoccidial efficacy compared in 5 field trials, commercial facilities
- Salinomycin**
 Kaemmerer, K.; and Fink, J., 1978, Deutsche Tierarztl. Wchnschr., v. 85 (6), 216-220
 monensin-natrium, lasalocid, salinomycin, influence of high dosages on heart of chickens
- Salinomycin**
 Karlsson, T.; and Reid, W. M., 1978, Avian Dis., v. 22 (3), 487-495
 Eimeria tenella, broiler chicks, effect of anticoccidials in feed on development of immunity to coccidiosis
- Salinomycin (Coxistac)**
 McQuiston, T. E.; and McDougald, L. R., 1979, Ztschr. Parasitenk., v. 59 (2), 107-113
 Eimeria tenella, surgical ligation of chick ceca used to study role of absorption and extraintestinal transport in action of anticoccidial drugs
- Salinomycin (Coxistac)**
 Migaki, T. T.; and Babcock, W. E., 1979, Poultry Science, v. 58 (2), 481-482
 salinomycin, anticoccidial, safety in broiler chickens compared with monensin
- Salinomycin (Coxistac)**
 Migaki, T. T.; Chappel, L. R.; and Babcock, W. E., 1979, Poultry Science, v. 58 (5), 1192-1196
 Eimeria spp., chicks (exper.), salinomycin, monensin, lasalocid, efficacy in battery trials
- Salinomycin**
 Morrison, W. D.; Ferguson, A. E.; and Leeson, S., 1979, Poultry Science, v. 58 (5), 1160-1166
 Eimeria spp., chicks (exper.), salinomycin and stenorol compared with other anticoccidials, efficacy and effect on chick performance
- Salinomycin + 3 Nitro-4 hydroxyphenylarsonic acid**
 Morrison, W. D.; Ferguson, A. E.; and Leeson, S., 1979, Poultry Science, v. 58 (5), 1160-1166
 Eimeria spp., chicks (exper.), salinomycin and stenorol compared with other anticoccidials, efficacy and effect on chick performance
- Salt.** See Sodium chloride.
- Samorin.** See Isometamidium.
- Sanasil.** See Sulfadoxine.
- Sansalid.** See Diuredosan.
- Santonin** -- Ascarel (with Cascara sagrada); Sodium santoninate.
- Sodium santoninate + Cascara sagrada (Ascarel)**
 Oldham, R. R.; et al., 1971, South. Med. J., v. 64 (4), 480-482
 possible santonin poisoning (hemolytic crisis) in young child treated for worms with ascarel
- Santonin**
 Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
 Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility
-

Saquadil. See Diaveridine or Sulfaquinoxaline.

Sarcocystis acridine

Krotov, A. I.; Cherniaeva, A. I.; and Budanova, I. S., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 165-168

Alveococcus multilocularis, white mice, effect of thiabendazole, sarcocystis acridine, levamisole, and mebendazole on larval cyst development

SB-EDTA. See Sodium-antimonyl-ethylene-diamino-tetracetate.

Schistocide T-109. See 2,8-Dimethyl-5- β -diethyl-amino-ethylamino-thiochromone.

Schistosomicides

Abdou, N. A.; et al., 1978, *Egypt. J. Pharm. Sc.*, v. 17 (2), 1976, 153-159

synthesis of some organosulphur compounds structurally related to certain antibilharzial drugs, to be screened for possible activity

Schistosomicides

Coutinho, A., 1969, *Rev. Inst. Med. Trop. S. Paulo*, v. 11 (5), 363-376

Schistosoma mansoni, human, current aspects of chemical and surgical therapy, review

Schistosomicides

Korolkovas, A.; et al., 1975, *Rev. Farm. e Bioquim. Univ. Sao Paulo*, v. 13 (2), 217-225

latentiation of 9 potential naphthylazo derivatives as schistosomicides, although the compounds were ineffective in trials with mice, the latentiation method used in the trials showed potential for wider application in drug testing

Schistosomicides

McMahon, J. E., 1978, *Trop. and Geogr. Med.*, v. 30 (2), 161-174

schistosomiasis, human, factors affecting chemotherapy, review

Schistosomicides

Woolhouse, N. M., 1979, *Biochem. Pharmacol.*, v. 28 (16), 2413-2418

antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Scolaban. See Bunamidine.

SDDS. See 2-Sulfamoyl-4,4'-diaminodiphenyl-sulfone.

Secnidazole -- (Hydroxy-2-propyl)-1-methyl-2-nitro-5-imidazole; 14.539 RP; PM-185184.

(Hydroxy-2-propyl)-1-methyl-2-nitro-5-imidazole (R.P. 14539)

da Cunha, A. S.; et al., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (5), 342-348

Entamoeba histolytica, human intestinal infections, clinical trials with R.P. 14539

Secnidazole (14.539 RP)

Rocha, A. M.; and Sarraff Neto, A., 1977, *Rev. Brasil. Med.*, v. 34 (6), 365-366

trichomoniasis, human urogenital, successful therapy of sexual partners with secnidazole

Selectomycin. See Spiramycin.

Selenourea

Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Selenium disulfide

Ogata, M.; et al., 1978, *Bull. Azabu Vet. Coll.*, v. 3 (2), 291-295

Cheyletiella sp. from cat, description; dermatitis of cat and her owners, selenium disulfide treatment of cat, case report: Tokyo, Japan

Senna -- Helmintia-P (with Phenothiazine, Piperazine, Tin, and Vernonia anthelmintica); *Sennae folia*.

Helmintia-P

Matta, S. C.; and Ahluwalia, S. S., 1979, *Indian Vet. J.*, v. 56 (7), 616-617

helminths, poultry, helmintia-P, sonex

Sennae folia. See Senna.

Septamycin

Hildebrandt, J.; Meingassner, J. G.; and Mieth, H., 1978, *Zentralbl. Vet.-Med., Reihe B.*, v. 25 (3), 186-193

Eimeria tenella, kidney cell cultures, chickens (exper.), septamycin, activity largely confined to first generation schizont

Septran. See Sulfamethoxazole or Trimethoprim.

Septrin. See Sulfamethoxazole or Trimethoprim.

Sevin. See Carbaryl.

SHAM. See Salicylhydroxamic acid.

Silver chloride

Sherkov, Sh.; et al., 1978, *Vet. Sbirka*, v. 76 (6), 39-41

coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat

Silver nitrate

Becker, J. H. R., 1979, *South African J. Surg.*, v. 17 (1), 43-45

Echinococcus, human, surgical evacuation of hepatic cyst using a cryogenic cone, sterilization of cavity with silver nitrate, prevents spillage of cyst fluid and possible anaphylactic shock

Simplotan. See Tinidazole.

Sintomycin. See Chloramphenicol.

Sisomicin, 6'-hydroxy analogue of
Davies, D. H.; et al., 1978, *J. Med. Chem.*,
v. 21 (2), 189-193
Trichomonas vaginalis, Entamoeba histolytica,
6'-hydroxy analogue of sisomicin, antiprotozoal
activity demonstrated in laboratory
trials; no activity against Histomonas meleagridis
and no anthelmintic activity observed

SN-9584
Schmidt, L. H.; et al., 1977, *Antimicrob. Agents and Chemotherapy*, v. 11 (5), 826-843
Plasmodium falciparum in Aotus trivirgatus, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

SN-10274
Schmidt, L. H.; et al., 1977, *Antimicrob. Agents and Chemotherapy*, v. 11 (5), 826-843
Plasmodium falciparum in Aotus trivirgatus, activities of various 4-aminoquinolines against chloroquine-resistant and -susceptible strains, observations confirm cross-resistance among 4-aminoquinolines but indicate that some derivatives may be therapeutically effective against infections refractory to maximally tolerated doses of chloroquine

Sodium antimony dimethylcysteine tartrate (NaP)
Katz, N.; and Pellegrino, J., 1974, *Rev. Inst. Med. Trop. S. Paulo*, v. 16 (6), 346-353
Schistosoma mansoni, sodium antimony dimethylcysteine tartrate, animal and human trials, toxic side effects and electrocardiographic changes in humans

Sodium antimony gluconate. See Antimony sodium gluconate.

Sodium-antimonyl-ethylene-diamino-tetracetate -- SB-EDTA.

SB-EDTA
Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Sodium antimonyl gluconate. See Antimony sodium gluconate.

Sodium arsenite
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Sodium azide
Brotherton, J., 1978, *Arzneimittel-Forsch.*, v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Sodium chloride -- Salt.

Sodium chloride
Bachinskii, V. P.; and Suspitsina, K. T., 1979, *Veterinariia, Moskva* (5), 50
trichodinosis, trout, treatment with sodium chloride in solution

Salt
Banerji, S. R.; Singh, U. N.; and Tiwari, S., 1978, *Current Sc., Bangalore*, v. 47 (8), 283-284 [Letter]
Trichodina [sp.] on exterior of Cyprinus carpio, severe infestation, copper sulphate effective treatment; quick lime, common salt, potassium permanganate, glacial acetic acid, and formalin were not effective: nursery ponds, Patna

Sodium chloride
Eslami, A.; Ahrari, H.; and Saadatzadeh, H., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (3), 307-308
Echinococcus granulosus, scolical activity of cetrimide compared to that of sodium chloride; findings suggest that cetrimide can be used successfully during human hydatid surgery

Sodium chloride
Imada, R.; and Muroga, K., 1979, *Bull. Japan. Soc. Scient. Fish. (Nippon Suisan Gakkaishi)*, v. 45 (1), 25-29
Pseudodactylogyrus microrchis on Anguilla anguilla (gills), trichlorfon, formalin, and sodium chloride baths compared, 2 trichlorfon baths effective and harmless

Sodium and Zinc chlorides, saturated solution
Kunstyr, I.; and Ammerpohl, E., 1978, *Lab. Animals*, v. 12 (2), 95-97
Spiro-nucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite

Sodium chloride
Ricetti, R. V., 1975, *Rev. Fac. Med. Vet. e Zootec. Univ. S. Paulo*, v. 12, 259-268
Cysticercus cellulosae-infected swine carcasses, sodium chloride treatment, length of storage necessary to render meat safe for consumption, comparison with refrigeration

Sodium chloride
Ricetti, R. V.; Panetta, J. C.; and Barbuto, O. J. M., 1977, *Rev. Fac. Med. Vet. e Zootec. Univ. S. Paulo*, v. 14 (1), 113-122
Cysticercus cellulosae in swine meat fragments, effect of refrigeration temperature and salt on viability

Sodium dihydroacetate
Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

- Sodium dimethyldithiocarbamate dihydrate
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- Sodium fluoride
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- Sodium fluoride
Jones, L. D.; and Cook, J., 1978, *Vet. and
Human Toxicol.*, v. 20 (2), 81-82
sodium fluoride, swine, anthelmintic dose,
no significant increase in fluoride concen-
tration in tissues
- Sodium fluoride
Saxena, S. N., 1974, *U. P. Vet. J.*, v. 2 (4),
132-137
Ascarops strongylina, piglets (exper.),
critical trials of efficacy of carbon di-
sulphide, thiabendazole, and sodium fluo-
ride against mature worms
- Sodium fluoride
Venkateswara Rao, P.; et al., 1977, *Riv.
Parassitol.*, Roma, v. 38 (1), 13-21
Prosthogonimus sp. cercariae, cercaricidal
effect of certain common fertilizers, am-
monium sulphate may be cercaricide of choice
- Sodium fluosilicate
Iakubovskii, M. V.; and Zen'kov, A. V., 1977,
Vet. Nauka--Proizvod., Trudy, Minsk, v. 15,
72-78
nematodes, swine raising complexes, combined
control measures, sanitation, anthelmintics,
suiverm most effective
- Sodium fluosilicate
Klenov, A. P., 1971, *Nauch. Trudy, Nauchno-
Issled. Vet. Inst.*, v. 9, 152-155
[Bothriocephalus], white amur, effectiveness
of various anthelmintics
- Sodium fluosilicate
Shnaidmiller, A. P., 1977, *Sborn. Nauch.
Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet.
Inst.* (29), 81-85
ascariasis, trichocephalosis, swine, indus-
trial use of sodium fluosilicate in feed
(indirect method)
- Sodium fluosilicate
Slepnev, N. K., 1971, *Nauch. Trudy, Nauchno-
Issled. Vet. Inst.*, v. 9, 108-112
ascariasis, trichuriasis, oesophagostomiasis,
swine, comparative effectiveness of various
anthelmintics
- Sodium fluosilicate
Tropin, M. I., 1977, *Sborn. Nauch. Rabot
SibNIVI* (28), 133-135
ascariasis, swine, sodium fluosilicate added
to food
- Sodium fluosilicate
Tropin, M. I., 1977, *Sborn. Nauch. Rabot
SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst.*
(29), 86-89
ascariasis, swine, sodium fluosilicate
as effective preimaginal anthelmintic in
feed, controlled and critical tests
- Sodium hydroxide
Uvaliev, I. U.; and Baigaziev, K. K., 1979,
Vestnik Sel'skokhoz. Nauki Kazakhstana (1),
75-78
besnoitiosis, bovine, disinfection of ani-
mals or hides by sodium hydroxide solution
spray; disinfection of premises by sodium
hydroxide or chlorine solution sprays
- Sodium hydroxide
Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-
230
Encephalitozoon cuniculi, survival of spores
after exposure to various temperatures and
disinfectants; growth-inhibition effect of
drugs in cell cultures
- Sodium hypochlorite
Kunstyr, I.; and Ammerpohl, E., 1978, *Lab.
Animals*, v. 12 (2), 95-97
Spiroucleus muris, faecal cysts, resistance
to physical and chemical factors tested,
data may be useful for control of infection
in rodents and for cryopreservation of
parasite
- Sodium hypochlorite
Sherkov, Sh.; et al., 1978, *Vet. Sbirka*, v. 76
(6), 39-41
coccidiosis, calves, disinfectants against
oocysts, sulfaquinoxaline as chemoprophyl-
actic, elancoban-100 as coccidiostat
- Sodium iodoacetate
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- Sodium lauryl sulfate -- SPA-S-222 (with Mepar-
tricin).
- Sodium lauryl sulfate + Mepartricin (= SPA-S-222)
Imparato, E.; et al., 1976, *Quad. Clin. Ostet.
e Ginec.*, v. 31 (4), 225-239
Trichomonas vaginalis, human vaginal tricho-
moniasis, evaluation of mepartricin as oral
therapy using nimorazole and clotrimazole as
reference drugs, best results obtained with
mepartricin-sodium lauryl sulfate
- Sodium lauryl sulfate + Mepartricin (= SPA-S-222)
Ruggeri, E., 1976, *Quad. Clin. Ostet. e Ginec.*,
v. 31 (4), 211-223
Trichomonas vaginalis, vaginal trichomoniasis,
oral mepartricin showed significant
therapeutic superiority when compared in
clinical trials with patients who received
nimorazole therapy orally or with patients
treated with a vaginal cream containing
amphotericin B and tetracycline
- Sodium malonate
Brotherton, J., 1978, *Arzneimittel-Forsch.*,
v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential
trichomonacides using Coulter Counter
- Sodium N-phenylglycinamide-p-arsenodithioglycol-
late. See Tryparsamide.

Sodium salicylate

Duncanson, F. P.; Abelmann, W. H.; and Pan, C., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 577-578

Trypanosoma cruzi, growth in vitro unaffected by sodium salicylate

Sodium santoninate. See Santonin.

Sodium stibogluconate. See Antimony sodium gluconate.

Sodium sulfadimethylpyrimidine. See Sulfamethazine.

Sodium thiacetarsamide. See Arsenamide.

Somonil. See Methidathion.

Sonex. See *Embelia ribes* extract or Nicotine or Pomegranate.

Spartrix. See Carnidazole.

Spectinomycin + Lincomycin (= Lincospectin)

Bischofova, N.; and Zajicek, D., 1976, *Veterinarstvi*, v. 26 (11), 506-507

Balantidium coli, pigs, large-scale fattening farms, lincospectin treatment

Spiramycin -- Rovamycin; Selectomycin.

Spiramycin (Selectomycin)

Brunnthaler, F., 1977, *Prakt. Tierarzt*, v. 58 (11), 849-851

coccidiosis, dogs, treatment with various drugs, best results with amprolium

Rovamycin

Gelle, P.; et al., 1975, *Rev. Franc. Gynec. et Obst.*, v. 70 (5), 329-333

toxoplasmosis, survey of pregnant women before and after delivery, non-immune women had more frequent abortions, of women who acquired infection during pregnancy those treated with rovamycin had fewer stillborn infants and infants free of infection than those who received no treatment

Spiramycin

Monnier, J. C.; et al., 1975, *Rev. Franc. Gynec. et Obst.*, v. 70 (5), 325-328

toxoplasmosis, survey of pregnant women using the complement fixation and immunofluorescence tests, those found to have evidence of infection were treated with spiramycin, all had normal deliveries and infants free of infection

Spiramycin

Ohshima, S.; Hoshino, M.; and Tanaka, H., 1977, *Kiseichugaku Zasshi (Japan. J. Parasitol.)*, v. 26 (3), 127-131

Toxoplasma, 11 strains, susceptibility to 6 drugs, mice

Spiramycin -- Continued.

Spiramycin + Sulfamethoxyypyridazine

Thiermann, E.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens compared during acute and late infections; pyrimethamine + sulfamethoxyypyridazine was most effective

Spiramycin (Rovamycin)

Trzaska, B.; and Leinweber-Noiszewska, H., 1974, *Ann. Acad. Med. Stetinensis*, suppl. 10, 19-25

Entamoeba histolytica, acute infection in Polish sailor who acquired disease in West Africa, therapy with bemarsal, mexaform and spiramycin resulted in relapse and chronic infection, apparent cure with metronidazole; pathology and clinical aspects of amoebiasis

Spiramycin

Waller, T., 1979, *Lab. Animals*, v. 13 (3), 227-230

Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Spirotrypan

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

SQ 21,704

Gemmell, M. A.; Johnstone, P. D.; and Oudemans, G., 1978, *Research Vet. Sc.*, v. 25 (1), 109-110

Echinococcus granulosus, *Taenia hydatigena*, dogs, antibiotic of streptothricin family (SQ 21,704), significant activity against *T. hydatigena*, evidence for similar activity against *E. granulosus* was less well marked

Squash seeds, defatted

Klenov, A. P., 1971, *Nauch. Trudy, Nauchno-Issled. Vet. Inst.*, v. 9, 152-155

[*Bothriocephalus*], white amur, effectiveness of various anthelmintics

Stanneous oxide. See Tin.

Stannotaen. See Tin.

Starbar GX-118. See Phosmet.

Starogyn. See Broxyquinoline.

Statyl. See Methyl benzoate.

Steclin. See Tetracycline.

Stenorol. See Halofuginone.

Sterosan. See Chlorquinaldol.

Stibocaptate -- Antimony dimercapto-succinate;
Astiban.

Astiban

Abdulla, W. A.; et al., 1977, Egypt. J. Bilharz., v. 4 (2), 109-115
S[chistosoma] mansoni, prophylactic activity, antischistosomal drugs, albino mice, most effective within 2 weeks post exposure

Stibocaptate (Astiban)

Denham, D. A.; Suswillo, R. R.; and Roger, R., 1978, J. Helminth., v. 52 (3), 227-228
Brugia pahangi, stibocaptate: killed 3rd stage larvae in vitro but had no effect on microfilariae, no larvae developed in infected treated mosquitoes, neither micro- nor macrofilaricidal in either Meriones unguiculatus or cats but did affect embryogenesis

Stibocaptate (Astiban)

Duke, B. O. L., 1977, Tropenmed. u. Parasitol., v. 28 (4), 447-455
Onchocerca volvulus, chimpanzees, pentamidine, stibocaptate, nifurtimox, 3 other compounds, macro- and microfilaricidal action, toxicity

Astiban (Antimony dimercapto-succinate)

El-Kholy, Z. A.; et al., 1979, Biochem. Pharmacol., v. 28 (21), 3171-3172
Schistosoma mansoni-infected and normal mice, effect of astiban on β -glucuronidase activity in liver, spleen, kidney, and bladder homogenates

Astiban

Erasmus, D. A.; and Davies, T. W., 1979, Exper. Parasitol., v. 47 (1), 91-106
Schistosoma mansoni, S. haematobium, calcareous corpuscles in vitelline cells, morphological observations, X-ray microanalysis, effect of drug treatment

Astiban

Magzoub, M., 1971, Sudan Med. J., v. 9 (3), 178-182
Schistosoma mansoni, untreated worms and worms treated with ambilhar or astiban, electron microscopy of cuticle, subcuticular region, and gut; possibility that egg formation is interrupted by either treatment

Stibophen -- Fouadin; Fuadin.

Stibophen (Fouadin)

Abdel Samad, M. M.; et al., 1977, Tropenmed. u. Parasitol., v. 28 (4), 554-559
Schistosoma mansoni, mice, liver monoamine oxidase activity during course of infection and after chemotherapy, may be useful index for progression or regression of liver fibrosis

Stibophen -- Continued.

Fouadin

Akhound-Zadeh, H., 1976, Rev. Internat. Serv. Sante Armees, v. 49 (5), 421-426
chronic cutaneous leishmaniasis, soldier with severe ulcers that did not heal despite 8 years of therapy with various anti-leishmanial drugs, chronicity thought to be result of immuno-deficiency, ulcers finally cured after additional therapy with monomycine: Iran (had travelled to Khouzistan)

Stibophen

Campbell, W. C.; Bartels, E.; and Cuckler, A. C., 1978, J. Parasitol., v. 64 (1), 69-77
Schistosoma mansoni, mice, simple and rapid assay suitable for routine screening of compounds for antischistosome activity, reduction in severity of hepatic lesions used as chief criterion of efficacy

Stibophen (Fuadin)

Ghazal, A.; Ismail, M.; and Sharabi, F. M., 1978, Egypt. J. Pharm. Sc., v. 17 (1), 1976, 43-52
Schistosoma mansoni-infected mice, tissue histamine content before and after treatment with several antischistosomal drugs

Fuadin

Loehr, H.; and Wolf, H., 1978, Deutsche Med. Wchnschr., v. 103 (10), 424-427
visceral leishmaniasis in child apparently acquired while traveling in Yugoslavia, manifestations of severe septic temperature elevations, diagnostic difficulties with diagnosis finally by immunoserologic means, improvement in condition after fuadin therapy: Germany

Fouadin

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Stibophen (Fuadin)

Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Fouadin

Youssef, A. H., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 147-157
Dipetalonema evansi, camels, filarial orchitis and possible significance as prevalent reproductive disease; surgical treatment and use of neosulversan, fouadin, and neguvon, histopathology of gonads: Egypt

Stilbamidine diisethionate

Kinnamon, K. E.; and Rane, D. S., 1978, Internat. J. Parasitol., v. 8 (6), 515-523
Trypanosoma rhodesiense, mice, greater than 1 year protection from lethal infections by prophylactic drugs and active immunity

Stilbazium iodide -- Monopar.

Stilbazium iodide

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, Rev. Brasil. Med., v. 31 (11), 791-794

human soil-transmitted nematodes, laboratory trials testing ovicidal and larvicidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Monopar

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Stirofos. See Tetrachlorvinphos.

St. John's wort, common, blossoms and stems

Slepnev, N. K., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 107-109

Ascaris suum, in vitro testing of anthelmintic activity of water extracts of some plants

Streptomycin

Evans, W. S., 1978, Canad. J. Zool., v. 56 (5), 1210-1211

Hymenolepis microstoma, effect of streptomycin and penicillin on growth and differentiation in vitro

Streptomycin sulphate

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Streptozotocin (WR 139 502)

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160

Trypanosoma rhodesiense, mice, active in screening of antitumor compounds for efficacy against infection

Strike Insect Strips. See Dichlorvos.

Strongid-P granules or paste. See Pyrantel.

Strongid-T. See Pyrantel.

Styrylacetic acid

Christow, C. P., 1974, Rev. Biol. Trop., v. 21 (2), 1973, 187-195

Trichomonas vaginalis, in vitro, reproduction inhibited by anti-amino acids (8-indolylacrylic acid, styrylacetic acid)

Suiverm. See Piperazine or Thiabendazole.

Sulfachloropyrazine -- N'-(6-Chloro-2 pyrazinyl)-sulfanilamide; Esb₃; Sulphachloropyrazine.

Sulphachloropyrazine (Esb₃)

Manuel, M. F.; and de Leon, M. L., 1978, Philippine J. Vet. Med., v. 17 (1-2), 129-141
Eimeria tenella, White Leghorn chicks (exper.), comparative efficacy of Esb₃ and Abi-Zet₅₀

Esb₃

Zaprianov, M. Ia., 1978, Vet. Med. Nauki, v. 15 (4), 103-107

Eimeria kofoidi in *Alectoris graeca cypritis* (nat. and exper.), amprolium, DOT-soluble, and Esb₃ tested

Sulfadiazine -- Belmet (with Sulfamerazine sodium and Sulfamethazine sodium); Gynben vaginal cream (with Diethylstilbestrol and Diiodo-hydroxyquin); Sulfadiazine sodium; 2-Sulfanilamidopyrimidine; Sulphadiazine; Tribriksen (with Trimethoprim); WR 7,557.

Gynben vaginal cream

Aguiar, L. M.; Bastos, A. da C.; and Salvatore, C. A., 1977, Rev. Brasil. Med., v. 34 (3), 151-152

vaginal trichomoniasis, humans, gynben vaginal cream, excellent antipruritic effect and good tolerance

Sulfadiazine + Trimethoprim (= Tribriksen)

Brunthaler, F., 1977, Prakt. Tierarzt, v. 58 (11), 849-851

coccidiosis, dogs, treatment with various drugs, best results with amprolium

Sulfadiazine

Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397

Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances

Sulfadiazine

Eke, R. A., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 1074-1075

Plasmodium falciparum, possible chloroquine-resistant strain, recrudescence of infection in 42-year-old hospital worker after chloroquine therapy, radical cure with sulfadiazine and pyrimethamine: Nigeria

Sulfadiazine

Elslager, E. F.; et al., 1979, J. Med. Chem., v. 22 (10), 1247-1257

Plasmodium spp., antimalarial activity of 2,4-diamino-6-(2-naphthylsulfonyl)quinazoline and related 2,4-diamino-6-[(phenyl and naphthyl)sulfinyl and sulfonyl]quinazolines

Sulfadiazine

Korolkovas, A.; and Pellegrino, J., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (4), 219-223

Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active

Sulfadiazine -- Continued.

Sulfadiazine-Trimethoprim

McKenzie, R. A.; Green, P. E. ; and Wood, A. D., 1978, Austral. Vet. J., v. 54 (2), 86-88
Angiostrongylus cantonensis in captive *Macropus rufogriseus* (surface of brain beneath leptomeninges, cerebellar folium, meninges), clinical symptoms, pathology, treatment with trimethoprim-sulfadiazine ineffective, case report: Brisbane

Sulfadiazine

McLeod, R.; et al., 1979, Am. J. Med., v. 67 (4), 711-714

Toxoplasma gondii, immunosuppressed man, brain abscesses, sulfadiazine, pyrimethamine, and calcium leukovorin, case report

Sulfadiazine sodium + Sulfamethazine sodium + Sulfamerazine sodium (=Belmet)

Manuel, M. F.; and Neri, R. A., 1975, Philippine J. Vet. Med., v. 14 (1), 106-116

Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfadiazine

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351

Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Sulfadiazine + Pyrimethamine

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351

Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Sulfadiazine + Trimethoprim

Neal, R. A., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 341-351

Leishmania tropica, 7 Latin-American and 2 Asiatic isolates, course of infection in hamsters, anti-folic reductase drugs compared with paromomycin and sodium stibogluconate

Sulfadiazine + Pyrimethamine

de Paillerets, F.; et al., 1975, Semaine Hop. Paris, v. 51 (43), 2589-2592 (Ann. Pediat., Paris, v. 22 (11), 801-804)

toxoplasmosis, infant, congenital infection, pyrimethamine-sulfadiazine therapy supplemented with folic acid

Sulfadiazine (WR 7,557)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947

sporozoite-induced *Plasmodium berghei* in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice.

Sulfadiazine

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 808-818

Plasmodium falciparum, *P. vivax*, various drug-resistant and drug-susceptible strains in *Aotus trivirgatus griseimembra*, capacity of sulfadiazine to enhance activities of WR-158,122 and WR-159,412

Sulfadiazine -- Continued.

Sulphadiazine (WR-7557) + 2,4-Diamino-6-(2-naphthylsulphonyl)quinazoline (WR-158122)

Wise, D. L.; Gresser, J. D.; and McCormick, G. J., 1979, J. Pharm. and Pharmacol., v. 31 (4), 201-204

dual antimalarial system, sustained release of ³H-labelled WR-7557 and ¹⁴C-labelled WR-158122 in biodegradable carrier, rhesus monkeys, mice

Sulfadiazine sodium. See Sulfadiazine.

Sulfadimethoxine -- Madribon; Pacprim (with Sulfisomidine, Sulfisoxazole and Trimethoprim); Sulfanilamide complex (with Sulfisomidine, Sulfisoxazole and Trimethoprim).

Sulfadimethoxine (Madribon)

Brunnthaler, F., 1977, Prakt. Tierarzt, v. 58 (11), 849-851

coccidiosis, dogs, treatment with various drugs, best results with amprolium

Sulfadimethoxin

Kogan, G. F., 1975, Vet. Nauka--Proizvod., Trudy, Minsk, v. 13, 196-198

nosematosis, bees, fumagillin had good therapeutic effect, metronidazole, sulfadimethoxin and enteroseptol showed no substantial effect

Sulfadimethoxine + Trimethoprim + Sulfisomidine + Sulfisoxazole (= Pacprim; Sulfanilamide complex)

Lee, H. S.; Lee, H. B.; and Moon, M. H., 1979, Korean J. Animal Sc., v. 21 (4), 281-288

Besnoitia besnoiti, goats (exper.), antimony potassium tartrate, pacprim

Sulfadimethoxine (Madribon)

Pandey, N. N.; and Mishra, S. S., 1978, Indian Vet. J., v. 55 (2), 144-148

Babesia bigemina, indigenous cow calves, berenil and acriflavin effective, sulfadimethoxine ineffective: India

Sulfadimezine. See Sulfamethazine.

Sulfadoxine -- Fansil; Fansidar (with Pyrimethamine); Sanasil; 4-Sulfanilamido-5,6-dimethoxyypyrimidine; Sulformethoxine; Sulphadoxine.

Sulfadoxine + Pyrimethamine (Fansidar)

Al Tawil, N., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (3), 409-413

Plasmodium falciparum, semi-immune humans, clearance of asexual parasitaemia with single dose sulfadoxine-pyrimethamine, comparison with standard dose of chloroquine over 3 days: Laos

Fansil + Pyrimethamine

Botelho, A., 1974, South African Med. J., v. 48 (31), 1351-1352

malarias, humans, clinical trials using sulfones and sulphonamides with a pyrimidine derivative

Sulfadoxine -- Continued.

- Sulfadoxine + Pyrimethamine (= Fansidar)
Chongsuphajaisiddhi, T.; et al., 1979, South-east Asian J. Trop. Med. and Pub. Health, v. 10 (1), 132-137
falciparum malaria, children, chloroquine resistance, efficacy of quinine and fansidar, clinical study: Thailand
- Sanasil (Fanasil)
Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances
- Sulfadoxine + Pyrimethamine (= Fansidar)
Doberstyn, E. B.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 15-17
Plasmodium vivax, humans, pyrimethamine alone or combined with sulfadoxine is not effective therapy for acute episode: Thailand
- Sulfadoxine + Pyrimethamine (= Fansidar)
Ferraroni, J. J.; and Hayes, J., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 909-911
Plasmodium falciparum outbreak among indigenous indian tribe, 3 cases resistant to chloroquine responded favorably to fansidar therapy: Uauaris, Territory of Roraima, Brazil
- Sulfadoxine + Pyrimethamine (= Fansidar)
Fogh, S.; Jepsen, S.; and Effersøe, P., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (2), 228-229
Plasmodium falciparum, chloroquine resistant strain in non-immune male, treatment with fansidar terminated infection: Danish tourist to Kenya
- Sulphormethoxine
Neves, J.; and Moura, H. B., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (2), 126-130
Plasmodium falciparum, P. vivax, prevalence survey in hospital patients, discussion of changes in prevalence with introduction of chloroquine resistant strains of P. falciparum, treatment trials with various malarial drugs: Brazil
- Sulfadoxine + Pyrimethamine (= Fansidar)
Ponnampalam, J. T., 1978, J. Trop. Med. and Hyg., v. 81 (10), 198-203
human malarias, comparative study of prophylaxis using chloroquine and a combination of sulfadoxine and pyrimethamine: residents of rubber estate in central Malaysia
- Sulphadoxine + Pyrimethamine (= Fansidar)
Rumans, L. W.; Dennis, D. T.; and Atmosoedjono, S., 1979, Lancet, London (8142), v. 2, 580-581 [Letter]
Plasmodium falciparum, fansidar-resistant malaria in case also resistant to chloroquine: Indonesia
- Sulphormethoxine (Fanasil)
Vincke, I. H., 1970, Ann. Soc. Belges Med. Trop. Parasitol., v. 50 (3), 339-358
Plasmodium berghei berghei, mice, action of pyrimethamine and sulphormethoxine on pre-erythrocytic and sporogonous cycles

Sulfadoxine -- Continued.

- Sulfadoxin + Trimethoprim
Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures
-
- Sulfaguandinine (Resulfon)
Brunnthaler, F., 1977, Prakt. Tierarzt, v. 58 (11), 849-851
coccidiosis, dogs, treatment with various drugs, best results with amprolium
- Sulfaguandinine
Dubey, J. P.; Weisbrode, S. E.; and Rogers, W. A., 1978, J. Am. Vet. Med. Ass., v. 173 (2), 185-191
Isospora ohioensis-like organism, dog (small intestine, cecum, colon, terminal ileum, villous epithelium, lamina propria, intestinal glands), description of asexual and sexual stages, pathology, attempted treatment with sulfaguandinine unsuccessful, case report: Ohio
- Sulfalene -- Kelfizine; 2-Methoxy-3-sulfanilamidopyrazine; Pyrikelfizine (with Pyrimethamine); Sulfamethoxyypyrazine; Sulfamethoxyypyrazine-longum.
- 2-Methoxy-3-sulfanilamidopyrazine (Kelfizine)
Chrusciel, T. L.; et al., 1970, Acta Parasitol. Polon., v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances
- Sulfamethoxyypyrazine (Kelfizine)
Herman, Z.; Sokoła, A.; and Szaflarski, J., 1970, Acta Parasitol. Polon., v. 18 (42-50), 483-486
Toxoplasma gondii, mice, comparison of pyrimethamine and sulfamethoxyypyrazine used separately or in combination and of a long-acting preparation of sulfamethoxyypyrazine
- Sulfamethoxyypyrazine-longum
Herman, Z.; Sokoła, A.; and Szaflarski, J., 1970, Acta Parasitol. Polon., v. 18 (42-50), 483-486
Toxoplasma gondii, mice, comparison of pyrimethamine and sulfamethoxyypyrazine used separately or in combination and of a long-acting preparation of sulfamethoxyypyrazine
- Sulfamethoxyypyrazine + Pyrimethamine (= Pyrikelfizine)
Herman, Z.; Sokoła, A.; and Szaflarski, J., 1970, Acta Parasitol. Polon., v. 18 (42-50), 483-486
Toxoplasma gondii, mice, comparison of pyrimethamine and sulfamethoxyypyrazine used separately or in combination and of a long-acting preparation of sulfamethoxyypyrazine

Sulfalene -- Continued.

Sulfalene

Williams, R. L.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (2, pt. 1), 226-231
Plasmodium falciparum, humans with chloroquine-resistant infection (exper.), acetylator phenotype does not influence therapeutic response to sulfalene or sulfalene combined with pyrimethamine

Sulfamerazine -- Belmet (with Sulfadiazine sodium and Sulfamethazine sodium); Sulfamerazine sodium; Supronal (with Sulfatolamide).

Sulfamerazine sodium + Sulfadiazine sodium + Sulfamethazine sodium (=Belmet)
 Manuel, M. F.; and Neri, R. A., 1975, *Philippine J. Vet. Med.*, v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfamerazine

Nishimura, T., 1977, *J. Tokyo Univ. Fish.*, v. 63 (2), 71-79
 sulfamonomethoxine, rainbow trout, toxicity compared with sulfamerazine

Supronal

Quintero Monasterios, R.; and Castro R., G., 1972, *Rev. Obst. y Ginec. Venez.*, v. 32 (3), 483-488
 human urogenital toxoplasmosis, statistics of cases studied because of secondary sterility, clinical management, therapy with daraprim and supronal

Sulfamerazine sodium. See Sulfamerazine.

Sulfameter -- Bayrena; 5-Methoxy-2-sulfanilamidepyrimidine.

5-Methoxy-2-sulfanilamidepyrimidine (Bayrena)
 Chrusciel, T. L.; et al., 1970, *Acta Parasitol. Polon.*, v. 18 (27-41), 393-397
Toxoplasma gondii, mice, therapeutic effect of bayrena and kelfizine alone or in combination with dimethylsulphoxide, and of several other antibacterial, antiviral, and antiprotozoan substances

Sulfamethazine -- Abi-Zet₅₀ (with Pyrimethamine); Belmet (with Sulfadiazine sodium and Sulfamerazine sodium); Sodium sulfadimethylpyrimidine; Sulfadimezine; Sulphadimidine; Sulfamethazine sodium; Sulmet; Sulphacombine (with Diaveridine).

Sulphadimidine + Diaveridine (=Sulphacombine)
 Danek, J.; et al., 1978, *Biol. a Chem. Zivoc. Vyroby, Vet.*, v. 14 (v. 20) (2), 151-169
Eimeria spp., rabbits (nat. and exper.), sulphacombine, controlled test, subacute toxicity, no negative effect on followed indicators

Sulfamethazine -- Continued.

Sulfamethazine

Korolkovas, A.; and Pellegrino, J., 1978, *Rev. Inst. Med. Trop. S. Paulo*, v. 20 (4), 219-223
Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active

Sulfadimezine

Krylov, V. F., 1978, *Veterinariia, Moskva* (10), 68-69
Eimeria tenella strain resistant to phar-mococcid after 35 laboratory passages in chickens, cross-resistance only to rigecoccin

Sulfadimidine + Pyrimethamine (=Abi-Zet₅₀)

Manuel, M. F.; and de Leon, M. L., 1978, *Philippine J. Vet. Med.*, v. 17 (1-2), 129-141
Eimeria tenella, White Leghorn chicks (exper.), comparative efficacy of Esb₃ and Abi-Zet₅₀

Sodium sulfadimethylpyrimidine (Sulmet)

Manuel, M. F.; and Neri, R. A., 1975, *Philippine J. Vet. Med.*, v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfamethazine sodium + Sulfadiazine sodium + Sulfamerazine sodium (=Belmet)

Manuel, M. F.; and Neri, R. A., 1975, *Philippine J. Vet. Med.*, v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulphadimidine

Oduye, O. O.; and Bobade, P. A., 1979, *J. Small Animal Practice*, v. 20 (3), 181-184
Isospora canis, kennelled dogs, outbreak of haemorrhagic diarrhoea, sulphadimidine treatment

Sulphadimidine + Diaveridine (=Sulphacombine)

Strakova, J.; Sevcik, B.; and Dvorak, M., 1978, *Biol. a Chem. Zivoc. Vyroby, Vet.*, v. 14 (v. 20) (2), 171-180
 coccidiosis, chicks, sulphacombine, acute and subacute toxicity studies

Sulfamethazine sodium. See Sulfamethazine.

Sulfamethoxazole -- Ciplin (with Trimethoprim); Co-trimoxazole (with Trimethoprim); Eusaprim (with Trimethoprim); 5-Methyl-3-sulfanilamido-isoxazole; Sulphamethoxazol; Septran (with Trimethoprim); Septrin (with Trimethoprim).

Sulfamethoxazole-Trimethoprim

Deeg, H. J.; et al., 1979, *Transplantation*, v. 28 (3), 243-246
 effect of trimethoprim-sulfamethoxazole on hematological recovery after total body irradiation and autologous marrow transplantation studied in dogs, results show that drug can be given safely and probably prevents very early cases of *Pneumocystis carinii* pneumonia

Sulfamethoxazole -- Continued.

Co-trimoxazole

Geddes, A. M.; Ball, A. P.; and Farrell, I. D., 1979, *J. Antimicrob. Chemother.*, v. 5, suppl. B, 221-230
co-trimoxazole for treatment of serious infections, review including information on *Pneumocystis carinii*, malaria, and toxoplasmosis

Sulfamethoxazole

Grossman, P. L.; and Remington, J. S., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (3), 445-455
Toxoplasma gondii, in vitro and in vivo in mice, effects of trimethoprim and sulfamethoxazole alone and in combination

Sulfamethoxazole-Trimethoprim

Hughes, W. T., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (3), 333-335
Pneumocystis carinii, immunosuppressed rats, trimethoprim-sulfamethoxazole has limited rather than lethal effect, protection is afforded only during period of administration

Sulfamethoxazole + Trimethoprim

Hughes, W. T.; et al., 1978, *J. Pediat.*, St. Louis, v. 92 (2), 285-291
Pneumocystis carinii, children, comparison of pentamidine isethionate and trimethoprim combined with sulfamethoxazole (TMP-SMZ) in treating *Pneumocystis pneumonia*, results show that TMP-SMZ is as effective as pentamidine, has minimal side effects, offers oral administration and is more readily available

Sulfamethoxazole

Kluge, R. M.; Spaulding, D. M.; and Spain, A. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (6), 975-978
Pneumocystis carinii in steroid-conditioned rats, combination of pentamidine with trimethoprim-sulfamethoxazole, data suggest that combination therapy is no more effective than trimethoprim-sulfamethoxazole alone and may be, in fact, harmful, trimethoprim by itself has no place in treatment of pneumocystosis

Sulfamethoxazole

Korolkovas, A.; and Pellegrino, J., 1978, *Rev. Inst. Med. Trop. S. Paulo*, v. 20 (4), 219-223
Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active

Sulphamethoxazol + Trimethoprim (= Septrin)

Kouba, K.; Nevarilova, A.; and Rajlichova, J., 1978, *Ceskoslov. Epidemiol., Mikrobiol., Immunol.*, v. 27 (3), 175-178
toxoplasmosis, human, therapy with septrin, poor results with allergic side effects

Sulfamethoxazole-Trimethoprim

Larter, W. E.; et al., 1978, *J. Pediat.*, St. Louis, v. 92 (5), 826-828
Pneumocystis carinii, trimethoprim-sulfamethoxazole treatment of pneumonitis in children

Sulfamethoxazole -- Continued.

Sulfamethoxazole

Lichtenwalner, D. M.; et al., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (5), 579-583
trimethoprim, sulfamethoxazole, rapid assay for determination of levels in serum by spectrofluorometry

Sulphamethoxazole

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511
Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in in vitro screens to test wide range of compounds for chemotherapeutic activity

Sulfamethoxazole + Trimethoprim (= Septrin)

Mancinella, J. E.; et al., 1975, *Semana Med.* (4914), an. 82, v. 147 (9), 230-235
Trichomonas, human vaginal infections, therapeutic action of sulfamethoxazole combined with trimethoprim analyzed

Sulphamethoxazole + Trimethoprim

Norrby, R.; et al., 1975, *Scand. J. Infect. Dis.*, v. 7 (1), 72-75
Toxoplasma gondii, humans, clinical and serological data on patients treated with trimethoprim-sulphamethoxazole

Sulfamethoxazole + Trimethoprim (= Co-trimoxazole; Eusaprim)

Norrby, R.; and Eilard, T., 1976, *Scand. J. Infect. Dis.*, v. 8 (4), 275-276
toxoplasmosis, recurrent infection in woman treated with co-trimoxazole, normal clinical response to each course of therapy, no evidence of impaired immunity

Sulphamethoxazole + Trimethoprim (= Septran; = Ciplin)

Shashindran, C. H.; et al., 1978, *Brit. J. Dermat.*, v. 98 (6), 699-700
human pediculosis capitis, successful systemic (oral) therapy using combination of trimethoprim and sulphamethoxazole without additional external application of insecticides; drugs when used separately were not effective

Sulfamethoxazole-Trimethoprim

Stevenson, D. K.; Christie, D. L.; and Haas, J. E., 1978, *Pediatrics*, *Am. Acad. Pediat.*, v. 61 (6), 864-866
trimethoprim-sulfamethoxazole, child, hepatic injury

Sulfamethoxazole + Trimethoprim (= Septrin)

Szaflarski, J.; Sokola, A.; and Herman, T. S., 1974, *Acta Parasitol. Polon.*, v. 22 (22-34), 261-263
Toxoplasma gondii, mice (exper.), trimethoprim and sulfamethoxazole alone and in combination, concluded that trimethoprim has no therapeutic effect but potentiates action of sulfamethoxazole

Sulfamethoxazole + Trimethoprim

Thiermann, E.; et al., 1978, *Am. J. Trop. Med. and Hyg.*, v. 27 (4), 747-750
Toxoplasma gondii, mice, 5 treatment regimens compared during acute and late infections; pyrimethamine + sulfamethoxyypyridazine was most effective

Sulfamethoxazole -- Continued.

Sulfamethoxazole + Trimethoprim (= Co-trimoxazole)
Westerman, E. L.; and Christensen, R. P., 1979,
Ann. Int. Med., v. 91 (3), 413-414

Isospora belli, immunosuppressed woman con-
currently infected with Giardia lamblia,
severe diarrhea, rapid remission with co-
trimoxazole, case report

Sulfamethoxazole + Trimethoprim

Wolff, L. J.; and Baehner, R. L., 1978, Am. J.
Dis. Child., v. 132 (5), 525-526

Pneumocystis carinii, clinical trials
evaluating prophylactic value of a 2-week,
high-dose course of trimethoprim-sulfamethoxa-
zole to prevent pneumonia in children with
cancer who are receiving their first 100
days of intensive immunosuppressive chemo-
therapy

Sulfamethoxazole+Trimethoprim

Yoshida, Y.; et al., 1977, Kiseichugaku Zasshi
(Japan. J. Parasitol.), v. 26 (6), 367-375

Pneumocystis carinii pneumonia, rats, com-
parative efficacy of pyrimethamine+sulfa-
monomethoxine vs. trimethoprim+sulfamethoxa-
zole

Sulfamethoxyprazine. See Sulfalene.

Sulfamethoxyprazine-longum. See Sulfalene.

Sulfamethoxyprazine + Pyrimethamine

Thiermann, E.; et al., 1977, Rev. Med. Chile,
v. 105 (7), 433-435

Toxoplasma gondii, mice, experimental trials
comparing efficacy of clindamycin with that
of pyrimethamine combined with sulfamethoxy-
prazine, combination drug cured 100% of
mice while mice treated with clindamycin sur-
vived during treatment but 50% died from
severe infections after therapy had been dis-
continued

Sulfamethoxyprazine (Lederkyn)

Thiermann, E.; et al., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens
compared during acute and late infections;
pyrimethamine + sulfamethoxyprazine was
most effective

Sulfamethoxyprazine + Clindamycin

Thiermann, E.; et al., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens
compared during acute and late infections;
pyrimethamine + sulfamethoxyprazine was
most effective

Sulfamethoxyprazine + Pyrimethamine

Thiermann, E.; et al., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens
compared during acute and late infections;
pyrimethamine + sulfamethoxyprazine was
most effective

Sulfamethoxyprazine + Spiramycin

Thiermann, E.; et al., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (4), 747-750

Toxoplasma gondii, mice, 5 treatment regimens
compared during acute and late infections;
pyrimethamine + sulfamethoxyprazine was
most effective

Sulfamonomethoxine -- 4-Methoxy-6-sulfanilamido-
pyrimidine monohydrate.

Sulfamonomethoxine + Pyrimethamine

Manuel, M. F.; Morales, E. G.; and Tavela,
E., [1977], Philippine J. Vet. Med., v. 15
(1-2), 1976, 87-95

Leucocytozoon caulleryi, White Leghorn
cockerels, superior prophylactic value of
sulfamonomethoxine + pyrimethamine admin-
istered in feed under field conditions, no
detrimental effects on growth rate or blood
picture

Sulfamonomethoxine

Matsui, T.; et al., 1977, Kiseichugaku Zasshi
(Japan. J. Parasitol.), v. 26 (4), 235-239

Isospora rivolta, cats, mice (all exper.),
sulfamonomethoxine

Sulfamonomethoxine

Nishimura, T., 1977, J. Tokyo Univ. Fish.,
v. 63 (2), 71-79

sulfamonomethoxine, rainbow trout, toxicity
compared with sulfamerazine

Sulfamonomethoxine

Ohshima, S.; Hoshino, M.; and Tanaka, H.,
1977, Kiseichugaku Zasshi (Japan. J. Parasi-
tol.), v. 26 (3), 127-131

Toxoplasma, 11 strains, susceptibility to
6 drugs, mice

Sulfamonomethoxine

Toriumi, T.; et al., 1978, Scient. Rep. Fac.
Agric. Okayama Univ. (52), 49-53

Leucocytozoon caulleryi, chickens (exper.),
sulfamonomethoxine and halofuginone in feed
prevented infection

Sulfamonomethoxine

Waki, S., 1976, Kiseichugaku Zasshi (Japan. J.
Parasitol.), v. 25 (6), 441-446

Plasmodium berghei, mice, protective immunity
induced by repeated infections followed by
radical chemotherapy with sulfamonomethoxine

Sulfamonomethoxine+Pyrimethamine

Yoshida, Y.; et al., 1977, Kiseichugaku Zasshi
(Japan. J. Parasitol.), v. 26 (6), 367-375

Pneumocystis carinii pneumonia, rats, com-
parative efficacy of pyrimethamine+sulfa-
monomethoxine vs. trimethoprim+sulfamethoxa-
zole

2-Sulfamoyl-4,4'-diaminodiphenylsulfone -- SDDS.

2-Sulfamoyl-4,4'-diamino diphenylsulfone
Kumar, P. S.; Kumar, R.; and Mohapatra, L. N.,
1978, Indian J. Med. Research, v. 67, 908-917
Toxoplasma gondii, rabbits treated with 2-
sulfamoyl-4,4'-diamino diphenylsulfone, de-
termination of minimum curative dose, haemag-
glutinating antibody response in primary and
challenge infection, immunity to challenge
infection, schedule for raising high titre
serum

2-Sulfamoyl-4,4'-diaminodiphenylsulfone (SDDS)
Ohshima, S.; Hoshino, M.; and Tanaka, H.,
1977, Kiseichugaku Zasshi (Japan. J. Parasit-
ol.), v. 26 (3), 127-131
Toxoplasma, 11 strains, susceptibility to
6 drugs, mice

SDDS

Werner, H.; et al., 1977, Tropenmed. u. Parasit-
tol., v. 28 (4), 528-532
Toxoplasma gondii, latent infected mice, sub-
stantial reduction in brain cysts obtained by
administration of hyperimmune serum, pyrimeth-
amine, and SDDS in various combinations;
effectiveness of therapy varied with parasite
strain

Sulfanilamide complex. See Sulfadimethoxine or
Sulfisomidine or Sulfisoxazole or Trimethoprim.

4-Sulfanilamido-5,6-dimethoxyypyrimidine. See
Sulfadoxine.

2-Sulfanilamidopyrimidine. See Sulfadiazine.

Sulfanitran -- Novostat-W (with Aklomide);
Polystat (with Dibutyltin dilaurate, Dinsed,
and Roxarsone).

Polystat

McDougald, L. R.; and McQuiston, T. E., 1978,
Avian Dis., v. 22 (4), 765-770
Eimeria spp., turkeys, coccidiosis manage-
ment, innate (age) resistance and acquired
immunity vs. anticoccidial medication

Sulfanitran + Aklomide (=Novostat-W)

Manuel, M. F.; and Neri, R. A., 1975, Philip-
pine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels,
efficacy of 7 water-soluble coccidiostats

Sulfaphenazole

Banerjee, N. C.; et al., 1979, Indian Poultry
Gaz., v. 63 (1), 19-21
sulfaphenazole, blood level, biological half
life, volume distribution, and tissue dis-
persion in poultry, possible public health
hazard

Sulfaquinoxaline -- Darvisul (with Diaveridine);
Embazin; Nitryl (with p-Dimethylaminobenzo-
nitrile); Noxal; Pancoxin (with Amprolium and
Ethopabate); Pancoxin plus (with Amprolium,
Ethopabate and Pyrimethamine); N'-(2-Quinox-
aliny) sulfanilamide; Saquadil (with Diaver-
idine); Sulfaquinoxaline sodium; Sulpha-
quinoxaline; Supacox (with Amprolium, Etho-
pabate and Pyrimethamine).

Embazin

Aggarwal, C. K.; et al., 1978, Indian Vet. J.,
v. 55 (12), 952-957
bifuran, embazin, effect of coccidiostats,
antibiotics, and litters (built up vs. fresh)
on broiler chick performance

Sulphaquinoxaline

Gupta, R. C.; and Sud, S. C., 1978, Indian J.
Animal Research, v. 12 (2), 91-96
sulphaquinoxaline, poultry, concentration
levels in blood and urine

Sulphaquinoxaline

Long, P. L.; and Millard, B. J., 1978, Parasit-
ology, v. 76 (1), 1-9
Eimeria grenieri in Numida meleagris (intes-
tine, caeca) (nat. and exper.), life cycle,
reproduction rate, pathogenicity (severe de-
pression of body weight gain), immunity to
reinfection, treatment with sulphaquinoxaline
in drinking water and robenidine in food:
Britain

Sulphaquinoxaline + Amprolium + Ethopabate
(= Pancoxin)

Long, P. L.; and Millard, B. J., 1978, Avian
Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on
oocyst output of various treatment regimens

Sulphaquinoxaline + Amprolium + Ethopabate +
Pyrimethamine (= Supacox)

Long, P. L.; and Millard, B. J., 1978, Avian
Path., v. 7 (3), 373-381
coccidiosis, broiler chickens, effect on
oocyst output of various treatment regimens

Sulphaquinoxaline

McHardy, N., 1978, Ann. Trop. Med. and Parasit-
tol., v. 72 (6), 501-511
Theileria parva- and T. annulata-infected
bovine lymphoblastoid cell cultures used in
in vitro screens to test wide range of
compounds for chemotherapeutic activity

Sulfaquinoxaline

McManus, E. C.; et al., 1979, Exper. Parasitol.,
v. 47 (1), 13-23
Eimeria tenella, chickens (exper.), t-butyl-
aminoethanol alone or in synergistic combina-
tion with sulfaquinoxaline and pyrimethamine,
anticoccidial efficacy, specific reversal of
toxicity for parasite and host by choline and
dimethylaminoethanol

Sulfaquinoxaline

McManus, E. C.; and Rogers, E. F., 1979, Exper.
Parasitol., v. 48 (2), 235-238
Eimeria tenella, chickens, synergistic in-
teraction of sulfaquinoxaline and t-butyl-
aminoethanol

Sulfaquinoxaline -- Continued.

Sulfaquinoxaline

Manuel, M. F.; and Neri, R. A., 1975, Philippine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfaquinoxaline sodium (Noxal)

Manuel, M. F.; and Neri, R. A., 1975, Philippine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfaquinoxaline + Diaveridine (=Darvisul liquid)

Manuel, M. F.; and Neri, R. A., 1975, Philippine J. Vet. Med., v. 14 (1), 106-116
Eimeria tenella, White Leghorn cockerels, efficacy of 7 water-soluble coccidiostats

Sulfaquinoxaline + p-Dimethylaminobenzonitrile (= Nitryl)

Mladenovic, Z.; Movsesijan, M.; and Borojevic, D., 1978, Vet. Glasnik, v. 32 (10), 829-834
Eimeria spp., chickens (exper.), mixed infections, cycostat, nitryl, and stenorol

Pancoxin plus

Mørch, J., 1978, Nord. Vet. Med., v. 30 (6), 253-259
Eimeria spp., chickens (exper.), efficacy of coccidiostats in feed, better production efficiency of medicated groups

Sulfaquinoxaline

Osweiler, G. D.; and Green, R. A., 1978, Vet. and Human Toxicol., v. 20 (3), 190-191
sulfaquinoxaline, death of pups, tentative diagnosis of vitamin K antagonism from drug therapy, case report

Sulfaquinoxaline

Ryley, J. F.; and Hardman, L., 1978, J. Parasitol., v. 64 (5), 878-881
Eimeria acervulina, E. mivati, speciation studies (cross-immunity and drug resistance studies), some immunological relationship was demonstrated but the failure of the 2 organisms to interbreed in the drug resistance studies lends support to status of E. mivati as distinct species

Pancoxin

Schindler, P.; et al., 1979, Poultry Science, v. 58 (1), 23-27
Eimeria spp., broiler chicken pen trials, arprinocid in feed highly effective prophylaxis, comparison with halofuginone, monensin, nicarbazin, and pancoxin: England; France; Germany

Pancoxin

Sherkov, Sh., 1977, Vet. Sbirka, v. 75 (3), 35-38
E[imeria] tenella, pancoxin, chickens raised under conditions of high and low temperatures; influence of thiamine on development of coccidiosis

Sulfaquinoxaline -- Continued.

Sulfaquinoxaline

Sherkov, Sh.; et al., 1978, Vet. Sbirka, v. 76 (6), 39-41
coccidiosis, calves, disinfectants against oocysts, sulfaquinoxaline as chemoprophylactic, elancoban-100 as coccidiostat

Pancoxin plus

Sherkov, S. N.; Leitch, B.; and Kokash, L., [1977], Egypt. J. Vet. Sc., v. 13 (1), 1976, 37-43
Sarcocystis tenella, kittens (exper.), development in intestines, life cycle; attempted parasite suppression using statyl and pancoxin plus

Sulphaquinoxaline

Shirley, M. W., 1978, Ztschr. Parasitenk., v. 57 (1), 83-87
Eimeria mivati and E. mivati var. diminuta strains differing in sensitivity to sulphaquinoxaline and electrophoretic mobility of lactate dehydrogenase crossed; electrophoretic variation of enzymes a further marker for genetic studies

Pancoxin plus

Stoianov, P.; et al., 1978, Vet. Med. Nauki, v. 15 (8), 105-114
E[imeria] tenella, broiler chickens, varied temperature and moisture regimes, blood biochemistry, host resistance, efficacy of pancoxin plus

Sulfaquinoxaline + Pyrimethamine + Amprolium + Ethopabate (= Pancoxin Plus)

Voeten, A. C.; et al., 1978, Tijdschr. Diergeneesk., v. 103 (23), 1284-1289
coccidiosis, broilers, anticoccidials, floor pen trials

Sulphaquinoxaline + Diaveridine (= Saquadil)

Williams, R. B., 1978, Vet. Parasitol., v. 4 (2), 193-197
Isospora mayuri and Eimeria colchici in Pavo cristatus controlled by sulphaquinoxaline and diaveridine in drinking water

Sulfaquinoxaline sodium. See Sulfaquinoxaline.

Sulfathiazole -- Norsulfazole.

Norsulfazole

Avakian, A. A.; et al., 1978, Veterinariia, Moskva (11), 76-77
coccidiosis, chickens (broilers), prophylactic control by various preparations, pharmacococci recommended, treatment economics: Krymsk oblast

Norsulfazole

Evplov, N. N.; and Nazarov, V. G., 1977, Veterinariia, Moskva (6), 65-66
Eimeria spp., calves, cnemococcide effective; compared with biomycin and norsulfazole: Belgorodsk oblast

Sulfathiazole -- Continued.

Sulfathiazole

- Korolkovas, A.; and Pellegrino, J., 1978, Rev. Inst. Med. Trop. S. Paulo, v. 20 (4), 219-223
Schistosoma mansoni, mice and hamsters, exper. drug trials with 30 latent forms of 1,4-naphthylenediamine and naphthoquinone, 4 drugs found to be active

Sulfatolamide -- Supronal (with Sulfamerazine).

Supronal

- Quintero Monasterios, R.; and Castro R., G., 1972, Rev. Obst. y Ginec. Venez., v. 32 (3), 483-488
human urogenital toxoplasmosis, statistics of cases studied because of secondary sterility, clinical management, therapy with daraprim and supronal

Sulfene. See Bithionol.

- 2,2'-Sulfinylbis(4-chloro-6-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

- 2,2'-Sulfinylbis(4,6-dichlorophenol). See Bithionol.

Sulfirame -- Monosulfiram; Tetmosol; Tetraethylthiuram monosulphide.

Monosulfiram (Tetmosol)

- Belda, W., 1975, Rev. Brasil. Clin. e Terap., v. 4 (8), 309-312
human scabies, increasing incidence, successful management with monosulfiram: Brazil

Monosulfiram (Tetmosol)

- Jardim, M. L.; and Motta, G. M. A., 1976, Rev. Brasil. Med., v. 33 (10), 361-362
scabies, human, therapeutic trials with monosulfiram, 95% cure rate: Pernambuco, Brazil

Monosulfiram (Tetraethylthiuram monosulphide; Tetmosol)

- Monaco, D.; and Lima, E., jr., 1975, Rev. Brasil. Med., v. 32 (10), 687-690
Sarcoptes scabiei, humans, clinical trials with monosulfiram, good results, few side effects: Estado de Sao Paulo, Brasil

Tetraethylthiuram monosulphide (Tetmosol)

- Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia muscili, Myocoptes musculinus, conventional mouse colony, acaricides

Tetmosol

- Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Sulfirame -- Continued.

Monosulfiram (Tetmosol)

- Robinson, D. D., 1976, Rev. Brasil. Med., v. 33 (5), 167-169
human scabies and pediculosis, clinical trials using monosulfiram to treat a rural population

Tetmosol

- Vihan, V. S.; Sinha, N. K.; and Sahni, K. L., 1979, Indian Vet. Med. J., v. 3 (3), 197-198
Sarcoptes scabiei, sheep, tetmosol and benzene hexachloride compared

- Sulfisomidine + Sulfadimethoxine + Trimethoprim + Sulfisoxazole (= Pacprim; Sulfanilamide complex)
Lee, H. S.; Lee, H. B.; and Moon, M. H., 1979, Korean J. Animal Sc., v. 21 (4), 281-288
Besnoitia besnoiti, goats (exper.), antimony potassium tartrate, pacprim

Sulfisoxazole -- Gantrisin; Pacprim (with Sulfadimethoxine, Sulfisomidine and Trimethoprim); Sulfanilamide complex (with Sulfadimethoxine, Sulfisomidine and Trimethoprim); Sulfisoxazole diolamine; Sulphafurazole.

Sulfisoxazole diolamine (Gantrisin)

- Jaffe, J. J.; et al., 1978, J. Parasitol., v. 64 (2), 193-197
Brugia pahangi-infected Aedes aegypti treated with sulfisoxazole and methotrexate singly or in combination, average number of infective larvae recovered was half of that recovered from controls and many larvae recovered were small and sluggish, most likely mode of action is inhibition of synthesis de novo of dihydrofolate in either parasite or more likely in mosquito host (leading to folate-related nutritional deficiencies inimical to normal filarial larval development)

Sulphafurazole + Trimethoprim

- Kyllerman, M.; and Strannegard, O., 1979, Arch. Dis. Childhood, v. 54 (4), 326-327
[Letter]
toxoplasmosis, 5-year-old boy, hemiplegia, disease most likely acquired from cat which had high serum levels of toxoplasma antibodies, trimethoprim-sulphafurazole treatment

- Sulfisoxazole + Sulfadimethoxine + Trimethoprim + Sulfisomidine (= Pacprim; Sulfanilamide complex)
Lee, H. S.; Lee, H. B.; and Moon, M. H., 1979, Korean J. Animal Sc., v. 21 (4), 281-288
Besnoitia besnoiti, goats (exper.), antimony potassium tartrate, pacprim

Sulfisoxazole diolamine. See Sulfisoxazole.

Sulfonamide

- Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230
Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

- 2,2'-Sulfonylbis(4-chloro-6-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives
- Sulfonylbis(2,3,6-trichlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan.
J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives
- Sulformethoxine. See Sulfadoxine.
- Sulfoxide
Bankov, D., 1976, Vet. Med. Nauki, v. 13 (10), 28-36
cestodes of sheep, drug trials; Stilesia globipunctata, tested several diagnostic methods with unfavorable results
- Sulfur ointment
Fletcher, K. C., 1978, J. Am. Vet. Med. Ass., v. 173 (9), 1231-1232
Notoedres cati in Uncia uncia (skin), intense pruritus with resulting alopecia, lime-sulfur solution and sulfur ointment healed all lesions without complications: zoo
- Sulfur powder
Needham, J. R., 1978, J. Inst. Animal Techn., v. 29 (1), 1-15
Myobia musculi, Myocoptes musculinus, conventional mouse colony, acaricides
- Sulmet. See Sulfamethazine.
- Sulph- See also Sulf-
- Sulphachloropyrazine. See Sulfachloropyrazine.
- Sulphacombine. See Diaveridine or Sulfamethazine.
- Sulphadiazine. See Sulfadiazine.
- Sulphadimidine. See Sulfamethazine.
- Sulphadoxine. See Sulfadoxine.
- Sulphafurazole. See Sulfisoxazole.
- Sulphamethoxazol. See Sulfamethoxazole.
- Sulphaquinoxaline. See Sulfaquinoxaline.
- Sulphene. See Bithionol.
- Sumithion. See Fenitrothion.
- Sumithion 50 EC. See Fenitrothion.
- Supacox. See Amprolium or Ethopabate or Pyrimethamine or Sulfaquinoxaline.
- Superphosphate. See Calcium phosphate.
- Supona. See Chlorfenvinphos.
- Supone. See Chlorfenvinphos.
- Suposan. See Chlorfenvinphos.
- Supronal. See Sulfamerazine or Sulfatolamide.
- Suramin -- Antrypol; Bayer 205; Germanin; Naganol; Suramin sodium; Suraminum natricum.
- Suramin
Buyst, H., 1975, Ann. Soc. Belge Med. Trop., v. 55 (2), 95-104
Trypanosoma rhodesiense, human, specific treatment with suramin and mel B, adjuvant antimalarial treatment with chloroquine and proguanil; modifications of sleeping sickness therapy advocated on physio-pathological and epidemiological grounds: Luangwa Valley, Zambia
- Suramin
Buyst, H., 1977, Ann. Soc. Belge Med. Trop., v. 57 (4-5), 201-212
sleeping sickness in children, epidemiologic data, age distribution, various therapeutic regimens, usefulness of antimalarial therapy administered simultaneously: Isoka, Zambia
- Suramin
Ercoli, N., 1978, Proc. Soc. Exper. Biol. and Med., v. 157 (3), 397-401
Trypanosoma venezuelense, rats, comparison of "true" and "false" prophylaxis using pentamidine, suramin and 98/202
- Suramin (Antrypol)
Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested
- Suramin-Tryparsamide complex
Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44
Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested
- Suramin sodium
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Suramin -- Continued.

Suramin

Gutteridge, W. E.; Dave, D.; and Richards, W. H. G., 1979, *Biochim. et Biophys. Acta*, v. 582 (3), 390-401

Kinetoplastida spp., Plasmodium spp., conversion of dihydroorotate to orotate, mechanism of reaction different in these 2 groups of protozoa, possible target of chemotherapeutic attack

Suramin

Jaffe, J. J.; and Chrin, L. R., 1978, *J. Parasitol.*, v. 64 (4), 661-668

Brugia pahangi-infected and normal *Aedes aegypti*, methylenetetrahydrofolate dehydrogenase (MTHFD) and reductase (MTHFR) activity, change in folate metabolism with advanced infections; suramin inhibited MTHFR activity but not MTHFD; MTHFR activity detected in crude extracts of adult parasites differed from that in mosquitoes

Suramin

James, D. M., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (5), 471-476

Trypanosoma congolense, *T. brucei*, rats, mice, prophylactic activity of various trypanocides complexed with dextran, comparison with uncomplexed drugs and with suramin-complexed drugs

Suramin sodium

Laemmler, G.; and El-Gendi, A. Y. I., 1978, *Ztschr. Parasitenk.*, v. 58 (1), 55-73

Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Suramin

McHardy, N., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (6), 501-511

Theileria parva- and *T. annulata*-infected bovine lymphoblastoid cell cultures used in vitro screens to test wide range of compounds for chemotherapeutic activity

Suramin

Shibuya, T.; et al., 1978, *Japan. J. Exper. Med.*, v. 48 (5), 411-418

Litomosoides carinii in *Sigmodon hispidus*, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Naganol (Suraminum natricum; Bayer 205)

Shien, Y. S., 1979, *J. Chinese Soc. Vet. Sc.*, v. 5 (1), 19-22

Trypanosoma evansi, goats (exper.), immunosuppression of response to *Brucella abortus* vaccinations, reversal after naganol treatment

Suramin (Germanin)

Tanaka, H.; et al., 1977, *Japan. J. Exper. Med.*, v. 47 (4), 315-317

Litomosoides carinii-infected cotton rats, improved method for intrapleural injection of anti-filarial drugs to evaluate macrofilaricidal action

Suramin -- Continued.

Suramin

Thylefors, B.; and Rolland, A., 1979, *Bull. World Health Organ.*, v. 57 (3), 479-480

Onchocerca volvulus, suramin-treated patients, increased incidence of optic atrophy

Suramin

Traub, N.; et al., 1978, *East African Med. J.*, v. 55 (10), 477-481

Trypanosoma brucei rhodesiense, human congenital, fatal infection in mother, infant successfully treated with suramin and mel-B, immunoglobulin levels at diagnosis, during treatment, and post-treatment, case reports: Zambia

Suramin + Glycerol

Van Der Meer, C.; Versluijs-Broers, J. A. M.; and Opperdoes, F. R., 1979, *Exper. Parasitol.*, v. 48 (1), 126-134

Trypanosoma brucei brucei, rats, treatment with salicylhydroxamic acid + glycerol and suramin + glycerol

Suramin

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Suramin

Worms, M. J.; and Hawking, F., 1978, *Tr. Roy. Soc. Trop. Med. and Hyg.*, v. 72 (5), 548-549

Litomosoides carinii-infected *Sigmodon hispidus*, effects of suramin, homidium bromide, quinapyramine, diminazene, and isometamidium after observation period of more than 5 weeks

Suramin sodium. See Suramin.

Suraminum natricum. See Suramin.

Sylvinite

Antonenko, A. E., 1975, *Vet. Nauka--Proizvod.*, Trudy, Minsk, v. 13, 121-126

nematode eggs, coccidian oocysts, non-litter pig manure, substances tested as disinfectants

Synanthic. See Oxfendazole.

Systemex. See Oxfendazole.

TAC. See Pararosaniline.

Taeniafugin

Corba, J.; et al., 1977, Veterinarstvi, v. 27 (11), 516-517

Moniezia sp., sheep, Taeniafugin treatment

Taktic. See Amitraz.

TAP

Slin'ko, V. G., 1979, Veterinaria, Moskva (5), 44-46

Sarcoptes suis, pigs, association with necrosis of ear helix, possible secondary bacterial infection; treatment with chlorophos or TAP

Tartar emetic. See Antimony potassium tartrate.

Task. See Dichlorvos.

Taurine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Ta-Verm. See Piperazine.

TBS. See Bromsalans.

Teclozan -- Am 13,146; N,N-Bis (dichloroacetyl)-N,N-bis(2-ethoxy-ethyl)-1,4-bis (aminomethyl) benzene; Falmonox; Win 13.146.

Teclozan (Falmonox; Win. 13.146)

Fleiss, J. R., 1973, Rev. Brasil. Med., v. 30 (7), 461-469

amoebiasis, human intestinal, teclozan for both diagnostic and therapeutic purposes in instances of diagnostic problems and persistent chronic infections, case reports

Falmonox (Teclozan; Am 13,146)

Zafar, M. H.; et al., 1976, Rawal Med. J., v. 5 (4), 140-142

Entamoeba histolytica, human, amoebic dysentery, clinical trials with teclozan, 94% cure rate with few side effects

Tego

Waller, T., 1979, Lab. Animals, v. 13 (3), 227-230

Encephalitozoon cuniculi, survival of spores after exposure to various temperatures and disinfectants; growth-inhibition effect of drugs in cell cultures

Telmezan

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176

Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Telmin. See Mebendazole.

Telmin RLT Sheep Drench. See Mebendazole.

Temephos -- Abate; Lypor 20; Tetramethyl 4,4'-(phenylenethiophenylene) di-(phosphorothionate); 0,0'-Thiodi-p-phenylene) 0,0,0', 0'-tetramethyl phosphorothioate.

Abate

Cole, M. M.; et al., 1973, J. Econom. Entom., v. 66 (1), 118-119

Pediculus h. humanus, strain from Burundi, resistance to malathion and 6 other insecticides

Temephos (Lypor 20)

Kettle, P. R.; and Lukies, J. M., 1979, N. Zealand Vet. J., v. 27 (4), 78-79

Linognathus vituli, cattle, pour-on formulations of phosmet, methidathion, chlorpyrifos, and temephos: Kaitoke, near Upper Hutt, New Zealand

Terenol. See Resorantel.

Terephthalic acid -- WR-74,106.

WR-74,106

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737

Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

Teroxalene -- A-16,612.

A-16,612

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46

Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Terradoxyn. See Doxycycline.

Terragluclin. See Oxytetracycline.

Terramycin. See Oxytetracycline.

Terramycin 100. See Oxytetracycline.

- Terramycin injectable long acting. See Oxytetracycline.
- Terramycin injectable solution. See Oxytetracycline.
- Terramycin/LA. See Oxytetracycline.
- Terramycin/LA (T-200). See Oxytetracycline.
- Testosterone propionate
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- Tetmosol. See Sulfirame.
- Tetraacetyl-ethylenediamine
Kolesnikov, V. I., 1977, *Sborn. Nauch. Rabot SibNIVI* (28), 143-146
Fasciola hepatica, rabbits, new anthelmintics tested, phenacetine highly effective
- 4,4',6,6'-Tetrabromo-2,2'-biphenyldiolmono(dihydrogenphosphate). See Bromophenophos.
- Tetracap. See Tetrachloroethylene.
- meso-Tetra-(p-carboxyphenyl)-porphine (TPPC.)
Meshnick, S. R.; et al., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis
- Tetrachlor- See also Tetrachloro-
- 3,5,3',5'-Tetrachlor-2,2'-dioxydiphenylsulfide
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,5,3',5'-Tetrachlor-2,2'-dioxydiphenylsulfone
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- 3,5,3',5'-Tetrachlor-2,2'-dioxydiphenylsulfoxide
Kozhokaru, A. F.; and Topaly, V. P., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 178-183
Fasciola hepatica total and mitochondrial lipids, ox brain total lipids, and ox heart mitochondrial lipids as sources of bimolecular phospholipid membranes in which proton conductivity induced by aromatic sulfides, sulfoxides, and sulfones correlated with their fasciolicidal effects and permitted toxicity evaluation
- Tetrachlorethylene. See Tetrachloroethylene.
- Tetrachloro- See also Tetrachloro-
- Tetrachloroethylene -- Tetracap; Tetrachloroethylene.
- Tetracap (Tetrachlor ethylene)
Cherian, Z.; Jose, M. P.; and Jayakumar, K. M., 1977, *Kerala J. Vet. Sc.*, v. 8 (1), 71-72
ancylostomiasis in mongrel dogs, clinical treatment trials: decaris highly effective without toxicity, tetracap reduced severity of infection, furoxone had no effect
- Tetrachloroethylene
Freire, J. dos S.; Bezerra, V. C.; and Prado, A. A., 1973, *Rev. Patol. Trop.*, v. 2 (2), 155-158
human ancylostomiasis, comparative therapeutic trials using jonit and tetrachloroethylene; tetrachloroethylene gave higher cure rates: Brazil
- Tetrachloroethylene
Goldsmith, R. S., 1978, *South. Med. J.*, v. 71 (12), 1513-1515
Metagonimus yokogawai, American woman traveling in the Orient, case report, chronic diarrhea treated unsuccessfully with hexylresorcinol, cure with tetrachloroethylene: California
- Tetrachlorethylene
Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, *Rev. Brasil. Med.*, v. 31 (11), 791-794
human soil-transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective
- Tetrachlorethylene
Migasena, S.; Suntharasamai, P.; and Harinatsuta, T., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (2), 199-200
Necator americanus, humans, treatment trials comparing efficacy of mebendazole, tetrachlorethylene and pyrantel pamoate: Bangkok
- Tetrachlorethylene
Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (1), 55-58
Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (Mesocricetus auratus)

Tetrachloroethylene -- Continued.

Tetrachlorethylene

Soelberg Sørensen, P.; and Melgaard, B., 1971, Scand. J. Infect. Dis., v. 3 (1), 65-69
Necator americanus, Ancylostoma duodenale, human hookworm anemia, comparative therapeutic study using tetrachlorethylene and iron singly and in combination, use of tetrachloroethylene alone was recommended for mass therapy with the additional use of iron when objective clinical signs of anemia were present: Koraput, Orissa State, India

Tetrachloroethylene

Swartzwelder, J. C.; et al., 1972, Rev. Biol. Trop., v. 20 (2), 295-307
tetrachloroethylene administered prophylactically to population of coffee plantation to reduce incidence of hookworm, piperazine hexahydrate administered simultaneously reduced incidence of Ascaris lumbricoides during initial phase of treatment only: Costa Rica

Tetrachlorethylene

Vaughan, J. P.; et al., 1977, Trop. and Geogr. Med., v. 29 (4), 369-373
hookworm and iron deficiency anemia in adults and children, therapy with tetrachlorethylene and oral iron, poor response to therapy by children suggests that malaria was causative factor: Tanzania

N-Tetrachloroethylthio-4-cyclohexene-1,2-dicarboximide

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Tetrachloroisophthalonitril

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

3,5,3',4'-Tetrachlorosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Tetrachlorvinphos -- 2-Chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate; Gardona; Rabon; Shell SD-8447; Stirofos.

Stirofos

Ahrens, E. H.; and Cocke, J., 1978, J. Econom. Entom., v. 71 (5), 764-765
Amblyomma maculatum, cattle, comparative test with insecticide-impregnated ear tags; longevity test to determine whether ranchers could tag cattle in early spring resulted in only marginally satisfactory control

Tetrachlorvinphos -- Continued.

Gardona

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory tests of insecticides

Stirofos

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Stirofos

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586
Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Stirofos

Hall, R. D.; Townsend, L. H., jr.; and Turner, E. C., jr., 1978, J. Econom. Entom., v. 71 (2), 315-318
Ornithonyssus sylviarum, laboratory and field tests to compare effectiveness of organophosphorous, carbamate, and synthetic pyrethroid acaricides, carbaryl most toxic to mites, ectiban permethrin and SD-43775 also effective; mites displayed tolerance to malathion

Rabon

Miller, R. W.; and Gordon, C. H., 1973, J. Econom. Entom., v. 66 (1), 135-138
rabon, feeding to dairy cows over extended periods, no adverse effects on general health or reproductive performance, negligible milk and tissue residues

Tetrachlorvinphos (Rabon, Gardona)

Noblet, G. P., 1978, Am. J. Vet. Research, v. 39 (6), 1033-1036
gastrointestinal nematodes, beef cattle, tetrachlorvinphos ineffective feed additive

Stirofos

Rawlins, S. C.; and Mansingh, A., 1979, J. Econom. Entom., v. 72 (3), 423-427
Boophilus microplus, bioassays of acaricidal residues on grass surfaces, greenhouse and pasture studies

Stirofos (Rabon)

Wilson, N. L.; Huston, J. E.; and Davis, D. I., 1978, Southwest. Vet., v. 31 (3), 197-199
Haematobia irritans, cattle, stirofos impregnated ear tags gave excellent control, weight gain not influenced by treatment: McCulloch County, Texas

Tetracosactide

Najar, H. C. F., 1978, Rev. Brasil. Med., v. 35 (3), 205-207
human scabies resulting in severe pruritis, clinical trials of milian ointment alone and combined with tetracosactide; combined therapy more successful with total eradication in 5 days

Tetracycline -- Achromycin; Codrinal (with β -Toluenesulfonyl- β -methoxy-ethyl urethane sodium); Contramibial (with Chloroquine and Diiodo-hydroxyquin); Steclin; Tetracycline hydrochloride; Tetracyn; WR 6,527.

Tetracycline

Austerman, J. W., 1979, Vet. Med. and Small Animal Clin., v. 74 (7), 954
haemobartonellosis, dog, case report, tetracycline

Tetracycline

Boggs, C. H.; and Chakravorty, R. C., 1979, Virginia Med., v. 106 (9), 674-675
Entamoeba histolytica, 56-year-old male, necrotizing amebic colitis and perforated colon, case report, successful treatment with surgery, metronidazole, tetracycline: Virginia

Tetracycline hydrochloride

Fuchs, P., 1978, Deutsche Med. Wchnschr., v. 103 (3), 97-98
Entamoeba histolytica, humans, pathogenicity, efficacy and toxicity of various drugs, recommended treatment for various forms of amoebiasis

Tetracycline HCl

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Contramibial

Ingelet, B., 1974, Rev. Infirm. Afrique Noire (26), 17-19
human amoebic colitis, clinical trials testing contramibial given orally over a 4-day period, good therapeutic results

Tetracycline hydrochloride (Tetracycline)

Kazim, M.; Puri, S. K.; and Dutta, G. P., 1979, Chemotherapy, v. 25 (4), 222-226
Plasmodium gallinaceum, chicks (exper.), minocycline and doxycycline, blood schizonticidal activity compared with that of known antibiotics, both more effective than oxytetracycline and tetracycline in controlling acute infection

Tetracycline (Tetracyn)

Lee, K. K.; et al., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (2), 217-220
Naegleria fowleri, in vitro susceptibility to selected antimicrobial agents singly and in combination

Tetracycline

Markowitz, S. M.; et al., 1978, Am. J. Path. (436), v. 92 (3), 733-743
Acanthamoeba castellanii, mice (exper.), pretreated with methylprednisolone or tetracycline, increased host mortality due to depressed host immunity; potentially pathogenic role for naturally occurring Acanthamoeba sp. in immunosuppressed humans

Tetracycline -- Continued.

Steclin

Mishra, A. K.; and Sharma, N. N., 1979, Trop. Animal Health and Prod., v. 11 (4), 222-226
Anaplasma marginale, calves (exper.), comparative efficacy of several drugs

Tetracycline + Amphotericin B

Moggian, G.; Tamburini, E.; and Visona, E., 1975, Quad. Clin. Ostet. e Ginec., v. 30 (4), 143-154
T[richomonas] vaginalis, human vulvovaginitis, clinical trials comparing mepartricin with amphotericin B combined with tetracycline in topical creams, good results

Tetracycline (WR 6,527)

Rane, D. S.; and Kinnamon, K. E., 1979, Am. J. Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in mice, development of high volume tissue schizonticidal drug screen based upon mortality of infected mice

Tetracycline + Amphotericin B

Ruggeri, E., 1976, Quad. Clin. Ostet. e Ginec., v. 31 (4), 211-223
Trichomonas vaginalis, vaginal trichomoniasis, oral mepartricin showed significant therapeutic superiority when compared in clinical trials with patients who received nimorazole therapy orally or with patients treated with a vaginal cream containing amphotericin B and tetracycline

Codrinal

Singh, J.; and Hussain, O., 1978, Indian Vet. J., v. 55 (1), 56-60
Eimeria tenella, chicks (exper.), amprolium provided better protection than codrinal, both drugs interfered to some extent with development of immunity

Achromycin

Sinha, R. P.; and Dubey, R. K., 1978, Indian Vet. J., v. 55 (5), 372-376
Anaplasma marginale outbreak in non-pre-immune Jersey cattle imported from United States and Denmark to farm in Bihar, clinical symptoms and pathological findings, epizootiological factors responsible for outbreak (high ambient temperature, stress of vaccination for rinderpest virus, presence of tick vectors), control achieved through chemotherapy of sick and healthy animals, removal of vectors, and housing in cool sheds; outbreaks in exotic herds could be avoided if cattle were imported in early winter: India

Tetracycline

Sucharit, S.; et al., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (1), 55-59
Brugia pahangi in Aedes togoi, effect of tetracycline phosphate complex on parasite development, distribution within host, and size and mosquito mortality

Tetracycline

Thong, Y. H.; Rowan-Kelly, B.; and Ferrante, A., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (3), 336-337
Naegleria fowleri-infected mice, amphotericin B potentiated by tetracycline shows promise in treating meningoencephalitis

Tetracycline -- Continued.

Tetracycline

Watson, A. D. J.; Farrow, B. R. H.; and Hoskins, L. P., 1978, *Austral. Vet. Practitioner*, v. 8 (3), 129-132
Haemobartonella felis, cats (nat. and exper.), thiacetarsamide sodium therapy; prednisolone, tetracycline, and chloramphenicol given sequentially

Tetracycline

Wilson, A. J.; et al., 1979, *Austral. Vet. J.*, v. 55 (2), 71-73
Anaplasma marginale, cattle and splenectomised calves (exper.), long-acting formulation of tetracycline compared with standard tetracycline and imidocarb dipropionate

Tetracycline hydrochloride. See Tetracycline.

Tetracyn. See Tetracycline.

Tetraethylthiuram disulfide. See Disulfiram.

Tetraethylthiuram monosulphide. See Sulfirame.

Tetrafinol. See Carbon tetrachloride.

 Δ^8 -Tetrahydrocannabinol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^8 -tetrahydrocannabinol and other cannabinoids

 Δ^9 -Tetrahydrocannabinol

Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, *Antimicrob. Agents and Chemotherapy*, v. 16 (5), 674-679
Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

trans-1,4,5,6-Tetrahydro-2-(3-hydroxystyryl)-1-methyl pyrimidine. See Oxantel.

trans-1,4,5,6-Tetrahydro-2-(trans-3-hydroxystyryl)-1-methyl pyrimidine. See Oxantel.

(--)-2,3,5,6-Tetrahydro-6-phenylimidazo(2,1-b)thiazol. See Tetramisole.

1-2,3,5,6-Tetrahydro-6-phenylimidazo (2,1-b)thiazole hydrochloride. See Tetramisole.

Tetramethyl 4,4'-(phenylenethiophenylene) di(phosphorothionate). See Temephos.

meso-Tetra-(p-N-methylpyridyl)-porphine

Meshnick, S. R.; et al., 1978, *J. Pharmacol. and Exper. Therap.*, v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Tetramethylthionine chloride. See Methylene blue.

Tetramisole -- Ascaridil; Brot; Citarin;

Citarin-L; Citarin-L spot on; Concurat; Decaris; Drofenit; Galinid; Ketrax; Laevamisole; Levamisole; Levamisole chlorhydrate; Levamisole hydrochloride; Levamisole phosphate; Levasole; Levasole Injection; Levasole tablets; Levo-tetramisole; Nemicide; Nemicide-L 15; Nilverm; Nilzan (with Oxyclozanide); R-8299; Ripercol; Ripercol-L; (--)-2,3,5,6-Tetrahydro-6-phenylimidazo(2,1-b)thiazol; 1-2,3,5,6-Tetrahydro-6-phenylimidazo (2,1-b)thiazole hydrochloride; dl-Tetramisole; L-Tetramisole; Tetramisole chlorhydrate; Tetramisole cyclamate; Tetramisole hydrochloride; Tetramisole, Laevo isomer; Tetramisole sulfamidine; Tetramisolum; Tramisol; L-Tramisol.

Citarin

Abdel Rahman, M. S.; et al., 1978, *Vet. Med. J.*, Giza, v. 24 (24), 1976, 39-48
 gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt

Concurat

Abdel Rahman, M. S.; et al., 1978, *Vet. Med. J.*, Giza, v. 24 (24), 1976, 39-48
 gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt

Tetramisole (Nilverm)

Altaif, K. I., 1979, *Trop. Animal Health and Prod.*, v. 11 (4), 241-245
 helminths, Awassi sheep, tetramisole, rafoxanide, body weight gains, wool growth: Iraq

Tetramisole hydrochloride (R-8299)

Amato Neto, V.; et al., 1969, *Rev. Inst. Med. Trop. S. Paulo*, v. 11 (4), 294-297
Ascaris lumbricoides, human, clinical trials testing efficacy of tetramisole administered orally: Brazil

Levamisole (Nilverm)

Anderson, N.; and Lord, V., 1979, *Austral. Vet. J.*, v. 55 (4), 158-162
Ostertagia ostertagi, *Trichostrongylus axei*, cattle, fenbendazole, oxfendazole, and levamisole compared

L-Tetramisole (Levamisole)

Arambulo, P. V. III; Lacuata, A. Q.; and Ocampo, W. G., 1975, *Philippine J. Vet. Med.*, v. 14 (1), 156-162
Dirofilaria immitis-infected dogs, clinical trials of L-tetramisole against microfilariae

Tetramisole -- Continued.

Levamisole hydrochloride

Atwell, R. B.; et al., 1979, Austral. Vet. J., v. 55 (6), 292-294 [Letter]
 Dirofilaria immitis, dogs, acute haemolytic anaemia suspected to have been induced by levamisole hydrochloride

Levamisole hydrochloride

Atwell, R. B.; and Baldock, C., 1979, J. Small Animal Practice, v. 20 (5), 299-302
 Dirofilaria immitis, dogs, oral treatment with levamisole hydrochloride, pilot trial

Levamisole hydrochloride (Levasole tablets)

Atwell, R. B.; Carlisle, C.; and Robinson, S., 1979, Austral. Vet. J., v. 55 (11), 531-534

Dirofilaria immitis adults, dogs, levamisole hydrochloride: Brisbane area

Tetramisole

Baqui, A.; and Ansari, J. A., 1976, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 25 (5), 409-414

Setaria cervi, white rats, hetrazan, thiazabendazole, tetramisole

Levamisole

Bennet, E. M.; Behm, C.; and Bryant, C., 1978, Internat. J. Parasitol., v. 8 (6), 463-466
 Mesocostoides corti, mice (infected, injected with dead larvae previous to infection, or irradiated), effects of mebendazole and levamisole alone or together on tetrathyridia, concluded that anthelmintic efficacy of mebendazole depends on its anthelmintic activity supplemented by host's immune response and that levamisole stimulates the latter

Levamisole

Bernberg, H. C.; Clarke, V. de V.; and Gelfand, M., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (2), 233-234
 Dipetalonema perstans-like unshathed microfilarial infection, humans, combined treatment with levamisole and mebendazole: Rhodesia

Nilverm (Ripercol; Citarin)

Bezubik, B.; Sinski, E.; and Swietlikowski, M., 1970, Acta Parasitol. Polon., v. 18 (27-41), 441-451
 gastrointestinal nematodes, Strongyloides papillosus, Dictyocaulus filaria, Moniezia sp., seasonal dynamics in different age classes of sheep in relation to nilverm treatment

Tetramisole cyclamate (Brot)

Biagi F., F.; and Navarrete C., E., 1969, Rev. Inst. Med. Trop. S. Paulo, v. 11 (1), 57-61
 Ascaris lumbricoides, humans, clinical trials testing efficacy of tetramisole, good results without significant side effects

Tetramisole (Decaris)

Bomb, B. S.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 110
 mixed hookworm and roundworm infection, man, treatment with tetramisole resulted in transient optic neuritis

Tetramisole -- Continued.

Tetramisole

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
 antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Levamisole (Nemicide)

Burnie, N., 1979, Vet. Rec., v. 104 (25), 585-586 [Letter]
 Aelurostrongylus larvae, cat (feces), paroxysmal coughing, levamisole, case history

Levamisole

Butler, P. G., 1978, J. Trop. Med. and Hyg., v. 81 (11), 221-224
 Leishmania tropica, human chronic form, levamisole, good therapeutic response in clinical trials: Saudi Arabia

Levamisole HCl (Ripercol-L)

Buys, S. B.; and van der Made, H. N., 1977, J. South African Vet. Ass., v. 48 (4), 284 [Letter]
 toxicity of levamisole in psittacine birds, reactions to parenteral administration not much more drastic than from oral dosing of tetramisole

Levamisole

Callinan, A. P. L.; and Cummins, L. J., 1979, Austral. Vet. J., v. 55 (8), 370-373
 nematodes, cattle (nat. and exper.), efficacies of various anthelmintics against adult and larval stages: western Victoria

Levamisole phosphate (Ripercol L)

Calverley, A. H., 1978, Proc. 23. Ann. Conv. Am. Ass. Equine Practitioners (Vancouver, British Columbia, Dec. 3-7, 1977), 363-365
 heaves caused by lungworms, horses, levamisole phosphate

Levamisole (Nilverm)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
 Haemonchus contortus, Trichostrongylus colubriformis, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Tetramisole

Chaia, G.; and da Cunha, A. S., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (2), 152-160
 intestinal helminths, children, efficacy of anthelmintic therapy: Vila Sao Vicente, Belo Horizonte

Levamisole hydrochloride (Ripercol)

Chaikin, R. J., 1979, Canine Pract., v. 6 (3), 32, 35-37
 Dirofilaria immitis, dogs, efficacy of levamisole as a simultaneous microfilaricide/adulticide

Decaris (Tetramisole)

Cherian, Z.; Jose, M. P.; and Jayakumar, K. M., 1977, Kerala J. Vet. Sc., v. 8 (1), 71-72
 ancylostomiasis in mongrel dogs, clinical treatment trials: decaris highly effective without toxicity, tetracap reduced severity of infection, furoxone had no effect

Tetramisole -- Continued.

Nilverm (Tetramisol hydrochloride)

Chroust, K., 1976, Acta Vet. Brno, v. 45 (4), 251-262
gastrointestinal nematodes, lungworms, sheep, economic aspects of treatment with nilverm

Levamisole hydrochloride

Coles, G. C.; Briscoe, M. G.; and Simpkin, K. G., 1979, Vet. Rec., v. 105 (20), 470
Haemonchus contortus, Trichostrongylus colubriformis, benzimidazole resistant and susceptible strains, lambs, levamisole activity

Tetramisole chlorhydrate

Costa, J. O.; et al., 1977, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 29 (2), 171-178
Strongylidae, calves, chlorhydrate of tetramisole, weight gain in treated and control groups not influenced by Strongylidae infection, low worm burden in both groups

Levamisole

Craig, T. M.; and Bell, R. R., 1978, Am. J. Vet. Research, v. 39 (6), 1037-1038
gastrointestinal nematodes, calves, natural infection, fenbendazole compared with levamisole, controlled experiment, varied results

Levamisole (Nilverm)

Cummins, L. J.; and Callinan, A. P. L., 1979, Vet. Rec., v. 104 (4), 77-78
O[stertagia] ostertagi in cattle, levamisole, combined treatment with human chorionic gonadotrophin did not improve anthelmintic activity against inhibited 4th stage larvae and lessened effectiveness against adults and developing larvae: western Victoria, Australia

Levamisole

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Nilverm

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Levamisole

Del Principe, D.; et al., 1979, Research Commun. Chem. Path. and Pharmacol., v. 25 (2), 307-317
levamisole, in vitro effect on oxygen consumption and survival of platelets

Levamisole (Nilverm)

Donald, A. D.; et al., 1979, Vet. Parasitol., v. 5 (2-3), 205-222
gastrointestinal nematodes with major emphasis on Ostertagia ostertagi, beef cattle, levels of infection and effects on live-weight gain, effects of pasture type (phalaris vs. lucerne) and stocking rate, effects of anthelmintic treatment, 4-year experiment: Canberra, Australia

Tetramisole -- Continued.

Levamisole (Nemicide L 15)

Dorchies, P.; Franc, M.; and Ducos de Lahitte, J., 1979, Rev. Med. Vet., Toulouse, v. 130 (3), 417-418, 421-425
nematodes, horses, levamisole administered per os and intramuscularly, toxicity; efficacy compared with thiabendazole

Levamisole

Downey, N. E., 1978, Vet. Rec., v. 103 (19), 427-428
gastrointestinal nematodes, calves, efficacy of albendazole, compared with levamisole: Ireland

Nilverm

Dvorakova, L.; and Rachac, L., 1978, Veterinarstvi, v. 28 (9), 398-399
Echinuria uncinata, domestic ducks, epizootic occurrence, prevalence, necrosis of proventriculus, nilverm and mebendazole, preventive measures: South Bohemia region

Levamisol (Ketrax)

El Boulaqi, H. A.; et al., 1979, Acta Trop., v. 36 (1), 85-90
Ascaris lumbricoides, human, levamisol treatment, degenerative changes in intestine and reproductive system of worms, worms not expelled by treatment produced only non-viable eggs

Levamisol

Enigk, K.; Dey-Hazra, A.; and Batke, J., 1975, Acta Parasitol. Polon., v. 23 (26-40), 367-372
helminths of geese, treatment (Amidostomum anseris with mebendazol, fenbendazol, levamisol, and pyrantel tartrat; Trichostrongylus tenuis with mebendazol and fenbendazol; Drepanidotaenia lanceolata and Hymenolepis setigera with mebendazol)

Levamisole hydrochloride

Ensley, P. K., 1978, J. Am. Vet. Med. Ass., v. 173 (9), 1246-1248
Paronchocerca ciconiarum in Leptoptilos crumeniferus (blood, right ventricle, pulmonary artery), levamisole hydrochloride, treatment of microfilaremia: San Diego Zoo, shipped from East Africa

Nilverm

Epel'dimov, L. S.; and Plotnikova, L. M., 1977, Sborn. Nauch. Rabot SibNIVI (28), 140-142
[Trichostrongylus], rabbits infected with sheep species as models for anthelmintic study, tests of nilverm, banminth-C, cupric carbonate

Tetramisole (Nilverm)

Fetisov, V. I.; and Maksina, T. P., 1977, Veterinariia, Moskva (8), 67-70
dictyocaulosis, cattle, tetramisole in feed, dosage at therapeutic level has no side effect, no significant anthelmintic action against muelleriasis in sheep

Tetramisole -- Continued.

Tetramisole (Ascaridil)
Fleury, G. C.; Correa, M. O. A.; and Amato Neto, V., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (4), 288-292

Trichostrongylus colubriformis, young child (feces), tetramisole, morphology: first report of this human parasitism in Brazil

Levamisole (Decaris)
Gatti, F.; et al., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 19-31
intestinal nematodes, African schoolchildren, trimetrial administration of levamisole compared with other anthelmintics: Kimwenza, Republique Democratique du Congo

Nilverm (Tetramisole)
Gaur, S. N. S.; and Dutt, S. C., 1979, Indian Vet. J., v. 56 (1), 62-64
Mecistocirrus digitatus, cattle, comparative efficacy of thibendole, nilverm, and helmetaq

Tetramisole
Georgieva, D., 1978, Vet. Sbirka, v. 76 (4), 32-33
anthelmintic treatment, lambs, weight gains

Tetramisole (Nilverm)
Gorodovich, N. M., 1979, Veterinariia, Moskva (2), 44-46
[Mecistocirrus], cattle, tetramisole

Tetramizole chlorhydrate
Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, Rev. Brasil. Med., v. 31 (11), 791-794
human soil-transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Levamisole phosphate (Levasole)
Haigh, J. C., 1979, J. Zoo Animal Med., v. 10 (3), 103-105
levamisole phosphate, Anas platyrhynchos and mixed group of waterfowl, effectiveness in reducing fecal egg count, toxicity

Levamisole
Heath, D. D.; and Lawrence, S. B., 1979, Internat. J. Parasitol., v. 9 (1), 73-76
Taenia crassiceps, rats, mebendazole, single large oral treatment markedly more effective in killing cysts than same amount of drug divided into daily smaller doses, levamisole promoted vigorous host cellular response but did not enhance action of mebendazole

Tetramisole
Iakubovskii, M. V., 1977, Veterinariia, Moskva (8), 72-73
ascariasis, trichuriasis, oesophagostomiasis, swine-breeding farms, tetramisole treatment

Tetramisole
Iakubovskii, M. V., 1979, Veterinariia, Moskva (2), 41-42
ascariasis, trichocephaliasis, oesophagostomiasis, swine, comparative effectiveness and economic value of various drugs: Minsk oblast

Tetramisole -- Continued.

Nilverm
Iakubovskii, M. V.; and Zen'kov, A. V., 1977, Vet. Nauka--Proizvod., Trudy, Minsk, v. 15, 72-78
nematodes, swine raising complexes, combined control measures, sanitation, anthelmintics, suiverm most effective

Citarin-L spot on (Levamisole)
Inderbitzin, F.; and Eckert, J., 1977, Prakt. Tierarzt, v. 58 (8), 579-584
Dictyocaulus viviparus, cattle, 2 formulations of citarin-L spot on (117-V and 111-V), drug trials

Levamisole
Jancloes, M. F.; Cornet, P.; and Thienpont, P., 1979, Trop. and Geog. Med., v. 31 (1), 111-121
intestinal nematodes, human, mass therapy with single oral doses of levamisole: Kisantu area, Republic of Zaire

Nilverm (Tetramisole)
Kadyrov, N. T., 1978, Veterinariia, Moskva (7), 57-58
Delafondia vulgaris, horses, anthelmintics tested; preventive dehelminthization every 15 days recommended for horses on pasture

Levamisole
Kerboeuf, D.; Hubert, J.; and Le Stang, J. P., 1979, Rec. Med. Vet., v. 155 (2), 143-148
Ostertagia ostertagi, cattle, serum pepsinogen levels in relation to worm burden and anthelmintic treatments

Levamisole
Koehler, P.; and Bachmann, R., 1978, Molec. Pharm., v. 14 (1), 155-163
Ascaris suum muscle tissue, comparison of effects of levamisole, thiabendazole, chloroquine, and praziquantel on electron transport in Ascaris muscle submitochondrial particles

Levamisole
Krotov, A. I.; Cherniaeva, A. I.; and Budanova, I. S., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 165-168
Alveococcus multilocularis, white mice, effect of thiabendazole, sarcocystin acridine, levamisole, and mebendazole on larval cyst development

Levamisole
Krubwa, F.; et al., 1974, Ann. Soc. Belge Med. Trop., v. 54 (3), 167-176
intestinal nematodes, heavily infected native population, effect of various regimens of levamisole administered over 1-year period on the prevalence and intensity of infection, results differed but levamisole seemed ideal anthelmintic for mass campaigns: Ndjili, Kinshasa, Zaire

Tetramisolium
Kurkela, P.; and Kaantee, E., 1978, Zentralbl. Vet.-Med., Reihe B, v. 25 (1), 81-87
parasites of domestic Rangifer tarandus tarandus in enclosure conditions, tetramisolium, good results, effective against Oedemagena tarandi larval forms, no harmful side effects: North of Finland

Tetramisole -- Continued.

Levamisole (Citarin-L)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Levamisole (Nilverm)

Le Jambre, L. F., 1979, Austral. Vet. J., v. 55 (2), 65-67
Ostertagia circumcincta, *O. trifurcata*, sheep (exper.), effectiveness of levamisole, thiabendazole, albendazole, and oxfendazole against levamisole-resistant strains

Levamisole (Nilverm)

Le Jambre, L. F.; and Martin, P. J., 1979, Vet. Sc. Commun., v. 3 (2), 153-158
 levamisole resistant *Ostertagia circumcincta* and *O. trifurcata*, sheep, cross resistant to morantel tartrate but not to naphthalophos

Levamisole

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Austral. Vet. J., v. 54 (12), 570-574
Trichostrongylus colubriformis, linear dose responses of selected and unselected strains to thiabendazole, levamisole, and morantel tartrate

Levamisole

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Internat. J. Parasitol., v. 8 (6), 443-447
Ostertagia circumcincta, development of simultaneous resistance to thiabendazole, morantel tartrate, and levamisole, multiple selection associated with increase in *O. trifurcata* in population and increase in larval inhibition

Levamisole (Ripercol L)

Lensink, B. M.; Rijpstra, A. C.; and Erken, A. H. M., 1979, Zool. Garten, n. F., v. 49 (2), 121-126
Ollulanus tricuspis in *Panthera tigris tigris* (vomitus, stomach), clinical symptoms in mother and offspring, treatment with various anthelmintics, complete recovery achieved with levamisole: Artis-Zoo, Amsterdam

Levamisole

Levi, G. C.; et al., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 392-396
 ascariasis, human, clinical trials comparing piperazine, tetramisole (or levamisole) and pyrantel pamoate: Brazil

Tetramisole hydrochloride

Levi, G. C.; et al., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (6), 392-396
 ascariasis, human, clinical trials comparing piperazine, tetramisole (or levamisole) and pyrantel pamoate: Brazil

Nilverm

Lipova, E.; and Zajicek, D., 1979, Veterinarstvi, v. 29 (3), 125-126
Strongyloides papillosus, calves, intensity of infection, clinical aspects, nilverm and nematin treatment

Tetramisole -- Continued.

Nilzan

McEwan, A. D.; and Oakley, G. A., 1978, Vet. Rec., v. 102 (14), 314-315
 panacur, thiabendazole, and nilzan with dye marker added, oral dosing of cattle showed evidence of rumen by-pass, reduced drug efficacy probably resulting from closure of oesophageal groove

Levamisole hydrochloride (Nemicide)

McEwan, A. D.; Oakley, G. A.; and Robinson, M., 1979, Vet. Rec., v. 105 (1), 15-16
Dictyocaulus viviparus, calves (exper.), lung lesions more severe with fenbendazole than with levamisole and certain types more severe in treated calves vs. controls, may be direct result of drug action

Levamisole

Macfarlane, D. G.; and Bacon, P. A., 1978, Brit. Med. J. (6110), v. 1, 407-408
 levamisole-induced vasculitis in patients with immunological disorders

Levamisole

McMahon, J. E., 1979, Ann. Trop. Med. and Parasitol., v. 73 (5), 465-472
Wuchereria bancrofti, humans, preliminary screening of levamisole and amodiaquine for antifilarial activity; some trials combined with diethylcarbamazine

Levotetramisole (Decaris)

Malik, G. Q.; et al., 1976, Rawal Med. J., v. 5 (11), 90-92
 human intestinal helminths, clinical trials with decaris, suggested use for mass therapy: Multan

Nilverm

Mamaev, N. Kh.; and Davudov, D. M., 1978, Veterinariia, Moskva (8), 69
 [Strongylata], sheep, nilverm granules mixed with feed

Levamisole HCl (Tramisole)

Marti, O. G.; Stewart, T. B.; and Hale, O. M., [1979], J. Parasitol., v. 64 (6), 1978, 1028-1031
 gastrointestinal nematodes, pigs raised under similar management conditions, comparative efficacy of fenbendazole, dichlorvos, and levamisole HCl

Levamisole hydrochloride

Martin, P. J.; and Le Jambre, L. F., 1979, Vet. Sc. Commun., v. 3 (2), 159-164
Ostertagia circumcincta, *O. trifurcata*, larval paralysis as in vitro assay of levamisole and morantel tartrate resistance

Tetramisole

de Melo, H. J. H., 1977, Arq. Escola Vet. Univ. Fed. Minas Gerais, v. 29 (3), 269-277
 helminths, weaned Zebu calves reared extensively on Jaragua grass, different schemes of treatment, effect of anthelmintic treatment on host growth seems to be conditioned to environmental conditions, especially nutrition

Tetramisole -- Continued.

Levamisole

Meltzer, R. S.; et al., 1979, Am. J. Med. Sc., v. 277 (1), 91-98

Strongyloides stercoralis, woman, development of hyperinfection syndrome while on high-dose corticosteroids and following splenectomy, central nervous system involvement, antemortem diagnosis, thiabendazole, levamisole, and mebendazole therapy: Memorial Sloan-Kettering Cancer Center, New York (had traveled in Italy and Sicily)

Levamisole (Decaris)

Miller, M. J.; et al., 1978, South. Med. J., v. 71 (2), 137-140

ascariasis, children throughout the world, levamisole vs. known anthelmintics, levamisole recommended for mass chemotherapy: Iran, Brazil, Mississippi, Louisiana

Tetramisole-hydrochloride

Mirck, M. H.; and Bergsma, R. F., 1979, Tijdschr. Diergeneesk., v. 104 (15-16), 633-634
anthelmintic containing '5-nitro-diphenol' as active principle found actually to contain tetramisole-hydrochloride, unsuitable systemic anthelmintic for horses, brief review

Levamisole

Moenz, M.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (5), 897-904

Ascaris lumbricoides, human, levamisole, analysis of combined results of 10 clinical studies with 1,734 patients

Tetramisole

Mourot, D.; et al., 1979, J. Pharm. Sc., v. 68 (6), 796-797

tetramisole, percent concentration in anthelmintic veterinary formulations, high-pressure liquid chromatography analysis

Levamisole (Ketrax)

Nagaty, H. F.; Mohajeri, M.; and Elahi, R., 1978, J. Trop. Med. and Hyg., v. 81 (10), 195-197

Ascaris lumbricoides, humans, clinical trials with levamisole: Khorasan, Iran

Levamisole

Narasimham, M. V. V. L.; et al., 1978, South-east Asian J. Trop. Med. and Pub. Health, v. 9 (4), 571-575

Wuchereria bancrofti, human, comparison of levamisole, levamisole + mebendazole, and diethylcarbamazine: Calicut, Kerala, India

Nilverm (Tetramisole)

Oripov, A. O., 1978, Veterinariia, Moskva (4), 74-76

Strongylata, sheep, various anthelmintics in granular form tested for mass dehelminthization, no harmful effects, nilverm most effective

Nilverm + Phenasal

Oripov, A. O.; Bekirov, R. E.; and Dzumaev, Z., 1978, Veterinariia, Moskva (12), 60

helminths, dogs, phenasal and nilverm given in feed (sausage form)

Tetramisole -- Continued.

Nilverm

Petrov, Iu. F., 1978, Veterinariia, Moskva (5), 64-66

[Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Levamisole + Diethylcarbamazine

Prod'hon, J.; et al., 1979, Med. Trop., v. 39 (6), 631-635

Onchocerca volvulus, humans, evaluation of 2 therapeutic schemes using diethylcarbamazine combined with levamisole, recommendations for optimum therapy: Haute Volta

Levamisole

Prod'hon, J.; Moreau, J. P.; and Mongin, C., 1979, Med. Trop., v. 39 (6), 619-628

Onchocerca volvulus, humans, mass therapy trials, diethylcarbamazine and levamisole, administered separately or combined: Haute Volta

Levamisole

Rao, Y. V. B. G., 1976, Indian Vet. J., v. 53 (10), 776-777

Capillaria obsignata, chickens (exper.), critical tests with methyridine, pyrantel tartrate, and levamisole

Tetramisole

Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, Ann. Trop. Med. and Parasitol., v. 72 (1), 55-58

Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (*Mesocricetus auratus*)

Levamisole (Nemicide)

Remfry, J., 1978, Lab. Animals, v. 12 (4), 213-218

helminth infections in imported *Macaca mulatta*, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Citarin-L spot-on (Levamisole)

da Rosa, C. E. Q.; Arnoni, J. V.; and Bastos Filho, A. F., 1976, Vet. Med. Rev. (2), 218-222

gastrointestinal nematodes, cattle, levamisole (injected form compared with externally applied form, citarin-L spot-on)

Levamisole

Sanchez Moreno, M.; and Barrett, J., 1979, Parasitology, v. 78 (1), 1-5

Hymenolepis diminuta, adults, monoamine oxidase, occurrence and properties, inhibition by several anthelmintics

Tetramisole

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426

Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Tetramisole -- Continued.

- Levamisole hydrochloride (Nilverm)
Sangster, N. C.; et al., 1979, Research Vet. Sc., v. 27 (1), 106-110
Trichostrongylus colubriformis, Ostertagia circumcincta, Merino and crossbred sheep, field observations and preliminary critical trials showed varying degrees of drug resistance to levamisole hydrochloride, morantel tartrate, and thiabendazole; differences in infectivity and drug efficacy between breeds: Australia
- Levamisole
Santiago, M. A. M.; et al., 1978, Rev. Centro Cien. Rurais, v. 8 (1), 83-87
Ancylostoma sp., Toxocara sp., dogs, levamisole by dermal application
- Levamisole (Nilverm)
Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1977, Rev. Centro Cien. Rurais, v. 7 (4), 421-422
Trichostrongylus colubriformis, levamisole-resistant strain in lambs
- Levamisole (Nilverm, Ripercol L)
Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1978, Rev. Centro Cien. Rurais, v. 8 (3), 257-261
Trichostrongylus colubriformis, levamisole resistant strain, sheep, dl-tetramisole, thiabendazole: Itaquí county, RS
- dl-Tetramisole
Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1978, Rev. Centro Cien. Rurais, v. 8 (3), 257-261
Trichostrongylus colubriformis, levamisole resistant strain, sheep, dl-tetramisole, thiabendazole: Itaquí county, RS
- Levamisole
Santos, L. E.; et al., 1977, Rev. Fac. Med. Vet. e Zootec. Univ. S. Paulo, v. 14 (1), 149-169
treated and untreated Guzera heifers compared for weight gains, hemoglobin levels, and helminth infestation: region of Sertãozinho, Sao Paulo, Brazil
- Levamisole
Scheinberg, M. A.; et al., 1978, Brit. Med. J. (6110), v. 1, 408
levamisole induced cutaneous necrotising vasculitis, case report
- Tramisol (Ripercol)
Schmidt, R. L.; et al., 1979, J. Wildlife Management, v. 43 (2), 461-467
Protostrongylus, domesticated, captive, or free-ranging Ovis c. canadensis, evaluation of 5 drugs
- Levamisole
Schock, R. C.; and Cooper, R., 1978, Mod. Vet. Pract., v. 59 (5), 439-443
nematode parasitism of captive birds, review; safety, efficacy, and dosage of levamisole as broad-spectrum dewormer

Tetramisole -- Continued.

- Galimid
Selim, M. K.; et al., 1972, Vet. Med. J., Giza, v. 20 (20), 125-130
Capillaria obsignata, pigeons and chickens, treatment with galimid effective and non-toxic, also effective against Ascaridia galli and Heterakis gallinae: Cairo, Egypt
- Nilverm
Shchelkanov, K. G.; and Epel'dimov, L. S., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 76-80
Dictyocaulus, sheep, divezid, nilverm, and cyazone, changes in lung tissue of sheep and some tissues of parasite resulting from drug action
- Levamisole
Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies
- Tetramisole
Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies
- Levamisole chlorhydrate
da Silva, D. J.; da Cunha, P. G.; and de Campos, B. do E. S., 1977, Bol. Indust. Animal, Sao Paulo, v. 34 (1), 55-67
nematodes, calves, levamisole chlorhydrate, 0-4 applications, analysis of weight gains
- Levamisole (Ripercol L)
Smith, J. P., 1979, Feline Pract., v. 9 (2), 14, 16, 18
intestinal parasites, cats, efficacy of parenteral aqueous levamisole
- Levamisole (Nilverm)
Smith, W. D.; and Christie, M. G., 1979, J. Comp. Path., v. 89 (1), 141-150
Haemonchus contortus, lambs (exper.), factors influencing degree of host resistance after immunization with attenuated larvae
- Tetramisole, Laevo isomer
de Souza, D. W. C.; Neves, J.; and de Lemos, M. S., 1972, Rev. Inst. Med. Trop. S. Paulo, v. 14 (1), 67-72
A[scaris] lumbricoides, humans, pyrantel pamoate and laevo isomer of tetramisole, clinical trials, satisfactory results: Brazil
- Levamisole (Tramisol)
Stockdale, P. H. G.; and Harries, W. N., 1979, Canad. Vet. J., v. 20 (9), 223-226
gastrointestinal nematodes, cattle, anthelmintic treatment upon entry into feedlot, no improvement in rate of weight gain or feed conversion efficiency: Alberta
- Levamisole (L-Tramisol)
Stokhof, A. A.; and Wolvekamp, W. T. C., 1978, Tijdschr. Diergeneesk., v. 103 (2), 1121-1129
Dirofilaria immitis, dogs, 4 case reports, chemotherapy: Netherlands (imported from United States, South America, or South Africa)

Tetramisole -- Continued.

Levamisole (Decaris)
Szabo, M.; et al., 1977, Therap. Hungar., v. 25 (4), 173-174
decaris, effect on phagocytosis of polymorphonuclear leukocytes varied with concentration

Levamisole
Thienpont, D.; et al., 1977, Tijdschr. Diergeneesk., v. 102 (19), 1123-1128
Toxocara vitulorum, suckling calves, differential diagnosis from *T. canis*, transmammmary transmission, successful treatment with levamisole: Belgium, imported from France

Tetramisole
Tongson, M. S.; et al., 1978, Philippine J. Vet. Med., v. 17 (1-2), 85-100
intestinal nematodes, beef cattle, frequency of deworming determined that will effect the most economical weight gains: Philippines

Levamisole
Turk, J. L.; and Parker, D., 1979, Internat. Arch. Allergy and Applied Immunol., v. 58 (2), 237-240
levamisole in guinea pigs is a potent skin sensitizer

Levamisole
Ueno, H.; and Chibana, T., 1978, Japan Agric. Research Quart., v. 12 (3), 152-156
Stephanofilaria okinawaensis, cattle, distribution, clinical signs, chemotherapy, intermediate host determined

Nilverm
Velichkin, P. A.; et al., 1977, Veterinariia, Moskva (7), 17-19
[Heterakis], [Ascaridia], control on poultry farms by phenothiazine, piperazine adipate, nilverm, and other control measures

Nilverm (Tetramisole; Citarin; Drogenit; Levamisole)
Veselova, T. P., 1979, Veterinariia, Moskva (5), 66-68
nilverm, pharmacological characteristics, effects on various organ systems

Nilverm (Tetramisole)
Vihan, V. S.; and Sahni, K. L., 1979, Indian J. Animal Research, v. 13 (1), 15-18
gastrointestinal nematodes, lambs, efficacy of helatac, nilverm, thiabendole, and banmynth-II: Makhdoom, Mathura (U. P.)

Levamisole
Wagner, E. D.; and Rexinger, D. D., 1978, Am. J. Trop. Med. and Hyg., v. 27 (1, pt. 1), 203-205
Ascaris lumbricoides, *Trichuris trichiura*, mebendazole and levamisole tested for possible ovicidal effects in vivo, human: Sierra Leone; Nassau, Bahamas

Tetramisole sulfamidine
Wang, P. C.; Ling, C. L.; and Chen, C. R., 1975, Tung Wu Hsueh Pao (Acta Zool. Sinica), v. 21 (2), 199-204
Stephanurus dentatus, pigs, seasonal infestation of soil, viability of larvae in different soil types, tetramisole sulfamidine, other control measures: Fukien

Tetramisole -- Continued.

Levamisole (Ripercol)
Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thia-bendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxylin are also useful chemical alternatives

Levamisole (Ketrax)
Willett, W. C.; Kilama, W. L.; and Kihamia, C. M., 1979, Am. J. Pub. Health, v. 69 (10), 987-991
Ascaris-infected children, levamisole, effect on growth rate; results consistent with a causal association between ascariasis and malnutrition: Ubiri village near Lushoto, Tanzania

Levamisole
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Levo-tetramisole (Decaris)
Zaman, V.; and Loh, Y. P., 1974, Singapore Med. J., v. 15 (2), 147-148
Necator americanus, humans, clinical trials testing efficacy of pyrantel embonate, levo-tetramisole, and bephenium hydroxynaphthoate: Singapore

dl-Tetramisole. See Tetramisole.

L-Tetramisole. See Tetramisole.

Tetramisole chlorhydrate. See Tetramisole.

Tetramisole cyclamate. See Tetramisole.

Tetramisole hydrochloride. See Tetramisole.

Tetramisole, Laevo isomer. See Tetramisole.

Tetramisole sulfamidine. See Tetramisole.

Tetramisolum. See Tetramisole.

meso-Tetraphenyl-porphine

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

meso-Tetra-(4-pyridyl)-porphine

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Tetra-sodium-meso-tetra-(p-sulfonato-phenyl)-porphine

Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis

Thenium -- Ancaris (with Piperazine phosphate); N,N-Dimethyl-N-2-phenoxyethyl-N-2'-thenylammonium)-p-chlorobenzenesulfonate; Thenium closylate.

Thenium closylate

Corwin, R. M.; and Miller, T. A., 1978, Am. J. Vet. Research, v. 39 (2), 263-265
Toxocara canis, young dogs and weaned pups, thenium closylate, piperazine phosphate, efficacy when administered alone or in combination, critical controlled trials

Thenium closylate + Piperazine phosphate (= Ancaris)

Remfry, J., 1978, Lab. Animals, v. 12 (4), 213-218
helminth infections in imported Macaca mulatta, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Thenium closylate. See Thenium.

2- α -Thenoyl-amino-5-nitrothiazole. See Atrican.

Theseinacid

Christow, C., 1978, Biochem. and Exper. Biol., v. 14 (2), 177-180
Trichomonas foetus, effect of certain B₁₂ antagonists upon growth

Thiabendazole -- BB-04 (with Piperazine hexahydrate); Equizole; Equizole A (with Piperazine); Gamaverm (with Piperazine); Mintesol; Mintezol; Mintezole; Minzolum; Omnizole; Omnizole Wormer Paste; Suiverm (with Piperazine); Thiabendazole hydrochloride; Thiabendole; Thiabenzole; 2-(4'-Thiazolyl)benzimidazole; Thibendole; Thibenzole; Thibenzole Veterinary Liquid; Tresaderm (with Dexamethasone and Neomycin).

Thiabendazole -- Continued.

Thiabendazole

Abdel Rahman, M. S.; et al., 1978, Vet. Med. J., Giza, v. 24 (24), 1976, 39-48
gastro-intestinal nematodes, sheep, comparative field trials with 6 anthelmintics: Egypt

Thiabendazole

Anosa, V. O., 1977, Trop. Animal Health and Prod., v. 9 (1), 11-17
Haemonchus contortus outbreak, Nigerian dwarf ewes and lambs, haematological observations before and after treatment with thiabendazole: Nigeria

Thiabendazole

Arguedas, J. A.; et al., 1975, Texas Rep. Biol. and Med., v. 33 (2), 265-268
Strongyloides stercoralis, mass treatment of a community with thiabendazole reduced prevalence, 2-year follow-up showed that rate remained low: Costa Rica

Thiabendazole lotion

Azulay, R. D.; et al., 1975, Rev. Brasil. Med., v. 32 (12), 800-802
scabies, human, clinical trials, efficacy of thiabendazole vs. benzyl benzoate as topical lotions

Thiabendazole

Baqui, A.; and Ansari, J. A., 1976, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 25 (5), 409-414
Setaria cervi, white rats, hetrazan, thiabendazole, tetramisole

Thiabendazole

Botero R., D., 1978, Ann. Rev. Pharmacol. and Toxicol., v. 18, 1-15
antiparasitic drugs in current use for human intestinal protozoa and helminths, brief review of pharmacology, secondary effects, toxicity and contraindications

Thiabendazole

Callinan, A. P. L.; and Cummins, L. J., 1979, Austral. Vet. J., v. 55 (8), 370-373
nematodes, cattle (nat. and exper.), efficacies of various anthelmintics against adult and larval stages: western Victoria

Thiabendazole + Piperazine hexahydrate (= BB-04)

Camillo-Coura, L.; et al., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (2), 103-113
human Ascaris lumbricoides, Trichuris trichiura, Ancylostomidae, treatment trials with piperazine combined with thiabendazole in silicone base (BB-04), mixed results

Thiabendazole (Thibenzole)

Campbell, N. J.; et al., 1978, Austral. Vet. J., v. 54 (1), 23-25
Haemonchus contortus, Trichostrongylus colubriformis, benzimidazole resistant strains, sheep, efficacy of 6 non-benzimidazole anthelmintics and thiabendazole, controlled test

Thiabendazole -- Continued.

Thiabendazole

Casali, A. J.; and De Costa, E. A., 1977, Bol. Chileno Parasitol., v. 32 (3-4), 66-70

T[richinella] spiralis, outbreak in 16 members of family group who had eaten raw pork, successful therapy with thiabendazole: Provincia de San Luis, Republica Argentina

Thiabendazole

Chaia, G.; and da Cunha, A. S., 1970, Rev. Inst. Med. Trop. S. Paulo, v. 12 (2), 152-160
intestinal helminths, children, efficacy of anthelmintic therapy: Vila Sao Vicente, Belo Horizonte

Thiabendazole (Thibenzole)

Chevis, R. A. F.; and Kelly, J. D., 1978, N. Zealand Vet. J., v. 26 (9), 218-220
Fasciola hepatica, rats and sheep (both exper.), mebendazole, parbendazole, cambendazole, thiabendazole, anthelmintic activity, molecular structure-activity analyses

Thiabendazole

Coles, G. C.; and Briscoe, M. G., 1978, Vet. Rec., v. 103 (16), 360-361 [Letter]
Fasciola hepatica eggs, LD₅₀ values of 7 benzimidazoles determined and compared with values for *Haemonchus contortus* eggs

Thiabendazole

Daniyarov, I. A.; et al., 1978, Veterinariia, Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics and dyes tested, none effective

Thiabendazole (Thibenzole)

Darvill, F. M.; Arundel, J. H.; and Brown, P. B., 1978, Austral. Vet. J., v. 54 (12), 575-584
Ostertagia, *Trichostrongylus*, maiden ewes, effect of pre-lambing and post-lambing thiabendazole treatment on pasture contamination by ewes and on bodyweights of ewes and lambs

2-(4¹-Thiazolyl)-benzimidazole (Thiabendazole)
Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

Thiabendazole

DePaoli, A.; and Johnsen, D. O., 1978, Vet. Path., v. 15 (1), 31-39
Strongyloides larvae in *Hylobates* lar, necropsy study, clinical signs, pathology, thiabendazole, high morbidity and mortality probably resulting from autoinfection and the hyperinfective syndrome

Thiabendazole -- Continued.

Thiabendazole (Thibenzole)

Donald, A. D.; et al., 1979, Vet. Parasitol., v. 5 (2-3), 205-222
gastrointestinal nematodes with major emphasis on *Ostertagia ostertagi*, beef cattle, levels of infection and effects on live-weight gain, effects of pasture type (*phalaris* vs. *lucerne*) and stocking rate, effects of anthelmintic treatment, 4-year experiment: Canberra, Australia

Thiabendazole (Thibenzole)

Dorchies, P.; Franc, M.; and Ducos de Lahitte, J., 1979, Rev. Med. Vet., Toulouse, v. 130 (3), 417-418, 421-425
nematodes, horses, levamisole administered per os and intramuscularly, toxicity; efficacy compared with thiabendazole

Thiabendazole (Omnizole)

Drudge, J. H.; Lyons, E. T.; and Tolliver, S. C., 1979, Am. J. Vet. Research, v. 40 (4), 590-594
large and small strongyles, horses, critical tests with 6 benzimidazoles, drug resistance

Thiabendazole (Thibenzole)

Evans, J. W.; and Green, P. E., 1978, Austral. Vet. J., v. 54 (9), 454-455 [Letter]
Platynosomum concinnum, cats, anthelmintics, drug trials

Thiabendazole

Fabiyi, J. P.; and Offiong, S. A., 1979, Vet. Rec., v. 104 (15), 348
Syngamus trachea in *Numida meleagris* *galeata*, severe outbreak after excessively wet weather, thiabendazole: Vom, Nigeria

Tresaderm

Faulk, R. H.; and Schwirck, S., 1978, Vet. Med. and Small Animal Clin., v. 73 (3), 307-308
Otodectes cynotis, dogs, cats (ears of both), tresaderm, clinical trial, highly effective

Thiabendazole + Piperazine

Fernandes, J. L.; and Garcia, E., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (3), 155-158
human intestinal parasites, clinical trials with combined piperazine and thiabendazole: Sao Paulo

Thiabendazole + Piperazine hexahydrate (= Gama-
verm)

Ferriolli Filho, F.; et al., 1971, Rev. Soc. Brasil. Med. Trop., v. 5 (4), 209-212
Ascaris lumbricoides, *Strongyloides stercoralis*, *Ancylostomidae*, mixed human infections, clinical trials testing efficacy of piperazine and thiabendazole in combination therapy

Thiabendazole

Fink, A. I.; MacKay, C. J.; and Cutler, S. S., 1978, Tr. Am. Ophth. Soc., v. 76, 108-115
sicca complex and cholestatic jaundice in 2 family members after thiabendazole treatment for pinworms

Thiabendazole -- Continued.

Thiabendazole

Friedman, P. A.; and Platzer, E. G., 1978, *Biochim. et Biophys. Acta*, v. 544 (3), 605-614
benzimidazoles and benzimidazole derivatives, interaction with bovine brain tubulin, implications for mode of anthelmintic action

Thiabendazole (Mintesol)

Gateff, C.; Lemarinier, G.; and Labusquiere, R., 1972, *Ann. Soc. Belge Med. Trop.*, v. 52 (2), 103-112

intestinal helminths, African schoolchildren, systematic thiabendazole therapy over 8-month period, increased weight gains with cost of therapy less than if given nutritional supplements, favorable impact on epidemiology: Yaounde, Cameroun

Thiabendazole (Mintezol)

Gatti, F.; et al., 1972, *Ann. Soc. Belge Med. Trop.*, v. 52 (1), 19-31

intestinal nematodes, African schoolchildren, trimestrial administration of levamisole compared with other anthelmintics: Kimwenza, Republique Democratique du Congo

Thiabendazole (Thiabendazole)

Gaur, S. N. S.; and Dutt, S. C., 1979, *Indian Vet. J.*, v. 56 (1), 62-64

Mecistocirrus digitatus, cattle, comparative efficacy of thiabendazole, nilverm, and helmatac

Thiabendazole

Gibson, T. E.; and Everett, G., 1971, *J. Comp. Path.*, v. 81 (4), 493-498

Ostertagia and *Trichostrongylus* spp., lambs treated with thiabendazole and moved to clean pasture had lowest worm burden

Thiabendazole

Gill, G. V.; and Bell, D. R., 1979, *Brit. Med. J.* (6190), v. 2, 572-574

Strongyloides stercoralis, former Far East prisoners of war, symptoms, diagnostic difficulties, thiabendazole: Britain

Thiabendazole (Mintezol)

Glickman, L.; et al., 1979, *Am. J. Trop. Med. and Hyg.*, v. 28 (1), 29-35

Toxocara canis, 2-year-old boy with concurrent ocular and visceral toxocariasis, case report, parasite-specific antibodies in serum and aqueous humor, response to treatment with prednisone and thiabendazole

Thiabendazole

Gordon, W., 1978, *South African Med. J.*, v. 54 (19), 767 [Letter]

sandworm, human, topical thiabendazole in anhydrous wool fat and paraffin: South Africa

Thiabendazole

Goulart, E. G.; de Arruda, M. E.; and Jourdan, M. C., 1974, *Rev. Brasil. Med.*, v. 31 (11), 791-794

human soil transmitted nematodes, laboratory trials testing ovicidal and larvacidal effects of selected anthelmintic drugs; prophylactic treatment of organic fertilizer or contaminated soil by these drugs seemed to be ineffective

Thiabendazole -- Continued.

Thiabendazole

Grell, G. A. C.; Watty, E. I.; and Muller, R. L., 1978, *Brit. Med. J.* (6150), v. 2, 1464
Syngamus [sp.] in woman who travelled extensively in Caribbean Islands, recovery after thiabendazole treatment, case report: Dominica, West Indies

Thiabendazole + Piperazine (=Equizole A)

Grieve, R. B.; Moore, B. G.; and Bradley, R. E., 1979, *Am. J. Vet. Research*, v. 40 (1), 139-141

gastrointestinal parasites, horses and ponies, critical test evaluation of butamisol, compared with efficacy of piperazine-thiabendazole

Thiabendazole

Gudimenko, I. I., 1970, *Nauchn. Trudy, Nauchno-Issled. Vet. Inst.*, v. 8, 100-104

trichuriasis, oesophagostomiasis, ascariasis, swine, testing thiabendazole, dithiazanine iodide, dipterex, hygromycin-B, and bubulin

Thiabendazole

Hall, C. A.; et al., 1978, *Research Vet. Sc.*, v. 25 (3), 364-367

Haemonchus contortus, *Trichostrongylus colubriformis*, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Thiabendazole

Hall, C. A.; Campbell, N. J.; and Richardson, N. J., 1978, *Research Vet. Sc.*, v. 25 (3), 360-363

Haemonchus contortus, *Trichostrongylus colubriformis*, levels of benzimidazole resistance recorded from an egg hatch test procedure

Thiabendazole

Hennessy, D. R.; and Prichard, R. K., 1979, *Austral. Vet. J.*, v. 55 (6), 298 [Letter]

deaths in cattle after administration of large doses of thiabendazole

Thiabendazole (Equizole)

Holt, P. E.; Brown, A.; and Brown, B., 1978, *Vet. Rec.*, v. 102 (18), 404-405

Strongyloides [sp.] in *Lampropeltis getulus holbrooki*, clinical symptoms, unsuccessful treatment with thiabendazole, mixed infection with flagellates and *Ochetosomatidae* sp., case report

Thiabendazole (Equizole)

Holt, P. E.; Cooper, J. E.; and Needham, J. R., 1979, *J. Small Animal Practice*, v. 20 (5), 269-286

Angusticaecum spp., *Tachygonetria* sp., and *Atractis dactyluris* in *Testudo*, treatment: Britain

Thiabendazole

Holt, P. E.; Cooper, J. E.; and Needham, J. R., 1979, *Vet. Rec.*, v. 104 (10), 213-214

Strongyloides larvae, snakes, pathology, treatment with thiabendazole, 3 case reports

Thiabendazole -- Continued.

Thiabendazole

Hopkins, T. J.; and Rafferty, M., 1978, Vet.-Med. Nachr. (2), 160-168
nematodes, sheep, febantel, drug trials, cross-resistance of thiabendazole-resistant strains of *Haemonchus contortus* and *Trichostrongylus colubriformis*: Australia

Thiabendazole (Thibenzole)

Horton, G. M. J., 1977, J. Animal Sc., v. 45 (6), 1453-1457
Trichostrongylus colubriformis, lambs (exper.), feed utilization, calcium and phosphorus metabolism and serum protein fractions, before and after treatment with thiabendazole

Suiverm

Iakubovskii, M. V.; and Zen'kov, A. V., 1977, Vet. Nauka--Proizvod., Trudy, Minsk, v. 15, 72-78
nematodes, swine-raising complexes, combined control measures, sanitation, anthelmintics, suiverm most effective

Thiabendazole

Ireland, C. M.; et al., 1979, Biochem. Pharmacol., v. 28 (17), 2680-2682
relative effectiveness of several benzimidazole carbamates and related compounds on assembly of sheep brain microtubules in vitro and on infections of *Nematospiroides dubius* in mice

Thiabendazole (Thibenzole)

Johnstone, I. L.; et al., 1979, Austral. J. Exper. Agric. and Animal Husb. (98), v. 19, 303-311
parasites, sheep, effect of four control schemes in two environments on wool production and liveweight gains

Thiabendazole (Thibenzole)

Johnstone, I. L.; Coote, B. G.; and Smart, K. E., 1979, Austral. J. Exper. Agric. and Animal Husb. (99), v. 19, 414-418
pre- and/or post-lambing anthelmintic treatment, effect on lamb birth weight and liveweight gain, fecal egg counts, pasture contamination: New South Wales

Thiabendazole

Koehler, P.; and Bachmann, R., 1978, Molec. Pharm., v. 14 (1), 155-163
Ascaris suum muscle tissue, comparison of effects of levamisole, thiabendazole, chloroquine, and praziquantel on electron transport in *Ascaris* muscle submitochondrial particles

Thiabendazole

Krotov, A. I.; Cherniaeva, A. I.; and Budanova, I. S., 1976, Med. Parazitol. i Parazitarn. Bolezni, v. 45 (2), 165-168
Alveococcus multilocularis, white mice, effect of thiabendazole, sarcocystin acridine, levamisole, and mebendazole on larval cyst development

Thiabendazole (Thibenzole)

Laemmler, G.; and El-Gendi, A. Y. I., 1978, Ztschr. Parasitenk., v. 58 (1), 55-73
Ancylostoma caninum in *Mastomys natalensis*, efficacy of various anthelmintics against third stage larvae

Thiabendazole -- Continued.

Thiabendazole (Thibenzole)

Le Jambre, L. F., 1979, Austral. Vet. J., v. 55 (2), 65-67
Ostertagia circumcincta, *O. trifurcata*, sheep (exper.), effectiveness of levamisole, thiabendazole, albendazole, and oxfendazole against levamisole-resistant strains

Thiabendazole

Le Jambre, L. F.; Martin, P. J.; and Webb, R. F., 1979, Austral. Vet. J., v. 55 (4), 163-166
Haemonchus contortus, thiabendazole resistance in field populations, use of egg hatch assay to detect low but significant levels of resistance: Northern Tablelands of New South Wales

Thiabendazole

Le Jambre, L. F.; Royal, W. M.; and Martin, P. J., 1979, Parasitology, v. 78 (2), 107-119
Haemonchus contortus, thiabendazole resistance is inherited as an autosomal and semi-dominant trait

Thiabendazole

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Austral. Vet. J., v. 54 (12), 570-574
Trichostrongylus colubriformis, linear dose responses of selected and unselected strains to thiabendazole, levamisole, and morantel tartrate

Thiabendazole

Le Jambre, L. F.; Southcott, W. H.; and Dash, K. M., 1978, Internat. J. Parasitol., v. 8 (6), 443-447
Ostertagia circumcincta, development of simultaneous resistance to thiabendazole, morantel tartrate, and levamisole, multiple selection associated with increase in *O. trifurcata* in population and increase in larval inhibition

Thiabendazole

Loria Cortes, R.; and Saborio Ruiz, M., 1974, Rev. Columb. Pediat. y Puericult., v. 28 (6), 409-413
Necator americanus, prenatal infection in 1 month-old infant presenting with intestinal bleeding, successful thiabendazole therapy: Costa Rica

Thiabendazole

Lumbreras, H.; et al., 1972, Rev. Peruana Med. Trop., v. 1 (2), 84-86
human uncinariasis, technique of Harada-Mori used to evaluate action of thiabendazole against viability of parasite eggs, inhibition and retardation of egg hatching was demonstrated

Thiabendazole (Thibenzole)

MacArthur, J. A.; and Wood, M., 1978, Lab. Animals, v. 12 (3), 141-143
Aspiculuris tetraptera, *Syphacia obvelata*, naturally infected mouse colony continuously medicated with thiabendazole in diet, anthelmintic effect, effect on growth and breeding

Thiabendazole -- Continued.

Thiabendazole

- Macchioni, G.; et al., 1978, Clin. Vet., Milano, v. 101 (4), 185-190
 Dicrocoelium dendriticum, sheep, efficacy of thiabendazole at various dosage rates, post mortem counts of adult parasites

Thiabendazole

- McEwan, A. D.; and Oakley, G. A., 1978, Vet. Rec., v. 102 (14), 314-315
 panacur, thiabendazole, and nilzan with dye marker added, oral dosing of cattle showed evidence of rumen by-pass, reduced drug efficacy probably resulting from closure of oesophageal groove

Thiabendazole

- MacKay, R. J.; and Urquhart, K. A., 1979, Equine Vet. J., v. 11 (2), 110-112
 Dictyocaulus arnfieldi, presumptive diagnosis in 8 horses with eosinophilic bronchitis, infection confirmed in companion donkey, thiabendazole treatment, usefulness of cytology of tracheobronchial secretions in differential diagnosis

Thiabendazole (Thibendole)

- Mallick, K. P.; Mishra, S. S.; and Sahai, B. N., 1977, Indian J. Animal Research, v. 11 (2), 77-80
 Haemonchus spp., goats (exper.), thiabendazole, pyrantel tartrate, krimmos, comparative efficacy, blood picture before and after treatment, krimmos not effective

Thiabendazole (Thibenzole)

- Mechow, A., 1977, Prakt. Tierarzt, v. 58 (11), 818, 822-825
 increased milk yield in dairy cows treated with thiabendazole at the beginning of lactation

Thiabendazole (Thibenzole)

- Melbourne, C. P., 1978, J. Zoo Animal Med., v. 9 (4), 146-148
 Trichuris sp. and 5 strongyloid genera, giraffes, camels, fenbendazole and thiabendazole: Longleat Safari Park, Great Britain

Thiabendazole

- Meltzer, R. S.; et al., 1979, Am. J. Med. Sc., v. 277 (1), 91-98
 Strongyloides stercoralis, woman, development of hyperinfection syndrome while on high-dose corticosteroids and following splenectomy, central nervous system involvement, antemortem diagnosis, thiabendazole, levamisole, and mebendazole therapy: Memorial Sloan-Kettering Cancer Center, New York (had traveled in Italy and Sicily)

Thiabendazole (Mintezol)

- Merdivenci, A.; Sengul, M.; and Baydemir, M., 1976, Turk Hijiy. ve Deneysel Biyol. Dergisi, v. 36 (2), 224-237
 Ascaris lumbricoides, Enterobius vermicularis, children, thiabendazole and mebendazole compared, both successful

Thiabendazole -- Continued.

Thiabendazole

- Molet, B.; and Kremer, M., 1975, Ann. Soc. Belge Med. Trop., v. 55 (5), 431-433
 "trichocephales", morphological alterations of eggs after thiabendazole therapy

Mintezole

- Nozais, J. P.; Heroin, P.; and Doucet, J., 1975, Rev. Med. Cote Ivoire (38), v. 11, 2. trimestre, 55-57
 strongyloidiasis in man with resulting severe urticarial type dermatitis, case report, cure with mintezole: Cote Ivoire

Thiabendazole

- Ogunsusi, R. A., 1978, Research Vet. Sc., v. 25 (2), 251-252
 Haemonchus spp. and Trichostrongylus spp., adult Yankasa sheep, thiophanate and thiabendazole at increased dosages highly effective in removing all stages of development including inhibited stages

Thiabendazole (Thibenzole)

- Owen, D.; and Turton, J. A., 1979, Lab. Animals, v. 13 (2), 115-118
 Syphacia obvelata, laboratory mice and rats, thiabendazole and piperazine citrate in feed and water

Thiabendazole

- Petrov, Iu. F., 1978, Veterinariia, Moskva (5), 64-66
 [Tetrameres], [Streptocara], ducks, effectiveness of various anthelmintics

Thiabendazole (Thibenzole)

- Pfister, K., 1978, Schweiz. Arch. Tierh., v. 120 (2), 89-99
 gastro-intestinal strongyles in lambs free of or also infected with Moniezia spp., thiabendazole administered periodically, changes in daily weight gain

Thiabendazole (Omnizole paste)

- Pluimers, E. J., 1979, Vet. Quart., v. 1 (2), 82-89
 gastrointestinal nematodes, Dutch dairy cattle, milk production increase following treatment with thiabendazole

Thiabendazole

- Pouplard, L., 1978, Vet. Rec., v. 103 (19), 434 [Letter]
 dairy cows, anthelmintic treatment with thiabendazole, increased milk production: Belgium

Thiabendazole

- Prichard, R. K.; et al., 1978, Vet. Rec., v. 102 (17), 382
 Ostertagia ostertagi arrested 4th stage larvae, incorporation of C¹⁴-labeled thiabendazole no lower than that of adult worms, increased tolerance may be due to lower energy demands of arrested larvae, higher efficacy can be achieved by persistently high anthelmintic concentrations in host

Thiabendazole -- Continued.

Thiabendazole

Prichard, R. K.; Hennessy, D. R.; and Steel, J. W., 1978, *Vet. Parasitol.*, v. 4 (4), 309-315

Haemonchus contortus, *Trichostrongylus colubriformis*, sheep, *Ostertagia ostertagi*, cattle, 4 benzimidazoles, mode of action and pharmacokinetic behavior, implications for prolonged administration as a new concept for increasing spectrum and effectiveness of anthelmintics

Thiabendazole

Prichard, R. K.; Kelly, J. D.; and Thompson, H. G., 1978, *Vet. Parasitol.*, v. 4 (3), 243-255
Haemonchus contortus, *Trichostrongylus colubriformis*, sheep, thiabendazole, fenbendazole, concentrations of anthelmintics or their radiolabelled metabolites in parasite tissues after administration to host, differences between amount of each anthelmintic incorporated by susceptible and resistant parasite strains and between the two parasites, effect of route of administration on anthelmintic concentration in parasite tissue and host plasma

Thiabendazole

Pujman, V. F.; and Hanusova, D., 1970, *J. Wildlife Dis.*, v. 6 (3), 163-166
 erythrocyte variations of normal vs. parasitized, mature vs. immature *Perdix perdix* and *Phasianus colchicus colchicus*, normalization of *Syngamus trachealis*-infected pheasant erythrocyte after addition of thiabendazole to feed

Thiabendazole

Ray, D. K.; Bhopale, E. K.; and Shrivastava, V. B., 1978, *Ann. Trop. Med. and Parasitol.*, v. 72 (1), 55-58
Ancylostoma ceylanicum, efficacy of 7 anthelmintics tested using an experimental patent infection in the golden hamster (*Mesocricetus auratus*)

Thiabendazole

Reinhardt, P., 1978, *Monatsh. Vet.-Med.*, v. 33 (23), 898-901
Dicrocoelium dendriticum, sheep, efficacy of various anthelmintics compared

Thiabendazole (Thibenzole)

Remfry, J., 1978, *Lab. Animals*, v. 12 (4), 213-218
 helminth infections in imported *Macaca mulatta*, incidence, pathogenicity, and treatment: imported from northern India to Primate Quarantine Unit, Oxford University

Thiabendazole (Mintezol)

Rougereau, A.; Lasnier, H.; and N'Diaye, A. M., 1977, *Med. et Nutrition*, v. 13 (6), 473-475
 oxyures, mice, thiabendazole, drug efficacy highest in hosts on a hyperproteinic diet

Thiabendazole hydrochloride

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Thiabendazole -- Continued.

Thiabendazole

Sanchez Moreno, M.; and Barrett, J., 1979, *Parasitology*, v. 78 (1), 1-5
Hymenolepis diminuta, adults, monoamine oxidase, occurrence and properties, inhibition by several anthelmintics

Thiabendazole

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, *Rev. Iber. Parasitol.*, v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Thiabendazole (Thibenzole)

Sangster, N. C.; et al., 1979, *Research Vet. Sc.*, v. 27 (1), 106-110
Trichostrongylus colubriformis, *Ostertagia circumcincta*, Merino and crossbred sheep, field observations and preliminary critical trials showed varying degrees of drug resistance to levamisole hydrochloride, morantel tartrate, and thiabendazole; differences in infectivity and drug efficacy between breeds: Australia

Thiabendazole (Thibenzole)

Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1978, *Rev. Centro Cien. Rurais*, v. 8 (1), 35-38
Haemonchus contortus, thiabendazole-resistant strain, sheep, fenbendazole

Thiabendazole (Thibenzole)

Santiago, M. A. M.; da Costa, U. C.; and Benevenga, S. F., 1978, *Rev. Centro Cien. Rurais*, v. 8 (3), 257-261
Trichostrongylus colubriformis, levamisole resistant strain, sheep, dl-tetramisole, thiabendazole: Itaqui county, RS

Thiabendazole

Sargent, R. G.; et al., *South. Med. J.*, v. 66 (2), 276-278
Trichuris trichiura, humans, clinical trials with thiabendazole

Thiabendazole (Mintezole)

Sastry, S. C.; Jaya Kumar, K.; and Lakshminarayana, V., 1978, *J. Trop. Med. and Hyg.*, v. 81 (2-3), 32-35
 human dracunculosis, treatment of 400 cases with thiabendazole, drug appears highly effective without major side effects: South India

Thiabendazole (Thibenzole)

Saxena, S. N., 1974, *U. P. Vet. J.*, v. 2 (4), 132-137
Ascarops strongylina, piglets (exper.), critical trials of efficacy of carbon disulphide, thiabendazole, and sodium fluoride against mature worms

Thiabendazole (Thiabendazole)

Schillhorn Van Veen, T. W., 1978, *Vet. Rec.*, v. 102 (16), 364-365
Haemonchus contortus, lambs (abomasum, faeces), casualties of young lambs following prolonged rainy season, further casualties at end of following dry season associated with inhibited *H. contortus* larvae suggest chronic haemonchosis syndrome (lambs had been previously treated with thiabendazole and rafoxanide); high pasture infection: Shika, near Zaria

Thiabendazole -- Continued.

Thiabendazole

Schmidt, R. L.; et al., 1979, J. Wildlife Management, v. 43 (2), 461-467
Protostrongylus, domesticated, captive, or free-ranging *Ovis c. canadensis*, evaluation of 5 drugs

Thiabendazole

Schumaker, J. D.; et al., 1978, Ann. Int. Med., v. 89 (5, pt. 1), 644-645
Strongyloides stercoralis, hyperinfected anephric patient on hemodialysis, successful use of thiabendazole, pharmacokinetic information

Thiabendazole

Shedivtsova, A., 1976, Med. Parazitol. i Parazit. Bolezni, v. 45 (2), 173-176
Nippostrongylus braziliensis, migratory phase, white mice, 16 anthelmintics tested, model for larval nematode treatment studies

Thiabendazole

Simpkin, K. G.; and Coles, G. C., 1978, Research Vet. Sc., v. 25 (2), 249-250
Haemonchus and *Trichostrongylus colubriformis* in sheep (exper.), instability of egg resistance to benzimidazoles, cross resistance between drugs (thiabendazole, cambendazole, mebendazole, parabendazole, oxbendazole)

Thiabendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 226-229
Ancylostoma caninum, dogs, comparative efficacy of disophenol, fenbendazole, pyrantel pamoate, and thiabendazole, clinical trials

Thiabendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1977, Haryana Vet., v. 16 (1), 5-7
Ancylostoma caninum, dogs, pyrantel pamoate, fenbendazole, thiabendazole, clinical trials, comparative efficacy

Thiabendazole

Singh, H.; Singh, R. P.; and Bali, M. K., 1978, Haryana Agric. Univ. J. Research, v. 8 (1), 55-58
Ancylostoma sp., thiabendazole, fenbendazole, ancylosol, and pyrantel pamoate, in vitro effects on development of eggs and infective larvae

Thiabendazole

Slepnev, N. K., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 108-112
ascariasis, trichuriasis, oesophagostomiasis, swine, comparative effectiveness of various anthelmintics

Thiabendazole (Thiabendazole)

Stockdale, P. H. G.; and Harries, W. N., 1979, Canad. Vet. J., v. 20 (9), 223-226
gastrointestinal nematodes, cattle, anthelmintic treatment upon entry into feedlot, no improvement in rate of weight gain or feed conversion efficiency: Alberta

Thiabendazole -- Continued.

thiabendazole (Minzolum)

Stumpf, J.; Landgraf, H.; and Undeutsch, K., 1978, Therapiewoche, v. 28 (42), 8010-8015
trichinellosis, humans, severe infections treated with thiabendazole, clinical course, side effects

Thiabendazole (Omnizole)

Tangredi, B. P., 1978, Vet. Med. and Small Animal Clin., v. 73 (2), 215-216
Capillaria sp., young caller ducks (intestine), case history, pathology, thiabendazole, good results: pond in Centre Island, Long Island, New York

Thiabendazole

Tidball, J. S.; Aguas, J. P.; and Aldis, J. W., 1978, Southeast Asian J. Trop. Med. and Pub. Health, v. 9 (1), 33-40
Capillaria philippinensis, humans (stools), 32 cases, history of eating raw fish, thiabendazole, mebendazole, life cycle discussed: San Antonio and San Narciso, Zambales Province, Western Luzon, Philippines

Thiabendazole

Tongson, M. S.; et al., 1978, Philippine J. Vet. Med., v. 17 (1-2), 85-100
intestinal nematodes, beef cattle, frequency of deworming determined that will effect the most economical weight gains: Philippines

Thiabendazole

Van Arman, G. G.; and Campbell, W. C., 1975, Texas Rep. Biol. and Med., v. 33 (2), 303-311
anti-inflammatory activity of thiabendazole assayed in laboratory animals, these same properties may contribute to clinical response after thiabendazole treatment of various human parasitic infections

Thiabendazole + Piperazine (= Gamaverm)

Velho Soli, A. S.; et al., 1972, Rev. Soc. Brasil. Med. Trop., v. 6 (4), 207-210
human helminths, clinical trials using combined piperazine and thiabendazole: Brazil

Thiabendazole (Thiabendazole)

Vihan, V. S.; and Sahni, K. L., 1979, Indian J. Animal Research, v. 13 (1), 15-18
gastrointestinal nematodes, lambs, efficacy of helatac, nilverm, thiabendazole, and banminth-II: Makhdoom, Mathura (U. P.)

Thiabendazole (Thibenzole)

Webb, R. F.; Jackson, A. R. B.; and McCully, C. H., 1978, Austral. Vet. J., v. 54 (10), 501-502 [Letter]
Haemonchus contortus, sheep (nat. and exper.), efficiency of various anthelmintics against field populations resistant to thiabendazole, results confirm the usefulness of levamisole, naphthalophos, and rafoxanide for this purpose, haloxon and nitroxynil are also useful chemical alternatives

Thiabendazole (Omnizole Wormer Paste)

Wescott, R.; and Shelton, T., 1979, West. Veterinarian, v. 17 (2), 11-14
internal parasites, cattle, prevalence on 5 ranches, effect of types of pasture, management practices, age of host, and treatment with thiabendazole: Butte County, Idaho

Thiabendazole -- Continued.

Thiabendazole (Thibenzole Veterinary Liquid; Mintezol)
Whiting, D. A., 1976, South African Med. J., v. 50 (8), 253-255
creeping eruption, humans, successful topical application of thiabendazole: South Africa

Thiabendazole
Wilson, C. G.; et al., 1979, Xenobiotica, v. 9 (6), 343-351
thiabendazole, inhibition of metabolism in the rat by ethoxyquin and desmethylimipramine; simultaneous administration of ethoxyquin (but not desmethylimipramine) potentiated anthelmintic effect of thiabendazole on *Nematospiroides dubius*

Thiabendazole + Ethoxyquin
Wilson, C. G.; et al., 1979, Xenobiotica, v. 9 (6), 343-351
thiabendazole, inhibition of metabolism in the rat by ethoxyquin and desmethylimipramine; simultaneous administration of ethoxyquin (but not desmethylimipramine) potentiated anthelmintic effect of thiabendazole on *Nematospiroides dubius*

Thiabendazole (Mintezol)
Yalcinkaya, F., 1976, Turk Hijiy. ve Deneysel Biyol. Dergisi, v. 35 (2-3), 1975, 101-106
helminths, human, thiabendazole

Thiabendazole
Yalcinkaya, F., 1978, Turk Hijiy. ve Tecrubu Biyol. Dergisi, v. 37 (3)-38 (1-3), 1977-1978, 253-261
Toxocara canis, mice (exper.), thiabendazole for larva migrans, early administration effective

Thiabendazole
Zavala-Velazquez, J.; Bolio-Cicero, A.; and Pinzon-Cantarell, J., 1976, SPM Salud Pub. Mexico, v. 18 (4), 735-742
Ancylostoma spp. causing creeping eruption in humans, clinical and epidemiological study, thiabendazole therapeutically effective: Yucatan, Mexico

Thiabendazole hydrochloride. See Thiabendazole.

Thiabendole. See Thiabendazole.

Thiabendole. See Thiabendazole.

Thiacetarsamide sodium. See Arsenamide.

Thiacoccide. See Amprolium.

2-(4'-Thiazolyl)benzimidazole. See Thiabendazole.

2-(4-Thiazolyl)-5-carbamoisopropoxybenzimidazole. See Cambendazole.

2-(4-Thiazolyl)-5-isopropoxycarbonylaminobenzimidazole. See Cambendazole.

4-[p-(2-Thiazolylsulfamyl)-phenylazo]-1-naphthylamine
Korolkovas, A.; and Barata, M. A. L., 1972, Rev. Farm. e Bioquim. Univ. Sao Paulo, v. 10 (1), 113-124
preparation and testing of 6 long-acting schistosomicidal resins

Thiabendole. See Thiabendazole.

Thiabendole. See Thiabendazole.

Thiabendole Veterinary Liquid. See Thiabendazole.

Thimerosal -- Merthiolate.

Merthiolate
Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Thiobis(2-acetoxy-3,5-dichlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Thiobis(2-acetoxy-3-nitro-5-chlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Thiobis(2-acetoxy-3,5,6-trichlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Thiobis(4-bromo-6-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Thiobis(4-bromophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Thiobis(2-carboxymethoxy-3,5-dichlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2'-Thiobis(4-chloro-6-bromophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

- 2,2'-Thiobis (4-chloro-6-nitrophenol)
Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice
- 2,2'-Thiobis (4-chloro-6-nitrophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis (4-chlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis (4,6-dibromophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis (4,6-dichlorophenol). See Bithiolenol.
- 4,4'-Thiobis (2,6-dichlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis (4-fluorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- Thiobis (2-methoxy-3-nitro-5-chlorobenzene)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- 2,2'-Thiobis (3,4,6-trichlorophenol)
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives
- Thiodiphenylamin. See Phenothiazine.
- 0,0'-(Thiodi-p-phenylene) 0,0,0',0'-tetramethyl phosphorothioate. See Temephos
- 5-Thio-D-glucose
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)
- 6-Thioguanine
Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214
Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo
- Thioguanine
Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection
- 6-Thioguanosine
Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856
Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs
- Thiophanate -- A 445; Diethyl 4,4'-O-phenylene bis (3-thioallophanate); Nemafox.
- Thiophanate
Baines, D. M.; et al., 1979, Vet. Rec., v. 105 (4), 81-82
nematodes, pigs, thiophanate given as 14-day low level in-feed anthelmintic, field studies
- Thiophanate (Nemafox)
Baines, D. M.; and Dalton, S. E., 1978, Vet. Rec., v. 103 (24), 527-530
nematodes of sheep and cattle, experimental studies and field trials with thiophanate administered at low daily dosages in various regimes, effects during and after medication, good results; shorter term/higher dosage regimes also show considerable promise
- Thiophanate (Nemafox)
Bali, M. K.; et al., 1978, Trop. Animal Health and Prod., v. 10 (1), 61-62
Haemonchus, Oesophagostomum, Trichostrongylus, and Trichuris spp. in pica-affected camels, thiophanate, good results against all but Trichuris sp.
- Nemafox
Bali, M. K.; and Singh, R. P., 1977, Haryana Agric. Univ. J. Research, v. 7 (4), 230-232
Haemonchus contortus, in vitro larvicidal effects of fenbendazole, banminth II, and nemafox
- Thiophanate (Nemafox)
Chandrasekharan, K.; et al., 1978, Kerala J. Vet. Sc., v. 9 (1), 167-170
gastrointestinal helminths in calves, efficacy of thiophanate, clinical trial
- Thiophanate
Dalton, S. E., 1978, Vet. Rec., v. 103 (7), 131-134
nematodes of sheep (nat. and exper.), thiophanate in low daily dosage reduced faecal egg output, egg hatchability, worm burdens, and pasture contamination, increased lamb weight gain, effective against Haemonchus contortus and Nematodirus spathiger, partially effective against Trichostrongylus colubriformis; comparison with phenothiazine treatment

Thiophanate -- Continued.

Thiophanate (Nemafax)

Duncan, J. L.; et al., 1979, Vet. Rec., v. 105 (19), 444-445
gastrointestinal nematodes including inhibited *Ostertagia ostertagi* larvae, calves, thiophanate

Thiophanate (Nemafax)

Fabiyi, J. P.; Oluyede, D. A.; and Negedu, J. O., 1979, Vet. Rec., v. 105 (16), 375
Cooperia punctata, *C. pectinata*, *Haemonchus placei*, cattle, thiophanate treatment: Jos Plateau, Vom, Nigeria

Thiophanate

Hall, C. A.; et al., 1978, Research Vet. Sc., v. 25 (3), 364-367
Haemonchus contortus, *Trichostrongylus colubriformis*, resistant strains selected with thiabendazole, dose response lines for 8 benzimidazole anthelmintics and thiophanate

Thiophanate (A 445, Nemafax)

Niec, R.; et al., 1979, Rev. Med. vet., Buenos Aires, v. 60 (1), 19-21
gastrointestinal nematodes, calves, thiophanate

Thiophanate

Ogunsusi, R. A., 1978, Research Vet. Sc., v. 25 (2), 251-252
Haemonchus spp. and *Trichostrongylus* spp., adult Yankasa sheep, thiophanate and thiabendazole at increased dosages highly effective in removing all stages of development including inhibited stages

Thiopropine

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Thiosemicarbazones

Foltinova, P.; et al., 1978, Acta Fac. Rerum Nat. Univ. Comenianae, Microbiol. (6), 35-50
Trypanosoma cruzi, trypanocidal effect of various thiosemicarbazones compared with standard anti-trypanosomes, benzazon VII proved effective in vitro (cultured crithidial forms) and in exper. infected mice and compared favorably with nitrofurazone and lampit

Thiosinamine -- Allyl thiourea.

Allyl thiourea

Ho, Y. H.; and Yang, H. C., 1974, Tung Wu Hsueh Pao (Acta Zool. Sinica), v. 20 (3), 243-262
Schistosoma japonicum, egg formation and chemical nature of egg shell, histological and histochemical study, morphological changes in egg formation following treatment of infected mice with thiourea compounds

Thiosinamine -- Continued.

Thiosinamine (Allyl thiourea)

Pellegrino, J.; and Machado, A., 1972, Rev. Brasil. Pesqui. Med. e Biol., v. 5 (1-2), 43-45
Schistosoma mansoni, mice and hamsters treated with thiosinamine, evidence of suppression of parasite egg laying process

Thiotepa

Kinnamon, K. E.; Steck, E. A.; and Rane, D. S., 1979, Antimicrob. Agents and Chemotherapy, v. 15 (2), 157-160
Trypanosoma rhodesiense, mice, inactive in screening of antitumor compounds for efficacy against infection

Thiourea

Ho, Y. H.; and Yang, H. C., 1974, Tung Wu Hsueh Pao (Acta Zool. Sinica), v. 20 (3), 243-262
Schistosoma japonicum, egg formation and chemical nature of egg shell, histological and histochemical study, morphological changes in egg formation following treatment of infected mice with thiourea compounds

2-(4-Thiozoly)-5-isopropylcarbonylaminobenzimidazole. See Cambendazole.

DL-Threo- α -2'-piperidyl-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridinemethanol. See α -(2-Piperidyl)-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridinemethanol.

DL-Threo- α -(2-piperidyl)-2-trifluoromethyl-6-(4-trifluoromethylphenyl)-4-pyridinemethanol phosphate. See α -(2-Piperidyl)-2-(4-trifluoromethylphenyl)-6-trifluoromethyl-4-pyridinemethanol.

Thymol

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

Thymol

Singh, Y., 1977, Indian Bee J., v. 36 (1-4), 1974, 16
Nosema apis, honey bees, formaldehyde (some improvement), thymol (good results): apiary at Jeolikote, India

Thymol iodide

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

Thymol palmitate

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

liberal. See Ornidazole.

Tifatol -- CGA 50439; 2-(2',4'-Dimethyl-phenyl-imino)-3-methyl-4-thiazolin.

CGA 50439 (Tifatol)
Gothe, R.; and Guizar, R. C., 1979, Zentralbl. Vet.-Med., Reihe B, v. 26 (4), 290-303
Argas walkerae, chickens (exper.), in vivo model for evaluating detaching potential of various acaricides

Tiguvon. See Fenthion.

Tiguvon Spot On. See Fenthion.

Tin -- Helmintha-P (with Phenothiazine, Piperazine, Senna, and Vernonia anthelmintica); Stannotaen; Stannous oxide; Tin arsenite.

Tin arsenite
Bogdanov, V. R.; et al., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 90-91
cestodes, pathomorphology resulting from action of various anthelmintics

Tin arsenite
Bogdanov, V. R.; Koshkina, N. G.; and Logachev, E. D., 1977, Sborn. Nauch. Rabot SibNIVI, Sibirsk. Nauchno-Issled. Vet. Inst. (29), 92-93
Dipylidium caninum, various anthelmintics, in vitro action on surface tissues and inactivation of enzymes

Stannotaen
Haak, W.; et al., 1972, Ang. Parasitol., v. 13 (4), 200-207
cestodes of humans, recommended drug therapy

Helmintha-P
Matta, S. C.; and Ahluwalia, S. S., 1979, Indian Vet. J., v. 56 (7), 616-617
helminths, poultry, helmintha-P, sonex

Tin arsenite. See Tin.

Tindurin. See Pyrimethamine.

Tinidazole -- Amoebicide 2004 (with Chlorphenoxamide); CP 12,574; Ethyl (2-(2-methyl-5-nitro-1-imidazolyl)ethyl) sulfone; 1-[2-(Ethylsulfonyl)ethyl]-2-methyl-5-nitroimidazole; Facizine; Fasigyn; Fasigyn 500; Fasigyne; Pletil; Simplotan; Tinigyn.

Tinidazole (Fasigyn)
Akashi, E.; et al., 1976, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 30 (7), 581-587
Trichomonas vaginitis, human, tinidazole, metronidazole, comparison clinical studies

Tinidazole -- Continued.

Tinidazole
Alonso, P., 1979, Chemotherapy, v. 25 (1), 40-47
tinidazole, free-living ciliates highly resistant, probably due to their aerobic condition, morphological and physiological alterations, comparison with parasitic protozoa

Tinidazole
Alonso, R. E., 1976, Semana Med. (4956), an. 83, v. 149 (7), 234-237
intestinal parasites, school children, clinical studies with tinidazole and mebendazole: Institute Aguirre 'Patronato de la Infancia'

Tinidazole
Alton, K. B.; and Patrick, J. E., 1979, J. Pharm. Sc., v. 68 (5), 599-601
tinidazole in human plasma, quantitative determination by high-performance liquid chromatographic assay

Tinidazole
Amato Neto, V.; et al., 1972, Rev. Soc. Brasil. Med. Trop., v. 6 (3), 147-149
Giardia lamblia, humans, therapeutic trials showed tinidazole to be useful antiparasitic treatment: Brazil

Tinidazole (Fasigyn)
Aure T., M., 1974, Rev. Obst. y Ginec. Venez., v. 34 (4), 437-440
human vaginal trichomoniasis, clinical trials comparing efficacy of tinidazole with metronidazole, concluded that drugs are of equal value in treatment: Venezuela

Tinidazole
Baranski, M. C.; et al., 1975, Rev. Soc. Brasil. Med. Trop., v. 9 (1), 15-18
Giardia lamblia, trials testing the therapeutic efficacy of tinidazole, adults and children both achieved cure rate of over 96%: Brasil

Tinidazole (Pletil)
Baranski, M. C.; and da Silva, A. F., 1976, Rev. Brasil. Clin. e Terap., v. 5 (11), 417-420
human intestinal amoebiasis, clinical trials testing the therapeutic effectiveness of tinidazole, recommended for individual and mass therapy because of ease of administration and virtual absence of side effects

Tinidazole (Fasigyn)
Blas Bava, A.; and Nino, R. F., 1974, Semana Med. (4876), an. 81, v. 145 (54), 2534-2539
Giardia lamblia, humans, clinical trials testing 2 dosage schedules of tinidazole, clinically and parasitologically effective with only mild side effects

Tinidazole
Brotherton, J., 1978, Arzneimittel-Forsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Tinidazole (Pletil)
Canzonieri, C. J.; et al., 1975, Rev. Brasil. Clin. e Terap., v. 4 (12), 479-482
Trichomonas vaginalis, Giardia lamblia, Entamoeba histolytica, humans, clinical trials testing efficacy of tinidazole therapy

Tinidazole -- Continued.

Tinidazole

Cervantes, L. F.; et al., 1975, Rev. Gastroenterol. Mexico, v. 40 (4), 185-193
human amoebiasis with hepatic abscess, clinical diagnosis confirmed by X-ray, laboratory studies and liver scan, therapy with tinidazole resulted in 93% cure rate: Mexico

Tinidazole (Fasigyne)

Crepin, G.; et al., 1976, Rev. Franc. Gyn. et Obst., v. 71 (11), 689-691
human urogenital trichomoniasis, treatment trials with tinidazole, good results

Tinidazole (Fasigyn)

Danzig, S.; and Hatchuel, W. L. F., 1977, South African Med. J., v. 52 (18), 708 [Letter]
Giardia lamblia, children, single dose tinidazole: Alexandra Health Clinic, Bergvlei, Tvl.

Fasigyne (CP 12,574)

Dellenbach, P.; and Muller, P., 1974, Rev. Franc. Gynec. et Obst., v. 70 (5), 357-359
human uro-genital trichomoniasis, therapeutic trials with fasigyne, recommended regimen includes treating both sexual partners, drug of choice

Tinidazole

Diez T., A.; et al., 1972, Rev. Obst. y Ginec. Venez., v. 32 (3), 433-438
human vaginal trichomoniasis, clinical trials comparing efficacy of tinidazole with that of metronidazole: Venezuela

Tinidazole (Fasigyn)

El-Masry, N. A.; Farid, Z.; and Miner, W. F., 1978, Am. J. Trop. Med. and Hyg., v. 27 (1, pt. 1), 201-202
Giardia lamblia, 53 of 55 heavily infected patients cured by single dose tinidazole: Egypt

Tinidazole

Fabio, P. F.; et al., 1978, J. Med. Chem., v. 21 (3), 273-276
Entamoeba histolytica, diaminoanthraquinone bisamidines, laboratory trials comparing activity against cecal form in rats and hepatic form in golden hamsters with activity of known amoebicides

Tinidazole (Fasigyn)

Farahmandian, I.; Sheiban, F.; and Sanati, A., 1978, J. Trop. Med. and Hyg., v. 81 (7), 139-140
Giardia lamblia, humans, evaluation of efficacy of single dose therapy with tinidazole, test findings of high cure rate with low side effects observed, recommended for both individual and mass therapy: Iran

Amoebicide 2004

Ferreira Tolsa, O., 1978, Semana Med. Mexico (1203), an. 25, v. 95 (4), 85-88
Entamoeba histolytica, school children, mass therapy with amoebicide 2004, well tolerated, good results

Tinidazole -- Continued.

Tinidazole

Forsgren, A.; and Forssman, L., 1979, Brit. J. Vener. Dis., v. 55 (5), 351-353
Trichomonas vaginalis, 35-year-old woman, symptomatic metronidazole-resistant vaginitis for 10 years, some resistance also to tinidazole and ornidazole: Sweden

Tinidazole

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Tinidazole

Hackett, L. P.; and Dusci, L. J., 1979, J. Chromatography, v. 175 (2), 347-349
metronidazole, tinidazole, detection and quantitation in human plasma using high-performance liquid chromatography

Tinidazole (Fasigyn)

Hatchuel, W., 1975, South African Med. J., v. 49 (45), 1879-1881
Entamoeba histolytica, patients with amoebic liver abscesses, tinidazole vs. metronidazole, equally efficacious

Fasigyne

Hervet, E.; and Aoun, A., 1978, Rev. Franc. Gynec. et Obst., v. 73 (6), 487-488
Trichomonas vaginalis, urogenital infection in pregnant women, treatment with fasigyne

Tinidazole

Hokanishi, H.; et al., 1976, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 30 (6), 515-518
human vaginal trichomoniasis, single dose therapy with tinidazole, clinical studies

Tinidazole

Islam, N.; and Hasan, M., 1978, J. Trop. Med. and Hyg., v. 81 (1), 20-22
Entamoeba histolytica, therapeutic efficacy of metronidazole and tinidazole compared in persons with hepatic infections, side effects

Tinidazole (Fasigyn)

Jokipii, A. M. M.; and Jokipii, L., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 758-761
Giardia lamblia, humans, tinidazole, comparative evaluation of 2 dosage regimens

Tinidazole (Fasigyn)

Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167
Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria

Fasigyne

Kaufmann, E., 1977, Rev. Franc. Gynec. et Obst., v. 72 (1), 33-35
Trichomonas vaginalis, round strain of parasite thought to be particularly virulent and resistant to various trichomonacides, therapy trials with fasigyne 500, varying treatment results

Tinidazole -- Continued.

Tinidazole (Facizine)

Kholodovskaia, I. V.; Minasova, G. S.; and Khokhlov, A. P., 1979, *Vestnik Dermat. i Venereol.* (4), 58-60
trichomoniasis, human vaginal infections, clinical trials with tinidazole

Tinidazole

Kunii, K.; et al., 1976, *Rinsho Fujinka Sanka* (Clin. Gynec. and Obst.), v. 30 (3), 237-240
human vaginal trichomoniasis, single dose therapy with tinidazole, clinical studies

Tinidazole

Lanbeck, K.; and Lindstrom, B., 1979, *J. Chromatography* (Biomed. Appl.), v. 162 (1), 117-121
assay of metronidazole and tinidazole in plasma and metronidazole in feces by high-performance liquid chromatography

Tinidazole

Levi, G. C.; de Avila, C. A.; and Amato Neto, V., 1977, *Rev. Inst. Med. Trop. S. Paulo*, v. 19 (6), 422-424
giardiasis, humans, efficacy of various drugs, comparative study, side-effects

Tinidazole

Lindmark, D. G.; and Mueller, M., 1976, *Antimicrob. Agents and Chemotherapy*, v. 10 (3), 476-482
metronidazole and 11 other nitroimidazoles, antitrichomonad activity against *Tritrichomonas foetus* and *Trichomonas vaginalis*, mutagenic action in Salmonella test, reducibility of nitro group by *T. foetus* homogenates, results underscore role of reduction of nitro group in antitrichomonad and mutagenic activity of nitroimidazoles

Tinidazole (Tinigyn)

Lovgren, T.; and Salmela, I., 1978, *Acta Path. et Microbiol. Scand.*, v. 86B (3), 155-158
Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents

Tinidazole

Manica, D.; and Peixoto, S., 1972, *Rev. Brasil. Clin. e Terap.*, v. 1 (5), 317-320
human trichomoniasis, clinical trials comparing efficacy of tinidazole and metronidazole

Tinidazole

Matsuda, S., 1978, *Rinsho Fujinka Sanka* (Clin. Gynec. and Obst.), v. 32 (2), 141-147
tinidazole, metronidazole, absorption, excretion, comparative study

Tinidazole

Meingassner, J. G.; and Thurner, J., 1979, *Antimicrob. Agents and Chemotherapy*, v. 15 (2), 254-257
Trichomonas vaginalis, isolation of strain resistant to metronidazole and other 5-nitroimidazoles

Tinidazole -- Continued.

Tinidazole (Fasigyn)

Merchan Lopez, D.; et al., 1977, *Rev. Obst. y Ginec. Venez.*, v. 37 (2), 267-273
human genital trichomoniasis, clinical trials testing efficacy of tinidazole, considered to be drug of choice for urogenital infections: Venezuela

Tinidazole (Fasigyn)

Merdivenci, A.; Baydemir, M.; and Sengul, M., 1976, *Turk Hijiy. ve Deneysel Biyol. Dergisi*, v. 36 (2), 238-247
Giardia intestinalis, children, tinidazole and nitrimidazine compared, both successful

Tinidazole (Fasigyn)

Nachbaur, J.; and Joly, H., 1978, *J. Chromatography* (Biomed. Appl.), v. 145 (2), 325-327
rapid assay of tinidazole in human plasma by high-performance liquid chromatography

Fasigyn (Tinidazole)

Nitzulescu, V.; Gherman, I.; and Popescu, A., 1977, *Rev. Pediat. (Pediat.)*, Bucuresti, v. 26 (2), 189-192
giardiasis, human, therapy with tinidazole, good results

Tinidazole

Orozco Hoyos, M.; Franco Marin, S.; and Soto, H., 1975, *Temas Escogidos Gastroenterol.*, v. 18, 253-262
amoebiasis, human acute rectocolitis, tinidazole

Tinidazole (Fasigyn)

Quaderi, M. A.; et al., 1978, *J. Trop. Med. and Hyg.*, v. 81 (1), 16-19
human hepatic amoebiasis, clinical diagnostic features, treatment trials with tinidazole, 100% cure rate at 2-month follow-up, some side effects: Bangladesh

Tinidazole (Fasigyn)

Rosemann, G. W.; and Vaughan, J., 1973, *South African Med. J.*, v. 47 (28), 1222-1224
Trichomonas vaginalis, human vaginal infections, tinidazole, single dose therapy, clinical trials

Tinidazole (Fasigyn)

Salles, A. de A.; et al., 1977, *Rev. Ginec. e Obst.*, Sao Paulo, v. 134 (3-4), 59-64
Trichomonas vaginalis, identification of "still" forms of parasites after therapy with single dose tinidazole, possible drug resistance in persons thought to be cured, need for treatment of sexual partners as additional precaution

Tinidazole

Sandront-Degee, M.; Werbrouck-Navette, J.; and Lambotte, R., 1975, *Rev. Med. Liege*, v. 30 (17), 560-562
Trichomonas vaginalis, humans, therapeutic trials comparing action of tiberl with metronidazole, fasigyn and tinidazole

Fasigyn

Sandront-Degee, M.; Werbrouck-Navette, J.; and Lambotte, R., 1975, *Rev. Med. Liege*, v. 30 (17), 560-562
Trichomonas vaginalis, humans, therapeutic trials comparing action of tiberl with metronidazole, fasigyn and tinidazole

Tinidazole -- Continued.

Tinidazole

Schenone, H.; et al., 1979, Bol. Chileno Parasitol., v. 34 (1-2), 2-6
Entamoeba histolytica and/or Giardia lamblia, children, clinical trials with varying doses of tinidazole: ciudad de Santiago

Tinidazole (Fasigyn)

Serup, J.; and Jensen, R. H., 1978, Ugeskr. Laeger, v. 140 (25), 1483-1484

Trichomonas vaginalis, women with vaginal infection and their sexual partners, controlled therapeutic trials, ornidazole vs. tinidazole as single oral dose

Tinidazole

Shimotomai, K.; et al., 1976, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 30 (9), 751-756

human vaginal trichomoniasis, oral use of tinidazole, clinical trials

Tinidazole

Shinohara, H.; and Sakai, H., 1975, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 29 (8), 635-638

human trichomoniasis vaginitis, combined oral and vaginal tinidazole

Tinidazole (Pletil)

Solano, J. N. de R.; Merino, C. G.; and Barrera, R. R., 1976, Rev. Brasil. Clin. e Terap., v. 5 (9), 353-365

human intestinal amoebiasis, successful clinical trials using tinidazole

Tinidazole (Fasigyn; Fasigyn 500)

Soyka, E., 1975, Therap. Umschau, v. 32 (9), 599-602

Trichomonas vaginalis, women, vaginal infections, single dose therapy with tinidazole compared with metronidazole: Switzerland

Tinidazole (Fasigyn)

Sucharit, P.; et al., 1979, Southeast Asian J. Trop. Med. and Pub. Health, v. 10 (4), 556-561

Trichomonas vaginalis, women with vaginal infections, single dose therapy; in vitro sensitivity studies with parasites cultured from human vaginal infections: Bangkok, Thailand

Tinidazole

Tazawa, T.; et al., 1974, Rinsho Fujinka Sanka (Clin. Gynec. and Obst.), v. 28 (9), 637-640

Trichomonas vaginalis vaginitis, combined oral and vaginal tinidazole therapy, patients' husbands also treated

Tinidazole

Valasco Chiriboga, A.; and Marquez Marino, G., 1975, Rev. Colomb. Obst. y Ginec., v. 26 (5), 291-298

human Trichomonas vaginalis vaginitis, comparative clinical trials using tinidazole and metronidazole; tinidazole recommended as drug of choice: Colombia

Tinidazole (Fasigyn)

do Vale, N., 1975, Rev. Brasil. Clin. e Terap., v. 4 (8), 305-308

human vaginal trichomoniasis, oral therapy with tinidazole, efficacy and simplicity of therapy makes it drug of choice for use in Public Health Clinics for control of venereal disease: Brazil

Tinidazole -- Continued.

Tinidazole

Valencia Torres, L., 1975, Temas Escogidos Gastroenterol., v. 18, 245-252
amoebiasis, human acute and chronic proctocolitis, clinical experiences with tinidazole

Tinidazole (Fasigyn)

Villamayor, J. H.; and Vivas, J., 1978, Semana Med. (5023), an. 85, v. 152 (6), 175-178
trichomoniasis, human, vulvo-vaginitis, clinical trials with tinidazole given as single dose

Tinidazole (Simplotan)

Walther, H., 1977, Therap. Gegenw., v. 116 (5), 969-975

trichomoniasis, human urogenital infections in sexual partners, simplotan, clinical trials

Tinidazole

Winkelmann, E.; Raether, W.; and Gebert, U., 1978, Arzneimittelforsch., v. 28 (10), 1682-1684

activity of 16 novel 5-nitroimidazoles against protozoa in mice and golden hamsters, compared with metronidazole and tinidazole, structure-activity relationships

Tinigyn. See Tinidazole.

Tioctilate (Octylthiobenzoate)

Pipyn, P.; Beaudet, C.; and Verstraete, W., 1979, Experientia, v. 35 (4), 480-481
tioctilate, biodegradability

Tioxidazole -- Methyl-6-n-propoxybenzothiazole-2-carbamate; Sch 21480.

Tioxidazole (Sch 21480)

Panitz, E.; et al., 1978, Experientia, v. 34 (6), 733

tioxidazole, new anthelmintic with broad spectrum activity against gastrointestinal roundworm infections

Tixantone. See Lucanthone. β -Toluenesulfonyl- β -methoxy-ethyl urethane sodium -- Codrinal (with Tetracycline).

Codrinal

Singh, J.; and Hussain, O., 1978, Indian Vet. J., v. 55 (1), 56-60

Eimeria tenella, chicks (exper.), amprolium provided better protection than codrinal, both drugs interfered to some extent with development of immunity

2-(p-Toluidino)-2-thiazoline

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and B. pahangi/patei hybrid, 23 anthelmintics tested in laboratory hosts (Aedes aegypti, Meriones unguiculatus, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

2-(o-Tolylydrazino)-2-thiazoline

Denham, D. A.; et al., 1978, Tr. Roy. Soc. Trop. Med. and Hyg., v. 72 (6), 615-618

Brugia pahangi and *B. pahangi/patei* hybrid, 23 anthelmintics tested in laboratory hosts (*Aedes aegypti*, *Meriones unguiculatus*, cats) and in vitro, concluded that insect and in vitro tests are of little value as primary screens

m-Tolyl methylcarbamate -- Tsumacide.

Tsumacide

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (1), 142-144

Boophilus microplus, five strains, susceptibility to acaricides: Jamaica; St. Kitts; Trinidad; Guyana

Toxaphene -- Altik (with Dioxathion); Camphechlor; Cooper-Tox; Coopertox [of Atef, M.; and ElSay, A., 1976]; Linton-X.

Toxaphene (Coopertox)

Atef, M.; and El-Say, A., 1976, J. Egypt. Vet. Med. Ass., v. 35 (3), 73-84

Hyalomma dromedarii, *Rhipicephalus s. sanguineus*, laboratory tests (immersion technique) with delnav, supona, toxaphene, BHC

Toxaphene (Camphechlor)

Crampton, P. L.; and Gichanga, M. M., 1979, Bull. Entom. Research, v. 69 (3), 427-439

economically important Ixodidae from major cattle-raising areas, survey of resistance to organochlorine and organophosphorus acaricides: Kenya

Toxaphene

Drummond, R. O.; et al., 1973, J. Econom. Entom., v. 66 (1), 130-133

Boophilus annulatus, *B. microplus*, laboratory tests of insecticides

Toxaphene

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886

laboratory-reared *Haematobia irritans*, susceptibility to topically applied insecticides

Toxaphene (Cooper-Tox)

Fisher, W. F.; et al., 1979, Southwest. Entom., v. 4 (3), 249-253

Psoroptes cuniculi, rabbits, phosmet, toxaphene, and 10 experimental alkyl amines compared

Toxaphene

Gladney, W. J.; Price, M. A.; and Graham, O. H., 1977, J. Med. Entom., v. 13 (4-5), 579-586

Amblyomma maculatum, cattle, efficacy of various insecticides applied as sprays, ear smears and dusts, or in slow-release devices, field tests

Camphechlor

Hamant, C. A., 1977, Rhodesian Vet. J., v. 8 (4), 67-70

ticks, resistance survey of field strains to commonly used ixodocides, changeover from arsenic to dioxathion dipping of cattle, dramatic improvement in tick control, necessitates complete overhaul of dipping facilities and retraining of personnel: Tribal Trust Lands of Rhodesia

Toxaphene -- Continued.

Toxaphene

Hamant, C. A.; and Matthewson, M. D., 1977, Rhodesian Vet. J., v. 8 (4), 71-73

Boophilus decoloratus, possible spread of organophosphate-resistant strain, cattle, case history, implications for control of ticks and tick-borne diseases: Rhodesia

Toxaphene + Dioxathion (= Altik)

Keating, M. I., 1978, Bull. Animal Health and Prod. Africa, v. 26 (4), 285-292

excretion of toxaphene and dioxathion in milk of dairy cows

Toxaphene

Lourens, J. H. M., 1979, J. Econom. Entom., v. 72 (5), 790-793

Amblyomma variegatum, *A. lepidum*, baseline data on susceptibility to organochlorine acaricides, genetic basis for resistance in *A. variegatum*

Toxaphene

Lourens, J. H. M.; and van de Klashorst, G., 1979, Ztschr. Ang. Entom., v. 87 (3), 230-238

Rhipicephalus appendiculatus, organochlorine susceptible and tolerant populations: East Africa

Toxaphene

Lourens, J. H. M.; and Tatchell, R. J., 1979, Bull. Entom. Research, v. 69 (2), 235-242

Rhipicephalus evertsi evertsi, identification and inheritance of resistance factors to organochlorine acaricides, experimental hybrids between susceptible and resistant strains

Toxaphene (Linton-X)

Meleney, W. P.; and Roberts, I. H., 1979, J. Med. Entom., v. 16 (1), 52-58

Psoroptes ovis, cattle, acaricides, dipping, spraying, or spray-dipping trials

Toxaphene

Rechav, Y.; and Whitehead, G. B., 1978, J. Econom. Entom., v. 71 (1), 149-151

Amblyomma hebraeum, field trials with pheromone-toxaphene mixtures applied to demarcated areas on cattle for tick control: near East London, South Africa

Toxaphene

Wright, F. C.; and Riner, J. C., 1979, Southwest. Entom., v. 4 (1), 40-45

Psoroptes ovis, *P. cuniculi*, 10 acaricides evaluated using 'tea-bag' technique

Toyomycin. See Chromomycin A₃.

Tramisol. See Tetramisole.

L-Tramisol. See Tetramisole.

Tremerad. See Clioxanide.

Tresaderm. See Dexamethasone or Neomycin or Thiabendazole.

Triamcinolone -- Aristocort; Triamcinolone diacetate.

Triamcinolone diacetate (Aristocort)
Nozik, R. A., 1977, Tr. Am. Acad. Ophth. and Otolaryngol., v. 83 (5), 811-818
toxoplasmic retinochoroiditis, humans, periorbital corticosteroid injections administered alone or in conjunction with systemic antimicrobial agents

Triamcinolone acetonide (Kenalog)
Nozik, R. A., 1977, Tr. Am. Acad. Ophth. and Otolaryngol., v. 83 (5), 811-818
toxoplasmic retinochoroiditis, humans, periorbital corticosteroid injections administered alone or in conjunction with systemic antimicrobial agents

Triamcinolone diacetate. See Triamcinolone.

DL-1,2,4,-Triazole-3-alanine
Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744
trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Tribrisen. See Sulfadiazine or Trimethoprim.

2,4,6-Tribromoimidazole
Brotherton, J., 1978, Arzneimittelforsch., v. 28 (10), 1665-1672
trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

3,5,4'-Tribromosalicylanilide. See Bromsalans.

3,5,5'-Tribromosalicyl-o-toluidide
Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Tribromsalan. See Bromsalans.

Tribromsalicylanide. See Bromsalans.

Tribulus pterocarpus herb
Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, Scientia Pharm., v. 47 (2), 114-118
Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

Tricandil. See Mepartricin.

Trichlorfon -- Arpalit; Bayer 2349; Bayer 15 922; Bilarcil; Bubulin; Chlorophos; ComBot; Combot; 0,0-Dimethyl-(1-hydroxy-2,2,2-trichloroethyl)-phosphate; 0,0-Dimethyl 2,2,2-trichloro-1-hydroxyethyl phosphate; Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate; 0,0-Dimethyl-2,2,2-trichloro-1-hydroxyethyl-phosphorous; Dipterex; Dixol; DTHP; Foschlor; Hypocid; Hypodermacide; Hypodermin; Hypodermin-chlorophos; L. 13/59; Metrifonate; Metrifonatum; Metriphosphate; Neguvon; Noevar; Ricifon; Trichlorofon; 2,2,2-Trichlorohydroxyethyl phosphoric acid; Trichlorophone; Trichlorphon; Trichlorphone.

Metrifonate
Abaru, D. E.; and McMahon, J. E., 1978, Tropenmed. u. Parasitol., v. 29 (2), 175-177
Onchocerca volvulus, Wuchereria bancrofti, humans, metrifonate demonstrated partial microfilaricidal activity against former but neither macro- nor microfilaricidal activity against latter: Tanga region, Tanzania

Trichlorfon (Dipterex, Neguvon)
Abdel Rahman, M. S.; El Gendi, A. Y. I.; and Moursi, H. S. A., 1977, J. Egypt. Vet. Med. Ass., v. 37 (2), 55-69
nematodes and cestodes, dogs, effectiveness of trichlorfon, oxinithiophos, and carbaryl compared: Giza and Cairo Governorates, Egypt

Neguvon
Alcaino, H. A.; and Calderon, M. T., 1976, Arch. Med. Vet., Valdivia, v. 8 (1), 40-41
Cheyletiella yasguri, dogs, neguvon

Metrifonate (Bilarcil)
Arap Siongok, T. K.; et al., 1978, J. Infect. Dis., v. 138 (6), 856-858
Schistosoma haematobium, children, mass treatment with single oral dose of metrifonate, effect on egg output: Coast Province, Kenya

Chlorophos
Azimov, Sh. A.; Enileeva, N. Kh.; and Ishmirzaev, B. R., 1977, Veterinariia, Moskva (7), 56-57
Gastrophilus spp. in horses, chlorophos granules in feed for control: Uzbekistan

Trichlorfon + Pyrantel pamoate
Bentley, O. E.; et al., 1978, Vet. Med. and Small Animal Clin., v. 73 (1), 70-73
horses, pyrantel pamoate + trichlorfon, field trials, results show that drugs are safe for horses when preceded by a feeding of grain or a complete horse ration, mild side effects: Kentucky; Alabama; Texas; Kansas

Metrifonate
Beulig, W.; et al., 1978, Monatsh. Vet.-Med., v. 33 (5), 181-184
metrifonate, sheep, toxicity

Trichlorfon -- Continued.

Trichlorphon (Metrifonate; Chlorophos)
Beulig, W.; Das, M.; and Bjamba, B., 1977,
Monatsh. Vet.-Med., v. 32 (23), 914-916
Dermacentor nuttalli, sheep, trichlorphon,
spray and pour-on methods evaluated, results
show that pour-on most effective and eco-
nomical; not more than 4 consecutive appli-
cations permissible, more applications cause
decline in cholinesterase activity

Trichlorfon

Bonin, W., 1977, Berl. u. Munchen. Tierarztl.
Wchnschr., v. 90 (2), 34-37
Amblyomma hebraeum, Psoroptes cuniculi,
Melophagus ovinus, Dermanyssus gallinae,
heptenophos, rapid mode of action, broad
range of efficacy, short residual effect
and effective as a vapour poison, compared
with other standard drug preparations

Trichlorfon (Noevar)

Boulard, C. F., 1979, Vet. Parasitol., v. 5
(4), 379-387
Hypoderma-infected or uninfected calves,
treatment with fenthion or trichlorfon,
blood histamine levels, circulating antibody
titers to Hypoderma lineatum antigen in
infected calves; blood histamine levels in
guinea pigs after injection of ground-up
Hypoderma lineatum larvae or application of
fenthion

Metrifonate (Neguvon)

Brandrup, F.; Andersen, K. E.; and Kristensen,
S., 1979, Ugeskr. Laeger, v. 141 (15), 1015-
1017
Cheyletiella yasguri infection in dogs and
their owners, removal of dogs from homes and
therapy with metrifonate baths exterminated
mite populations: Denmark

Metrifonate

Burchard, G. D.; Albiez, E. J.; and Bierther,
M., 1979, Tropenmed. u. Parasitol., v. 30 (1),
97-102
onchocerciasis, humans, electron microscopic
studies of skin and of microfilariae after
treatment with metrifonate: Liberia

Dipterex

Chaia, G.; et al., 1975, Rev. Inst. Med. Trop.
S. Paulo, v. 17 (5), 298-306
Dermatobia hominis, life cycle maintained
under laboratory conditions, infection of
rats for study of chemotherapeutics

Neguvon

da Costa, U. C.; and Saraiva, D., 1977, Rev.
Centro Cien. Rurais, v. 7 (4), 417-419
Myocoptes musculus, white mice [in Eng-
lish title; "camundongos brancos (Wistar)"
in Portuguese text], treatment with asuntol
and neguvon

Bubulin

Daniyarov, I. A.; et al., 1978, Veterinariia,
Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics
and dyes tested, none effective

Trichlorfon -- Continued.

Chlorophos

Daniyarov, I. A.; et al., 1978, Veterinariia,
Moskva (2), 64-65
Echinococcus spp., sheep, 28 anthelmintics
and dyes tested, none effective

Neguvon

Delak, M., 1977, Vet. Arhiv, Zagreb, v. 47
(5), 231-238
Gasterophilus larvae, horses, percutaneous
application of trichlorphon and neguvon

Trichlorphon

Delak, M., 1977, Vet. Arhiv, Zagreb, v. 47
(5), 231-238
Gasterophilus larvae, horses, percutaneous
application of trichlorphon and neguvon

Neguvon

Derylo, A., 1978, Polskie Pismo Entom., v. 48
(2), 253-259
Hypoderma bovis, cattle, incidence prior to
and during control with neguvon, seasonal
dynamics

Neguvon

Derylo, A.; and Haliniarz, A., 1977, Med. Wet.,
v. 33 (10), 624-626
Hypoderma bovis, cattle, neguvon, economic
losses before and after treatment: Lublin,
Poland

Trichlorphon (Neguvon)

Dorrestein, G. M.; and Van Bronswijk, J. E. M.
H., 1979, Vet. Parasitol., v. 5 (4), 389-398
Trixacarus caviae as cause of mange in Cavia
porcellus (nat. and exper.), clinical symp-
toms, pathology, treatment; papular urti-
caria in humans associated with mangy guinea-
pigs: The Netherlands

Trichlorfon (ComBot paste)

Drudge, J. H.; Lyons, E. T.; and Tolliver,
S. C., 1978, Am. J. Vet. Research, v. 39 (9),
1419-1421
nematodes, bots, horses, febantel, activity
of paste formulation alone or with tri-
chlorfon paste, critical tests

Trichlorfon (Comboto liquid) + Febantel

Drudge, J. H.; Lyons, E. T.; and Tolliver,
S. C., 1979, J. Equine Med. and Surg., v. 3
(3), 135-140
parasites, horses, critical tests with
febantel alone or in combination with tri-
chlorfon

Trichlorfon

Drummond, R. O.; et al., 1973, J. Econom. En-
tom., v. 66 (1), 130-133
Boophilus annulatus, B. microplus, laboratory
tests of insecticides

Chlorophos

Evstaf'ev, M. N., 1978, Veterinariia, Moskva
(11), 70-72
Hypoderma bovis, cattle, insecticides tes-
ted, aerosol method of application more use-
ful for large, specialized farms than for
individual treatment: Tiimensk oblast

Trichlorfon -- Continued.

Hypodermin-chlorophos

Evstaf'ev, M. N., 1978, Veterinariia, Moskva (11), 70-72

Hypoderma bovis, cattle, insecticides tested, aerosol method of application more useful for large, specialized farms than for individual treatment: Tiumensk oblast

Neguvon

Evstaf'ev, M. N., 1978, Veterinariia, Moskva (11), 70-72

Hypoderma bovis, cattle, insecticides tested, aerosol method of application more useful for large, specialized farms than for individual treatment: Tiumensk oblast

Neguvon

Fadzil, M., 1977, Vet.-Med. Nachr. (1), 44-52
Stephanofilaria kaeli, cattle, incidence in different ecological areas, role of Staphylococcus aureus and S. albus in setting up the inflammation; neguvon: malaysischen Halbinsel

Neguvon

Fadzil, M., 1977, Vet. Med. Rev. (1), 44-52
Stephanofilaria kaeli in cattle, prevalence, temperature and humidity in endemic areas favor vectors, Staphylococcus play important role in setting up inflammation, treatment with neguvon gave excellent results: west coast of Peninsular Malaysia

Trichlorfon

Frazar, E. D.; and Schmidt, C. D., 1979, J. Econom. Entom., v. 72 (6), 884-886
laboratory-reared Haematobia irritans, susceptibility to topically applied insecticides

Metrifonate

Fuglsang, H.; and Anderson, J., 1977, Tropenmed. u. Parasitol., v. 28 (4), 439-446

Onchocerca volvulus, human, single dose of metrifonate, microfilaricidal effects, side effects (better tolerated than diethylcarbamazine): Cameroon

Metrifonate

Fuglsang, H.; and Anderson, J., 1978, Tropenmed. u. Parasitol., v. 29 (2), 168-174

Onchocerca volvulus, humans, treatment trials with metrifonate using 3 different regimens: Cameroon

Chlorophos

Getta, G. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 72-75

hypodermatosis, cattle, insecticides, repellents

Hypodermin-chlorophos

Getta, G. I., 1977, Sborn. Nauch. Rabot SibNIVI (28), 72-75

hypodermatosis, cattle, insecticides, repellents

Chlorophos

Goncharov, A. P.; and Kudriavikh, V. I., 1978, Veterinariia, Moskva (8), 30-31

Haematopinus suis, pig farm, chlorophos in single dose for prophylaxis, two doses for treatment

Trichlorfon -- Continued.

Metrifonate

Gonzalez Barranco, D., 1977, Rev. Invest. Salud Pub., Mexico, v. 37 (1), 37-45

Onchocerca volvulus, humans, brief review of in vitro and in vivo effects of several drugs studied for possible therapeutic use, dosage recommendations for some drugs, special emphasis on studies with metrifonate

Bubulin

Gudimenko, I. I., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 100-104

trichuriasis, oesophagostomiasis, ascariasis, swine, testing thiabendazole, dithiazanine iodide, dipterex, hygromycin-B, and bubulin

Dipterex

Gudimenko, I. I., 1970, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 8, 100-104

trichuriasis, oesophagostomiasis, ascariasis, swine, testing thiabendazole, dithiazanine iodide, dipterex, hygromycin-B, and bubulin

Dipterex

Gudimenko, I. I., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 112-116

trichuriasis, ascariasis, oesophagostomiasis, swine, dipterex tested

Dipterex

Gudimenko, I. I., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 137-140

[Oesophagostomum], [Trichocephalus], [Ascariids], pregnant and lactating sows, dipterex treatment

Neguvon

Guler, S., 1977, Vet. Fak. Dergisi, Ankara Univ., v. 24 (3-4), 441-449

Demodex canis, dogs, neguvon as bath, good therapeutic agent

Chlorophos

Iakubovskii, M. V.; and Zen'kov, A. V., 1977, Vet. Nauka--Proizvod., Trudy, Minsk, v. 15, 72-78

nematodes, swine raising complexes, combined control measures, sanitation, anthelmintics, suiverm most effective

Ricifon (Chlorophos)

Iamov, V. Z., 1977, Veterinariia, Moskva (9), 64-67

[Hypoderma], cattle, ricifon effective, no toxic effects; toxicity tested in white mice; tissues of calves and cow's milk tested for residues

Chlorophos

Iamov, V. Z.; and Kolesnik, N. V., 1978, Veterinariia, Moskva (11), 68-70

Hypoderma bovis larval stages, esterases, molecular forms, effect of inhibitors, including chlorophos

Trichlorfon -- Continued.

Trichlorfon

- Imada, R.; and Muroga, K., 1979, Bull. Japan. Soc. Scient. Fish. (Nippon Suisan Gakkaishi), v. 45 (1), 25-29
Pseudodactylogyrus microrchis on Anguilla anguilla (gills), trichlorfon, formalin, and sodium chloride baths compared, 2 trichlorfon baths effective and harmless

Metriphonate

- James, M. F. M.; and Jewsbury, J. M., 1978, Brit. Med. J. (6110), v. 1, 442 [Letter]
human urinary schistosomiasis, depression of blood cholinesterase activity during therapy with metriphonate, cautions regarding use of suxamethonium during course of schistosomal therapy

Trichlorophone (Metriphonate, Bilarcil)

- Kale, O., 1978, Tropenmed. u. Parasitol., v. 29 (2), 163-167
Onchocerca volvulus, humans, small-scale trials of 6 known parasiticides, none showed any evidence of substantial activity against microfilariae or adult worms: Western Nigeria

Neguvon

- Kamyszek, F., 1977, Med. Wet., v. 33 (6), 354-356
Sarcoptes scabiei, pigs, alugan, neguvon

Dipterex

- Kashnikov, A. A., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 140-142
macracanthorhynchosis, swine, action of various anthelmintics

Trichlorfon

- Kawatsu, H., 1978, Bull. Japan. Soc. Scient. Fish. (Nippon Suisan Gakkaishi), v. 44 (12), 1315-1319
Diplozoon nipponicum in crucian carp (gills), hypochromic microcytic anemia, hematological characteristics, incidence in relation to season and host size, effective treatment with trichlorfon (DEP): basin of the river Asakawa

Trichlorfon (Neguvon)

- Keller, H.; and Mueller, R., 1979, Berl. u. Munchen. Tierarztl. Wchnschr., v. 92 (4), 63-65
dichlorvos, trichlorfon, horses, decrease in plasma cholinesterase activity, concluded that relaxation of horses with succinylcholine should not be carried out within 10 days after exposure to organophosphorus type anthelmintics

Trichlorfon

- Kolacz, J., 1977, Med. Wet., v. 33 (3), 157-159
Hypoderma sp., cattle, trichlorfon: Poland

Dixol (Trichlorophone)

- Koltai, L., 1979, Magy. Allat. Lapja, v. 34 (2), 87-91
Hypoderma, cattle, trichlorophone 'pour on', economic losses and cost of control: Hungary

Trichlorfon -- Continued.

Neguvon

- Kristensen, S.; Brandrup, F.; and Andersen, K. E., 1978, Dansk Vet.-Tidsskr., v. 61 (8), 369-375
Cheyletiella yasguri, dogs and humans exposed to dogs, case histories, neguvon successful in dogs

Chlorophos

- Kruchinina, M. P., 1977, Sborn. Nauch. Rabot SibNIVI (28), 129-132
hypodermatosis, cattle, chlorophos in feed, large scale trials

Dipterex

- Kumar, A.; et al., 1977, Indian J. Poultry Sc., v. 12 (3), 48-50
Eomenacanthus stramineus, chicks (exper.), sumithion, malathion, dipterex, comparative efficacy, haematology and blood glucose levels

Chlorophos

- Mamaev, N. Kh.; Golin, P. I.; and Omarov, M. V., 1978, Veterinariia, Moskva (11), 73-74
Hypoderma, B[oophilus] calcaratus, cattle, chlorophos treatment, prophylaxis against piroplasmosis: Dagestan

Dixol

- Merenyi, L., 1978, Magy. Allat. Lapja, v. 100, v. 33 (2), 132-134
Hypoderma, control, treatment with Dixol

Chlorophos

- Meshkov, S.; Iotov, I.; and Sirachev, D., 1977, Vet. Sbirka, v. 75 (5), 32-33
piroplasmosis, berenil, cattle, control of Boophilus calcaratus with chlorophos

Hypodermin-chlorophos

- Metelitsa, V. K.; et al., 1977, Veterinariia, Moskva (9), 68-70
[Hypoderma], cattle, dosing apparatus for applying hypodermin-chlorophos in measured doses to several animals, economic benefits of treatment

Neguvon

- Mihai, M.; and Badescu, C., 1973, Lucrar. Stiint. Inst. Agron. N. Balcescu, s. C. Med. Vet., v. 15, 1972, 225-229
neguvon, alterations of free amino acids in animal tissues

Trichlorfon (Neguvon)

- Miller, B. E.; et al., 1978, J. Med. Entom., v. 14 (6), 651-661
flea control on rodents and rabbits, evaluation of 7 organophosphates as oral systemics, open-field and enclosure tests: southeastern New Mexico

Foschlor

- Moczon, T., 1976, Bull. Acad. Polon. Sc., Cl. II, s. Sc. Biol., v. 24 (5), 289-292
Fasciola hepatica miracidia, inhibitory effect of pesticides on enzyme activity

Trichlorfon -- Continued.

Chlorophos

Mordasov, P. M.; and Kisliakova, Z. I., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 162-165
[Hypoderma], calves, early treatment with hypodermin + chlorophos or chlorophos, weight gain

Hypodermin

Mordasov, P. M.; and Kisliakova, Z. I., 1973, Vet. Nauka--Proizvod., Trudy, Minsk, v. 11, 162-165
[Hypoderma], calves, early treatment with hypodermin + chlorophos or chlorophos, weight gain

Chlorophos

Mordasov, P. M.; Kisliakova, Z. I.; and Bukin, V. A., 1971, Nauch. Trudy, Nauchno-Issled. Vet. Inst., v. 9, 130-134
[Hypoderma], cattle, early treatment with chlorophos poured on, high therapeutic effect

Trichlorfon (Comboto)

Morrow, G. L., 1978, Vet. Med. and Small Animal Clin., v. 73 (11), 1388, 1393
strongyles, ascarids, bots, horses, febantel and trichlorfon paste formulations, drug trials

Trichlorfon + Mebendazole

Muyllle, E.; Oyaert, W.; and Rogiers, M., 1979, Vlaams Diergeneesk. Tijdschr., v. 48 (4), 279-282
Gasterophilus intestinalis larvae, horses, mebendazole + trichlorfon paste, endoscopic assessment of efficacy

Metrifonate (Dipterex)

Niemandt, S.; and Murahwa, S., 1975, South African Med. J., v. 49 (45), 1860 [Letter]
Schistosoma haematobium, schoolchildren, trials with metrifonate, annual treatment may reduce prevalence in endemic areas: Rhodesia

Neguvon + Asuntol

Oba, M. S. P.; Ogassawara, S.; and Costa, A. J. S., 1977, Arq. Inst. Biol., Sao Paulo, v. 44 (1-2), 95-97
Chorioptes bovis var. bovis, bovines, clinical aspects, cure with neguvon + asuntol: Municipio de Sao Carlos, Sao Paulo

Neguvon (Metrifonatum)

Olsson, T., 1977, Svensk Vet.-Tidn., v. 29 (20), 795-800
Ascaris suum, slaughter swine, treatment with neguvon or mascyl

Metrifonate (Trichlorfon; Bilarcil; Neguvon)

Omer, A. H. S.; and Teesdale, C. H., 1978, Ann. Trop. Med. and Parasitol., v. 72 (2), 145-150
Schistosoma mansoni, S. haematobium, humans, single or mixed infections including some patients passing S. mansoni eggs in urine, efficacy of metrifonate, results suggest that site of infection rather than species of parasite renders parasite more susceptible to metrifonate and may further clarify mode of action of metrifonate: Khartoum, Sudan

Trichlorfon -- Continued.

Metrifonate (Trichlorphon)

Ong, T. M., 1978, Mutation Research, v. 55 (1), 43-70
hycanthone and other antischistosomal drugs, general properties, teratogenicity, carcinogenicity, mutagenicity, and other genetically related activities, review

Neguvon

Oproiu, V.; et al., 1977, Rev. Crest. Animal-elor, v. 27 (7), 47-49
Hypoderma bovis, cattle, neguvon, ruelene, trichlorphon, curative treatment in spring, prophylactic treatment in autumn more efficient

Trichlorphon

Oproiu, V.; et al., 1977, Rev. Crest. Animal-elor, v. 27 (7), 47-49
Hypoderma bovis, cattle, neguvon, ruelene, trichlorphon, curative treatment in spring, prophylactic treatment in autumn more efficient

Trichlorphone

Pellegrino, J.; Pereira, L. H.; and Mello, R. T., 1977, Rev. Inst. Med. Trop. S. Paulo, v. 19 (1), 43-46
Schistosoma mansoni, mice, chemoprophylactic activity of 17 known schistosomicidal agents compared

Neguvon

Petrov, D.; Milushev, I.; and Monov, M., 1978, Vet. Med. Nauki, v. 15 (4), 33-39
Oestrus ovis, sheep, neguvon, ranide, and dovenix compared; use of ranide economically justified only for mixed infection with Fasciola hepatica

Chlorophos (Hypodermacide)

Potemkin, V. I.; and Nadiradze, O. Z., 1977, Veterinariia, Moskva (9), 67-68
[Hypoderma], cattle, chlorophos (Hypodermacide), effective, low toxicity

Chlorophos

Puchkova, E. A., 1977, Veterinariia, Moskva (7), 19-22
D[ermanyssus] gallinae, C[imex] lectularius, lice, chickens on industrial scale farms, control, sevin, dicresil, chlorophos, carbophos; other complex sanitation measures

Metrifonate (Bilarcil)

Pugh, R. N. H., 1978, Ann. Trop. Med. and Parasitol., v. 72 (5), 495-497
Schistosoma haematobium, human, concurrent single-dose therapy with metrifonate and niri-dazole: Malumfashi District, Nigeria

Neguvon

Rak, H., 1976, J. Vet. Fac. Univ. Tehran, v. 32 (1-4), 77-96
ectoparasites, veterinary practice, berco-tox, asuntol 50, alon, bolfo, alugan, opigal, gamatox, tetmosol, neguvon: Iran

Trichlorfon -- Continued.

Chlorophos

Rastegaev, Iu. M., 1978, Vestnik Sel'skokhoz. Nauki Kazakhstana (3), 73-76
myiasis, helminths, horses, chlorophos, amidophos, economic effectiveness of treatment: northern Kazakhstan

Trichlorphon

Rawlins, S. C.; and Mansingh, A., 1978, J. Econom. Entom., v. 71 (6), 956-960
Boophilus microplus, 6 Jamaican strains, patterns of resistance to acaricides

Metrifonate

Reiner, E.; et al., 1978, Comp. Biochem. and Physiol., v. 60C (2), 155-157
Metastrongylus apri, cholinesterase, kinetic properties with respect to substrate hydrolysis and inhibition by organophosphorus compounds

Trichlorphon (Metrifonate)

Ribbeck, R.; Schroeder, E.; and Schumann, H., 1979, Monatsh. Vet.-Med., v. 34 (10), 383-384
Lucilia sericata, dogs, cats, myiasis, surgical removal and drug treatment

Hypocid

Riha, J.; Minar, J.; and Hradil, M., 1976, Veterinarstvi, v. 26 (10), 456-458
Hypoderma bovis, H. lineatum, cattle, treatment with hypocid, economic importance of control, effect on heifer weight gain

Hypocid (Trichlorfon)

Riha, J.; Minar, J.; and Matouskova, O., 1977, Vet. Med., Praha, v. 50, v. 22 (4), 193-200
Hypoderma bovis, heifers, hypocid preparation, highly effective and profitable, as measured by weight gains

Metrifonate (Bayer 2349)

Salazar-Mallen, M.; Gonzalez-Barranco, D.; and del Carmen Montes H., M., 1971, Rev. Inst. Med. Trop. S. Paulo, v. 13 (5), 363-368
Onchocerca volvulus, human, trials with metrifonate attempting to establish maximum microfilaricidal activity with minimum side effects

Dipterex

Sanchez Moreno, M.; Monteoliva, M.; and Hermoso, R., 1978, Rev. Iber. Parasitol., v. 38 (1-2), 415-426
Ascaris lumbricoides, in vitro, anthelmintics and pesticides, effects on motility

Metrifonate

Shibuya, T.; et al., 1978, Japan. J. Exper. Med., v. 48 (5), 411-418
Litomosoides carinii in Sigmodon hispidus, screening filaricides for human filariasis, evaluation of intrathoracic injection method

Trichlorfon (Comboto)

Shmidl, J. A.; et al., 1979, Vet. Med. and Small Animal Clin., v. 74 (12), 1796-1799
safety evaluation of concurrent administration of febantel and trichlorfon in paste and liquid forms to horses, no significant toxic effects

Trichlorfon -- Continued.

Chlorophos

Shoshkov, D.; and Kolarova, V., 1977, Vet. Med. Nauki, v. 14 (8), 91-98
Argulus foliaceus, Dactylogyrus extensus, carp, neguvon, chlorophos, good results

Neguvon

Shoshkov, D.; and Kolarova, V., 1977, Vet. Med. Nauki, v. 14 (8), 91-98
Argulus foliaceus, Dactylogyrus extensus, carp, neguvon, chlorophos, good results

Chlorophos

Slin'ko, V. G., 1979, Veterinariia, Moskva (5), 44-46
Sarcoptes suis, pigs, association with necrosis of ear helix, possible secondary bacterial infection; treatment with chlorophos or TAP

Trichlorfon

Smith D. L., 1976, Manitoba Entom., v. 10, 5-8
Hypoderma spp., calves, weight gains, no significant difference between treated and untreated calves in response to control of cattle grubs with insecticides (trichlorfon and crufomate): Manitoba

Chlorophos

Sverba, V. A.; and Shemchuk, V. R., 1978, Veterinariia, Moskva (10), 69-71
Sinergasilus major, white amur, copper sulfate and ferrous sulfate mixture, chlorophos, carbophos, formula for estimating concentrations in relation to temperature and other factors in aquaria or ponds

Chlorophos

Talanov, G. A.; and Nikolaev, P. I., 1978, Veterinariia, Moskva (9), 59-60
Parascaris, [Strongylata], Gasterophilus, horses, use of chlorophos and amidophos: Yakutsk ASSR

Metrifonate (Trichlorofon, Dipterex)

Tanaka, H.; et al., 1977, Japan. J. Exper. Med., v. 47 (4), 315-317
Litomosoides carinii-infected cotton rats, improved method for intrapleural injection of anti-filarial drugs to evaluate macrofilaricidal action

Chlorophos

Ternovoi, V. I., 1978, Veterinariia, Moskva (9), 62-63
Oestrus ovis, sheep, chlorophos in intranasal irrigation

Arpalit (Trichlorofon; DTHP)

Tesik, I., 1978, Biol. a Chem. Zivoc. Vyroby, Vet., v. 14 (v. 20) (4), 379-384
ectoparasites of reptiles, arpalit spray, toxicity to snakes and tortoise tested, TMB-4 as antidote

Neguvon

Ueno, H.; and Chibana, T., 1978, Japan Agric. Research Quart., v. 12 (3), 152-156
Stephanofilaria okinawaensis, cattle, distribution, clinical signs, chemotherapy, intermediate host determined

Trichlorfon -- Continued.

Metrifonate

Woolhouse, N. M., 1979, *Biochem. Pharmacol.*, v. 28 (16), 2413-2418
antischistosomal drugs, biochemical and pharmacological effects in relation to mode of action

Trichlorfon

Wright, F. C.; and Riner, J. C., 1979, *Southwest. Entom.*, v. 4 (1), 40-45
Psoroptes ovis, *P. cuniculi*, 10 acaricides evaluated using 'tea-bag' technique

Neguvon

Yagi, A. I.; Abdel Razig, M. T.; and Osman, O. M., 1975, *Bull. Animal Health and Prod. Africa*, v. 23 (2), 181-184
Menacanthus cornutus, poultry, neguvon solution for spraying and dipping

Neguvon

Youssef, A. H., 1976, *J. Egypt. Vet. Med. Ass.*, v. 35 (3), 147-157
Dipetalonema evansi, camels, filarial orchitis and possible significance as prevalent reproductive disease; surgical treatment and use of neosulversan, fouadin, and neguvon, histopathology of gonads: Egypt

Neguvon

Zarzara, C., 1979, *Rev. Crest. Animalelor*, v. 29 (5), 42-45
Cimex lectularius, lindatox-20, neguvon, and carbetox 37, laboratory tests; carbetox 37 for control in poultry sheds, application methods

Chlorophos

Zharikov, I. S.; Litviak, V. S.; and Nikolaenko, G. V., 1973, *Vet. Nauka--Proizvod.*, Trudy, Minsk, v. 11, 119-121
chlorophos, cattle given anthelmintic dose, blood picture, intoxication

1,1,1-Trichloro-2,2-bis(p-methoxyphenyl) ethane.
See Methoxychlor.

3,4,4'-Trichlorocarbanilide

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Trichlorofon. See Trichlorfon.

2,4,4'-Trichloro-2'-hydroxydiphenyl ether

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

2,2,2-Trichlorohydroxyethyl phosphoric acid.
See Trichlorfon.

N-Trichloromethyl-thio-4-cyclohexene-1,2-dicarboximide. See Captan.

Trichlorophen -- Trichlosal (with Phenasal).

Trichlorophen

Bankov, D., 1976, *Vet. Med. Nauki*, v. 13 (10), 28-36
cestodes of sheep, drug trials; *Stilesia globipunctata*, tested several diagnostic methods with unfavorable results

Trichlosal (=Phenasal + Trichlorophene)

Grinenko, N. V.; et al., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (1), 101-103
H[ymenolepis] nana, patients of different age groups, dichlosal or trichlosal in divided doses, 5 or 7 day courses

Trichlorophene

Grinenko, N. V.; et al., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasal, trichlorophene, dichlosal, and trichlosal tested in graded doses

Trichlosal

Grinenko, N. V.; et al., 1976, *Med. Parazitol. i Parazitar. Bolezni*, v. 45 (2), 176-178
H[ymenolepis] nana, white mice, phenasal, trichlorophene, dichlosal, and trichlosal tested in graded doses

Trichlorophene

Lopatin, B. V.; and Bekhli, A. F., 1979, *Khimiko-Farm. Zhurnal*, v. 13 (9), 103-105
trichlorophene, photometric method of quality control in production

Trichlorophen

Rusak, L. V.; and Kovchur, V. N., 1972, *Parazitologiya*, Leningrad, v. 6 (1), 85-87
Hymenolepis nana, glycogen content in parasite tissues decreased after treatment of infected mice with aminoacrichine, phenasal, or trichlorophen, implications for mechanism of drug action

Trichlorophone. See Trichlorfon.

3,5,4'-Trichlorosalicylanilide

Sakamoto, T., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 115-128
Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

3,5,4'-Trichlorosalicylanilide

Sakamoto, T.; and Gemmell, M. A., 1975, *Japan. J. Vet. Research*, v. 23 (3), 81-94
Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Trichlorophon. See Trichlorfon.

Trichlorophone. See Trichlorfon.

Trichlosal. See Niclosamide or Trichlorophen.

Trichomycin

Lovgren, T.; and Salmela, I., 1978, Acta Path. et Microbiol. Scand., v. 86B (3), 155-158
Trichomonas vaginalis, in vitro sensitivity to 7 chemotherapeutic agents

Trichopol. See Metronidazole.

Tricocel. See Oxantel.

Trifluoperazine

Hillman, G. R.; Gibler, A. M.; and Anderson, J. W., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 992-997

Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

**3-Trifluoromethyl-4,4'-dichlorocarbani-
lilide**

Sakamoto, T., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 115-128

Echinococcus multilocularis, various anthelmintics, scolicidal effects in vitro and/or therapeutic effects in mice

**3-Trifluoromethyl-4,4'-dichlorocarbani-
lilide**

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolicidal effect of salicylanilide and bisphenol derivatives

Trigonella foenum-greekum seeds

Abdulla, W. A.; Kadry, H.; and Mahran, S. G., 1979, Scientia Pharm., v. 47 (2), 114-118

Ascaridia galli, Ascaris vitulorum, in vitro anthelmintic activity of some Egyptian plants; only Nerium oleander caused death of worms

Trimagill. See Trimonil.

Trimelarsen. See Melarsonyl potassium.

Trimethoprim -- Ciplin (with Sulfamethoxazole);

Co-trimoxazole (with Sulfamethoxazole); 2,4-Diamino-5-(3,4,5-trimethoxybenzyl)-pyrimidine; Eusaprim (with Sulfamethoxazole); Pacprim (with Sulfadimethoxine, Sulfisomidine, and Sulfisoxazole); Septran (with Sulfamethoxazole); Septrin (with Sulfamethoxazole); Sulfanilamide complex (with Sulfadimethoxine, Sulfisomidine, and Sulfisoxazole); Tribissen (with Sulfadiazine); WR 5,949.

Trimethoprim

Brotherton, J., 1978, Arzneimittelforsch., v. 28 (10), 1665-1672

trichomonads, in vitro testing of potential trichomonacides using Coulter Counter

Trimethoprim + Sulfadiazine (Tribissen)

Brunnthaler, F., 1977, Prakt. Tierarzt, v. 58 (11), 849-851

coccidiosis, dogs, treatment with various drugs, best results with amprolium

Trimethoprim -- Continued.**Trimethoprim-Sulfamethoxazole**

Deeg, H. J.; et al., 1979, Transplantation, v. 28 (3), 243-246

effect of trimethoprim-sulfamethoxazole on hematological recovery after total body irradiation and autologous marrow transplantation studied in dogs, results show that drug can be given safely and probably prevents very early cases of Pneumocystis carinii pneumonia

Co-trimoxazole

Geddes, A. M.; Ball, A. P.; and Farrell, I. D., 1979, J. Antimicrob. Chemother., v. 5, suppl. B, 221-230

co-trimoxazole for treatment of serious infections, review including information on Pneumocystis carinii, malaria, and toxoplasmosis

Trimethoprim

Grossman, P. L.; and Remington, J. S., 1979, Am. J. Trop. Med. and Hyg., v. 28 (3), 445-455

Toxoplasma gondii, in vitro and in vivo in mice, effects of trimethoprim and sulfamethoxazole alone and in combination

Trimethoprim

Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338

Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice

Trimethoprim-Sulfamethoxazole

Hughes, W. T., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (3), 333-335

Pneumocystis carinii, immunosuppressed rats, trimethoprim-sulfamethoxazole has limited rather than lethal effect, protection is afforded only during period of administration

Trimethoprim + Sulfamethoxazole

Hughes, W. T.; et al., 1978, J. Pediat., St. Louis, v. 92 (2), 285-291

Pneumocystis carinii, children, comparison of pentamidine isethionate and trimethoprim combined with sulfamethoxazole (TMP-SMZ) in treating Pneumocystis pneumonia, results show that TMP-SMZ is as effective as pentamidine, has minimal side effects, offers oral administration and is more readily available

Trimethoprim

Kluge, R. M.; Spaulding, D. M.; and Spain, A. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 975-978

Pneumocystis carinii in steroid-conditioned rats, combination of pentamidine with trimethoprim-sulfamethoxazole, data suggest that combination therapy is no more effective than trimethoprim-sulfamethoxazole alone and may be, in fact, harmful, trimethoprim by itself has no place in treatment of pneumocystosis

Trimethoprim + Sulphamethoxazol (= Septrin)

Kouba, K.; Nevarilova, A.; and Rajlichova, J., 1978, Ceskoslov. Epidemiol., Mikrobiol., Immunol., v. 27 (3), 175-178

toxoplasmosis, human, therapy with septrin, poor results with allergic side effects

Trimethoprim -- Continued.

Trimethoprim + Sulphafurazole
Kyllerman, M.; and Strannegard, O., 1979,
Arch. Dis. Childhood, v. 54 (4), 326-327
[Letter]
toxoplasmosis, 5-year-old boy, hemiplegia,
disease most likely acquired from cat which
had high serum levels of toxoplasma anti-
bodies, trimethoprim-sulphafurazole treatment

Trimethoprim-Sulfamethoxazole
Larter, W. E.; et al., 1978, J. Pediat., St.
Louis, v. 92 (5), 826-828
Pneumocystis carinii, trimethoprim-sulfa-
methoxazole treatment of pneumonitis in
children

Trimethoprim + Sulfisomidine + Sulfadimethoxine +
Sulfisoxazole (= Pacprim; Sulfanilamide complex)
Lee, H. S.; Lee, H. B.; and Moon, M. H., 1979,
Korean J. Animal Sci., v. 21 (4), 281-288
Besnoittia besnoiti, goats (exper.), antimony
potassium tartrate, pacprim

Trimethoprim
Lichtenwalner, D. M.; et al., 1979, Antimicrob.
Agents and Chemotherapy, v. 16 (5), 579-583
trimethoprim, sulfamethoxazole, rapid assay
for determination of levels in serum by
spectrofluorometry

Trimethoprim-Sulphadiazine
McKenzie, R. A.; Green, P. E.; and Wood, A. D.,
1978, Austral. Vet. J., v. 54 (2), 86-88
Angiostrongylus cantonensis in captive
Macropus rufogriseus (surface of brain be-
neath leptomeninges, cerebellar folium,
meninges), clinical symptoms, pathology,
treatment with trimethoprim-sulphadiazine
ineffective, case report: Brisbane

Trimethoprim + Sulfamethoxazole (= Septrin)
Mancinella, J. E.; et al., 1975, Semana Med.
(4914), an. 82, v. 147 (9), 230-235
Trichomonas, human vaginal infections, thera-
peutic action of sulfamethoxazole combined
with trimethoprim analyzed

Trimethoprim
Neal, R. A., 1972, Rev. Inst. Med. Trop. S.
Paulo, v. 14 (6), 341-351
Leishmania tropica, 7 Latin-American and 2
Asiatic isolates, course of infection in
hamsters, anti-folic reductase drugs compared
with paromomycin and sodium stibogluconate

Trimethoprim + Sulfadiazine
Neal, R. A., 1972, Rev. Inst. Med. Trop. S.
Paulo, v. 14 (6), 341-351
Leishmania tropica, 7 Latin-American and 2
Asiatic isolates, course of infection in
hamsters, anti-folic reductase drugs compared
with paromomycin and sodium stibogluconate

Trimethoprim + Sulphamethoxazole
Norrby, R.; et al., 1975, Scand. J. Infect.
Dis., v. 7 (1), 72-75
Toxoplasma gondii, humans, clinical and
serological data on patients treated with
trimethoprim-sulphamethoxazole

Trimethoprim -- Continued.

Trimethoprim + Sulfamethoxazole (= Co-trimoxa-
zole; Eusaprim)
Norrby, R.; and Eilard, T., 1976, Scand. J.
Infect. Dis., v. 8 (4), 275-276
toxoplasmosis, recurrent infection in woman
treated with co-trimoxazole, normal clinical
response to each course of therapy, no evi-
dence of impaired immunity

Trimethoprim (WR 5,949)
Rane, D. S.; and Kinnamon, K. E., 1979, Am. J.
Trop. Med. and Hyg., v. 28 (6), 937-947
sporozoite-induced Plasmodium berghei in
mice, development of high volume tissue
schizonticidal drug screen based upon mor-
tality of infected mice

Trimethoprim + Sulphamethoxazole (= Septran;
= Ciplin)
Shashindran, C. H.; et al., 1978, Brit. J. Der-
mat., v. 98 (6), 699-700
human pediculosis capitis, successful system-
ic (oral) therapy using combination of trime-
thoprim and sulphamethoxazole without addi-
tional external application of insecticides;
drugs when used separately were not effective

Trimethoprim-Sulfamethoxazole
Stevenson, D. K.; Christie, D. L.; and Haas, J.
E., 1978, Pediatrics, Am. Acad. Pediat., v. 61
(6), 864-866
trimethoprim-sulfamethoxazole, child, hepatic
injury

Trimethoprim + Sulfomethoxazole (= Septrin)
Szaflarski, J.; Sokola, A.; and Herman, Z. S.,
1974, Acta Parasitol. Polon., v. 22 (22-34),
261-263
Toxoplasma gondii, mice (exper.), trimetho-
prim and sulfomethoxazole alone and in combi-
nation, concluded that trimethoprim has no
therapeutic effect but potentiates action of
sulfomethoxazole

Trimethoprim + Sulfamethoxazole
Thiermann, E.; et al., 1978, Am. J. Trop.
Med. and Hyg., v. 27 (4), 747-750
Toxoplasma gondii, mice, 5 treatment regimens
compared during acute and late infections;
pyrimethamine + sulfamethoxyypyridazine was
most effective

Trimethoprim + Sulfadoxin
Waller, T., 1979, Lab. Animals, v. 13 (3), 227-
230
Encephalitozoon cuniculi, survival of spores
after exposure to various temperatures and
disinfectants; growth-inhibition effect of
drugs in cell cultures

Trimethoprim + Sulfamethoxazole (= Co-trimoxazole)
Westerman, E. L.; and Christensen, R. P., 1979,
Ann. Int. Med., v. 91 (3), 413-414
Isospora belli, immunosuppressed woman con-
currently infected with Giardia lamblia,
severe diarrhea, rapid remission with co-
trimoxazole, case report

Trimethoprim -- Continued.

Trimethoprim + Sulfamethoxazole

Wolff, L. J.; and Baehner, R. L., 1978, Am. J. Dis. Child., v. 132 (5), 525-526

Pneumocystis carinii, clinical trials evaluating prophylactic value of a 2-week, high-dose course of trimethoprim-sulfamethoxazole to prevent pneumonia in children with cancer who are receiving their first 100 days of intensive immunosuppressive chemotherapy

Trimethoprim+Sulfamethoxazole

Yoshida, Y.; et al., 1977, Kiseichugaku Zasshi (Japan. J. Parasitol.), v. 26 (6), 367-375

Pneumocystis carinii pneumonia, rats, comparative efficacy of pyrimethamine+sulfamonomethoxine vs. trimethoprim+sulfamethoxazole

Trimonil (Trimagill)

Karnaky, K. J., 1972, Southwest. Med., v. 53 (1), 10-12

Trichomonas vaginalis and other vaginal infections, women, micronized form of trimonil instilled vaginally, restores normal pH, thus is useful prophylactically and therapeutically

Trinoin. See Bromsalans.

6,6,9-Tri nor-6-oxo Δ^{6a-10a} -tetrahydrocannabinol Pringle, H. L.; Bradley, S. G.; and Harris, L. S., 1979, Antimicrob. Agents and Chemotherapy, v. 16 (5), 674-679

Naegleria fowleri, susceptibility to Δ^9 -tetrahydrocannabinol and other cannabinoids

Triostam. See Antimony sodium gluconate.Triostib. See Antimony sodium gluconate.

Tripiperazine tetrabithionol

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Tris (p-aminophenyl) carbonium. See Pararosaniline.Tris (p-aminophenyl) carbonium pamoate. See Pararosaniline. β, β, β -Tris-(4-chlorophenyl)-propionic acid-N'-methylpiperazide. See Hetolin.

2,2'-Trithiobis(4-chloro-6-nitrophenol)

Sakamoto, T.; and Gemmell, M. A., 1975, Japan. J. Vet. Research, v. 23 (3), 81-94

Echinococcus granulosus in vitro, scolical effect of salicylanilide and bisphenol derivatives

Trodax. See Nitroxynil.Trolene. See Ronnel.Trypaflavine. See Acriflavine.Trypanidium. See Isometamidium.

Trypan blue

Irvin, A. D.; and Young, E. R., 1978, Research Vet. Sc., v. 25 (2), 211-214

Babesia spp., drug inhibition of hypoxanthine uptake in vitro could be used as primary screen for babesicidal drugs but drugs showing in vitro activity are not necessarily active in vivo

Trypanblue

Nikol'skii, S. N.; Nikiforenko, V. I.; and Pozov, S. A., 1977, Veterinariia, Moskva (4), 71-75

Piroplasma jakimovi, cattle, morphological and biological comparison with *P. bigeminum*, epizootiology (*Ixodes ricinus* as main vector; frequent association with leptospirosis), treatment: Siberia

Trypan blue

Vulchovski, Ia., 1977, Vet. Med. Nauki, v. 14 (3), 79-87

Babesia, cattle, efficacy of trypan blue, trypaflavin, acaprin, berenil, and pyrodis (berenil and pyrodis superior to other drugs)

Trypanocides

Lausi, L., 1976, Semana Med. (4952), an. 83, v. 149 (3), 84-100; (4960), an. 83, v. 149 (11), 356-365

Chagas disease, humans, extensive therapeutic review, current concepts

Trypanocides

Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579

Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Trypanocides

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Tryparsamide -- Sodium N-phenylglycinamide-p-arsenodithioglucolate.

Tryparsamide

Gill, B. S., 1972, Ann. Soc. Belge Med. Trop., v. 52 (1), 33-44

Trypanosoma evansi, rats, chemoprophylactic trials, 8 compounds tested

Tryparsamide

Meshnick, S. R.; et al., 1978, J. Exper. Med., v. 148 (2), 569-579

Trypanosoma brucei brucei, attempt to develop new trypanocidal drugs based on inability of bloodstream form to decompose hydrogen peroxide, experiments with porphyrins, naphthoquinones, and arsenicals in vitro and in vivo, possible mechanisms of combination of agents

Tsumacide. See m-Tolyl methylcarbamate.

Tubercidin -- 4-Aminopyrazolo-(2,3-d)-pyrimidine beta-D-ribofuranoside; 7-Deazaadenosine.

Tubercidin

Dewey, V. C.; Kidder, G. W.; and Nolan, L. L., 1978, Biochem. Pharmacol., v. 27 (10), 1479-1485

Crithidia fasciculata, mechanism of inhibition of growth by adenosine and adenosine analogs

Tubercidin

Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130

Echinococcus granulosus, scolicalidal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Tubercidin -- Continued.

Tubercidin

Senft, A. W.; and Crabtree, G. W., 1977, Biochem. Pharmacol., v. 26 (20), 1847-1856

Schistosoma mansoni, inhibition of adenine and guanine nucleotide synthesis by purine analogs in intact worms in vitro, implications in development of new anti-schistosomal drugs

Tubercidin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

d-Tubocurarine

Hillman, G. R.; Gibler, A. M.; and Anderson, J. W., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 992-997

Schistosoma mansoni, anticholinergic drugs as inhibitors of labeling of parasite by a fluorescent derivative of acetylcholine, scanning microfluorimetric system

Turingin. See Bacillus thuringiensis.

Tyrothricin

Williamson, J.; and Scott-Finnigan, T. J., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (5), 735-744

trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

- U. K. 5066
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5378
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5574
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5704
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5876
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5925
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5585-11
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- U. K. 5778/11
Pellegrino, J.; Mello, R. T.; and Pereira, L. H., 1976, Rev. Inst. Med. Trop. S. Paulo, v. 18 (3), 149-151
Schistosoma mansoni, mice, 8 pyrazinoquinolines
- Umbethion. See Coralox.
- 10-Undecen-1-yl-isothiuronium. See Undecenyl-pseudothiourea.

Undecenyl-pseudothiourea -- 10-Undecen-1-yl-isothiuronium.

Undecenyl-pseudothiourea
Payares, G.; and Ercoli, N., 1978, Exper. Parasitol., v. 45 (1), 1-7
Schistosoma mansoni, drug-immobilized cercariae have reduced virulence but are not dead, cercariae become avirulent only when flame cell is affected, no protection against reinfection in mice injected with immobilized cercariae of reduced virulence

Uden. See Propoxur

Urea
Venkateswara Rao, P.; et al., 1977, Riv. Parasitol., Roma, v. 38 (1), 13-21
Prosthogonimus sp. cercariae, cercaricidal effect of certain common fertilizers, ammonium sulphate may be cercaricide of choice

Urea stibamine
Naik, S. R.; et al., 1979, Tr. Roy. Soc. Trop. Med. and Hyg., v. 73 (1), 61-65
Leishmania donovani, human, 24 sporadic cases of kala-azar, epidemiology, clinical features, complications and associations, hematological and biochemical findings, diagnosis, treatment: north-western India

Uredofos. See Diuredosan.

Uredofos, Disodium salt. See Diuredosan.

Uroporphyrin I
Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, Biochem. Pharmacol., v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism

Ursovermit. See Rafoxanide.

Vagilen. See Metronidazole.

Valbazen. See Albendazole.

Vankin. See Pyrvinium.

Vansil. See Oxamniquine.

Vapam. See Metham sodium.

Vapona. See Dichlorvos.

Vapona Strips. See Dichlorvos.

Vaporett dog collars. See Dichlorvos.

Vendex. See Hexakis (β , β -dimethylphenethyl) distannoxane.

Vermex. See Piperazine.

Vermiculine

Fuskova, A.; Fuska, J.; and Kettner, M., 1978, *Folia Microbiol.*, v. 23 (5), 389-393
Trichomonas foetus, in vitro inhibition by vermiculine, mode of action

Vermox. See Mebendazole.

Vernonia anthelmintica -- Helmintha-P (with Phenothiazine, Piperazine, Senna, and Tin).

Helmintha-P

Matta, S. C.; and Ahluwalia, S. S., 1979, *Indian Vet. J.*, v. 56 (7), 616-617
 helminths, poultry, helmintha-P, sonex

Vidarabine -- Adenine arabinoside.

Adenine arabinoside

Pfefferkorn, E. R.; and Pfefferkorn, L. C., 1978, *J. Parasitol.*, v. 64 (3), 486-492
Toxoplasma gondii, lack of adenosine kinase is biochemical basis for resistance to adenine arabinoside in mutant

Vinblastine

Williamson, J.; and Scott-Finnigan, T. J., 1978, *Antimicrob. Agents and Chemotherapy*, v. 13 (5), 735-744
 trypanocidal activity of antitumor antibiotics and other metabolic inhibitors, microtest for rapid preliminary assay in vitro, parasite motility and infectivity for mice are indexes respectively of respiration and glycolysis and of cell division, implications of results for combination chemotherapy and deposit prophylaxis (with polyanions)

Vinblastine sulfate

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Vincristine sulfate

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Vinylphosphate -- 0,0-Dimethyl-0,1,2,4,5-trichlorphenyl (2-chlorvinyl) phosphate.

Vinylphosphate

Frolov, B. A.; et al., 1978, *Veterinariia, Moskva* (11), 75-76
Dermanyssus gallinae, *Argas persicus*, chickens, acaricides tested

N-Vinyl phthalimide

Sakamoto, T.; and Gemmell, M. A., 1979, *Mem. Fac. Agric. Kagoshima Univ.* (24), v. 15, 125-130
Echinococcus granulosus, scolical effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro

Virazole. See Ribavirin.

VUAgT-71

Privora, M.; Rupes, V.; and Cerny, V., 1970, *Folia Parasitol.*, v. 17 (1), 81-84
Dermapteron marginatus, laboratory trials testing six insecticides

Warbex. See Famphur.

Wopell

Srivastava, G. C.; and Kumar, J., 1979, Indian Vet. Med. J., v. 3 (2), 85-87
Raillietina tetragona, poultry, wopell ex-pels segments but not scolices

Wormguard. See Parabendazole.

Wormwood. See Absinthium.

WR 5 990

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR 6 007

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR 6 025

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR 6 028

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR-7,930

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta

WR 29 597

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR-77,135

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-91,808

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-98,057

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-98,057

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

WR-99,210

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR-99,682

Schmidt, L. H., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 718-737
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, methods employed in search for new blood schizonticidal drugs

WR 127 854

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

WR-131,834

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

WR-135,403

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

WR 136 479

Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

- WR-143,803
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-146,459
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-148,763
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-148,946
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-149,809
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-150,726
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-151,312
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-158,121
Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines
- WR-162,878
Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines
- WR-165,355
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-165,533
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-173,060
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in Aotus trivirgatus griseimembra and human volunteers, P. cynomolgi in Macaca mulatta, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds
- WR-175,039
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and P. vivax in Aotus trivirgatus griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for P. berghei and these same compounds

WR-175,412

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in *Aotus trivirgatus* griseimembra and human volunteers, *P. cynomolgi* in *Macaca mulatta*, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-177,602

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus* griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

WR-178,919

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus* griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-178,979

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in *Aotus trivirgatus* griseimembra and human volunteers, *P. cynomolgi* in *Macaca mulatta*, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-180,117

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus* griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-180,153

Schmidt, L. H., 1979, Am. J. Trop. Med. and Hyg., v. 28 (5), 793-807
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus* griseimembra, strains resistant to chloroquine, quinine, or pyrimethamine, antimalarial properties of selected 2,4-diamino-6-substituted quinazolines

WR-181,018

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in *Aotus trivirgatus* griseimembra and human volunteers, *P. cynomolgi* in *Macaca mulatta*, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-182,231

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 420-435
Plasmodium falciparum and *P. vivax* in *Aotus trivirgatus* griseimembra, antimalarial activities of various 4-pyridinemethanols with special attention to WR-172,435 and WR-180,409, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-183,544

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus* griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

WR-183,545

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus* griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

WR-183,546

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus* griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

WR-183,606

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in *Aotus trivirgatus* griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with *P. vivax* in *Aotus trivirgatus* and *P. cynomolgi* in *Macaca mulatta*

WR-185,020

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in *Aotus trivirgatus* griseimembra and human volunteers, *P. cynomolgi* in *Macaca mulatta*, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

WR-190,420

Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 14 (3), 292-314
Plasmodium falciparum in *Aotus trivirgatus* griseimembra and human volunteers, *P. cynomolgi* in *Macaca mulatta*, antimalarial activities of various 9-phenanthrenemethanols with special attention to WR-122,455 and WR-171,669, includes some [apparently unpublished] results of other workers for *P. berghei* and these same compounds

- WR 192 515
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR-199,426
Schmidt, L. H.; et al., 1978, Antimicrob. Agents and Chemotherapy, v. 13 (6), 1011-1030
Plasmodium falciparum in Aotus trivirgatus griseimembra, pilot appraisals of activities of 12 4-quinolinemethanols, further appraisal of mefloquine with P. vivax in Aotus trivirgatus and P. cynomolgi in Macaca mulatta
- WR 201 678
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 206 027
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 208 442
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 210 810
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 077
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 663
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 666
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 672
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 784
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 211 819
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 212 293
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 213640
Gutteridge, W. E.; Cover, B.; and Gaborak, M., 1978, Ann. Trop. Med. and Parasitol., v. 72 (4), 329-338
Trypanosoma cruzi, rapid, simple primary screen to test compounds for activity as potential trypanocides using infected A/JAX inbred mice
- WR 214 198
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 215 300
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 215 732
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 216 837
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 217 271
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)
- WR 218 574
Kinnamon, K. E.; et al., 1978, Am. J. Trop. Med. and Hyg., v. 27 (4), 751-757
Leishmania donovani-Mesocricetus auratus model, antileishmanial activity of lepidines (6-methoxy-4-methyl-8-aminoquinoline derivatives)

- Xexy p-hydroxy benzoate
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Yoghurt
Gunston, K. D.; and Fairbrother, P. F., 1975, South African Med. J., v. 49 (16), 675-676
Trichomonas vaginalis, human, unsuccessful trials using yoghurt to reduce vaginal infestations
- Yomesan. See Niclosamide.
- Zanil. See Oxyclozanide.
- Zectran. See 4-Dimethylamino-3,5-xylol methyl-carbamate.
- Zinc and Sodium chlorides, saturated solution
Kunstyr, I.; and Ammerpohl, E., 1978, Lab. Animals, v. 12 (2), 95-97
Spironucleus muris, faecal cysts, resistance to physical and chemical factors tested, data may be useful for control of infection in rodents and for cryopreservation of parasite
- Zinc-conjugated manganese ethylene bisdithiocarbamate
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Zinc ethylene bisdithiocarbamate
Sakamoto, T.; and Gemmell, M. A., 1979, Mem. Fac. Agric. Kagoshima Univ. (24), v. 15, 125-130
Echinococcus granulosus, scolicedal effect of 65 antibiotic, antineoplastic, cytostatic, and other agents in vitro
- Zinc hematoporphyrin
Meshnick, S. R.; Chang, K. P.; and Cerami, A., 1977, Biochem. Pharmacol., v. 26 (20), 1923-1928
Trypanosoma brucei, T. congolense, heme lysis of bloodstream forms, T. brucei, lytic effect of porphyrins, in vitro and in vivo (mice) studies, mechanism of action believed to be homolytic cleavage of intracellular H₂O₂ to form hydroxyl radicals which can react with vital cell components and kill the organism
- Zn hematoporphyrin D
Meshnick, S. R.; et al., 1978, J. Pharmacol. and Exper. Therap., v. 207 (3), 1041-1050
Trypanosoma brucei brucei, mice, rats, rabbits, evaluation of trypanocidal activity of series of porphyrins and metalloporphyrins, role of zinc in porphyrin-induced lysis
- Zoalen. See Dinitolmide.
- Zoalene. See Dinitolmide.
- Zoamix. See Dinitolmide.
- Zoecon 515. See Methoprene.
- Zolone. See Phosalone.



**United States Department of Agriculture
Agricultural Research Service
Washington, D.C. 20250**

**OFFICIAL BUSINESS
Penalty for Private Use, \$300**

**Postage and Fees Paid
U.S. Department of Agriculture
AGR-101**

